

1 **Transient climate simulations of the deglaciation 21-9**
2 **thousand years before present; PMIP4 Core experiment**
3 **design and boundary conditions.**

4 **Ruza F. Ivanovic¹, Lauren J. Gregoire¹, Masa Kageyama², Didier M. Roche^{2,3}, Paul**
5 **J. Valdes⁴, Andrea Burke⁵, Rosemarie Drummond⁶, W. Richard Peltier⁶, Lev**
6 **Tarasov⁷**

7 [1]{School of Earth & Environment, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT,
8 United Kingdom}

9 [2]{[Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-](#)
10 [UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France](#)}

11 [3]{Earth and Climate Cluster, Faculty of Earth and Life Sciences, Vrije Universiteit
12 Amsterdam, Amsterdam, the Netherlands}

13 [4]{School of Geographical Sciences, University of Bristol, University Road, Bristol, BS8 1SS,
14 United Kingdom}

15 [5]{Department of Earth and Environmental Sciences, Irvine Building, University of St.
16 Andrews, St. Andrews, KY16 9AL, United Kingdom}

17 [6]{Department of Physics, University of Toronto, 60 St George Street, Toronto, Ontario,
18 Canada M5S 1A7}

19 [7]{Department of Physics and Physical Oceanography, Memorial University of
20 Newfoundland and Labrador, St. John's, NL, Canada A1B 3X7}

21 Correspondence to: Ruza F. Ivanovic (r.ivanovic@leeds.ac.uk)

22 **Abstract**

23 The last deglaciation, which marked the transition between the last glacial and present
24 interglacial periods, was punctuated by a series of rapid (centennial and decadal) climate
25 changes. Numerical climate models are useful for investigating mechanisms that underpin the
26 [climate change](#) events, especially now that some of the complex models can be run for multiple

1 millennia. We have set up a Paleoclimate Modelling Intercomparison Project (PMIP) working
2 group to coordinate efforts to run transient simulations of the last deglaciation, and to facilitate
3 the dissemination of expertise between modellers and those engaged with reconstructing the
4 climate of the last 21 thousand years. Here, we present the design of a coordinated Core
5 [simulation experiment](#) over the period 21-9 thousand years before present (ka) with time varying
6 orbital forcing, greenhouse gases, ice sheets, and other geographical changes. A choice of two
7 ice sheet reconstructions is given, [but no and we make recommendations for prescribing](#) ice
8 [sheet or iceberg](#) meltwater [should be prescribed \(or not\)](#) in the Core [simulation experiment](#).
9 Additional *focussed* simulations will also be coordinated on an ad-hoc basis by the working
10 group, for example to investigate [more thoroughly](#) the effect of ice [sheet and iceberg](#) meltwater-
11 [and on climate system evolution, and to examine](#) the uncertainty in other forcings. Some of
12 these *focussed* simulations will [focus on target](#) shorter durations around specific events [to in](#)
13 [order to understand them in more detail and](#) allow the more computationally expensive models
14 to take part.

15 1 Introduction

16 1.1 Climate evolution over the last deglaciation

17 The last deglaciation is a period of major climate change, when Earth transitioned from its last
18 full glacial state, to the current interglacial climate. The *Last Glacial Maximum* (LGM) marked
19 the culmination of the last glacial cycle when vast ice sheets covered large regions of the
20 Northern Hemisphere, stretching over North America and Eurasia (e.g. Boulton et al., 2001;
21 Dyke et al., 2002; Peltier et al., 2015; Svendsen et al., 2004; Tarasov et al., 2012), and the
22 Antarctic Ice Sheet expanded to the edge of the continental shelf (Argus et al., 2014; Briggs et
23 al., 2014; Lambeck et al., 2014 and references therein). Changes in the ice sheets resulted in a
24 total sea level rise of ~115-130 m between LGM and the late Holocene (Lambeck et al., 2014;
25 Peltier and Fairbanks, 2006) depending upon the time assumed to correspond to the LGM, and
26 ~100 m from 21 ka to 9 ka (the period of focus for this manuscript).

27 Historically, [the EPILOG group](#) defined the LGM as having occurred 23-19 ka (21 ka centre
28 point), when climate was generally cool and ice sheets were more or less at their largest, based
29 on ice core and sea level records (Mix et al., 2001). It represents the time of maximum terrestrial

1 ice volume. More recently, the last sea level lowstand has been found to have occurred either
2 around 26 ka (Peltier and Fairbanks, 2006) or 21 ka (Lambeck et al., 2014) with relatively stable
3 (low) sea level between those dates. Nearly all ice sheets were at or close to their maximum
4 extent between 26 ka and 19 ka (Clark et al., 2009).

5 During the LGM, global annual mean surface temperatures are estimated to have been around
6 4.0 ± 0.8 °C colder than today (Annand and Hargreaves, 2013). The Earth began warming
7 towards its present state from around 19 ka (Fig. 1h; Buizert et al., 2014; Jouzel et al., 2007),
8 as summer insolation at northern high latitudes and global atmospheric greenhouse gas
9 concentrations gradually increased (Fig. 1c-f; Bereiter et al., 2015; Berger, 1978; Loulergue et
10 al., 2008; Marcott et al., 2014). By 9 ka, although the northern ice sheets had not quite retreated
11 (or disappeared) to their present day configuration, most of the Northern Hemisphere
12 deglaciation had taken place (Clark et al., 2012; Lambeck et al., 2014; Peltier et al., 2015;
13 Tarasov et al., 2012; Figures 1g and 2), with both surface air temperatures (Fig. 1h-i) and
14 atmospheric greenhouse gases (Fig. 1d-f) approaching present day values. However, much of
15 Antarctica remained heavily glaciated well into the Holocene, with the majority of its deglacial
16 ice melting/gloss taking place between 12 and 6 ka (Argus et al., 2014; Briggs et al., 2014;
17 Mackintosh et al., 2014). Antarctica's total contribution to post-glacial eustatic sea level is
18 poorly constrained, but recent studies have not supported LGM contributions greater than about
19 15 m eustatic sea level equivalent (Bentley et al., 2014; Briggs et al., 2014; Golledge et al.,
20 2013; Mackintosh et al., 2011; Philippon et al., 2006; Whitehouse et al., 2012), emphasising
21 the dominance of North American and Eurasian Ice Sheet dynamics in the global sea level
22 record during the last deglaciation (Argus et al., 2014; Lambeck et al., 2014; Peltier et al., 2015).
23 It should be noted that there is some controversy over whether deglacial ice sheet
24 reconstructions close the global sea level budget (Clark and Tarasov, 2014), with a potential
25 LGM shortfall of 'missing ice'.

26 The last deglaciation is not only an interesting case study for understanding multi-millennial
27 scale processes of deglaciation, but also provides the opportunity to study shorter and more
28 dramatic climate changes. Superimposed over the gradual warming trend (EPICA Community
29 Members, 2004; Jouzel et al., 2007; Petit et al., 1999; Stenni et al., 2011) are several abrupt
30 climate transitions lasting from a few years to a few centuries (examples of which are given

1 below) and it remains a challenge to reconstruct or understand the chain of events surrounding
2 these instances of rapid cooling and warming.

3 *Heinrich Event 1* (approx. 16.8 ka; Hemming, 2004) occurred during the relatively cool
4 Northern Hemisphere Heinrich Stadial 1 (~18-14.7 ka). It was characterised by the release of a
5 vast number of icebergs from the North American and Eurasian ice sheets into the open North
6 Atlantic, where they melted. The existence of these iceberg ‘armadas’ is evidenced by a high
7 proportion of ice rafted debris in North Atlantic sediments between 40° N and 55° N,
8 predominantly of Laurentide (Hudson Strait) provenance (Hemming, 2004 and references
9 therein). There are several competing theories for the cause of Heinrich Event 1. There is a
10 substantial body of evidence ~~to suggest~~suggesting that it occurred during or was precursory to
11 a period of Atlantic Meridional Overturning Circulation (AMOC) slow down (e.g. Hall et al.,
12 2006; Hemming, 2004; McManus et al., 2004) and weak North Atlantic Deep Water (NADW)
13 formation (e.g. Keigwin and Boyle, 2008; Roberts et al., 2010) under a relatively cold, Northern
14 Hemisphere surface climate (Shakun et al., 2012). Even though the interpretation of a cause
15 and effect link between Heinrich Event 1 and the diminished strength of the AMOC remains
16 rather compelling (e.g. Kageyama et al., 2013), it is increasingly being suggested that the
17 melting icebergs might not have caused the recorded AMOC slow down, but may have provided
18 a positive feedback to amplify or prolong AMOC weakening and widespread North Atlantic
19 cooling (e.g. Álvarez-Solas et al., 2011; Barker et al., 2015),~~whilst also causing mid-latitude~~
20 Atlantic sea surface warming through northward expansion of the subtropical gyre.

21 During the subsequent 14.2-14.7 ka interval, Northern Hemisphere temperatures are seen to
22 have risen by as much as 14.4 ± 1.9 °C in just a few decades (Buizert et al., 2014; Goujon et
23 al., 2003; Kindler et al., 2014; Lea et al., 2003; Severinghaus and Brook, 1999), with a dramatic
24 shift in some components of Greenland climate taking place in as little as one to three years
25 (Steffensen et al., 2008). This abrupt event is termed the *Bølling Warming* or *Bølling Transition*
26 (Severinghaus and Brook, 1999). At roughly the same time (~14.6 ka), there was a rapid jump
27 in global sea level of 12-22 metres in around 350 years or less, known as *Meltwater Pulse 1a*
28 (MWP1a; Deschamps et al., 2012). It is not known exactly which ice mass(es) contributed this
29 40 mm yr^{-1} (or greater) flux of water to the oceans (e.g. Lambeck et al., 2014; Peltier, 2005).
30 Some ~~older~~ studies have mainly attributed it to a southern source (Bassett et al., 2005, 2007;
31 Carlson, 2009; Clark et al., 1996, 2002; Weaver et al., 2003), whereas more recent work has

1 suggested that at most, less than 4.3 metres eustatic sea level equivalent of meltwater could
2 have come from Antarctica (Argus et al., 2014; Bentley et al., 2010, 2014; Briggs et al., 2014;
3 Golledge et al., 2012, 2013, 2014; Licht, 2004; Mackintosh et al., 2011, 2014; Whitehouse et
4 al., 2012) and that Northern Hemisphere ice was the primary contributor (Aharon, 2006;
5 Gregoire et al., 2012; Keigwin et al., 1991; Marshall and Clarke, 1999; Peltier, 2005; Tarasov
6 et al., 2012; Tarasov and Peltier, 2005). Exactly how the Bølling Warming and MWP1a are
7 linked, or what triggered either, remains uncertain.

8 Ice core records of δD indicate that from around 14.5 ka to 12.8 ka, the general trend of
9 increasing Southern Hemisphere warming, temporarily stalled (Jouzel et al., 2007; ice core
10 chronology from Veres et al., 2013) for a period known as the *Antarctic Cold Reversal* (Jouzel
11 et al., 1995). Southern Hemisphere cooling is thought to have been relatively widespread,
12 extending from the South Pole to the southern mid-latitudes, with glacial readvance (or stall in
13 glacial retreat) recorded to have peaked 13.0-14.2 ka in Patagonia (García et al., 2012; Kaplan
14 et al., 2011; Strelin et al., 2011) and ~13.0 ka in New Zealand (Putnam et al., 2010; Rother et
15 al., 2014). There are several hypotheses for the cause of the Antarctic Cold Reversal. For
16 example, some have linked it to a change in ocean circulation induced by the delivery of
17 Antarctic ice melt to the Southern Ocean (Menviel et al., 2010, 2011), or possibly as a bipolar
18 response to AMOC recovery and Northern Hemisphere warming during the Bølling Warming
19 (Menviel et al., 2011; Stocker, 1998). Using a CMIP5 level coupled atmosphere-ocean model,
20 Peltier and Vettoretti (2014) and Vettoretti and Peltier (2015) have recently shown that ice core
21 inferred southern hemisphere cooling and northern hemisphere warming could have been
22 caused by a nonlinear salt oscillator mechanism. Others have argued that a change in Southern
23 Hemisphere winds and ocean circulation is the explanation; for example, a simultaneous
24 northward migration of the southern Subtropical Front and northward expansion of cold water
25 originating in the Southern Ocean (Putnam et al., 2010). The ongoing disagreement over the
26 timing, duration and extent of the Antarctic Cold Reversal means that its cause is difficult to
27 pin down.

28 The next event of particular interest is the *Younger Dryas cooling*, when Northern Hemisphere
29 temperatures are thought to have dropped by several degrees at 12.8-11.7 ka and most
30 prominently in high latitudes (Buizert et al., 2014; Heiri et al., 2007; Lea et al., 2003; Liu et al.,
31 2012; Simonsen et al., 2011; Steffensen et al., 2008). The event presents a conceptual paradox;

1 the magnitude of the cooling is difficult to reconcile with rising atmospheric CO₂
2 (approximately +10 ppm compared to the earlier Bølling period ~ 14.5 ka; Bereiter et al., 2015)
3 and increasing boreal summer insolation (Berger and Loutre, 1991). It is possible that changes
4 in the atmospheric hydrological cycle, such as a shift in source moisture region, could be partly
5 responsible for the δ¹⁸O signal, requiring a smaller temperature anomaly to match the records
6 (Liu et al., 2012). For the climate cooling itself, a rerouting of North American freshwater
7 discharge to the Arctic and/or Atlantic Oceans might have caused a reduction in NADW
8 formation (Broecker et al., 1989; Condron and Winsor, 2012; Tarasov and Peltier, 2005).
9 Simulating this period within the context of the preceding climate evolution,⁻ could be key to
10 understanding exactly what the surface climate and deep ocean changes were during the
11 Younger Dryas, and how these relate to contemporaneous proxy records (e.g. Buizert et al.,
12 2014).

13 In this description, we have sought to capture some of the last deglaciation's main climatic
14 events, but there are others that could shape the focus of further study in the working group.
15 For example, early on in the period there is evidence of around 10 m sea level rise taking place
16 in 500-800 years around 20-19 ka (Clark et al., 2004; Clark and Mix, 2002; De Deckker and
17 Yokoyama, 2009; Yokoyama et al., 2001a, 2001b). Whilst the event itself remains somewhat
18 controversial (Cabioc'h et al., 2003; Hanebuth et al., 2000, 2009; Peltier and Fairbanks, 2006;
19 Shennan and Milne, 2003), it could be the expression of accelerating deglacial ice melt
20 following the Last Glacial Maximum. More recently, the Barbados record of relative sea level
21 history indicates that following the Younger Dryas cooling episode, there may have been
22 another meltwater pulse (Fairbanks, 1989; Peltier and Fairbanks, 2006), referred to as
23 Meltwater Pulse 1b. Significant debate surrounds the magnitude and timing of Meltwater Pulse
24 1b (Bard et al., 1996; Cabioc'h et al., 2003; Cutler et al., 2003; Edwards et al., 1993; Shennan,
25 1999; Stanford et al., 2011) and even its existence, because similar to the 19 ka event, it is not
26 seen in all sea level records spanning the interval (e.g. Bard et al., 1996, 2010; Hanebuth et al.,
27 2000). However, evidence of rapid Antarctic retreat around the time of the event could provide
28 a possible cause for this late deglacial rapid sea level rise (Argus et al., 2014).

1.2 Transient modelling of the last deglaciation

Transient modelling of the last deglaciation is valuable for examining dynamic and threshold behaviours (Braconnot et al., 2012) endemic to the Earth's non-stationary climate system, especially ice-ocean-atmosphere interactions. It is the best tool for reaching a comprehensive understanding of complex and interrelating climate processes with specific regard to chains of events.

Such simulations are useful for examining the effect of temporally varying climate forcings across the globe and in different environmental systems: what geographical patterns arise and how are they connected, how do these vary through time from seasonal to millennial time scales, and how long does it take before a change in forcing is manifested in a climate response? The spatial coherency of specific events can be investigated to identify processes for simultaneous change as well as lead/lag mechanisms. For example, Roche et al. (2011) investigated patterns of spatial variability in the deglaciation as caused by long-term changes in orbital parameters, atmospheric greenhouse gas concentrations, and ice sheet extent/topography. The results indicated a simultaneous onset of hemispheric warming in the North and South, showing that obliquity forcing was the main driver of the early deglacial warming. In the same investigation, it was found that sea-ice covered regions were the first parts of the world to exhibit significant rises in temperature, implying that a better knowledge of sea-ice evolution could be key to fully understanding the trigger for widespread deglaciation and warming feedbacks. A further example of the insights available into lead-lag relationships provided by long, transient climate simulations under glacial boundary conditions is provided by the previously referenced Dansgaard-Oeschger oscillation-related analyses of Peltier and Vettoretti (2014) and Vettoretti and Peltier (2015), which appear to mimic the Heinrich Stadial 1 to Bølling transition.

Through comparison to geological timeseries data, transient simulations enable the 'fingerprinting' of specific climate processes to find out what mechanisms [in the model] can cause recorded climate signals. Comparing complex, global-scale models to combined geological records can provide multiple 'fingerprints' in different variables from different archives and in different locations to help narrow down plausible scenarios. For example, Menzel et al. (2011) ran a suite of simulations, varying oceanic meltwater fluxes through the last deglaciation in order to identify which freshwater-forcing scenarios reproduce the Atlantic

1 Ocean circulation state implied by sedimentary records of AMOC strength/depth and
2 ventilation age (Gherardi et al., 2005; McManus et al., 2004 with ages shifted as per Alley,
3 2000; Thornalley et al., 2011) as well as the Northern Hemisphere surface climate (Alley, 2000;
4 Bard, 2002; Bard et al., 2000; Heiri et al., 2007; Lea et al., 2003; Martrat et al., 2004, 2007). It
5 was argued that such climate simulations could be used to improve constraints on the timing,
6 duration, magnitude, and location of meltwater inputs to the global ocean.

7 Liu et al. (e.g. 2009) used climate ‘fingerprinting’ to identify possible mechanisms for the
8 abrupt Bølling Warming Event, finding that in their model, a forced cessation of freshwater
9 inputs to the North Atlantic (representing ice sheet melt) superimposed on a steady increase in
10 atmospheric CO₂ caused an abrupt resumption in the strength of the AMOC (almost matching
11 a record produced by McManus et al., 2004). This in turn induced a rapid warming in Northern
12 Hemisphere surface climate (close to records from Bard et al., 2000; Cuffey and Clow, 1997;
13 and Waelbroeck et al., 1998) and an increase in tropical rainfall over the Cariaco Basin
14 (comparable to Lea et al., 2003), whilst Antarctic surface temperatures remained relatively
15 stable (similar to Jouzel et al., 2007). Using a suite of simulations from the same model, Otto-
16 Bliesner et al. (2014) went on to suggest that a combination of rapid strengthening of NADW
17 seen by Liu et al. (e.g. 2009) and rising greenhouse gas concentrations was responsible for
18 increased African humidity around 14.7 ka, matching the model output to a range of regional
19 climate proxies (including deMenocal et al., 2000; Tierney et al., 2008; Tjallingii et al., 2008;
20 Verschuren et al., 2009; Weijers et al., 2007).

21 Thus, climate proxy fingerprinting can be useful for understanding the spatial coherency of
22 climatic changes and their underlying mechanisms. However, correlation between model and
23 geological data does not guarantee that the correct processes have been simulated; there is
24 always the problem of *equifinality*, whereby the same end state can be reached by multiple
25 means. In a process sense, this may be particularly uncertain when a model does not reproduce
26 the full chain of events that led to a distinguishable climatic signal. For example, mechanisms
27 for many of the major changes in oceanic freshwater inputs proposed by Liu et al. (2009) and
28 Men viel et al. (2011) have not yet been directly simulated (e.g. by dynamic ice sheet models).
29 In both studies, they are imposed as model boundary conditions. Further simulations with
30 different forcing scenarios and from a range of models would help to address such uncertainties.

1 Transient simulations of the last deglaciation also provide necessary boundary conditions for
2 modelling a variety of Earth System components that may not be interactively coupled to the
3 climate model being used. For example, Gregoire et al. (2015) drove a dynamic ice sheet model
4 with climate data produced by a similar set of simulations to Roche et al. (2011). Using a low
5 resolution GCM, individual climate forcings – including orbit, greenhouse gases, and meltwater
6 fluxes – were isolated so that their relative contribution to melting the modelled North American
7 ice sheets could be examined. The work concluded that the last deglaciation was primarily
8 driven by changes in Northern Hemisphere insolation, causing around 60% of the North
9 American Ice Sheet melt, whilst increasing CO₂ levels were responsible for most of the
10 remaining changes (Gregoire et al., 2015). The sufficiency of these two forcings for North
11 American glaciation/deglaciation had previously also been identified with fully coupled
12 glaciological and energy balance climate models (Tarasov and Peltier, 1997). Gregoire et al
13 (2012) were also able to highlight a ~~possible~~ ‘saddle-collapse’ mechanism, whereby gradual
14 warming trends could result in abrupt ice sheet melting events, ~~such as MWP1a and the 8.2 kyr~~
15 ~~Event,~~ when a threshold in ice mass balance was crossed. ~~The opening of the ice free corridor~~
16 ~~between the Cordilleran and Laurentide ice sheets has long been built into the ICE NG, Tarasov~~
17 ~~and Peltier and Tarasov et al. sequence of models as geological inferences indicate that it,~~
18 ~~which could have~~ occurred ~~around the same time as during~~ MWP1a- ~~and the 8.2 kyr event..~~

19 A further example is given by Liu et al. (2012), who carried out an asynchronous (or ‘offline’)
20 coupling between simulated sea surface temperatures and an isotope-enabled atmospheric
21 model to investigate the Younger Dryas cooling event (~12 ka). The results revised the
22 presupposed Greenland temperatures at this time by 5 °C, demonstrating that changes in
23 moisture source must be an important consideration for the robust interpretation of Greenland
24 ice core δ¹⁸O records and our understanding of high-latitude climate sensitivity. More recently,
25 the same methodology was applied to understanding Chinese cave records of the East Asian
26 Summer Monsoon 21-0 ka (Liu et al., 2014), not only to better interpret what the speleothem
27 δ¹⁸O tells us about regional hydroclimate variability, but also to understand the wider
28 teleconnections controlling those patterns.

29 In addition, there are now transient simulations of the last deglaciation from climate models
30 that have been interactively coupled with dynamic ice sheet models (Bonelli et al., 2009;
31 Heinemann et al., 2014) and isotope systems (Caley et al., 2014). Furthermore, a fast Earth

1 System Model of Intermediate Complexity (EMIC) that includes an interactive ice sheet model
2 has been used to look at Earth System dynamics (the role of orbital cycles, aeolian dust,
3 subglacial regolith properties, the carbon cycle, and atmospheric trace gases) on much longer,
4 glacial-interglacial timescales >120 ka and encompassing the last deglaciation (Bauer and
5 Ganopolski, 2014; Brovkin et al., 2012; Ganopolski et al., 2010; Ganopolski and Calov, 2011).
6 However, the older, uncoupled climate-ice sheet model approach discussed above remains
7 useful because it enables a wider suite of models to be employed than would otherwise be
8 feasible due to limited computational efficiency (e.g. of state-of-the-art, high
9 resolution/complexity models) or software engineering capability. It may also allow for the
10 same Earth System component model (e.g. of ice sheets or $\delta^{18}\text{O}$) to be driven by multiple
11 climate models, in order to examine the range of responses and assess [climate] model
12 performance.

13 With sufficient computational power to make long simulations of the last deglaciation a feasible
14 undertaking, it is timely to coordinate new efforts to ensure that a framework exists to (i) utilise
15 the cutting edge science in climate modelling and palaeoclimate reconstruction, and (ii) robustly
16 intercompare simulations run with different models by different groups and palaeoclimatic data.

17 **1.3 Establishing a new PMIP working group**

18 For [more than](#) twenty years, the Paleoclimate Modeling Intercomparison Project (PMIP) has
19 been internationally coordinating multi-model simulations with complex climate models in
20 order to evaluate model performance and better understand [past] climate changes (Braconnot
21 et al., 2007, 2012; PMIP website, 2007). Currently entering its fourth phase, PMIP is a growing
22 organisation that continues to contribute towards other coordinated efforts to understand present
23 day climate change; including the Coupled Model Intercomparison Project (Taylor et al., 2011a,
24 CMIP; e.g. 2011b) and the Intergovernmental Panel on Climate Change's (IPCC) Assessment
25 Reports (e.g. the Fifth Assessment Report; Flato et al., 2013; Masson-Delmotte et al., 2013). It
26 encompasses a broad range of models, from very fast, lower resolution EMICS, through a range
27 of coupled GCMs to the latest generation of higher resolution and complexity Earth System
28 Models. Thus, the main challenges for the fourth Phase of PMIP include: designing experiments
29 that are suitable for all of its participants; addressing sufficiently fundamental questions to be
30 of interest to the EMIC community; defining adequately focused scope for the feasible

1 participation of the latest generation of ESMs; and prescribing flexible model setups that can
2 be implemented in this range of models, whilst maintaining the ability to robustly compare
3 results. [In addition, a continuing challenge for PMIP is to assemble suitable palaeoclimatic](#)
4 [datasets for comparison to model results.](#)

5 One of the most recent working groups to be established in PMIP is the Last Deglaciation
6 Working Group. With the aim of coordinating transient simulations of the last deglaciation, the
7 challenge of including the full range of PMIP models is at the forefront of our experiment
8 design. The experiment will be partitioned into three phases (Fig. 1b and Sect. 4), which will
9 form milestones for managing its long duration (12 thousand years) as well as for scheduling
10 any shorter, alternative simulations to the Core.

11 The aim of this paper is to outline the model setup for the transient Core [simulation](#)
12 [of experiment for](#) the last deglaciation, specifically for the sub-period of 21-9 ka. Prescribed
13 boundary conditions include orbital parameters, atmospheric trace gases and ice sheets. In
14 association with the ice sheet reconstructions, we also provide bathymetric, orographic and
15 [land-sea mask evolution-, as well as make recommendations for freshwater forcing \(or global](#)
16 [ocean salinity changes\) through the period.](#)

17 **1.4 Approach**

18 One of the roles of PMIP has been to systematically study the ability of climate models to
19 retrodict different past climates for which there are ‘observational’ data from geological
20 archives (e.g. Braconnot et al., 2000, 2007, 2012; Haywood et al., 2010; Joussaume et al., 1999;
21 Kageyama et al., 2006; Kohfeld and Harrison, 2000; Masson-Delmotte et al., 2006; Otto-
22 Bliesner et al., 2009; Weber et al., 2007). In this vein, many palaeoclimate model
23 intercomparison projects have been designed to facilitate the robust comparison of results from
24 the same ‘experiment’ (i.e. simulation set) across a range of different models, usually taking a
25 prescriptive approach to model setup to ensure that any differences observed in the results are
26 attributable to differences in model structure and not to differences in chosen ‘boundary
27 conditions’ and climate forcings. However, as Schmidt et al. (2011) point out, the choice of one
28 particular configuration from a range of plausible boundary conditions and forcings is often
29 arbitrary and does not account for uncertainties in the data used for developing the
30 forcings/boundary conditions. Moreover, in designing the PMIP last deglaciation experiment,

1 we have attempted to strike a balance between establishing a framework within which to assess
2 model differences and performance, and taking the opportunity to utilise the full range of PMIP
3 climate models (Earth System, General Circulation and Intermediate Complexity) to examine
4 uncertainties in deglacial forcings, trigger-mechanisms and dynamic feedbacks. In short, when
5 we do not precisely know the climate forcing for an event, or the temporal evolution of model
6 boundary conditions, it is more efficient to compare the results from models that use different
7 forcings with geological and palaeoclimatic data than to run one scenario with all models and
8 all scenarios with all models. The aim is to use the results of the comparison to narrow down
9 the range of uncertainty in the forcings/boundary conditions and reach a better understanding
10 of underlying climate mechanisms.

11 Consequently, forcings/boundary conditions that are relatively well established (atmospheric
12 trace gases and orbital parameters) are tightly constrained in the Core experiment design. Others
13 are given with multiple precisely described possibilities to choose from (ice sheet
14 reconstructions) and the remainder (e.g. freshwater/salinity, aerosols and vegetation) are left to
15 the discretion of individual participants, although we recommend. Recommendations will be
16 made for the latter grouping of forcings/boundary conditions; for example, freshwater/global
17 salinity fluxes that are consistent with the provided ice sheet evolutions, and the use of
18 preindustrial aerosol and/or vegetation values when they are not model prognostics; but a
19 flexible approach is advantageous not only scientifically (i.e. for examining the climatic
20 response to uncertain forcings, see above), but also practically (for accommodating the wide
21 range of participating models). Further to this, it will be left to the expert user to decide how
22 often to make manual updates to those boundary conditions that cannot evolve automatically in
23 the model, such as bathymetry, orography and land sea mask. This is also necessary because of
24 the specific technical and resource requirements associated with setting up and running each
25 participant model.

26 In addition to the Core, we will also coordinate additional a series of experiments that are
27 designed to:

28 (i) explore uncertainties in the boundary conditions and climate forcings.
29 (ii) test specific hypotheses for mechanisms of climate change and to explain individual
30 events.

1 (iii) focus on shorter time periods (for example, abrupt events) and thus include
2 computationally expensive models for which a twelve thousand year simulation is
3 unfeasible.

4 These optional simulations will be referred to as *focussed* experiments, and participants are
5 encouraged to contribute towards the design and coordination of these simulations within the
6 working group [\(dedicated Wiki page to coordinate these here: <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglab:index>\)](#).

8 The start date for the experiment has been chosen to be in line with PMIP's historical definition
9 of the LGM; 21 ka (Abe-Ouchi et al., 2015; e.g. Braconnot et al., 2000; Kohfeld and Harrison,
10 2000). However, we are aware that some groups may prefer to begin their simulations from the
11 earlier date of 26 ka (around the last sea level lowstand; Clark et al., 2009; Lambeck et al.,
12 2014; Peltier and Fairbanks, 2006) and both orbital and atmospheric trace gas parameters will
13 be provided from this earlier date. Although the working group's focus will at least initially be
14 21-9 ka, boundary conditions for the Core [simulation simulations](#) will be provided from 21 ka
15 to the preindustrial (26 ka to the preindustrial for orbital insolation and trace gases).

16 The following is not meant to be an exhaustive review of climate forcing reconstructions
17 through the last deglaciation. Instead, our intention is to consolidate the current knowledge in
18 a practical experiment design for a range of climate models. Within this coordinated context,
19 the aim is to explore the forcings and underlying feedback mechanisms for the rapid climate
20 events that punctuated the gradual warming and deglaciation of the Earth.

21 The paper is structured so that Sect. 2 outlines the model boundary conditions and climate
22 forcings for the Core [simulation experiment](#). Section 3 presents how we will ensure the feasible
23 participation of a range of climate models with different complexity and computational
24 efficiency, as well as the plan to run additional, targeted, hypothesis- and sensitivity-led
25 simulations. Section 4 discusses the three phases of the long Core experiment.

26 **2 Core [simulation experiment](#) (21 ka to 9 ka)**

27 The Core [simulation for simulations](#) of the last deglaciation will focus on the period from 21 ka
28 to 9 ka, although there will also be the option to spin up the simulation with time-evolving
29 orbital and trace gas parameters from 26 ka and all boundary conditions will be available from
30 21 ka to the preindustrial. Recommendations for the initialisation state at 21 ka are summarised

1 in Table 1 and described below (Sect. 2.1). Prescribed boundary conditions include insolation
2 via the Earth's astronomical parameters (Sect. 2.2), atmospheric trace gases (Sect. 2.3), ice
3 sheets (Sect. 2.4), meltwater fluxes (Sect. 2.5), and orography/bathymetry (Sect. 2.6), as
4 summarised in Table 2. Boundary condition data for the Core [simulationexperiment](#) are
5 provided on the PMIP wiki; <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglab:bc:core>
6 (PMIP Last Deglaciation Working Group, 2015).

7 **2.1 Last Glacial Maximum spinup**

8 There is a choice of two possibilities for starting the last deglaciation Core
9 [simulation](#)[simulations](#). Either the simulation should be initialised from the end of a spun-up,
10 PMIP-compliant LGM (21 ka) simulation, or a simulation with transient orbital and trace gas
11 forcing should be run from an earlier time period (orbital and trace gas parameters will be
12 provided from 26 ka onwards). Whichever method is applied, we require that it is
13 comprehensively documented along with information on the model's state of spinup at 21 ka
14 (e.g. timeseries of surface climates, maximum strength of the North Atlantic Meridional
15 Overturning Circulation stream function, net radiation at the top of the atmosphere [etc.](#)).

16 **2.1.1 Equilibrium-type spinup (21 ka)**

17 For setting up an equilibrium-type spinup, please make sure to use the following constraints,
18 which may differ from other PMIP 21 ka simulation protocols:

- 19 - Insolation should be set so that eccentricity is 0.018994, obliquity is 22.949°,
20 perihelion–180° is 114.42°, the date of the vernal equinox is 21st March at noon,
21 and the solar constant is the same as for the preindustrial (e.g. 1365 W m⁻², as in the
22 PMIP3-CMIP5 preindustrial experiment). These are consistent with previous PMIP
23 LGM boundary conditions (PMIP LGM Working Group, 2010).
- 24 - Prescribed atmospheric trace gases should be as follows: CO₂ at [188190](#) ppm, CH₄
25 at 375 ppb, N₂O at 200 ppb (Fig. 3), with CFCs at 0 and O₃ at the PMIP3-CMIP5
26 preindustrial value (e.g. 10 DU). This is to be compatible with the time-evolving
27 boundary conditions for the Core [simulation](#)[simulations](#) (Sect. 2.3). Note that the
28 LGM atmospheric CO₂ and CH₄ concentrations have changed slightly from earlier
29 LGM experiments (e.g. PMIP3, which used 185 ppm and 350 ppb, respectively);

PMIP LGM Working Group, 2010). However, N₂O remains at 200ppb, which is more representative of the longer glacial period than the 187 ppb concentration recorded at 21 ka (Fig. 3c). These updates are in line with the latest ice core age model, (AICC2012; Veres et al., 2013) [and records](#) (Bereiter et al., 2015; Schilt et al., 2010), [which is, which are](#) also used for the transient forcings described below (Sect. 2.3).

- Prescribed ice sheets should use either the GLAC-1D or ICE-6G_C reconstruction at 21 ka (see Sect. 2.4). The associated topography and coastlines should be used as per the chosen ice sheet reconstruction: [Beyond maintaining consistency with the coastlines](#), it is optional whether or not to implement the associated bathymetry- [and participants should adapt the bathymetry according to their model's capabilities \(for example, depending on whether the spatial resolution allows for it or makes this a useful adaptation\)](#). These data will be provided with the ice sheet reconstructions. Whichever ice sheet reconstruction is chosen for the LGM spinup should be carried through to the Core transient simulation.
- Global ocean salinity should be +1 psu, compared to preindustrial, to account for the increased terrestrial ice mass at the LGM (PMIP LGM Working Group, 2015).
- Any other boundary conditions should be set to be consistent with the Core transient simulation to follow (Sect. 2.2-2.7).

On the freshwater budget, PMIP advises groups to ‘carefully check the fresh water budget in their LGM experiments in order to avoid unnecessary drifts of the ocean salinity. It can be necessary to route the snow which has fallen in excess on the ice sheets to the ocean. Given the change in coastlines, it is also sometimes necessary to relocate the large river estuaries on the coast’ (PMIP LGM Working Group, 2015). Tarasov and Peltier (2006) [providesprovide](#) a glaciological example of the possible re-routings for North America. [As they become available, routing maps for the Last Glacial Maximum continents will be provided on the last deglaciation PMIP Wiki \(address above\)](#).

The integration time required for spinning up the LGM climate state should be decided on a case-by-case basis by the user. Groups may choose to initialise their equilibrium-type simulation from other PMIP LGM runs. However, [please be careful.caution is advised](#). Some

1 of the boundary conditions for ~~the PMIP4 CMIP6 (not finalised at the time of writing)~~ and
2 previous PMIP LGM simulations are different to the setup outlined here, specifically in terms
3 of ice sheets and trace gases concentrations, and therefore need to be adapted to match these
4 requirements. ~~Please also~~The protocol for the PMIP4-CMIP6 (being finalised at the time of
5 writing) is currently compatible with the LGM spin-up described here. Therefore, provided that
6 either the ICE-6G C or GLAC-1D ice sheet reconstruction is used for both the LGM spin-up
7 and transient run, the PMIP4-CMIP6 LGM simulation can be used to initialise transient
8 simulations of the last deglaciation without alteration. Please provide timeseries data for the
9 diagnosis of model [dis]equilibrium at 21 ka (introduction to Sect. 2.1).

10 2.1.2 Transient orbital and trace gas parameters (26-21 ka)

11 If this is the preferred option to initialise the Core, it is recommended that the simulation is
12 setup as per Sect. 2.1.1, but with time-evolving orbital and trace gas parameters instead of fixed
13 ones. Specifically for orbit, the eccentricity, obliquity, perihelion–180° and date of the vernal
14 equinox values listed above should be replaced with their transient equivalents, as per Berger
15 (1978). For the atmospheric trace gases, carbon dioxide, methane and nitrous oxide values
16 should be replaced with the transient equivalents provided on the PMIP Wiki
17 (<https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglab:core>) and according to
18 LüthiBereiter et al. (2015), Loulergue et al. (2008) and Schilt et al. (2010), respectively, on the
19 AICC2012 chronology (Veres et al., 2013); Fig. 3.

20 In this case, all other boundary conditions should remain fixed in line with the LGM
21 equilibrium-type experiment design until 21 ka, when the fully transient Core simulation
22 begins:simulations begin. This transient spin-up can be initialised from a spun-up previous
23 LGM, cold ocean, preindustrial, or observed present day ocean simulation.

24 2.2 Insolation (21-9 ka)

25 As per Sect. 2.1, the solar constant should be fixed to the established preindustrial conditions
26 (e.g. 1365 W m⁻²) throughout the run, which is the PMIP preindustrial experiment setup (PMIP
27 LGM Working Group, 2015). However, the orbital parameters should be time-evolving through
28 the deglaciation to follow Berger (1978); e.g. Fig. 1c.

2.3 Atmospheric trace gases (21-9 ka)

For the deglaciation, CFCs should be fixed at 0, and O₃ should be set to PMIP3-CMIP5 preindustrial values (e.g. 10 DU), as used for the LGM. When a model is not running with dynamic atmospheric chemistry, the remaining trace gases should be time-evolving, with CO₂ following Lüthi et al. (2015), CH₄ following Loulergue et al. (2008) and N₂O following Schilt et al. (2010), all adjusted to the AICC2012 chronology (Veres et al., 2013); Fig. 1d-f.

The atmospheric CO₂ concentrations provided by Bereiter et al. (2015) is a composite dataset, combining previous Antarctic ice core records and composites (for the period 26-0 ka: Ahn and Brook, 2014; Lüthi et al., 2008; MacFarling Meure et al., 2006; Marcott et al., 2014; Rubino et al., 2013; Siegenthaler et al., 2005) on the AICC2012 timescale of Veres et al. (2013) Temporally higher resolution CO₂ data from the West Antarctic Ice Sheet Divide has been provided by Marcott et al. , spanning 23-9 ka ('WDC' on Fig. 3a). However, the newer data are consistently offset from other Antarctic ice core data by ~4 ppm and the cause for this remains unresolved. Furthermore, although the data encompasses the last deglaciation (and the period we are focussing on; 21-9 ka), it would not be easily spliced into a longer record (e.g. for groups wishing to run their simulations through to the present day). This is why the higher resolution data will not be used for the Core, reverting to the older record from Lüthi et al. However, it to produce a high resolution record that is consistent with the other, lower resolution trace gas records used in this experiment (CH₄ and N₂O as discussed above). Groups are free to decide on the temporal resolution of trace gas model inputs based on these records and if lower resolution is employed, the method used to smooth or create a spline through the data should be fully documented. Exploring the influence of CO₂ resolution on the climate system may form the basis of a coordinated additional simulation, which will be optional for participant groups. Other sensitivity type simulations could also be coordinated to assess the influence of timing in the CO₂ records on climate and ice sheet evolution, addressing age model uncertainty. The details of the setup for such focussed simulations (also discussed in Sect. 3) will be discussed and determined at a later date.

It is noted that the N₂O value from Schilt et al. (2010) and Veres et al. (2013) does not match the previously defined LGM N₂O concentration (Sect. 2.1.1); 187 ppb compared to 200 ppb (Fig. 3c). This is because the N₂O record is highly variable during the last glacial lowstand (26-21 ka), with a range of ~33 ppb (183-216 ppb) and a mean of 201 ppb. Thus 200 ppb seems a

1 reasonably representative N₂O concentration for the spinup phase of the simulation, although
2 the Core [simulations](#) will start with the more chronologically accurate value of 187
3 ppb.

4 **2.4 Ice sheet reconstructions (21-9 ka)**

5 For the Core experiment, ice sheet extent and topography should be prescribed from one of two
6 possible reconstructions: ICE-6G_C (Fig. 2a and Fig. 4a) and GLAC-1D (Fig. 2b and Fig. 4b).

7 The ICE-6G_C reconstruction is fully published (Argus et al., 2014; Peltier et al., 2015), and
8 the reader is directed to this literature for further information. The GLAC-1D reconstruction is
9 combined from different sources (Briggs et al., 2014; Tarasov et al., 2012; Tarasov and Peltier,
10 2002) and whilst it is mostly published, there are some new components; therefore, a short
11 description follows. The Eurasian and North American components are from Bayesian
12 calibrations of a glaciological model (Tarasov et al., 2012; this study), the Antarctic component
13 is from a scored ensemble of 3344 glaciological model runs (Briggs et al., 2014) and the
14 Greenland component is the hand-tuned glaciological model of Tarasov and Peltier (2002)
15 [updated to the GICC05 age chronology](#) (Rasmussen et al., 2006). All four of the GLAC-1D
16 ice sheet components employ dynamical ice sheet models that have been constrained with
17 relative sea level data. Where available, they have also been constrained by geologically-
18 inferred deglacial ice margin chronologies, pro-glacial lake levels, ice core temperature
19 profiles, present-day vertical velocities, past ice thickness, and present day ice configuration.
20 Details of exactly how these constraints were derived and applied are given in the relevant
21 references above. The four components (North American, Eurasia, Antarctica and Greenland)
22 were combined under Glacial Isostatic Adjustment (GIA) post-processing for a near-
23 gravitationally self-consistent solution (Tarasov and Peltier, 2004), which was tested against
24 complete Glacial Isostatic Adjustment solutions ([Tarasov, pers. comm. 2014](#)). The topography
25 in the global combined solution was adjusted in Patagonia and Iceland following ICE-5G
26 (Peltier, 2004), but the changes in these ice caps are not reflected in the ice mask.

27 Both datasets include ice extent and topography at intervals of 1,000 years or less through the
28 deglaciation. [Ice-Specifically, the ICE-6G_C reconstruction is provided at 1,000-year intervals](#)
29 [for the period spanning 26-21 ka and 500-year intervals for 21-0 ka. For GLAC-1D, the data](#)

1 [are at 100-year intervals 21-0 ka. In both reconstructions, ice](#) extent is provided as a fractional
2 ice mask [for ICE-6G_C and a binary ice mask in GLAC-1D.](#)

3 The two reconstructions incorporate similar constraints for North American ice sheet extent
4 (i.e. Dyke, 2004). For Eurasia, ICE-6G_C follows the ice extent provided by Gyllencreutz et
5 al. (2007), whereas GLAC-1D uses data from Hughes et al. (2015). The reconstructions only
6 differ slightly in their ice extent evolution (Figures 2 and 4), for example the Barents Sea
7 deglaciates earlier in GLAC-1D than in ICE-6G_C (Fig. 2). The main differences between the
8 reconstructions are in the shape and volume of individual ice sheets. In particular, the North
9 American Ice Sheet reaches an elevation of 4000 m in ICE-6G_C, but is only 3500 m high in
10 GLAC-1D. Similarly, the shape and thickness of the Barents Sea Ice Sheet are not the same in
11 the two reconstructions. The ICE-6G_C dataset is [been](#) provided at [both](#) 1 degree [horizontal](#)
12 [resolution](#) and 10 minute horizontal resolution, GLAC-1D is provided at 1 degree [\(longitude\)](#)
13 [× 0.5 degree \(latitude\)](#) horizontal resolution.

14 Ice surface elevation (topography) should be implemented as an anomaly from present day
15 topography and added to the model's present day topography after regridding onto the model
16 resolution, following the [previous](#) LGM experimental protocol (PMIP LGM Working Group,
17 2010, 2015). Land surface properties will need to be adjusted for changes in ice extent. Where
18 ice retreats, land surface should be initialised as bare soil if a dynamic vegetation model is used,
19 otherwise use prescribed vegetation (see Sect. 2.7) with appropriate consideration of soil
20 characteristics. Where ice is replaced by ocean, it is advised to follow the procedure for
21 changing coastlines described in Sect. 2.7. Inland lakes can be prescribed based on the ice sheet
22 and topography reconstructions, but this is not compulsory. It is also optional whether to include
23 changes in river routing basins [\(i.e. catchments\)](#) and outlets, which can [either](#) be calculated
24 from the provided topography and land-sea mask data (see Sect. [2.6](#)), [or can be manually](#)
25 [set to follow routing maps, which will be provided on the last deglaciation PMIP Wiki.](#)

26 Groups are free to choose how often to update ice extent and elevation. This could be done at
27 regular intervals (e.g. the [sub](#)-1000 year time slices provided) or at specific times during the
28 deglaciation, as was done in the TraCE-21 ka experiment (Liu et al., 2009). Changes in ice
29 extent can have a large impact on climate through ice albedo changes and feedbacks. We thus
30 recommend that when possible, ice sheets are not updated at times of abrupt regional or global
31 climate change, particularly the events that the working group will focus on, as this could

1 artificially introduce stepped shifts in climate. Groups are also advised to consider that ice sheet
2 associated boundary conditions (ice extent and elevation, land-sea mask, bathymetry) may need
3 to be updated more often at times of rapid ice retreat. The timing and way in which land ice
4 changes are implemented must be documented.

5 Alternative ice sheet reconstructions or simulations can be used to test the sensitivity of climate
6 to this boundary condition. Simulations with coupled ice sheet-climate models are also
7 welcomed. Although these will not form part of the Core, for which ICE-6G_C or GLAC-1D
8 should be used, they will be coordinated as important supplementary *focussed* simulations.

9 **2.5 Ice meltwater**

10 The Core simulation willexperiment protocol is flexible on whether or not to include any
11 prescribed ice melt (i.e. freshwater fluxes) delivered from the ice sheets to the ocean. This may
12 seem controversial given the levels and how to do it. It is recommended to run at least one
13 version of terrestrialthe Core experiment with ice sheet melt and included, since around 110 m
14 of ice-volume equivalent sea-level rise knownis thought to have taken place during this
15 periodmelted 26-9 ka (e.g. Lambeck et al., 2014) and considering the historical importance
16 attached to the influence of [de]glacial freshwater fluxes on climate (e.g. Broecker et al., 1989;
17 Condron and Winsor, 2012; Ganopolski and Rahmstorf, 2001; Liu et al., 2009; Rahmstorf,
18 1995, 1996; Teller et al., 2002; Thornalley et al., 2010; Weaver et al., 2003). However,
19 considering the current uncertainty on exactly when and where ice melt entered the ocean
20 during the last deglaciation (e.g. discussion of MWP1a in Sect. 1.1), thisit is the best wayalso
21 important to ensure that the Core experiment is based on robust geological data. Furthermore,
22 there is an note the ongoing debate over the role ofextent to which catastrophic freshwater
23 fluxes in bringingbrought about abrupt deglacial climate change and; several alternative or
24 complementary mechanisms have been proposed (e.g. Adkins et al., 2005; Álvarez-Solas et al.,
25 2011; Barker et al., 2010, 2015; Broecker, 2003; Hall et al., 2006; Knorr and Lohmann, 2003,
26 2007; Roche et al., 2007; Rogerson et al., 2010; Thiagarajan et al., 2014). In light of this, and
27 because we are keen to see what the climate response to non freshwater forced scenarios will
28 be in the PMIP models, the decision has been made to have no prescribed freshwater fluxes in
29 the Core simulation. This experiment is thus designed to constitute a reference for experiments
30 in which fresh water fluxes will be introduced.

Moreover, a thorough investigation of the extent to which non-freshwater-forced climate evolution matches the geological records has merit in its own right; can abrupt deglacial changes be simulated without ice-meltwater, as has been proposed (e.g. discussion above)?²² To what extent can ‘observed’ patterns be attributed to better constrained forcings, such as atmospheric CO₂ and Earth’s orbit? To complete the investigation, freshwater flux scenarios will be targeted by opt-in focussed simulations that test specific ice-melt hypotheses as well as instances where/when the Core falls short of the ‘observed’ patterns. For example, routing of ice melt computed from GLAC-1D (Sect. 2.4) will be provided as a possible transient boundary condition. It is for all of these reasons that a flexible protocol is required.

Freshwater forcing scenarios consistent with the ice sheet reconstructions and which hence conserve salinity throughout the deglacial experiment are provided in two formats (the ‘melt-’ scenarios described below). In addition, there is the option to run without any ice meltwater (‘no-melt’) to provide a robust reference for simulations that include uncertain meltwater fluxes. Thus, at least one Core simulation should be run using one of the following ice sheet meltwater scenarios:

melt-uniform: a globally uniform freshwater flux (or salinity target) through time, designed to conserve ocean salinity based on changing terrestrial ice mass. Fluxes consistent with the ice sheet reconstructions are provided.

melt-routed: a distributed routing that is consistent with the geographic evolution of the ice sheet reconstructions (GLAC-1D and ICE-6G C; Sect. 2.4) and gives the flux through time at individual meltwater river outlets along the coast. Again, versions of this scenario are provided.

no-melt: no ice meltwater is included in the core; neither a globally integrated ocean salinity target (*melt-uniform*) nor a distributed routing at the coastlines (*melt-routed*) is implemented. This is best implemented as a sensitivity-type experiment to account for model-specificness and meltwater flux uncertainty when also implementing *melt-* scenarios in accompanying versions of the Core simulation.

Multiple Core simulations exploring more than one of these scenarios are welcomed.

1 Data for the *melt-* scenarios will be available from the PMIP last deglaciation Wiki. The data
2 for *melt-uniform* are available at the time of writing (following the respective ice volume
3 changes from ICE-6G_C and GLAC1-D; Fig. 1g), data for *melt-routed* will be made available
4 as they are produced (anticipated by April/May 2016). These *melt-* scenarios represent a ‘best-
5 estimate’ approach to resolving the yet unknown geographically- and temporally-precise
6 freshwater fluxes of the last deglaciation, and they are also consistent with the ice sheet
7 reconstructions employed in the core. As such, they provide robust and justifiable boundary
8 conditions for simulations that will be assessed against palaeoclimate reconstructions.

9 However, participants do not have to use the [recommended] versions of *melt-uniform* or *melt-*
10 *routed* that are consistent with ICE-6G_C and GLAC-1D, and can instead use their own
11 scenarios to explore uncertainty in the ice sheet meltwater flux forcing. This is because the
12 working group aims to use the full suite of PMIP climate models to examine forcing/boundary
13 condition uncertainty (see discussion of model intercomparison project approaches in Sect.
14 1.4). Please note that in some ice melt (including *no-melt*) scenarios, global water budget may
15 not be balanced through time (as is also true for *no-melt*). Therefore, it is advised to also use at
16 least one scenario that falls within geological constraints (such as the ICE-6G_C or GLAC-1D
17 consistent scenarios for *melt-uniform* and *melt-routed*).

18 Regardless of which scenario is employed, it is important that meltwater fluxes are prescribed
19 as time-evolving model boundary conditions; rather than as step-wise adjustments at the same
20 time as the ice sheets are updated, for example. Unless they are intentional conditions of the
21 scenario, there should be no sudden jumps in the freshwater being applied. Furthermore, we
22 invite participants to upload the boundary condition data for other freshwater flux scenarios
23 along with appropriate documentation as/when they become available, and to contribute
24 towards the coordination of *focussed* experiments (see Sect. 3) that will test specific hypotheses
25 associated with model and climate sensitivity to the location, duration and magnitude of
26 freshwater fluxes.

27 **2.6 Topography, bathymetry, coastlines and rivers**

28 Changes in the ice sheets and their glacial eustatic and isostatic influence affected continental
29 topography and ocean bathymetry, which in turn shifted the coordinates of river mouths and
30 the coastal outline throughout the deglaciation. Hence time-varying topographic, bathymetric

1 and land-sea mask fields that match the chosen ice sheet from Sect. 2.4 (i.e. ICE-6G_C or
2 GLAC-1D) should be used; [these are provided within the ice sheet reconstruction datasets](#).

3 Topography should be updated at the same time as the model's ice sheet is updated; this is
4 mainly implicit to implementing the ice sheet reconstruction because the major orographic
5 changes through the deglaciation relate directly to ice sheet evolution. This said, due to glacial
6 isostatic adjustment components in the ice sheet reconstructions, there is evolution in
7 continental topography that is not directly the lowering/heightening of the ice surface, and it is
8 up to individuals whether they incorporate this or mask only the changes in ice sheet orography.

9 Ocean bathymetry will be provided, [but is an optional. When deemed possible, this](#) boundary
10 condition [to vary should be varied](#) through time. [Coastlines, Where differences in the land-sea](#)
11 [mask require extra land to fill up coastal regions, or land to be cut away into ocean as sea level](#)
12 [rises \(see next paragraph on the other hand, coastlines\), the model must be changed accordingly,](#)
13 [because it is important to adequately represent the changing land-sea mask; for example, in](#)
14 [order to include overlying grounded ice.](#)

15 [Following on from this, coastlines](#) will need to be varied according to changes in global sea
16 level (and each model's horizontal grid resolution). It will be left to the discretion of participants
17 to decide how often to update either boundary condition, and when deciding on their frequency
18 it is recommended that groups consider the implications for opening/closing seaways and their
19 effect on ocean circulation and climate. Furthermore, the frequency need not be regular and
20 may instead focus on key 'events' in the marine [gateway] realm. However, whenever possible
21 and foreseeable, groups are encouraged to avoid making stepwise changes to model boundary
22 conditions that would interfere with signals of abrupt climate change; particularly those events
23 that the working group aims to focus on (e.g. Heinrich Event 1, the Bølling Warming, MWP1a,
24 the Younger Dryas etc.) unless the forcing (e.g. opening of a gateway) is assumed to be linked
25 with the event.

26 If groups wish, model river networks can be remapped to be consistent with this and updated
27 on the same timestep as the ice sheet reconstruction, either manually or by the model. However,
28 it is appreciated that the technical challenges associated with such a methodology would be
29 impractical for many. Therefore, following the recommendation of the PMIP3 LGM Working
30 Group (2010) [and Kageyama et al. \(in prep.\)](#), 'river pathways and basins should be at least
31 adjusted so that fresh water is conserved at the Earth's surface and care should be taken that

1 rivers reach the ocean' at every timestep that the bathymetry is adjusted; for example, when sea
2 levels were lower, some river mouths may need to be displaced towards the [new] coastline to
3 make sure they reach the ocean.

4 **2.7 Vegetation, land surface and other forcings**

5 In this section, recommendations are made for last deglaciation vegetation, land surface and
6 aerosol (dust) parameters in the model.

7 There are three recommended options for setting up the Core simulation's experiment's
8 vegetation and land surface parameters, they can either be: (i) computed using a dynamical
9 vegetation model (e.g. coupled to the atmospheric component of the model); (ii) prescribed to
10 match the CMIP5 preindustrial setup (Taylor et al., 2011a, 2011b) with fixed vegetation types
11 and fixed plant physiology (including leaf area index); or (iii) prescribed to match the CMIP5
12 preindustrial setup (Taylor et al., 2011a, 2011b) with fixed vegetation types and interactive
13 plant physiology if running with an enabled carbon cycle. If prescribing vegetation and land
14 surface, i.e. using option (ii) and (iii), groups should be aware that coastal land will be emerged
15 compared to preindustrial because of the increased terrestrial ice volume and associated lower
16 eustatic sea level (with the maximum during the early stages of the Core). Therefore,
17 vegetation/land surface will need to be interpolated onto the emerged land from preindustrial
18 grid cells, for example using nearest neighbour methods.

19 For models with prognostic aerosols, the parameters for dust [forcing] can be computed
20 dynamically. Alternatively, it is recommended that Core simulations fix the associated
21 parameters according to the CMIP5 preindustrial simulation (Taylor et al., 2011a, 2011b),with
22 no temporal variation., with no temporal variation. Examining the influence of different
23 transient aerosol scenarios (for those models that do not include prognostic dust, for example)
24 could constitute a further suite of sensitivity simulations for comparison with the Core

25 It has already been described that for the LGM (i.e. the very start of the Core simulation), groups
26 are recommended to adjust the global freshwater budget by +1 psu to account for the increased
27 [terrestrial] ice volume (Sect. 2.1.1). If salinity is reset at any subsequent point (e.g. to correct
28 for model drifts or to account for ice volume changes), this must be documented.

- 1 There is no last deglaciation protocol for setting up other forcings, transient or fixed in time.
- 2 For all simulations, groups are required to fully document their methods, including experiment
- 3 design and especially when different or with additional components to the setup described here.

4 **3 Coordinating further simulations**

5 As already [alluded to](#) [discussed](#), we are faced with the challenge of designing an experiment that
6 is suitable to be run with a wide range of models, from the more computationally efficient class
7 of intermediate complexity models, to state-of-the-art Earth System Models. One particular
8 difficulty is enabling the most complex and highest resolution climate models to participate in
9 this 12 thousand year long experiment when for some, even the integration to reach the LGM
10 spinup state demands a huge amount of computational resource. There is no easy solution and
11 our approach will be to augment the Core [simulation](#) [simulations](#) with shorter *focussed*
12 simulations that target specific questions, mechanisms and time periods. Whilst the most
13 computationally expensive models (e.g. the latest generation of Earth System Models) may not
14 [feasibly](#) be able to participate in the Core, they will be included in the shorter subset of *focussed*
15 simulations. Similarly, alternative full-deglaciation simulations can be coordinated for the less
16 computationally expensive models in the working group (e.g. low resolution General
17 Circulation Models, and Earth System Models of Intermediate Complexity).

18 One line of investigation relating to meltwater inputs from ice sheets and icebergs is to carry
19 out a suite of sensitivity simulations examining different injection sites. These simulations
20 would help to address some of the uncertainty [that led to the exclusion of](#) [in](#) freshwater [fluxes](#)
21 [from the Core](#) [flux scenarios](#). For example, geochemical evidence suggests that smaller and
22 more localised discharges of freshwater than have traditionally been considered in climate
23 models may have an important influence on ocean circulation (e.g. Hall et al., 2006), implying
24 that precise freshwater fluxes are needed in the models to examine their effect. Certainly, others
25 have shown that the location of injection is a controlling factor on the impact of freshwater
26 delivery to the ocean, not just laterally (e.g. Condron and Winsor, 2012; Smith and Gregory,
27 2009), but also in terms of depth (e.g. Roche et al., 2007).

28 A set of coordinated simulations exploring a range of uncertainty in the freshwater forcing
29 (location, depth, duration, magnitude, and physical characteristics such as temperature and
30 density) would be well suited for the *focussed* experiments, thus building on the [meltwater free](#)

1 Core. Core simulations, which may themselves indicate interesting avenues for investigation;
2 partly the purpose of a flexible meltwater approach.

3 However, freshwater is not the only issue and other *focussed* experiments could include the
4 influence of timing in greenhouse gas records, record, differences in ice sheet reconstructions
5 (e.g. the PMIP3 merged ice sheet from Abe-Ouchi et al., 2015; ICE-6G_C; GLAC-1D) or
6 simulations with [coupled] ice-sheet models, the relative importance of different forcings (e.g.
7 insolation vs. trace gases vs. ice sheet evolution), sensitivity to dust-forcing scenarios, the
8 influence of changes in tidal energy dissipation (Schmittner et al., 2015), event-specific
9 hypothesis testing, and shorter-term variability within the climate system.

10 Based on on-going discussions, it is likely that the first sets of *focussed* simulations will be:

- 11 • Sensitivity and hypothesis-driven, investigating simulations that compare results from
12 uniformly distributed meltwater fluxes to results from river-routed meltwater fluxes to
13 examine the impact of the regional specificity of freshwater forcing upon climate system
14 evolution.
- 15 • Sensitivity simulations that are free from ice meltwater fluxes to provide information
16 on what climate evolution was caused by processes other than freshwater fluxes to the
17 ocean.
- 18 • A hypothesis-driven investigation of the possible mechanisms for preconditioning the
19 glacial ocean for the relatively cool Heinrich Stadial 1 and ensuing catastrophic iceberg
20 discharge (Barker et al., 2015).
- 21 • Sensitivity experiments examining the role of trace gas forcing resolution on climate
22 evolution; for example, smoothing the record provided by Bereiter et al. (2015).

23 We have described the plans for *focussed* simulations to highlight the depth of the working
24 group's aims and to properly contextualise the Core simulations, but the purpose of
25 this manuscript is to outline the model setup for the Core simulation. The experiment. The
26 design for subsequent *focussed* simulations will be described at a later date on the PMIP Last
27 Deglaciation Working Group Wiki
28 (<https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglag:index>) and we welcome
29 contributions to the discussion of what further simulations to coordinate there.

1 **4 Working group phases**

2 The experiment will be split into three phases that are designed to run seamlessly into each
3 other (Fig. 1a). Phase one begins at the LGM (21 ka) and will finish at the abrupt Bølling
4 Warming event, which is where Phase 2 picks up, encompassing the Bølling Warming. Phase
5 3 begins at the start of the Younger Dryas cooling and is currently planned to continue through
6 to the end of the Core [simulation experiment](#) at 9 ka.

7 Perhaps most importantly, this affords near-future milestones for managing the ultimate
8 completion of the long full deglacial simulation across all participant groups. It will provide a
9 timetabled framework for beginning and continuing the longer simulations; for scheduling
10 shorter, event- or challenge-specific transient simulations by more computationally expensive
11 models (see discussion in Sect. 3); and for the analysis and publication of results as the
12 milestones are reached. Another motivation is to ensure that the experiment design for later
13 periods of the last deglaciation is updated according to knowledge gained from simulations of
14 the preceding time period; for example, changes in ocean and climate states, which have
15 previously been shown to have a strong influence on climate trajectories (e.g. Kageyama et al.,
16 2010; Timm and Timmermann, 2007). [This is particularly important for setting up shorter,](#)
17 [event-specific focussed simulations, but it is not planned to be explicitly used to influence the](#)
18 [Core.](#) Splitting the period into phases also provides the opportunity to update model boundary
19 conditions and climate forcing data with cutting edge palaeoclimate reconstructions, as they
20 emerge during the lifespan of the multi-model experiment. However, care will be taken to
21 ensure that these are physically consistent between phases. [–, and these updates will not](#)
22 [compromise the Core simulations described in this manuscript. This is so as not to disadvantage](#)
23 [more computationally efficient models that may have already completed simulating the full 21–](#)
24 [9 ka \(or beyond\) period. Instead, the information will be incorporated into focussed versions of](#)
25 [the last deglaciation simulations; possibly spun-off sub-periods that do not have to start again](#)
26 [at the LGM.](#)

27 Each phase will encompass at least one distinguishable climate event; Heinrich Stadial 1 and
28 Heinrich Event 1 in Phase 1 following on from the LGM; MWP1a, the Bølling Warming and
29 the Antarctic Cold Reversal in Phase 2; and the Younger Dryas cooling in Phase 3 (Fig. 1b).
30 As outlined in Sect. 3, simulations of these shorter events can be coordinated in the *focussed*
31 simulations. This is to engage the higher complexity/resolution models, which are unable to run

1 longer simulations, but can use the wider framework of the working group to provide valuable
2 knowledge on rapid climate changes known to have taken place in the last 21 ka.

3 **5 Summary**

4 The last deglaciation presents a host of exciting opportunities to study the Earth System and in
5 particular, to try to understand a range of abrupt climate changes that occurred over just a few
6 years to centuries within the context of more gradual trends. Numerical climate models provide
7 useful tools to investigate the mechanisms that underpin the events of this well-studied time
8 period, especially now that technological and scientific advances make it possible to run multi-
9 millennium simulations with some of the most complex models. Several recent modelling
10 studies have begun this task, but many questions and untested hypotheses remain. Therefore,
11 under the auspices of the Paleoclimate Modelling Intercomparison Project (PMIP), we have set
12 up an initiative to coordinate efforts to run transient simulations of the last deglaciation, and to
13 facilitate the dissemination of expertise between modellers and those engaged with
14 reconstructing the climate of the last 21 thousand years.

15 The first step has been to design a [single](#), Core [simulationexperiment](#) suitable for a range of
16 PMIP models; from relatively fast and coarse resolution Earth System Models of Intermediate
17 Complexity, to new generations of the more complex and higher resolution General Circulation
18 and Earth System Models. The setup for this Core [simulationexperiment](#), is based on an
19 approach that tries to combine a traditional Model Intercomparison Project method of strictly
20 prescribing boundary conditions across all models, and the philosophy of utilising the breadth
21 of participants to address outstanding uncertainty in the climate forcings, model structure and
22 palaeoclimate reconstructions. Accordingly, we have made recommendations for the
23 initialisation conditions for the simulation and have stated our minimum requirements for the
24 transient experiment design, as summarised in Table 1 and 2, respectively.

25 However, there are some uncertainties that the Core is not designed to deal with directly [or](#)
26 [exhaustively](#); two examples discussed in this manuscript being the [effect of trace gas record](#)
27 [resolution and the](#) influence of ice melt on the oceans and climate, [and the effect of timing in](#)
28 [the trace gas records respectively](#). We know that the Core [simulation](#)[simulations](#) will not tackle
29 all of our questions, and [is](#)[are](#) likely to give rise to others. Therefore, additional *focussed*
30 simulations will also be coordinated on an ad-hoc basis by the working group. Many of these

1 will build on and be centred around the Core; often taking shorter snapshots in time, thus
2 including the most computationally expensive models in the experiment, or presenting twelve-
3 thousand year alternatives to the Core for faster models to contribute. Not all simulations will
4 be suitable for all models, but the aim is that taken as a whole, the experiment can utilise the
5 wide range of PMIP model strengths and hence minimise individual weaknesses.

6 Essentially, the Core [simulationexperiment](#) has been designed to be inclusive, taking into
7 account the best compromise between uncertainties in the geological data and model
8 limitations. The hypothesis-driven *focussed* experiments will go further than the Core to target
9 the questions that remain. It is hoped that this exciting initiative will improve our individual
10 efforts, providing new opportunities to drive the science forwards towards understanding this
11 fascinating time period, specific mechanisms of rapid climate warming, cooling and sea level
12 change, and Earth's climate system more broadly.

13 **Author Contributions**

14 RFI and LJG lead the PMIP Last Deglaciation Working Group, for which AB, MK, DMR and
15 PJV act as the advisory group. RFI, LJG, MK, DMR, PJV and AB collaboratively designed the
16 working group's aims, structure, Core [simulationexperiment](#) and additional experiments in
17 consultation with the wider community. RD, WRP and LT provided the ice sheet
18 reconstructions, plus associated boundary conditions. RFI and LJG collated these and all other
19 boundary condition data for the simulations. RFI and LJG wrote the manuscript and produced
20 the figures with contributions from all authors.

21 **Acknowledgments**

22 RFI is funded by a NERC Independent Research Fellowship [#NE/K008536/1]. Data
23 processing for boundary condition preparation was carried out using the computational facilities
24 of the Palaeo@Leeds modelling group, University of Leeds, UK. All authors would like to
25 thank everyone who has taken the time to discuss the Working Group's aims and experiments
26 with us. We are especially grateful to Jean-Yves Peterschmitt (LSCE, France) for archiving the
27 boundary conditions, Emilie Capron (BAS, UK) for help with the ice core data and Bette Otto-
28 Bliesner (NCAR, USA) for useful comments on an earlier version of this manuscript. [Specific](#)
29 [thanks also go to Anders Carlson, Eric Wolff, Andreas Schmittner, Shawn Marshall and an](#)

1 [anonymous reviewer for valuable comments on the manuscript, and to Jeremy Fyke for editorial](#)
2 [handling.](#)

3 **References**

4 Abe-Ouchi, A., Saito, F., Kageyama, M., Braconnot, P., Harrison, S. P., Lambeck, K., Otto-
5 Bliesner, B. L., Peltier, W. R., Tarasov, L., Peterschmitt, J.-Y. and Takahashi, K.: Ice-sheet
6 configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments, *Geosci Model Dev*
7 *Discuss*, 8(6), 4293–4336, doi:10.5194/gmdd-8-4293-2015, 2015.

8 Adkins, J. F., Ingersoll, A. P. and Pasquero, C.: Rapid climate change and conditional instability
9 of the glacial deep ocean from the thermobaric effect and geothermal heating, *Quat. Sci. Rev.*,
10 24(5–6), 581–594, doi:10.1016/j.quascirev.2004.11.005, 2005.

11 Aharon, P.: Entrainment of meltwaters in hyperpycnal flows during deglaciation superfloods in
12 the Gulf of Mexico, *Earth Planet. Sci. Lett.*, 241(1–2), 260–270,
13 doi:10.1016/j.epsl.2005.10.034, 2006.

14 Ahn, J. and Brook, E. J.: Siple Dome ice reveals two modes of millennial CO₂ change during
15 the last ice age, *Nat. Commun.*, 5, 3723, doi:10.1038/ncomms4723, 2014.

16 Alley, R. B.: The Younger Dryas cold interval as viewed from central Greenland, *Quat. Sci.*
17 *Rev.*, 19(1–5), 213–226, doi:10.1016/S0277-3791(99)00062-1, 2000.

18 Álvarez-Solas, J., Montoya, M., Ritz, C., Ramstein, G., Charbit, S., Dumas, C., Nisancioglu,
19 K., Dokken, T. and Ganopolski, A.: Heinrich event 1: an example of dynamical ice-sheet
20 reaction to oceanic changes, *Clim Past*, 7(4), 1297–1306, doi:10.5194/cp-7-1297-2011, 2011.

21 Annan, J. D. and Hargreaves, J. C.: A new global reconstruction of temperature changes at the
22 Last Glacial Maximum, *Clim Past*, 9(1), 367–376, doi:10.5194/cp-9-367-2013, 2013.

23 Argus, D. F., Peltier, W. R., Drummond, R. and Moore, A. W.: The Antarctica component of
24 postglacial rebound model ICE-6G_C (VM5a) based on GPS positioning, exposure age dating
25 of ice thicknesses, and relative sea level histories, *Geophys. J. Int.*, ggu140,
26 doi:10.1093/gji/ggu140, 2014.

27 Bard, E.: Climate shock- Abrupt changes over millenial time scales, *Phys. Today*, 55, 32–38,
28 2002.

1 Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G. and Rougerie, F.:
2 Deglacial sea-level record from Tahiti corals and the timing of global meltwater discharge,
3 *Nature*, 382(6588), 241–244, doi:10.1038/382241a0, 1996.

4 Bard, E., Rostek, F., Turon, J.-L. and Gendreau, S.: Hydrological Impact of Heinrich Events in
5 the Subtropical Northeast Atlantic, *Science*, 289(5483), 1321–1324,
6 doi:10.1126/science.289.5483.1321, 2000.

7 Bard, E., Hamelin, B. and Delanghe-Sabatier, D.: Deglacial Meltwater Pulse 1B and Younger
8 Dryas Sea Levels Revisited with Boreholes at Tahiti, *Science*, 327(5970), 1235–1237,
9 doi:10.1126/science.1180557, 2010.

10 Barker, S., Knorr, G., Vautravers, M. J., Diz, P. and Skinner, L. C.: Extreme deepening of the
11 Atlantic overturning circulation during deglaciation, *Nat. Geosci.*, 3(8), 567–571,
12 doi:10.1038/ngeo921, 2010.

13 Barker, S., Chen, J., Gong, X., Jonkers, L., Knorr, G. and Thornalley, D.: Icebergs not the
14 trigger for North Atlantic cold events, *Nature*, 520(7547), 333–336, doi:10.1038/nature14330,
15 2015.

16 Bassett, S. E., Milne, G. A., Mitrovica, J. X. and Clark, P. U.: Ice Sheet and Solid Earth
17 Influences on Far-Field Sea-Level Histories, *Science*, 309(5736), 925–928,
18 doi:10.1126/science.1111575, 2005.

19 Bassett, S. E., Milne, G. A., Bentley, M. J. and Huybrechts, P.: Modelling Antarctic sea-level
20 data to explore the possibility of a dominant Antarctic contribution to meltwater pulse IA, *Quat.*
21 *Sci. Rev.*, 26(17–18), 2113–2127, doi:10.1016/j.quascirev.2007.06.011, 2007.

22 Bauer, E. and Ganopolski, A.: Sensitivity simulations with direct shortwave radiative forcing
23 by aeolian dust during glacial cycles, *Clim Past*, 10(4), 1333–1348, doi:10.5194/cp-10-1333-
24 2014, 2014.

25 Bentley, M. J., Fogwill, C. J., Brocq, A. M. L., Hubbard, A. L., Sugden, D. E., Dunai, T. J. and
26 Freeman, S. P. H. T.: Deglacial history of the West Antarctic Ice Sheet in the Weddell Sea
27 embayment: Constraints on past ice volume change, *Geology*, 38(5), 411–414,
28 doi:10.1130/G30754.1, 2010.

1 Bentley, M. J., Ó Cofaigh, C., Anderson, J. B., Conway, H., Davies, B., Graham, A. G. C.,
2 Hillenbrand, C.-D., Hodgson, D. A., Jamieson, S. S. R., Larter, R. D., Mackintosh, A., Smith,
3 J. A., Verleyen, E., Ackert, R. P., Bart, P. J., Berg, S., Brunstein, D., Canals, M., Colhoun, E.
4 A., Crosta, X., Dickens, W. A., Domack, E., Dowdeswell, J. A., Dunbar, R., Ehrmann, W.,
5 Evans, J., Favier, V., Fink, D., Fogwill, C. J., Glasser, N. F., Gohl, K., Golledge, N. R.,
6 Goodwin, I., Gore, D. B., Greenwood, S. L., Hall, B. L., Hall, K., Hedding, D. W., Hein, A. S.,
7 Hocking, E. P., Jakobsson, M., Johnson, J. S., Jomelli, V., Jones, R. S., Klages, J. P.,
8 Kristoffersen, Y., Kuhn, G., Leventer, A., Licht, K., Lilly, K., Lindow, J., Livingstone, S. J.,
9 Massé, G., McGlone, M. S., McKay, R. M., Melles, M., Miura, H., Mulvaney, R., Nel, W.,
10 Nitsche, F. O., O'Brien, P. E., Post, A. L., Roberts, S. J., Saunders, K. M., Selkirk, P. M.,
11 Simms, A. R., Spiegel, C., Stolldorf, T. D., Sugden, D. E., van der Putten, N., van Ommen, T.,
12 Verfaillie, D., Vyverman, W., Wagner, B., White, D. A., Witus, A. E. and Zwart, D.: A
13 community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last
14 Glacial Maximum, *Quat. Sci. Rev.*, 100, 1–9, doi:10.1016/j.quascirev.2014.06.025, 2014.

15 Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H.,
16 Kipfstuhl, S. and Chappellaz, J.: Revision of the EPICA Dome C CO₂ record from 800 to
17 600 kyr before present, *Geophys. Res. Lett.*, 42(2), 2014GL061957,
18 doi:10.1002/2014GL061957, 2015.

19 Berger, A.: Long-Term Variations of Daily Insolation and Quaternary Climatic Changes, *J.*
20 *Atmospheric Sci.*, 35(12), 2362–2367, doi:10.1175/1520-
21 0469(1978)035<2362:LTVODI>2.0.CO;2, 1978.

22 Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quat.*
23 *Sci. Rev.*, 10(4), 297–317, doi:10.1016/0277-3791(91)90033-Q, 1991.

24 Bonelli, S., Charbit, S., Kageyama, M., Woillez, M.-N., Ramstein, G., Dumas, C. and Quiquet,
25 A.: Investigating the evolution of major Northern Hemisphere ice sheets during the last glacial-
26 interglacial cycle, *Clim Past*, 5(3), 329–345, doi:10.5194/cp-5-329-2009, 2009.

27 Boulton, G. S., Dongelmans, P., Punkari, M. and Broadgate, M.: Palaeoglaciology of an ice
28 sheet through a glacial cycle:: the European ice sheet through the Weichselian, *Quat. Sci. Rev.*,
29 20(4), 591–625, doi:10.1016/S0277-3791(00)00160-8, 2001.

1 Braconnot, P., Joussaume, S., de Noblet, N. and Ramstein, G.: Mid-Holocene and Last Glacial
2 Maximum African monsoon changes as simulated within the Paleoclimate Modelling
3 Intercomparison Project, *Glob. Planet. Change*, 26(1–3), 51–66, doi:10.1016/S0921-
4 8181(00)00033-3, 2000.

5 Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J.-Y., Abe-Ouchi,
6 A., Crucifix, M., Driesschaert, E., Fichefet, T., Hewitt, C. D., Kageyama, M., Kitoh, A., Laîné,
7 A., Loutre, M.-F., Marti, O., Merkel, U., Ramstein, G., Valdes, P., Weber, S. L., Yu, Y. and
8 Zhao, Y.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial
9 Maximum – Part 1: experiments and large-scale features, *Clim Past*, 3(2), 261–277,
10 doi:10.5194/cp-3-261-2007, 2007.

11 Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-Ouchi,
12 A., Otto-Bliesner, B. and Zhao, Y.: Evaluation of climate models using palaeoclimatic data,
13 *Nat. Clim. Change*, 2(6), 417–424, doi:10.1038/nclimate1456, 2012.

14 Briggs, R. D., Pollard, D. and Tarasov, L.: A data-constrained large ensemble analysis of
15 Antarctic evolution since the Eemian, *Quat. Sci. Rev.*, 103, 91–115,
16 doi:10.1016/j.quascirev.2014.09.003, 2014.

17 Broecker, W. S.: Does the Trigger for Abrupt Climate Change Reside in the Ocean or in the
18 Atmosphere?, *Science*, 300(5625), 1519–1522, doi:10.1126/science.1083797, 2003.

19 Broecker, W. S., Kennett, J. P., Flower, B. P., Teller, J. T., Trumbore, S., Bonani, G. and Wolfli,
20 W.: Routing of meltwater from the Laurentide Ice Sheet during the Younger Dryas cold
21 episode, *Nature*, 341(6240), 318–321, doi:10.1038/341318a0, 1989.

22 Brovkin, V., Ganopolski, A., Archer, D. and Munhoven, G.: Glacial CO₂ cycle as a succession
23 of key physical and biogeochemical processes, *Clim Past*, 8(1), 251–264, doi:10.5194/cp-8-
24 251-2012, 2012.

25 Buizert, C., Gkinis, V., Severinghaus, J. P., He, F., Lecavalier, B. S., Kindler, P., Leuenberger,
26 M., Carlson, A. E., Vinther, B., Masson-Delmotte, V., White, J. W. C., Liu, Z., Otto-Bliesner,
27 B. and Brook, E. J.: Greenland temperature response to climate forcing during the last
28 deglaciation, *Science*, 345(6201), 1177–1180, doi:10.1126/science.1254961, 2014.

1 Cabioch, G., Banks-Cutler, K. A., Beck, W. J., Burr, G. S., Corrège, T., Lawrence Edwards, R.
2 and Taylor, F. W.: Continuous reef growth during the last 23 cal kyr BP in a tectonically active
3 zone (Vanuatu, SouthWest Pacific), *Quat. Sci. Rev.*, 22(15–17), 1771–1786,
4 doi:10.1016/S0277-3791(03)00170-7, 2003.

5 Caley, T., Roche, D. M. and Renssen, H.: Orbital Asian summer monsoon dynamics revealed
6 using an isotope-enabled global climate model, *Nat. Commun.*, 5, doi:10.1038/ncomms6371,
7 2014.

8 Carlson, A. E.: Geochemical constraints on the Laurentide Ice Sheet contribution to Meltwater
9 Pulse 1A, *Quat. Sci. Rev.*, 28(17–18), 1625–1630, doi:10.1016/j.quascirev.2009.02.011, 2009.

10 Clark, P. U. and Mix, A. C.: Ice sheets and sea level of the Last Glacial Maximum, *Quat. Sci.*
11 *Rev.*, 21(1–3), 1–7, doi:10.1016/S0277-3791(01)00118-4, 2002.

12 Clark, P. U. and Tarasov, L.: Closing the sea level budget at the Last Glacial Maximum, *Proc.*
13 *Natl. Acad. Sci.*, 111(45), 15861–15862, doi:10.1073/pnas.1418970111, 2014.

14 Clark, P. U., Alley, R. B., Keigwin, L. D., Licciardi, J. M., Johnsen, S. J. and Wang, H.: Origin
15 of the first global meltwater pulse following the Last Glacial Maximum, *Paleoceanography*,
16 11(5), 563–577, doi:10.1029/96PA01419, 1996.

17 Clark, P. U., Mitrovica, J. X., Milne, G. A. and Tamisiea, M. E.: Sea-Level Fingerprinting as a
18 Direct Test for the Source of Global Meltwater Pulse IA, *Science*, 295(5564), 2438–2441,
19 doi:10.1126/science.1068797, 2002.

20 Clark, P. U., McCabe, A. M., Mix, A. C. and Weaver, A. J.: Rapid Rise of Sea Level 19,000
21 Years Ago and Its Global Implications, *Science*, 304(5674), 1141–1144,
22 doi:10.1126/science.1094449, 2004.

23 Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J.
24 X., Hostetler, S. W. and McCabe, A. M.: The Last Glacial Maximum, *Science*, 325(5941), 710–
25 714, doi:10.1126/science.1172873, 2009.

26 Clark, P. U., Shakun, J. D., Baker, P. A., Bartlein, P. J., Brewer, S., Brook, E., Carlson, A. E.,
27 Cheng, H., Kaufman, D. S., Liu, Z., Marchitto, T. M., Mix, A. C., Morrill, C., Otto-Bliesner,
28 B. L., Pahnke, K., Russell, J. M., Whitlock, C., Adkins, J. F., Blois, J. L., Clark, J., Colman, S.
29 M., Curry, W. B., Flower, B. P., He, F., Johnson, T. C., Lynch-Stieglitz, J., Markgraf, V.,

1 McManus, J., Mitrovica, J. X., Moreno, P. I. and Williams, J. W.: Global climate evolution
2 during the last deglaciation, Proc. Natl. Acad. Sci., 109(19), E1134–E1142,
3 doi:10.1073/pnas.1116619109, 2012.

4 Condron, A. and Winsor, P.: Meltwater routing and the Younger Dryas, Proc. Natl. Acad. Sci.,
5 109(49), 19928–19933, doi:10.1073/pnas.1207381109, 2012.

6 Cuffey, K. M. and Clow, G. D.: Temperature, accumulation, and ice sheet elevation in central
7 Greenland through the last deglacial transition, J. Geophys. Res., 102(C12), 26383–26,396,
8 doi:10.1029/96JC03981, 1997.

9 Cutler, K. B., Edwards, R. L., Taylor, F. W., Cheng, H., Adkins, J., Gallup, C. D., Cutler, P.
10 M., Burr, G. S. and Bloom, A. L.: Rapid sea-level fall and deep-ocean temperature change since
11 the last interglacial period, Earth Planet. Sci. Lett., 206(3–4), 253–271, doi:10.1016/S0012-
12 821X(02)01107-X, 2003.

13 De Deckker, P. and Yokoyama, Y.: Micropalaeontological evidence for Late Quaternary sea-
14 level changes in Bonaparte Gulf, Australia, Glob. Planet. Change, 66(1–2), 85–92,
15 doi:10.1016/j.gloplacha.2008.03.012, 2009.

16 deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L. and Yarusinsky,
17 M.: Abrupt onset and termination of the African Humid Period:: rapid climate responses to
18 gradual insolation forcing, Quat. Sci. Rev., 19(1–5), 347–361, doi:10.1016/S0277-
19 3791(99)00081-5, 2000.

20 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson, G.
21 M., Okuno, J. 'ichi and Yokoyama, Y.: Ice-sheet collapse and sea-level rise at the Bolling
22 warming 14,600 years ago, Nature, 483(7391), 559–564, doi:10.1038/nature10902, 2012.

23 Dyke, A. S.: An outline of North American deglaciation with emphasis on central and northern
24 Canada, in Quaternary Glaciations-Extent and Chronology - Part II: North America, vol.
25 Volume 2, Part 2, pp. 373–424, Elsevier. [online] Available from:
26 https://www.lakeheadu.ca/sites/default/files/uploads/53/outlines/2014-15/NECU5311/Dyke_2004_DeglaciationOutline.pdf (Accessed 20 October 2015), 2004.

1 Dyke, A. S., Andrews, J. T., Clark, P. U., England, J. H., Miller, G. H., Shaw, J. and Veillette,
2 J. J.: The Laurentide and Innuitian ice sheets during the Last Glacial Maximum, *Quat. Sci. Rev.*,
3 21(1–3), 9–31, doi:10.1016/S0277-3791(01)00095-6, 2002.

4 Edwards, R. L., Beck, J. W., Burr, G. S., Donahue, D. J., Chappell, J. M. A., Bloom, A. L.,
5 Druffel, E. R. M. and Taylor, F. W.: A Large Drop in Atmospheric $^{14}\text{C}/^{12}\text{C}$ and Reduced
6 Melting in the Younger Dryas, Documented with ^{230}Th Ages of Corals, *Science*, 260(5110),
7 962–968, doi:10.1126/science.260.5110.962, 1993.

8 EPICA Community Members: Eight glacial cycles from an Antarctic ice core, *Nature*,
9 429(6992), 623–628, doi:10.1038/nature02599, 2004.

10 Fairbanks, R. G.: A 17,000-year glacio-eustatic sea level record: influence of glacial melting
11 rates on the Younger Dryas event and deep-ocean circulation, *Nature*, 342(6250), 637–642,
12 doi:10.1038/342637a0, 1989.

13 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P., Driouech,
14 F., Emori, S., Eyring, V., Forest, C., Glecker, P., Guilyardi, E., Jackob, C., Kattsov, V., Reason,
15 C. and Rummukainen, M.: Evaluation of Climate Models, in *Climate Change 2013: The*
16 *Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of*
17 *the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G. K. Plattner,
18 M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, pp. 741–
19 866, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
20 [online] Available from: http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter09_FINAL.pdf (Accessed 20 October 2015), 2013.

22 Ganopolski, A. and Calov, R.: The role of orbital forcing, carbon dioxide and regolith in 100
23 kyr glacial cycles, *Clim Past*, 7(4), 1415–1425, doi:10.5194/cp-7-1415-2011, 2011.

24 Ganopolski, A. and Rahmstorf, S.: Rapid changes of glacial climate simulated in a coupled
25 climate model, *Nature*, 409(6817), 153–158, doi:10.1038/35051500, 2001.

26 Ganopolski, A., Calov, R. and Claussen, M.: Simulation of the last glacial cycle with a coupled
27 climate ice-sheet model of intermediate complexity, *Clim Past*, 6(2), 229–244, doi:10.5194/cp-
28 6-229-2010, 2010.

1 García, J. L., Kaplan, M. R., Hall, B. L., Schaefer, J. M., Vega, R. M., Schwartz, R. and Finkel,
2 R.: Glacier expansion in southern Patagonia throughout the Antarctic cold reversal, *Geology*,
3 40(9), 859–862, doi:10.1130/G33164.1, 2012.

4 Gherardi, J.-M., Labeyrie, L., McManus, J. F., Francois, R., Skinner, L. C. and Cortijo, E.:
5 Evidence from the Northeastern Atlantic basin for variability in the rate of the meridional
6 overturning circulation through the last deglaciation, *Earth Planet. Sci. Lett.*, 240(3–4), 710–
7 723, doi:10.1016/j.epsl.2005.09.061, 2005.

8 Golledge, N. R., Fogwill, C. J., Mackintosh, A. N. and Buckley, K. M.: Dynamics of the last
9 glacial maximum Antarctic ice-sheet and its response to ocean forcing, *Proc. Natl. Acad. Sci.*,
10 109(40), 16052–16056, doi:10.1073/pnas.1205385109, 2012.

11 Golledge, N. R., Levy, R. H., McKay, R. M., Fogwill, C. J., White, D. A., Graham, A. G. C.,
12 Smith, J. A., Hillenbrand, C.-D., Licht, K. J., Denton, G. H., Ackert Jr., R. P., Maas, S. M. and
13 Hall, B. L.: Glaciology and geological signature of the Last Glacial Maximum Antarctic ice
14 sheet, *Quat. Sci. Rev.*, 78, 225–247, doi:10.1016/j.quascirev.2013.08.011, 2013.

15 Golledge, N. R., Menziel, L., Carter, L., Fogwill, C. J., England, M. H., Cortese, G. and Levy,
16 R. H.: Antarctic contribution to meltwater pulse 1A from reduced Southern Ocean overturning,
17 *Nat. Commun.*, 5, doi:10.1038/ncomms6107, 2014.

18 Goujon, C., Barnola, J.-M. and Ritz, C.: Modeling the densification of polar firn including heat
19 diffusion: Application to close-off characteristics and gas isotopic fractionation for Antarctica
20 and Greenland sites, *J. Geophys. Res. Atmospheres*, 108(D24), 4792,
21 doi:10.1029/2002JD003319, 2003.

22 Gregoire, L. J., Payne, A. J. and Valdes, P. J.: Deglacial rapid sea level rises caused by ice-
23 sheet saddle collapses, *Nature*, 487(7406), 219–222, doi:10.1038/nature11257, 2012.

24 Gregoire, L. J., Valdes, P. J. and Payne, A. J.: The relative contribution of orbital forcing and
25 greenhouse gases to the North American deglaciation, *Geophys. Res. Lett.*, 42(22),
26 2015GL066005, doi:10.1002/2015GL066005, 2015.

27 Gyllencreutz, R., Mangerud, J., Svendsen, J.-I. and Lohne, Ø.: DATED—a GIS-based
28 reconstruction and dating database of the Eurasian deglaciation, *Appl. Quat. Res. Cent. Part*
29 *Glaciat. Terrain Geol. Surv. Finl. Spec. Pap.*, 46, 113–120, 2007.

1 Hall, I. R., Moran, S. B., Zahn, R., Knutz, P. C., Shen, C.-C. and Edwards, R. L.: Accelerated
2 drawdown of meridional overturning in the late-glacial Atlantic triggered by transient pre-H
3 event freshwater perturbation, *Geophys. Res. Lett.*, 33(16), L16616,
4 doi:10.1029/2006GL026239, 2006.

5 Hanebuth, T., Stattegger, K. and Grootes, P. M.: Rapid Flooding of the Sunda Shelf: A Late-
6 Glacial Sea-Level Record, *Science*, 288(5468), 1033–1035,
7 doi:10.1126/science.288.5468.1033, 2000.

8 Hanebuth, T. J. J., Stattegger, K. and Bojanowski, A.: Termination of the Last Glacial
9 Maximum sea-level lowstand: The Sunda-Shelf data revisited, *Glob. Planet. Change*, 66(1–2),
10 76–84, doi:10.1016/j.gloplacha.2008.03.011, 2009.

11 Haywood, A. M., Dowsett, H. J., Otto-Bliesner, B., Chandler, M. A., Dolan, A. M., Hill, D. J.,
12 Lunt, D. J., Robinson, M. M., Rosenbloom, N., Salzmann, U. and Sohl, L. E.: Pliocene Model
13 Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment
14 1), *Geosci. Model Dev.*, 3(1), 227–242, doi:10.5194/gmd-3-227-2010, 2010.

15 Heinemann, M., Timmermann, A., Elison Timm, O., Saito, F. and Abe-Ouchi, A.: Deglacial
16 ice sheet meltdown: orbital pacemaking and CO₂ effects, *Clim Past*, 10(4), 1567–1579,
17 doi:10.5194/cp-10-1567-2014, 2014.

18 Heiri, O., Cremer, H., Engels, S., Hoek, W. Z., Peeters, W. and Lotter, A. F.: Lateglacial
19 summer temperatures in the Northwest European lowlands: a chironomid record from
20 Hijkemeer, the Netherlands, *Quat. Sci. Rev.*, 26(19–21), 2420–2437,
21 doi:10.1016/j.quascirev.2007.06.017, 2007.

22 Hemming, S. R.: Heinrich events: Massive late Pleistocene detritus layers of the North Atlantic
23 and their global climate imprint, *Rev. Geophys.*, 42(1), RG1005, doi:10.1029/2003RG000128,
24 2004.

25 Hughes, A. L. C., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J. and Svendsen, J. I.: The last
26 Eurasian ice sheets – a chronological database and time-slice reconstruction, DATED-1,
27 *Boreas*, doi:10.1111/bor.12142, 2015.

28 Joussaume, S., Taylor, K. E., Braconnot, P., Mitchell, J. F. B., Kutzbach, J. E., Harrison, S. P.,
29 Prentice, I. C., Broccoli, A. J., Abe-Ouchi, A., Bartlein, P. J., Bonfils, C., Dong, B., Guiot, J.,

1 Herterich, K., Hewitt, C. D., Jolly, D., Kim, J. W., Kislov, A., Kitoh, A., Loutre, M. F., Masson,
2 V., McAvaney, B., McFarlane, N., de Noblet, N., Peltier, W. R., Peterschmitt, J. Y., Pollard,
3 D., Rind, D., Royer, J. F., Schlesinger, M. E., Syktus, J., Thompson, S., Valdes, P., Vettoretti,
4 G., Webb, R. S. and Wyputta, U.: Monsoon changes for 6000 years ago: Results of 18
5 simulations from the Paleoclimate Modeling Intercomparison Project (PMIP), *Geophys. Res.*
6 Lett.

7 Lett., 26(7), 859–862, doi:10.1029/1999GL900126, 1999.

7 Jouzel, J., Vaikmae, R., Petit, J. R., Martin, M., Duclos, Y., Stievenard, M., Loria, C., Toots,
8 M., Mélières, M. A., Burckle, L. H., Barkov, N. I. and Kotlyakov, V. M.: The two-step shape
9 and timing of the last deglaciation in Antarctica, *Clim. Dyn.*, 11(3), 151–161,
10 doi:10.1007/BF00223498, 1995.

11 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster,
12 B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S., Leuenberger,
13 M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A.,
14 Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J. P., Stenni, B.,
15 Stocker, T. F., Tison, J. L., Werner, M. and Wolff, E. W.: Orbital and Millennial Antarctic
16 Climate Variability over the Past 800,000 Years, *Science*, 317(5839), 793–796,
17 doi:10.1126/science.1141038, 2007.

18 Kageyama, M., Braconnot, P., Peterchmitt, J.-Y., Harrison, S. P., Jungclaus, J. H., Otto-
19 Bliesner, B., Brierley, C., Crucifix, M., Haywood, A. M., Ivanovic, R. F., Lunt, D. J., Phipps,
20 S. J., Roche, D. M., Schmidt, G. A., Zhang, Q. and Zhou, Y.: Overview of the PMIP4-CMIP6
21 experiments, *Geosci. Model Dev.*, in prep.

22 Kageyama, M., Laîné, A., Abe-Ouchi, A., Braconnot, P., Cortijo, E., Crucifix, M., de Vernal,
23 A., Guiot, J., Hewitt, C. D., Kitoh, A., Kucera, M., Marti, O., Ohgaito, R., Otto-Bliesner, B.,
24 Peltier, W. R., Rosell-Melé, A., Vettoretti, G., Weber, S. L. and Yu, Y.: Last Glacial Maximum
25 temperatures over the North Atlantic, Europe and western Siberia: a comparison between PMIP
26 models, MARGO sea–surface temperatures and pollen-based reconstructions, *Quat. Sci. Rev.*,
27 25(17–18), 2082–2102, doi:10.1016/j.quascirev.2006.02.010, 2006.

28 Kageyama, M., Paul, A., Roche, D. M. and Van Meerbeeck, C. J.: Modelling glacial climatic
29 millennial-scale variability related to changes in the Atlantic meridional overturning

1 circulation: a review, *Quat. Sci. Rev.*, 29(21–22), 2931–2956,
2 doi:10.1016/j.quascirev.2010.05.029, 2010.

3 Kageyama, M., Merkel, U., Otto-Bliesner, B., Prange, M., Abe-Ouchi, A., Lohmann, G.,
4 Ohgaito, R., Roche, D. M., Singarayer, J., Swingedouw, D. and X Zhang: Climatic impacts of
5 fresh water hosing under Last Glacial Maximum conditions: a multi-model study, *Clim Past*,
6 9(2), 935–953, doi:10.5194/cp-9-935-2013, 2013.

7 Kaplan, M. R., Strelin, J. A., Schaefer, J. M., Denton, G. H., Finkel, R. C., Schwartz, R.,
8 Putnam, A. E., Vandergoes, M. J., Goehring, B. M. and Travis, S. G.: In-situ cosmogenic ^{10}Be
9 production rate at Lago Argentino, Patagonia: Implications for late-glacial climate chronology,
10 *Earth Planet. Sci. Lett.*, 309(1–2), 21–32, doi:10.1016/j.epsl.2011.06.018, 2011.

11 Keigwin, L. D. and Boyle, E. A.: Did North Atlantic overturning halt 17,000 years ago?,
12 *Paleoceanography*, 23(1), PA1101, doi:10.1029/2007PA001500, 2008.

13 Keigwin, L. D., Jones, G. A., Lehman, S. J. and Boyle, E. A.: Deglacial meltwater discharge,
14 North Atlantic Deep Circulation, and abrupt climate change, *J. Geophys. Res. Oceans*, 96(C9),
15 16811–16826, doi:10.1029/91JC01624, 1991.

16 Kindler, P., Guillevic, M., Baumgartner, M., Schwander, J., Landais, A. and Leuenberger, M.:
17 Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Clim Past*, 10(2),
18 887–902, doi:10.5194/cp-10-887-2014, 2014.

19 Knorr, G. and Lohmann, G.: Southern Ocean origin for the resumption of Atlantic thermohaline
20 circulation during deglaciation, *Nature*, 424(6948), 532–536, doi:10.1038/nature01855, 2003.

21 Knorr, G. and Lohmann, G.: Rapid transitions in the Atlantic thermohaline circulation triggered
22 by global warming and meltwater during the last deglaciation, *Geochem. Geophys. Geosystems*, 8(12), Q12006, doi:10.1029/2007GC001604, 2007.

24 Kohfeld, K. . and Harrison, S. .: How well can we simulate past climates? Evaluating the models
25 using global palaeoenvironmental datasets, *Quat. Sci. Rev.*, 19(1–5), 321–346,
26 doi:10.1016/S0277-3791(99)00068-2, 2000.

27 Lambeck, K., Rouby, H., Purcell, A., Sun, Y. and Sambridge, M.: Sea level and global ice
28 volumes from the Last Glacial Maximum to the Holocene, *Proc. Natl. Acad. Sci.*,
29 doi:10.1073/pnas.1411762111, 2014.

1 Lea, D. W., Pak, D. K., Peterson, L. C. and Hughen, K. A.: Synchronicity of Tropical and High-
2 Latitude Atlantic Temperatures over the Last Glacial Termination, *Science*, 301(5638), 1361–
3 1364, doi:10.1126/science.1088470, 2003.

4 Licht, K. J.: The Ross Sea's contribution to eustatic sea level during meltwater pulse 1A,
5 *Sediment. Geol.*, 165(3–4), 343–353, doi:10.1016/j.sedgeo.2003.11.020, 2004.

6 Liu, Z., Otto-Bliesner, B. L., He, F., Brady, E. C., Tomas, R., Clark, P. U., Carlson, A. E.,
7 Lynch-Stieglitz, J., Curry, W., Brook, E., Erickson, D., Jacob, R., Kutzbach, J. and Cheng, J.:
8 Transient Simulation of Last Deglaciation with a New Mechanism for Bølling-Allerød
9 Warming, *Science*, 325(5938), 310–314, doi:10.1126/science.1171041, 2009.

10 Liu, Z., Carlson, A. E., He, F., Brady, E. C., Otto-Bliesner, B. L., Briegleb, B. P., Wehrenberg,
11 M., Clark, P. U., Wu, S., Cheng, J., Zhang, J., Noone, D. and Zhu, J.: Younger Dryas cooling
12 and the Greenland climate response to CO₂, *Proc. Natl. Acad. Sci.*,
13 doi:10.1073/pnas.1202183109, 2012.

14 Liu, Z., Wen, X., Brady, E. C., Otto-Bliesner, B., Yu, G., Lu, H., Cheng, H., Wang, Y., Zheng,
15 W., Ding, Y., Edwards, R. L., Cheng, J., Liu, W. and Yang, H.: Chinese cave records and the
16 East Asia Summer Monsoon, *Quat. Sci. Rev.*, 83, 115–128,
17 doi:10.1016/j.quascirev.2013.10.021, 2014.

18 Louergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola,
19 J.-M., Raynaud, D., Stocker, T. F. and Chappellaz, J.: Orbital and millennial-scale features of
20 atmospheric CH₄ over the past 800,000 years, *Nature*, 453(7193), 383–386,
21 doi:10.1038/nature06950, 2008.

22 Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud,
23 D., Jouzel, J., Fischer, H., Kawamura, K. and Stocker, T. F.: High-resolution carbon dioxide
24 concentration record 650,000–800,000 years before present, *Nature*, 453(7193), 379–382,
25 doi:10.1038/nature06949, 2008.

26 MacFarling Meure, C., Etheridge, D., Trudinger, C., Steele, P., Langenfelds, R., van Ommen,
27 T., Smith, A. and Elkins, J.: Law Dome CO₂, CH₄ and N₂O ice core records extended to 2000
28 years BP, *Geophys. Res. Lett.*, 33(14), L14810, doi:10.1029/2006GL026152, 2006.

1 Mackintosh, A., Golledge, N., Domack, E., Dunbar, R., Leventer, A., White, D., Pollard, D.,
2 DeConto, R., Fink, D., Zwart, D., Gore, D. and Lavoie, C.: Retreat of the East Antarctic ice
3 sheet during the last glacial termination, *Nat. Geosci.*, 4(3), 195–202, doi:10.1038/ngeo1061,
4 2011.

5 Mackintosh, A. N., Verleyen, E., O'Brien, P. E., White, D. A., Jones, R. S., McKay, R., Dunbar,
6 R., Gore, D. B., Fink, D., Post, A. L., Miura, H., Leventer, A., Goodwin, I., Hodgson, D. A.,
7 Lilly, K., Crosta, X., Golledge, N. R., Wagner, B., Berg, S., van Ommen, T., Zwart, D.,
8 Roberts, S. J., Vyverman, W. and Masse, G.: Retreat history of the East Antarctic Ice Sheet
9 since the Last Glacial Maximum, *Quat. Sci. Rev.*, 100, 10–30,
10 doi:10.1016/j.quascirev.2013.07.024, 2014.

11 Marcott, S. A., Bauska, T. K., Buizert, C., Steig, E. J., Rosen, J. L., Cuffey, K. M., Fudge, T.
12 J., Severinghaus, J. P., Ahn, J., Kalk, M. L., McConnell, J. R., Sowers, T., Taylor, K. C., White,
13 J. W. C. and Brook, E. J.: Centennial-scale changes in the global carbon cycle during the last
14 deglaciation, *Nature*, 514(7524), 616–619, doi:10.1038/nature13799, 2014.

15 Marshall, S. J. and Clarke, G. K. C.: Modeling North American Freshwater Runoff through the
16 Last Glacial Cycle, *Quat. Res.*, 52(3), 300–315, doi:10.1006/qres.1999.2079, 1999.

17 Martrat, B., Grimalt, J. O., Lopez-Martinez, C., Cacho, I., Sierro, F. J., Flores, J. A., Zahn, R.,
18 Canals, M., Curtis, J. H. and Hodell, D. A.: Abrupt Temperature Changes in the Western
19 Mediterranean over the Past 250,000 Years, *Science*, 306(5702), 1762–1765,
20 doi:10.1126/science.1101706, 2004.

21 Martrat, B., Grimalt, J. O., Shackleton, N. J., Abreu, L. de, Hutterli, M. A. and Stocker, T. F.:
22 Four Climate Cycles of Recurring Deep and Surface Water Destabilizations on the Iberian
23 Margin, *Science*, 317(5837), 502–507, doi:10.1126/science.1139994, 2007.

24 Masson-Delmotte, V., Kageyama, M., Braconnot, P., Charbit, S., Krinner, G., Ritz, C.,
25 Guilyardi, E., Jouzel, J., Abe-Ouchi, A., Crucifix, M., Gladstone, R. M., Hewitt, C. D., Kitoh,
26 A., LeGrande, A. N., Marti, O., Merkel, U., Motoi, T., Ohgaito, R., Otto-Bliesner, B., Peltier,
27 W. R., Ross, I., Valdes, P. J., Vettoretti, G., Weber, S. L., Wolk, F. and Yu, Y.: Past and future
28 polar amplification of climate change: climate model intercomparisons and ice-core constraints,
29 *Clim. Dyn.*, 26(5), 513–529, doi:10.1007/s00382-005-0081-9, 2006.

1 Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolsk, A., González Rouco, J.
2 F., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T., Osborn, T., Otto-Bliesner, B., Quinn,
3 T., Ramesh, R., Rojas, M., Shao, X. and Timmermann, A.: Information from Paleoclimate
4 Archives, in Climate Change 2013: The Physical Science Basis. Contribution of Working
5 Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change,
6 edited by T. F. Stocker, D. Qin, G. K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels,
7 Y. Xia, V. Bex, and P. M. Midgley, pp. 383–464, Cambridge University Press, Cambridge,
8 United Kingdom and New York, NY, USA. [online] Available from:
9 http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter05_FINAL.pdf
10 (Accessed 20 October 2015), 2013.

11 McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L. D. and Brown-Leger, S.: Collapse
12 and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes,
13 *Nature*, 428(6985), 834–837, doi:10.1038/nature02494, 2004.

14 Men viel, L., Timmermann, A., Timm, O. E. and Mouchet, A.: Climate and biogeochemical
15 response to a rapid melting of the West Antarctic Ice Sheet during interglacials and implications
16 for future climate, *Paleoceanography*, 25(4), PA4231, doi:10.1029/2009PA001892, 2010.

17 Men viel, L., Timmermann, A., Timm, O. E. and Mouchet, A.: Deconstructing the Last Glacial
18 termination: the role of millennial and orbital-scale forcings, *Quat. Sci. Rev.*, 30(9–10), 1155–
19 1172, doi:10.1016/j.quascirev.2011.02.005, 2011.

20 Mix, A. C., Bard, E. and Schneider, R.: Environmental processes of the ice age: land, oceans,
21 glaciers (EPILOG), *Quat. Sci. Rev.*, 20(4), 627–657, doi:10.1016/S0277-3791(00)00145-1,
22 2001.

23 Monnin, E., Steig, E. J., Siegenthaler, U., Kawamura, K., Schwander, J., Stauffer, B., Stocker,
24 T. F., Morse, D. L., Barnola, J.-M., Bellier, B., Raynaud, D. and Fischer, H.: Evidence for
25 substantial accumulation rate variability in Antarctica during the Holocene, through
26 synchronization of CO₂ in the Taylor Dome, Dome C and DML ice cores, *Earth Planet. Sci.*
27 *Lett.*, 224(1–2), 45–54, doi:10.1016/j.epsl.2004.05.007, 2004.

28 Otto-Bliesner, B. L., Schneider, R., Brady, E. C., Kucera, M., Abe-Ouchi, A., Bard, E.,
29 Braconnot, P., Crucifix, M., Hewitt, C. D., Kageyama, M., Marti, O., Paul, A., Rosell-Melé,
30 A., Waelbroeck, C., Weber, S. L., Weinelt, M. and Yu, Y.: A comparison of PMIP2 model

1 simulations and the MARGO proxy reconstruction for tropical sea surface temperatures at last
2 glacial maximum, *Clim. Dyn.*, 32(6), 799–815, doi:10.1007/s00382-008-0509-0, 2009.

3 Otto-Bliesner, B. L., Russell, J. M., Clark, P. U., Liu, Z., Overpeck, J. T., Konecky, B.,
4 deMenocal, P., Nicholson, S. E., He, F. and Lu, Z.: Coherent changes of southeastern equatorial
5 and northern African rainfall during the last deglaciation, *Science*, 346(6214), 1223–1227,
6 doi:10.1126/science.1259531, 2014.

7 Peltier, W. R.: Global glacial isostasy and the surface of the Ice-Age Earth: The ICE-5G (VM2)
8 model and GRACE, *Annu. Rev. Earth Planet. Sci.*, 32(1), 111–149,
9 doi:10.1146/annurev.earth.32.082503.144359, 2004.

10 Peltier, W. R.: On the hemispheric origins of meltwater pulse 1a, *Quat. Sci. Rev.*, 24(14–15),
11 1655–1671, doi:10.1016/j.quascirev.2004.06.023, 2005.

12 Peltier, W. R. and Fairbanks, R. G.: Global glacial ice volume and Last Glacial Maximum
13 duration from an extended Barbados sea level record, *Quat. Sci. Rev.*, 25(23–24), 3322–3337,
14 doi:10.1016/j.quascirev.2006.04.010, 2006.

15 Peltier, W. R. and Vettoretti, G.: Dansgaard-Oeschger oscillations predicted in a comprehensive
16 model of glacial climate: A “kicked” salt oscillator in the Atlantic, *Geophys. Res. Lett.*, 41(20),
17 2014GL061413, doi:10.1002/2014GL061413, 2014.

18 Peltier, W. R., Argus, D. F. and Drummond, R.: Space geodesy constrains ice age terminal
19 deglaciation: The global ICE-6G_C (VM5a) model, *J. Geophys. Res. Solid Earth*,
20 2014JB011176, doi:10.1002/2014JB011176, 2015.

21 Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J.-M., Basile, I., Bender, M.,
22 Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M.,
23 Lipenkov, V. Y., Lorius, C., PÉpin, L., Ritz, C., Saltzman, E. and Stievenard, M.: Climate and
24 atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*,
25 399(6735), 429–436, doi:10.1038/20859, 1999.

26 Philippon, G., Ramstein, G., Charbit, S., Kageyama, M., Ritz, C. and Dumas, C.: Evolution of
27 the Antarctic ice sheet throughout the last deglaciation: A study with a new coupled climate—
28 north and south hemisphere ice sheet model, *Earth Planet. Sci. Lett.*, 248(3–4), 750–758,
29 doi:10.1016/j.epsl.2006.06.017, 2006.

1 PMIP Last Deglaciation Working Group: PMIP3 Last Deglaciation experiment design, [online]
2 Available from: <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglac:index> (Accessed 9
3 April 2015), 2015.

4 PMIP LGM Working Group: PMIP3-CMIP5 Last Glacial Maximum experiment design,
5 [online] Available from: <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:design:21k:final>
6 (Accessed 9 April 2015), 2010.

7 PMIP LGM Working Group: PMIP-CMIP6 Last Glacial Maximum experiment design, [online]
8 Available from: <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:cmip6:design:21k:index>
9 (Accessed 9 April 2015), 2015.

10 PMIP website: Paleoclimate Model Intercomparison Project, [online] Available from:
11 <http://pmip.lsce.ipsl.fr/> (Accessed 13 November 2014), 2007.

12 Putnam, A. E., Denton, G. H., Schaefer, J. M., Barrell, D. J. A., Andersen, B. G., Finkel, R. C.,
13 Schwartz, R., Doughty, A. M., Kaplan, M. R. and Schlüchter, C.: Glacier advance in southern
14 middle-latitudes during the Antarctic Cold Reversal, *Nat. Geosci.*, 3(10), 700–704,
15 doi:10.1038/ngeo962, 2010.

16 Rahmstorf, S.: Bifurcations of the Atlantic thermohaline circulation in response to changes in
17 the hydrological cycle, *Nature*, 378(6553), 145–149, doi:10.1038/378145a0, 1995.

18 Rahmstorf, S.: On the freshwater forcing and transport of the Atlantic thermohaline circulation,
19 *Clim. Dyn.*, 12(12), 799–811, doi:10.1007/s003820050144, 1996.

20 Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen,
21 H. B., Siggaard-Andersen, M.-L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M.,
22 Röhlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E. and Ruth, U.: A new Greenland
23 ice core chronology for the last glacial termination, *J. Geophys. Res. Atmospheres*, 111(D6),
24 D06102, doi:10.1029/2005JD006079, 2006.

25 Roberts, N. L., Piotrowski, A. M., McManus, J. F. and Keigwin, L. D.: Synchronous Deglacial
26 Overturning and Water Mass Source Changes, *Science*, 327(5961), 75–78,
27 doi:10.1126/science.1178068, 2010.

1 Roche, D. M., Renssen, H., Weber, S. L. and Goosse, H.: Could meltwater pulses have been
2 sneaked unnoticed into the deep ocean during the last glacial?, *Geophys. Res. Lett.*, 34(24),
3 n/a–n/a, doi:10.1029/2007GL032064, 2007.

4 Roche, D. M., Renssen, H., Paillard, D. and Levavasseur, G.: Deciphering the spatio-temporal
5 complexity of climate change of the last deglaciation: a model analysis, *Clim Past*, 7(2), 591–
6 602, doi:10.5194/cp-7-591-2011, 2011.

7 Rogerson, M., Colmenero-Hidalgo, E., Levine, R. C., Rohling, E. J., Voelker, A. H. L., Bigg,
8 G. R., Schönfeld, J., Cacho, I., Sierro, F. J., Löwemark, L., Reguera, M. I., Abreu, L. de and
9 Garrick, K.: Enhanced Mediterranean-Atlantic exchange during Atlantic freshening phases,
10 *Geochem. Geophys. Geosystems*, 11, 22 PP., doi:201010.1029/2009GC002931, 2010.

11 Rother, H., Fink, D., Shulmeister, J., Mifsud, C., Evans, M. and Pugh, J.: The early rise and late
12 demise of New Zealand's last glacial maximum, *Proc. Natl. Acad. Sci.*, 111(32), 11630–11635,
13 doi:10.1073/pnas.1401547111, 2014.

14 Rubino, M., Etheridge, D. M., Trudinger, C. M., Allison, C. E., Battle, M. O., Langenfelds, R.
15 L., Steele, L. P., Curran, M., Bender, M., White, J. W. C., Jenk, T. M., Blunier, T. and Francey,
16 R. J.: A revised 1000 year atmospheric $\delta^{13}\text{C}$ -CO₂ record from Law Dome and South Pole,
17 Antarctica, *J. Geophys. Res. Atmospheres*, 118(15), 8482–8499, doi:10.1002/jgrd.50668, 2013.

18 Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J., Louergue,
19 L., Schüpbach, S., Spahni, R., Fischer, H. and Stocker, T. F.: Atmospheric nitrous oxide during
20 the last 140,000 years, *Earth Planet. Sci. Lett.*, 300(1–2), 33–43,
21 doi:10.1016/j.epsl.2010.09.027, 2010.

22 Schmidt, G. A., Jungclaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J.,
23 Delaygue, G., Joos, F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J.,
24 Shindell, D. T., Solanki, S. K., Steinhilber, F. and Vieira, L. E. A.: Climate forcing
25 reconstructions for use in PMIP simulations of the last millennium (v1.0), *Geosci Model Dev*,
26 4(1), 33–45, doi:10.5194/gmd-4-33-2011, 2011.

27 Schmittner, A., Green, J. a. M. and Wilmes, S.-B.: Glacial ocean overturning intensified by
28 tidal mixing in a global circulation model, *Geophys. Res. Lett.*, 42(10), 2015GL063561,
29 doi:10.1002/2015GL063561, 2015.

1 Severinghaus, J. P. and Brook, E. J.: Abrupt Climate Change at the End of the Last Glacial
2 Period Inferred from Trapped Air in Polar Ice, *Science*, 286(5441), 930–934,
3 doi:10.1126/science.286.5441.930, 1999.

4 Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto-Bliesner, B.,
5 Schmittner, A. and Bard, E.: Global warming preceded by increasing carbon dioxide
6 concentrations during the last deglaciation, *Nature*, 484(7392), 49–54,
7 doi:10.1038/nature10915, 2012.

8 Shennan, I.: Global meltwater discharge and the deglacial sea-level record from northwest
9 Scotland, *J. Quat. Sci.*, 14(7), 715–719, doi:10.1002/(SICI)1099-
10 1417(199912)14:7<715::AID-JQS511>3.0.CO;2-G, 1999.

11 Shennan, I. and Milne, G.: Sea-level observations around the Last Glacial Maximum from the
12 Bonaparte Gulf, NW Australia, *Quat. Sci. Rev.*, 22(14), 1543–1547, doi:10.1016/S0277-
13 3791(03)00088-X, 2003.

14 Siegenthaler, U., Stocker, T. F., Monnin, E., Lüthi, D., Schwander, J., Stauffer, B., Raynaud,
15 D., Barnola, J.-M., Fischer, H., Masson-Delmotte, V. and Jouzel, J.: Stable Carbon
16 Cycle–Climate Relationship During the Late Pleistocene, *Science*, 310(5752), 1313–
17 1317, doi:10.1126/science.1120130, 2005.

18 Simonsen, S. B., Johnsen, S. J., Popp, T. J., Vinther, B. M., Gkinis, V. and Steen-Larsen, H. C.:
19 Past surface temperatures at the NorthGRIP drill site from the difference in firn diffusion of
20 water isotopes, *Clim Past*, 7(4), 1327–1335, doi:10.5194/cp-7-1327-2011, 2011.

21 Smith, R. S. and Gregory, J. M.: A study of the sensitivity of ocean overturning circulation and
22 climate to freshwater input in different regions of the North Atlantic, *Geophys. Res. Lett.*,
23 36(15), L15701, doi:10.1029/2009GL038607, 2009.

24 Spahni, R., Chappellaz, J., Stocker, T. F., Louergue, L., Hausmann, G., Kawamura, K.,
25 Flückiger, J., Schwander, J., Raynaud, D., Masson-Delmotte, V. and Jouzel, J.: Atmospheric
26 Methane and Nitrous Oxide of the Late Pleistocene from Antarctic Ice Cores, *Science*,
27 310(5752), 1317–1321, doi:10.1126/science.1120132, 2005.

1 Stanford, J. D., Hemingway, R., Rohling, E. J., Challenor, P. G., Medina-Elizalde, M. and
2 Lester, A. J.: Sea-level probability for the last deglaciation: A statistical analysis of far-field
3 records, *Glob. Planet. Change*, 79(3–4), 193–203, doi:10.1016/j.gloplacha.2010.11.002, 2011.

4 Steffensen, J. P., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Fischer, H.,
5 Goto-Azuma, K., Hansson, M., Johnsen, S. J., Jouzel, J., Masson-Delmotte, V., Popp, T.,
6 Rasmussen, S. O., Röhlisberger, R., Ruth, U., Stauffer, B., Siggaard-Andersen, M.-L.,
7 Sveinbjörnsdóttir, Á. E., Svensson, A. and White, J. W. C.: High-Resolution Greenland Ice
8 Core Data Show Abrupt Climate Change Happens in Few Years, *Science*, 321(5889), 680–684,
9 doi:10.1126/science.1157707, 2008.

10 Stenni, B., Buiron, D., Frezzotti, M., Albani, S., Barbante, C., Bard, E., Barnola, J. M., Baroni,
11 M., Baumgartner, M., Bonazza, M., Capron, E., Castellano, E., Chappellaz, J., Delmonte, B.,
12 Falourd, S., Genoni, L., Iacumin, P., Jouzel, J., Kipfstuhl, S., Landais, A., Lemieux-Dudon, B.,
13 Maggi, V., Masson-Delmotte, V., Mazzola, C., Minster, B., Montagnat, M., Mulvaney, R.,
14 Narcisi, B., Oerter, H., Parrenin, F., Petit, J. R., Ritz, C., Scarchilli, C., Schilt, A., Schüpbach,
15 S., Schwander, J., Selmo, E., Severi, M., Stocker, T. F. and Udisti, R.: Expression of the bipolar
16 see-saw in Antarctic climate records during the last deglaciation, *Nat. Geosci.*, 4(1), 46–49,
17 doi:10.1038/ngeo1026, 2011.

18 Stocker, T. F.: The Seesaw Effect, *Science*, 282(5386), 61–62,
19 doi:10.1126/science.282.5386.61, 1998.

20 Strelin, J. A., Denton, G. H., Vandergoes, M. J., Ninnemann, U. S. and Putnam, A. E.:
21 Radiocarbon chronology of the late-glacial Puerto Bandera moraines, Southern Patagonian
22 Icefield, Argentina, *Quat. Sci. Rev.*, 30(19–20), 2551–2569,
23 doi:10.1016/j.quascirev.2011.05.004, 2011.

24 Svendsen, J. I., Alexanderson, H., Astakhov, V. I., Demidov, I., Dowdeswell, J. A., Funder, S.,
25 Gataullin, V., Henriksen, M., Hjort, C., Houmark-Nielsen, M., Hubberten, H. W., Ingólfsson, O.,
26 Jakobsson, M., Kjær, K. H., Larsen, E., Lokrantz, H., Lunkka, J. P., Lyså, A., Mangerud, J.,
27 Matiouchkov, A., Murray, A., Möller, P., Niessen, F., Nikolskaya, O., Polyak, L., Saarnisto, M.,
28 Siegert, C., Siegert, M. J., Spielhagen, R. F. and Stein, R.: Late Quaternary ice sheet history
29 of northern Eurasia, *Quat. Sci. Rev.*, 23(11–13), 1229–1271,
30 doi:10.1016/j.quascirev.2003.12.008, 2004.

1 Tarasov, L. and Peltier, W. R.: Terminating the 100 kyr ice age cycle, *J. Geophys. Res.*
2 *Atmospheres*, 102(D18), 21665–21693, doi:10.1029/97JD01766, 1997.

3 Tarasov, L. and Peltier, W. R.: Greenland glacial history and local geodynamic consequences,
4 *Geophys. J. Int.*, 150(1), 198–229, doi:10.1046/j.1365-246X.2002.01702.x, 2002.

5 Tarasov, L. and Peltier, W. R.: A geophysically constrained large ensemble analysis of the
6 deglacial history of the North American ice-sheet complex, *Quat. Sci. Rev.*, 23(3–4), 359–388,
7 doi:10.1016/j.quascirev.2003.08.004, 2004.

8 Tarasov, L. and Peltier, W. R.: Arctic freshwater forcing of the Younger Dryas cold reversal,
9 *Nature*, 435(7042), 662–665, doi:10.1038/nature03617, 2005.

10 Tarasov, L. and Peltier, W. R.: A calibrated deglacial drainage chronology for the North
11 American continent: evidence of an Arctic trigger for the Younger Dryas, *Quat. Sci. Rev.*, 25(7–
12 8), 659–688, doi:10.1016/j.quascirev.2005.12.006, 2006.

13 Tarasov, L., Dyke, A. S., Neal, R. M. and Peltier, W. R.: A data-calibrated distribution of
14 deglacial chronologies for the North American ice complex from glaciological modeling, *Earth*
15 *Planet. Sci. Lett.*, 315–316, 30–40, doi:10.1016/j.epsl.2011.09.010, 2012.

16 Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: An Overview of CMIP5 and the Experiment
17 Design, *Bull. Am. Meteorol. Soc.*, 93(4), 485–498, doi:10.1175/BAMS-D-11-00094.1, 2011a.

18 Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: A Summary of the CMIP5 Experiment Design,
19 [online] Available from: http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf
20 (Accessed 13 November 2014b), 2011.

21 Teller, J. T., Leverington, D. W. and Mann, J. D.: Freshwater outbursts to the oceans from
22 glacial Lake Agassiz and their role in climate change during the last deglaciation, *Quat. Sci.*
23 *Rev.*, 21(8–9), 879–887, doi:10.1016/S0277-3791(01)00145-7, 2002.

24 Thiagarajan, N., Subhas, A. V., Southon, J. R., Eiler, J. M. and Adkins, J. F.: Abrupt pre-
25 Bolling-Allerod warming and circulation changes in the deep ocean, *Nature*, 511(7507), 75–
26 78, doi:10.1038/nature13472, 2014.

27 Thornalley, D. J. R., McCave, I. N. and Elderfield, H.: Freshwater input and abrupt deglacial
28 climate change in the North Atlantic, *Paleoceanography*, 25(1), PA1201,
29 doi:10.1029/2009PA001772, 2010.

1 Thornalley, D. J. R., Barker, S., Broecker, W. S., Elderfield, H. and McCave, I. N.: The
2 Deglacial Evolution of North Atlantic Deep Convection, *Science*, 331(6014), 202–205,
3 doi:10.1126/science.1196812, 2011.

4 Tierney, J. E., Russell, J. M., Huang, Y., Damsté, J. S. S., Hopmans, E. C. and Cohen, A. S.:
5 Northern Hemisphere Controls on Tropical Southeast African Climate During the Past 60,000
6 Years, *Science*, 322(5899), 252–255, doi:10.1126/science.1160485, 2008.

7 Timm, O. and Timmermann, A.: Simulation of the Last 21 000 Years Using Accelerated
8 Transient Boundary Conditions*, *J. Clim.*, 20(17), 4377–4401, doi:10.1175/JCLI4237.1, 2007.

9 Tjallingii, R., Claussen, M., Stuut, J.-B. W., Fohlmeister, J., Jahn, A., Bickert, T., Lamy, F. and
10 Röhl, U.: Coherent high- and low-latitude control of the northwest African hydrological
11 balance, *Nat. Geosci.*, 1(10), 670–675, doi:10.1038/ngeo289, 2008.

12 Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi, M., Svensson, A., Vinther, B. and Wolff, E. W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Clim Past*, 9(4), 1733–1748, doi:10.5194/cp-9-1733-2013, 2013.

17 Verschuren, D., Sinninghe Damsté, J. S., Moernaut, J., Kristen, I., Blaauw, M., Fagot, M.,
18 Haug, G. H., Geel, B. van, Batist, M. D., Barker, P., Vuille, M., Conley, D. J., Olago, D. O.,
19 Milne, I., Plessen, B., Eggermont, H., Wolff, C., Hurrell, E., Ossebaar, J., Lyaruu, A., Plicht, J.,
20 van der, Cumming, B. F., Brauer, A., Rucina, S. M., Russell, J. M., Keppens, E., Hus, J.,
21 Bradley, R. S., Leng, M., Mingram, J. and Nowaczyk, N. R.: Half-precessional dynamics of
22 monsoon rainfall near the East African Equator, *Nature*, 462(7273), 637–641,
23 doi:10.1038/nature08520, 2009.

24 Vettoretti, G. and Peltier, W. R.: Interhemispheric air temperature phase relationships in the
25 nonlinear Dansgaard-Oeschger oscillation, *Geophys. Res. Lett.*, 42(4), 2014GL062898,
26 doi:10.1002/2014GL062898, 2015.

27 Waelbroeck, C., Labeyrie, L., Duplessy, J. C., Guiot, J., Labracherie, M., Leclaire, H. and
28 Duprat, J.: Improving past sea surface temperature estimates based on planktonic fossil faunas,
29 *Paleoceanography*, 13(3), 272–283, doi:10.1029/98PA00071, 1998.

1 Weaver, A. J., Saenko, O. A., Clark, P. U. and Mitrovica, J. X.: Meltwater Pulse 1A from
2 Antarctica as a Trigger of the Bølling-Allerød Warm Interval, *Science*, 299(5613), 1709–1713,
3 doi:10.1126/science.1081002, 2003.

4 Weber, S. L., Drijfhout, S. S., Abe-Ouchi, A., Crucifix, M., Eby, M., Ganopolski, A.,
5 Murakami, S., Otto-Bliesner, B. and Peltier, W. R.: The modern and glacial overturning
6 circulation in the Atlantic ocean in PMIP coupled model simulations, *Clim Past*, 3(1), 51–64,
7 doi:10.5194/cp-3-51-2007, 2007.

8 Weijers, J. W. H., Schefuß, E., Schouten, S. and Damsté, J. S. S.: Coupled Thermal and
9 Hydrological Evolution of Tropical Africa over the Last Deglaciation, *Science*, 315(5819),
10 1701–1704, doi:10.1126/science.1138131, 2007.

11 Whitehouse, P. L., Bentley, M. J. and Le Brocq, A. M.: A deglacial model for Antarctica:
12 geological constraints and glaciological modelling as a basis for a new model of Antarctic
13 glacial isostatic adjustment, *Quat. Sci. Rev.*, 32, 1–24, doi:10.1016/j.quascirev.2011.11.016,
14 2012.

15 Yokoyama, Y., Esat, T. M. and Lambeck, K.: Last glacial sea-level change deduced from
16 uplifted coral terraces of Huon Peninsula, Papua New Guinea, *Quat. Int.*, 83–85, 275–283,
17 doi:10.1016/S1040-6182(01)00045-3, 2001a.

18 Yokoyama, Y., De Deckker, P., Lambeck, K., Johnston, P. and Fifield, L. K.: Sea-level at the
19 Last Glacial Maximum: evidence from northwestern Australia to constrain ice volumes for
20 oxygen isotope stage 2, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 165(3–4), 281–297,
21 doi:10.1016/S0031-0182(00)00164-4, 2001b.

22

1 Tables

2 Table 1. Summary of recommended model boundary conditions to spin up the last deglaciation
3 Core [simulationexperiment](#) (pre 21 ka); see text for details. Participants are not required to
4 follow the recommendation for these boundary conditions, but must document the method used,
5 including information on the simulation's state of spinup at the point when the Core is started.
6 Data are available from PMIP Last Deglaciation Working Group Wiki:
7 <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglag:index>. Boundary condition group
8 headings are in bold.

Spinup type	Boundary condition	Description
Last Glacial Maximum (LGM; 21 ka)	Insolation Solar constant Eccentricity Obliquity Perihelion-180° Vernal equinox	Preindustrial (e.g. 1365 W m ⁻²) 0.018994 22.949° 114.42° Noon, 21 st March
	Trace gases Carbon dioxide (CO ₂) Methane (CH ₄) Nitrous oxide (N ₂ O) Chlorofluorocarbon (CFC) Ozone (O ₃)	188190 ppm 375 ppb 200 ppb 0 Preindustrial (e.g. 10 DU)
	Ice sheets, orography and coastlines	21 ka data from either: - ICE-6G_C (references in text) - GLAC-1D (references in text)
	Bathymetry	Keep consistent with the coastlines, using either: - Data associated with the ice sheet - Preindustrial bathymetry
	Global ocean salinity	+ 1 psu, relative to preindustrial
Transient orbit and trace gases (26-21 ka)	Orbital parameters	All orbital parameters should be transient, as per Berger (1978) 26-21 ka
	Trace gases Carbon dioxide (CO ₂) Methane (CH ₄) Nitrous oxide (N ₂ O)	Adjusted to the AICC2012 (Veres et al., 2013) Transient, as per LüthiBereiter et al. (20082015) Transient, as per Loulergue et al. (2008) Transient, as per Schilt et al. (2010)
	All others	As per LGM (21 ka) spinup type.

1 **Table 2.** Summary of required model boundary conditions for the last deglaciation Core
 2 [simulation experiment](#) 21-9 ka; optional boundary conditions are labelled as such. Data are
 3 available from PMIP Last Deglaciation Working Group Wiki:
 4 <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:degl:index>. See text for details. Boundary
 5 condition group headings are in bold.

Boundary condition	Description
Initial conditions (pre 21 ka)	Recommended (optional) to use either: <ul style="list-style-type: none"> - Last Glacial Maximum (LGM; 21 ka) equilibrium simulation, including +1 psu global ocean salinity - Transient orbit and trace gases (26-21 ka) and all other boundary conditions fixed as per equilibrium LGM See Table 1 for details. The method must be documented, including information on the state of spinup
Insolation	
Solar constant	Preindustrial (e.g. 1365 W m ⁻²)
Orbital parameters	Transient, as per Berger (1978)
Trace gases	Adjusted to the AICC2012 age model (Veres et al., 2013):
Carbon dioxide (CO ₂)	Transient, as per Lüthi-Bereiter et al. (2015)
Methane (CH ₄)	Transient, as per Loulergue et al. (2008)
Nitrous oxide (N ₂ O)	Transient, as per Schilt et al. (2010)
Chlorofluorocarbon (CFC)	0
Ozone (O ₃)	Preindustrial (e.g. 10 DU)
Ice sheet	Transient, with a choice of either : <ul style="list-style-type: none"> - ICE-6G_C reconstruction (references in text) - GLAC-1D reconstruction (references in text) How often to update the ice sheet is optional
Orography and coastlines	Transient. To be consistent with the choice of ice sheet. Orography is updated on the same timestep as the ice sheet. It is optional how often the land-sea mask is updated, but ensure consistency with the ice sheet reconstruction is maintained
Bathymetry	Keep consistent with the coastlines and otherwise use either: <ul style="list-style-type: none"> - Transient data associated with the chosen ice sheet; it is optional how often the bathymetry is updated- - Preindustrial bathymetry
River routing	Ensure that rivers reach the coastline It is recommended (optional) to use one of the following: <ul style="list-style-type: none"> - Preindustrial configuration for the model - Transient routing provided with the GLAC-1D-ice sheet reconstructions - Manual/model calculation of river network to match topography
Freshwater fluxes	No land ice or iceberg meltwater fluxes to the ocean At participant discretion. Three options are: <i>melt-uniform</i> , <i>melt-routed</i> and <i>no-melt</i> (see text). It is recommended (optional) to run at least one Core simulation with a scenario consistent with the chosen ice sheet reconstruction to conserve salinity (e.g. as provided). See text for full details (Sect. 2.5)
Other (optional)	
Vegetation and land cover	Prescribed preindustrial cover or dynamic vegetation model
Aerosols (dust)	Prescribed preindustrial distribution or prognostic aerosols

1 **Figures**

2 **Figure 1.** The last deglaciation; forcings and events. **(a)** The three phases of the Core
3 [simulationexperiment](#) (Sect. 4). **(b)** Climate events/periods discussed in the text; Last Glacial
4 Maximum (LGM; 23-19 ka as according to the EPILOG definition; Mix et al., 2001), Heinrich
5 Stadial 1 (HS1), Heinrich Event 1 (H1), Bølling Warming (BW) and Meltwater Pulse 1a
6 (MWP1a), Antarctic Cold Reversal (ACR) and the Younger Dryas cooling (YD). **(c)** June
7 insolation at 60° N and December insolation at 60° S (Berger, 1978). **(d)** Atmospheric carbon
8 dioxide concentration (recent composite of EPICA Dome C, Vostok, Taylor Dome, Siple Dome
9 and West Antarctic Ice Sheet Divide records, Antarctica; Bereiter et al., 2015); black dashed
10 line shows preindustrial concentration. **(e)** Atmospheric methane concentration (EPICA Dome
11 C, Antarctica; Loulergue et al., 2008); green dashed line shows preindustrial concentration. **(f)**
12 Atmospheric nitrous oxide concentration (Talos Dome, Antarctica; Schilt et al., 2010); brown
13 dashed line shows preindustrial concentration. **(g)** Volume of the ice sheets according to the
14 ICE-6G_C reconstruction (solid lines; Argus et al., 2014; Peltier et al., 2015) and the GLAC-
15 1D reconstruction (dashed lines; Briggs et al., 2014; Tarasov et al., 2012; Tarasov and Peltier,
16 2002). [Associated meltwater scenarios melt-uniform and melt-routed \(see Sect. 2.5\) are](#)
17 [consistent with these; all ice mass loss shown is supplied as freshwater to the ocean.](#) **(h)**
18 Greenland temperature reconstruction with $\pm 1 \sigma$ shaded (averaged GISP2, NEEM and NGRIP
19 records; Buizert et al., 2014). **(i)** Antarctic δD (EPICA Dome C; Jouzel et al., 2007). **(d)-(f)** and
20 **(h)-(i)** are given on the AICC2012 timescale (Veres et al., 2013).

21 **Figure 2.** Northern Hemisphere ice sheet elevation at 21, 18, 15, 12 and 9 ka; **(a)** ICE-6G_C
22 reconstruction at 10 arcminute horizontal resolution, elevation is plotted where the fractional
23 ice mask is more than 0.5 (Peltier et al., 2015); **(b)** GLAC-1D reconstruction at 1° [\(longitude\)](#)
24 [× 0.5° \(latitude\)](#) horizontal resolution, elevation is plotted where the [binaryfractional](#) ice mask
25 is [onemore than 0.5](#) (Briggs et al., 2014; Tarasov [et al., 2012; Tarasov](#) and Peltier, 2002;
26 [Tarasov et al., 2012](#)[this study](#)).

27 **Figure 3.** Atmospheric trace gases through the last deglaciation from Antarctic ice cores. **(a)**
28 [CarbonCore experiment carbon](#) dioxide according to a [recent](#) composite record from EPICA
29 Dome C (EDC), [West Antarctic Ice Sheet Divide \(WDC\)](#), Vostok [and](#), Taylor Dome [and](#) Siple
30 Dome [\(thick black line; Bereiter et al., 2015\), adjusted to, which was produced on](#) the
31 AICC2012 chronology (Veres et al., 2013). [Also shown for comparison is an older composite](#)

1 [record from EDC, Vostok and Taylor Dome](#) (thin blue line; Lüthi et al., 2008, adjusted to the
2 AICC2012 chronology). [The, as well as the](#) original EDC CO₂ record (green line; Monnin et
3 al., 2004) and [more](#) the recent, higher resolution [West Antarctic Ice Sheet Divide](#) (WDC) CO₂
4 record (dark red line; Marcott et al., 2014); [which were incorporated into the newer composite](#)
5 [by Bereiter et al.](#) (2015) [are shown for comparison.](#) (b) Methane according to the EPICA Dome
6 C (EDC) record (Loulergue et al., 2008), shown both on the original EDC1 chronology (green
7 line; Spahni et al., 2005) and adjusted to the more recent AICC2012 chronology [for the Core](#)
8 [experiment](#) (thick black line; Veres et al., 2013). (c) Nitrous oxide according to the Talos Dome
9 (TALDICE) record (Schilt et al., 2010), adjusted to the AICC2012 chronology [for the Core](#)
10 [experiment](#) (thick black line; Veres et al., 2013). For comparison, the earlier EPICA Dome C
11 (EDC) record on the EDC1 chronology is also shown (green line; Spahni et al., 2005). The
12 nearest measured N₂O concentration to 21 ka is from 21.089 ka; hence the small offset between
13 the slightly earlier concentration (187 ppb) used for the Core and the interpolated value plotted
14 at 21 ka. For (a)-(c) 21 ka concentrations according to the AICC2012 age model (red dots) are
15 shown in contrast to previous PMIP3 LGM concentrations (blue dots; PMIP LGM Working
16 Group, 2010). If using an equilibrium-type spinup for the start of [the](#) transient Core simulation
17 at 21 ka (Sect. 2.1.1), use [188190](#) ppm CO₂, 375 ppb CH₄ and 200 ppb N₂O.

18 **Figure 4.** Southern Hemisphere ice sheet elevation at 21, 12 and 9 ka; (a) ICE-6G_C
19 reconstruction at 10 arcminute horizontal resolution, ice elevation is plotted where the fractional
20 ice mask is more than 0.5 (Argus et al., 2014; Peltier et al., 2015); (b) GLAC-1D reconstruction
21 at 1° [\(longitude\) × 0.5° \(latitude\)](#) horizontal resolution, [ice](#) elevation is plotted where [the](#)
22 [binaryfractional](#) ice mask is [1more than 0.5](#) (Briggs et al., 2014; Tarasov et al., 2012; Tarasov
23 and Peltier, 2002).