



1 **PMCAMx-2015 evaluation over Europe against** 2 **AERONET and MODIS aerosol** 3 **optical depth measurements** 4

5 **Antigoni Panagiotopoulou^{1,2}, Panagiotis Charalambidis^{1,3}, Christos Fountoukis¹,**
6 **Christodoulos Pilinis³, Spyros N. Pandis^{1,2,4}**

7 ¹Institute of Chemical Engineering Sciences (ICE-HT/FORTH), Platani, P.O. Box 1414, Patras, 26504,
8 Greece

9 ²Department of Chemical Engineering, University of Patras, University Hill, Patras, 26504, Greece

10 ³Department of Environment, University of the Aegean, University Hill, Mytilene, 81100, Greece

11 ⁴Department of Chemical Engineering, Carnegie Mellon University, Pittsburgh, PA 15213, USA

12

13 *Correspondence to:* Spyros N. Pandis (spyros@chemeng.upatras.gr)

14

15 **Abstract.** The ability of the chemical transport model (CTM) PMCAMx to reproduce aerosol optical
16 depth (AOD) measurements by the Aerosol Robotic Network (AERONET) and the Moderate
17 Resolution Imaging Spectroradiometer (MODIS) over Europe during a photochemically active period is
18 evaluated. Periods with high dust levels are excluded so the analysis focuses on the ability of the model
19 to simulate the mostly secondary aerosol and its interactions with water. PMCAMx reproduces the
20 monthly mean MODIS and AERONET AOD values over the Iberian Peninsula, the British Isles, central
21 Europe, and Russia with fractional bias less than 15% and fractional error less than 30%. However, the
22 model overestimates the AOD over northern Europe probably due to an overestimation of organic
23 aerosol and sulfates. On the other end, PMCAMx underestimates the monthly mean MODIS AOD over
24 the Balkans, the Mediterranean, and the South Atlantic. These errors are probably due to an
25 underestimation of sulfates. Sensitivity tests indicate that the evaluation results of the monthly mean
26 AODs are quite sensitive to the relative humidity (RH) fields used by PMCAMx, but are not sensitive to
27 the simulated size distribution and the black carbon mixing state.



1 Introduction

Atmospheric aerosols are suspensions of solid and/or liquid particles in air that scatter and absorb light. The aerosol optical depth (AOD) is defined as the integrated extinction coefficient over the entire atmospheric column and is a measure of the total aerosol loading (King et al., 1999; Kokhanovsky, 2008; Vijayarachavan et al., 2008; Hidy et al., 2009). Calculations of AOD require knowledge of the aerosol vertical profile, including the particulate matter size distribution, chemical composition, and microphysical state (Seinfeld and Pandis, 2006).

Aerosol properties can be retrieved from ground-based measurements as well as from satellite earth observations (Holben et al., 1998; Levy et al., 2007a, b; Kokhanovsky, 2008; Levy et al., 2010; Duncan et al., 2014; Hu et al., 2014). Global observations of high spatial coverage are provided by satellites (King et al., 1999; Vijayarachavan et al., 2008; Hidy et al., 2009) and more limited spatial coverage by ground-based stations. Regarding temporal coverage, satellite observations are sparse when compared against ground measurements. Ground-based measurements of AOD are direct measurements while satellite AOD measurements are indirect, resulting from inversion procedures and exhibiting larger uncertainties. The magnitude of the satellite AOD uncertainties is higher over land where the surface reflectance cannot be neglected and it must be retrieved simultaneously with the aerosol properties (Levy et al., 2007a, 2007b, 2010). The satellite inversion procedure is simpler over water since the surface contribution is small and the detected signal is mostly due to aerosol reflectance (Shi et al., 2011; Anderson et al., 2013; Schutgens et al., 2013).

Chemical transport models (CTMs) are valuable tools for the study of the impact of pollutant emissions, the development of air quality improvement strategies, studies of aerosol radiative forcing, visibility, and global climate change. Uncertainties of the CTM's input data, including meteorological fields, emission inventories, and boundary conditions as well as weaknesses in representation of atmospheric processes may lead to weak model performance (Kinne et al., 2003, 2006). CTMs have



1 been used in the past to provide AOD predictions either globally (Chin et al., 2002, 2004; Lee et al.,
2 2010; Johnson et al., 2012; Meij et al., 2012; Pozzer et al., 2012; Yu et al., 2012) or over specific
3 regions like Asia (Han et al., 2010; Park et al., 2011), United States (Roy et al., 2007), and Europe
4 (Jeuken et al., 2001; Hodzic et al., 2006; Meij et al., 2007; Tombette et al., 2008; Myhre et al., 2009;
5 Carnevale et al., 2011; Im et al., 2014). Model evaluation often relies on in-situ ground measurements
6 but also measurements from airborne platforms. These in-situ measurements cover by necessity a
7 limited part of the modeling domain. Comparisons against remote sensing data have been used to close
8 that gap.

9 Jeuken et al. (2001) compared the TM3 CTM AOD predictions with the ATSR-2 radiometer
10 AOD retrievals during a 1997 summer episode over Europe. Model errors (neglecting organics and
11 mineral aerosol) in the vertical distributions of sulfate, ammonium, and nitrate, in the hygroscopic
12 growth, and in the optical parameters led to an average AOD (at 550 nm) underestimation by 0.17-0.19.
13 Hodzic et al. (2006) used the CHIMERE model to simulate AOD at 865 nm over Europe during August
14 2003. The model generally reproduced AOD within a factor of 2 and with correlation coefficients
15 ranging from 0.4 to 0.6 in comparison with POLDER and AERONET. Sporadic aerosol emissions due
16 to forest fires or dust events led to regional AOD underestimations. Meij et al. (2007) used the
17 mesoscale TAPOM model to investigate AOD over Milan, Italy during June 2001. Simulated and
18 observed AODs by AERONET, MODIS and MISR (Multi-angle Imaging Spectroradiometer) differed
19 by a factor of 2 or 3 in days with cirrus clouds and Saharan dust but showed good agreement in clear
20 sky days. A finer model resolution gave a more detailed AOD distribution pattern and improved by 15%
21 the agreement with the AOD observations. Tombette et al. (2008) compared the Polair3D estimated
22 AOD against AERONET measurements over Europe for 2001. The black carbon (BC) mixing state had
23 almost no effect on the estimated single scattering albedo (SSA) but the aerosol water content
24 influenced significantly both the SSA and the AOD. Myhre et al. (2009) used the global Oslo CTM2 to



1 predict AODs at 550 nm focusing on specific European regions (Adriatic Sea, Black Sea, and Po
2 Valley). Comparisons against AOD measurements from AERONET, MODIS, and MISR were
3 presented for a short period during late summer-early autumn of 2004. The model underestimated AOD
4 around Venice against AERONET because of organic carbon underestimation. Carnevale et al. (2011)
5 implemented the TCAM CTM to simulate AODs during 2004 over Italy. In general, TCAM was found
6 to underestimate MODIS AODs. Analysis of the extinction coefficient showed that the submicron
7 inorganic aerosol played a key role. Im et al. (2014) simulated air pollution over Europe using the
8 WRF-CMAQ modeling system for 2008. The model underestimated AERONET AOD measurements
9 by 3-22% on average. AOD underestimations were attributed to underestimation of either the
10 anthropogenic emissions or the natural and re-suspended dust emissions.

11 The PM₁ composition predictions of PMCAMx have been evaluated over Europe in May 2008
12 (Fountoukis et al., 2011). PMCAMx performance against airborne measurements was as good as its
13 performance against the hourly ground measurements. More than 94% of the organic aerosol (OA)
14 hourly values and more than 82% of the sulfate ones were reproduced within a factor of 2. However, the
15 evaluation was limited in space. The model performance was only evaluated against ground
16 measurements which were taken at 4 stations located in the Netherlands, Greece, Ireland, and Germany
17 as well as against airborne measurements from 15 flights in North-Western Europe.

18 One of the limitations of the previous evaluation exercises is that errors in dust emissions,
19 transport, and removal often dominate the overall results. In the present work MODIS and AERONET
20 AODs are filtered to exclude periods with high dust levels and to focus on the rest of the anthropogenic
21 and biogenic aerosol components. A period with high photochemical activity is selected so that the
22 emphasis is on secondary aerosol components.

23 In the present study we provide a first time evaluation of the ability of PMCAMx (Murphy and
24 Pandis, 2009; Fountoukis et al., 2011) to reproduce AOD observations over Europe. The objective of



1 this work is to identify weaknesses and strengths of PMCAMx and its inputs, by taking advantage of the
2 wide spatial coverage of MODIS and the temporal coverage of AERONET.

3 4 **2 PMCAMx description**

5 PMCAMx-2015 is a three-dimensional CTM that employs the framework of CAMx (Environ,
6 2003) simulating the processes of horizontal and vertical advection, horizontal and vertical dispersion,
7 wet and dry deposition as well as gas, aqueous, and aerosol-phase chemistry. Three detailed aerosol
8 models are employed: inorganic aerosol growth (Gaydos et al., 2003; Koo et al., 2003), aqueous-phase
9 chemistry (Fahey and Pandis, 2001) as well as OA formation and chemical aging (Murphy and Pandis,
10 2009). The specific modules utilize a sectional approach that dynamically models the evolution of the
11 aerosol size distribution. Ten size sections covering particle diameters from 40 nm to 40 μm are used.
12 PMCAMx treats both primary and secondary organic aerosol as semivolatile and photochemically
13 reactive employing the volatility basis set (Murphy and Pandis, 2009).

14 The PMCAMx European modeling domain in this application is a region of 5,400x5,832 km^2
15 with 36x36 km^2 grid resolution and 14 vertical layers extending up to approximately 6 km. Simulations
16 were performed on a polar stereographic map projection. Horizontal wind components, vertical
17 diffusivity, temperature, pressure, water vapor, clouds, and rainfall were provided by the Weather
18 Research and Forecasting meteorological model (WRF) (Skamarock et al., 2008). Anthropogenic gas
19 emissions are from the European emissions database GEMS while elemental carbon and organic carbon
20 emissions are from the Pan European Carbonaceous Aerosol Inventory (Kulmala et al., 2011). Biogenic
21 emissions were calculated by the MEGAN v2.04 model (Guenther et al., 2006). The marine aerosol
22 emission model developed by O' Dowd et al. (2008) was employed for the estimation of mass fluxes for
23 both accumulation and coarse mode, including the organic aerosol fraction. Emissions from wildfires
24 are taken from IS4FIRES (Sofiev et al., 2009).



1 To limit the effect of the initial conditions on the results, the first six days of each simulation
2 were excluded from the analysis. Concentrations of the major PM_{2.5} aerosol components at the
3 boundaries of the domain (Table S1 in Supplementary Information) are based on measurements of
4 typical background concentrations in sites close to the domain boundaries (Zhang et al., 2007; Seinfeld
5 and Pandis, 2006). All concentrations given here are under ambient temperature and pressure
6 conditions.

7

8 **2.1 AOD prediction by PMCAMx**

9 The size and chemically resolved concentrations of aerosol particles are simulated by PMCAMx
10 for every computational cell. Inorganic aerosol water concentration is calculated online by the
11 thermodynamic equilibrium model ISORROPIA (Nenes et al., 1998). Taking into account all the
12 vertical layers, we calculate the PMCAMx AOD at 550 nm as the sum of the extinction coefficients at
13 each layer:

14

$$15 \quad \text{AOD} = \sum_{i=1}^{14} b_{ext,i} \Delta z_i \quad (1)$$

16 where $b_{ext,i}$ is the extinction coefficient of layer i and Δz_i is the corresponding layer thickness. Assuming
17 that the particles are homogeneous spheres and that all particles in each size bin have the same
18 composition (internal mixture), the aerosol extinction coefficient ($b_{ext,i}$) for layer i is:

19

$$20 \quad b_{ext,i} = \sum_{j=1}^{10} \frac{\pi D_j^2}{4} N_j Q_{ext,j}(m_j, D_j) \quad (2)$$

21 where D_j is the mean diameter of size bin j , N_j is the aerosol number concentration for bin j and $Q_{ext,j}$ is
22 the extinction efficiency of a single particle having a complex refractive index m_j . The extinction
23 efficiency for bin j is estimated as the sum of the scattering, $Q_{scat,j}$, and absorption, $Q_{abs,j}$, efficiencies:



$$Q_{ext,j} = Q_{scat,j} + Q_{abs,j} \quad (3)$$

1
2 Aerosol scattering and absorption efficiencies ($Q_{scat,j}$, $Q_{abs,j}$) are calculated using Mie theory (Seinfeld
3 and Pandis, 2006) and mass concentrations provided for each size bin by PMCAMx, including the
4 concentrations of particulate water. The complex refractive, m_j , index of a homogeneous sphere is
5 estimated using the volume weighted average of the individual refractive indices (Pilinis and Pandis,
6 1995). Sulfate and ammonium are assumed to have a real refractive index of 1.53, which is the value of
7 ammonium sulfate (GEISA, 2011; NASA, 2006). Nitrate is assumed to have a real refractive index of
8 1.56, similar to the value of ammonium nitrate (NASA, 2006). Sodium and chloride have a real
9 refractive index of 1.5 (GEISA, 2011; NASA, 2006). Dust is assumed to have a complex refractive
10 index of $1.53-0.0055i$ (GEISA, 2011). OA is assumed to be non-absorbing with a refractive index of 1.5
11 (Nessler et al., 2005; Fierz Schmidhauser et al., 2010). Biomass burning was minimal during the period
12 of interest (Crippa et al., 2014), so this simplifying assumption regarding the OA absorptivity has little
13 effect on the predicted AOD. The black carbon refractive index has the largest uncertainties (Bond and
14 Bergstrom, 2005) and we use a value of $1.75-0.44i$ (GEISA, 2011). In the base case BC is assumed to
15 be internally mixed with the other components in each size range. The sensitivity of the model
16 predictions to this assumption is discussed in a subsequent section.

17 18 **3 MODIS and AERONET data**

19 The cloud screened and quality assured Level 2 AERONET direct AOD measurements are used
20 for the PMCAMx evaluation. AERONET applies the Beer Lambert Bouguer law to measure AOD from
21 direct sun observations (Holben et al., 1998) therefore it is considered to be the ground truth with AOD
22 uncertainties of 0.01 – 0.02 (Eck et al., 1999). In this work only the AOD values corresponding to
23 Angstrom Exponent values greater than 0.9 are employed in an effort to exclude periods with high dust
24 levels (Schuster et al., 2006). This filter rejected 29% of AODs over land and 28% over water. The
25 geographical distribution of the corresponding AERONET stations is depicted in Fig. 1 and the number



1 of stations in each region is shown in Table 1. Some AERONET stations in the domain of interest did
2 not have available Level 2 AOD data for the period of interest while all data from three stations
3 (OHP_OBSERVATOIRE in South France, FORTH_CRETE in Crete, Greece, and ATHENS_NOA in
4 Athens, Greece) have been excluded after the dust coarse particle rejection filtering.

5 The polar-orbiting MODIS monitors global aerosol properties from two satellites: Terra and
6 Aqua (Salomonson et al., 1989). MODIS employs 36 channels from 0.412 to 14.2 μm , has a wide swath
7 of 2,330 km, and observes every part of the globe at least once daily. The default resolution for aerosol
8 retrieval is 10x10 km^2 (Levy et al., 2009). Each data set retrieved by MODIS is associated with a
9 Quality Assurance Confidence (QAC) flag which ranges from 0 (no confidence) to 3 (highest
10 confidence). For increased spatial coverage we use the union of Terra and Aqua MODIS AOD
11 retrievals with $\text{QAC} \geq 1$. We employ the MODIS Level 2 Collection 5.1 aerosol datasets. AOD
12 retrievals are provided at seven wavelengths (470, 550, 660, 870, 1,200, 1,600, 2,100 nm) over water
13 surface and four wavelengths (470, 550, 660, 2,100 nm) over land. In this study we focus on the 550 nm
14 values. Figure 2 presents the geographical distribution of the available MODIS AOD measurements
15 during the period of interest (1-29 May 2008) over Europe. The average number of retrievals is 12 ± 9 .
16 The maximum number of retrievals is 65 in areas in the North Atlantic.

17 Dust emissions from the Sahara are not included in the PMCAMx emissions used here and the
18 focus of this study is on periods and regions in which Saharan dust does not contribute significantly to
19 the AOD. To exclude periods with high dust levels and to focus on the rest of the anthropogenic and
20 biogenic aerosol components, MODIS AODs are filtered. Over water we employ the dust coarse
21 particle rejection filter of Barnaba and Gobbi (2004). According to this filter, AOD values greater than
22 0.3 also corresponding to coarse mode fraction higher than 0.3 are assumed to be dust-influenced
23 periods. Over land we only use the AOD values which correspond to Angstrom Exponent values



1 exceeding 0.9 (Schuster et al., 2006). The above filters discard 16% of MODIS AOD values over land
2 and 0.4% over water.

3 The evaluation of the MODIS AODs at 550 nm for the land algorithm was performed following
4 the approach of Remer et al. (2005) and Levy et al. (2007b). The collocated data were sorted according
5 to the AERONET measurements. The resulting data were partitioned into bins of 100 points and then
6 averaged. At higher optical depths since the data became sparser we employed 25 points for each bin.
7 The regression results of the collocated AODs, prior to binning had a slope of 1.05. 73% of the 8,331
8 collocated points fall within the expected error envelope denoted by the dashed lines in Fig. S1. These
9 results indicate that the mean MODIS AOD over land in the region and period of interest was retrieved
10 with the expected accuracy. The highest quality flag QAC = 3 provides the closest match but including
11 the QAC = 2 and 1 retrievals results in only a minor reduction of accuracy while increasing significantly
12 the size of the dataset (Table S2).

13 Previous studies have shown that MODIS AOD retrievals have an expected error of $\pm(0.05 +$
14 $0.15AOD_{AERONET})$ over land and $\pm(0.03 + 0.05AOD_{AERONET})$ over water (Chu et al., 2002; Remer et al.,
15 2005; Levy et al., 2007a,b, 2010; Anderson et al., 2013). Table S3 summarizes the values of the
16 expected MODIS AOD uncertainties for the various regions in our modeling domain during May 2008,
17 based on the monthly mean values of AERONET AOD. The MODIS-AERONET AOD differences for
18 this period are consistent with the expected uncertainty of the MODIS retrievals (Fig. S1).

19 20 **4 Evaluation of PMCAMx AOD predictions**

21 The dust-screened monthly mean AODs for Europe during May 2008 retrieved by MODIS and
22 predicted by PMCAMx are shown in Fig. 3. The PMCAMx AODs have been calculated for exactly the
23 same periods as the MODIS retrievals to allow the direct comparison of the two.

24 The MODIS retrievals show high AOD values (> 0.25) over England, South Ireland, North Italy,
25 South Poland, East Romania, Greece, and North Atlantic. Low AOD values (< 0.1) were retrieved over



1 East France, Belgium, Sweden, and North Russia. PMCAMx predicts high AODs over England, South
2 Ireland, North Italy, and central Atlantic and low AODs over North Sweden, East Russia, North and
3 South Atlantic.

4

5 **4.1 Overall evaluation**

6 The difference between PMCAMx and MODIS monthly mean AODs is depicted in Fig. 4.
7 PMCAMx AODs are higher than those of MODIS over England, Ireland, France, Germany, central and
8 South Italy, North and East Europe, central, North and West Russia, West Balkans, and central Atlantic.
9 On the other hand PMCAMx predicts lower AODs than MODIS over parts of Russia, North Italy,
10 central and South Balkans, South Poland, North and South Atlantic, and the African coast of the
11 Mediterranean. On a domain average basis PMCAMx predicts an AOD equal to 0.14 while MODIS
12 retrieved 0.16. Detailed comparisons for each region can be found in Table 2. 94% of the monthly mean
13 AOD values fall inside the expected MODIS error envelope over land (Fig. 5a). Over the whole domain
14 the PMCAMx monthly mean AODs have a mean error of 0.05 and a fractional bias of -16% compared
15 to the MODIS monthly mean AODs (Table 2).

16 PMCAMx AODs were also compared against the AERONET values for the simulation period.
17 Once more the comparisons were done for the grid cells of the AERONET stations and corresponding
18 measurement periods. The PMCAMx monthly mean AODs had a mean error of 0.03 and a fractional
19 bias of 4% compared to the AERONET monthly mean AODs (Table 1). The comparison of the
20 PMCAMx with AERONET monthly mean AODs is summarized in Fig. 5b for the 50 AERONET
21 stations which are employed in the present study.

22

23

24



1 **4.2 Regional evaluation**

2 *Spain and Portugal:* The relatively low AOD levels (0.11 for the 8 AERONET stations and 0.14 for
3 MODIS) are reproduced well by PMCAMx (0.12 for the AERONET sites and 0.12 for the periods of
4 the MODIS retrievals). The monthly mean PMCAMx AOD predictions have a mean error of 0.02
5 (AERONET) and 0.04 (MODIS) (Tables 1 and 2). The model shows little bias (5%) compared to the
6 AERONET stations and a small tendency towards underprediction (-15%) compared to MODIS. 83%
7 of the monthly mean PMCAMx AODs are within the expected MODIS error envelope. Sulfate and
8 organic aerosol are the major components of dry fine PM in Spain and Portugal (Table 3).

9
10 *Russia, Belarus, and Ukraine:* PMCAMx reproduces well (0.14 predicted versus 0.15 measured) the
11 average AOD observations at the 5 AERONET stations in this region (2 in West Russia, 1 in Belarus, 1
12 in Ukraine, and 1 in Crimea) (Table 1). The model has a similar good performance against the MODIS
13 retrievals (0.12 predicted versus 0.13 retrieved) (Table 2). As a result, the monthly mean PMCAMx
14 AOD predictions have a low mean error of 0.02 (AERONET) and 0.04 (MODIS). PMCAMx shows a
15 slight tendency towards underprediction (-11%) compared to AERONET and no bias (<1%) compared
16 to MODIS. 92% of the monthly mean PMCAMx AODs are within the expected MODIS error envelope.
17 Sulfates and organic aerosol predominate in this region and it appears that PMCAMx performs
18 reasonably well in this ground-level measurement poor region.

19
20 *United Kingdom (UK) and Ireland:* This area was relatively polluted during the simulation period with
21 high levels of nitrates, sulfates, and organic aerosol (Table 3). PMCAMx reproduces the relatively high
22 average MODIS (0.23 predicted versus 0.21 retrieved) and AERONET (0.24 predicted versus 0.25
23 measured in the station of Chibolton). The monthly mean PMCAMx AODs have a mean error of 0.04
24 compared to MODIS with a small tendency towards overprediction (14%). 90% of the monthly mean
25 PMCAMx AODs fall within the expected MODIS error envelope.



1 *Balkans*: The Balkans according to PMCAMx had some of the highest sulfate levels in the domain
2 during the simulation period (Table 3). The model underpredicts the AOD both against MODIS (0.14
3 predicted versus 0.19 retrieved) and the two AERONET stations (0.15 predicted versus 0.21 measured).
4 The corresponding fractional biases are -24% against MODIS and -33% against AERONET. However,
5 80% of the monthly mean PMCAMx AODs fall within the expected MODIS error envelope. Given that
6 most of the predicted AOD is due to the sulfate these results suggest that the PMCAMx underprediction
7 is probably due to their underestimation.

8
9 *Central Europe*: PMCAMx showed a small tendency towards overprediction of the moderate AODs in
10 this region compared to both AERONET (12%) and MODIS (13%). For example, overpredictions were
11 evident over France and Germany (Fig. 4). The corresponding fractional errors on a monthly average
12 basis were 22% against AERONET and 30% against MODIS. Sulfate and organic aerosol are the major
13 fine PM components in central Europe and they are probably slightly overestimated by PMCAMx in
14 this region during the simulation period. Errors in the relative humidity fields could also explain parts
15 of these discrepancies.

16
17 *East Europe*: PMCAMx slightly overpredicted the AODs in this region compared to both AERONET
18 (24%) and MODIS (25%). 82% of the monthly mean PMCAMx AODs fell within the expected MODIS
19 error envelope. PMCAMx predicts more frequently AODs > 0.1 than measured by AERONET during
20 the corresponding period of measurements probably because of an overestimation of sulfates and
21 organic aerosol.

22
23 *North Europe*: PMCAMx reproduces the low pollution levels in this area with a mean AOD of 0.12
24 compared to 0.08 by the 4 AERONET stations. The absolute monthly mean errors are low: 0.04
25 (AERONET) and 0.06 (MODIS). However, there is significant fractional positive bias compared to



1 both AERONET (36%) and MODIS (47%). 54% of the monthly mean PMCAMx AODs fall outside the
2 expected MODIS error envelope. 53% of the hourly PMCAMx AODs are greater than 0.1 while only
3 18% of the AERONET values are greater than 0.1. Sulfates and organic aerosol are the dominant fine
4 PM components in this region and at least one of them is overestimated by the model.

5
6 *Turkey and Northern Africa:* There are only two AERONET stations in this area and PMCAMx
7 underpredicts by 30% the corresponding moderate AOD measurements. However, the model
8 performance appears to be much better against the MODIS retrievals covering a much bigger area and
9 the underprediction drops to 13%. 79% of the PMCAMx AODs fall inside the expected MODIS error
10 envelope. Sulfates and organic aerosol are the major fine PM components in these regions and they are
11 probably slightly underestimated by the model.

12
13 *Mediterranean Sea:* PMCAMx exhibits a tendency towards underprediction (-24%) against MODIS and
14 58% of the monthly mean PMCAMx AODs fall inside the expected MODIS error envelope. The major
15 discrepancies are evident in the southern part of the Mediterranean especially close to the African coast.
16 These suggest that dust may be partially responsible for the errors even after the filtering of the data.
17 The model performance is better in the eastern Mediterranean (Fig. 4). Sulfates dominated the AOD in
18 the Mediterranean during the simulation period according to PMCAMx.

19
20 *South Atlantic:* The PMCAMx AOD predictions are significantly lower (-45%) compared to the
21 MODIS retrievals in this region with 74% of the monthly mean PMCAMx AODs falling outside the
22 expected MODIS error envelope. Sulfate and sea-salt dominated the predicted AOD in this region in
23 May 2008 and there is evidence that they may be underpredicted. However, errors in relative humidity
24 or cloud contamination could be also responsible for these discrepancies (Anderson et al., 2013).



1 *North Atlantic:* The model performance is much better in the North than in the South Atlantic. The
2 mean AOD error is 0.04 compared to MODIS with a tendency towards underprediction (-21%). 53% of
3 the monthly mean PMCAMx AODs fall inside the expected MODIS error envelope. There is one
4 AERONET station in this area (in Helgoland around 50 km from the coast of Germany) and PMCAMx
5 predicts an average AOD equal to 0.16 compared to the 0.11 measured. Sulfates, organic aerosol, and
6 sea-salt were the major fine PM components in North Atlantic during May 2008 (Table 3).

7
8 *Black Sea:* PMCAMx exhibits a tendency towards underprediction (-18%) versus MODIS in this
9 relatively polluted region. 66% of the PMCAMx AODs fall within the expected MODIS error envelope.
10 Sulfates were the major fine PM components in the Black Sea during the simulation period.

11
12 The results of the PMCAMx-MODIS comparison for the various regions are summarized in Fig. 6.
13 These results suggest that the variability of the MODIS retrievals exceeds that of the PMCAMx
14 predictions for almost all areas. This could be due to the spatial resolution of the model inputs including
15 the emissions, but also to other reasons including missing short-term air pollution sources in the
16 inventory, potential cloud contamination of the retrievals, etc.

18 **5 Sensitivity analysis of the predicted AODs**

19 There are various possible sources of bias in the PMCAMx predictions of AOD other than the
20 concentration and composition of aerosol. We explore here the role of the relative humidity calculated
21 by the WRF model, the role of the mixing state of BC, as well as that of the predicted aerosol size
22 distribution.

23 In the first test the absolute humidity was increased uniformly by 5%, while maintaining the
24 maximum relative humidity in cloud-free regions at 99%. The PMCAMx monthly mean AOD increased
25 on average by 13% (Fig. S2). The increases ranged from 7% in Turkey and Northern Africa to 31% in



1 the North Atlantic. This AOD change can explain a significant part of the base case discrepancies which
2 cause a fractional error of PMCAMx 22% versus AERONET and 33% versus MODIS.

3 In another test the diameter of all particles was increased by 20%. 72% of the PMCAMx
4 monthly mean AOD values changed by less than 0.01. The average increase of the monthly mean AOD
5 was 1% (ranging from 0.3% in the Black Sea to 4% in the UK and Ireland).

6 In a third sensitivity test we assumed that BC was always externally mixed with the other
7 components in each size range, forming pure BC spheres. 73% of the PMCAMx monthly mean AOD
8 values changed by less than 0.01 in this test. The average change of the monthly mean AOD was
9 negligible ($< 0.5\%$).

10

11 **6 Conclusions**

12 Previous evaluations of the ability of the 3-D CTM PMCAMx to reproduce the aerosol levels in
13 Europe, the US, and Mexico City have been based on comprehensive chemical composition
14 measurements at a few ground sites and limited data from a few flights. In this study we expand these
15 efforts by using the MODIS retrievals of AOD over Europe during a photochemically active period
16 (May 2008). Given the uncertainty of the MODIS AOD retrievals we compliment the evaluation using
17 the corresponding AERONET observations in the same region which are more accurate but cover only
18 specific areas. We exclude periods during which the different areas are strongly affected by dust
19 (mainly from the Sahara) in an effort to focus on the other primary and secondary anthropogenic and
20 biogenic aerosol components.

21 The details of the evaluation results differ, as expected, depending on the use of either the
22 MODIS or the AERONET results. These differences are due to the different spatial coverage of the
23 datasets, but also to the MODIS AOD retrieval uncertainties. The major conclusion is that PMCAMx
24 can reproduce the observed AODs for this period with little bias (-16% for MODIS and +4% for



1 AERONET). The corresponding fractional errors are 33% against MODIS and 22% against
2 AERONET. These results are consistent with those of Fountoukis et al. (2001) who compared the
3 PMCAMx predictions for the same period against ground measurements of fine PM composition in four
4 sites and airborne measurements from several flights over central and northern Europe.

5 The AOD performance of PMCAMx against the MODIS retrievals is excellent (absolute
6 fractional bias less than 15% and fractional error less than 35%) in the Iberian Peninsula, UK/Ireland,
7 central Europe, Russia-Belarus-Ukraine, Turkey-northern Africa. It is good (absolute fractional bias less
8 than 25% and fractional error less than 35%) in East Europe, the Balkans, and over the Mediterranean,
9 the North Atlantic, and the Black Sea. Finally, its performance is average (absolute fractional bias less
10 than 50% and fractional error less than 55%) in the relatively clean area of North Europe and the South
11 Atlantic. The performance is more or less similar against AERONET with the exception of a few areas
12 with only one or two AERONET stations. The average performance against the AERONET
13 measurements is considered using the above criteria excellent and against MODIS it is the borderline
14 between good and excellent.

15 The above PMCAMx performance suggests that overall the model does a good job in
16 reproducing the fine aerosol sulfate, organics, nitrate, and sea-salt levels over Europe during the
17 evaluation period. Its major weaknesses appear to be potential overpredictions of sulfate and/or organics
18 over North and East Europe, underprediction of sulfate over the Balkans, and underprediction of fine
19 sodium chloride, sulfates, or organics in the southern Mediterranean and South Atlantic. However, these
20 discrepancies are quite sensitive to the relative humidity fields predicted by WRF. In a sensitivity test
21 the average predicted AOD increased by 13% (ranging from 7 to 31% depending on the area) for a
22 uniform 5% change in RH. On the other hand, the details of the fine PM size distribution and the black
23 carbon mixing state have a very small effect on the AOD predictions.

24



1 **Code availability**

2 PMCAMx is the research version of the publicly available CAMx (www.camx.org). The Fortran source
3 code of CAMx (Version 6.20 was posted on March 23, 2015) and a User's Guide both prepared by
4 ENVIRON can be downloaded through the above website. The PMCAMx code is used as testbed for
5 testing of different hypotheses, algorithms, etc. The version used in this paper (PMCAMx-2015) as well
6 as the most current version can be obtained upon request by contacting Prof. S. Pandis
7 (spyros@chemeng.upatras.gr).

8

9 *Acknowledgements.* The NASA MODIS team is acknowledged for preparing and making available
10 MODIS observations. The AERONET team is acknowledged for establishing and maintaining the
11 AERONET sites used in this study. Funding was provided by the FP7 ERC IDEAS project
12 ATMOPACS.

13

14

15

16 **References**

17 Anderson, J. C., Wang, J., Zeng, J., Leptoukh, G., Petrenko, M., Ichoku, C. and Hu, C.: Long-term
18 statistical assessment of Aqua-MODIS aerosol optical depth over coastal regions: bias
19 characteristics and uncertainty sources, *Tellus B*, 65, 1-22, 2013.

20 Barnaba, F. and Gobbi, G. P.: Aerosol seasonal variability over the Mediterranean region and relative
21 impact of maritime, continental and Saharan dust particles over the basin from MODIS data in the
22 year 2001, *Atmos. Chem. Phys.*, 4, 2367-2391, 2004.

23 Bond, T. C. and Bergstrom, W.: Light absorption by carbonaceous particles: An investigative review,
24 *Aerosol. Sci. & Tech.*, 40, 27 – 67, 2005.

25 Carnevale, C., Finzi, G., Mannarini, G., Pisoni, E. and Volta, M.: Comparing mesoscale chemistry-
26 transport model and remote-sensed Aerosol Optical Depth, *Atmos. Envir.*, 45, 289-295, 2011.

27 Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B., Duncan, B., Martin, R., Logan, J., Higurashi,
28 A. and Nakajima, T.: Tropospheric aerosol optical thickness from the GOCART model and
29 comparisons with satellite and sun photometer measurements, *J. Atmos. Sci.*, 59, 461-483, 2002.

30 Chin, M., Chu, A., Levy, R., Remer, L., Kaufman, Y., Holben, B., Eck, T., Ginoux, P. and Gao, Q.:
31 Aerosol distribution in the Northern Hemisphere during ACE-Asia: Results from global model,
32 satellite observations and sun photometer measurements, *J. Geoph. Res.*, 109, 1-15, 2004.



- 1 Chu, D., Kaufman, J. Y., Ichoku, C., Remer, L., Tanre, D. and Holben, N. B.: Validation of MODIS
2 aerosol optical depth retrieval over land, *Geophys. Res. Lett.*, 29, MOD02-1 to MOD02-4,
3 doi:10.1029/2001GL013205, 2002.
- 4 Crippa, M., Canonaco, F., Lanz, V. A., Äijälä, M., Allan, J. D., Carbone, S., Capes, G., Ceburnis, D.,
5 Dall'Osto, M., Day, D. A., De-Carlo, P. F., Ehn, M., Eriksson, A., Freney, E., Hildebrandt, R. L.,
6 Hillamo, R., Jimenez, J. L., Junninen, H., Kiendler-Scharr, A., Kortelainen, A.-M., Kulmala, M.,
7 Laaksonen, A., Mensah, A. A., Mohr, C., Nemitz, E., O'Dowd, C., Ovadnevaite, J., Pandis, S. N.,
8 Petäjä, T., Poulain, L., Saarikoski, S., Sellegri, K., Swietlicki, E., Tiitta, P., Worsnop, D. R.,
9 Baltensperger, U. and Prévôt, A. S. H.: Organic aerosol components derived from 25 AMS data
10 sets across Europe using a consistent ME-2 based source apportionment approach, *Atmos. Chem.*
11 *Phys.*, 14, 6159–6176, doi:10.5194/acp-4-6159-2014, 2014.
- 12 Duncan, B. N., Prados, A. I., Lamsal, L. N., Liu, Y., Streets, D. G., Gupta, P., Hilsenrath, E., Kahn, R.
13 A., Nielsen, J. A., Beyersdorf, A. J., Burton, S. P., Fiore, A. M., Fishman, J., Henze, D. K.,
14 Hostetler, C. A., Krotkov, N. A., Lee, P., Lin, M., Pawson, S., Pfister, G., Pickering, K. E., Pierce,
15 B. R., Yoshida, Y. and Ziem, L. D.: Satellite data of atmospheric pollution for U.S. air quality
16 applications: Examples of applications, summary of data end-user resources, answers to FAQs,
17 and common mistakes to avoid, *Atmos. Env.*, 94, 647-662, 2014.
- 18 Eck, T. F., Holben, B. N., Reid, J. S., Dubovik, O., Smirnov, A., O'Neill, N. T., Slutsker, I. and Kinne,
19 S.: Wavelength dependence of the optical depth of biomass burning, urban, and desert dust
20 aerosols, *J. Geophys. Res.*, 104, 31333–31349, doi: 10.1029/1999JD900923, 1999.
- 21 Environ: User's guide to the comprehensive air quality model with extensions (CAMx), version 4.02,
22 Report, ENVIRON Int. Corp., Novato, Calif., 2003.
- 23 Fahey, K. and Pandis, S.N.: Optimizing model performance: variable size resolution in cloud chemistry
24 modeling, *Atmos. Environ.*, 35, 4471-4478, 2001.
- 25 Fierz, S. R., Zieger, P., Vaishya, A., Monahan, C., Bialek, J., Dowd, O. D. C., Jennings, S. G.,
26 Baltensperger, G. and Weingartner, E.: Light scattering enhancement factors in the marine
27 boundary layer (Mace Head, Ireland), *J. Geophys. Res.*, 115, D20204,
28 doi:20210.21029/22009JD013755, 2010.
- 29 Fountoukis, C., Racherla, P. N., Denier van der Gon, H. C. A., Polymeneas, P., Charalampidis, P. E.,
30 Pilinis, C., Wiedensohler, A., Dall'Osto, M., O'Dowd, C. and Pandis, S. N.: Evaluation of a three-
31 dimensional chemical transport model (PMCAMx) in the European domain during the EUCAARI
32 May 2008 Campaign, *Atmos. Chem. Phys.*, 11, 10331-10347, 2011.
- 33 Gaydos, T., Koo, B. and Pandis, S. N.: Development and application of an efficient moving sectional
34 approach for the solution of the atmospheric aerosol condensation/evaporation equations, *Atmos.*
35 *Environ.*, 37, 3303-3316, 2003.



- 1 GEISA (Gestion et Etude des Informations Spectroscopiques Atmosphériques: Management and Study
2 of Atmospheric Spectroscopic Information) 2011: <http://www.pole-ether.fr/>
- 3 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I. and Geron, C.: Estimates of global
4 terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from
5 Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, 2006.
- 6 Han, X., Zhang, M., Han, Z., Xin, J., Wang, L., Qiu, J. and Liu, Y.: Model analysis of aerosol optical
7 depth distributions over East Asia, *Scienc. China Earth Scienc.*, 53, 1079-1090, 2010.
- 8 Hidy, G. M., Brook, J. R., Chow, J. C., Green, M., Husar, R. B., Lee, C., Scheffe, R. D., Swanson, A.
9 and Watson, J. G.: Remote sensing of particulate pollution from space: Have we reached the
10 promised land? *J. Air & Waste Manag. Assoc.*, 59, 1130-1139, 2009.
- 11 Hodzic, A., Vautard, R., Chepfer, H., Goloub, P., Menut, L., Chazette, P., Deuz, J. L., Apituley, A. and
12 Couvert, P.: Evolution of aerosol optical thickness over Europe during the August 2003 heat wave
13 as seen from CHIMERE model simulations and POLDER data, *Atmos. Chem. Phys.*, 6, 1853-
14 1864, 2006.
- 15 Holben, B. N., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A.,
16 Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I. and Smirnov, A.: AERONET-A
17 federated instrument network and data archive for aerosol characterization, *Remote Sens.*
18 *Environ.*, 66, 1-16, 1998.
- 19 Hu, X., Waller, L. A., Lyapustin, A., Wang, Y. and Liu, Y.: 10-year spatial and temporal trends of
20 PM_{2.5} concentrations in the southeastern US estimated using high-resolution satellite data, *Atmos.*
21 *Chem. Phys.*, 14, 6301-6314, 2014.
- 22 Im, U., Daskalakis, N., Markakis, K., Vrekoussis, M., Hjorth, J., Myriokefalitakis, S.,
23 Gerasopoulos, E., Kouvarakis, G., Richter, A., Burrows, J., Pozzoli, A., Unal, A., Kindap, T. and
24 Kanakidou, M.: Simulated air quality and pollutant budgets over Europe in 2008, *Scienc. Total*
25 *Environ.*, 470-471, 270-281, 2014.
- 26 Jeuken, A., Veeffkind, P., Dentener, F., Metzger, S. and Robles-Gonzales, C.: Simulation of the aerosol
27 optical depth over Europe for August 1997 and a comparison with observations, *J. Geophys. Res.*,
28 106, 28,295-28,311, 2001.
- 29 Johnson, M. S., Meskhidge, N. and Kiliyanpillakil, V. P.: A global comparison of GEOS-Chem-
30 predicted and remotely-sensed mineral dust aerosol optical depth and extinction profiles, *J.*
31 *Advances Model. Earth Syst.*, 4, 1-15, 2012.
- 32 King, M. D., Kaufman, Y. J., Tanre, D. and Nakajima, T.: Remote sensing of tropospheric aerosols
33 from space: Past, present and future, American Meteorological Society, 1999.
- 34 Kinne, S., Lohmann, U., Feichter, J., Timmreck, C., Schulz, M., Ghan, S., Easter, R., Chin, M., Ginoux,
35 P., Takemura, T., Tegen, I., Koch, D., Herzog, M., Penner, J., Pitari, G., Holben, B., Eck, T.,
36 Smirnov, A., Dubovik, O., Slutsker, I., Tanre, D., Torres, O., Mishchenko, M., Geogdzhayev, I.,



- 1 Chu, D. A. and Kaufman, Y.: Monthly averages of aerosol properties: A Global comparison
2 among models, satellite data and AERONET ground data, *J. Geophys. Res*, 108, 3-1 to 3-42,
3 2003.
- 4 Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Bernsten, T., Berglen, T.,
5 Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D.,
6 Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Huang, P.,
7 Isaksen, I., Iversen, T., Koch, D., Kirkevåg, A., Kloster, S., Krol, M., Kristjansson, E., Lauer, A.,
8 Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari,
9 G., Reddy, S., Seland, O., Stier, P., Takemura, T. and Tie, X.: An AeroCom initial assessment –
10 optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–
11 1834, 2006.
- 12 Kokhanovsky, A. A.: *Aerosol Optics: Light absorption and scattering by particles in the atmosphere*,
13 Springer and Praxis Publishing, UK, pp. 6, 2008.
- 14 Koo, B., Pandis, S. N. and Ansari, A.: Integrated approaches to modeling the organic and inorganic
15 atmospheric aerosol components, *Atmos. Environ.*, 37, 4757–4768, 2003.
- 16 Kulmala, M., Asmi, A., Lappalainen, H. K., Baltensperger, U., Brenguier, J.-L., Facchini, M. C.,
17 Hansson, H. C., Hov, O., O'Dowd, C. D., Pöschl, U., Wiedensohler, A., Boers, R., Boucher,
18 O., de Leeuw, G., Denier van der Gon, H. A. C., Feichter, J., Krejci, R., Laj, P., Lihavainen, H.,
19 Lohmann, U., McFiggans, G., Mentel T., Pilinis, C., Riipinen, I., Schulz, M., Stohl, A.,
20 Swietlicki, E., Vignati, E., Alves, C., Amann, M., Ammann, M., Arabas S., Artaxo, P., Baars, H.,
21 Beddows, D. C. S., Bergström R., Beukes, J. P., Bilde, M., Burkhardt, J. F., Canonaco, F., Clegg,
22 S. L., Coe, H., Crumeyrolle, S., D'Anna, B., Decesari, S., Gilardoni, S., Fischer, M., Fjaeraa, A.
23 M., Fountoukis, C., George, C., Gomes, L., Halloran, P., Hamburger, T., Harrison, R. M.
24 Herrmann, H., Hoffmann, T., Hoose, C., Hu, M., Hyvärinen, A., Hörrak, U., Iinuma, Y., Iversen,
25 T., Josipovic, M., Kanakidou, M., Kiendler-Scharr, A., Kirkevåg, A., Kiss, G., Klimont, Z.,
26 Kolmonen, P., Komppula, M., Kristjánsson, J.-E., Laakso, L., Laaksonen, A., Labonnote, L.,
27 Lanz, V. A., Lehtinen, K. E. J., Rizzo, L. V., Makkonen, R., Manninen, H. E. McMeeking,
28 G, Merikanto, J., Minikin, A., Mirme, S., Morgan, W. T., Nemitz, E., O'Donnell, D., Panwar, T.
29 S., Pawlowska, H., Petzold, A., Pienaar, J. J., Pio, C., Plass-Duelmer, C., Prévôt, A. S. H., Pryor,
30 S., Reddington, C. L., Roberts, G., Rosenfeld, D., Schwarz, J., Seland, O., Sellegri, K., Shen, X.
31 J., Shiraiwa, M., Siebert, H., Sierau, B., Simpson, D., Sun, J., Y., Topping, D., Tunved,
32 P., Vaattovaara, P., Vakkari, V., Veefkind, J. P., Visschedijk, A., Vuollekoski, H., Vuolo,
33 R., Wehner, B., Wildt, J., Woodward, S., Worsnop, D. R., van Zadelhoff J.-L., Zardini, A., A.,
34 Zhang, K., van Zyl, P. G., Kerminen, V. M., Carslaw, K. S. and Pandis, S. N.: General
35 Overview: EUCAARI-integrating aerosol research from nano to global scales, *Atmos. Chem.*
36 *Phys.*, 11, 13061-13143, 2011.



- 1 Lee, Y. H. and Adams, J. P.: Evaluation of aerosol distributions in the GISS-TOMAS global aerosol
2 microphysics model with remote-sensing applications, *Atmosph. Chem. Phys.*, 10, 2129-2144,
3 2010.
- 4 Levy, R. C., Remer, L. A. and Dubovic, O.: Global aerosol optical properties and application to
5 Moderate Resolution Imaging Spectroradiometer aerosol retrieval over land, *J. Geophys Res.*,
6 112, D13210, 2007a.
- 7 Levy, R. C., Remer, L. A., Mattoo, S., Vermote, E. F. and Kaufman, Y. J.: Second-generation
8 operational algorithm: Retrieval of aerosol properties over land from inversion of Moderate
9 Resolution Imaging Spectroradiometer spectral reflectance, *J. Geophys. Res.*, 112, D13211,
10 2007b.
- 11 Levy, R. C., Remer, L. A., Tanre, D., Mattoo, S. and Kaufman, Y. J.: Algorithm for remote sensing of
12 tropospheric aerosol over dark targets from MODIS: Collections 005 and 051: Revision 2.
13 Product ID: MOD04/MYD04, Feb 2009.
- 14 Levy, R. C., Remer, L. A., Kleidman, R. G., Mattoo, S., Ichoku, C., Kahn, R. and Eck, T. F.: Global
15 evaluation of the Collection 5 MODIS dark-target aerosol products over land, *Atmos. Chem.*
16 *Phys.*, 10, 10399-10420, 2010.
- 17 Meij, A., Wagner, S., Cuvelier, C., Dentener, F., Gobron, N., Thunis, P. and Schaap, M.: Model
18 evaluation and scale issues in chemical and optical aerosol properties over the greater Milan area
19 (Italy) for June 2001, *Atmosph. Res.*, 85, 243-267, 2007.
- 20 Meij, A., Pozzer, A., Pringle, K. J., Tost, H. and Lelieveld, J.: EMAC model evaluation and analysis of
21 atmospheric aerosol properties and distribution with a focus on the Mediterranean region,
22 *Atmosph. Res.*, 114-115, 38-69, 2012.
- 23 Murphy, B.N. and Pandis, S. N.: Simulating the formation of semivolatile primary and secondary
24 organic aerosol in a regional chemical transport model, *Environ. Scienc. & Technol.*, 43, 4722-
25 4728, 2009.
- 26 Myhre, G., Berglen, T. F., Hoyle, C. R., Christopher, S. A., Coe, H., Crosier, J., Formenti, P., Haywood,
27 J. M., Johnsrud, M., Jones, T. A., Loeb, N., Osborne, S. and Remer, L. A.: Modelling of chemical
28 and physical aerosol properties during the ADRIEX aerosol campaign, *Quarterly J. Royal Meteor.*
29 *Societ.*, 135, 53-66, 2009.
- 30 NASA Global Aerosol Climatology Project (GACP) 2006: <http://gacp.giss.nasa.gov/>
- 31 Nenes, A., Pandis, S. N. and Pilinis, C.: ISORROPIA: a new thermodynamic equilibrium model for
32 multiphase multicomponent inorganic aerosols, *Aquat. Geochem.*, 4, 123-152, 1998.
- 33 Nessler, R., Weingartner, E. and Baltensperger, U.: Adaptation of dry nephelometer measurements to
34 ambient conditions at the Jungfraujoch, *Environ. Sci. Technol.*, 39, 2219-2228, 2005.
- 35 O' Dowd, C. D., Langmann, B., Varghese, S., Scannel, C., Ceburnis, D. and Facchini, M. C.: A
36 combined organic-inorganic sea-spray source function, *Geophys. Res. Lett.*, 35, 1-5, 2008.



- 1 Park, R. S., Song, C. H., Han, K. M., Park, M. E., Lee, S.-S., Kim, S.-B. and Shimizu, A.: A study on
2 the aerosol optical properties over East Asia using a combination of CMAQ-simulated aerosol
3 optical properties and remote-sensing data via a data assimilation technique, *Atmos. Chem. Phys.*,
4 11, 12275-12296, doi:10.5194/acp-11-12275-2011, 2011.
- 5 Pilinis, C. and Pandis, S. N.: Physical, Chemical and Optical Properties of Atmospheric Aerosols. The
6 handbook of Environmental Chemistry, Airborn Particulate Matter, Springer, pp. 99 –124, 1995.
- 7 Pozzer, A., Meij, A., Pringle, K. J., Tost, H., Doering, U. M., Van, Aardenen J. and Lelieveld, J.:
8 Distributions and regional budgets of aerosols and their precursors simulated with the EMAC
9 chemistry-climate model, *Atmosph. Chem. Phys.*, 12, 961-987, 2012.
- 10 Remer, L. A., Kaufman, Y. J., Tanre, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R. R., Ichoku C.,
11 Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E. and Holben, B. N.: The MODIS aerosol
12 algorithm, products, and validation, *J. Atmos. Sci.*, 62, 947-973, 2005.
- 13 Roy, B., Matur, R., Gilliland, A. and Howard, S.: A comparison of CMAQ-based aerosol properties
14 with IMPROVE, MODIS and AERONET data, *J. Geophys. Res.*, 112, 1-17, 2007.
- 15 Salomonson, V. V., Barnes, W., Maymon, P. W. and Montgomery, H. E.: MODIS: Advanced Facility
16 Instrument for Studies of the Earth as a System, *IEEE Trans. Geosc. Remote Sens.*, 27, 145-153,
17 1989.
- 18 Shi, Y., Zhang, J., Reid, J. S., Holben, B., Hyer, E. J. and Curtis, C.: An analysis of the collection 5
19 MODIS over-ocean aerosol optical depth product for its implication in aerosol assimilation,
20 *Atmos. Chem. Phys.*, 11, 557-565, 2011.
- 21 Schutgens, N. A. J., Nakata, M. and Nakajima, T.: Validation and empirical correction of MODIS AOT
22 and AE over ocean, *Atmos. Meas. Tech.*, 6, 2455-2475, 2013.
- 23 Schuster, G. L., Dubovik, O. and Holben, B. N.: Angstrom exponent and bimodal aerosol size
24 distributions, *J. Geophys. Res.*, 111, 1-14, 2006.
- 25 Seinfeld, J. H and Pandis, S. N.: Atmospheric Chemistry and Physics: From Air Pollution to Climate
26 Change, John Wiley and Sons, U.S.A., 2006.
- 27 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X., Wang,
28 W. and Powers, J. G.: A Description of the Advanced Research WRF Version 3, NCAR Technical
29 Note, available at: http://www.mmm.ucar.edu/wrf/users/docs/arw_v3.pdf, 2008.
- 30 Sofiev, M., Vankevich, R., Lotjonen, M., Prank, M., Petukhov, V., Ermakova, T., Koskinen, J. and
31 Kukkonen, J.: An operational system for the assimilation of the satellite information on wild-land
32 fires for the needs of air quality modeling and forecasting, *Atmosph. Chem. Phys.*, 9, 6833-6847,
33 2009.
- 34 Tombette, M., Chazette, P., Sportisse, B. and Roustan, Y.: Simulation of aerosol optical properties over
35 Europe with a 3-D size-resolved aerosol model: comparisons with AERONET data, *Atmosph.*
36 *Chem. Phys.*, 8, 7115-7132, 2008.



1 Yu, F., Luo, G. and Ma, X.: Regional and global modeling of aerosol optical properties with a size,
2 composition and mixing state resolved particle microphysics model, *Atmosph. Chem. Phys.*, 12,
3 5719-5736, 2012.

4 Vijayaraghavan, K., Snell, H. E. and Seigner, C.: Practical aspects of using satellite data in air quality
5 modeling, *Environ. Scienc. Techn.*, 42, 8187-8192, 2008.

6 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami,
7 A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., De-Carlo, P., Salcedo,
8 D., Onasch, T. B., Jayne, J. T., Miyoshi, T., Shimo, A., Hatakeyama, N., Takegawa, N., Kondo,
9 Y., Schneider, J., Drewnick, F., Weimer, S., Demerjian, K. L., Williams, P. I., Bower, K. N.,
10 Bahreini, R., Cottrell, L., Griffin, R. J., Rautianen, J. and Worsnop, D. R.: Ubiquity and
11 dominance of oxygenated species in organic aerosols in anthropogenically influenced Northern
12 Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007GL029979, 2007.

13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36



1 **Table 1.** Error metrics for the evaluation of PMCAMx against AERONET monthly mean AODs.

Region	Number of AERONET stations	Mean AERONET AOD	Mean PMCAMx AOD	Mean Error	Mean Bias	Fractional Error	Fractional Bias
UK/Ireland	1	0.25	0.24	0.01	-0.01	0.04	-0.04
Central Europe	25	0.16	0.17	0.03	0.02	0.22	0.12
North Europe	4	0.08	0.12	0.04	0.04	0.36	0.36
Spain and Portugal	8	0.11	0.12	0.02	0.01	0.20	0.05
East Europe	2	0.11	0.14	0.03	0.03	0.24	0.24
Balkans	2	0.21	0.15	0.06	-0.06	0.33	-0.33
Russia, Belarus, and Ukraine	5	0.15	0.14	0.02	-0.02	0.16	-0.11
Turkey and Northern Africa	2	0.17	0.12	0.05	-0.05	0.30	-0.30
Mediterranean Sea	-	-	-	-	-	-	-
North Atlantic Ocean	1	0.11	0.16	0.05	0.05	0.37	0.37
South Atlantic Ocean	-	-	-	-	-	-	-
Black Sea	-	-	-	-	-	-	-
Domain	50	0.15	0.15	0.03	0.001	0.22	0.04

2
 3
 4

$$\text{Mean Error} = \frac{1}{N} \sum_{i=1}^N |P_i - O_i| \quad \text{Mean Bias} = \frac{1}{N} \sum_{i=1}^N (P_i - O_i)$$

5
 6
 7

$$\text{Fractional Error} = \frac{2}{N} \sum_{i=1}^N \frac{|P_i - O_i|}{P_i + O_i} \quad \text{Fractional bias} = \frac{2}{N} \sum_{i=1}^N \left(\frac{P_i - O_i}{P_i + O_i} \right)$$

8 where P_i are predicted values by PMCAMx, O_i the AERONET retrievals and N the number of stations.



1 **Table 2.** Error metrics for the evaluation of PMCAMx against MODIS monthly mean AODs.
 2

Region	Mean MODIS AOD	Mean PMCAMx AOD	Mean Error	Mean Bias	Fractional Error	Fractional Bias
UK and Ireland	0.21	0.23	0.04	0.02	0.22	0.14
Central Europe	0.16	0.17	0.05	0.01	0.30	0.13
North Europe	0.09	0.14	0.06	0.05	0.53	0.47
Spain and Portugal	0.14	0.12	0.04	-0.03	0.28	-0.15
East Europe	0.13	0.15	0.05	0.03	0.35	0.25
Balkans	0.19	0.14	0.05	-0.04	0.28	-0.24
Russia, Belarus, and Ukraine	0.13	0.12	0.04	0.01	0.30	-0.01
Turkey and Northern Africa	0.16	0.14	0.05	-0.03	0.31	-0.13
Mediterranean Sea	0.18	0.14	0.04	-0.04	0.25	-0.24
North Atlantic Ocean	0.17	0.14	0.04	-0.03	0.30	-0.21
South Atlantic Ocean	0.16	0.10	0.06	-0.06	0.45	-0.45
Black Sea	0.17	0.14	0.04	-0.03	0.22	-0.18
Domain	0.16	0.14	0.05	-0.02	0.33	-0.16

26



1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24

Table 3. Monthly predicted mean ground-level concentration in $\mu\text{g}/\text{m}^3$ of the major $\text{PM}_{2.5}$ components.

Region	SO_4^{2-}	OA	EC	Cl^-	Na^+	NH_4^+	NO_3^-	H_2O	Crustal
UK and Ireland	3.6	3.4	0.5	0.6	0.8	2.4	3.8	33.3	0.7
Central Europe	3.0	3.4	0.5	0.2	0.4	1.4	1.3	10.1	0.6
North Europe	2.2	2.3	0.2	0.2	0.4	0.9	0.6	5.1	0.4
Spain and Portugal	1.6	1.2	0.2	0.1	0.2	0.7	0.5	11.7	0.3
East Europe	2.9	3.1	0.4	0.1	0.4	1.2	0.8	9.2	0.6
Balkans	3.9	2.7	0.3	0.1	0.3	1.4	0.3	4.9	0.6
Russia, Belarus, and Ukraine	2.5	2.1	0.3	0.03	0.2	0.9	0.2	3.9	0.5
Turkey and Northern Africa	2.7	2.2	0.2	0.2	0.4	1.0	0.6	8.2	0.5
Mediterranean Sea	4.2	2.4	0.3	1.3	1.4	1.2	0.3	11.3	0.7
North Atlantic Ocean	2.1	1.8	0.2	1.0	1.0	1.0	1.0	24.2	0.4
South Atlantic Ocean	1.5	1.1	0.06	0.8	0.8	0.5	0.3	8.8	0.3
Black Sea	3.8	2.8	0.3	0.6	0.7	1.3	0.4	9.7	0.7
Domain	2.4	1.9	0.2	0.5	0.6	0.9	0.5	10	0.5



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28





Figure 1. Geographical distribution of the 50 AERONET stations used in the present study.



1
2
3



4
5
6
7
8
9
10
11
12
13

Figure 2. Geographical distribution of the number of available AOD retrievals from MODIS over Europe during May 2008. White color denotes no retrievals. Land is partitioned into 8 regions including the United Kingdom and Ireland, central Europe, North Europe, Spain and Portugal, East Europe, Balkans, Russia/Belarus/Ukraine, Turkey, and Northern Africa. The sea is partitioned into 4 regions: the Mediterranean, North Atlantic, South Atlantic, and Black Sea.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36





Figure 3. Monthly mean AODs from PMCAMx and MODIS ($QAC \geq 1$) during May 2008. White color denotes no AOD retrieval. A coarse particle rejection filter has been employed. The PMCAMx AODs correspond to the periods of the MODIS retrievals.



31 **Figure 4.** Difference of the PMCAMx from MODIS ($QAC \geq 1$) monthly mean AODs during May
32 2008. Positive means that PMCAMx overpredicts AOD compared to MODIS. There were not enough
33 dust-screened AOD retrievals for the model evaluation in the white areas.
34
35
36
37
38




Figure 5. a) Comparison of the PMCAMx predictions with MODIS ($QAC \geq 1$) monthly mean AODs. The different colors indicate density. The dashed red lines denote the ocean expected error envelope and the dotted lines denote the land envelope which describes MODIS AOD uncertainties with respect to AERONET. The solid red line is the 1:1 line. **b)** Comparison of the PMCAMx with AERONET monthly mean AODs. The PMCAMx values correspond to the periods of measurement for the 50 AERONET stations.




Figure 6. a) Box plots of the PMCAMx and MODIS ($QAC \geq 1$) monthly mean AODs for land. The central mark is the median, the edges of the box are the 25th and 75th percentiles, the whiskers extend to the extreme data points considered to be not outliers, and the outliers are plotted individually by the red marks. Points are drawn as outliers if they are larger than $Q3 + 1.5(Q3 - Q1)$ or smaller than $Q1 - 1.5(Q3 - Q1)$, where $Q1$ and $Q3$ are the first and third quartiles, respectively. **b)** Box plots of the PMCAMx and MODIS ($QAC \geq 1$) monthly mean AODs for water.