

Dear Dr Williams,

Please find attached our revised manuscript with changes from the original version highlighted in red. We found the reviewers' comments to be very helpful and have responded to these comments below.

-----

Reviewer 1: Dáithí Stone

**Reviewer's summary:**

This paper describes the experimental setup of an atmospheric modelling system for examination of extreme weather over the land territories of Australia and New Zealand in the context of anthropogenic climate change. It is well designed, well described in this paper, and various aspects of the output of the modelling system are adequately summarised. I recommend the paper for publication. I have some minor comments and suggested edits below, but I do not consider any of them to be required.

**1. Reviewer's comment (general):** You examine DJF and JJA values, and some SONDJF values. The onset/cessation seasons for temperature and (I think) rainfall occur during the SON and MAM seasons, and I believe extreme early/late onsets/cessations can be at least as important e.g. for water resources and agriculture. Have you done these analyses for those seasons and are you able to summarise them? It probably does not have to be in the sort of detail done for DJF and JJA, but could just highlight any cases where e.g. the model might happen to be rather late (as proxied by the mean during the onset).

*Authors' response:* This additional analysis has been completed and is included within the supplementary material (see new Figures S11–S13). Overall, the weather@home model is seen to adequately resolve the onset/cessation seasons for temperature and precipitation.

**2. Reviewer's comment (technical):** page 1, lines 6-7 "more robust estimates of uncertainty" than what? You are using a single modelling system, so I am not sure how you can e.g. robustly estimate the uncertainty due to approximations in model design.

*Authors' response:* The manuscript has been revised to address this point (see page 1, line 7).

**3. Reviewer's comment (technical):** page 1, lines 12-13. Where or how? I do not see the URL of any data portal listed in the document, for instance.

*Authors' response:* The relevant information for accessing the data has been added under the section *Data and code availability*. Members of the research community wishing to access the data are encouraged to contact the authors directly. An online data portal (with public URL) will become available in the future.

**4. Reviewer's comment (technical):** page 1, line 16. I suppose wildfire might be considered a "climate-related" process, but I usually think of it as an ecological process.

*Authors' response:* The lead author (Mitchell Black) is currently completing a PhD thesis that is using the weather@home ANZ framework to investigate the influence of anthropogenic climate change on wildfire risk. This material will be published in due course.

**5. Reviewer's comment (technical):** page 1, lines 16-20. Are there any reports that document such "loss" and "tasks"?

*Authors' response:* Citations have been included within the revised manuscript (see page 1 of revised manuscript, lines 17-19).

**6. Reviewer's comment (technical):** page 2, line 1. Reisinger et alii (2014) (IPCC AR5 WGII Ch25) might be an appropriate reference here, as it is in this chapter of the IPCC AR5 that trends in Australian/New Zealand climate are assessed specifically

*Authors' response:* This citation has been included within the revised manuscript (see page 2, line 3).

**7. Reviewer's comment (technical):** page 2, line 10 "Distinguishing between internal" -> "Distinguishing between the responses to internal"

*Authors' response:* Text revised accordingly (see page 2, line 12)

**8. Reviewer's comment (technical):** page 2, line 12. While the magnitude of the response is more or less the same at regional and global scales?

*Authors' response:* This sentence has been revised to allow the message to be clearer. (see page 2, lines 13-16).

**9. Reviewer's comment (technical):** page 2, lines 20-22 Are these "drivers" or mechanisms? For instance, atmospheric blocking over Australia is a manifestation of climate variability over Australia, not a driver thereof.

*Authors' response:* This sentence has been revised to remove this confusion between drivers and mechanisms (see page 2, lines 23-27).

**10. Reviewer's comment (technical):** page 2, line 31 "respond to anthropogenic" -> "respond to the absence of anthropogenic" This discussion concerns estimation of the counterfactual climate, correct?

*Authors' response:* This sentence has been revised accordingly (see page 2, line 35).

**11. Reviewer's comment (technical):** page 6, line 9 Might the observations have any deficiencies?

*Authors' response:* A sentence has been added to section *Observational datasets* (page 6, lines 11-13) to acknowledge potential limitations of the observational datasets used in this study. However, as indicated in the revised text, these datasets are nevertheless well regarded for use in model evaluation studies.

**13. Reviewer's comment (technical):** page 6, lines 10-11 How do DJF and JJA project onto the wet, dry, onset, and/or cessation seasons over these regions?

*Authors' response:* As indicated for point 1 above, additional analysis has been undertaken to examine the model's ability to resolve the onset/cessation seasons for temperature and precipitation. Overall, the model appears to adequately resolve this timing for each region. The supplementary material has been updated to include these additional figures and a brief summary of these results are included in the main body of manuscript. (see page 6, lines 22-23).

**14. Reviewer's comment (technical):** page 8, lines 3-4 This is not clear to me for Tmin. The uncertainty on the median should be  $\sim\sqrt{75}\sim 8.7$  times smaller than the range of the whiskers: are the respect Tmin median values larger than this? I cannot tell from the plot.

*Authors' response:* Figures 8-10 have been updated to make the box plots clearer.

**15. Reviewer's comment (technical):** page 8, line 27 Maybe a 360-day calendar?

*Authors' response:* Manuscript updated accordingly.

**16. Reviewer's comment (technical):** page 8, lines 32-33 How is "model uncertainty" estimated with just the single model?

*Authors' response:* This has been corrected to indicate 'sampling uncertainty', not 'model uncertainty'.

**17. Reviewer's comment (technical):** page 9, lines 4-5 I am not sure, the observations lie within the spread of the simulations. page 9, lines 9-10 The observations lie within the spread of the simulations for 11d, and are pretty much bang on for 11e.

*Authors' response:* Indeed, this section of text was incorrect as the observed values fell within the model spread. This section of text has been updated to remove these incorrect statements (see page 9, lines 17-22).

**18. Reviewer's comment (technical):** page 9, lines 30-31 Re the sea ice extent, does this make sense for the Antarctic, where the recent trend has been toward slightly larger extent?

*Authors' response:* This point was investigated during the beta-testing phase of the weather@home Australia-New Zealand experiment. During this testing counterfactual climates were simulated using: 1.) sea ice extent corresponding to the year of maximum sea ice extent in the Southern Hemisphere from the OSTIA records (1985–2014), and 2.) sea ice extent corresponding to the year of minimum sea ice extent in

the Southern Hemisphere. While not shown here, the results of these experiments identified that the choice of counterfactual sea ice extent had negligible impact on the resulting climates of Australia and New Zealand. Therefore, the weather@home Australia-New Zealand experiments presented in this manuscript used the maximum observed sea ice extent as a proxy for the counterfactual sea ice extent, for both hemispheres. This is consistent with the methodology of the existing weather@home projects – that is, weather@home Europe (Massey et al. 2015) and weather@home Pacific North-West (Mote et al. 2015).

**19. Reviewer's comment (technical):** page 9, lines 32-33 You did not mention when describing the simulations if anything is done concerning land use/cover change. Is this included and, if so, how is this treated in the counterfactual simulations?

*Authors' response:* The weather@home ANZ setup uses the MOSES1.0 land surface scheme. The surface type is fixed and is the same between the historical and counterfactual climate simulations. The manuscript has been revised to include this information (see page 4, paragraph 1).

**20. Reviewer's comment (technical):** page 10, line 18 "quantified" -> "characterised" I do not see any reason to believe that these 10 estimates can be assumed to be uniformly sampled from a plausible posterior distribution. Rather than producing a posterior distribution, I think you are testing robustness against plausible alternative estimates.

*Authors' response:* Manuscript updated accordingly.

**21. Reviewer's comment (technical):** page 10, lines 29-30 Where and/or how?

*Authors' response:* The relevant information for accessing the data has been added under section *Data and code availability* (page 12, paragraph 1).

**22. Reviewer's comment (technical):** "allows extreme events" -> "allows certain types of extreme weather events"

*Authors' response:* Manuscript updated accordingly.

-----

Reviewer 2: Andrew Ciavarella

**Reviewer's summary:**

This well written paper introduces a unique and valuable resource for event attribution over the Australia and New Zealand region, building on the success of equivalent systems focussing on different regions.

With regards reproducibility I have found some omissions in the model and experimental description that can be easily addressed. The system description is otherwise clear and well motivated.

A decent attempt to explore model uncertainty is made through the construction of 10 estimates of the counterfactual boundary conditions while the limitations of the system are also discussed. The basic validation conducted is probably sufficient given



that the studies that are performed with such a system generally require bespoke validation and are in fact already in print [Herring et al., 2015]. Altogether I found no issues that I consider major and so I would recommend this work for publication.

**1. Reviewer's comment (specific):** [p2] Use of phrase “internal ... climate forcings” While it is not unreasonable to refer to major modes of variability as forcing a regional climate akin to an external forcings it would be better to use a phrase like “internal climate variability”.

*Authors' response:* Manuscript updated accordingly

**2. Reviewer's comment (specific):** [p2.7, also p2.20,23] “chaotic natural variability” is referring to variability generated internal to the climate system while in attribution we normally retain the phrase “natural variability” to refer to externally forced variability (i.e. by solar and volcanic forcing). “chaotic internal variability” would be better and is probably adequately distinguished from the major modes of internally generated variability in the context of this sentence. Later [p2.30] “natural forcing” is used in the normal sense so there is clear room for confusion.

*Authors' response:* Manuscript updated accordingly

**3. Reviewer's comment (specific):** Page 2.19-24 emphasises the importance of several major modes of internal variability in addition to ENSO (SAM, position of storm tracks, blocking) which are important in the Australia / New Zealand region but only ENSO is addressed in the remainder of the paper. I am not asking that additional validation be done for these other factors but could the authors comment on the relative importance of ENSO with respect to these others? For e.g. some reference to the literature to give an idea of the share of variance explained in inter-seasonal or monthly regional means of the two variables presented? It appears Risbey et al., 2009 Figs. 15 & 16 (already cited by the authors) may contain enough to go on.

*Authors' response:* As Reviewer 1 correctly identified, the original body of text confused drivers of internal climate variability (e.g., ENSO) from manifestations of climate variability (e.g., blocking). The text was revised to address this point. As the text now stands, three drivers of climate variability are identified – ENSO, Southern Annular Mode and the Indian Ocean Dipole. While each of these three modes of variability have an important influence on Australian temperature and precipitation extremes (e.g., Risbey et al. 2009, Min et al. 2013), ENSO was examined in this manuscript as it is the mode of variability that has been examined in a number of event attribution studies to date (e.g., King et al. 2013; Lewis and Karoly 2013; Christidis et al. 2013) and there are a number of planned studies wanting to specifically investigate the role of ENSO (e.g., Black and Karoly, 2016; Karoly et al. 2016). For brevity, the analysis for IOD and SAM was not presented in the current manuscript but will feature in an upcoming publication. Analysis of blocking within the weather@home ANZ model is provided in the recent publication by Grose et al. (2015).

**4. Reviewer's comment (specific):** [p2.28, also p9.22, Fig. 13 caption] Use of “observed” climate in reference to historical climate. In a modelling context we would avoid using “observed” to refer to experiments with anthropogenic and natural

forcings present as these are simply not observations but simulations whose climate is intended to reproduce an observed climate, but which may not do so. I advise substituting this with “anthropogenically forced”, “historical” or similar. Elsewhere [p5.3] “historical ... climate scenario” is used.

*Authors' response:* The manuscript has been updated to replace ‘observed’ with ‘historical’

**5. Reviewer's comment (specific):** [p4.33] Further initial condition perturbations are applied by a range of start conditions with difference large scale circulation and soil moisture patterns. No details or references are given for the selection criteria or pattern generation. Please elaborate. Are the patterns physical / taken from a control?

*Authors' response:* For the weather@home ANZ experiment, each model simulation is initialised using a restart file created in a control simulation for the previous year. For each experiment, 100 unique restart files are created (each with a different atmospheric state and soil moisture pattern/profile).

For example, consider the weather@home experiment for the year 2015. For this experiment, the model is run 100 times for the preceding year (2014) and the resulting restart files from these experiments are used to initialise the 2015 simulations. Therefore, there are essentially 100 ‘groups’ of simulations created for the year 2015, with members of each ‘group’ containing the same restart file. For members within each ‘group’, slightly different initial condition perturbations are applied to the three-dimensional potential profile of the restart file (as described within the manuscript).

The manuscript has revised to clarify this (see page 5).

**6. Reviewer's comment (specific):** [p5.4] Lower boundary conditions are taken from a daily analysis product (OSTIA) while previous versions of the weather@home experimental setup [Massey et al., 2015, Mote et al., 2015] used interpolation from a monthly observational dataset (HadISST1). Was there a good reason for this new choice? For instance was it felt that the daily analysis provides more faithful sub-monthly variability than interpolation from monthly data?

*Authors' response:* All recent weather@home experiments have transitioned to using sea surface temperatures and sea ice extent from the OSTIA dataset (e.g., see Schaller et al. 2016, Mitchell et al. 2016). As the reviewer correctly identified, these daily fields provide a more faithful estimate of sub-monthly variability than is achieved from interpolation of monthly data.

**7. Reviewer's comment (specific):** [5.10] Halocarbon prescription: experimental progenitor [Massey et al. 2015] prescribe a single halocarbon value designed to give the radiative forcing corresponding to the presence of all (6 AR4 recommended) represented species. Is the same manner of prescription used in these experiments or are the AR5 individual species concentrations separately prescribed?

*Authors' response:* As per Massey et al. (2015), a single halocarbon value is used to give the radiative forcing corresponding to the presence of all represented species.

The manuscript has been updated to include this information (see page 5, paragraph 2).

**8. Reviewer's comment (specific):** [5.11] GHG concentrations and aerosol emissions. Could you be explicit about the concentrations that are prescribed post-2005? Specifically, is one of the RCP scenarios followed (for e.g. from <http://www.pik-potsdam.de/~mmalte/rcps/index.htm>)?

*Authors' response:* Post-2005, the greenhouse gas concentrations and aerosol emissions follow the RCP 8.5 scenario. The manuscript has been updated to include this information (see page 5, paragraph 2).

**9. Reviewer's comment (specific):** Section 2 does not mention if land use changes are prescribed. I can see (from Massey et al., 2015, section 2.2.4) that fractions of surface types are specified. Is this specification fixed or time dependent? Are the prescribed fraction of the natural simulations representative of preindustrial conditions? Section 2 I could not gather from this or Massey et al., 2015 what land surface scheme the models use.

*Authors' response:* The model uses the MOSES 1.0 land surface scheme with fixed surface type. There is not change in surface type between the historical and counterfactual climate scenarios. The manuscript has been updated to include this information (see page 4, paragraph 1).

**10. Reviewer's comment (specific):** Section 2 Spin down. Is there a spin down period allowed prior to creation of the experimental initial conditions or to the analysis period? Has any continued drift in climate variables such as soil moisture been seen over the 29 year historical experiment?

*Authors' response:* As indicated above, each model year is initialised from a restart file output from a model simulation for the previous year. Therefore, there is a 12-month spin-down period allowed. Continuous integration of the model over an extended period (1985–2014) has not revealed any continued drift in soil moisture.

**11. Reviewer's comment (specific):** [p6.9] After taking 29 year means “any differences between the obs. and model output may be interpreted as model deficiencies”. This is not strictly true as, for example, even the means of output from two members of the same model will still be subject to “standard error” which will decrease as  $1/\text{SQRT}(n)$  for  $n$  data points. In comparing the difference of two such means the errors will also add in quadrature. Nevertheless I would estimate that the discussed biases depicted in the figures 2 – 4 easily stand out from this level of noise. Also I acknowledge that p6.15 says that this can be regarded as “an indication of model bias”.

*Authors' response:* As the reviewer indicates, the biases depicted in Figures 2–4 easily stand out from sampling noise and can be regarded as an indication of model bias. The purpose of these figures is to provide a general overview of model performance; we feel that these images are able to portray this required information in a simple and adequate manner.

**12. Reviewer's comment (specific):** [p7.16] A comparison of time series variances, power spectra or quantile plots would provide a more objective measure of agreement than simply eye-balling that the obs. time series sits mostly inside the envelope of the models, especially given that the objective measure provided (correlation coefficients) will not allow an assessment of overall amplitude of the series. However given the intended brevity of the validation and later closer focus on daily data we can probably make do with this.

*Authors' response:* We thank the author for these suggestions. Given the brevity of the paper we too feel that the level of information currently presented is appropriate.

**13. Reviewer's comment (specific):** [p7.18] Precisely what series are the correlation coefficients between? Is it obs. and model median? This should go into caption to Fig. 5 also.

*Authors' response:* The correlation coefficients are calculated between the observation and model medians. The manuscript has been updated to include this, including the caption of Figure 5.

**14. Reviewer's comment (specific):** [Figures 5 - 7] Can you confirm that the p-values are for a one-sided test?

*Authors' response:* The p-values are for a two-sided test. The figure captions have been updated to include this information.

**15. Reviewer's comment (specific):** [p7.21] ENSO as driver of "natural climate variability" would better be "internal climate variability", again avoiding confusion with solar and volcanically forced variability.

*Authors' response:* Manuscript updated accordingly.

**16. Reviewer's comment (specific):** [p9.30] What year or period are the "pre-industrial" GHG, ozone & aerosol levels taken from?

*Authors' response:* The manuscript has been updated to include this information.

**17. Reviewer's comment (specific):** [p10.1] "cannot be known" could better be phrased as "cannot be observed", which is indisputable. We may dispute whether the counterfactual world is knowable.

*Authors' response:* Manuscript updated accordingly.

**18. Reviewer's comment (specific):** [p11.7-10] Use of an atmosphere only model is here portrayed as a limitation but the atmosphere only approach simply allows us to frame a different event attribution question than that provided by coupled experiments. Namely, we ask for the likelihood of an event subject to the lower boundary forcing provided by the precise phases of the various modes of oceanic (and

cryospheric) variability at the time of the event, which is not possible with a coupled model.

*Authors' response:* The reviewer has raised an important point – the atmosphere only model is not necessarily a limitation. The text has been revised accordingly (see page 11, paragraph 3).

**19. Reviewer's comment (technical corrections):** [p6.12] Could insert the word “daily” to be completely clear (sentence could be interpreted as maximum of seasonal average over 29 years, 75 members.). Ditto the caption to Figure 2.

*Authors' response:* Manuscript updated accordingly.

**20. Reviewer's comment (technical corrections):** [p7.14] “individual years” would better be phrased as “specific years”.

*Authors' response:* Manuscript updated accordingly.

**21. Reviewer's comment (technical corrections):** [p9.26] Insert word “lower” before “boundary conditions” to distinguish from lateral.

*Authors' response:* Manuscript updated accordingly.

**22. Reviewer's comment (technical corrections):** [p9.32] “boundary conditions common to both” is incorrect here, should be “forcings common to both”.

*Authors' response:* Manuscript updated accordingly.

**23. Reviewer's comment (technical corrections):** [p11.7] Unnecessary comma after “weather@home”

*Authors' response:* Manuscript updated accordingly.

**24. Reviewer's comment (technical corrections):** [Fig S1] “summertime” should be “wintertime” if genuinely June – August.

*Authors' response:* Manuscript updated accordingly.

**25. Reviewer's comment (technical corrections):** [p14.19 and elsewhere] Massey et al. “2014” should be “2015”.

*Authors' response:* Manuscript updated accordingly.

## References

- Black, M.T. and Karoly, D.J. (2016). Climate change was an important driver of southern Australia's warmest October on record [in “Explaining Extreme Events of 2015 from a Climate Perspective”], *Bulletin of the American Meteorological Society*, under review.
- Christidis, N et al. (2013). An attribution study of the heavy rainfall over eastern Australia in March 2012 [in “Explaining Extreme Events of 2012 from a Climate Perspective”], *Bulletin of the American Meteorological Society*, 94, S58-S61.
- Grose, M. R. et al. (2015) Attribution of exceptional mean sea level pressure anomalies south of Australia in August 2014 [in “Explaining Extreme Events of 2014 from a Climate Perspective”], *Bulletin of the American Meteorological Society*, 96, S158–S162.
- Karoly, D.J. et al. (2016). The roles of climate change and El Niño in the record low rainfall in October 2015 in Tasmania, Australia [in “Explaining Extreme Events of 2015 from a Climate Perspective”], *Bulletin of the American Meteorological Society*, under review.
- King, A et al. (2013). Limited Evidence of Anthropogenic Influence on the 2011-12 Extreme Rainfall over Southeast Australia [in “Explaining Extreme Events of 2012 from a Climate Perspective”], *Bulletin of the American Meteorological Society*, 94, S55-S58.
- Lewis, S and Karoly, D.J (2013). Anthropogenic contributions to Australia's record summer temperatures of 2013, *Geophysical Research Letters*, 40, 3705-3709.
- Mitchell, D. et al. (2016). Attributing human mortality during extreme heat waves to anthropogenic climate change. *Environmental Research Letters*, doi: 10.1088/1748-9326/11/7/074006
- Schaller, N. et al. (2016). The human influence on climate in the winter 2013/2014 floods in southern England. *Nature Climate Change*, doi: 10.1038/NCLIMATE2927

# The weather@home regional climate modelling project for Australia and New Zealand

Mitchell T. Black<sup>1</sup>, David J. Karoly<sup>1</sup>, Suzanne M. Rosier<sup>2</sup>, Sam M. Dean<sup>2</sup>, Andrew D. King<sup>1</sup>, Neil R. Massey<sup>3</sup>, Sarah N. Sparrow<sup>3,4</sup>, Andy Bowery<sup>4</sup>, David Wallom<sup>4</sup>, Richard G. Jones<sup>2,5</sup>, Friederike E. L. Otto<sup>3</sup>, and Myles R. Allen<sup>3</sup>

<sup>1</sup>School of Earth Sciences and ARC Centre of Excellence for Climate System Science, The University of Melbourne, Melbourne, Australia.

<sup>2</sup>National Institute of Water and Atmospheric Research, Wellington, New Zealand

<sup>3</sup>Environmental Change Institute, Oxford University, Oxford, United Kingdom

<sup>4</sup>Oxford e-Research Centre, Oxford University, Oxford, United Kingdom

<sup>5</sup>Met Office Hadley Centre, Exeter, United Kingdom

*Correspondence to:* M.T. Black (mtblack@student.unimelb.edu.au)

**Abstract.** A new climate modelling project has been developed for regional climate simulation and the attribution of weather and climate extremes over Australia and New Zealand. The project, known as *weather@home Australia-New Zealand*, uses public volunteers' home computers to run a moderate-resolution global atmospheric model with a nested regional model over the Australasian region. By harnessing the aggregated computing power of home computers, weather@home is able to generate an unprecedented number of simulations of possible weather under various climate scenarios. This combination of large ensemble sizes with high spatial resolution allows extreme events to be examined with ~~more robust estimates of uncertainty~~ **well-constrained estimates of sampling uncertainty**. This paper provides an overview of the weather@home Australia-New Zealand project, including initial evaluation of the regional model performance. The model is seen to be capable of resolving many climate features that are important for the Australian and New Zealand regions, including the influence of El Niño-Southern Oscillation on driving natural climate variability. To date, 75 model simulations of the ~~observed~~ **historical** climate have been successfully integrated over the period 1985–2014 in a time-slice manner. In addition, multi-thousand member ensembles have also been generated for the years 2013, 2014 and 2015 under climate scenarios with and without the effect of human influences. All data generated by the project is freely available to the broader research community.

## 1 Introduction

Extreme weather and climate-related events often have a serious impact on our economy, environment and society. This is particularly true in Australia and New Zealand where recurring heat waves, floods, droughts and wildfires have resulted in the loss of life, property and livelihoods (e.g., Steffen et al., 2013). In the aftermath of these events the scientific community is often faced with the task of quantifying the link to different causal factors, including human-induced climate change (e.g., Arblaster et al., 2015). The delivery of such information in a timely, clear and reliable manner is an ongoing challenge. Therefore,

developing a capacity to research extremes and understand their causes continues to be crucial for predicting and managing their impacts.

There is clear evidence that the climate has changed as a result of human influence (Stocker et al., 2014; Reisinger et al., 2014) and that some aspects of extremes have changed across the globe as a result (Seneviratne et al., 2012). However, this does not imply that the occurrence of every recently observed extreme weather or climate-related event was the result of human influence on the climate system, as such events may still have occurred (however unlikely) in the absence of such an influence (e.g., Stott et al., 2013). This can be understood by recognising that our climate is a complex, chaotic system that is influenced by internal climate variability and external forcings. Processes that generate internal climate variability include atmosphere-ocean teleconnections, such as El Niño-Southern Oscillation (ENSO), as well as chaotic natural internal variability. Meanwhile, external forcings of climate can be either natural, such as explosive volcanic eruptions, or anthropogenic, such as greenhouse gas emissions from the burning of fossil fuels.

Distinguishing between the responses to internal and external climate forcings becomes increasingly difficult when moving from global to regional scales, due to a lower signal-to-noise ratio (Károly and Wu, 2005; Stott et al., 2011). ~~That is to say, the noise associated with internal climate variability is greater at regional scales than at the global scale. That is to say, analysis on smaller spatial scales offers less opportunity to reduce the magnitude of natural variability through spatial averaging or other techniques.~~ This is particularly true for Australia, which is recognised as having one of the most variable climates in the world (e.g., Nicholls et al., 1997). The variability in Australia's climate is driven by a number of factors, in particular the year-to-year variations in sea surface temperatures in both the Pacific and Indian Oceans (Risbey et al., 2009). ENSO represents the variations in sea-surface temperatures and atmospheric patterns across the Pacific Ocean, with warm (El Niño) conditions producing below-average rainfall, above-average temperatures and often drought over much of northern and eastern Australia (McBride and Nicholls, 1983; Holland, 1986; Jones and Trewin, 2000). The reverse is true during cool (La Niña) conditions. Although New Zealand's climate is not usually affected as strongly by ENSO as are parts of Australia, there is nevertheless a significant influence (e.g., Gordon, 1986). In addition to ENSO there are a number of other drivers of natural internal climate variability for Australia and New Zealand, including the Southern Annular Mode (Hendon et al., 2007), ~~variations in storm tracks (Frederiksen and Frederiksen, 2007) and atmospheric blocking (Pook and Gibson, 1999)~~ and the Indian Ocean Dipole (Saji et al., 1999). Therefore, any assessment of extreme weather and climate-related events needs to consider the interplay of both natural internal climate variability and forced external changes, such as the warming effect caused by increased greenhouse gas emissions.

In light of this challenge, an emerging field of climate science (known as event attribution) is seeking to quantify how the risk of weather and climate-related extremes has changed as a consequence of particular forcings acting on the climate system (Allen, 2003; Stott et al., 2004; Pall et al., 2011; NASEM, 2016). This is typically achieved by comparing the probability of such events under the current (observed/historical) climate against that for counterfactual worlds in which particular forcing factors (such as human-induced climate change) are absent. We of course are unable to observe a world in which either anthropogenic or natural forcing is absent, therefore physically based climate models are required to estimate how the climate would respond to the absence of anthropogenic and natural forcings (Hegerl and Zwiers, 2011).



Undertaking event attribution studies of extreme weather events is typically restricted by two important modelling requirements: ensemble size and model resolution. Extreme weather events are, by definition, rare and therefore very large ensembles of climate model simulations are needed in order to study the event with a high degree of confidence. Meanwhile, as many extreme events occur at a regional or local scale, the model must have sufficient resolution to realistically capture the event.

5 Due to these requirements, such an undertaking would be computationally expensive and typically beyond the capability of conventional computing resources. However, these demands may be met through the aggregated power of distributed computing projects. Proposed by Allen (1999) and launched in 2003, *climateprediction.net* became the largest climate modelling experiment to date by running climate models on volunteers' home computers. While the project was originally focussed on running low-resolution global coupled atmosphere-ocean (Frame et al., 2009) and medium-resolution atmosphere-only models

10 (Pall et al., 2011), a recent advancement (known as *weather@home*) involves running the global atmosphere-only model with a nested higher-resolution regional model to generate very large ensembles of model simulations (Massey et al., 2015). This regional model configuration has been implemented and evaluated over Europe (Massey et al., 2015) and the Western United States (Li et al., 2015; Mote et al., 2015) and successfully used in a number of event attribution studies (e.g., Peterson et al., 2013; Herring et al., 2014, 2015; Schaller et al., 2016). Given the success of these existing *weather@home* regional climate

15 modelling projects it was decided to implement a regional configuration over Australia and New Zealand.

The primary purpose of this paper is to provide a description and basic evaluation of the *weather@home* Australia-New Zealand modelling setup. This is achieved by comparing the regional model output with observations from the recent past over regions of Australia and New Zealand. Because the modelling setup is intended to be used for event attribution studies, particular focus is given to an assessment of how well the model represents 1) mean spatial fields and interannual climate

20 variability, 2) regional teleconnections to ENSO and 3) the distribution of daily variables at regional and local scales. For the purpose of this study we have restricted our analysis to only consider temperature and precipitation as these are the variables most commonly assessed in event attribution studies.

The remainder of this paper is structured as follows: Section 2 describes the model setup and summarises the experimental design for the representation of the ~~observed~~historical climate, while Section 3 provides details on the evaluation of the system.

25 As a thorough comparison with observations is beyond the scope of this paper, we provide some illustrative comparisons of both temporal and spatial patterns. Section 4 describes how the counterfactual climate scenarios are constructed for the purpose of undertaking event attribution studies. The main conclusions are given in Section 5, including plans for future improvements.

## 2 Model description

*Weather@home* Australia-New Zealand uses the the Hadley Centre Atmospheric General Circulation Model 3P (HadAM3P; Massey et al., 2015) with an embedded regional model (HadRM3P; Jones et al., 2004) over the Australasian CORDEX domain (Figure 1). The HadAM3P/HadRM3P model formulation is based on the atmospheric component of the HadCM3 general circulation model (Gordon et al., 2000) with a number of improvements with respect to the calculation of clouds and convection, and a more realistic coupling of vegetated surfaces with the soil (Massey et al., 2015). HadAM3P/HadRM3P is a grid-point

30

model which solves equations of motion, radiative transfer and dynamics explicitly on the same scale as the grid. HadAM3P is integrated with a 15 minute timestep, has 19 vertical levels and has a regular latitude-longitude grid (1.25° longitude by 1.875° latitude) with regular poles. HadRM3P has a 5 minute timestep, has 19 vertical levels and uses a rotated grid (0.44° longitude by 0.44° latitude) with an artificial north pole at 60.31° N, 141.38° E for the Australia-New Zealand configuration.

5 This allows the region of interest to lie about the equator of the rotated grid, thus ensuring that each grid box in the nested region has approximately the same area. HadAM3P/HadRM3P are run in an interleaved manner: HadAM3P first runs for a full model day, providing the lateral boundary conditions to HadRM3P, which then also runs for one full model day. The coupling is strictly one-way, meaning that there is no feedback from the regional model to the global model. There is a four-point buffer zone around the perimeter of the regional model, where the lateral boundary conditions are relaxed to values temporarily

10 interpolated from 6-hourly output from HadAM3P. **The land surface scheme incorporated within the model is MOSES 1.0 (Met Office Surface Exchange Scheme, Cox et al. 1999), with fixed surface types (one vegetation type and one soil type per grid box).** Further details of the HadAM3P/HadRM3P configuration are provided by Massey et al. (2015), but with the European region replacing the Australasian region.

Weather@home is able to generate very large ensembles of climate model simulations by harnessing spare CPU time on

15 a network of volunteers' personal computers. This distributed computing capacity is made possible by the Berkeley Open Infrastructure for Network Computing (BOINC; Anderson, 2004) open source infrastructure. Each volunteer signs up for the weather@home project via the BOINC client software, which automatically downloads the climate model setup to the volunteer's computer. Individual workunits are then received from the BOINC server and run when the computer is idle. The workunit contains all necessary configuration inputs needed by the climate model to run the experiment for one model

20 year (December–November), under a specific climate scenario. After the completion of each model month the output is post-processed to retain only a selection of key meteorological variables. This is required in order to minimise file size for data transfer and storage. A complete listing of these output variables is provided as supplementary material. Following this post-processing stage the final results are returned to a server hosted at the Tasmanian Partnership for Advanced Computing in Hobart, Australia. On average, it takes a standard home computer around 4–5 days to integrate over the model year. At the

25 completion of the model year an additional file (the restart file) is returned that represents the final state of the atmosphere. This final state can then be incorporated as the initial conditions for a new workunit describing the next year of the climate scenario. Therefore, this allows the system to run for a year at a time, in a time-slice manner, to generate an extended timeseries of climate model integrations. As this resubmission process is not automated, the generation of these continuous model runs is somewhat restricted by the need for project scientists to manage restart files and work unit regeneration. Therefore, typical

30 event attribution studies will only generate multi-thousand ensemble members for a single model year of interest.

In order to represent the range of internal variability that is possible with the model, a perturbation is applied to the initial conditions of each workunit. These perturbations are applied to the global climate model in the form of slight changes to the three-dimensional potential temperature field. The initial condition perturbations were generated by calculating the next-day differences within a 1-year integration of the global model and then multiplying by five global scaling factors (1.1, 1.2, 1.3, 1.4

35 and 1.6) (see Massey et al. (2015) for details). This resulted in the generation of 1740 different initial condition perturbations.

In the case of an extended (multi-year) model experiment, these perturbations are only applied to the initial condition of the first model year; no perturbation is applied thereafter so as to allow for the continuous integration of the model under its specific climate scenario. Further initial condition perturbations are also applied to the first year of the model integration using a range of starting conditions with different large-scale circulations and soil moisture amounts. **That is to say, sets of model simulations are initialised using 100 different restart files taken from control simulations for the preceding year.** Although all of the initial condition perturbations are only applied to the global model, they immediately affect the regional simulations through the previously described transfer of lateral boundary conditions at the end of the first global model day.

As HadAM3P and HadRM3P are both atmosphere-only models, they require specified forcings at the boundary between the atmosphere and ocean. These lower boundary conditions come in the form of prescribed sea surface temperature (SST) and sea ice fraction (SIF) fields. For the historical 1985–2014 climate scenario used in this paper for the purpose of model evaluation, both the SST and SIF fields were sourced from the UK Met Office Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) dataset (Donlon et al., 2012). OSTIA provides global, daily fields with a spatial resolution of  $0.05^\circ$  latitude  $\times$   $0.05^\circ$  longitude. In order for these fields to be defined per grid box of the global climate model, they are regridded to the HadAM3P resolution of  $1.875^\circ$  latitude  $\times$   $1.25^\circ$  longitude using an area-weighted averaging method. Any discrepancy between the HadAM3P and OSTIA land-sea masks is resolved by taking the mean of surrounding ocean grid points. In addition to these lower boundary conditions, the model also requires the atmospheric composition to be specified. The concentrations of greenhouse gases ( $\text{CO}_2$ ,  $\text{CH}_4$  and  $\text{N}_2\text{O}$ ), ozone, halocarbons, sulphur species and solar anomalies are all prescribed to follow the recommendations outlined by the Coupled Model Intercomparison Project Phase 5 (CMIP5; Taylor et al., 2012). **The halocarbon gases (CFC113, CFC11, CFC12, HCFC22, HFC124 and HFC134A) are represented as a single value per timepoint in the timeseries, which produces the equivalent radiative forcing as if all six gases were modelled. Post-2005, the greenhouse gas concentrations and aerosol emissions follow the RCP 8.5 scenario.** An overview of the model boundary conditions for the counterfactual climate scenarios is presented in Section 4.

### 3 Model evaluation

In order to establish how well weather@home represents the ~~observed~~ **historical** climate over Australia and New Zealand, we generate 75 model simulations for each year over the period December 1985 to November 2014. While the modelling setup is capable of generating much larger ensemble sizes, this would be unnecessary for characterising the climatology and overall distribution of climate variables over a 29-year period. Therefore, computing resources were directed towards generating very large ensembles of climate simulations for the individual years of 2013, 2014 and 2015, for subsequent use in event attribution studies that are beyond the scope of the current paper.

The spatial fields of the regional model output are separated into six regions (Figure 1) for subsequent examination. Australia has been broken up into five established regions based on distinct climatic zones: northern Australia (NAUS), central Australia (CAUS), eastern Australia (EAUS), southwest Australia (SWAUS) and southeast Australia (SEAUS) (see CSIRO and Bureau of Meteorology (2015) for details). Meanwhile, the North and South Islands of New Zealand (NZ) are treated as a single region.

While each of the six regions could have been further broken down for sub-regional detail, it would be impractical to present such a mass of information here.

### 3.1 Observational datasets

Evaluation of weather@home is undertaken by comparing the regional model output to two observational data sets: for evaluation over Australia we use the Australian Water and Availability Project dataset (AWAP; Jones et al., 2009) and for New Zealand we use the Virtual Climate Station Network dataset (VCSN; Tait et al., 2006). The AWAP dataset provides daily and monthly gridded fields of rainfall and temperature extending back to 1911 on a  $0.05^\circ \times 0.05^\circ$  grid, and is highly regarded for studying trends and variability over Australia (e.g., Risbey et al., 2013; Min et al., 2013; King et al., 2013; Perkins and Alexander, 2013). For the purpose of this study we have masked the AWAP data over inland regions of Australia where there is low station density. Analogous to AWAP, VCSN provides high resolution ( $0.05^\circ$  latitude  $\times$   $0.05^\circ$  longitude) estimates of daily rainfall and temperature over New Zealand extending back to 1972. **While these observational datasets may be subject to uncertainties in and of themselves, they have nevertheless been found to be appropriate for use in model evaluation studies (e.g., Bennett et al., 2014; Moise et al., 2015).**

In order to compare model output and observations, a remapping of the observational datasets onto the HadRM3P model grid is required. For temperature this is achieved using bilinear interpolation. For precipitation, a conservative remapping scheme is used to ensure that the total amount of precipitation in the remapped data is the same as in the original data.

### 3.2 Climatological mean fields and inter-annual variability

We first examine weather@home's ability to correctly represent seasonal mean fields of temperature and precipitation. By averaging over the 29-year period we are attempting to reduce the internal atmospheric model variability about the mean state. Therefore, any differences between the observations and model output may be interpreted as model deficiencies. We have separated our analysis into seasons and for brevity are only showing results for the Austral summer (December–February) and winter (June–August). **While not shown here, the weather@home model is able to adequately resolve the onset/cessation seasons for temperature and precipitation (see supplementary Figures S11–S13).**

The spatial fields of seasonal average **daily** maximum temperature ( $T_{\max}$ ) and minimum temperature ( $T_{\min}$ ) are shown in Figure 2 and Figure 3, respectively. In each of these figures, the top panels show the mean field from HadRM3P, averaged over the 75 ensemble members for the 29-year period, while the middle panels show the mean for the observational datasets. The bottom panels show the difference between the model and observations and can be interpreted as an indication of model bias. Overall, the model is able to capture the large-scale spatial patterns of temperature very well, including the regions of warmest temperature over northern parts of Australia and the persistently cooler temperatures over southeast Australia and New Zealand, associated with topography. For  $T_{\max}$ , HadRM3P is capable of representing mean summertime values to within  $\pm 1^\circ$  C over most parts of Australia and over the North Island of New Zealand (Figure 2e). In winter, the model underestimates  $T_{\max}$  at almost every land grid point across the model domain (Figure 2f). For  $T_{\min}$ , the model overestimates summertime values at most locations, particularly over southwest and southeast Australia and the northern and eastern parts of New Zealand

(Figure 3e). In winter, the model overestimates T<sub>min</sub> in the north and east of Australia and in parts of New Zealand, while it underestimates temperatures to the west (Figure 3f). The prominent negative bias in temperatures along the western coastline of the South Island of New Zealand in both seasons may be the result of two features: an inability of the model to correctly resolve temperature in this region of complex topography, as well as possible limitations in the VCSN network due to a lack of stations at high elevations.

The simulated patterns of seasonal average precipitation (Figure 4) clearly demonstrate weather@home's ability to capture both seasonal variations and, at least to some extent, the influence of topography. Over Australia, the regional model is able to capture the strong summertime monsoon rainfall over the northern parts of the continent, as well as the rainfall associated with onshore moisture transport along the eastern seaboard (Figure 4a). There is a distinctly different rainfall distribution over Australia in winter, with the highest rainfall restricted to the southern parts of the continent, including regions of topography, where rainfall is often associated with the passage of frontal systems (Figure 4b). Over New Zealand, the model is able to resolve the strong rainfall gradient along the South Island, reflecting the region's complex topography. Overall, weather@home tends to underestimate rainfall in both seasons over Australia and New Zealand, with the exception of parts of southwest and eastern Australia in summer (Figure 4e), and parts of the South Island of New Zealand in winter (Figure 4f). The prominent differences in wintertime rainfall along the southern and western coastlines of Australia (model  $\leq 50\%$  of observations; Figure 4f) suggest that the model may not be able to fully capture the influence of local land sea breezes and/or the influence of passing frontal systems at those locations. Meanwhile, the prominent differences over northern Australia in JJA (Figure 4f) are an artefact of expressing the differences as percentages; the actual rainfall values for both observations and the model output are both small over this region, meaning that any resulting small differences equates to a large percentage.

Next, we assess HadRM3P's capacity to represent interannual variability. For each of the regional clusters identified in Figure 1, we compare timeseries of annual variations of seasonal average temperature and precipitation from the model and the observational datasets (Figures 5–7). Here, we express these timeseries as anomalies relative to the period mean. The solid lines show the median of the 75 ensemble members, while the shading represents the 5th–95th percentile range. Meanwhile, the dashed line represents the corresponding observational dataset. For brevity, we only show timeseries for summer here (winter timeseries are included as supplementary material).

There is general agreement between the interannual variability captured by weather@home and the observational datasets (Figures 5–7). For each of T<sub>max</sub>, T<sub>min</sub> and precipitation there are individual-specific years that correspond to peaks/troughs in both the model estimates and observations, and the overall shapes of the curves are similar. In addition, the observations lie within the model ensemble range for each year with only a few exceptions (e.g., northern Australian rainfall; Figure 7a). Because sea surface temperatures are the only source of interannual variability that is common to both the weather@home simulations and the observational records, the agreement between the timeseries (as represented by the correlation coefficients between the median values for the model simulations and observations) in Figures 5–7 highlights the importance of sea surface temperatures on driving the climates of Australia and New Zealand.

### 3.3 Response to ENSO

Given that ENSO is an important driver of **natural****internal** climate variability for Australia and New Zealand (e.g., Nicholls et al., 1997), we assess weather@home’s ability to correctly simulate ENSO teleconnections. This is achieved by comparing regional model output against observations for the different phases of ENSO: La Niña, neutral and El Niño. La Niña (El Niño) events were defined when the average Nino-3.4 index was at or below (above)  $-1^{\circ}\text{C}$  ( $+1^{\circ}\text{C}$ ) anomaly for at least three months in the September–February period. Neutral events were defined as periods when the average Nino-3.4 index did not go beyond  $\pm 1^{\circ}\text{C}$  anomaly in any month of September–February. These criteria allowed an equal number of events to be selected for each of the three ENSO phases: La Nina (1988/1989, 1998/1999, 2007/2008, 2010/2011), neutral (1992/1993, 1993/1994, 2003/2004, 2012/2013) and El Niño (1994/1995, 1997/1998, 2002/2003, 2009/2010). Furthermore, these criteria allowed the events to be relatively evenly spread across the period of available model simulations (1985–2014). The events were grouped according to their ENSO phase for subsequent analysis.

Figures 8–10 show the distributions of temperature and rainfall, averaged over September–February, for the different phases of ENSO. For the purpose of model evaluation we present results for each of the six study regions. The model-derived distributions are shown as box and whisker plots; each box represents the median and first and third quartiles, while the whiskers extend to the 5th and 95th percentiles. Meanwhile, the corresponding values calculated from the observational dataset are represented as dots. It is worth noting that while we only have four examples of observed atmospheric response to each of the La Niña, neutral and El Niño forcings, we have  $4 \times 75$  examples from weather@home. This large number of model simulations allows us to reduce the influence of internal chaotic variability by averaging in the modelled ensemble and thus, differences between the median values of the box plots are likely to be representative of forced responses to the observed teleconnections.

Overall, weather@home is able to correctly represent the response of temperature and rainfall to changes in the phase of ENSO. As ENSO changes from the La Niña to El Niño phase, there is a warming shift in the distributions of Tmax over each of the Australian regions (Figure 8a–e). Meanwhile, the model is able to capture the reverse relationship for Tmax over New Zealand (Figure 8f). The response of Tmin to changes in the phase of ENSO is less pronounced over Australia (Figure 9a–e), while conditions continue to be warmer over New Zealand during the La Niña phase (Figure 9f). For precipitation, there is a shift towards higher rainfall totals over each of the study regions during La Niña conditions (Figure 10). Weather@home seems able to capture the observed non-linear ENSO-precipitation relationship despite many global coupled and atmosphere-only models failing to do so (King et al., 2015b). While the limited observations prevent us from determining if the magnitude of these shifts are suitably represented by the model, they do suggest that the directions of these shifts are correct.

### 3.4 Daily variability

Because the weather@home setup is specifically designed for use in the attribution of extreme weather events, it is important that the model is able to correctly represent the distribution of daily values of temperature and precipitation at regional and local scales. Such an assessment needs to consider not only the model’s ability to correctly resolve the mean state, but also the

tails of the distributions where the extreme events lie. By identifying any limitations of the model, a bias correction approach may be used to correct for systematic errors.

Here, we compare the distributions of daily Tmax, Tmin and precipitation from the ensemble of weather@home regional simulations against the distribution of these variables in the observational dataset. By way of example, results for the SEAUS region are presented in Figure 11 for both summer (DJF) and winter (JJA) in the form of quantile-quantile plots. The corresponding plots for the other regions are presented as supplementary material. For brevity, we do not intend to provide a thorough assessment of the model's representation of daily fields over each of the defined study domains. Rather, we highlight how the large ensemble provided by weather@home allows us to systematically identify biases in the modelled distribution.

Figure 11 is constructed by extracting the daily fields of Tmax, Tmin and precipitation from the regional model simulations and calculating area averages over the SEAUS region. As the models uses a 360-day calendar, and there are 2175 model simulations (75 model realisations for each of the 29 years), this results in a total sample size of 195,750 daily values for each season. This large sample size provides a thorough sampling of physically plausible climate states represented by the model and thus, allows the tails of the distribution to be resolved with confidence. The solid blue line in Figure 11 identifies the percentile values when considering all of the 2175 model runs together, while the envelope shows the 5th to 95th percentile range for values at each percentile when considering each model run separately. Therefore, the range of this envelope provides an assessment of both sampling uncertainty and internal variability.

Figure 11 shows that the weather@home regional model provides an adequate representation of the distribution of daily summertime Tmax, Tmin and precipitation averaged over SEAUS, when compared against the AWAP observational dataset. That is to say, the solid blue line is almost directly overlying the 1:1 line of agreement (shown in black). The model is capable of not only correctly resolving the mean state, but also the tails of the distribution (represented by the 1st and 99th percentiles). The large spread in the envelope indicates that the model is capable of representing a wide range of temperatures and precipitation rates and therefore, in order to fully sample internal variability of the model, a large ensemble is necessary.

The relatively high resolution of the weather@home regional model, and the large number of model simulations, allows the performance of the model to also be assessed at much more local scales. By way of example, quantile-quantile plots have also been generated for the coastal city of Melbourne and the inland city of Mildura by extracting and examining the corresponding nearest model grid point (see supplementary material). Overall, the weather@home setup provides sufficient model resolution and ensemble sizes to allow the model to be assessed (and where appropriate, bias corrected) for subsequent use in event attribution at both the regional and local scales.

#### 4 Creating counterfactual climate scenarios

In order to quantify how human-induced climate change has altered the likelihood of extreme weather and climate related events, large ensembles of model simulations are required under two distinct scenarios: under current (observed/historical) climate and under a counterfactual (natural) climate as might have been without human influence on atmospheric composition.



Up to this point we have only considered the weather@home model under the observed climate scenario. Therefore, a brief description of the counterfactual climate scenarios is provided here.

The key differences between the observed and counterfactual climate scenarios are the **lower** boundary conditions used to drive the weather@home model. As outlined in Section 2, simulations for the **observedhistorical** climate are driven by **observedhistorical** SSTs and sea-ice from the OSTIA dataset, as well as present day atmospheric composition (well-mixed greenhouse gases, ozone and aerosols). For the counterfactual climate the model is driven by different atmospheric composition and different sea ice and SST specifications; the atmosphere has prescribed pre-industrial (**1850**) levels of greenhouse gases, ozone and aerosols, the sea ice extent corresponds to the year of maximum sea ice extent in each hemisphere of the OSTIA record, and SSTs are modified to remove estimates of anthropogenic warming. Meanwhile, **boundary conditionsforcings** common to both scenarios are the natural forcing factors, such as changes in volcanic aerosols and solar irradiance. **There is no change in the land surface types between the historical and counterfactual climate scenarios.**

As the true climate conditions for the ‘world without humans’ cannot be **knownobserved**, weather@home simulations are run under ten alternative realisations of the counterfactual climate scenario. These alternative realisations are derived from different estimates of the underlying SST warming (delta-SST) due to human influence, which are separately calculated from ten available Coupled Model Intercomparison Project Phase 5 (CMIP5) models (Taylor et al. 2012; see supplementary material for details). Monthly-average delta-SST estimates are calculated for each of the CMIP5 models by calculating the difference between the decadal-average (1996–2005) SSTs from the “historical” simulations (which include both anthropogenic and natural forcings) and the corresponding “natural” simulations. The resulting patterns and magnitudes of warming are seen to differ across the ten delta-SST estimates (Figure 12). These delta-SST patterns are then subtracted from the **observedhistorical** OSTIA SSTs to provide the lower boundary conditions for each of the respective counterfactual realisations.

The use of multiple realisations of the counterfactual scenario allows us to account for some of the uncertainty in our estimates of a world without anthropogenic influence. By way of example, Figure 13 shows weather@home model estimates of summertime daily Tmax at Melbourne for 2014/2015, under the **observedhistorical** and counterfactual climate scenarios. For each of these scenarios the model has been run thousands of times and the daily values of Tmax have been extracted from the nearest model grid point to Melbourne. The return time curve for the **observedhistorical** climate scenario (shown in red) is positioned to the left of the curves for the respective counterfactual scenarios (shown in shades of grey). This suggests that anthropogenic climate change has shifted the distribution of Melbourne summertime maximum temperatures towards warmer conditions. However, the extent of this shift varies when considering each of the separate counterfactual scenarios. The multiple realisations of the counterfactual climate scenario allow for uncertainty to be **characterised** and communicated in any resulting attribution statement. More detailed examples of event attribution studies performed using the weather@home Australia-New Zealand system can be found in the 2015 special issue of the Bulletin of the American Meteorological Society investigating extreme events of 2014 (e.g., Black et al., 2015; Grose et al., 2015; King et al., 2015a; Rosier et al., 2015). The model evaluation undertaken in each of these studies was tailored to the region of interest and builds upon the general model evaluation in this paper.



## 5 Discussion and conclusions

The weather@home Australia-New Zealand climate modelling setup has been described and briefly evaluated. By harnessing spare computing power of volunteers' home computers, weather@home is capable of generating very large ensembles of regional climate model simulations over Australia and New Zealand. This provides a unique tool for undertaking attribution studies of extreme weather and climate events in the region. To date, 75 model simulations have been successfully integrated over the period 1985–2014 in a time-slice manner, while multi-thousand member ensembles have also been generated under both ~~observed~~historical and counterfactual climate scenarios for the years 2013, 2014 and 2015. All of this model output is freely available to the research community.

The weather@home regional model is seen to be capable of resolving many climate features that are important for the Australia and New Zealand regions. This is reflected in the model's ability to provide a good representation of temperature and precipitation, both spatially and temporally. Results presented here suggest that the model is capable of correctly simulating ENSO teleconnections, which is a key requirement given the importance of ENSO on driving natural climate variability in the region. The reasonably high resolution of the regional model, and the large ensemble size achieved through the distributed computing setup, **allows certain types of extreme weather events** to be examined with confidence at regional and local scales. While the model is seen to exhibit varying degrees of bias in temperature and precipitation for different regions, this bias may be corrected through a simple scaling and offset approach (e.g., Sippel and Otto, 2014; Mera et al., 2015), or through more complicated approaches such as quantile mapping (e.g., Bergaoui et al., 2015) or ensemble re-sampling techniques (e.g., Sippel et al., 2016).

Despite the strengths of weather@home it is important to recognise some of the limitations of the project. ~~Because the modelling setup relies on the aggregated power of volunteers' home computers, restriction on machine CPU means that the regional modelling setup can only be achieved using atmosphere-only models. Therefore, climate realisations using high-resolution coupled ocean-atmosphere models are not possible at this point in time.~~ **Under the current model configuration, land surface data (e.g., vegetation roughness and type) are fixed and are the same for the historical and counterfactual climate scenarios.** Furthermore, weather@home only uses a single atmospheric model, meaning that any resulting attribution statement can only be made within the context of that specific modelling setup. In order to test the dependence of the model simulation on physical parameterization, future work will employ a perturbed physics approach whereby perturbations will be applied to components of atmospheric and surface physics.

Other areas of current and future work involve generating larger ensembles for the recent past (1985–2015), as well as sets of future simulations under varying projections of climate change. In addition, the model will be driven with idealised SSTs for different phases of ENSO, under both current and counterfactual climate scenarios, so as to provide a novel framework for assessing the relative roles of ENSO and anthropogenic climate change on recent extreme weather events. Overall, the weather@home Australia-New Zealand modelling setup provides a unique modelling resource and greatly enhances Australia and New Zealand's capacity for researching extremes and understanding their causes.

## 6 Data and code availability

All data generated by the weather@home Australia-New Zealand project is hosted on a server at the Tasmanian Partnership for Advanced Computing, Hobart. This data is freely available to the broader research community and can be accessed by contacting the authors of this manuscript. The HadAM3P and HadRM3P models are both available from the UK Met Office as part of the Providing REgional Climates for Impacts Studies (PRECIS) program. Access to standard versions of the software is dependent on attendance at a PRECIS training workshop after which all source and other materials is made available (<http://www.metoffice.gov.uk/research/applied/applied-climate/precis/obtain>). As a program for supporting developing countries this workshop is free for officially categorised developing countries and attracts a charge for other country participants. The code to manage and embed these models within the Weather@Home project is specific to their utilisation within the BOINC environment and we consider not within the scope of this publication.

*Author contributions.* Mitchell Black's contribution towards this work was performed as part of his PhD project. The weather@home Australia-New Zealand project was initiated by Myles Allen and David Karoly and was set up with the assistance of Richard Jones, Neil Massey, Andy Bowery, Mitchell Black, Suzanne Rosier, Sam Dean, Sarah Sparrow and Friederike Otto. All results were plotted and analysed by Mitchell Black with advice from David Karoly and Andrew King. The paper was written in its final form by Mitchell Black with input from all contributing authors.

*Acknowledgements.* The authors would like to thank Dáithí Stone and Andrew Ciavarella for their assistance in improving the original manuscript. M. T. Black, D. J. Karoly and A. D. King have been supported by funding from the ARC Centre of Excellence for Climate System Science (Grant CE110001028). Weather@home ANZ is a collaboration among the University of Oxford, the UK Met Office, the ARC Centre of Excellence for Climate System Science in Australia, NIWA in New Zealand, the University of Melbourne, the University of Tasmania and the Tasmanian Partnership for Advanced Computing. We thank the volunteers who donated their computing time to run weather@home.


## References

- Allen, M.: Do-it-yourself climate prediction, *Nature*, 401, 642–642, doi:10.1038/44266, 1999.
- Allen, M.: Liability for climate change, *Nature*, 421, 891–892, doi:10.1038/421891a, 2003.
- Anderson, D.: BOINC: A system for public-resource computing and storage, in: 5th IEEE/ACM International Workshop on Grid Computing, 5 8 November, Pittsburgh, P.A., 2004.
- Arblaster, J., Jubb, I., Braganza, K., Alexander, L., Karoly, D., and Colman, R.: Weather extremes and climate change, Tech. rep., Australian Climate Change Science Programme, 2015.
- Bennett, J. C., Grose, M. R., Corney, S. P., White, C. J., Holz, G. K., Katzfey, J. J., Post, D. A., and Bindoff, N. L.: Performance of an empirical bias-correction of a high-resolution climate dataset, *International Journal of Climatology*, 34, 2189–2204, doi:10.1002/joc.3830, 2014.
- 10 Bergaoui, K., Mitchell, D., Zaaoul, R., McDonnell, R., Otto, F., and Allen, M.: The contribution of human-induced climate change to the drought of 2014 in the southern Levant region [in “Explaining Extremes of 2014 from a Climate Perspective”], *Bulletin of the American Meteorological Society*, 96, S66–S69, doi:10.1175/BAMS-D-15-00129.1, 2015.
- Black, M. T., Karoly, D. J., and King, A. D.: The contribution of anthropogenic forcing to the Adelaide and Melbourne, Australia, heat waves of January 2014, *Bulletin of the American Meteorological Society*, 96, S145–S148, doi:10.1175/BAMS-D-15-00097.1, <http://journals.ametsoc.org/doi/10.1175/BAMS-D-15-00097.1>, 2015.
- 15 Cox, M. P., Betts, A. R., Bunton, B. C., Essery, H. R. L., Rowntree, R. P., and Smith, J.: The impact of new land surface physics on the GCM simulation of climate and climate sensitivity, *Climate Dynamics*, 15, 183–203, doi:10.1007/s003820050276, <http://dx.doi.org/10.1007/s003820050276>, 1999.
- CSIRO and Bureau of Meteorology: Climate Change in Australia Information for Australia’s Natural Resource Management Regions: Technical Report, Tech. rep., CSIRO and Bureau of Meteorology, Australia, 2015.
- 20 Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system, *Remote Sensing of Environment*, 116, 140–158, doi:10.1016/j.rse.2010.10.017, 2012.
- Frame, D. J., Aina, T., Christensen, C. M., Faull, N. E., Knight, S. H. E., Piani, C., Rosier, S. M., Yamazaki, K., Yamazaki, Y., and Allen, M. R.: The climateprediction.net BBC climate change experiment: design of the coupled model ensemble., *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences*, 367, 855–70, doi:10.1098/rsta.2008.0240, 2009.
- 25 Frederiksen, J. S. and Frederiksen, C. S.: Interdecadal changes in southern hemisphere winter storm track modes, *Tellus, Series A: Dynamic Meteorology and Oceanography*, 59, 599–617, doi:10.1111/j.1600-0870.2007.00264.x, 2007.
- Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments, *Climate Dynamics*, 30 16, 147–168, doi:10.1007/s003820050010, 2000.
- Gordon, N. D.: The Southern Oscillation and New Zealand Weather, *Monthly Weather Review*, 114, 371–387, 1986.
- Grose, M. R., Black, M. T., Risbey, J. S., and Karoly, D. J.: Attribution of exceptional mean sea level pressure anomalies south of Australia in August 2014, *Bulletin of the American Meteorological Society*, 96, S158–S162, doi:10.1175/BAMS-D-15-00116.1, 2015.
- Hegerl, G. and Zwiers, F.: Use of models in detection and attribution of climate change, *Wiley Interdisciplinary Reviews: Climate Change*, 35 2, 570–591, doi:10.1002/wcc.121, 2011.
- Hendon, H. H., Thompson, D. W. J., and Wheeler, M. C.: Australian rainfall and surface temperature variations associated with the Southern Hemisphere Annular Mode, *Journal of Climate*, 20, 2452–2467, doi:10.1175/JCLI4134.1, 2007.


- Herring, S., Hoerling, M., Peterson, T., and Stott, P.: Explaining extreme events of 2013 from a climate perspective, *Bulletin of the American Meteorological Society*, 95, S1–96, 2014.
- Herring, S. C., Hoerling, M. P., Kossin, J. P., Peterson, T. C., and Stott, P. A.: Explaining Extreme Events of 2014 from a Climate Perspective, *Bulletin of the American Meteorological Society*, 96, S1–S172, 2015.
- 5 Holland, G. J.: Interannual Variability of the Australian Summer Monsoon at Darwin: 1952–82, *Monthly Weather Review*, 114, 594–604, 1986.
- Jones, D., Wang, W., and Fawcett, R.: High-quality spatial climate data-sets for Australia, *Australian Meteorological and Oceanographic Journal*, 58, 233–248, 2009.
- Jones, D. A. and Trewin, B. C.: On the relationships between the El Niño-Southern Oscillation and Australian land surface temperature, *International Journal of Climatology*, 20, 697–719, 2000.
- 10 Jones, R., Noguera, M., Hassell, D., Wilson, S., Jenkins, G., and Mitchell, J.: Generating high resolution climate change scenarios using PRECIS, Tech. rep., Met Office Hadley Centre, 2004.
- Karoly, D. J. and Wu, Q.: Detection of Regional Surface Temperature Trends, *Journal of Climate*, 18, 4337–4343, doi:10.1175/JCLI3565.1, 2005.
- 15 King, A., Black, M., Karoly, D., and Donat, M.: Increased likelihood of Brisbane, Australia, G20 heat event due to anthropogenic climate change, *Bulletin of the American Meteorological Society*, 96, S141–S144, doi:DOI:10.1175/BAMS-D-15-00098.1, 2015a.
- King, A. D., Alexander, L. V., and Donat, M. G.: The efficacy of using gridded data to examine extreme rainfall characteristics: a case study for Australia, *International Journal of Climatology*, 33, 2376–2387, doi:10.1002/joc.3588, 2013.
- King, A. D., Donat, M. G., Alexander, L. V., and Karoly, D. J.: The ENSO-Australian rainfall teleconnection in reanalysis and CMIP5, *Climate Dynamics*, 44, 2623–2635, doi:10.1007/s00382-014-2159-8, 2015b.
- 20 Li, S., Mote, P. W., Rupp, D. E., Vickers, D., Mera, R., and Allen, M.: Evaluation of a regional climate modeling effort for the western United States using a superensemble from weather@home, *Journal of Climate*, pp. 7470–7488, doi:10.1175/JCLI-D-14-00808.1, 2015.
- Massey, N., Jones, R., Otto, F. E. L., Aina, T., Wilson, S., Murphy, J. M., Hassell, D., Yamazaki, Y. H., and Allen, M. R.: Weather@home – development and validation of a very large ensemble modelling system for probabilistic event attribution, *Quarterly Journal of the Royal Meteorological Society*, doi:10.1002/qj.2455, 2015.
- 25 McBride, J. L. and Nicholls, N.: Seasonal relationships between Australian rainfall and the Southern Oscillation, *Monthly Weather Review*, 111, 1998–2004, 1983.
- Mera, R., Massey, N., Rupp, D. E., Mote, P., Allen, M., and Frumhoff, P. C.: Climate change, climate justice and the application of probabilistic event attribution to summer heat extremes in the California Central Valley, *Climatic Change*, doi:10.1007/s10584-015-1474-3, 30 2015.
- Min, S. K., Cai, W., and Whetton, P.: Influence of climate variability on seasonal extremes over Australia, *Journal of Geophysical Research Atmospheres*, 118, 643–654, doi:10.1002/jgrd.50164, 2013.
- Moise, A., Wilson, L., Grose, M., Whetton, P., Watterson, I., Bhend, J., Bathols, J., Hanson, L., Erwin, T., Bedin, T., Heady, C., and Rafter, T.: Evaluation of CMIP3 and CMIP5 Models over the Australian Region to Inform Confidence in Projections, *Australian Meteorological and Oceanographic Journal*, 65, 19–53, 2015.
- 35 Mote, P. W., Allen, M. R., Jones, R. G., Li, S., Mera, R., Rupp, D. E., Salahuddin, A., and Vickers, D.: Superensemble regional climate modeling for the western US, *Bulletin of the American Meteorological Society*, doi:10.1175/BAMS-D-14-00090.1, 2015.

- NASEM: Attribution of Extreme Weather Events in the Context of Climate Change, Tech. rep., The National Academies of Sciences, Engineering and Medicine, Washington, D.C., doi:10.17226/21852, <http://www.nap.edu/catalog/21852>, 2016.
- Nicholls, N., Drosowsky, W., and Lavery, B.: Australian rainfall variability and change, *Weather*, 52, 66–72, doi:10.1002/j.1477-8696.1997.tb06274.x, 1997.
- 5 Pall, P., Aina, T., Stone, D. A., Stott, P. A., Nozawa, T., Hilberts, A. G. J., Lohmann, D., and Allen, M. R.: Anthropogenic greenhouse gas contribution to flood risk in England and Wales in autumn 2000., *Nature*, 470, 382–5, doi:10.1038/nature09762, 2011.
- Perkins, S. E. and Alexander, L. V.: On the measurement of heat waves, *Journal of Climate*, 26, 4500–4517, doi:10.1175/JCLI-D-12-00383.1, 2013.
- Peterson, T. C., Hoerling, M. P., Stott, P. A., and Herring, S. C.: Explaining extreme events of 2012 from a climate perspective, *Bulletin of the American Meteorological Society*, 94, S1–S74, doi:10.1175/BAMS-D-13-00085.1, 2013.
- 10 Pook, M. and Gibson, T.: Atmospheric blocking and storm tracks during SOP-1 of the FROST Project, *Australian Meteorological Magazine*, Special Ed, 51–60, 1999.
- Reisinger, A., Kitching, R. L., Chiew, F., Hughes, L., Newton, P. C. D., Schuster, S. S., Tait, A., and Whetton, P.: Australasia, in: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*, edited by Barros, V. R., Field, C. B., Dokken, D. J., Mastrandrea, M. D., Mach, K. J., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., pp. 1371–1438, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
- 15 Risbey, J. S., Pook, M. J., McIntosh, P. C., Wheeler, M. C., and Hendon, H. H.: On the remote drivers of rainfall variability in Australia, *Monthly Weather Review*, 137, 3233–3253, doi:10.1175/2009MWR2861.1, 2009.
- Risbey, J. S., McIntosh, P. C., and Pook, M. J.: Synoptic components of rainfall variability and trends in southeast Australia, *International Journal of Climatology*, 33, 2459–2472, doi:10.1002/joc.3597, 2013.
- Rosier, S., Dean, S., Stuart, S., Carey-Smith, T., Black, M., and Massey, N.: Extreme rainfall in early July 2014 in Northland, New Zealand -was there an anthropogenic influence?, *Bulletin of the American Meteorological Society*, 96, S136–S140, doi:10.1175/BAMS-D-15-00105.1, 2015.
- 25 Saji, N. H., Goswami, B. N., Vinayachandran, P. N., and Yamagata, T.: A dipole mode in the tropical Indian Ocean., *Nature*, 401, 360–363, doi:10.1038/43854, <http://www.ncbi.nlm.nih.gov/pubmed/16862108>, 1999.
- Schaller, N., Kay, A. L., Lamb, R., Massey, N. R., van Oldenborgh, G. J., Otto, F. E. L., Sparrow, S. N., Vautard, R., Yiou, P., Bowery, A., Crooks, S. M., Huntingford, C., Ingram, W. J., Jones, R. G., Legg, T., Miller, J., Skeggs, J., Wallom, D., Weisheimer, A., Wilson, S., and Allen, M. R.: The human influence on climate in the winter 2013/2014 floods in southern England., *Nature Climate Change*, accepted, doi:10.1038/nclimate2927, 2016.
- 30 Seneviratne, S., Nicholls, N., Easterling, D., Goodess, C., Kanae, S., Kossin, J., Luo, Y., Marengo, J., McInnes, K., Rahimi, M., Reichstein, M., Sorteberg, A., Vera, C., and Zhang, X.: Changes in Climate Extremes and their Impacts on the Natural Physical Environment, Tech. rep., Intergovernmental Panel on Climate Change, 2012.
- Sippel, S. and Otto, F. E. L.: Beyond climatological extremes - assessing how the odds of hydrometeorological extreme events in South-East Europe change in a warming climate, *Climatic Change*, 125, 381–398, doi:10.1007/s10584-014-1153-9, 2014.

- Sippel, S., Otto, F. E. L., Forkel, M., Allen, M. R., Guillod, B. P., Heimann, M., Reichstein, M., Seneviratne, S. I., Thonicke, K., and Mahecha, M. D.: A novel bias correction methodology for climate impact simulations, *Earth System Dynamics*, 7, 71–88, doi:10.5194/esdd-6-1999-2015, <http://www.earth-syst-dynam-discuss.net/6/1999/2015/>, 2016.
- Steffen, W., Hughes, L., and Karoly, D. J.: The critical decade: extreme weather, Tech. Rep. April, The Climate Commission, 2013.
- 5 Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., eds.: *Climate Change 2013 - The Physical Science Basis*, Cambridge University Press, Cambridge, doi:10.1017/CBO9781107415324, 2014.
- Stott, P. A., Stone, D. A., and Allen, M. R.: Human contribution to the European heatwave of 2003, *Nature*, 432, 610–614, doi:<http://dx.doi.org/10.1038/nature03089>, 2004.
- Stott, P. A., Christidis, N., and Betts, R. A.: Changing return periods of weather-related impacts: the attribution challenge, *Climatic Change*, 109, 263–268, doi:10.1007/s10584-011-0265-8, 2011.
- 10 Stott, P. A., Allen, M., Christidis, N., Dole, R., Hoerling, M., Huntingford, C., Pall, P., Perlwitz, J., and Stone, D.: Attribution of weather and climate-related extreme events, in: *Climate Science for Serving Society: Research, Modeling and Prediction Priorities*, edited by Asrar, G. and Hurrell, J., pp. 307–337, Springer Science and Business Media, Dordrecht, 2013.
- Tait, A., Henderson, R., Turner, R., and Zheng, X.: Thin plate smoothing spline interpolation of daily rainfall for New Zealand using a climatological rainfall surface, *International Journal of Climatology*, 26, 2097–2115, doi:10.1002/joc.1350, 2006.
- 15 Taylor, K. E., Stouffer, R. J., and Meehl, G. a.: An Overview of CMIP5 and the Experiment Design, *Bulletin of the American Meteorological Society*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, <http://journals.ametsoc.org/doi/abs/10.1175/BAMS-D-11-00094.1>, 2012.




**Figure 1.** Domain and elevation of terrain (metres) used in the weather@home regional model simulations. Land areas have been separated into six regions for subsequent evaluation: northern Australia (NAUS), central Australia (CAUS), eastern Australia (EAUS), southwest Australia (SWAUS), southeast Australia (SEAUS) and New Zealand (NZ). The coastal city of Melbourne and the inland city of Mildura are identified by the asterisk symbols.




**Figure 2.** Seasonal average daily maximum temperatures ( $^{\circ}$  C) for 1985–2014 for DJF (left), and JJA (right), from the weather@home regional model HadRM3P (top), and the observational datasets (middle; AWAP over Australia and VCSN over New Zealand). The bottom panels show the difference between HadRM3P and the corresponding observational dataset.







**Figure 3.** As in Figure 2, but showing seasonal average minimum temperature ( $^{\circ}$  C).




**Figure 4.** As in Figure 2, but showing seasonal average precipitation (mm/day). The difference fields are expressed as percentages relative to the observational datasets.




**Figure 5.** Time series of summertime (December–February) average maximum temperature for the respective study regions (as labelled) for 1986–2014. Ensemble-mean values from weather@home simulations are shown by the solid line (5th–95th percentile shaded envelope) while the observations (AWAP over Australia and VCSN over New Zealand) are shown by the dashed line. The time series are given as anomalies relative to the mean of the entire period. The bias between the medians for the model and observations is indicated, along with the Pearson correlation coefficient ( $r$ ; calculated using a two-sided test) and  $p$ -value for testing non-correlation.




**Figure 6.** As in Figure 5, but showing average minimum temperature.




**Figure 7.** As in Figure 5, but showing average precipitation.



**Figure 8.** September–February average maximum temperature for the respective study regions (as labelled) during different phases of ENSO: La Niña (LN), neutral (NU) and El Niño (EN). Observed values are plotted as colored circles while values from the weather@home HadRM3P simulations are shown as box-and-whisker plots. The boxes show the median and interquartile range while the whiskers extend to the 5th and 95th percentiles. See text for details.




**Figure 9.** As in Figure 8, but showing minimum temperature.




**Figure 10.** As in Figure 8, but showing precipitation.






**Figure 11.** Quantile-quantile plots showing distributions of daily maximum temperature (a, b), minimum temperature (c, d) and precipitation (e, f), averaged over southeast Australia. Distributions are shown for December–February (a,c,e) and June–August (b, d, f). The solid blue line shows the percentile values for the entire ensemble of model simulations, while the blue envelope shows the 5th to 95th percentile range of values for individual ensemble members.



**Figure 12.** Estimated sea surface temperature response pattern ( $^{\circ}\text{C}$ ) to anthropogenic forcing, calculated from 10 different CMIP5 models (as labelled). The temperature responses are calculated for each month (January–December) but are shown here as annual averages.



**Figure 13.** Return periods of daily December–February maximum temperature at Melbourne, Australia, for historical climate conditions (red) and various counterfactual climate conditions (grey).