

Dear Executive Editor and Referees,

We received your comments and suggestions for our discussion paper "MATRIX-VBS: implementing organic aerosol volatility in an aerosol microphysics model" (now: "MATRIX-VBS (v1.0): an aerosol microphysics model including organic aerosol volatility") on GMDD. We thank you very much for your efforts in evaluating our submission.

We have used the comments received to guide our revision of the discussion paper. Almost all of the advice received were incorporated into the revised paper. Point-by-point answers to all questions raised are listed below, together with highlighted changes in the revised manuscript.

We hereby resubmit the revised discussion paper to be considered for publication in *Geoscientific Model Development*. We confirm that all authors listed on the manuscript concur with submission in its revised form. Should you have any remaining questions, we will be happy to address them.

Sincerely,

C. Y. Gao, K. Tsigaridis and S. E. Bauer

Executive Editor

...In particular, please note that for your paper, the following requirement has not been met in the Discussions paper: • "The main paper must give the model name and version number (or other unique identifier) in the title." Please add a version number for MATRIX-VBS in the title upon your revised submission to GMD.

Now the title reads: "MATRIX-VBS (v1.0): an aerosol microphysics model including organic aerosol volatility".

Concerning your statements in the Code Availability Section, I like to inquire, if the box model, which is subject to your publication, might also be available independent of the full GISS ModelE Earth System Model.

Yes, the box model is available on request. We added this information in the code availability section: "In addition, the box model used here is available on request".

REFEREE #1

- 1. Figures 6 and 7: In Figures 6 and 7, for each population, the median diameters of the number, surface-area, and volume distributions are all the same. For example, in Figure 6 for “T=120hr new”, the median diameter for BOC is 80 nm in each of the three distributions. The median diameter of a mode is only the same for number, surface area, and volume distributions if the particles are in that mode are all exactly the same size. Yet, the modes representing the populations in Figures 6 and 7 are shown having a finite width, which means that the median diameter of the surface-area median diameter should be larger than the number-median diameter, and the volume median diameter larger than that for surface area. I’m guessing that MATRIX is only simulating 2 moments per population (though this is never explicitly stated) and an assumption is made about a 3rd moment is made in order to get the modal width in Figures 6 and 7. This 3rd-moment assumption is fine, but this 3rd-moment assumption needs to be consistently used such that the median diameters shift between the three distributions.**

Thank you for pointing this out. We discovered a bug in the plotting of the size distribution which was not affecting the model results. Now Figures 6 and 7 have been updated and the discussion has been modified accordingly. In addition, it is true that MATRIX only simulates 2 moments per population, and each population has a fixed width. Although we never explicitly mention the term “moments” in the manuscript, the fact that we simulate number and mass concentrations per population is explicitly mentioned in the manuscript, including the abstract. However, to make this point even clearer, we expanded an explicit sentence about it in section 2.1: “MATRIX is designed to resolve the aerosol temporal evolution and represent the mixing states of a user-selected set of aerosol populations, which are modes of different composition as listed in Table 2, tracking two moments each, number and mass, while keeping the width of the distribution fixed”.

- 2. The following information is either missing from the paper or shows up later than ideal:
2.1 Enthalpy of vaporization that drives the temperature dependence of C*s.**

The enthalpy of vaporization is calculated using equation 12 from Epstein et al., 2010. This is now included in Table 1.

bin	Enthalpy of vaporization
-2	153
-1	142
0	131
1	120
2	109
3	98
4	87
5	76
6	65

2.2 How many moments tracked per population? I'm guessing 2 since that what I remember from previous MATRIX papers, but it needs to be explicitly stated here. Which moments? Number and mass (volume)?

Yes, two moments, number and mass. See also answer to point 1.

2.3 In Figures 6 and 7, it looks like the moments are converted to modes. If there are just 2 moments, I'm guessing you assumed a fixed width (this looks to be the case). What width did you use or what did you assume about a 3rd moment?

We included the fixed widths for the modes at the bottom of Table 2: “*The sigma values for all populations are 1.80, except for AKK, which has a sigma of 1.60, and for SSC and MXX, which both have a sigma of 2.00.”

Also see comment about the number, surface area, and volume median diameters of the modes incorrectly being the same.

Thank you, this was a plotting bug, please see answer to point 1.

3. Several things should be explicitly stated in the methods but aren't discussed until later in the paper:

3.1 the duration of the simulations (10 days)


This is already stated in section 4, description of the simulations. We changed “ten” to “10”.

3.2 are emission continuous?

Yes. We added the word “continuously” in the emissions description in section 4. The sentence now reads: “Semi-volatile POA, sulfate in the accumulation mode, and black carbon, are emitted continuously in the OCC, ACC, BC1 populations, respectively, shown in Figure 1 as yellow circles.”

3.3 oxidant concentrations? What is the diurnal cycle of temperature and oxidant concentrations?

We did not consider a diurnal cycle of temperature, for simplicity. Inserted in page 5 line 23: “All parameters and emissions are held constant throughout the simulations”. We also changed the row titles in Table 3 to read “Fixed parameters”. The oxidant concentrations change with time, as calculated by photochemistry and the diurnal variability of solar zenith angle which affects photolysis rates. The mean calculated OH radical concentrations are already mentioned in the results section, but we selected to not show a plot of them, for clarity. They are shown below.



3.4 What are the gas-phase chemical rate constants?

Gas phase organics are oxidized by OH radicals with a rate constant of $1 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ (Donahue et al., 2006). We added this information in section 2.2.

3.5 Is condensation/evaporation to each population calculated through kinetic mass transfer, or are the populations and gas-phase assumed to be in instantaneous equilibrium? Instantaneous equilibrium might work ok here where all of the populations interacting with the gas-phase organics are essentially all accumulation mode (and thus the populations have similar equilibrium timescales); however, this assumption will likely fail when the authors begin to consider organic uptake to the nucleation and Aitken modes, and the equilibration time will vary between modes.

Yes, we used instantaneous equilibrium, as described in Donahue et al. (2006), which is based on the Pankow (1994) theory. This is now mentioned in the revised manuscript in sections 2.2 and 3.3.

4. Other comments Title: “an evolving organic aerosol volatility” sounds awkward to me. What about “an organic aerosol volatility scheme” or removing the work “an” from the current title?

We modified the title to “MATRIX-VBS (v.1): an aerosol microphysical model including organic aerosol volatility”.

5. P1 L21: Can you define the low- and high-volatility ranges?

We added our definition in the VBS framework description: “We classify organics as Murphy et al., 2014 does: low-volatility organics are in bins 10^{-2} to 10^{-1} $\mu\text{g m}^{-3}$ (M2 and M1 in Table 1), semi-volatile organics are in bins 10^0 to 10^2 $\mu\text{g m}^{-3}$ (M0, P1, P2), and intermediate-volatility organics are in bins 10^3 to 10^6 $\mu\text{g m}^{-3}$ (P3, P4, P5, and P6). “ We also modified the terms accordingly throughout the paper.

We also modified the abstract: “Emitted semi-volatile primary organic aerosols evaporate almost completely in the intermediate volatility range, while they remain in the particle phase in the low volatility range.”

6. P1 L22: “The *final* volatility distribution. . .”?

At any point in time the volatility distribution would depend on those factors. We modified the sentence to read “The volatility distribution at any point in time depends on...”

7. P4 L27: This sentence is confusing. I thought that in the old scheme, non-volatile SOA condensed onto modes too (see P3 L21), but this sentence says coagulation is the only process involving organics.

Non-volatile SOA do not condense onto any modes (aerosol populations), they are directly emitted into those, same as other primary aerosols, e.g. BC, as already explicitly stated in the lines mentioned by the reviewer: “In the original version of the MATRIX model, organics only contribute to particle growth and mix with other aerosol species via coagulation. Primary organic aerosols are emitted only as non-volatile particulate organic matter, and do not exist in the gas phase or interact with other aerosol populations”. Page 3 line 21 does not say non-volatile SOA condensed onto modes, it says “The semi-volatile nature of biogenic SOA is not represented in the VBS framework in this work”, so we are not certain why the reviewer was confused here.

8. Figures 2 and 3: I think it would make more sense to have the black line showing the mass in the old scheme in the “aerosol phase” panels rather than the “total” panels since the old scheme was only tracking the aerosol mass. I view the aerosol mass in the new scheme and the old scheme’s aerosol mass as the apples-to-apples comparison.

The reason why we decided to plot the original in the total rather than the aerosol phase is to directly visualize the Shrivastava et al. (2008) emission factors impact on the total organics, which takes into account IVOC emissions. We do agree though that for an apples-to-apples the line is probably better to be presented in the aerosol phase, which is what we now show.

9. P7 L30: “...total gas-phase *concentration* reaches. . .”

Corrected.

10. P8 L6: “...would expect *higher* gas-phase *concentrations* due to. . .”

Corrected.

11. Copy editing: The paper would benefit from having compound adjectives being properly hyphenated. Also commas appear in some places they shouldn’t and don’t appear in some places they should.

When gas-phase is a compound adjective it's hyphenated, when it's not, for instance "in the gas phase" then it's not hyphenated. Same applies for aerosol-phase. In addition, after acceptance for publication and during the production phase, there is a copy-editing process that should take care of any remaining issues on that front. Taken from the GMD website (<http://www.geoscientific-model-development.net/production.html>): "Where applicable, the article undergoes the copy-editing process (spelling, grammar, sentence structure) and the changes are incorporated into the text and again converted into a PDF."

REFEREE #2

- 1. Page 3, line 3-6: This sentence is talking about the role of semi-volatile organics but mentions new particle formation - I would suggest moving the reference to new particle formation to the previous sentence about very low volatility organics, and including some additional references, i.e.: “It has been established that highly oxidised, low-volatility organics play a key role in new particle formation (Metzger et al., 2010; Riccobono et al., 2014; Kirkby et al., 2016) and particle growth (Trostl et al., 2016), while the range. . .” [this is alluded to in your Methods section but not clear here]**

Now the sentence reads: “It has been established that the highly oxidized, very low volatility organics play a key role in particle formation (Metzger et al., 2010; Paasonen et al., 2013; Riccobono et al., 2014; Kirkby et al., 2016) and particle growth (Trostl et al., 2016),”

- 2. Page 4, line 1-14: There are a few more details that would be useful to include in this model description. Specifically: Clarify whether this model tracks both aerosol number and aerosol mass? [I assume both from Figures 6 and 7] What are the “aerosol populations” – they seem to be specific modes of particular composition (perhaps refer the reader to Table 2 at this point)? What are the diameter ranges of these populations? (this also becomes relevant when interpreting Figures 6 and 7)**

Please see response to referee #1 2.2, the two moments, number and mass, are clarified in page 3, line 8. The reference to Table 2 for the clarifying what aerosol populations are is included as well. Now the sentence reads: “MATRIX is designed to resolve the aerosol temporal evolution and represent the mixing states of a user-selected set of aerosol populations, which are modes of different composition as listed in Table 2, tracking two moments each, number and mass, while keeping the width of the distribution fixed.”

Populations have a characteristic size at emission time, and can grow or shrink to the whole size distribution range, contrary to other modal models where mass and number move from one mode to another. A notable exception relevant to the work presented here is the move of aerosol number and mass from BC1 to BC2; this, however, happens with compositional criteria, not size assumptions. For details, see Bauer et al., 2008.

- 3. Page 5, lines 25 - 30: Can you expand on what these mass-based emission factors represent? (enough for the reader to understand without having to refer to Shrivastava et al., 2008) To what is the additional factor of 1.5 applied?**

The emission factors represent the volatility distribution of aerosols at emission time. Current emission inventories assume POA are nonvolatile, thus in order to include a volatility distribution we use these factors. In addition, the current emission inventories do not include the intermediate volatility organic compounds (IVOCs), the sum of which makes up the additional factor of 1.5 of the original POA, based on Shrivastava et al. (2008). This was described in detail in paragraph 30 of Shrivastava et al. 2008, and it is already written in the text: “Adding up the 9 factors from each bin listed in Table 1, we obtain a total factor of 2.5, which means the new scheme’s organics emission is 2.5 times that of the organics emissions in the original scheme. The additional multiplication factor of 1.5 is applied to the emission to account for missing sources of volatile organics in the IVOC volatility regime in the inventories (Shrivastava et al. 2008).”

- 4. While the VBS scheme is outlined in the Model Description, it would be useful to include a brief description of how this influences, and what controls, the gas-particle partitioning among the different populations.**

Added at the end of model development section: “The amount of gas-phase species partitioned onto each aerosol population is based on the surface area of that population, in addition to the mass of that population and the volatility of species, and equilibrium partitioning is assumed.”

- 5. Reaction with OH is mentioned as a mechanism for chemical aging of the organic species, is the OH concentration taken from GISS ModelE? How often are the concentrations updated? How does this process occur (does it occur at a specified reaction rate for each of the volatility bins of organic species?)**

No, the OH radical concentration is calculated in the box model by using the same chemical mechanism and time step (30 minutes) as in the global model. The aging is a reaction which is taken into account in the gas-phase chemical mechanism like any other chemical reaction in the model. We added the following at the beginning of section 2: “A time step of 30 minutes is used, for consistency with the global model.”

- 6. Where do the products of the oxidation of biogenic organics go? Into the lowest volatility bin? Or are they treated entirely separately?**

They are treated separately, as already mentioned in page 3 21-25: “The semi-volatile nature of biogenic SOA is not represented in the VBS framework in this work. Instead, biogenic SOA are treated as non-volatile, as in the original MATRIX version, and are produced with a 10% constant yield from terpenes emissions without any requirement for oxidation before the OA is formed (Lathière et al., 2005; Tsigaridis et al., 2014). The inclusion of semi-volatile biogenic SOA will be parameterized in the same way as in the VBS framework presented here in the future.”

- 7. Some of the information in Section 4 seems like it would be more appropriate in the model (i.e., description of the treatment of BC) / VBS description (i.e., description of the emission rates for the VBS species), rather than the discussion of the simulations?**

We moved the description of BC in the model description, in section 2.1, and the VBS emissions factors in section 2.2. In addition, we added a short sentence here to account for the BC conversion from BC1 to BC2 that was moved elsewhere: “Condensation of VBS species on BC1 can increase the non-absorbing shell of that population, leading to formation of BC2, as described above.”

- 8. Page 6, lines 19 – 20: this currently sounds like each emission factor is 2.5 (which is not the case?), perhaps rephrase this to clarify what you mean.**

We agree that the presence of 2.5 here was confusing. We rephrased the sentence: “... derived from the Shrivastava et al. (2008) mass-based emission factors.”

- 9. In a couple of places, compounds that are low-, intermediate-, and high-volatility are referred to (e.g., Page 7, lines 13-15); in terms of the volatility bins used here, how are these categories defined?**

Added our definition at the description for VBS: “We classify organics as Murphy et al., 2014 does: low-volatility organics are in bins 10^{-2} to 10^{-1} $\mu\text{g m}^{-3}$ (M2 and M1 in Table 1), semi-volatile organics are in bins 10^0 to 10^2 $\mu\text{g m}^{-3}$ (M0, P1, P2), and intermediate-volatility organics are in bins 10^3 to 10^6 $\mu\text{g m}^{-3}$ (P3, P4, P5, and P6).”

10. Page 22, Figure 2: Refer to Table 1 or Figure 1 to explain the legend entries and define OCAR

Fixed Figure 2 legend.

11. Page 24/25, Figure 4/5: The layout of the columns doesn't quite match the description in the caption.

Corrected. Legends now read:

“Figure 4. Temporal evolution of organic aerosol mass concentration in each organics-containing population from the new scheme (first column for January, third column for July), and the old scheme (second column for January, fourth column for July). “

“Figure 5. Temporal evolution of organic aerosol mass concentration fraction in each organics-containing population from the new scheme (first column for January, third column for July), and the old scheme (second column for January, fourth column for July).”

12. General comments: On several occasions, quantities are referred to as “about” ##, it would be better to either refer to a more specific value or refer to this same value as “approximate”.

Replaced all “about” to “approximately.”

13. Specific suggestions:

Page 1, line 21: should this be “high volatility” rather than “high volatile”?

Corrected.

Page 2, line 21: insert “the” between “with” and “hydroxyl”

Corrected.

Page 2, line 27: remove “it” from between “space” and “also”

Removed.

Page 2, line 28/29: specify that the “it” being referred to here is the 2D-VBS?

Inserted.

Page 3, line 23: replace “terpenes emissions” with “emissions of monoterpenes” [if you are only referring to monoterpenes here]

The model version we used contains both monoterpenes and higher terpenes. No changes made.

Page 4, line 21: remove the word “ones” from between “volatile” and “and”

Removed.

Page 5, line 11: insert a hyphen into “nonvolatile”

Inserted.

Page 5, line 12-14: rephrase this sentence, it is currently not clear what you mean – and be consistent with writing numerical values as words or numbers, i.e. “eight” v. “8”

Rephrased, now the sentence reads: “Now each of the 8 organic-containing populations carry 9 additional semi-volatile VBS species listed in Table 1. Together with the 5 original tracers, we now have up to 14 available tracers per population, depending on whether they carry organic aerosols or not, with the original organics tracer (OCAR) representing the non-volatile biogenic OA, as it did in the original mechanism.”

Page 5, line 18: you could replace “of different” with “and”?

Replaced.

Page 5, line 19: you could move “(Mexico City) to the end of this description (after “tropics”) to avoid confusion

Moved.

Page 5, line 23: you could separate this into two sentences, with the first ending after “Table 3” and the second beginning with “Here we do not include. . . .”

Separated.

Page 6, line 30: replace “least one” with “least volatile”

Replaced.

Page 7, line 11: should this say “more oxidized” rather than “less oxidized”?

Corrected.

Page 7, line 18: replace “same with that” as “same as that”

Replaced.

Page 8, line 19: replace “much more” with “higher concentrations”

Replaced.

Page 8, line 20: replace “is also very different” with “are also very different”

Replaced.

Page 9, line 18: replace “effects” with “affects” Page 9, line 29: replace “reason of this” with “reason for this”

Replaced.

REFEREE #3

1. Page 3 lines 21-25: This paragraph needs to be moved to the model description

Moved to the end of the model development section.

2. Section 2.1: I suggest to describe more in detail here the aerosol populations used by the MATRIX model.

We are now referring the description to Table 2: “MATRIX is designed to resolve the aerosol temporal evolution and represent the mixing states of a user-selected set of aerosol populations, which are modes of different composition as listed in Table 2, tracking two moments each, number and mass, while keeping the width of the distribution fixed.”

3. Section 2.2: I suggest to improve the description of the VBS framework used.

3.1 Important information that affect the presented results are missing here: For instance, the description of the ageing mechanism (i.e., the oxidation rate constants used, the reduction in volatility after each oxidation step, and the oxygen mass added per oxidation).

Please see response to Referee #1 3.4: Gas phase organics are oxidized by OH radicals with a rate constant of $10^{-11} \text{ cm}^3 \text{ s}^{-1}$, which is stated in section 3.3.1 of Donahue et al., 2006. The reduction in volatility after each oxidation step is already stated in the VBS framework description in section 2.2 of the paper, which is one bin per oxidation step.

We do not explicitly track oxygen because that's for 2D VBS, we use OM:OC=1.6 which can be modified as needed. For simplicity, we chose to use a constant value for all volatility bins, although when we will do global modeling studies this value will be different for each bin. Choosing to use a higher OM:OC as volatility decreases would have no significant effect on results here other than a higher organic mass with decreasing volatility. Following referee 1's suggestion, we added enthalpy of vaporization in Table 1.

3.2 Furthermore, how do you describe the formation of SOA from traditional VOCs in the model? Do you use a narrower volatility distribution (e.g., up to 10^3)?

As already mentioned in the text: “The semi-volatile nature of biogenic SOA is not represented in the VBS framework in this work. Instead, biogenic SOA are treated as non-volatile, as in the original MATRIX version, and are produced with a 10% constant yield from terpenes emissions without any requirement for oxidation before the OA is formed (Lathièrè et al., 2005; Tsigaridis et al., 2014). The inclusion of semi-volatile biogenic SOA will be 25 parameterized in the same way as in the VBS framework presented here in the future.”

3.3 Do you use aerosol yields for the oxidation of VOCs (if so please report them)?

Other than the 10% from biogenic VOCs that form non-volatile SOA, which is already mentioned in the manuscript (see previous comment as well), no. All VBS-related aerosols are coming from particulate emissions that evaporated, not from gaseous precursors that got oxidized.

3.4 Another issue is how do you perform the partition between the two phases. Do you assume instant equilibrium?

Yes. See also comment 3.5 from reviewer #1.

3.5 Do you account for the temperature dependence of saturation concentrations? If so, what are the enthalpies of vaporization used for each of your organic species?

Yes. The enthalpy of vaporization is calculated using equation 12 from Epstein et al., 2010. This is now included in Table 1. See also comment 2.1 from reviewer #1.

3.6 Finally, since you use terms such as “high volatility range”, “intermediate volatility range”, and “low volatility range” in your results I would recommend to define these terms here in respect to the effective saturation concentration.

We modified and added our definition in the VBS framework description. See also comment 5 from reviewer #1.

4. Page 5, lines 1-4: Which is the extra population compared to the “14 populations” configuration? Furthermore, according to Table 2, the aerosol populations ACC, BC1, BC2, and BCS do not contain organics. Please explain what do you mean that these populations “are set to contain organics as semi-volatile VBS species”.

Thanks for spotting a typo in the manuscript. The original mechanism had 16 populations, not 14. This is now fixed in the revised manuscript. The one population that was eliminated is DBC (dust/BC mixtures), which we know from past work is almost always negligible. We also modified the sentence quoted by the reviewer to “could contain organics as semi-volatile VBS species”. The new mechanism allows for organics to be present to more populations than the original mechanism does, namely the populations mentioned by the reviewer.

5. Page 6 lines 1-2: Why the overestimation of biomass burning emission factors is not an issue for your experiments? Your experiments include forested areas during summer (where you have high emissions from open biomass burning) and highly populated areas during winter (where you have high biomass burning emissions from residential heating).

This statement was not accurate. Hodzic et al. (2015) did not show that Shrivastava et al. (2008) overestimated biomass burning, instead they showed that Shrivastava et al. (2015) had the overestimation. The line has been removed.

6. Page 7 lines 1-12: Most of this discussion belongs to the model description.

Lines abridged and combined in the VBS part of the model description.

7. Page 11 line 30: You did not only include the volatility of the organics, but also their reactivity (by ageing with OH).

Yes. Now the sentence reads: “... how the inclusion of semi-volatility of organics and their reactivity affected ...”

8. Table 2: What is the size of the mode (Aitken, accumulation, or coarse) used for the aerosol populations OCC, BC1, BC2, BCS, BOC, OCS, and MXX? Also, does the OCC has sulfate? Because in that case it seems to be identical to the OCS.

Populations can grow or shrink to the whole size distribution range, contrary to other modal models. See also reply 2 of reviewer #2. OCC could be the same in terms of composition, but has different origin: OCC is directly emitted and can be sulfate-free, while OCS comes from

coagulation of OCC with ACC and a) is a secondary particle, rather than a primary, and b) will always contain sulfate.

- 9. Figure 2: “P2” and “OCAR” seem to have very similar color. Please change the color of “P2”. Furthermore, it would be nice to add for comparison the “OCAR” dashed line in the “Aerosol phase” column as well.**

Following the recommendation of reviewer #2, we moved OCAR to the aerosol phase column. This also makes the line much more visible, and its black color does not get confused with the grey P2.

- 10. Figure 4: The label is wrong. It writes that the second column is “July new” and the third for “January old” which is not the case.**

Legend is fixed. See also comment 11 from reviewer #2.

References:

Bauer, S. E., Wright, D., Koch, D., Lewis, E. R., McGraw, R., Chang, L.-S., Schwartz, S. E. and Ruedy, R.: MATRIX (Multiconfiguration Aerosol TRacker of mIXing state): an aerosol microphysical module for global atmospheric models, *Atmos. Chem. Phys. Discuss.*, 8(3), 9931–10003, doi:10.5194/acpd-8-9931-2008, 2008.

Donahue, N. M., Robinson, a. L., Stanier, C. O. and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40(8), 2635–2643, doi:10.1021/es052297c, 2006.

Epstein, S. A., Riipinen, I., and Donahue, N. M.: A Semi-Empirical Correlation between Enthalpy of Vaporization and Saturation Concentration for Organic Aerosol, *Environ. Sci. Technol.*, 44, 743–748, doi:10.1021/es902497z, 2010.

Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C., Jimenez, J. L., Madronich, S., and Park, R. J.: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmos. Chem. Phys. Discuss.*, 15, 32413-32468, doi:10.5194/acpd-15-32413-2015, 2015.

Murphy, B. N., Donahue, N. M., Robinson, A. L., and Pandis, S. N.: A naming convention for atmospheric organic aerosol, *Atmos. Chem. Phys.*, 14, 5825-5839, doi:10.5194/acp-14-5825-2014, 2014.

Pankow, J. F.: An absorption-model of gas-particle partitioning of organic compounds in the atmosphere, *Atmos. Environ.*, 28, 185–188, 1994.

Shrivastava, M. K., Lane, T. E., Donahue, N. M., Pandis, S. N. and Robinson, A. L.: Effects of gas particle partitioning and aging of primary emissions on urban and regional organic aerosol concentrations, *J. Geophys. Res. Atmos.*, 113(18), doi:10.1029/2007JD009735, 2008.

Shrivastava, M., Easter, R., Liu, X., Zelenyuk, A., Singh, B., Zhang, K., Ma, P-L, Chand, D., Ghan, S., Jimenez, J. L., Zhang, Q., Fast, J., Rasch, P., and Tiitta, P.: Global transformation and fate of SOA: Implications of low volatility SOA and gasphase fragmentation reactions, *J. Geophys. Res.-Atmos.*, 120, 4169–4195, doi:10.1002/2014JD022563, 2015.

MATRIX-VBS (v1.0): an aerosol microphysics model including organic aerosol volatility

C. Y. Gao^{1,2}, K. Tsigaridis^{3,2}, S. E. Bauer^{3,2}

¹Department of Earth and Environmental Sciences, Columbia University, New York, NY, 10027, USA

²NASA Goddard Institute for Space Studies, New York, NY, 10025, USA

³Center for Climate System Research, Columbia University, New York, NY, 10025, USA

Correspondence to: K. Tsigaridis (kostas.tsigaridis@columbia.edu)

Abstract. The gas-particle partitioning and chemical aging of semi-volatile organic aerosol are presented in a newly developed box model scheme, where its effect on the growth, composition and mixing state of particles is examined. The volatility-basis set (VBS) framework is implemented into the aerosol microphysical scheme MATRIX (Multiconfiguration Aerosol TRacker of mIXing state), which resolves mass and number aerosol concentrations and in multiple mixing-state classes. The new scheme, MATRIX-VBS, has the potential to significantly advance the representation of organic aerosols in Earth system models by improving upon the conventional representation as non-volatile particulate organic matter, often with also an assumed fixed size distribution. We present results from idealized cases representing Beijing, Mexico City, a Finnish and a Southeast U.S. forest, and investigate the evolution of mass concentrations and volatility distributions for organic species across the gas and particle phases, as well as assessing their mixing state among aerosol populations. Emitted semi-volatile primary organic aerosols evaporate almost completely in the intermediate volatility range, while they remain in the particle phase in the low volatility range. Their volatility distribution at any point in time depends on the applied emission factors, oxidation by OH radicals, and temperature. We also compare against parallel simulations with the original scheme, which represented only the particulate and non-volatile component of the organic aerosol, examining how differently the condensed phase organic matter is distributed across the mixing states in the model. The results demonstrate the importance of representing organic aerosol as a semi-volatile aerosol, and explicitly calculating the partitioning of organic species between the gas and particulate phases.

Keywords: organic aerosols, volatility-basis set, aerosol mixing state, box model

1 Introduction

Atmospheric aerosols play a key role in the Earth system with great impacts on global air quality, public health and climate (Boucher et al., 2013; Myhre et al., 2013; Seinfeld and Pandis, 2016). One contribution to the large uncertainty in aerosol radiative forcing is organic aerosol (OA), which is ubiquitous in the

Deleted: implementing an evolving organic aerosol volatility and its evolution in

Deleted: 5

Deleted: e

Deleted: high

Deleted: e

Deleted: low volatility

Deleted: least-volatile

atmosphere and contribute to a large portion of submicron particulate mass in various regions around the world (Zhang et al., 2007; Jimenez et al., 2009). Advancements in measurement techniques greatly improved our understanding of the evolution of OA and lifetime in the atmosphere at the process level (Jimenez et al., 2009). However, OA processes in models still remain poorly constrained. Measurements imply that OA concentrations are potentially underestimated in current models (Tsigaridis et al., 2014). Such a discrepancy hints at large uncertainties in the prediction of aerosol-radiation interactions, their hygroscopicity, aerosol-cloud interactions and their overall impact on climate (Petters and Kreidenweis, 2007).

Missing sources of secondary organic aerosol (SOA) in models have been suggested to be the main cause of the underestimated OA formation (Heald et al., 2005; Volkamer et al., 2006; Hodzic et al., 2010; Spracklen et al., 2011). More recently, studies have sought to investigate the underestimation of organic aerosol mass within more advanced model frameworks, which are capable of resolving semi-volatile primary organic aerosol (POA) and including secondary organic aerosol (SOA) from a wider set of precursors including intermediate-volatility organic compounds (IVOCs). The volatility-basis set was developed (Donahue et al., 2006) to provide a relatively simple framework whereby models can represent the overall behavior of the myriad of compounds that constitute organic aerosol and their precursors. The approach involves considering OA as being composed of a number of representative species, each with a particular volatility, spanning a spectrum in vapor pressures from highly volatile (which essentially remains in the gas phase) to very low vapor pressure species which partition readily into the particle phase. VBS then captures the chemical aging of the organic species in the gas-phase, with the hydroxyl radical oxidizing them and producing the adjacent lower volatility class as a product. This method has been used extensively in regional studies (Robinson et al., 2007; Shrivastava et al., 2008; Murphy and Pandis, 2009; Tsimpidi et al., 2010; Hodzic et al., 2010; Fountoukis et al., 2011; Tsimpidi et al., 2011; Bergström et al., 2012; Athanasopoulou et al., 2013; Zhang et al., 2013; Fountoukis et al., 2014) but less so in global models (Pye and Seinfeld, 2010; Jathar et al., 2011; Jo et al., 2013; Tsimpidi et al., 2014; Hodzic et al., 2015). Other studies have used the 2D-VBS (Donahue et al., 2011; Murphy et al., 2011), an approach that in addition to the volatility space, also resolves that of chemical composition, by tracking the amount of oxygenation in the representative organic compounds. However, the 2D-VBS is not implemented in global models, due to its large amount of tracers and the large number of free parameters that are involved in the parameterization.

The inclusion of semi-volatile organics is important for accounting for the total mass of organics in the particulate phase, since an increase in particulate organic matter may not be the result of chemically produced low-volatility species, but simply be reflecting a temperature-driven increase in the partitioning of semi-volatile organic aerosol into the particle phase. It has been established that the highly oxidized, very low volatility organics play a key role in particle formation (Metzger et al., 2010; Paasonen et al., 2013; Riccobono

Deleted:

Deleted: intermediate volatility

Deleted: e

Deleted: on top of

Deleted: it

Deleted: it

Deleted: low volatility

et al., 2014; Kirkby et al., 2016) and particle growth (Tröstl et al., 2016), while the range of volatilities contributing to aerosol growth increases with aerosol size (Pierce et al., 2011; Yu, 2011). Semi-volatile organics also affect aerosol size and mixing state, as well as their impact on climate, due to changes in cloud condensation nuclei (CCN) formation rates (Petters et al., 2006; Riipinen et al., 2011; Scott et al., 2015).
5 hygroscopicity (Petters and Kreidenweis, 2007) and optical properties (Mehre et al., 2013). Since OA emissions are on the rise from developing countries (Lamarque et al., 2010) and no Earth system model considers anthropogenic OA as semi-volatile as measurements suggest, it is important to include and constraint semi-volatile organics to ultimately reduce uncertainties in aerosol radiative forcing and make climate model simulated aerosol changes more realistic.

Deleted: o

Deleted: in particle growth

10 The objective of this study is to further develop an aerosol microphysics model by including a more advanced representation of organic aerosol, including semi-volatile primary OA and an evolving OA volatility during chemical aging in the gas phase in its calculations. This objective was achieved by implementing the VBS framework in the aerosol microphysical scheme MATRIX (Bauer et al., 2008), which represents major aerosol processes such as nucleation, condensation (excluding organics in its original version) and coagulation, and
15 explicitly tracks the mixing state of different aerosol populations. As many traditional chemistry-climate models do (Tsigaridis et al., 2014), MATRIX treats POA and SOA as non-volatile (Bauer et al., 2008). By coupling MATRIX with VBS, POA are treated as condensable semi-volatile organic compounds. These can partition among different aerosol populations based on their volatility and aerosol population size distribution, capturing particle growth via condensation of low-volatility organic vapors, thus providing a more physically-
20 based calculation of aerosol microphysics.

Deleted: e

Deleted: -

Deleted: low volatility

2 Model description

A box model is used for this study. The gas-phase chemical mechanism CBM-IV (Gery et al., 1989), as used in the NASA GISS ModelE (Shindell et al., 2001; Shindell et al., 2003), is coupled to the MATRIX aerosol microphysics scheme, utilizing the Kinetic Pre-Processor KPP (Sandu and Sander, 2006) to solve the
25 differential equations of the gas-phase chemistry scheme. [A time step of 30 minutes is used, for consistency with the global model.](#)

Deleted: The semi-volatile nature of biogenic SOA is not represented in the VBS framework in this work. Instead, biogenic SOA are treated as non-volatile, as in the original MATRIX version, and are produced with a 10% constant yield from terpenes emissions without any requirement for oxidation before the OA is formed (Lathièrre et al., 2005; Tsigaridis et al., 2014). The inclusion of semi-volatile biogenic SOA will be parameterized in the same way as in the VBS framework presented here in the future.*

2.1 MATRIX box model

MATRIX (Multiconfiguration Aerosol TRacker of mIXing state; Bauer et al., 2008) is an aerosol microphysical model based on the Quadrature Method of Moments scheme (McGraw, 1997) in the NASA
30 GISS ModelE Earth System Model, which can be used either as a module within the global model or as a stand-alone box-model. Here, the stand-alone box model is used for development. The design of the code is

such that the box-model code can be used as-is in the global model, without any changes, allowing for seamless transition and maximum portability. MATRIX is designed to resolve the aerosol temporal evolution and represent the mixing states of a user-selected set of aerosol populations, which are modes of different composition as listed in Table 2, tracking two moments each, number and mass, while keeping the width of the distribution fixed. It describes new particle formation, particle growth through condensation with explicit treatment of sulfuric acid condensation and lumped treatment of the NH₄-NO₃-H₂O system, as well as coagulation of particles among different populations. Each aerosol population has its own set of aerosol components, which may be primary (from direct aerosol emissions), secondary (formed by nucleation or condensation of gas-phase components onto existing primary particles), or mixed (from any constituent, following condensation on primary aerosols or coagulation between primary/secondary/mixed populations). Black carbon is uniquely treated in MATRIX, in order to separate the coated (via condensation) from the mixed (via coagulation) populations. It is emitted in BC1, which can grow (blue arrow in Figure 1) with inorganic and organic coating, and as its coating volume fraction reaches 5%, it would be moved in the BC2 population, shown as the orange circle.

2.2 VBS framework

The volatility-basis set approach is introduced to the original model; it is an organic aerosol volatility parameterization that separates semi-volatile organic compounds into logarithmically-spaced bins of effective saturation concentrations, which are used for gas-particle partitioning and photochemical aging (Donahue et al., 2006). The scheme groups organic compounds into nine surrogate VBS species according to their effective saturation concentrations (C*) at 298 K, which are separated by factors of ten, ranging from 10⁻² to 10⁶ μg m⁻³. We classify organics as [Murphy et al., 2014](#) does: low-volatility organics are in bins 10⁻² to 10⁻¹ μg m⁻³ (M2 and M1 in Table 1), semi-volatile organics are in bins 10⁰ to 10² μg m⁻³ (M0, P1, P2), and intermediate-volatility organics are in bins 10³ to 10⁶ μg m⁻³ (P3, P4, P5, and P6). Low-volatility organics partition almost exclusively to the particulate phase, the semi-volatile species are present in both the gas and aerosol phase, and intermediate-volatility organics are the most volatile ones in the framework and remain almost exclusively in the gas phase. Equilibrium partitioning is assumed for all volatility bins. Gas phase organics can become chemically aged by the extremely reactive hydroxyl radicals (°OH) during daytime with a rate constant of 10⁻¹¹ cm³ s⁻¹, and as they become more oxidized, their volatility decreases and they move down to the adjacent bin with a factor of 10 lower volatility (Donahue et al., 2006). Parameters and names used to represent them in this study are listed in Table 1.

The emission rates for the VBS species were derived from the POA emission rate in the global model for the corresponding gridbox and month, which were distributed in the volatility space by using mass-based emission factors from [Shrivastava et al. 2008](#) (Table 1). Adding up the 9 factors from each bin listed in Table

Deleted: -

Deleted:

Formatted: Font: 12 pt

Formatted: Heading 2, Line spacing: single

Deleted: Shrivastava

Deleted: 2008

Deleted: least-volatile

Formatted: Not Superscript/ Subscript

Deleted: ³

Formatted: Not Superscript/ Subscript

Deleted: , and P3

Deleted: intermediate volatility

Deleted: ⁴

Formatted: Not Superscript/ Subscript

Deleted: M

Deleted: M

Deleted: M

Deleted: Species in the 10² μg m⁻³ bin

Deleted: Least-volatile

Deleted: are the least volatile ones and

Deleted:

Deleted: while species in the 10⁶ μg m⁻³ bin

Deleted: intermediate volatility

Deleted: These compounds

Deleted: while in the gas-phase following reaction with °OH radicals, which results in oxidizing to a species

Deleted:

1, we obtain a total factor of 2.5, which means the new scheme's organics emission is 2.5 times that of the organics emissions in the original scheme. The additional multiplication factor of 1.5 is applied to the emission to account for missing sources of volatile organics in the IVOC volatility regime in the inventories (Shrivastava et al. 2008).

3 Model development

In the original version of the MATRIX model, organics only contribute to particle growth and mix with other aerosol species via coagulation. Primary organic aerosols are emitted only as non-volatile particulate organic matter, and do not exist in the gas phase or interact with other aerosol populations. Implementing the VBS scheme adds these missing processes. Before this development, there were 8 alternative configurations of MATRIX available to the user, each representing a distinct set of aerosol populations whose number, composition and interactions by coagulation vary. A 9th configuration with 15 selected aerosol populations is created for this study (Table 2), in which 8 of the 16 populations, ACC, OCC, BC1, BC2, OCS, BOC, BCS, and MXX, could contain organics as semi-volatile VBS species. We only included semi-volatile organics in 8 populations, so that we can examine the BC-OA-sulfate-nitrate system first, before adding them into the nucleation population AKK and the dust and sea salt populations (DD1, DS1, DD2, DS2, SSA, SSC). Through coagulation, the 15 donor populations grow or mix and are placed into recipient populations, based on the donor population composition, as described in Bauer et al. (2008). In a future stage, organics will also be implemented in the AKK mode to present nanoparticle growth and we will include an additional nucleation scheme that considers the dependence of new particle formation that involve organics (Kirkby et al., 2016; Tröstl et al., 2016).

Previously, each aerosol population carried up to 5 tracers – sulfate, black carbon, non-volatile organics, dust and sea salt. Now each of the 8 organic-containing populations carry 9 additional semi-volatile VBS species listed in Table 1. Together with the 5 original tracers, we now have up to 14 available tracers per population, depending on whether they carry organic aerosols or not, with the original organics tracer (OCAR) representing the non-volatile biogenic OA, as it did in the original mechanism. This newly coupled model MATRIX-VBS treats POA as semi-volatile gas-phase species, which then partition into and out of the particulate phase. The amount of gas-phase species partitioned onto each aerosol population is based on the surface area of that population, in addition to the mass of that population and the volatility of species, and equilibrium partitioning is assumed.

The semi-volatile nature of biogenic SOA is not represented in the VBS framework in this work. Instead, biogenic SOA are treated as non-volatile, as in the original MATRIX version, and are produced with a 10% constant yield from terpenes emissions without any requirement for oxidation before the OA is formed

Deleted: 1

Deleted: eight

Deleted: ninth

Formatted: Superscript

Deleted: which builds on top of configuration 1 in Bauer et al., 2008,

Deleted: eight

Deleted: 14

Deleted: are set to

Deleted: eight

Deleted: fifteen

Deleted: .

Deleted: eight

Deleted:

Deleted: .

Deleted: The addition to

Deleted: countto

Deleted: s

Deleted: .

(Lathièrè et al., 2005; Tsigaridis et al., 2014). The inclusion of semi-volatile biogenic SOA will be parameterized in the same way as in the VBS framework presented here in the future.

4 Simulations

To test the newly developed model's behavior, we simulated idealized cases representative of four different locations and environments: one very polluted city (Beijing), another cleaner yet still very polluted city at high altitude and closer to the tropics (Mexico City), a very clean Finnish forest (Hyytiälä), and an anthropogenically-affected forest in the Southeast U.S. (Centreville, Alabama). The experiments are performed for a winter (January) and a summer month (July) for 10 days, and initial conditions and emission rates for each location were extracted from a GISS ModelE simulation (similar setup as described in Mezuman et al., 2016) for the year 2006, listed in Table 3. All parameters and emissions are held constant throughout the simulations. Here we do not include deposition and dilution, for simpler mass-balance calculations. Semi-volatile POA, sulfate in the accumulation mode, and black carbon, are emitted continuously in the OCC, ACC, BC1 populations, respectively, shown in Figure 1 as yellow circles. Condensation of VBS species on BC1 can increase the non-absorbing shell of that population, leading to formation of BC2, as described above.

The four organic-containing populations described above can coagulate (black arrows in Figure 1) with themselves and each other and form three additional organic-containing mixed populations, BOC, OCS and BCS, shown as green circles. This schematic includes seven of the eight organic-containing populations in the model.

5 Results and discussion

The temporal evolution of the total organics mass concentrations from the new scheme and the old scheme are presented in Figure 2 for January and Figure 3 for July in the four locations under study. They show large changes in organics concentrations between the old scheme (black line on the right column) and the new one (colors). The organics in the new scheme are represented and distributed by organic tracers of different volatility, whose saturation concentration C^* ranges from the least volatile $10^{-2} \mu\text{g m}^{-3}$ ("M2" in Figures 2 and 3) to the most volatile $10^6 \mu\text{g m}^{-3}$ ("P6" in Figures 2 and 3). They are distributed between the gas and aerosol phases by gas-particle partitioning, whereas the organics in the original scheme are only represented by one nonvolatile organic aerosol tracer ("OCAR").

As mentioned in the model description, the emission rates for organics in the each of the volatility bin in the new scheme were derived from the Shrivastava et al. (2008) mass-based emission factors. Consequently, since there is no deposition and dilution in the simulations, the new scheme's organics total mass concentrations

Deleted: of different

Deleted: (Mexico City)

Deleted: ten

Deleted: ,

Deleted: il

Deleted: with no

Deleted: The emission rates for the VBS species were derived from the POA emission rate in the global model for the corresponding gridbox and month, which were distributed in the volatility space by using mass-based emission factors from Shrivastava et al. 2008 (Table 1). Adding up the 9 factors from each bin listed in Table 1, we obtain a total factor of 2.5, which means the new scheme's organics emission is 2.5 times that of the organics emissions in the original scheme. The additional multiplication factor of 1.5 is applied to the emission to account for missing sources of volatile organics in the IVOC volatility regime in the inventories (Shrivastava et al. 2008).

Deleted: Hodzic et al. (2015) showed that Shrivastava et al. (2008) overestimates biomass burning, which could affect the emission factors we used, however, it is not an issue for the idealized experiments in the present study.

Deleted: Black carbon is uniquely treated in MATRIX, in order to separate the coated (via condensation) from the mixed (via coagulation) populations. It is emitted in BC1, which can grow (blue arrow in Figure 1) with inorganic and organic coating, and as its coating volume fraction reaches 5%, it would be moved in the BC2 population, shown as the orange circle.

Deleted: '

Deleted: '

Deleted: , of 2.5, from Shrivastava et al. 2008

(shown in color in the right columns of Figures 2 and 3) always adds up to 2.5 times that of the old scheme (shown as dash-dotted lines) throughout the simulations in both January and July.

5.1 Winter

In January, the total mass concentration for organics in Beijing, Centreville, Hyytiälä and Mexico City at the end of 10 days are approximately 115 $\mu\text{g}/\text{m}^3$, 16 $\mu\text{g}/\text{m}^3$, 13 $\mu\text{g}/\text{m}^3$, and 65 $\mu\text{g}/\text{m}^3$, respectively. Organic VBS species partition between the gas and aerosol phases within their corresponding volatility bin. The more volatile the species, the more it partitions into the gas phase. The concentration evolution of VBS species in the gas phase from the four locations are shown in the left column of Figure 2 for January. From top to bottom in each panel, volatility decreases from the most volatile species (“P6”) to the least volatile (“M2”). Although semi-volatile organics are emitted in the aerosol phase, in the intermediate volatility range from P6 to P3 bins, the species are so volatile that they evaporate and partition into the gas phase almost completely. ↓

In all four locations, almost all species in the intermediate volatility range are in the gas phase, those in the semi-volatile range partition between the gas and aerosol phases, and those in the low volatility range are in the aerosol phase in January. This is especially true for Beijing and Hyytiälä, where the volatility distributions are very similar (in relative terms), where the total concentration of gas-phase species is higher than the sum of all aerosol-phase species. In Centreville, the total amount of gas-phase species is approximately the same as that of the aerosol-phase species, whereas in Mexico City there are more species in the aerosol than in the gas phase. In Centreville and Mexico City, the species show a diurnal variability, which will be explained later.

Aging can help explain the similar volatility distributions in Beijing and Hyytiälä. The $\cdot\text{OH}$ concentration in both locations are low in January: Beijing’s mean $\cdot\text{OH}$ is approximately 10^5 molecules/ cm^3 and Hyytiälä’s mean $\cdot\text{OH}$ is approximately 10^4 molecules/ cm^3 . Low $\cdot\text{OH}$ concentrations limit the aging of intermediate-volatility organics and their ability to move to the lower volatility bins, thus the volatility distributions do not change drastically, something that is also evident by the lack of a daily cycle. On the other hand, much higher mean $\cdot\text{OH}$ concentrations in Centreville ($2 \cdot 10^6$ molecules/ cm^3) and Mexico City ($5 \cdot 10^6$ molecules/ cm^3) provide more oxidation power, making oxidation a significant pathway in aerosol evolution. The higher mean $\cdot\text{OH}$ concentrations also explain the diurnal variability of both gas-phase and aerosol-phase mass concentrations we see in the two locations, because $\cdot\text{OH}$ is only produced during daytime and has very low concentrations during night. Since Mexico City has slightly higher $\cdot\text{OH}$ concentration than Centreville, its total gas-phase concentration reaches a dynamic equilibrium after approximately 4 days, whereas Centreville’s total gas-phase continues to rise approaching equilibrium at a slower pace.

Looking at the total of the organics (right column of Figure 2), it is not surprising that the very polluted Beijing has the highest concentration of total organics while the cleanest location, Hyytiälä, has the lowest; what is interesting, however, is that organics at these locations share similar volatility distributions. By the end

Deleted: about

Deleted: one

Deleted: high volatility

Deleted: 4

Deleted: Volatile organics are oxidized in the gas phase by the extremely reactive hydroxyl radicals ($\cdot\text{OH}$) during daytime, and as they become more oxidized, their volatility decreases and they move down to the adjacent lower volatility bin. This oxidation process is called “aging” in the VBS framework. For example, species from the P6 bin are emitted in the aerosol phase, but because they are so volatile, they evaporate immediately into the gas phase. During daytime, when OH radicals are abundant, they age, become less volatile, and are placed in the lower volatility bin P5, which has a C^* one order of magnitude lower than its precursor. Similarly, as gas-phase species in all volatility bins age, they get oxidized, become less volatile, and move further down the volatility space. At intermediate volatilities species are becoming less volatile, and they no longer completely evaporate after emission, but are present in both the gas and the aerosol phases, following partitioning. However, they continue to become even more less oxidized following oxidation by OH , until they have such low volatility that they almost exclusively partition into the aerosol phase (Figure 2 middle column).

Deleted: high volatility

Deleted: intermediate volatility

Deleted: low volatility

Deleted: least-volatile

Deleted:

Deleted: about

Deleted: with

Deleted:

Deleted: about

Deleted: about

Deleted: high volatility

Deleted: intermediate volatility

Deleted:

Deleted:

Deleted:

Deleted: about

Deleted:

of the 10-day simulations in the new scheme, the volatility distributions in Beijing and Hyytiälä are very similar to the emission factors distribution among the volatility, with factor differences of less than 0.1. This behavior is, again, a result of the low $\cdot\text{OH}$ concentrations in the two locations, and the low oxidation rate that limits the change in volatility distribution. Volatility is also temperature dependent, which is also relevant to the total aerosols present. In Beijing, we would expect higher gas-phase concentration due to the higher temperatures. However, the larger amount of aerosols moves the partitioning point towards the aerosol phase, which offsets the temperature difference in the colder Hyytiälä case, and gives us similar results.

On the other hand, the volatility distributions in Centreville and Mexico City are very different from the applied emission factor distribution, except the two bins in the low volatility range, M2 and M1. Due to the high concentrations of $\cdot\text{OH}$, both sites have low gas-phase organics concentrations because the intermediate-volatility gases are more efficiently oxidized and their less-volatile products partition into the aerosol phase. Therefore, the relative amount of organics from the intermediate volatility range no longer resembles the applied emission factors. The organics in the intermediate volatility range from P6 to P3, are totaling factors of approximately 0.38 and 0.15 in Centreville and Mexico City, respectively, which are, in sharp contrast to the factors of 0.4, 0.5, 0.8 applied to each of the respective bins.

5.2 Summer

The total mass concentration of organics in Beijing and Mexico City at the end of 10 days in July are approximately $130 \mu\text{g}/\text{m}^3$ and $67 \mu\text{g}/\text{m}^3$, very similar to the amounts in January. However, Centreville and Hyytiälä have higher concentrations of organics than they did in January, with $90 \mu\text{g}/\text{m}^3$ and $43 \mu\text{g}/\text{m}^3$. The volatility distributions for the four locations in July (Figure 3) are, also very different from that of January. Organics are all very low in the intermediate volatility and semi-volatile ranges, and they are all high in the low volatility ranges, with less than 10% of the total organics in the gas phase in all four locations. This behavior means that at all locations oxidation is very strong, stronger than any place during January. This sharp change in behavior is caused by the difference in $\cdot\text{OH}$ concentrations during the two months. July's concentrations are much higher than those in January because $\cdot\text{OH}$ production is increased due to increased photolysis in the summer. The mean $\cdot\text{OH}$ concentration is approximately $1.5 \cdot 10^7$ molecules/ cm^3 in Beijing and Hyytiälä, and it is approximately $1 \cdot 10^7$ molecules/ cm^3 in Centreville and $2 \cdot 10^7$ molecules/ cm^3 in Mexico City. More $\cdot\text{OH}$ leads to faster oxidation of the gas-phase organics and the consequent partitioning of the less-volatile oxidation products into the aerosol phase. This is evident in Figure 3, where the gas-phase concentrations in all four locations are very low. In all cases, dynamic equilibrium was reached after just two days. They also exhibit a strong diurnal variability, as expected from the fast $\cdot\text{OH}$ oxidation, which decreases with decreasing volatility.

Deleted: more

Deleted: est

Deleted: high volatility

Deleted: intermediate volatility

Deleted: -

Deleted: high volatility

Deleted: high volatility

Deleted: 4

Deleted: a

Deleted: around

Deleted: is

Deleted: about

Deleted: much more

Deleted: is

Deleted: high

Deleted: intermediate volatility

Deleted: low volatility

Deleted: least-volatile

Deleted: about

Deleted: about

Deleted:

Deleted:

5.3 Mixing state

The temporal evolution of total organic aerosol mass concentration per population is shown in Figure 4 (absolute amounts) and Figure 5 (relative amounts). The first and third columns are results from the new scheme with condensing and coagulating organics for January and July, respectively, while the second and fourth columns are results from the old scheme with only coagulating organics from January and July, respectively. The organic aerosol mass concentrations in Figure 4 correspond to the aerosol-phase concentrations in Figures 2 and 3 (middle column), except they are now separated by population, whereas in the earlier two figures they were separated by mass tracers representing volatility. At a first glance, the population with the highest organic mass is BOC for January and July in both schemes. BOC is the population that contains OC, BC, and sulfate, and is the end result of coagulation of all populations in our idealized cases. However, in the old scheme, populations OCC and OCS also have significant amounts of organics. This is because in the new scheme the emitted populations are ACC, BC1 and OCC, and organics that are emitted in the OCC population can condense on and/or coagulate with other populations, including being lost by evaporation and then repartitioning to other populations. Thus there is an additional loss mechanism of organics from those populations in the new scheme. In addition, there is competition between the ACC and BC1 populations in both schemes, and in the new scheme, aerosol-phase organics in the OCC population could either coagulate with the ACC population to form OCS, or they could coagulate with the BC1 population to form BOC. This competition determines how much OCS and BOC are formed, and it affects how much gas-phase organics from the OCC population could condense on the two populations and the distribution of organics among the populations. Since partitioning adds a loss mechanism to OCC, part of the evaporated mass will go to BOC, making it larger, and a more efficient scavenger of other particles. As a result, most organics coagulate with and condense on the BOC population and/or the OCS population, and together with the emitted OCC population, hold the most organics and dominate the mass fractions. There is some similarity between the January and July results between the new and the old schemes (Figure 5). This similarity means that the distribution of organics among aerosol populations is not significantly affected by season. This is consistent with a study by Bauer et al. (2013), where they found that the mixing state distribution is rather a characteristic of a region and not so much of a season, although the total (absolute) amounts by season may vary. By the end of the simulations, most locations have more organics present in the BOC population, except those in Centreville. The reason for this is sulfate; from the sulfate and black carbon emissions listed in Table 3, we can calculate the sulfate to black carbon ratio in Centreville to be 2:1, higher than the corresponding ratios in all other locations. This high ratio helps the ACC population to survive the competition against BC1 for coagulation with OCC. This leads to higher OCS formation, which is available for gas-phase organics to condense on, thus coagulation and condensation both bring more organics in the

Deleted:

Deleted:

Deleted: e

Deleted:

Deleted: of

Deleted:

OCS population during the first half of the simulation. These results show that the sulfate to black carbon ratio is important for the mixing state by delaying the inevitable BOC domination. Also, comparing the distribution fraction in Figure 5, volatile organics create rather different mixing states as those created by coagulation alone in the original scheme, meaning that the semi-volatility did alter mixing state significantly.

5 5.4 Size distribution

Another important factor on the evolution of aerosols is their size distribution. Shown in Figures 6 and 7 are the January size distributions from Mexico City and Centreville. The first row shows number concentration, the second row surface area, and the third row volume. The first two columns are results from the new and old schemes after 24 hours of simulation, and the right two columns are after 120 hours. The total number concentration, surface area and volume from the eight populations are shown as dotted lines. Note also that these plots show the total aerosol size distribution per population, which includes the contribution of species other than organics.

The size distributions in July are very similar to January in all locations, therefore only January is shown here. Beijing, Hyytiälä and Mexico City exhibit somewhat similar size distributions (with different absolute amounts), just as their mass fractions. The size distribution is dominated by OCC, OCS and BOC in the first 3 to 4 days, but later only by BOC. On the other hand, Centreville, similar to its mixing state, is different in size distribution of different aerosol populations from the other three locations. Therefore, only size distributions of Mexico City and Centreville are shown here.

In the new scheme for Mexico City after 24 hours of simulation, the number concentration has two modes. OCC has even smaller size as Aitken mode sulfate AKK does, as a result of the evaporation of organics, but its number concentration is higher. OCS and BOC have started to form from coagulation of OCC with ACC and BC1, and their diameter, number concentration, surface area and volume are very similar, almost overlapping, with BOC slightly smaller in diameter. After 120 hours of simulation, OCC's number concentration has decreased significantly, from $4 \cdot 10^7 \text{ m}^{-3}$ to $1 \cdot 10^7 \text{ m}^{-3}$. This is because OCC is semi-volatile, it has evaporated and condensed onto other populations, and at the same time its loss due to coagulation with other populations has increased, due to the increase of their number concentration and decrease in size. OCS size grew very slightly, but BOC grew significantly, with peaks of surface area and volume both increasing approximately one order of magnitude. Its peak surface area increased from $1.5 \cdot 10^5 \text{ } \mu\text{m}^2 \text{ m}^{-3}$ to $9 \cdot 10^5 \text{ } \mu\text{m}^2 \text{ m}^{-3}$, and its peak volume grew from approximately $2 \cdot 10^4 \text{ } \mu\text{m}^3 \text{ m}^{-3}$ to $2 \cdot 10^5 \text{ } \mu\text{m}^3 \text{ m}^{-3}$. BOC's growing large surface area is another reason why it has so much organics and dominates the mass concentration: the greater the surface area, the more gas-phase species are able to condense. This matches the mixing state results (Figure 5), where we saw after 24 hours ACC, OCC and BOC have high mass fractions, whereas after 120 hours OCC and OCS are negligible, and more than 90% of the total organic aerosol mass is in the BOC population.

Deleted: 2
Deleted: 6
Deleted: 0.5
Deleted: 6
Deleted: about
Deleted: 3
Deleted: 3
Deleted: 6
Deleted: about
Deleted: 1
Deleted: 5
Deleted: 1
Deleted: 6
Deleted:

In the old scheme, after 24 hours OCC has higher number concentration (peaking at $5 \cdot 10^7 \text{ m}^{-3}$) and size than in the new scheme, and higher surface area and volume, due to its greater number and diameter. OCS and BOC are both fewer in number (peaks are $1 \cdot 10^7 \text{ m}^{-3}$ and $1.5 \cdot 10^7 \text{ m}^{-3}$ lower in the old scheme) but slightly greater in diameter than they are in the new scheme. Later, after 120 hours, OCC decreases in number to a peak at $3.5 \cdot 10^7 \text{ m}^{-3}$, due to coagulation with ACC and BC1 to form more OCS and BOC. Therefore, OCS and BOC increased in number and size, with BOC seeing greater growth (the peak of number concentration increased from $7 \cdot 10^6 \text{ m}^{-3}$ to $2.3 \cdot 10^7 \text{ m}^{-3}$, the peak of surface area increased from $1.5 \cdot 10^5 \text{ } \mu\text{m}^2 \text{ m}^{-3}$ to $1.1 \cdot 10^6 \text{ } \mu\text{m}^2 \text{ m}^{-3}$ and the peak of volume increased from $3 \cdot 10^4 \text{ } \mu\text{m}^3 \text{ m}^{-3}$ to $3.1 \cdot 10^5 \text{ } \mu\text{m}^3 \text{ m}^{-3}$). For OCS we calculated more modest increases of approximately 50% in number, surface area and volume concentration peaks; the number concentration from $4 \cdot 10^6 \text{ m}^{-3}$ to $7 \cdot 10^6 \text{ m}^{-3}$, surface area from $1 \cdot 10^5 \text{ } \mu\text{m}^2 \text{ m}^{-3}$ to $1.5 \cdot 10^5 \text{ } \mu\text{m}^2 \text{ m}^{-3}$ and volume from $2 \cdot 10^4 \text{ } \mu\text{m}^3 \text{ m}^{-3}$ to $4 \cdot 10^4 \text{ } \mu\text{m}^3 \text{ m}^{-3}$, as seen in the new scheme as well. However, BOC's growth in the old scheme is even greater than that in the new scheme. This slightly accelerated growth slows down at later hours (not shown), because BOC dominates faster in the new scheme than in the old one (Figure 4).

The Centreville size distributions tells a different story. In the early stages with the new scheme, OCS has greater number concentration and size than BOC does; OCS's peak number concentration is $0.5 \cdot 10^7 \text{ m}^{-3}$, more than double than that of BOC, while its peak surface area and volume are $1 \cdot 10^5 \text{ } \mu\text{m}^2 \text{ m}^{-3}$ and $2 \cdot 10^5 \text{ } \mu\text{m}^3 \text{ m}^{-3}$, whereas those of BOC are negligible. Later, OCS still outgrows BOC in number, but barely exceeds in surface area and is not greater in volume. BOC shifts to greater diameters, therefore it has greater volume than OCS does after 120 hours. As for the old scheme, OCC does not decrease in number from 24 hours to 120 hours as it does in Mexico City, but its number increases from $1.7 \cdot 10^7 \text{ m}^{-3}$ to $2.5 \cdot 10^7 \text{ m}^{-3}$. This means that in that period of time coagulation loss is less than the amount of OCC emitted, which is what was also seen earlier for the mass concentrations (Figure 2). At 120 hours, OCS has again higher number concentration than BOC does, but only slightly (peak number concentration difference is approximately $1 \cdot 10^6 \text{ m}^{-3}$) and not as much as in the case of Mexico City, and the latter's surface area and volume continue to be greater than those of the former due to its increasing diameter.

6 Conclusions

Organic aerosol volatility calculations were implemented into a new aerosol microphysics scheme, MATRIX-VBS. Results from idealized cases in Beijing, Centreville, Hyytiälä and Mexico City during summer and winter using the new scheme were compared against the original scheme and showed how the inclusion of semi-volatility of organics and their reactivity affected aerosol mass concentration, as well as their mixing state and size distribution. Emission factors, OH oxidation, temperature and total aerosol levels are the key factors determining organics' volatility distribution and mass concentration. The mixing state is affected by particle size and concentration, which determines coagulation and condensation pathways. Results from the

- Deleted: much
- Deleted: 1.2
- Deleted: 2
- Deleted: 6
- Deleted: 6
- Deleted: 1.2
- Deleted: 7
- Deleted: 6-fold,
- Deleted: 3
- Deleted: 6
- Deleted: 1.8
- Deleted: 3
- Deleted: 4.3
- Deleted: 1
- Deleted: 5
- Deleted: 1.6
- Deleted: 6
- Deleted: changes, with
- Deleted: all increasing by less than 50%
- Deleted: 2
- Deleted: 3
- Deleted: 3
- Deleted: 4
- Deleted: 0.8
- Deleted: 5
- Deleted: 1
- Deleted: 5
- Deleted: 2
- Deleted: 2
- Deleted: 6
- Deleted: 2
- Deleted: 6
- Deleted: 0.5
- Deleted: 5
- Deleted: , but not in size
- Deleted: surface area and
- Deleted: 4
- Deleted: 6
- Deleted: 7
- Deleted: 6
- Deleted: less than

new scheme showed different mixing state distribution from the original scheme.

Going forward, the new scheme will be simplified, and we will reduce the number of tracers needed, in order to simplify the model and save computational resources, without losing the essential information needed for volatility. The simplified version of the box model will then be implemented in the NASA GISS ModelE

5 Earth System Model. While this study is purely theoretical, we will evaluate MATRIX-VBS after its implementation into GISS ModelE. We will gain even better understanding of how semi-volatile organics are altering aerosol mixing state, how meteorological conditions and pollution levels influence organics' volatility distribution, as well as their mixing state in the real world, and what implications these processes have on the climate system.

10

Code Availability

This model development is part of GISS ModelE Earth System Model, which is publicly available. [In addition, the box model used here is available on request.](#)

15 Acknowledgements

We thank the NASA Modeling, Analysis, and Prediction program, which supports the GISS ModelE development.

References

- 5 Athanasopoulou, E., Vogel, H., Vogel, B., Tsimpidi, A. P., Pandis, S. N., Knote, C., and Fountoukis, C.: Modeling the meteorological and chemical effects of secondary organic aerosols during an EUCAARI campaign, *Atmos. Chem. Phys.*, 13, 625–645, doi:10.5194/acp-13-625-2013, 2013.
- 10 Bauer, S. E., Wright, D., Koch, D., Lewis, E. R., McGraw, R., Chang, L.-S., Schwartz, S. E. and Ruedy, R.: MATRIX (Multiconfiguration Aerosol TRacker of mIXing state): an aerosol microphysical module for global atmospheric models, *Atmos. Chem. Phys. Discuss.*, 8(3), 9931–10003, doi:10.5194/acpd-8-9931-2008, 2008.
- 15 Bauer, S. E., Ault, A. and Prather, K. A.: Evaluation of aerosol mixing state classes in the GISS modelE-MATRIX climate model using single-particle mass spectrometry measurements, *J. Geophys. Res. Atmos.*, 118, 9834–9844, doi:10.1002/jgrd.50700, 2013.
- 20 Bergström, R., Denier van der Gon, H. A. C., Prévôt, A. S. H., Yttri, K. E., and Simpson, D.: Modelling of organic aerosols over Europe (2002–2007) using a volatility basis set (VBS) framework: application of different assumptions regarding the formation of secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 8499–8527, doi:10.5194/acp-12-8499-2012, 2012.
- 25 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S.K., Sherwood, S., Stevens, B., and Zhang, X.Y.: Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 571–657 pp., doi:10.1017/CBO9781107415324, 2013.
- 30 Donahue, N. M., Robinson, a. L., Stanier, C. O. and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40(8), 2635–2643, doi:10.1021/es052297c, 2006.
- 35 Donahue, N. M., Epstein, S. A., Pandis, S. N. and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics, *Atmos. Chem. Phys.*, 11(7), 3303–3318, doi:10.5194/acp-11-3303-2011, 2011.
- [Epstein, S. A., Riipinen, I. and Donahue, N. M.: A Semi-Empirical Correlation between Enthalpy of Vaporization and Saturation Concentration for Organic Aerosol, *Environ. Sci. Technol.*, 44, 743–748, doi:10.1021/es902497z, 2010.](#)
- 40 Fountoukis, C., Racherla, P. N., Denier van der Gon, H. A. C., Polymeneas, P., Charalampidis, P. E., Pilinis, C., Wiedensohler, A., Dall'Osto, M., O'Dowd, C., and Pandis, S. N.: Evaluation of a three-dimensional chemical transport model (PMCAMx) in the European domain during the EUCAARI May 2008 campaign, *Atmos. Chem. Phys.*, 11, 10331–10347, doi:10.5194/acp-11-10331-2011, 2011.
- 45 Fountoukis, C., Megaritis, A. G., Skyllakou, K., Charalampidis, P. E., Pilinis, C., Denier van der Gon, H. A. C., Crippa, M., Canonaco, F., Mohr, C., Prévôt, A. S. H., Allan, J. D., Poulain, L., Petäjä, T., Tiitta, P., Carbone, S., Kiendler-Scharr, A., Nemitz, E., O'Dowd, C., Swietlicki, E., and Pandis, S. N.: Organic aerosol concentration and composition over Europe: insights from comparison of regional model predictions with

- aerosol mass spectrometer factor analysis, *Atmos. Chem. Phys.*, 14, 9061-9076, doi:10.5194/acp-14-9061-2014, 2014.
- 5 Gery, M.W., Whitten, G. Z., Killus, J. P., and Dodge, M. C.: A photochemical kinetics mechanism for urban and regional scale computer modeling, *J. Geophys. Res.*, 94, 12925–12956, doi:10.1029/JD094iD10p12925, 1989.
- 10 Heald, C. L., Jacob, D. J., Park, R. J., Russell, L. M., Huebert, B. J., Seinfeld, J. H., Liao, H. and Weber, R. J.: A large organic aerosol source in the free troposphere missing from current models, *Geophys. Res. Lett.*, 32(18), 1–4, doi:10.1029/2005GL023831, 2005.
- 15 Hodzic, A., Jimenez, J. L., Madronich, S., Canagaratna, M. R., Decarlo, P. F., Kleinman, L. and Fast, J.: Modeling organic aerosols in a megacity: Potential contribution of semi-volatile and intermediate volatility primary organic compounds to secondary organic aerosol formation, *Atmos. Chem. Phys.*, 10(12), 5491–5514, doi:10.5194/acp-10-5491-2010, 2010.
- 20 Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C., Jimenez, J. L., Madronich, S., and Park, R. J.: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmos. Chem. Phys. Discuss.*, 15, 32413-32468, doi:10.5194/acpd-15-32413-2015, 2015.
- Jathar, S. H., Farina, S. C., Robinson, a. L. and Adams, P. J.: The influence of semi-volatile and reactive primary emissions on the abundance and properties of global organic aerosol, *Atmos. Chem. Phys.*, 11(15), 7727–7746, doi:10.5194/acp-11-7727-2011, 2011.
- 25 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, a S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, a C., Docherty, K. S., Ulbrich, I. M., Grieshop, a P., Robinson, a L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. a, Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, a, Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. a, Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, a, Miyoshi, T., Hatakeyama, S., Shimo, a, Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, a M., Williams, L. R., Wood, E. C., Middlebrook, a M., Kolb, C. E., Baltensperger, U. and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere., *Science*, 326(5959), 1525–1529, doi:10.1126/science.1180353, 2009.
- 35 Jo, D. S., Park, R. J., Kim, M. J., and Spracklen, D. V.: Effects of chemical aging on global secondary organic aerosol using the volatility basis set approach, *Atmos. Environ.*, 81, 230– 244, 2013.
- 40 Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C. and Almeida, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533(7604), 521–526, doi:10.1038/nature17953, 2016.
- 45 Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liou, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K. and Van Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application, *Atmos. Chem. Phys.*, 10(15), 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- 50 Lathière, J., Hauglustaine, D. A., De Noblet-Ducoudré, N., Krinner, G. and Folberth, G. A.: Past and future changes in biogenic volatile organic compound emissions simulated with a global dynamic vegetation model, *Geophys. Res. Lett.*, 32(20), L20818, doi:10.1029/2005GL024164, 2005.

- McGraw, R.: Description of Aerosol Dynamics by the Quadrature Method of Moments, *Aerosol Sci. Technol.*, 27(2), 255–265, doi:10.1080/02786829708965471, 1997.
- 5 Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S. H., Weingartner, E., Riipinen, I., Kulmala, M., Spracklen, D. V., Carslaw, K. S., and Baltensperger, U.: Evidence for the role of organics in aerosol particle formation under atmospheric conditions, *P. Natl. Acad. Sci. USA*, 107, 6646–6651, doi:10.1073/pnas.0911330107, 2010.
- 10 Mezuman, K., Bauer, S. E., and Tsigaridis, K.: Evaluating secondary inorganic aerosols in three dimensions, *Atmos. Chem. Phys.*, 16, 10651-10669, doi:10.5194/acp-16-10651-2016, 2016.
- 15 Murphy, B. N. and Pandis, S. N.: Simulating the formation of semivolatile primary and secondary organic aerosol in a regional chemical transport model, *Environ. Sci. Technol.*, 43(13), 4722–4728, doi:10.1021/es803168a, 2009.
- Murphy, B. N., Donahue, N. M., Fountoukis, C. and Pandis, S. N.: Simulating the oxygen content of ambient organic aerosol with the 2D volatility basis set, *Atmos. Chem. Phys.*, 11(15), 7859–7873, doi:10.5194/acp-11-7859-2011, 2011.
- 20 Murphy, B. N., Donahue, N. M., Robinson, A. L., and Pandis, S. N.: A naming convention for atmospheric organic aerosol, *Atmos. Chem. Phys.*, 14, 5825-5839, doi:10.5194/acp-14-5825-2014, 2014.
- 25 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 659-740 pp., doi:10.1017/CBO9781107415324, 2013.
- 30 Paasonen, P., Asmi, A., Petäjä, T., Kajos, M. K., Äijälä, M., Junninen, H., Holst, T., Abbatt, J. P. D., Arneth, A., Birmili, W., van der Gon, H. D., Hamed, A., Hoffer, A., Laakso, L., Laaksonen, A., Richard Leaitch, W., Plass-Dülmer, C., Pryor, S. C., Räsänen, P., Swietlicki, E., Wiedensohler, A., Worsnop, D. R., Kerminen, V.-M. and Kulmala, M.: Warming-induced increase in aerosol number concentration likely to moderate climate change, *Nat. Geosci.*, 6(6), 438–442, doi:10.1038/ngeo1800, 2013.
- Petters, M. D., Prenni, A. J., Kreidenweis, S. M., DeMott, P. J., Matsunaga, A., Lim, Y. B., and Ziemann, P. J.: Chemical aging and the hydrophobic-hydrophilic conversion of carbonaceous aerosol, *Geophys. Res. Lett.*, 33, L24806, doi:10.1029/2006GL027249, 2006.
- 40 Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, doi:10.5194/acp-7-1961-2007, 2007.
- 45 Pierce, J. R., Riipinen, I., Kulmala, M., Ehn, M., Petäjä, T., Junninen, H., Worsnop, D. R. and Donahue, N. M.: Quantification of the volatility of secondary organic compounds in ultrafine particles during nucleation events, *Atmos. Chem. Phys.*, 11(17), 9019–9036, doi:10.5194/acp-11-9019-2011, 2011.
- Pye, H. O. T. and Seinfeld, J. H.: A global perspective on aerosol from low-volatility organic compounds, *Atmos. Chem. Phys.*, 10(9), 4377–4401, doi:10.5194/acp-10-4377-2010, 2010.
- 50

Formatted: Font: Times New Roman, 12 pt, Font color: Auto

Deleted: Mezuman, K., Bauer, S. E., and Tsigaridis, K.: Evaluating Secondary Inorganic Aerosols in 3-Dimensions, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-199, in review, 2016.

Formatted: Font: Times New Roman, 12 pt, Font color: Auto

Deleted: ¶

Deleted:

5 [Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A., Kürten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petäjä, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H., Sipilä, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tomé, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R., and Baltensperger, U.: Oxidation products of biogenic emissions contribute to nucleation of atmospheric particles, *Science*, 344, 717–721, 2014.](#)

10 [Riipinen, I., Pierce, J. R., Yli-Juuti, T., Nieminen, T., Häkkinen, S., Ehn, M., Junninen, H., Lehtipalo, K., Petäjä, T., Slowik, J., Chang, R., Shantz, N. C., Abbatt, J., Leaitch, W. R., Kerminen, V. -M., Worsnop, D. R., Pandis, S. N., Donahue, N. M., and Kulmala, M.: Organic condensation: a vital link connecting aerosol formation to cloud condensation nuclei \(CCN\) concentrations, *Atmos. Chem. Phys.*, 11, 3865–3878, doi:10.5194/acp-11-3865-2011, 2011.](#)

15 [Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. a, Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R. and Pandis, S. N.: Rethinking organic aerosols: semivolatile emissions and photochemical aging., *Science*, 315\(5816\), 1259–1262, doi:10.1126/science.1133061, 2007.](#)

20 [Sandu, A. and Sander, R.: Technical note: Simulating chemical systems in Fortran90 and Matlab with the Kinetic PreProcessor KPP-2.1, *Atmos. Chem. Phys.*, 6, 187-195, doi:10.5194/acp-6-187-2006, 2006.](#)

25 [Scott, C. E., Spracklen, D. V., Pierce, J. R., Riipinen, I., D'Andrea, S. D., Rap, A., Carslaw, K. S., Forster, P. M., Artaxo, P., Kulmala, M., Rizzo, L. V., Swietlicki, E., Mann, G. W., and Pringle, K. J.: Impact of gas-to-particle partitioning approaches on the simulated radiative effects of biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 15, 12989-13001, doi:10.5194/acp-15-12989-2015, 2015.](#)

30 [Seinfeld, J. H. and Pandis, S. N. : Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, third edition, John Wiley & Sons Inc., Hoboken, New Jersey, 2016.](#)

35 [Shindell, D. T., Grenfell, J. L., Rind, D., Grewe, V. and Price, C.: Chemistry-climate interactions in the Goddard Institute for Space Studies general circulation model: 1. Tropospheric chemistry model description and evaluation, *J. Geophys. Res.*, 106\(D8\), 8047, doi:10.1029/2000JD900704, 2001.](#)

40 [Shindell, D. T., Faluvegi, G. and Bell, N.: Preindustrial-to-present-day radiative forcing by tropospheric ozone from improved simulations with the GISS chemistry-climate GCM, *Atmos. Chem. Phys. Discuss.*, 3\(4\), 3939–3989, doi:10.5194/acpd-3-3939-2003, 2003.](#)

45 [Shrivastava, M. K., Lane, T. E., Donahue, N. M., Pandis, S. N. and Robinson, A. L.: Effects of gas particle partitioning and aging of primary emissions on urban and regional organic aerosol concentrations, *J. Geophys. Res. Atmos.*, 113\(18\), doi:10.1029/2007JD009735, 2008.](#)

50 [Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W., Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G., Rap, a. and Forster, P.: Aerosol mass spectrometer constraint on the global secondary organic aerosol budget, *Atmos. Chem. Phys.*, 11\(23\), 12109–12136, doi:10.5194/acp-11-12109-2011, 2011.](#)

[Tröstl, J., Chuang, W., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F. and Wagner, R.: The role of low-volatility organic compounds in initial particle growth in the atmosphere, *Nature*, 533\(7604\), 527–531, doi:10.1038/nature18271, 2016.](#)

Formatted: Justified

Deleted: ¶

- 5 Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S. E., Bellouin, N., Benedetti, A., Bergman, T., Berntsen, T. K., Beukes, J. P., Bian, H., Carslaw, K. S., Chin, M., Curci, G., Diehl, T., Easter, R. C., Ghan, S. J., Gong, S. L., Hodzic, A., Hoyle, C. R., Iversen, T., Jathar, S., Jimenez, J. L., Kaiser, J. W., Kirkevåg, A., Koch, D., Kokkola, H., H Lee, Y., Lin, G., Liu, X., Luo, G., Ma, X., Mann, G. W., Mihalopoulos, N., Morcrette, J. J., Müller, J. F., Myhre, G., Myriokefalitakis, S., Ng, N. L., O'donnell, D., Penner, J. E., Pozzoli, L., Pringle, K. J., Russell, L. M., Schulz, M., Sciare, J., Seland, Shindell, D. T., Sillman, S., Skeie, R. B., Spracklen, D., Stavrou, T., Steenrod, S. D., Takemura, T., Tiitta, P., Tilmes, S., Tost, H., Van Noije, T., Van Zyl, P. G., Von Salzen, K., Yu, F., Wang, Z., Wang, Z., Zaveri, R. A., Zhang, H., Zhang, K., Zhang, Q. and Zhang, X.: The AeroCom evaluation and intercomparison of organic aerosol in global models, *Atmos. Chem. Phys.*, 14(19), 10845–10895, doi:10.5194/acp-14-10845-2014, 2014.
- 10 Tsimpidi, a. P., Karydis, V. a., Pozzer, a., Pandis, S. N. and Lelieveld, J.: ORACLE (v1.0): module to simulate the organic aerosol composition and evolution in the atmosphere, *Geosci. Model Dev.*, 7(6), 3153–3172, doi:10.5194/gmd-7-3153-2014, 2014.
- 15 Tsimpidi, A. P., Karydis, V. A., Zavala, M., Lei, W., Molina, L., Ulbrich, I. M., Jimenez, J. L. and Pandis, S. N.: Evaluation of the volatility basis-set approach for the simulation of organic aerosol formation in the Mexico City metropolitan area, *Atmos. Chem. Phys.*, 10(2), 525–546, doi:10.5194/acp-10-525-2010, 2010.
- 20 Tsimpidi, A. P., Karydis, V. A., Zavala, M., Lei, W., Bei, N., Molina, L. and Pandis, S. N.: Sources and production of organic aerosol in Mexico City: Insights from the combination of a chemical transport model (PMCAMx-2008) and measurements during MILAGRO, *Atmos. Chem. Phys.*, 11(11), 5153–5168, doi:10.5194/acp-11-5153-2011, 2011.
- 25 Volkamer, R., Jimenez, J. L., San Martini, F., Dzepina, K., Zhang, Q., Salcedo, D., Molina, L. T., Worsnop, D. R. and Molina, M. J.: Secondary organic aerosol formation from anthropogenic air pollution: Rapid and higher than expected, *Geophys. Res. Lett.*, 33(17), doi:10.1029/2006GL026899, 2006.
- 30 Yu, F.: A secondary organic aerosol formation model considering successive oxidation aging and kinetic condensation of organic compounds: Global scale implications, *Atmos. Chem. Phys.*, 11(3), 1083–1099, doi:10.5194/acp-11-1083-2011, 2011.
- 35 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, a., Middlebrook, a. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimo, a., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M. and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34(13), 1–6, doi:10.1029/2007GL029979, 2007.
- 40 Zhang, Q. J., Beekmann, M., Drewnick, F., Freutel, F., Schneider, J., Crippa, M., Prevot, A. S. H., Baltensperger, U., Poulain, L., Wiedensohler, A., Sciare, J., Gros, V., Borbon, A., Colomb, A., Michoud, V., Doussin, J. F., Denier Van Der Gon, H. A. C., Haefelin, M., Dupont, J. C., Siour, G., Petetin, H., Bessagnet, B., Pandis, S. N., Hodzic, A., Sanchez, O., Honoré, C. and Perrussel, O.: Formation of organic aerosol in the Paris region during the MEGAPOLI summer campaign: Evaluation of the volatility-basis-set approach within the CHIMERE model, *Atmos. Chem. Phys.*, 13(11), 5767–5790, doi:10.5194/acp-13-5767-2013, 2013.
- 45

1 Table 1. Naming convention and parameters used in the VBS implementation described here.

Parameter	9 Virtual VBS Species								
C* $\mu\text{g m}^{-3}$ at 298K	10^{-2}	10^{-1}	10^0	10^1	10^2	10^3	10^4	10^5	10^6
Name of volatility bins	M2	M1	M0	P1	P2	P3	P4	P5	P6
Mass-based emission factors applied to POA emissions (Shrivastava et al., 2008)	0.03	0.06	0.09	0.14	0.18	0.30	0.40	0.50	0.80
Enthalpy of vaporization¹	153	142	131	120	109	98	87	76	65

2
3 ¹: [enthalpy of vaporization is calculated using Eqn.12 from Epstein et al. 2010.](#)

Deleted: log(C*)

Deleted: OC

Deleted: OC

Deleted: OC

Deleted: OC

Deleted: OC

Deleted: OC

Deleted: OC

Deleted: OC

Formatted Table

Formatted: Superscript

Deleted: in VBSxnp, x is m for minus and p for plus, n is the absolute value of log(C*), and p is g for the gas-phase and a for the sum across all populations of the aerosol-phase concentration.

Table 2. Aerosol population chemical composition in MATRIX.

Population abbreviation	Description	Composition (constituents other than NH ⁺ ₄ , NO ⁻ ₃ , and H ₂ O)
AKK	sulfate (Aitken mode)	SO ₄ ²⁻
ACC	sulfate (accumulation mode)	SO ₄ ²⁻
OCC	organic carbon	OC, SO ₄ ²⁻
BC1	fresh black carbon (<5% coating)	BC, SO ₄ ²⁻
BC2	aged (by condensation) black carbon (>5% coating)	BC, SO ₄ ²⁻
BCS	aged (by coagulation) black carbon	BC, SO ₄ ²⁻
BOC	black and organic carbon	BC, OC, SO ₄ ²⁻
OCS	organic carbon and sulfate	OC, SO ₄ ²⁻
SSA	sea salt (accumulation mode)	sea salt, SO ₄ ²⁻
SSC	sea salt (coarse mode)	sea salt, SO ₄ ²⁻
DD1	dust (accumulation mode; <5% coating)	mineral dust, SO ₄ ²⁻
DD2	dust (coarse mode; <5% coating)	mineral dust, SO ₄ ²⁻
DS1	dust (accumulation mode; >5% coating)	mineral dust, SO ₄ ²⁻
DS2	dust (coarse mode; >5% coating)	mineral dust, SO ₄ ²⁻
MXX	mixed (all components)	BC,OC, mineral dust, sea salt, SO ₄ ²⁻

*The sigma values for all populations are 1.80, except for AKK, which has a sigma of 1.60, and for SSC and MXX, which both have a sigma of 2.00.

- Deleted: fixed width
- Deleted: μm
- Deleted: SSA and
- Formatted: English (United States)
- Deleted: a
- Deleted: fixed widths
- Deleted: μm
- Formatted: English (United States)

Table 3. Conditions of each location used in the simulations, taken from the GISS ModelE for January and July 2006.

January 2006		Units	Beijing	Centreville	Hyytiälä	Mexico City
Fixed parameters	Temperature	K	270	279	260	283
	Pressure	hPa	1007	996	1009	797
	RH	%	46.8	77.7	79.5	62.5
Gaseous emissions	NO _x		216.5	92.4	169.7	148.7
	CO		6943.3	1199.3	557.3	2308.4
	Alkenes		4.3	0.3	0.1	1.3
	Paraffin	pptv/hr	8.2	2.1	0.6	10.5
	Terpenes		1.8	26.3	9.4	25.8
	Isoprene		1.3	23.8	0.0	0.0
	SO ₂		555.8	191.7	24.1	538.7
	NH ₃		181.3	24.2	50.7	63.3
Aerosol emissions	sulfate		0.06	0.02	0.003	0.05
	black carbon	μg/m ³ /hr	0.09	0.01	0.008	0.03
	organics*		0.19	0.03	0.02	0.11
July 2006		Units	Beijing	Centreville	Hyytiälä	Mexico City
Fixed parameters	Temperature	K	304	303	292	289
	Pressure	hPa	986	995	998	800
	RH	%	59.8	61.8	77.8	83.1
Gaseous emissions	NO _x		281.3	124.3	200.9	165.3
	CO		8111.9	1749.9	630.5	2276.1
	Alkenes		5.0	0.5	0.1	1.3
	Paraffin	pptv/hr	9.6	2.7	0.7	10.7
	Terpenes		36.9	145.4	87.6	44.9
	Isoprene		916.1	795.5	47.2	0.0
	SO ₂		653.7	206.5	26.8	549.5
	NH ₃		211.7	38.7	58.1	63.3
Aerosol emissions	sulfate		0.06	0.02	0.002	0.05
	black carbon	μg/m ³ /hr	0.10	0.01	0.01	0.03
	organics*		0.21	0.03	0.07	0.11

Deleted: P

Deleted: P




Figure 1. Schematic showing coagulation pathways among organic-containing aerosol populations as colored circles, with 9 organic VBS species condensed as grey outer circles. In yellow are the emitted donor aerosol populations, and green are the mixed recipient populations. OCC has a semi-transparent yellow core because it is actually emitted as the VBS species that can serve as condensation medium for gaseous VBS species, represented by the grey outer circles. In orange is population BC2, which contains >5% coating of sulfate and organics, which is formed rapidly from the growth of population BC1, which has <5% sulfate/organics coating.





Figure 2. Temporal evolution of the mass concentration of semi-volatile organics in the gas phase (left column), aerosol phase (across all populations; middle column) and total (right column) using the new scheme (refer to Table 1 for legend) for January. The total of non-volatile organics from the original scheme (OCAR) is shown in black dash-dotted lines in the aerosol phase column (middle). OCAR from the old scheme is exactly 2.5 times smaller from the total organic species in the new scheme.

5



Deleted:

Deleted: right





Figure 3. Same as Figure 2, for July.



Deleted:




Figure 4. Temporal evolution of organic aerosol mass concentration in each organics-containing population from the new scheme (first column for January, third column for July), and the old scheme (second column for January, fourth column for July).

5

- Deleted: left two columns,
- Deleted: first
- Deleted: second
- Deleted: right two columns,
- Deleted: third





Figure 5. Temporal evolution of organic aerosol mass concentration fraction in each organics-containing population from the new scheme (first column for January, [third](#) column for July), and the old scheme ([second](#), [fourth](#) column for January, [fourth](#) column for July).



- Deleted: 1
- Deleted: left two columns,
- Deleted: secon
- Deleted: right two columns, third

5





Figure 6. Organics-containing aerosol populations (except MXX) and AKK (Aitken mode sulfate) size distributions for Mexico City in January. Top row: number concentration, middle row: surface area, bottom row: volume. Total of all populations in dotted black lines.






Figure 7. Same as Figure 6 for Centreville.

