

## Anonymous Referee #1

We would like to thank Reviewer #1 for the very valuable comments and suggestions to improve the manuscript.

The authors have driven the atmospheric model WRF with analysis fields of the operational forecasting system of the ECMWF to simulate the period July to August 2013 in a mid-latitude belt (20°N to 65°N) around the globe with two grid sizes: 0.12° and 0.03°. Both simulations make use of sea surface temperatures from the Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) data. In the 0.03° simulation, parameterisation for deep convection is turned off. In order to evaluate the belt simulations, they are compared to various reference datasets (analysis fields of the operational forecasting system of the ECMWF, EOB-S, CMORPH) and in several sub-domains. In addition, the authors ask specific questions about the added value in the 0.03° simulation.

### General Comments

The paper combines two relatively new innovations, convection permitting simulations and belt simulations. The strengths and weaknesses of both innovations are largely unknown and hence, the paper is worth to be published. However there are two major issues that need to be clarified first:

#### Scientific quality

The solution of a local area model is partly predominated by its lateral boundary conditions (LBCs). The larger the model domain becomes, the weaker becomes the coupling to its LBCs and the larger become large-scale deviations from its driving data in the interior of the model. Kida et al. (1991) and Paegle et al. (1996) are often cited in this context. More recently, Becker et al. (2015) demonstrated that a local area model creates artificial flows to compensate those large-scale deviations in order to achieve physical consistency with the LBCs along the lateral boundaries and that an increase of the model domain does not change this – the artificial flows simply become more complex.

In our configuration, the model is driven only by northern and southern LBCs and the SSTs. This set up makes it easier to differentiate between effects due to LBC (which should be small) and internal model physics.

In contrast, in the study of Becker et al. (2015), the secondary circulation patterns were detected in a 41 year COSMO-CLM downscaling effort using ECHAM5 simulations on a T63 grid as LBCs. Therefore, there are 2 major differences to our study: 1) we do not have a 41 year climatology, as this is simply not possible, and 2) the latitude-belt domain does not have any boundaries in west-east direction preventing deflections from the boundaries. It is also challenging to derive whether the T63 model or the high-resolution LAM is more accurate because partly the internal circulation, which was argued to be induced over the Alpine high mountain range, may be more accurate in the LAM rather than in the T63 simulations. Therefore, the results of Becker et al. (2015) are hardly applicable to our study so that we included it in our introduction as a motivation for performing latitude-belt simulations. Also in the study of Zagar et al. (2013) [Žagar, N., L. Honzak, R. Žabkar, G. Skok, J. Rakovec, and A. Cegljar (2013), Uncertainties in a regional climate model in the midlatitudes due to the nesting

technique and the domain size, *J. Geophys. Res. Atmos.*, 118, 6189–6199, doi:10.1002/jgrd.50525], the location of lateral boundaries in west-east direction disturbed the model performance. The smaller the model domain was, the larger was the influence of the LBCs. As we do have a very large model domain where almost 50% are water surfaces, the SST forcing plays a stronger role compared to the influence of the LBCs. In order to reduce the effect of the LBCs further, we are studying here the performance of a latitude belt simulation what is to our knowledge for the first time.

The study of Kida et al. (1991) describes an alternative way to nest LAMs into coarser resolution models by applying the lateral boundaries in wavenumber space. A prerequisite is that the LAM model to be nested in the coarser model has to be a spectral model like the ALADIN and AROME models from Météo France. As WRF is not a spectral model, this method cannot be used in our case.

In the study of Paegle et al. (1996) the model was forced towards the coarser resolution simulation by applying nudging. Our intention is not to have a time-space interpolator but that the model develops its own balance for process studies and for detecting errors in model physics.

Both of these studies are not applicable, because the purpose of our study is to investigate the behavior of a latitude-belt configuration at two different resolutions which are hardly affected by LBCs in west-east direction. This type of simulation can be seen as an ensemble member of a seasonal forecast system initialized by a global model and forced by observed and simulated SSTs.

As it is well known and subject of many ongoing studies, there is predictive skill up to the seasonal scale due to the memory of the Earth system with respect to ocean circulations, soil moisture distribution, and vegetation properties. Additionally, model performance should improve on the convection permitting scale because land-atmosphere interaction is better represented particularly in heterogeneous terrain, orographic effects are simulated more accurately, and the parameterization of deep convection, which is subject of severe model errors, is turned off (Rotach et al. 2009, Wulfmeyer et al. 2011).

Based on these considerations, we disagree with the reviewers that the comparison of the latitude-belt simulations with our two different resolutions cannot be compared with observations. Deviations with respect to ECMWF analyses should degrade slower and deviate with less rms in the high-resolution model even when forced only with northern and southern LBCs but with SSTs.

Therefore, we consider this study as a first steps towards the analysis of the predictive skill of seasonal ensemble members and partly of future latitude belt dynamical downscaling runs for regional climate simulations. This prospect is currently extensively discussed in the regional climate and seasonal forecast communities for the development of next generation seasonal forecast and regional climate models.

We clarified the first paragraph of the abstract and it reads now:

“Increasing computational resources and the demands of impact modelers, stake holders and society envision seasonal and climate simulations at the convection permitting resolution. So far

such a resolution is only achieved with limited area model whose results are impacted by zonal and meridional boundaries. Here we present the set-up of a latitude-belt domain that reduces disturbances originating from the western and eastern boundaries and therefore allows for studying the impact of model resolution and physical parameterization. The Weather Research and Forecasting (WRF) model coupled to the NOAA land surface model was operated during July and August 2013 at two different horizontal resolutions, namely  $0.03^\circ$  (HIRES) and  $0.12^\circ$  (LOWRES). Both simulations were forced by ECMWF operational analysis data at the northern and southern domain boundaries, and the high-resolution Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) data at the sea surface. The simulations are compared to the operational ECMWF analysis for the representation.”

In the presented belt simulations, there are no western and eastern boundaries and hence, the decoupling becomes an important factor.

This is correct and the analysis of the performance of the model system under these conditions is the goal of our study.

This can be seen in principal in Fig. 9: the model creates significant anomalies in MSLP (low and high pressure systems are created that do not exist and vice versa) in the Atlantic region, but from time to time (e.g. around July 27 and August 10 to 15) the influence of the driving data becomes dominant. The fact that there is some coupling to the LBCs at all comes from the location of the sub-domain: the Atlantic region touches the northern boundary. In the interior of the model domain, the coupling might be much smaller.

We investigated a possible coupling effect from the northern boundaries and reduced the averaging domain to  $60W-10W$  and  $40N-55N$  (1000km away from the northern boundary), the result is very similar to that observed in Figure 10. Additionally we also investigated the 500hPa geopotential height, but again, the behavior is nearly the same independent of the selected domain across the Atlantic. Thus we are confident that the influence of the northern boundary can be neglected compared to SST forcing and internal dynamics.

A short paragraph was added to the discussion section on page 17, line 29:

“Referring to Fig. 10, the potential influence of the northern boundaries was investigated by slightly varying the domain. When selecting a much smaller domain between  $60^\circ W - 10^\circ W$  and  $40^\circ N - 55^\circ N$ , the curve progression of the MSLP and 500-hPa geopotential height is very similar to the behavior shown in Fig. 10 (not shown here). This indicates that the influence of the northern boundaries on the development of the simulation compared to the SST is not significant- especially as the meridional wind speed is very weak in this area.”

Hence, the simulations may be affected by large-scale decoupling to such a large extent that the entire evaluation in its current stage is flawed.

We disagree that this configuration does not allow for evaluating our model runs. The model is not only driven at the LBs but also by the SSTs. Furthermore, some predictive skill of the model is kept up to the seasonal scale. Therefore, we argue that a model with better physics and resolution will demonstrate a better performance and it should be possible to identify problems in model physics. This is an advantage of this model run.

The description of Fig. 10 now reads as follows on page 13, line 3:

“In addition, Figure 10 shows the time series of the averaged MSLP over the North Atlantic between 40°N and 65°N and 60°W and 10°E (white rectangle named Atlantic in Fig. 1. During the first ~10 days, the HIRES simulation (red line) agrees well with the ECMWF analysis while the LOWRES simulation show slightly lower pressure values. After this period, the LOWRES simulation shows considerably lower MSLP compared to the ECMWF analysis while the HIRES simulation is much closer the ECMWF analysis until day 18 of the forecast where both simulations miss the development of a depression. Both simulations are able to capture the pressure drop after 25 days of forecast but the HIRES simulation shows a better agreement with the ECMWF analysis. In the further course, both WRF simulation overestimate the strength of the high-pressure situation with being closer to the analysis again after 45 days. Overall, the LOWRES simulation shows a tendency to even further overestimate the strength of low and high pressure systems. The mean bias of the HIRES simulation during July is 1.6 hPa while it is -0.8 hPa for the LOWRES simulation. In August, the bias of the HIRES simulation stays the same while for the LOWRES simulation it now turns into a positive bias of 2.2hPa. The root mean square error during July is 4.5 hPa and 4.65 hPa for the HIRES and LOWRES simulation, respectively. It further reduces to 3.5 hPa (HIRES) and 3.65 hPa (LOWRES) during August 2013.”

Shifts in time/space between modelled and observed phenomena are limiting the applicability of traditional statistical analysis. Biases and other error measures are showing the summary effect of large-scale decoupling and model deficiencies (which should be the only focus in a model evaluation study).

We agree that this is not a traditional analysis but new model configurations need new ideas of model evaluation. We do not see it as a disadvantage that decoupling takes place but as an advantage to disentangle errors due to boundaries (strongly reduced here), model physics, and model resolution. The model is still forced by SST data and some predictive skill remains up to the seasonal scale (see above).

A common approach to overcome this mismatch is to extend the simulation period to multiple decades and evaluate statistical measures in a climatological way (as it is done for climate models, for instance). However, in the face of high computational costs, this might not be feasible.

As pointed out in the computational setup section on page 7 ff., this is currently impossible due to limited computing and storage resources. This model run was a pioneering, special project at HLRS in order to demonstrate the power of corresponding, future model configurations.

For the purpose of model evaluation it would be enough to demonstrate that the simulations are lying within the bandwidth of possible realistic developments. The climatological year-to-year variability (on a monthly basis), which could be derived from ERA-Interim, or extended ensemble forecast data (e.g. <http://www.ecmwf.int/en/forecasts/datasets/set-vi>) – as the forecast model runs without ingestion of observation – could be used to define such space of possible developments.

This is a great idea. We followed your suggestion and compared the 500 hPa geopotential height anomalies with the biases of the WRF forecasts. The anomalies in Figure 2 show maximum values of 60 gpm during the two month period. Allowing a factor of 2 to identify the space for possible developments and comparing the values shown in Figure 9, the HIRES simulation lies

mostly within this range, while the deviations of the LOWRES simulation are larger. Figure 9 also suggests that the LOWRES simulation starts to drift away earlier from the ECMWF analysis. According to the suggestion of Referee #2, we added an additional plot (Figure 6) showing whether both WRF simulations are within  $\pm 2$  standard deviations of the operational ECMWF analysis in terms of the mean sea level pressure field and 500hPa geopotential height. Both simulations mostly stay well within  $\pm 1$  standard deviations of the ECMWF operational model with advantages, especially during August, of the HIRES simulation.

The following was added to the manuscript on page 11, line 25:

“To further assess the quality of the simulation, Fig. 6 shows the confidence of the WRF simulation biases expressed in terms of ECMWF standard deviations for MSLP and 500-hPa geopotential height indicating that the bias mostly stays within  $\pm 2$  standard deviations of the ECMWF analysis for both variables. The mean value of the deviation expressed in terms of ECMWF standard deviations for the MSLP is 0.22 and 0.36 (HIRES) and 0.29 and 0.43 (LOWRES) for July and August, respectively. For the deviations of the 500 hPa geopotential height the values are 0.21 and 0.24 (HIRES) and 0.21 and 0.31 (LOWRES), respectively.”

Because of the large-scale decoupling that makes the simulations partly incomparable to observational or observation based data, the investigation of added value is flawed, too.

We do not agree with this statement (see above) due to remaining internal forcing by the SSTs and remaining predictive skill up to the seasonal scale.

In addition, the asked questions about added value are way too generally expressed. With one coarse and one fine resolved simulation of the same model, no robust conclusion on added value can be drawn. In such a case, the added value analysis is limited to this specific case.

We agree that based on these model runs, identification of issues with model physics are challenging. However, for the above mentioned reasons we are convinced that our analyses are still valid for the time period of the model runs and confirm an improved performance of the model running on the convection-permitting scale. The benefit is shown by a reduction of the classical scores like bias, RMSE, correlation and Pearson Skill Score of the HIRES simulation compared to the LOWRES experiment. From our results it is clear that a better representation of the terrain and land-use heterogeneity and the possibility of waiving the application of a convection parametrization leads to a better precipitation forecast as shown by the traditional scores and PSS. Most of the applied convection schemes are developed for scales of  $\sim 50$ km and appear do not work properly on resolution at  $0.12^\circ$  as also indicated in the studies of Prein et al. (2015a, Clim Dyn.) and Warrach-Sagi et al. (2013, Clim:Dyn.)

As we are able to trace model errors back to problems with model physics, the results form a basis for more detailed and more extended future studies.

Unfortunately it was not possible to perform longer term simulations. With increasing computational performance, longer simulations can be performed in the future.

By the way, for demonstrating added value, it is not enough to show that biases on a monthly basis are reduced, because monthly biases are the result of multiple processes and phenomena that may take place at the same time and also in sequences. So, a reduction of a monthly bias can be the result of enlarged process and phenomena related biases that are simply cancelling out each other. Hence, demonstrating added value includes a thorough investigation of the underlying processes and phenomena plus a demonstration that these processes and phenomena are more properly captured by the finer resolved model.

While admitting that the evaluation of the models is difficult, a reduction of bias during the simulated time period is important and significant.

When increasing the horizontal resolution of the model, several ambient conditions are improved: 1) the representation of the terrain is much more realistic as compared to the 12 km run. The publication of Prein et al (2015), which was cited by you below, points towards that a resolution increase from 0.44 to 0.11 degree still suffers from the windward-lee effect. Only a further increase to the CP scale gives a chance to considerably improve precipitation (e.g. Prein et al, 2015, Rev. Geophys) 2) The land-use cover, soil texture and its variability is also much more realistically represented on the higher resolution which is absolutely necessary 3) The high-resolution SST combined with better resolved coast lines will improve coastal effects.

As the RMSE (PSS) of mean sea level pressure, geopotential height and especially precipitation are considerably reduced (increased), we are convinced that atmospheric processes and phenomena are more properly captured by the simulation on a convection permitting resolution.

Prein, A. F., W. Langhans, G. Fosser, A. Ferrone, N. Ban, K. Goergen, M. Keller, M. Tölle, O. Gutjahr, F. Feser, et al. (2015), A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges, Rev. Geophys., 53, 323–361.  
doi:[10.1002/2014RG000475](https://doi.org/10.1002/2014RG000475).

To solve this issue, the authors could either include such a thorough process and phenomena based analysis or should put more effort on the model evaluation and its problematic (see above) and do not announce added value in such a prominent way.

We agree that thorough process and phenomena based analysis would strengthen the evaluation of our simulations.

The difficulty is to obtain suitable observations for the whole model domain which allow a fair comparison of e.g. diurnal cycles of temperature, wind, and precipitation. At first glance, the ECMWF data availability chart for conventional observations (see e.g. <http://www.ecmwf.int/en/forecasts/charts/monitoring/dcover?time=2017020100,0,2017020100&obs=synop-ship>) shows a nice coverage, but a closer inspection reveals that most of the stations only report in 3h or 6h intervals. If considering wind and (hourly) precipitation observations, the station density dramatically reduces. From the ECMWF analysis or ERA-Interim data, no diurnal cycles can be displayed since only 6 hourly data are available. Therefore, we would like to keep this suggestion for future studies when suitable model and observational data sets are available.

Since large-scale decoupling plays such an important role, it needs to be an integrative part of the discussion (Section 5) and summary (Section 6).

Yes we agree with your suggestion. The discussion section was reordered and the following was added to address the decoupling on page 16, line 13:

“As the simulations are only driven by high-resolution SST data and no zonal lateral boundaries are applied, this can be isolated to the applied model configurations. The Pacific and North Atlantic are the most sensitive areas with respect to the development of storms (Fig. 7a,b), thus small differences in temperatures due to the applied model physics can lead to different spatial and temporal evolutions of storm systems.”

The summary was also reordered and now contains information on the decoupling starting on page 19, line 13.

### **Presentation quality**

Reference data, error measures, including an explanation why these reference datasets and error measures are selected and how data from different grids is remapped onto a common evaluation grid is missing in the experimental setup (Section 2). Instead this information is (partly) given at other places, for instance at the beginning of the result section (Section 4). Having an evaluation concept in section 2 summarising all of this would increase the readability of the manuscript.

We agree. The paragraph dealing with the observational data set was moved to the experimental setup section on page starting now on page 6, line 33.

### **Specific Comments**

The following study would nicely fit into the introduction section:

Prein, A. F., A. Gobiet, H. Truhetz, K. Keuler, K. Goergen, C. Teichmann, C. Fox Maule, E. van Meijgaard, M. Déqué, G. Nikulin, R. Vautard, A. Colette, E. Kjellström, and D. Jacob (2015), Precipitation in the EURO-CORDEX 0.11° and 0.44° simulations: High resolution, high benefits?, *Climate Dynamics*, 46, 383–412, 10.1007/s00382-015- 2589-y

This paper was added in the introduction as a further motivation to go to the convection-permitting scale because in this work the remaining deficiencies of models running on grids with convection parameterizations are suffering from severe errors such as the windward-lee effect in orographic terrain (Wulfmeyer et al. 2011) making their input almost useless for most end users such as hydrologists. This reference was added to the third paragraph of the introduction on page 2, line 28. We also added the reference of Prein et al. (2015, *Rev. Geophys*) to the introduction on page 3 line 10 being in favor for the necessity of convection permitting scale simulations.

Page 3, lines 14, 15: It is unclear how errors in the large scale circulation patterns can be traced back to the applied physics schemes, especially in the light of large-scale decoupling.

As we did not apply zonal LBCs, the main atmospheric flow in this direction is not disturbed by different physics between the LAM. Thus, deviations in the atmospheric flow can be related

to the applied model as especially the CP resolution is much closer to reality in terms of terrain, coast lines, and land use than the driving model.

We corrected the sentence on page 3, line 35 and it now reads: “As the general circulation is west-east oriented and lateral forcing is only applied at the northern and southern boundaries, e.g. errors in the large scale circulation patterns can be traced back to the applied model with its specific physics schemes. The model physics of the coarser resolution model (ECMWF) providing the lateral boundaries in south-north direction only plays a minor role.”

Page 3, lines 22, 23: It is not clear how large uncertainties over the Atlantic and Pacific can be explained by differences in model physics, especially in the light of large-scale decoupling which is also active when model physics are identical (simply extending the model domain of a local area model by some grid cells into one direction gives different results; see Becker et al., 2015).

Both areas are the most active regions in terms of the Jet stream and tropical storms. Thus small differences e.g. in the temperature fields between WRF and ECMWF can lead to an amplification of the development of weather systems potentially leading to phase shifts and different storm tracks. In case a classic LAM approach is used, the internal variability of the nested model is strongly influenced by the boundary conditions. This is not the case if a latitude-belt is applied.

The paragraph on page 4, line 9 was modified and it reads now:

“Their results indicate a strong influence of the zonal LBCs on the internal model variability due to different model physics and the applied nesting technique. In case the model domain is made smaller and smaller, the RCM does not have the chance to develop its own internal variability and the results are mainly driven by the LBCs. This means that the analysis of the model errors is giving more insights into the applied model in case of a latitude-belt set-up.”

Page 5, line 14: What is the advantage of using OSTIA instead of SST from the operational ECMWF analyses? OSTIA is given on a daily resolution and need to be interpolated in time (see page 5 line 34 to page 6, line 3), while SST from ECMWF is already on a 6 h basis. I am not an SST expert, but a short literature research brought up a paper from Seo et al. (2014) which demonstrates the importance of sub-daily SST variability to properly capture the onset and intensity of Madden–Julian oscillation (MJO) convection in the Indian Ocean in a coupled WRF-ocean model. Seo, H., A. C. Subramanian, A. J. Miller, and N. R. Cavanaugh (2014), Coupled Impacts of the Diurnal Cycle of Sea Surface Temperature on the Madden–Julian Oscillation. *J. Climate*, 27, 8422–8443, doi: 10.1175/JCLI-D-14-00141.1.

The major advantage of OSTIA is the native resolution of  $1/20^\circ$  (5km) while the SST data from the operational analysis would be on the same resolution of as the driving data ( $0.125^\circ$  in our case). The higher resolution of the SST data becomes especially important in coastal regions (e.g. Himada, S., Ohsawa, T., Kogaki, T., Steinfeld, G., and Heinemann, D. (2015), Effects of sea surface temperature accuracy on offshore wind resource assessment using a mesoscale model. *Wind Energy*, 18, 1839–1854. doi: 10.1002/we.1796).

The ECMWF applies the daily OSTIA SST data set at initial time and the SST is kept constant during the operational 10-day forecast (section 8.9 of <http://www.ecmwf.int/sites/default/files/elibrary/2013/9245-part-iv-physical-processes.pdf>).



This also means that the SST data from ECMWF are constant throughout the day (see chapter 12 in <http://www.ecmwf.int/sites/default/files/elibrary/2013/9243-part-ii-data-assimilation.pdf>).

The study of Seo et al. is very interesting, however our study region covers only a small part of the tropics and thus we assume it is feasible to use constant SST data throughout the day. Also this study applies a very coarse resolution together with the necessary convection parametrization which is well known to deteriorate the quality of precipitation forecasts.

The paragraph on page 6, starting line 17 now reads:

“Forcing data at the northern and southern boundaries were provided by 6 hourly ECMWF operational analysis data on model 15 levels and are blended with the default linear decay over five grid points into the WRF model grid. Sea Surface temperatures were provided by the high-resolution Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) data (Donlon et al., 2012) with a resolution of 5km. As this study only contains a small part of the tropics, it appears practicable to use more or less constant SST data for each day. As they are only available in daily intervals, these data were linearly interpolated to the 6 h intervals of the ECMWF analysis. This interpolation was performed by using version 1.7.0 of the Climate Data Operators 20 (CDO;<https://code.zmaw.de/projects/cdo>).

Page 6, lines 11 to 17: the 0.03° and 0.12° simulations make use of pnetcdf. A discussion about pnetcdf is missing here. Does pnetcdf solve the problem?

PNetCDF only reduces the amount of computing time that is spent for I/O. The model results are the same no matter which version of NetCDF is applied. The I/O rates on the system used for this study is around 7GB/s with PNetCDF while it is only around 500MB/s with serial NetCDF. The CDF5 format convection is only available when using PNetCDF. Serial NetCDF does not offer this capability.

The paragraph on page 9, line 5 was enhanced and now reads:

“If even a higher number of grid points is planned to use, one has to take care about the NetCDF limitations in the presently used CDF-2 format. This convention only allows 2<sup>32</sup>-4 bytes per array which can be too small for future experiments so that the new CDF-5 standard has to be considered. This feature is available from PNetCDF version 1.6.0 onwards.

In order to have the possibility to apply such a large domain latitude-belt simulation on the CP scale, we modified the source code by exchanging the second argument of the `nf_create` function from `NF_64BIT_OFFSET` to `NF_64BIT_DATA` in `frame/module_bdywrite.F`. A similar change was performed in `external/io_pnetcdf/wrf_io.F90` In the `NFMPI_CREATE` function, the third argument has to be replaced by `NF_64BIT_DATA`. In the `NFMPI_OPEN` function, `NF_NOWRITE` has to be replaced by `NF_WRITE`.”

Page 8, line 25: What is “good agreement”? What biases are acceptable? (These questions should be tackled in an evaluation concept in section 2.)

We agree that a more precise definition is necessary. Following the study of Kotlarski et al, who evaluated a 20-year forecasting ensemble, a mean sea level pressure bias of 3 hPa is acceptable. For temperature, mean deviations of 3 °C are tolerable in homogenous terrain while for precipitation a relative difference of 100 % is acceptable. Where precipitation amounts are low like in Africa, even a difference of more than 100% is acceptable.

We added this short paragraph to the verification data strategy section on page 7, line 18.

“Following the study of Kotlarski et al. (2014), who evaluated a 20-year forecast ensemble, a mean sea level pressure bias of 15.3 hPa is acceptable. For temperature, mean deviations of up to 3°C are tolerable in homogeneous terrain while for precipitation relative differences of 100% are reasonable. In case of very low precipitation amounts like in North Africa, relative deviations of more than 100% are tolerable.”

Kotlarski, S., Keuler, K., Christensen, O. B., Colette, A., Déqué, M., Gobiet, A., Goergen, K., Jacob, D., Lüthi, D., van Meijgaard, E., Nikulin, G., Schär, C., Teichmann, C., Vautard, R., Warrach-Sagi, K., and Wulfmeyer, V.: Regional climate modeling on European scales: a joint standard evaluation of the EURO-CORDEX RCM ensemble, *Geosci. Model Dev.*, 7, 1297-1333, doi:10.5194/gmd-7-1297-2014, 2014.

Page 11, lines 7 to 8: There must be something fundamentally going wrong with the model in this specific region. Maybe it is related to the initialisation of the soil.

We investigated the soil moisture content in this area. The ECMWF analysis fields and the temporal evolution of the WRF soil moisture data is very similar.

The authors are encouraged to contact WRF experts (e.g. Walter Immerzeel, University of Utrecht, or the CORDEX-South-Asia or CORDEX-Central-Asia communities) that are operating the model in this region. There is also a new reference dataset for temperature available. It is called WFDEI (Weedon et al., 2010; 2011) and can be downloaded from <ftp://rfddata:forceDATA@ftp.iiasa.ac.at>

Weedon, G.P., Gomes, S., Viterbo, P., Österle, H., Adam, J.C., Bellouin, N., Boucher, O., and Best, M., 2010. The WATCH Forcing Data 1958-2001: a meteorological forcing dataset for land surface- and hydrological models. *WATCH Tech. Rep. 22*, 41p (available at [www.eu-watch.org/publications](http://www.eu-watch.org/publications)).

Weedon, G.P., Gomes, S., Viterbo, P., Shuttleworth, W.J., Blyth, E., Österle, H., Adam, J.C., Bellouin, N., Boucher, O., and Best, M., 2011. Creation of the WATCH Forcing data and its use to assess global and regional reference crop evaporation over land during the twentieth century. *J. Hydrometeorol.* 12, 823-848, doi: 10.1175/2011JHM1369.1.

This dataset is based on ERA-Interim, but it corrects temperatures in a way that it is consistent with the observed behaviours of glaciers in the Himalayan region. It might be a more reliable reference dataset than the ECMWF analysis fields.

The WATCH data set is based on ERA-40 downscaled to a 0.5° grid to match the CRU land mask. We compare our model with the ECMWF operational analysis which already considers a lot of 2-m temperatures which are used in the 4DVAR analysis (see e.g. <http://www.ecmwf.int/en/forecasts/charts/monitoring/dcover?time=2016122200,0,2016122200&obs=synop-ship>). To complement 2-m observations, ECMWF also applies assimilation of satellite based surface temperature observations. At this time, the 4DVAR was performed on an outer loop T799 grid with 137 levels (cycle 38r2) while the ERA-Interim analysis is performed on a T255 grid with only 62 levels. Assuming a better resolved terrain and

underlying land-use data set in the operational HTESSEL land-surface model, the operational analysis should be superior to even a corrected ERA-40 analysis on a 0.5° grid.

The large temperature deviations over the Steppe regions are also observed in a study of Zhang et al. (2014, J. Hydromet). They conclude that the soil hydraulic parameters used in the NOAH LSM are inappropriate in steppe regions. Currently these variables are read in from tables so it is difficult to adjust these values for specific regions in case of a large model domain covering different climate regimes. The regions showing large biases of the 2-m temperatures (California, eastern Canada, and China/Mongolia) also exhibit large deviations of the upper soil temperatures compared to the ECMWF analysis fields.. As a 4DVAR also includes requires a forecast model, in this case the ECMWF operational model with different physics compared to WRF, this can also lead to differences between the WRF simulations and the analysis in case the observations density is low.

Another possible factor are the values for the background albedo. ECMWF uses a climatological value while the WRF model offers monthly varying albedos. A closer look into both fields revealed that in these specific areas, the ECMWF albedo values are higher by about 5-10% leading to ~50-100W/m<sup>2</sup> more solar radiation absorbed by the ground.

A reference to the study of Zhang et al. is given in the discussion on page 18, line 16.

“As pointed out by Zhang et al. (2014), the soil hydraulic parameters used in the NOAH LSM show some deficiencies in desert and steppe regions over Inner Mongolia. As our study already makes use of an improved version of the thermal roughness length calculation over land (Chen and Zhang, 2009), it appears that a more proper description of the canopy resistance over the desert steppe can be beneficial. At present, the WRF model system unfortunately does not offer the possibility of latitude or region varying parameters for the land-surface models.”

Zhang, G., G. Zhou, F. Chen, M. Barlage, and L. Xue, 2014: A Trial to Improve Surface Heat Exchange Simulation through Sensitivity Experiments over a Desert Steppe Site. J. Hydrometeor., **15**, 664–684, doi: 10.1175/JHM-D-13-0113.1.

#### Technical Corrections

Page 3, line 16: Is there a reference for the storm systems affecting Europe?

With the citation of Rogers (1997) we added a reference to the storm track climatology over the Atlantic Ocean on page 4, line 3.

Page 3, line 33: typo – “This study is organised . . .”

This was corrected on page 4, line 1314.

Page 4, line 14: Euro-CORDEX should be referenced by Jacob et al. (2014).

Jacob, D., J. Petersen, B. Eggert, A. Alias, O. B. Christensen, L. M. Bouwer, A. Braun, A. Colette, M. Déqué, G. Georgievski, E. Georgopoulou, A. Gobiet, L. Menut, G. Nikulin, A. Haensler, N. Hempelmann, C. Jones, K. Keuler, S. Kovats, N. Kröner, S. Kotlarski, A. Kriegsmann, E. Martin, E. van Meijgaard, C. Moseley, S. Pfeifer, S. Preuschmann, C. Radermacher, K. Radtke, D. Rechid, M. Rounsevell, P. Samuelsson, S. Somot, J.-F. Soussana, C. Teichmann, R. Valentini, R. Vautard, B. Weber, and P. Yiou (2014), EURO-

CORDEX: New high-resolution climate change projections for European impact research, *Regional Environmental Change*, 14, 563–578, 10.1007/s10113-013-0499-2.

This reference is added to the introduction on page 5, line 14.

Page 5, line 18: Are there any references for the studies that have shown the spin-up time for NOAH's land surface model?

E.g. Angevine et al. 2014 [Angevine, W. M., Bazile, E., Legain, D., and Pino, D.: Land surface spinup for episodic modeling, *Atmos. Chem. Phys.*, 14, 8165-8172, doi:10.5194/acp-14-8165-2014, 2014.] performed spin-up experiments using the NOAH LSM. In their study, WRF was driven with coarse resolution ERA-Interim data. Their study indicates that already after one day, the spatial structure of the soil moisture is clearly visible.

We also investigated the soil moisture over Europe. Only minor differences are visible in the first three layers for about 3 weeks between WRF and ECMWF. At the end of July, the WRF soil moisture starts to deviate from ECMWF, probably as a result of different precipitation amounts. The soil moisture in the 4<sup>th</sup> layer is very homogenous although the WRF model shows lower values compared to ECMWF. The 4<sup>th</sup> layer of the ECMWF analysis covers the depth between 100cm and 255cm, while the 4<sup>th</sup> layer of the NOAH model covers the depth between 100cm and 200cm. Therefore the observed slow decay of the ECMWF soil moisture can be explained.

The following was added to the manuscript on page 6, line 24:

“Soil moisture and temperature were initialized from the ECMWF operational analysis. The Hydrology land-surface model HTESSEL (Balsamo et al., 2009) assimilates ASCAT soil moisture data since 2012 (Albergel et al., 2012). A brief comparison of the analyzed ECMWF soil moisture and HIREs soil moisture data over Europe revealed no major differences between both data sets during the first 17 forecast days. The absolute soil moisture content in the three topmost layers is between 0.25 and 0.3 m<sup>3</sup>/m<sup>3</sup> and the differences between HIREs and ECMWF vary around 0.05 m<sup>3</sup>/m<sup>3</sup>. This is very promising especially as ECMWF assimilates ASCAT soil moisture data since 2012 (Albergel et al., 2012). Thus it appears feasible to waive a separate spin-up run for this two month period. Afterwards, the soil moisture shows a different behavior most probably due to different evapotranspiration and precipitation patterns.”

Page 9, line 1 to 2: Is there a reference to the analysis on tropical storms of JMA?

A link to the JMA website (<http://www.jma.go.jp/en/typh/>) was added on page 12, line 13.

Page 9, line 5: Is there a reference to this jet stream north of the Tibetan Plateau which is typical for the summer monsoon?

A reference to Xie et al. (2015, *Journal of Climate*) was added on page 12, line 16.

Page 9, line 10: up to now, it was not clear that RMSE is used at all. It is also not clear which RMSE is used (the RMSE of monthly means or on a daily basis or whatsoever)

Thanks for the comment. As we do not show the RMSE for the sake of brevity, we clarified the sentence that the RMSE is not shown here.

RMSE and standard deviation are calculated on a daily basis. This was clarified in the Verification data strategy section on page 7, line 13. "Standard deviation, bias and RMSE are calculated on a daily basis by comparing the 12Z time steps for each day. The scores are finally averaged over the two month period."

For the verification of precipitation, we also added a sentence on page 8, line 2:

"The RMSE and biases with respect to the CMORPH and E-OBS data sets are calculated from the two month accumulated precipitation over the whole observation domain."

Page 10, line 8: typo – "... due to small ..."

This typo was corrected.

Page 10, line 19: typo – "... caused by the inaccurate ..."

This typo was corrected on page 14, line 1

Page 15, line 29 to 30: cumulus parameterisation is also changed in LOWRES. Is GRIMS also active in LOWRES?

Yes, GRIMS is also active in the LOWRES simulation. We ensured that the description of the model setup on page 5, line 32 is clear on that point.

Figure 2: colours of the shades are too intensive, continents can only hardly be seen. Also the structure of the plot should be consistent with Figure 4 (anomalies should be shaded and the climatology should be in solid lines).

The shaded fields and contour lines were swapped so that the anomalies are shaded and the climatology is shown by contour lines. Also the color of the geophysical border in Figures 2 and 4 has been changed to blue for a better distinction.

## **Anonymous Referee #2**

We would like to thank Reviewer #2 for the valuable comments and suggestions to improve the manuscript.

The authors investigate the benefits of convection permitting modeling by employing the WRF modeling system in a channel configuration over the Northern Hemisphere (between 20° N and 65°N) at resolutions of 0.12° (LOWRES) and 0.03° (HIRES), respectively. In HIRES the deep convection parameterization scheme is turned off. The necessarily short integration period covers the summer of 2013 (July and August). This period was notable for exhibiting a strongly positive phase of the North Atlantic Oscillation and generally weaker subtropical highs over the Atlantic and Pacific basins. The driving data is from the ECMWF operational analysis data at the northern and southern boundaries and the OSTIA 5km SST data set at the sea surface. The authors then compare their results to reanalysis and gridded combined data products (e.g., E-OBS, CMORPH). They aim to answer the three questions: 1) What is the benefit of a CP resolution with respect to the spatial representation of large-scale features in comparison to coarse resolution? 2) Does the higher resolution lead to an improvement of surface variables such as 10m windspeed and 2m temperatures? 3) What is the benefit of the CP resolution with respect to the spatial distribution and amount of precipitation?

Given large channel domain and the fact that convective permitting simulations are still relatively rare this study can potentially make a useful contribution to our understanding of how and why simulations at these grid spacings are useful. However, the experiment is not designed in such a way that it can answer the questions as they are posed. The shortcomings are detailed below as are suggests for improvement. I will focus on what I see as the two major issues that must be addressed before the manuscript can move forward. There are likely more specific comments but these can be addressed in the next revision.

### **General comments**

Due to the channel set up the model simulations are largely “free”.

This is not entirely correct. The simulations are still driven by observed SST data and thus the simulation is only partly free. The relative area covered by the SSTs is increasing with the latitude belt configuration in contrast to an LAM; therefore, increasing the domain and removing the west-east boundaries has also considerable advantages with respect to model performance.

In other words internal variability can account for much of the difference that we see between the simulations and the reference data sets. In fact there is little reason to assume that they would in anyway resemble each other. In the absence of nudging the one-to-one comparison of the model simulations with the reference fields for the large-scale circulation is doomed to fail.

Of course one cannot expect that the model reproduces the observation (ECMWF analysis in this case) one-to-one since in contrast to WRF, ECMWF uses sophisticated data assimilation of a huge set of observations to keep the analysis close to the real observations. However, comparing these simulations with high-resolution (re-)analysis and observational data is the only way to specify how the model is performing on different resolutions and different physics.

As both Reviewers address a similar point here, we are adding the response to reviewer 1:

The purpose of our study is to investigate the behavior of a latitude-belt configuration at two different resolutions which are affected by LBCs in west-east direction. This type of simulation can be seen as an ensemble member of a seasonal forecast system initialized by a global model and forced by observed and simulated SSTs.

As it is well known and subject of many ongoing studies, there is predictive skill up to the seasonal scale due to the memory of the Earth system with respect to ocean circulations, soil moisture distribution, and vegetation properties. Additionally, model performance should improve on the convection permitting scale because land-atmosphere interaction is better represented particularly in heterogeneous terrain, orographic effects are simulated more accurately, and the parameterization of deep convection, which is subject of severe model errors, is turned off (Rotach et al. 2009, Wulfmeyer et al. 2011).

Based on these considerations, we disagree with the reviewers that the comparison of the latitude-belt simulations with our two different resolutions cannot be compared with observations or analyses. Deviations with respect to ECMWF analyses should degrade slower and deviate with less rms in the high-resolution model even when forced only with northern and southern EBCs but with SSTs.

Therefore, we consider this study as a first steps towards the analysis of the predictive skill of seasonal ensemble members and partly of future latitude belt dynamical downscaling runs for regional climate simulations. This prospect is currently extensively discussed in the regional climate and seasonal forecast communities for the development of next generation seasonal forecast and regional climate models.

We clarified this in the first paragraph of the abstract and it reads now:

“Increasing computational resources and the demands of impact modelers, stake holders and society envision seasonal and climate simulations at the convection permitting resolution. So far such a resolution is only achieved with limited area model whose results are impacted by zonal and meridional boundaries. Here we present the set-up of a latitude-belt domain reduces disturbances originating from the western and eastern boundaries and therefore allows for studying the impact of model resolution and physical parameterization. The Weather Research and Forecasting (WRF) model coupled to the NOAH land surface model was operated during July and August 2013 at two different horizontal resolutions, namely 0.03° (HIRES) and 0.12° (LOWRES). Both simulations were forced by ECMWF operational analysis data at the northern and southern domain boundaries, and the high-resolution Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) data at the sea surface. The simulations are compared to the operational ECMWF analysis for the representation.”

This was also clarified in introduction on page 4, line 15:

“The goal of this study is to evaluate the benefit of a unique convection permitting latitude-belt simulation over a 2 month period which is not disturbed by zonal lateral boundaries. Due to its high computational demand it is investigated in comparison with a lower resolution set-up as currently applied in seasonal global forecasts. The simulations were not performed in forecast mode and without data assimilation as the study is considered as a pilot study for future convection permitting seasonal forecasting. In this study, we would like to answer whether a convection permitting latitude-belt simulation for a two month period improves the model performance compared to a commonly applied coarser resolution of 0.12°.”

This is illustrated most clearly through examination of the Figures 4 and 5. The dominant anomalies in the large-scale circulation for 2013 are a weakening of the subtropical highs over the ocean basins and a strengthening of the low-pressure anomalies over the Eurasia. The so-called model biases wipe out this weakening over the subtropical highs and intensify the low-pressure anomalies over the Eurasia. From there the rest of the comparisons are uninformative at best.

Based on the arguments above, we disagree with this particular statement and hope that these are convincing for the reviewers.

Therefore, I would suggest the authors focus on whether this type of simulation is fit for purpose.

Yes, absolutely, the simulation fits our purposes and other research teams working on regional climate and seasonal simulations. Zagar et al. (2013) showed that large disturbances can

occur in case of classical LAM simulations, which is still to date a great matter of concern. Our configuration can be the starting point for reducing basic problems of LAMs.

Žagar, N., L. Honzak, R. Žabkar, G. Skok, J. Rakovec, and A. Ceglar (2013), Uncertainties in a regional climate model in the midlatitudes due to the nesting technique and the domain size, *J. Geophys. Res. Atmos.*, 118, 6189–6199, [doi:10.1002/jgrd.50525](https://doi.org/10.1002/jgrd.50525).

In other words, can the model perform the task for which it is intended and does the HIRES simulation perform this task better, or more accurately than the LOWRES simulation?

See above.

One way the authors could do this would be to show, via hatching for example, areas where the modeled field falls outside the +/- 2 standard deviation confidence bounds of the observations. Given that these simulations are basically single realizations of internal variability, weakly constrained by the lower and north/south boundaries this is a more fair and appropriate comparison.

Thank you for your suggestion. A new Fig. 6 was included indicating that the mean sea level pressure bias stays well within 2 standard deviation of the ECMWF analysis. Additionally, the same type of plots for the 500 hPa geopotential height has been included. Here, the beneficial influence of the higher resolution is clearly visible especially during August.

The following was added to the manuscript on page 11, line 25:

“To further assess the quality of the simulation, Fig. 6 shows the confidence of the WRF simulation biases expressed in terms of ECMWF standard deviations for MSLP and 500-hPa geopotential height indicating that the bias mostly stays within  $\pm 2$  standard deviations of the ECMWF analysis for both variables. The mean value of the deviation expressed in terms of ECMWF standard deviations for the MSLP is 0.22 and 0.36 (HIRES) and 0.29 and 0.43 (LOWRES) for July and August, respectively. For the deviations of the 500-hPa geopotential height the values are 0.21 and 0.24 (HIRES) and 0.21 and 0.31 (LOWRES), respectively.”

Another solution would be to re-run the experiments, but constrain the flow so that expectation could be that the model, in the absence of internal model errors, would reproduce the temporal evolution of the weather over the course of July and August 2013.

Re-running these experiments is simply not possible as this implies blocking half of the current HPC system for almost three days to finish in reasonable time including post processing all data. Also disk space is an issue and is currently not available on the system.

The other issue relates to added value. Given the issues described above and the fact that this is a single model, case study experiment, it is very, very, difficult to convincingly argue for added value. Rather than focus on added value using such measure as Taylor diagrams and RMSE, the authors could perhaps focus more on processes that are more accurately captured in the HIRES simulation. Examples are diurnal cycles of winds and precipitation, blocking associated with heat waves, etc.



In general, it is a good idea to compare diurnal cycles. The difficulty is to obtain suitable observations for the whole model domain which allow a fair comparison. It is especially difficult to obtain observations for wind measurements and hourly precipitation. At first glance, the ECMWF data availability chart for conventional observations (see e.g. <http://www.ecmwf.int/en/forecasts/charts/monitoring/dcover?time=2017020100,0,2017020100&obs=synop-ship>) shows a nice coverage, but a closer inspection reveals that most of the stations only report in 3h or 6h intervals. If considering wind and (hourly) precipitation observations, the station density dramatically reduces. From the ECMWF analysis or ERA-Interim data, no diurnal cycles can be displayed since only 6 hourly data are available. Therefore, we would like to keep this suggestion for future studies when suitable model and observational data sets are available.

The authors should not focus on spatial comparisons as there is little reason to expect high spatial correlation between the simulations and the reference data other than that due to the fact there are climatological patterns that the simulations will somewhat follow. If the authors are really set on showing added value then I would recommend they use something like the Perkins skill score which assesses the similarity of two pdfs (Perkins et al. 2007). This metric is quite a bit more informative than the approaches shown used in the manuscript, which rely heavily on visual inspection.

We do not claim that a high spatial correlation can be expected. Nevertheless, the model should be able to remain close to large scale patterns for the reasons mentioned above. Therefore we consider it as essential to include spatial comparisons here, which also helps to detect model errors.

We followed your suggestion and calculated the Perkins skill score in addition to the other error measures. The result for the averaged August MSLP reveals a value of 0.86 for the LOWRES and 0.91 for the HIRES simulation. For the precipitation over Europe during the whole period, the Perkins score is 0.75 for the LOWRES and 0.84 for the HIRES simulation.

The following sentences were added to the results section on page 11, line 13:

“The PSS of the MSLP over Europe for the HIRES simulation is 0.91 for July and 0.9 for August 2013 whereas the LOWRES simulation yields values of 0.92 and 0.86 for July and August, respectively. Perkins et al. (2007) and Devis et al. (2013) suggest that a PSS of 0.7 indicates a reasonable model performance when compared to reference data sets. Therefore the achieved scores indicate a good performance of both WRF simulations over Europe with better results in the HIRES simulation on longer time scales.”

The following sentence was added on page 15, line 28:

“The PSS during the two month period yields a value of 0.75 for the LOWRES simulation and a value of 0.84 for the HIRES simulation.”

We also added a short paragraph on page 11, line 14.:

“Perkins et al. (2007) and Devis et al. (2013) suggest that a PSS of 0.7 indicates a reasonable model performance when compared to reference data sets. Therefore the achieved scores indicate a good performance of both WRF simulations over Europe with better results in the HIRES simulation on longer forecast lead times.”

Devis, A., N. P. M. van Lipzig, and M. Demuzere (2013), A new statistical approach to downscale wind speed distributions at a site in northern Europe, *J. Geophys. Res. Atmos.*, 118, 2272–2283, doi:10.1002/jgrd.50245.

Perkins, S. E., Pitman, A. J., Holbrook, N. J., & McAneney, J. (2007). Evaluation of the AR4 climate models' simulated daily maximum temperature, minimum temperature, and precipitation over Australia using probability density functions. *Journal of climate*, 20(17), 4356-4376.

### **Specific comments**

The abstract is much too long and without critical insight. The abstract should not just be a laundry list of the results but a brief exposition of key findings. The reader should immediately grasp why this paper is of interest. The contribution this study is making should come through in the abstract.

The abstract was modified so that the intention of our study and the main results of our study are more clearly visible.

Page 5 L3-20: The authors go on about how important soil moisture is but then choose not to spin up soil moisture? This is confusing if, as the authors claim, only 10-14 days are required for spin up. Given that there was a heat wave over Europe in 2013 having the correct soil moisture field would be critical to get the proper atmospheric circulation.

We agree that this is an important topic. With this large domain, we start with an ECMWF soil moisture analysis. While admitting that a spin up run would have been better we found in many of our studies that due to similar physics, ECMWF and NOAH soil moisture is quite similar and that the required spin up time is short.

The following was added to the manuscript on page 6, line 24:

“Soil moisture and temperature were initialized from the ECMWF operational analysis. The Hydrology land-surface model HTESSEL (Balsamo et al., 2009) assimilates ASCAT soil moisture data since 2012 (Albergel et al., 2012). A brief comparison of the analyzed ECMWF soil moisture and HIRES soil moisture data over Europe revealed no major differences between both data sets during the first 17 forecast days. The absolute soil moisture content in the three topmost layers is between 0.25 and 0.3 m<sup>3</sup>/m<sup>3</sup> and the differences between HIRES and ECMWF vary around 0.05 m<sup>3</sup>/m<sup>3</sup>. This is very promising especially as ECMWF assimilates ASCAT soil moisture data since 2012 (Albergel et al., 2012). Thus it appears feasible to waive a separate spin-up run for this two month period. Afterwards, the soil moisture shows a different behavior most probably due to different evapotranspiration and precipitation patterns.”

Page 8 L14: “low pressure” should be replaced with “negative bias”

This was replaced.

Page 8 L19: How is the standard deviation calculated? On mean daily values? Something else? This lack of clarity on calculations appears in other areas of the manuscript as well.

Standard deviation and RMSE values were calculated on a daily basis at the 12Z time steps and finally averaged over the month. This was clarified in the Verification data strategy now starting on page 6.

Page 8 L26: Delete “significantly”. Unless describing the result of a hypothesis test this term should not be used in such a context. There are other areas of the manuscript where this is used.

As we did not perform any statistical significance test, this word is replaced by “considerably”.

Figures

As stated in the general comments the figures could benefit from inclusion of confidence bounds from the reference data.

We included the new Figure 6 to display confidence bounds with respect to the reference data of MSLP and 500hPa geopotential height.

Figure 9 can be removed, as there is no reason to expect these simulations to match the temporal march of the reanalysis.

Due to the inclusion of the new Figure 6, the Figure you refer to is now Figure 10. We decided to keep Figure 10 as this figure nicely shows that the general trend of the HIRES simulation is in a better agreement with ECMWF compared to the LOWRES simulation.

# Continuous high resolution mid-latitude belt simulations for July-August 2013 with WRF

Thomas Schwitalla<sup>1</sup>, Hans-Stefan Bauer<sup>1</sup>, Volker Wulfmeyer<sup>1</sup>, and Kirsten Warrach-Sagi<sup>1</sup>

<sup>1</sup>Institute of Physics and Meteorology, University of Hohenheim, Garbenstrasse 30, 70599 Stuttgart, Germany

Correspondence to: Thomas Schwitalla (thomas.schwitalla@uni-hohenheim.de)

**Abstract.** ~~The impact of a convection permitting (CP) northern hemisphere~~

~~Increasing computational resources and the demands of impact modelers, stake holders and society envision seasonal and climate simulations at the convection permitting resolution. So far such a resolution is only achieved with limited area model whose results are impacted by zonal and meridional boundaries. Here we present the set-up of a latitude-belt simulation with the domain that reduces disturbances originating from the western and eastern boundaries and therefore allows for studying the impact of model resolution and physical parameterization. The Weather Research and Forecasting (WRF) model was investigated during the coupled to the NOAH land surface model was operated during July and August 2013. For this application, the WRF model together with the NOAH land surface model (LSM) was applied 2013 DIFaddend at two different horizontal resolutions, namely 0.03° (HIRES) and 0.12° (LOWRES). The set-up as a latitude-belt domain avoids disturbances that originate from the western and eastern boundaries and therefore allows to study the impact of model resolution and physical parameterizations on the results.~~ Both simulations were forced by ECMWF operational analysis data at the northern and southern domain boundaries, and the high-resolution Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) data at the sea surface.

The simulations are compared to the operational ECMWF analysis for the representation of large scale features. To compare analyze the simulated precipitation, the operational ECMWF forecast, the CPC MORPHing (CMORPH), and the ENSEMBLES gridded observation precipitation data set (E-OBS) were used as references.

~~Compared to the operational high-resolution ECMWF analysis, both simulations are able to capture the large scale circulation pattern though the strength of the Pacific high is considerably overestimated in the LOWRES simulation. Major differences between ECMWF and WRF occur during July 2013 when the lower resolution simulation shows a significant negative bias over the North Atlantic which is not observed in the CP simulation. The analysis indicates deficiencies in the applied combinations of cloud microphysics and convection parametrization on the coarser grid scale in subpolar regions. The overall representation of the 500 hPa geopotential height surface is also improved by the CP simulation compared to the LOWRES simulation apart across Newfoundland where the geopotential height is higher than in the LOWRES simulation due to a northward shift of the location of the Atlantic high pressure system.~~

~~Both simulations show higher wind speeds in the boundary layer by about  $1.5 \text{ m s}^{-1}$  compared to the the ECMWF analysis. Due to the higher surface evaporation, this results in a moist bias of  $0.5 \text{ g kg}^{-1}$  at 925 hPa in the planetary boundary layer compared to the ECMWF analysis. Major differences between ECMWF and WRF occur in the simulation of the 2-m~~

temperatures over the Asian desert and steppe regions. They are significantly higher in WRF by about 5 K both during day and night time presumably as a result of different soil hydraulic parameters used in the NOAA land surface model for steppe regions.

5 The precipitation of the HRES simulation shows a better spatial agreement with CMORPH especially over mountainous terrain. The overall bias reduces from 80 mm at the coarser resolution to 50 mm in the HRES simulation and the Analyzing pressure, geopotential height, wind and temperature fields as well as precipitation revealed: 1) a benefit from the higher resolution concerning the reduction of monthly biases, root mean square error is reduced by about 35% when compared to the CMORPH precipitation analysis. The precipitation distribution agrees much better with the CMORPH data than and an improved Pearson Skill Score 2) deficiencies in the physical parametrizations leading to notable biases in distinct regions like  
10 the Polar Atlantic for the LOWRES simulation which tends to overestimate precipitation, mainly caused by the convection parametrization. Especially over Europe the CP resolution reduces the precipitation bias by about 30% to 20 mm as a result of a better terrain representation and due to the avoidance of the convection parameterization., the North Pacific and Inner Mongolia for both resolutions.

15 In summary, the application of a latitude-belt on a convection permitting resolution shows promising results beneficial for future seasonal forecasting.

## 1 Introduction

On longer time scales like seasonal, decadal, and climate predictions, global General Circulation Models (GCM) are commonly applied with a typical horizontal resolution in the range of 1–2° (Taylor et al., 2012, e.g.). Since it is often desired to have higher resolutions over a region of interest to better represent the land-surface interaction, more and more regional climate models  
20 (RCMs) covering only a subregion of the globe are still applied at a resolution-resolutions between 0.1° and 0.5°.

In the Coordinated Regional Downscaling Experiment CORDEX (<http://www.cordex.org>; Giorgi et al., 2009), several RCMs are applied with grid distances of 0.44° for different continental scale regions around the globe at affordable computing power.

As this resolution may still suffer still suffers from a too coarse horizontal resolution, e.g., the EURO-CORDEX project (<http://www.euro-cordex.net/>) focuses on regional climate simulations for Europe at 0.11° resolution. Studies of Kotlarski et al. (2014) and  
25 Vautard et al. (2013) Evaluation studies of Kotlarski et al. (2014), Vautard et al. (2013), and Prein et al. (2015a) indicated that increasing the resolution from 0.44° to 0.11° results in beneficial or detrimental effects with respect to the simulation of 2-m temperatures with biases in the range of  $\pm 2$  K. However, Kotlarski et al. (2014) show a large model variability with respect to convective precipitation during the summer season over Europe. Further, Prein et al. (2015a) show a strong windward-lee effect in low mountain regions in these coarse RCM simulations.

30 Heikkilä et al. (2011) applied the Weather Research and Forecasting model (WRF) over Norway at 0.33° and 0.11° showing a superior performance of the 0.11° domain with respect to precipitation and 2-m temperatures. Warrach-Sagi et al. (2013) performed a 20 year simulation with the WRF model over Europe at 0.33° and 0.11° resolution where the focus was set on precipitation in Germany. Their study shows an overestimation of precipitation and a higher wet day frequency than ob-

served. The 0.11° simulation shows the windward-lee effect in the low mountain ranges in Germany also observed in a study of Schwitalla et al. (2008) who performed simulations at 7 km horizontal resolution using the MM5 model ~~Grell et al. (1995)~~ [\(Grell et al., 1995\)](#). Due to the application of a convection parameterization, convection ~~was~~ is triggered too early with underestimated peak precipitation rates.

5 As 0.11° resolution can still be too coarse to resolve orographic precipitation, Warrach-Sagi et al. (2013) applied the WRF model with a resolution of 0.0367° during the Convective and Orographically-induced Precipitation Study (COPS; Wulfmeyer et al., 2011) period in Summer 2007. Their study demonstrated a significant improvement with respect to the spatial distribution of precipitation when applying a convection permitting (CP) resolution due to the better resolved terrain and explicit treatment of deep convection. A better spatial distribution of precipitation was also observed in studies of Bauer et al. (2011), Prein et al. 10 (2013), ~~Warrach-Sagi et al. (2013), and Piere et al. (2015)~~ [Piere et al. \(2015\)](#), and [Prein et al. \(2015b\)](#) who clearly identified the benefit of performing convection permitting simulations. ~~seales~~ [Recently, Miyamoto et al. \(2013\) performed global high resolution simulations down to the km-scale for a few hours due to limited computational resources. Their results clearly indicate a benefit from increasing the resolution to ~ 3 km or less. Also Palmer \(2013\) recommended that running climate models on very high horizontal resolution is necessary to represent even the large scale features like Rossby waves.](#)

15 RCMs are either driven by coarser scale models, GCMs, or coarser scale reanalysis data like ERA-Interim (Dee et al., 2011). Therefore the numerical solution of a RCM is driven by the lateral boundary conditions (LBCs) given by the driving model. As the inflow boundaries at coarser grid scales may be imperfect, this can deteriorate the results of the RCM. Laprise et al. (2008) suggested that RCMs require a large model domain to capture all the fine scale features especially in the upper troposphere in mid-latitudes. Schwitalla et al. (2011) evaluated the performance of a limited area WRF set up on a large scale 20 driven precipitation event in summer 2007. The WRF model showed superior performance with respect to the representation of precipitation compared to the smaller domain operational numerical weather prediction (NWP) model of the German Meteorological Service.

Diaconescu and Laprise (2013) tested the effect of different domain sizes on large scale features with simulations at ~ 0.5° horizontal resolution. When RCMs are driven by LBCs containing errors, RCMs can reduce errors in the large scale circulation 25 by applying a large model domain. Problems still can occur as the driving models often contain different physics schemes than the LAM leading to inconsistencies at the boundaries which can penetrate into the model domain [\(Žagar et al., 2013\)](#). [Becker et al. \(2015\) compared a 41-year simulation over Europe using a limited area model and observed secondary circulation patterns when compared to the driving global climate model. These circulations appear to be the result of having domain boundaries in west-east direction.](#)

30 An option to partially overcome the necessity to apply ~~boundary conditions~~ LBCs from a coarser LAM or GCM are channel or latitude-belt simulations. With this type of simulations, it is only required to apply LBCs on the northern and southern boundaries. A typical application is a tropical channel covering an area between 30° S and 30° N. Due to computational constraints, these simulation often have a resolution between 20–30 km [\(e.g. Coppala et al., 2012; Evan et al., 2013; Fonseca et al., 2015\)](#) [\(Coppala et al.,](#) One idea of this special type of simulations is to allow storm systems to cross a whole ocean basin without being truncated 35 by domain boundaries. As the general circulation is west-east oriented [and lateral forcing is only applied at the northern and](#)

southern boundaries, e.g. errors in the large scale circulation patterns can be traced back to the applied model with its specific physics schemes.

Europe is frequently affected by storm systems transiting from Newfoundland towards Europe (Rogers, 1997, e.g.). By applying RCMs, western LBCs can destroy certain features of these storms before they reach Ireland and ~~the~~ Western Europe.

5 Žagar et al. (2013) performed one of the first higher resolution latitude belt simulation covering the northern hemisphere between 35° N and 70° N. They applied the WRF model for a three month period covering January-March 2009. The horizontal resolution was 0.25° which, at this time, was very close to the horizontal resolution of the ECMWF operational model used to force the lateral boundaries. To show the benefit of such a latitude belt, they performed additional LAM simulations with different west-east stretching domain sizes. Their results ~~show the largest uncertainties over the Atlantic and Pacific due to an imperfect nesting of WRF, which~~ indicate a strong influence of the zonal LBCs on the internal model variability due to different model physics and the applied nesting technique. In case the model domain is made smaller and smaller, the RCM does not have the chance to develop its own internal variability and the results are mainly driven by the LBCs. This means that the model physics between WRF and the driving model differ analysis of the model errors is giving more insights into the applied model in case of a latitude-belt set-up.

15 The goal of this study is to evaluate the ~~added value of a convection permitting simulation without any deterioration by LBCs in west-east direction. So far no forecasts on the convection permitting scale in a latitude belt configuration have been performed.~~

~~We are investigating whether a very high-resolution latitude belt domain improves the long-term skill with respect to the large-scale circulation and especially precipitation. In our work, we are addressing the following questions:-~~

- 20
- ~~- What is the benefit of a CP resolution with respect to the spatial representation of large scale features in comparison to coarse resolution?~~
  - ~~- Does the higher resolution lead to an improvement of surface variables such as 10-m wind speed and 2-m temperatures?~~
  - ~~- What is the benefit of the CP resolution regarding the spatial distribution and amount of precipitation?~~

25 benefit of a unique convection permitting latitude-belt simulation over a 2 month period which is not disturbed by zonal lateral boundaries. Due to its high computational demand it is investigated in comparison with a lower resolution set-up as currently applied in seasonal global forecasts. The simulations were not performed in forecast mode and without data assimilation as the study is considered as a pilot study for future convection permitting seasonal forecasting. In this study, we would like to answer whether a convection permitting latitude-belt simulation for a two month period improves the model performance compared to

30 a commonly applied coarser resolution of 0.12°.

This study is organized as follows: Section 2 gives an overview about the technical details ~~and~~, the experimental set-up, and verification strategy followed by a review of the climatological weather situation during the simulation period in section 3. In section 4, a comparison of the large scale circulation against ECMWF operational analysis followed by a comparison of

2-temperatures, 10-m wind speeds and precipitation will be ~~performed~~shown. Section 5 provides a discussion of the results - ~~The and the~~ final section 6 summarizes our results.

## 2 Experimental setup

### Model setup

5 For the experiment, the limited area WRF model (Skamarock et al., 2008) version 3.6.1 was applied. Due to the greater variety of physics options, the fully compressible non-hydrostatic Advanced Research WRF (ARW) is used in this study. In contrast to the most commonly applied limited area grids, a latitude belt was selected for this study. This latitude belt covers the northern hemisphere between 20° N and 65° N and is shown in Figure 1. This is the typical latitude range for weather systems affecting Europe.

10 Two configurations with a latitude-longitude grid are selected: a simulation with 0.12° resolution where convection was parametrized (hereafter named LOWRES) and a convection permitting configuration consisting of 12000×1500 grid cells with a horizontal resolution of 0.03° (hereafter named HIRES). The reason to choose a 0.12° resolution is that the current resolution of the ECMWF operational model is similar and it is also similar to the resolution applied in the EURO-CORDEX experiment (~~Giorgi et al., 2009, e.g.~~)(Jacob et al., 2014, e.g.).

15 Both simulations were performed as dynamical downscaling of the ECMWF analysis with 57 vertical levels, of which 14 levels were within the first 1500 m above ground level, and the model top was set to 10 hPa. The numerical time step was 10 s in the HIRES simulation and 40 s in the LOWRES simulation in order to avoid Courant-Friedrichs-Lewy (CFL) criteria violations in the northern part of the model domain. In addition, the *eps<sub>sm</sub>* parameter ( $\beta$  in the study of Dudhia, 1995) was set to 0.5. This parameter biases the average in vertical wind speed for sound wave computation leading to an increased stability  
20 when the terrain slope is steep.

WRF-ARW offers multiple physics parametrizations. The surface layer above the ground is parametrized by the revised MM5 surface layer scheme of Jimenéz et al. (2012) and is combined with the YSU boundary layer scheme of Hong (2007). The YSU is widely used and extensively evaluated in the WRF community (Nolan et al., 2009; Schwitalla et al., 2011; Shin and Hong, 2011; Milovac et al., 2016, e.g.).

25 Cloud microphysics are parametrized by the Morrison two-moment scheme (Morrison et al., 2009) which includes prognostic variables for liquid and frozen hydrometeors and their corresponding number concentrations. The Morrison scheme is a full 2-moment scheme which is beneficial to represent summertime convection where frozen particles can collect liquid water. This scheme was used during summertime convective precipitation events as shown in studies by Schwitalla and Wulfmeyer (2014) and Bauer et al. (2015a).

30 ~~For the 0.12° simulation, the Kain-Fritsch (KF) cumulus scheme (Kain, 2004) together with the default trigger function was applied.~~The RRTMG longwave and shortwave schemes of Iacono et al. (2008) were applied to parametrize radiation transport. In addition to cloud water, cloud ice and snow, RRTMG interacts with rain water. Shallow convection was parametrized by the GRIMS scheme of Hong et al. (2013) in both simulations. At the lower boundary, the WRF model is coupled to the



land surface model (LSM) NOAH (Chen and Dudhia, 2001; Ek et al., 2003). For the 0.12° simulation, the Kain-Fritsch (KF) cumulus scheme (Kain, 2004) together with the default trigger function was applied in addition. The different physics options are summarized in Table 1.

The representation of the soil texture is crucial when performing simulations on higher resolution. Studies of Warrach-Sagi et al. (2008, 2013), and Acs et al. (2010) indicated that the global FAO (Food and Agricultural Organization of UNO) soil texture data set, which has a resolution of 5' (approx. 10 km), shows significant deviations from high-resolution soil databases. When approaching the convection permitting scale, a soil texture data set at the corresponding resolution is required since the texture determines the soil moisture. We used a modified soil texture data set from Milovac et al. (2014) which is derived from the Harmonized World Soil Database (HWSD) between 60° N and 60° S available at 1 km resolution. Land cover is described by the 20-category MODIS data set from the International Geosphere Biosphere Programme (IGBP) program available at 30'' resolution.

The model set-up applied in this study is very similar to a seasonal forecast system and can be seen as a single realization of a such a forecast ensemble with different resolutions. The operational model of ECMWF was upgraded to cycle 38r2 in 2013. This upgrade includes a change in vertical resolution from 91 to 137 levels with a densification especially in the lowest 1500 m above ground. This increase in vertical resolution becomes more and more important with increasing horizontal resolution. The simulations were performed for a 2 month period starting at 01 July 2013 00 UTC.

Forcing data at the northern and southern boundaries were provided by ~~6h~~ 6 hourly ECMWF operational analysis data on model levels. ~~The LBCs and~~ are blended with the default linear decay over five grid points into the WRF model grid. Sea Surface temperatures were provided by the high-resolution Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) data ~~Donlon et al. (2012)~~ (Donlon et al., 2012) with a resolution of ~ 5km. As this study only contains a small part of the tropics, it appears practicable to use more or less constant SST data for each day. As they are only available in daily intervals, these data were linearly interpolated to the 6 h intervals of the ECMWF analysis. This interpolation was performed by using version 1.7.0 of the Climate Data Operators (CDO;https://code.zmaw.de/projects/cdo).

Soil moisture and temperature were initialized from the ECMWF operational analysis. The Hydrology land-surface model HTESSSEL (Balsamo et al., 2009) ~~of ECMWF operational system includes seven different soil textures which are in a better accordance with the soil textures used in the NOAH LSM as compared to e.g. assimilates ASCAT soil moisture data since 2012 (Albergel et al., 2012).~~ A brief comparison of the analyzed ECMWF soil moisture and HIRES soil moisture data over Europe revealed no major differences between both data sets during the first 17 forecast days. The absolute soil moisture content in the three topmost layers is between 0.25 and 0.3 m<sup>3</sup> m<sup>-3</sup> and the differences between HIRES and ECMWF vary around 0.05m<sup>3</sup> m<sup>-3</sup>. This is very promising especially as ECMWF assimilates ASCAT soil moisture data since 2012 (Albergel et al., 2012). Thus it appears feasible to waive a separate spin-up run for this two month period. Afterwards, the soil moisture shows a different behavior most probably due to different evapotranspiration and precipitation patterns.

### Verification data strategy

Obtaining consistent observations of wind, temperature, and moisture at different altitudes, which are on a comparable resolution to 0.12°, is currently very challenging. They are only available for a few countries and are not homogeneous. Satellite derived products like integrated water vapor, radiation data, and cloud products are not available at the required resolution and spatial and temporal coverage. Therefore the operational ECMWF analysis provides the basis to analyze the results of both WRF simulations with respect to the large scale patterns apart from precipitation where more suitable data sets are available.

The ECMWF analysis is generated by a four-dimensional variational data assimilation system (4DVAR; Rabier et al., 2000). It combines a model background field from a previous forecast with high-resolution observations in order to obtain a high quality gridded analysis field. The 4DVAR at ECMWF includes several different observation types like surface measurements, radio soundings, satellite radiances and aircraft measurements. Regridding of WRF pressure level output to the ECMWF grid at 0.125° was performed with an MPI-compiled version of the Earth System Modelling Framework (ESMF) RegridWeightGen tool<sup>1</sup> within the NCAR Command Language (NCL) framework using bilinear interpolation. ECMWF reanalysis data (Dee et al., 2011), available on a 0.75° grid, are applied for evaluations within CORDEX (Vautard et al., 2013; Warrach-Sagi et al., 2013, e.g.).

Standard deviation, bias and RMSE were calculated on a daily basis using the *dim\_rmsd\_n* and *dim\_stat\_4\_n* functions of NCL by comparing the 12Z time steps for each day. The scores are finally averaged over the two month period. In order to show that both WRF simulations do not deviate too far from the operational analysis, the MSLP and 500 hPa geopotential height bias are expressed in terms of standard deviations of the analysis fields during each month to verify that the bias of both WRF simulations stays within  $\pm 1$  standard deviations.

Following the study of Žagar et al. (2013) where the old TESSEL (Viterbo et al., 1999) was available at ECMWF. Previous studies showed that NOAA's soil moisture and temperatures spin up within 10–14 days in Europe when initialized with HTESSEL. A soil moisture spin-up run was not performed as the main focus. Kotlarski et al. (2014), who evaluated a 20-year forecast ensemble, a mean sea level pressure bias of 3 hPa is acceptable. For temperature, mean deviations of up to 3°C are tolerable in homogeneous terrain while for precipitation relative differences of 100% are reasonable. In case of very low precipitation amounts like in North Africa, relative deviations of more than 100% are tolerable.

For the verification of precipitation, the CPC MORPHing technique (CMORPH; Joyce et al., 2004) data set was applied. It is an almost global precipitation analysis, based on low orbit microwave satellite data. In version 1.0, this product is bias corrected and also uses surface precipitation data where available (blended product). The daily precipitation analysis is available on a 0.25°×0.25° grid. Studies of e.g. Liu et al. (2015) and Stampoulis et al. (2013) show a reasonable correlation of the CMORPH precipitation analysis with ground stations in different regions around the globe. A study of Gebremichael et al. (2014) indicated that although weaknesses in satellite derived precipitation products exist over the mountains, CMORPH performs considerably better than products from the Tropical Rainfall Measuring Mission (TRMM) mission (Huffman et al., 2007). In order to compare the observations with the simulations, the HIRES and LOWRES data were interpolated to the CMORPH grid using the ESMF regridding routines.

A recent study of Skok et al. (2016) showed a superior performance of the E-OBS data (Haylock et al., 2008) over Europe compared to the CMORPH observations. Therefore, the E-OBS precipitation data set was selected for comparisons in Europe.

---

<sup>1</sup><https://www.earthsystemcog.org/projects/regridweightgen/>

To perform the interpolation to the corresponding observation grid, the ESMF tools within NCL have been applied in the same way as for the interpolation of the WRF output to the ECMWF grid. The RMSE and biases with respect to the CMORPH and E-OBS data sets are calculated from the two month accumulated precipitation over the whole observation domain.

5 In addition to the traditional scores, the Perkins Skill Score (PSS, Perkins et al., 2007) was applied for the validation of the mean sea level pressure and European precipitation data. The PSS measures the overlap of two probability density functions (PDF) by calculating the cumulative minimum value of each binned variable. In case both PDFs show an ideal overlap, the PSS would be equal to 1. If there is no overlap between both PDFs then the PSS would be zero.

10 As both WRF simulations are performed without the aid of data assimilation, it cannot be anticipated that they represent single extreme weather events. Nevertheless both simulations are expected to reproduce the large scale weather situation with advantages when applying a convection permitting resolution. The main goal of this study is to compare the different resolutions on a rather short time scale of two months relative to each other rather than specifying whether a certain bias is acceptable.

### Computational aspects

The WRF model simulations were performed at the High Performance Computing Center Stuttgart (HLRS) on the Cray XC40 (<http://www.hlrs.de/systems/cray-xc40-hazel-hen/>). At the time when the simulations were performed, the system consisted of approx. 4000 compute nodes each equipped with 2 Intel 12-Core CPUs with 2.5 GHz clock frequency. The model was compiled with version 14.7 of the Portland Group compiler, Cray MPI 4.3.2 and parallel NetCDF 1.5.0. The total number of cores was partitioned in such a way that each node was filled with four MPI tasks and six OpenMP threads so that in total 14000 MPI tasks were used.

20 The Lustre file system was configured so that 128 object storage targets (OSTs) were used for writing into a single NetCDF file of 92GB size for the HIRES simulation. Further testing revealed that it was not beneficial to use more MPI tasks as this deteriorated the I/O rate which was in the range of ~ 6–7 GByte/s. The total data amount including restart files for the HIRES simulation is about 300TB.

25 The necessary input fields from the ECMWF analysis are about 454 and required high-resolution SST are data about 465 GB in size. It has to be noted that due to limitations in the WPS code of the WRF model system, each GRIB2 file has to be smaller than 2 GB as otherwise this file cannot be fully read in by the ungrib program. The required high-resolution SST data are about 10 GB in total. As they are only available in daily intervals, these data have to be interpolated to the 6 h intervals of the ECMWF analysis. This interpolation was performed by using version 1.7.0 of the Climate Data Operators (CDO;). The time required to download and postprocess-process these data is about 3 days.

30 As the Metgrid interpolation program requires a lot of memory and does not support Parallel NetCDF, the input fields had to be splitted using a value of 102 for `io_form_metgrid` in `namelist.wps` so that each MPI task writes its own small NetCDF file. These files were ingested into the Real program using `io_form_input=102` in the `namelist.input` and the required wall time for Real was about 24 h. The used `namelist.input` for the HIRES simulation is shown in the Appendix.

For the HIRES simulation, 3500 compute nodes (84000 Cores) were used for 3.5 days wall time resulting in  $\sim 0.15$  s for each model time step giving a speed up of 66 compared to real time. The  $0.12^\circ$  simulation was performed on 120 compute nodes and was finished within 31 hours wall time.

If such a high-resolution simulation is considered for operational applications, users have to reduce the output frequency considerably as otherwise the time for writing the files becomes the prevailing process. If even a higher number of grid points is planned to use, one has to take care about the NetCDF limitations in the commonly used CDF-2 format. This convention only allows  $2^{32}-4$  bytes per array which can be too small for future experiments so that the new CDF-5 standard has to be considered. [This feature is available from PNetCDF version 1.6.0 onwards.](#)

[In order to have the possibility to run a large domain latitude-belt simulation on the CP scale, we modified the source code by exchanging the second argument of the `nf\_create` function from `NF\_64BIT\_OFFSET` to `NF\_64BIT\_DATA` in `frame/module\_bdywrite.F`. A similar change was performed in `external/io\_pnetcdf/wrf\_io.F90`. In the `NFMPI\_CREATE` function, the third argument has to be replaced by `NF\_64BIT\_DATA`. In the `NFMPI\_OPEN` function, `NF\_NOWRITE` has to be replaced by `NF\_WRITE`.](#)

Another alternative is to use NetCDF4 with HDF5 support but due to the applied compression, this may require the same time for writing although the file sizes may be considerably smaller than with classic NetCDF. Further information about technical challenges can be found in [\(Bauer et al., 2015b, e.g.\)](#) [\(Bauer et al., 2015b, e.g.\)](#).

### 3 Seasonal statistics

In order to classify the meteorological conditions of summer 2013, climatologies from the ERA-Interim analysis (Dee et al., 2011) were analyzed. Figure 2 displays the mean 500 hPa geopotential [\(solid lines\)](#) together with the anomaly of July and August 2013 compared to 1979–2012 [\(contour lines shaded\)](#). It shows the [subtropical-Azores](#) high with a geopotential height of more than 5900 gpm over the Central Atlantic which is in accordance with the climatological mean of 1979–2012.

Over the northern mid-latitudes, a positive anomaly of the 500 hPa geopotential height is observed over Newfoundland, western United States and the northern Pacific while especially over Europe the 500 hPa geopotential height is significantly higher than the climatological mean. In connection with lower geopotential values over Greenland this leads to stronger wind speeds in the mid-troposphere and thus changes the circulation [pattern-patterns](#) compared to the climatology. The stronger gradient in the 500 hPa geopotential between the northern Pacific and East Asia [support-supports](#) the transport of warm and moist air masses towards North and East Asia, especially as the SSTs are higher than the climatological average during both months (Fig. 3). The location of the jet stream is similar during the simulation period as compared to the climatology. The most remarkable difference is the considerably increased wind speed east of Newfoundland and over Central Asia while the wind speeds are weaker over the Pacific (not shown).

The mid-troposphere [Azores High high](#) extends further towards Central Europe and is also visible at the surface. Fig. 4 shows the average MSLP (contour lines) for the two month period together with the corresponding anomalies (shaded). The MSLP anomaly reaches 3 hPa over Central Europe while at the same time the MSLP bias is negative between Greenland and Iceland

leading to higher low level wind speeds than normal. July and August 2013 were characterized by a strong positive [Northern North Atlantic](#) Oscillation (NAO) index of 0.7 and 1.0 in July and August, respectively. Apparently, the positive SST anomalies (Fig. 3) around 2.5°C over the the [Central-central](#) Atlantic and the northern Pacific are responsible for the lower MSLP over the North Atlantic and the Eastern Pacific as compared to the climatology. Particularly in Europe, there was a heat wave in July leading to dry conditions in Western Europe (Dong et al., 2014). The precipitation amounts over [Central and Northern central and northern](#) Europe were less than 50% of the climatological mean as indicated by the E-OBS (Haylock et al., 2008) precipitation data set (not shown).

## 4 Results

~~Obtaining consistent observations of wind, temperature, and humidity at different altitudes, which are on a comparable resolution to 0.12°, is currently very challenging. They are only available for a few countries and are not homogeneous. Satellite derived products like integrated water vapor, radiation data, and cloud products are often available on grids with resolutions coarser than 0.5°. Also they are mostly only available for a certain region depending on the satellite coverage. Therefore we decided to use the operational ECMWF analysis to compare the results of both WRF simulations with respect to the large scale patterns apart from precipitation where suitable data sets are available.~~

~~The ECMWF analysis is generated by a four-dimensional variational data assimilation system (4DVAR;?). It combines a model background field from a previous forecast with high-resolution observations in order to obtain a high-quality gridded analysis field. The 4DVAR at ECMWF includes several different observation types like surface measurements, radio soundings, satellite radiances and aircraft measurements. ECMWF reanalysis data (Dec et al., 2011), available on a 0.75° grid are widely used for verification of RCMs (Vautard et al., 2013; Warrach-Sagi et al., 2013; Katragkou et al., 2015, e.g.) as they can easily be obtained. As this resolution is too coarse for this high-resolution simulations, we decided to use the operational ECMWF analysis for comparison. The regridding of the WRF output to the ECMWF grid at 0.125° for was performed with an MPI-compiled version of the Earth System Modelling Framework (ESMF) RegridWeightGen tool<sup>2</sup> within the NCAR Command Line (NCL) framework.~~

~~For the verification of precipitation, the CPC MORPHing technique (CMORPH; Joyce et al., 2004) data set was applied. It is an almost global precipitation analysis, based on low-orbit microwave satellite data. In version 1.0, this product is bias-corrected and also uses surface precipitation data where available (blended product). The daily precipitation analysis is available on a 0.25°×0.25° grid. Studies of e.g. Liu et al. (2015) and Stampoulis et al. (2013) show a reasonable correlation of the CMORPH precipitation analysis with ground stations in different regions around the globe. In order to compare the observations with the simulations, the HRES and LOWRES data were interpolated to the CMORPH grid. For Europe, the E-OBS precipitation data set (Haylock et al., 2008) was selected. A recent study of Skok et al. (2016) showed a superior performance of the E-OBS data over Europe compared to the CMOPRH observations.~~

---

<sup>2</sup>

~~As both WRF simulations are performed without data assimilation, it cannot be expected that they represent single extreme weather events. Nevertheless both simulations are expected reproduce to the large scale weather situation reasonably well with advantages when applying a convection permitting resolution.~~

#### 4.1 Large scale circulation

5 Figure 5 shows the comparison of the averaged MSLP at 12 UTC for July and August 2013. In July the ECMWF model shows a strong high pressure system over the Eastern Pacific and a well defined high pressure system over the Atlantic. This is a typical situation during ~~the~~ summer over the northern hemisphere (see Fig. 4).

During July, both WRF simulations are able to capture the general features compared to the ECMWF analysis. Larger differences occur over the Pacific Ocean and over the Central Atlantic where the high pressure systems are located. The intensity of the Pacific high is significantly overestimated in both simulations (Fig. 5g,i) and its location is slightly shifted to the south showing a dipole structure. This behavior was also observed in a study of Cassano et al. (2011) who performed month long simulations using WRF over the polar and subpolar region.

15 The PSS of the MSLP over Europe for the HIRES simulation is 0.91 for July and 0.9 for August 2013 whereas the LOWRES simulation yields values of 0.92 and 0.86 for July and August, respectively. Perkins et al. (2007) and Devis et al. (2013) suggest that a PSS of 0.7 indicates a reasonable model performance when compared to reference data sets. Therefore the achieved scores indicate a good performance of both WRF simulations over Europe with better results in the HIRES simulation on longer forecast lead times.

In both simulations the Atlantic high pressure system extends further to the north towards the Azores ~~islands~~ and also the intensity is overestimated as compared to the ECMWF analysis. The LOWRES simulation shows a ~~low pressure area negative bias~~ east of Greenland. This is not ~~simulated-seen~~ in the HIRES simulation which is in a better accordance with the ECMWF analysis.

20 During August 2013 (right column of Fig. 5) the Pacific high is still overestimated with a bias of more than 5 hPa (Fig. 5h,j). ~~The and the~~ high pressure system over the Atlantic shows a different shape compared to the ECMWF analysis (Fig. 5b,d,f). The strong negative bias over Central Asia is the result of too high 2-m temperatures (see later in section 4.3).

25 To further assess the quality of the simulation, Fig. 6 shows the confidence of the WRF simulation biases expressed in terms of ECMWF standard deviations for MSLP and 500 hPa geopotential height indicating that the bias mostly stays within  $\pm 2$  standard deviations of the ECMWF analysis for both variables. The mean value of the deviation expressed in terms of ECMWF standard deviations for the MSLP is 0.22 and 0.36 (HIRES) and 0.29 and 0.43 (LOWRES) for July and August, respectively. For the deviations of the 500 hPa geopotential height the values are 0.21 and 0.24 (HIRES) and 0.21 and 0.31 (LOWRES),  
30 respectively.

Both WRF simulations show different sensitive regions compared to ECMWF as indicated by the MSLP standard deviation shown in Figure 7 for the different months. The LOWRES experiment exhibits an unrealistically large variability over the Hudson Bay in July associated with a stronger variability of the 850 hPa wind speeds (not shown). The large standard deviation over the North Atlantic shown by the HIRES simulation can be explained by a higher internal variability due to the higher

resolution. Such variability cannot be expected in the ECMWF analysis due to its coarser resolution which does not resolve the ~~belonging corresponding~~ high-resolution dynamical processes. ~~Nevertheless the overall location of the high standard deviation areas in the HRES simulation is in good agreement with the ECMWF analysis.~~

~~The variability of the MSLP over the Western Pacific is significantly overestimated by both WRF experiments. The LOWRES experiment seems to have an even higher tendency to develop tropical storms as seen by the simulated corridor of higher standard deviation. The very high standard deviation over the Aleutian Islands may be related to the higher resolution as these islands consist of volcanoes with elevations up to 2000 m. As they are only partially resolved in the ECMWF model and especially its 4DVAR system running at even coarser resolution, this can explain the different behavior in this region. In combination with higher resolution SSTs and a better represented landmask this also contributes to higher sensitivities.~~

10 Especially in August, the LOWRES simulation tends to exaggerate the development of tropical storms as indicated by the large standard deviation of more than 12 hPa south of Japan. Apparently this is related to the stronger pressure gradient over the West Pacific (Fig. 5). According to the analysis of the Japanese Meteorological Agency (JMA, <http://www.jma.go.jp/en/typh/>), only two tropical storms were present during August in the West Pacific north of 20° N.

15 Figure 8 shows the mean 300 hPa wind speed of both WRF simulations compared with the ECMWF analysis for 12 UTC. During both months, the ECMWF analysis (Fig. 8) shows a well defined subtropical jet stream north of the Tibetan Plateau with an average wind speed of 30 m s<sup>-1</sup> over Central Asia which is typical for the monsoon season ([Xie et al., 2015, e.g.](#)). Also the polar jet over the Pacific and Newfoundland is clearly visible. Compared to the climatology from 1979–2012, the position of the subtropical and polar jet is very similar, with considerably higher wind speeds along the subtropical jet.

20 The large scale structure is captured in both WRF simulations while the HRES simulation shows a weaker maximum over Central Asia as compared to ECMWF. ~~They Both WRF simulations~~ tend to overestimate the intensity of the subtropical jet over the North Pacific but the HRES simulation has a lower RMSE as compared to the LOWRES experiment (~~not shown~~). At the 200 hPa level (not shown here), the wind maximum over Central Asia is simulated more accurately in the HRES simulation. This indicates a possible influence of the better resolved terrain over Asia. As the surface low over the Tibetan plateau is deeper than observed, this can induce a force which moves the subtropical jet ~~further~~ to the north deforming ~~the subtropical jet it~~ as  
25 shown by the reddish colors over the northwest of China and Mongolia in Fig. 8. In addition, the better representation of the Pamir and Tien Shan Mountains in the higher resolution model also play a role in terms of blocking ~~the backward motion southward movement~~ of the jet.

To complement the results for the large scale circulation, Figure 9 shows the mean 500 hPa geopotential height of the ECMWF analysis at 12 UTC time steps. Here a wave like structure with 5–6 stationary waves is visible during both months in  
30 the analysis indicated by the alternating reddish and red colors.

In July, the general features agree in both WRF simulations and the differences partially reflect the the displacement of the low pressure systems shown in Fig. 5. The LOWRES simulation simulates high geopotential over Mongolia and the West Pacific in July 2013 as compared to the ECMWF analysis. This bias further increases in August 2013 exceeding 100 gpm over the North Pacific and Newfoundland (Fig. 9d) as a result of the even stronger displacement of the pressure systems similar to

the results of Cassano et al. (2011). The HIRES simulation also simulates high geopotential at 500 hPa over the Atlantic but the differences over the West Pacific ~~remains are~~ much smaller in August 2013 compared to the LOWRES simulation.

In addition, Figure 10 shows the time series of the averaged MSLP over the North Atlantic between 40° N and 65° N and 60° W and 10° E (white rectangle named Atlantic in Fig. 1). During the first ~ 10 days, the HIRES simulation (red line) agrees well with the ECMWF analysis while the LOWRES simulation show slightly lower pressure values. After this period, the LOWRES simulation shows considerably lower MSLP compared to the ECMWF analysis while the HIRES simulation is much closer the ECMWF analysis until day 18 of the forecast where both simulations miss the development of a depression. Both simulations are able to capture the pressure drop after 25 days of forecast but the HIRES simulation shows a better agreement with the ECMWF analysis. In the ~~further course following~~, both WRF simulation overestimate the strength of the high-pressure situation with being closer to the analysis again after 45 days. Overall, the LOWRES simulation shows a tendency to even further overestimate the strength of low and high pressure systems.

The mean bias of the HIRES simulation during July is 1.6 hPa while it is -0.8 hPa for the LOWRES simulation. In August, the bias of the HIRES simulation stays the same while for the LOWRES simulation it now turns into a positive bias of 2.2 hPa. The root mean square error during July is 4.5 hPa and 4.65 hPa for the HIRES and LOWRES simulation, respectively. It further reduces to 3.5 hPa (HIRES) and 3.65 hPa (LOWRES) during August 2013.

## 4.2 Temperature and moisture in the lower troposphere

The moisture availability in the boundary layer is an important factor for the development of convection and precipitation. As an example, Figure 11a shows the mean 925 hPa water vapor mixing ratio of the ECMWF analysis at 12 UTC. The areas with high moisture ~~availability content~~ over India during the monsoon season and the low amount of water vapor over continental Africa and the African west coast can be recognized by the greenish and blueish colors.

From Figure 11 it is seen that WRF estimates a higher moisture content over the central Pacific with a strong bias of ~ 1.5 g kg<sup>-1</sup>. The same holds for the Gulf of Mexico and the Western Atlantic. There are only minor differences in the moisture content at 925 hPa north of 45° N due to small differences in the MSLP field. Both simulations show similar RMSE values of ~ 2.4 g kg<sup>-1</sup> with an improvement of about 5% in the HIRES simulation (not shown). The ~~highest largest~~ deviations from the ECMWF analysis occur at the east coast of Canada, the North Atlantic, and the West Pacific which are the regions with the highest pressure deviations as these are the ares where the low pressure systems develop. The higher moisture values at the North American east coast can be related to the transportation of humidity from the Gulf of Mexico due to the more intense high pressure system over the Atlantic leading to a stronger southwesterly flow in the lower troposphere. This behavior is sustained at 850 hPa where a similar pattern is observed.

The upper panel of Figure 12 shows the mean temperature of the ECMWF analysis at 925 hPa. The warm air masses transported from the desert towards the Atlantic due to Passat winds can be identified. The LOWRES simulation (Fig. 12b) shows a very strong positive temperature bias exceeding 3 K over Europe, North Africa and the Northwest Pacific. The HIRES simulation also shows a positive temperature bias but it is less pronounced as in the LOWRES simulation and the bias over the Northwest pacific is significantly reduced by about 2 K.



The temperature bias over Newfoundland is caused by the ~~the~~-inaccurate position of the Atlantic high pressure systems which extends too far to the west (see Fig. 5). Due to the different wind direction, warmer air masses from the Gulf of Mexico are advected towards Canada. Another interesting feature is the strong overestimation of 925 hPa temperatures in both WRF simulations at the west coast of California. This is due to an overestimation of wind speeds associated with the stronger pressure gradient which dries out the air coming from Cascade Mountain range.

~~The temperature bias is even higher in the LOWRES simulation (Fig. 12b) because of even stronger winds in the boundary layer.~~In general, the LOWRES simulation shows an even higher temperature bias exceeding 5 K over the North Pacific (Fig. 12b). Note that the average RMSE in the HIRES simulation over Europe is very small with around 3 K at 925 hPa and 850 hPa (not shown). Further the LOWRES simulation does not simulate the tongue of cold air extending from the central North Atlantic towards the west of the Canary islands (indicated by the warm bias south of the Azores in Fig. 12b). The LOWRES simulation tends to overestimate the boundary layer wind speeds in combination with the spatial shift of the high pressure system. Due to the strong high pressure over the Atlantic and the resulting subsidence, warm air masses are transported from the African desert towards the Canary Islands. In addition the insufficient representation of the terrain in the LOWRES simulation ~~as e.g. the mountains are not represented by the LOWRES simulation. This leads~~ can lead to a different circulation pattern than in the HIRES simulation.

### 4.3 Surface fields

Figure 13 shows the mean 2-m temperature for the 12 UTC time steps. The LOWRES simulation shows hardly any bias over the western half of the model domain during July 2013 while the bias considerably grows in August 2013. The HIRES simulation (lower row of Fig.13) shows hardly any bias over the ocean except over the Mediterranean where the model exhibits a cold bias of  $\sim 1-2$  K. In August, both WRF simulations show a similar temperature bias as the pressure gradient at the east coast is very similar.

The 2-m temperatures over the Tarim basin north of the Tibetan Plateau are significantly overestimated as shown in Figs. 13c-f. The simulated skin temperatures (TSK) of both WRF simulations are  $\sim 6$  K higher than in the ECMWF analysis. As the 2-m temperatures are calculated based on the TSK and the second lowest model level, this leads to higher values. In addition, the warm bias over Africa during daytime turns into a cold bias during night time (not shown).

The 10-m wind speeds show a weak bias over the continents (Fig. 14) while larger deviations occur over the Ocean due to the different location of the low and high pressure systems. Especially in the West Pacific, the LOWRES simulation shows a large bias of about  $5 \text{ m s}^{-1}$  during both months while the HIRES simulation is closer to the ECMWF analysis. The deviations in the Atlantic are the results of the slightly larger extent of the high pressure system (see Fig. 5).

### 4.4 Precipitation

The upper panel of Figure 15 shows the CMORPH accumulated precipitation for the 2 month period over land~~only~~. Precipitation in most regions is between 50 mm and 300 mm for the 2 months. The precipitation peaks in the summer monsoon

dominated Southeast Asia and India. The precipitation in the Southern United States is dominated by moist air mass inflow from the Gulf of Mexico in August 2013 (see Fig. 11).

Overall, precipitation amounts are overestimated in both WRF simulations (Fig. 15b,c) apart from the [west-coast-western part](#) of the United States.

5 The LOWRES simulation (Fig. 15b) shows an even stronger overestimation of precipitation in this regions related to the required convection parametrization which is responsible for over 90% of the total precipitation. [Also the LOWRES simulation It also](#) shows a tendency to simulate more widespread precipitation of lower intensities [Schwitalla et al. \(2008\)\(Schwitalla et al., 2008\)](#).

The precipitation maximum over the Korean peninsula caused by the East Asian Monsoon and the maximum over Mexico due to the North American Monsoon are also [well](#) captured in the HIRES experiment. For the whole model domain, the mean estimated precipitation from CMORPH during the two month period is 137 mm, the HIRES experiment simulates 186 mm, and the LOWRES experiment predicts 219 mm within the two months. The variance in both simulations is notably higher as given by the CMORPH analysis (161 mm) and the RMSE is 188 mm for the HIRES simulation and 207 mm for the LOWRES experiment. The average precipitation amount from the operational ECMWF forecasting system (Fig. 15d) is similar to the LOWRES simulation, although the average of 186 mm is more closer to the HIRES simulation.

15 Although both WRF simulations show a positive precipitation bias, it is seen from Fig. 16 that the shape of the precipitation distribution is better represented in the HIRES simulation. Especially the secondary peak in the precipitation amounts of 100 mm is not visible in the LOWRES simulation. The positive benefit of the high-resolution is also seen in the scatter plot displayed in Fig. 17. The regression line is showing a systematic bias of  $\sim 70-80$  mm in both simulations, however the LOWRES regression line has a different slope pointing to an increasing bias with increasing precipitation intensities [which is](#)  
20 [a typical systematic error when applying a convection parametrization \(Schwitalla et al., 2008\)](#).

As also Central Europe together with the Alpine region and the Spanish dry region is of interest in terms of natural disasters caused by droughts and heavy precipitation (Gobiet et al., 2014, e.g.), Fig. 18 displays the accumulated precipitation over Europe for the 2 month period. The E-OBS analysis shows high precipitation amounts induced by orography over southwestern Norway, Central United Kingdom and the Alps with values higher than 175 mm. The low precipitation amount over the Iberian Peninsula with values lower than 20 mm is clearly visible and well simulated by the HIRES experiment. Compared to the mean precipitation of 87.7 mm, the LOWRES simulation overestimates the total precipitation over Europe by 55% while the HIRES simulation only shows an overestimation of 25% in this region. Especially the low precipitation amounts over Spain and Sweden are much better represented compared to the LOWRES simulation (Fig. 18b). [The PSS during the two month period yields a value of 0.75 for the LOWRES simulation and a value of 0.84 for the HIRES simulation. For a qualitative](#)  
30 [comparison, the ECMWF precipitation forecast is shown in addition \(Fig. 18d\)](#).

The precipitation over the Alps is considerably overestimated by almost 100% and also the precipitation amounts over Spain are too high due to the application of a cumulus scheme. In addition, the overestimation due to an inaccurate representation of the terrain is clearly visible in the United Kingdom and southern Scandinavia. Compared to the observation and the HIRES simulation, the LOWRES experiment does not simulate the rain shadow area over Sweden. Although both WRF simulations  
35 show a positive bias, the precipitation distribution is much better represented by the HIRES simulation (Fig. 19).

Summarizing the statistical results, Fig. 20 shows a Taylor diagram for the spatial distribution of precipitation. The different verification regions are marked by the white rectangles shown in Fig. 1. This Taylor diagram combines information about the spatial correlation (azimuth angle) with the normalized centered root mean square error (RMSE, blue circles) and normalized standard deviation (dashed black circles). A perfect model would be at the point marked REF.

- 5 On the global scale, over Europe, and East Asia an indication for applying a CP resolution is given by the lower RMSE and standard deviations. Over Central Asia and North America the benefit is not as clear as the correlation of the LOWRES simulation is better and the bias is not reduced by the higher resolution. Over Africa, the correlation is almost similar but the HIRES simulation tends to slightly underestimate the amount of precipitation.

## 5 Discussion

- 10 The results from section 4 indicate that both simulations show deficiencies in the representation of the large scale and precipitation patterns. However, the convection permitting resolution helps to reduce biases associated with a reduction of the RMSE.

Especially over the northern Pacific, the lower resolution simulation shows major deficiencies with an overstrong subtropical Pacific high showing a MSLP bias of more than 5 hPa. As the simulations are mainly driven by high-resolution SST data and no zonal lateral boundaries are applied, the reasons can be traced back to the applied model configurations. The Pacific and  
15 North Atlantic are the most sensitive areas with respect to the development of storms (Fig. 7a,b), thus small differences in temperatures due to the applied model physics can lead to different spatial and temporal evolutions of storm systems.

- At coarser model resolutions, ~~this problem a strong MSLP bias of more than 10 hPa~~ was also observed in a study of Cassano et al. (2011) and is significantly reduced when a higher horizontal resolution is applied. Studies of Pai Mazumder et al. (2012) over Siberia and Efstathiou et al. (2013) over Greece also observed a tendency to overestimate the intensity of high and  
20 ~~low pressure systems when applying the YSU PBL scheme. The different location of the polar jet over the North Atlantic can result in the transport of warmer and moist air masses from the Central Atlantic towards the north. This can enhance convection over the Atlantic having a strong influence on the simulation of precipitation over Europe. It also intensifies the cyclogenesis which can lead to more severe storms over Europe and also more precipitation over the eastern part of the United States~~

The variability of the MSLP over the Western Pacific is considerably overestimated by both WRF experiments. The LOWRES  
25 experiment seems to have an even higher tendency to develop tropical storms as seen by the simulated corridor of higher standard deviation. The very high standard deviation over the Aleutian Islands may be related to the higher resolution as these islands consist of volcanoes with elevations up to 2000 m. As they are only partially resolved in the ECMWF model and especially its 4DVAR system running at even coarser resolution, this can explain the different behavior in this region. In combination with higher resolution SSTs and a better represented landmask this also contributes to higher sensitivities.

- 30 Although the strong pressure and geopotential height bias over the East Pacific exist in both simulations, Figure 6 gives an indication that on longer time scales the CP resolution has a beneficial impact on the simulation of the large scale patters. The bias expressed in terms of ECMWF standard deviations is considerably smaller the longer the forecast period is.

As the LOWRES simulation shows a totally different large scale pattern in July 2013 compared to the HIRES simulation, the question arises whether the strong negative pressure bias over the North Atlantic simulation is caused by the combination of the applied physics scheme at this particular resolution.

5 A study of Kotlarski et al. (2014) revealed a similar result when comparing the large scale circulation during the summer months averaged over a 20 year period with the same physics combination as for the LOWRES simulation [using a common LAM approach](#). If a different microphysics scheme as e.g. the WSM6 (Hong and Lim, 2006) is applied, the strong sea level pressure bias is not present anymore as seen in the CRP-GL configuration in Kotlarski et al. (2014). As the strong negative pressure bias is ~~also~~ not visible in the HIRES simulation. ~~This, this~~ points towards an unfavorable combination of the Kain-Fritsch convection parameterization with the Morrison microphysics scheme at this particular resolution over the subpolar  
10 regions.

~~Cassano et al. (2011) performed a simulation with exchanging the default Goddard microphysics scheme (Tao and Simpson, 1993) by the Morrison 2-moment scheme combined with the Grell-Devenji convection parameterization (Grell and Dévényi, 2002). The precipitation bias over the Arctic is increased by about 50% by applying this physics combination.~~

The Morrison scheme uses a fixed cloud droplet concentration of  $250 \text{ cm}^{-3}$ . This concentration is adjusted at every model  
15 time step and is set to this constant value at the end of a vertical loop. The ice nucleation follows a formula of Rasmussen et al. (2002) which is primarily designed for mid-latitudes. WRF offers another switch based on observations from the Arctic but this is an on-off switch [for the whole model domain](#). Especially the fixed particle concentration can lead to a more intense formation of optically thick clouds reducing the solar irradiation. E.g. in the Polar-WRF model (Bromwich et al., 2013), the cloud droplet concentration is reduced to  $50 \text{ cm}^{-3}$  to produce fewer liquid water droplets. ~~This points out that~~

20 [The different location of the polar jet over the North Atlantic can result in the transport of warmer and moist air masses from the Central Atlantic towards the north. This can enhance convection over the Atlantic having a strong influence on the simulation of precipitation over Europe. It also intensifies the cyclogenesis which can lead to more severe storms over Europe and also more precipitation over the eastern part of the United States.](#)

[Cassano et al. \(2011\) performed a simulation in which the exchanged the default Goddard microphysics scheme \(Tao and Simpson, 1993\) by the Morrison 2-moment scheme combined with the Grell-Devenji convection parameterization \(Grell and Dévényi, 2002\). The precipitation bias over the Arctic is increased by about 50% by applying this physics combination. Therefore](#) it may be necessary to either adjust these parameters according to the latitude when performing simulations in sub-polar regions or to apply a different combination of cloud microphysics and convection parametrizations at coarser [grid](#) resolutions.

[Referring to Fig. 10, the potential influence of the northern boundaries was investigated by slightly varying the domain. When selecting a much smaller domain between  \$60^\circ \text{ W}\$ – \$10^\circ \text{ W}\$  and  \$40^\circ \text{ N}\$ – \$55^\circ \text{ N}\$ , the curve progression of the MSLP and 500 hPa geopotential height is very similar to the behavior shown in Fig. 10 \(not shown here\). This indicates that the influence of the northern boundaries on the development of the simulation compared to the SST is not significant- especially as the meridional wind speed is very weak in this area.](#)

The general appearance of the boundary layer humidity fields is comparable with the ECMWF analysis ~~–Both but both~~  
35 simulations show a positive water vapor bias of  $0.6 \text{ g kg}^{-1}$  ~~for the HIRES simulation (HIRES)~~ and  $0.4 \text{ g kg}^{-1}$  ~~for the~~

~~LOWRES simulation and a strong moisture bias of more than  $2 \text{ g kg}^{-1}$  over Newfoundland as this is a sensitive region with respect to the development of low pressure systems. Although the overall bias seems to be fairly small, this can have a meaningful influence on the initiation of convection (LOWRES) during the two month period. This is associated with a temperature bias up to the middle troposphere over the West Coast of the US, East Canada and Inner Mongolia.~~

5 ~~The 925 hPa and In the regions where the 2-m temperatures exhibit a large positive bias in both simulations whereas the bias of the LOWRES simulation is even higher especially over the ocean. The large deviation over the Taclamacan and Gobi desert is the result of higher surface temperatures. Compared to the analyzed surface temperatures from ECMWF, the simulated surface temperatures are significantly higher temperature bias is very large (West Coast of the US, East Canada, Mongolia, Inner Mongolia and Xinjiang region), the background albedo used in the WRF simulations. A possible explanation could be the different albedo in the WRF simulation and is 5–10% lower compared to the ECMWF albedo leading to higher absorption of radiation by the surface resulting in higher surface and soil temperatures (Branch et al., 2014, e.g.). The main difference between both albedo data sets is that the ECMWF model. The albedo has a climatological, fixed albedo throughout the year while the albedo used in the WRF model is 0.04 smaller than in the ECMWF analysis thus allowing a 4% higher absorption of radiation and thus higher temperatures (Branch et al., 2014, e.g.). Further, as simulations is time varying. As these regions are~~  
10 ~~covered with snow in winter, this can explain the higher albedo in these areas.~~

15 ~~As pointed out by Zhang et al. (2014), the soil hydraulic parameters used in the NOAH LSM show some deficiencies in desert and steppe regions over Inner Mongolia. As our study already makes use of an improved version of the thermal roughness length calculation over land (Chen and Zhang, 2009), it appears that a more proper description of the canopy resistance over the desert steppe can be beneficial. At present, the WRF model system unfortunately does not offer the possibility of latitude or region varying parameters for the land-surface models.~~

A major advantage of the HIRES simulation is that the precipitation distribution is much better represented compared to the LOWRES simulation. This is especially true for Europe where the simulations were verified against E-OBS data. Here, the HIRES simulation is much closer to the observed precipitation distribution although it also tends to produce spurious precipitation amounts. A reason for the overestimation of precipitation over Asia in the LOWRES simulation are the higher  
25 wind speeds at 10 m over eastern China and the Pacific (Fig. 14) leading to higher evaporation and thus a higher moisture availability (see also Fig. 11). ~~As also the location of the subtropical high is changed, this can also lead to different precipitation patterns. The Together with the applied convection parametrization this can enhance the precipitation amounts considerably. In addition, the~~ LOWRES simulation shows a similar overestimation of precipitation over India, Bangladesh, and Myanmar ~~as compared to the ECMWF operational model, probably related to the convection parameterization. This is not a big surprise as~~  
30 ~~both models use a convection parametrization although the applied schemes differ.~~

Over Central Asia, the benefit of the CP resolution is not clearly visible. This can be related to possible weaknesses in the CMORPH analysis over very complex terrain even when corrected with in-situ observations (Skok et al., 2016, e.g.) (Gebremichael et al., 2016). Another factor influencing the Indian Monsoon can be the role of aerosols but this is beyond the scope of this study. Over North America, the HIRES simulation shows a slightly worse correlation compared to the LOWRES experiment due to an overesti-

mation of precipitation over the eastern United States (Fig. 15b) which apparently can be connected to a moist inflow bias at 925 hPa (Fig. 11b).

The overestimation of precipitation over the eastern part of the United States in both WRF simulations is related to the shift of the pressure system. This shift allows a moist inflow from the Atlantic and the Gulf of Mexico leading to higher precipitation amounts which are not simulated by the operational ECMWF forecasts.

On the global scale, over Europe, and East Asia an indication for applying a CP resolution is given by the lower RMSE and standard deviations. Over Central Asia and North America the benefit is not as clear as the correlation of the LOWRES simulation is better and the bias is not reduced by the higher resolution. Here, a better resolved terrain might deteriorate the situation with inaccurate soil hydraulic parameters. Over Africa, the correlation is almost similar but the HIRES simulation tends to slightly underestimate the amount of precipitation. Following Perkins et al. (2007) and Devis et al. (2013), the improved PSS clearly points towards the application of the CP resolution.

## 6 Summary

Two ~~latitude belt simulations with WRF~~ latitude-belt simulations with the WRF-ARW model between 20° N and 65° N, ~~one have been evaluated for July and August 2013. One simulation was performed at 0.12° resolution, one at typical for currently applied RCMs and the other simulation was performed on~~ the convection permitting scale with a resolution of 0.03° ~~were analyzed for July and August 2013. Such high-resolution simulations require computing resources of 84000 cores for 3.5 days and therefore are often limited to shorter periods or even case studies.~~ Meridional boundaries were provided every six hours by the ECMWF operational analysis. The lower boundary forcing is provided by daily high-resolution SST data from the OSTIA project interpolated to six hourly intervals.

Nevertheless, they are undisturbed by ~~lateral boundary conditions at the western and eastern domain boundaries as in limited area model~~ LBCs in west-east direction as in commonly applied LAM applications and therefore allow for new insights into model resolution dependence of the results. Further, since the results now depend on model physics and resolution only, the results can be assessed with respect to the model performance itself rather than the domain size and inconsistencies of model physics at the ~~meridional-zonal~~ boundaries. This is important since as e.g. Eurasia and North America are characterized by the impact of the polar and subtropical jets and the sea surface temperatures of the Atlantic and Pacific Gyres, namely the Gulfstream and Kuroshio current on the general atmospheric circulation.

The simulations were compared to ECMWF operational analyses data at 0.12° resolution and to observational precipitation data sets of CMORPH and E-OBS at 0.25° resolution, since ~~precipitation data is not available worldwide~~ precipitation data sets are not available at higher resolution.

The objective of the study was to ~~answer the questions posed in the introduction. The results are as follows:-~~

What is the benefit of a CP resolution with respect to the spatial representation of large scale features in comparison to coarse resolution? evaluate the performance and benefit of applying a convection-permitting resolution when performing latitude-belt simulations over an extended forecast period.

A benefit of the higher model resolution can be seen in ~~a reduced bias and RMSE of the pressure and 500 hPa geopotential height fields during both months~~ the pressure fields of both months in terms of a bias reduction. The spatial distribution ~~of the errors is not impacted by the resolution. The 925 hPa temperature shows an added value in the Pacific and Atlantic Ocean, but no added value is seen in the 925 hPa humidity.~~

5 ~~Does the higher resolution lead to an improvement of surface variables such as 10-m wind speed and 2-m temperatures?~~ between both WRF simulations do greatly differ over the Atlantic during the first month while there is a considerable improvement at longer forecast periods at higher resolution over the Pacific indicated by bias reduction and improvement of the Perkins Skill Score. The CP resolution also seems to be able to better capture the middle and upper tropospheric features like e.g. the location of the subpolar jet stream. Also the temperature and moisture bias is slightly reduced in the lower troposphere although  
10 an average bias of  $0.5 \text{ g kg}^{-3}$  and 1.5 K still remains.

Over land both simulations show the same biases in the ~~2m-temperature and 10m-~~ 2-m temperature and 10-m wind speeds. This indicates that the biases are subject to the physical parameterization schemes of WRF, namely those describing the sub-grid scale land-atmosphere feedback processes. ~~Concerning the 10m wind speed HRES shows an added value over the oceans and are inferior to the applied resolution.~~

15 ~~What is the benefit of the CP resolution regarding the spatial distribution and amount of precipitation?~~ HRES shows an added value ~~The CP resolution shows an enhancement~~ concerning the precipitation amount in the whole domain except Northern America, where LOWRES compares better with the studied precipitation data sets. ~~In Europe HRES~~ Over Europe, the CP resolution results in an improved pdf distribution of precipitation amounts indicated by a higher Perkins Skill Score and considerably lower bias and RMSE. Concerning the spatial correlation ~~no added value was gained from the~~  
20 ~~HRES simulation, however this is expected when comparing to the coarse observational data sets.~~

~~All in all the study reveals that a high resolution improves the general circulation, but is not sufficient to tackle the biases in long term simulations concerning the surface variables. Though computing resources are still growing, LAM will still be required for climate simulations. This study, there is no clear signal for a major improvement of the higher resolution but one~~ has to bear in mind the rather coarse resolution of the precipitation data sets.

25 This study also showed that the physical parameterizations need to be assessed to provide more accurate simulations of the climate and also to provide less biased surface variables to the impact models as required by the society. Namely the land-atmosphere feedback and interactions need to be investigated in a synergy of novel high resolution observational data (e.g. from the Surface-Atmosphere-Boundary-Layer-Exchange experiment, Wulfmeyer and Coauthors, 2015) seamless model applications down to LES and new evaluation techniques (Wulfmeyer et al., 2016, e.g.) to improve the physical parameterization  
30 schemes on the applied model resolution.

As computing resources are continuously growing, the potential for an increase of the simulation period for such CP latitude-belt simulations is given.

## Code availability

To download the WRF source code, users need to register on the following website:

<http://www2.mmm.ucar.edu/wrf/users/download/wrf-regist.php>. Apart from the default required NetCDF and MPI libraries, users need to install the PNetCDF libraries version 1.5.0 or higher from the Argonne National Laboratory

5 (<https://trac.mcs.anl.gov/projects/parallel-netcdf>).

## Appendix A: namelist.input used for the WRF simulations

The following namelist.input was used for both simulations. For the LOWRES simulation, the time step, grid resolution, and number of grid cells need to be adjusted. In addition, `cu_physics` needs to be set to zero.

```
&time_control
10  run_days           = 0,
    run_hours         = 0,
    run_minutes       = 0,
    run_seconds       = 0,
    start_year        = 2013
15  start_month       = 07
    start_day         = 01
    start_hour        = 0
    start_minute      = 0
    start_second      = 0
20  end_year          = 2013
    end_month         = 09
    end_day           = 01
    end_hour          = 0
    end_minute        = 0
25  end_second        = 0
    interval_seconds  = 21600,
    input_from_file   = . true .
    history_interval  = 30,
    frames_per_outfile = 1
30  restart           = . false .,
    restart_interval  = 720,
    override_restart_timers = . true .
```



```

io_form_history           = 11,
io_form_restart          = 11,
io_form_input            = 102,
io_form_boundary         = 11,
5 io_form_auxinput1      = 11,
debug_level              = 0,
nocolons= . true .
io_form_auxinput4        = 11
auxinput4_inname         = "wrflowinp_d<domain>"
10 auxinput4_interval     = 360
auxhist23_outname='wrfpress_d<domain>_<date>'
io_form_auxhist23 = 11
auxhist23_interval = 30,
frames_per_auxhist23 = 1
15 diag_print=1,
auxhist2_outname='afwa_d<domain>_<date>'
io_form_auxhist2 = 11
auxhist2_interval = 15,
frames_per_auxhist2 = 1
20 use_netcdf_classic =. true .
/

&diags
p_lev_diags              = 1
25 num_press_levels      = 7
press_levels             = 92500, 85000, 70000, 50000, 30000, 20000, 10000
use_tot_or_hyd_p         = 2
/

&domains
30 time_step              = 10
time_step_fract_num      = 0
time_step_fract_den      = 1
max_dom                  = 1
s_we                     = 1
35 e_we                   = 12000

```

```

s_sn                = 1
e_sn                = 1500
s_vert              = 1
e_vert              = 57
5 eta_levels = 1.000,0.997,0.993,0.989,0.983,0.972,0.962,0.952
,0.942,0.932,0.917,0.903,0.889,0.875,0.852,0.826,0.799,0.771,
0.748,0.725,0.7,0.678,0.653,0.628,0.590,0.557,0.515,0.480,
0.445,0.410,0.375,0.340,0.305,0.280,0.25,0.219,0.191,0.174,
0.157,0.142,0.128,0.114,0.102,0.091,0.080,0.070,0.061,0.052
10 ,0.044,0.037,0.030,0.024,0.018,0.013,0.008,0.003,0.000,
num_metgrid_levels = 138,
p_top_requested     = 1000,
dx                  = 3335.324,
dy                  = 3335.324,
15 grid_id          = 1,
parent_id           = 1,
i_parent_start      = 1,
j_parent_start      = 1,
parent_grid_ratio    = 1,
20 parent_time_step_ratio = 1,
feedback            = 1,
smooth_option        = 0,
use_surface          = . false .,
sfcfcp_to_sfcfcp    = . false .
25 use_adaptive_time_step = . false .
step_to_output_time = . true .
target_cfl           = 1.3,
max_step_increase_pct = 50,
starting_time_step   = -1,
30 max_time_step     = 15,
min_time_step        = 1,
/

&physics
35 sst_update         = 1,

```

```

mp_physics           = 10
ra_lw_physics       = 4
ra_sw_physics       = 4
radt                = 3
5  sf_sfclay_physics = 1
   sf_surface_physics = 2,
   bl_pbl_physics     = 1,
   bldt               = 0,
   topo_wind          = 1
10  cu_physics        = 0,
   cudt               = 0,
   kfeta_trigger      = 2,
   isfflx              = 1,
   ifsnow              = 1,
15  icloud             = 1,
   surface_input_source = 1,
   num_soil_layers     = 4,
   mp_zero_out         = 0,
   sf_urban_physics    = 0,
20  maxiens            = 1,
   maxens              = 3,
   maxens2              = 3,
   maxens3              = 16,
   ensdim               = 144,
25  slope_rad          = 0,
   topo_shading         = 0,
   num_land_cat         = 21,
   iz0tln               = 1,
   shcu_physics         = 3
30  sf_ocean_physics    = 0
   usemonalb = . true .
   do_radar_ref = 1,
   hail_opt     = 1,
/

```

35

```

&afwa
afwa_diag_opt=1
afwa_severe_opt=1
afwa_ptype_opt=1
5 afwa_radar_opt=1
afwa_vis_opt=1
afwa_cloud_opt=1
/

10 &dynamics
w_damping = 1,
diff_opt = 1,
km_opt = 4,
gwd_opt = 0,
15 diff_6th_opt = 2,
diff_6th_factor = 0.12,
base_temp = 290.
damp_opt = 3,
zdamp = 5000.,
20 dampcoef = 0.2,
khdif = 0,
kvdif = 0,
non_hydrostatic = . true .,
moist_adv_opt = 1,
25 scalar_adv_opt = 1,
epssm = 0.5
/

30 &bdy_control
spec_bdy_width = 5,
spec_zone = 1,
relax_zone = 4,
specified = . true .,
35 nested = . false .,

```

```
periodic_x = .true .
```

```
/
```

```
&namelist_quilt
```

```
5 nio_tasks_per_group = 0,
```

```
nio_groups = 1,
```

```
/
```

*Author contributions.* The experiment including setting up the simulations and modifying the code was carried out by T. Schwitalla. The analysis of the simulations was performed in a collaborative effort and the manuscript was prepared by T. Schwitalla with contributions by  
10 all co-authors.

*Acknowledgements.* We are grateful to the High-Performance Computing Center Stuttgart (HLRS) and to Cray Inc. for providing the tremendous amount of computing time required for this simulation on XC40 system. Special thanks goes to U. Küster, T. Beisel, and T. Bönisch from HLRS and to S. Andersson and S. Dieterich from Cray Inc. . We are also grateful to the ECMWF for providing operational analysis data.

## References

- Acs, F., Horvath, A., Breuer, H., and Rubel, F.: Effect of soil hydraulic parameters on the local convective precipitation, *Meteorol. Z.*, 19, 143–153, 2010.
- Albergel, C., de Rosnay, P., Gruhier, C., Munoz Sabater, J., Hasenauer, S., Isaksen, L., Kerr, Y., and Wagner, W.: Evaluation of remotely  
5 sensed and modelled soil moisture products using global ground-based in situ observations, *Remote Sensing of Environment*, 118, 215–226, doi:10.1016/j.rse.2011.11.017, 2012.
- Balsamo, G., Viterbo, P., Beljaars, A., van den Hurk, B., Hirschi, M., Betts, A., and Scipal, K.: A Revised Hydrology for the ECMWF Model: Verification from Field Site to Terrestrial Water Storage and Impact in the Integrated Forecast System, *J. Hydrometeor.*, 10, 623–641, 2009.
- Bauer, H.-S., Weusthoff, T., Dorninger, M., Wulfmeyer, V., Schwitalla, T., Gorgas, T., Arpagaus, M., and Warrach-Sagi, K.: Predictive skill  
10 of a subset of models participating in D-PHASE in the COPS region, *Q. J. R. Meteorol. Soc.*, 137, 287–305, 2011.
- Bauer, H.-S., Schwitalla, T., Wulfmeyer, V., Bakhshaii, A., Ehret, U., Neuper, M., and Caumont, O.: Quantitative precipitation estimation based on high-resolution numerical weather prediction and data assimilation with WRF - a performance test, *Tellus A*, 67, 2015a.
- Bauer, P., Thorpe, A., and Brunet, G.: The quiet revolution of numerical weather prediction, *Nature*, 525, 47–55, doi:doi:10.1038/nature14956, 2015b.
- 15 Becker, N., Ulbrich, U., and Klein, R.: Systematic large-scale secondary circulations in a regional climate model, *Geophysical Research Letters*, 42, 4142–4149, doi:10.1002/2015GL063955, <http://dx.doi.org/10.1002/2015GL063955>, 2015GL063955, 2015.
- Branch, O., Warrach-Sagi, K., Wulfmeyer, V., and Cohen, S.: Simulation of semi-arid biomass plantations and irrigation using the WRF-NOAH model; a comparison with observations from Israel, *Hydrol. Earth Syst. Sc.*, 18, 1761–1783, doi:10.5194/hess-18-1761-2014, 2014.
- 20 Bromwich, D. H., Otieno, F. O., Hines, K. M., Manning, K. W., and Shilo, E.: Comprehensive evaluation of polar weather research and forecasting model performance in the Antarctic, *J. Geophys. Res.-Atmos.*, 118, 274–292, doi:10.1029/2012JD018139, 2013.
- Cassano, J. J., Higgins, M. E., and Seefeldt, M. W.: Performance of the Weather Research and Forecasting Model for Month-Long Pan-Arctic Simulations, *Mon. Wea. Rev.*, 139, 3649–3488, doi:10.1175/MWR-D-10-05065.1, 2011.
- Chen, F. and Dudhia, J.: Coupling an advanced land-surface/hydrology model with the Penn State NCAR MM5 modeling system. Part I:  
25 Model implementation and sensitivity, *Mon. Wea. Rev.*, 129, 569–585, 2001.
- Chen, F. and Zhang, Y.: On the coupling strength between the land surface and the atmosphere: From viewpoint of surface exchange coefficients, *Geophys. Res. Lett.*, 36, doi:10.1029/2009GL037980, 110404, 2009.
- Coppala, E., Giorgi, F., Mariotti, L., and Bi, X.: RegT-Band: a tropical band version of RegCM4, *Clim. Res.*, 52, 115–133, 2012.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P.,  
30 Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.
- Devis, A., van Lipzig, N. P. M., and Demuzere, M.: A new statistical approach to downscale wind speed distributions at a site in northern  
35 Europe, *Journal of Geophysical Research: Atmospheres*, 118, 2272–2283, doi:10.1002/jgrd.50245, 2013.
- Diaconescu, E. P. and Laprise, R.: Can added value be expected in RCM-simulated large scales?, *Clim. Dyn.*, 41, 1769–1800, doi:10.1007/s00382-012-1649-9, 2013.

- Dong, B., Sutton, R., and Shaffrey, L.: The 2013, hot, dry, summer in western Europe, *Bull. Amer. Meteor. Soc.*, 95, S62–S66, in "Explaining Extremes of 2013 from a Climate Perspective", 2014.
- Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system, *Remote Sens. Environ.*, 116, 140 – 158, doi:<http://dx.doi.org/10.1016/j.rse.2010.10.017>, Advanced Along Track Scanning Radiometer(AATSR) Special Issue, 2012.
- 5     Dudhia, J.: Reply to comment on "A nonhydrostatic version of the Penn State NCAR mesoscale model: Validation tests and simulations of an Atlantic cyclone and cold front" by J. Steppeler., *Mon. Wea. Rev.*, 123, 2573–2575, 1995.
- Efstathiou, G., Zoumakis, N., Melas, D., Lolis, C., and Kassomenos, P.: Sensitivity of {WRF} to boundary layer parameterizations in simulating a heavy rainfall event using different microphysical schemes. Effect on large-scale processes, *Atmos. Res.*, 132-133, 125–143, doi:<http://dx.doi.org/10.1016/j.atmosres.2013.05.004>, 2013.
- 10     Ek, M. B., Mitchell, K. E., Lin, Y., Rogers, E., Grummann., P., Koren, V., Gayno, G., and Tarpley, J. D.: Implementation of NOAA land surface model advances in the National Centers for Environmental Prediction operational Mesoscale Eta Model, *J. Geophys. Res.*, 108, doi:10.1029/2002JD003296, 2003.
- Evan, S., Rosenlof, K. H., Dudhia, J., Hassler, B., and Davis, S. M.: The representation of the TTL in a tropical channel version of the WRF model, *J. Geophys. Res.-Atmos.*, 118, 2835–2848, doi:10.1002/jgrd.50288, 2013.
- 15     Fonseca, R. M., Zhang, T., and Yong, K.-T.: Improved simulation of precipitation in the tropics using a modified BMJ scheme in the WRF model, *Geosci. Model Dev.*, 8, 2915–2928, doi:10.5194/gmd-8-2915-2015, 2015.
- Gebremichael, M., Bitew, M. M., Hirpa, F. A., and Tesfay, G. N.: Accuracy of satellite rainfall estimates in the Blue Nile Basin: Lowland plain versus highland mountain, *Water Resources Research*, 50, 8775–8790, doi:10.1002/2013WR014500, 2014.
- 20     Giorgi, F., Jones, C., and Asrar, G. R.: Addressing climate information needs at the regional level: the CORDEX framework, *WMO Bull.*, 58, 2009.
- Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., and Stoffel, M.: 21st century climate change in the European Alps- A review, *Science of the Total Environment*, 493, 1138–1151, doi:10.1016/j.scitotenv.2013.07.050, <http://dx.doi.org/10.1016/j.scitotenv.2013.07.050>, 2014.
- 25     Grell, G. A. and Dévényi, D.: A generalized approach to parameterizing convection combining ensemble and data assimilation techniques, *Geophys. Res. Lett.*, 29, 38–1–38–4, doi:10.1029/2002GL015311, 2002.
- Grell, G. A., Dudhia, J., and Stauffer, R. R.: A description of the fifth-generation Penn State/NCAR mesoscale model (MM5), NCAR technical Note TN-398+STR, NCAR, Boulder/CO, 122pp, 1995.
- Haylock, M. R., Hofstra, N., Klein Tank, A. M. G., Klok, E. J., Jones, P. D., and New, M.: A European daily high-resolution gridded data set of surface temperature and precipitation for 1950-2006, *J.G.R.: Atmospheres*, 113, doi:10.1029/2008JD010201, 2008.
- 30     Heikkilä, U., Sandvik, A., and Sorteberg, A.: Dynamical downscaling of ERA-40 in complex terrain using the WRF regional climate model, *Clim. Dyn.*, 37, 1551–1564, doi:10.1007/s00382-010-0928-6, 2011.
- Hong, S.-Y.: Stable Boundary Layer Mixing in a Vertical Diffusion Scheme, the Korea Meteor. Soc., Fall conference, Seoul, Korea, Oct. 25-26., 2007.
- 35     Hong, S. Y. and Lim, J.-O. J.: The WRF Single-Moment 6-Class Microphysics Scheme (WSM5), *J. Korean Meteor. Soc.*, 42, 129–151, 2006.
- Hong, S.-Y., Park, H., Cheong, H.-B., Kim, J.-E., Koo, M.-S., Jang, J., Ham, S., Hwang, S.-O., Park, B.-K., Chang, E.-C., and Li, H.: The Global/Regional Integrated Model system (GRIMs), *Asia-Pacific Journal of Atmospheric Sciences*, 49, 219–243, 2013.


- Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P., and Stocker, E. F.: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales, *Journal of Hydrometeorology*, 8, 38–55, doi:10.1175/JHM560.1, 2007.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models., *J. Geophys. Res.*, 113, 2008.
- Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A., Colette, A., D  qu   M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S., Kr  ner, N., Kotlarski, S., Kriegsmann, A., Martin, E., van Meijgaard, E., Moseley, C., Pfeifer, S., Preuschmann, S., Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsen, P., Somot, S., Soussana, J.-F., Teichmann, C., Valentini, R., Vautard, R., Weber, B., and Yiou, P.: EURO-CORDEX: New high-resolution climate change projections for European impact research, *Regional Environmental Change*, 14, 563–578, doi:10.1007/s10113-013-0499-2, 2014.
- Jimen  z, P., Dudhia, J., Fidel Gonz  lez-Rouc, F., Navarro, J., Mont  vez, J., and Garcia-Bustamante, E.: A Revised Scheme for the WRF Surface Layer Formulation, *Mon. Wea. Rev.*, 140, 898–918, 2012.
- Joyce, R. J., Janowiak, J. E., Arkin, P. A., and Xie, P.: CMORPH: A Method that Produces Global Precipitation Estimates from Passive Microwave and Infrared Data at High Spatial and Temporal Resolution, *J. Hydrometeorol.*, 5, 487–503, 2004.
- Kain, J. S.: The Kain-Fritsch Convective Parameterization: An update, *J. Appl. Meteorol.*, 43, 170–181, 2004.
- Katragkou, E., Garc  a-D  ez, M., Vautard, R., Sobolowski, S., Zanis, P., Alexandri, G., Cardoso, R. M., Colette, A., Fernandez, J., Gobiet, A., Goergen, K., Karacostas, T., Knist, S., Mayer, S., Soares, P. M. M., Pytharoulis, I., Tegoulis, I., Tsikerdekis, A., and Jacob, D.: Regional climate hindcast simulations within EURO-CORDEX: evaluation of a WRF multi-physics ensemble, *Geosci. Model Dev.*, 8, 603–618, 2015.
- Kotlarski, S., Keuler, K., Christensen, O. B., Colette, A., D  qu   M., Gobiet, A., Goergen, K., Jacob, D., L  thi, D., van Meijgaard, E., Nikulin, G., Sch  r, C., Teichmann, C., Vautard, R., Warrach-Sagi, K., and Wulfmeyer, V.: Regional climate modeling on European scales: a joint standard evaluation of the EURO-CORDEX RCM ensemble, *Geosci. Model Dev.*, 7, 1297–1333, 2014.
- Laprise, R., de Elia, R., Caya, D., Biner, S., Lucas-Picher, P., Diaconescu, E., Leduc, M., Alexandru, A., and Separovic, L.: Challenging some tenets of Regional Climate Modelling, *Meteorol. Atmos. Phys.*, 100, 3–22, doi:10.1007/s00703-008-0292-9, 2008.
- Liu, Z., Duan, Z., Jian, J., and Zhu, A.: Evaluation of Three Satellite Precipitation Products TRMM 3B42, CMORPH, and PERSIANN over a Subtropical Watershed in China., *Advances in Meteorology*, 2015, 13pp, 2015.
- Milovac, J., Ingwersen, J., and Warrach-Sagi, K.: Soil texture forcing data for the whole world for the Weather Research and Forecasting (WRF) Model of the University of Hohenheim (UHOH) based on the Harmonized World Soil Database (HWSD) at 30 arc-second horizontal resolution, doi:10.1594/WDCC/WRF\_NOAH\_HWSD\_world\_TOP\_SOILTYP, 2014.
- Milovac, J., Warrach-Sagi, K., Behrendt, A., Sp  th, F., Ingwersen, J., and Wulfmeyer, V.: Investigation of PBL schemes combining the WRF model simulations with scanning water vapor DIAL measurements, *J. Geophys. Res.-Atmos.*, 121, doi:10.1002/2015JD023927, 2016.
- Miyamoto, Y., Kajikawa, Y., Yoshida, R., Yamaura, T., Yashiro, H., and Tomita, H.: Deep moist atmospheric convection in a subkilometer global simulation, *Geophysical Research Letters*, 40, 4922–4926, doi:10.1002/grl.50944, 2013.
- Morrison, H., Thompson, G., and Tatarskii, V.: Impact of Cloud Microphysics on the Development of Trailing Stratiform Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes, *Mon. Wea. Rev.*, 137, 991–1007, 2009.




- Nolan, D. S., Stern, D. P., and Zhang, J. A.: Evaluation of Planetary Boundary Layer Parameterizations in Tropical Cyclones by Comparison of In Situ Observations and High-Resolution Simulations of Hurricane Isabel (2003). Part II: Inner-Core Boundary Layer and Eyewall Structure, *Mon. Wea. Rev.*, 137, 3675–3698, 2009.
- Pai Mazumder, D., Henderson, D., and Mölders, N.: Evaluation of WRF-forecasts over Siberia: Air mass formation, clouds and precipitation., *The Open Atmospheric Science Journal*, 6, 93–110, 2012.
- 5 Palmer, T. N.: Climate extremes and the role of dynamics, *Proceedings of the National Academy of Sciences*, 110, 5281–5282, doi:10.1073/pnas.1303295110, 2013.
- Perkins, S. E., Pitman, A. J., Holbrook, N. J., and McAneney, J.: Evaluation of the AR4 Climate Models Simulated Daily Maximum Temperature, Minimum Temperature, and Precipitation over Australia Using Probability Density Functions, *Journal of Climate*, 20, 4356–4376, doi:10.1175/JCLI4253.1, 2007.
- 10 Piere, A., von Hardenberg, J., Parodi, A., and Provenzale, A.: Sensitivity of precipitation statistics to resolution, microphysics, and convective parameterization: A case study with the high-resolution WRF climate model over Europe, *J. Hydrometeorol.*, 16, 1857–1872, doi:10.1175/JHM-D-14-0221.1, 2015.
- Prein, A. F., Gobiet, A., Suklitsch, M., Truhetz, H., Awan, N. K., Keuler, K., and Georgievski, G.: Added value of convection permitting seasonal simulations, *Climate Dynamics*, 41, 2655–2677, doi:10.1007/s00382-013-1744-6, 2013.
- 15 Prein, A. F., Gobiet, A., Truhetz, H., Keuler, K., Goergen, K., Teichmann, C., Maule, C. F., van Meijgaard, E., Džur, M., Nikulin, G., Vautard, R., Colett, A., Kjellström, E., and Jacob, D.: Precipitation in the EURO-CORDEX 0.11° and 0.44° simulations: High resolution, high benefits?, *Clim. Dyn.*, 46, 383–412, doi:10.1007/s00382-015-2589-y, 2015a.
- Prein, A. F., Langhans, W., Fosser, G., Ferrone, A., Ban, N., Goergen, K., Keller, M., Tölle, M., Gutjahr, O., Feser, F., Brisson, E., Kollet, S., Schmidli, J., van Lipzig, N. P. M., and Leung, R.: A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges, *Reviews of Geophysics*, 53, 323–361, doi:10.1002/2014RG000475, 2014RG000475, 2015b.
- 20 Rabier, F., Järvinen, H., Klinker, E., Mahfouf, J.-F., and Simmons, A.: The ECMWF operational implementation of four-dimensional variational assimilation. I: Experimental results with simplified physics, *Quarterly Journal of the Royal Meteorological Society*, 126, 1143–1170, doi:10.1002/qj.49712656415, <http://dx.doi.org/10.1002/qj.49712656415>, 2000.
- 25 Rasmussen, R. M., Geresdi, I., Thompson, G., Manning, K., and Karplus, E.: Freezing Drizzle Formation in Stably Stratified Layer Clouds: The Role of Radiative Cooling of Cloud Droplets, Cloud Condensation Nuclei, and Ice Initiation., *J. Atmos. Sci.*, 59, 837–860, 2002.
- Rogers, J. C.: North Atlantic Storm Track Variability and Its Association to the North Atlantic Oscillation and Climate Variability of Northern Europe, *Journal of Climate*, 10, 1635–1647, doi:10.1175/1520-0442(1997)010<1635:NASTVA>2.0.CO;2, 1997.
- Schwitalla, T. and Wulfmeyer, V.: Radar data assimilation experiments using the IPM WRF Rapid Update Cycle, *Meteorol. Z.*, 23, 79–102, 30 2014.
- Schwitalla, T., Bauer, H.-S., Wulfmeyer, V., and Zängl, G.: Systematic errors of QPF in low-mountain regions as revealed by MM5 simulations, *Meteorol. Z.*, 17, 903–919, 2008.
- Schwitalla, T., Bauer, H.-S., Wulfmeyer, V., and Aoshima, F.: High-resolution simulation over central Europe: assimilation experiments during COPS IOP 9c, *Q. J. R. Meteorol. Soc.*, 137, 156–175, 2011.
- 35 Shin, H. H. and Hong, S.-Y.: Intercomparison of Planetary Boundary-Layer Parametrizations in the WRF Model for a Single Day from CASES-99, *Boundary-Layer Meteorology*, 139, 261–281, doi:10.1007/s10546-010-9583-z, 2011.

- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D., Barker, D. O., Duda, M. G., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 3, NCAR Technical Note TN-475+STR, NCAR, Boulder/CO., [http://www.mmm.ucar.edu/wrf/users/docs/arw\\_v3.pdf](http://www.mmm.ucar.edu/wrf/users/docs/arw_v3.pdf), 2008.
- 5 Skok, G., Žagar, N., Honzak, L., Žabkar, R., Rakovec, J., and Ceglar, A.: Precipitation intercomparison of a set of satellite- and raingauge-derived datasets, ERA Interim reanalysis, and a single WRF regional climate simulation over Europe and the North Atlantic, *Theoretical and Applied Climatology*, 123, 217–232, doi:10.1007/s00704-014-1350-5, <http://dx.doi.org/10.1007/s00704-014-1350-5>, 2016.
- Stampoulis, D., Anagnostou, E., and Nikolopoulos, E.: Assessment of High-Resolution Satellite-Based Rainfall Estimates over the Mediterranean during Heavy Precipitation Events, *J. Hydrometeorol.*, 14, 1500–1514, 2013.
- Tao, W. K. and Simpson, J.: The Goddard cumulus ensemble model. Part I: Model description., *Terr. Atmos. Oceanic Sci.*, 4, 19–54, 1993.
- 10 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *Bulletin of the American Meteorological Society*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, 2012.
- Vautard, R., Gobiet, A., Jacob, D., Belda, M., Colette, A., Déqué, M., Fernández, J., García-Díez, M., Goergen, K., Güttler, I., Halenka, T., Karacostas, T., Katragkou, E., Keuler, K., Kotlarski, S., Mayer, S., Meijgaard, E., Nikulin, G., Patarčić, M., Scinocca, J., Sobolowski, S., Suklitsch, M., Teichmann, C., Warrach-Sagi, K., Wulfmeyer, V., and Yiou, P.: The simulation of European heat waves from an ensemble of regional climate models within the EURO-CORDEX project, *Climate Dynamics*, 41, 2555–2575, doi:10.1007/s00382-013-1714-z, 2013.
- 15 Viterbo, P., Beljaars, A., Mahfouf, J.-F., and Teixeira, J.: The representation of soil moisture freezing and its impact on the stable boundary layer, *Quarterly Journal of the Royal Meteorological Society*, 125, 2401–2426, doi:10.1002/qj.49712555904, 1999.
- Žagar, N., Honzak, L., Žabkar, R., Skok, G., Rakovec, J., and Ceglar, A.: Uncertainties in a regional climate model in the midlatitudes due to the nesting technique and the domain size, *J. Geophys. Res.*, 118, 6189–6199, 2013.
- 20 Warrach-Sagi, K., Wulfmeyer, V., Grasselt, R., Ament, F., and Simmer, C.: Streamflow simulations reveal the impact of the soil parameterization, *Meteorologische Zeitschrift*, 17, 751–762, 2008.
- Warrach-Sagi, K., Schwitalla, T., Wulfmeyer, V., and Bauer, H.-S.: Evaluation of a climate simulation in Europe based on the WRF-NOAH Model System: precipitation in Germany, *Clim. Dynam.*, 41, 755–774, doi:10.1007/S00382-013-1727-7, 2013.
- Wulfmeyer, V. and Coauthors: New Concepts for Studying Land-Surface-Atmosphere Feedback Based on a new Lidar Synergy and Grey Zone Simulations, *Geophysical Research Abstracts*, 17, <http://meetingorganizer.copernicus.org/EGU2015/EGU2015-5054.pdf>, 2015.
- 25 Wulfmeyer, V., Behrendt, A., Kottmeier, C., Corsmeier, U., Barthlott, C., Craig, G. C., Hagen, M., Althausen, D., Aoshima, F., Arpagaus, M., Bauer, H.-S., Bennett, L., Blyth, A., Brandau, C., Champollion, C., Crewell, S., Dick, G., DiGirolamo, P., Dorninger, M., Dufournet, Y., Eigenmann, R., Engelmann, R., Flamant, C., Foken, T., Gorgas, T., Grzeschik, M., Handwerker, J., Hauck, C., Höller, H., Junkermann, W., Kalthoff, N., Kiemle, C., Klink, S., König, M., Krauss, L., Long, C. N., Madonna, F., Mobbs, S., Neining, B., Pal, S., Peters, G., Pigeon, G., Richard, E., Rotach, M. W., Russchenberg, H., Schwitalla, T., Smith, V., Steinacker, R., Trentmann, J., Turner, D. D., van Baelen, J., Vogt, S., Volker, H., Weckwerth, T., Wernli, H., Wieser, A., and Wirth, M.: The Convective and Orographically Induced Precipitation Study (COPS): The Scientific Strategy, the Field Phase, and research Highlights, *Q. J. R. Meteorol. Soc.*, 137, 3–30, 2011.
- 30 Wulfmeyer, V., Muppa, S. K., Behrendt, A., Hammann, E., Späth, F., Sorbjan, Z., Turner, D. D., and Hardesty, R. M.: Determination of Convective Boundary Layer Entrainment Fluxes, Dissipation Rates, and the Molecular Destruction of Variances: Theoretical Description and a Strategy for Its Confirmation with a Novel Lidar System Synergy, *J. Atmos. Sci.*, 73, 667–692, doi:10.1175/JAS-D-14-0392.1, 2016.
- Xie, Z., Du, Y., and Yang, S.: Zonal Extension and Retraction of the Subtropical Westerly Jet Stream and Evolution of Precipitation over East Asia and the Western Pacific, *Journal of Climate*, 28, 6783–6798, doi:10.1175/JCLI-D-14-00649.1, 2015.



Zhang, G., Zhou, G., Chen, F., Barlage, M., and Xue, L.: A trial to improve surface heat exchange simulation through sensitivity experiments over a desert steppe site., *J. Hydrometeor.*, 15, 664–684, 2014.




**Figure 1.** Model domain of the latitude belt simulation. The white rectangles denote the domains used for verification of precipitation.




**Figure 2.** ERA-INTERIM 500 hPa geopotential height climatology for July and August 2013 (shaded black solid lines). The contour lines shaded areas show the anomaly during July and August 2013 compared to the period 1979–2012.



**Figure 3.** ERA-INTERIM sea surface temperature (SST) anomalies in July 2013 (upper panel) and August 2013 (lower panel) compared to the climatological period 1979–2012.




**Figure 4.** Mean sea level pressure climatology for July and August 2013 (black solid lines) from ERA-INTERIM. The shaded areas show the anomaly during July and August 2013 compared to the period 1979–2012.




**Figure 5.** Average mean sea level pressure at 12 UTC for July 2013 (left column) and August 2013 (right column). The first row shows the ECMWF analysis followed by the LOWRES simulation (c,d) and the Hires simulation (e,f). The two lowermost rows show the mean bias for the LOWRES (g,h) and the Hires simulations (i,j) compared to the ECMWF analysis for the two different months.







**Figure 6.** Model biases expressed in absolute terms 2-ECMWF standard deviations for the LOWRES simulation (upper row) and the HIRES simulation (bottom row) for the 12 UTC time steps. Left column: July 2013. Right column: August 2013. First two rows: Mean sea level pressure. Bottom two rows: 500 hPa geopotential height.




**Figure 7.** Mean sea level pressure standard deviation at 12 UTC for July 2013 (left column) and August 2013 (right column). The top row shows the ECMWF analysis, the middle row shows the LOWRES simulation and the bottom row displays the HIRES simulation.




**Figure 8.** Mean 300 hpa wind velocities for July 2013 (left column) and August 2013 (right column). The top row shows the ECMWF analysis, the middle row shows the LOWRES simulation and the bottom row displays the HIRES simulation.




**Figure 9.** Mean 500 hPa geopotential height and mean differences between the WRF simulations and ECMWF analysis for July 2013 (left column) and August 2013 (right column). The top row show the ECMWF analysis, the middle row show the LOWRES simulation and the bottom row display the HIRES simulations.




**Figure 10.** Time series of the MSLP averaged between 40° N and 65° N, and 60° W and 10° E.



**Figure 11.** Average 925 hPa water vapor mixing ratio for 12 UTC of the ECMWF analysis (a) and the LOWRES simulation (b). The bottom panel shows the HIRES simulation.




**Figure 12.** Mean 925 hPa temperature for 12 UTC of the ECMWF analysis (a). (b) and (c) show the deviation of the LOWRES and HIRES simulation from the ECMWF analysis, respectively.




**Figure 13.** Mean 2-m temperature at 12 UTC in July (left column) and August (right column). The top row shows the ECMWF analysis, the middle row displays the LOWRES simulation and the bottom row represents the HIRES simulation. Reddish colors indicate a warm bias of the WRF simulations.







**Figure 14.** 10-m wind speed bias in July (left column) and August 2013 (right column) for the LOWRES simulation (upper row) and the HIRES simulation (lower row).




**Figure 15.** 2 month accumulated precipitation. (a) shows the CMORPH analysis, (b) shows the LOWRES simulation and (c) displays the HIRES simulation. (d) displays the accumulated precipitation from the operational ECMWF 12 h forecast started at 00 UTC and 12 UTC each day.



**Figure 16.** Histogram of the 2 month accumulated precipitation by using the CMORPH data over land points only. The filled blue bars denote the CMORPH data set and the cross-hatched bars denote the LOWRES (left) and HIRES simulation data (right).




(a)




(b)


**Figure 17.** Scatter plot of the accumulated precipitation over the two month period including regression lines. The left panel shows the LOWRES simulation vs. the CMORPH data. The red line would be the perfect result. The right panel shows the HIRES simulation vs. the CMORPH data.



**Figure 18.** 2 month accumulated precipitation over Europe. (a) shows the E-OBS data set, (b) shows the LOWRES simulation, and (c) displays the HIRES simulation, and (d) shows the ECMWF forecast.



**Figure 19.** Histogram of the 2 month accumulated precipitation over Europe using the E-OBS data set. The filled blue bars denote the observation data set and the cross-hatched bars denote the HIRES (left column) and LOWRES simulation data (right column).



**Figure 20.** Taylor diagram of the accumulated precipitation over land points. The simulations were plotted against CMORPH observations, except for Europe where the E-OBS data set is the reference.

**Table 1.** Physics parameterizations used in the WRF simulations.

Parameterization	Scheme	Reference
Cloud microphysics	Morrison 2-moment	Morrison et al. (2009)
Radiation	RRTMG	Iacono et al. (2008)
PBL	YSU	Hong (2007)
Shallow convection	GRIMS	Hong et al. (2013)
Cumulus parameterization	KF-ETA	Kain (2004)
Surface layer	MM5 scheme	Jiménez et al. (2012)
Land Surface	NOAH LSM	Ek et al. (2003)