

The TOMCAT global chemical transport model v1.6: Description of chemical mechanism and model evaluation

Sarah A. Monks^{1,2,3}, Stephen R. Arnold¹, Michael J. Hollaway¹, Richard J. Pope^{1,4}, Chris Wilson^{1,4}, Wuhu Feng^{1,5}, Kathryn M. Emmerson⁶, Brian J. Kerridge⁷, Barry L. Latter⁷, Georgina M. Miles⁷, Richard Siddans⁷, and Martyn P. Chipperfield¹

¹Institute for Climate and Atmospheric Science, University of Leeds, UK.

²Chemical Sciences Division, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA.

³Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA..

⁴National Centre for Earth Observation, University of Leeds, UK.

⁵National Centre for Atmospheric Science, University of Leeds, UK.

⁶CSIRO Oceans and Atmosphere Flagship, Aspendale, Australia.

⁷Remote Sensing Group, STFC Rutherford Appleton Laboratory, Harwell Oxford, UK.

Correspondence to: Sarah Monks (sarah.monks@noaa.gov)

Abstract. ~~The~~ This paper documents the tropospheric chemical mechanism scheme used in the TOMCAT 3-D chemical transport model ~~has been updated with the emissions and chemical degradation~~. The current scheme includes a more detailed representation of hydrocarbon chemistry than previously included in the model, with the inclusion of the emission and oxidation of ethene, propene, ~~toluene, butane~~ ~~butane, toluene~~ and monoterpenes. The full tropospheric chemical mechanism is described

5 and the model is evaluated against a range of surface, balloon, aircraft and satellite measurements. The model is generally able to capture the main spatial and seasonal features of high and low concentrations of carbon monoxide (CO), ozone (O₃), volatile organic compounds (VOCs) and reactive nitrogen. However, model biases are found ~~in some species~~, some of which are common to chemistry models and some that are specific to TOMCAT and warrant further investigation.

Simulated O₃ is found to generally lie within the range of ozonesonde observations and shows good agreement with surface sites. The most notable ~~exceptions to this are during winter at high latitudes, when O₃ is underestimated, and during summer over North America, when O₃ is overestimated. Global Ozone Monitoring Experiment-2~~ of these biases are 1) a negative bias in Northern Hemispheric (~~GOME-2~~) comparisons suggest that TOMCAT sub-column tropospheric O₃ in DJF may also be underestimated outside of the Arctic, particularly near tropical regions.

TOMCAT CO is negatively biased during winter and spring in the Northern Hemisphere (NH) when compared to ground-based observations and MOPITT (Measurements Of Pollution In The Troposphere) satellite data. In contrast, CO is positively biased winter and spring CO, and a positive bias in Southern Hemispheric (SH) CO throughout the year ~~in the Southern Hemisphere (SH)~~. The negative bias in the NH is a common feature in chemistry models and TOMCAT lies well within the range of biases found in other models, while the TOMCAT SH positive bias is at the upper range of positive biases reported in other models.

Using two simulations with different boundary conditions highlighted the sensitivity of model performance to the chosen 20 emission dataset when simulating VOCs, nitrogen oxides (NO_x) and peroxyacetyl nitrate (PAN). VOC measurements show

winter/spring negative biases in C2-C3, 2) a positive bias in NH O_3 in summer and a negative bias at high latitudes during SH winter and 3) a negative bias in NH winter C2 and C3 alkanes and alkenes, which is likely driven by underestimated anthropogenic emissions. TOMCAT is able to capture the seasonal minima and maxima of PAN and HNO_3 . However, comparisons to an aircraft climatology show that PAN may be overestimated in winter and HNO_3 may be overestimated in winter and 5 spring in regions over North America. The model showed different biases in NO_x , depending on location, with evidence of underestimated Asian emissions contributing to negative model biases over China and underestimated fire emissions contributing to negative biases in the SH.

10 TOMCAT global mean tropospheric hydroxyl radical (OH) concentrations are higher than estimates inferred from observations of methyl chloroform, but similar to, or lower than, multi-model mean concentrations reported in recent model intercomparison studies. TOMCAT shows peak OH concentrations in the tropical lower troposphere, unlike other models, which show peak concentrations in the tropical upper troposphere. This is likely to affect the lifetime and transport of important trace gases and warrants further investigation.

1 Introduction

15 Atmospheric chemistry plays a central role in air quality and climate change, which can have a negative effect on humans on a global-scale. Air pollution has been estimated to have caused over 3 million deaths worldwide in 2010 and this rate is estimated to double by 2050 due to projected increases in emissions (Lelieveld et al., 2015). Increases in anthropogenic emissions have led to higher atmospheric concentrations of greenhouse gases, such as methane (CH_4) and ozone (O_3), contributing significantly to the observed rise in global mean surface temperature (Stocker et al., 2013). Chemical processing, emissions, and transport determine the concentrations and distribution of pollutants within the atmosphere and the impact that they have on society. 20 Reactive gases, such as volatile organic compounds (VOCs) and nitrogen oxides (NO_x), influence air quality and climate as they result in the formation of O_3 and aerosols. Other gases such as carbon monoxide (CO), which may not directly affect the climate, can have secondary impacts by influencing the lifetime of gases such as CH_4 (Berntsen et al., 2005).

25 Atmospheric chemistry models help to inform our understanding of how atmospheric chemistry affects climate and air quality on a global- or regional-scale. These models can be used to simulate the temporal and spatial evolution of important short-lived pollutants, taking into account the main physical and chemical processes that act on trace constituents in the troposphere (emissions, chemistry, transport and deposition). The chemical and dynamical complexity and the spatial resolution of such models is a compromise between model accuracy and computational efficiency. Atmospheric chemistry models are often run as chemical transport models (CTMs), where transport is constrained by reanalysis products that assimilate meteorological observations. This allows the simulated chemical fields to provide context for measurements, which are often 30 limited spatially and temporally. They can also be used to further understand the impacts of new atmospheric processes that have been identified by measurements (e.g., Lelieveld et al., 2008). CTMs are of particular use in investigating the impacts of natural and anthropogenic emissions on atmospheric burdens of pollutants that are important for air quality and climate reasons and for source-receptor studies for policy-making purposes (e.g., Sanderson et al., 2008; Fiore et al., 2009).

The TOMCAT CTM is a three-dimensional (3-D) global Eulerian model that has been used for a wide range of tropospheric and stratospheric chemistry studies. For example, it has been used to investigate the impacts of O₃ on crop yields (Holloway et al., 2012), fire emissions on Arctic interannual variability (Monks et al., 2012) and to identify the main sources of peak summertime O₃ in the Mediterranean (Richards et al., 2013). In the stratosphere the model has been used to study issues such as ozone depletion (e.g., Chipperfield et al., 2015) and the impact of solar variability (e.g., Dhomse et al., 2013). TOMCAT is also the host model for the GLOMAP aerosol module (Mann et al., 2010).

This paper summarises ~~recent updates to the~~ ~~the current~~ tropospheric chemical mechanism, ~~documenting the current full chemical scheme~~ ~~scheme used in TOMCAT~~ (Section 2). ~~The scheme gives a more detailed representation of hydrocarbon chemistry than previously included in the model, with the inclusion of the emission and oxidation of ethene, propene, butane, toluene and monoterpenes. Alkenes have the greatest potential for forming O₃ (Saunders et al., 2003), and previously, isoprene was the only alkene treated in the TOMCAT model. In addition, a more extensive VOC scheme makes it possible to couple the TOMCAT tropospheric chemistry to the formation of secondary organic aerosol in future versions of the GLOMAP aerosol model (Mann et al., 2010).~~ Key gas-phase species simulated by the latest version of the model are shown and evaluated using a range of observations. The model simulations that are evaluated are described in Section 2.2 and the observations that are used are described in Section 3. The observational platforms that are used include surface, satellite, aircraft and balloon sounding measurements. The model results and comparisons with observations are shown in Section 4 and focus on annual, seasonal and monthly mean simulated concentrations. The chemical species that are discussed include CO, O₃, VOCs, reactive nitrogen (NO_y) and the hydroxyl radical (OH).

2 The TOMCAT model

The TOMCAT model is an Eulerian offline 3-D global CTM and is described by Chipperfield (2006). The model has a flexible horizontal and vertical resolution and the vertical domain can be varied depending on the problem being studied. Typical horizontal resolutions range from 5.6° x 5.6° for multidecadal stratospheric studies to 1.2° x 1.2° for short case studies. The model uses a σ - p coordinate system, with near-surface levels following the terrain (σ) and higher levels ($\sim >100$ hPa) using pressure levels (p). The model extends from the surface to ~ 10 hPa for tropospheric simulations, as used in this study. ~~The global mean pressure levels are shown in Figure 1. Model meteorology is forced by winds, temperature and humidity fields from the European Centre for Medium-Range Weather Forecasts (ECMWF) reanalyses (Dee et al., 2011). These data are read in every 6 hours and~~ ~~interpolated to the linearly interpolated in time to the model time-step and to the~~ TOMCAT grid. To avoid inconsistencies between horizontal and vertical winds after this interpolation, the vertical motion is diagnosed from horizontal divergence instead of using analysed vertical velocities. Large-scale tracer advection in the meridional, zonal and vertical direction is based on the Prather (1986) scheme, which conserves mass and maintains tracer gradients (Chipperfield, 2006). Sub-grid scale transport (boundary layer mixing and convective transport) is treated in the model using the Holtslag and Bolville (1993) and Tiedtke (1989) schemes. There is also an option to run the model using archived convective mass fluxes (Feng et al., 2011). Wilson et al. (2014) used sulphur hexafluoride (SF₆) to evaluate model tracer transport and showed that the

model is able to reproduce seasonal transport timescales and patterns along with the location of the intertropical convergence zone. However, they also noted that the model inter-hemispheric transport is somewhat slow, resulting in an interhemispheric gradient in SF₆ that was 18% too large.

5 Natural and anthropogenic surface emissions are read into the model on a 1°x1° resolution and regredded online to the model grid. The model is usually provided with monthly mean emissions and a temporal interpolation is performed online to the model time step. Isoprene emissions are emitted with and then have a diurnal cycle imposed online to account for the dependence of emissions on daylight. Lightning emissions of NO_x are coupled to convection in the model and therefore vary in space and time according to the seasonality and spatial pattern of convective activity (Stockwell et al., 1999). Aircraft emissions of NO_x
10 ~~are based on estimated aircraft movements for the year 2002 (Lamarque et al., 2010) and were calculated for the European QUANTIFY project (<http://www.pa.op.dlr.de/quantify/>). They are available on 25 vertical levels from the surface to 14.5 km and are regredded to the TOMCAT vertical levels online.~~

15 Dry deposition velocities are weighted by prescribed fixed land cover fields and seasonally varying sea-ice fields from the NCAR community land model (CLM) (Oleson et al., 2010). The 16 CLM land types were regredded onto the model resolution and reclassified into the TOMCAT's five land types (Forest~~forest~~, grass/shrub/crop, bare ground, sea-ice and water). Chemical species' deposition velocities were then determined based upon time of day, season and were weighted by the proportion of the grid box covered by each land type. Wet deposition is parameterised according to the proportionality of the removal rate to the concentration of the species and is dependent on convection rates, precipitation and the solubility of gases. The scheme has been shown to perform well within the TOMCAT model with a 4% bias compared to Radon observations (Giannakopoulos
20 et al., 1999).

2.1 Tropospheric chemistry scheme

~~TOMCAT O_x–HO_x–NO_x–CO–CH₄ and C1–C3 alkane hydrocarbon chemistry was previously described by Arnold et al. (2005). TOMCAT has since been extended to also include the The previously documented TOMCAT tropospheric chemical mechanism included odd oxygen (O_x), reactive nitrogen (NO_y), carbon monoxide, methane, ethane, propane, acetaldehyde, acetone
25 and formaldehyde chemistry (Arnold et al., 2005). TOMCAT also includes oxidation of isoprene based on the Mainz Isoprene Mechanism (MIM) scheme (Pöschl et al., 2000). The implementation of this scheme into TOMCAT is described by Young (2007). Isoprene is one of the largest single sources of VOCs to the atmosphere, accounting for around a third of total natural and anthropogenic VOC emissions (Guenther et al., 2006). After emission, isoprene is highly reactive and can influence O₃ concentrations both regionally (Chameides et al., 1988) and globally (Wang and Shallcross, 2000). O₃, OH,
30 CO and PAN from the condensed MIM scheme were found to agree within 10% of the concentrations calculated from a more explicit representation of isoprene chemistry in the Master Chemical Mechanism (Pöschl et al., 2000). However, significant uncertainties still exist in the representation of isoprene chemistry in models and chemical mechanisms will likely evolve in the future (e.g., Archibald et al., 2010, 2011; Squire et al., 2015). Most recently, the TOMCAT model has been updated chemistry has been expanded to include the emission and destruction of some C2–C7 unsaturated and aromatic hydrocarbons (ethene, propene, toluene and butane) based on the Extended Tropospheric Chemistry scheme (ExtTC) (Folberth et al., 2006). Biogenic~~

emission and chemistry of, and monoterpenes based on the MOZART-3 chemical mechanism (Kinnison et al., 2007) has also been added. The model now includes [The extended chemistry scheme results in an increase in the global burden of CO₂](#).
5 O₃, PAN, HO₂ and a small decrease in OH in summer (see supplementary material). Whilst this scheme has been used in the TOMCAT model for some scientific studies (e.g., Richards et al., 2013; Emmons et al., 2015), the expanded scheme was not fully documented. This is the purpose of this study.

The current model chemistry scheme has a total of 79 species and they are listed in Table 2, identifying whether they are emitted or undergo dry or wet deposition. A few shorter-lived species are grouped into families for transport processes, which 10 are also identified in Table 2 [16 of which are emitted \(see Table 2\)](#), and approximately 200 chemical reactions. The bimolecular, termolecular and photolysis reactions are shown in Tables 3, 4 and 5, respectively. The chemical reactions are implemented via a [integrated in the model with a 15-minute chemical timestep using a](#) software package, ASAD (Carver et al., 1997), which allows the use of input files that contain the information listed in Tables 2. The package allows fractional products and the use of families, which can be used for grouping very short-lived species together for transportation (see Table 2 for 15 species treated as families). The bimolecular and termolecular kinetic rates are mostly taken from the 2005 International Union of Pure and Applied Chemistry recommendations (<http://iupac.pole-ether.fr/>) and the Leeds Master Chemical Mechanism (MCM, 2004). Simplified ethene, propene and butane chemistry is based on von Kuhlmann (2001), with reaction rates taken from IUPAC (Atkinson et al., 2006). Ethane chemistry adds 1 bimolecular reaction (R135–R136, Table 3) and 1 termolecular reaction (R17–19, Table 5) to TOMCAT. Oxidation of ethane by OH forms PrpeOO, a peroxy radical, which continues in the 20 propene oxidation chain (von Kuhlmann, 2001). Propane oxidation adds 6 bimolecular reactions (R137–145, Table 3) and 1 termolecular reaction to TOMCAT (R17–19, Table 4) and butane adds 10 bimolecular reactions (R123–134, Table 3) and 5 photolysis reactions (R40a–46b, Table 5). Ethene, propene and butane emissions are emitted into the respective compounds, with no lumping of higher alkenes/alkanes. Toluene is emitted into a generic aromatic compound, AROM, which produces AROMO₂ (peroxy radicals) and AROMOOH (hydroperoxides). Including AROM adds 11 bimolecular reactions to the model 25 (R146–R156, Table 3) and 2 photolysis reactions (R48a–R48b, Table 5), with reaction rates taken from Folberth et al. (2006). Including ethene, propene, butane and toluene will account for missing sources of carbon in the model and produces peroxy radicals that are important for O₃ production von Kuhlmann (2001). Alkenes are particularly efficient at producing O₃, with photochemical ozone creation potentials of 100 and 105 for ethene and propene, respectively, compared to 8.8 and 18.3 for ethane and propane, respectively Saunders et al. (2003). The lumped monoterpene compound (C₁₀H₁₆) is treated 30 as alpha-pinene, with emissions made up of the sum of alpha-pinene, beta-pinene, limonene, myrcene, ocimene, sabinene, and delta-3-carene. Including monoterpene chemistry adds 6 bimolecular reactions (R116–R122, Table 3) and one photolysis reaction (R26–27, Table 5) to the TOMCAT extended scheme. Monoterpenes are the second largest biogenic source of VOCs, after isoprene, and play an important role in OH, NO₃, O₃ and aerosol chemistry (e.g., Atkinson and Arey, 2003; Fuentes et al., 2000). Accounting for monoterpenes also allows the coupling of the atmospheric chemistry model, TOMCAT, to the aerosol model, 35 GLOMAP.

Photolysis rates are calculated online [based on Hough \(1988\)](#) at each chemical timestep based on the two-stream method of Hough (1988), which considers [both](#) direct and scattered radiation. [Within TOMCAT, this scheme is](#) The scheme has total of

203 wavelength intervals from 120 nm to 850 nm, though only wavelengths above 175 nm are used for stratosphere-troposphere studies. These wavelength intervals are the same as those employed in the TOMCAT stratospheric chemistry scheme (Chipperfield et al., 2005). The top of the atmosphere solar flux spectrum is fixed in time and there is no account of, for example, the 11-yr solar cycle in the standard model. This photolysis scheme is coupled with the TOMCAT model by using the model temperature and ozone concentration profiles. The scheme is also supplied with surface albedo, aerosol concentrations and monthly mean climatological cloud fields and total column ozone and temperature profiles. The bimolecular and termolecular kinetic rates are mostly taken from the International Union of Pure and Applied Chemistry (Atkinson et al., b) and the Leeds Master Chemical Mechanism (MCM, 2004). The bimolecular, termolecular and photolysis reactions are listed in Tables 3-5. This scheme was first used in this manner by Arnold et al. (2005). Previously, an offline approach was used where photolysis rates were calculated offline and then read in to the model (e.g. Law et al. (1998)). Where possible, photochemical data is taken from Sander (2011) for species which are also relevant for the stratosphere. Otherwise photochemical data is generally taken from IUPAC (Atkinson et al., 2004, 2006). The UV absorption cross-sections for methyl hydroperoxide (MeOOH), which are from JPL (Sander, 2006), are used for the hydroperoxides produced from the oxidation of butane (BtOOH), toluene (AROMOOH) and monoterpene (TERPOOH). For the photolysis of ONIT, which represents organic nitrates produced from higher alkanes (currently only butane), cross-sections for methyl nitrate are used based on IUPAC recommendations (Atkinson et al., 2006). ONIT can be an important reservoir of reactive nitrogen (von Kuhlmann, 2001). Stratospheric concentrations of O₃ and NO_y calculated offline by the 2-D Cambridge model (Law and Pyle, 1993) are read in by TOMCAT in the absence of stratospheric chemistry. Hydrogen, carbon dioxide, oxygen and nitrogen are kept at fixed global mean volume mixing ratios in the model. Water vapour is calculated from the specific humidity field read in from the meteorological input data.

Heterogeneous chemistry is known to affect the global concentrations of O₃, OH and NO_x in the troposphere (Jacob, 2000). One important reaction is that of dinitrogen pentoxide (N₂O₅) with water (H₂O) on the surface of aerosols to form nitric acid (HNO₃). HNO₃ is highly soluble and is therefore efficiently lost through wet deposition, making this an important loss channel for NO_x from the atmosphere. This is important in the troposphere when there is no sunlight, allowing time for the formation of N₂O₅. TOMCAT can be run coupled to the GLOMAP aerosol module (Mann et al., 2010), which can then calculate the available aerosol surface area for use in the heterogeneous chemistry calculation (e.g., Breider et al., 2010). When TOMCAT is run without the coupled GLOMAP scheme uncoupled to GLOMAP there is an option to account for heterogeneous uptake of N₂O₅ using prescribed monthly mean aerosol concentrations that have been calculated for number density and radius for 5 different aerosol types (sulphate, black carbon, organic carbon, sea salt and dust). Currently this data is taken from a previous GLOMAP run for the year 2000 by the GLOMAP model (Mann et al., 2010). In this simplified scheme, uptake coefficients for sulphate, black carbon, organic carbon and sea salt the uptake coefficients are based on Evans and Jacob (2005) and the uptake coefficient for dust, with the exception of dust, which is based on Mogil et al. (2006), with the (See Table 6). The overall uptake coefficient varying varies as a function of temperature, humidity and aerosol composition. Similarly, computationally cheap TOMCAT-GLOMAP 'aerosol-only' experiments can be run using specified fields of oxidants. Uptake of N₂O₅ on cloud surfaces is currently not included due to the use of climatological clouds in the model. Code exists to take account of HO₂ uptake but is currently not used in the model as it requires evaluation and testing within the model.

2.2 Model simulations and set-up and emissions

5 ~~Two simulations have~~ A simulation has been performed using the ~~new~~ current chemical mechanism scheme for the year 2000 (RUN) ~~or year~~ 2008 (RUN) ~~(both~~ with a 1-year spin-up), ~~which differ slightly in their inputs. Using two simulations for~~ different years and with different set-ups allows insight into whether a model bias is systematic or possibly due to the model boundary conditions, such as emissions. Both simulations use ~~The model uses~~ 31 vertical levels (surface to 10 hPa) and a horizontal resolution of $2.8^\circ \times 2.8^\circ$. ~~Each model run uses~~ ERA-Interim meteorology ~~and emissions for the specific year of the~~ 10 run and uses offline aerosols is used to drive the model. Offline aerosol concentrations are used for N_2O_5 uptake.

15 ~~RUN~~ uses anthropogenic and biomass burning emission estimates for the year 2000. The anthropogenic and ship emissions are from a dataset that was created for the IPCC Fifth Assessment Report (AR5) (Lamarque et al., 2010). The biomass burning emissions are taken from the Global Fire Emissions Database (GFED) version 3.1 (van der Werf et al., 2010). Oceanic CO and soil NO_x emissions are from the POET emission inventory (Granier et al., 2005) and biogenic emissions of volatile organic compounds (VOCs) were calculated offline by the Model of Emissions of Gases and Aerosol from Nature version 2.1 (Guenther et al., 2012) within the NCAR Community Climate Model (Lawrence et al., 2011). The MEGAN emissions are described and used in Scott et al. (2016). Due to the long lifetime of methane (~ 10 years), a very long spin-up time would be required in order to simulate a realistic atmospheric methane distribution using online emissions and loss. To avoid this long spin-up, a common method in CTMs is to use a fixed methane field from offline sources. In this TOMCAT simulation, CH_4 20 emissions are from Wilson et al. (2016), with tropospheric surface concentrations being scaled to match a realistic global mean concentration of 1800 ppbv. This results in realistic model concentrations of CH_4 , whilst the spatial distribution of high/low emission regions is maintained.

25 ~~RUN~~ This run uses emissions that were chosen for the POLARCAT (POLar study using Aircraft, Remote Sensing, surface measurements and models of Climate, chemistry, Aerosols, and Transport) ~~Model Intercomparison Project (POLMIP)~~ (Emons et al., 2015). Monthly mean anthropogenic and ship emissions are based on the Streets v1.2 inventory (Zhang et al., 2009), which was updated with ~~the latest~~ several recent regional inventories in 2008 for the POLARCAT campaign. ~~Monthly mean~~ This included Asian emissions from Zhang et al. (2009), North America emissions from USNEI 2002 and CAC 2005 and European emissions from EMEP 2006 database (<http://www.ceip.at>). Where regional inventories were unavailable emissions were taken from the EDGAR 3.2FT2000 database. These emissions have no seasonal cycle and are therefore the same each 30 month. Monthly-varying biogenic emissions are from the MACC (Monitoring Atmospheric Composition and Climate) project (MACCcity), which provides simulated VOCs calculated offline by the Model of Emissions of Gases and Aerosols from Nature (MEGAN) v2.1 (Guenther et al., 2012). ~~0 for a reference year 2000 (Guenther et al., 2006)~~, Oceanic CO and VOC emissions, and soil NO_x are from the POET inventory. For 2008, daily biomass burning emissions are taken from the Fire INventory from NCAR (FINN) (Wiedinmyer et al., 2011). ~~Aircraft emissions of NO_x are based on estimated aircraft movements for the year 2002 (Lamarque et al., 2010) and were calculated for the European QUANTIFY project (<http://www.pa.op.dlr.de/quantify/>).~~ They are provided on 25 vertical levels from the surface to 14.5 km and are regressed to the TOMCAT vertical levels online.

Surface CH₄ is set to equal zonal mean concentrations calculated from NOAA/ESRL/GMD surface observations observations 5 for the year 2000 (Meinshausen et al., 2011).

The emissions for both runs are shown in Table 1. Some differences exist between the two sets of emissions, with RUN having higher total biogenic emissions and RUN having higher total anthropogenic and fire emissions. There is also the option to use an emission file for CH₄ and then scale the surface global mean concentration to a suitable value for the year in question. As already mentioned, lightning NO_x emissions are also included but are dealt with online depending on the convection in the 10 model.

3 Observations

3.1 Satellite data

Simulated CO is compared on a global scale to CO distributions retrieved from the satellite instrument, MOPITT (Measurements Of Pollution In The Troposphere) version 6. MOPITT is a nadir-viewing instrument on-board the NASA Terra satellite 15 and retrieves CO concentrations globally at a horizontal resolution of \sim 22 km by measuring infrared radiances in the CO absorption band (Deeter et al., 2010). The Terra satellite has an overpass time at the equator of 10:30 local time (LT). Version 6 uses an *a priori* based on climatological output from the CAM-Chem model for 2000 to 2009 (Deeter, 2013). It has increased sensitivity to lower tropospheric CO by using both near-infrared and thermal infrared wavelengths (Deeter et al., 2011). As MOPITT is a nadir-viewing instrument, it is more sensitive to certain altitudes, therefore averaging kernels (AKs) that contain 20 information about the instrument's varying sensitivities at different altitudes are used, along with the *a priori*, to apply the same vertical sensitivity to the TOMCAT CO profiles. This allows a more accurate comparison between the observed and simulated CO. Data where the degrees of freedom signal (DOFS) are less than 1 have been removed from the model and satellite columns to identify retrievals where the satellite sensitivity is low.

Satellite O₃ is taken from Global Ozone Monitoring Experiment-2 (GOME-2) aboard EUMETSAT's Metop-A polar orbiting 25 satellite. GOME-2 is a nadir-viewing instrument with an approximate local equator crossing time of 09:30 LT. It has a spectral range of 240–790 nm and the pixel sizes are between 40 km and 80 km along and across track, respectively (Miles et al., 2015b). The data comes from the Rutherford Appleton Laboratory and is based on an optimal estimation algorithm (Rodgers, 1976). Miles et al. (2015b) describes how the GOME-2 retrievals are quality controlled prior to use, with data being removed where geometric cloud fraction is greater than 0.2 and the solar zenith angle is less than 80°. For optimal comparisons, the 30 GOME-2 AKs are applied to the TOMCAT data, as described in Miles et al. (2015a). The model and satellite data is matched in space and time by choosing the closest model grid box to the satellite pixel, to within 3 hours of the satellite daylight overpass time (6-hourly model output is being used).

For nitrogen dioxide (NO₂), we use data from the Ozone Monitoring Instrument (OMI) aboard NASA's EOS-Aura polar orbiting satellite. It has an approximate equator crossing of 13:30 LT (Boersma et al., 2007) and is a nadir-viewing instrument with a spectral range of 270–500 nm. The pixel sizes are between 16–23 km and 24–135 km along and across track, respectively, depending on the viewing zenith angle (Boersma et al., 2007). The tropospheric column NO₂ data, known as the DOMINO

5 product (v2.0) (Boersma et al., 2011), was downloaded from the Tropospheric Emissions Monitoring Internet Service (TEMIS; <http://www.temis.nl/airpollution/no2.html>). The retrieval of OMI tropospheric column NO₂ is based on Differential Optical Absorption Spectroscopy (DOAS), as discussed by Eskes and Boersma (2003). OMI retrievals have been quality controlled and data is only used where they have geometric cloud cover less than 20 % and good quality data flags. The product also uses the algorithm of Braak (2010) to remove OMI pixels affected by row anomalies. Studies have shown the DOMINO product to
10 have small biases against other independent observational data with some evidence of a small low bias over oceans (Irie et al., 2012; Boersma et al., 2008). The product has also been used in model evaluation studies previously (e.g., Huijnen et al., 2010; Pope et al., 2015). For the TOMCAT comparisons, AKs are applied following Boersma et al. (2011). The model and satellite
data is matched in space and time by choosing the closest model grid box to the satellite pixel, to within 3 hours of the satellite
daylight overpass time (6-hourly model output is being used).

15 **3.2 Surface data**

We take O₃ measurements at the surface over the U.S. from the United States Environmental Protection Agency (EPA) Clean Air Status and Trends Network (CASTNET) database. They provide hourly mean concentrations from continuous ozone monitoring instruments that have undergone a large amount of quality assurance. Here we use ~~an average of data available for the years 2000 to 2008 at~~ data from 44 sitescovering the whole which cover large parts of the U.S., excluding highly urbanised
20 sites as identified by Sofen et al. (2016). The model output is interpolated to the location of each station both horizontally and vertically.

Observations of CO, VOCs, peroxyacetyl nitrate (PAN) and some O₃ measurements are taken from the World Data Centre for Greenhouse Gases (WDCGG; <http://ds.data.jma.go.jp/gmd/wdcgg/>; see Figure 2 for locations). Most of the surface O₃ and CO measurements are provided by the National Oceanic and Atmospheric Administration (NOAA). NOAA CO is from
25 flask samples that have been analysed using gas chromatography (Novelli et al., 1998) and O₃ is measured by ultraviolet (UV) light absorption at 254 nm (Oltmans and Levy, 1994). The O₃ measurements at Cape Verde are provided by the University of York and were made using a UV light absorption instrument (Read et al., 2008). CO at Minamitorishima is from continuous measurements made by the Japan Meteorological Agency (JMA) using a gas chromatography (Watanabe et al., 2000). PAN measurements at Zugspitze and Schauinsland are provided to the WDCGG by the German Federal Environment Agency
30 (UBA) and were made using a commercial gas chromatograph (GC) analyser (Pandey Deolal et al., 2014). VOC measurements of ethene, ethane, propene, propane, toluene and butane made using gas chromatography at Hohenpeissenberg were provided by the German Meteorological Service (DWD) (Plass-Dülmer et al., 2002). All NO_x measurements were made using Chemiluminescence and are provided by DWD at Hohenpeissenberg (Mannschreck et al., 2004), UBA at Zugspitze, Empa (Swiss Federal Laboratories for Materials Science and Technology) at Jungfraujoch, Payerne and Rigi (Zellweger et al., 2003), and by RIVM (Netherlands National Institute for Public Health and the Environment) at Kollumerwaard.

5 3.3 Ozonesonde data climatology

Simulated O₃ profiles are compared to ozonesonde data from a climatology, which uses 17 years of ozone balloon soundings made between 1995 and 2011 (Tilmes et al., 2012). The data is available as profiles between 1000 hPa and 10 hPa at 42 stations, covering large parts of the globe. The model output is interpolated to the longitude and latitude of each station location. The site locations are shown in Figure 2b. The ozonesondes tend to measure concentrations around 10 ppbv higher over eastern 10 U.S. and around 5 ppbv lower over Europe compared to independent observational data from aircraft and surface data (Tilmes et al., 2012). For comparison to TOMCAT, both the model and the observations have been averaged into 3 different altitude and latitude bands for comparison.

3.4 Aircraft climatology ARCTAS data

~~Measurements from several aircraft campaigns have been compiled into regional mean profiles calculated over domains covered by the aircraft (Emmons et al., 2000) . The so-called climatology includes campaigns that were conducted between 1983 and 2001, covering several months of the year and both hemispheres, providing an insight into the temporal and spatial distribution of some species that are not routinely measured. Whilst this climatology is useful for this purpose, it is noted that aircraft campaigns often target pollution plumes that may not be captured by global models, in addition to the model generally not being representative of the year in which the observations were collected. This means that differences in meteorology and emissions may play a role in some of the discrepancies. For the TOMCAT comparisons we limit the data to the campaigns that occurred in the most recent~~ We use observations taken aboard the DC-8 aircraft between 29 June to 10 years (1992–2001) July 2008 during the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARTCAS) campaign (Jacob et al., 2010) . At this time the aircraft was based at Cold Lake, Canada and flew over large parts of North America and the Arctic. For the comparisons the model monthly mean data for July is used and is average over a region contained by the 20 25 maximum and minimum longitude and latitude range of the aircraft during the campaign.

3.5 OH estimates

OH is difficult to measure due to its very short lifetime (~1s) and low concentrations, and even with vast improvements to *insitu* measurement techniques (Heard and Pilling, 2003) , they do not provide a global picture. A common method to estimate OH is by using measurements of methyl chloroform (CH₃CCl₃, MCF), for which the primary loss channel is 30 through reaction with OH. Accurate determination of OH from MCF relies on accurate estimation of emissions and the use of models, introducing possible biases. These measurements are frequently used to estimate the global mean OH concentration (e.g., Krol et al., 1998; Prinn et al., 2001; Montzka et al., 2011) and can offer some insight into the regional distribution of OH (e.g., Krol and Lelieveld, 2003; Patra et al., 2014) . We use published estimates of global mean OH to discuss possible biases in simulated OH in Section 4.2.

4 Results

5 4.1 Simulated distributions of CO, O₃ and OH

Figure 3 shows annual mean surface and zonal mean concentrations of CO, O₃ and OH from ~~RUN~~^{the} ~~TOMCAT~~ 2008 simulation. CO is emitted directly from natural and anthropogenic sources and produced in the atmosphere from chemical destruction of VOCs (Logan et al., 1981). Direct emission at the Earth's surface and secondary production in the troposphere from VOCs (most notably CH₄) are estimated to be of equal importance in terms of total global tropospheric CO sources (Duncan et al., 2007). High concentrations due to direct emission of CO from fossil fuel burning can be seen in Figure 3a in the densely populated regions of North America, Central Europe and Asia. Large concentrations are also seen over regions with high rates of biomass burning, such as South America and Africa. Both at the surface and throughout the troposphere higher background concentrations of CO are seen in the Northern Hemisphere (NH) due to larger emissions.

O₃ is important in the troposphere as it is a major source of OH, the primary oxidising agent in the troposphere, and is an air pollutant and greenhouse gas (Monks et al., 2015a). It is not directly emitted but produced from photochemical reactions involving NO_x, VOCs and CO, and is transported from the stratosphere to the troposphere (Lelieveld and Dentener, 2000). The atmospheric burden of O₃ is controlled by a balance between these sources and loss through chemical reactions and deposition (Stevenson et al., 2006). Figure 3c shows the highest concentrations at the surface lie within the NH extra-tropical region due to the proximity to large emissions of NO_x and VOCs, and photochemical production. Some of the highest concentrations of O₃ are found downwind of regions with high NH anthropogenic emissions (identified by CO in Figure 3a). This is due to production of O₃ being greater downwind of source regions away from very high NO_x concentrations that can titrate O₃ in urban environments (Monks et al., 2015a). Low O₃ over the central Pacific Ocean and northern South America is also seen in the model, ~~most likely due to abundance of water vapour in the tropical regions limiting O₃ production through the reaction of O(¹D) with H₂O. Zonally averaged O₃ in Figure 3d shows the highest concentrations of~~. In the tropics, lower O₃ in stratosphere. Low concentrations are seen at ~~higher altitudes in the tropics due to the uplift of 100–300 hPa due to a higher tropopause in this region and the uplift of air with low O₃ concentrations within deep convection, whilst within deep tropical convection. At around 20–40 S/N, evidence of the downward transport of stratospheric O₃ to the troposphere can be seen in the extra-tropics due to the by the~~ Brewer-Dobson circulation (Butchart, 2014) ~~can be seen~~. The overall features of TOMCAT O₃ are consistent with multi-model results from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) (Young et al., 2013) and those observed by satellite (Ziemke et al., 2011). TOMCAT simulates an annual mean tropospheric burden of ~~336 Tg in RUN and 331 Tg in RUN~~, which agrees well with the present day ACCMIP multi-model mean tropospheric ozone burden of 337±23 Tg (Young et al., 2013).

OH is the dominant radical responsible for the removal of pollutants such as NO_x and VOCs from the atmosphere, initiating the production of O₃ (Gligorovski et al., 2015) and aerosols (e.g., Carlton et al., 2009). OH is produced in the troposphere when O₃ is photolysed to produce O(¹D) and subsequent reaction with H₂O. It is therefore produced in large quantities in the tropics, where there are large concentrations of H₂O and a high incidence of solar radiation. This can be seen in TOMCAT in

Figure 3e-f with high concentrations of OH occurring between 50°N and 50°S. The spatial distribution of TOMCAT surface OH is broadly similar to multi-model surface OH from the ACCMIP study shown by Voulgarakis et al. (2013).

5 4.2 Evaluation of OH

~~OH is difficult to measure due to its very short lifetime (~1s) and low concentrations, and even with vast improvements to *in situ* measurement techniques (Heard and Pilling, 2003), they do not provide a global picture. A common method to estimate OH is by using measurements of methyl chloroform (CH_3CCl_3 , MCF), for which the primary loss channel is through reaction with OH. Accurate determination of OH from MCF relies on accurate estimation of emissions and the use of models, introducing possible biases. These measurements are frequently used to estimate the global mean OH concentration (e.g., Krol et al., 1998; Prinn et al., 2001; Montzka et al., 2011) and can offer some insight into the regional distribution of OH (e.g., Krol and Lelieveld, 2003; Patra et al., 2014).~~

TOMCAT global mean airmass-weighted tropospheric OH was calculated using a climatological tropopause (see definition in Figure 4) following Lawrence et al. (2001). TOMCAT has an annual mean tropospheric OH concentration of ~~1.09 $\times 10^6$ molecules/cm³ in RUN and 1.08 $\times 10^6$ molecules/cm³ in RUN~~. Concentrations of global mean tropospheric OH calculated from MCF observations have been estimated to be 0.94×10^6 molecules/cm³ by Prinn et al. (2001), 1.0×10^6 molecules/cm³ by Krol et al. (2003) and 0.98×10^6 molecules/cm³ by Bousquet et al. (2005). These estimates indicate that the TOMCAT global mean OH may be slightly high. However, a recent inverse modelling study calculated a global mean OH concentration of 1.06×10^6 molecules/cm³, highlighting uncertainties in using MCF observations to calculate OH (Wang et al., 2008). In addition to this, concentrations reported by model intercomparison studies are also higher than those reported based on observations. The POLARCAT Model Intercomparison Project (POLMIP) found a multi-model mean value of $1.08 \pm 0.6 \times 10^6$ molecules/cm³ when using 8 models (including a previous version of TOMCAT). The multi-model mean was the same whether a climatological tropopause was used, as done here, or when the 150 ppb O₃ contour line was used. Voulgarakis et al. (2013) found a multi-model mean concentration of $1.17 \pm 0.1 \times 10^6$ molecules/cm³, when using a subset of 12 ACCMIP models, and Naik et al. (2013) found a multi-model mean of $1.11 \pm 0.2 \times 10^6$ molecules/cm³, when using all 16 ACCMIP models. Both of these ACCMIP concentrations were calculated using a tropopause of 200 hPa. However, Voulgarakis et al. (2013) found little difference in the resulting concentrations of OH when using different methods of defining the tropopause (200 hPa, 150 ppbv O₃ contour and the climatological tropopause, as used here).

Whilst comparing the global mean OH concentration in TOMCAT to those reported in the literature is very useful, it is also important to consider the regional distribution of OH in TOMCAT. Figure 4 shows TOMCAT OH ~~from both runs~~ averaged into 9 regional subsections defined by Lawrence et al. (2001), along with OH from Spivakovsky et al. (2000) (referred to as the Spivakovsky dataset) and the multi-model mean OH from the ACCMIP study (Naik et al., 2013). Patra et al. (2011) used the Spivakovsky dataset in a recent multi-model intercomparison project, but revised the concentrations down by 8% to match more recent measurements of MCF. This highlights that quantitative comparison of TOMCAT OH with the Spivakovsky dataset is limited due to observational and modelling uncertainties. However, the Spivakovsky dataset is still valuable for estimating the regional distribution of OH.

The largest concentrations of OH are found in the tropics for the Spivakovsky dataset and for the ACCMIP and TOMCAT simulations. However, the ACCMIP models have the highest OH concentrations between 500 hPa and 250 hPa, Spivakovsky 5 has the highest concentrations between 750 hPa and 500 hPa and TOMCAT has the highest concentrations between the surface and 750 hPa. Large differences in the spatial distribution of simulated OH has recently been identified in models, highlighting uncertainties in the ability of current models to accurately simulate OH concentrations and distributions (Emmons et al., 2015; Monks et al., 2015b). TOMCAT was shown to have lower photolysis rates in the upper troposphere and higher photolysis rates 10 in the lower troposphere compared to other models, with model differences in clouds and water vapour in the POLMIP models being identified as possible reasons for differences in the OH (Emmons et al., 2015; Monks et al., 2015b).

In addition to this, Patra et al. (2014) found that the NH to SH ratio of OH, inferred from observations of MCF, is equal to 0.97. TOMCAT has an annual NH/SH ratio of 1.37~~in both runs~~. Naik et al. (2013) found a NH/SH ratio of 1.28 ± 0.1 for the ACCMIP models, which is also higher than that estimated from observations, indicating that this is a common feature in global models.

15 TOMCAT OH results in a chemical methane lifetime of ~~7.6 yrs in RUN and 7.9 yrs in RUN~~. Voulgarakis et al. (2013) found an ACCMIP multi-model mean methane lifetime of 9.3 ± 0.9 , with a minimum of 7.1 years and a maximum of 13.9 years. This indicates TOMCAT has a methane lifetime that is generally shorter than other models. As the majority of methane oxidation occurs in the tropics near the surface (Lawrence et al., 2001; Bloss et al., 2005), the short methane lifetime is likely due to TOMCAT having a higher concentration of OH in this region compared to other models.

20 4.3 Evaluation of carbon monoxide

As mentioned in Section 4.1, CO is emitted from a wide range of natural and anthropogenic sources and can provide insight into model emissions and subsequent transport of sources due to its lifetime of several months. Figure 5 shows retrieved CO from MOPITT (see Section 3) at 500 hPa during April and October 2008 along with simulated CO from ~~RUN~~TOMCAT with the MOPITT averaging kernels applied.

25 In April, both the model and the satellite show higher CO concentrations in the NH compared to the Southern Hemisphere (SH) due to a longer CO lifetime at this time of year in conjunction with higher anthropogenic emissions in the NH. MOPITT observes concentrations around 10–30 ppbv larger than simulated in the NH mid-latitudes and Arctic (Figure 5c). This negative model bias is a well-known problem with current CTMs during winter and spring, with models having a 15 to 50 ppbv negative bias against MOPITT at 500 hPa in April in the NH (Shindell et al., 2006) and 5 to 40 ppbv negative bias against Arctic surface 30 stations in Spring (Monks et al., 2015b). The model shows the best agreement in the NH tropics at this time of year.

TOMCAT CO concentrations in the SH in April are around 10–15 ppbv larger than observed. Shindell et al. (2006) found good agreement between a 26-model ensemble mean at 500 hPa compared to MOPITT, with individual models showing both negative and positive biases of between -15 ppbv and +15 ppbv, showing that the TOMCAT bias at this time of year is at the high end of the multi-model positive bias range.

35 The model negative bias in the NH and positive bias in the SH leads to a simulated interhemispheric gradient that is too low (see Figure 5c), which is a common feature in chemistry models (Shindell et al., 2006). Several inverse modelling studies have

suggested that wintertime CO emissions in the NH need to be increased in order to better match observations of CO (Pétron et al., 2004; Kopacz et al., 2010; Fortems-Cheiney et al., 2011). Transport errors in the model could also play a role, however, they are unlikely to cause such widespread biases of this magnitude in background CO. In addition to this, as mentioned in 5 Section 4.2, the OH in TOMCAT is most likely too high at the surface, particularly in the tropics, and the NH/S_H OH ratio in TOMCAT and other chemistry models is higher than estimates based on observations, suggesting that there is too much OH in the NH or too little in the SH. This is likely to influence the lifetime of simulated CO and will contribute to the NH and 10 SH biases. Strode et al. (2015) showed that by lowering the NH/S_H OH ratio of current state-of-the-art models, simulation simulations of CO can be improved. The cause of the lower simulated NH/S_H OH ratio in models is still unclear and may be linked to emission biases, where higher emissions of CO and VOCs in the NH may reduce OH concentrations, reducing the NH/S_H OH ratio.

In October, the interhemispheric gradient in CO is no longer as clear due to longer CO lifetimes in the SH and shorter lifetimes in the NH. This time of year is characterised by peak fire emissions in the SH (van der Werf et al., 2010). For this reason, 15 high concentrations of CO are seen by MOPITT over South America and there is a shift in the biomass burning emissions further south over Africa, resulting in higher CO over the Southern Ocean. TOMCAT also shows higher concentrations over the Southern Ocean due to the influence of fire emissions compared to April. However, fire emission location errors are clearly contributing to a mismatch between the CO plumes in the model and those seen by MOPITT. Total column CO over this region suggests that emissions from fires may be too large in the tropics, particularly over tropical Asia (not shown), and the fires are located too far north in Africa and too far west in South America, resulting in too much CO being transported out over the 20 oceans in the tropics (see Figure 5d,e). Naik et al. (2013) also showed that the ACCMIP multi-model annual mean simulated CO at 500 hPa was 2–45 ppbv too high compared to MOPITT in this region supporting a high bias in CO fire emissions across different emission inventories in the SH and tropics at this time of year. Outside of the 10°S–30°N region, the zonal mean CO shows much better agreement between TOMCAT and MOPITT than seen in April (see Figure 5f).

Figure 6 compares simulated and measured CO at 14 different surface observatories that are located at several different 25 latitudes and longitudes for the year 2008 (see Figure 2a for station locations). TOMCAT generally captures the seasonal cycle, with high correlations values found at most stations (see r values in Fig. 6). However, the amplitude of the seasonal cycle is often less pronounced in the model due to biases that exist in the first half of the year in some regions. In agreement with MOPITT comparisonsthe MOPITT comparison results, the model shows a large negative bias in winter and spring in the NH, with particularly large biases at stations located at higher latitudes (Alert and Mace Head). This has been documented at Arctic 30 surface sites previously (Shindell et al., 2008; Monks et al., 2015b). At latitudes $>25^{\circ}\text{N}$ the model has a normalised mean bias (NMB) of between -17.0 % and -38.1 % for RUN and -19.2 % and -33.1 % for RUN. The winter/spring model bias is smaller at locations closer to the NH tropics resulting in lower NMB, with the best overall model agreement found at Key Biscayne in Florida (NMB of 3.6% and 4.8%) -14.6 % and -22.4 %. In the SH, the model has a tendency to overestimate CO concentrations The model performs the best near the tropics, particularly in the austral summer. Overall, the model has NMBs in the SH of with NMBs of between 4.7 % - 8.5 % and 37.3 % for RUN and -0.8 % and 30.3 %, with the best model performance at Easter Island in the Pacific Ocean, and the worst at Cape Grim, Australia. In the SH, RUN has consistently lower NMBs, most likely

due to lower emissions. ~~8.2 %~~. In the mid to high SH latitudes the model overestimates CO concentrations throughout the year, with the largest biases occurring during the austral summer (NMBs of 21.2 % to 28.5 %).

5 The 26 multi-model study by Shindell et al. (2006) found that models have a negative bias of between 20–80 ppbv at Alert in the Arctic during winter/spring and a more persistent positive bias throughout the year of up to 20–25 ppbv at Cape Grim, exhibiting a ~~similar~~ transition from a negative bias in the NH to a positive bias in the SH ~~that is similar~~ to that found in TOMCAT. TOMCAT is within the bias range at Alert, with a winter negative bias of up to ~50 ppbv, and at the upper end of the bias range at Cape Grim, with up to ~25 ppbv at Cape Grim. ~~This and the MOPITT comparisons shows~~ ~~The surface and~~
10 ~~MOPITT comparisons show~~ that these model biases exist at the surface and throughout the free troposphere, and are generally consistent with biases found in other chemical transport models.

4.4 Evaluaton of ozone

Ozonesonde data is compared to simulated O₃ in Figure 7. The data has been separated into three different altitude and latitude bands. The model overestimates O₃ at higher NH and SH latitudes in the highest altitude band (NMB of 22 % to 43.2 %),
15 possibly due to too much downward mixing of stratospheric O₃ in the model at these altitudes. TOMCAT also overestimates O₃ at the surface in the tropics (NMB of 14.4 % to 16.7 %), but the model lies within the range of observations. Elsewhere, the model has a negative bias (NMB of -1.2 % to -24.6%), but lies within the range of observations at several times of the year. Most of the negative bias in the higher latitudes is being driven by wintertime underestimates in O₃ in both the SH and NH. Young et al. (2013) ~~compared~~ ~~found that~~ the multi-model ACCMIP mean O₃ ~~to ozonesonde data and found that it~~ is also
20 negatively biased in the SH during the winter months, ~~but when compared to the same data, suggesting that this is a common~~ feature in chemistry models. However, they found that the ACCMIP models overestimated O₃ in the NH high latitudes during winter.

~~This~~ ~~The~~ low TOMCAT bias in wintertime O₃ can also be seen in surface data located at high latitudes ~~in the SH at Arrival Heights~~ (see Figure 8), ~~where~~ TOMCAT has a negative O₃ bias of ~10–15 ppbv during the ~~NH boreal winter at Heimaey, Iceland and during the SH austral winter at Arrival Heights and South Pole~~ (NMB of ~~-19.7 % to -26.3 %~~). ~~All three of these~~ stations are located near the poles suggesting ~~-21.1 %~~. ~~This suggests~~ that the model may have difficulties reproducing O₃ photochemistry in the winter in remote, dark and cold regions or the model may deposit too much O₃ onto snow/ice covered surfaces. Whilst most models in the POLMIP study were also negatively biased at the Summit observatory in the Arctic during winter, TOMCAT simulated some of the lowest concentrations (Monks et al., 2015b). Outside of the poles, the model simulates
30 concentrations of O₃ that ~~agree well with observations at this time of the year (see Figure 8), with the worst agreement occurring~~ ~~in the NH summer, where the model are in much better agreement with the observations (NMB of 0.7 % to 15.2 %)~~. ~~In the~~ NH during the summer TOMCAT tends to overestimate concentrations. This is a common feature in models in the NH during summer, which has been identified to be paticularly pronounced over Eastern U.S. (e.g., Ellingsen et al., 2008; Fiore et al., 2009; Yu et al., 2010). O₃ at the surface is also compared ~~at to data from~~ 44 EPA CASTNET stations located in the U.S. (Figure 9). This high ~~summer~~ bias over the U.S. is clearly evident, with a large mean bias (MB) of ~~26.8/32.3/28~~ ppbv (NMB of ~~89.9/107.6~~ = ~~97.8~~ %). The best agreement is seen in winter (MB < ~~4.6~~ = ~~2.7~~ ppbv, NMB < ~~14.6~~ = ~~-8~~ %). ValMartin et al. (2014)

showed that model summertime O₃ biases could be reduced from 44 % to 28 % over the U.S. and from 25 % to 14 % over Europe when improvements were made to ~~the a~~ coupled land-atmosphere model's deposition scheme. This suggests that using 5 a more sophisticated deposition scheme coupled to a land model may improve TOMCAT simulations of summertime O₃.

~~RUN_{sub-column} Sub-column~~ O₃ between 0–6 km (up to ~500 hPa) is compared to GOME-2 retrievals in Figure 10. MB errors that are greater than the satellite error are highlighted with green polygons. In DJF, GOME-2 measures the highest concentrations of O₃ (~25 DU) in regions near O₃ precursor emissions and enough sunlight to initiate photochemistry at this 10 time of year (e.g. India, China and northern Africa; Figure 10b). TOMCAT shows negative MBs of up to -10 DU in several regions, with some of the larger biases being co-located with high observed O₃ concentrations (see Figure 10d). Comparisons to ozonesondes (see Figure 7) further support this and show that the model O₃ may be biased low (by 5–10 ppbv) in the tropical 15 region at this time of year at altitudes between 750 hPa and 450 hPa (although the model does lie within the ozonesonde observed ranges). In JJA, the model bias is much smaller with very few significant MBs being highlighted (see Figure 10c). There is evidence that the model overestimates O₃ at this time of year over South-East U.S., in agreement with the CASTNET 20 model-observation comparisons, and some evidence that O₃ is also overestimated near Cape Verde off the coast of Africa, as seen in Figure 8.

4.5 Evaluation of VOCs

~~Global maps of simulated propene (C₃H₆, Figure ??), ethene (C₂H₄, Figure ??), propane (C₃H₈, Figure ??) and ethane (C₂H₆, Figure ??) are shown averaged over 2–4 km and 5–8 km for both model runs. The lifetimes of these VOCs vary from a few hours to a few months, with propene having the shortest lifetime, followed by ethene, propane and then ethane (Rudolph et al., 1989). Due to the shorter lifetime of alkenes (C₂H₄, C₃H₆), the concentrations are mostly elevated near source regions due to limited long-range transport. When the lifetimes of the alkenes are longer due to lower OH (e.g. in winter), higher concentrations can be seen further from sources. In contrast, the longer lifetime of alkanes results in much more transport throughout the globe, with the highest concentrations being located in the NH due to higher anthropogenic emissions.~~

~~25 Overlaid on the maps are mean observed concentrations from several aircraft campaigns collected between 1992 and 2001 (Emmons et al., 2000). The spatial distribution of high and low concentrations of the alkanes and alkenes seems to be well represented by the model. However, it is noticeable that simulated concentrations, particularly of alkanes, are consistently lower than observed, suggesting a negative bias in the model. As these VOCs are emitted and not formed, this indicates that emissions are probably too low. A negative model bias in ethane and propane at several surface sites located in the NH has 30 been shown previously to exist in several models (Emmons et al., 2015).~~

In Figure 11, measurements of ethene, ethane, propene, propane, toluene and butane are compared to simulated concentrations at the mountain site Hohenpeissenberg, Germany. The observations show a ~~distinct seasonal cycle~~ seasonal cycle that is particularly pronounced for ethene, ethane, propene and toluene, with peak concentrations in winter and spring, when OH concentrations are lower and the lifetimes of VOCs are longer, and a minimum in summer. For ethane and propane, the model captures the seasonal ~~cycle (r=0.88–0.98 transitions (r-values of 0.94 and 0.99, respectively)~~ transitions (r-values of 0.94 and 0.99, respectively), but shows a much smaller amplitude due to large negative biases, particularly in winter (NMB of ~~-29.2 % to -80.5 %~~ -31.3 % and -25.8 %,

respectively). This can also be seen at high latitudes over North America throughout the troposphere when compared to the ARCTAS data in Figure 12. It has been suggested previously that the current anthropogenic emissions of ethane are too low in global models (Franco et al., 2016; Tilmes et al., 2016), which is likely to explain the negative bias in TOMCAT ethane. Propane is also shown to be too low throughout the tropospheric column (Fig. 12) and has also been found to be biased low in the NH across different models, suggesting a similar problem with emissions (Emmons et al., 2015; Tilmes et al., 2015). For ethene and propene, the seasonal cycle is not well captured by the model due to enhancements in summer ($r=0.13$ – 0.25 – 0.70). The model simulates summer minimum concentrations in JJA elsewhere in the NH (see Figure ??–??). Therefore, this is likely due to incorrect local emissions or at this time of year (from biogenic or fire sources) or difficulties capturing local turbulent transport at this mountain site, which is a common problem in models (Zhang et al., 2008; Feng et al., 2011). Similar to ethane and propane, the model also shows negative biases that are particularly large in winter, suggesting underestimated anthropogenic emissions in the NH. For toluene and butane, the model captures the seasonal cycle well ($r=0.90$ – 0.95 – 0.91 and 0.97 , respectively), but some large positive biases are found (toluene: NMB of $25.3/302$ %, butane: NMB of $91/155.3$ %). For all the VOCs, the two simulations show large differences in concentrations and biases, with RUN generally simulating lower VOC concentrations for all the VOCs at this measurement site. This results in lower NMBs in toluene and butane in RUN but higher NMBs for the C2–C3 alkanes/alkenes. This is likely due to the fact that RUN and RUN emit different magnitudes of anthropogenic VOCs (see Table 1). In addition to this, RUN has larger biogenic C2–C3 alkene emissions, resulting in higher concentrations compared to RUN near biogenic emissions sources (e.g. over the Amazon) (see Figures ??–??). This highlights that regional model performance for VOCs is very dependent on the emission dataset chosen, consistently throughout the year (NMB of 270.5 % and of 175.6 %, respectively).

4.6 Evaluation of reactive nitrogen

Oxides of nitrogen (NO_y) are important atmospheric pollutants and are key in the production of O₃. In addition, speciation of NO_y is dependent on oxidative capacity, organic chemistry and heterogeneous chemistry. Hence, evaluation of speciated NO_y is a valuable test of several inter-related aspects of model chemistry. Here we use observations of nitrogen oxide (NO), NO₂, nitric acid (HNO₃) and peroxyacetyl nitrate (PAN) to evaluate the model NO_y.

In Figure 13 2008 measurements of NO_x (NO + NO₂) from two European observatories are compared to simulated concentrations at several European locations at a range of altitudes (see Figure 2c for locations). The observations generally show a minimum in summer and a maximum in winter, with the model showing variable skill in capturing the seasonal cycle depending on the surface station ($r=0.08$ – 0.89). Both TOMCAT simulations underestimate NO_x concentrations (NMB well suggesting that the model is able to reproduce seasonal changes in photochemistry (r -values of -30 %– 0.89 and -101.1 %), with RUN having the lowest overall bias due to larger concentrations 0.94 , respectively). However, TOMCAT overestimates the concentrations at Hohenpeissenberg throughout the year (NMB=78.9 %) but underestimates concentrations at Payerne (NMB=–28.4 %). The model shows only marginally higher concentrations at Payerne compared to Hohenpeissenberg. The observations show that this difference is larger in reality, suggesting a higher gradient in concentrations between the two different stations both horizontally and vertically. The largest

biases occur in winter, resulting in a smaller seasonal cycle amplitude. As NO_x is very short-lived, it is difficult for global models to reproduce observations due to coarse horizontal resolutions (Huijnen et al., 2010), which may partly explain the negative biases seen in the model and vertical resolutions, which is likely to affect the models ability to capture concentration gradients (Huijnen et al., 2010). The model is able to capture the changes in NO_2 with altitude and captures the magnitude well in the middle troposphere (see Fig 12).

Figure 14 shows 2008 DJF and JJA satellite OMI NO₂ column for 2008 and alongside the TOMCAT MB from RUN. Due to the short lifetime of NO_2 , high concentrations are located near emission regions. In the NH, high concentrations are seen over Asia, North America and Europe, near some of the largest anthropogenic emission sources. In both seasons the model simulates concentrations that are too high over parts of Europe. This is in contrast to the surface comparisons shown in Figure 13, where the model underestimated NO_x at several locations at altitudes below 3 km. However, most of these sites were located further west of the region where OMI shows a positive bias in the model likely to be linked to emissions due to the short lifetime of NO_2 . Due to the location near the Baltic and North Seas, this could indicate that ship emissions are too large in this region. Large negative biases in NO_2 near China are seen in the model in the NH winter. This has been seen in several models previously when comparing to OMI and is thought to be due to anthropogenic emissions that are too low (Emmons et al., 2015). In contrast, TOMCAT has a positive model bias in this region during summer, most likely due to fire emissions that are the FINN fire emissions being too high, which has also been seen in multiple models being compared to OMI (Emmons et al., 2015).

In the SH, OMI observes the largest concentrations over the high biomass burning regions of South America, Africa and Australia. In these regions the model shows NO_2 concentrations that are too low during both seasons, suggesting emissions from fires FINN fire emissions are too low in the SH. This is in contrast to CO satellite comparisons, which suggested fire emissions are too high in this region (see Section 4.3). This therefore indicates that emission factors used to calculate fire emissions need to be further evaluated in the tropics and the SH.

PAN is formed from the oxidation of VOCs and reaction of the peroxyacetyl radical with NO_2 . It is an important NO_x reservoir and acts to transport and supply NO_x to regions remote from emission sources, where it can contribute to O_3 production (e.g., Hudman et al., 2004). Figure ?? shows TOMCAT PAN peaks in the lower troposphere in spring and peaks in the upper troposphere in summer, which is in agreement with the GEOS-Chem model (Fischer et al., 2014) and upper tropospheric satellite observations from MIPAS-E (Moore and Remedios, 2010). The aircraft climatology shows that TOMCAT captures the spatial variability of PAN well (see Figure ??). However, TOMCAT may overestimate concentrations of PAN over North America in winter. Pope et al. (2016) also found that TOMCAT PAN may be too high against aircraft data in the Arctic and over North America in winter. They also found that TOMCAT PAN overestimated upper tropospheric MIPAS satellite PAN in winter and spring in parts of the NH. A high spring bias in PAN has also been shown to be present in the GEOS-Chem CTM when compared to aircraft data (Fischer et al., 2014).

Figure 15 shows surface PAN comparisons at two locations in Europe at two different altitudes, with concentrations peaking in the spring. Both model simulations show a summer maximum and a Figure 15 shows PAN comparisons at the Mountain site, Schauinsland, Germany. Observations at this location show concentrations that peak in April, with a winter minimum.

TOMCAT concentrations peak later in the year in June but capture the strong drop off in concentrations leading towards a winter minimum ($r=0.8-0.85$), with a ~ 2 month lag in the peak concentrations. At Schauinsland (1205 m), both of the model simulations lie within the range of observations. At Zugspitze, which has lower observed concentrations, both model simulations fall outside of the range of observations during certain times of the year. RUN simulates the highest concentrations and as a result, overestimates PAN in summer at both stations (NMB=43–77.5%). RUN has much better overall agreement with observations due to lower simulated concentrations (0.82). Simulated concentrations show reasonable agreement with the observations (NMB=−4.0–17.46.9%). RUN also showed better agreement against the aircraft climatology in the NH due to lower concentrations. As already mentioned, the VOC emissions differ between the runs, which will affect PAN production. Differences in model meteorology between the years may also play a role.

15 HNOPAN during the summer months shows reasonable agreement with aircraft data shown in Figure 12. However, HNO_3 is compared to the aircraft climatology in Figure ???. The spatial variability in HNO_3 is generally captured overestimated in the model where observations exist. However, both RUN and RUN simulate HNO_3 concentrations that are higher than those from the aircraft climatology over parts of North America during DJF and MAM, possibly due to too much production or not enough washout. In an Arctic model intercomparison project (POLMIP), TOMCAT had some of the highest concentrations of PAN 20 and HNO_3 compared to other models (Emmons et al., 2015), suggesting that TOMCAT NO_y production may be more efficient than other models, is higher and/or loss may be underestimated in TOMCAT in the mid to high latitude NH regions. This is important for lower compared to other models. The observations at Schauinsland suggest that TOMCAT may do a better job in simulating PAN over Europe at lower latitudes. In addition to this, Pope et al. (2016) found that TOMCAT PAN overestimated upper tropospheric MIPAS PAN at altitudes above 200 hPa in winter and spring. Due to the importance of reactive nitrogen in 25 O_3 production and warrants further investigation there is a need for further investigation and evaluation of these species in the future when more recent observations become available.

5 Summary

The TOMCAT This paper describes the TOMCAT 3-D chemical transport model has been updated with the Extended Tropospheric Chemistry scheme, adding 's tropospheric chemistry scheme. The current scheme has a more detailed representation of 30 hydrocarbon chemistry compared to the previously documented version found in Arnold et al. (2005). The current scheme includes the degradation of ethene, propene, toluene and butane. Monoterpene chemistry has also been added based on the Extended Tropospheric Chemistry scheme and monoterpene chemistry based on MOZART-3 chemistry. Two model simulations, which differ in their boundary conditions but both use the new chemistry scheme, are used to evaluate the model. A one-year 35 simulation for the year 2008 is used to document model performance against a range of surface, satellite, aircraft and balloon measurements. The model is generally able to capture the main spatial and seasonal features of high and low concentrations of CO , O_3 , VOCs and reactive nitrogen. However, several negative and positive biases are present in TOMCAT during certain times of the year and at certain locations. Some of these biases are prevalent in current state-of-the-art chemistry models, but some biases that are specific to TOMCAT are also highlighted.

TOMCAT global mean tropospheric OH ($1.07\text{--}1.08\text{--}1.08 \times 10^6$ molecules/cm³) is higher than estimates inferred from MCF observations ($0.94\text{--}1.0 \times 10^6$ molecules/cm³). However, this is a common feature across chemistry models and the TOMCAT global mean OH is at the lower end of concentrations reported in previous multi-model intercomparison projects ($1.08\text{--}1.17 \times 10^6$ molecules/cm³). TOMCAT has the highest concentrations (in molec./cm³) of OH in the lower tropical troposphere, which is in contrast to the ACCMIP multi-model mean OH, which has the highest OH concentrations in the tropical upper troposphere. ~~However, observationally-constrained~~ Observationally-constrained OH shows the highest ~~concentration~~ concentrations of OH in the middle tropical troposphere suggesting that TOMCAT has too much OH at the surface in the tropics. Further to this, TOMCAT has a higher NH/SH OH ratio (1.37) compared to the ratio inferred from MCF observations (0.98), which is again a common feature in chemistry models, with TOMCAT being at the upper limit of the multi-model mean value calculated from the ACCMIP models (1.28 ± 0.1). This suggests that simulated OH in current chemistry models is largely uncertain ~~and more work is needed to understand the cause of the lower simulated NH/SH OH ratio in models. One possibility~~ could be underestimated emissions in the NH which may be contributing to OH concentrations being too high in this region.

TOMCAT CO is negatively biased during winter and spring in the NH when compared to MOPITT and surface observations. In contrast, CO is positively biased throughout the year in the SH. The negative bias in the NH is a common feature in chemistry models and TOMCAT lies well within the range of biases found in other models. The TOMCAT SH positive bias is at the upper range of positive biases reported in other models, with some models reporting negative biases. Underestimated emissions in the NH are thought to play a role in the negative NH CO bias, whilst comparisons with MOPITT suggest that TOMCAT fire emissions may be too high in the SH. ~~Biases could also be reduced by lowering the NH, contributing to the model positive bias. OH biases could also play a role in the CO bias, in particular in the NH where near-surface OH is around 50 % larger than methyl chloroform-constrained OH estimates. Lower OH concentrations in the model at the surface would lead to an increase in CO concentrations in the NH and would also reduce NH/SH OH ratio, where a lower OH concentration in the NH will increase the lifetime of CO and/or a higher OH concentration in the SH would decrease the lifetime of CO~~ SH OH ratio.

TOMCAT is able to capture the seasonality of O₃ in most locations, with the model lying within the range of observations made during balloon soundings during most times of the year. The notable exceptions to this are: 1) at high latitudes during winter conditions, where TOMCAT simulates O₃ that is negatively biased by up to 15 ppbv when compared to both surface and ozonesonde measurements and 2) in the NH during summer, where TOMCAT is positively biased by up to ~~32~~ 28 ppbv over North America when compared to surface sites. GOME-2 satellite data shows that model performance is better in JJA compared to ~~DJADJF~~, where the model underestimates O₃ by up to 10 DU in regions with high observed O₃ concentrations near Asia and Africa.

~~Comparison of simulated VOCs, NO_x, PAN and HNO₃ to observations highlights the sensitivity of model performance to model boundary conditions and emissions, with a large range in concentrations between RUN and RUN. Comparison with an aircraft climatology shows that the model generally captures the spatial and temporal variability of these species. However some biases are found. VOC aircraft and VOC surface measurements show large negative biases in simulated winter/spring C2–C3 alkanes and alkenes, which is likely driven by underestimated anthropogenic emissions. This has been seen previously~~

5 for ethane and propane in several models in the NH. In contrast, RUN showed very good agreement with the seasonality and magnitude of surface measurements of toluene and butane in Europe.

TOMCAT is also able to capture the seasonal minima and maxima of PAN that vary with altitude. However, TOMCAT has a 2 month lag in peak PAN concentrations when compared to mountain site observations made in Europe. The aircraft climatology also shows that wintertime PAN may be overestimated in winter and HNO_3 may be overestimated in winter and spring in 10 some regions over North America. Whilst the aircraft climatology is useful for general comparisons it is noted that the model simulations do not match the observations in time and therefore some discrepancies are expected. For NO captures the rapid decline in PAN concentrations between summer and winter at a European mountain site but simulates peak PAN concentrations in June rather April. TOMCAT is able to capture the seasonal cycle of NO_x , the amplitude of simulated NO_2 at European sites is much smaller than observed due to large negative model biases, particularly in winter well at two European surface sites but has trouble capturing the concentrations, overestimating them at the lower altitude Payerne site and underestimating them at the higher altitude site of Hohenpeissenberg. This is likely to be at least partly due to the very short lifetime of NO_x and the 5 coarse model grid. Satellite Tropospheric satellite OMI NO_2 showed regional differences in TOMCAT biases, with negative biases existing over China in DJF (possibly due to anthropogenic emissions) and South America and Africa (possibly due to fire emissions), and positive biases over Europe in DJF and JJA. The biases over Asia have been shown to exist in several other models when using the same emissions as used for RUN here. Further to this, models have been shown previously to vary widely in the simulation of species such as HNO_3 , PAN and acetaldehyde. Therefore, observations of such these species that 10 are collected continuously throughout the year at several locations globally would be valuable in evaluating chemical transport models in the future and understanding model biases in O_3 .

6 Code Availability

TOMCAT/SLIMCAT (www.see.leeds.ac.uk/tomcat) is a UK community model. It is available to UK (or NERC-funded) researchers who normally access the model on common facilities or who are helped to install it on their local machines. As it 15 is a complex research tool, new users will need help to use the model optimally. We do not have the resources to release and support the model in an open way. Any potential user interested in the model should contact Martyn Chipperfield. The model updates described in this paper are included in the standard model library.

Acknowledgements. Funding for this work was provided by the NERC EurEx project (NE/H020241/1) and TOMCAT model simulations were performed on the UK Archer HPC system. We would like to thank the many providers of observational data that has been used in this 20 paper. Specifically, the World Ozone and Ultraviolet Data Center (WOUDC), the National Oceanic and Atmospheric Administration Earth System Research Laboratory, Global Monitoring Division (NOAA ESRL, GMD) and the Southern Hemisphere ADDitional OZonesondes (SHADOZ) for data that was used in the ozonesonde climatology. The MOPITT team that provided CO satellite data and L.K. Emmons for providing code to process the MOPITT data. The TEMIS website and KNMI OMI satellite group for use of tropospheric NO_2 column data. The ARCTAS DC-8 team who provided aircraft measurements. The United States Environmental Protection Agency (EPA) Clean Air Status

25 and Trends Network (CASTNET) for O₃ surface measurements. The Global Atmosphere Watch (GAW) program for the use of the World Data Centre for Greenhouse Gases (WDCGG), which provided surface measurements of CO, O₃, NO_x, PAN and VOCs. WDCGG providers whose data was used includes NOAA/ESRL, University of York, UBA, Empa, RIVM and DWD.

References

- Archibald, A. T., Cooke, M. C., Utembe, S. R., Shallcross, D. E., Derwent, R. G., and Jenkin, M. E.: Impacts of mechanistic changes on HO_x formation and recycling in the oxidation of isoprene, *Atmos. Chem. Phys.*, 10, 8097–8118, doi:10.5194/acp-10-8097-2010, <http://www.atmos-chem-phys.net/10/8097/2010/>, 2010.
- Archibald, A. T., Levine, J. G., Abraham, N. L., Cooke, M. C., Edwards, P. M., Heard, D. E., Jenkin, M. E., Karunaharan, A., Pike, R. C., Monks, P. S., Shallcross, D. E., Telford, P. J., Whalley, L. K., and Pyle, J. A.: Impacts of HO_x regeneration and recycling in the oxidation of isoprene: Consequences for the composition of past, present and future atmospheres, *Geophys. Res. Lett.*, 38, n/a–n/a, doi:10.1029/2010GL046520, 105804, 2011.
- Arnold, S. R., Chipperfield, M. P., and Blitz, M. A.: A three-dimensional model study of the effect of new temperature-dependent quantum yields for acetone photolysis, *J. Geophys. Res.*, 110, doi:10.1029/2005JD005998, 2005.
- Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review, *Atmos. Environ.*, 37, Supplement 2, 197–219, doi:[http://dx.doi.org/10.1016/S1352-2310\(03\)00391-1](http://dx.doi.org/10.1016/S1352-2310(03)00391-1), 2003.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: IUPAC Subcommittee for Gas Kinetic Data Evaluation. Summary of evaluated kinetic data and photochemical data for atmospheric chemistry, <http://www.iupac-kinetic.ch.cam.ac.uk/>, accessed in 2004, a.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: IUPAC Subcommittee for Gas Kinetic Data Evaluation. Summary of evaluated kinetic data and photochemical data for atmospheric chemistry, <http://www.iupac-kinetic.ch.cam.ac.uk/>, accessed in 2005, b.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: IUPAC Subcommittee for Gas Kinetic Data Evaluation. Summary of evaluated kinetic data and photochemical data for atmospheric chemistry, <http://www.iupac-kinetic.ch.cam.ac.uk/>, accessed in 2006, c.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II - gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, doi:10.5194/acp-6-3625-2006, 2006.
- Berntsen, T., Fuglestvedt, J., Joshi, M., Shine, K., Stuber, N., Ponater, M., Sausen, R., Hauglustaine, D., and Li, L.: Response of climate to regional emissions of ozone precursors: sensitivities and warming potentials, *Tellus B*, 57, 283–304, doi:10.1111/j.1600-0889.2005.00152.x, 2005.
- Bloss, W. J., Evans, M. J., Lee, J. D., Sommariva, R., Heard, D. E., and Pilling, M. J.: The oxidative capacity of the troposphere: Coupling of field measurements of OH and a global chemistry transport model, *Faraday Discuss.*, 130, 425–436, doi:10.1039/B419090D, 2005.
- Boersma, K., Jacob, D., Bucsela, E., Perring, A., Dirksen, R., van der A, R., Yantosca, R., Park, R., Wenig, M., Bertram, T., and Cohen, R.: Validation of OMI tropospheric NO₂ observations during INTEX-B and application to constrain emissions over the eastern United States and Mexico, *Atmos. Environ.*, 42, 4480 – 4497, doi:<http://dx.doi.org/10.1016/j.atmosenv.2008.02.004>, 2008.

- Boersma, K. F., Eskes, H. J., Veefkind, J. P., Brinksma, E. J., van der A, R. J., Sneep, M., van den Oord, G. H. J., Levelt, P. F., Stammes, P., Gleason, J. F., and Bucsela, E. J.: Near-real time retrieval of tropospheric NO₂ from OMI, *Atmos. Chem. Phys.*, 7, 2103–2118, 30 doi:10.5194/acp-7-2103-2007, 2007.
- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905–1928, doi:10.5194/amt-4-1905-2011, 2011.
- Bousquet, P., Hauglustaine, D. A., Peylin, P., Carouge, C., and Ciais, P.: Two decades of OH variability as inferred by an inversion of 35 atmospheric transport and chemistry of methyl chloroform, *Atmos. Chem. Phys.*, 5, 2635–2656, doi:10.5194/acp-5-2635-2005, 2005.
- Braak, R.: Row Anomaly Flagging Rules Lookup Table, KNMI Technical Document TN-OMIE-KNMI-950, 2010.
- Breider, T. J., Chipperfield, M. P., Richards, N. A. D., Carslaw, K. S., Mann, G. W., and Spracklen, D. V.: Impact of BrO on dimethylsulfide 40 in the remote marine boundary layer, *Geophys. Res. Lett.*, 37, doi:10.1029/2009GL040868, l02807, 2010.
- Butchart, N.: The Brewer-Dobson circulation, *Rev. Geophys.*, 52, 157–184, doi:10.1002/2013RG000448, 2014.
- Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmos. Chem. Phys.*, 9, 4987–5005, doi:10.5194/acp-9-4987-2009, 2009.
- Carver, G., Brown, P., and Wild, O.: The ASAD atmospheric chemistry integration package and chemical reaction database, *Comput. Phys. 5 Commun.*, 105, 197–215, 1997.
- Chameides, W. L., Lindsay, R. W., Richardson, J., and Kiangs, C. S.: The role of biogenic hydrocarbons in urban photochemical smog: Atlanta as a case study, *Science*, 241, 1473–1475, doi:10.5194/acp-9-4987-2009, 1988.
- Chipperfield, M. P.: New version of the TOMCAT/SLIMCAT off-line chemical transport model: Intercomparison of stratospheric tracer 5 experiments, *Q. J. R. Meteorol. Soc.*, 132, 1179–1203, doi:10.1256/qj.05.51, 2006.
- Chipperfield, M. P., Dhomse, S. S., Feng, W., McKenzie, R. L., Velders, G., and Pyle, J. A.: Quantifying the ozone and ultraviolet benefits 10 already achieved by the Montreal Protocol, *Nat. Commun.*, 6, doi:10.1038/ncomms8233, 2015.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Mor- 15 crette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.
- Deeter, M. N.: MOPITT (Measurements of Pollution in the Troposphere) Version 6 Product User's Guide, 2013.
- Deeter, M. N., Edwards, D. P., Gille, J. C., Emmons, L. K., Francis, G., Ho, S.-P., Mao, D., Masters, D., Worden, H., Drummond, J. R., and 20 Novelli, P. C.: The MOPITT version 4 CO product: Algorithm enhancements, validation, and long-term stability, *J. Geophys. Res.*, 115, D07 306, 2010.
- Deeter, M. N., Worden, H. M., Gille, J. C., Edwards, D. P., Mao, D., and Drummond, J. R.: MOPITT multispectral CO retrievals: Origins 25 and effects of geophysical radiance errors, *J. Geophys. Res.*, 116, doi:10.1029/2011JD015703, 2011.
- Dhomse, S. S., Chipperfield, M. P., Feng, W., Ball, W. T., Unruh, Y. C., Haigh, J. D., Krivova, N. A., Solanki, S. K., and Smith, A. K.: Stratospheric O₃ changes during 2001–2010: the small role of solar flux variations in a chemical transport model, *Atmos. Chem. Phys.*, 13, 10 113–10 123, doi:10.5194/acp-13-10113-2013, 2013.
- Duncan, B. N., Logan, J. A., Bey, I., Megretskiaia, I. A., Yantosca, R. M., Novelli, P. C., Jones, N. B., and Rinsland, C. P.: Global budget of 20 CO, 1988–1997: Source estimates and validation with a global model, *J. Geophys. Res.*, 112, doi:10.1029/2007JD008459, d22301, 2007.

- Ellingsen, K., Gauss, M., Van Dingenen, R., Dentener, F. J., Emberson, L., Fiore, A. M., Schultz, M. G., Stevenson, D. S., Ashmore, M. R., Atherton, C. S., Bergmann, D. J., Bey, I., Butler, T., Drevet, J., Eskes, H., Hauglustaine, D. A., Isaksen, I. S. A., Horowitz, L. W., Krol, M., Lamarque, J. F., Lawrence, M. G., van Noije, T., Pyle, J., Rast, S., Rodriguez, J., Savage, N., Strahan, S., Sudo, K., Szopa, S., and Wild, O.: Global ozone and air quality: a multi-model assessment of risks to human health and crops, *Atmos. Chem. Phys. Discuss.*, 8, 2163–2223, doi:10.5194/acpd-8-2163-2008, 2008.
- Emmons, L. K., Hauglustaine, D. A., Müller, J.-F., Carroll, M. A., Brasseur, G. P., Brunner, D., Staehelin, J., Thouret, V., and Marenco, A.: Data composites of airborne observations of tropospheric ozone and its precursors, *J. Geophys. Res.*, 105, 20497–20538, doi:10.1029/2000JD900232, 2000.
- Emmons, L. K., Arnold, S. R., Monks, S. A., Huijnen, V., Tilmes, S., Law, K. S., Thomas, J. L., Raut, J.-C., Bouarar, I., Turquety, S., Long, Y., Duncan, B., Steenrod, S., Strode, S., Flemming, J., Mao, J., Langner, J., Thompson, A. M., Tarasick, D., Apel, E. C., Blake, D. R., Cohen, R. C., Dibb, J., Diskin, G. S., Fried, A., Hall, S. R., Huey, L. G., Weinheimer, A. J., Wisthaler, A., Mikoviny, T., Nowak, J., Peischl, J., Roberts, J. M., Ryerson, T., Warneke, C., and Helmig, D.: The POLARCAT Model Intercomparison Project (POLMIP): overview and evaluation with observations, *Atmos. Chem. Phys.*, 15, 6721–6744, doi:10.5194/acp-15-6721-2015, 2015.
- Eskes, H. J. and Boersma, K. F.: Averaging kernels for DOAS total-column satellite retrievals, *Atmos. Chem. Phys.*, 3, 1285–1291, doi:10.5194/acp-3-1285-2003, 2003.
- Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N_2O_5 hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, *Geophys. Res. Lett.*, 32, doi:10.1029/2005GL022469, 109813, 2005.
- Feng, W., Chipperfield, M. P., Dhomse, S., Monge-Sanz, B. M., Yang, X., Zhang, K., and Ramonet, M.: Evaluation of cloud convection and tracer transport in a three-dimensional chemical transport model, *Atmos. Chem. Phys.*, 11, 5783–5803, doi:10.5194/acp-11-5783-2011, 2011.
- Fiore, A. M., Dentener, F. J., Wild, O., Cuvelier, C., Schultz, M. G., Hess, P., Textor, C., Schulz, M., Doherty, R. M., Horowitz, L. W., MacKenzie, I. A., Sanderson, M. G., Shindell, D. T., Stevenson, D. S., Szopa, S., Van Dingenen, R., Zeng, G., Atherton, C., Bergmann, D., Bey, I., Carmichael, G., Collins, W. J., Duncan, B. N., Faluvegi, G., Folberth, G., Gauss, M., Gong, S., Hauglustaine, D., Holloway, T., Isaksen, I. S. A., Jacob, D. J., Jonson, J. E., Kaminski, J. W., Keating, T. J., Lupu, A., Marmer, E., Montanaro, V., Park, R. J., Pitari, G., Pringle, K. J., Pyle, J. A., Schroeder, S., Vivanco, M. G., Wind, P., Wojcik, G., Wu, S., and Zuber, A.: Multimodel estimates of intercontinental source-receptor relationships for ozone pollution, *J. Geophys. Res.*, 114, doi:10.1029/2008JD010816, d04301, 2009.
- Fischer, E. V., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Millet, D. B., Mao, J., Paulot, F., Singh, H. B., Roiger, A., Ries, L., Talbot, R., Dzepina, K., and Pandey Deolal, S.: Atmospheric peroxyacetyl nitrate (PAN): a global budget and source attribution, *Atmos. Chem. Phys.*, 14, 2679–2698, doi:10.5194/acp-14-2679-2014, 2014.
- Folberth, G. A., Hauglustaine, D. A., Lathière, J., and Brocheton, F.: Interactive chemistry in the Laboratoire de Météorologie Dynamique general circulation model: model description and impact analysis of biogenic hydrocarbons on tropospheric chemistry, *Atmos. Chem. Phys.*, 6, 2273–2319, doi:10.5194/acp-6-2273-2006, 2006.
- Fortems-Cheiney, A., Chevallier, F., Pison, I., Bousquet, P., Szopa, S., Deeter, M. N., and Clerbaux, C.: Ten years of CO emissions as seen from Measurements of Pollution in the Troposphere (MOPITT), *J. Geophys. Res.*, 116, doi:10.1029/2010JD014416, d05304, 2011.
- Franco, B., Mahieu, E., Emmons, L., Tzompa-Sosa, Z., Fischer, E., Sudo, K., Bovy, B., Conway, S., Griffin, D., Hannigan, J., and Strong, K.: Evaluating ethane and methane emissions associated with the development of oil and natural gas extraction in North America, *Atmos. Chem. Phys.*, 11, p.044 010, 2016.

- Fuentes, J. D., Gu, L., Lerdau, M., Atkinson, R., Baldocchi, D., Bottenheim, J. W., Ciccioli, P., Lamb, B., Geron, C., Guenther, A., Sharkey, T. D., and Stockwell, W.: Biogenic Hydrocarbons in the Atmospheric Boundary Layer: A Review, *Bull. Amer. Meteor. Soc.*, 81, 1537–1575, doi:10.1175/1520-0477(2000)081<1537:BHITAB>2.3.CO;2, 2000.
- 30 Giannakopoulos, C., Chipperfield, M. P., Law, K. S., and Pyle, J. A.: Validation and intercomparison of wet and dry deposition schemes using ^{210}Pb in a global three-dimensional off-line chemical transport model, *J. Geophys. Res.*, 104, 23 761–23 784, doi:10.1029/1999JD900392, 1999.
- Gligorovski, S., Strekowski, R., Barbat, S., and Vione, D.: Environmental Implications of Hydroxyl Radicals ($\cdot\text{OH}$), *Chem. Rev.*, 115, 35 13 051–13 092, doi:10.1021/cr500310b, 2015.
- Granier, C., Lamarque, J., Mieville, A., Muller, J., Olivier, J., Orlando, J., Peters, J., Petron, G., Tyndall, G., and Wallens, S.: POET, <http://www.aero.jussieu.fr/projet/ACCENT/POET.php>, 2005.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, doi:10.5194/gmd-5-1471-2012, 2012.
- 5 Heard, D. E. and Pilling, M. J.: Measurement of OH and HO_2 in the Troposphere, *Chem. Rev.*, 103, 5163–5198, doi:10.1021/cr020522s, 2003.
- Holloway, M. J., Arnold, S. R., Challinor, A. J., and Emberson, L. D.: Intercontinental trans-boundary contributions to ozone-induced crop yield losses in the Northern Hemisphere, *Biogeosciences*, 9, 271–292, doi:doi:10.5194/bg-9-271-2012, 2012.
- 10 Holtslag, A. and Bolville, B.: Local versus nonlocal boundary layer diffusion in a global climate model, *J. Climate*, 6, 1825–1842, doi:[http://dx.doi.org/10.1175/1520-0442\(1993\)006<1825:LVNBLD>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(1993)006<1825:LVNBLD>2.0.CO;2), 1993.
- Hough, A.: The calculation of photolysis rates for use in global tropospheric modelling studies, *AERE Rep.*, R-13259, 1988.
- 15 Hudman, R. C., Jacob, D. J., Cooper, O. R., Evans, M. J., Heald, C. L., Park, R. J., Fehsenfeld, F., Flocke, F., Holloway, J., Hübner, G., Kita, K., Koike, M., Kondo, Y., Neuman, A., Nowak, J., Oltmans, S., Parrish, D., Roberts, J. M., and Ryerson, T.: Ozone production in transpacific Asian pollution plumes and implications for ozone air quality in California, *J. Geophys. Res.*, 109, doi:10.1029/2004JD004974, d23S10, 2004.
- Huijnen, V., Eskes, H. J., Poupkou, A., Elbern, H., Boersma, K. F., Foret, G., Sofiev, M., Valdebenito, A., Flemming, J., Stein, O., Gross, A., Robertson, L., D'Isidoro, M., Kioutsioukis, I., Friese, E., Amstrup, B., Bergstrom, R., Strunk, A., Vira, J., Zyryanov, D., Maurizi, A., Melas, D., Peuch, V.-H., and Zerefos, C.: Comparison of OMI NO_2 tropospheric columns with an ensemble of global and European regional air quality models, *Atmos. Chem. Phys.*, 10, 3273–3296, doi:10.5194/acp-10-3273-2010, 2010.
- Irie, H., Boersma, K. F., Kanaya, Y., Takashima, H., Pan, X., and Wang, Z. F.: Quantitative bias estimates for tropospheric NO_2 columns retrieved from SCIAMACHY, OMI, and GOME-2 using a common standard for East Asia, *Atmos. Meas. Tech.*, 5, 2403–2411, doi:10.5194/amt-5-2403-2012, 2012.
- 20 Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131 – 2159, doi:[http://dx.doi.org/10.1016/S1352-2310\(99\)00462-8](http://dx.doi.org/10.1016/S1352-2310(99)00462-8), 2000.
- Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A., Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the

- Troposphere from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results, *Atmospheric Chemistry and Physics*, 10, 5191–5212, doi:10.5194/acp-10-5191-2010, 2010.
- 30 Kinnison, D. E., Brasseur, G. P., Walters, S., Garcia, R. R., Marsh, D. R., Sassi, F., Harvey, V. L., Randall, C. E., Emmons, L., Lamarque, J. F., Hess, P., Orlando, J. J., Tie, X. X., Randel, W., Pan, L. L., Gettelman, A., Granier, C., Diehl, T., Niemeier, U., and Simmons, A. J.: Sensitivity of chemical tracers to meteorological parameters in the MOZART-3 chemical transport model, *J. Geophys. Res.*, 112, D20302, 2007.
- Kopacz, M., Jacob, D. J., Fisher, J. A., Logan, J. A., Zhang, L., Megretskaya, I. A., Yantosca, R. M., Singh, K., Henze, D. K., Burrows, J. P.,
35 Buchwitz, M., Khlystova, I., McMillan, W. W., Gille, J. C., Edwards, D. P., Eldering, A., Thouret, V., and Nedelev, P.: Global estimates of CO sources with high resolution by adjoint inversion of multiple satellite datasets (MOPITT, AIRS, SCIAMACHY, TES), *Atmos. Chem. Phys.*, 10, 855–876, doi:10.5194/acp-10-855-2010, 2010.
- Krol, M. and Lelieveld, J.: Can the variability in tropospheric OH be deduced from measurements of 1,1,1-trichloroethane (methyl chloroform)?, *J. Geophys. Res.*, 108, doi:10.1029/2002JD002423, 4125, 2003.
- Krol, M., van Leeuwen, P. J., and Lelieveld, J.: Global OH trend inferred from methylchloroform measurements, *J. Geophys. Res.*, 103, 10 697–10 711, doi:10.1029/98JD00459, 1998.
- 5 Krol, M. C., Lelieveld, J., Oram, D. E., Sturrock, G. A., Penkett, S. A., Brenninkmeijer, C. A. M., Gros, V., Williams, J., and Scheeren, H. A.: Continuing emissions of methyl chloroform from Europe, *Nature*, 421, 131–135, doi:doi:10.1038/nature01311, 2003.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- 10 Law, K. S. and Pyle, J. A.: Modeling trace gas budgets in the troposphere: 1. Ozone and odd nitrogen, *Journal of Geophysical Research: Atmospheres*, 98, 18 377–18 400, doi:10.1029/93JD01479, 1993.
- Law, K. S., Plantevin, P. H., Shallcross, D. E., Rogers, H. L., Pyle, J. A., Grouhel, C., Thouret, V., and Marenco, A.: Evaluation of modeled O₃ using Measurement of Ozone by Airbus In-Service Aircraft (MOZAIC) data, *Journal of Geophysical Research: Atmospheres*, 103, 15 721–25 737, doi:10.1029/98JD01482, 1998.
- 15 Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in Version 4 of the Community Land Model, *J. Adv. Model. Earth Sy.*, 3, doi:10.1029/2011MS000045, m03001, 2011.
- Lawrence, M. G., Jöckel, P., and von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmos. Chem. Phys.*, 1, 37–49, 20
20 doi:10.5194/acp-1-37-2001, 2001.
- Lelieveld, J. and Dentener, F. J.: What controls tropospheric ozone?, *J. Geophys. Res.*, 105, 3531–3551, doi:10.1029/1999JD901011, 2000.
- Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzeveld, L., Harder, H., Lawrence, M. G., Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation capacity sustained by a tropical forest, *Nature*, 452, 737–740, doi:10.1038/nature06870, 2008.
- 25 Lelieveld, J., Evans, J., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367–371, doi:10.1038/nature15371, 2015.
- Logan, J., Prather, M., Wofsy, S., and McElroy, M.: Tropospheric chemistry: A global perspective, *J. Geophys. Res.*, 86, 7210–7254, doi:10.1029/JC086iC08p07210, 1981.

- Mann, G., Carslaw, K. S., Spracklen, D. V., Ridley, D., Manktelow, P. T., Chipperfield, M. P., Pickering, S., , and Johnson, C. E.: Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model, *Geosci. Model Dev.*, 3, 519–551, 2010.
- Mannschreck, K., Gilge, S., Plass-Duelmer, C., Fricke, W., and Berresheim, H.: Assessment of the applicability of NO-NO₂-O₃ photostationary state to long-term measurements at the Hohenpeissenberg GAW Station, Germany, *Atmos. Chem. Phys.*, 4, 1265–1277, doi:10.5194/acp-4-1265-2004, 2004.
- MCM: Master Chemical Mechanism v3.1, <http://mcm.leeds.ac.uk/MCM/>, accessed: February 2004, 2004.
- Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J.-F., Matsumoto, K., Montzka, S. A., Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., and van Vuuren, D. P.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Clim. Chang.*, 109, 213–241, doi:10.1007/s10584-011-0156-z, 2011.
- Miles, G., Siddans, R., Kerridge, B., and Latter, B.: RAL Ozone Profile Algorithm Product User Guide Version 1.1, 2015a.
- Miles, G. M., Siddans, R., Kerridge, B. J., Latter, B. G., and Richards, N. A. D.: Tropospheric ozone and ozone profiles retrieved from GOME-2 and their validation, *Atmos. Meas. Tech.*, 8, 385–398, doi:10.5194/amt-8-385-2015, 2015b.
- Mogili, P. K., Kleiber, P. D., Young, M. A., and Grassian, V. H.: N₂O₅ hydrolysis on the components of mineral dust and sea salt aerosol: Comparison study in an environmental aerosol reaction chamber, *Atmos. Environ.*, 40, 7401–7408, doi:<http://dx.doi.org/10.1016/j.atmosenv.2006.06.048>, 2006.
- Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G. E., Stevenson, D. S., Tarasova, O., Thouret, V., von Schneidemesser, E., Sommariva, R., Wild, O., and Williams, M. L.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmos. Chem. Phys.*, 15, 8889–8973, doi:10.5194/acp-15-8889-2015, 2015a.
- Monks, S. A., Arnold, S. R., and Chipperfield, M. P.: Evidence for El Niño-Southern Oscillation (ENSO) influence on Arctic CO interannual variability through biomass burning emissions, *Geophys. Res. Lett.*, 39, doi:10.1029/2012GL052512, 2012.
- Monks, S. A., Arnold, S. R., Emmons, L. K., Law, K. S., Turquety, S., Duncan, B. N., Flemming, J., Huijnen, V., Tilmes, S., Langner, J., Mao, J., Long, Y., Thomas, J. L., Steenrod, S. D., Raut, J. C., Wilson, C., Chipperfield, M. P., Diskin, G. S., Weinheimer, A., Schlager, H., and Ancellet, G.: Multi-model study of chemical and physical controls on transport of anthropogenic and biomass burning pollution to the Arctic, *Atmos. Chem. Phys.*, 15, 3575–3603, doi:10.5194/acp-15-3575-2015, 2015b.
- Montzka, S. A., Krol, M., Dlugokencky, E., Hall, B., Jöckel, P., and Lelieveld, J.: Small Interannual Variability of Global Atmospheric Hydroxyl, *Science*, 331, 67–69, doi:10.1126/science.1197640, 2011.
- Moore, D. P. and Remedios, J. J.: Seasonality of Peroxyacetyl nitrate (PAN) in the upper troposphere and lower stratosphere using the MIPAS-E instrument, *Atmos. Chem. Phys.*, 10, 6117–6128, doi:10.5194/acp-10-6117-2010, 2010.
- Naik, V., Voulgarakis, A., Fiore, A. M., Horowitz, L. W., Lamarque, J.-F., Lin, M., Prather, M. J., Young, P. J., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R., Eyring, V., Faluvegi, G., Folberth, G. A., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., van Noije, T. P. C., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R., Shindell, D. T., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., and Zeng, G.: Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 5277–5298, doi:10.5194/acp-13-5277-2013, 2013.
- Novelli, P., Masarie, K., and Lang, P.: Distributions and recent changes of carbon monoxide in the lower troposphere, *J. Geophys. Res.*, 103, 19 015–19 033, 1998.

- Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzeck, E., Lawrence, P. J., Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddema, J., Heald, C. L., Hoffman, F., Lamarque, J.-F., Mahowald, N., Niu, G.-Y., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stockli, R., Wang, A., Yang, Z.-L., Zeng, X., and Zeng, X.: Technical Description of version 4.0 of the Community Land Model(CLM), Ncar technical note ncar/tn-478+str, NationalCenter for Atmospheric Research, Boulder, CO, 2010.
- Oltmans, S. J. and Levy, H.: Surface ozone measurements from a global network, *Atmos. Environ.*, 28, 9 – 24, doi:[http://dx.doi.org/10.1016/1352-2310\(94\)90019-1](http://dx.doi.org/10.1016/1352-2310(94)90019-1), 1994.
- Pandey Deolal, S., Henne, S., Ries, L., Gilge, S., Weers, U., Steinbacher, M., Staehelin, J., and Peter, T.: Analysis of elevated springtime levels of Peroxyacetyl nitrate (PAN) at the high Alpine research sites Jungfraujoch and Zugspitze, *Atmos. Chem. Phys.*, 14, 12 553–12 571, doi:10.5194/acp-14-12553-2014, 2014.
- Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Belikov, D., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield, M. P., Corbin, K., Fortems-Cheiney, A., Fraser, A., Gloor, E., Hess, P., Ito, A., Kawa, S. R., Law, R. M., Loh, Z., Maksyutov, S., Meng, L., Palmer, P. I., Prinn, R. G., Rigby, M., Saito, R., and Wilson, C.: TransCom model simulations of CH₄ and related species: linking transport, surface flux and chemical loss with CH₄ variability in the troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 11, 12 813–12 837, doi:10.5194/acp-11-12813-2011, 2011.
- Patra, P. K., Krol, M. C., Montzka, S. A., Arnold, T., Atlas, E. L., Lintner, B. R., Stephens, B. B., Xiang, B., Elkins, J. W., Fraser, P. J., Ghosh, A., Hintsa, E. J., Hurst, D. F., Ishijima, K., Krummel, P. B., Miller, B. R., Miyazaki, K., Moore, F. L., Mühle, J., O'Doherty, S., Prinn, R. G., Steele, L. P., Takigawa, M., Wang, H. J., Weiss, R. F., Wofsy, S. C., and Young, D.: Observational evidence for interhemispheric hydroxyl-radical parity, *Nature*, 513, 219–223, doi:10.1038/nature13721, 2014.
- Pétron, G., Granier, C., Khattatov, B., Yudin, V., Lamarque, J.-F., Emmons, L., Gille, J., and Edwards, D. P.: Monthly CO surface sources inventory based on the 2000–2001 MOPITT satellite data, *Geophys. Res. Lett.*, 31, doi:10.1029/2004GL020560, 121107, 2004.
- Plass-Dülmer, C., Michl, K., Ruf, R., and Berresheim, H.: C2–C8 Hydrocarbon measurement and quality control procedures at the Global Atmosphere Watch Observatory Hohenpeissenberg, *J. Chromatogr. A*, 953, 175–197, doi:[http://dx.doi.org/10.1016/S0021-9673\(02\)00128-0](http://dx.doi.org/10.1016/S0021-9673(02)00128-0), 2002.
- Pope, R. J., Chipperfield, M. P., Savage, N. H., Ordóñez, C., Neal, L. S., Lee, L. A., Dhomse, S. S., Richards, N. A. D., and Keslake, T. D.: Evaluation of a regional air quality model using satellite column NO₂: treatment of observation errors and model boundary conditions and emissions, *Atmos. Chem. Phys.*, 15, 5611–5626, doi:10.5194/acp-15-5611-2015, 2015.
- Pope, R. J., Richards, N. A. D., Chipperfield, M. P., Moore, D. P., Monks, S. A., Arnold, S. R., Glatthor, N., Kiefer, M., Breider, T. J., Harrison, J. J., Remedios, J. J., Warneke, C., Roberts, J. M., Diskin, G. S., Huey, L. G., Wisthaler, A., Apel, E. C., Bernath, P. F., and Feng, W.: Intercomparison and evaluation of satellite peroxyacetyl nitrate observations in the upper troposphere - lower stratosphere, *Atmos. Chem. Phys. Discuss.*, 2016, 1–44, doi:10.5194/acp-2016-386, 2016.
- Pöschl, U., von Kuhlmann, R., Poisson, N., and Crutzen, P. J.: Development and intercomparison of condensed isoprene oxidation mechanisms for global atmospheric modeling, *J. Atmos. Chem.*, 37, 29–52, 2000.
- Prather, M.: Numerical advection by conservation of second-order moments., *J. Geophys. Res.*, 91, 6671–6681, 1986.
- Prinn, R. G., Huang, J., Weiss, R. F., Cunnold, D. M., Fraser, P. J., Simmonds, P. G., McCulloch, A., Harth, C., Salameh, P., O'Doherty, S., Wang, R. H. J., Porter, L., and Miller, B. R.: Evidence for Substantial Variations of Atmospheric Hydroxyl Radicals in the Past Two Decades, *Science*, 292, 1882–1888, doi:10.1126/science.1058673, 2001.

- 30 Ravishankara, A. R., Dunlea, E. J., Blitz, M. A., Dillon, T. J., Heard, D. E., Pilling, M. J., Strekowski, R. S., Nicovich, J. M., and Wine, P. H.: Redetermination of the rate coefficient for the reaction of O(¹D) with N₂, *Geophys. Res. Lett.*, 29, 35–1–35–4, doi:10.1029/2001GL014850, 2002.
- Read, K., Mahajan, A., Carpenter, L., Evans, M., Faria, B., Heard, D., Hopkins, J., Lee, J., Moller, S., Lewis, A., Mendes, L., McQuaid, J., Oetjen, H., Saiz-Lopez, A., Pilling, M., and Plane, J.: Extensive halogen-mediated ozone destruction over the tropical Atlantic Ocean, *Nature*, 453, 1232–1235, doi:10.1038/nature07035, 2008.
- Richards, N. A. D., Arnold, S. R., Chipperfield, M. P., Miles, G., Rap, A., Siddans, R., Monks, S. A., and Hollaway, M. J.: The 35 Mediterranean summertime ozone maximum: global emission sensitivities and radiative impacts, *Atmos. Chem. Phys.*, 13, 2331–2345, doi:doi:10.5194/acp-13-2331-2013, 2013.
- Rodgers, C. D.: Retrieval of atmospheric temperature and composition from remote measurements of thermal radiation, *Rev. Geophys.*, 14, 609–624, doi:10.1029/RG014i004p00609, 1976.
- Rudolph, J., Khedim, A., and Wagenbach, D.: The seasonal variation of light nonmethane hydrocarbons in the Antarctic troposphere, *J. Geophys. Res.*, 94, 13 039–13 044, doi:10.1029/JD094iD10p13039, 1989.
- Sander, S.P., e. a.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies Evaluation Number 15. JPL Publication 06-2, 5 Tech. rep., Jet Propulsion Laboratory, Pasadena, USA, 2006.
- Sander, S.P., e. a.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies Evaluation Number 17, JPL Publication 10-6, Tech. rep., Jet Propulsion Laboratory, Pasadena, USA, 2011.
- Sanderson, M. G., Dentener, F. J., Fiore, A. M., Cuvelier, C., Keating, T. J., Zuber, A., Atherton, C. S., Bergmann, D. J., Diehl, T., Doherty, R. M., Duncan, B. N., Hess, P., Horowitz, L. W., Jacob, D. J., Jonson, J.-E., Kaminski, J. W., Lupu, A., MacKenzie, I. A., 10 Mancini, E., Marmer, E., Park, R., Pitari, G., Prather, M. J., Pringle, K. J., Schroeder, S., Schultz, M. G., Shindell, D. T., Szopa, S., Wild, O., and Wind, P.: A multi-model study of the hemispheric transport and deposition of oxidised nitrogen, *Geophys. Res. Lett.*, 35, doi:10.1029/2008GL035389, 117815, 2008.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM 15 v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmospheric Chemistry and Physics*, 3, 161–180, doi:10.5194/acp-3-161-2003, <http://www.atmos-chem-phys.net/3/161/2003/>, 2003.
- Scott, C. E., Monks, S. A., Spracklen, D. V., Arnold, S. R., Forster, P. M., Rap, A., Äijälä, M., Artaxo, Carslaw, K. S., Chipperfield, M. P., Ehn, M., Gilardoni, Heikkinen, L., Kulmala, M., Petäjä, T., Reddington, C. L. S., Rizzo, L., Swielicki, E., Vignati, E., and Wilson, C.: submitted to *Nature Commun.*, 2016.
- Shindell, D. T., Faluvegi, G., Stevenson, D. S., Krol, M. C., Emmons, L. K., Lamarque, J.-F., Pétron, G., Dentener, F. J., Ellingsen, K., 20 Schultz, M. G., Wild, O., Amann, M., Atherton, C. S., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Lawrence, M. G., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Strahan, S. E., Sudo, K., Szopa, S., Unger, N., van Noije, T. P. C., and Zeng, G.: Multimodel simulations of carbon monoxide: Comparison with observations and projected near-future changes, *J. Geophys. Res.*, 111, doi:10.1029/2006JD007100, d19306, 2006.
- 25 Shindell, D. T., Chin, M., Dentener, F., Doherty, R. M., Faluvegi, G., Fiore, A. M., Hess, P., Koch, D. M., MacKenzie, I. A., Sanderson, M. G., Schultz, M. G., Schulz, M., Stevenson, D. S., Teich, H., Textor, C., Wild, O., Bergmann, D. J., Bey, I., Bian, H., Cuvelier, C., Duncan, B. N., Folberth, G., Horowitz, L. W., Jonson, J., Kaminski, J. W., Marmer, E., Park, R., Pringle, K. J., Schroeder, S., Szopa, S.,

- Takemura, T., Zeng, G., Keating, T. J., and Zuber, A.: A multi-model assessment of pollution transport to the Arctic, *Atmos. Chem. Phys.*, 8, 5353–5372, doi:10.5194/acp-8-5353-2008, 2008.
- 30 Sofen, E. D., Bowdalo, D., Evans, M. J., Apadula, F., Bonasoni, P., Cupeiro, M., Ellul, R., Galbally, I. E., Grgzdienė, R., Luppo, S., Mimouni, M., Nahas, A. C., Saliba, M., and Tørseth, K.: Gridded global surface ozone metrics for atmospheric chemistry model evaluation, *Earth System Science Data*, 8, 41–59, doi:10.5194/essd-8-41-2016, 2016.
- 35 Spivakovskiy, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B. A., Horowitz, L. W., Fusco, A. C., Brenninkmeijer, C. A. M., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Three-dimensional climatological distribution of tropospheric OH: Update and evaluation, *J. Geophys. Res.*, 105, 8931–8980, 2000.
- Squire, O. J., Archibald, A. T., Griffiths, P. T., Jenkin, M. E., Smith, D., and Pyle, J. A.: Influence of isoprene chemical mechanism on modelled changes in tropospheric ozone due to climate and land use over the 21st century, *Atmos. Chem. Phys.*, 15, 5123–5143, doi:10.5194/acp-15-5123-2015, 2015.
- 40 Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F., Lawrence, M. G., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., 5 Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.*, 111, doi:10.1029/2005JD006338, d08301, 2006.
- 50 Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., eds.: IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 10 Stockwell, D., Giannakopoulos, C., Plantevin, P.-H., Carver, G., Chipperfield, M., Law, K., Pyle, J., Shallcross, D., and Wang, K.-Y.: Modelling NO_x from lightning and its impact on global chemical fields, *Atmos. Environ.*, 33, 4477 – 4493, doi:[http://dx.doi.org/10.1016/S1352-2310\(99\)00190-9](http://dx.doi.org/10.1016/S1352-2310(99)00190-9), 1999.
- 15 Strode, S. A., Duncan, B. N., Yegorova, E. A., Kouatchou, J., Ziemke, J. R., and Douglass, A. R.: Implications of carbon monoxide bias for methane lifetime and atmospheric composition in chemistry climate models, *Atmos. Chem. Phys.*, 15, 11 789–11 805, doi:10.5194/acp-15-11789-2015, 2015.
- Sukhodolov, T., Rozanov, E., Ball, W. T., Bais, A., Tourpali, K., Shapiro, A. I., Telford, P., Smyshlyaev, S., Fomin, B., Sander, R., Bossay, S., Bekki, S., Marchand, M., Chipperfield, M. P., Dhomse, S., Haigh, J. D., Peter, T., and Schmutz, W.: Evaluation of simulated photolysis rates and their response to solar irradiance variability, *J. Geophys. Res.*, 121, 6066–6084, doi:10.1002/2015JD024277, 2015JD024277, 2016.
- 20 Tiedtke, M.: A comprehensive mass flux scheme for cumulus parameterization in large-scale models, *Mon. Weather Rev.*, 117, 1779–1800, 1989.
- Tilmes, S., Lamarque, J.-F., Emmons, L. K., Conley, A., Schultz, M. G., Saunois, M., Thouret, V., Thompson, A. M., Oltmans, S. J., Johnson, B., and Tarasick, D.: Technical Note: Ozonesonde climatology between 1995 and 2011: description, evaluation and applications, *Atmos. Chem. Phys.*, 12, 7475–7497, doi:10.5194/acp-12-7475-2012, 2012.
- 25 Tilmes, S., Lamarque, J.-F., Emmons, L. K., Kinnison, D. E., Ma, P.-L., Liu, X., Ghan, S., Bardeen, C., Arnold, S., Deeter, M., Vitt, F., Ryerson, T., Elkins, J. W., Moore, F., Spackman, J. R., and Val Martin, M.: Description and evaluation of tropospheric chemistry and aerosols in the Community Earth System Model (CESM1.2), *Geosci. Model Dev.*, 8, 1395–1426, doi:10.5194/gmd-8-1395-2015, 2015.

- Tilmes, S., Lamarque, J.-F., Emmons, L. K., Kinnison, D. E., Marsh, D., Garcia, R. R., Smith, A. K., Neely, R. R., Conley, A., Vitt, F., Val Martin, M., Tanimoto, H., Simpson, I., Blake, D. R., and Blake, N.: Representation of the Community Earth System Model (CESM1) CAM4-chem within the Chemistry-Climate Model Initiative (CCMI), *Geosci. Model Dev.*, 9, 1853–1890, doi:doi:10.5194/gmd-9-1853-2016, 2016.
- Tyndall, G. S., Cox, R. A., Granier, C., Lesclaux, R., Moortgat, G. K., Pilling, M. J., Ravishankara, A. R., and Wallington, T. J.: Atmospheric chemistry of small organic peroxy radicals, *J. Geophys. Res.*, 106, 12 157–12 182, doi:10.1029/2000JD900746, 2001.
- ValMartin, M., Heald, C. L., and Arnold, S. R.: Coupling dry deposition to vegetation phenology in the Community Earth System Model: Implications for the simulation of surface O₃, *Geophys. Res. Lett.*, 41, 2988–2996, doi:10.1002/2014GL059651, 2014.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11 707–11 735, doi:10.5194/acp-10-11707-2010, 2010.
- von Kuhlmann, R.: Photochemistry of Tropospheric Ozone, its Precursors and the Hydroxyl Radical: A 3D-Modeling Study Considering Non-Methane Hydrocarbons, Ph.D. thesis, Johannes Gutenberg-Universitat Mainz, Mainz, Germany, 2001.
- Voulgarakis, A., Naik, V., Lamarque, J.-F., Shindell, D. T., Young, P. J., Prather, M. J., Wild, O., Field, R. D., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Folberth, G. A., Horowitz, L. W., Josse, B., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Stevenson, D. S., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Analysis of present day and future OH and methane lifetime in the ACCMIP simulations, *Atmos. Chem. Phys.*, 13, 2563–2587, doi:10.5194/acp-13-2563-2013, 2013.
- Wang, J. S., McElroy, M. B., Logan, J. A., Palmer, P. I., Chameides, W. L., Wang, Y., and Megretskaya, I. A.: A quantitative assessment of uncertainties affecting estimates of global mean OH derived from methyl chloroform observations, *J. Geophys. Res.*, 113, doi:10.1029/2007JD008496, d12302, 2008.
- Wang, K.-Y. and Shallcross, D.: Modelling terrestrial biogenic isoprene fluxes and their potential impact on global chemical species using a coupled LSM–CTM model, *Atmospheric Environment*, 34, 2909 – 2925, doi:[http://dx.doi.org/10.1016/S1352-2310\(99\)00525-7](http://dx.doi.org/10.1016/S1352-2310(99)00525-7), 2000.
- Watanabe, F., Uchino, O., Joo, Y., Aono, M., Higashijima, K., Hirano, Y., Tsuboi, K., and Suda, K.: Interannual variation of growth rate of atmospheric carbon dioxide concentration observed at the JMA's three monitoring stations: Large increase in concentration of atmospheric carbon dioxide in 1998, *J. Meteorol. Soc. Jpn.*, 78, 673—682, 2000.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, doi:10.5194/gmd-4-625-2011, 2011.
- Wilson, C., Chipperfield, M. P., Gloor, M., and Chevallier, F.: Development of a variational flux inversion system (INVICAT v1.0) using the TOMCAT chemical transport model, *Geosci. Model Dev.*, 7, 2485–2500, doi:doi:10.5194/gmd-7-2485-2014, 2014.
- Wilson, C., Gloor, M., Gatti, L. V., Miller, J. B., Monks, S. A., McNorton, J., Bloom, A. A., Basso, L. S., and Chipperfield, M. P.: Contribution of regional sources to atmospheric methane over the Amazon Basin in 2010 and 2011, *Global Biogeochem. Cycles*, doi:10.1002/2015GB005300, 2016.
- Young, P.: The influence of biogenic isoprene emissions on atmospheric chemistry: A model study for present and future atmospheres, Ph.D. thesis, University of Cambridge, 2007.
- Young, P. J., Archibald, A. T., Bowman, K. W., Lamarque, J.-F., Naik, V., Stevenson, D. S., Tilmes, S., Voulgarakis, A., Wild, O., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Horowitz, L. W., Josse, B.,

- Lee, Y. H., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R. B., Shindell, D. T., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Pre-industrial to end 21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 2063–2090, doi:10.5194/acp-13-2063-2013, 2013.
- 1070 Yu, S., Mathur, R., Sarwar, G., Kang, D., Tong, D., Pouliot, G., and Pleim, J.: Eta-CMAQ air quality forecasts for O₃ and related species using three different photochemical mechanisms (CB4, CB05, SAPRC-99): comparisons with measurements during the 2004 ICARTT study, *Atmos. Chem. Phys.*, 10, 3001–3025, doi:10.5194/acp-10-3001-2010, 2010.
- Zellweger, C., Forrer, J., Hofer, P., Nyeki, S., Schwarzenbach, B., Weingartner, E., Ammann, M., and Baltensperger, U.: Partitioning of reactive nitrogen (NO_y) and dependence on meteorological conditions in the lower free troposphere, *Atmos. Chem. Phys.*, 3, 779–796, doi:10.5194/acp-3-779-2003, 2003.
- 1075 Zhang, K., Wan, H., Zhang, M., and Wang, B.: Evaluation of the atmospheric transport in a GCM using radon measurements: sensitivity to cumulus convection parameterization, *Atmos. Chem. Phys.*, 8, 2811–2832, doi:10.5194/acp-8-2811-2008, 2008.
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, doi:10.5194/acp-9-5131-2009, 2009.
- 1080 Ziemke, J. R., Chandra, S., Labow, G. J., Bhartia, P. K., Froidevaux, L., and Witte, J. C.: A global climatology of tropospheric and stratospheric ozone derived from Aura OMI and MLS measurements, *Atmos. Chem. Phys.*, 11, 9237–9251, doi:10.5194/acp-11-9237-2011, 2011.




Figure 1. Global annual mean pressure levels from the TOMCAT model.

As Figure ??, but for nitric acid (HNO_3 ; pptv).




Figure 2. Location of WDCGG surface observatories and ozonesonde release sites used to evaluate the model for a) CO, b) O₃, and c) PAN, NO_x and VOCs.




Figure 3. Concentrations of TOMCAT simulated annual surface mean and annual zonal mean CO (a,b), O₃ (c,d) and OH (e,f) ~~from TOMCAT simulation RUN~~.




Figure 4. Regional [annual](#) mean OH concentrations ($\times 10^6$ molecules/cm 3) split into subsections as recommended by Lawrence et al. (2001). a) OH estimated from methyl chloroform observations from Spivakovsky et al. (2000), b) ACCMIP multi-model mean simulated OH concentrations from Naik et al. (2013) and [c](#)) TOMCAT-simulated OH concentrations for [e\) RUN](#) and [d\) RUN](#). The air-mass-weighted global mean tropospheric OH (OH_{gm}) is indicated above each plot for panels b [-d](#) and [c](#). In TOMCAT, the troposphere was defined as the area below a climatological tropopause ($p = 300 - 215(\cos(lat))^2$) (as discussed in Lawrence et al. (2001)) and for ACCMIP it was defined as below 200 hPa. The colours in [panel c](#) [-d](#) are scaled according to the difference from [panel a](#), with the darkest blue representing the largest negative differences and the darkest red representing the largest positive differences.




Figure 5. April and October 2008 monthly mean 500 hPa CO concentrations (ppbv) observed by MOPITT (a,c) and simulated by TOMCAT (b,d) during April and October 2008. The zonal mean concentrations at 500 hPa are also shown (e,f; data only shown when there is 25% coverage in a given latitude band). MOPITT Averaging kernels have been applied to the TOMCAT fields.




Figure 6. Observed (blue line) and simulated (red line) monthly mean CO (ppbv) at a range of WDCGG (blue line) several surface observatory sites located throughout the globe. The observations (blue line) are shown as an average arranged by latitude from north to south, with Pearson correlation (black solid line) and as minimum and maximum concentrations normalised mean bias (black dashed lines) of all available data between 2000 and 2008. The sites simulated monthly mean data are arranged by latitude from north to south printed on each panel.




Figure 7. Median O_3 concentrations (ppbv) taken from the Tilmes et al. (2012) ozonesonde climatology compared to [TOMCAT simulated](#) concentrations [from RUN and RUN](#). The data is [average averaged](#) over three latitude ranges (left to right) and three pressure level ranges (top to bottom), where the error bars show the 25th and 75th percentiles of the observed concentrations and Nstat gives the number of sonde release sites located within each latitude range.




Figure 8. Observed Monthly mean 2008 observed and simulated O_3 (ppbv) at several WDCGG surface observatories sites located throughout the globe. The observations panels are shown as an average arranged by latitude from north to south, with Pearson correlation (black solid line) and as minimum and maximum concentrations normalised mean bias (black dashed lines NMB) of all available data between 2000 the observed and 2008. The sites simulated monthly mean data are arranged by latitude from north to south printed on each panel.




Figure 9. Scatter plots of seasonal mean observed and simulated [2008](#) O₃ concentrations (ppbv) at CASTNET EPA monitoring stations located in North America. [The observations are a mean of 2000 to 2008 available data.](#)




Figure 10. GOME-2 sub-column O₃ (0-6 km, DU) on the TOMCAT $2.8^\circ \times 2.8^\circ$ grid for a) June-July-August 2008 (JJA) and b) December-January-February 2008 (DJF). c) and d) show the difference in concentrations between TOMCAT [RUN](#) and GOME-2. The green polygons are where the mean bias (MB) is greater than the satellite error.




Figure 11. Maps of simulated seasonal ~~Monthly~~ mean concentrations of propene ~~2008~~ observed and simulated VOCs (C_3H_6 ; pptv) averaged over two ~~at the European high~~ altitude ~~bands~~ observatory, Hohenpeissenberg. Pearson correlation ($2-4\text{ km}$) and $5-8\text{ km}$ normalised mean bias (NMB), with overlaid circles coloured according to concentrations from ~~between the aircraft climatology of~~ ~~Emmons et al. (2000)~~ observed and simulated monthly mean data are printed on each panel.




Figure 12. As Figure ?? Comparison of simulated and observed concentrations of CO, but for propane (O_3 , C_2H_6 , C_3H_8 ; pptv), NO_2 , PAN and HNO_3 for the ARCTAS July 2008 flights (7 flights).

As Figure ??, but for ethane (C_2H_6 ; pptv).

Observed and simulated VOCs (pptv) at the European high altitude observatory, Hohenpeissenberg. The observations are shown as an average (black solid line) and as minimum and maximum concentrations (black dashed lines) of all available data between 2000 and 2008.




Figure 13. Observed Monthly mean 2008 observed and simulated NO_2 (ppbv) at several European surface observatories/sites located in Europe. The observations are shown as an average Pearson correlation (black solid line) and as minimum and maximum concentrations normalised mean bias (black dashed lines NMB) of all available data between 2000 the observed and 2008 simulated monthly mean data are printed on each panel.




Figure 14. OMI₂₀₀₈ tropospheric column NO₂ column ($\times 10^{15}$ molecules cm^{-2}) from OMI on the TOMCAT model $2.8^\circ \times 2.8^\circ$ grid for a) June-July-August (2008) and b) January-February-December (2008) December-January-February, e) and d) are along with the TOMCAT OMI tropospheric column NO₂ mean bias (MB) for the same periods (c,d). The green polygons are where the $|\text{MB}| >$ satellite error.

Maps of simulated seasonal mean concentrations of peroxyacetyl nitrate (PAN; pptv) averaged over two altitude bands (2–4 km and 5–8 km), with overlaid circles coloured according to concentrations from the aircraft climatology of Emmons et al. (2000).




Figure 15. Observed and simulated PAN (pptv) at two European high altitude observatories. The observations are shown as averages (black solid line) and as minimum and maximum concentrations (black dashed lines) of all available data between 2000 and 2008.

Table 1. TOMCAT annual global emissions [used in the model simulations presented in this paper](#) (Tg(species)/year).

Species	Anthropogenic	Fires	Biogenic	Ocean	Soil	Total
CO	331.62	76.57	20.01		1023.47	
Ethene	6.81	2.84	16.70	1.40		27.75
Ethane	6.34	1.67	0.14	0.98		9.14
Propene	3.04	1.57	6.10	1.52		12.23
Propane	5.68	0.38	0.02	1.30		7.37
Toluene	25.34	10.66	0.26			36.26
Butane	12.38	0.60				12.98
Formaldehyde	2.99	4.13	4.03			11.15
Acetone	0.54	1.86	28.58			30.98
Acetaldehyde	2.00	4.55	11.20			17.75
Methanol	0.93	5.38	159.87			166.18
Isoprene	0.80	525.84				526.64
Monoterpenes	0.28	97.10				97.37
NO _x	107.73	19.41			16.31	143.46

Table 2: Chemical species treated in the tropospheric chemistry scheme of the TOMCAT CTM. If the species are emitted, dry deposited or wet deposited a Y is in the relevant column. The family column indicates which short-lived species are grouped together for advection and chemistry. TOMCAT abbreviations: Me=CH₃, Et=C₂H₅, Pr=C₃H₇, MACR=lumped species (methacrolein, methyl vinyl ketone and other C4 carbonyls), HACET=hydroxyacetone, MGLY=methylglyoxal, NALD=nitrooxy acetaldehyde, TERP=generic terpene compound, AROM=generic aromatic compound, MEK=Methyl ethyl ketone, Prpe=C₃H₇O, ONIT=organic nitrate, S=stratospheric tracer (TOMCAT species 39–43).

	<u>TOMCAT</u> Species	Family	Dry Deposited?	Wet Deposited?	Emitted?
1	O(³ P)	Ox			
2	O(¹ D)	Ox			
3	O ₃	Ox	Y		
4	NO	NOx	Y		
5	NO ₃	NOx	Y	Y	
6	NO ₂	NOx	Y		Y
7	N ₂ O ₅		Y	Y	
8	HO ₂ NO ₂		Y	Y	
9	HONO ₂		Y	Y	
10	OH				
11	HO ₂			Y	
12	H ₂ O ₂		Y	Y	
13	CH ₄				Y
14	CO		Y		Y
15	HCHO		Y	Y	Y
16	MeOO			Y	
17	H ₂ O				
18	MeOOH		Y	Y	
19	HONO		Y	Y	
20	C ₂ H ₆				Y
21	EtOO				
22	EtOOH		Y	Y	
23	MeCHO		Y		Y
24	MeCO ₃				
25	PAN		Y		
26	C ₃ H ₈				Y
27	n-PrOO				
28	i-PrOO				
29	n-PrOOH		Y	Y	
30	i-PrOOH		Y	Y	
31	EtCHO		Y		
32	EtCO ₃				
33	Me ₂ CO		Y		Y
34	MeCOCH ₂ OO				
35	MeCOCH ₂ OOH		Y	Y	

Continued on next page

Table 2 – continued from previous page

	TOMCAT Species	Family	Dry Deposited?	Wet Deposited?	Emitted?
36	PPAN		Y		
37	MeONO ₂				
38	O(³ P)S	Sx			
39	O(¹ D)S	Sx			
40	O ₃ S	Sx	Y		
41	NOXS		Y		
42	HNO ₃ S		Y	Y	
43	NOYS		Y	Y	
44	C ₅ H ₈				Y
45	C ₁₀ H ₁₆				Y
46	TERPOOH		Y	Y	
47	ISO ₂				
48	ISOOH		Y	Y	
49	ISON		Y	Y	
50	MACR		Y		
51	MACRO ₂				
52	MACROOH		Y	Y	
53	MPAN		Y		
54	HACET		Y	Y	
55	MGLY		Y	Y	
56	NALD		Y		
57	HCOOH		Y	Y	
58	MeCO ₃ H		Y	Y	
59	MeCO ₂ H		Y	Y	
60	MeOH		Y	Y	Y
61	TERPO ₂				
62	C ₂ H ₄				Y
63	C ₂ H ₂				Y
64	C ₄ H ₁₀				Y
65	C ₃ H ₆				Y
66	AROM				Y
67	MEK				
68	MeCOCOME		Y	Y	
69	BtOO				
70	PrpeOO				
71	AROMO ₂				
72	MEKOO				
73	BtOOH		Y	Y	
74	PrpeOOH		Y	Y	
75	AROMOOH		Y	Y	

Continued on next page

Table 2 – continued from previous page

	TOMCAT Species	Family	Dry Deposited?	Wet Deposited?	Emitted?
76	MEKOOH		Y	Y	
77	ONIT				
78	EtCO ₃ H				
79	EtCO ₂ H				

Table 3: TOMCAT gas-phase bimolecular reactions. T is the model grid-box temperature in [kelvins](#)[kelvin](#). Reaction rate references 1: Atkinson et al. (a), 2: Atkinson et al. (b), 3: Atkinson et al. (c), 4: MCM (2004), 5: Tyndall et al. (2001), 6: Ravishankara et al. (2002), 7: Pöschl et al. (2000), 9: Kinnison et al. (2007), 10: Folberth et al. (2006).

	Reactants	Products	k	Reference
1	$\text{HO}_2 + \text{NO}$	$\rightarrow \text{OH} + \text{NO}_2$	$3.60 \times 10^{-12} \exp\left(\frac{270}{T}\right)$	2
2	$\text{HO}_2 + \text{NO}_3$	$\rightarrow \text{OH} + \text{NO}_2$	4.00×10^{-12}	2
3	$\text{HO}_2 + \text{O}_3$	$\rightarrow \text{OH} + \text{O}_2$	$2.03 \times 10^{-16} \left(\frac{T}{300}\right)^{4.57} \exp\left(\frac{693}{T}\right)$	2
4	$\text{HO}_2 + \text{HO}_2$	$\rightarrow \text{H}_2\text{O}_2$	$2.20 \times 10^{-13} \exp\left(\frac{600}{T}\right)$	2
5	$\text{HO}_2 + \text{MeOO}$	$\rightarrow \text{MeOOH}$	$3.80 \times 10^{-13} \exp\left(\frac{780}{T}\right)$	2
6	$\text{HO}_2 + \text{MeOO}$	$\rightarrow \text{HCHO}$	$3.80 \times 10^{-13} \exp\left(\frac{780}{T}\right)$	2
7	$\text{HO}_2 + \text{EtOO}$	$\rightarrow \text{EtOOH}$	$3.80 \times 10^{-13} \exp\left(\frac{900}{T}\right)$	2
8	$\text{HO}_2 + \text{MeCO}_3$	$\rightarrow \text{MeCO}_3\text{H}$	$2.08 \times 10^{-13} \exp\left(\frac{980}{T}\right)$	2
9	$\text{HO}_2 + \text{MeCO}_3$	$\rightarrow \text{MeCO}_2\text{H} + \text{O}_3$	$1.04 \times 10^{-13} \exp\left(\frac{980}{T}\right)$	2
10	$\text{HO}_2 + \text{MeCO}_3$	$\rightarrow \text{OH} + \text{MeOO}$	$2.08 \times 10^{-13} \exp\left(\frac{980}{T}\right)$	2
11	$\text{HO}_2 + \text{n-PrOO}$	$\rightarrow \text{n-PrOOH}$	$1.51 \times 10^{-13} \exp\left(\frac{1300}{T}\right)$	4
12	$\text{HO}_2 + \text{i-PrOO}$	$\rightarrow \text{i-PrOOH}$	$1.51 \times 10^{-13} \exp\left(\frac{1300}{T}\right)$	4
13	$\text{HO}_2 + \text{EtCO}_3$	$\rightarrow \text{O}_2 + \text{EtCO}_3\text{H}$	$3.05 \times 10^{-13} \exp\left(\frac{1040}{T}\right)$	4
14	$\text{HO}_2 + \text{EtCO}_3$	$\rightarrow \text{O}_3 + \text{EtCO}_2\text{H}$	$1.25 \times 10^{-13} \exp\left(\frac{1040}{T}\right)$	4
15	$\text{HO}_2 + \text{MeCOCH}_2\text{OO}$	$\rightarrow \text{MeCOCH}_2\text{OOH}$	$1.36 \times 10^{-13} \exp\left(\frac{1250}{T}\right)$	4
16	$\text{MeOO} + \text{NO}$	$\rightarrow \text{HO}_2 + \text{HCHO} + \text{NO}_2$	$2.95 \times 10^{-12} \exp\left(\frac{285}{T}\right)$	2
17	$\text{MeOO} + \text{NO}$	$\rightarrow \text{MeONO}_2$	$2.95 \times 10^{-15} \exp\left(\frac{285}{T}\right)$	2
18	$\text{MeOO} + \text{NO}_3$	$\rightarrow \text{HO}_2 + \text{HCHO} + \text{NO}_2$	1.30×10^{-12}	2
19	$\text{MeOO} + \text{MeOO}$	$\rightarrow \text{MeOH} + \text{HCHO}$	$1.03 \times 10^{-13} \exp\left(\frac{365}{T}\right)$	4
20	$\text{MeOO} + \text{MeOO}$	$\rightarrow \text{HO}_2 + \text{HO}_2 + \text{HCHO} + \text{HCHO}$	$1.03 \times 10^{-13} \exp\left(\frac{365}{T}\right)$	2
21	$\text{MeOO} + \text{MeCO}_3$	$\rightarrow \text{HO}_2 + \text{HCHO} + \text{MeOO}$	$1.80 \times 10^{-12} \exp\left(\frac{500}{T}\right)$	2
22	$\text{MeOO} + \text{MeCO}_3$	$\rightarrow \text{MeCO}_2\text{H} + \text{HCHO}$	$2.00 \times 10^{-13} \exp\left(\frac{500}{T}\right)$	2
23	$\text{EtOO} + \text{NO}$	$\rightarrow \text{MeCHO} + \text{HO}_2 + \text{NO}_2$	$2.60 \times 10^{-12} \exp\left(\frac{380}{T}\right)$	2
24	$\text{EtOO} + \text{NO}_3$	$\rightarrow \text{MeCHO} + \text{HO}_2 + \text{NO}_2$	2.30×10^{-12}	2
25	$\text{EtOO} + \text{MeCO}_3$	$\rightarrow \text{MeCHO} + \text{HO}_2 + \text{MeOO}$	$4.40 \times 10^{-13} \exp\left(\frac{1070}{T}\right)$	2
26	$\text{MeCO}_3 + \text{NO}$	$\rightarrow \text{MeOO} + \text{CO}_2 + \text{NO}_2$	$7.50 \times 10^{-12} \exp\left(\frac{290}{T}\right)$	2
27	$\text{MeCO}_3 + \text{NO}_3$	$\rightarrow \text{MeOO} + \text{CO}_2 + \text{NO}_2$	4.00×10^{-12}	4
28	$\text{n-PrOO} + \text{NO}$	$\rightarrow \text{EtCHO} + \text{HO}_2 + \text{NO}_2$	$2.90 \times 10^{-12} \exp\left(\frac{350}{T}\right)$	2
29	$\text{n-PrOO} + \text{NO}_3$	$\rightarrow \text{EtCHO} + \text{HO}_2 + \text{NO}_2$	2.50×10^{-12}	4
30	$\text{i-PrOO} + \text{NO}$	$\rightarrow \text{Me}_2\text{CO} + \text{HO}_2 + \text{NO}_2$	$2.70 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	2
31	$\text{i-PrOO} + \text{NO}_3$	$\rightarrow \text{Me}_2\text{CO} + \text{HO}_2 + \text{NO}_2$	2.50×10^{-12}	4
32	$\text{EtCO}_3 + \text{NO}$	$\rightarrow \text{EtOO} + \text{CO}_2 + \text{NO}_2$	$6.70 \times 10^{-12} \exp\left(\frac{340}{T}\right)$	2
33	$\text{EtCO}_3 + \text{NO}_3$	$\rightarrow \text{EtOO} + \text{CO}_2 + \text{NO}_2$	4.00×10^{-12}	4
34	$\text{MeCOCH}_2\text{OO} + \text{NO}$	$\rightarrow \text{MeCO}_3 + \text{HCHO} + \text{NO}_2$	$2.80 \times 10^{-12} \exp\left(\frac{300}{T}\right)$	5
35	$\text{MeCOCH}_2\text{OO} + \text{NO}_3$	$\rightarrow \text{MeCO}_3 + \text{HCHO} + \text{NO}_2$	2.50×10^{-12}	4
36	$\text{NO} + \text{NO}_3$	$\rightarrow \text{NO}_2 + \text{NO}_2$	$1.80 \times 10^{-11} \exp\left(\frac{110}{T}\right)$	2
37	$\text{NO} + \text{O}_3$	$\rightarrow \text{NO}_2$	$1.40 \times 10^{-12} \exp\left(\frac{-1310}{T}\right)$	2
38	$\text{NO}_2 + \text{O}_3$	$\rightarrow \text{NO}_3$	$1.40 \times 10^{-13} \exp\left(\frac{-2470}{T}\right)$	2

Continued on next page

Table 3 – continued from previous page

Reactants	Products	<i>k</i>	Reference
39 NO ₃ + HCHO	→ HONO ₂ + HO ₂ + CO	$2.00 \times 10^{-12} \exp\left(\frac{-2440}{T}\right)$	2
40 NO ₃ + MeCHO	→ HONO ₂ + MeCO ₃	$1.40 \times 10^{-12} \exp\left(\frac{-1860}{T}\right)$	2
41 NO ₃ + EtCHO	→ HONO ₂ + EtCO ₃	$3.46 \times 10^{-12} \exp\left(\frac{-1862}{T}\right)$	4
42 NO ₃ + Me ₂ CO	→ HONO ₂ + MeCOCH ₂ OO	3.00×10^{-17}	2
43 N ₂ O ₅ + H ₂ O	→ HONO ₂ + HONO ₂	2.50×10^{-22}	2
44 O(³ P) + O ₃	→ O ₂ + O ₂	$8.00 \times 10^{-12} \exp\left(\frac{-2060}{T}\right)$	2
45 O(¹ D) + CH ₄	→ OH + MeOO	1.05×10^{-10}	2
46 O(¹ D) + CH ₄	→ HCHO + H ₂	7.50×10^{-12}	2
47 O(¹ D) + CH ₄	→ HCHO + HO ₂ + HO ₂	3.45×10^{-11}	2
48 O(¹ D) + H ₂ O	→ OH + OH	2.20×10^{-10}	2
49 O(¹ D) + N ₂	→ O(³ P) + N ₂	$2.10 \times 10^{-11} \exp\left(\frac{115}{T}\right)$	6
50 O(¹ D) + O ₂	→ O(³ P) + O ₂	$3.20 \times 10^{-11} \exp\left(\frac{67}{T}\right)$	2
51 OH + CH ₄	→ H ₂ O + MeOO	$1.85 \times 10^{-12} \exp\left(\frac{-1690}{T}\right)$	2
52 OH + C ₂ H ₆	→ H ₂ O + EtOO	$6.90 \times 10^{-12} \exp\left(\frac{-1000}{T}\right)$	2
53 OH + C ₃ H ₈	→ n-PrOO + H ₂ O	$7.60 \times 10^{-12} \exp\left(\frac{-585}{T}\right)$	2
54 OH + C ₃ H ₈	→ i-PrOO + H ₂ O	$7.60 \times 10^{-12} \exp\left(\frac{-585}{T}\right)$	2
55 OH + CO	→ HO ₂	1.44×10^{-13}	2
56 OH + EtCHO	→ H ₂ O + EtCO ₃	$5.10 \times 10^{-12} \exp\left(\frac{405}{T}\right)$	2
57 OH + EtOOH	→ H ₂ O + MeCHO + OH	8.01×10^{-12}	4
58 OH + EtOOH	→ H ₂ O + EtOO	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	4
59 OH + H ₂	→ H ₂ O + HO ₂	$7.70 \times 10^{-12} \exp\left(\frac{-2100}{T}\right)$	2
60 OH + H ₂ O ₂	→ H ₂ O + HO ₂	$2.90 \times 10^{-12} \exp\left(\frac{-160}{T}\right)$	2
61 OH + HCHO	→ H ₂ O + HO ₂ + CO	$5.40 \times 10^{-12} \exp\left(\frac{135}{T}\right)$	1
62 OH + HO ₂	→ H ₂ O	$4.80 \times 10^{-11} \exp\left(\frac{250}{T}\right)$	2
63 OH + HO ₂ NO ₂	→ H ₂ O + NO ₂	$1.90 \times 10^{-12} \exp\left(\frac{270}{T}\right)$	2
64 OH + HO ₂ NO ₂	→ H ₂ O + NO ₃	1.50×10^{-13}	2
65 OH + HONO	→ H ₂ O + NO ₂	$2.50 \times 10^{-12} \exp\left(\frac{260}{T}\right)$	2
66 OH + MeOOH	→ H ₂ O + HCHO + OH	$1.02 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	2
67 OH + MeOOH	→ H ₂ O + MeOO	$1.89 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	2
68 OH + MeONO ₂	→ HCHO + NO ₂ + H ₂ O	$4.00 \times 10^{-13} \exp\left(\frac{-845}{T}\right)$	2
69 OH + Me ₂ CO	→ H ₂ O + MeCOCH ₂ OO	$8.80 \times 10^{-12} \exp\left(\frac{-1320}{T}\right)$	2
70 OH + Me ₂ CO	→ H ₂ O + MeCOCH ₂ OO	$1.70 \times 10^{-14} \exp\left(\frac{420}{T}\right)$	2
71 OH + MeCOCH ₂ OOH	→ H ₂ O + MeCOCH ₂ OO	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	4
72 OH + MeCOCH ₂ OOH	→ OH + MGLY	8.39×10^{-12}	4
73 OH + MeCHO	→ H ₂ O + MeCO ₃	$4.40 \times 10^{-12} \exp\left(\frac{365}{T}\right)$	2
74 OH + NO ₃	→ HO ₂ + NO ₂	2.00×10^{-11}	2
75 OH + O ₃	→ HO ₂ + O ₂	$1.70 \times 10^{-12} \exp\left(\frac{-940}{T}\right)$	2
76 OH + OH	→ H ₂ O + O(³ P)	$6.31 \times 10^{-14} \left(\frac{T}{300}\right)^{2.6} \exp\left(\frac{945}{T}\right)$	2
77 OH + PAN	→ HCHO + NO ₂ + H ₂ O	3.00×10^{-14}	2
78 OH + PPAN	→ MeCHO + NO ₂ + H ₂ O	1.27×10^{-12}	4
79 OH + n-PrOOH	→ n-PrOO + H ₂ O	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	4

Continued on next page

Table 3 – continued from previous page

Reactants	Products	<i>k</i>	Reference
80 OH + n-PrOOH	→ EtCHO + H ₂ O + OH	1.10×10^{-11}	4
81 OH + i-PrOOH	→ i-PrOO + H ₂ O	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	4
82 OH + i-PrOOH	→ Me ₂ CO + OH	1.66×10^{-11}	4
83 O(³ P) + NO ₂	→ NO + O ₂	$5.50 \times 10^{-12} \exp\left(\frac{188}{T}\right)$	2
84 OH + C ₅ H ₈	→ ISO ₂	$2.70 \times 10^{-11} \exp\left(\frac{390}{T}\right)$	2
85 ^a OH + C ₅ H ₈	→ MACR + HCHO + MACRO ₂ + MeCO ₃	$3.33 \times 10^{-15} \exp\left(\frac{-1995}{T}\right)$	2
86 ^a OH + C ₅ H ₈	→ MeOO + HCOOH + CO + H ₂ O ₂	$3.33 \times 10^{-15} \exp\left(\frac{-1995}{T}\right)$	2
87 ^a OH + C ₅ H ₈	→ HO ₂ + OH	$3.33 \times 10^{-15} \exp\left(\frac{-1995}{T}\right)$	2
88 NO ₃ + C ₅ H ₈	→ ISON	$3.15 \times 10^{-12} \exp\left(\frac{-450}{T}\right)$	2
89 NO + ISO ₂	→ NO ₂ + MACR + HCHO + HO ₂	$2.43 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4,7
90 NO + ISO ₂	→ ISON	$1.12 \times 10^{-13} \exp\left(\frac{360}{T}\right)$	4,7
91 HO ₂ + ISO ₂	→ ISOOH	$2.05 \times 10^{-13} \exp\left(\frac{1300}{T}\right)$	4,7
92 ISO ₂ + ISO ₂	→ MACR + MACR + HCHO + HO ₂	2.00×10^{-12}	7
93 OH + ISOOH	→ MACR + OH	1.00×10^{-10}	7
94 OH + ISON	→ HACET + NALD	1.30×10^{-11}	7
95 OH + MACR	→ MACRO ₂	$1.30 \times 10^{-12} \exp\left(\frac{610}{T}\right)$	2
96 OH + MACR	→ MACRO ₂	$4.00 \times 10^{-12} \exp\left(\frac{380}{T}\right)$	2
97 ^a O ₃ + MACR	→ MGLY + HCOOH + HO ₂ + CO	$2.13 \times 10^{-16} \exp\left(\frac{-1520}{T}\right)$	2
98 ^a O ₃ + MACR	→ OH + MeCO ₃	$2.13 \times 10^{-16} \exp\left(\frac{-1520}{T}\right)$	2
99 ^a O ₃ + MACR	→ MGLY + HCOOH + HO ₂ + CO	$3.50 \times 10^{-16} \exp\left(\frac{-2100}{T}\right)$	2
100 ^a O ₃ + MACR	→ OH + MeCO ₃	$3.50 \times 10^{-16} \exp\left(\frac{-2100}{T}\right)$	2
101 ^a NO + MACRO ₂	→ NO ₂ + MeCO ₃ + HACET + CO	$1.27 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4,7
102 ^a NO + MACRO ₂	→ MGLY + HCHO + HO ₂	$1.27 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4,7
103 HO ₂ + MACRO ₂	→ MACROOH	$1.83 \times 10^{-13} \exp\left(\frac{1300}{T}\right)$	4,7
104 ^a MACRO ₂ + MACRO ₂	→ HACET + MGLY + HCHO + CO	1.00×10^{-12}	4,7
105 ^a MACRO ₂ + MACRO ₂	→ HO ₂	1.00×10^{-12}	4,7
106 OH + MPAN	→ HACET + NO ₂	2.90×10^{-11}	2
107 OH + MACROOH	→ MACRO ₂	3.00×10^{-11}	7
108 OH + HACET	→ MGLY + HO ₂	3.00×10^{-12}	2,7
109 OH + MGLY	→ MeCO ₃ + CO	1.50×10^{-11}	2,7
110 NO ₃ + MGLY	→ MeCO ₃ + CO + HONO ₂	$3.46 \times 10^{-12} \exp\left(\frac{-1860}{T}\right)$	4
111 OH + NALD	→ HCHO + CO + NO ₂	$4.40 \times 10^{-12} \exp\left(\frac{365}{T}\right)$	2,7
112 OH + MeCO ₃ H	→ MeCO ₃	3.70×10^{-12}	4,7
113 OH + MeCO ₂ H	→ MeOO	$4.00 \times 10^{-13} \exp\left(\frac{200}{T}\right)$	8
114 OH + HCOOH	→ HO ₂	4.50×10^{-13}	2
115 MeOH + OH	→ HCHO + HO ₂	$2.85 \times 10^{-12} \exp\left(\frac{-345}{T}\right)$	3
116 OH + C ₁₀ H ₁₆	→ TERPO ₂	$1.20 \times 10^{-11} \exp\left(\frac{444}{T}\right)$	9
117 O ₃ + C ₁₀ H ₁₆	→ OH + MEK + HO ₂	$1.00 \times 10^{-15} \exp\left(\frac{-732}{T}\right)$	9
118 NO ₃ + C ₁₀ H ₁₆	→ ISON + MACR	$1.20 \times 10^{-12} \exp\left(\frac{490}{T}\right)$	9
119 ^a NO + TERPO ₂	→ Me ₂ CO + HO ₂ + NO ₂	$2.10 \times 10^{-12} \exp\left(\frac{180}{T}\right)$	9
120 ^a NO + TERPO ₂	→ MACR + MACR	$2.10 \times 10^{-12} \exp\left(\frac{180}{T}\right)$	9

Continued on next page

Table 3 – continued from previous page

Reactants	Products	<i>k</i>	Reference
121 ^a	$\text{HO}_2 + \text{TERPO}_2 \rightarrow \text{TERPOOH}$	$7.50 \times 10^{-13} \exp\left(\frac{700}{T}\right)$	9
122 ^a	$\text{OH} + \text{TERPOOH} \rightarrow \text{TERPO}_2$	$3.80 \times 10^{-12} \exp\left(\frac{200}{T}\right)$	9
123	$\text{C}_4\text{H}_{10} + \text{OH} \rightarrow \text{BtOO} + \text{H}_2\text{O}$	$9.10 \times 10^{-12} \exp\left(\frac{-405}{T}\right)$	3
124 ^a	$\text{BtOO} + \text{NO} \rightarrow \text{NO}_2 + \text{MEK} + \text{HO}_2 + \text{EtOO}$	$1.27 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4
125 ^a	$\text{BtOO} + \text{NO} \rightarrow \text{ONIT} + \text{MeCHO}$	$1.27 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4
126	$\text{BtOO} + \text{HO}_2 \rightarrow \text{BtOOH}$	$1.82 \times 10^{-13} \exp\left(\frac{1300}{T}\right)$	4
127 ^a	$\text{BtOO} + \text{MeOO} \rightarrow \text{MEK} + \text{HCHO} + \text{HO}_2 + \text{MeCHO}$	1.25×10^{-13}	4
128 ^a	$\text{BtOO} + \text{MeOO} \rightarrow \text{MeOH} + \text{EtOO}$	1.25×10^{-13}	4
129 ^a	$\text{BtOOH} + \text{OH} \rightarrow \text{BtOO} + \text{MEK} + \text{OH} + \text{H}_2\text{O}$	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	4
130	$\text{MEK} + \text{OH} \rightarrow \text{MEKOO}$	$1.30 \times 10^{-12} \exp\left(\frac{-25}{T}\right)$	3
131	$\text{MEKOO} + \text{NO} \rightarrow \text{MeCHO} + \text{MeCO}_3 + \text{NO}_2 + \text{ONIT}$	$2.54 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4
132	$\text{MEKOO} + \text{HO}_2 \rightarrow \text{MEKOHH}$	$1.82 \times 10^{-13} \exp\left(\frac{1300}{T}\right)$	4
133	$\text{MEKOHH} + \text{OH} \rightarrow \text{MeCOCOMe} + \text{OH} + \text{OH}$	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	4
134	$\text{ONIT} + \text{OH} \rightarrow \text{MEK} + \text{NO}_2 + \text{H}_2\text{O}$	1.60×10^{-12}	3
135 ^a	$\text{C}_2\text{H}_4 + \text{O}_3 \rightarrow \text{HCHO} + \text{HO}_2 + \text{OH} + \text{CO}$	$4.55 \times 10^{-15} \exp\left(\frac{-2580}{T}\right)$	3
136 ^a	$\text{C}_2\text{H}_4 + \text{O}_3 \rightarrow \text{H}_2 + \text{CO}_2 + \text{HCOOH}$	$4.55 \times 10^{-15} \exp\left(\frac{-2580}{T}\right)$	3
137 ^a	$\text{C}_3\text{H}_6 + \text{O}_3 \rightarrow \text{HCHO} + \text{MeCHO} + \text{OH} + \text{HO}_2$	$1.83 \times 10^{-15} \exp\left(\frac{-1880}{T}\right)$	3
138 ^a	$\text{C}_3\text{H}_6 + \text{O}_3 \rightarrow \text{EtOO} + \text{MGLY} + \text{CH}_4 + \text{CO}$	$1.83 \times 10^{-15} \exp\left(\frac{-1880}{T}\right)$	3
139 ^a	$\text{C}_3\text{H}_6 + \text{O}_3 \rightarrow \text{MeOH} + \text{MeOO} + \text{HCOOH}$	$1.83 \times 10^{-15} \exp\left(\frac{-1880}{T}\right)$	3
140 ^a	$\text{C}_3\text{H}_6 + \text{NO}_3 \rightarrow \text{ONIT}$	$4.60 \times 10^{-13} \exp\left(\frac{-1155}{T}\right)$	3
141 ^a	$\text{PrpeOO} + \text{NO} \rightarrow \text{MeCHO} + \text{HCHO} + \text{HO}_2 + \text{NO}_2$	$1.27 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4
142 ^a	$\text{PrpeOO} + \text{NO} \rightarrow \text{ONIT}$	$1.27 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	4
143	$\text{PrpeOO} + \text{HO}_2 \rightarrow \text{PrpeOOH}$	$1.50 \times 10^{-13} \exp\left(\frac{1300}{T}\right)$	4
144	$\text{PrpeOOH} + \text{OH} \rightarrow \text{PrpeOO} + \text{H}_2\text{O}$	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	4
145	$\text{PrpeOOH} + \text{OH} \rightarrow \text{HACET} + \text{OH}$	2.44×10^{-11}	4
146 ^a	$\text{AROM} + \text{OH} \rightarrow \text{AROMO}_2 + \text{HO}_2$	$1.81 \times 10^{-12} \exp\left(\frac{338}{T}\right)$	10
147 ^a	$\text{AROMO}_2 + \text{NO} \rightarrow \text{MGLY} + \text{NO}_2 + \text{MeCO}_3 + \text{CO}$	$1.35 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	10
148 ^a	$\text{AROMO}_2 + \text{NO} \rightarrow \text{HO}_2$	$1.35 \times 10^{-12} \exp\left(\frac{360}{T}\right)$	10
149 ^a	$\text{AROMO}_2 + \text{NO}_3 \rightarrow \text{MGLY} + \text{NO}_2 + \text{MeCO}_3 + \text{CO}$	1.20×10^{-12}	10
150 ^a	$\text{AROMO}_2 + \text{NO}_3 \rightarrow \text{HO}_2$	1.20×10^{-12}	10
151 ^a	$\text{AROMO}_2 + \text{HO}_2 \rightarrow \text{AROMOOH}$	$1.90 \times 10^{-13} \exp\left(\frac{-1300}{T}\right)$	10
152 ^a	$\text{AROMO}_2 + \text{MeOO} \rightarrow \text{MGLY} + \text{CO} + \text{MeCO}_3 + \text{MeOH}$	1.15×10^{-13}	10
153 ^a	$\text{AROMO}_2 + \text{MeOO} \rightarrow \text{HO}_2 + \text{HCHO}$	1.15×10^{-13}	10
154 ^a	$\text{AROMOOH} + \text{OH} \rightarrow \text{AROMO}_2$	$1.90 \times 10^{-12} \exp\left(\frac{190}{T}\right)$	10
155 ^a	$\text{AROMOOH} + \text{OH} \rightarrow \text{OH} + \text{H}_2\text{O}$	$4.61 \times 10^{-18} \exp\left(\frac{253}{T}\right)$	10
156 ^a	$\text{AROMOOH} + \text{OH} \rightarrow \text{MeCO}_3 + \text{CO} + \text{HO}_2 + \text{OH}$	$4.19 \times 10^{-17} \exp\left(\frac{696}{T}\right)$	10
157	$\text{HO}_2 + \text{O}_3\text{S} \rightarrow \text{HO}_2 + \text{O}_2$	$2.03 \times 10^{-16} \left(\frac{T}{300}\right)^{4.57} \exp\left(\frac{693}{T}\right)$	2
158	$\text{OH} + \text{O}_3\text{S} \rightarrow \text{OH} + \text{O}_2$	$1.70 \times 10^{-12} \exp\left(\frac{-940}{T}\right)$	2
159	$\text{O}^{(1)\text{D}}\text{S} + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}$	2.20×10^{-10}	2
160	$\text{O}^{(1)\text{D}}\text{S} + \text{N}_2 \rightarrow \text{O}^{(3)\text{P}}\text{S} + \text{N}_2$	$2.10 \times 10^{-11} \exp\left(\frac{115}{T}\right)$	6

Continued on next page

Table 3 – continued from previous page

	Reactants	Products	<i>k</i>	Reference
161	$O(^1D)S + O_2$	$\rightarrow O(^3P)S + O_2$	$3.20 \times 10^{-11} \exp(\frac{67}{T})$	2

1085

a: Reactions are split between multiple lines.

Table 4: TOMCAT gas-phase termolecular and thermal decomposition reactions. Rate constant $k = \left(\frac{k_0[M]}{1+k_0[M]/k_\infty}\right) F_c^{(1+\log\frac{k_0[M]}{k_\infty})^2-1}$, where k_0 is the low pressure limit, k_∞ is the high pressure limit and M is the number density in molecules/cm³. $F_c = f$ when $f < 1$ else $F_c = \exp(-T/f)$. Low pressure limit $k_0 = k_1 \left(\frac{T}{300}\right)^{\alpha_1} \exp\left(-\frac{\beta_1}{T}\right)$ and high pressure limit $k_\infty = k_2 \left(\frac{T}{300}\right)^{\alpha_2} \exp\left(-\frac{\beta_2}{T}\right)$. Reaction rate references 1: Atkinson et al. (b), 2: MCM (2004), 3: Pöschl et al. (2000), 4: Atkinson et al. (c).

	Reactants	Products	f	k_1	α_1	β_1	k_2	α_2	β_2	Reference
1 ^a	$\text{HO}_2 + \text{HO}_2 + \text{M}$	$\rightarrow \text{H}_2\text{O}_2 + \text{O}_2 + \text{M}$	0.00	1.90×10^{-33}	0.00	-980.0	$0.00 \times 10^{+00}$	0.00	0.0 ¹	1
2	$\text{HO}_2 + \text{NO}_2 + \text{M}$	$\rightarrow \text{HO}_2\text{NO}_2 + \text{M}$	0.60	1.80×10^{-31}	-3.20	0.0	4.70×10^{-12}	0.00	0.0	1
3	$\text{HO}_2\text{NO}_2 + \text{M}$	$\rightarrow \text{HO}_2 + \text{NO}_2 + \text{M}$	0.60	4.10×10^{-05}	0.00	10650.0	$4.80 \times 10^{+15}$	0.00	11170.0	1
4	$\text{MeCO}_3 + \text{NO}_2 + \text{M}$	$\rightarrow \text{PAN} + \text{M}$	0.30	2.70×10^{-28}	-7.10	0.0	1.20×10^{-11}	-0.90	0.0	1
5	$\text{PAN} + \text{M}$	$\rightarrow \text{MeCO}_3 + \text{NO}_2 + \text{M}$	0.30	4.90×10^{-03}	0.00	12100.0	$5.40 \times 10^{+16}$	0.00	13830.0	1
6	$\text{N}_2\text{O}_5 + \text{M}$	$\rightarrow \text{NO}_2 + \text{NO}_3 + \text{M}$	0.35	1.30×10^{-03}	-3.50	11000.0	$9.70 \times 10^{+14}$	0.10	11080.0	1
7	$\text{NO}_2 + \text{NO}_3 + \text{M}$	$\rightarrow \text{N}_2\text{O}_5 + \text{M}$	0.35	3.60×10^{-30}	-4.10	0.0	1.90×10^{-12}	0.20	0.0	1
8	$\text{O}({}^3\text{P}) + \text{O}_2 + \text{M}$	$\rightarrow \text{O}_3 + \text{M}$	0.00	5.70×10^{-34}	-2.60	0.0	$0.00 \times 10^{+00}$	0.00	0.0	1
9	$\text{OH} + \text{NO} + \text{M}$	$\rightarrow \text{HONO} + \text{M}$	1420.00	7.40×10^{-31}	-2.40	0.0	3.30×10^{-11}	-0.30	0.0	1
10	$\text{OH} + \text{NO}_2 + \text{M}$	$\rightarrow \text{HONO}_2 + \text{M}$	0.40	3.30×10^{-30}	-3.00	0.0	4.10×10^{-11}	0.00	0.0	1
11	$\text{OH} + \text{OH} + \text{M}$	$\rightarrow \text{H}_2\text{O}_2 + \text{M}$	0.50	6.90×10^{-31}	-0.80	0.0	2.60×10^{-11}	0.00	0.0	1
12	$\text{EtCO}_3 + \text{NO}_2 + \text{M}$	$\rightarrow \text{PPAN} + \text{M}$	0.30	2.70×10^{-28}	-7.10	0.0	1.20×10^{-11}	-0.90	0.0	2
13	$\text{PPAN} + \text{M}$	$\rightarrow \text{EtCO}_3 + \text{NO}_2 + \text{M}$	0.36	1.70×10^{-03}	0.00	11280.0	$8.30 \times 10^{+16}$	0.00	13940.0	1
14	$\text{MACRO}_2 + \text{NO}_2 + \text{M}$	$\rightarrow \text{MPAN} + \text{M}$	0.30	2.70×10^{-28}	0.00	11280.0	$8.30 \times 10^{+16}$	0.00	13940.0	3
15	$\text{MPAN} + \text{M}$	$\rightarrow \text{MACRO}_2 + \text{NO}_2 + \text{M}$	0.30	4.90×10^{-03}	0.00	12100.0	$5.40 \times 10^{+16}$	0.00	13830.0	3
16	$\text{O}({}^3\text{P}) + \text{O}_2 + \text{M}$	$\rightarrow \text{O}_3 + \text{M}$	0.00	5.70×10^{-34}	-2.60	0.0	$0.00 \times 10^{+00}$	0.00	0.0	1
17 ^b	$\text{C}_2\text{H}_4 + \text{OH} + \text{M}$	$\rightarrow \text{PrpeOO} + \text{M}$	0.48	2.87×10^{-29}	-3.10	0.0	3.00×10^{-12}	-0.85	0.0	4
18 ^b	$\text{C}_2\text{H}_4 + \text{OH} + \text{M}$	$\rightarrow \text{PrpeOO} + \text{M}$	0.48	2.87×10^{-29}	-3.10	0.0	3.00×10^{-12}	-0.85	0.0	4
19 ^b	$\text{C}_2\text{H}_4 + \text{OH} + \text{M}$	\rightarrow	0.48	2.87×10^{-29}	-3.10	0.0	3.00×10^{-12}	-0.85	0.0	4
20	$\text{C}_3\text{H}_6 + \text{OH} + \text{M}$	$\rightarrow \text{PrpeOO} + \text{M}$	0.50	8.00×10^{-27}	-3.50	0.0	3.00×10^{-11}	-1.00	0.0	4

a: Reaction rate is dependent on H_2O so k is weighted by factor of $1 + 1.4E - 21[\text{H}_2\text{O}] \exp(2200/T)$, where $[\text{H}_2\text{O}]$ is in molecules cm⁻³.

b: Reactions are split between multiple lines.

Table 5: TOMCAT photolysis reactions.

Reaction	Reactants	Products
1	EtOOH + $h\nu$	\rightarrow MeCHO + HO ₂ + OH
2	H ₂ O ₂ + $h\nu$	\rightarrow OH + OH
3a	HCHO + $h\nu$	\rightarrow HO ₂ + HO ₂ + CO
3b	HCHO + $h\nu$	\rightarrow H ₂ + CO
5	HO ₂ NO ₂ + $h\nu$	\rightarrow HO ₂ + NO ₂
6	HONO ₂ + $h\nu$	\rightarrow OH + NO ₂
7a	MeCHO + $h\nu$	\rightarrow MeOO + HO ₂ + CO
7b	MeCHO + $h\nu$	\rightarrow CH ₄ + CO
9	MeOOH + $h\nu$	\rightarrow HO ₂ + HCHO + OH
10	N ₂ O ₅ + $h\nu$	\rightarrow NO ₃ + NO ₂
11	NO ₂ + $h\nu$	\rightarrow NO + O(³ P)
12a	NO ₃ + $h\nu$	\rightarrow NO + O ₂
12b	NO ₃ + $h\nu$	\rightarrow NO ₂ + O(³ P)
14	O ₂ + $h\nu$	\rightarrow O(³ P) + O(³ P)
15a	O ₃ + $h\nu$	\rightarrow O ₂ + O(¹ D)
15b	O ₃ + $h\nu$	\rightarrow O ₂ + O(³ P)
17	PAN + $h\nu$	\rightarrow MeCO ₃ + NO ₂
18	HONO + $h\nu$	\rightarrow OH + NO
19	EtCHO + $h\nu$	\rightarrow EtOO + HO ₂ + CO
20	Me ₂ CO + $h\nu$	\rightarrow MeCO ₃ + MeOO
21	n-PrOOH + $h\nu$	\rightarrow EtCHO + HO ₂ + OH
22	i-PrOOH + $h\nu$	\rightarrow Me ₂ CO + HO ₂ + OH
23	MeCOCH ₂ OOH + $h\nu$	\rightarrow MeCO ₃ + HCHO + OH
24	PPAN + $h\nu$	\rightarrow EtCO ₃ + NO ₂
25	MeONO ₂ + $h\nu$	\rightarrow HO ₂ + HCHO + NO ₂
26a	TERPOOH + $h\nu$	\rightarrow OH + HO ₂ + MACR + MACR
26b	TERPOOH + $h\nu$	\rightarrow TERPOOH + Me ₂ CO
28	ISOOH + $h\nu$	\rightarrow OH + MACR + HCHO + HO ₂
29	ISON + $h\nu$	\rightarrow NO ₂ + MACR + HCHO + HO ₂
30	MACR + $h\nu$	\rightarrow MeCO ₃ + HCHO + CO + HO ₂
31	MPAN + $h\nu$	\rightarrow MACRO ₂ + NO ₂
32a	MACROOH + $h\nu$	\rightarrow OH + HO ₂ + OH + HO ₂
32b	MACROOH + $h\nu$	\rightarrow HACET + CO + MGLY + HCHO
34	HACET + $h\nu$	\rightarrow MeCO ₃ + HCHO + HO ₂
35	MGLY + $h\nu$	\rightarrow MeCO ₃ + CO + HO ₂ +
36	NALD + $h\nu$	\rightarrow HCHO + CO + NO ₂ + HO ₂
37	MeCO ₃ H + $h\nu$	\rightarrow MeOO + OH
38a	O ₃ S + $h\nu$	\rightarrow O ₂ + O(¹ D)S
38b	O ₃ S + $h\nu$	\rightarrow O ₂ + O(³ P)S
40a	BtOOH + $h\nu$	\rightarrow MEK + MEK + EtOO + MeCHO

Continued on next page

Table 5 – continued from previous page

Reaction	Reactants	Products
40b	BtOOH + $h\nu$	\rightarrow HO ₂ + HO ₂
40c	BtOOH + $h\nu$	\rightarrow OH + OH + OH
43	MEK + $h\nu$	\rightarrow MeCO ₃ + EtOO
44	MeCOCOMe + $h\nu$	\rightarrow MeCO ₃ + MeCO ₃
45	MEKO ₂ H + $h\nu$	\rightarrow MeCO ₃ + MeCHO + OH
46a	ONIT + $h\nu$	\rightarrow NO ₂ + MEK + HO ₂ + EtOO
46b	ONIT + $h\nu$	\rightarrow MeCHO + ONIT
48a	AROMOOH + $h\nu$	\rightarrow OH + Me ₂ CO + HO ₂ + CO
48b	AROMOOH + $h\nu$	\rightarrow MeCO ₃ + AROMOOH

Table 6. List of γ values used in TOMCAT for heterogeneous uptake of N_2O_5 by aerosol.

Aerosol Type	Reaction Probability (T=temperature (K), RH= relative humidity (%))
Sulphate	$\gamma = \alpha \times 10^\beta$ $\alpha = 2.79 \times 10^{-4} + 1.3$ $\times 10^{-4} \times RH - 3.43$ $\times 10^{-6} \times RH^2 + 7.52$ $\times 10^{-8} \times RH^3$ $\beta = 4 \times 10^{-2} \times (T - 294)$ ($T \geq 282\text{K}$) $\beta = -0.48$ ($T < 282\text{K}$)
Organic Carbon	$\gamma = RH \times 5.2 \times 10^{-4}$ ($RH < 57\%$)
Black Carbon	$\gamma = 0.005$
Sea Salt	$\gamma = 0.005$ ($RH < 62\%$) $\gamma = 0.03$ ($RH \geq 62\%$)
Dust	$\gamma = 0.02$

Table 7. Model diagnostics compared to previously published values

Diagnostic	TOMCAT	Published Values	Reference
O_3 Burden (Tg) ^a	331	337 ± 23	Young et al. (2013)
OH concentration ($\times 10^6$ molecules/cm ³) ^b	1.08	0.94–1.06	Krol and Lelieveld (2003), Prinn et al. (2001), Bousquet et al. (2005), Wang et al. (2008)
CH_4 lifetime (yrs)	7.9	9.3 ± 0.9	Voulgarakis et al. (2013)

a: Annual mean; b: Mass-weighted annual mean.