

GMD Manuscript, response to editor and reviewer comments

Editor Comment 1: A. Kerkweg

Dear authors,

In my role as Executive editor of GMD, I would like to bring to your attention our Editorial version 1.1:

<http://www.geosci-model-dev.net/8/3487/2015/gmd-8-3487-2015.html>

This highlights some requirements of papers published in GMD, which is also available on the GMD website in the 'Manuscript Types' section:

http://www.geoscientific-model-development.net/submission/manuscript_types.html

In particular, please note that for your paper, the following requirement has not been met in the Discussions paper:

- "The main paper must give the model name and version number (or other unique identifier) in the title."

Please add a version number for your model in the title upon your revised submission to GMD.

Yours,

Astrid Kerkweg

Thank you for bringing this requirement to our attention. Since our model is a modification to UVic v 2.9, we have chosen to remove this distinction from the title. Therefore, we have revised our title to read:

A Cloud Feedback Emulator (CFE v1.0) for an intermediate complexity model

Response to Anonymous Referee #1

We thank Anonymous Referee #1 for his/her thoughtful and insightful comments on our manuscript. We have responded to the comments below (in red).

General comments:

The paper "The University of Victoria Cloud Feedback Emulator (UVic-CFE): cloud radiative feedbacks in an intermediate complexity model" describes and evaluates a new method for applying GCM-derived cloud feedbacks to intermediate complexity models. The new method is able to capture the spread in TOA radiative feedbacks between the original GCMs, implying that the tool is generally efficient. Given that cloud feedback plays an essential role in determine the magnitude of global warming, this method is expected to be useful in improving intermediate complexity models that are in lack of cloud feedbacks. Therefore, I suggest the paper be published after addressing my following comments.

1. The cloud masking effect has not been removed from cloud radiative effect when cloud feedback is calculated. This may result in a systematic bias to the TOA radiations. Consider an assumptive situation that there is no change in cloud properties under global warming, then no cloud terms need to be added up to Eq. (6), and the $OLW_{cloud}(t)$ term in Eq. (15) should be zero. However, the cloud longwave radiative effect in GCM would still change due to changes in water vapor and temperature (cloud masking effect), leading to a non-zero value in Eq. (15). Therefore, an additional term is needed to compensate the cloud masking effect (this may be done with radiative kernels, Soden et al. 2008, doi: 10.1175/2007JCLI2110.1).

Because the cloud masking effect is evident in the 4xCO₂ GCM simulations, we choose to include it in our OLW cloud feedback. The OLW parameterization of UVic (Eq. 6) lacks the full cloud dynamics of the GCMs, so by assessing our OLW cloud feedback with the full cloud radiative effect (CRE) anomalies, we aim to incorporate the full OLW cloud changes of the GCMs, including cloud-masking effects. That said, the Referee is correct in pointing out the differences in the CRE and kernel methods, so we make note of potential issues as suggested by the reviewer (section 2.3.2).

2. The calculation of "TOA feedbacks" is inaccurate, so I suggest the authors to either calculate the TOA feedbacks with the standard method, or to replace "TOA feedbacks" with another phrase. In this paper, climate forcing is included in the "TOA feedbacks" (Page 9, line 6), so the TOA feedbacks in Fig. 6 ($_{OLW}/m^2/K$) is much larger than that calculated by Andrews et al. 2012 (-1.08 W/m²/K).

Since the exact greenhouse gas forcing is unknown for each of the GCMs, we favor our current method of calculating TOA "feedbacks". Therefore, we have replaced the phrase "TOA feedbacks" with "TOA radiative-temperature response" so as not to confuse readers with the standard method.

3. The cloud rapid adjustment has not been removed from the cloud feedback (Zelinka et al. 2013, doi:10.1175/JCLI-D-12-00555.1). This may be partially responsible to the loss of spread in UVic-CFE simulations (Fig. 6). I expect that the UVic-CFE would be more accurate after the above comments are addressed.

Yes, by calculating our cloud radiative effect using CRE anomalies instead of the 'Gregory' slope method, we are neglecting influence of cloud rapid adjustment. However, we are unable to include them as part of the forcing (as suggested by Zelinka et al., 2013), so we must include them in the cloud feedback. We make note of this issue in the revised text (discussion) and its possible influence on the spread in UVic-CFE.

Specific comments:

Page 1, Line 29. "The relative magnitude and net effect of these feedbacks depends on cloud altitude. . . High clouds, on the other hand, radiate at much colder temperatures than the surface, which can make the longwave effect dominate and lead to net warming"

The authors may also discuss the effect of cloud optical depth here. The net cloud radiative effect of high clouds could be either positive (high thin clouds) or negative (high thick clouds) depending on the cloud optical depth.

Yes, the referee is correct in noting the importance of optical depth. We now include a statement on cloud optical depth and its relationship with cloud radiative effect.

Page 3, Line 30. It is worth discussing whether the cloud radiative effect is included when the empirical parameters in Eq. (6) were calculated.

The original parameterization of Thompson and Warren (1982) is for clear-sky outgoing LW fluxes. Therefore, there is no cloud feedback within this parameterization. We now make note of this in our description of Eq. (6).

Page 7, Line 17. Please provide more details for the 4xCO₂ and LGM experiment design.

Our experimental design is described in the second paragraph of section 2.4. Is there a particular aspect of the experimental design that you would like to see further described?

Figure 4 and 5. There are some white pixels surrounded by blue/green pixels (for example, Central tropical Pacific Ocean in Fig. 4a,d). Are these white pixels induced by missing values? Is it possible to eliminate them?

These white pixels appear to be due to an improper image rendering of the original postscript image. We have corrected these images by first saving the file to a jpeg.

Technical comments

Page 3, Line 25. How is the variable f calculated?

The variable f is as an adjustment factor that is necessary to correct for a radiative imbalance that arises in our estimate of atmospheric albedo from the CERES observational data. Therefore, this adjustment factor directly affects global mean surface air temperature. We ran a series of control simulations with different f -values to tune our control simulation to observational estimates of -13.9 C ($f=0.95$). We have rephrased our description in section 2.2 to address this confusion.

Figure 6, 8. Figure legend: "UVIC control" -> "UVic control", to be consistent with the figure description.

Fixed

Response to Anonymous Referee #2

We thank Anonymous Referee #2 for his/her thoughtful and insightful comments on our manuscript. We have responded to the comments below (in red).

From the reference to the paper by Weaver and co-authors, UVic needs a choice between options, like humidity transport or diffusive. It also looks as if the atm wind can be prescribed or sensitive to SAT and

density. Evidently, such options impact the feedbacks in UVic. I suggest a table to summarize these options, and the feedback concerned.

The referee is correct in noting that UVic contains options regarding atmospheric transport/diffusion and atm winds. In the first paragraph of section 2.1 (4th sentence), we have stated that we are using UVic version 2.9, which includes the atmospheric heat diffusion feedback (diffusion as a function of global mean surface air temperature). This is a feedback that the latest version of UVic includes and we make note that it has been shown to improve the latitudinal temperature gradient for the Last Glacial Maximum (when compared with proxy data; Fyke and Eby, 2012).

To isolate the effect of cloud feedbacks in our emulator, we choose to prescribe atm wind stress (no SAT feedback). However, large differences in the surface boundary conditions at the LGM (ice sheets) have been shown to greatly impact wind stress anomalies in LGM simulations (Muglia and Schmittner, 2016). The optional wind-SAT feedback would not capture these changes; therefore, we apply wind stress anomalies as diagnosed from the LGM GCM results (see end of section 2.4).

To be consistent in our model design, we also prescribe modern wind stress for our 4xCO₂ simulations. Wind stress anomalies across the CMIP5 4xCO₂ experiments are small; therefore, we use the prescribed wind stress fields of the control UVic 2.9 model. Upon reviewing our manuscript, we discussed the use of LGM wind stress anomalies at the end of section 2.4, but did not note our wind stress boundary condition for the 4xCO₂ simulations. We have added an additional sentence at the end of section 2.4 that further discusses the prescribed wind stress for the 4xCO₂ simulations.

In the same views, the way cloud feedback, as approximated by the atm-albedo could be explicitly described for clarity. The description of some feedback-loops would be of great help. For instance, atm-albedo -> SAT-> OLW etc How is the ocean dynamics impacted? What changes are observed concerning the thermohaline circulation, the thermocline etc What about the sea-ice extent?

We provide a description of the nature of cloud feedback loops (through their impact on SW and LW radiative balance) in our introduction.

Regarding ocean dynamics and sea-ice, we have chosen to concentrate this manuscript on how our linear parameterization of cloud feedbacks helps capture the change in radiative balance that would otherwise be missing in UVic (or similar EMIC). We have focused our discussion of radiative feedbacks on surface air temperature evolution and climate sensitivity, by association with the 4xCO₂ experiments of the CMIP5 coordinated framework. Therefore, we have not included any analysis on the impacts of these radiative changes on ocean dynamics in this study.

Some comments are already included in the RESULTS sections that could be related more closely to the UVic extended results others that the averaged global results directly as support to the comparison with the seven GCMs.

See responses to comments from other reviewer/editors.

Editor Comment 2: K. Gierens

I have a few comments/questions concerning several equations in the manuscript. Please consider them for your revised version.

1) Eq. 3: a better explanation of f is needed. How does the choice of f guarantee that α_{atm} remains bounded by 0 and 1? Also it should be stated that S in equation 2 is identical to $SW_{\text{in,TOA}}$ in eq. 4.

For explanation of f , see response to Referee #1. We have rephrased our description in section 2.2 to address this confusion.

The variable f does not guarantee that α_{atm} is bounded by 0 and 1. However, the editor is correct in noting that such a limit should be in place. We set this limit in our calculation of α_{plt} and α_{sfc} . Under low-light conditions (winter, high-latitudes), the denominators of equation 4 and 5 ($SW_{\text{in,TOA}}$) can become small relative to the numerator, resulting in a value of $\alpha_{\text{plt}} > 1$ or $\alpha_{\text{sfc}} > 1$ across a latitudinal band. Therefore, we limit α_{plt} and α_{sfc} to be between 0 and 1. If α_{plt} and α_{sfc} are greater than 1, we assign them with a value from the next closest month

in time where α_{plf} and α_{sfc} are appropriately defined. However, we note that these are months with low incoming light ($SW_{\text{in,TOA}}$), so the effect of α_{plf} and α_{sfc} on local radiative balance is negligible. In short, we do limit α_{atm} through a limit on α_{plf} and α_{sfc} . With these limits, α_{plf} is bounded by 0 and 1. We have added a sentence explaining this in our description of new α_{atm} with CERES data (section 2.2).

We have adjusted S in eq. 2 so that it now uses the consistent variable " $SW_{\text{in,TOA}}$ " as in eq. 4.

2) Eqs. 11 and 12: The argument that albedo values are not additive leads you to formally consider the ratio $\alpha_{\text{atm,perturbed}}/\alpha_{\text{atm,CERES}}$ in eq. 11, however it is necessary to subtract one from this ratio. Mathematically, we then have the difference of the albedo values back, since $(\alpha_{\text{atm,perturbed}}/\alpha_{\text{atm,CERES}}) - 1 = (\alpha_{\text{atm,perturbed}} - \alpha_{\text{atm,CERES}})/\alpha_{\text{atm,CERES}}$.

In eq. 12 this expression is then multiplied by $\alpha_{\text{atm,CERES}}$, and the simple difference of the albedo values returns back. So this argumentation seems to add unnecessary complexity.

Yes, the editor is correct in noting that the 1 is mathematically unnecessary. However, in defining the atm albedo feedback this way (centered around zero), it is easier to demonstrate when the feedback is positive and when it is negative. By association, we feel the plots of the atm albedo feedback are more illustrative when centered around zero.

3) Eqs. 12 and 15: I wonder whether these equations are used at every timestep. If so, how do you distinguish climatological temperature variations from diurnal and seasonal temperature variations? Should a feedback not work only on the long climatological time scales? Furthermore, are these equations applied to each grid point independently or are they averaged over, e.g., latitude zones?

As stated in section 2.3, these equations are used at every timestep. We assess these cloud radiative feedbacks ($\alpha_{\text{atm}}FB$ and $OLW_{\text{cloud}}FB$) over a 12-month climatology to incorporate any seasonality in the feedbacks (e.g. monsoon impacts, etc). In addition, $\alpha_{\text{atm}}FB$ and $OLW_{\text{cloud}}FB$ are applied at each grid cell to incorporate the spatial patterns in the cloud feedbacks that are unique to each source GCM. We have attempted to clarify this in our revised manuscript.

4) Page 11, line 4: Why do you write $F_{2\times\text{CO}_2} = F_{4\times\text{CO}_2}/2$ when there is a logarithmic relation between radiative fluxes and the CO_2 concentration? Is this close to linear because the absolute change is very small?

Yes, there is a logarithmic relationship between CO_2 radiative forcing and concentration. In UVic, it looks like this:

$$F_{\text{CO}_2} = \text{CO}_2\text{FOR} * \ln \frac{[\text{CO}_2 (\text{ppm})]}{280 \text{ ppm}},$$

where CO_2FOR is the CO_2 radiative forcing term (5.35 W m^{-2}), equivalent to 3.71 W m^{-2} for a doubling of CO_2 .

The forcing of $4\times\text{CO}_2$ (1120 ppm) is mathematically equivalent to $2\times$ the forcing of a doubling of CO_2 (560 ppm). Therefore, $F_{2\times\text{CO}_2} = F_{4\times\text{CO}_2}/2$

A Cloud Feedback Emulator (CFE, version 1.0) for an intermediate complexity model ~~The University of Victoria Cloud Feedback Emulator (UVic-CFE): cloud radiative feedbacks in an intermediate complexity model~~

5 Ullman, D.J.^{1,*} and Schmittner, A.¹

¹College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, USA

*Now at Northland College, Ashland, WI, USA

Correspondence to: David J. Ullman (dullman@northland.edu)

Abstract. The dominant source of inter-model differences in comprehensive global climate models (GCMs) are cloud radiative effects on Earth's energy budget. Intermediate complexity models, while able to run more efficiently, often lack cloud feedbacks. Here, we describe and evaluate a method for applying GCM-derived shortwave and longwave cloud feedbacks from 4xCO₂ and Last Glacial Maximum experiments to the University of Victoria Earth System Climate Model. The method generally captures the spread in top-of-the-atmosphere radiative feedbacks between the original GCMs, which impacts the magnitude and spatial distribution of surface temperature changes and climate sensitivity. These results suggest that the method is suitable to incorporate multi-model cloud feedback uncertainties in ensemble simulations with a single intermediate complexity model.

10
15

1 Introduction

The predominant trade-off in climate modeling is that of systematic complexity versus computational expense. While comprehensive global climate models (GCMs) attempt to resolve the complex interactions between earth systems, their computational expense limits the exploration of parametric uncertainty. Conversely, more simplified models, such as earth system models of intermediate complexity (EMICs), can be employed for large-ensemble analysis of parametric variability but their reliance on fixed boundary conditions or generalized parameterizations of earth processes may not capture all important feedbacks driving system dynamics.

20

One of the largest sources of intermodel spread in GCM-based climate projections is the magnitude and direction of radiative cloud feedbacks (Soden and Held, 2006; Dufrense and Bony, 2008; Tomassini et al., 2013; Vial et al., 2013). Clouds affect climate through their impacts on both shortwave (solar radiation mostly in the visible part of the spectrum) and longwave (terrestrial, infrared radiation) fluxes and therefore determine the sensitivity of GCMs to changes in radiative forcing (Andrews et al., 2012; Sherwood et al., 2014). Because clouds are more reflective than most other surfaces, an increase in clouds will reduce the amount of shortwave energy absorbed by the Earth and lead to cooling. Conversely, clouds

25

ability to absorb upward longwave fluxes and re-radiate them back down causes warming at the surface (Hartmann and Short, 1980). The relative magnitude and net effect of these feedbacks depends on cloud altitude. For low clouds, which radiate longwave fluxes at a similar temperature as the surface, shortwave effects dominate and their net effect is cooling. High clouds, on the other hand, radiate at much colder temperatures than the surface, which can make the longwave effect dominate and lead to net warming (Hartmann et al., 1992). However, the cloud liquid water content and associated optical depth of high and low clouds likely also plays a role in the absorption and reflection of incoming shortwave fluxes (Tselioudis et al., 1992). Therefore, the net cloud feedback may be positive or negative feedback depending on whether low versus high cloud cover and cloud optical depth responds more to local and global temperature change. For state of the art GCMs, the spread in GCM cloud feedbacks is primarily driven by model differences in low cloud cover changes (Sherwood et al., 2014). In addition, the spread in GCM cloud feedbacks manifests in both the global mean as well as regional variability (Tomassini et al., 2013; Vial et al., 2013). This spatial variability likely has a profound impact on the magnitude of climate response to perturbations (Marvel et al., 2015).

Since EMICs use simplified atmospheric components, the cloud radiative forcing is typically fixed (Plattner et al., 2001; Joos et al., 2001; Driesschaert, 2005; Crucifix et al., 2002; Weaver et al., 2001). Therefore, the uncertainties in cloud feedbacks demonstrated in GCMs are typically neglected in the non-interactive cloud schemes of EMICs. Schmittner et al. (2011), e.g., hypothesized that their estimate of climate sensitivity, determined using the University of Victoria (UVic) EMIC and paleoclimate observations, resulted in a too narrow probability distribution due to the neglect of cloud feedback uncertainties. Here we describe and evaluate a new method for diagnosing and applying cloud feedbacks of state-of-the-art GCMs into an EMIC, thereby creating a computationally less-expensive emulator of more complex models.

20 **2 Methods**

2.1 Model Description

The UVic Earth System Climate Model (Weaver et al., 2001) is an EMIC with a 3-dimensional ocean general circulation model coupled to a dynamic-thermodynamic sea ice model, a 2-dimensional single-layer energy-moisture balance atmosphere, and a dynamic land (Meissner et al., 2003) and vegetation model (Cox, 2001). Surface wind speeds used in the calculations of air-sea exchange and atmospheric transport of heat and moisture are prescribed in the model, thereby limiting variability in the atmospheric model. The model conserves heat and moisture without the need for a flux correction (Weaver et al., 2001). We employ version 2.9 of UVic (Eby et al., 2013), in which atmospheric heat diffusion varies with changes in global mean surface air temperature; this modification has been shown to improve the latitudinal temperature gradient for the Last Glacial Maximum when compared with high-latitude proxy data (Fyke and Eby, 2012). All model components have a horizontal grid resolution of 1.8° latitude by 3.6° longitude, with 19 vertical levels in the ocean model increasing from 50 m thickness in the surface level to 590 m thickness in the deepest gridcell.

The net radiative balance (NETRAD) at the top of the atmosphere (TOA) is the difference between the net shortwave radiation (SW_{TOA}) and the outgoing longwave radiation (LW_{TOA}):

$$NETRAD = SW_{TOA} - LW_{TOA} \quad (1)$$

5

Clouds impact SW_{TOA} through prescribed monthly fields of atmospheric albedo (α_{atm}):

$$SW_{TOA} = SW_{in,TOA} - SW_{in,TOA} \cdot \alpha_{atm} - SW_{in,TOA} \cdot (1 - \alpha_{atm}) \cdot \alpha_{sfc} \cdot \tau^2 \quad (2)$$

10 where S is the flux of incoming (incident) solar radiation energy per unit area ($W\ m^{-2}$) at the top of the atmosphere (with seasonal and latitudinal variation), τ is a constant atmospheric transmission coefficient (0.77), and α_{sfc} is the surface albedo. The second term of Eq. (2) represents the proportion of incoming SW radiation that is immediately reflected by clouds, while the third term represents the portion that is reflected by the surface, which passes through the atmosphere twice. During its first (downward) pass $S \cdot (1 - \alpha_{atm}) \cdot (1 - \tau)$ is absorbed by the atmosphere. During its second (upward) pass,
 15 $S \cdot (1 - \alpha_{atm}) \cdot (1 - \tau) \cdot \alpha_{sfc} \cdot (1 - \tau)$ is absorbed. All variables except for τ vary over space and time, but while α_{sfc} is allowed to evolve with changes in surface model components (sea ice, snow cover, vegetation, etc.), α_{atm} is a fixed boundary condition at monthly resolution to resolve seasonal changes in regional cloud cover. In the control version of UVic, α_{atm} is estimated with the following relationship:

$$20 \quad \alpha_{atm} = \frac{f \cdot \alpha_{plt} - \alpha_{sfc}}{1 - \alpha_{sfc} \cdot \tau^2} \quad (3)$$

where

$$\alpha_{plt} = \frac{SW_{out,TOA}}{SW_{in,TOA}} \quad (4)$$

25

$$\alpha_{sfc} = \frac{SW_{out,sfc}}{SW_{in,sfc}} \quad (5)$$

where the planetary (α_{plt}) and surface albedo (α_{sfc}) are calculated using the incoming and outgoing shortwave satellite observational measurements at the surface and top of the atmosphere from the Earth Radiation Budget Experiment (ERBE;
 30 Barkstrom, 1984; Barkstrom and Smith, 1986; Ramanathan et al., 1989). This α_{atm} relationship is directly derived from Eq. (2) so as to be internally consistent with the radiative balance relationship from the UVic model. The variable f in Eq. (3) is a

constant planetary albedo adjustment factor to account for radiative imbalances that arise in the implementation of the derived α_{atm} .

The outgoing longwave radiation (*OLW*) is parameterized in UVic using an empirical relationship (Thompson and Warren, 1982; Weaver et al., 2001) that ~~depends~~ determines clear-sky *OLW* as a function of on surface relative humidity (*rh*) and temperature (*SAT*):

$$\begin{aligned} OLW = & c_{00} + c_{01}rh + c_{02}rh^2 + (c_{10} + c_{11}rh + c_{12}rh^2)SAT \\ & + (c_{20} + c_{21}rh + c_{22}rh^2)SAT^2 + (c_{30} + c_{31}rh + c_{32}rh^2)SAT^3 \\ & + \Delta F_{2xCO_2} \ln \frac{[CO_2]_t}{[CO_2]_o} \end{aligned} \quad (6)$$

where the final term adjusts *OLW* for a change in the atmospheric CO₂ concentrations. The value of $\Delta F_{2xCO_2} = 5.35 \text{ W m}^{-2}$ is selected as the radiative forcing associated with 3.71 W m^{-2} (IPCC, 2001). The constants (c_{xx}) are provided by Thompson and Warren (1982). Since this was originally estimated as a clear-sky is an empirical relationship, the effect of clouds on the *OLW* radiative balance is not explicitly ~~but implicitly~~ included.

2.2 CERES update to Atmospheric Albedo boundary conditions

Because of discontinuities in satellite coverage, missing data, and poor resolution, the Clouds and the Earth's Radiant Energy System (CERES, Wielicki et al., 1996) was launched in late 1999 to better observe the earth's radiative balance (Fasullo and Trenberth, 2008). The CERES experiment uses an updated satellite architecture and provides higher spatial resolution observations over a longer time domain (2000-2013 for CERES compared with 1985-1989 for ERBE), thereby providing more robust modern climatology on the impact of clouds on atmospheric albedo (Wielicki et al., 1996). In addition, the duration of the ERBE experiment between 1985 to 1989 spans a somewhat large El Niño event (1987), which may bias the equatorial Pacific toward enhanced cloudiness in the calculation of atmospheric albedo climatology using the ERBE data (Cess et al., 2001).

In this paper, we use the climatology (2000-2013) of CERES surface and top of the atmosphere shortwave fluxes to better estimate α_{atm} boundary conditions in UVic (using Eq. (3)). Under low-light conditions (winter, high-latitudes), satellite-derived estimates of incoming SW are small, which occasionally results in values of α_{plt} and α_{sfc} that are greater than 1. Therefore, we limit α_{plt} and α_{sfc} to values less 1, which ensures that α_{atm} is within appropriate limits.

An ensemble of control simulations was performed using the new CERES-based estimates of α_{atm} with varying values of the *f* parameter in Eq. (3). From the resulting equilibrium simulations, a value of $f=0.95$ in Eq. (3) was selected in order to match 20th century global mean temperature data estimates of $\sim 13.9 \text{ }^\circ\text{C}$ (NOAA, 2016) in a UVic control simulation.

The-This final estimate of CERES-based α_{atm} ~~is~~ was smoothed and regridded to the UVic grid.

Figure 1 compares the annual mean values of α_{atm} as derived from the ERBE and CERES datasets. In the tropics, the ERBE-based estimates of α_{atm} generally match those of the CERES-based values (Figure 1). In the high latitudes, however, the ERBE-based α_{atm} values are generally higher than the CERES-based values. Such differences are likely related to improvement in sampling orbit of the CERES satellite and the associated reduction in zenith angle-dependent biases, which may result in large errors in the top of the atmosphere flux measurements in the ERBE data (Loeb et al., 2009). As such, the use of CERES-based estimates of α_{atm} provides an improvement in UVic, particularly at high latitudes.

2.3 Innovations

With the use of CERES-based α_{atm} estimates, the UVic model now includes an updated effect of clouds on the Earth's shortwave radiative balance. However, the control UVic model design does not incorporate any change in the shortwave or longwave radiative effect of clouds due to changes in temperature. This lack of cloud feedbacks may significantly limit the ability of UVic to capture global temperature in perturbed simulations. Here, we provide a simple method of diagnosing cloud radiative forcings from GCM results of the Coupled Model Intercomparison Project 5 (CMIP5) and Paleoclimate Model Intercomparison Project 3 (PMIP3) archives (Braconnot et al., 2011; Taylor et al., 2012) and incorporate the associated shortwave and longwave cloud feedbacks into UVic for both 4 times CO₂ (4xCO₂) and Last Glacial Maximum (LGM) climate simulations. Reanalysis of satellite observations suggests that the range of CMIP5 models present widespread agreement with cloud data, both in spatial extent and vertical distribution, across the historical record (Norris et al., 2016). We have selected model output from 7 GCMs: CCSM4 (abbreviated as CCSM), CNRM-CM5 (CNRM), GISS-E2-R (GISS), IPSL-CM5A-LR (IPSL), MIROC-ESM (MIROC), MPI-ESM-P (MPI), and MRI-CGCM3 (MRI). These models were chosen because they have results for both 4xCO₂ and LGM simulations and all of the relevant variables for calculating shortwave and longwave cloud feedbacks (see below). The following innovations demonstrate how we employ UVic as a cloud feedback emulator (~~UVic~~-CFE [version 1.0](#); henceforth CFE) of the full AOGCMs.

2.3.1 Shortwave Cloud Feedbacks in UVic

Since UVic incorporates the shortwave impact of clouds through atmospheric albedo, we assess the shortwave cloud feedback as the change in α_{atm} due to the change in temperature in each of the GCM simulations. Albedo anomalies are not mathematically additive; therefore, we first calculate α_{atm} for each perturbed state (4xCO₂, LGM) by adding GCM anomalies of each of the individual fluxes to the CERES observations:

$$SW_{in,TOA,GCM} = (SW_{in,TOA,perturbed} - SW_{in,TOA,control}) + SW_{in,TOA,CERES} \quad (7)$$

$$SW_{out,TOA,GCM} = (SW_{out,TOA,perturbed} - SW_{out,TOA,control}) + SW_{out,TOA,CERES} \quad (8)$$

$$SW_{in,sfc,GCM} = (SW_{in,sfc,perturbed} - SW_{in,sfc,control}) + SW_{in,sfc,CERES} \quad (9)$$

$$SW_{out,sfc,GCM} = (SW_{out,sfc,perturbed} - SW_{out,sfc,control}) + SW_{out,sfc,CERES} \quad (10)$$

5 For each of the variables, we have calculated a 12-month climatology (separate averaging for each month) that is assessed over the final 10 years of the 150 year transient 4xCO₂ simulations, the final 100 years of the LGM equilibrium simulations, and the final 100 years of the equilibrium control simulations. The anomaly-perturbed values of each of the shortwave fluxes (Eq. (7)-(10)) are then used to calculate an $\alpha_{atm,perturbed}$ for each of the perturbed GCM simulations using Eq. (3)-(5).

Again, because albedo values are not additive, we calculate the albedo anomaly as the ratio of the atmospheric
10 albedo of the GCM perturbed state to CERES-derived atmospheric albedo. Therefore, the α_{atm} feedback ($\alpha_{atm}FB$) is this albedo anomaly divided by the change in temperature:

$$\alpha_{atm}FB = \frac{[\alpha_{atm,perturbed}/\alpha_{atm,CERES}] - 1}{SAT_{perturbed} - SAT_{control}} \quad (11)$$

15 The subtraction of 1 in the numerator is necessary such that when there is no change in α_{atm} ($\alpha_{atm,perturbed} = \alpha_{atm,CERES}$), then there is no atmospheric albedo feedback. This α_{atm} feedback is calculated as a 12-month climatology at each gridcell for each of the 7 GCMs that are sampled in this analysis (Figure 2, 3). Positive (negative) values for this atmospheric albedo feedback indicate a negative (positive) shortwave cloud feedback since increases in temperature cause an increase (decrease) in atmospheric albedo, which cools (warms) the surface. The magnitude of these atmospheric albedo feedbacks varies
20 considerably among the GCMs and between perturbed climate states (4xCO₂ versus LGM), which is consistent with the large spread in cloud shortwave feedbacks found in previous studies (Tomassini et al., 2013; Vial et al., 2013). For example, GISS-E2-R shows a strongly positive atmospheric albedo feedback from the 4xCO₂ results, while IPSL-CM5A-LR generally shows a strongly negative atmospheric albedo feedback, particularly in the tropics (Figure 2).

The innovation to UVic is the application of these GCM-diagnosed α_{atm} feedbacks to the shortwave radiative
25 balance. First, we calculate a SAT climatology from a long-term control simulation of UVic that uses $\alpha_{atm,CERES}$ as the control atmospheric albedo. Then at each timestep (t) of a model simulation, we calculate the difference in surface air temperature from this control monthly-climatology, and perturb atmosphere albedo at each gridcell using the GCM-derived $\alpha_{atm}FB$ of Eq. (11):

$$30 \quad \alpha_{atm}(t) = [\alpha_{atm}FB \cdot [SAT(t) - SAT_{ctl}] + 1] \cdot \alpha_{atm,CERES} \quad (12)$$

The above calculation is done at every timestep and each gridcell allowing for spatially- and seasonallymonthly-specific atmospheric albedo feedbacks as diagnosed from the GCMs.

2.3.2 Shortwave-Longwave Cloud Feedbacks in UVic

Because UVic lacks a longwave cloud feedback in the calculation of OLW , we provide an additional term to Eq. (6), which now includes the OLW due to changes in the cloud longwave effect in the GCM simulations. First, we diagnose the outgoing longwave radiation at the top of the atmosphere from the GCM output:

$$OLW_{cloud} = OLW_{total} - OLW_{clearsky} \quad (13)$$

The outgoing longwave cloud feedback is therefore the cloud longwave forcing anomaly divided by temperature anomaly:

$$OLW_{cloud}FB = \frac{OLW_{cloud,perturbed} - OLW_{cloud,control}}{SAT_{perturbed} - SAT_{control}} \quad (14)$$

as diagnosed from results of the GCM perturbed simulations. These outgoing longwave cloud feedbacks are calculated as monthly climatologies at each gridcell, and are assessed separately for both the 4xCO₂ and LGM perturbed states (Figure 4, 5). Again, $OLW_{cloud}FB$ values are assessed using the 12-month climatologies assessed over the final 10 years of the 150-year transient 4xCO₂ simulations, the final 100 years of the LGM equilibrium simulations, and the final 100 years of the equilibrium control simulations. We note that by calculating the OLW cloud radiative effect using the total OLW minus clear-sky OLW (Eq. 13), we are implicitly including the effects of cloud masking and rapid cloud adjustments (Zelinka et al., 2013). Including both of these effects has been shown to reduce both LW and SW cloud feedbacks relative to a more explicit cloud radiative kernel method (Zelinka et al., 2012; Zelinka et al., 2013). Both effects may limit the magnitude of the total cloud feedback.

Most models show more areas of positive $OLW_{cloud}FB$. This indicates a negative climate feedback since increasing temperatures lead to more OLW , which cools the surface. Again, the outgoing longwave cloud feedbacks vary considerable between models and climate state. The largest variability in OLW cloud feedbacks between models exists in the tropics, which is consistent with prior results suggesting that model differences in convective mixing and resulting cloud height greatly impacts the magnitude and direction of cloud feedbacks (Sherwood et al., 2014). Generally, the OLW cloud feedback is stronger in magnitude for the LGM state (Figure 5) than for the 4xCO₂ state.

Similar to the inclusion of the atmospheric albedo feedbacks in UVic, we multiply the outgoing longwave cloud feedback by the temperature difference from the long-term control UVic simulation:

$$OLW_{cloud}(t) = OLW_{cloud}FB \cdot [SAT(t) - SAT_{ctl}] \quad (15)$$

This OLW_{cloud} term is calculated at each timestep [and gridcell](#) in the model and is added to the OLW parameterization (Eq. (6)) as an additional cloud longwave feedback term.

2.4 Numerical Experiments

5 To estimate how well our [UVic-CFECFE](#) cloud feedback emulator ([UVic-CFECFE](#)) captures the original cloud radiative effects from the GCMs, we present an ensemble of [UVic-CFECFE](#) control and perturbed experiments (4xCO₂ and LGM) that use the α_{atm} and OLW_{cloud} feedbacks diagnosed from each of the 7 GCMs employed in this analysis. Because our diagnosed cloud feedbacks differ between the 4xCO₂ and LGM climate states (Figures 2-5), we ran 2 separate preindustrial control simulations for each ensemble member: one with 4xCO₂ cloud feedbacks (ctl4x) and one with LGM cloud feedbacks
10 (ctlLGM). Indeed, the inclusion of these cloud feedbacks in the control climate state leads to slight differences in control global mean temperature, indicating that separate controls are necessary in the calculation of resulting radiative feedbacks. Therefore, we present the results from 28 separate [UVic-CFECFE](#) simulations: 4 simulations (ctl4x, ctlLGM, 4xCO₂, LGM) for each of the 7 GCM-derived cloud feedbacks.

Preindustrial control and LGM simulations with each of the GCM-derived cloud feedbacks were run to extended
15 equilibrium (>2000 years) to be certain of minimal model drift (global mean SAT trend < 0.04 °C per 100 years). Both 4xCO₂ and LGM simulations follow the CMIP5/PMIP3 protocol (Braconnot et al., 2011; Taylor et al., 2012) as closely as possible as these are the boundary conditions used in the original GCM simulations. Our 4xCO₂ simulations using modern boundary conditions, an instantaneous increase in atmospheric CO₂ concentration to 1120 ppm, and a simulation length of 150 years, starting from the end of the preindustrial control simulation (ctl4x). Our LGM simulations have reduced
20 greenhouse gas concentrations (atmospheric CO₂ = 185 ppm; radiative forcing adjusted for appropriate CH₄/N₂O concentrations; Schmittner et al., 2011), altered orbital state, full glacial ice sheet extent/topography (Peltier, 2004), modified river pathways, and +1 PSU (Practical Salinity Unit) increase in mean ocean salinity. In addition, we apply LGM surface wind stress anomalies that are diagnosed from the LGM GCM results (Muglia and Schmittner, 2015). [Wind stress anomalies at the end of the CMIP5 4xCO₂ simulations are small; therefore, we use the prescribed wind stress fields of the control UVic
25 \[2.9 model \\(from NCEP reanalysis\\) in our 4xCO₂ simulations.\]\(#\)](#)

3 Results

3.1 Assessment of GCM-diagnosed cloud feedbacks

Across the historical record with a warming climate, the cloud trends in CMIP5 models have been shown to be in agreement with satellite observations, with robust reductions in cloudiness across the mid-latitude and tropics, as well as an increase in
30 cloud top height at all latitudes (Norris et al., 2016). Our calculated 4xCO₂ atmospheric albedo feedbacks are consistent with

these observations, generally showing a reduction in α_{atm} in the mid-latitudes and tropics (Figure 2). Only one model (GISS) shows an increase in α_{atm} across the 4xCO₂ simulations. Most of the 4xCO₂ GCM-diagnosed α_{atm} feedbacks seem to suggest an increase in α_{atm} in the high-latitudes with warming (particularly over the Southern Ocean), which is likely related to a poleward shift in the storm tracks due to warming (Lu et al., 2007; Norris et al., 2016).

5 The 4xCO₂ GCM-derived OLW_{cloud} feedbacks are also most prominent in the tropics with considerable variability in the location, magnitude and direction of peak feedback (Fig. 4). However, all models show a negative OLW_{cloud} feedback across the equatorial Pacific and a positive OLW_{cloud} feedback over the Indonesia Archipelago, South America and off the equator. Outside of the tropics, most models show positive OLW_{cloud} feedbacks in the mid-latitudes and slight negative feedbacks in the polar regions. These data are consistent with observations of increased cloud top height (Norris et al., 2016),
10 as regions with enhanced cloudiness (increased α_{atm} , Figure 2) also typically show decreased OLW (Figure 4).

For the LGM, GCM-derived cloud feedbacks are less coherent. Nearly all models show large changes in the tropical α_{atm} feedback, particularly across the equatorial Pacific and Indonesian Archipelago (Figure 3). Such changes may be suggestive of changes in the position of the Intertropical Convergence Zone (ITCZ) associated changes in deep convective cloud systems that are specific to each model (Braconnot et al., 2007; Arbuszewski et al., 2013). In addition, nearly all
15 GCM-derived feedbacks show a reduction in α_{atm} over the North Atlantic (note that LGM cooling indicates that direction of feedback change is opposite that shown in Figure 3), which may be indicative of a shift in the position of the Gulf Stream seen in some models (Otto-Bliesner et al., 2006). The prominent feature in the LGM GCM-derived OLW_{cloud} feedback is a large reduction in the tropics (green-blue-purple colors in Figure 5), which is likely related to the reduction in tropical convection due to lower sea surface temperatures (Yin and Battisti, 2001). However, this spatial extent and magnitude of
20 reduction in OLW_{cloud} for the LGM vary appreciably among the GCMs.

3.2 Radiative balance in ~~UVie-CFECFE~~ 4xCO₂ simulation

To compare the global radiative balance of ~~UVie-CFECFE~~ with that of the GCMs, we calculate the total change in TOA shortwave and longwave fluxes per global mean surface temperature change from the final 10 years of the 150-year 4xCO₂ simulations (relative to the control simulation) and compare the raw GCM results with our cloud feedback-forced ~~UVie-CFECFE~~ simulations (Figure 6). The changes in longwave fluxes include the CO₂ forcing, which may differ by ~15%
25 between models (Andrews et al., 2012). Because the forcing is included in the longwave fluxes, the flux/temperature ratios shown in Fig. 6 are not a true “feedback,” strictly speaking; therefore, we use the term “radiative-temperature response.” However, variations in the forcings are presumably relatively small compared to variations in feedbacks. The shortwave flux/temperature ratios in Fig. 6 are true feedbacks and consistent with numbers reported previously (Tomassini et al., 2013).
30

In general, the spread of TOA shortwave and longwave ~~feedbacks~~ radiative-temperature response in the 4xCO₂ ~~UVie-CFECFE~~ simulations matches that of the original GCM results (Figure 6) and is consistent with previous work

(Tomassini et al. 2013). For instance, the IPSL model exhibits the largest positive shortwave ~~feedback~~ and largest negative longwave ~~feedback~~ radiative-temperature response in the GCM results, which is also captured in our UVie-CFECFE simulations (Figure 2, 4). Conversely, the GISS model is the only simulation to show a negative shortwave ~~feedback~~ and positive longwave ~~feedback~~ radiative temperature response, which is consistent with the UVie-CFECFE results. All other GCM and UVie-CFECFE simulations have positive shortwave ~~feedbacks~~ and negative longwave ~~feedbacks~~ radiative-temperature response that are both smaller in magnitude than the IPSL-based simulations.

While the relative magnitude of the UVie-CFECFE radiative-temperature response feedback results captures that of the original GCM results, the absolute magnitude of the radiative-temperature response feedbacks is generally slightly reduced in UVie-CFECFE. We also present the results from a control 4xCO₂ UVic simulation, without the implementation of any cloud feedbacks (grey bar, Figure 6). Here, the TOA shortwave radiative-temperature response feedback is ~0.40 W m⁻² °C⁻¹ and the TOA longwave radiative-temperature response feedback is ~-0.03 W m⁻² °C⁻¹, while the average radiative-temperature response feedbacks from the GCMs are ~0.87 W m⁻² °C⁻¹ and ~-0.55 W m⁻² °C⁻¹, respectively. Therefore, the application of α_{atm} and OLW-cloud feedbacks in UVie-CFECFE are prominent drivers in the spread of total TOA shortwave and longwave radiative-temperature response feedbacks. In general, the GCMs show a greater reduction in global surface albedo with increasing temperature compared to the UVie-CFECFE (not shown). Therefore, the differences in surface albedo processes between the GCMs and UVie-CFECFE, likely explains some of the reduction in TOA shortwave radiative-temperature response feedback magnitude in the UVie-CFECFE simulations.

3.3 Radiative balance in UVie-CFECFE LGM simulations

For the UVie-CFECFE LGM simulations, we calculate TOA shortwave and longwave radiative-temperature response feedbacks at equilibrium conditions, averaged over the last 100 years of the LGM and ctlLGM experiments. Note that in this case the shortwave fluxes include forcing from prescribed ice sheets and therefore are not strictly speaking feedbacks. UVie-CFECFE generally captures the spread of the shortwave and longwave radiative-temperature response feedbacks from the GCMs although it is slightly reduced (Figure 6). The total imbalance seems to be smaller in UVie-CFECFE compared with most GCMs indicating that UVie-CFECFE is closer to equilibrium, perhaps because it was integrated longer. Thus a larger remaining imbalance could contribute to the larger spread in the GCMs compared with UVie-CFECFE.

The absolute magnitude of the radiative-temperature response feedbacks is mostly reduced in the UVie-CFECFE relative to the GCM simulations. Similar to the 4xCO₂ results, the IPSL-based simulations present the strongest shortwave and longwave radiative-temperature response feedbacks. Conversely, the CNRM-based UVie-CFECFE simulation shows enhanced shortwave and longwave radiative-temperature response feedbacks relative to those of the GCM, suggesting that non-cloud processes or differences in the forcings are likely important for this model.

3.4 Effect of ~~UVic-CFECFE~~ on modeled temperature evolution and spatial distribution

As expected, the incorporation of cloud feedbacks into ~~UVic-CFECFE~~ has a direct impact on modeled surface temperature anomalies in perturbed experiments. For the 4xCO₂ experiments, global mean surface air temperature anomalies at the end of the 150-year simulation range from +3.9 °C (GISS) to +8.8 °C (IPSL), where the control UVic simulation without cloud feedbacks results in a final anomaly of +5.1°C (Figure 7). Only two ~~UVic-CFECFE~~ simulations (GISS and MRI) result in a year 150 temperature anomaly that is less than the UVic control, confirming that the 4xCO₂ net cloud feedbacks are generally positive (see above) and consistent with the analysis of the individual models themselves (Vial et al., 2013; Tomassini et al., 2013).

The spatial variability in GCM cloud feedbacks (Figure 2, 4) is also expressed in the 4xCO₂ zonal mean temperature anomalies (Figure 7). All models show the effects of strong polar amplification by the end of the 4xCO₂ simulations, but the addition of cloud feedbacks to ~~UVic-CFECFE~~ appears to enhance this polar amplification in most cases. In addition, the change in temperature due to cloud feedbacks is not uniform for all models. For example, the CCSM-driven simulation presents some of the largest temperature anomalies in the southern high-latitudes but relatively reduced anomalies at the low-latitudes, resulting in an overall global anomaly that is similar to the that of the control UVic simulation (Figure 7).

For the LGM simulations, the global mean temperature change at the end of the simulation ranges from -4.1 °C (CCSM) to -8.2 °C (CNRM), whereas the control UVic simulation has a cooling of 5.7°C (Figure 7). Nearly half of the UVic simulations show enhanced global mean cooling (CNRM, IPSL, and MRI) relative to the UVic control (Figure 7), while the other four simulations show reduced cooling (CCSM, GISS, MIROC, and MPI). Again, zonal mean temperature anomalies at the LGM show that enhanced cloud feedbacks lead to enhanced polar amplification, but spatial differences in the magnitude of feedbacks may impact regional temperature change. For example, the CNRM-based simulation shows the strongest cooling in the southern high latitude, whereas the IPSL-based simulation has the largest cooling in the northern high latitudes (Figure 7).

3.5 Using ~~UVic-CFECFE~~ to estimate climate sensitivity

Intermodel spread in GCM cloud feedbacks has been shown to have a large impact on the modelled sensitivity to perturbation in greenhouse gas radiative forcing (Fasullo and Trenberth, 2012; Andrews et al., 2012; Sherwood et al., 2014). To estimate the effect of the cloud feedbacks in ~~UVic-CFECFE~~ on global climate, we calculate effective equilibrium climate sensitivity ($\Delta T_{2xCO_2,eff}$) from the 150-year 4xCO₂ simulations by regressing the global net downward heat flux at the TOA onto the change in temperature. The slope of this regression is the climate response parameter (α) and the intercept is the 4xCO₂ forcing (F_{4xCO_2}) specific to each model (Gregory et al., 2004). These values can be used to estimate the effective equilibrium climate sensitivity to a doubling of CO₂ by dividing the implied global 2xCO₂ forcing ($F_{2xCO_2} = F_{4xCO_2}/2$) by α (Gregory et al., 2004). We calculate $\Delta T_{2xCO_2,eff}$ for both the raw GCM model output as well as the associated ~~UVic-CFECFE~~ simulations.

With the introduction of cloud feedbacks, UVie-CFECFE is able to capture much of the intermodel variability in climate sensitivity (Figure 8). The seven GCMs sampled in this analysis show values of $\Delta T_{2xCO_2,eff}$ ranging from 2.15 °C (GISS) to 4.10 °C (IPSL), which agrees well with Andrews et al. (2012) for those models that were used in both studies. In the UVie-CFECFE simulations, $\Delta T_{2xCO_2,eff}$ values range from 2.34 °C (GISS) to 7.00 °C (IPSL). Again, the IPSL-based UVie-CFECFE simulation is a noticeable outlier, while all of the values of $\Delta T_{2xCO_2,eff}$ in UVie-CFECFE are more comparable to the values from the raw GCM output and the magnitude relative to each of the models is generally the same (Figure 8). However, most of the UVie-CFECFE simulations show elevated $\Delta T_{2xCO_2,eff}$ relative to their GCM counterpart (Figure 8). The $\Delta T_{2xCO_2,eff}$ in the 4xCO₂ control UVic simulation (grey bar, figure 8) is 3.63 °C, a value that is higher than most of the GCM results, suggesting that the control UVic climate sensitivity without explicit cloud feedbacks may already be higher than that of most of the sampled GCMs. This suggests that the control UVic model's clear sky (without explicit clouds) feedbacks are larger than those of most GCMs. Adding the mostly positive cloud feedbacks thus makes the UVic model's climate sensitivities considerably larger than those of the GCMs. Clear-sky feedbacks in the UVic model could be tuned by e.g. varying the coefficients of eq. (6) if a better match with individual GCM's climate sensitivity was desired.

4 Discussion and Conclusions

The cloud feedbacks (α_{atm} and OLW_{cloud} feedbacks) derived from the GCMs and employed in UVie-CFECFE are generally consistent between climate states (4xCO₂ vs LGM) for each GCM, with some notable exceptions. For example, the 4xCO₂ α_{atm} feedbacks (Figure 2) are generally consistent between models in showing a prominent negative feedback across the southern ocean, with CCSM being the only model with a positive α_{atm} feedback. However, for the LGM, the CCSM-derived α_{atm} feedback is negative along with all other models in general (Figure 3). In addition, the α_{atm} feedbacks across the equatorial Pacific are not always consistent between climate states, with the CNRM-, GISS-, MIROC-, and MPI-based fields showing a pronounced difference in the direction of the α_{atm} feedback (Figure 2, 3). Similarly, the OLW_{cloud} feedbacks across the equatorial Pacific and North Pacific differ in magnitude and direction between the climate states in nearly all models (Figure 4, 5). These differences likely arise due to shifts in the ITCZ and Gulf Stream between climate states (Otto-Bliesner et al., 2006; Braconnot et al., 2007; Arbuszewski et al., 2013), and they suggest that such cloud feedbacks are not universal to all climate states. As such, the cloud feedbacks derived from the GCMs should only be applied to a consistent climate state experiment when using UVie-CFECFE.

In general, the application of GCM-derived cloud feedbacks to UVie-CFECFE captures the changes in TOA radiative balance of the original GCMs, for both the 4xCO₂ and LGM experiments. Differences in total radiative feedbacks between each GCM and the associated UVie-CFECFE may exist for several reasons. First, the derivation of the cloud feedbacks are parameterized from the original GCM results and therefore may not be a perfect representation of the full complexity of cloud radiative forcing in each GCM. This is particularly the case for the shortwave cloud feedback, which is

applied using a calculation of the α_{atm} feedback, which uses an assumption of a global mean atmospheric transmissivity (Eq. (3)). The OLW-cloud feedbacks, on the other hand, are a direct calculation of the longwave cloud feedbacks from each GCM.

Second, total TOA feedbacks in [UVic-CFECFE](#) may not perfectly match those of the source GCMs because the resulting feedbacks are still partly controlled by the control radiative balance code of the UVic model. Other components of the Earth system, apart from clouds, impact the shortwave and longwave radiative balance in UVic, which may feedback on the simulated climate in a different manner than in the GCMs. For instance, the total TOA shortwave feedbacks include the effect of surface albedo change. Therefore, differences in vegetation and sea ice dynamics and their effect on surface albedo in the GCMs relative to UVic may help explain some of the differences in the shortwave feedbacks. Similarly, the longwave feedback in UVic is in part controlled by the SAT-based parameterization of OLW in Eq. 6, which may be different from the clear-sky feedbacks in the GCMs

Third, the ratios of TOA flux and temperature changes shown in [Figure- 6](#) include forcings (greenhouse gas for both 4xCO₂ and LGM and surface albedo for LGM). Therefore, differences in the forcings would also impact the total TOA “feedbacks”. The forcings differ between the GCMs but are constant among the [UVic-CFECFE](#) experiments. [In addition, our method of estimating cloud feedbacks neglects the effects of cloud masking and cloud rapid adjustment \(Zelinka et al., 2013\),](#) which may explain some of the loss of spread in [UVic-CFECFE](#) compared with the GCMs.

However, despite the potential for differences in total radiative feedbacks, our results suggest that a simple parameterization of cloud shortwave and longwave feedbacks may be applied to UVic to generally capture dominant inter-model spread in total radiative feedbacks. This result confirms that cloud feedbacks dominate the multi-model uncertainty in GCM radiative balance (Soden and Held, 2006; Dufrense and Bony, 2008; Tomassini et al., 2013; Vial et al., 2013). The addition of GCM-derived cloud feedbacks to the UVic leads to only small increases in computational expense, while capturing an important component of the Earth’s radiative balance that is otherwise lacking in the default UVic model. Indeed, the inclusion of cloud feedbacks leads to a large spread in surface air temperature anomalies for both the 4xCO₂ and LGM experiments (Figure 7). In addition, spatial variability in the cloud feedbacks (Figure 2-5) leads to some differences in the latitudinal distribution of this temperature change (Figure 7), suggesting that certain regional cloud changes may be important on the global scale. Differences in equator-pole temperature contrast do to cloud feedbacks in [UVic-CFECFE](#) could impact ocean heat transport in the model.

The application of cloud feedbacks in [UVic-CFECFE](#) provides an important source of inter-model uncertainty that is present in CMIP5/PMIP3. Recent model-data comparisons suggest that the state-of-the-art CMIP5 simulations capture important cloud feedbacks across the observational record (Norris et al., 2016), providing assurance that the feedbacks in [UVic-CFECFE](#) are also within the range of observations. However, as model physics of cloud dynamics and spatial distribution continue to improve in future GCM simulations, the GCM cloud radiative effects can again be applied in [UVic-CFECFE](#) ensemble analyses to emulate the multi-model uncertainty in cloud feedbacks.

Finally, we confirm that the cloud feedbacks in each of the GCMs plays a prominent role in determining the resulting climate sensitivity of each simulation (Fasullo and Trenberth, 2012; Andrews et al., 2012; Sherwood et al., 2014). By incorporating cloud feedbacks into ~~UVic-CFECFE~~, we generally capture the relative spread $\Delta T_{2x,eff}$ of the GCMs (Figure 8). The absolute magnitude of $\Delta T_{2x,eff}$ is typically larger in our ~~UVic-CFECFE~~ simulations relative to each of the GCMs. Since net cloud feedbacks are generally positive in CMIP5 (Vial et al., 2013; Tomassini et al., 2013), the addition of these radiative feedbacks may require a revision of the overall radiative balance in ~~UVic-CFECFE~~. Specifically, future versions of CFE may consider the effects of cloud masking and rapid adjustment in the cloud feedback parameterization (Zelinka et al., 2013). Conversely, the full radiative balance may be adjusted through an enhanced OLW parameterization by slight modification to the constants in Eq. (6). This method of has been applied to UVic to effectively adjust $\Delta T_{2x,eff}$ (Schmittner et al., 2011). The ~~UVic-CFECFE~~ is currently being applied to a study of climate sensitivity using paleoclimate reconstructions (Ullman et al., in prep.).

Code and Data Availability

~~UVic-CFECFE v1.0~~ model code, associated cloud feedback input files, and other relevant data files are available as a Supplement to this manuscript. See the README file in the Supplement for description of contents.

Acknowledgements

This work was supported by a grant from the National Science Foundation's Paleoclimate Perspectives on Climate Change (P2C2) program (award number 1204243).

References

- Andrews, T., Gregory, J.M., Webb, M.J., and Taylor, K.E.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere-ocean climate models, *Geophysical Research Letters*, 39, L09712, doi:10.1029/2012GL051607, 2012.
- Arbuszewski, J., deMenocal, P.B., Cléroux, C., Bradtmiller, L., and Mix, A.: Meridional shifts of the Atlantic intertropical convergence zone since the Last Glacial Maximum, *Nature Geoscience*, 6, 959-962, doi:10.1038/NGEO1961, 2013.
- Barkstrom, B.R.: The earth radiation budget experiment (ERBE), *Bulletin of the American Meteorological Society*, 65, 1170-1185, 1984.
- Barkstrom, B.R., and Smith, G.L.: The earth radiation budget experiment: Science and implementation, *Reviews of Geophysics*, 24, 379-390, 1986.
- Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J.Y., Abe-Ouchi, A., Crucifix, M., Driesschaert, E., Fichefet, T., Hewitt, C.D., and Kageyama, M.: Results of PMIP2 coupled simulations of the Mid-Holocene and

Last Glacial Maximum–Part 2: feedbacks with emphasis on the location of the ITCZ and mid-and high latitudes heat budget, *Climate of the Past*, 3, 279-296, 2007.

- Braconnot, P., Harrison, S.P., Otto-Bliesner, B., Abe-Ouchi, A., Jungclaus, J., and Petterschmitt, J.-Y.: The Paleoclimate Modeling Intercomparison Project contribution to CMIP5, *CLIVAR Exchanges*, 16, 15–19, 2011.
- 5 Braconnot, P., Harrison, S.P., Kageyama, M., Bartlein, P.J., Masson-Delmotte, V., Abe-Ouchi, A., Otto-Bliesner, B., and Zhao, Y.: Evaluation of climate models using palaeoclimatic data, *Nature Climate Change*, 2, 417-424, 2012.
- Cess, R.D., Zhang, M., Wang, P.H., and Wielicki, B.A.: Cloud structure anomalies over the tropical Pacific during the 1997/98 El Nino, *Geophysical research letters*, 28, 4547-4550, 2001.
- Cox, P.M.: *Description of the TRIFFID dynamic global vegetation model*. Technical Note 24, Hadley Centre, United Kingdom Meteorological Office, Bracknell, UK., pp. 1-16, 2001.
- 10 Crucifix, M., M. F. Loutre, P. Tulkens, T. Fichefet, and A. Berger: Climate evolution during the Holocene: A study with an Earth system model of intermediate complexity, *Climate Dynamics*, 19, 43–60, 2002.
- Driesschaert, E.: Climate change over the next millennia using LOVECLIM, a new Earth system model including the polar ice sheets, Ph.D. thesis, 214 pp., Univ. Catholique de Louvain, Louvain-la-Neuve, Belgium, 2005.
- 15 Dufresne, J.L. and Bony, S.: An assessment of the primary sources of spread of global warming estimates from coupled atmosphere-ocean models, *Journal of Climate*, 21, 5135-5144, 2008.
- Eby, M., Weaver, A.J., Alexander, K., Zickfeld, K., Abe-Ouchi, A., Cimatoribus, A.A., Crespin, E., Drijfhout, S.S., Edwards, N.R., Eliseev, A.V., Feulner, G., Fichefet, T., Forest, C.E., Goosse, H., Holden, P.B., Joos, F., Kawamiya, M., Kicklighter, D., Kienert, H., Matsumoto, K., Mokhov, I.I., Monier, E., Olsen, S.M., Pedersen, J.O.P., Perrette, M., Philippon-Berthier, G., Ridgwell, A., Schlosser, A., Schneider von Deimling, T., Shaffer, G., Smith, R.S., Spahni, R., Sokolov, A.P., Steinacher, M., Tachiiri, K., Tokos, K., Yoshimori, M., Zeng, N., and Zhao, F.: Historical and idealized climate model experiments: an intercomparison of Earth system models of intermediate complexity, 9, 1111-1140, doi:10.5194/cp-9-1111-2013, 2013.
- 20 Fasullo, J.T. and Trenberth, K.E.: The annual cycle of the energy budget. Part I: Global mean and land-ocean exchanges, *Journal of Climate*, 21, 2297-2312, 2008.
- 25 Fyke, J. and Eby, M.: Comment on “Climate sensitivity estimated from temperature reconstructions of the Last Glacial Maximum”, *Science*, 337, 1294-1294, 2012.
- Gregory, J.M., Ingram, W.J., Palmer, M.A., Jones, G.S., Stott, P.A., Thorpe, R.B., Lowe, J.A., Johns, T.C., and Williams, K.D.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophysical Research Letters*, 31, L03205, doi:10.1029/2003GL018747, 2004.
- 30 Hartmann, D.L. and Short, D.A.: On the use of earth radiation budget statistics for studies of clouds and climate, *Journal of the Atmospheric Sciences*, 37, 1233-1250, 1980.
- Hartmann, D.L., Ockert-Bell, M.E., and Michelsen, M.L.: The effect of cloud type on Earth's energy balance: Global analysis, *Journal of Climate*, 5, 1281-1304, 1992.

- IPCC Working Group I: Houghton, J.T., Ding, Y., Griggs, D.J., Noguer, M., van der Linden, P.J., Dai, X., Maskell, K., and Johnson, C.A., eds.: *Climate Change 2001: The Scientific Basis, Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2001.
- 5 Joos, F., Prentice, I.C., Sitch, S., Meyer, R., Hooss, G., Plattner, G-K., Gerber, S., and Hasselmann, K.: Global warming feedbacks on terrestrial carbon uptake under the IPCC emission scenarios, *Global Biogeochemical Cycles*, 15, 891–907, 2001
- Loeb, N.G., Wielicki, B.A., Doelling, D.R., Smith, G.L., Keyes, D.F., Kato, S., Manalo-Smith, N., and Wong, T.: Toward optimal closure of the Earth's top-of-atmosphere radiation budget, *Journal of Climate*, 22, 748-766, 2009.
- 10 Lu, J., Vecchi, G.A., and Reichler, T.: Expansion of the Hadley cell under global warming, *Geophysical Research Letters*, 34, L06805, doi:10.1029/2006GL028443, 2007.
- Meissner, K.J., Weaver, A.J., Matthews, H.D., and Cox, P.M.: The role of land surface dynamics in glacial inception: a study with the UVic Earth System Model, *Climate Dynamics*, 21, 515-537, 2003.
- Muglia, J., and Schmittner, A.: Glacial Atlantic overturning increased by wind stress in climate models, *Geophysical Research Letters*, 42, 9862-9868, doi:10.1002/2015GL064583, 2015.
- 15 NOAA National Centers for Environmental Information, State of the Climate: Global Analysis for Annual 2015, published online January 2016, retrieved on July 6, 2016 from <http://www.ncdc.noaa.gov/sotc/global/201513>.
- Norris, J.R., Allen, R.J., Evan, A.T., Zelinka, M.D., O'Dell, C.W., and Klein, S.A.: Evidence for climate change in the satellite cloud record, *Nature*, 536, 72-75, 2016.
- Otto-Bliesner, B.L., Brady, E.C., Clauzet, G., Tomas, R., Levis, S., and Kothavala, Z.: Last glacial maximum and Holocene 20 climate in CCSM3. *Journal of Climate*, 19, 2526-2544, 2006.
- Peltier, W.R.: Global glacial isostasy and the surface of the ice-age Earth: the ICE-5G (VM2) model and GRACE. *Annu. Rev. Earth Planet. Sci.*, 32, 111-149, 2004.
- Plattner, G.-K., Joos, F., Stocker, T.F., and Marchal, O.: Feedback mechanisms and sensitivities of ocean carbon uptake under global warming, *Tellus*, 53B, 564–592, 2001.
- 25 Ramanathan, V., Cess, R.D., Harrison, E.F., Minnis, P., Barkstrom, B.R., Ahmad, E., and Hartmann, D.: Cloud-Radiative Forcing and Climate: Results from the Earth Radiation Budget Experiment, *Science*, 243, 57-63, 1989.
- Schmittner, A., Urban, N.M., Shakun, J.D., Mahowald, N.M., Clark, P.U., Bartlein, P.J., Mix, A.C., and Rosell-Melé, A.: Climate sensitivity estimated from temperature reconstructions of the Last Glacial Maximum, *Science*, 334, 1385-1388, 2011.
- 30 Sherwood, S.C., Bony, S., and Dufresne, J.L.: Spread in model climate sensitivity traced to atmospheric convective mixing, *Nature*, 505, 37-42, 2014.
- Soden, B.J. and Held, I.M.: An assessment of climate feedbacks in coupled ocean-atmosphere models, *Journal of Climate*, 19, 3354-3360, 2006.

- Taylor, K.E., Stouffer, R.J., and Meehl, G.A.: An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, 93, 485-498, 2012.
- Thompson, S.L. and Warren, S.G.: Parameterization of outgoing infrared radiation derived from detailed radiative calculations, *Journal of the Atmospheric Sciences*, 39, 2667-2680, 1982.
- 5 [Tselioudis, G., Rossow, W.B., and Rind, D.: Global patterns of cloud optical thickness variation with temperature, *Journal of Climate*, 5, 1484-1495, 1992.](#)
- Tomassini, L., Geoffroy, O., Dufresne, J-L., Idelkadi, A., Cagnazzo, C., Block, K., Mauritsen, T., Giorgetta, M., and Quaas, J.: The respective roles of surface temperature driven feedbacks and tropospheric adjustment to CO₂ in CMIP5 transient climate simulations, *Climate Dynamics*, 41, 3103-3126, doi:10.1007/s00382-013-1682-3, 2013.
- 10 Ullman, D.J., Schmittner, A., and Urban, N.M.: A new estimate of climate sensitivity using Last Glacial Maximum model-data constraints that includes parametric, feedback, and proxy uncertainties, in prep.
- Vial, J., Dufresne, J.L., and Bony, S., 2013: On the interpretation of inter-model spread in CMIP5 climate sensitivity estimates, *Climate Dynamics*, 41, 3339-3362, 2013.
- Weaver, A.J., Eby, M., Wiebe, E., Bitz, C.M., Duffy, P.B., Ewen, T.L., Fanning, A.F., Holland, M.M., MacFadyen, A., 15 Matthews, H.D., Meissner, K.J., Saenko, O., Schmittner, A., Wang, H., and Yoshimori, M.: The UVic Earth System Climate Model: model description, climatology, and applications to past, present and future climates, *Atmosphere-Ocean*, 39, 361-428, 2001.
- Wielicki, B.A., Barkstrom, B.R., Harrison, E.F., Lee III, R.B., Louis Smith, G., and Cooper, J.E.: Clouds and the Earth's Radiant Energy System (CERES): An earth observing system experiment, *Bulletin of the American Meteorological Society*, 77, 853-868, 1996.
- 20 Yin, J.H. and Battisti, D.S.: The importance of tropical sea surface temperature patterns in simulations of Last Glacial Maximum climate, *Journal of Climate*, 14, 565-581, 2001.
- [Zelinka, M.D., Klein, S.A., and Hartmann: Computing and partitioning cloud feedbacks using cloud property histograms. Part I: Cloud radiative kernels, *Journal of Climate*, 25, 3736-3754.](#)
- 25 [Zelinka, M.D., Klein, S.A., Taylor, K.E., Andrews, T., Webb, M.J., Gregory, J.M., and Forster, P.M.: Contributions of different cloud types to feedbacks and rapid adjustments in CMIP5, *Journal of Climate*, 26, 5007-5027.](#)




Figure 1: Comparison of annual-averaged atmospheric albedo (α_{atm}) as calculated using Eq. (3) and the climatology of ERBE (left) and CERES (right) data.




Figure 2: Maps of annual-mean atmospheric albedo feedback term ($\alpha_{atm}FB$), as calculated using Eq. (11) and the $4\times\text{CO}_2$ results of the 7 CMIP5 models discussed in the text. Units are albedo fraction change per $^{\circ}\text{C}$.




Figure 3: Maps of annual-mean atmospheric albedo feedback term ($\alpha_{atm}FB$), as calculated using Eq. (11) and the LGM results of the 7 PMIP3 models discussed in the text. Units are albedo fraction change per $^{\circ}C$. Note that because the LGM represents a period of global cooling (Braconnot et al., 2012), the direction of change in α_{atm} is opposite that shown in these figures.




Figure 4: Maps of annual-mean outgoing longwave feedback term (OLW_{cloud}^{FB}), as calculated using Eq. (14) and the $4xCO_2$ results of the 7 CMIP5 models discussed in the text. Units are $W m^{-2} °C^{-1}$.






Figure 5: Maps of annual-mean outgoing longwave feedback term ($OLW_{cloud}FB$), as calculated using Eq. (14) and the LGM results of the 7 PMIP3 models discussed in the text. Units are $W m^{-2} °C^{-1}$. Note that because the LGM represents a period of global cooling (Braconnot et al., 2012), the direction of change in OLW_{cloud} is opposite that shown in these figures.




5 Figure 6: Comparison of 4xCO₂ (top) and LGM (bottom) top-of-the-atmosphere feedbacks calculated from raw CMIP5/PMIP3 output from each of the 7 GCMs (CMIP5/PMIP3) and from UVic simulations using GCMs-derived cloud feedbacks (UVic). Shortwave feedbacks are shown on the left, longwave feedbacks on the right. Positive values designate an increased forcing *TO* the climate system with increased temperature (i.e. positive feedback). Feedbacks from the UVic control simulation without cloud feedbacks is shown in grey.


Global Mean SAT anomaly, 4xCO₂



Global Mean SAT anomaly, LGM



Zonal Mean SAT anomaly, 4xCO₂



Zonal Mean SAT anomaly, LGM





Figure 7: Global mean surface air temperature anomalies for the 4xCO₂ (upper left) and LGM (upper right) UVic-CFE simulations. Zonal mean surface air temperature anomalies from the UVic-CFE simulations, averaged over the last 10 years of the 4xCO₂ simulations (lower left) and the last 100 years of the LGM simulation (lower right).



5

Figure 8: Comparison of effective equilibrium climate sensitivity ($\Delta T_{2xCO_2,eff}$) calculated from raw CMIP5 output from each of the 7 GCMs (CMIP5) and from UVic simulations using GCMs-derived cloud feedbacks (UVic). $\Delta T_{2xCO_2,eff}$ from the UVic control simulation without cloud feedbacks is shown in grey.

10