



A global wetland methane emissions and uncertainty dataset for atmospheric chemical transport models.

A. Anthony Bloom¹, Kevin Bowman¹, Meemong Lee¹, Alexander J. Turner², Ronny Schroeder³, John
5 R. Worden¹, Richard Weidner¹, Kyle C. McDonald^{1,3}, Daniel J. Jacob².

¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

²School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA

³The City College of New York, NY, USA

10 *Correspondence to:* A. Anthony Bloom (abloom@jpl.nasa.gov)

Abstract. Wetland emissions remain one of the principal sources of uncertainty in the global atmospheric methane (CH₄) budget, largely due to poorly constrained process controls on CH₄ production in waterlogged soils. Process-based estimates of global wetland CH₄ emissions and their
15 associated uncertainties can provide crucial prior information for model-based top-down CH₄ emission estimates. Here we construct a global wetland CH₄ emission model ensemble for use in atmospheric chemical transport models. Our 0.5°×0.5° resolution model ensemble is based on satellite-derived surface water extent and precipitation re-analyses, nine heterotrophic respiration simulations (eight carbon cycle models and a data-constrained terrestrial carbon cycle analysis) and three temperature
20 parameterizations for the period 2009-2010; an extended ensemble subset – based solely on precipitation and the data-constrained terrestrial carbon cycle analysis – is derived for the period 2001-2015. We incorporate the mean of the full and extended model ensembles into GEOS-Chem and compare model against surface measurements of atmospheric CH₄; model performance (site-level and zonal mean anomaly residuals) compares favourably against published wetland CH₄ emissions
25 scenarios. We find that uncertainties in carbon decomposition rates and wetland extent together account for more than 80% of the primary uncertainty in the timing, magnitude and seasonal variability of wetland CH₄ emissions, although uncertainty in the temperature CH₄:C dependence is a significant



contributor to seasonal variations in mid-latitude wetland CH₄ emissions. The combination of satellite, carbon cycle models and temperature dependence parameterizations provides a physically informed structural a priori uncertainty critical for top-down estimates of wetland CH₄ fluxes: specifically, our ensemble can provide enhanced information on the prior CH₄ emissions uncertainty and the error covariance structure, as well as a means for using posterior flux estimates and their uncertainties to quantitatively constrain global wetland CH₄ emission biogeochemical process controls.

1 Introduction

10 Methane (CH₄) is a potent greenhouse gas, with a global warming potential of more than 25 times that of CO₂ on a 100-year time horizon (Myhre et al., 2013). The global CH₄ budget and growth rate remain poorly understood, largely due to poorly resolved evolution of atmospheric CH₄ sources and sinks (Nisbet et al., 2014). Wetland CH₄ emissions are the largest natural source of atmospheric CH₄, amounting to roughly 20 – 40% of global CH₄ emissions (Ciais et al., 2013). The large disparities between a range of top-down and bottom up wetland CH₄ estimates (Kirschke et al., 2013; Melton et al., 2013) arise from large uncertainties in the timing, distribution and the underlying processes controlling net wetland CH₄ production.

In wetland soils, CH₄ is produced by the decomposition of organic matter in anaerobic (oxygen depleted) environments. The dominant processes controlling the seasonal and inter-annual variations include carbon availability (soil C substrate) and decomposition rate, wetland inundation extent, and temperature (Yvon-Durocher et al., 2014). Other important controls on wetland CH₄ emissions include the presence of macrophytes (Laanbroek 2010), organic C decomposition rates (Miyajima et al., 1997) and soil pH (Singh et al., 2000), amongst other factors. The link between terrestrial carbon-water cycling and wetland CH₄ emissions is of particular interest from a terrestrial greenhouse gas emissions standpoint: inter-annual variations in terrestrial carbon cycling (Le Quéré et al., 2013) can affect wetland CH₄ emissions on seasonal-to-century timescales (Hodson et al., 2011). The role of carbon cycle dynamics in global wetland CH₄ emissions is increasingly recognized: temporal variations in gross primary production influence short-term carbon supply (such as carbon inputs from root exudates



and fine litter), as well as long-lived carbon stores (such as wood litter turnover or soil organic C) in wetland soils (Riley et al., 2011; Bloom et al., 2012; Melton et al., 2013). The combined response of CO₂ and CH₄ fluxes to climatic variability remains poorly characterized. For example, increasing temperatures in boreal ecosystems could lead to higher carbon uptake, increased respiration and drier
5 soils (Watts et al., 2014), and it is currently unclear whether these processes amount to an amplifying or dampening effect on boreal CH₄ emissions. From a greenhouse gas balance standpoint, quantifying the global-scale process links between terrestrial carbon cycling and wetland CH₄ emissions is crucial to characterizing the combined terrestrial biosphere CO₂ and CH₄ flux response to climatic variability.

Quantification of regional wetland CH₄ emissions remains challenging. While wetland CH₄
10 emissions are relatively well constrained on a global scale (Kirschke et al., 2013), regional CH₄ fluxes are difficult to detect, due to their comparatively diffuse nature – relative to anthropogenic point sources – and the scarcity of direct measurements of wetland CH₄ emissions. From a bottom-up perspective, challenges in wetland CH₄ modelling stem from order-of-magnitude uncertainties on wetland CH₄ emissions factors and their spatio-temporal dependence on biogeochemical process controls.
15 Nonetheless, for top-down CH₄ emission estimates, prior knowledge of wetland CH₄ emissions and their associated uncertainty is critical in the formulation of Bayesian atmospheric CH₄ inversions. Atmospheric inversions combine CH₄ measurements from surface, aircraft and satellites (Wecht et al., 2014b; Jacob et al., 2016) and the prior probability on the magnitude and uncertainty characteristics of CH₄ emissions (Bousquet et al., 2011; Pison et al., 2013; Fraser et al., 2014; Turner et al., 2015): these
20 CH₄ inversions typically formulate wetland CH₄ emission uncertainty as a spatio-temporally uncorrelated and normally distributed CH₄ prior. However, inter-model similarities reveal significant levels of emergent correlations in the timing, magnitude and spatial variability of wetland CH₄ emissions. For example, the Wetland CH₄ Inter-comparison of Models Project (WETCHIMP) model ensemble (Melton et al., 2013) reveals varying levels of spatial and temporal agreement between
25 models; these correlations stem from large-scale patterns in biogeochemical process controls (such as temperature, inundation and carbon cycling). Given the relatively large WETCHIMP CH₄ emission uncertainties (model range is typically 150-300% of model mean over major wetland areas, and greater elsewhere), this prior ‘biogeochemical covariance’ can potentially amount to a critical constraint on



atmospheric CH₄ inversions: such a covariance structure can be incorporated in an atmospheric inversion cost-function (Michalak et al., 2005) or as a means for improving attribution of posterior CH₄ fluxes to wetland CH₄ emissions (Wecht et al., 2014a).

Here we propose a process-informed wetland CH₄ emission ensemble based on multiple
5 terrestrial biosphere models, wetland extent scenarios and CH₄:C temperature dependencies. In contrast to a conventional process-based model inter-comparison approach, our ensemble statistics are derived by exhaustively combining a range of temperature, carbon and wetland extent parameterizations. An advantage of our approach is that it provides a prior probability distribution of biogeochemical process control uncertainty: our ensemble can be further constrained – based top-down CH₄ emission estimates
10 – to quantify the combined probability distribution of carbon models, CH₄:C temperature dependencies and wetland extent scenarios.

We formulate a full (2009-2010) and extended (2001-2015) estimate of wetland CH₄ emission magnitude and its associated biogeochemical covariance structure, based on knowledge of the global wetland CH₄ source and the primary biogeochemical process controls. We validate and compare the
15 wetland CH₄ emissions ensemble against a suite of regional studies, and we use a global atmospheric chemical transport model (GEOS-Chem, Bey et al., 2001) to evaluate the CH₄ emissions ensemble relative to existing wetland CH₄ emission models (sections 2 and 3). Finally, we summarize the strengths and limitations of our wetland emissions ensemble and outline its potential applications in global atmospheric inversion frameworks (section 4).

20

2. Wetland CH₄ model ensemble

The wetland CH₄ emissions ensemble provides CH₄ fluxes and associated uncertainty estimates based on four wetland extent parameterizations, nine terrestrial biosphere models of heterotrophic
25 respiration and three CH₄:C temperature parameterization. Global monthly 0.5°×0.5° emissions and their associated uncertainty structure span 2009-2010 (full ensemble, henceforth FE); we also evaluate a subset of the model ensemble spanning 2001-2015 (extended ensemble, henceforth EE). We validate FE



and EE emissions against a range of regional CH₄ emission estimates, and we test the updated GEOS-Chem model against 104 surface CH₄ measurement sites.

2.1 Wetland CH₄ emissions & uncertainty

5

We derive wetland CH₄ emissions F ($\text{mg CH}_4 \text{ m}^{-2} \text{ day}^{-1}$) at time t and location x as:

$$F(t, x) = s A(t, x) R(t, x) q_{10}^{\frac{T(t, x)}{10}} \quad (1)$$

10 where $A(t, x)$ is the wetland extent fraction, $R(t, x)$ is the total C heterotrophic respiration at time for a unit area at time t , $q_{10}^{T(t, x)/10}$ is the temperature dependence of the ratio of C respired as CH₄ (where q_{10} represents the relative CH₄:C respiration for a 10°C increase) and s is a global scale factor. This empirical parameterization provides first order constraints on the role of carbon, water and temperature variability on the global spatial and temporal variability of wetland CH₄ emissions. Variants of the
15 equation 1 parameterization have been used within a range of wetland CH₄ emission models (e.g., Hodson et al., 2011, Pickett-Heaps et al., 2011, Melton et al., 2013 amongst others).

In our approach, wetland CH₄ emissions statistics within each 0.5°×0.5° gridcell are derived based on a 108-member ensemble of wetland CH₄ emission simulations. The 108-member FE is based on 3 CH₄:C temperature dependencies, 9 heterotrophic respiration configurations and 4 wetland extent
20 scenarios (3×9×4 = 108); the six-member EE ensemble is a subset of FE, based on data availability during 2001-2015 (see table 1 for summary). For each ensemble member, the magnitude and uncertainty of s is optimized to match global estimates of the annual wetland CH₄ source (Kirshke et al., 2013).

The heterotrophic respiration configurations are derived from 8 terrestrial biosphere models used
25 in the Multi-scale Synthesis and Terrestrial Model Intercomparison Project (MsTMIP BG1 simulations, see Huntzinger et al., 2013 and Wei et al., 2014 for model and experiment details) and the global CARbon DATa-MODEl fraMework (CARDAMOM) terrestrial carbon analysis (Bloom et al., 2016). V1.0 outputs from the MsTMIP are available for the period 1900-2010 (Huntzinger et al., 2015); the



CARDAMOM analysis was extended to span 2001-2015 based on the Bloom et al., (2016) methodology (see Appendix A for details). Since MsTMIP and CARDAMOM respiration estimates vary intrinsically as a function of temperature, q_{10} only accounts for the temperature dependence of the fraction of C respired as CH_4 . We prescribe three CH_4 :C temperature dependencies (table 1) which are

5 broadly equivalent to a $\pm 50\%$ range on the CH_4 : CO_2 temperature dependence reported by Yvon-Durocher et al., (2014).

Here we use two spatial ($i = 1,2$) and two temporal ($j = 1,2$) wetland extent parameterizations approaches to represent the uncertainty associated with the role of hydrology on wetland CH_4 emissions. We parameterize wetland extent as the product of a static extent constraint $w_i(x)$ and a

10 normalized time-varying scale factor $h_{ij}(x,t)$, in the following manner

$$A(x, t) = w_i(x)h_{ij}(x, t), \quad (2)$$

$w_1(x)$ is the sum of all GLOBCOVER wetland and freshwater land cover types (all flooded, water-logged, and inland water body land-cover types; Bontemps et al., 2011) and $w_2(x)$ the Global

15 Wetland and Lakes Database (GLWD) maximum recorded wetland and freshwater body extent map by Lehner & Doll (2004).

For $h_{ij}(x,t)$, we use (a) the Surface Water Microwave Product Series (SWAMPS) multi-satellite surface water product (Schroeder et al., 2015; $j=1$), and (b) monthly ERA-interim precipitation

20 ($j=2$): for $i = 1$ ($i = 2$), $h_{ij}(x,t)$ is normalized such that mean (maximum) $h_{ij}(x,t)$ is equal to 1. We note that the two hydrological proxies provide contrasting advantages and disadvantages. Satellite-retrieved surface water extent provides an observation-based constraint on the spatial and temporal extent of wetlands and freshwater bodies. While our temporal scaling of static wetland and freshwater extent mitigates the role of spatial biases in satellite-retrieved inundation, vegetation cover remains a major

25 confounding variable in satellite-constrained wetland extent (Schroeder et al., 2015). Moreover, satellites cannot directly observe subsurface soil saturation, even though these soils amount to significant CH_4 fluxes to the atmosphere (Turetsky et al., 2014). On the other hand, precipitation does not provide a direct constraint on the wetland and freshwater extent; however, it provides an aggregate



constraint on ecosystem hydrological variability and wall-to-wall coverage across the globe. We henceforth refer F as “wetland CH_4 emissions”; however, we recognize that lakes, rivers and reservoirs account for ~20% of the total wetland and freshwater body extend (Lehner and Döll, 2004). We discuss the implications of including non-wetland freshwater bodies in $w_i(x)$ section 4.

5 For each of the 108 FE ensemble configurations ($c = 1 - 108$), and 6 EE ensemble configurations ($c = 1-6$), we derive s_c such that:

$$s_c = \frac{G}{\sum_x \sum_t F_{x,t,c} A_x \frac{\Delta t}{n}} \quad (3)$$

10 where $F_{x,t,c}$ are the c^{th} ensemble member fluxes at grid-cell x and time t , A_x is the area of grid-cell x , Δt is the timestep (1 month), n is the number of years, and G is the global total CH_4 emitted from wetlands. We derive s_c such that each ensemble members are scaled to $G = 175 \text{ Tg CH}_4 \text{ yr}^{-1}$ throughout the duration of FE (2009-2010): this is consistent with the Kirschke et al., (2013) mean 2000-2009 top-down wetland CH_4 emission estimates ($175 \text{ Tg CH}_4 \text{ yr}^{-1}$).

15 We derive the CH_4 flux ensemble mean and 5th – 95th percentile ranges at location x and time t based on all 108 $F_{x,t,c}$ ensemble members. For the percentile range calculations, we propagate the global mean wetland CH_4 emission uncertainty by Kirschke et al., (2013) (minimum-to-maximum range = $142 - 208 \text{ Tg CH}_4 \text{ yr}^{-1}$, or $175 \text{ Tg CH}_4 \text{ yr}^{-1} \pm 19\%$). We create an expanded ensemble (FE_{exp}) by randomly perturbing all 108 model s_c values by a factor of $U(0.81, 1.19)$ 1000 times, where $U()$ denotes
20 a random number sampled from a continuous uniform distribution spanning the bracketed numbers. We use the expanded ensemble FE_{exp} to derive the ensemble’s spatiotemporal error covariance structure: the quantitative derivation and qualitative interpretation of the error covariance structure is fully described in appendix B.

We attribute the uncertainty of the timing and magnitude of $F(x,t)$ (maximum CH_4 emission
25 month, mean CH_4 emissions and CH_4 emission variability) to carbon decomposition, wetland extent and $\text{CH}_4:\text{C}$ temperature dependence uncertainty; the derivation of the “primary uncertainty” within each



zonal band (i.e. the dominance of carbon, water or temperature as the dominant source of uncertainty) is fully described in Appendix C.

2.2 GEOS-Chem atmospheric CH₄ simulations

5

We evaluate the FE and EE wetland CH₄ emission means against the World Data Centre for Greenhouse Gases (WDCGG) CH₄ measurement sites by incorporating these into the 4°×5° resolution GEOS-Chem atmospheric chemical and transport model (version 10.01; acmg.seas.harvard.edu/geos). We benchmark the FE and EE runs against GEOS-Chem simulations with the GEOS-Chem wetland CH₄ emission inventory (Pickett-Heaps et al., 2011; henceforth GC) and the Bloom et al., (2012) satellite-constrained wetland emissions (henceforth BL), as these emission estimates have been in a range of atmospheric chemical transport model simulations (Fraser et al., 2013; Turner et al., 2015; Wilson et al., 2016 amongst others). We perform each GEOS-Chem forward run for the period 2009-2010 with a four-year (2005-2009) spin-up period. The non-wetland CH₄ sources in GEOS-Chem consist of biofuel, fossil fuel, livestock, waste, Rice (EDGAR v4.2; European Commission, 2011), fires (Global Fire Emissions Database version 4; van der Werf et al., 2010), soil C sinks and termites (Fung et al., 1991). The non-wetland CH₄ fluxes are the same in each run, with the exception of rice source in run BL (as global wetland and rice emissions are treated as one source by Bloom et al., 2012). While model CH₄ surface concentrations are strongly influenced by wetland CH₄ magnitude, timing and distribution (Bloom et al., 2012, Meng et al., 2015), comparisons between GEOS-Chem outputs and surface CH₄ measurement may also be affected by errors in non-wetland CH₄ emissions and in transport. However, Wecht et al. (2012) and Turner et al. (2015) show that the GEOS-Chem emissions and transport provide an unbiased representation of the observed latitudinal background. The global inversion of Turner et al. (2015) using GEOS-Chem emissions as prior further shows no large errors in non-wetland emissions that would confound the analysis presented here.

For each of the four runs (FE, EE, GC and BL), we use the Wecht et al., (2014a) 1st Jan 2005 initial conditions for atmospheric CH₄ concentrations in GEOS-Chem. For each simulation, we performed a four-year spin-up period (2005-2009) using 2009 emissions to reduce the potential



inconsistency between initial conditions and the global distribution of wetland CH₄ emissions; this spin-up ensures that the relative variations in Jan 1st 2009 CH₄ concentrations for each run are broadly consistent with each emission scenario. We save GEOS-Chem atmospheric CH₄ concentrations every 3 hours. We compare mean monthly GEOS-Chem output against all WDCGG sites below 500m altitude (104 sites with monthly 2009-2010 data in total); this minimizes the topographic mismatch due to vertical CH₄ representation in the model. For each site, the nearest 4°×5° GEOS-Chem grid-cell is used for comparison.

3. Results, Comparison and Validation

10

Mean full ensemble (FE) global wetland emissions are largely accounted for by three high-latitude regions, three tropical regions, and sub-tropical southeast Asia (Figure 1a). North America, Scandinavia and Siberia median (5th – 95th percentiles) CH₄ fluxes amount to 8% (3 – 30%), 2% (0 – 6%) and 2% (1 – 6%) of global emissions; Amazon wetland emissions amount (30%; 21 – 38%) account for the largest tropical emission source, followed by the Indonesian archipelago (13%; 7 – 22%), and central Africa (12%; 7 – 24%); subtropical southeast Asia emissions account for 5% (1 – 10%). High-latitude (>50°N) and tropical emissions amount to 11% (4 – 29%) and 68% (43 – 84%) of global wetland CH₄ emissions, respectively. High-latitude FE emissions exhibit a peak at 60°N (Figure 1b) similar to the GEOS-Chem wetland CH₄ emissions inventory (Pickett-Heaps et al., 2011; GC) and further north than the Bloom et al., (2012) emissions (BL). Tropical emissions for all three emission datasets peak within 0° – 5°S. A comparison between zonal mean emissions (Figure 2) reveals differences of less than 1 Tg/yr/°lat between FE and the extended ensemble (EE) s; the FE zonal mean is comparable to the BL in the near-equatorial tropics and significantly (with respect to the FE model ensemble 90% confidence range) lower everywhere else; the FE zonal mean is comparable to GC in high-latitude and temperate regions, but significantly lower than GC in the tropics and southern hemisphere.

25

All CH₄ emission models show similar patterns in the temporal distribution of CH₄ emissions in high-latitude and temperate regions (with CH₄ emissions peaking between July and September, Figure 3).



- We note that the larger CH₄ fluxes in the BL emissions over Asia and Oceania are due to rice paddy CH₄ emissions. The CH₄ emission models exhibit 1-month differences in the timing of maximum seasonal CH₄ emission across the high-latitudes (generally between June and August). In tropical South America 0° – 20°S latitudes, FE and EE emissions peak March –May, which is comparable to BL (March); and overall earlier than GC (5°S – 20°S emission peak in September). There is a considerable disagreement between northern tropical Africa emission variability amongst all models. Subtropical Asia FE and EE emissions (20°N – 30°N) peak in June-August, earlier than BL emissions (August-September) and comparable to GC emissions (June).
- 10 We compare mean FE and EE (2009-2010) wetland emissions against a range of independent wetland CH₄ regional emission estimates (Figure 4). Emissions from Siberian wetlands (Glagolev et al., 2011) Hudson bay lowlands (Pickett-Heaps et al., 2011), and Amazon river basin (Melack et al., 2004) are within 25th – 75th percentile estimates of FE and EE wetland CH₄ emissions; Chang et al., (2014)(May-September 2012) wetlands are higher (2.1 Tg CH₄ yr⁻¹) but within the 5th -95th percentile range of FE and EE wetland CH₄ emission estimates. BL (2009-2010) and GC (2009-2010) estimates are also within 5th – 95th percentile ranges. With the exception of Amazon river basin estimates, the FE and EE emission estimate uncertainty is larger than the Melton et al., (2013) wetland CH₄ emission model (WETCHIMP 1993-2004) range. We note the temporal mismatch between the wetland emission estimates shown in Figure 4: however, we expect inter-annual variation in wetland CH₄ emissions (e.g. Bloom et al., 2010; Melton et al., 2013) to be substantially smaller than the FE and EE estimate uncertainty.

On a zonal basis, the “primary uncertainty” – i.e. the dominant source of uncertainty within each band – in mean CH₄ emissions and the timing of maximum CH₄ emissions is almost completely dominated by carbon decomposition and wetland extent uncertainties (Figure 5). Seasonal variability of CH₄ emissions is also largely dominated by carbon and extent uncertainties, although the temperature CH₄:C dependence is the dominant source of uncertainty in temperate latitudes. At latitudes > 20°N, wetland extent is the dominant source of uncertainty in mean CH₄ emissions, while temperature CH₄:C



dependence accounts for less than 10% of the primary uncertainty attribution. At all latitudes $> 10^{\circ}\text{S}$, carbon decomposition accounts for the vast majority of the primary uncertainty in the timing of wetland CH_4 emissions.

5 We summarizing the FE_{exp} global error covariance structure as an error correlation matrix between mean monthly 2009-2010 emissions across boreal & arctic ($>55^{\circ}\text{N}$) temperate ($23^{\circ}\text{N} - 55^{\circ}\text{N}$), tropical ($23^{\circ}\text{S} - 23^{\circ}\text{N}$) and southern hemisphere ($<23^{\circ}\text{S}$) latitudes (Figure 6); the error correlation matrix quantitatively summarizes similarities in the spatial and temporal patterns between ensemble members, relative to the ensemble mean (see appendix B for description and interpretation). The FE_{exp} error
10 correlation matrix highlights positively correlated ensemble member CH_4 emissions within each region, with larger correlations (generally $r>0.8$) between emissions separated by 1-2 months; tropical emissions exhibiting the largest overall temporal correlations ($r>0.5$). Tropical and boreal & arctic CH_4 emissions are overall anti-correlated, however no temporal patterns emerge between these anti-correlations.

15

Mean 2009-2010 observed and GEOS-Chem forward model run CH_4 concentrations (with FE, EE, BL and GC wetland emissions) are broadly consistent on a latitudinal basis. The observed and modelled zonal atmospheric CH_4 concentration anomaly (relative to mean global 2009-2010 CH_4 concentrations) is shown in Figure 7 (zonal profile root-mean-square errors – RMSE – are 6.4 ppb, 6.6 pbb, 8.4 ppb, 9.2
20 ppb for FE, EE, BL and GC relative to the observed CH_4 anomaly zonal profile). Within the primary wetland CH_4 emission latitudes ($10^{\circ}\text{S} - 80^{\circ}\text{N}$; Figure 2), all mean CH_4 model estimates are within the mean standard deviation of observed CH_4 , except GC at $>60^{\circ}\text{N}$ and all models at 80°N . The median site-level correlation (Pearson's r) between observed and model de-trended CH_4 concentrations for FE (0.79; 5th – 95th percentiles = -0.15 - 0.97) is highest, followed by EE (0.77; 5th - 95th percentiles = -
25 0.24 - 0.96), BL (0.75; 5th - 95th percentiles = -0.08 - 0.99) and GC (0.72; 5th - 95th percentiles = -0.24 - 0.93). The median RMSE between observed and model de-trended CH_4 concentrations for FE (10.98 ppb; 5th - 95th percentiles = 2.81 - 53.11 ppb) and EE (11.61ppb 5th - 95th percentiles = 4.01 - 51.69) are lower than BL (12.29 ppb; 5th - 95th percentiles = 2.35 - 49.31 ppb) and GC (median =



12.95 ppb; 5th - 95th percentiles = 5.06 - 50.98). FE and EE improvements (relative to GC and BL Pearson's r and RMSE) are primarily in northern hemisphere high-latitudes latitudes ($>50^{\circ}\text{N}$; Figure 8). FE Pearson's r and RMSE suggest better performance than GC in southern hemisphere extra-tropical latitudes ($<23^{\circ}\text{S}$); BL is comparable to FE and outperforms EE across southern hemisphere latitudes.

5

4. Discussion

4.1 Model limitations

10 Densely vegetated wetland areas are likely to amount to a large component of the global wetland CH_4 sources; high-carbon density (and high temperatures in the case of tropical wetlands) result in high CH_4 emissions under inundated conditions. However, satellite-derived observations of surface water area (Schroeder et al., 2015) are ill-equipped to observe densely vegetated wetland areas, as the passive microwave sensors become increasingly sensitive to vegetation moisture within high biomass ecosystems (Sippel et al., 1994). For example, estimates of Amazon river basin wetland CH_4 emissions range between 16% - 29% (5th - 95th percentiles) of the global wetland emissions source; high biomass density in this region (Saatchi et al., 2011) may be a significant source of inundation area bias. Therefore, while we incorporate prior information on the mean and maximum wetland extent to scale the satellite-derived inundation fraction, we anticipate that errors in seasonal and inter-annual inundation variability are likely to be larger within densely vegetated wetland areas. We are optimistic that current and upcoming missions such as SMAP and BIOMASS (Entekhabi et al., 2010; Le Toan et al., 2011) combined with data integration approaches (Schroeder et al., 2015; Fluet-Chouinard et al., 2015) can potentially provide additional constraints required to extend current inundation datasets and to improve current surface inundation detection capabilities.

25

The MsTMIP model ensemble provides a first-order estimate of the magnitude and variability of C decomposition within each $0.5^{\circ}\times 0.5^{\circ}$ grid-cell. Here we highlight 4 potentially major sources of error: (a) differences in aerobic:anaerobic turnover rates of major (labile and recalcitrant) C pools (b)



systematic differences in wetland and non-inundated area carbon uptake within each $0.5^\circ \times 0.5^\circ$ grid-cell, (c) systematic differences in dead organic matter C stocks and accumulation between wetland and non-inundated areas, and (d) lateral flows of C into (or out of) wetland areas. Top-down estimates of seasonal and inter-annual terrestrial CO_2 fluxes (e.g. Liu et al., 2014) could be used to independently assess the validity of heterotrophic respiration from the MSTMIP models and CARDAMOM. In turn, top-down CH_4 and CO_2 flux retrievals, and range of in-situ and regional-scale CH_4 flux estimates (Schriel-Uijl et al., 2011; Chang et al., 2014; Budishchev et al., 2014; amongst others) can be combined to assess whether our empirical parameterization is able capture regional, seasonal and inter-annual wetland CH_4 emission variability and their link to the broader terrestrial carbon cycle. Finally, in succession to eddy covariance tower site analyses of CO_2 respiration dependence on temperature (Mahecha et al., 2010), we anticipate that CH_4 eddy covariance measurements will provide critical site-level constraints on the temperature dependence of wetland CH_4 emissions.

Rice paddies likely amount to <20% of wetland CH_4 emissions, and the majority of rice paddy areas are implicitly excluded from our analysis: GLOBCOVER distinguishes between natural and irrigated water bodies, and GLWD explicitly excludes rice paddy extents in China (which alone accounts for a large portion of global rice paddy CH_4 emissions). However, satellite-based inundation fraction retrievals are unable to distinguish the temporal variability of co-located agriculture and natural wetland inundation extent; moreover $0.5^\circ \times 0.5^\circ$ carbon cycle model resolution may be insufficient to resolve spatial differences in wetland and agricultural C cycling. Inadvertent inclusion of co-located rice CH_4 emissions is therefore a potential source of bias in our approach. We note that the distinction between wetland and rice CH_4 emissions has yet to be consistently addressed in global wetland CH_4 emission quantification efforts (see Bloom et al., 2010; Hodson et al., 2011; Melton et al., 2013, and references therein).

25

CH_4 production in non-wetland freshwater bodies, such as very small ponds (Holgerson and Raymond 2016), lakes (Wik et al., 2016) and rivers (Bastviken et al., 2011) is potentially a significant – albeit highly uncertain – term in the global CH_4 budget (Kirshke et al., 2013; Bridgman et al., 2013). Our



approach implicitly accounts for non-wetland freshwater body emissions, since their extent is incorporated in grid-cell scaling factors (see Eq. 2). We recognise the challenge in explicitly distinguishing between wetlands and non-wetland freshwater body CH₄ emissions, as well as the associated physical and biogeochemical process controls: the spatial and temporal characterization of wetland and non-wetland freshwater extent remains challenging from the current spatial resolution (~25km) of surface inundation retrievals (Prigent et al., 2007; Schroeder et al., 2015). Equally, from a carbon perspective, 0.5°×0.5° carbon cycle model resolution is insufficient to resolve spatial variability in wetland and non-wetland freshwater body extent (Lehner and Döll, 2004). Contingent on future resolution enhancements in surface inundation and carbon cycle models, we recommend further investigation on the adequate distinction and estimation of non-wetland freshwater CH₄ emissions for atmospheric CH₄ chemical transport modelling applications.

By constraining global emission estimates to the Kirshke et al., (2013) model range, our approach does not challenge the global annual CH₄ source and uncertainty (175 Tg CH₄ yr⁻¹; range = 142 – 206 Tg CH₄ yr⁻¹ or ±19%); rather, it places constraints on spatial and temporal wetland CH₄ source variability. Since the global uncertainty is substantially smaller than regional, zonal and grid scale uncertainties (Figures 1,2 and 4), we highlight that new or improved constraints on the global wetland CH₄ source are unlikely to substantially influence our regional or grid-scale CH₄ flux confidence range estimates.

4.2 Applications

Based on comparisons against measured CH₄ concentrations and a range of regional and global CH₄ emission estimates (Figures 2-4, 7-8), we have shown that the FE and EE wetland CH₄ emission ensembles robustly represent the global magnitude and uncertainty of wetland CH₄ emissions. The ensemble configurations of inundation extent, carbon decomposition and temperature dependence have together provided a characterization of the dominant source of uncertainty in global wetland CH₄ estimates (Figure 5). The approach outlined here provides a framework for producing prior emission estimates and associated uncertainty. The error covariance structure – along with the CH₄ observing system capabilities (Wecht et al., 2014a) – can be used to devise an optimal strategy for spatially and/or



- temporally aggregating CH₄ fluxes in an atmospheric inversion framework. Retrieved CH₄ flux from assimilating atmospheric CH₄ observations in an inverse modelling framework (e.g. Fraser et al., 2013) could in turn provide a quantitative constraint on the wetland ensemble: the FE and EE model members can be treated as an ensemble of probable biogeochemical process hypotheses that can be weighted
- 5 against atmospheric constraints. In contrast to conventional wetland CH₄ emission estimates (Riley et al., 2011; Pickett-Heaps et al., 2011) and model inter-comparisons (Melton et al., 2013), top-down CH₄ flux estimates can constrain the joint probability distribution of FE and EE carbon models, wetland extent parameterizations, and temperature dependencies.
- 10 We anticipate extensions of the FE beyond the 2009-2010 time period, contingent on the extensions of the MsTMIP and SWAMPS dataset beyond 2010 and 2012 respectively. In light of continued satellite CH₄ retrievals from GOSAT (Parker et al., 2011; Butz et al., 2011) and upcoming satellite CH₄ measurement from TROPOMI on-board ESA Sentinel 5 precursor (Veefkind et al., 2012), we anticipate that the FE and EE datasets will provide key process-based prior knowledge in future atmospheric CH₄
- 15 inversions.

Appendix A: CARDAMOM extension

- 20 CARDAMOM heterotrophic respiration was derived from the Bloom et al., (2016) global terrestrial C cycle 1°×1° analysis. CARDAMOM retrieved C state and process variables for the period 2001-2010 were used to run the ecosystem carbon balance model DALEC2 (Bloom & Williams 2015) to span 2001-2015. The 2011-2015 ERA-interim meteorological drivers and MODIS burned area were obtained as described by Bloom et al., (2016). The CARDAMOM output consists of 4000 heterotrophic
- 25 respiration realisations at each monthly time-step: for each time-step, we use the median CARDAMOM heterotrophic respiration output. We downscale the data to a 0.5°×0.5° resolution using a nearest neighbour interpolation.



Appendix B: Error correlation structure

We derive the model ensembles' space-time $n \times n$ error correlation matrix \mathbf{M} as follows:

$$M_{ij} = \text{cor}(\mathbf{A}_{i,*} | \mathbf{A}_{j,*}) \quad (\text{A1})$$

where n corresponds to the number of space and time wetland CH_4 emission aggregations, and i, j span $1-n$. $A_{(i,m)}$ and $A_{(j,m)}$ correspond to the total CH_4 flux for model m within the i th and j th space-time aggregations (i.e. total wetland CH_4 emissions within a given time & area); $\mathbf{A}_{i,*}$ and $\mathbf{A}_{j,*}$ are $1 \times N$ vectors, where N is the number of models within the ensemble. the “cor()” operator denotes the Pearson's correlation coefficient between the two bracketed vectors. For Figure 6, $A_{1,m}$ we aggregate model wetland CH_4 emissions for each month across four zonal bands: Boreal & Arctic ($>55^\circ\text{N}$) Temperate ($23^\circ\text{N} - 55^\circ\text{N}$), Tropical ($23^\circ\text{S} - 23^\circ\text{N}$) and Southern Hemisphere ($<23^\circ\text{S}$). **Interpretation:** a perfect correlation between the i th and j th indices ($M_{ij} = 1$) indicates that models are consistently over- or under-predicting CH_4 emissions at times-and-locations i and j relative to the ensemble mean; a perfect anti-correlation ($M_{ij} = -1$) indicates that models consistently over-predicting CH_4 emissions at time-and-location i consistently under-predict CH_4 emissions at time-and-location j (relative to the ensemble mean) and vice versa.

Appendix C: Primary process uncertainty

We quantify the primary process uncertainty of wetland CH_4 emission state variables ($s = 1-3$; 1. maximum emission month, 2. mean CH_4 emissions and 3. seasonal variability (standard deviation)) to wetland emission controls ($e = 1-3$; 1. model carbon decomposition, 2. CH_4 :C temperature dependence and 3. wetland extent parameterization) at location x as follows:

$$R_{x,s,e} = \sum_{c=1}^N \frac{\max(\mathbf{M}_{x,s,m_c}) - \min(\mathbf{M}_{x,s,m_c})}{N} \quad (\text{A2})$$



where $R_{x,s,e}$ is the mean range of state variable s across the ensemble given a fixed emission control e ; $\mathbf{M}_{x,s,*}$ is a vector of all ensemble member state variables s at location x ; \mathbf{m}_c denotes the indices of ensemble subset driven by c th emission control e ; N are the number of configurations for each e (the ensemble configuration details are show in Table 1). For example, $R_{100,1,1}$ is the mean range of seasonal CH₄ variability ($s=3$) for a fixed carbon model configuration ($e=1$) at the 100th gridcell ($x=100$). We attribute the zonal primary uncertainty of state variable s to emission control e as:

$$P_{z,s,e} = \frac{\sum_{x_z} r_{x_z,s,e} F_{x_z}}{\sum_{x_z} F_{x_z}} \times 100\% \quad (\text{A3})$$

where x_z are the pixels x within 5° zonal band z , F_{x_z} is the mean 2009-2010 area-integrated CH₄ flux (Eq. 1 in main text). $r_{x_z,s,e} = 1$ if $R_{x_z,s,e} = \min(\mathbf{R}_{x_z,s,*})$ otherwise $r_{x_z,s,e} = 0$; the “min()” function denotes the minimum element of the bracketed vector; i.e. e is the largest source of uncertainty when the mean range in state variable s is smallest for a fixed e . $P_{z,s,e}$ denotes the percentage of zonal band z where emission control e is the greatest source of uncertainty for each s .

15

Data availability

The full ensemble (FE), extended ensemble (EE) presented here are currently available upon request and will become publicly available through the NASA DAAC. MsTMIP monthly 0.5°×0.5° datasets were obtained from nacp.ornl.gov/MsTMIP.shtml. ERA-interim datasets were obtained from apps.ecmwf.int/datasets/data/interim-full-mnth. CARDAMOM 2001-2010 heterotrophic respiration outputs are available at datashare.is.ed.ac.uk/handle/10283/875; 2011-2015 heterotrophic extensions outputs are available upon request. Inundation datasets were obtained from wetlands.jpl.nasa.gov. The GLWD dataset was obtained from gcmd.gsfc.nasa.gov. The GLOBCOVER dataset was obtained from due.esrin.esa.int. The WDCGG data was obtained from ds.data.jma.go.jp/gmd/wdcgg.

25

Acknowledgments



Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. Funding for this study was provided through a NASA Carbon Monitoring System Grant #NNH14ZDA001N-CMS.

- 5 *Funding for the Multi-scale synthesis and Terrestrial Model Intercomparison Project (MsTMIP; <http://nacp.ornl.gov/MsTMIP.shtml>) activity was provided through NASA ROSES Grant #NNX10AG01A. Data management support for preparing, documenting, and distributing model driver and output data was performed by the Modeling and Synthesis Thematic Data Center at Oak Ridge National Laboratory (ORNL; <http://nacp.ornl.gov>), with funding through NASA ROSES Grant*
- 10 *#NNH10AN681. Finalized MsTMIP data products are archived at the ORNL DAAC (<http://daac.ornl.gov>).*

References

- 15 Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., and Enrich-Prast, A.: Freshwater methane emissions offset the continental carbon sink, *Science*, 331, p. 50, doi: 10.1126/science.1196808, 2011.

- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated
- 20 meteorology: Model description and evaluation, *J. Geo-phys. Res.*, 106, 23 073–23 096, doi:10.1029/2001JD000807, 2001.

- Bloom, A. A., Palmer, P. I., Fraser, A., Reay, D. S., and Frankenberg, C.: Large-Scale Controls of Methanogenesis Inferred from Methane and Gravity Spaceborne Data, *Science*, 327, 322–325,
- 25 doi:10.1126/Science.1175176, 2010.



Bloom, A. A., Palmer, P. I., Fraser, A., and Reay, D. S.: Seasonal variability of tropical wetland CH₄ emissions: the role of the methanogen-available carbon pool, *Biogeosciences*, 9, 2821– 2830, doi:10.5194/bg-9-2821-2012, 2012.

5 Bloom, A. A. and Williams, M.: Constraining ecosystem carbon dynamics in a data-limited world: integrating ecological "common sense" in a model–data fusion framework, *Biogeosciences*, 12, 1299–1315, doi:10.5194/bg-12-1299-2015, 2015.

Bloom, A. A., Exbrayat, J.-F., van der Velde, I. R., Feng, L., Williams, M. The decadal state of the
10 terrestrial carbon cycle: Global retrievals of terrestrial carbon allocation, pools, and residence times. *P. Natl. Acad. Sci.*, 113, 1285–1290, doi: 10.1073/pnas.1515160113, 2016.

Bontemps, S., Defourny, P., Bogaert, E. V., Arino, O., Kalogirou, V., and Perez, J. R.: Globcover
Products Description and Validation Report, Tech. rep., ESA, 2011.

15

Bousquet, P., Ringeval, B., Pison, I., Dlugokencky, E. J., Brunke, E.-G., Carouge, C., Chevallier, F.,
Fortems-Cheiney, A., Frankenberg, C., Hauglustaine, D. A., Krummel, P. B., Langenfelds, R. L.,
Ramonet, M., Schmidt, M., Steele, L. P., Szopa, S., Yver, C., Viovy, N., and Ciais, P.: Source
20 attribution of the changes in atmospheric methane for 2006–2008, *Atmos. Chem. Phys.*, 11, 3689–3700,
doi:10.5194/acp-11-3689-2011, 2011.

Budishchev, A., Mi, Y., van Huissteden, J., Belelli-Marchesini, L., Schaepman-Strub, G., Parmentier, F.
J. W., Fratini, G., Gallagher, A., Maximov, T. C., and Dolman, A. J.: Evaluation of a plot-scale methane
emission model using eddy covariance observations and footprint modelling, *Biogeosciences*, 11, 4651–
25 4664, doi:10.5194/bg-11-4651-2014, 2014.

Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-
M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G., Wunch, D., Wennberg, P., Deutscher, N., Griffith,



D., Macatangay, R., Messerschmidt, J., Notholt, J., and Warneke, T.: Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophys. Res. Lett.*, 38, L14812, doi:10.1029/2011GL047888, 2011.

Bridgham, S. D., Cadillo-Quiroz, H., Keller, J. K., and Zhuang, Q.: Methane emissions from wetlands: 5 biogeochemical, microbial, and modeling perspectives from local to global scales, *Glob. Change Biol.*, 19, 1325–1346, doi:10.1111/gcb.12131, 2013.

Chang, R. Y.-W., Miller, C. E., Dinardo, S. J., Karion, A., Sweeney, C. S., Daube, B. C., Henderson, J. M., Mountain, M. E., Eluszkiewicz, J., Miller, J. B., Bruhwiler, L. M. P., and Wofsy, S.C.: Methane 10 emissions from Alaska in 2012 from CARVE airborne observations, *P. Natl. Acad. Sci.*, 111, 16694–16699, doi:10.1073/pnas.1412953111, 2014.

Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., and Thornton, P.: Carbon and Other 15 Biogeochemical Cycles, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: T. F. Stocker, Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., p. 1535, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

20

Entekhabi, D., Njoku, E.G., O'Neill, P.E., Kellogg, K.H., Crow, W.T., Edelstein, W.N., Entin, J.K., Goodman, S.D., Jackson, T.J., Johnson, J. and Kimball, J.: The soil moisture active passive (SMAP) mission. *Proceedings of the IEEE*, 98(5), pp.704-716, doi: 10.1109/JPROC.2010.2043918, 2010.

25 European Commission: Emission Database for Global Atmospheric Research (EDGAR), release version 4.2, Tech. rep., Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL), available at: <http://edgar.jrc.ec.europa.eu> (last access: 1 December 2014), 2011.



Fluet-Chouinard, E., Lehner, B., Rebelo, L. M., Papa, F., & Hamilton, S. K.: Development of a global inundation map at high spatial resolution from topographic downscaling of coarse-scale remote sensing data. *Remote Sensing of Environment*, 158, 348-361. doi:10.1016/j.rse.2014.10.015, (2015)

5 Fraser, A., Palmer, P. I., Feng, L., Boesch, H., Cogan, A., Parker, R., Dlugokencky, E. J., Fraser, P. J., Krummel, P. B., Langenfelds, R. L., O'Doherty, S., Prinn, R. G., Steele, L. P., van der Schoot, M., and Weiss, R. F.: Estimating regional methane surface fluxes: the relative importance of surface and GOSAT mole fraction measurements, *Atmos. Chem. Phys.*, 13, 5697-5713, doi:10.5194/acp-13-5697-2013, 2013.

10

Fraser, A., Palmer, P. I., Feng, L., Bösch, H., Parker, R., Dlugokencky, E. J., Krummel, P. B., and Langenfelds, R. L.: Estimating regional fluxes of CO₂ and CH₄ using space-borne observations of XCH₄: XCO₂, *Atmos. Chem. Phys.*, 14, 12883-12895, doi:10.5194/acp-14-12883-2014, 2014.

15 Fung, I., John, J., Lerner, J., Matthews, E., Prather, M., Steele, L. P., and Fraser, P. J.: Three-dimensional model synthesis of the global methane cycle, *J. Geophys. Res.*, 96, 13033, doi:10.1029/91jd01247, 1991.

Glagolev, M., Kleptsova, I., Filippov, I., Maksyutov, S., and Machida, T.: Regional methane emission
20 from West Siberia mire landscapes, *Environ. Res. Lett.*, 6, 045214, doi:10.1088/1748-9326/6/4/045214, 2011.

Hodson, E. L., Poulter, B., Zimmermann, N. E., Prigent, C., and Kaplan, J. O.: The El Niño-Southern
25 Oscillation and wetland methane interannual variability, *Geophys. Res. Lett.*, 38, L08810, doi:10.1029/2011GL046861, 2011.

Holgerson, M. A. and Raymond, P. A.: Large contribution to inland water CO₂ and CH₄ emissions from very small ponds. *Nature Geoscience*, 9, 222-226, doi:10.1038/ngeo2654, 2016.



Huntzinger, D. N., Schwalm, C., Michalak, A. M., Schaefer, K., King, A. W., Wei, Y., Jacobson, A., Liu, S., Cook, R. B., Post, W. M., Berthier, G., Hayes, D., Huang, M., Ito, A., Lei, H., Lu, C., Mao, J., Peng, C. H., Peng, S., Poulter, B., Ricciuto, D., Shi, X., Tian, H., Wang, W., Zeng, N., Zhao, F., and
5 Zhu, Q.: The North American Carbon Program Multi-Scale Synthesis and Terrestrial Model Intercomparison Project – Part 1: Overview and experimental design, *Geosci. Model Dev.*, 6, 2121-2133, doi:10.5194/gmd-6-2121-2013, 2013.

Huntzinger, D. N., Schwalm, C. R., Wei, Y., Cook, R. B., Michalak, A. M., Schaefer, K., Jacobson, A.
10 R., Arain, M. A., Ciais, P., Fisher, J. B., Hayes, D. J., Huang, M., Huang, S., Ito, A., Jain, A. K., Lei, H., Lu, C., Maignan, F., Mao, J., Parazoo, N., Peng, C., Peng, S., Poulter, B., Ricciuto, D. M., Tian, H., Shi, X., Wang, W., Zeng, N., Zhao, F., and Zhu, Q.: NACP MsTMIP: Global 0.5-deg Terrestrial Biosphere Model Outputs (version 1) in Standard Format, Data set, available at: <http://daac.ornl.gov>, from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA,
15 doi:10.3334/ORNLDAAAC/1225, 2016.

Jacob, D. J., Turner, A. J., Maasackers, J. D., Sheng, J., Sun, K., Liu, X., Chance, K., Aben, I., McKeever, J., and Frankenberg, C.: Satellite observations of atmospheric methane and their value for
20 2016. quantifying methane emissions, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-555, in review,

Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B.,
25 Lamarque, J.-F., Langenfelds, R. L., Le Quéré, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R.,



Voulgarakis, A., van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6, 813–823, doi:10.1038/ngeo1955, 2013.

Laanbroek, H. J.: Methane emission from natural wetlands: interplay between emergent macrophytes
5 and soil microbial processes. A mini-review, *Ann. Botany*, 105, 141–153, doi:10.1093/aob/mcp201, 2010.

Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters,
G. P., van der Werf, G., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C.,
10 Enright, C., Friedling-stein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F.,
Klein Goldewijk, K., Levis, S., Levy, P., Lo-mas, M., Poulter, B., Raupach, M. R., Schwinger, J.,
Sitch, S., Stocker, B. D., Viovy, N., Zaehle, S., and Zeng, N.: The global carbon budget 1959–2011,
Earth Syst. Sci. Data, 5, 165–185, doi:10.5194/essd-5-165-2013, 2013.

15 Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs, and
wetlands, *J. Hydrol.*, 296, 1–22, doi:10.1016/j.jhydrol.2004.03.028, 2004.

Le Toan, T., Quegan, S., Davidson, M.W.J., Balzter, H., Paillou, P., Papathanassiou, K., Plummer, S.,
Rocca, F., Saatchi, S., Shugart, H. and Ulander, L.: The BIOMASS mission: Mapping global forest
20 biomass to better understand the terrestrial carbon cycle. *Remote sensing of environment*, 115(11),
pp.2850-2860, doi:10.1016/j.rse.2011.03.020, 2011.

Liu, J., Bowman, K. W., Lee, M., Henze, D. K., Bousserrez, N., Brix, H., Collatz, G. J., Menemenlis, D.,
Ott, L., Pawson, S., Jones, D., and Nassar, R.: Carbon monitoring system flux estimation and
25 attribution: impact of ACOS-GOSAT XCO₂ sampling on the inference of terrestrial biospheric sources
and sinks, *Tellus B*, 66, 22486, doi:10.3402/tellusb.v66.22486, 2014.



Mahecha, M. D., Reichstein, M., Carvalhais, N., Lasslop, G., Lange, H., Seneviratne, S. I., Vargas, R., Ammann, C., Arain, M. A., Cescatti, A., Janssens, I. A., Migliavacca, M., Montagnani, L., and Richardson, A. D.: Global convergence in the temperature sensitivity of respiration at ecosystem level, *Science*, 329, 838–840, doi:10.1126/science.1189587, 2010.

5

Melack, J. M., Hess, L. L., Gastil, M., Forsberg, B. R., Hamilton, S. K., Lima, I. B. T., and Nova, E. M. L. M.: Regionalization of methane emissions in the Amazon basin with microwave remote sensing, *Global Change Biol.*, 10, 530–544, 10.1111/j.1365-2486.2004.00763.x, 2004.

- 10 Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753-788, doi:10.5194/bg-10-753-2013, 2013.
- 15

Meng, L., Paudel, R., Hess, P. G. M., and Mahowald, N. M.: Seasonal and interannual variability in wetland methane emissions simulated by CLM4Me' and CAM-chem and comparisons to observations of concentrations, *Biogeosciences*, 12, 4029-4049, doi:10.5194/bg-12-4029-2015, 2015.

20

Michalak, A. M., Hirsch, A., Bruhwiler, L., Gurney, K. R., Peters, W., and Tans, P. P.: Maximum likelihood estimation of covariance parameters for Bayesian atmospheric trace gas surface flux inversions, *J. Geophys. Res.*, 110(D24), D24107, doi:10.1029/2005JD005970, 2005.



Miyajima, T., Wada, E., Hanba, Y. T., and Vijarnsorn, P.: Anaerobic mineralization of indigenous organic matters and methanogenesis in tropical wetland soils, *Geochimica et Cosmochimica Acta*, 61, 3739–3751, doi:10.1016/S0016-7037(97)00189-0, 1997.

- 5 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

Nisbet, E. G., Dlugokencky, E. J., and Bousquet, P.: Methane on the Rise—Again, *Science*, 343, 493–495, doi:10.1126/science.1247828, 2014.

15

Parker, R., Boesch, H., Cogan, A., Fraser, A., Feng, L., Palmer, P.I., Messerschmidt, J., Deutscher, N., Griffith, D.W.T., Notholt, J., Wennberg, P.O., and Wunch, D.: Methane observations from the Greenhouse Gases Observing SATellite: Comparison to ground-based TCCON data and model calculations, *Geophys. Res. Lett.*, 38, L15807, doi:10.1029/2011GL047871, 2011.

20

Pickett-Heaps, C. A., Jacob, D. J., Wecht, K. J., Kort, E. A., Wofsy, S. C., Diskin, G. S., Worthy, D. E. J., Kaplan, J. O., Bey, I., and Drevet, J.: Magnitude and seasonality of wetland methane emissions from the Hudson Bay Lowlands (Canada), *Atmos. Chem. Phys.*, 11, 3773–3779, doi:10.5194/acp-11-3773-2011, 2011.

25



- Pison, I., Ringeval, B., Bousquet, P., Prigent, C., and Papa, F.: Stable atmospheric methane in the 2000s: key-role of emissions from natural wetlands, *Atmos. Chem. Phys.*, 13, 11609-11623, doi:10.5194/acp-13-11609-2013, 2013.
- 5 Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., Mahowald, N. M., and Hess, P.: Barriers to predicting changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in CESM, *Biogeosciences*, 8, 1925–1953, doi:10.5194/bg-8-1925-2011, 2011.
- 10 Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *P. Natl. Acad. Sci.*, 108, 9899–9904, doi:10.1073/pnas.1019576108, 2011.
- 15 Schrier-Uijl, A.-P., Veraart, A. J., Leffelaar, P. A., Berendse, F., and Veenendaal, E. M.: Release of CO₂ and CH₄ from lakes and drainage ditches in temperate wetlands, *Biogeochemistry*, doi:10.1007/s10533-010-9440-7, 2011.
- Schroeder, R., McDonald, K.C., Chapman, B.D., Jensen, K., Podest, E., Tessler, Z.D., Bohn, T.J. and
20 Zimmermann, R.: Development and Evaluation of a Multi-Year Fractional Surface Water Data Set Derived from Active/Passive Microwave Remote Sensing Data. *Remote Sensing*, 7(12), 16688-16732, doi:10.3390/rs71215843, 2015.
- Singh, S., Kulshreshtha, K., and Agnihotri, S.: Seasonal dynamics of methane emission from wetlands,
25 *Chemosphere - Global Change Science*, 2, 39–46, doi:10.1016/S1465-9972(99)00046- X, 2000.



Sippel, S. J., Hamilton, S. K., Melack, J. M., and Choudhury, B. J.: Determination of inundation area in the Amazon River floodplain using the SMMR 37 GHz polarization difference. *Remote Sensing of Environment*, 48(1), 70-76, doi:10.1016/0034-4257(94)90115-5, 1994.

5

Turetsky, M. R., Kotowska, A., Bubier, J., Dise, N. B., Crill, P., Hornibrook, E. R. C., Minkinen, K., Moore, T. R., Myers-Smith, I. H., Nykänen, H., Olefeldt, D., Rinne, J., Saarnio, S., Shurpali, N., Tuittila, E.-S., Waddington, J. M., White, J. R., Wickland, K. P., and Wilmking, M.: A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands, *Global Change Biol.*, 20, 2183–2197, doi:10.1111/gcb.12580, 2014.

Turner, A. J., Jacob, D. J., Wecht, K. J., Maasackers, J. D., Lundgren, E., Andrews, A. E., Biraud, S. C., Boesch, H., Bowman, K. W., Deutscher, N. M., Dubey, M. K., Griffith, D. W. T., Hase, F., Kuze, A., Notholt, J., Ohyama, H., Parker, R., Payne, V. H., Sussmann, R., Sweeney, C., Velasco, V. A.,
15 Warneke, T., Wennberg, P. O., and Wunch, D.: Estimating global and North American methane emissions with high spatial resolution using GOSAT satellite data, *Atmos. Chem. Phys.*, 15, 7049-7069, doi:10.5194/acp-15-7049-2015, 2015.

Van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D.
20 C., Defries, R. S., Jin, Y. and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997-2009), *Atmos. Chem. Phys.*, 10(23), 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Veefkind, J. P., Aben, I., McMullan, K., Forster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de
25 Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120, 70–83, 2012.



Watts, J. D., Kimball, J. S., Bartsch, A., and McDonald, K. C.: Surface water inundation in the boreal-Arctic: potential impacts on regional methane emissions. *Environmental Research Letters* 9, 075001, doi: 10.1088/1748-9326/9/7/075001 (2014).

5

Wecht, K. J., Jacob, D. J., Frankenberg, C., Jiang, Z., and Blake, D. R.: Mapping of North American methane emissions with high spatial resolution by inversion of SCIAMACHY satellite data, *J. Geophys. Res.-Atmos.*, 119, 7741– 7756, doi:10.1002/2014jd021551, 2014a.

10 Wecht, K.J., D.J. Jacob, M.P. Sulprizio, G.W. Santoni, S.C. Wofsy, R. Parker, H. Bösch, and J.R. Worden, Spatially resolving methane emissions in California: constraints from the CalNex aircraft campaign and from present (GOSAT, TES) and future (TROPOMI, geostationary) satellite observations, *Atm. Chem. Phys.*, 14, 8175- 8184, 2014b.

15 Wei, Y., Liu, S., Huntzinger, D. N., Michalak, A. M., Viovy, N., Post, W. M., Schwalm, C. R., Schaefer, K., Jacobson, A. R., Lu, C., Tian, H., Ricciuto, D. M., Cook, R. B., Mao, J., and Shi, X.: The North American Carbon Program Multi-scale Synthesis and Terrestrial Model Intercomparison Project: Part 2 - Environmental Driver Data. *Geosci. Model Dev.*, 7, 2875-2893, doi:[10.5194/gmd-7-2875-2014](https://doi.org/10.5194/gmd-7-2875-2014), 2014.

20


Wik, M., Thornton, B. F., Bastviken, D., Uhlback, J., and Crill, P. M.: Biased sampling of methane release from northern lakes: A problem for extrapolation, *Geophys. Res. Lett.*, 43, doi: 10.1002/2015GL066501, 2016.

25 Wilson, C., Gloor, M., Gatti, L. V., Miller, J. B., Monks, S. A., McNorton, J., Bloom, A. A., Basso, L. S., and Chipperfield, M. P.: Contribution of regional sources to atmospheric methane over the Amazon Basin in 2010 and 2011, *Global Biogeochem. Cycles*, 30, 400–420, doi:10.1002/2015GB005300, 2016.



Yvon-Durocher, G., Allen, A. P., Bastviken, D., Conrad, R., Gudas, C., St-Pierre, A., Thanh-Duc, N., and del Giorgio, P. A.: Methane fluxes show consistent temperature dependence across microbial to ecosystem scales, *Nature*, 507, 488–491, doi:10.1038/nature13164, 2014.


- 5 Ziehn, T., Scholze, M., and Knorr, W.: On the capability of monte carlo and adjoint inversion techniques to derive posterior parameter uncertainties in terrestrial ecosystem models, *Global Biogeochem. Cy.*, 26, GB3025, doi:10.1029/2011GB004185, 2012.



10 **Figure 1.** Top row: 2009-2010 full model ensemble (FE, left) and extended model ensemble (EE, right) mean wetland CH₄ emissions; Middle row: 2009-2010 emissions from the GEOS-Chem wetland



CH₄ emissions inventory (GC, left) model and satellite-constrained estimates by Bloom et al., (2012) (BL, right). **Bottom row:** Mean 2009-2010 FE 5th – 95th percentile range (left) and uncertainty factor (5th – 95th percentile range normalized by mean 2009-2010 emissions, right).



5

Figure 2. Top: Zonal profile of full ensemble (FE, red) extended ensemble (EE, black dashed line) and mean wetland CH₄ emissions from 108 ensemble members (grey). **Bottom:** mean FE (red), mean EE (black dashed line) and 90% FE confidence range (pink), GEOS-Chem wetland emission inventory (GC) and the Bloom et al., (2012) emissions (BL).

10




Figure 3. Seasonal mean zonal profiles for this study (full ensemble: FE; extended ensemble: EE) Bloom et al., (2012) wetland emissions (BL) and GEOS-Chem wetland emissions inventory (GC) for North & South America (left column; 180°W – 35°W); Europe & Africa (center column; 35°W – 55°E) and Asia & Oceania (right column; 55°E – 180°E). The black dotted line denotes the maximum emissions month within each 5° latitude bin.




Figure 4. Comparison between mean annual regional wetland CH₄ emission estimates (1. Glagolev et al., 2011; 2. Picket Heaps et al., 2011; 3. Chang et al., 2014; 4. Melack et al., 2004) and global wetland emission datasets by Bloom et al., 2012 emissions (BL), the GEOS-Chem wetland CH₄ emission inventory (GC); this study (full ensemble: FE; extended ensemble: EE), and the range of WETCHIMP models (Melton et al., 2013). The regions (1-4) are shown in the inset map.




Figure 5. Primary uncertainty attribution of maximum CH₄ emissions month (left), magnitude (center) and seasonal variability (right), to carbon decomposition, temperature CH₄:C dependence (q_{10}) and wetland extent parameterization, within 5° latitude bins. The derivation of primary uncertainties is described in Appendix C.




Figure 6. Full ensemble (FE) spatial and temporal error covariance, summarized as monthly error correlation across boreal & arctic ($>55^{\circ}\text{N}$) temperate ($23^{\circ}\text{N} - 55^{\circ}\text{N}$), tropical ($23^{\circ}\text{S} - 23^{\circ}\text{N}$) and southern hemisphere ($<23^{\circ}\text{S}$) latitudes. A correlation between two location-and-time indices indicates the degree to which models consistently over- or under-predict wetland CH_4 emissions relative to the ensemble mean. The non-zero off-diagonal correlation patterns emerge as a function of varying biogeochemical commonalities across ensemble members, such as wetland CH_4 dependencies on temperature, carbon availability and wetland extent. Negative correlations between tropical and northern hemisphere extratropical (i.e. temperate, boreal and arctic) wetlands emerge as a function of a global constraint on wetland CH_4 emissions ($175 \text{ Tg } \text{CH}_4 \text{ yr}^{-1} \pm 19\%$).




Figure 7. Mean 2009-2010 CH₄ measurements and model CH₄ zonal anomalies (ΔCH_4), relative to the mean 2009-2010 global CH₄ concentration. The black dots denote mean WDCGG network observed CH₄ concentrations within 5° latitude bins; the grey envelope denotes the mean 2009-2010 standard deviation across all sites within 5° latitude bins. The coloured symbols and error bars denote the GEOS-Chem equivalent model concentrations statistics based on the FE and EE ensembles (this study), Bloom et al., (2012) (BL) and the GEOS-Chem emission inventory (GC) wetland CH₄ emissions datasets.




Figure 8. Symbol colours denote the monthly de-trended CH₄ model-observation Pearson's r correlation (left column) and RMSE (right column) for the FE (top-row) and EE (bottom-row) wetland CH₄ emissions (CH₄ observations are from the WDCGG measurement site network). The y-axis denotes the difference between FE/EE and model runs with Bloom et al., (2012) wetland CH₄ emissions (BL) and the GEOS-Chem wetland CH₄ emissions inventory (GC).



Tables

Table 1: Wetland CH₄ model ensemble configurations

Parameter	Description		Ensemble configurations
s	Global scaling factor ^(*)		N/A (all configurations are scaled to 175 Tg CH ₄ yr ⁻¹)
A	Wetland extent	<i>Spatial Extent</i>	2 spatial extent parameterization (scaled using GLOBCOVER and GLWD)
		<i>Temporal Variability</i>	SWAMPS inundation extent ^(**)
			ERA-interim precipitation
R	Heterotrophic respiration		8 MsTMIP terrestrial C models ^(**)
			CARDAMOM terrestrial C cycle analysis
$q_{10(c)}$	Temperature-dependent CH ₄ respiration fraction.		3 CH ₄ :C temperature parameterizations: $q_{10(c)} = [1, 2, 3]$

^(*)Global scaling factor is derived such that the global annual CH₄ emissions amount to 175 Tg CH₄ yr⁻¹.

5 ^(**) These datasets are only used in the 2009-2010 “Full Ensemble” (FE).