

Corresponding Author response to Anonymous Referee #1

Comment : I suggest a modification of title according to this last remark:
The Interactions between Soil-Biosphere-Atmosphere land surface model with Multi-Energy Balance option (ISBA-MEB) in SURFEX – Part 2: Introduction of litter formulation and model evaluation for forest sites.

Answer : In accordance with both referee suggestions, we have modified the title to : “The Interactions between Soil-Biosphere-Atmosphere (ISBA) land surface model Multi-Energy Balance (MEB) option in SURFEXv8 - Part 2: Introduction of a litter formulation and model evaluation for local scale forest sites”

Comment : P3 l90: “the” in excess

Answer : corrected

Comment : P4 l92: reference when introduce DIF option

Answer : the reference has been added

Comment : P5 l126: “that” in excess

Answer : corrected

Comment : P5 l155: is there a condition in residual term such $res \geq 0$?

Answer : We did not apply such a condition. We assume (as seems to be the case generally in the literature as far as we can tell) in the correction method that it is H and LE that are underestimated compared to the available energy $R_n - G$, but indeed, we understand the relevance of the comment. To explore the sensitivity to this condition, we plotted (see fig1.png) the adjusted sensible heat flux just as it appears in Figs.6-8 for the three sites. Blue curves are plotted using the correction method of the paper and green curves are plotted by only computing the corrections when $res \geq 0$, otherwise the turbulent heat flux are not adjusted. Fig1.png shows that the differences are quite small between these two methods and mostly occur during nighttime when the fluxes are relatively weak. We checked the statistics and differences are slight and don't change our conclusions.

Comment : P7 l212: reference when introduce ECOCLIMAP database

Answer : The reference has been added

Comment : P9 l272: ‘is’ in excess

Answer : corrected

Comment : P9 l274: is it possible to precise “veg” default value for forests

Answer : the parenthesis is replaced by : “which is constant in time for forests and varies between 0.95 and 0.99 as a function of the forest cover type”

Comment : P9 l287: suppose fig. 3c instead of fig. 3B

Answer : Indeed this is an error and has now been corrected.

Comment : P11 I359-361: Fig2 don't present very clearly that both MEB simulations simulate less ground evaporation compare to ISBA, even for MEBL. I suggest to moderate affirmation or link comment to another figure or table.

Answer : Thank you for this comment, there was a little confusion here. We removed the reference to the figure and better explained that this sentence refers to summer time. We replace these lines by "During this period, both MEB simulations simulate considerably less ground evaporation (about 4 % of summer LE) compared to the standard ISBA simulation which simulates 25 % of summer LE as ground evaporation."

Comment : P13 I 431: ad "at different depths" after soil temperature

Answer : added

Comment : P16 I523: I suggest justifying here why only MEBL is considered here.

Answer : We changed this sentence to : For improved clarity, only the MEBL and the ISBA models are compared here, since the previous evaluation showed the consistent improvement of using the litter option when using MEB for forests.

Comment : Legend Table 3: change "indicates that figures come from" by "indicates that values come from"

Answer : changed

Comment : Figure 2: precise if it is partitioning for a specific year or mean of many years

Answer : The years have been added in the legend

Comment : Figure 4 and 5: there is no unit on Y-axis

Answer : this has been done, the figure has been modified accordingly

Comment : Legend figure 4: "indicated" and not "indciated"

Answer : corrected

Comment : Figure 7: a,b,c indication are missing

Answer : this has been added

Comment : Figure 9: For total WG, please precise soil thickness used to calculus

Answer : we modified the first part of the legend into : "The total soil water content calculated over the root depth indicated in Table.2 "

Comment : Figure 10: G RMSE in legend, H RMSE in Y-axis legend. Need to be the same. Litter thickness is in 10-2m, not in m.

Answer : "H" has been replaced by "G" in the text and the units have been corrected

Comment : Figure 14: Precise H and LE in Y-axis title

Answer : corrected

Corresponding Author response to Anonymous Referee #2

Comment : Change the title to: "The Interactions between Soil-Biosphere-Atmosphere land surface model with a Multi-Energy Balance (ISBA-MEB) option in SURFEXv8 - Part 2: Model evaluation for local scale forest sites"

Answer : In accordance with both referee suggestions, we have modified the title to : "The Interactions between Soil-Biosphere-Atmosphere (ISBA) land surface model Multi-Energy Balance (MEB) option in SURFEXv8 - Part 2: Introduction of a litter formulation and model evaluation for local scale forest sites"

Comment : P3. L83: Remove "from kilometer resolution" or change to "at the resolutions of kilometers"

Answer : The text has been modified to : "This is essential since ISBA is used within the SURFEX platform in various configurations at resolutions ranging **from several to just under 10 kilometers** at the regional scale, such as within the operational mesoscale numerical weather prediction model AROME, (Seity et al., 2011) and the operational distributed hydrological model system SIM, (Habets et al., 2008), **to resolutions ranging from tens to hundreds of kilometers** in global scale models, such as within the global climate models CNRM-CM5.1 (Voldoire et al., 2013) and CNRM-ESM1 (Séférián et al., 2015)."

Comment : P4. L92: What DIF stands for? Is it ISBA-DF in Boone et al. 2000? Also correct it in P7 L203.

Answer : A reference has been added. DIF stands for the diffusive option from *Decharme et al. 2011* but is replaced by DF for more clarity.

Comment : P4. L 93: change to " The prognostic soil temperature is represented by Tg for Ng soil layers"

Answer : This has been changed

Comment : P4. L 93: Change to " soil volumetric water content and water content equivalent of frozen water"

Answer : This has been changed

Comment : P4. 104: Are you referring to the part 1 of this article? Boone et al. 2017a? Correction should be done in reference section and also line 835.

Answer : This is right, date has been corrected

Comment : P4. 108: change to "water content equivalent of snow"

Answer : This has been changed

Comment : P4. 118: change “models” to “simulates”

Answer : This has been changed

Comment : P4. 125: Change to “litter water content equivalent of ice”

Answer : This has been changed

Comment : P6 168: Correct T to Tg

Answer : This has been corrected

Comment : P6 191 Correction “assess”

Answer : This has been corrected

Comment : P7 210: A_{gs}??

Answer : We clarify what the A_{gs} definition by changing the sentence to “The canopy resistance formulation is based on the A_{gs} (leaf net assimilation of CO₂ - leaf conductance to water vapour) model (Calvet et al. 1998, Givelin et al. 2006) which simulates photosynthesis and its coupling to the stomatal conductance in response to atmospheric CO₂”

Comment : P7 L 12 : Define ECOCLIMAP

Answer : The appropriate reference has been added

Comment : P8 230: Add “leaf area index (LAI)”

Answer : It was defined previously on P4.L116, so we opted not to re-define it here

Comment : P9 L 270 : Change R_{net} to R_n as it is in eq 1

Answer : This is done

Comment : P9 L 273 : Remove “is”

Answer : corrected

Comment : P9 279: Is these averages based on the results of the all models at all three sites? For which variable? S_{wnet}?

Answer : It is true that this sentence was not really clear, it has been changed to : “Despite these differences, results are very similar for the S_{wnet} calculation, as seen in Table 4. The models perform well with relatively low values of RMSE (< 10 W m⁻²) and annual error (AE < 8 W m⁻²) for all three sites.” With these changes, we hope that it is now more clear that this sentence is about S_{wnet} and that it is valid for each model and site and not the average.

Comment : P9 287: correct Fig 3b to Fig 3c and a,b,c to the figures

Answer : corrected

Comment : P9 294: Based on table 4, RMSE is higher than 8 Wm⁻² for Barbeau!
Also, is this average AE for all the models and sites? It does not seem to be correct.
Check your calculations please.

The values of RMSE and AE are not quite similar across the models as it is mentioned in the text.

Answer : Indeed, this is a mistake: the maximum value for RMSE and AE are changed to 10 Wm⁻². It is true that the values are not similar in terms of relative difference (e.g. 5.8 and 9.3 W m⁻²) but for such small values, we think it is more pertinent to consider the absolute differences. We clarify this by modifying the text to : “quite similar in terms of absolute errors”.

Comment : P10. 304. Define RCA, replace “one” with “1”

Answer : The sentence is modified to : “ A ratio of 1 is used for forests within the default version of the original two-source model in the RCA (Samuelsson et al. 2011) dual-energy budget LSM”

Comment : P10. 317. Except with ISBA for Le Bray!

Answer : this has been added. The text is now worked as : “Except with ISBA at Le Bray, the simulations tend to slightly underestimate”

Comment : P10 327. Do you mean good results for the sensible heat flux?

Answer : We mean good results for LE. This has been clarified by changing the sentence to : “For the Mediterranean forest at Puechabon, most of the net radiation is converted into sensible heat flux (Fig.6a) which leads to low values of LE in accordance with observations (fig.6b)”

Comment : P 16. Can you also include a figure for Rnet comparison?

Answer : Please find attached the figure named fig2.png for an Rn comparison. However, we think that this figure does not add much since there are mostly minor differences between the two models. The main information in fig2.png is that the average over all sites is slightly biased and that RMSE is higher, compared to the three local sites (due to more uncertainties in albedo, LAI ...): this is already explained in the text. We can add this figure, but since we feel it does not add a lot of additional information and since the paper already includes a lot of figures, we can let this reviewer the editor decide.

Comment : P19 548: New paragraph “ In terms of prospective”

Answer : A new paragraph has been designated

Comment : P19. 549: MEB or MEBL? Define SIM

Answer : corrected and the definition of SIM has been added “within the SIM chain (SAFRAN-ISBA-MODCOU, Habets et al., 2008)”

Comment : Section 5.3: In world wide assessment, is it possible to report also the results with MEB? It would be interesting for readers to see how including litter layer will affect the results.

Answer : We plot on fig5.png the results for RMSE calculations for H (left panel) and LE (right panel). It shows that the litter option almost always performed better for both H and LE. We chose not to show this result mainly for 2 reasons: first, because the paper already includes many figures and we have tried to include those that are the most essential, and second, because the first part of the validation already showed the necessity of the litter representation so that the litter option can be considered for forest as the default version. That being said, if this reviewer think that it is necessary, we propose to add this figure (fig5.png) in the appendix. We propose to refer to it in the text (if it is added) by adding the text: “ As mentioned in Section 5.1, significant improvement is obtained with MEB compared to ISBA, and even more improvement with MEBL. An example of the improvement in H and LE between MEB and MEBL is shown in Appendix C Fig.16. But because of the consistently best behavior with MEBL verses ISBA compared to MEB, MEBL has become the default option for forests.”

Comment : Table 1. nedle? You mean needle?

Answer : yes, this has been corrected

Comment : Table 2: Correction required : “Mean annual temperature” and “annual rainfall “

Answer : corrected

Comment :Change the number of Fig 11 to Fig 2. Change order of Table 1 and Table 2. Try to number figures and tables in the order that they are mentioned in the text.

Answer : These modifications have been done

Comment : Figures 4b and 5b: Hard to see the lines. Can you possible change the scale of the yaxis ? also add the units in on Y-axis. Shaded are is not shown well in the figures.

Answer : We have plotted both turbulent flux components with the same scale so as the reader can easily understand that the energy from the G mostly goes into H. fig3.png and fig4.png shows figures 4 and 5 of the paper, respectively, with changes to the yaxis scale. We think that the new figure could bias the reader in terms of the relative weigh of H and LE and that it doesn't add much information since the LE fluxes are quite similar between the models.

Comment : Figure 9: what is the total depth of the soil for WG (better to change it to wg). Can you specify on the figure?

Answer : the total depth is the root depth indicated in Table 3. This information has been added in the legend.

Comment : Figure 10: Correct the Y-axis title to G RMSE also litter thickness should be in cm on X-axis title

Answer : This change has been done

The Interactions between Soil-Biosphere-Atmosphere (ISBA) land surface model Multi-Energy Balance (MEB) option in ~~SURFEX~~ SURFEXv8 - Part 2: ~~Model~~ Introduction of a litter formulation and model evaluation for local scale forest sites

Adrien Napoly¹, Aaron Boone¹, Patrick Samuelsson², Stefan Gollvik², Eric Martin³, Roland Seferian¹, Dominique Carrer¹, Bertrand Decharme¹, and Lionel Jarlan⁴

¹CNRM UMR 3589, Météo-France/CNRS, Toulouse, France

²Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

³IRSTEA, U-R RECOVER, Aix en Provence, FRANCE

⁴Centre d'études Spatiales de la Biosphère (CESBIO), Toulouse, France

Correspondence to: Adrien Napoly (adr.napoly@gmail.com)

Abstract. Land surface models (LSMs) need to balance a complicated trade-off between computational cost and complexity in order to adequately represent the exchanges of energy, water and matter with the atmosphere and the ocean. Some current generation LSMs use a simplified or composite canopy approach that generates recurrent errors in simulated soil temperature and turbulent fluxes. In response to these issues, a new version of the Interactions between the Surface Biosphere Atmosphere (ISBA) land surface model has recently been developed which explicitly solves the transfer of energy and water from the upper canopy and the forest floor which is characterized as a litter layer. The so-called Multi Energy Balance (MEB) version of ISBA is first evaluated for three well-instrumented contrasting local scale sites, and sensitivity tests are performed to explore the behavior of new model parameters. Second, ISBA-MEB is benchmarked against observations from 42 forested sites from the global micro-meteorological network (FluxNet) for multiple annual cycles.

It is shown that ISBA-MEB outperforms the composite version of ISBA in improving the representation of soil temperature, ground, sensible and to a lesser extent latent heat fluxes. Both versions of ISBA give comparable results in terms of simulated latent heat flux because of the similar formulations of the water uptake and the stomatal resistance. However, MEB produces a better agreement with the observations of sensible heat flux than the previous version of ISBA for 87.5 % of the simulated years across the 42 forested FluxNet sites. Most of this improvement arises owing to the improved simulation of the ground conduction flux, which is greatly improved using MEB, especially owing to the forest litter parameterization. It is also shown that certain processes are also modeled more realistically (such as the partitioning of evapotranspiration into transpiration and ground evap-

oration), even if certain statistical performances are neutral. The analyses demonstrate that shading effect of the vegetation, the explicit treatment of turbulent transfer for the canopy and ground, and the insulating thermal and hydrological effects of the forest floor litter turn out to be essential for simulating the exchange of energy, water and matter across a large range of forest types and climates.

25 1 Introduction

The land surface model (LSM) is one of the key parametrization schemes of atmospheric models used for numerical weather prediction and climate simulations. It is used to compute the turbulent fluxes of heat, water and momentum, along with the radiative fluxes at the surface-atmosphere interface. In addition, the current generation of LSMs are used to compute flux exchanges of certain
30 chemical species (such as carbon dioxide and biogenic organic volatile carbon) and emissions of particles (such as aerosols from dust or biomass burning). The interaction soil biosphere atmosphere model (ISBA) is part of the SURFace EXternalisée platform (SURFEX) developed in recent years at Météo-France (Masson et al., 2013) and a suite of international partners. It is used to either a suit of surface schemes in either offline or coupled mode with an atmospheric model and/or a hydro-
35 logical model. ISBA has benefited from continuous improvements since its first version (Noilhan and Planton, 1989), in particular for the carbon cycle (Calvet et al., 1998; Gibelin et al., 2006), soil mass and heat transfer (~~Boone et al., 1999; ?~~)([Boone et al., 1999](#); [Decharme et al., 2011](#)), snowpack processes (Boone and Etchevers, 2001; Decharme et al., 2016), sub-grid hydrology (Decharme and Douville, 2006), and radiative transfer through the canopy (Carrer et al., 2013).

40 In order to remain consistent with the aforementioned developments and to respond to both current and future user demands, a multi energy balance (MEB) approach has been developed (Boone et al., 2017). This improvement consists in representing the surface as a so-called multi-source model (i.e. a separation of the canopy and the surface soil layer) in contrast to the standard ISBA soil-vegetation composite model. This new model has been designed to better represent certain processes
45 for forested areas, and to incorporate new modeled processes which can benefit from a more accurate representation of the soil-vegetation continuum. In particular, partitioning of the incoming energy into turbulent and ground heat fluxes are expected to be more realistic with the explicit vegetation scheme as well as the partitioning of latent heat flux into its different components. The carbon cycle should also benefits from a more conceptually accurate representation of the surface (Carrer et al., 2013) since it is strongly linked to leaf temperature via assimilation of atmosphere Carbon. Moreover, snow melt during spring is very sensitive to the snow net radiation budget, which is impacted by the presence of tall vegetation, and interception and loss by the canopy during the winter season (Rutter et al., 2009).

The surface in forested regions beneath the canopy is often covered by a layer of dead leaves or
55 needles, branches, fruits and other organic material which can be characterized as a litter layer. The

explicit inclusion of such a layer is neglected for the most part in LSMs used in global scale models, or it is only partly taken into account. For example, Sakaguchi and Zeng (2009) implemented a specific resistance due to litter for soil evaporation. The inclusion of litter-related processes has been shown to have a substantial impact on hydrological (Putuhena and Cordery, 1996; Guevara-
60 Escobar et al., 2007) and thermal processes (Andrade et al., 2010). Litter has an important insulating capacity as its thermal diffusivity is small compared to soil (Riha et al., 1980), which leads to a strong impact on the soil temperature diurnal cycle and ground conduction flux. Changes in this flux can then impact the energy available for radiative and turbulent fluxes. The presence of litter also alters ground evaporation by covering the soil with a high porosity layer which essentially prevents
65 liquid water capillary rise (Schaap and Bouten, 1997). It also constitutes another water interception reservoir for rainfall and canopy drip prior to infiltration and runoff (e.g., Gerrits et al., 2007). Some models have introduced parameterizations for litter (Ogée and Brunet, 2002; Gonzalez-sosa et al., 1999; Wilson et al., 2012; Haverd and Cuntz, 2010; Sakaguchi and Zeng, 2009), but the approach can be very different from one to another depending on their complexity (only modifying or adding
70 a ground resistance, modeling the litter using an explicit single or multi-layer model). In addition, each of the models were validated against observations made for a single well-instrumented site.

The goal of the current study is to evaluate the impact of a new parametrization of the soil-litter-vegetation-atmosphere continuum at the local scale. In the first part of this study, an in-depth evaluation for three forest sites in France is carried out. They have been selected in order to represent
75 a range of forest types and climates. The main goal is to understand the effect of both the explicit canopy layer and the litter layer on the available energy partitioning (latent, sensible, ground), soil temperatures and soil water content. The second part of this study consists in a benchmark study using 42 sites from the Fluxnet network (Baldochi et al., 2001). The objective is twofold: (1) to evaluate the performance of the MEB options against the classic composite soil-vegetation version
80 of the ISBA model over a wide range of forest species and climates; and (2) to analyze whether the general improvements seen at the three well-instrumented and documented local scale sites in France can be extended to other climates and forest types.

This is essential since ~~, as part of the SURFEX platform,~~ ISBA is used within the SURFEX platform in various configurations ~~from kilometer resolutions at resolutions ranging from several~~
85 to just under 10 kilometers at the regional scale ~~such as,~~ such as within the operational mesoscale numerical weather prediction model AROME, (Seity et al., 2011) ~~, or~~ and the operational distributed hydrological model system SIM, (Habets et al., 2008), ~~and in to resolutions ranging from tens to~~ hundreds of kilometers in global scale models, such as within the global climate models CNRM-CM5.1 (~~Voldoire et al., 2013b~~) (Voldoire et al., 2013a) and CNRM-ESM1 (Séférian et al., 2015).

90 2 Model

2.1 The standard ISBA composite soil-vegetation model

The standard ISBA model uses a single composite soil-vegetation surface energy budget which means that the ~~the~~ properties of the soil and vegetation are aggregated within each grid-cell point (Noilhan and Planton, 1989; Noilhan and Mahfouf, 1996). In this study, the so-called multi-layer soil diffusion (~~DFDF~~, [Decharme et al., 2011](#)) option is used, along with the explicit multi-layer snow scheme. The prognostic ~~heat-storage-variables-consist-in-the-temperatures,~~ [soil temperature is represented by](#) T_g , for N_g soil layers. The snowpack uses the snow enthalpy as a prognostic variable, from which both the snow liquid water content and temperature can be diagnosed for N_n layers. The hydrological prognostic variables are the canopy reservoir for water interception, W_r (kg m^{-2}), the snow liquid water content, W_n , for N_n snow layers, and the soil volumetric water ~~and liquid water equivalent ice contents~~ [content and water content equivalent of frozen water](#), w_g and w_{gf} ($\text{m}^3 \text{m}^{-3}$), respectively, for $N_{g,l}$ active hydrological soil layers (where $N_{g,l} \leq N_g$). Further details describing the soil and snowpack representations as well as other prognostic variables and model assumptions are detailed in Decharme et al. (2016).

105 2.2 The ISBA-MEB model

The multi energy balance model, ISBA-MEB, treats up to three fully coupled distinct surface energy budgets (i.e. the snow surface, the bulk vegetation canopy and the ground which is characterized as either a soil surface or litter layer: see Section 2.3). The reader is referred to Boone et al. (2017) for an extended description of the various assumptions of the MEB approach, its full set of governing equations and its numerical aspects. Compared to the classic ISBA approach, there is one additional prognostic heat storage variable which is the vegetation temperature, T_v . The new hydrological prognostic variable is the ~~liquid water equivalent~~ [water content equivalent of](#) snow stored within the vegetation canopy, W_{rn} (kg m^{-2}).

2.3 Explicit model for litter in ISBA-MEB

115 Two methods are generally used to represent the effect of a litter layer within LSMs. The first method consists in adding a specific ground resistance in order to reduce the soil evaporation due to the presence of a litter layer. For example, Sakaguchi and Zeng (2009) and Park et al. (1998) have used this method in their models with the resistance depending on the leaf area index (LAI) of the litter or its thickness. The second method utilizes an explicit model to represent the effects of the litter. 120 The implementation of this method turns out to be quite variable among different LSMs, although generally it accounts for both thermal and hydrological effects. In the current study, a model was introduced which ~~models-simulates~~ the first order effects of a litter layer on the surface energy and water budgets while minimizing the number of parameters and the complexity. To this end, a

single bulk-layer approach, which is based on the model of (Schaap and Bouten, 1997), has been developed. The aforementioned study has also inspired similar approaches by (Ogée and Brunet, 2002) and (Haverd and Cuntz, 2010). The method is relatively simple: a single explicit litter layer is inserted between the vegetation layer and the upper soil layer. With this approach, three additional state equations have to be solved: i) the litter average temperature, T_l (K), ii) the litter liquid water content, W_l (kg m^{-2}), and iii) the litter ~~liquid water equivalent ice content~~ water content equivalent of ice, W_{lf} (kg m^{-2}). Two noteworthy assumptions have been made: first, it is assumed that ~~that~~ there is no evaporation directly from the soil in the presence of litter, and second, water movement by capillarity rise from the soil upwards into litter is neglected. The energy budget is computed for the litter layer using prescribed values of thermal conductivity, heat capacity and litter thickness. The water budget is computed in a similar manner as for the vegetation canopy layer. Finally, frozen water in the litter is modeled using the same simple freezing model method as for the soil in ISBA (Boone et al., 2000). The model is fully described in Appendix A.

3 Data

3.1 Energy balance closure

Energy balance closure is a well-known issue when turbulent fluxes are computed with the eddy covariance technique. The closure, δ , is defined as the ratio of turbulent fluxes to available energy at the surface:

$$\delta = \frac{H + LE}{R_n - G - S} \quad (1)$$

where S , R_n , G , H and LE (W m^{-2}) represent the vegetation heat storage, the net radiation, ground conduction (heat) flux, the sensible heat flux and the latent heat flux, respectively. The basic idea is that closure should be as close as possible to unity when observations are compared with LSM output since such models close the energy budget (to a high degree) by design. As it turns out, the mean closure imbalance for FluxNet sites is typically 20 % (Wilson et al., 2002). According to Twine et al. (2000), the net radiation is probably the most accurately measured component of the energy balance and closure can be reasonably done by adjusting H and LE . The Bowen ratio closure method is often adopted to correct the non-closure. It consists in assuming that the Bowen ratio (ratio of the sensible to the latent heat fluxes) is well estimated by the eddy covariance system, so that the turbulent fluxes can be adjusted while conserving the Bowen ratio to ensure closure (Blyth et al., 2010; Zheng et al., 2014; Er-Raki et al., 2008). The adjusted sensible and latent heat fluxes can be computed as:

$$H_{adj} = H + \text{res} \times \frac{H}{H + LE} \quad (2)$$

$$LE_{adj} = LE + \text{res} \times \frac{LE}{H + LE} \quad (3)$$

where H_{adj} and LE_{adj} (W m^{-2}) represent the adjusted sensible and latent heat fluxes, respectively, and the residual energy is defined as

$$\text{res} = R_n - G - S - H - LE \quad (4)$$

160 Among the terms in Eq. 4, the energy stored in the canopy, S , is generally not measured. Even though it is relatively small compared to the net radiation, it has been shown to be non-negligible in some forest canopies (Oliphant et al., 2004). One possible method which can be used to address this issue is to use the simulated residual (Blyth et al., 2010). In the current study, tests were done using both the model simulated S and neglecting it. It was found that incorporating this term had
165 very little influence on the adjusted fluxes. In the analysis that follows, we neglected this term and both the original and the adjusted fluxes are shown.

3.2 Local scale evaluation over France

Three well-instrumented forest sites have been used for model evaluation which cover a range in climate, soils and vegetation characteristics. Their location is shown in Fig. 1 and their main characteristics are summarized in Table 1. At each site, measurements of sensible and latent heat fluxes
170 were made using the eddy covariance technique, and the ground heat flux was measured with flux plates. Soil temperature (T_g) and volumetric moisture content (w_g) profiles were also available for two of the sites (Le Bray and Barbeau).

The Barbeau site is located in the Barbeau National Forest which is approximately 60 km from
175 Paris (48.4°N, 2.7°E, France, elevation 90 m). This site consists mainly of a deciduous broadleaf mature oak forest which has an average height of 27 m. The climate is temperate, with a mean annual temperature of 10.7 °C and the annual rainfall is 680 kg m^{-2} . The soil texture is loam in the upper soil and clay loam at the lower portion of the soil profile (Prévost-Bouré et al., 2009).

The Le Bray site is located about 20 km from Bordeaux, France (44.7°N, -0.7°E, elevation 62 m).
180 It is a maritime pine forest classified as evergreen needleleaf with an average height of 18 m with a dense grass understory which comprises about half of the total LAI at its maximum. The mean annual temperature is 12.9 °C and the mean annual precipitation is 997 kg m^{-2} . The soil is a sandy and hydromorphic podzol with a layer of compacted sand which starts at a depth between 0.4 and 0.8 m, which constitutes the limit of root penetration. The site also has a water table whose depth
185 fluctuates during the year and, at times, can reach the surface. This is the only site where specific measurements for litter were made, and a near constant litter thickness of 0.05 m was observed (Ogée and Brunet, 2002).

The Puéchabon site is located roughly 35 km northwest of Montpellier in the Puéchabon State Forest (43.7°N, 3.5°E, elevation 270 m). Vegetation is largely dominated by a dense over-story of holm
190 oak (evergreen broadleaf forest) with a mean height of 5.5 m. The climate is typical Mediterranean with mean annual temperature of 13.5 °C and a mean annual precipitation of 872 kg m^{-2} . The soil

texture is homogeneous down to about 0.5 m depth and it can be described as silty clay loam. The parent rock is limestone (Rambal et al., 2014)

3.3 Benchmark

195 The second part of the study uses observations from a subset of the FluxNet sites to ~~asses-~~assess systematically or benchmark the suite of ISBA versions developed at Météo-France. The FluxNet database has been used for LSM evaluation by several widely used LSM models (Stöckli et al., 2008; Blyth et al., 2010; Ukkola et al., 2015). Many sites are available within the FluxNet database, however in the current study, only those sites and years with an energy balance closure of 20 %
200 (or less) before adjustment of the turbulent fluxes are retained. Ground heat flux was assumed to represent 3 % of net radiation when it is not available (this is close to the average value over the three French sites). After screening, 42 forested sites remain which give a total of 179 years of observations (see Fig. 2 for the locations of the sites used in the current study). The method used to gap fill missing atmospheric forcing data is described in Vuichard and Papale (2015). The three sites
205 from the first part of this study have been removed in this analysis.

4 Model set up

~~The-All the~~ simulations are performed using SURFEXv8. ~~We use~~ the diffusive soil (DIFDF) option meaning that the soil heat and mass transfers are solved on a multi-layer grid (Decharme et al., 2011). A recently developed multi-layer canopy radiative transfer scheme (Carrer et al., 2013) has
210 been incorporated into MEB in order to determine the fraction of incoming radiation intercepted by the canopy layer, transmitted to the ground and reflected. This model uses structural organization of the canopy and spectral properties of the leaves as well as the albedo from the soil and the vegetation for two different spectral bands (visible and near-infrared). ~~Finally, the estimation of the stomatal resistance is made with-~~The canopy resistance formulation is based on the A-gs ~~formulation-(leaf~~
215 net assimilation of CO₂ - leaf conductance to water vapour) model (Calvet et al., 1998; Gibelin et al., 2006) which simulates photosynthesis and its coupling to the stomatal conductance in response to atmospheric CO₂. Parameters of the stomatal resistance scheme are vegetation type dependent and were chosen according to the ECOCLIMAP database (Faroux et al., 2013).

Three simulations were performed for each site in order to assess the impact of the new canopy
220 and litter layers on the simulated fluxes, soil temperature and soil moisture. The models were forced with atmospheric data at half-hourly time steps. In the first simulation, the reference model (i.e. using a single composite vegetation and surface-soil layer) was used and it is hereafter referred to as ISBA. The second simulation with the explicit canopy layer corresponds to the ISBA model using the MEB approach (referred as MEB herein for simplicity). The last simulation was carried out with
225 both the explicit canopy layer and the explicit forest litter layer and it is referred to as MEBL.

4.1 Local scale

The pedotransfer functions of Clapp and Hornberger (1978) are used for the soil water flow. The soil hydraulic conductivity at saturation, the so-called b -exponent and the matric potential at saturation are computed with the continuous formulations of Noilhan et al. (1995) using the textural properties at each site. The wilting point, field capacity and the water content at saturation were computed using the observations of soil volumetric water content and/or following the recommendations of the site principal investigators (Table 2). An hydraulic conductivity exponential profile was used to model the packing of the soil with depth. Soil organic content, which affects the hydrological and thermal coefficients as in Decharme et al. (2016), is modeled using the input data from the Harmonized World Soil Database (HWSD, Nachtergaele and Batjes, 2012). Leaf are index (LAI), canopy height, vegetation type, total soil and root depths were set to values found in the corresponding literature for each site (Table 2). Litter thickness, Δz_l , which is one of the key parameters of the litter module, varies in space and time. This study assumes it is constant in time based on estimates made at each site. This is an important approximation, but a literature review reveals that measured litter thickness generally varies from one to ten centimeters (see Table 3). Because of the uncertainties related to specification of this parameter, a series of sensitivity tests are done and the results are presented in Sec. 5.2.

At the Le Bray site, the water table has a significant influence on the seasonal soil wetness. Measurements of its depth were available and allowed a simple parametrization to be developed. It consists in a strong relaxation towards saturation in soil layers below the observed water table depth. Thus, soil moisture within the saturated zone is very close to the observed values in the saturated or nearly saturated layers, while soil moisture above this zone is permitted to freely evolve.

For each site, simulated turbulent heat fluxes were compared (~~Figs. 7a-b; 6a-b; 8a-b~~) to both the original and the adjusted-observed values assuming that model results should fall inside the area delimited by these two curves. Moreover, since a proper evaluation of each flux component of the energy balance has to be done with a closed energy budget (in order to be consistent with the model which imposes closure by design), the scores are computed with the adjusted flux values.

4.2 Benchmark

The ECOCLIMAP land cover and the HWSD soil data bases were used to prescribe most of the needed parameters as was done for the local scale evaluation for the three French sites (in the previous section). This is also consistent with the method used for spatially distributed offline and fully coupled (with the atmosphere) simulations with ISBA. However, note that soil texture, canopy height and vegetation type were chosen in agreement with literature values for each site where available, therefore they superseded the ECOCLIMAP values for these parameters. The default thickness of

260 the litter layer is set to three centimeters based on the sensitivity test results presented in Section 5.2 and in overall agreement with literature values (Table 3).

Initial conditions can have a significant impact on both the sensible and the latent heat fluxes, but they were not known thus a spin-up period was used for each site. Sites were initialized with saturated soil water content conditions and the first available year was repeated at least ten times
265 until a predefined convergence criteria was achieved. Only the latent and sensible heat fluxes are evaluated since the ground heat flux, soil temperature and soil water content were not available at each FluxNet site.

5 Results and discussion

The MEB model has been compared to both observations and the standard ISBA composite vegetation model for several well-instrumented contrasting (in terms of forest type and climate) sites
270 in France, and to a sub-set of the FluxNet sites for multiple annual cycles using a benchmarking application. The results of these evaluations and of several sensitivity tests are given in this section.

5.1 Evaluation for three well-instrumented forested sites

5.1.1 Net radiation

275 The total net radiation (R_{net}) is analyzed in terms of the net longwave (LW_{net}) and short-wave (SW_{net}) components. The SW_{net} primarily depends on the prescribed albedo for the soil, α_g , and the vegetation, α_v . In ISBA, the net shortwave radiation is simply defined as $SW_{net} = (1 - \alpha_{eff}) SW \downarrow$, where the effective albedo is simply $\alpha_{eff} = veg \alpha_v + (1 - veg) \alpha_g$ and veg is the vegetation cover fraction (which is constant for forests in time for forests and varies between 0.95 and
280 0.99 according to the forest cover). For the MEB simulations, the scheme from Carrer et al. (2013) is used which uses the soil and vegetation albedo for two spectral bands (which are aggregated to all-wavelength values for ISBA), along with the LAI (the notion of veg is not applicable to MEB) in order to obtain an estimate of the vegetation optical thickness. Despite these differences, results are very close similar for the SW_{net} calculation, as seen in Table 4. The models perform well with
285 relatively low values of average RMSE (< 6.1 RMSE (< 10 W m⁻²) and average annual error (AE < 8 W m⁻²) for all three sites.

Since MEB explicitly uses the shortwave radiation transmitted through the canopy for the ground net radiation computation, it can be compared with the photosynthetically active radiation (PAR) measurements below the canopy within the Barbeau and Puéchabon forests (Fig. 3). Comparison
290 with MEB results revealed that the transmission coefficient for the PAR is overestimated. The radiative transfer scheme parameter which exerts the most control over this process is the clumping index, and it has been adjusted for these two sites. However, the shortwave radiation transmission remains slightly over-estimated at the Barbeau forest during the first three months of the year in-

cluding this adjustment (Fig. 3bc). The main reason is that during this period, the LAI is very low
295 but the trunks and branches intercept solar radiation (the forest is relatively old and tall, so that this
effect can be quite significant) which is not taken into account by the model. This could be done by
including a stem area index (SAI), but currently the default minimum LAI is used as a proxy for
this effect.

The simulated LW_{net} , which depends on the explicit contributions from the soil, vegetation and
300 snow in MEB, and the composite soil-vegetation layer and snow in ISBA, was quite similar in terms
of absolute errors among the model versions and led to a fairly good comparison with measurements.
The annual RMSE is less ~~than 8~~ or equal than 10 W m^{-2} for each site and ~~run~~ on average, and the
~~annual~~ AE is less than ~~3-10~~ W m^{-2} (Table 4). This is due, in part, to the use of the same values
of emissivity for soil, snow and vegetation in ISBA and MEB. It is also due to the fact that $LW \downarrow$
305 is the same for both models. Note that another parameter which can have an impact on LW_{net} is
the ratio of the roughness length for momentum to that for heat, which is commonly referred to
as $k B^{-1} = \ln(z_0/z_{0h})$. In the default version of ISBA, the roughness length ratio is defined as
 $z_0/z_{0h} = 10$ (which corresponds to $k B^{-1} \approx 2.3$), and this value has been determined as a value
which works well generally (although for local scale applications, values can be prescribed to range
310 from 1 to on the order of 1×10^2). But note that there is evidence that $k B^{-1}$ is actually closer to
1 for tall canopies (e.g., Yang and Friedl, 2003) which corresponds to a roughness length ratio of
approximately 2.7. A ratio of ~~one~~ 1 is used for forests within the default version of the original
two-source model in the RCA (Samuelsson et al., 2011) dual-energy budget LSM. Thus currently,
a ratio of one is used for MEB since it is closer to $k B^{-1} \approx 1$, in contrast to 10 for ISBA. The
315 larger roughness ratio in ISBA can be seen as a way of compensating for an under-estimated diurnal
cycle of the surface radiative temperature owing to a composite soil-vegetation heat capacity (which
is larger than that for a bulk vegetation layer as in MEB). This implies that the explicit canopy
representation in MEB allows us to use a slightly more realistic value of roughness length ratio for
forest canopies.

320 5.1.2 Latent heat flux

At Le Bray and Puéchabon, the simulated values of total LE are relatively close between the three
simulations and in fairly good agreement with the adjusted measurements. Above the forest, most of
the evapotranspiration comes from canopy transpiration (Fig. 4), which is modulated to a large extent
for the three sites by the root zone soil moisture. All three of the simulations use the same scheme
325 for the soil water and the stomatal resistance. ~~The simulations tend~~ Except with ISBA at Le Bray,
the simulations tend to slightly underestimate LE (negative AE) for both these sites and models
on an annual basis. One possible explanation is that the formulation of the stomatal resistance was
originally adapted for low vegetation types, such as crops (Calvet et al., 1998). At Barbeau, the LE
underestimation (bias) is larger. The likely reason is owing to the presence of a water table that was

330 not explicitly measured, but strong evidence for its existence can be seen in the measured soil water
(especially in winter and spring: Fig. 5e). Without an explicit groundwater parameterization, it is
not possible to model this water table accurately and development of such a scheme was beyond the
scope of the current study. But these very wet observed soil conditions resulted in mostly unstressed
conditions during this time frame, and this high level of soil water was not modeled by ISBA or
335 MEB.

For the Mediterranean forest at Puéchabon, most of the net radiation is converted into sensible
heat flux (Fig. 6a) which leads to ~~good results for the three simulations (Table 4)~~ low values of LE
in accordance with observations (fig. 6b) . However, the R^2 for LE is still improved by 14 % for
MEBL compared to ISBA (Table 4), which is related to the improved sensible heat flux simulation
340 (see Section 5.1.3 for more details). The RMSE scores are not very different between the three
simulations, although the results are slightly improved with MEB and MEBL (the RMSE is 7 %
lower with MEB and 10 % with MEBL compared to ISBA). The main difference between the three
simulations for this site is the ground evaporation. It is very small for ISBA, but this is largely due
to the class-dependent prescription of a vegetation fraction, veg , of 0.99 (Table 2). This severely
345 limits the baresoil evaporation for this particular class. This high value was defined for this class
within ISBA to avoid excessive baresoil evaporation in the composite scheme (amounting to a tuning
parameter) resulting from the use of an aggregated surface roughness (resulting in a relatively large
roughness for the ground) and to model in a very simplistic way the shading effect. For MEB, the
 veg parameter does not exist, and instead the partitioning between E_g and E_v is controlled more by
350 physical processes (largely by the annual cycle of LAI). The MEB E_g turns out to be significantly
larger than that for ISBA for this site, but MEBL produces an E_g which is closer to that of ISBA.

The two other sites have a higher annual evapotranspiration than the Mediterranean site and a
larger ground evaporation, especially for ISBA. At Le Bray, evapotranspiration is almost the same for
the three simulations (Fig. 4a), with a slight underestimation between September and April (Fig. 7b).
355 Ground evaporation was measured at this site from March 14 to April 6 in 1998 Lamaud et al. (2001),
leading to an estimate of the cumulative evaporation from the ground of approximately 7.2 kg m^{-2}
during this period. For each of the three available years of this study (2006-2008), MEBL simulates
an evaporation from the litter of around 6.5 kg m^{-2} during this period, whereas ISBA evaporates
 16 kg m^{-2} from the ground. This shows the ability of the litter model to limit ground evaporation
360 during this period.

At Barbeau, the interpretation of the results is much more complex because of the deciduous
broadleaf nature of the forest. In winter, the LAI is very low and latent heat flux is dominated by
the ground evaporation. In spring, when net radiation increases but vegetation is not fully developed,
MEB significantly overestimates LE owing to a large ground evaporation (Fig. 8b), in contrast to
365 MEBL, which limits surface evaporation to 1.5 times less owing to the litter layer. In summer, the
ISBA simulation appears to better simulate LE (in terms of total amount). However, taking a closer

look at the partitioning of evapotranspiration reveals that transpiration and evaporation from the canopy reservoir are almost the same but ground evaporation is very different compared to MEB and MEBL. ~~Both~~ During this period, both MEB simulations simulate considerably less ground evaporation (about 4% of summer LE) compared to the standard ISBA simulation which simulates 25 % of annual-summer LE as ground evaporation as shown in Fig. 4 (especially due to the summertime contribution). This value is certainly over-estimated since at this time of the year, very little energy is expected to reach the ground (the leaf area index-LAI is around $5 \text{ m}^2 \text{ m}^{-2}$). For example, during July, ISBA and MEBL evaporate 33.4 kg m^{-2} and 4.5 kg m^{-2} directly from the ground, respectively, whereas the total energy reaching the soil, is only equivalent to a maximum possible evaporation of 8.1 kg m^{-2} based on PAR measurements. Second, in the composite ISBA scheme, the roughness length, z_0 , used for ground evaporation is the same as that for transpiration, and it is computed using an inverse-log averaging between soil and vegetation values weighted by the fraction cover Noilhan et al. (1995). The forest cover in ISBA uses a fraction cover of 0.95, thus the effective surface roughness is dominated by the vegetation roughness and is nearly 2 orders of magnitude larger than a typical soil roughness (0.01 m, which is close to the MEB value) leading to a relatively low aerodynamic resistance. Thus, the ISBA model simulates a reasonable LE which is comparable to MEB owing to a weak transpiration compensating an excessive bare soil evaporation. This is one of the known biases which MEB was intended to reduce.

5.1.3 Sensible heat flux

For all of the sites, more significant differences occur for the simulated H compared to LE ; with the most significant differences occurring with MEBL (see Table 4). At Le Bray and Puéchabon, the observed H is in better agreement between the MEBL and MEB runs with an average RMSE over these two sites of 67.7 W m^{-2} with the ISBA, 49.5 W m^{-2} with MEB and 48 W m^{-2} for MEBL. In addition, the R^2 are improved with average of 0.82 for ISBA, 0.89 for MEB and 0.90 for MEBL. At Barbeau during summer, the MEB and MEBL simulations constantly over-estimate H . As shown in the previous section, LE is underestimated in particular for MEB and MEBL. As net radiation and ground heat flux (section 5.1.4) are well simulated, H tends to be over-estimated. ISBA produces fairly good results despite overestimations of the ground heat flux and ground evaporation.

The improvement for the two MEB simulations are mainly related to three processes;

- The use of an explicit canopy layer which intercepts most of the downward solar radiation thereby reducing the net radiation at the ground surface. This leaves more energy available for turbulent fluxes in contrast to ISBA which is directly connected to forest floor and can more easily propagate energy into the ground by conduction.
- The use of a lower roughness length ratio (momentum to heat: z_0/z_{0h}) which leads to an increase in the roughness length for heat and water vapor and therefore the turbulent fluxes.

- The presence of a litter layer limits the penetration of energy into the ground because of its low thermal diffusivity which leads to a reduction of the ground heat flux (amplitude) and thus to more available energy for H and LE .

405 5.1.4 Ground heat flux

A substantial effect of both MEB and MEBL is the ground heat flux reduction. The explicit representation of the canopy induces a shading effect due to the leaves, stems and branches, so that less energy reaches the ground. But in addition, the explicit representation of the litter layer also modifies the ground heat flux by acting as a buffer for the top soil layer due to its low thermal diffusivity. It can be seen in Fig. 9 that for each site, MEB and MEBL had improved simulations of G in terms of amplitude and phase with regards to ISBA. The mean AE of the three experiments remains very low (Table 4), even for the ISBA model. This demonstrates that the two MEB experiments reduce the amplitude and improve the phasing of this flux while its daily average was reasonably well represented with the standard ISBA model. The mean RMSE averaged over all years and sites is 11.7 W m⁻² with the explicit litter, 24.3 W m⁻² without it and 47.1 W m⁻² using ISBA (the scores are summarized in Table 4).

The ISBA overestimation of the flux amplitude often represents several 10's of W m⁻² during both daytime and nighttime (Figs 7c; 6c; 8c), which reduces the amount of energy available for turbulent fluxes. For all sites, improvements of the ground heat flux are the same from one season to another which means that using a constant thickness for the litter layer is likely adequate. The MEBL overestimates the reduction of G from June to September only for the Le Bray site (Fig. 7c). However, the presence of a very dense understory at this site (accounting for half of the LAI over this period) makes the radiative transfer modeling much more complicated and could potentially lead to an underestimation of the incoming radiation at the forest floor. The understory contribution to the shortwave fluxes is simply represented using a bulk overstory-understory aggregated LAI . In contrast to the other two sites, the effect of shading and litter at Puéchabon are not sufficient to reduce the G significantly (to be more in line with observed values) no matter what the season. The parameters of the radiative transfer scheme have been calibrated through the modification of the clumping index so that the energy passing through the canopy is well modeled (Fig 3b). Sensitivity tests have been done for which the ground albedo and soil texture were modified over realistic ranges, but this did not resolve the issue, and so the over-estimated G is likely due to other causes (such as the need to better adjust the turbulent transfer parameters to better correspond to the vegetation at this site). At any rate, the MEBL G is still superior to that simulated using MEB and ISBA for this site. Finally, at Barbeau, the MEBL simulation produces good results for G (Fig. 8c) except for a short period in winter when the shortwave radiation transmission is also over-estimated (see Fig. 3c).

5.1.5 Soil temperature

A good description of the soil thermal characteristics is needed to model temperatures at different depths, along with a correspondingly good estimate of the surface soil heat flux. The soil characteristics (thermal conductivity and heat capacity) are calculated based on the input soil texture (sand and clay fractions) and organic matter contained in the soil. The soil temperature simulation statistics in Table 5 were computed for the three model configurations using available measurements at each site [and at different depth](#). MEBL and MEB have considerably lower RMSE of soil temperatures for each site compared to ISBA. The RMSE of the first measured depth of each site (between 0.04 and 0.10 m) is 10 to 20 % lower for MEB compared to the ISBA model, and 20 to 60 % lower between MEBL and ISBA. Moreover, R^2 is also improved with an average value for all of the sites of 0.96 for MEBL, 0.94 with MEB and 0.90 for ISBA. The good agreement between simulated and measured temperatures and the improvements with MEB are consistent with the improvement of the ground heat flux shown in the previous section (Fig. 9). Note, however, that the improvement of the simulated temperature at Puéchabon is not as significant as for the other sites. This was expected since, as shown in the previous section, the G was over-estimated for this site: the RMSE is reduced with the MEB and MEBL experiments, but it remains high (3.8 and 3.3 K, respectively). In general, there are relatively low AE values for each site and soil depth ($AE < 1.3$ K), but a seasonally varying AE remains. The general effect of MEB is to reduce the mean annual temperature by 1 to 2 K (resulting in a shift in the annual cycle), while MEBL dampens the annual cycle amplitude. Indeed, the litter reduces both the energy loss during the cold season and gains in heat during the summer season. The composite monthly average diurnal cycles of the uppermost available measured temperature for each site is shown in Fig. 10. The relatively low AE calculated over the entire year (Table 5) masks a negative AE during the cold season (November to March) and a positive AE during the warm season (May to September). At all sites, both MEB simulations improve the mean daily values during the warm season, but only MEBL reduces the negative bias that occurs during the cold season. So, when looking at the composite diurnal cycles, it is obvious that the use of MEBL versus MEB improves the performance in terms of amplitude and phase all year long even if a relatively small bias remains. The ISBA model daytime overestimation can be as high as 10 K, whereas with the MEBL model it does not exceed 3 K.

5.1.6 Soil moisture

The observed and simulated soil water content time series for an annual cycle are displayed in Fig. 5. Soil volumetric water content measures are available at 30 minute time steps at Le Bray and Barbeau. At Puéchabon, no half-hourly measurements were available but a reference curve, which corresponds to a model-derived interpolation of discrete soil water storage measurement (Lempereur et al., 2015), has been added to the figure. A fairly good overall agreement between simulated and observed total

soil water content is found. Moreover, the three experiments lead to very similar results, which is consistent with the fact that evapotranspiration has been found to be the least impacted flux between ISBA, MEB and MEBL.

More significant differences can be seen in terms of the top soil water content. In particular, for the Le Bray and Barbeau forests, at the end and beginning of the year, respectively, ISBA and MEB simulate unrealistic drops in the liquid soil water content (Fig. 5, b and f). These drops are caused by the underestimation of the soil temperature during these periods which makes the water freeze. These drops are not present in the MEBL simulation. In fact, the near-surface soil temperature stays above 0 deg C owing to an insulating effect from the litter so that the water remains liquid as seen in the measurements. At Barbeau, all the three model approaches underestimate the total soil moisture during the first two thirds of the year (Fig. 5e) which is mainly related to the presence of a water table which is not explicitly modeled. MEB and MEBL are more sensitive than ISBA to this process since they simulate more transpiration than ISBA (although MEBL is slightly wetter since E_g is always less in MEBL compared to MEB).

5.2 Sensitivity tests

There is uncertainty with respect to the definition of several key model parameters which are not usually available in the observations so that several sensitivity tests have been undertaken. Three parameters have been tested; i) the extinction coefficient for the long wave transmission through the canopy (Boone et al., 2017), ii) the leaf geometry which modulates the solar radiation transmitted through the canopy (Carrer et al., 2013), and iii) the litter thickness which controls the impact of litter on both the hydrological and thermal regimes (see Appendix A).

For all three of the sites, the long wave transmission coefficient has a very weak influence on each of the simulations for a reasonable range of values (0.3 to 0.7, 0.5 being the default value which is based on a value used by Versegny et al. (1993). The RMSE calculated with observed and simulated total long wave up radiation varies less than 2 W m^{-2} over this range for the 3 sites on an annual basis.

The litter layer thickness values tested range between 0.01 and 0.10 m based on values from the literature (Table 3). The MEBL default values based on recommendations from the site principal investigators are shown in Table 1. The G is the most sensitive to these changes and RMSE values for these tests are shown on Fig. 11 for Le Bray and Barbeau. Note that these tests were not done for Puéchabon since G was overestimated despite the litter layer (which implies that optimizing litter thickness for these site to improve fluxes would likely result as a compensating error related to another process). Minimum errors values are reached for thicknesses above approximately 0.03 m which is consistent with the recommended site values and those from several other studies (see Table 3). Note that these thickness values also produce the lowest H and LE RMSE (not shown).

The clumping index was tested since it is the key parameter of the radiative transfer controlling the transmission and absorption of incoming short wave radiation through the canopy (Carrer et al., 2013). Three configurations were tested for the MEBL configuration; i) the default value, ii) a clumping index increased by 50 % (a more closed canopy) and iii) an index decreased by 50 %. This parameter alters the computation of the upwelling shortwave radiation, and the values of the turbulent and ground heat flux since it changes the partitioning of available energy between the canopy and the litter layer. However, the H and LE RMSE only changed by 3 to 8 % depending on site, thus for now, the default clumping index values are used.

In summary, owing to these sensitivity tests, the default extinction coefficient for long wave transmission is retained as a constant value of 0.5. A default constant value of litter thickness is defined as 0.03 m since it is both not widely observed and it tends to be a threshold for which the effect of litter on G becomes notable and positive (and errors begin to slowly increase slightly above this threshold) for several contrasting sites. One could even define a slightly larger value based on Fig. 11 (0.04 m for example), but a first order approximation is required as opposed to a tuned value based on just two sites. Also, the layer should be relatively thin to insure a robust diurnal cycle (as opposed to using values on the high side from these tests or based on the literature). The default clumping index used in the shortwave radiative transfer led to an over-estimation of the below-canopy radiation for the two sites where measurements of below canopy PAR were available, and slight improvements were found by adjusting this parameter (to values still within its realistic range). But in fact, the impact of tuning the clumping index was found to have only a relatively small impact on the simulated turbulent heat fluxes with the MEBL simulation so that the default value is considered to be robust enough for now. As a final note, if measured values are available or can be readily determined for the longwave transmission coefficient, litter thickness, and shortwave radiation transmission (for example, at local scale well-instrumented sites for studies oriented towards detailed process analysis), they can be used to replace the aforementioned default values.

5.3 Worldwide skill score assessment

The comparison of modeling results with field measurements from over 42 Fluxnet sites (Fig. 2) reveals that the results from the benchmark are consistent with those from the three local French sites. Only For improved clarity, only the MEBL and the ISBA models are considered here compared here, since the previous evaluation showed the consistent improvement of using the litter option when using MEB for forests.

MEBL and ISBA generally performed well in simulating the R_{net} , but both tended to slightly underestimate daytime values with a mean bias of 10 W m^{-2} corresponding to approximately 9 % of the average net radiation. Note that the R_{net} was slightly better simulated for the three local scale sites (Sec. 5.1), but this results in part, because the benchmark uses values from the ECOCLIMAP database (which likely introduces some error relative to using observed values

of albedo for example). Scatter plot of the statistical metrics (RMSE, R^2 and AE) obtained for the sensible and the latent heat fluxes are shown in Fig. 12 and Fig. 13, respectively. Each black dot corresponds to the average score over all available years for a given site. The shaded area corresponds to the sites and years for which the MEBL model outperforms the ISBA model and the dashed lines represent the average over all sites. Finally, the percentage of sites for which MEBL was superior to ISBA for a particular score is given in the title of each panel within the aforementioned figures.

The sensible heat flux AE (Fig. 12c) is quite similar for the two model approaches as evidenced by the relatively low scatter (most of the values are within -20 to 20 W m^{-2}). This is consistent with previous conclusions for the detailed local scale run analysis (Table 4) and confirms that the influence of the multi-source model is more significant for diurnal cycles than for daily average results. However, a very significant improvement of the sensible heat flux R^2 is observed and for 93.5 % of the sites for MEBL. This shows the ability of MEBL, as demonstrated in Sec. 5.1, to better represent the phase of turbulent fluxes owing to the explicit representation of the canopy layer and especially the more realistic lower value of the heat capacity of vegetation with respect to ISBA.

The amplitude of the sensible heat flux diurnal cycle is also improved as suggested by the improved RMSE values (Fig. 12a) over 90.5 % of the sites. This better amplitude performance is related to both the canopy shading and litter insulation effects which lower the diurnal amplitude of the ground heat flux and provide more energy for the turbulent fluxes. The recent so called PLUMBER experiment Best et al. (2015) found that for H , most of the LSMs they studied (including ISBA) were not able to beat the one variable non-linear regression that uses instantaneous downward short-wave radiation and observed surface fluxes. The results in this study present a possible reason for this (since most of the LSMs do not include an explicit litter layer), therefore we recommend that the ground heat flux should be better investigated and could be contributing to the generally poor performance of simulating H among LSMs for the Fluxnet sites.

The latent heat flux, LE , results between the two models are more similar than for H . Based on the results from Sec. 5.1, this was expected since the vapor flux is mainly controlled by the stomatal resistance parametrization and soil hydrology (which are the same for MEBL and ISBA). Fig. 13a shows that the differences in RMSE between MEBL and ISBA are relatively small between the two simulations. However, an improvement in the R^2 coefficient is obtained (Fig. 13b), with 73.3 % of the sites showing an improvement with MEBL. Note that a negative AE is obtained for most of the sites. This result is also consistent with the local evaluation over the three French sites (Table 4 and as shown in Fig. 4). Since both the ISBA and MEBL approaches have similar errors in this respect, it is assumed that this is likely caused by an aspect of ISBA which is common among the two models, such as root-zone water uptake, hydrology, or stomatal resistance for example. In a general, a very good consistency is found between the analysis in Sec. 5.1 for highly instrumented sites and the more general benchmark application in this section. This is a positive result for a model which is designed for weather forecast and climate simulations at a global scale.

Finally, it is of interest to determine if the results are conditioned by certain key physiographic parameters, notably the LAI . The differences in RMSE between MEBL and ISBA are shown in Fig. 14 for different ranges of LAI as box plots. Each box is plotted using the 10, 25, 50, 75 and 90 percentiles of the normalized difference in RMSE calculated over one-month periods for each site that satisfies the closure condition adopted for this study (i.e. less than 20 % imbalance). In Fig. 14a, the improvement of RMSE is significant for forested sites with a relatively low LAI ($LAI < 2$) which represents 19 % of the cases, moderate for medium LAI ($2 < LAI < 4$), 46 % of the cases, and weak for high LAI ($LAI > 4$), 35 % of the cases. This shows that the gains of the multi source energy budget are more significant when the canopy is relatively sparse or open (corresponding to a relatively low LAI), so that the surface fluxes have significant contributions from both the ground and the canopy. When there is a medium LAI range, the same conclusion applies to a lesser extent. Thus as LAI becomes large, the MEBL and ISBA results converge (since the fluxes are dominated by the canopy). Fig. 14 is consistent with previous results in that the simulated LE is similar between MEBL and ISBA so that RMSE differences are low on average.

6 Conclusions

This study is the second of a set of two papers which describe the introduction of the new multi energy balance (MEB) approach within the interactions between the soil biosphere atmosphere model (ISBA) as part of the SURFEX platform. Two new explicit bulk layers have been implemented, one for the vegetation canopy and the other for a litter layer. This paper describes a two-part local-scale offline evaluation of both the bulk canopy scheme (MEB) and the combined bulk canopy-litter layer (MEBL) approaches, and the results are also compared to the standard composite-vegetation ISBA model. The model parameterization governing the litter layer is also presented. The evaluation is done by investigating the ability of the models to simulate the fluxes above (sensible and latent heat flux) and below (ground heat flux) different forest canopies and the ability of the different approaches to reproduce the observed soil temperatures and soil water content.

In the first part of this study, an evaluation over three well-instrumented forested sites in France was done using observed forest characteristics, and turbulent, radiative and heat conduction (ground flux) measurements with a particular attention paid to the energy balance closure issue. The mid-latitude forest sites were contrasting in terms of both vegetation type (needleleaf, broadleaf) and climate (Mediterranean, temperate). In the second part of this study, a statistical evaluation was done using the framework of a benchmark platform based upon 42 sites scattered throughout the world from the Fluxnet network.

In terms of the model evaluation for the three French forested sites, the standard ISBA model was found to underestimate the amplitude of the sensible heat flux, H , and overestimate that of the ground heat flux, G , the latter of which is in agreement with an overestimation of the soil temperature

amplitude. Also, latent heat flux is generally well simulated despite being slightly underestimated.

615 The simulation with the two new explicit layers greatly reduced the G and soil temperature errors owing mainly to shading effect of the canopy layer and the low thermal diffusivity of the litter layer. The relatively low thermal conductivity of the litter greatly reduced the ground heat flux and soil temperature amplitudes, in addition to modifying their phase in better agreement with the measurements: the RMSE decreased from 47.1 to 10.0 W m^{-2} on average. As a result, more energy was
620 available for the turbulent heat fluxes. Since latent heat flux was generally water limited, most of this excess energy went into H : the average RMSE for the three well-instrumented sites decreased from 62.1 to 50 W m^{-2} with the new parameterizations. This result might have importance beyond those for ISBA, since the results of the recent PLUMBER (Best et al., 2015) LSM inter-comparison and evaluation experiment highlighted the general poor performance of LSM-estimated H (including
625 ISBA) compared to a subset of the Fluxnet observations. Since net radiation and latent heat fluxes were generally reasonably well simulated, this implies that one of the main sources of error in H is likely related to the ground heat flux simulation based on the energy budget equation. This study showed that at least for ISBA (using the MEBL option), the inclusion of litter lead to significantly improved ground conduction which resulted in better sensible heat fluxes.

630 In terms of temporal dynamics, the main differences in latent heat flux between ISBA and MEBL occur during spring for the deciduous forest site where the litter layer acts to significantly limit soil evaporation, whereas ISBA and MEB (without explicit litter) overestimate evapotranspiration due to strong ground evaporation (owing to a relatively low LAI combined with large incoming radiation and generally low to unstressed soil conditions). And despite the overall similar total annual evap-
635 otranspiration simulated by ISBA and MEBL, the partitioning between canopy evapotranspiration, E_v , and ground evaporation, E_g , is more realistic with MEBL than ISBA. This is mainly related to the conceptual design of the composite vegetation, which uses a single aggregated roughness length for both vegetation and soil (which leads to a slight underestimate of canopy roughness but a significant overestimate of ground roughness and thus relatively large E_g compared to the explicit bulk
640 canopy approach of MEB). In addition, the structure of MEB enables the model to directly use the radiation transmitted through the canopy when computing the ground surface energy budget. Measurements from two of the three well-instrumented forest sites indicate that MEBL simulates the downwelling shortwave radiation relatively well, and this sets a theoretical limit on the amount of energy available for the turbulent fluxes at the surface (which again, is often lower than the energy
645 available for E_g in the composite ISBA scheme). The more physically-based vegetation canopy in MEB also implies that there is no longer a dependence on the ISBA parameter veg (vegetation cover fraction), which was constant for forests and tuned (to values between 0.95 and 0.99) to prevent excessive E_g .

The main conclusions made from analysis of the three well-instrumented sites were found to be
650 consistent with the results of a statistical benchmark analysis over a subset of 42 forested sites from

the Fluxnet network. The sensible heat flux RMSE was improved for 87.5 % of these sites with the new parameterizations (MEBL). The selected sites encompassed a wide range of climate and several different forest land-cover types, thus it is a necessary test before implementing MEBL in regional to global scale applications. The benchmark also showed that the impact of the explicit
655 treatment of the canopy and litter layer was more significant for relatively open canopies (low to medium LAI values), whereas for closed canopies (high LAI), all three of the model approaches simulation results converge. This is not overly surprising since in the limit as a canopy becomes tall and quite dense, the composite scheme resembles a vegetation canopy (the soil contribution becomes significantly less). But again, ISBA tends to simulate considerably more baresoil evaporation in the
660 peak growing season (maximum LAI) than MEBL, so there is error compensation which is masked to a large extent when looking at the total evapotranspiration.

In terms of prospectives, evaluation of MEB/MEBL is ongoing, and offline spatially distributed applications within ~~SIM (Habets et al., 2008)~~ the SIM chain (SAFRAN-ISBA-MODCOU, Habets et al., 2008) and at the global scale are being tested and compared to ISBA, along with an examination of
665 the improvements obtained in high latitudes (owing, in part, to a more detailed representation of snow interception processes). Also, additional local scale tests are being done for both tropical (since there were relatively few such sites in Fluxnet) and semi-arid forests (owing to the impact of MEB for the latter as shown herein). Finally, experiments are being prepared for fully coupled land-atmosphere simulation tests in both mesoscale and climate model applications at Météo-France.

670 7 Code Availability

The MEB code is a part of the ISBA LSM and is available as open source via the surface modelling platform called SURFEX, which can be downloaded at <http://www.cnrm-game-meteo.fr/surfex/>. SURFEX is updated at a relatively low frequency (every 3 to 6 months) and the developments presented in this paper are available starting with SURFEX version 8.0. If more frequent updates are needed,
675 or if what is required is not in Open-SURFEX (DrHOOK, FA/LFI formats, GAUSSIAN grid), you are invited to follow the procedure to get a SVN account and to access real-time modifications of the code (see the instructions at the previous link).

Acknowledgements. This work is a contribution to the ongoing efforts to improve the SURFEX platform. The authors would like to thank the teams of the three French eddy flux towers involved in this study: Le Bray,
680 Barbeau and Puéchabon, for making their data available, and to all of the contributors to the Fluxnet data base. This study was fully funded by a Météo-France grant.

References

- Andrade, J. A. V., Abreu, F. M. G. d., and Madeira, M. A. V.: Influence of litter layer removal on the soil thermal regime of a pine forest in a Mediterranean climate, *Rev. Bras. Ciênc. Solo*, 34, 1481–1490, 2010.
- 685 Arunachalam, A. and Arunachalam, K.: Influence of gap size and soil properties on microbial biomass in a subtropical humid forest of north-east India, *Plant Soil*, 223, 187–195, 2000.
- Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., et al.: FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities, *B. Am. Meteorol. Soc.*, 82, 2415–2434, 2001.
- 690 Best, M., Abramowitz, G., Johnson, H., Pitman, A., Balsamo, G., Boone, A., Cuntz, M., Decharme, B., Dirmeyer, P., Dong, J., et al.: The plumbing of land surface models: benchmarking model performance, *J. Hydrometeorol.*, 16, 1425–1442, 2015.
- Blyth, E., Gash, J., Lloyd, A., Pryor, M., Weedon, G. P., and Shuttleworth, J.: Evaluating the JULES land surface model energy fluxes using FLUXNET data, *J. Hydrometeorol.*, 11, 509–519, 2010.
- 695 Boone, A. and Etchevers, P.: An intercomparison of three snow schemes of varying complexity coupled to the same land-surface model: Local scale evaluation at an Alpine site, *J. Hydrometeorol.*, 2, 374–394, 2001.
- Boone, A., Calvet, J., and Noilhan, J.: Inclusion of a third soil layer in a land surface scheme using the force-restore method, *Journal of Applied Meteorology*, 38, 1611–1630, 1999.
- Boone, A., Masson, V., Meyers, T., and Noilhan, J.: The influence of the inclusion of soil freezing on simulations by a soil-vegetation-atmosphere transfer scheme, *J. Appl. Meteorol.*, 9, 1544–1569, 2000.
- 700 Boone, A., Samuelsson, P., Gollvik, S., Napoly, A., Brun, E., Decharme, B., and Co-Authors: The Interactions between Soil-Biosphere-Atmosphere (ISBA) land surface model Multi-Energy Balance (MEB) option in SURFEX, *Geosci. Model Dev.*, 2017.
- Bulcock, H. and Jewitt, G.: Modelling canopy and litter interception in commercial forest plantations in South Africa using the Variable Storage Gash model and idealised drying curves, *Hydrol. Earth Syst. Sc.*, 16, 4693–4705, 2012.
- 705 Calvet, J.-C., Noilhan, J., Roujean, J.-L., Bessemoulin, P., Cabelguenne, M., Olioso, A., and Wigneron, J.-P.: An interactive vegetation SVAT model tested against data from six contrasting sites, *Agr. For. Meteorol.*, 92, 73–95, 1998.
- 710 Carrer, D., Roujean, J.-L., Lafont, S., Calvet, J.-C., Boone, A., Decharme, B., Delire, C., and Gastellu-Etchegorry, J.-P.: A canopy radiative transfer scheme with explicit FAPAR for the interactive vegetation model ISBA-A-gs: Impact on carbon fluxes, *J. Geophys. Res.*, 118, 888–903, 2013.
- Choudhury, B. J. and Monteith, J. L.: A four-layer model for the heat budget of homogeneous land surfaces, *Q. J. Roy. Meteor. Soc.*, 114, 373–398, 1988.
- 715 Clapp, R. and Hornberger, G.: Empirical equations for some soil hydraulic properties, *Water Resour. Res.*, 14, 601–604, 1978.
- De Vries, D. A.: Thermal properties of soils, *Physics of plant environment*, 1963.
- Decharme, B. and Douville, H.: Introduction of a sub-grid hydrology in the ISBA land surface model., *Clim. Dynam.*, 26, 65–78, 2006.
- 720 Decharme, B., Boone, A., Delire, C., and Noilhan, J.: Local evaluation of the Interaction between Soil Biosphere Atmosphere soil multilayer diffusion scheme using four pedotransfer functions, *J. Geophys. Res.*, 116, 2011.

- Decharme, B., Brun, E., Boone, A., Delire, C., Moigne, P. L., and Morin, S.: Impacts of snowpack properties and soil organic carbon content on characteristics and soil temperature profiles simulated by the ISBA land surface model, *Cryosphere*, 10, 853–877, doi:10.5194/tc-10-853-2016, 2016.
- 725 Er-Raki, S., Chehbouni, A., Hoedjes, J., Ezzahar, J., Duchemin, B., and Jacob, F.: Improvement of FAO-56 method for olive orchards through sequential assimilation of thermal infrared-based estimates of ET, *Agr. Water Manage.*, 95, 309–321, 2008.
- Farouki, T. O., ed.: *Thermal Properties of Soils*, vol. 11, Clausthal-Zellerfeld, Germany: Trans Tech, 1986.
- Faroux, S., Tchuente, A. T. K., Roujean, J.-L., Masson, V., Martin, E., and Moigne, P. L.: ECOCLIMAP-
730 II/Europe: a twofold database of ecosystems and surface parameters at 1 km resolution based on satellite information for use in land surface, meteorological and climate models, *Geosci. Model Dev.*, 6, 563–582, doi:10.5194/gmd-6-563-2013, 2013.
- Gerrits, A., Savenije, H., Hoffmann, L., and Pfister, L.: New technique to measure forest floor interception: an application in a beech forest in Luxembourg, *Hydrol. Earth Syst. Sc.*, 11, 695–701, doi:10.5194/hess-11-
735 695-2007, 2007.
- Giard, D. and Bazile, E.: Implementation of a new assimilation scheme for soil and surface variables in a global NWP model, *Monthly Weather Review*, 128, 997–1015, 2000.
- Gibelin, A.-L., Calvet, J. C., Roujean, J.-L., Jarlan, L., and Los, S. O.: Ability of the land surface model ISBA-A-gs to simulate leaf area index at the global scale: Comparison with satellites products, *J. Geophys. Res.*,
740 111, doi:10.1029/2005JD006691, d18102, 2006.
- Gonzalez-sosa, E., Braud, I., Jean-Louis, T., Michel, V., Pierre, B., and Jean-Christophe, C.: Modelling heat and water exchanges of fallow land covered with plant-residue mulch, *Agr. For. Meteorol.*, 97, 151–169, 1999.
- Guevara-Escobar, A., Gonzalez-Sosa, E., Ramos-Salinas, M., and Hernandez-Delgado, G.: Experimental analysis of drainage and water storage of litter layers, *Hydrol. Earth Syst. Sc.*, 11, 1703–1716, 2007.
- 745 Habets, F., Boone, A., Champeaux, J., Etchevers, P., Franchisteguy, L., Leblois, E., Ledoux, E., Moigne, P. L., Martin, E., Morel, S., et al.: The SAFRAN-ISBA-MODCOU hydrometeorological model applied over France, *J. Geophys. Res.*, 113, doi:10.1029/2007JD008548, 2008.
- Haila, Y., Niemelä, J., et al.: Leaf litter and the small-scale distribution of carabid beetles (Coleoptera, Carabidae) in the boreal forest, *Ecography*, 22, 424–435, 1999.
- 750 Haverd, V. and Cuntz, M.: Soil–Litter–Iso: A one-dimensional model for coupled transport of heat, water and stable isotopes in soil with a litter layer and root extraction, *J. Hydrol.*, 388, 438–455, 2010.
- Kostel-Hughes, F., Young, T. P., and Carreiro, M. M.: Forest leaf litter quantity and seedling occurrence along an urban-rural gradient, *Urban Ecosys.*, 2, 263–278, doi:10.1023/A:1009536706827, 1998.
- Lamaud, E., Ogée, J., Brunet, Y., and Berbigier, P.: Validation of eddy flux measurements above the understorey
755 of a pine forest, *Agr. For. Meteorol.*, 106, 187–203, 2001.
- Lempereur, M., Martin-StPaul, N. K., Damesin, C., Joffre, R., Ourcival, J.-M., Rocheteau, A., and Rambal, S.: Growth duration is a better predictor of stem increment than carbon supply in a Mediterranean oak forest: implications for assessing forest productivity under climate change, *New Phytologist*, 207, 579–590, 2015.
- Marsh, D. and Pearman, P.: Effects of habitat fragmentation on the abundance of two species of Leptodactylid
760 frogs in an Andean montane forest, *Conserv. Biol.*, 11, 1323–1328, 1997.

- Masson, V., Moigne, P. L., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouyssel, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Khatib, R. E., Essaouini², K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafont, S., Lebeaupin, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Muktari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of Earth surface variables and fluxes, *Geosci. Model Dev.*, 6, 929–960, doi:10.5194/gmd-6-929-2013, 2013.
- 765 Meekins, J. F. and McCarthy, B. C.: Effect of environmental variation on the invasive success of a nonindigenous forest herb, *Ecol. Appl.*, 11, 1336–1348, doi:10.1890/1051-0761(2001)011[1336:EOEVOT]2.0.CO;2, 2001.
- 770 Nachtergaele, F. and Batjes, N.: Harmonized world soil database, FAO Rome, Italy, 2012.
- Noilhan, J. and Mahfouf, J.-F.: The ISBA land surface parameterisation scheme, *Global Planet. Change*, 13, 145–159, 1996.
- Noilhan, J. and Planton, S.: A simple parameterization of land surface processes for meteorological models, *Mon. Wea. Rev.*, 117, 536–549, 1989.
- 775 Noilhan, J., and Lacarrere, P.: GCM grid-scale evaporation from mesoscale modeling, *J. Climate*, 8, 206–223, 1995.
- Ogée, J. and Brunet, Y.: A forest floor model for heat and moisture including a litter layer, *J. Hydrol.*, 255, 212–233, 2002.
- Oliphant, A., Grimmond, C., Zutter, H., Schmid, H., Su, H.-B., Scott, S., Offerle, B., Randolph, J., and Ehman, J.: Heat storage and energy balance fluxes for a temperate deciduous forest, *Agr. For. Meteorol.*, 126, 185–201, 2004.
- 780 Park, H.-T., Hattori, S., and Tanaka, T.: Development of a numerical model for evaluating the effect of litter layer on evaporation, *Journal of Forest Research*, 3, 25–33, 1998.
- Prévost-Bouré, N. C., Ngao, J., Berveiller, D., Bonal, D., Damesin, C., Dufrière, E., Lata, J.-C., Le Dantec, V., Longdoz, B., Ponton, S., et al.: Root exclusion through trenching does not affect the isotopic composition of soil CO₂ efflux, *Plant Soil*, 319, 1–13, 2009.
- 785 Putuhena, W. M. and Cordery, I.: Estimation of interception capacity of the forest floor, *J. Hydrol.*, 180, 283–299, 1996.
- Rambal, S., Lempereur, M., Limousin, J., Martin-StPaul, N., Ourcival, J., and Rodríguez-Calcerrada, J.: How drought severity constrains gross primary production (GPP) and its partitioning among carbon pools in a *Quercus ilex* coppice?, *Biogeosciences*, 11, 6855–6869, 2014.
- 790 Riha, S. J., McInnes, K., Childs, S., and Campbell, G.: A finite element calculation for determining thermal conductivity, *Soil Science Society of America Journal*, 44, 1323–1325, 1980.
- Rutter, N., Essery, R., Pomeroy, J., Altimir, N., Andreadis, K., Baker, I., Barr, A., Bartlett, P., Boone, A., Deng, H., Douville, H., Dutra, E., Elder, K., Ellis, C., Feng, X., Gelfan, A., Goodbody, A., Gusev, Y., Gustafsson, D., Hellström, R., Hirabayashi, Y., Hirota, T., Jonas, T., Koren, V., Kuragina, A., Lettenmaier, D., Li, W.-P., Luce, C., Martin, E., Nasonova, O., Pumpanen, J., Pyles, R. D., Samuelsson, P., Sandells, M., Schädler, G., Shmakin, A., Smirnova, T. G., Stähli, M., Stöckli, R., Strasser, U., Su, H., Suzuki, K., Takata, K., Tanaka, K., Thompson, E., Vesala, T., Viterbo, P., Wiltshire, A., Xia, K., Xue, Y., and Yamazaki, T.: Evaluation of forest snow processes models (SnowMIP2), *J. Geophys. Res.*, 114, doi:10.1029/2008JD011063, 2009.
- 800

- Sakaguchi, K. and Zeng, X.: Effects of soil wetness, plant litter, and under-canopy atmospheric stability on ground evaporation in the Community Land Model (CLM3. 5), *J. Geophys. Res.*, 114, 2009.
- Samuelsson, P., Jones, C., Willén, U., Ullerstig, A., Gollvik, S., Hansson, U., Jansson, C., Kjellström, E., Nikulin, G., and Wyser, K.: The Rossby Centre Regional Climate Model RCA3: Model description and performance, *Tellus A*, 63, 1–3, doi:10.1111/j.1600-0870.2010.00478.x, 2011.
- 805 Sato, Y., Kumagai, T., Kume, A., Otsuki, K., and Ogawa, S.: Experimental analysis of moisture dynamics of litter layers - the effects of rainfall conditions and leaf shapes, *Hydrol. Process.*, 18, 3007–3018, 2004.
- Schaap, M. and Bouten, W.: Forest floor evaporation in a dense Douglas fir stand, *J. Hydrol.*, 193, 97–113, 1997.
- 810 Séférian, R., Gehlen, M., Bopp, L., Resplandy, L., Orr, J., Marti, O., Dunne, J., Christian, J., Doney, S., Ilyina, T., et al.: Inconsistent strategies to spin up models in CMIP5: implications for ocean biogeochemical model performance assessment, 2015.
- Seity, Y., Brousseau, P., Malardel, S., Hello, G., Bénard, P., Bouttier, F., Lac, C., and Masson, V.: The AROME-France convective-scale operational model, *Monthly Weather Review*, 139, 976–991, 2011.
- 815 Stöckli, R., Lawrence, D., Niu, G.-Y., Oleson, K., Thornton, P., Z.-L., Bonan, G., Denning, A., and Running, S.: Use of FLUXNET in the Community Land Model development, *Journal of Geophysical Research*, 113, 2008.
- Twine, T. E., Kustas, W., Norman, J., Cook, D., Houser, P., Meyers, T., Prueger, J., Starks, P., and Wesely, M.: Correcting eddy-covariance flux underestimates over a grassland, *Agr. For. Meteorol.*, 103, 279–300, 2000.
- 820 Ukkola, A., Pitman, A., Decker, M., Kauwe, M. D., Abramowitz, G., Kala, J., and Wang, T.: Modelling evapotranspiration during precipitation deficits: identifying critical processes in a land surface model, *Hydrol. Earth Syst. Sc.*, 12, 10 789–10 825, 2015.
- Versegny, D. L., McFarlane, N. A., and Lazare, M.: CLASS – A Canadian land surface scheme for GCMs, II. Vegetation model and coupled runs, *Int. J. Climatol.*, 13, 347–370, 1993.
- 825 Voldoire, A., Sanchez-Gomez, E., Méliá, D. S., B. Decharme, C. C., Sénési, S., Valcke, S., Beau, I., Alias, A., Chevallier, M., Déqué, M., Deshayes, J., Douville, H., E. Fernandez, G. M., Maisonnave, E., Moine, M. P., Planton, S., D. Saint-Martin, S. S., Tyteca, S., Alkama, R., Belamari, S., A. Braun, L. C., and Chauvin, F.: The CNRM-CM5.1 global climate model: description and basic evaluation, *Clim. Dynam.*, 40, 2091–2121, doi:10.1007/s00382-011-1259-y, 2013a.
- 830 Voldoire, A., Sanchez-Gomez, E., Méliá, D. S., Decharme, B., Cassou, C., Sénési, S., Valcke, S., Beau, I., Alias, A., Chevallier, M., et al.: The CNRM-CM5. 1 global climate model: description and basic evaluation, *Clim. Dynam.*, 40, 2091–2121, 2013b.
- Vorobeichik, E.: On the methods for measuring forest litter thickness to diagnose the technogenic disturbances of ecosystems, *Russian Journal of Ecology*, 28, 230–234, 1997.
- 835 Vuichard, N. and Papale, D.: Filling the gaps in meteorological continuous data measured at FLUXNET sites with ERA-interim reanalysis, *Earth System Science Data Discussions*, 8, 23–55, 2015.
- Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., Bernhofer, C., Ceulemans, R., Dolman, H., Field, C., et al.: Energy balance closure at FLUXNET sites, *Agr. For. Meteorol.*, 113, 223–243, 2002.

- 840 Wilson, T., Meyers, T., Kochendorfer, J., Anderson, M., and Heuer, M.: The effect of soil surface litter residue on energy and carbon fluxes in a deciduous forest, *Agr. For. Meteorol.*, 161, 134–147, 2012.
- Wu, H., Peng, C., Moore, T., Hua, D., Li, C., Zhu, Q., Peichl, M., Arain, M., and Guo, Z.: Modeling dissolved organic carbon in temperate forest soils: TRIPLEX-DOC model development and validation, *Geosci Model Dev Discuss*, 6, 3473–3508, 2013.
- 845 Yang, R. and Friedl, M. A.: Determination of roughness lengths for heat and momentum over Boreal forests, *Bound.-Lay. Meteorol.*, 107, 581–603, 2003.
- Zheng, D., Van Der Velde, R., Su, Z., Booi, M. J., Hoekstra, A. Y., and Wen, J.: Assessment of roughness length schemes implemented within the Noah land surface model for high-altitude regions, *J. Hydrometeorol.*, 15, 921–937, 2014.
- 850 Zhu, J., Matsuzaki, T., Lee, F., and Gonda, T.: Effect of gap size created by thinning on seedling emergence, survival and establishment in a coastal pine forest, *Forest Ecol. Manag.*, 182, 339–354, 2003.

Appendix A: Description of the litter layer model

Forest litter is represented using a single model layer which generally ranges in thickness from 0.01 to 0.10 m, and in the absence of ancillary data, the default value is 0.03 m. When this option
 855 is active, an additional layer is added to the soil for the thermal and energy budget computations with litter-specific thermal properties. This means that the numerical solution method is identical to that presented in the companion paper by Boone et al. (2017). In terms of hydrology, an additional reservoir is added which uses a relatively simple bucket-type scheme with a litter-specific maximum water storage capacity. The model physics and governing equations are reviewed herein.

860 A1 Prognostic equations

The energy budget for the snow-free litter layer can be expressed as:

$$C_l \frac{\partial T_l}{\partial t} = R_{nl} - H_l - LE_l - G_{g,1} + L_f \Phi_l \quad (\text{A1})$$

where T_l is the litter temperature (K), Δz_l (m) is the thickness of the litter layer, and C_l ($\text{J K}^{-1} \text{m}^{-2}$) is the effective heat capacity of the litter. $R_{n,l}$, H_l , LE_l , $G_{g,1}$ represent the net radiation, sensible
 865 heat flux, latent heat flux and ground conduction flux from the litter layer, respectively. Note that when litter is present, $R_{n,l}$, H_l , LE_l correspond to the ground fluxes in Boone et al. (2017). The additional terms added owing to a snow cover are described in Boone et al. (2017), and are therefore not repeated here.

The liquid water content of the litter layer evolves following:

$$870 \quad \frac{\partial W_l}{\partial t} = P_l - E_l - D_l - \Phi_l \quad (0 < W_l < W_{l,max}) \quad (\text{A2})$$

where P_l is the sum of the rates of the rainfall passing through the canopy, canopy drip and ground-based snow melt, E_l represents the litter evaporation rate, D_l is the drainage rate from the litter to

the soil (all in $\text{kg m}^{-2} \text{ s}^{-1}$). The maximum liquid water content in the litter reservoir is defined as $W_{l,max} = w_{l,max} \Delta z_l \rho_w$ (kg m^{-2}). The default value for the maximum holding capacity of the litter layer, $w_{l,max}$, is $0.12 \text{ m}^3 \text{ m}^{-3}$ (Putuhena and Cordery, 1996). All water in excess of this maximum value is then partitioned between infiltration and surface runoff by the ISBA soil hydrological model. The liquid water equivalent of ice contained in the litter layer is governed by:

$$\frac{\partial W_{lf}}{\partial t} = \Phi_l - E_{lf} \quad (\text{A3})$$

where E_{lf} represents the sublimation of ice contained within the litter layer.

880 A2 Phase Change

The phase change rate, Φ_l ($\text{kg m}^{-2} \text{ s}^{-1}$), is defined as:

$$\Phi_l = \frac{1}{\tau_{ice}} \left\{ \delta_f \min \left[\frac{\rho_i C_i \Delta z_l (T_l - T_f)}{L_f}, W_{lf} \right] + (1 - \delta_f) \min \left[\frac{\rho_i C_i \Delta z_l (T_f - T_l)}{L_f}, W_l \right] \right\} \quad (\text{A4})$$

where L_f represents the latent heat of fusion (J kg^{-1}), ρ_i is the density of ice (here defined as 920 kg m^{-3}), the freezing point temperature is $T_f = 273.15 \text{ K}$, and C_i is the specific heat capacity of ice ($2.106 \times 10^3 \text{ J K}^{-1} \text{ kg}^{-1}$). The delta function $\delta_f = 1$ if energy is available for melting (i.e. $T_l - T_f > 0$), otherwise it is $\delta_f = 0$. τ_{ice} is a parameter which represents the characteristic time scale for phase changes (Giard and Bazile, 2000). The updated temperature is first computed from Eq. A1 with $\Phi_l = 0$, then the phase change is computed as an adjustment to T_l , W_l and W_{lf} as in Boone et al. (2000).

890 A3 Energy Fluxes

It is assumed that litter below the canopy is spatially homogeneous so that it intercepts all of the incoming radiation. Thus, the net radiation R_{nl} for the litter layer is the same that for the first soil layer in the standard model:

$$R_{n,g} = SW_{net,g} + LW_{net,g} \quad (\text{A5})$$

895 where $SW_{net,g}$ and $LW_{net,g}$ correspond to the net shortwave and longwave radiation (W m^{-2}) as in Boone et al. (2017). Note that currently, the soil emissivity and albedo values are used for the litter for spatially distributed simulations pending the development of global datasets of these parameters for litter or the development of an appropriate model to estimate them. For local scale simulations, the values can be defined based on observations.

900 The below-canopy sensible heat flux, H_l (W m^{-2}), is computed the same way that for the top soil layer in the ISBA model as:

$$H_l = \rho_a \frac{(T_l - T_c)}{R_{ag-c}} \quad (\text{A6})$$

where R_{ag-c} is the aerodynamic resistance between the ground and the canopy air space which is based on Choudhury and Monteith (1988). T_l and T_c ($J\ kg^{-1}$) are thermodynamic variables which are linearly related to temperature (Boone et al., 2017). The latent heat flux is partitioned between evaporation and sublimation in the litter layer:

$$LE_l = (1 - p_{lf}) LE_l + p_{lf} LE_{lf} \quad (A7)$$

where p_{lf} is the fraction of frozen water in the litter layer and

$$LE_l = L_v \rho_a \frac{[h_{ul} q_{sat}(T_l) - q_c]}{R_{ag-c}} \quad (A8a)$$

$$910 \quad LE_{lf} = L_s \rho_a \frac{[h_{ulf} q_{sat}(T_l) - q_c]}{R_{ag-c}} \quad (A8b)$$

where the specific humidity of the canopy air space is represented by q_c . The specific humidity at saturation over liquid water is represented by q_{sat} ($kg\ kg^{-1}$). Note, it would be more accurate to use the specific humidity at saturation over ice in Eq. A8b, but this complicates the linearization and this effect is neglected for now (Boone et al., 2017). The surface humidity factors for liquid and frozen water are represented by h_{ul} and h_{ulf} , respectively. They are computed as the relative humidity following (Noilhan and Planton, 1989):

$$h_{ul} = \frac{1}{2} \left[1 - \cos \left(\pi \frac{W_l}{W_{l,max}} \right) \right] \quad (A9)$$

Note that h_{ulf} is computed by replacing W_l and $W_{l,max}$ by the values for the liquid water equivalent ice content. The maximum liquid holding capacity is modified for ice following (Boone et al., 2000).

Finally, the ground conduction flux ($W\ m^{-2}$) between the litter layer and the underlying soil is computed as:

$$G_{g,1} = \frac{T_l - T_{g,1}}{(\Delta z_l / \lambda_l) + (\Delta z_{g,1} / \lambda_{g,1})} \quad (A10)$$

where λ_l and $\lambda_{g,1}$ are the litter and first soil layer thermal conductivities respectively and $\Delta z_{g,1}$ is the thickness of the first soil layer.

A4 Water interception and fluxes

The water intercepted by the litter layer corresponds to the sum of the rain passing through the canopy and the drip from the canopy.

$$P_l = (1 - \sigma_v) P_r + D_c \quad (A11)$$

930 Note that when snow is present, melt from the snowpack is also included in this term (Boone et al., 2017). The fraction of the total rainfall P_r ($kg\ m^{-3}\ s^{-1}$) intercepted by the canopy is modulated by σ_v , which depends on the Leaf Area Index of the canopy. Finally, the canopy drip is represented by

D_c . Further details on canopy interception and drip are given in Boone et al. (2017). The drainage from the litter is simply computed as in Noilhan and Mahfouf (1996):

$$935 \quad D_l = \max(0, W_l - W_{l,max}) \quad (A12)$$

and the amount of D_l which can infiltrate into the soil is limited by Darcy's law, with any residual contributing to surface runoff. Note that for simplicity, a gravitational drainage type formulation is not used for litter, but rather a tipping bucket following Ogée and Brunet (2002).

A5 Thermal properties

940 The litter thermal conductivity, λ_l ($\text{W m}^{-1} \text{K}^{-1}$), is computed according to De Vries (1963) as:

$$\lambda_l = 0.1 + 0.03 \left(\frac{W_l}{\rho_w \Delta z_l} \right) \quad (A13)$$

The effective heat capacity of the litter, \mathcal{C}_l ($\text{J m}^{-2} \text{K}^{-1}$), is computed using

$$\mathcal{C}_l = \Delta z_l \rho_{ld} C_{ld} + W_l C_w + W_{lf} C_i \quad (A14)$$

where the specific heat capacity of liquid water is $C_w = 4.218 \times 10^3$ ($\text{J K}^{-1} \text{kg}^{-1}$). The dry density of the litter is defined as ρ_{ld} . Ogée and Brunet (2002) used a value of dry litter density of 45 kg m^{-3} for a pine forest. Meekins and McCarthy (2001) measured a litter density of 46 kg m^{-3} in a deciduous forest and Kostel-Hughes et al. (1998) estimated values varying between 27 to 38 kg m^{-3} for oak forests. Currently ECOCLIMAP doesn't distinguish between different types of deciduous trees, thus by default, ρ_{ld} is assigned a value of 45 kg m^{-3} . As a proxy for the specific heat of litter, we use the specific heat capacity of organic material from Farouki (1986) which is $C_{ld} = 1.926 \times 10^3$ $\text{J kg}^{-1} \text{K}^{-1}$. Currently, constant values for ρ_{ld} and C_{ld} are used for spatially distributed applications or on the local scale, unless observational data are available.

Appendix B: Statistical Scores

This three scores are defined respectively as:

$$955 \quad R = \frac{\overline{(x_m - \bar{x}_m)(x_o - \bar{x}_o)}}{\sigma_m \sigma_o} \quad SDV = \frac{\sigma_m}{\sigma_o} \quad cRMSE = \frac{\sqrt{[(x_m - \bar{x}_m) - (x_o - \bar{x}_o)]^2}}{\sigma_m} \quad (B1)$$

where the overbar represents the average, x_m and x_o are the modeled and observed datasets respectively and σ is the standard deviation defined as:

$$\sigma = \sqrt{\overline{(x - \bar{x})^2}} \quad (B2)$$

The cRMSE difference between MEBL and ISBA is defined as:

$$960 \quad \Delta_{cRMSE} = \frac{cRMSE(ISBA) - cRMSE(MEBL)}{\min[cRMSE(ISBA), cRMSE(MEBL)]} \times 100 \quad (B3)$$

where a positive value corresponds to an improvement using MEBL.

Measured litter thickness reported for various sites. SiteCountryMain coverThicknessMassRéféréncetypecmkg m⁻²

Le Bray FranceMaritime Pine52.6Ogée and Brunet (2002)needleconstantOak Ridge USAdeciduous oak40.6Wilson et al. (2012)leafconstantKyushu Univ.JapanC.japonica5.21.7Sato et al. (2004)needleMarch 2002Kyushu Univ.JapanL.edulis8.62.1Sato et al. (2004)leafMarch 2002NetherlandDouglas fir5Schaap and Bouten (1997)needleRussiaTaiga4.2Vorobeichik (1997)leafJuly 1990RussiaTaiga5.4Vorobeichik (1997)needleJuly 1990CSIRSouth AfricaEucalyptus grandis3.82.3Bulecock and Jewitt (2012)leavesmarch 2011CSIRSouth AfricaAcacia mearnsii1.82.4Bulecock and Jewitt (2012)nedlemarch 2011CSIRSouth AfricaPinus patula4.53.3Bulecock and Jewitt (2012)nedlemarch 2011TumbarumbaAustraliaEucalyptus3Haverd and Cuntz (2010)leavesCanadaWhite pine2.5Wu et al. (2013)needleEcuadorOcotea infrafoveolata1.8 (average)Marsh and Pearman (1997)leafMawphlangIndiaQuercus griffithii1.2 (average)Arunachalam and Arunachalam (2000)leafAoyamaJapanPinus thunbergii4.5Zhu et al. (2003)needleSeirseminenFinlandSpruce3.5Haila et al. (1999)needle

Table 1. Characteristics of the three French forest sites.

Site	Barbeau	Puéchabon	Bray
Year	2013	2006	2006
Localisation	Paris	Montpellier	Bordeaux
Main vegetation type	Sessile oak	Green oak	Maritime pine / Grass
Climate	Temperate	Mediterranean	Maritime
Forest type	Deciduous broadleaf	Evergreen broadleaf	Evergreen needleleaf
Mean <u>annual</u> temp (°C)	10.7	13.5	12.9
<u>Annual</u> Rainfall (mm)	680	872	997

Table 2. *The main model parameters for each site. Literature indicates that figures come from studies cited in the text and estimated means that values were provided by the principal investigators of each site. The main model parameters for each site. Literature indicates that values come from studies cited in the text and estimated means that values were provided by the principal investigators of each site.*

Site	Barbeau	Puechabon	Bray	Source
Soil Parameters				
Sand (%)	41	14	98	literature
Clay (%)	39	40	2	literature
W_{sat}	0.36 - 0.48	0.06	0.42 - 0.42	estimated
W_{fc}	0.15 - 0.35	0.046	0.17 - 0.16	estimated
W_{wilt}	0.05	0.018	0.04 - 0.03	estimated
SOC 0-30 cm (kg m ⁻²)	5	5	4.8	HWSD
SOC 70-100 cm (kg.m ⁻²)	4	5.5	9	HWSD
Root Depth (m)	1.2	5.0	0.8	literature
Soil albedo (vis / nir)	0.03 / 0.13	0.05 / 0.2	0.1 / 0.18	estimated
Vegetation Parameters				
LAI (m ² m ⁻²)	0.5 - 6.0	2.4	2.0-4.0	literature
Height (m)	27	5.5	18	literature
Vegetation albedo (vis / nir)	0.05 / 0.20	0.03 / 0.17	0.04 / 0.17	estimated
Vegetation fraction	0.95	0.99	0.95	ECOCLIMAP
Litter Parameters				
Thickness (m)	0.03	0.01	0.05	estimated

Table 3. *Measured litter thickness reported for various sites.*

<u>Site</u> <u>Référence</u>	<u>Country</u>	<u>Main cover</u> <u>type</u>	<u>Thickness</u> <u>cm</u>	<u>Mass</u> <u>kg m⁻²</u>
<u>Le Bray</u> <u>Ogée and Brunet (2002)</u>	<u>France</u>	<u>Maritime Pine</u> <u>needle</u>	<u>5</u> <u>constant</u>	<u>2.6</u>
<u>Oak Ridge</u> <u>Wilson et al. (2012)</u>	<u>USA</u>	<u>deciduous oak</u> <u>leaf</u>	<u>4</u> <u>constant</u>	<u>0.6</u>
<u>Kyushu Univ.</u> <u>Sato et al. (2004)</u>	<u>Japan</u>	<u>C.japonica</u> <u>needle</u>	<u>5.2</u> <u>March 2002</u>	<u>1.7</u>
<u>Kyushu Univ.</u> <u>Sato et al. (2004)</u>	<u>Japan</u>	<u>L.edulis</u> <u>leaf</u>	<u>8.6</u> <u>March 2002</u>	<u>2.1</u>
<u>Schaap and Bouten (1997)</u>	<u>Netherland</u>	<u>Douglas fir</u> <u>needle</u>	<u>5</u>	
<u>Vorobeichik (1997)</u>	<u>Russia</u>	<u>Taiga</u> <u>leaf</u>	<u>4.2</u> <u>July 1990</u>	
<u>Vorobeichik (1997)</u>	<u>Russia</u>	<u>Taiga</u> <u>needle</u>	<u>5.4</u> <u>July 1990</u>	
<u>CSIR</u> <u>Bulcock and Jewitt (2012)</u>	<u>South Africa</u>	<u>Eucalyptus grandis</u> <u>leaves</u>	<u>3.8</u> <u>march 2011</u>	<u>2.3</u>
<u>CSIR</u> <u>Bulcock and Jewitt (2012)</u>	<u>South Africa</u>	<u>Acacia mearnsii</u> <u>needle</u>	<u>1.8</u> <u>march 2011</u>	<u>2.4</u>
<u>CSIR</u> <u>Bulcock and Jewitt (2012)</u>	<u>South Africa</u>	<u>Pinus patula</u> <u>needle</u>	<u>4.5</u> <u>march 2011</u>	<u>3.3</u>
<u>Tumbarumba</u> <u>Haverd and Cuntz (2010)</u>	<u>Australia</u>	<u>Eucalyptus</u> <u>leaves</u>	<u>3</u>	
<u>Wu et al. (2013)</u>	<u>Canada</u>	<u>White pine</u> <u>needle</u>	<u>2.5</u>	
<u>Marsh and Pearman (1997)</u>	<u>Ecuador</u>	<u>Ocotea infrafoveolata</u> <u>leaf</u>	<u>1.8 (average)</u>	
<u>Mawphlang</u> <u>Arunachalam and Arunachalam (2000)</u>	<u>India</u>	<u>Quercus griffithii</u> <u>leaf</u>	<u>1.2 (average)</u>	
<u>Aoyama</u> <u>Zhu et al. (2003)</u>	<u>Japan</u>	<u>Pinus thunbergii</u> <u>needle</u>	<u>4.5</u>	
<u>Seirseminen</u> <u>Haila et al. (1999)</u>	<u>Finland</u>	<u>Spruce</u> <u>needle</u>	<u>3-5</u>	

Table 4. The root mean square error, RMSE, square correlation coefficient, R^2 , and annual error, AE, for the three French sites for the three different experiments.

Site	Flux	MEBL			MEB			ISBA		
		RMSE	R^2	AE	RMSE	R^2	AE	RMSE	R^2	AE
Bray	SW_{net}	3.0	1.0	-0.1	3.0	1.0	-0.1	3.1	1.0	0.5
	LW_{net}	7.1	0.95	2.3	6.0	0.97	2.1	7.1	0.95	-0.2
	H	52.4	0.84	5.7	52.5	0.83	4.7	63.1	0.77	-2.1
	LE	59.2	0.54	-1.2	60.8	0.5	-1.5	58.9	0.55	0.8
	G	9.6	0.75	-3.3	21.6	0.55	-3.4	42.1	0.36	-3.0
Puechabon	SW_{net}	6.4	0.97	-1.3	6.4	0.97	-1.3	6.4	0.97	-2.5
	LW_{net}	7.2	0.98	-0.9	5.8	0.99	-1.3	9.3	0.98	4.2
	H	43.7	0.95	4.1	46.4	0.94	2.7	72.3	0.87	-2.2
	LE	37.3	0.66	-5.5	38.6	0.62	-4.1	41.3	0.58	-7.3
	G	15.8	0.81	0.8	29.2	0.78	0.5	56.9	0.48	1.4
Barbeau	SW_{net}	8.7	0.96	-5.9	8.7	0.96	-5.9	9.4	0.97	-7.6
	LW_{net}	9.8	0.98	-8.8	10.0	0.99	-9.7	8.7	0.99	-7.2
	H	54.1	0.8	23.1	63.8	0.67	15.9	51.1	0.8	17.2
	LE	53.4	0.76	-15.1	59.6	0.66	-7.3	48.6	0.78	-10.2
	G	4.7	0.75	-3.2	22.0	0.55	-3.3	42.4	0.36	-2.8

Table 5. The root mean square error, RMSE, square correlation coefficient, R^2 and annual error, AE, computed with available soil temperatures of each site for the three different experiments

Site	Depth (m)	MEBL			MEB			ISBA		
		RMSE	R^2	AE	RMSE	R^2	AE	RMSE	R^2	AE
Bray	0.04	1.9	0.98	-1.0	2.9	0.96	-1.3	3.6	0.93	0.1
	0.32	2.3	0.97	-0.9	2.8	0.97	-1.2	3.0	0.96	0.2
	1.0	3.0	0.94	-1.1	3.5	0.91	-1.2	3.8	0.89	0.2
Puechabon	0.10	3.3	0.91	-0.9	3.8	0.92	-1.2	4.2	0.89	0.9
Barbeau	0.04	1.4	0.98	-0.4	2.8	0.93	-0.9	3.9	0.89	0.5
	0.08	1.4	0.98	-0.4	2.7	0.93	-0.9	3.7	0.91	0.5
	0.16	1.4	0.97	-0.5	2.5	0.94	-0.9	3.2	0.92	0.4
	0.32	1.3	0.98	-0.6	2.2	0.95	-1.0	2.7	0.94	0.3




Figure 1. The location of the three well-instrumented forested sites in France.




Figure 2. The forested sites from the Fluxnet network. Selected sites (shown) have a maximum energy imbalance at or below 20%. The circles indicate the location of sites retained for this study.

~~The forested sites from the Fluxnet network. Selected sites (shown) have a maximum energy imbalance at or below 20%. The circles indicate the location of sites retained for this study.~~




Figure 3. The modelled (thin dashed line) and observed (thick dashed line) incoming short wave radiation transmitted through the canopy at Puéchabon and Barbeau . The observations are based on below-canopy *PAR* measurements. The total incoming short wave radiation is also plotted (full black line) as a reference.




Figure 4. The partitioning of latent heat flux for each site and model option into: transpiration (black), ground/litter evaporation (white) and evaporation from the canopy (grey). Values for Le Bray (2006), Puechabon (2006) and Barbeau (2013) are shown in panels a, b and c, respectively.

The modelled (thin dashed line) and observed (thick dashed line) incoming short wave radiation transmitted through the canopy at Puechabon and Barbeau. The observations are based on below-canopy PAR measurements. The total incoming short wave radiation is also plotted (full black line) as a reference.




Figure 5. The total soil water content calculated over the root depth indicated in Table 2 (left: panels a, c and e) and the near-surface volumetric soil water content (right: panels b, d and f) at each site. Observations are in black (for Puechabon, the black curve corresponds to the output from a site-specific calibrated reference model from Lempereur et al. (2015), see text). Results for MEBL are in red, for MEB in blue and for ISBA in green.





Figure 6. The monthly diurnal cycle composite at Le Bray. MEBL is in red, MEB in blue, ISBA in green, measurements are indicated by a dashed black line and adjusted measurements are represented using a solid black curve. As a visual aid, the area between the latter two curves is shaded. Ideally, model results fall within this area.



Puechabon:

Figure 7. As in Fig 7 except for Le Bray.




Figure 8. As in Fig.7 except for Barbeau.




Figure 9. The Taylor diagram for ground heat flux, G , at Le Bray (panel a), Puechabon (b), and Barbeau (c). MEBL is represented by a circle, MEB by a square, ISBA by a triangle and measurements are indicated using a star. Three scores are represented in each diagram; the correlation coefficient (R) is the angle of the polar plot, the normalized standard deviation (NSDV) is the radial distance from the origin, and the normalized centered root mean square error (cRMSE) is proportional to the distance with the reference (star).




Figure 10. Monthly average soil temperature (K) diurnal cycle (at 0.04 m soil depth) composites at Le Bray (panel a), Puechabon (b) and Barbeau (c). MEBL is in red, MEB in blue, ISBA in green and the observations are shown in black.

The total soil water content (left: panels a, c and e) and the near-surface volumetric soil water content (right: panels b, d and f) at each site. Observations are in black (for Puechabon, the black curve corresponds to the output from a site-specific calibrated reference model from Lempereur et al. (2015), see text). Results for MEBL are in red, for MEB in blue and for ISBA in green.




Figure 11. The G RMSE computed at Le Bray and Barbeau for different values of the litter thickness.




Figure 12. Scatter plots of RMSE, R^2 and AE ~~dee~~ for sensible heat flux (H) for each site and year. The abscissa corresponds to the ISBA model and the ordinate to the MEB-L model. Points falling within the grey shaded area imply a better statistical score for MEBL. Each dot corresponds to one site. The intersection of the dashed lines corresponds to the average over all 42 sites. The percentages at the top of each panel correspond to the percent of sites which have better ~~statities~~ statistics for MEBL.




Figure 13. As in Fig.12 except for latent heat flux (~~LELE~~)




Figure 14. Boxplots of improvement in RMSE between MEBL and ISBA. Each RMSE value is computed with over a month-long period which satisfies the energy budget closure condition. The sensible heat flux is shown in panel a, and the latent heat flux is in panel b. The box plots contain 6, 13, 19, 27, 21, and 14 % of all considered months (values) for ranges in LAI increasing from 0 to 1 $\text{m}^2 \text{m}^{-2}$, to 5-8 $\text{m}^2 \text{m}^{-2}$, respectively.