

## ***Interactive comment on “Locally-orthogonal unstructured grid-generation for general circulation modelling on the sphere\*” by Darren Engwirda***

### **Anonymous Referee #1**

Received and published: 11 January 2017

I thank the author for his answers. The author clearly answered to my previous questions. I only have a few issues left.

1 Yes you are right about Voronoi-Delaunay grid staggering. I was too fast and apologize. I indeed was referring to what you call well-centered.

2 My main concern is still about the novelty of the method. You say:

In the current work, it's shown that a combination of Frontal-Delaunay refinement and hill-climbing optimisation is an effective strategy able to produce very high-quality well-centred Voronoi-Delaunay grids even when complex, highly non-uniform grid sizing constraints are imposed. I believe this to be a new result of benefit to the unstruc-

C1

tured oceanic/atmospheric modelling communities. Public availability of the associated JIGSAW-GEO grid-generator is also thought to be a further benefit to the community.

and:

I do not believe that the methods presented in the present work are preexisting. The hybrid Frontal-Delaunay surface meshing technique described here, able to guarantee worst-case bounds on element quality and sizing conformance are, in my view, new. I am not aware of another algorithm with the same properties able to produce smoothly varying Voronoi-Delaunay grids with very high mean element quality (similar to advancing front type schemes), while also guaranteeing worst-case bounds on element angles and conformance (à la standard Delaunay-refinement techniques). Existing methods for unstructured oceanic/atmospheric modelling appear to either lack provable worst-case bounds [Jacobsen et al., 2013], or generally produce grids with somewhat lower overall quality [Lambrechts et al., 2008]. The combination of the Frontal-Delaunay scheme with a coupled hill-climbing optimisation strategy to generate 'well-centred' grids is also, in my view, new.

EVERY mesh generator (edge, face, volume) has a main engine (Delaunay, Frontal, Octree, coupled) and an optimization phase that follows [1], so there is nothing new to that. The facts that you apply it to oceanic/atmospheric communities or that it is publicly available do not make these techniques new. I have added a list of references on high quality surface mesh generation that present the same high quality based on the same techniques [2,3,4,5] on top of the coupled Delaunay advancing front variants which are not fully referenced. Feel free to include them or not.

Nevertheless, I completely agree with the fact that the application of these techniques to the oceanic/atmospheric communities is new and interesting and therefore, the paper should be published.

[1] Frey,P.J. and George,P.L., Meshing, applications to finite elements, Hermes, Paris, 1999 [2] J. Tristano, S. Owen, S. Canann, Advancing front surface mesh generation

C2

in parametric space using a Riemannian surface definition, in: IMR, 1998, pp. 429–445. [3] D. Ryppl, P. Krysl, Triangulation of 3D surfaces, Eng. Comput. 13 (1997) 87–98. [4] C. Lee, Automatic metric advancing front triangulation over curved surfaces, Eng. Comput. 17 (1) (2000) 48–74. 642–667. [5] Löhner R. Regridding surface triangulations. Journal of Computational Physics 1996; 126:1–10.

---

Interactive comment on Geosci. Model Dev. Discuss., doi:10.5194/gmd-2016-296, 2016.