

**The Land Surface, Snow and Soil moisture Model Intercomparison Program (LS3MIP):
aims, set-up and expected outcome** (original title)

Reply to reviewers

Bart van den Hurk et al, 13 July 2016

We thank the reviewers, editor and representatives from the various organisational units of the CMIP panel for their constructive comments on the LS3MIP documentation paper. Although the LS3MIP protocol as described in this paper can be regarded as a reference document guiding the implementation of the experiment, many questions on technical and scientific aspects still arise as the experiment is being set up and critically evaluated. We therefore welcome the editors suggestion to provide a version number to the experimental description, as future redesigns or reconsiderations may be likely.

Reply to the editor

We have changed the title of the manuscript into “LS3MIP (v1.0) contribution to CMIP6: The Land Surface, Snow and Soil moisture Model Intercomparison Program – aims, set-up and expected outcome”

Reply to reviewer Paul Dirmeyer

- *Objectives section: An obvious "omission" is anything to do directly with vegetation or the carbon cycle, which will probably stir up questions in the minds of readers. The authors should declare the territory of this MIP up front, presaging Fig 3, that the focus is on the "GEWEXy" bits, especially the water cycle. State explicitly that there are other MIPs (e.g., LUMIP) that are concerned with the vegetation aspect of the land surface (can say "...as described later..." as this does get addressed eventually with Figs 2 and 3 on p.6).*

Good point, also commented on by other reviewers. In the “objectives” section we added a paragraph explaining the link to LUMIP: “While vegetation, carbon cycle, soil moisture, snow, surface energy balance and land-atmosphere interaction are all intimately coupled in the real world, LS3MIP focuses – necessarily – on the physical subdomain in this complex system. Interactions with vegetation and carbon cycle are included in the analyses wherever this is possible without losing this essential focus. In the complementary experiments Land Use MIP (LUMIP; see Lawrence et al. submitted) and C4MIP (Jones et al, 2016) vegetation, the terrestrial carbon cycle and land management are the central topics of analysis. LS3MIP and LUMIP share some model experiments

and analyses (see below) to allow addressing the complex interactions at the land surface and yet remain able to focus on well-posed hypotheses and research approaches.”

- *Highly relevant work on snow-climate coupling should be cited (Xu and Dirmeyer 2011) especially regarding the albedo versus delayed hydrologic effects (Xu and Dirmeyer 2013), the latter also at line 191.*

Sorry we missed these references. We’ve added a sentence in the introduction section “Temporal dynamics of the snow-atmosphere coupling during various phases of snow depletion (Xu and Dirmeyer 2011, 2012) are crucial for a proper representation of the timing and atmospheric response to snow melt.” and added a reference near line 191

- *Para lines 176-192: This is rather redundant with the first paragraph of the introduction. Probably needs to be in only one place. Same para: Is WCRP the only "customer" for this project? Seems that the potential audience is much broader - should be stated here. Along with the next paragraph, gives short shrift to the broader impacts of LS3MIP.*

Apart from removing some redundant references to WCRP, we’ve added a sentence describing the potential revenues of LS3MIP: “LS3MIP is geared to extend and consolidate available data, models and theories to support human awareness and resilience to highly variable environmental conditions in a large ensemble of sectoral domains, including disaster risk reduction, food security, public safety, nature conservation and societal wellbeing.”

- *L297: Which NCEP reanalysis? There are 3 separate unique NCEP reanalyses.*

This can be found in the documentation of the CRU-NCEP data set and is of minor relevance for this paper

- *Para lines 427-439: Really need to avoid prescribing surface soil layer moisture as well, because it can cause highly unrealistic Bowen ratios where net radiation is high, (cf. GLACE2 experiment "S"; Koster et al. 2006)*

Yes, we are aware of this, and have added a reference to this notion. As we made clear in the manuscript, a standardization of this approach is difficult, and should be carefully tested by the modelling groups.

- *L510-12: What is the protocol for ensemble construction? Are there suggestions of a prioritized list of preferred approaches like there were for GLACE-1 and -2?*

Good point. We’ve added the phrase “The procedure to initialize the land surface states in the ensemble members is left to the participant, but should allow to generate sufficient

spread that can be considered representative for the climate system under study. Koster et al. (2006) proposed a preference hierarchy of methods depending on the availability of initialization fields, and LS3MIP will follow this proposal.”

- *Tier 2 experiments in LFMIP: The mean AOGCM climatology of SST will certainly differ from that of an AMIP run based on observed SST, introducing two differences between the experiments, not one. What are the implications?*

The implications will be substantial, but also a systematic biases in SSTs is an inherent part of the analysis of the role of SSTs on land-atmosphere coupling.

- *L699-702: It seems agricultural areas in general should be a focus.*

Indeed, added as such

- Other minor corrections and suggestions for citations: changed as suggested

Reply to reviewer Gab Abramowitz

- *For this to work as a stand-alone paper, I feel like a little more contextualisation of the divisions between CMIP6 related projects might make sense. Why, for example, is there so little carbon cycle discussion (noting it's not circled in Figure 2) in an experiment that is ostensibly about all things land surface (“LS3MIP fills a major gap by considering systematic land biases and land feedbacks”)? The carbon cycle is clearly relevant for a water resources discussion when CO₂ is rapidly increasing. If there is a clear science rationale for the dividing line between another CMIP6 project (say, C4MIP) that investigates the land component of the carbon cycle it really should spelled out in detail here. C4MIP is only mentioned in passing and isn't shown on the diagram of LandMIPs (Figure 3). I would have thought it evident that the carbon cycle affects the water cycle, and that its effect is not limited to “impacts of snow and soil moisture processes . . . on terrestrial carbon exchanges” (L219-220). Alternatively, if there are historical institutional and/or political reasons for such a division I think that needs to be laid bare in a journal article describing the rationale for a science program.*

This point was also made by Paul Dirmeyer. We've stressed the complementarity of LS3MIP with LUMIP and C4MIP, as indicated in the new text (see reply to Paul Dirmeyer above).

- *As a description of what LS3MIP participants will produce and why, this document is clear in its motivation and detail, and is well thought out. What's less clear, to me at least, is how we can meaningfully evaluate the model output that this experiment will produce. I understand that analysis of CMIP data is not coordinated in the way that the production of simulations is, but nevertheless the production protocol significantly affects what can or cannot be investigated.*

Indeed, the manuscript primarily focuses on the experimental protocol, and gives examples of analyses and important research questions that can be addressed with these experiments. As such it does not describe so much the dynamics of the research network that is active in the planning, execution and analysis of LS3MIP.

I've added a paragraph on this in the "time line/participation" section: "The organisational structure of LS3MIP consistently relies on active participation of modelling groups. Coordination structures are put in place for the collection and dissemination of data and model results (Eyring et al. 2015), and for the organisation of meetings and seminars (by the core team members of LS3MIP, first five authors of this manuscript). Different from earlier experiments such as GSWP2 and GLACE1/2, no central "analysis group" is put in place that is responsible for the analyses as proposed in this manuscript. The execution and publication of analyses is considered to be a community effort of participating researchers, under coordination of the core LS3MIP team members, for instance in order to avoid duplication of efforts and coordinate the production of scientific papers."

- *One of the stated objectives of this work is to "diagnose systematic biases and processlevel deficiencies in the land modules of current Earth System Models". This requires an ability to 'ground-truth' a sufficient subset of model states and fluxes, at high temporal scales, to be able to categorically identify and quantify the fidelity of process representation. At this point in time, as I understand it, we don't come close to having this kind of observational data collection at gridded scales (despite the many products described on p17/18). While this experiment (laudably) uses multiple gridded driving data sets in Land-Hist2, this very real uncertainty, together with the significant disagreement amongst the multiple historical gridded evapotranspiration products that are available (as an example), means that we are usually unable to categorically describe the cause of differences between a model simulation and evaluation products. This problem is even tougher in the coupled environment. Essentially I don't think we can use this approach for model diagnosis, unless model problems are extreme. It is essentially a confirmation holism problem, well described in the broader climate modelling context by Lenhard and Winsberg (2010). It is clearly also problematic when we try to "quantify the associated uncertainties" with the land surface in climate projections – another stated objective. How do the authors propose we get around this issue?*

This is an interesting and well posed issue: the complexity of the true climate system will not allow a comprehensive analysis of all its relevant interactions and dynamics given the limited ability of models and observations to capture these. Personally I am not a believer of "reducing uncertainty" as a key role of climate (model) research,

but am convinced that within the limits of “understandability” valuable statements on plausibility of processes or events to occur can be derived from well designed model experiments. It goes too far to devote an extensive discussion on this issue in this manuscript, but we included a reference to Lenhard and Winsberg in the discussion section: “Within the limits to which complex models such as ESMs can be evaluated with currently available observational evidence (see e.g. the interesting philosophical discussion on climate model evaluation by Lenhard and Winsberg; 2010), it will lead to enhanced understanding of the contribution of land surface treatment to overall climate model performance...”

- *A partial antidote to the problem outlined in (2). Despite the glaring scale mismatch for model application, using a broad collection of site-based data sets to thoroughly understand the fidelity of process representation might well help regional and global scale applications. Lines 354-356 indicate that some forcing from single sites will be included in LS3MIP, but there is very little detail, presumably because the authors felt this spatial scale was not especially relevant for global scale simulations. My feeling is that if we really want to diagnose process level deficiencies and provide the means to quantify uncertainty, this really needs to be the starting point, since it's the only context in which we can meaningfully understand the uncertainties in both the forcing and evaluation data. This is not to suggest that model biases / errors at this scale necessarily translate directly to larger scales, of course. The results in Best et al (2015) and Haughton et al (2016) illustrate the power of the constraint that observational data provide at these scales. Do the authors have any reason to believe, if we had “true” gridded forcing and evaluation data at global scales, that the benchmarking results from these papers would not still be evident at gridded scales? If there is any doubt, I think a comprehensive set of site-based experiments would be very useful as part of LS3MIP, at least as its objectives currently stand. Again, I'm not sure of the extent to which the experimental protocol is already fixed, but if not, this may be a useful addition.*

Although we do agree with this notion, the exact point the reviewer wants to make is not clear. The experimental design is not particularly geared towards either local or global evaluation, but indeed analyses of larger scale interactions have a stronger emphasis than process evaluation at the local scale. However, also analysis using in situ observations must be put in the broader context in order to gain insight and inspiration for model development, and the “holistic view” described by Lenhard and Winsberg similarly applies to in situ data. We felt there is not a very clear message from this statement that we could include in the revised version of the manuscript.

- *L319 / Figure 5: are these the PLUMBER sites from Best et al 2015? If so, a simple reference gives readers enough information to get a lot more from this figure.*

Indeed, it was the PALS data set that was used here. We've indicated that in the figure caption.

- *L393-394: How is the choice to “represent the ensemble spread efficiently and reliably” going to be made? Evans et al (2013)? Global temperature trend? Could be controversial!*

We are aware of the controversy but have not yet made a decision on how this choice will be made. The reference to Evans et al is added for inspiration.

- *L501-509: this seems a little vague - are periods for extremes analysis part of LS3MIP or not? If so, which periods, why?*

At this point in time it is very difficult to be more specific: early results should give inspiration to zooming in on particular episodes.

- Other minor text suggestions and citations have been included as suggested.

Reply to review by Ron Stouffer

- *In the Introduction, this paper needs to clearly state what is its focus and what is found in the other strongly related GMD CMIP6 papers. The split between the physical climate and the carbon MIPs needs to be made much clearer and early in the paper.*

This comment is also made by Paul Dirmeyer and Gab Abramowitz. We've added a paragraph on the LS3MIP focus and links to LUMIP and C4MIP in particular (see comments above)

- *Page 11, Lines 389-402 – You may want to note here that these runs will be performed sometime in the future after the ESM data is available in the CMIP6 archive. This could be a year or 2 or more in the future.*

Pointed out in a comment

- *Page 12, line 405 – Is there an interaction between LFMIP and FAFMIP? It seems there should be and it should be noted in this section.*

We did make a reference to FAFMIP in this section, but plans for coordinated analyses have not yet been discussed with the FAFMIP panel: “This set-up is similar to the Flux Anomaly Forced MIP (FAFMIP, Gregory et al. 2016), where the role of ocean-atmosphere interaction at climate time scales is diagnosed by idealised surface perturbation experiments.”. However, a stronger link with OMIP is included in the new version, to cross-reference offline generated freshwater fluxes: “Interactions with the Ocean MIP (OMIP; Griffies et al. 2016) are arranged by the use of terrestrial freshwater fluxes produced in the LMIP simulations as a boundary condition for the forced ocean-

only simulations in OMIP, in addition to the forcing provided by (Dai and Trenberth 2002)."

- *Page 14, line 523 – "A perfect boundary condition" – several studies have shown that prescribed SSTs are less than perfect since it breaks the atmosphere-ocean coupling and feedbacks. This issue distorts the variability in models forced by SSTs relative predicted SSTs. I assume the land surface will have even larger issues since it has much smaller heat capacity. Reword.*

Also Paul Dirmeyer made a comment on the notion that prescribed SSTs are not necessarily perfect. We've rephrased it as "pseudo-observed boundary condition" experiment.

- *In the table "direction" should be changed to "Positive direction". Just to be very clear.*

See reply above

Reply to Anonymous Referee #3

We thank the reviewer for the overall positive assessment of the experimental set-up and its description, and his/her recommendation to have the paper published in GMD.

- *Data volume estimates for the requested ESM model output are currently missing and it is recommended to add the information for instance in table 1. This is easy to compute if the cost of 1-year of output (mandatory/extra) is made available. The information can be very helpful to plan storage of the output and runs throughput.*

Although this is a valuable comment, we cross-checked a few other CMIP6 papers in GMD, and none of them provide these estimates. I understand the CMIP coordination panel is preparing a paper describing the planned data exchange and storage, and I would expect that document to act as a reference for resource planning by the modelling groups.

- *Links to other projects such as the PRIMAVERA-H2020 <https://www.primaverah2020.eu> or CRESCENDO-H2020 is also worth mentioning.*

These projects are well known to us. We did make cross-references to a number of earlier experiments, but chose to confine ourselves to those experiments that have a direct relation with the LS3MIP protocol and analyses. We are aware that projects like CRESCENDO (and also others) will be used to carry out the simulations and analyses mentioned in LS3MIP (and other MIPs)

- *There is no mention to the reproducibility of the results and whether the data repository will facilitate for instance re-run the Land experiment series with another model at a later stage.*

We don't have a lot of experience with reproducibility, but generally outcomes are pretty sensitive to computer platforms, initialization, subtle configuration settings

etc that make direct reproducibility limited. Later participation to the experiment by other modelling groups is encouraged and facilitated by the infrastructure. A comment on this is added in the “Data Availability” section: “This infrastructure makes it possible to carry out the experiments in a distributed manner, and to allow later participation of additional modelling groups.”

LS3MIP (v1.0) contribution to CMIP6: The Land Surface, Snow and Soil moisture Model
Intercomparison Program-(LS3MIP) – ÷
aims, set-up and expected outcome

Met opmaak: Engels (V.S.)

Bart van den Hurk¹, Hyungjun Kim², Gerhard Krinner³, Sonia I. Seneviratne⁴, Chris Derksen⁵,
Taikan Oki², Hervé Douville⁶, Jeanne Colin⁶, Agnès Ducharne⁷, Frederique Cheruy⁷, Nicholas
Viovy⁸, Michael Puma⁹, Yoshihide Wada¹⁰, Weiping Li¹¹, Binghao Jia¹², Andrea Alessandri¹³,
Dave Lawrence¹⁴, Graham P. Weedon¹⁵, Richard Ellis¹⁶, Stefan Hagemann¹⁷, Jiafu Mao¹⁸,
Mark G Flanner¹⁹, Matteo Zampieri²⁰, Rachel Law²¹, and Justin Sheffield^{22,23}

Met opmaak: Engels (V.S.)

¹ KNMI The Netherlands (corresponding author: hurkvd@knmi.nl)

² Univ Tokyo Japan – MIROC6

³ LGGE Grenoble France – IPSL

⁴ ETH Zürich Switzerland

⁵ Environment and climate change Canada – CCCma

⁶ CNRM France – CNRM

⁷ UPMC France – IPSL

⁸ LSCE France – IPSL

⁹ NASA US – GISS

¹⁰ IIASA Laxenburg Austria

¹¹ CMA China – BCC

¹² Chinese Academy of Sciences – FGOALS

¹³ ENEA Italy – EC-Earth

¹⁴ NCAR Boulder – CESM

¹⁵ UK Metoffice

¹⁶ CEH Wallingford – hadGEM3

¹⁷ Max Planck Hamburg – MPI

¹⁸ Oak Ridge National Laboratory – ACME Land model

¹⁹ University of Michigan USA

²⁰ CMCC and University of Bologna – CMCC-CM2

²¹ CSIRO Australia – ACCESS

²² Princeton University USA

Met opmaak: Engels (V.S.)

²³ University of Southampton, UK

33
34
35
36
37
38
39

Revised Manuscript submitted to GMD

v17, 136 April-July 2016

40 Abstract

41 The Land Surface, Snow and Soil Moisture Model Intercomparison Project (LS3MIP) is
42 designed to provide a comprehensive assessment of land surface, snow, and soil moisture
43 feedbacks on climate variability and climate change, and to diagnose systematic biases in
44 the land modules of current Earth System Models (ESMs). The solid and liquid water stored
45 at the land surface has a large influence on the regional climate, its variability and
46 predictability, including effects on the energy, water and carbon cycles. Notably, snow and
47 soil moisture affect surface radiation and flux partitioning properties, moisture storage and
48 land surface memory. They both strongly affect atmospheric conditions, in particular
49 surface air temperature and precipitation, but also large-scale circulation patterns.
50 However, models show divergent responses and representations of these feedbacks as well
51 as systematic biases in the underlying processes. LS3MIP will provide the means to quantify
52 the associated uncertainties and better constrain climate change projections, which is of
53 particular interest for highly vulnerable regions (densely populated areas, agricultural
54 regions, the Arctic, semi-arid and other sensitive terrestrial ecosystems).

Met opmaak: Engels (V.S.)

55 The experiments are subdivided in two components, the first addressing systematic land
56 biases in offline mode ("LMIP", building upon the 3rd phase of Global Soil Wetness Project;
57 GSWP3) and the second addressing land feedbacks attributed to soil moisture and snow in
58 an integrated framework ("LFMIP", building upon the GLACE-CMIP blueprint).

60 Introduction

61 Land surface processes, including heat fluxes, snow, soil moisture, vegetation, turbulent
62 transfer and runoff, continue to be ranked highly on the list of the most relevant yet
63 complex and poorly represented features in state-of-the-art climate models. People live on
64 land, exploit its water and natural resources, and experience day-to-day weather that is
65 strongly affected by feedbacks with the land surface. The six Grand Challenges of the World
66 Climate Research Program (WCRP)¹ include topics governed primarily (Water Availability,
67 Cryosphere) or largely (Climate Extremes) by land surface characteristics.

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

68 Despite the importance of a credible representation of land surface processes in Earth
69 System Models (ESMs), a number of systematic biases and uncertainties persist. Biases in
70 hydrological characteristics (e.g. moisture storage in soil and snow, runoff, vegetation and
71 surface water bodies), partitioning of energy and water fluxes (Seneviratne et al. 2010),
72 definition of initial and boundary conditions at the appropriate spatial scale, feedback
73 strengths (Koster et al. 2004; Qu and Hall 2014) and inherent land surface related
74 predictability (Douville et al. 2007; Dirmeyer et al. 2013) are still subjects of considerable
75 research effort.

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

Gewijzigde veldcode

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

76 These biases and uncertainties are problematic, because they affect, among others, forecast
77 skill (Koster et al. 2010a), regional climate change patterns (Campoy et al. 2013; Seneviratne

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

¹ <http://www.wcrp-climate.org/grand-challenges>

78 et al. 2013; Koven et al. 2012), and explicable trends in water resources (Lehning 2013). In
79 addition, there is evidence of the presence of large-scale systematic biases in some aspects
80 of land hydrology in current climate models (Mueller and Seneviratne 2014) and the
81 terrestrial component of the carbon cycle (Anav et al. 2013; Mystakidis et al. 2016). Notably,
82 land surface processes can be an important reason for a direct link between the climate
83 models' temperature biases in the present period and in the future projections with
84 increased radiative forcings at the regional scale (Cattiaux et al. 2013).

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

85 For snow cover, a better understanding of the links with climate is critical for interpretation
86 of the observed dramatic reduction in springtime snow cover over recent decades (e.g.
87 (Derksen and Brown 2012; Brutel-Vuilmet et al. 2013), to improve the seasonal to
88 interannual forecast skill of temperature, runoff and soil moisture (e.g. Thomas et al. 2015;
89 Peings et al. 2011), and to adequately represent polar warming amplification in the Arctic
90 (e.g. Holland and Bitz, 2003). Snow-related biases in climate models may arise from the
91 snow-albedo feedback (Qu and Hall 2014; Thackeray et al. 2015a), but also from the energy
92 sink induced by snow melting in spring and the thermal insulation effect of snow on the
93 underlying soil (Koven et al. 2012; Gouttevin et al. 2012). Temporal dynamics of snow-
94 atmospheric coupling during various phases of snow depletion (Xu and Dirmeyer 2011,
95 2012) are crucial for a proper representation of the timing and atmospheric response to
96 snow melt. Phase 1 and 2 of the Snow Model Intercomparison Project (SnowMIP) (Etchevers
97 et al. 2004; Essery et al. 2009) provided useful insights in the capacity of snow models of
98 different complexity to simulate the snowpack evolution from local meteorological forcing
99 but did not explore snow-climate interactions. Because of strong snow/atmosphere
100 interactions, it remains difficult to distinguish and quantify the various potential causes for
101 disagreement between observed and modeled snow trends and the related climate
102 feedbacks.

Met opmaak: Lettertype: Calibri

103 Soil moisture plays a central role in the coupled land – vegetation – snow – water –
104 atmosphere system (Seneviratne et al., 2010; van den Hurk et al., 2011), where interactions
105 are evident at many relevant time scales: diurnal cycles of land surface fluxes, ~~(sub-)~~
106 seasonal and subseasonal predictability of droughts, floods, and hot extremes, annual cycles
107 governing the water buffer in dry seasons, and shifts in the climatology in response to
108 changing patterns of precipitation and evaporation. The representation of historical
109 variations in land water availability and droughts still suffer from large uncertainties, due to
110 model parameterizations, unrepresented hydrologic processes such as lateral groundwater
111 flow, lateral flows connected to reinfiltration of river water or irrigation with river water,
112 and/or atmospheric forcings (Sheffield et al. 2012; Zampieri et al. 2012); Trenberth et al.
113 2014; Greve et al. 2014; Clark et al. 2015). This also applies to the energy and carbon
114 exchanges between the land and the atmosphere (e.g. Mueller and Seneviratne 2014;
115 Friedlingstein et al. 2013).

Gewijzigde veldcode

Met opmaak: Nederlands (standaard)

Met opmaak: Nederlands (standaard)

Gewijzigde veldcode

Met opmaak: Nederlands (standaard)

Met opmaak: Nederlands (standaard)

Gewijzigde veldcode

Met opmaak: Nederlands (standaard)

Met opmaak: Nederlands (standaard)

116 It is difficult to generate reliable observations of soil moisture and land surface fluxes that
117 can be used as boundary conditions for modelling and predictability studies. Satellite
118 retrievals, in situ observations, offline model experiments (Second Global Soil Wetness
119 Project, GSWP2; Dirmeyer et al. 2006) and indirect estimates all have a potential to

120 generate relevant information but are largely inconsistent, covering different model
121 components, and suffer from methodological flaws (Mueller et al. 2013; Jiafu Mao et al.
122 2015). As a consequence, the pioneering work on deriving soil moisture related land-
123 atmosphere coupling strength (Koster et al. 2004) and regional/global climate responses in
124 both present and future climate (Seneviratne et al. 2006, 2013) has been carried out using
125 (ensembles of) modelling experiments. The second Global Land Atmosphere Coupling
126 Experiment (GLACE2; Koster et al., 2010a) measured the actual temperature and
127 precipitation skill improvement of using GSWP2 soil moisture initializations, which is much
128 lower than suggested by the coupling strength diagnostics. Limited quality of the initial
129 states, limited predictability and poor representation of essential processes determining the
130 propagation of information through the hydrological cycle in the models all play a role.

131 Altogether, there are substantial challenges concerning both the representation of land-
132 surface processes in current-generation ESMs and the understanding of related climate
133 feedbacks. The Land Surface, Snow and Soil moisture Model Intercomparison Project
134 (LS3MIP) is designed to allow the climate modelling community to make substantial
135 progress in addressing these challenges. It is part of the sixth phase of the Coupled Model
136 Intercomparison Project (CMIP6; Eyring et al. 2015). The following section further develops
137 the objectives and rationale of LS3MIP. The experimental design and analysis plan is
138 presented thereafter. The final discussion section describes the expected outcome and
139 impact of LS3MIP.

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

140

141 Objectives and rationale

142 The goal of the collection of LS3MIP experiments is to provide a comprehensive assessment
143 of land surface, snow, and soil moisture-climate feedbacks, and to diagnose systematic
144 biases and process-level deficiencies in the land modules of current ESMs. While vegetation,
145 carbon cycle, soil moisture, snow, surface energy balance and land-atmosphere interaction
146 are all intimately coupled in the real world, LS3MIP focuses – necessarily – on the physical
147 land surface in this complex system: interactions with vegetation and carbon cycle are
148 included in the analyses wherever possible without losing this essential focus. In the
149 complementary experiment Land Use MIP (LUMIP; see Lawrence et al. submitted) and
150 C4MIP (Jones et al. 2016) vegetation, the terrestrial carbon cycle and land management are
151 the central topics of analysis. LS3MIP and LUMIP share some model experiments and
152 analyses (see below) to allow to be addressed the complex interactions at the land surface
153 and yet remain able to focus on well-posed hypotheses and research approaches.

Gewijzigde veldcode

Met opmaak: Lettertype: Calibri

Met opmaak: Engels (V.S.)

Met opmaak: Lettertype: Calibri

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

154 LS3MIP will provide the means to quantify the associated uncertainties and better
155 constrain climate change projections, of particular interest for highly vulnerable regions
156 (including densely populated regions, the Arctic, agricultural areas, and some terrestrial
157 ecosystems).

158 The LS3MIP experiments collectively address the following objectives:

- evaluate the current state of land processes including surface fluxes, snow cover and soil moisture representation in CMIP DECK (Diagnostic, Evaluation and Characterization of Klima) experiments and CMIP6 historical simulations (Eyring et al. 2015), to identify the main systematic biases and their dependencies;
- estimate multi-model long-term terrestrial energy/water/carbon cycles, using the land modules of CMIP6 models under observation-constrained historical (land reanalysis) and projected future (impact assessment) climatic conditions considering land use/land cover changes;
- assess the role of snow and soil moisture feedbacks in the regional response to altered climate forcings, focusing on controls of climate extremes, water availability and high-latitude climate in historical and future scenario runs;
- assess the contribution of land surface processes to systematic Earth System model biases and the current and future predictability of regional temperature/precipitation patterns.

These objectives address each of the three CMIP6 overarching questions: 1) What are regional feedbacks and responses to climate change?; 2) What are the systematic biases in the current climate models?; and 3) What are the perspectives concerning the generation of predictions and scenarios?

LS3MIP encompasses a family of model experiments building on earlier multi-model experiments, particularly a) offline land surface experiments (GSWP2 and its successor GSWP3), b) the coordinated snow model intercomparisons SnowMIP phase 1 and 2 (Etchevers et al., 2002; Essery et al., 2009), and c) the coupled climate time-scale GLACE-type configuration (GLACE-CMIP, Seneviratne et al. 2013). Within LS3MIP the Land-only experimental suite is referred to as **LMIP (Land Model Intercomparison Project)** with the experiment ID **Land**, while the coupled suite is labelled as **LFMIP (Land Feedback MIP)**. A detailed description of the model design is given below, and a graphical display of the various components within LS3MIP is shown in Figure 13.




Figure 13: Structure of the “LandMIPs”. LS3MIP includes (1) the offline representation of land processes (LMIP) and (2) the representation of land-atmosphere feedbacks related to snow and soil moisture (LFMIP). Forcing associated with land use is assessed in LUMIP. Substantial links also exist to C4MIP (terrestrial carbon cycle). Furthermore, a land albedo testbed

experiment is planned within GeoMIP. From Seneviratne et al. (2014)




Figure 24: Relevance of LS3MIP for WCRP Core Projects and Grand Challenges²

As illustrated in Figure 24, LS3MIP is addressing multiple WCRP Grand Challenges and core projects ~~and is therefore relevant for a large fraction of the WCRP activities. It is initiated by two out of four WCRP core projects (CIC and GEWEX) and directly related to three WCRP Grand Challenges (Cryosphere in a Changing Climate, Changes in Water Availability, and Climate Extremes).~~ The LMIP experiment will provide better estimates of historical changes in snow and soil moisture at global scale, thus allowing the evaluation of changes in freshwater, agricultural drought, and streamflow extremes over continents, and a better understanding of the main drivers of these changes. The LFMIP experiments are of high relevance for the assessment of key feedbacks and systematic biases of land surfaces processes in coupled mode. (Dirmeyer et al. 2015), and are particularly focusing on two of the main feedback loops over land: the snow-albedo-temperature feedback involved in Arctic Amplification, and the soil moisture-temperature feedback leading to major changes in temperature extremes (Douville et al. 2016). In addition, LS3MIP will allow the exchange of data and knowledge across the snow and soil moisture research communities that address a common physical topic: terrestrial water in liquid and solid form. Snow and soil moisture dynamics are often interrelated (e.g. Hall et al. 2008; Xu and Dirmeyer 2012), ~~Hall et al. 2008~~ and jointly contribute to hydrological variability (e.g. Koster et al. 2010b).

LS3MIP will also provide relevant insights for other research communities ~~within WCRP~~, such as global reconstructions of land variables that are not directly observed for detection and attribution studies (Douville et al. 2013), estimates of freshwater inputs to the oceans (which are relevant for sea-level changes and regional impacts; Carmack et al. 2015), the assessment of feedbacks shown to strongly modulate regional climate variability relevant for regional climate information, as well as the investigation of land climate feedbacks on large-scale circulation patterns and cloud occurrence (Zampieri and Lionello 2011). This will thus also imply potential contributions to ~~the other WCRP Grand Challenges and core projects and to~~ programmes like the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP; Warszawski et al. 2014) and the International Detection and Attribution Group (IDAG). LS3MIP is geared to extend and consolidate available data, models and theories to support human awareness and resilience to highly variable environmental conditions in a

Met opmaak: Lettertype: Calibri

Gewijzigde veldcode

Met opmaak: Lettertype: Calibri

Met opmaak: Lettertype: +Hoofdstekst (Calibri)

² <http://wcrp-climate.org/index.php/grand-challenges>; status Dec 2015

large ensemble of sectoral domains, including disaster risk reduction, food security, public safety, nature conservation and societal wellbeing.




Figure 32: Embedding of LS3MIP within CMIP6. Adapted from Eyring et al. (2015)

Figure 32 illustrates the embedding of LS3MIP within CMIP6. LS3MIP fills a major gap by considering systematic land biases and land feedbacks. In this context, LS3MIP is part of a larger “LandMIP” series of CMIP6 experiments fully addressing biases, uncertainties, feedbacks and forcings from the land surface (Figure 13), which are complementary to similar experiments for ocean or atmospheric processes (Seneviratne et al. 2014). In particular, we note that while LS3MIP focuses on systematic biases in land surface processes (Land) and on feedbacks from the land surface processes on the climate system (LFMIP), the complementary Land Use MIP (LUMIP) experiment addresses the role of land use forcing on the climate system. The role of vegetation and carbon stores in the climate system is a point of convergence between LUMIP, C4MIP and LS3MIP, and the offline LMIP experiment will serve as land-only reference experiments for both the LS3MIP and LUMIP experiments. In addition, there will also be links to the C4MIP experiment with respect to impacts of snow and soil moisture processes (in particular droughts and floods) on terrestrial carbon exchanges and resulting feedbacks to the climate system.




Figure 3: Structure of the “LandMIPs”. LS3MIP includes (1) the offline representation of land processes (LMIP) and (2) the representation of land-atmosphere feedbacks related to snow and soil moisture (LFMIP). Forcing associated with land use is assessed in LUMIP. Substantial links also exist to C4MIP (terrestrial carbon cycle). Furthermore, a land albedo testbed experiment is planned within GeoMIP. From Seneviratne et al. (2014)

Experimental design

The experimental design of LS3MIP consists of a series of offline land-only experiments (LMIP) driven by a land surface forcing data set and a variety of coupled model simulations (LFMIP) (see Figure 4 and Table 1):




Figure 4: Schematic diagram for the experiment structure of LS3MIP. Tier 1 experiments are indicated with a heavy black outline, and complementary ensemble experiments are indicated with white hatched lines. Land-Altfor represents 3 alternative forcings for the Land-Hist experiment. For further details on the experiments and acronyms, see Table 1 and text.

(1) Offline land model experiments (“Land offline MIP”, experiment ID “Land”):

Offline simulations of land surface states and fluxes allow for the evaluation of trends and variability of snow, soil moisture and land surface fluxes, carbon stocks and vegetation dynamics, and climate change impacts. Within the CMIP6 program various Model Intercomparison Projects make use of offline terrestrial simulations to benchmark or force coupled climate model simulations: LUMIP focusing on the role of land use/land cover change, C4MIP to address the terrestrial component of the carbon cycle and its feedback to climate, and LS3MIP to provide soil moisture and snow boundary conditions.

Meteorological forcings, ancillary data (e.g., land use/cover changes, surface parameters, CO₂ concentration and nitrogen deposition) and documented protocols to spin-up and execute the experiments are essential ingredients for a successful offline land model experiment (Wei et al. 2014). The first Global Soil Wetness Project (GSWP; Dirmeyer et al. 1999), covering two annual cycles (1987 – 1988), established a successful template, which was updated and fine-tuned in a number of follow-up experiments, both with global (Dirmeyer et al. 2006; Sheffield et al. 2006) and regional (Boone et al. 2009) coverage.

Available data sets for meteorological forcing

Offline experiments will primarily use GSWP3³ (Tier 1) forcing (Kim et al., in preparation) with alternate forcing used in Tier 2 experiments.

The third Global Soil Wetness Project (GSWP3) provides meteorological forcings for the entire 20th century and beyond, making extensive use of the 20th Century Reanalysis (20CR) (Compo et al. 2011). In this reanalysis product only surface pressure and monthly sea-surface temperature and sea-ice concentration are assimilated. The ensemble uncertainty in

³ <http://hydro.iis.u-tokyo.ac.jp/GSWP3/>

the synoptic variability of 20CR varies with the time-changing observation network. High correlations for geopotential height (500 hPa) and air temperature (850 hPa) with an independent long record (1905-2006) of upper-air data were found (Compo et al. 2011), comparable to forecast skill of a state-of-the-art forecasting system at 3 days lead time.

GSWP3 forcing data are generated based on a dynamical downscaling of 20CR. A simulation of the Global Spectral Model (GSM), run at a T248 resolution (~50km) is nudged to the vertical structures of 20CR zonal and meridional winds and air temperature using a spectral nudging dynamical downscaling technique that effectively retains synoptic features in the higher spatial resolution (Yoshimura and Kanamitsu 2008). Additional bias corrections using observations, vertical damping (Hong and Chang 2012) and single ensemble member correction (Yoshimura and Kanamitsu 2013) are applied, giving considerable improvements.

Weedon et al. (2011) provide the meteorological forcing data for the EU Water and Global Change (WATCH) programme⁴, designed to evaluate global hydrological trends and impacts using offline modelling. The half-degree resolution, 3 hourly WATCH Forcing Data (WFD) was based on the ECMWF ERA-40 reanalysis and included elevation correction and monthly bias correction using CRU observations (and alternative GPCC precipitation total observations). WATCH hydrological modelling led to the WaterMIP study (Haddeland et al. 2011). The WFD stops in 2001, but within a follow-up project EMBRACE Weedon et al. (2014) generated the WFDEI dataset that starts in 1979 and was recently extended to 2014. The WFDEI was based on the WATCH Forcing Data methodology but used the ERA-Interim reanalysis (4D-var and higher spatial resolution than ERA-40) so that there are offsets for some variable in the overlap period with the WFD. The forcing consists of 3-hourly ECMWF ERA-Interim reanalysis data (WFD used ERA-40) interpolated to half degree spatial resolution. The 2m temperatures are bias-corrected in terms of monthly means and monthly average diurnal temperature range using CRU half degree observations. The 2m temperature, surface pressure, specific humidity and downwards long-wave radiation fluxes are sequentially elevation corrected. Short-wave radiation fluxes are corrected using CRU cloud cover observations and corrected for the effects of seasonal and interannual changes in aerosol loading. Rainfall and snowfall rates are corrected using CRU wet days per month and according to CRU or GPCC observed monthly precipitation gauge totals. The WFDEI data set is also used as forcing to the ISIMIP2.1 project, which focuses on historical validation of global water balance under transient land use change (Warszawski et al. 2014).

To support the Global Carbon Project⁵ (Le Quere et al. 2009) with annual updates of global carbon pools and fluxes, the offline modelling framework TRENDY⁶ applies an ensemble of terrestrial carbon allocation and land surface models. For this a forcing data set is prepared in which NCEP reanalysis data are bias corrected using the gridded in situ climate data from the Climate Research Unit (CRU), the so-called CRU-NCEP dataset⁷. This dataset is currently

⁴ <http://www.eu-watch.org/>

⁵ <http://www.globalcarbonproject.org/about/index.htm>


⁶ <http://dgvn.ceh.ac.uk/node/21>

⁷ Viovy N, Ciais P (2009) A combined dataset for ecosystem modelling, Available at:

325 available from 1901 to 2014 at 0.5 degrees horizontal spatial resolution and 6 hourly time-
 326 step. It is being updated annually.

327 The Princeton Global Forcing dataset⁸ (Sheffield et al. 2006) was developed as a forcing for
 328 land surface and other terrestrial models, and for analyzing changes in near surface climate.
 329 The dataset is based on 6-hourly surface climate from the NCEP-NCAR reanalysis, which is
 330 corrected for biases at diurnal, daily and monthly time scales using a variety of
 331 observational datasets. The data are available at 1.0, 0.5 and 0.25-degree resolution and 3-
 332 hourly time-step. The latest version (V2.2) covers 1901-2014, with a real-time extension
 333 based on satellite precipitation and weather model analysis fields. The reanalysis
 334 precipitation is corrected by adjusting the number of rain days and monthly accumulations
 335 to match observations from CRU and the Global Precipitation Climatology Project (GPCP).
 336 Precipitation is downscaled in space using statistical relationships based on GPCP and the
 337 TRMM Multi-satellite Precipitation Analysis (TMPA), and to 3-hourly resolution based on
 338 TMPA. Temperature, humidity, pressure and longwave radiation are downscaled in space
 339 with account for elevation. Daily mean temperature and diurnal temperature range are
 340 adjusted to match the CRU monthly data. Short- and long-wave surface radiation are
 341 adjusted to match satellite-based observations from the University of Maryland (Zhang et al.
 342 submitted) and to be consistent with CRU cloud cover observations outside of the satellite
 343 period. An experimental version (V3) assimilates station observations into the background
 344 gridded field to provide local-scale corrections (Sheffield et al., in preparation).

345 Figure 5 shows the performance in terms of correlation and standard deviation of the
 346 forcing data sets compared to daily observations from 20 globally distributed in-situ
 347 FLUXNET sites (Baldocchi et al. 2001). Although for precipitation intrinsic heterogeneity
 348 leads to significant differences with the in-situ observations, long- and short-wave
 349 downward radiation (not shown) and air temperature show variability characteristics similar
 350 to the observations.



351
 352 *Figure 5: Taylor diagram for evaluating the forcing datasets comparing to daily observations*
 353 *from FLUXNET sites, as used by Best et al. (2015): (a) 2m air temperature and (b)*
 354 *precipitation. Red, blue, and green dots indicate GSWP3, Watch Forcing Data (Weedon et al.*
 355 *2011) and Princeton forcing (Sheffield et al. 2006), respectively. Grey and orange dots*
 356 *indicate 20CR and its dynamically downscaled product (GSM248).*

357
 358 The participating modelling groups are invited to run a number of experiments in this land-
 359 only branch of LS3MIP.

Opmerking [BvdH1]: Question from Gab: are these the PLUMBER sites from Best et al 2015? If so, a simple reference gives readers enough information to get a lot more from this figure.

Gewijzigde veldcode

Met opmaak: Lettertype: Cursief

Met opmaak: Lettertype: Cursief

<http://dods.extra.cea.fr/data/p529viov/cruncep/readme.htm>

⁸ <http://hydrology.princeton.edu/data.php>

360

361 *Historical offline simulations: Land-Hist*

362 The Tier 1 experiments of the offline LMIP experiment consist of simulations using the
363 GSWP3 forcing data for a historical (1831-2014) interval. The land model configuration
364 should be identical to that used in the DECK and CMIP6 historical simulations for the parent
365 coupled model.

366 The atmospheric forcing will be prepared at a standard $0.5 \times 0.5^\circ$ spatial resolution at 3
367 hourly intervals and distributed with a package to regrid data to the native grids of the
368 GCMs. Also vegetation, soil, topography and land/sea mask data will be prescribed
369 following the protocol used for the CMIP6 DECK simulations. Spin-up of the land-only
370 simulations should follow the TRENDY protocol⁹ which calls for recycling of the climate
371 mean and variability from two decades of the forcing dataset (e.g., 1831-1850 for GSWP3,
372 1901-1920 for the alternative land surface forcings). Land use should be held constant at
373 1850 as in the DECK 1850 coupled control simulation (*piControl*). See discussion and
374 definition of “constant land-use” in Section 2.1 of LUMIP protocol paper (Lawrence et al.
375 submitted). CO₂ and all other forcings should be held constant at 1850 levels during spinup.
376 For the period 1850 to the first year of the forcing dataset, the forcing data should continue
377 to be recycled but all other forcings (land-use, CO₂, etc.) should be as in the CMIP6
378 historical simulation. Transient land use is a prescribed CMIP6 forcing and is described in
379 the LUMIP protocol (Lawrence et al. submitted).

380 Interactions with the Ocean MIP (OMIP; Griffies et al. 2016) are arranged by the use of
381 terrestrial freshwater fluxes produced in the LMIP simulations as a boundary condition for
382 the forced ocean-only simulations in OMIP, in addition to the forcing provided by (Dai and
383 Trenberth 2002).

Met opmaak: Lettertype: Calibri

Met opmaak: Lettertype: Calibri

384 Single site time series of in-situ observational forcing variables from selected reference
385 locations (from FLUXNET, Baldocchi et al. 2001) are supplied in addition to the forcing data
386 for additional site level validation.

387 Although Land-Hist is not a formal component of the DECK simulations which form the core
388 of CMIP6 (see Figure 3.2), the WCRP Working Group on Climate Modelling (WGCM)
389 recognized the importance of these land-only experiments for the process of model
390 development and benchmarking. A future implementation of a full or subset of this
391 historical run is proposed to become part of the DECK in future CMIP exercises and is
392 included as a Tier 1 experiment in LS3MIP. Land surface model output from this subset of
393 LMIP will also be used as boundary condition in some of the coupled climate model
394 simulations, described below.

395

396 *Historical simulations with alternative forcings*

397 Additional Tier 2 experiments are solicited where the experimental set-up is similar to the
398 Tier 1 simulations, but using 3 alternative meteorological forcing data sets that differ from

⁹ <http://dgvn.ceh.ac.uk/node/9>

GSWP3: the Princeton forcing (Sheffield et al. 2006), WFD and WFDEI combined (allowing for offsets as needed (Weedon et al. 2014) and the CRU-NCEP forcing (Wei et al. 2014) used in TRENDY (Sitch et al. 2015). These Tier 2 experiments cover the period 1901 – 2014. The model outputs will allow assessment of the sensitivity of land-only simulations to uncertainties in forcing data. Differences in the outputs compared to the primary runs with the GSWP3 forcing will help in understanding simulation sensitivity to the selection of forcing datasets. Kim (2010) utilized a similarity index (Ω ; Koster et al. 2000) to estimate the uncertainty derived from an ensemble of precipitation observation data sets relative to the the uncertainty from an ensemble of model simulations for evapotranspiration and runoff. The joint utilization of common monthly observations by the various forcing data sets leads to a high value of Ω when evaluated using monthly mean values. However, evaluation of dataset consistency of monthly variance leads to much larger disparities and considerably lower values of Ω (Figure 6). This uncertainty will propagate differently to other hydrological variables, such as runoff or evapotranspiration (Kim 2010).




Figure 6: Global distributions of the similarity index (Ω) for 2001-2010 of monthly mean (a, c) and (b, d) monthly variance (calculated from daily data from each data set) of 2m air temperature (top panels) and precipitation (bottom panels), respectively. Shown are global distributions and zonal means. After (Kim 2010).

Climate change impact assessment: Land-Future

A set of future land-only time slice simulations (2015-2100) will be generated via forcing data obtained from at least 2 future climate scenarios from the ScenarioMIP- (O'Neill et al. 2016) and will be executed at a later stage during CMIP6. Tentatively, Shared Socio-economic Pathway SSP5-8.5 and SSP4-3.7¹⁰ will be selected, run by 3 model realizations each. The models will be chosen based on the evaluation of the results from the Historical simulations from the CMIP6 Nucleus in order to represent the ensemble spread efficiently and reliably (Evans et al. 2013). To generate a set of ensemble forcing data for the future, a trend preserving statistical bias correction method will be applied to the 3-hourly surface meteorology variables (Table A4) from the scenario output (Hempel et al. 2013; Watanabe et al. 2014). Gridded forcings will be provided in a similar data format as the historical simulations.

Land-Future is a Tier 2 experiment in LS3MIP and focuses on assessment of climate change impact (e.g. shifts of the occurrence of critical water availability due to changing statistical distributions of extreme events) and on the assessment of the land surface analogue of climate sensitivity for various key land variables (Perket et al. 2014; Flanner et al. 2011).

¹⁰ <https://cmip.ucar.edu/scenario-mip/experimental-protocols>

436

437 (2) Prescribed land surface states in coupled models for land surface feedback assessment
438 (“Land Feedback MIP”, LFMIP):

439 Land surface processes do not act in isolation in the climate system. A tight coupling with
440 the overlying atmosphere takes place on multiple temporal and spatial scales. A systematic
441 assessment of the strength and spatial structure of land surface interaction at
442 subcontinental, seasonal time scales has been performed with the initial GLACE set-up
443 (GLACE1 and GLACE2 experiments; Koster et al. (2004)) in which essentially the spread in an
444 ensemble simulation of a coupled land-atmosphere model was compared to a model
445 configuration in which the land-atmosphere interaction was greatly bypassed by prescribing
446 soil conditions throughout the simulation in all members of the ensemble. Examination of
447 the significance of land-atmosphere feedbacks at the centennial climate time scale was later
448 explored at the regional scale in a single-model study (Seneviratne et al. 2006) and on global
449 scale in the GLACE-CMIP5 experiment in a small model ensemble (Seneviratne et al. 2013).

450 A protocol very similar to the design of GLACE-CMIP5 is followed in LFMIP. Parallel to a set
451 of reference simulations taken from the CMIP6 DECK, a set of forced experiments is carried
452 out where land surface states are prescribed from or nudged towards prescribed fields
453 derived from coupled simulations. The land surface states are prescribed or nudged at a
454 daily time scale. This set-up is similar to the Flux Anomaly Forced MIP (FAFMIP, Gregory et
455 al. 2016), where the role of ocean-atmosphere interaction at climate time scales is
456 diagnosed by idealised surface perturbation experiments.

Met opmaak: Lettertype: Calibri

457 While earlier experiments used model configurations with prescribed SST and sea ice
458 conditions, the Tier 1 experiment in LFMIP will be based on coupled AOGCM simulations
459 and comprise simulations for a historical (1980-2014) and future (2015-2100) time range.
460 The selection of the future scenario (from the ScenarioMIP experiment) will be based on the
461 choices made in the offline LMIP experiment (see above).

462 In GLACE-CMIP5 only soil moisture states were prescribed in the forced experiments. The
463 configuration of the particular land surface models may introduce the need to make
464 different selections of land surface states to be prescribed, for instance to avoid strong
465 inconsistencies in the case of frozen ground (soil moisture rather than soil water state
466 should be prescribed; *M. Hauser, ETH Zurich, personal communication*), melting snow, or
467 growing vegetation. Prescribing surface soil moisture only (experiment “S” in Koster et al.
468 2006) gave unrealistic values of the surface Bowen ratio. A standardization of this selection
469 is difficult as the implementation and consequences may be highly model specific. Here we
470 recommend to prescribe only the water reservoirs (soil moisture, snow mass). The disparity
471 of possible implementations is adding to the uncertainty range generated by the model
472 ensemble, similar to the degree to which implementation of land use, flux corrections or
473 downscaling adds to this uncertainty range. Participating modelling groups are encouraged
474 to apply various test simulations focusing both on technical feasibility and experimental
475 impact to evaluate different procedures to prescribe land surface conditions.

Met opmaak: Lettertype: Calibri

476 The earlier experience with GLACE-type experiments has revealed a number of technical
477 and scientific issues. Because in most GCMs the land surface module is an integral part of

the code describing the atmosphere, prescribing land surface dynamics requires a non-conventional technical interface, reading and replacing variables throughout the entire simulations. Many ~~participants to~~ LS3MIP participants have participated earlier in GLACE-type experiments, but for some the code adjustments will require a technical effort. Interpretation of the effect of the variety of implementations of prescribed land surface variables by the different modelling groups (see above) is helped by a careful documentation of the way the modelling groups have implemented this interface. Tight coordination and frequent exchange among the participating modelling groups on the technical modalities of the implementation of the required forcing methods will be ensured during the preparatory phase of LS3MIP in order to maximize the coherence of the modelling exercise and to facilitate the interpretation of the results.

By design, the prescribed land surface experiments do not fully conserve water and energy, similar to AMIP, nudged, and data assimilation experiments. A systematic addition or removal of water or energy can even emerge as a result of asymmetric land surface responses to dry and to wet conditions, e.g. when surface evaporation or runoff depend strongly non-linearly to soil moisture or snow states (e.g. Jaeger and Seneviratne 2011). Also, unrepresented processes (such as water extraction for irrigation or exchange with the groundwater) may lead to imbalances in the budget (Wada et al. 2012). This systematic alteration of the water and energy balance may not only perturb the simulation of present-day climate (e.g. Douville 2003; Douville et al. 2016) but may also interact with the projected climate change signal, where altered climatological soil conditions can contribute to the climate change induced temperature or precipitation signal or water imbalances can lead to imposed runoff changes that could affect ocean circulation and SSTs. Earlier GLACE-type experiments revealed that the problems of water conversion are often reduced when prescribed soil water conditions are taken as the median rather than the mean of a sample over which a climatological mean is calculated (Hauser et al. *subm*). In the analyses of the experiments this asymmetry and lack of energy/water balance closure will be examined and put in context of the climatological energy and water balance and its climatic trends.

To be able to best quantify the forcing that prescribing the land surface state represents, the increments of both snow and soil moisture imposed as a consequence of this prescription are required as an additional output. This will enable us to estimate the amplitude of implicit water and energy fluxes imposed by the forcing procedure.

Complementary experiments following an almost identical setup as LFMIP, but limiting the prescription of land surface variables to snow-related variables and thus leaving soil moisture free-running, are carried out in the framework of the ESM-SnowMIP (Earth System Model - Snow Module Intercomparison Project) carried out within the WCRP Grand Challenge “Melting Ice and Global Consequences”¹¹. ESM-SnowMIP being tightly linked to LS3MIP, these complementary experiments will allow separating effects of soil moisture and snow feedbacks.

¹¹ <http://www.climate-cryosphere.org/activities/targeted/esm-snowmip>

518 *Tier 1 experiments in LFMIP*

519 Similar to the set-up of GLACE-CMIP5 (Seneviratne et al. 2013), the core experiments of
520 LFMIP (tier 1) evaluate two different sets of prescribed land surface conditions (snow and
521 soil moisture):

- 522 • LFMIP-pdLC: the experiments comprise transient coupled atmosphere-ocean
523 simulations in which a selection of land surface characteristics is prescribed rather
524 than interactively calculated in the model. This “climatological” land surface forcing
525 is calculated as the mean annual cycle in the period 1980-2014 from the Historical
526 GCM simulations. The experiment aims at diagnosing the role of land-atmosphere
527 feedback at the climate time scale. Seneviratne et al. (2013) found a substantial
528 effect of changes in climatological soil moisture on projected temperature change in
529 a future climate, both for seasonal mean and daytime extreme temperature in
530 summer. Effects on precipitation are less clear, and the multi-model nature of
531 LS3MIP is designed to sharpen these quantitative effects. Also, LS3MIP will take a
532 potential damping (or amplifying) effect of oceanic responses on altered land surface
533 conditions into account, in contrast to GLACE-CMIP5. Experiments using this set-up
534 (i.e. coupled ocean) in a single-model study have shown that the results could be
535 slightly affected by the inclusion of an interactive ocean, although the effects were
536 not found to be large overall (Orth and Seneviratne submitted).
- 537 • LFMIP-rmLC: a prescribed climatology using a transient 30-yr running mean, where a
538 comparison to the standard CMIP6 runs allows diagnosis of shifts in the regions of
539 strong land-atmosphere coupling as recorded by e.g. Seneviratne et al. (2006), and
540 shifts in potential predictability related to land surface states (Dirmeyer et al. 2013).

541 Both sets of simulations cover the historical period (1850-2014) and extend to 2100, based
542 on a forcing scenario to be identified at a later stage. The procedure to initialize the land
543 surface states in the ensemble members is left to the participant, but should allow to
544 generate sufficient spread that can be considered representative for the climate system
545 under study. Koster et al. (2006) proposed a preference hierarchy of methods depending on
546 the availability of initialization fields, and LS3MIP will follow this proposal.

Met opmaak: Lettertype: Calibri

Met opmaak: Lettertype: Calibri

547 Output in high temporal resolution (daily, as well as sub-daily for some fields and time
548 slices) is required to address the role of land surface-climate feedbacks on climate extremes
549 over land.

550 Multi-member experiments are encouraged, but the mandatory tier 1 simulations are
551 limited to one realization for each of the two prescribed land surface time series described
552 above.

553

554 *Tier 2 experiments in LFMIP*

555 To analyse a number of additional features of land –atmosphere feedbacks, a collection of
556 tier 2 simulations is proposed in LS3MIP:

- 557 • *Simulations with observed SST*: The AOGCM simulations from Tier 1 are duplicated
558 with a prescribed SST configuration taken from the AMIP runs in the DECK (AGCM),

in order to isolate the role of the ocean in propagating and damping/reinforcing land surface responses on climate (Koster et al. 2000). Both the historic and running mean land surface simulations are requested (LFMIP-pdLC+SST and -rmLC+SST, respectively)

- *Simulations with observed SST and Land-hist output:* A “~~perfect~~ pseudo-observed boundary condition” set of experiments use the AMIP SSTs and the Land-Hist land boundary conditions generated by the land surface model used in the participating ESM, leading to simulations driven by surface fields that are strongly controlled by observed forcings. This will only cover the historic period (1901-2014) (LFMIP-PObs+SST). For this the land-only simulations in LMIP need to be interpolated to the native GCM grid, preserving land-sea boundaries and other characteristics.
- *Separate effects of soil moisture and snow, and role of additional land parameters and variables:* Additional experiments, in which only snow, snow albedo or soil moisture is prescribed will be conducted to assess the respective feedbacks in isolation, and have control on possible interactions between snow cover and soil moisture content. Also vegetation parameters and variables (e.g. leaf area index, canopy height and thickness) are considered. These experiments are not listed in Table 1, but will be detailed in a follow-up protocol to be defined later.
- *Fixed land use conditions:* in conjunction with the Land Use MIP (LUMIP) a repetition of the Tier 1 experiment under fixed 1850 land cover and land use conditions highlights the role of soil moisture in modulating the climate response to land cover and land use. (Not listed in Table 1).

(3) Prescribed land surface states derived from pseudo-observations (LFMIP-Pobs)

The use of LMIP (land-only simulations) to initialize the AOGCM experiments (LFMIP) allows a set of predictability experiments in line with the GLACE2 set-up (Koster et al. 2010a). The LFMIP-Pobs experiment is an extension to GLACE2 by (a) allowing more models to participate, (b) improving the statistics by extending the original 1986 – 1995 record to 1980 – 2014, (c) evaluating the quality of newly available land surface forcings, and (d) executing the experiments in AOGCM mode. Koster et al. (2010a) and van den Hurk et al. (2012) concluded that the forecast skill improvement from models using initial soil moisture conditions was relatively low. Possible causes for this low skill are the limited record length and limited quality of the (precipitation) observations used to generate the soil conditions. These issues are explicitly addressed in LFMIP-Pobs.

All LFMIP-Pobs experiments are Tier 2, which also gives room for additional model design elements such as the evaluation of various observational data sources (such as for SWE or snow albedo, using satellites derived, reanalysis and land surface model outputs). The predictability assessments include the evaluation of the contribution of snow cover melting and its related feedbacks to the underestimation of recent boreal polar warming by climate models.

The experimental protocol (number of simulations years, ensemble size, initialization, model configuration, output diagnostics) has a strong impact on the results of the experiment (e.g.

601 Guo and Dirmeyer 2013). This careful design of the LFMIP-Pobs experiment needed for a
602 succesfull implementation has currently not yet taken place. Therefore these experiments
603 are listed as Tier 2 in Table 1, with the comment that the detailed experimental protocol still
604 needs to be defined.
605

606 *Table 1: Summary of LS3MIP experiments. Experiments with specific treatment of subsets of*
607 *land surface features are not listed in this overview.*

Experiment ID and Tier	Experiment Description / Design	Config (L/A/O)*	Start	End	# Ens**	# Total Years***	Science Question and/or Gap Being Addressed	Synergies with other CMIP6 MIPs
Land-Hist (1)	Land only simulations	L	1850	2014	1	165	Historical land simulations	LUMIP, C4MIP, CMIP6 historical
Land-Hist-cruNcep Land-Hist-princeton Land-Hist-wfdei2 (2)	Land only simulations	L	1901	2014	3	342	As Land-Hist but with three different forcing data sets (Princeton forcing, CRU-NCEP, and WFDEI)	
Land-Future (2)	Land only simulations	L	2015	2100	6	516	Climate trend analysis	LUMIP, C4MIP, ScenarioMIP
LFMIP-pdLC (1)	Prescribed land conditions 1980-2014 climate	LAO	1980	2100	1	121	diagnose land-climate feedback including ocean response	ScenarioMIP
LFMIP-pdLC2 (2)	as LFMIP-pdLC with multiple model members	LAO	1980	2100	4	484	diagnose land-climate feedback including ocean response	ScenarioMIP
LFMIP-pdLC+SST (2)	Prescribed land conditions 1980-2014 climate; SSTs prescribed	LA	1980	2100	5	605	diagnose land-climate feedback over land	ScenarioMIP
LFMIP-Pobs+SST (2)	Land conditions from Land-hist; SSTs prescribed	LA	1901	2014	1	115	"perfect boundary condition" simulations	
LFMIP-rmLC (1)	Prescribed land conditions 30yr running mean	LAO	1980	2100	1	121	diagnose land-climate feedback including ocean response	ScenarioMIP
LFMIP-rmLC2 (2)	as LFMIP-rmLC with multiple model members	LAO	1980	2100	4	484	diagnose land-climate feedback including ocean response	ScenarioMIP
LFMIP-rmLC+SST (2)	Prescribed land conditions 30yr running mean; SSTs prescribed	LA	1980	2100	5	605	diagnose land-climate feedback over land	ScenarioMIP
LFMIP-Pobs (2) ^{ptbd}	Initialized pseudo-observations land	LAO	1980	2014	10	350	land-related seasonal predictability	CMIP6 historical

608 * Config L/A/O refers to land/atmosphere/ocean model configurations

609 ** # Ens refers to number of ensemble members.

610 *** # Total years is total number of simulation years.

611 ^{ptbd} experimental protocol needs to be detailed in a later stage

612

613 **Analysis strategy**

614 LS3MIP is designed to push the land surface component of climate models, observational
615 data sets and projections to a higher level of maturity. Understanding the propagation of
616 model and forecast errors and the design of model parameterizations is essential to realize
617 this goal. The LS3MIP steering group is a multi-disciplinary team (climate modellers, snow
618 and soil moisture model specialists, experts in local and remotely sensed data of soil
619 moisture and snow properties) that ensures that the experiment setups, model evaluations
620 and analyses/interpretations of the results are pertinent.

621 For both snow and soil moisture the starting point will be a careful analysis of model results
622 from on the one hand a) the DECK historic simulations (both the AMIP and the historical
623 coupled simulation) and b) on the other hand the (offline) LMIP historical simulations.

624 For the evaluation of snow representation in the models, large-scale high-quality datasets of
625 snow mass (SWE) and snow cover extent (SCE) with quantitative uncertainty characteristics
626 will be provided by the Satellite Snow Product Intercomparison and Evaluation Experiment
627 (SnowPEX¹²). Analysis within SnowPEX is providing the first evaluation of satellite derived
628 snow extent (15 participating datasets) and SWE derived from satellite measurements, land
629 surface assimilation systems, physical snow models, and reanalyses (7 participating
630 datasets). Internal consistency between products, and bias relative to independent
631 reference datasets are being derived based on standardized and consistent protocols. The
632 evaluation of variability and trends in terrestrial snow cover extent and mass was examined
633 previously for CMIP3 and CMIP5 by e.g. Brown and Mote (2009), Derksen and Brown (2012)
634 and Brutel-Vuilmet et al. (2013). While these assessments were based on single
635 observational datasets, and hence provide no perspective on observational uncertainty and
636 spread relative to multi-model ensembles, standardized multi-source datasets generated by
637 SnowPEX will allow assessment using a multi-dataset observational ensemble (e.g. Mudryk
638 et al. 2015). For snow albedo, multiple satellite-derived datasets are available, including 16-
639 day MODIS¹³ data from 2001 – present, the ESA GlobAlbedo product¹⁴, the recently updated
640 twice-daily APP-x¹⁵ product (1982 – 2011), and a derivation of the snow shortwave radiative
641 effect from 2001 – 2013 (Singh et al. 2015). Satellite retrievals of snow cover fraction in
642 forested and mountainous areas is an ongoing area of uncertainty which influences the
643 essential diagnostics related to climate sensitivity of snow cover (Thackeray et al. 2015b),

¹² <http://calvalportal.ceos.org/projects/snowpex>

¹³ <http://modis-atmos.gsfc.nasa.gov/ALBEDO/>

¹⁴ <http://www.globalbedo.org>

¹⁵ <http://stratus.ssec.wisc.edu/products/appx/appx.html>

644 feeding into essential diagnostics related to climate sensitivity of snow cover (Qu and Hall,
645 2014; Fletcher et al. 2012).

646 In the case of soil moisture, land hydrology and vegetation state, several observations-based
647 datasets will be used in the evaluation of the coupled DECK simulations and offline Land
648 experiments. Data considered will include the first multidecadal satellite-based global soil
649 moisture record (Essential Climate Variable Soil Moisture ECVSM) (Liu et al. 2012; Dorigo et
650 al. 2012), long-term (2002-2015) records of terrestrial water storage from the GRACE
651 satellite (Rodell et al. 2009; Reager et al. 2016; Kim et al. 2009), the multi-product LandFlux-
652 EVAL evapotranspiration synthesis (Mueller et al. 2013), multi-decadal satellite retrievals of
653 the Fraction of Photosynthetically Absorbed Radiation (FPAR, e.g. Gobron et al. 2010;
654 Zscheischler et al. 2015), and upscaled Fluxnet based products (Jung et al. 2010).

655 Several details of snow and soil moisture dynamical processes can be indirectly inferred
656 through the analysis of river discharge (Orth et al. 2013; Zampieri et al. 2015). Variables
657 simulated by the routing schemes included in the land surface models can be compared
658 with the station data available from the Global Runoff Database (GRDC¹⁶). Combined use of
659 in-situ discharge observations and terrestrial water storage changes observed by GRACE will
660 verify how the land surface simulations partition the terms in the water balance equation
661 (i.e., precipitation, evapotranspiration, runoff, and water storage changes)(Kim et al. 2009).

662 The coupled LS3MIP (LFMIP) simulations will be analyzed in concert with the control runs to
663 quantify various climatic effects of snow and soil moisture, detect systematic biases and
664 diagnose feedbacks. Anticipated analyses include:

- 665 • *Drivers of variability at multiple time scales:* comparison of simulations with
666 prescribed soil moisture and snow (LFMIP-pdLC) allows ~~the~~ quantification of the
667 impact of land surface state variability on variability of climate variables ~~such as, for~~
668 ~~instance,~~ temperature, relative humidity, cloudiness, precipitation and river
669 discharge at several time scales. The LFMIP-rmLC simulation allows evaluation of this
670 contribution on seasonal time scales, and changes of patterns of high/low land
671 surface impact in a future climate. In particular, a focus ~~will~~ be ~~set-put~~ on impacts
672 on climate extremes (temperature extremes, heavy precipitation events, see e.g.
673 Seneviratne et al. 2013) and the possible role of land-based feedbacks in amplifying
674 regional climate responses compared to changes in global mean temperature
675 (Seneviratne et al. 2016). A secondary focus will be on the impacts of snow and soil
676 moisture variability on the extremes of river discharge, which can be related to large-
677 scale floods and to non-local propagation of droughts signals. These aspects will be
678 analyzed in the context of water management and to quantify feedbacks of ~~the~~ river
679 discharge ~~to~~ the climate system (through the discharge in the oceans, Materia et
680 al. 2012; Carmack et al. 2015) and to the carbon cycle (through the methane
681 produced in flooded areas, Meng et al. 2015)).

¹⁶ <http://www.bafg.de/GRDC>

- *Attribution of model disagreement:* the multi-model set up of the experiment allows closer inspection of the effects of modeled soil moisture and snow (and related processes such as plant transpiration, photosynthesis, or snowmelt) ~~on~~^{to} calculated land temperature, precipitation, runoff, vegetation state, and gross primary production. The comparison of LFMIP-pdLC and LFMIP-rmLC will be useful to isolate ~~the~~ model disagreement in land surface feedbacks potentially induced by including coupling to a dynamic ocean despite similar land response to climate change.
- *Emergent constraints:* while the annual cycle of snow cover and local temperature (Qu and Hall 2014), and the relation between global mean temperature fluctuations and CO₂-concentration (Cox et al. 2013) provide observational constraints on snow-albedo and carbon-climate feedback respectively, similar emergent constraints may be defined to constrain (regional) soil moisture or snow related feedbacks with temperature or hydrological processes such as, for instance, the timing of spring onset which may be related to snowmelt, spring river discharge (Zampieri et al. 2015) and vegetation phenology (Xu et al. 2013). Use of appropriate observations and diagnostics as emergent constraints will reduce uncertainties in projections of mean climate and extremes (heat extremes, droughts, floods) (Hoffman et al. 2014). The analysis of amplitude and timing of seasonality of hydrological and ecosystem processes will provide additional diagnostics.
- *Attribution of model bias:* a positive relationship between model temperature bias in the current climate, and (regional) climate response can partly be attributed to the soil moisture-climate feedback, which acts on both the seasonal and climate time scale (Cheruy et al. 2014). A multi-model assessment of this relationship is enabled via LS3MIP. The comparison of AMIP-DECK, LFMIP-CA and LFMIP-LCA will be used to assess the impact of atmospheric-related errors in land boundary conditions on the AGCM biases.
- *Changes in feedback hotspots and predictability patterns:* land surface conditions don't exert uniform influence on the atmosphere in all areas of the globe: a distribution of strong interaction "hotspots" and areas of high potential predictability contributions from the land surface exists (e.g. Koster et al. 2004). These patterns may change in a future climate (e.g. Seneviratne et al. 2006). A multi-model assessment such as foreseen in LS3MIP allows mapping changes in these patterns, with implications for the occurrence of droughts, heat waves, irrigation limitations or river discharge anomalies and their predictability (Dirmeyer et al. 2013).
- *Snow shortwave radiative effect analysis:* The Snow Shortwave Radiative Effect (SSRE) can be diagnosed through parallel calculations of surface albedo and shortwave fluxes with and without model snow on the ground or in the vegetation canopy (Perket et al. 2014). This metric provides a precise, overarching measure of the snow-induced perturbation to solar absorption in each model, integrating over the variable influences of vegetation masking, snow grain size, snow cover fraction,

soot content, etc. SSRE is analogous to the widely-used cloud radiative effect diagnostic, and its time evolution provides a measure of snow albedo feedback in the context of changing climate (Flanner et al. 2011). We recommend that the diagnostic snow shortwave radiative effect (SSRE) calculation be implemented in standard LS3MIP simulations (Tiers 1 and 2). This will enable us to evaluate the integrated effect of model snow cover on surface radiative fluxes.

- *Complementary snow-related offline experiments:* Additional offline experiments with prescribed snow albedo or snow water equivalent are planned as a follow-up to LS3MIP within the ESM-SnowMIP¹⁷ initiative. This is aimed at improving our understanding of sources of coupled model biases (global offline and site scale experiments) in order to identify priority avenues for future model development.

Regarding the snow analyses, the initial geographical focus of LS3MIP is on the continental snow cover of both hemispheres, both in ice-free areas (Northern Eurasia and North America) and on the large ice sheets (Greenland and Antarctica). Effects of snow on sea ice, and the quality of the representation of snow on sea ice in climate models, will be explored later, but is of interest because of strong recent trends of Arctic sea ice decline and the potential amplifying effect of earlier spring snow melt over land.

For soil moisture, the geographical focus is on all land areas, with special interest in [agricultural](#) locations with strong land-atmosphere interaction (transition zones between wet and dry climates), extensive irrigation areas, and high interannual variability of warm season climate in densely populated areas.

The analyses are carried out on a standardized model output data set. A summary of the requested output data is given in tables in the Annex.

Table 2: Earth System Modelling groups participating ~~to~~ in LS3MIP

Model Name	Institute	Country
ACCESS	CSIRO/Bureau of Meteorology	Australia
ACME Land Model	U.S. Department of Energy	USA
BCC-CSM2-MR	BCC,CMA	China
CanESM	CCCma	Canada
CESM		USA
CMCC-CM2	Centro Euro-Mediterraneo sui Cambiamenti Climatici	Italy
CNRM-CM	CNRM-CERFACS	France
EC-Earth	SMHI and 26 other institutes	Sweden and 9 other European

¹⁷ <http://www.climate-cryosphere.org/activities/targeted/esm-snowmip>

		countries
FGOALS	LASG, IAP, CAS	China
GISS	NASA GISS	USA
IPSL-CM6	IPSL	France
MIROC6-CGCM	AORI, University of Tokyo / JAMSTEC / National Institute for Environmental Studies	Japan
MPI-ESM	Max Planck Institute for Meteorology (MPI- M)	Germany
MRI-ESM1.x	Meteorological Research Institute	Japan
NorESM	Norwegian Climate Service Centre	Norway
hadGEM3	Met Office	UK

747

748 Data availability

749 The offline forcing data for the Land-Hist experiments and output from the model
 750 simulations described in this paper will be distributed through the Earth System Grid
 751 Federation (ESGF) with digital object identifiers (DOIs) assigned. The model output required
 752 for LS3MIP is listed in the Annex. Model data distributed via ESGF will be freely accessible
 753 through data portals after registration. This infrastructure makes it possible to carry out the
 754 experiments in a distributed matter, and to allow later participation of additional modelling
 755 groups. Links to all forcings datasets will be made available via the CMIP Panel website¹⁸.
 756 Information about accreditation, data infrastructure, metadata structure, citation and
 757 acknowledging is provided by Eyring et al. (2015).

758

759 Time line ~~and~~ participating models and interaction strategy

Met opmaak: Engels (V.S.)

760 The offline land surface experiments (Land-Hist) are expected to be completed in early
 761 2017. Future time slices can only be performed when the Scenario-MIP results become
 762 available. All coupled LS3MIP simulations and their subsequent analyses will be timed after
 763 the completion of the DECK and historical 20th century simulations, expected by mid 2017.
 764 Table 2 lists the participating Earth System modelling groups.

765 The organisational structure of LS3MIP relies on active participation of modelling groups.
 766 Coordination structures are in place for the collection and dissemination of data and model
 767 results (Eyring et al. 2015), and for the organisation of meetings and seminars (by the core
 768 team members of LS3MIP, first six authors of this manuscript). Different from earlier
 769 experiments such as GSWP2 and GLACE1/2, no central “analysis group” is put in place that is
 770 responsible for the analyses as proposed in this manuscript. The execution and publication
 771 of analyses is considered to be a community effort of participating researchers, in order to
 772 avoid duplication of efforts and coordinate the production of scientific papers.

Met opmaak: Engels (V.S.)

Met opmaak: Lettertype: Calibri

Met opmaak: Engels (V.S.)

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

773

¹⁸ <http://www.wcrp-climate.org/index.php/wgcm-cmip/about-cmip>

774 **Discussion: expected outcome and impact of LS3MIP**

775 The treatment of the land surface in the current generation of climate models plays a critical
776 role in the assessment of potential effects of widespread changes in radiative forcing, land
777 use and biogeochemical cycles. The land surface both “receives” climatic variations (by its
778 atmospheric forcing) and “returns” these variations as feedbacks or land surface features
779 that are of high relevance to the people living on it. The strong coupling between land
780 surface, atmosphere, hydrosphere and cryosphere makes an analysis of its performance
781 characteristics challenging: the response and the state of the land surface strongly depend
782 on the climatological context, and metrics of interactions or feedbacks, which are all difficult
783 to define and observe (van den Hurk et al. 2011).

784 LS3MIP addresses these challenges by enhancing earlier diagnostic studies and experimental
785 designs. Within the limits to which complex models such as ESMs can be evaluated with
786 currently available observational evidence (see e.g. the interesting philosophical discussion
787 on climate model evaluation by Lenhard and Winsberg; 2010). It will lead to enhanced
788 understanding of the contribution of land surface treatment to overall climate model
789 performance; give inspiration on how to optimize land surface parameterizations or ~~their~~
790 forcing; support the development of better forecasting tools, where initial conditions affect
791 the trajectory of the forecast and can be used to optimize forecast skill; and, last but not
792 least, provide a better historical picture of the evolution of our vital water resources during
793 the recent century. In particular, LS3MIP will provide a solid benchmark for assessing water
794 and climate related risks and trends therein. Given the critical importance of changes in land
795 water availability and of impacts of changes in snow, soil moisture and land surface states
796 for the projected evolution of climate mean and extremes, we expect that LS3MIP will help
797 the research community make fundamental advances in this area.

798

799 **Acknowledgements**

800 The authors thank the CMIP panel of the WCRP Working Group on Climate Modelling for
801 their efforts in coordinating the CMIP6 enterprise. G.P.W. was supported by the Joint UK
802 DECC/Defra Met Office Hadley Climate Centre Programme (GA01101). J.M. is supported by
803 the Biogeochemistry-Climate Feedbacks Scientific Focus Area project funded through the
804 Regional and Global Climate Modeling Program in Climate and Environmental Sciences
805 Division (CESD) of the Biological and Environmental Research (BER) Program in the U.S.
806 Department of Energy (DOE) Office of Science. Oak Ridge National Laboratory is managed by
807 UT-BATTELLE for DOE under contract DE-AC05-00OR22725. Hanna Lee (NorESM) has
808 expressed intention to participate in LS3MIP when feasible, but has not contributed to this
809 manuscript.

810

811 **References**

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

Met opmaak: Lettertype: Calibri

Met opmaak: Lettertype: Calibri

Gewijzigde veldcode

Met opmaak: Engels (V.S.)

812 Anav, A., and Coauthors, 2013: Evaluating the Land and Ocean Components of the Global
813 Carbon Cycle in the CMIP5 Earth System Models. *J. Clim.*, **26**, 6801–6843,
814 doi:10.1175/JCLI-D-12-00417.1.

815 Baldocchi, D., and Coauthors, 2001: FLUXNET: A New Tool to Study the Temporal and Spatial
816 Variability of Ecosystem–Scale Carbon Dioxide, Water Vapor, and Energy Flux
817 Densities. *Bull. Am. Meteorol. Soc.*, **82**, 2415–2434, doi:10.1175/1520-
818 0477(2001)082<2415:FANTTS>2.3.CO;2.

819 Best, M. J., and Coauthors, 2015: The Plumbing of Land Surface Models: Benchmarking
820 Model Performance. *J. Hydrometeorol.*, **16**, 1425–1442, doi:10.1175/JHM-D-14-
821 0158.1.

822 Boone, A., and Coauthors, 2009: The AMMA Land Surface Model Intercomparison Project
823 (ALMIP). *Bull. Am. Meteorol. Soc.*, **90**, 1865–1880, doi:10.1175/2009BAMS2786.1.

824 Brown, R. D., and P. W. Mote, 2009: The Response of Northern Hemisphere Snow Cover to a
825 Changing Climate*. *J. Clim.*, **22**, 2124–2145, doi:10.1175/2008JCLI2665.1.

826 Brutel-Vuilmet, C., M. Ménégoz, and G. Krinner, 2013: An analysis of present and future
827 seasonal Northern Hemisphere land snow cover simulated by CMIP5 coupled climate
828 models. *The Cryosphere*, **7**, 67–80, doi:10.5194/tc-7-67-2013.

829 Campoy, A., A. Ducharne, F. Cheruy, F. Hourdin, J. Polcher, and J. C. Dupont, 2013: Response
830 of land surface fluxes and precipitation to different soil bottom hydrological
831 conditions in a general circulation model. *J. Geophys. Res. Atmospheres*, **118**,
832 10,725–10,739, doi:10.1002/jgrd.50627.

833 Carmack, E., and Coauthors, 2015: Fresh water and its role in the Arctic Marine System:
834 sources, disposition, storage, export, and physical and biogeochemical consequences
835 in the Arctic and global oceans. *J. Geophys. Res. Biogeosciences*, n/a – n/a,
836 doi:10.1002/2015JG003140.

837 Cattiaux, J., H. Douville, and Y. Peings, 2013: European temperatures in CMIP5: origins of
838 present-day biases and future uncertainties. *Clim. Dyn.*, **41**, 2889–2907,
839 doi:10.1007/s00382-013-1731-y.

840 Cheruy, F., J. L. Dufresne, F. Hourdin, and A. Ducharne, 2014: Role of clouds and land-
841 atmosphere coupling in midlatitude continental summer warm biases and climate
842 change amplification in CMIP5 simulations. *Geophys. Res. Lett.*, **41**, 6493–6500,
843 doi:10.1002/2014GL061145.

844 Clark, M. P., and Coauthors, 2015: Improving the representation of hydrologic processes in
845 Earth System Models. *Water Resour. Res.*, **51**, 5929–5956,
846 doi:10.1002/2015WR017096.

847 Compo, G. P., and Coauthors, 2011: The Twentieth Century Reanalysis Project. *Q. J. R.*
848 *Meteorol. Soc.*, **137**, 1–28, doi:10.1002/qj.776.

Met opmaak: Bibliografie, Zwevende regels voorkomen, Spatiëring tussen Aziatische en Latijnse tekst aanpassen, Spatiëring tussen Aziatische tekst en nummers aanpassen

Gewijzigde veldcode

849 [Cox, P. M., D. Pearson, B. B. Booth, P. Friedlingstein, C. Huntingford, C. D. Jones, and C. M.](#)
850 [Luke, 2013: Sensitivity of tropical carbon to climate change constrained by carbon](#)
851 [dioxide variability. *Nature*, **494**, 341–344, doi:10.1038/nature11882.](#)

852 [Dai, A., and K. E. Trenberth, 2002: Estimates of Freshwater Discharge from Continents:](#)
853 [Latitudinal and Seasonal Variations. *J. Hydrometeorol.*, **3**, 660–687,](#)
854 [doi:10.1175/1525-7541\(2002\)003<0660:EOFDfC>2.0.CO;2.](#)

855 [Derksen, C., and R. Brown, 2012: Spring snow cover extent reductions in the 2008–2012](#)
856 [period exceeding climate model projections. *Geophys. Res. Lett.*, **39**, L19504,](#)
857 [doi:10.1029/2012GL053387.](#)

858 [Dirmeyer, P. A., A. J. Dolman, and N. Sato, 1999: The Pilot Phase of the Global Soil Wetness](#)
859 [Project. *Bull. Am. Meteorol. Soc.*, **80**, 851–878, doi:10.1175/1520-](#)
860 [0477\(1999\)080<0851:TPPOTG>2.0.CO;2.](#)

861 [—, X. Gao, M. Zhao, Z. Guo, T. Oki, and N. Hanasaki, 2006: GSWP-2: Multimodel Analysis](#)
862 [and Implications for Our Perception of the Land Surface. *Bull. Am. Meteorol. Soc.*, **87**,](#)
863 [1381–1397, doi:10.1175/BAMS-87-10-1381.](#)

864 [—, S. Kumar, M. J. Fennessy, E. L. Altshuler, T. DelSole, Z. Guo, B. A. Cash, and D. Straus,](#)
865 [2013: Model Estimates of Land-Driven Predictability in a Changing Climate from](#)
866 [CCSM4. *J. Clim.*, **26**, 8495–8512, doi:10.1175/JCLI-D-13-00029.1.](#)

867 [Dirmeyer, P. A., C. Peters-Lidard, and G. Balsamo, 2015: Land-Atmosphere Interactions and](#)
868 [the Water Cycle. *Seamless prediction of the Earth system: from minutes to months*,](#)
869 [G. Brunet, S. Jones, and P.M. Ruti, Eds., Vol. No 1156 of, WMO, Geneva, p. Ch 15.](#)

870 [Dorigo, W., R. de Jeu, D. Chung, R. Parinussa, Y. Liu, W. Wagner, and D. Fernández-Prieto,](#)
871 [2012: Evaluating global trends \(1988–2010\) in harmonized multi-satellite surface soil](#)
872 [moisture. *Geophys. Res. Lett.*, **39**, n/a – n/a, doi:10.1029/2012GL052988.](#)

873 [Douville, Conil, Tyteca, and Voldoire, 2007: Soil moisture memory and West African](#)
874 [monsoon predictability: artefact or reality? *Clim. Dyn.*, **28**, 723–742,](#)
875 [doi:10.1007/s00382-006-0207-8.](#)

876 [Douville, H., 2003: Assessing the Influence of Soil Moisture on Seasonal Climate Variability](#)
877 [with AGCMs. *J. Hydrometeorol.*, **4**, 1044–1066, doi:10.1175/1525-](#)
878 [7541\(2003\)004<1044:ATIOSM>2.0.CO;2.](#)

879 [—, A. Ribes, B. Decharme, R. Alkama, and J. Sheffield, 2013: Anthropogenic influence on](#)
880 [multidecadal changes in reconstructed global evapotranspiration. *Nat. Clim Change*,](#)
881 [3, 59–62, doi:10.1038/nclimate1632.](#)

882 [—, J. Colin, E. Krug, J. Cattiaux, and S. Thao, 2016: Midlatitude daily summer temperatures](#)
883 [reshaped by soil moisture under climate change. *Geophys. Res. Lett.*, **43**, 812–818,](#)
884 [doi:10.1002/2015GL066222.](#)

885 [Essery, R., and Coauthors, 2009: SNOWMIP2: An Evaluation of Forest Snow Process](#)
886 [Simulations. *Bull. Am. Meteorol. Soc.*, **90**, 1120–1135,](#)
887 [doi:10.1175/2009BAMS2629.1.](#)

888 [Etchevers, P., and Coauthors, 2004: Validation of the energy budget of an alpine snowpack](#)
889 [simulated by several snow models \(SnowMIP project\). *Ann. Glaciol.*, **38**, 150–158,](#)
890 [doi:10.3189/172756404781814825.](#)

891 [Evans, J., F. Ji, G. Abramowitz, and M. Ekström, 2013: Optimally choosing small ensemble](#)
892 [members to produce robust climate simulations. *Environ. Res. Lett.*, **8**, 044050.](#)

893 [Eyring, V., S. Bony, G. A. Meehl, C. Senior, B. Stevens, R. J. Stouffer, and K. E. Taylor, 2015:](#)
894 [Overview of the Coupled Model Intercomparison Project Phase 6 \(CMIP6\)](#)
895 [experimental design and organisation. *Geosci Model Dev Discuss*, **2015**, 10539–](#)
896 [10583, doi:10.5194/gmdd-8-10539-2015.](#)

897 [Flanner, M. G., K. M. Shell, M. Barlage, D. K. Perovich, and M. A. Tschudi, 2011: Radiative](#)
898 [forcing and albedo feedback from the Northern Hemisphere cryosphere between](#)
899 [1979 and 2008. *Nat. Geosci*, **4**, 151–155, doi:10.1038/ngeo1062.](#)

900 [Fletcher, C. G., S. C. Hardiman, P. J. Kushner, and J. Cohen, 2009: The Dynamical Response to](#)
901 [Snow Cover Perturbations in a Large Ensemble of Atmospheric GCM Integrations. *J.*](#)
902 [Clim., **22**, 1208–1222, doi:10.1175/2008JCLI2505.1.](#)

903 [—, H. Zhao, P. J. Kushner, and R. Fernandes, 2012: Using models and satellite observations](#)
904 [to evaluate the strength of snow albedo feedback. *J. Geophys. Res. Atmospheres*,](#)
905 [117, D11117, doi:10.1029/2012JD017724.](#)

906 [Friedlingstein, P., M. Meinshausen, V. K. Arora, C. D. Jones, A. Anav, S. K. Liddicoat, and R.](#)
907 [Knutti, 2013: Uncertainties in CMIP5 Climate Projections due to Carbon Cycle](#)
908 [Feedbacks. *J. Clim.*, **27**, 511–526, doi:10.1175/JCLI-D-12-00579.1.](#)

909 [Gobron, N., A. Belward, B. Pinty, and W. Knorr, 2010: Monitoring biosphere vegetation](#)
910 [1998–2009. *Geophys. Res. Lett.*, **37**, n/a – n/a, doi:10.1029/2010GL043870.](#)

911 [Gouttevin, I., M. Menegoz, F. Dominé, G. Krinner, C. Koven, P. Ciais, C. Tarnocai, and J.](#)
912 [Boike, 2012: How the insulating properties of snow affect soil carbon distribution in](#)
913 [the continental pan-Arctic area. *J. Geophys. Res. Biogeosciences*, **117**, n/a – n/a,](#)
914 [doi:10.1029/2011JG001916.](#)

915 [Gregory, J. M., and Coauthors, 2016: The Flux-Anomaly-Forced Model Intercomparison](#)
916 [Project \(FAFMIP\) contribution to CMIP6: Investigation of sea-level and ocean climate](#)
917 [change in response to CO2 forcing. *Geosci Model Dev Discuss*, **2016**, 1–37,](#)
918 [doi:10.5194/gmd-2016-122.](#)

919 [Greve, P., B. Orlowsky, B. Mueller, J. Sheffield, M. Reichstein, and S. I. Seneviratne, 2014:](#)
920 [Global assessment of trends in wetting and drying over land. *Nat. Geosci*, **7**, 716–](#)
921 [721.](#)

922 [Griffies, S. M., and Coauthors, 2016: Experimental and diagnostic protocol for the physical](#)
923 [component of the CMIP6 Ocean Model Intercomparison Project \(OMIP\). *Geosci*](#)
924 [Model Dev Discuss, **2016**, 1–108, doi:10.5194/gmd-2016-77.](#)

925 [Guo, Z., and P. A. Dirmeyer, 2013: Interannual Variability of Land–Atmosphere Coupling](#)
926 [Strength. *J. Hydrometeorol.*, **14**, 1636–1646, doi:10.1175/JHM-D-12-0171.1.](#)

927 [Haddeland, I., and Coauthors, 2011: Multimodel Estimate of the Global Terrestrial Water](#)
928 [Balance: Setup and First Results. *J. Hydrometeorol.*, **12**, 869–884,](#)
929 [doi:10.1175/2011JHM1324.1.](#)

930 [Hall, A., X. Qu, and J. D. Neelin, 2008: Improving predictions of summer climate change in](#)
931 [the United States. *Geophys. Res. Lett.*, **35**, L01702, doi:10.1029/2007GL032012.](#)

932 [Hauser, M., R. Orth, and S. I. Seneviratne, subm: Role of soil moisture vs recent climate](#)
933 [change for heat waves in western Russia. *Geophys.Res.Lett.*,](#)

934 [Hempel, S., K. Frieler, L. Warszawski, J. Schewe, and F. Piontek, 2013: A trend-preserving](#)
935 [bias correction – the ISI-MIP approach. *Earth Syst Dynam.*, **4**, 219–236,](#)
936 [doi:10.5194/esd-4-219-2013.](#)

937 [Hoffman, F. M., and Coauthors, 2014: Causes and implications of persistent atmospheric](#)
938 [carbon dioxide biases in Earth System Models. *J. Geophys. Res. Biogeosciences*, **119**,](#)
939 [141–162, doi:10.1002/2013JG002381.](#)

940 [Holland, M. M., and C. M. Bitz, 2003: Polar amplification of climate change in coupled](#)
941 [models. *Clim. Dyn.*, **21**, 221–232, doi:10.1007/s00382-003-0332-6.](#)

942 [Hong, S.-Y., and E.-C. Chang, 2012: Spectral nudging sensitivity experiments in a regional](#)
943 [climate model. *Asia-Pac. J. Atmospheric Sci.*, **48**, 345–355, doi:10.1007/s13143-012-](#)
944 [0033-3.](#)

945 [van den Hurk, B., M. Best, P. Dirmeyer, A. Pitman, J. Polcher, and J. Santanello, 2011:](#)
946 [Acceleration of Land Surface Model Development over a Decade of Glass. *Bull. Am.*](#)
947 [Meteorol. Soc., **92**, 1593–1600, doi:10.1175/BAMS-D-11-00007.1.](#)

948 [—, F. Doblas-Reyes, G. Balsamo, R. Koster, S. Seneviratne, and H. Camargo Jr, 2012: Soil](#)
949 [moisture effects on seasonal temperature and precipitation forecast scores in](#)
950 [Europe. *Clim. Dyn.*, **38**, 349–362, doi:10.1007/s00382-010-0956-2.](#)

951 [Jaeger, E. B., and S. I. Seneviratne, 2011: Impact of soil moisture–atmosphere coupling on](#)
952 [European climate extremes and trends in a regional climate model. *Clim. Dyn.*, **36**,](#)
953 [1919–1939, doi:10.1007/s00382-010-0780-8.](#)

954 [Jiafu Mao, and Coauthors, 2015: Disentangling climatic and anthropogenic controls on](#)
955 [global terrestrial evapotranspiration trends. *Environ. Res. Lett.*, **10**, 094008.](#)

956 [Jones, C. D., and Coauthors, 2016: The C4MIP experimental protocol for CMIP6. *Geosci*](#)
957 [Model Dev Discuss, **2016**, 1–52, doi:10.5194/gmd-2016-36.](#)

958 [Jung, M., and Coauthors, 2010: Recent decline in the global land evapotranspiration trend](#)
959 [due to limited moisture supply. *Nature*, **467**, 951–954, doi:10.1038/nature09396.](#)

960 [Kim, H., 2010: *Role of rivers in the spatiotemporal variations of terrestrial hydrological*](#)
961 [circulations. University of Tokyo.](#)

962 [Kim, H., P. J.-F. Yeh, T. Oki, and S. Kanae, 2009: Role of rivers in the seasonal variations of](#)
963 [terrestrial water storage over global basins. *Geophys. Res. Lett.*, **36**, n/a – n/a,](#)
964 [doi:10.1029/2009GL039006.](#)

965 [Koster, R. D., M. J. Suarez, and M. Heiser, 2000: Variance and Predictability of Precipitation](#)
966 [at Seasonal-to-Interannual Timescales. *J. Hydrometeorol.*, **1**, 26–46,](#)
967 [doi:10.1175/1525-7541\(2000\)001<0026:VAPOPA>2.0.CO;2.](#)

968 [—, and Coauthors, 2004: Regions of Strong Coupling Between Soil Moisture and](#)
969 [Precipitation. *Science*, **305**, 1138–1140, doi:10.1126/science.1100217.](#)

970 [—, and Coauthors, 2006: GLACE: The Global Land–Atmosphere Coupling Experiment. Part](#)
971 [I: Overview. *J. Hydrometeorol.*, **7**, 590–610, doi:10.1175/JHM510.1.](#)

972 [Koster, R. D., and Coauthors, 2010a: Contribution of land surface initialization to](#)
973 [subseasonal forecast skill: First results from a multi-model experiment. *Geophys.*](#)
974 [Res. Lett., **37**, L02402, doi:10.1029/2009GL041677.](#)

975 [Koster, R. D., S. P. P. Mahanama, B. Livneh, D. P. Lettenmaier, and R. H. Reichle, 2010b: Skill](#)
976 [in streamflow forecasts derived from large-scale estimates of soil moisture and](#)
977 [snow. *Nat. Geosci.*, **3**, 613–616, doi:10.1038/ngeo944.](#)

978 [Koven, C. D., W. J. Riley, and A. Stern, 2012: Analysis of Permafrost Thermal Dynamics and](#)
979 [Response to Climate Change in the CMIP5 Earth System Models. *J. Clim.*, **26**, 1877–](#)
980 [1900, doi:10.1175/JCLI-D-12-00228.1.](#)

981 [Lawrence, D. M., and Coauthors, submitted: The Land-Use Model Intercomparison Project](#)
982 [\(LUMIP\) experimental design. *Geosci Model Dev.*](#)

983 [Lehning, M., 2013: Snow–atmosphere interactions and hydrological consequences. *Snow–*](#)
984 [Atmosphere Interact. Hydrol. Consequences, **55**, 1–3,](#)
985 [doi:10.1016/j.advwatres.2013.02.001.](#)

986 [Lenhard, J., and E. Winsberg, 2010: Holism, entrenchment, and the future of climate model](#)
987 [pluralism. *Stud. Hist. Philos. Sci. Part B*, **41**, 253–262.](#)

988 [Le Quere, C., M. R. Raupach, J. G. Canadell, and G. Marland et al., 2009: Trends in the](#)
989 [sources and sinks of carbon dioxide. *Nat. Geosci.*, **2**, 831–836, doi:10.1038/ngeo689.](#)

990 [Liu, Y. Y., W. A. Dorigo, R. M. Parinussa, R. A. M. de Jeu, W. Wagner, M. F. McCabe, J. P.](#)
991 [Evans, and A. I. J. M. van Dijk, 2012: Trend-preserving blending of passive and active](#)
992 [microwave soil moisture retrievals. *Remote Sens. Environ.*, **123**, 280–297,](#)
993 [doi:10.1016/j.rse.2012.03.014.](#)

994 [Materia, S., S. Gualdi, A. Navarra, and L. Terray, 2012: The effect of Congo River freshwater](#)
995 [discharge on Eastern Equatorial Atlantic climate variability. *Clim. Dyn.*, **39**, 2109–](#)
996 [2125, doi:10.1007/s00382-012-1514-x.](#)

997 [Meng, L., R. Paudel, P. G. M. Hess, and N. M. Mahowald, 2015: Seasonal and interannual](#)
998 [variability in wetland methane emissions simulated by CLM4Me' and CAM-chem and](#)
999 [comparisons to observations of concentrations. *Biogeosciences*, **12**, 4029–4049,](#)
1000 [doi:10.5194/bg-12-4029-2015.](#)

1001 [Mudryk, L. R., C. Derksen, P. J. Kushner, and R. Brown, 2015: Characterization of Northern](#)
1002 [Hemisphere Snow Water Equivalent Datasets, 1981–2010. *J. Clim.*, **28**, 8037–8051,](#)
1003 [doi:10.1175/JCLI-D-15-0229.1.](#)

1004 [Mueller, B., and S. I. Seneviratne, 2014: Systematic land climate and evapotranspiration](#)
1005 [biases in CMIP5 simulations. *Geophys. Res. Lett.*, **41**, 128–134,](#)
1006 [doi:10.1002/2013GL058055.](#)

1007 [—, and Coauthors, 2013: Benchmark products for land evapotranspiration: LandFlux-](#)
1008 [EVAL multi-data set synthesis. *Hydrol Earth Syst Sci*, **17**, 3707–3720,](#)
1009 [doi:10.5194/hess-17-3707-2013.](#)

1010 [Mystakidis, S., E. L. Davin, N. Gruber, and S. I. Seneviratne, 2016: Constraining future](#)
1011 [terrestrial carbon cycle projections using observation-based water and carbon flux](#)
1012 [estimates. *Glob. Change Biol.*, n/a – n/a, doi:10.1111/gcb.13217.](#)

1013 [O'Neill, B. C., and Coauthors, 2016: The Scenario Model Intercomparison Project](#)
1014 [\(ScenarioMIP\) for CMIP6. *Geosci Model Dev Discuss*, **2016**, 1–35, doi:10.5194/gmd-](#)
1015 [2016-84.](#)

1016 [Orth, R., and S. I. Seneviratne, submitted: Soil moisture and sea surface temperatures](#)
1017 [similarly important for land climate in the warm season in the Community Earth](#)
1018 [System Model. *J. Clim.*,.](#)

1019 [Orth, R., R. D. Koster, and S. I. Seneviratne, 2013: Inferring Soil Moisture Memory from](#)
1020 [Streamflow Observations Using a Simple Water Balance Model. *J. Hydrometeorol.*,](#)
1021 [14, 1773–1790, doi:10.1175/JHM-D-12-099.1.](#)

1022 [Peings, Y., H. Douville, R. Alkama, and B. Decharme, 2011: Snow contribution to springtime](#)
1023 [atmospheric predictability over the second half of the twentieth century. *Clim. Dyn.*,](#)
1024 [37, 985–1004, doi:10.1007/s00382-010-0884-1.](#)

1025 [Perket, J., M. G. Flanner, and J. E. Kay, 2014: Diagnosing shortwave cryosphere radiative](#)
1026 [effect and its 21st century evolution in CESM. *J. Geophys. Res. Atmospheres*, **119**,](#)
1027 [1356–1362, doi:10.1002/2013JD021139.](#)

1028 [Qu, X., and A. Hall, 2014: On the persistent spread in snow-albedo feedback. *Clim. Dyn.*, **42**,](#)
1029 [69–81, doi:10.1007/s00382-013-1774-0.](#)

1030 [Reager, J. T., A. S. Gardner, J. S. Famiglietti, D. N. Wiese, A. Eicker, and M.-H. Lo, 2016: A](#)
1031 [decade of sea level rise slowed by climate-driven hydrology. *Science*, **351**, 699–703,](#)
1032 [doi:10.1126/science.aad8386.](#)

1033 [Rodell, M., I. Velicogna, and J. S. Famiglietti, 2009: Satellite-based estimates of groundwater](#)
1034 [depletion in India. *Nature*, **460**, 999–1002, doi:10.1038/nature08238.](#)

1035 [Seneviratne, S. I., D. Luthi, M. Litschi, and C. Schar, 2006: Land-atmosphere coupling and](#)
1036 [climate change in Europe. *Nature*, **443**, 205–209, doi:10.1038/nature05095.](#)

1037 [—, T. Corti, E. L. Davin, M. Hirschi, E. B. Jaeger, I. Lehner, B. Orlowsky, and A. J. Teuling,](#)
1038 [2010: Investigating soil moisture–climate interactions in a changing climate: A](#)
1039 [review. *Earth-Sci. Rev.*, **99**, 125–161, doi:10.1016/j.earscirev.2010.02.004.](#)

1040 [—, and Coauthors, 2013: Impact of soil moisture-climate feedbacks on CMIP5 projections:](#)
1041 [First results from the GLACE-CMIP5 experiment. *Geophys. Res. Lett.*, **40**,](#)
1042 [2013GL057153, doi:10.1002/grl.50956.](#)

1043 [Seneviratne, S. I., B. Van den Hurk, D. M. Lawrence, G. Krinner, G. Hurtt, H. Kim, C. Derksen,](#)
1044 [and et al., 2014: Land Processes, Forcings, and Feedbacks in Climate Change](#)
1045 [Simulations: The CMIP6 “LandMIPs.” *GEWEX Newsl.*, 6–10.](#)

1046 [Seneviratne, S. I., M. G. Donat, A. J. Pitman, R. Knutti, and R. L. Wilby, 2016: Allowable CO₂](#)
1047 [emissions based on regional and impact-related climate targets. *Nature*, **529**, 477–](#)
1048 [483.](#)

1049 [Sheffield, J., G. Goteti, and E. F. Wood, 2006: Development of a 50-Year High-Resolution](#)
1050 [Global Dataset of Meteorological Forcings for Land Surface Modeling. *J. Clim.*, **19**,](#)
1051 [3088–3111, doi:10.1175/JCLI3790.1.](#)

1052 [—, E. F. Wood, and M. L. Roderick, 2012: Little change in global drought over the past 60](#)
1053 [years. *Nature*, **491**, 435–438, doi:10.1038/nature11575.](#)

1054 [Singh, D., M. G. Flanner, and J. Perket, 2015: The global land shortwave cryosphere radiative](#)
1055 [effect during the MODIS era. *The Cryosphere*, **9**, 2057–2070, doi:10.5194/tc-9-2057-](#)
1056 [2015.](#)

1057 [Sitch, S., and Coauthors, 2015: Recent trends and drivers of regional sources and sinks of](#)
1058 [carbon dioxide. *Biogeosciences*, **12**, 653–679, doi:10.5194/bg-12-653-2015.](#)

1059 [Thackeray, C. W., C. G. Fletcher, and C. Derksen, 2015a: Quantifying the skill of CMIP5](#)
1060 [models in simulating seasonal albedo and snow cover evolution. *J. Geophys. Res.*](#)
1061 [Atmospheres, **120**, 5831–5849, doi:10.1002/2015JD023325.](#)

1062 [—, —, and —, 2015b: Quantifying the skill of CMIP5 models in simulating seasonal](#)
1063 [albedo and snow cover evolution. *J. Geophys. Res. Atmospheres*, **120**, 5831–5849,](#)
1064 [doi:10.1002/2015JD023325.](#)

1065 [Thomas, J., A. Berg, and W. Merryfield, 2015: Influence of snow and soil moisture](#)
1066 [initialization on sub-seasonal predictability and forecast skill in boreal spring. *Clim.*](#)
1067 [*Dyn.*, 1–17, doi:10.1007/s00382-015-2821-9.](#)

1068 [Trenberth, K. E., A. Dai, G. van der Schrier, P. D. Jones, J. Barichivich, K. R. Briffa, and J.](#)
1069 [Sheffield, 2014: Global warming and changes in drought. *Nat. Clim Change*, **4**, 17–22.](#)

1070 [Wada, Y., L. P. Beek, F. C. Sperna Weiland, B. F. Chao, Y. Wu, and M. F. Bierkens, 2012: Past](#)
1071 [and future contribution of global groundwater depletion to sea-level rise. *Geophys.*](#)
1072 [*Res. Lett.*, **39**.](#)

1073 [Warszawski, L., K. Frieler, V. Huber, F. Piontek, O. Serdeczny, and J. Schewe, 2014: The Inter-](#)
1074 [Sectoral Impact Model Intercomparison Project \(ISI-MIP\): Project framework. *Proc.*](#)
1075 [*Natl. Acad. Sci.*, **111**, 3228–3232.](#)

1076 [Watanabe, S., and Coauthors, 2014: Application of performance metrics to climate models](#)
1077 [for projecting future river discharge in the Chao Phraya River basin. *Hydrol. Res.*](#)
1078 [*Lett.*, **8**, 33–38, doi:10.3178/hrl.8.33.](#)

1079 [Weedon, G. P., and Coauthors, 2011: Creation of the WATCH Forcing Data and Its Use to](#)
1080 [Assess Global and Regional Reference Crop Evaporation over Land during the](#)
1081 [Twentieth Century. *J. Hydrometeorol.*, **12**, 823–848, doi:10.1175/2011JHM1369.1.](#)

1082 [Weedon, G. P., G. Balsamo, N. Bellouin, S. Gomes, M. J. Best, and P. Viterbo, 2014: The](#)
1083 [WFDEI meteorological forcing data set: WATCH Forcing Data methodology applied to](#)
1084 [ERA-Interim reanalysis data. *Water Resour. Res.*, **50**, 7505–7514,](#)
1085 [doi:10.1002/2014WR015638.](#)

1086 [Wei, Y., and Coauthors, 2014: The North American Carbon Program Multi-scale Synthesis](#)
1087 [and Terrestrial Model Intercomparison Project – Part 2: Environmental driver data.](#)
1088 [*Geosci Model Dev*, **7**, 2875–2893, doi:10.5194/gmd-7-2875-2014.](#)

1089 [Xu, L., and P. Dirmeyer, 2011: Snow-atmosphere coupling strength in a global atmospheric](#)
1090 [model. *Geophys. Res. Lett.*, **38**, n/a – n/a, doi:10.1029/2011GL048049.](#)

1091 [—, and —, 2012: Snow–Atmosphere Coupling Strength. Part II: Albedo Effect Versus](#)
1092 [Hydrological Effect. *J. Hydrometeorol.*, **14**, 404–418, doi:10.1175/JHM-D-11-0103.1.](#)

1093 [Xu, L., and Coauthors, 2013: Temperature and vegetation seasonality diminishment over](#)
1094 [northern lands. *Nat. Clim Change*, **3**, 581–586.](#)

1095 [Yoshimura, K., and M. Kanamitsu, 2008: Dynamical Global Downscaling of Global Reanalysis.](#)
1096 [*Mon. Weather Rev.*, **136**, 2983–2998, doi:10.1175/2008MWR2281.1.](#)

1097 [—, and —, 2013: Incremental Correction for the Dynamical Downscaling of Ensemble](#)
1098 [Mean Atmospheric Fields. *Mon. Weather Rev.*, **141**, 3087–3101, doi:10.1175/MWR-](#)
1099 [D-12-00271.1.](#)

1100 [Zampieri M, and Lionello P, 2011: Anthropic land use causes summer cooling in](#)
1101 [-Central Europe. *Clim. Res.*, **46**, 255–268.](#)

1102 [Zampieri, M., E. Serpetzoglou, E. N. Anagnostou, E. I. Nikolopoulos, and A. Papadopoulos,](#)
1103 [2012: Improving the representation of river–groundwater interactions in land](#)
1104 [surface modeling at the regional scale: Observational evidence and parameterization](#)
1105 [applied in the Community Land Model. *J. Hydrol.*, **420–421**, 72–86,](#)
1106 [doi:10.1016/j.jhydrol.2011.11.041.](#)

1107 [Zampieri, M., E. Scoccimarro, S. Gualdi, and A. Navarra, 2015: Observed shift towards earlier](#)
1108 [spring discharge in the main Alpine rivers. *Better Underst. Links Stress. Hazard*](#)
1109 [*Assess. Ecosyst. Serv. Water Scarcity*, **503–504**, 222–232,](#)
1110 [doi:10.1016/j.scitotenv.2014.06.036.](#)

1111 [Zhang, Y., and Coauthors, submitted: A Climate Data Record \(CDR\) for the global terrestrial](#)
1112 [water budget: 1984-2010. *J. Clim.*.](#)

1113 [Zscheischler, J., R. Orth, and S. I. Seneviratne, 2015: A submonthly database for detecting](#)
1114 [changes in vegetation-atmosphere coupling. *Geophys. Res. Lett.*, **42**, 9816–9824,](#)
1115 [doi:10.1002/2015GL066563.](#)

1116 ▲ **Met opmaak:** Engels (V.S.)

1117

1118 | **Annex: output data tables requested for LS3MIP**

1119

1120 | *Table A1: Variable request table “LEday”: daily variables related to the energy cycle. Priority*
 1121 | *index (p*) in column 1 indicates 1: “Mandatory” and 2: “Desirable”. The dimension (dim.)*
 1122 | *column indicates T: time, Y: latitude, X: longitude, and Z: soil or snow layers. “Direction”*
 1123 | *identifies the direction of positive numbers.*

p*	name	standard_name (cf)	long_name (netCDF)	unit	direction	dim.
1	rss	surface_net_downward_shortwave_flux	Net shortwave radiation	W/m ²	Downward	TYX
1	rls	surface_net_downward_longwave_flux	Net longwave radiation	W/m ²	Downward	TYX
2	rsds	surface_downwelling_shortwave_flux_in_air	Downward short-wave radiation	W/m ²	Downward	TYX
2	rlsds	surface_downwelling_longwave_flux_in_air	Downward long-wave radiation	W/m ²	Downward	TYX
2	rsus	surface_upwelling_shortwave_flux_in_air	Upward short-wave radiation	W/m ²	Upward	TYX
2	rlus	surface_upwelling_longwave_flux_in_air	Upward long-wave radiation	W/m ²	Upward	TYX
1	hfls	surface_upward_latent_heat_flux	Latent heat flux	W/m ²	Upward	TYX
1	hfss	surface_upward_sensible_heat_flux	Sensible heat flux	W/m ²	Upward	TYX
1	hfds	surface_downward_heat_flux	Ground heat flux	W/m ²	Downward	TYX
1	hfdsn	surface_downward_heat_flux_in_snow	Downward heat flux into snow	W/m ²	Downward	TYX
2	hfmlt	surface_snow_and_ice_melt_heat_flux	Energy of fusion	W/m ²	Soild to Liquid	TYX
2	hfsbl	surface_snow_and_ice_sublimation_heat_flux	Energy of sublimation	W/m ²	Soild to Vapor	TYX
2	tau	surface_downward_stress	Momentum flux	N/m ²	Downward	TYX
2	hfrs	temperature_flux_due_to_rainfall_expressed_as_heat_flux_onto_snow_and_ice	Heat transferred to snowpack by rainfall	W/m ²	Downward	TYX
1	dtes	change_over_time_in_thermal_energy_content_of_surface	Change in surface heat storage	J/m ²	Increase	TYX
1	dtesn	change_over_time_in_thermal_energy_content_of_surface_snow_and_ice	Change in snow/ice cold content	J/m ²	Increase	TYX
1	ts	surface_temperature	Average surface temperature	K	-	TYX
2	tsns	surface_snow_skin_temperature	Snow Surface Temperature	K	-	TYX
2	tcs	surface_canopy_skin_temperature	Vegetation Canopy Temperature	K	-	TYX
2	tgss	surface_ground_skin_temperature	Temperature of bare soil	K	-	TYX
2	tr	surface_radiative_temperature	Surface Radiative Temperature	K	-	TYX
1	albs	surface_albedo	Surface Albedo	-	-	TYX
1	albsn	snow_and_ice_albedo	Snow Albedo	-	-	TYX

1	snc	surface_snow_area_fraction	Snow covered fraction	-	-	TYX
2	albc	canopy_albedo	Canopy Albedo	-	-	TYX
2	cnc	surface_canopy_area_fraction	Canopy covered fraction	-	-	TYX
1	tsl	soil_temperature	Average layer soil temperature	K	-	TZYX
1	tsnl	snow_temperature	Temperature profile in the snow	K	-	TZYX
1	tasmax	air_temperature_maximum	Daily Maximum Near-Surface Air Temperature	K	-	TYX
1	tasmin	air_temperature_minimum	Daily Minimum Near-Surface Air Temperature	K	-	TYX
2	clt	cloud_area_fraction	Total cloud fraction	-	-	TYX

1124

Table A2: Variable request table "LWday": daily variables related to the water cycle.

p*	name	standard_name (cf)	long_name (netCDF)	unit	direction	dim.
1	pr	precipitation_flux	Precipitation rate	kg/m ² /s	Downward	TYX
2	prra	rainfall_flux	Rainfall rate	kg/m ² /s	Downward	TYX
2	prsn	snowfall_flux	Snowfall rate	kg/m ² /s	Downward	TYX
2	prrc	convective_rainfall_flux	Convective Rainfall rate	kg/m ² /s	Downward	TYX
2	prnc	convective_snowfall_flux	Convective Snowfall rate	kg/m ² /s	Downward	TYX
1	prveg	precipitation_flux_onto_canopy	Precipitation onto canopy	kg/m ² /s	Downward	TYX
1	et	surface_evapotranspiration	Total Evapotranspiration	kg/m ² /s	Upward	TYX
1	ec	liquid_water_evaporation_flux_from_canopy	Interception evaporation	kg/m ² /s	Upward	TYX
1	tran	Transpiration	Vegetation transpiration	kg/m ² /s	Upward	TYX
1	es	liquid_water_evaporation_flux_from_soil	Bare soil evaporation	kg/m ² /s	Upward	TYX
2	eow	liquid_water_evaporation_flux_from_open_water	Open water evaporation	kg/m ² /s	Upward	TYX
2	esn	liquid_water_evaporation_flux_from_surface_snow	Snow Evaporation	kg/m ² /s	Upward	TYX
2	sbl	surface_snow_and_ice_sublimation_flux	Snow sublimation	kg/m ² /s	Upward	TYX
2	slbnosn	sublimation_amount_assuming_no_snow	Sublimation of the snow free area	kg/m ² /s	Upward	TYX
2	potet	water_potential_evapotranspiration_flux	Potential Evapotranspiration	kg/m ² /s	Upward	TYX
1	mrro	runoff_flux	Total runoff	kg/m ² /s	Out	TYX
2	mrros	surface_runoff_flux	Surface runoff	kg/m ² /s	Out	TYX
1	mrrob	subsurface_runoff_flux	Subsurface runoff	kg/m ² /s	Out	TYX
1	snm	surface_snow_and_ice_melt_flux	Snowmelt	kg/m ² /s	Solid to liquid	TYX
1	snrefr	surface_snow_and_ice_refreezing_flux	Re-freezing of water in the snow	kg/m ² /s	Liquid to solid	TYX
2	snmsl	surface_snow_melt_flux_into_soil_layer	Water flowing out of snowpack	kg/m ² /s	Out	TYX
2	qgwr	water_flux_from_soil_layer_to_groundwater	Groundwater recharge from soil layer	kg/m ² /s	Out	TYX
2	rivo	water_flux_from_upstream	River Inflow	m ³ /s	In	TYX
2	rivi	water_flux_to_downstream	River Discharge	m ³ /s	Out	TYX
1	dslw	change_over_time_in_water_content_of_soil_layer	Change in soil moisture	kg/m ²	Increase	TYX
1	dsn	change_over_time_in_surface_snow_and_ice_amount	Change in snow water equivalent	kg/m ²	Increase	TYX
1	dsw	change_over_time_in_surface_water_amount	Change in Surface Water Storage	kg/m ²	Increase	TYX
1	dcw	change_over_time_in_canopy_water_amount	Change in interception storage	kg/m ²	Increase	TYX

2	dgw	change_over_time_in_groundwater	Change in groundwater	kg/m ²	Increase	TYX
2	drivw	change_over_time_in_river_water_amount	Change in river storage	kg/m ²	Increase	TYX
1	rzwc	water_content_of_root_zone	Root zone soil moisture	kg/m ²	-	TYX
1	cw	canopy_water_amount	Total canopy water storage	kg/m ²	-	TYX
1	snw	surface_snow_amount	Snow Water Equivalent	kg/m ²	-	TZYX
1	snwc	canopy_snow_amount	SWE intercepted by the vegetation	kg/m ²	-	TYX
2	lwsnl	liquid_water_content_of_snow_layer	Liquid water in snow pack	kg/m ²	-	TZYX
1	sw	surface_water_amount_assuming_no_snow	Surface Water Storage	kg/m ²	-	TYX
1	mrlsl	moisture_content_of_soil_layer	Average layer soil moisture	kg/m ²	-	TZYX
1	mrsos	moisture_content_of_soil_layer	Moisture in top soil (10cm) layer	kg/m ²	-	TYX
1	mrsow	relative_soil_moisture_content_above_field_capacity	Total Soil Wetness	-	-	TYX
2	wtd	depth_of_soil_moisture_saturation	Water table depth	m	-	TYX
1	tws	canopy_and_surface_and_subsurface_water_amount	Terrestrial Water Storage	kg/m ²	-	TYX
2	mrlqso	mass_fraction_of_unfrozen_water_in_soil_layer	Average layer fraction of liquid moisture	-	-	TZYX
1	mrfsofr	mass_fraction_of_frozen_water_in_soil_layer	Average layer fraction of frozen moisture	-	-	TZYX
2	prrsn	mass_fraction_of_rainfall_onto_snow	Fraction of rainfall on snow.	-	-	TYX
2	prnsn	mass_fraction_of_snowfall_onto_snow	Fraction of snowfall on snow.	-	-	TYX
1	lqsn	mass_fraction_of_liquid_water_in_snow	Snow liquid fraction	-	-	TZYX
1	snd	surface_snow_thickness	Depth of snow layer	m	-	TYX
1	agesno	age_of_surface_snow	Snow Age	day	-	TYX
2	sootsn	soot_content_of_surface_snow	Snow Soot Content	kg/m ²	-	TYX
2	sic	sea_ice_area_fraction	Ice-covered fraction	-	-	TYX
2	sit	sea_ice_thickness	Sea-ice thickness	m	-	TYX
2	dfr	depth_of_frozen_soil	Frozen soil depth	m	Downward	TYX
2	dmlt	depth_of_subsurface_melting	Depth to soil thaw	m	Downward	TYX
2	tpf	permafrost_layer_thickness	Permafrost Layer Thickness	m	-	TYX
2	pflw	liquid_water_content_of_permafrost_layer	Liquid water content of permafrost layer	kg/m ²	-	TYX
			Aerodynamic conductance	m/s	-	TYX
2	ares	aerodynamic_resistance	Aerodynamic resistance	s/m	-	TYX
1	<u>nudgincw</u>	<u>nudging_increment_of_total_water</u>	<u>Nudging Increment of Water</u>	<u>kg/m2</u>	<u>Increase</u>	<u>TYX</u>
1	hur	relative_humidity	Relative humidity	%	-	TYX
1	hurmax	relative_humidity_maximum	Daily Maximum Near-Surface Relative Humidity	%	-	TYX

1126

1	hurmin	relative_humidity_minimum	Daily Minimum Near-Surface Relative Humidity	%	-	TYX
---	--------	---------------------------	---	---	---	-----

1127 *Table A3: Variable request table “LCmon”: monthly variables related to the carbon cycle.*

P*	name	standard_name (cf)	long_name (netCDF)	unit	direction	dim.
1	gpp	gross_primary_productivity _of_carbon	Gross Primary Production	Kg/m ² /s	Downward	TYX
1	npp	net_primary_productivity_of_carbon	Net Primary Production	Kg/m ² /s	Downward	TYX
1	nep	surface_net_downward_mass_flux _of_carbon_dioxide_expressed _as_carbon_due_to _all_land_processes_excluding _anthropogenic_land_use_change	Net Ecosystem Exchange	Kg/m ² /s	Downward	TYX
1	ra	plant_respiration_carbon_flux	Autotrophic Respiration	Kg/m ² /s	Upward	TYX
1	rh	heterotrophic_respiration _carbon_flux	Heterotrophic Respiration	Kg/m ² /s	Upward	TYX
1	fLuc	surface_net_upward_mass_flux _of_carbon_dioxide_expressed _as_carbon_due_to_emission_from _anthropogenic_land_use_change	Net Carbon Mass Flux into Atmosphere due to Land Use Change	Kg/m ² /s	Upward	TYX
1	cSoil	soil_carbon_content	Carbon Mass in Soil Pool	Kg/m ²	-	TYX
1	cLitter	litter_carbon_content	Carbon Mass in Litter Pool	Kg/m ²	-	TYX
1	cVeg	vegetation_carbon_content	Carbon Mass in Vegetation	Kg/m ²	-	TYX
1	cProduct	carbon_content_of_products_of _anthropogenic_land_use_change	Carbon Mass in Products of Land Use Change	Kg/m ²	-	TYX
2	cLeaf	leaf_carbon_content	Carbon Mass in Leaves	Kg/m ²	-	TYX
2	cWood	wood_carbon_content	Carbon Mass in Wood	Kg/m ²	-	TYX
2	cRoot	root_carbon_content	Carbon Mass in Roots	Kg/m ²	-	TYX
2	cMisc	miscellaneous_living_matter _carbon_content	Carbon Mass in Other Living Compartments on Land	Kg/m ²	-	TYX
2	fVegLitter	litter_carbon_flux	Total Carbon Mass Flux from Vegetation to Litter	Kg/m ² /s	-	TYX
2	fLitterSoil	carbon_mass_flux_into_soil _from_litter	Total Carbon Mass Flux from Litter to Soil	Kg/m ² /s	-	TYX
2	fVegSoil	carbon_mass_flux_into_soil _from_vegetation_excluding_litter	Total Carbon Mass Flux from Vegetation Directly to Soil	Kg/m ² /s	-	TYX
1	treeFrac	area_fraction	Tree Cover Fraction	%	-	TYX
1	grassFrac	area_fraction	Natural Grass Fraction	%	-	TYX
1	shrubFrac	area_fraction	Shrub Fraction	%	-	TYX
1	cropFrac	area_fraction	Crop Fraction	%	-	TYX
1	pastureFrac	area_fraction	Anthropogenic Pasture Fraction	%	-	TYX

1128

1	baresoilFrac	area_fraction	Bare Soil Fraction	%	-	TYX
1	residualFrac	area_fraction	Fraction of Grid Cell that is Land but Neither Vegetation-Covered nor Bare Soil	%	-	TYX
1	lai	leaf_area_index	Leaf Area Index	Kg/m ²	-	TYX

1129 *Table A4: Variable request table “L3hr”: 3-hourly variables to generate the atmospheric*
1130 *boundary conditions for the off-line simulation.*


p*	name	standard_name (cf)	long_name (netCDF)	unit	direction	dim.
1	rsds	surface_downwelling_shortwave_flux_in_air	Downward short-wave radiation	W/m ²	Downward	TYX
1	rlds	surface_downwelling_longwave_flux_in_air	Downward long-wave radiation	W/m ²	Downward	TYX
1	hus	specific_humidity	Near surface specific humidity	kg/kg	-	TYX
1	ta	air_temperature	Near surface air temperature	K	-	TYX
1	ps	surface_air_pressure	Surface Pressure	Pa	-	TYX
1	ws	wind_speed	Near surface wind speed	m/s	-	TYX
2	va	northward_wind	Near surface northward wind component	m/s	Northward	TYX
2	ua	eastward_wind	Near surface eastward wind component	m/s	Eastward	TYX
2	pr	precipitation_flux	Precipitation rate	kg/m ² /s	Downward	TYX
1	prra	rainfall_flux	Rainfall rate	kg/m ² /s	Downward	TYX
1	prsn	snowfall_flux	Snowfall rate	kg/m ² /s	Downward	TYX
2	prrc	convective_rainfall_flux	Convective Rainfall rate	kg/m ² /s	Downward	TYX
2	prsn	convective_snowfall_flux	Convective Snowfall rate	kg/m ² /s	Downward	TYX
1	clt	cloud_area_fraction	Total cloud fraction	-	-	TYX
2	co2c	mole_fraction_of_carbon_dioxide_in_air	Near surface CO2 concentration	-	-	TYX

1131

1132

1133 Figures

1134




1135

1136 *Figure 1: Structure of the “LandMIPs”. LS3MIP includes (1) the offline representation of land*
 1137 *processes (LMIP) and (2) the representation of land-atmosphere feedbacks related to snow*
 1138 *and soil moisture (LFMIP). Forcing associated with land use is assessed in LUMIP. Substantial*
 1139 *links also exist to C4MIP (terrestrial carbon cycle). Furthermore, a land albedo testbed*
 1140 *experiment is planned within GeoMIP. From Seneviratne et al. (2014)*

1141

1142

1143



1144

1145 *Figure 24: Relevance of LS3MIP for WCRP Core Projects and Grand Challenges¹⁹*

1146

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

¹⁹ <http://wcrp-climate.org/index.php/grand-challenges>; status Dec 2015




Figure 32: Embedding of LS3MIP within CMIP6. Adapted from Eyring et al. (2015)






Figure 3: Structure of the “LandMIPs”. LS3MIP includes (1) the offline representation of land processes (LMIP) and (2) the representation of land-atmosphere feedbacks related to snow and soil moisture (LFMIP). Forcing associated with land use is assessed in LUMIP. Substantial links also exist to C4MIP (terrestrial carbon cycle). Furthermore, a land albedo testbed experiment is planned within GeoMIP. From Seneviratne et al. (2014)

1159

1160



1161



1162

Figure 4: Schematic diagram for the experiment structure of LS3MIP. Tier 1 experiments are indicated with a heavy black outline, and complementary ensemble experiments are indicated with white hatched lines. Land-Altforc represents 3 alternative forcings for the Land-Hist experiment. For further details on the experiments and acronyms, see Table 1 and text.




Figure 5: Taylor diagram for evaluating the forcing datasets comparing to daily observations from FLUXNET sites, as used by (Best et al. 2015): (a) 2m air temperature and (b) precipitation. Red, blue, and green dots indicate GSWP3, Watch Forcing Data (Weedon et al. 2011) and Princeton forcing (Sheffield et al. 2006), respectively. Grey and orange dots indicate 20CR and its dynamically downscaled product (GSM248).




Figure 6: Global distributions of the similarity index (Ω) for 2001-2010 of monthly mean (a, c) and (b, d) monthly variance (calculated from daily data from each data set) of 2m air temperature (top panels) and precipitation (bottom panels), respectively. Shown are global distributions and zonal means. After (Kim 2010).

Gewijzigde veldcode