

# RESPONSE TO THE REFEREES COMMENTS

Title: ORCHIDEE-MICT (revision 4126), a land surface model for the high-latitudes: model description and validation

Author(s): Matthieu Guimberteau et al.

MS No.: gmd-2017-122

MS Type: Model evaluation paper

Iteration: Revised Submission

We would like to express our gratitude towards the two anonymous Referees for their constructive comments. We value very much their help in our effort toward a revised version of the manuscript which is at the end of this document. In the following, we write our point by point response.

- Reviewer's comments are in bold
- Modifications done in the new submitted version of the manuscript are in blue
- Figure and Table numbers, line numbers and pages all correspond to the initial manuscript version

## REFEREE #1

**This paper presents and evaluates the ORCHIDEE “high latitude” model version which is known as ORCHIDEE-MICT. The difference compared with the trunk version of ORCHIDEE is a vertically discretized soil carbon scheme and the coupling of this scheme to the soil thermal/hydrological properties, along with a representation of fire. What I really like about the paper is the coincident evaluation of so many variables (in order to correctly interpret interrelated biases) and the use of multiple datasets for the same variable when those are available (in order to give an idea of observational uncertainties). It is also great that two different forcing datasets were used, and I think it is a valuable conclusion that the uncertainty in forcing datasets needs to be taken into account to avoid “over-calibrating”.**

Clearly, this paper is extremely long. I think a suggestion of splitting it could be rightly met with the argument that all of the components are interacting so it would be difficult to split. However, depending on what the other reviewers or editors think, there could be a reasonable split into two linked papers that cover thermal/hydrological processes (in one paper) and carbon cycle processes (in another). Given the size of the work, it is well written so that it does not become too confusing to the reader. So for now (aside from the idea of splitting into two papers) I suggest only minor changes.

We agree that the paper is long but we think that splitting it into two linked papers would not be necessary, as mentioned by the reviewer #2 in his first general comment. Following the comments of this reviewer, we propose to remove the last paragraph on page 4 and replace it by a table of

content between the abstract and the introduction. Moreover, Figure 4 will be moved to the supplementary section, as you suggested in point 3. We propose also to put the description of the evaluation datasets (sections 5.2 to 5.4, P.12 to 20) into an appendix section after the conclusions for a better clarity of the paper.

**1. A theme that runs through the paper is the late response of LAI in the spring. The reason that is suggested (several times) in the paper is that this could be linked to the late persistence of snow cover. However, from my experience of such land surface models, they often don't incorporate a direct influence of snow on vegetation - perhaps this is incorporated in ORCHIDEE? But if so, can you make it more clear in the paper how the snow cover influences the vegetation in the model? The late LAI in spring also occurs in ORCHIDEE simulations where the snow does not stay too late (Chadburn et al., 2017), and I have heard that bud burst is simply triggered by the number of growing degree days and requires a rather large number to initiate bud burst. Therefore, I suggest you consider this alternative (possibly, more likely?) explanation, and recommend further study on the phenology scheme.**

The reviewer is right, the snow cover is not directly taken into account in the vegetation phenology models used for the different PFTs of ORCHIDEE. After the sentences on Page 26, Lines 13-15: "In all the basins, the LAI simulated by ORCHIDEE-MICT has a phase delay of up to one month compared to both products. This is due to a delay in the start of the growing season, which may be related to excessive persistence of the snow cover (Fig. 10)" in section 8.1, we thus detailed the arguments to support our hypothesis: "The phenological models in ORCHIDEE (detailed in MacBean et al., 2015, Appendix A) do not explicitly take into account this influence, unlike what is done in Van Wijk et al. (2003), who model the link of the start of the tussock tundra growing season to the soil thaw at 10-cm depth. However there is a first indirect link between the snow cover and the vegetation phenology through air temperature, which influences both the start of the growing season, determined in ORCHIDEE using growing degree days (GDD)-based phenological models for deciduous species, and the start of the snowmelt season. There is a second indirect link through snowmelt. While there is still a large amount of snow, the soil surface temperature is kept at zero degree Celsius or below and the soil cannot thaw. Only when snowmelt occurs and when the snow fraction is small enough, will the soil start thawing, thus increasing soil liquid water content. This impacts the start of the growing season for grasses and crops, which use both a GDD and a soil moisture thresholds, and also reduces water stress, thus favoring photosynthesis for all PFTs."



In the discussion part, section 10.3, after the sentences Lines 32-33 P.33: “The phase of simulated LAI in spring lags satellite observations, in particular for BONA and BOAS sub-regions (Fig. 14).”, we nevertheless acknowledge the reference to Chadburn et al. (2017), cited by the reviewer: “We argued that this lag is related to the late persistence of the snow cover. However a recent work (Chadburn et al., 2017) shows a late onset even in the absence of snow persistence on site simulations by ORCHIDEE. This calls for a revisit of the phenology-related thresholds in the high-latitudes, perhaps by introducing new PFTs (arctic C3 grass and shrub, and non-vascular plants), with their separate set of parameters calibrated, to better represent Arctic vegetation and their phenology (Druel et al., 2017).”.

**2. Another bias is the deep active layer, which really suggests that the soil properties should be better representative of the organic carbon content. The high organic content at the surface is quite well simulated (Figure 22), and this should have a great impact on the soil temperature. I would suggest that the problem might be the use of the linear weighted average for thermal conductivity. In terms of water and ice, the geometric mean is used - Equation 4 - so it might make sense to use that form for weighting the soil thermal conductivity as well, particularly the dry soil thermal conductivity which can be very low for an organic soil.**

We would like to note that the soil carbon concentrations used in the thermal and hydrological modules were prescribed from the empirical soil databases of NCSCD and HWSD (as mentioned on Page 8 Lines 15-16), similar to the treatment in other land surface models like CLM (Lawrence and Slater, 2008) and JULES (Chadburn et al., 2015). So the modeled SOC by the carbon cycle module did not yet feed back on the soil physical properties. Although the coupling between soil thermodynamics and the prognostically modeled SOC is readily achievable by changing a configuration in the model set-up, we tend to exclude the bias of the carbon cycle module in the physical processes in this study as a first step.

The simulated ALT is indeed generally overestimated compared to the site observations from CALM network and the regional ALT map for Yakutia. However, apart from the bias in the parameterization of soil thermal properties, the fact that we did not use the site-specific organic layer thickness for the CALM sites, and that we did not prescribe regionally a fixed thickness for organic layer (to mimic the insulating effect of moss layer), also contributed to the bias in modeled ALT. Now we conducted, following the comment by Reviewer #2, additional CALM site simulations in which site-specific organic layer thicknesses were prescribed. These site runs then produced significantly shallower ALTs compared to the previous regional simulation. Please refer to our second response to Reviewer #2 for details.

As for the averaging method for soil thermal conductivity, it makes sense indeed to use geometric mean, which will be tested for the next steps of model development. We added the following sentence after the sentence on Page 9 Line 2: “[Note that here we followed Lawrence and Slater \(2008\) to use linear weighting organic and mineral soil properties, while in some other models like JULES \(Chadburn et al., 2015\) and ISBA \(Decharme et al., 2016\), soil thermal conductivities are calculated as geometric averages of organic and mineral soils, consistent with the treatment for soil water and ice \(Eq. 4\). The geometric averaging method increases the effect of the organic fraction compared to arithmetic averages, and would be tested in ORCHIDEE-MICT in future developments.](#)”

**In the seasonal cycle of NEE is an unrealistic peak of emissions in spring, which can be partly due to the late LAI already discussed, but also partly because of soil decomposition starting too early. You talked about CO<sub>2</sub> trapping in the soil (P34 Line 16) but in fact it may be much more simply that the ground is thawing too quickly - again, due to the lack of thermal insulation from the organic layer. Although when the seasonal cycle of soil temperature is studied (Figure 6), this is not obviously the case, I think there might be a bias in the Russian dataset as it seemed to behave differently from other in-situ data. The problem is potentially with the removal of vegetation from the surfaces and site disturbance, which can result in the insulation of the ‘organic layer’ being removed (Frauenfeld et al., 2004). Certainly when comparing with in situ data in Chadburn et al. (2017), the soil in ORCHIDEE is thawing too early and likewise the soil respiration starts a bit too early in the spring.**

**I suggest adding discussion of the above points relating to the link between soil carbon and soil thermal properties.**

We would like first to note that the Russian dataset for soil temperatures between 0.2 and 3.2 m depths (that we used in this study) have been measured under natural vegetation (mainly grass) and undisturbed snow; although the grasses were regularly mowed to keep their height below 20 cm (as described in [http://nsidc.org/data/docs/fgdc/ggd251\\_soiltemp\\_fsul/](http://nsidc.org/data/docs/fgdc/ggd251_soiltemp_fsul/)), the main insulating effect of the organic matter near the ground surface has been kept. Therefore, bias of these measurements induced by human disturbance are generally very small.

Figure 6cd indeed shows a cold bias of soil temperature during spring; however, this figure is the average for all sites located in continuous permafrost region, while spatially, there is a warm bias in the Lena basin and a cold bias for the further eastern sites (please also see our second response to Reviewer #2), which partly offset each other in Fig. 6cd.

We agree that the spring peak in NEE may be partly explained by a too big soil respiration. But also note that the spring GPP is underestimated, especially in BOAS and BOEU (Fig. 15). To

address these issues, we modified the text on Page 34 Lines 15-17: “Interestingly the reasonable seasonal phase of simulated GPP (Fig. 14) can be contrasted with the larger lag of spring NEE uptake compared to inversions results (Fig. 19). This may be partly explained by the underestimated GPP at the beginning of the growing season (Fig. 14), and also by a possible too big soil respiration in spring. The moss/lichen surface coverage could be over 70% under the vast boreal forests (Porada et al., 2016), but we did not prescribe an additional moss layer in the regional simulations, which could lead to a too early thawing of the soil in spring.”

**3. These two issues that I have mentioned: The phenology and the organic soil thermal properties, both seem quite important to me, and worthy of being mentioned in the conclusion, along with the issue of snow thermal conductivity which is certainly too high. These issues are extensively discussed in the text but not mentioned in the conclusions. (In particular, the organic soil properties are the ‘new’ process that is included in the paper so it seems important to include them in the conclusion.) It seems to me that the other processes are appropriately discussed (at least, as far as my expertise goes: I can’t comment on fires or say much about forests.)**

We add these sentences in the conclusion:

- P. 34 L.27 “... in the ORCHIDEE-MICT land surface model. The effects of soil organic matter on soil thermal and hydraulic properties are incorporated.”
- P. 34 L.30 “Naturally, there remains significant room for improvement. The model appears to underestimate evapotranspiration and overestimate surface temperature, particularly in the southern portion of the boreal zone. Simulated phenology shows generally a delay in the onset of growing season. And the snow module underestimates the thermal insulation of snow.”

**I would like to suggest some kind of reduction in the text, as the same points are sometimes made a few times, but it is hard to envisage how to do this- I’m sure you have thought about the same thing! However, to shorten I suggest at least moving Figure 4 to the supplementary as it doesn’t contain observations and doesn’t seem so informative as the others.**

Yes, Figure 4 is moved to the supplementary section. We also propose to remove the last paragraph on page 4 and replace it by a table of content between the abstract and the introduction. We put the description of the evaluation datasets (sections 5.2 to 5.4) into an appendix section after the conclusion for a better clarity of the paper.

#### **Small comments:**

**P29 Lines 25: “SOC stocks simulated by the model fit the spatial pattern from observed inventory data” - this does not seem convincing to me looking at Figure 22, I think this statement should be more qualified e.g. ‘to some extent’ !**

Yes, we revised it as : “SOC stocks simulated by the model fit **to some extent** the spatial pattern from observed inventory data ...”.

**P28, line 27/8: just says “see 7” - should this be “see Figure 7”?**

Corrected now in the text.

---

#### **REFeree #2**

The manuscript presents a new version of the global land surface model ORCHIDEE, which aims at a more realistic representation of hydrological processes and carbon fluxes at high latitudes and is called ORCHIDEE-MICT. To this end, several new components are introduced, such as a vertical soil carbon profile, influence of soil carbon on soil thermal properties, and a revised scheme for plant water stress. The new model is thoroughly evaluated by comparing multiple variables, such as snow properties, soil moisture and temperature, runoff and evapotranspiration, GPP, NPP, biomass and soil carbon. Explanations for mismatches between model estimates and observations are provided. The paper is well written and of good scientific quality. I therefore recommend to publish it with some minor revisions (see below).

#### **General comments**

(1) I agree with reviewer #1 that the paper is quite long, but I do not think it is necessary to split the manuscript. Instead, I suggest replacing the last paragraph on page 4 with a table or schematic, similar to a table of contents, which lists the sections of the paper. This might help the reader to get a better overview of the paper.

Thank you for this good suggestion. We will remove the last paragraph on page 4 and replace it by a table of content between the abstract and the introduction. Moreover, Figure 4 is moved to the supplementary section, as suggested by the Reviewer #1. We propose also to put the description


of the evaluation dataset (sections 5.2 to 5.4) into an appendix section after the conclusion for a better clarity of the paper.

**(2) The simulated soil temperatures show, in general, a cold bias, only for one soil depth in combination with the GSWP3 forcing data the temperature is overestimated. However, ALT is significantly overestimated, which does not make much sense to me. Even if soil thermal conductivity was overestimated, leading to an overestimation of ALT, this does not explain why soil temperature is underestimated. Could you please explain this a bit more? Furthermore, I do not understand why spatial heterogeneity should lead to underestimated ALT in the field measurements (page 22, line 14)?**

Simulated ALT is indeed generally overestimated compared to the site observations from CALM network and the empirically-derived map for Yakutia (Beer et al., 2013). This seems inconsistent with the cold bias in soil temperature compared to the Russian meteorological stations' observations as shown in Fig. 6. However, we would like to clarify several points:

First, there is a mismatch for the locations of CALM sites for ALT and of Russian stations for soil temperature, the former mostly within the Arctic regions near the coast (for the Eurasian sites), and the latter more scattered throughout Russia. Therefore, the slight overestimation in peak summer-time soil temperature (and even underestimation with the CRUNCEP forcing) shown in Fig. 6cd for the Russian sites does not necessarily contradict with the significantly overestimated ALT compared to the CALM sites.

As for the regional maps for soil temperature and ALT, we acknowledge that Fig. 6ab may be a bit misleading if readers connect this figure to Fig. S3 which shows on the contrary a generally deeper ALT compared to Beer et al., (2013). This is because Fig. 6ab shows the annual mean temperature at 0.2 m depth, in which the larger cold bias in winter outweighs the warm bias in summer. To complement Fig. 6ab, we added maps for the maximum monthly soil temperature for the four depths in the Supplementary, also shown below, to facilitate a link between bias in soil temperature and bias in ALT. This figure shows an overestimation in maximum soil temperature below 0.8 m in the Lena basin, consistent with the deeper ALT for the same region shown in Fig. S3.




**Figure S4.** Maximum monthly soil temperature at different depths in **(a)** GSWP3 and **(b)** CRUNCEP-forced simulation (background maps), compared to the site observations (color filled circles), averaged over the period 1981-2000.

To clarify it, we revised the following sentences on Page 21 Lines 32-33: “Summer soil temperatures are higher in the GSWP3-forced simulation relative to those of CRUNCEP, and warmer than observations from the Russian meteorological stations in continuous permafrost region by 1~2 °C on average at 0.8 and 1.6 m depths (Fig. 5c,d). Spatially however, the bias in peak summer soil temperature varies for different regions, with a large warm bias for the Lena basin below 0.8 m, and some cold bias for the further eastern sites (Fig. S4). This is consistent with the overestimation of ALT for Yakutia (Fig. S5) compared to the empirically-derived map by Beer et al., (2013) (see Section 6.4). Differences between the two simulations...” And added one sentence at the end of the caption for Fig. 6: “...over the period 1981-2000. The spatial patterns of maximum monthly soil temperature are also shown in Fig. S4.”

Second, the mismatch between the local-scale ALT measurements at the CALM sites and the modeled ALT could be partly explained by the fact that we did not use the site information of the soil organic layer thickness in the calculation of soil physical properties, but used the gridded soil carbon database from NCSCD (Hugelius et al., 2013) for permafrost regions (as mentioned on Page 8 Lines 15-16), which was upscaled in the model to match the spatial resolution for each simulation (here 1° by 1°).

To further explore the impact of the site-specific organic layer thickness on modeled ALT, we conducted additional runs at the CALM sites, in which we assumed  $f_{i,\text{SOC}}$  in Eq. 9 equaling to one for the soil layers above the organic layer thickness at each site. While the other model inputs including climate and soil texture are the same as the previous northern hemisphere simulation. There are in total 69 sites that provide explicit organic layer thickness (<https://www2.gwu.edu/~calm/data/north.html>). Some sites, e.g. the Dot Lake in Alaska, also have a thick sphagnum layer above the decomposing organic soil; in such cases, we summed up their depths to derive a total organic layer thickness. The figure below shows the result, which is now added to the Supplementary.



**Figure S6.** (a) Scatter plot of modeled ALT forced by CRUNCEP compared with observed ALT from the CALM network, averaged over the period 1990-2007. The black circles represent the grid cells taken from the regional simulation (shown in Fig. 8b). The blue circles represent a subset of the CALM sites for which we performed additional runs using site-specific organic layer thicknesses, with the result shown by the red circles. (b) Illustration for the difference of the additional site simulations. Each grey arrow connects the same site, pointing from the blue circles using soil carbon content from NCSCD (upscaled at  $1^\circ$  by  $1^\circ$  resolution) to calculate soil physical properties, to the red circles using the organic layer thickness provided by the sites. See text for further information.

Accordingly, the following discussions were added on Page 22 Line 10: “...whereas CRUNCEP-forced output shows relatively better agreement with the observations. Apart from the uncertainty induced by climate forcing, the model-data mismatch may also arise from scale differences for the organic carbon content that is used to calculate soil physical properties for each grid cell. As mentioned in Section 4.2, the empirical SOC map from NCSCD (Hugelius et al., 2013) is prescribed for permafrost regions in the soil thermal and hydrological modules, which is upscaled



by the model to the target spatial resolution ( $1^\circ$  by  $1^\circ$  in this study). These SOC values thus do not represent the site-level soil conditions, aside from the uncertainty of the NCSCD database itself. To further investigate this impact, we conducted additional simulations for the sites that provide explicit organic layer thicknesses (in total 69 sites), forced by CRUNCEP. In these runs, we assumed pure organic soil, i.e.  $f_{i,\text{SOC}}$  in Eq. 9 equaling to one, for the soil layers above the site-specific organic layer thickness, while kept the SOC concentration unchanged below this thickness, i.e. from NCSCD. Note that the moss layer, vegetation mat, and/or organic root zone as described in some sites were all summed to derive a total organic layer thickness. The other configurations including climate forcing and soil texture were the same as the regional simulation. The result is displayed in Fig. S6, showing significantly shallower ALTs simulated by these site runs which better match the observations (Fig. S6a), with different magnitudes of ALT reductions among the sites (Fig. S6b).”

**(3) In the abstract, it is stated that the new processes put ORCHIDEE-MICT at the forefront of land surface modelling at high latitudes. However, I would expect more comparison at the process level to other models (e.g. JULES or JSBACH) to substantiate this statement, maybe through a short paragraph in the discussion. I would also like to know why the inclusion of an organic layer or a moss/lichen layer, which has been done in JULES (Chadburn et al, 2015, TC) and JSBACH (Porada et al, 2016, TC) was not considered? Could you please explain in this context the relation of ORCHIDEE-MICT to another ORCHIDEE version which is currently in review in GMD (Druel et al, 'Towards a more detailed representation of high-latitude vegetation in the global land surface model ORCHIDEE (ORC-HL-VEGv1.0)') ?**

The current model described in this study indeed lacks explicit representation of moss and lichen growth. For an inclusion of organic layer, however, the multi-layer structure of the model enables it to approximate the effect of an organic layer by assuming 100% organic soil above the prescribed organic layer thickness, as we did for the additional CALM site simulations. Actually, this method could also approximate the effect of moss/lichen on soil thermodynamics, assuming similar thermal properties of the moss/lichen layer to that of the organic layer, as the implementation in JULES (Chadburn et al., 2015, GMD). Therefore, the current ORCHIDEE-MICT is able to represent the insulating effect of moss/lichen and organic layer in a simplified way, given an input of their thickness.

However, for large-scale simulations to account for moss/lichen layer, what is indeed lacking in current ORCHIDEE-MICT is a prognostic modeling of moss/lichen surface cover, considering the lack of a gridded map for moss/lichen coverage, especially in the boreal forest understory (Chadburn et al., 2015, TC; Porada et al, 2016, TC). Chadburn et al. (2015) used functions of



temperature, moisture, light and snow to empirically determine the ground cover of moss to be used in the soil thermal module, but did not simulate the carbon cycle of mosses. Porada et al., (2016) modeled the productivity and expansion of moss/lichen, which then, combined with fire disturbance, determined the dynamic surface coverage of moss/lichen.

Druel et al. (2017, GMDD) implemented an explicit representation of the high-latitude vegetation types including shrubs, boreal grasses, and non-vascular plants that are missing in standard ORCHIDEE. Processes and parameters regarding the growth of these new PFTs were defined, and the main biogeochemical results were evaluated. At the moment, their work is in parallel with the recent developments in ORCHIDEE-MICT described in this study, but could be merged within ORCHIDEE-MICT in the next step.

To discuss these issues, we added a paragraph at the end of Section 10.2, Page 32 Line 15:

“Previous land surface modeling studies have shown the critical role of organic matter in soil thermodynamics in permafrost regions (e.g. Lawrence et al., 2008; Chadburn et al., 2015), while different parameterizations of such effects are implemented in different models. Most of the recent models, like CLM (Lawrence et al., 2008), JULES (Chadburn et al., 2015), ISBA (Decharme et al., 2016), and ORCHIDEE-MICT in this study, assume weighted combinations of organic soil and mineral soil in the calculation of soil physical parameters for each soil layer in the model. This structure is more flexible than a fixed thickness of organic layer or moss layer as the implementation in JSBACH (Ekici et al., 2014), since the former could approximate the latter by assuming 100% organic soil above the prescribed thickness. Note that for the insulating effect of moss/lichen layer, the same values of thermal properties to that of the organic soil are usually used in recent models (Chadburn et al., 2015; Porada et al., 2016). In this study, however, we did not apply a fixed moss layer in the thermal module for the regional simulations, due to the lack of a gridded map for moss/lichen ground covers especially on the boreal forest floor, and to the lack of a representation for dynamic moss/lichen coverage as in JULES (Chadburn et al., 2015) and JSBACH (Porada et al., 2016). This could partly explain the regionally overestimated ALT compared to the empirical map for Yakutia (Fig. S5). An explicit representation for non-vascular plants in ORCHIDEE (Druel et al., 2017) has been worked in parallel with this study at the moment, but would be incorporated in ORCHIDEE-MICT in the future developments.”

**(4) I agree with the authors that the new processes implemented in ORCHIDEE-MICT should improve the model performance at high latitudes. However, I did not find in the manuscript any comparison with the previous ORCHIDEE version in this regard. Could you please show with 2 or 3 examples how ORCHIDEE-MICT represents an improvement over the previous version, e.g. with respect to simulation of runoff, snow patterns, carbon fluxes etc.?**

ORCHIDEE-MICT is a new branch building on several separated former works with important processes for high latitudes for both the physical processes (e.g. Gouttevin et al., 2012a; Wang et al., 2013) and the carbon cycle (e.g. Koven et al., 2009). This paper is thus like the birth certificate of this new branch, demonstrating the effectiveness of all these combined processes in reproducing important observed variables. A comparison with the TRUNK, which does not yet incorporate all of these processes like the soil carbon discretization, would be unfair. An outcome of this paper will indeed be the integration of these high-latitude processes in the standard TRUNK version.

### **Specific comments**

**p.3,l.4: Is the correct buildup of soil carbon pools during the spin-up the only important factor for the correct short-term (100yr) prediction of soil carbon fluxes? I would argue that accurate representation of decomposition is at least equally important. Please explain shortly in the discussion why you did not revise the decomposition scheme.**

We fully agree with the reviewer that a good representation of SOC is highly important to represent the carbon fluxes to the atmosphere. At some points we wondered how we could improve the scheme in ORCHIDEE-MICT, and from a short literature review (well summarized in Manzoni and Porporato, 2009, SBB, Wutzler and Reichstein 2008, BG or more recently in Luo et al. 2016 GBC) there is still no consensus in the soil science community, and the different approach with their own underlying assumptions can hardly impact the model outputs. The representation of SOC decomposition in land surface models is still under debate and within this debate we choose to be conservative and keep the scheme used for decades now based on CENTURY (Parton et al., 1988). Nevertheless, some actions are ongoing in our group to improve the SOC decomposition scheme including the impact of priming, dissolved organic carbon, etc.

We add these sentences in the discussion on Page 33 Line 18: "Other studies (Koven et al., 2013; Burke et al., 2017a) further limited the rate of decomposition of SOC at depth, to reproduce the lack of oxygen inhibiting decomposition. [It should be noted also that the decomposition scheme of SOC is still based on Parton et al., \(1988\) as classically done in land surface models \(Friedlingstein et al., 2006\). Different approaches were proposed in models focusing only on SOC decomposition \(Manzoni and Porporato, 2009; Wutzler and Reichstein, 2008\) based on different assumptions \(substrate driven, decomposer driven, etc.\), but no clear consensus emerged up to date to revise the SOC decomposition scheme in land surface models \(Luo et al., 2016\).](#)"

**p.4,l.3: If transpiration is calculated per PFT, some averaging has to take place in order to calculate the energy balance per grid cell; Please explain.**

A transpiration flux is calculated for each PFT in each tile (forest and short vegetation). In each tile, a weighted spatial average of the different PFTs is performed. Then, the model calculates a weighted spatial average across the soil tiles to obtain a total representative flux of the grid-cell.

**p.5,I.13: To what extent does soil water content fluctuate in 11th soil layer? If significant changes occur, these are transferred with unlimited speed through the whole soil column down to 38m. This may lead to unrealistic dynamics of thermal properties.**

The soil water content fluctuates in a variation range given by the hydrodynamic parameters of the different soils prescribed in the model. Thus it can vary from 0.034 to 0.460 m<sup>3</sup>/m<sup>3</sup> (residual and saturated water content for a silt soil respectively). This corresponds to a maximum variation range of 68 to 920 mm for the 2-meter depth soil of the model. Yet, the most important variations of water content occurs in the superficial layers of the soil column that are submitted to precipitation events and infiltration. The non-linear decrease of the hydraulic conductivity with soil water content leads to reasonable values of gravitational drainage at the bottom and thus cannot lead to unlimited speed through the whole soil column.

**p.9,I.14: Why do you assume that the residual soil moisture and the Van Genuchten coefficients are independent of soil carbon content? Does soil carbon have no effect on soil texture? Please explain shortly.**

Soil organic matter may indeed contribute to the variation of the Van Genuchten equation coefficients (except for  $\alpha$ ) (Wang et al., CLEAN-Soil Air Water, 2014). However, we chose as a first step to keep the coefficients unchanged, and the resulted relationship of field capacity/wilting point versus SOC (Fig. S2) could capture the first-order characters in the observations by Hudson (1994). Modifications of the Van Genuchten equation coefficients would require an in-depth sensitivity study of soil hydrology in the model. This could be considered for further developments of the model.

**p.24,I.10: It is suggested that low speed of infiltration is the reason for the underestimation of soil water content in the deep soil. In addition to the mentioned deficiencies in the representation of infiltration into frozen soils, I would like to know whether the sensitivity of deep soil moisture to the parameterization of soil hydraulic conductivity was tested? I think the importance of the soil water deficit should be pointed out a bit more in sect. 7, since it has far-reaching effects such as reduced transpiration, increased surface temperature, and reduced productivity.**

In our parameterization of SOM in the model, the soil hydraulic conductivity can be affected by SOM through the increase of the porosity. However, we didn't perform any sensitivity tests of deep soil moisture to the parameterization of soil hydraulic conductivity. We point out the importance of the soil water deficit at the beginning of the paragraph "In the root zone" of sub-section 7.3 "Soil moisture", Page 24 Line 16: "The soil water deficit is of primary importance during spring and summer in the high latitudes because of its potential impacts on the vegetation transpiration, leading to a surface temperature increase and a reduction in the productivity. Yet, a soil water comparison between GLEAM and ORCHIDEE-MICT is difficult because ..."

**p.26,l.30 Why is peak GPP overestimated? Please explain.**

The CO<sub>2</sub> fertilization seems indeed too important in ORCHIDEE. We have at least two leads to improve this behavior. Kuppel et al. (2012) used FLUXNET daily observations to optimize several photosynthetic parameters like the maximum carboxylation rate V<sub>cmax</sub>. Given all the modifications brought to the model since this first FLUXNET optimization, mainly regarding the physics, it is probably time to recalibrate these photosynthetic parameters to get a more realistic GPP. Second, plants grown under elevated CO<sub>2</sub> show a photosynthetic downregulation (Sellers et al., 1996 ; Bounoua et al., 1999 ; Bounoua et al., 2010). This downregulation calibration is coded in the model to rectify V<sub>cmax</sub> under different atmospheric CO<sub>2</sub> levels to empirically limit the CO<sub>2</sub> fertilization effect, but this option was not activated in these simulations.

- Kuppel, S., Peylin, P., Chevallier, F., Bacour, C., Maignan, F., and Richardson, A. D.: Constraining a global ecosystem model with multi-site eddy-covariance data, *Biogeosciences*, 9, 3757-3776, DOI 10.5194/bg-9-3757-2012, 2012.
- Sellers, P. J., Bounoua, L., Collatz, G. J., Randall, D. A., Dazlich, D. A., Los, S. O., Berry, J. A., Fung, I., Tucker, C. J., Field, C. B., and Jensen, T. G.: Comparison of radiative and physiological effects of doubled atmospheric co<sub>2</sub> on climate, *Science*, 271, 1402-1406, DOI 10.1126/science.271.5254.1402, 1996.
- Bounoua, L., Collatz, G. J., Sellers, P. J., Randall, D. A., Dazlich, D. A., Los, S. O., Berry, J. A., Fung, I., Tucker, C. J., Field, C. B., and Jensen, T. G.: Interactions between vegetation and climate: Radiative and physiological effects of doubled atmospheric co<sub>2</sub>, *Journal of Climate*, 12, 309-324, Doi 10.1175/1520-0442(1999)012<0309:lbgacr>2.0.Co;2, 1999.
- Bounoua, L., Hall, F. G., Sellers, P. J., Kumar, A., Collatz, G. J., Tucker, C. J., and Imhoff, M. L.: Quantifying the negative feedback of vegetation to greenhouse warming: A modeling approach, *Geophysical Research Letters*, 37, Artn L23701, Doi 10.1029/2010gl045338, 2010.

**p.27,l.4: NPP is underestimated (Fig 16) due to water stress and lack of nitrogen fertilisation. GPP, however, is overestimated, which should lead to an underestimation of CUE. However, CUE is overestimated, and this is explained with a lack of nutrient limitation ('too much' nutrients), which is inconsistent with the lack of nitrogen ('too little' nutrients) mentioned above. Please explain.**

GPP is overestimated at the regional scale largely for BONA, slightly for BOEU and is correct for BOAS (see Fig. 15), but GPP is underestimated over the Campioli sites (see Fig. 17), where the CUE is computed. We can thus infer that the Campioli database is not a representative sample of our domain, however this database offers the advantage of having collocated GPP and NPP observations, and thus access to realistic CUE estimates.

Nevertheless the reviewer is right, as we use the nutrients argument in opposite manners to explain a too low NPP and a too high CUE over the Campioli sites. We thus removed the following sentences in section 8.2.2 on Page 27 Line 6: "... or due to a lack of N-deposition combined with soil fertility effects in modeled NPP" and on Page 27 Lines 18-19 "This high CUE bias can be expected, as ORCHIDEE-MICT omits the effects of low nutrient availability on CUE (Vicca et al. 2012)". We now just stick to our water stress hypothesis, to explain these low NPP and GPP, resulting in a high CUE.

**p.28,l.30: The respiration could also originate from a moss/lichen layer which may show some activity at low temperatures and under snow.**

Yes. We added these sentences in the text:

- at Line 27 P.28: "... (ii) insufficient snow insulation of soils (See Fig. 7); and (iii) the lack of the carbon cycle of mosses and lichens which could have respiration under winter low temperatures (Atanasiu, 1971)."
- at Lines 29-31 p.28: "...improve the snow insulating, prescribe an organic layer of insulating topsoil (e.g. mosses, O-horizons observed in boreal forests, see O'Donnell et al., 2011) into the thermal module, or explicitly represent the moss/lichen plants including their carbon cycle and physical effects (Porada et al., 2016; Druel et al., 2017)."

**p.29,l.6: Biomass is significantly overestimated (see also Fig S5), and, in my opinion, the difference between climate forcing data sets cannot explain this easily: Precipitation in CRUNCEP is lower than in GSWP3, but biomass is higher, which seems counterintuitive. Could you please explain?**

The higher biomass by CRUNCEP than by GSWP3 could be partly explained by the higher GPP (Fig. 15). Indeed, precipitation in CRUNCEP is lower than in GSWP3; however, the specific air humidity in CRUNCEP during summer is higher than in GSWP3 (Fig. S14). A lower air humidity leads to a higher atmospheric vapour pressure deficit (VPD). ORCHIDEE being VPD-dependent, the stomatal conductance of the vegetation decreases to prevent excessive water loss when VPD is high, and consequently reduces photosynthetic rates. To address it, we add the following paragraph on Page 26, below Line 31: “Interestingly, comparing GPP forced by the two climate datasets shows higher values by CRUNCEP than by GSWP3 (Fig. 14), despite a generally lower precipitation in CRUNCEP (Fig. S17). This could be explained by the higher specific air humidity during summer in CRUNCEP than in GSWP3 (Fig. S14). A low air humidity increases the atmospheric vapor pressure deficit (VPD) and the leaf to air vapour pressure difference; plants then partially close the stomata to constrain a potentially fast transpiration (Oren et al., 1999; McAdam et al., 2015), which leads to reduced photosynthetic rate. The photosynthesis module in ORCHIDEE largely follows Yin and Struik (2009) in which stomata conductance decreases with an increasing VPD, thus is able to simulate a lower GPP under dry air conditions. A recent study (Novick et al., 2016) showed that, between the two factors that impact plant water stress, i.e. soil moisture supply and atmospheric demand for water (reflected by VPD), the latter limits evapotranspiration to a greater extent than the former in relatively wet forested ecosystems. In spite of its importance, the effect of VPD on vegetation productivity has been far less studied than soil water availability (Konings et al., 2017), warranting further investigations in both observations and land surface models.”

Apart from a higher GPP, the higher biomass by CRUNCEP could also be because of the allocation scheme in ORCHIDEE. As detailed in Krinner et al., (2005), if LAI is above a PFT-specific maximum value (about 4 for boreal tree PFTs), carbon will not be allocated to leaves but to the sapwood which slowly converts to heartwood. Therefore, a higher LAI simulated by CRUNCEP leads to more carbon allocation to the wood for some areas, the turnover time of which is much longer than leaves. To address it, the paragraph on Page 29, Lines 13-14 is revised as: “Both model output and observation data are subject to the spatial uncertainties introduced by the use of satellite-derived land cover maps. We thus used the forest cover map as prescribed in the model for both data sets, and calculated latitudinal averages to compare with model results (Fig. S8). GSWP3-forced model output agrees well with observations averaged over the whole study region, while CRUNCEP-forced output overestimates biomass at all latitudes (Fig. S8a). For the sub-regions, the overestimation of biomass in BONA is consistent with that of GPP (Fig. 14), while biomass is more overestimated in BOEU compared with GPP, probably because of the lack of forest management and forest age structure for Europe. Comparing the two model results, CRUNCEP-forced biomass is much higher than GSWP3-forced biomass, which cannot solely be explained by the higher GPP by CRUNCEP (Fig. 14), indicating a bias in the allocation scheme in ORCHIDEE. As detailed in Krinner et al., (2005), the photosynthates are partitioned into leaves,

roots, sapwood, and carbohydrate reserve, dependent on soil moisture etc. If LAI is above a PFT-specific maximum value, carbon will not be allocated to leaves but to the sapwood, which slowly converts to heartwood. Therefore, a higher LAI forced by CRUNCEP leads to more carbon allocation to the wood for some areas, the turnover time of which is much longer than leaves. A new allocation scheme based on the pipe model was implemented in another branch of ORCHIDEE (Naudts et al., 2015), which provides a physiologically meaningful relationship between foliage, roots, and wood. This would be incorporated in ORCHIDEE-MICT in the future developments. Simulated total forest biomass for the whole domain is 95 PgC under GSWP3 forcing (165 PgC under CRUNCEP), close to estimates from forest inventory data in Pan et al. (2011) of 92.1 PgC. Somewhat lower estimates are derived by Avitabile et al. (2016) and Thurner et al. (2014), of 73 and 84 PgC, respectively.”

**p.32,I.23: I think biomass compares well to observations only for GSWP3 forcing (Fig 21).**

Yes, you are right. This is now corrected in the text: “ORCHIDEE-MICT performs generally well for biomass with GSWP3 forcing, including the latitudinal profile, ...”

**p.33,I.18: The model already has a cold bias, so even lower soil temperatures would be required for a more realistic (higher) soil carbon content. Why do you not mention explicit simulation of cryoturbation as a potential missing process?**

Actually, we take into account the cryoturbation effect in ORCHIDEE-MICT using a diffusion term (see Eq. 2 in Section 4.1).

**supplement: Please indicate if CRUNCEP is subtracted from GSWP3 or vice versa.**

This is added now in the title of the section of supplementary Figures and in the caption of Figure S9

---

# ORCHIDEE-MICT (~~revision 4126~~)(v8.4.1), a land surface model for the high-latitudes: model description and validation

Matthieu Guimberteau<sup>1\*</sup>, Dan Zhu<sup>1\*</sup>, Fabienne Maignan<sup>1</sup>, Ye Huang<sup>1</sup>, Chao Yue<sup>1</sup>, Sarah Dantec-Nédélec<sup>1</sup>, Catherine Ottlé<sup>1</sup>, Albert Jornet-Puig<sup>1</sup>, Ana Bastos<sup>1</sup>, Pierre Laurent<sup>1</sup>, Daniel Goll<sup>1</sup>, Simon Bowring<sup>1</sup>, Jinfeng Chang<sup>2</sup>, Bertrand Guenet<sup>1</sup>, Marwa Tifafi<sup>1</sup>, Shushi Peng<sup>1,3</sup>, Gerhard Krinner<sup>4</sup>, Agnès Ducharne<sup>5</sup>, Fuxing Wang<sup>6</sup>, Tao Wang<sup>7,8</sup>, Xuhui Wang<sup>1,9</sup>, Yilong Wang<sup>1</sup>, Zun Yin<sup>1</sup>, Ronny Lauerwald<sup>10,1,11</sup>, Emilie Joetzjer<sup>1,12</sup>, Chunjing Qiu<sup>1</sup>, Hyungjun Kim<sup>13</sup> and Philippe Ciais<sup>1</sup>

~~\* These authors contributed equally to this work~~

<sup>1</sup> Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA - CNRS - UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

<sup>2</sup> Sorbonne Universités (UPMC), CNRS-IRD-MNHN, LOCEAN/IPSL, 4 place Jussieu, 75005 Paris, France

<sup>3</sup> Sino-French Institute for Earth System Science, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

<sup>4</sup> CNRS, Univ. Grenoble Alpes, Institut des Géosciences de l'Environnement (IGE), 38000 Grenoble, France

<sup>5</sup> UMR 7619 METIS, Sorbonne Universités, UPMC, CNRS, EPHE, 4 place Jussieu, 75005 Paris, France

<sup>6</sup> Laboratoire de Météorologie Dynamique, Ecole Polytechnique, F 91128 Palaiseau, France

<sup>7</sup> Key Laboratory of Alpine Ecology and Biodiversity, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100085, China

<sup>8</sup> CAS Center for Excellence in Tibetan Plateau Earth Sciences, Chinese Academy of Sciences, Beijing 100085, China

<sup>9</sup> Laboratoire de Météorologie Dynamique, Université Pierre et Marie Curie, 75005 Paris, France

<sup>10</sup> Université Libre de Bruxelles, Belgium

<sup>11</sup> University of Exeter, Exeter, United Kingdom

<sup>12</sup> CNRS, Université Paul Sabatier, ENFA; UMR5174 EDB (Laboratoire Evolution et Diversité Biologique), 118 route de Narbonne, Toulouse F-31062, France

<sup>13</sup> Institute of Industrial Science, The University of Tokyo, Tokyo, Japan

*\* These authors contributed equally to this work*

~~matthieu.guimberteau@lsce.ipsl.fr~~ Correspondence to: [matthieu.guimberteau@lsce.ipsl.fr](mailto:matthieu.guimberteau@lsce.ipsl.fr) and [dan.zhu@lsce.ipsl.fr](mailto:dan.zhu@lsce.ipsl.fr)

**Abstract.** The high-latitude regions of the northern hemisphere are a nexus for the interaction between land surface physical properties and their exchange of carbon and energy with the atmosphere. At these latitudes, two carbon pools of planetary significance – those of the permanently frozen soils (permafrost), and of the great expanse of boreal forest – are vulnerable to destabilization in the face of currently observed climatic warming, the speed and intensity of which are expected to in-



crease with time. Improved projections of future Arctic and boreal ecosystem transformation require improved land surface models that integrate processes specific to these cold biomes. To this end, this study lays out relevant new parameterizations in the ORCHIDEE-MICT land surface model. These describe the interactions between soil carbon, soil temperature and hydrology, and their resulting feedbacks on water and CO<sub>2</sub> fluxes, in addition to a recently-developed fire module. Outputs from ORCHIDEE-MICT, when forced by two climate input data sets, are extensively evaluated against: (i) temperature gradients between the atmosphere and deep soils; (ii) the hydrological components comprising the water balance of the largest high-latitude basins, and (iii) CO<sub>2</sub> flux and carbon stock observations. The model performance is good with respect to empirical data, despite a simulated excessive plant water stress and a positive land surface temperature bias. In addition, acute model sensitivity to the choice of input forcing data suggests that the calibration of model parameters is strongly forcing-dependent. Overall, we suggest that this new model design is at the forefront of current efforts to reliably estimate future perturbations to the high-latitude terrestrial environment.

## Contents

	<b>1 Introduction</b>	<b>3</b>
	<b>2 ORCHIDEE model overview</b>	<b>4</b>
15	<b>3 High-latitude processes in the initial ORCHIDEE version</b>	<b>5</b>
	<b>4 New processes and parameterizations</b>	<b>7</b>
	<b>5 Simulation protocol, forcing and evaluation data sets</b>	<b>11</b>
	<b>6 Evaluation of the atmosphere-snow-soil continuum</b>	<b>21</b>
	<b>7 Evaluation of large-scale water storage and fluxes</b>	<b>24</b>
20	<b>8 Evaluation of the leaf area index, gross and net CO<sub>2</sub> fluxes</b>	<b>27</b>
	<b>9 Evaluation of carbon stocks</b>	<b>31</b>
	<b>10 Discussion</b>	<b>32</b>
	<b>11 Conclusions</b>	<b>37</b>
	<b>12 Appendix</b>	<b>38</b>

## 1 Introduction

In the high latitudes, the complex coupling between soil thermal and hydraulic processes, snowpack properties, and plant and soil carbon pools is of great importance. Snow accumulation and freezing of soil water lead to a net storage of water from October to April. Through the processes of snow melt and the onset of soil thaw in spring, water is made available for plant uptake and growth. Simultaneously however, much of this is ‘lost’ as runoff to rivers, causing peak discharge rates in May-June (Yang et al., 2003) and the flooding of large flatland areas from May to September (Papa et al., 2008; Biancamaria et al., 2009). In summertime, the peak in incident solar radiation causes a temperature maximum that increases water evaporative demand on the land surface. Many boreal and Arctic regions thus have a negative water balance in summer (Schulze et al., 1999), which may impose powerful constraints on plant growth. Siberian and Canadian boreal forests have thus been shown to experience water stress, with ratios of surface sensible to latent heat flux of up to ~2 (Jarvis et al., 1997; Baldocchi et al., 1997; Schulze et al., 1999) causing further heating of the near-surface atmosphere.

These large seasonal shifts of the high-latitude water balance – how water input from precipitation is shared between changes of water storage in snow, ice and soil moisture, and balanced against losses from evapotranspiration, sublimation and river discharge – can now be better assessed and evaluated using state-of-the-art observation datasets. In addition, in terms of realistic process representation, land surface models (LSMs) focusing on high-latitude phenomena require the inclusion of the following non-exhaustive series of pivotal hydrological and biogeochemical interactions:

1- A representation of permafrost physics and seasonal freeze-thaw cycles, which determine the soil hydrologic and thermal budgets and the volume and timing of lateral water flows to rivers.

2- The impact of winter snow acting as an insulating ‘barrier’ between soils and overlying air from Autumn to early Spring. These have subsequent effects on soil temperature and water content, feeding back onto snow thickness itself.

3- The seasonal mediation of plant water availability via snowmelt water, transpiration losses and the depth of the permafrost table (active layer thickness), which in turn determine the availability of the lateral water flows that feed rivers in the warmer months.

4- The limitations on plant productivity and biomass due to acute climatic conditions in high-latitude regions. These primarily involve biotically prohibitive cold temperatures from Autumn to late Spring, low soil moisture in dry-summer regions, and fire events, caused by hot and dry conditions.

5- The buildup of large soil carbon stocks under cold conditions through the slow burial of organic matter in the permafrost via cryoturbation and sedimentary soil formation processes (e.g. Hugelius et al., 2013; Tarnocai et al., 2009).

6- Feedbacks between high soil carbon concentrations and profiles of soil temperature, water and permafrost carbon content (e.g. Lawrence and Slater, 2008; Decharme et al., 2016).

We represent the above processes in an updated version of the ORCHIDEE LSM (ORganizing Carbon and Hydrology in Dynamic EcosystEms), known as ORCHIDEE-MICT (aMeliorated Interactions between Carbon and Temperature), which we describe in this study. Since the comprehensive description of the model ORCHIDEE by Krinner et al. (2005), the model went through major modifications and improvements; we present here the major ones linked to high-latitude processes. ORCHIDEE-

MICT is evaluated over the last two to three decades (depending on the variable) against empirically generated datasets. Against these, we are able to evaluate model performance regarding the distribution of permafrost and the effect of snow on soil thermics (mechanisms 1 and 2); the different components of the water cycle over a wide range of high-latitude basins (mechanism 3); plant primary productivity as constrained by high-latitude conditions (mechanisms 3 and 4), and replication of soil carbon stocks and feedback dynamics (mechanisms 5 and 6).

## 2 ORCHIDEE model overview

The starting point for our updated land surface model (ORCHIDEE-MICT) is ORCHIDEE-TRUNK revision 3976. As detailed in Section 3, its description of soil temperature and vertical water transport dynamics is based on coupled diffusion equations with identical vertical discretization (Wang et al., 2016a), and includes soil freezing, its effect on water infiltration, and phase change-induced heat sources and sinks in the soil column (Gouttevin et al., 2012b). The snow model described by Wang et al. (2013) is incorporated in this version, where snow is discretized into three layers of variable thickness, conductivity and density, accounting for snow liquid water content (Boone and Etchevers, 2001). In terms of large-scale hydrology, a river routing scheme including floodplains and their dynamics (d'Orgeval et al., 2008; Guimberteau et al., 2012) is coupled to simulated grid-cell runoff (Section 3.2), permitting the calculation of 'natural' river discharge (i.e. in the absence of dams or human water withdrawals).

The carbon cycle model includes half-hourly photosynthesis (GPP), daily allocation of GPP assimilates to autotrophic respiration and 8 plant biomass pools (foliage, roots, above/below-ground sapwood and heartwood, fruits and carbon reserves), and prognostic phenology (Botta et al., 2000). These pools are characterized by different turnover times, mortality rates and subsequent litter and soil carbon decomposition rates. Litter carbon is funneled between structural and metabolic fractions, and soil carbon between active, slow and passive pools, following Parton et al. (1987).

The model divides vegetation into 13 plant functional types (PFTs). Each PFT follows the same suite of equations but with PFT-specific parameter values and phenology functions (Krinner et al., 2005). PFT fractions are assigned to three soil tiles corresponding to bare soil, short vegetation (grass and crop PFTs) and forests (all tree PFTs). The soil moisture budget of each soil tile is calculated separately, but different PFTs in the same soil tile interact as they share the same soil moisture source. While transpiration is calculated separately for each PFT, and soil moisture for each soil tile, the energy budget of a grid cell with multiple PFTs is calculated using the area-weighted average of those PFTs. This in turn defines mean grid-cell land surface temperature, giving the upper boundary condition for the vertically discretized soil thermal module.

Temperature, water and carbon interactions described in ORCHIDEE revision 3976 are summarized in Fig. 1 by the black arrows. Air temperature and humidity impact phenology, photosynthesis, autotrophic respiration and the water and heat fluxes comprising the surface energy budget. Soil moisture in the root zone modulates photosynthesis and transpiration, which depends on wilting point and field capacity. In ORCHIDEE revision 3976, while soil carbon decomposition is impacted by soil water and temperature, soil carbon stocks themselves exert no feedback on the soil physical state.

The key model developments presented here in ORCHIDEE-MICT ([revision 4126v8.4.1](#)) thus include the feedback effects of soil organic carbon (SOC) concentration on both soil thermic and soil water dynamics (Fig. 1, red arrows). Because these SOC-affected soil physics alter the above and below-ground components of the carbon cycle, as well as plant transpiration via hydraulic stress, we can expect complex indirect effects on the energy, water and carbon budgets (Fig. 1). Note that several other updates were implemented in ORCHIDEE-TRUNK (revision 3976) and passed to ORCHIDEE-MICT ([revision 4126v8.4.1](#)), including a revised background albedo based on satellite observations, and updates of photosynthesis scheme. These will be described in an upcoming paper for the ORCHIDEE-TRUNK (version close to revision 3976) that will be used for CMIP6 exercise. In the following, we describe the parameterizations that define ORCHIDEE-MICT ([revision 4126v8.4.1](#)). ~~Section 3 briefly summarizes snow and soil freeze developments (Gouttevin et al., 2012a; Wang et al., 2013) that are incorporated already in ORCHIDEE-TRUNK revision 3976. In Section 4, we focus on new features of ORCHIDEE-MICT compared to revision 3976, including SOC vertical discretization and processes leading to SOC burial in frozen horizons (Section 4.1), and functions describing how the SOC content in each soil layer alters soil thermal and hydraulic properties (Section 4.2). The simulation protocol, forcing data and different observations used to evaluate ORCHIDEE-MICT are described in Section 5. Model outputs are evaluated against multiple observations for grid-based simulations forced by two climate datasets, CRUNCEP v7 and Global Soil Wetness Project Phase 3 (GSWP3) v0, at 1° resolution over the Northern Hemisphere over periods within the range 1960–2007. In the result sections, Section 6 evaluates snow depth and soil temperature against measurements from in-situ stations, and land skin temperature against satellite-derived dataset. Section 7 evaluates modeled components of the water budget and their sum, using remote-sensing based observations of total land water storage changes, snow water equivalent, and evapotranspiration, combined with in-situ river discharge data, to close the water budget at the scale of large northern hemisphere catchments. Sections 8 and 9 evaluate carbon fluxes and stocks, using observation-based gridded GPP, fire emissions, forest biomass and SOC, together with site-based measurements of NPP (net primary production) and GPP, and regional net CO<sub>2</sub> fluxes from atmospheric inversions. In the discussion (Section 10), we address some of the critical issues in the model performances for processes related to seasonal changes in water budgets, thermal gradients from atmosphere to soils, and carbon budgets.~~

## 25 3 High-latitude processes in the initial ORCHIDEE version

### 3.1 Soil freezing and snow processes

The soil freezing scheme developed by Gouttevin et al. (2012a) describes phase changes of soil water, simulating the latent heat exchanges involved in the freezing and melting of soil water, and subsequent changes in thermal and hydrological ground properties. Soil heat conductivity and heat capacity are dependent on soil ice content. The hydraulic conductivity of the soil is parameterized according to its liquid water content and decreases with the frozen soil fraction. Heat transfer through the

soil column is represented by a one-dimensional heat conduction equation, with latent heat acting as a source or sink term (Gouttevin et al., 2012a), in the following function:

$$c \frac{\partial T}{\partial t} = \frac{\partial}{\partial z} \left( \lambda \frac{\partial T}{\partial z} \right) + \rho_{ice} L \frac{\partial \theta_{ice}}{\partial t} \quad (1)$$

where  $c$  is volumetric soil heat capacity ( $\text{JK}^{-1} \text{m}^{-3}$ );  $T$  is soil temperature (K);  $\lambda$  is soil thermal conductivity ( $\text{J m}^{-1} \text{s}^{-1} \text{K}^{-1}$ );  $\rho_{ice}$  is ice density ( $\text{kg m}^{-3}$ );  $L$  is latent heat of fusion ( $\text{J kg}^{-1}$ );  $\theta_{ice}$  is volumetric ice content ( $\text{m}^3 \text{m}^{-3}$ );  $t$  is time (s) and  $z$  is depth (m). In ORCHIDEE-MICT, this equation is discretized on the 32 vertical layers of the model with a total soil depth of 38 m (Fig. S1). Note that the soil hydrology has only 11 layers up to 2 m, so the volumetric contents of water and ice below 2 m take the values of the bottom layer (i.e. the 11<sup>th</sup> layer).

Snowpack is represented by a 3-layer snow model of intermediate complexity, as described in Wang et al. (2013). This scheme was implemented to resolve the energy and water budgets inside the snowpack, accounting for thawing and refreezing of liquid water. The snow model produces prognostic snow temperature, density and SWE for the 3 snow layers. Modifications were recently implemented to represent snowpack sub-grid scale variability. A snow cover fraction over the grid cell was introduced as a function of SWE. This was used for improving albedo and surface temperature estimates. Although this snow cover fraction was calculated for glacier and vegetated-surface areas separately, it is not dependent on the vegetation cover. Additional modifications were implemented (uniformization of the energy budget calculation, update of the snow-covered vegetation albedo...) and will be described in the upcoming CMIP6 ORCHIDEE paper as mentioned in Section 2.

### 3.2 Soil hydrology and river routing

ORCHIDEE simulates soil water fluxes and storage through a multi-layer soil hydrology scheme described by de Rosnay et al. (2000, 2002) and Campoy et al. (2013). Soil moisture is redistributed in the column by solving the Richards equation for vertical unsaturated flow under the effect of root uptake. The hydraulic conductivity and diffusivity depend on soil moisture, following the Mualem-van Genuchten model (Mualem, 1976; Van Genuchten, 1980), and using parameters defined by Carsel and Parrish (1988). These variables depend on the dominant soil texture in each grid-cell, based on the 12 USDA texture classes provided at the  $0.08^\circ$  resolution from Reynolds et al. (2000). For frozen soils, the decrease of the hydraulic conductivity (Gouttevin et al., 2012a) reduces infiltration into the soil and drainage, and increases surface runoff. The 2m soil column is divided into 11 layers, with layer thickness increasing geometrically with depth (Fig. S1). The saturated hydraulic conductivity is modified according to the scheme in d'Orgeval et al. (2008). This decreases exponentially below a top-30 centimeter depth boundary to account for increased soil compaction, as suggested by Beven and Germann (1982), and increases above that boundary towards the soil surface due to the enhanced infiltration capacity afforded by vegetative roots, whose presence increases soil porosity in the root zone (Beven, 1984). The canopy throughfall rate and soil hydraulic conductivity govern the partitioning between surface runoff and soil infiltration. This partitioning involves a time-splitting procedure inspired by Green and Ampt (1911),

describing the propagation of the wetting front. The second physical factor contributing to total runoff is free gravitational drainage at the bottom of the soil.

The runoff routing module (Polcher, 2003; Ngo-Duc et al., 2005; Guimberteau et al., 2012) aggregates surface runoff and drainage produced at a 30-minute time step to calculate daily flow between grid cells and discharge to the ocean. Grid cells are subdivided into basins in which water is transferred through a series of linear reservoirs along the drainage network, derived from a 0.5 degree resolution data set (Vörösmarty et al., 2000; Oki et al., 1999). In a given basin, a ‘slow’ reservoir collects drainage water, while a ‘fast’ reservoir collects surface runoff, each with different linear response timescales. Corresponding outflows are transferred to the stream reservoir of the downstream basin. The process is fully detailed in Guimberteau et al. (2012).

The routing scheme includes a parameterization of floodplains (d’Orgeval et al., 2008; Guimberteau et al., 2012), the maximum extent of which is prescribed by the GLWD (Global Lakes and Wetlands Database) map (Lehner and Döll, 2004). In grid-cells with flooded areas, river discharge from upstream basins is diverted to a floodplain reservoir, which then feeds a delayed return flow back to the stream reservoir of the basin.

#### 4 New processes and parameterizations

##### 4.1 Soil carbon discretization

In ORCHIDEE-MICT, the three soil carbon pools (active, slow and passive) share a common 32-layer discretization scheme as that of soil temperature, to a maximum depth of 38 m. Carbon inflows to the soil pools from decomposed litter are partitioned along this depth using an exponential function that corresponds to the prescribed PFT root profile. Decomposition of soil carbon is calculated at each layer as a function of soil temperature, moisture, and texture (Koven et al., 2009; Zhu et al., 2016). Vertical mixing of soil carbon due to cryoturbation (mixing of soil layers induced by repeated freeze-thaw cycles) and bioturbation are accounted for by adding a diffusion term in the soil carbon equation:

$$\frac{\partial C_i(z,t)}{\partial t} = I_i(z,t) - g_i(z,t)C_i(z,t) + D \frac{\partial^2 C_i(z,t)}{\partial z^2} \quad (2)$$

where  $C_i(z,t)$  is carbon content of pool  $i$  at depth  $z$  and time  $t$  ( $\text{gCm}^{-3}$ );  $I_i(z,t)$  is carbon input ( $\text{gCm}^{-3} \text{d}^{-1}$ );  $g_i(z,t)$  is decomposition rate ( $\text{d}^{-1}$ );  $D$  is diffusive mixing rate, set as  $10^{-3} \text{m}^2 \text{yr}^{-1}$  through the active layer and decreases linearly to zero at 3 m in permafrost regions, to represent cryoturbative mixing (Koven et al., 2009), and set as  $10^{-4} \text{m}^2 \text{yr}^{-1}$  above 2 m in non-permafrost regions to represent bioturbation (Koven et al., 2013).

##### 4.2 SOM-dependent soil thermal and hydraulic parameters

Soil organic matter (SOM) significantly modifies soil thermal and hydraulic properties. SOM lowers thermal conductivity and increases heat capacity (e.g. Lawrence and Slater, 2008; Decharme et al., 2016), and increases soil porosity, which in

turn increase saturated hydraulic conductivity and available water capacity (e.g. Hudson, 1994; Morris et al., 2015). As a consequence, the presence of SOM modulates heat transfer from the surface through the soil column, typically leading to cooler soil temperature during summer. SOM-effected increases in soil water holding capacity also enhances plant available water and thus primary productivity (Krull et al., 2004) and transpiration. SOM impacts on soil thermics and hydraulics have previously been parameterized in the global LSMs CLM (Lawrence and Slater, 2008), JULES (Chadburn et al., 2015b) and ISBA (Decharme et al., 2016). In ORCHIDEE, SOM thermal insulation was previously investigated by Koven et al. (2009) but its parameterization was imbedded in a prior model version which employed bucket-type soil hydrology. This however, is not applicable to ORCHIDEE-MICT, which uses a new vertically discretized hydrology scheme and its coupling with the thermal module. In addition, the Koven et al. (2009) study did not include SOM effects on soil hydraulic properties, which are addressed in ORCHIDEE-MICT and described in detail below.

### Thermal conductivity and heat capacity

By default, soil thermal conductivity and heat capacity in ORCHIDEE are calculated in each soil layer as empirical functions of the 12 USDA soil texture classifications (see Table S1) and soil water and ice contents, following Wang et al. (2016a):

$$\lambda_i = Ke_i \lambda_{i,sat} + (1 - Ke_i) \lambda_{i,dry} \quad (3)$$

with:

$$\lambda_{i,sat} = \lambda_{i,solid}^{(1-\theta_{i,sat})} \lambda_{liq}^{\left(\frac{\theta_{i,sat} \theta_{i,liq}}{\theta_{i,liq} + \theta_{i,ice}}\right)} \lambda_{ice}^{\left(\frac{\theta_{i,sat} \theta_{i,ice}}{\theta_{i,liq} + \theta_{i,ice}}\right)} \quad (4)$$

$$c_i = c_{i,dry} + \theta_{i,liq} c_{liq} + \theta_{i,ice} c_{ice} \quad (5)$$

where  $\lambda_{i,sat}$  and  $\lambda_{i,dry}$  are saturated and dry thermal conductivities for layer  $i$ ;  $\lambda_{liq}$  and  $\lambda_{ice}$  are thermal conductivities of liquid water and ice, equaling to 0.57 and 2.2 respectively ( $W m^{-1} K^{-1}$ );  $\lambda_{i,solid}$  is thermal conductivity of soil solids (see Table S1);  $c_{liq}$  and  $c_{ice}$  are heat capacities of liquid water and ice, equaling to  $4.18 \cdot 10^6$  and  $2.11 \cdot 10^6$  respectively ( $JK^{-1}m^{-3}$ );  $c_{dry}$  is dry soil heat capacity depending on soil texture;  $\theta_{i,sat}$  is volumetric moisture content at saturation (porosity), and it varies with soil textures;  $\theta_{i,liq}$  and  $\theta_{i,ice}$  are prognostic volumetric liquid water and ice contents ( $m^3 m^{-3}$ ) that are computed by the soil hydrology model;  $Ke_i$  is the Kersten number given by:

For unfrozen soils:

$$Ke_i = \begin{cases} \log_{10}(S_r) + 1 \\ 0.7 \log_{10}(S_r) + 1 \\ 0 \end{cases} \quad \text{if } \begin{cases} S_r > 0.1 \\ 0.05 < S_r \leq 0.1 \\ S_r \leq 0.05 \end{cases} \quad (6)$$

with:

$$S_r = \frac{\theta_i}{\theta_{i,sat}} \quad (7)$$

5 For frozen soils:

$$Ke_i = S_r \quad (8)$$

where  $S_r$  is the degree of saturation.

To account for the impacts of organic carbon on soil thermal properties in ORCHIDEE-MICT, we follow Lawrence and Slater (2008) in assuming that soil physical properties are weighted averages of mineral soil (as the default values in standard ORCHIDEE) and pure organic soil, with the organic soil fraction  $f_{i,soc}$  calculated as:

$$f_{i,soc} = \min \left( 1, \frac{\rho_{i,soc}}{\rho_{soc,max}} \right) \quad (9)$$

where  $\rho_{i,soc}$  is the carbon content for layer  $i$  ( $\text{kgC m}^{-3}$ ), derived from observation-based soil organic carbon map from NCSCD (Hugelius et al., 2013) in permafrost regions and from HWSD (FAO, 2012) in non-permafrost regions, after linear vertical interpolation from their original soil horizons to fit ORCHIDEE-MICT vertical layers;  $\rho_{soc,max}$  equals to  $130 \text{ kgC m}^{-3}$ , a typical soil carbon density of peat (Lawrence and Slater, 2008). Therefore, the red-colored parameters in Eqs. 3-7 are calculated as:

$$P_i = (1 - f_{i,soc}) P_{\text{mineral}} + f_{i,soc} P_{\text{soc}} \quad (10)$$

where  $P_i$  represents different properties  $\lambda_{i,dry}$ ,  $\lambda_{i,solid}$ ,  $c_{i,dry}$ , and  $\theta_{i,sat}$ . The values of  $P_{\text{mineral}}$  for each soil texture and  $P_{\text{soc}}$  are listed in Table S1. Note that here we followed Lawrence and Slater (2008) to use linear weighting organic and mineral soil properties, while in some other models like JULES (Chadburn et al., 2015a) and ISBA (Decharme et al., 2016), soil thermal



conductivities are calculated as geometric averages of organic and mineral soils, consistent with the treatment for soil water and ice (Eq. 4). The geometric averaging method increases the effect of the organic fraction compared to arithmetic averages, and would be tested in ORCHIDEE-MICT in future developments.

#### Available water capacity

- 5 Plant available water capacity, defined as the difference in the amount of water held by each soil layer between field capacity ( $\theta_{fc}$ ) and permanent wilting point ( $\theta_{wp}$ ), determines the capacity of the soil to store and supply water for plants and is therefore an important aspect of soil fertility (Hudson, 1994). For mineral soils in ORCHIDEE,  $\theta_{fc}$  and  $\theta_{wp}$  are derived from measurements of the soil matric potential at field capacity and wilting point, based on the soil water retention curve described by van Genuchten equation (Van Genuchten, 1980):

$$10 \quad \theta = \frac{(\theta_{sat} - \theta_r)}{[1 + (\alpha (-\Psi))^n]^{(1 - \frac{1}{n})}} + \theta_r \quad (11)$$

where  $\psi$  is soil matric potential (kPa), and  $\psi = -33$  kPa (or -10 kPa for the three sandy soils, see Table S2) corresponds to field capacity  $\theta_{fc}$ , while  $\psi = -1500$  kPa corresponds to wilting point  $\theta_{wp}$  for all textures;  $\theta_r$  is the residual volumetric water content ( $\text{m}^3 \text{m}^{-3}$ );  $\alpha$  and  $n$  are empirical fitting coefficients, with their values for different soil textures listed in Table S2.

- 15 SOC has been shown to significantly increase water retention (Rawls et al., 2003). To parameterize this SOM effect, we assume that  $\theta_r$  and the coefficients in Eq. 11 do not change with carbon content, while porosity  $\theta_{sat}$  increases with organic carbon (Eq. 10). Therefore, both  $\theta_{fc}$  and  $\theta_{wp}$  increase under higher carbon contents, but  $\theta_{fc}$  increases faster, resulting in a higher available water capacity (Fig. S2), consistent with the patterns observed in Hudson (1994).

#### 4.3 Reformulation of soil hydric stress above the permafrost table

- It is known that reduced soil moisture availability decreases the rate of photosynthesis, but the parameterization of this photosynthetic stress differs amongst models (Medlyn et al., 2015). ORCHIDEE-MICT lacks a fully mechanistic plant hydraulic structure that calculates plant internal water movement via constraints from water potential ( $\psi$ ) and conductance of roots, stems and leaves. Instead, a stress factor, which ranges from 0 to 1, is calculated based on the relative moisture content at each soil layer. This factor is applied to stomatal conductance and mesophyll conductance, as well as the maximum RuBisCO activity rate ( $V_{cmax}$ ) and maximum electron transport rate ( $J_{max}$ ), in order to account for experimentally observed effects of drought on stomatal and non-stomatal photosynthetic limitation (Zhou et al., 2014). The stress factor ( $\gamma$ ) of water limitation is calculated as:

$$25 \quad \gamma_i = \frac{\theta_i - \theta_{wp}}{\theta_{wp} + \rho (\theta_{fc} - \theta_{wp})} \quad (12)$$

$$\gamma = \sum_{i=1}^{11} \gamma_i w_i \quad (13)$$

where  $\gamma_i$  is relative moisture content at each soil layer  $i$ , bounded between 0 and 1;  $\rho$  represents the fraction above which photosynthesis rate is not limited by soil moisture, and is set at 0.8;  $w_i$  is the weighting factor for each layer.

In the initial version of ORCHIDEE, the profile of  $w_i$  was assumed to be constant over time, although differed between tree and grass PFTs, with the highest value at 1.5 m depth for trees and 0.37 m depth for grasses. We considered this description inappropriate for the high latitudes, and in particular for permafrost regions, where trees develop shallow and lateral roots above the permanently frozen layer (Kajimoto et al., 2003). Thus, in ORCHIDEE-MICT,  $w_i$  is modified to be a dynamic profile which optimizes plant water use, in a manner inspired by the representation given in Beer et al. (2007):

$$w_i = \frac{\gamma_i}{\sum_{i=1}^{11} \gamma_i} \quad (14)$$

where if layer  $i$  is below modeled active layer thickness,  $w_i$  is set to zero, and the remaining  $w$  are re-normalized to one.

#### 4.4 Fires

The prognostic fire module SPITFIRE (SPread and InTensity of FIRE), which has been previously integrated to and calibrated for a standard version of ORCHIDEE (Thonicke et al., 2010; Yue et al., 2014), was merged into ORCHIDEE-MICT. Ignitions were re-calibrated using the GFED4s data set (<http://www.globalfiredata.org/data.html>) by using region-specific scaling factors (see Table S3) and the exclusion of cropland fires to ensure that simulated mean annual burned area for 1997-2013 was equal to that of the GFED4s data set. Note that this method only calibrated for mean annual regional burned area, and that simulated latitudinal distributions and grid cell spatial patterns of burned area and fire carbon emissions, and their interannual and seasonal variabilities, could be still compared with observation-based data. Deforestation and peatland fires are not explicitly simulated, but as both fire types rely on suitable weather conditions to occur, which could be partly captured by SPITFIRE (Yue et al., 2015), model simulations are expected to partially include these fire types.

### 5 Simulation protocol, forcing and evaluation data sets

#### 5.1 Simulation protocol and forcings

##### 5.1.1 Simulation set-up

Two separate runs using different climate forcing input data – CRUNCEP v7 (hereafter CRUNCEP) and GSWP3 – were performed with ORCHIDEE-MICT for the terrestrial Northern Hemisphere ( $> 30^\circ\text{N}$ ) at  $1^\circ$  spatial resolution. Both sets of runs

encompass the 20<sup>th</sup> and beginning of 21<sup>st</sup> century, and were preceded by separate spin-ups for each climate dataset, forced by fixed pre-industrial conditions of atmospheric CO<sub>2</sub> (286 ppm) and vegetation map. The dynamic vegetation model is deactivated throughout both runs. In order to accumulate soil organic carbon in the model, which requires substantial computing time before reaching near-equilibrium in the presence of the slow mixing processes described in Section 4.1, the spin-up procedure comprised three steps: 1) The full ORCHIDEE-MICT model was forced by looped climate fields over the period 1960-1990 for 100 years to reach equilibria for soil temperature, soil moisture, vegetation productivity, soil carbon inputs from dead plants, etc. We used the 1960-1990 loop, instead of pre-industrial climate, to approximate the higher Holocene temperatures relative to the ‘pre-industrial’ period that have been reconstructed in Marcott et al. (2013); 2) A soil carbon sub-model was run for 20,000 years, forced by the outputs from the preceding step; 3) The full ORCHIDEE-MICT model was run for 100 years, forced by looped 1901-1920 climate data, to approach to the pre-industrial equilibrium for physical variables, carbon fluxes, and fast carbon pools. A final transient simulation from 1861 to 2007 (using 1901-1920 climate loop for the period 1861-1900 due to the lack of climate forcing before the 20<sup>th</sup> century) was then run from the last year of spin-up stage 3, forced by historical climate forcing and land cover maps, and rising CO<sub>2</sub> concentrations, as detailed below.

### 5.1.2 Atmospheric forcing datasets

The use of two different forcing datasets represents a first step in documenting atmospheric forcing based uncertainty in model output. Runoff has been shown for instance to be particularly affected by differences in precipitation from different datasets (Fekete et al., 2004; Biancamaria et al., 2009), and by the methods to partition total precipitation volumes between rainfall and snowfall during the cold season (Haddeland et al., 2011). The bias of meteorological driver also impacts the carbon budget (Zhao et al., 2012). A description of the two datasets used follows.

#### 20 GSWP3 v0

This 3-hourly 0.5° global forcing product (1901-2007) was developed for the third phase of GSWP3 (<http://hydro.iis.u-tokyo.ac.jp/GSWP3/>). It is based on the 20<sup>th</sup> Century Reanalysis (20CR) version 2 performed with the NCEP land-atmosphere model (Compo et al., 2011). 20CR was dynamically downscaled to T248 (0.5°) resolution using the Global Spectral Model (GSM) by data assimilation using the spectral nudging technique (Yoshimura and Kanamitsu, 2008). Bias correction for precipitation, temperature and longwave and shortwave downward radiations were made using GPCC v6 (Global Precipitation Climatology Centre), CRU TS v3.21 (Climate Research Unit), and SRB (Surface Radiation Budget) datasets, respectively. Precipitation was partitioned into rainfall and snowfall referring to the ratio of the downscaled 20CR, and wind induced undercatch correction (Motoya et al., 2002) were applied separately. We upscaled the GSWP3 forcing for 1° spatial resolution.

#### CRUNCEP v7

30 This 6-hourly 0.5° global forcing product (1901-2015) is a combination of the annually-updated CRU TS v3.24 monthly climate dataset (New et al., 2000) and NCEP reanalysis (Kalnay et al., 1996). The latter is only used to generate diurnal and daily

anomalies added to CRU TS monthly means, after bi-linear interpolation to the  $0.5^\circ$  resolution of CRU, except for the precipitation which were linearly interpolated. A threshold of  $0^\circ\text{C}$  in 2-m temperature were used to partition the precipitation into rainfall and snowfall in the CRUNCEP forcing. Rainfall, cloudiness, relative humidity and temperature are taken from the CRU while the others fields (pressure, longwave radiation, windspeed) were directly derived from NCEP (see more details at: <https://vesg.ipsl.upmc.fr/thredds/fileServer/store/p529viov/cruncep/readme.html>). We upscaled the forcing to a  $1^\circ$  spatial resolution dataset, which can be found at [https://vesg.ipsl.upmc.fr/thredds/catalog/store/p529viov/cruncep/V7\\_1901\\_2015/catalog.html](https://vesg.ipsl.upmc.fr/thredds/catalog/store/p529viov/cruncep/V7_1901_2015/catalog.html).

### 5.1.3 Vegetation and soil texture map

The ESA CCI Land Cover map (Bontemps et al., 2013) was used to produce the PFT map for ORCHIDEE. The ESA CCI land cover product is given by 3 maps at a 300 m spatial resolution, corresponding to the years 2010, 2005 and 2000. These maps were derived from the interpretation of MERIS full and reduced resolutions and SPOT-Vegetation time series. Land cover was classified according to the 22 classes used in the UN-LCCS (land cover classification system) scheme, which was translated into PFT fractions used in ORCHIDEE, following the cross-walking method presented by Poulter et al. (2011, 2015). Historical land use maps from the Harmonized Global Land Use dataset (Chini et al., 2014) were incorporated to reconstruct PFT fractions since 1860, following Peng et al. (2017), which will be detailed in the upcoming ORCHIDEE-TRUNK paper for CMIP6.

For soil texture, we use the 12 USDA texture classes provided at a global  $0.08^\circ$  resolution from Reynolds et al. (2000) and upscaled these to the resolution of the atmospheric dataset ( $1^\circ \times 1^\circ$ ). Only the dominant texture type for a grid cell is used at the  $1^\circ$  resolution for defining soil hydraulic parameters (Carsel and Parrish, 1988) in ORCHIDEE-MICT.

## 5.2 Evaluation datasets ~~for the water budget~~

~~Table 1 gives an overview of the selected datasets to evaluate ORCHIDEE-MICT. The selected datasets used to evaluate ORCHIDEE-MICT are summarized in Table 1 and described in Appendix.~~ For the water budget evaluation, we selected six Arctic river basins (Fig. 2b) which are important contributors to total Arctic Ocean river inflow: the four largest Eurasian Arctic basins (Ob, Yenisei, Lena and Kolyma), the Mackenzie basin in northwestern Canada and the Yukon basin in Alaska. The four Eurasian basins (with the smaller Pechora and the Severnaya Dvina basins) drain about two-thirds of the Eurasian Arctic landmass (Peterson et al., 2002), while the Mackenzie is the largest North American river bringing freshwater to the Arctic Ocean (Woo and Thorne, 2003). We also evaluated the Volga basin (Fig. 2b), which is subject to snowfall events during the year but is not underlain by permafrost, in order to compare results with the high-permafrost Arctic basins (Fig. 2a and Table S4).

### 5.2.1 ~~Total terrestrial water storage~~

~~The change of total terrestrial water storage (TWS) is an integrated measure of the ability of a LSM to partition the cold season storage of water as snow and ice, and its decrease from losses to downstream river discharge, plant and soil evapotranspiration,~~

and soil moisture increase. Alkama et al. (2010) showed with the ISBA-TRIP model that seasonal variations in TWS over high-latitude basins resulted primarily from snow accumulation in the cold season, despite a wintertime underestimation of TWS values. Decreasing soil moisture contributed to a small decrease in springtime TWS, while exports with riverflow increased dramatically during the snowmelt period. The GRACE (Gravity Recovery And Climate Experiment) satellite mission permits estimation of monthly TWS variations through measurements of the Earth's gravitational field. Three solutions, at 1° resolution, based on spherical harmonic coefficients (Release 05), are obtained by different processing centers: CSR (Center for Space Research), JPL (Jet Propulsion Laboratory) and GFZ (GeoForschungsZentrum Potsdam) and provided in the GRACE Tellus dataset (Swenson and Wahr, 2006; Swenson, 2012; Landerer and Swenson, 2012). We use the ensemble mean of these solutions to reduce uncertainty in GRACE data (Sakumura et al., 2014). To compare TWS simulated by ORCHIDEE-MICT to GRACE data over the common time period between the two meteorological forcing data sets and GRACE data (July 2003–December 2007), we summed the water stocks simulated by ORCHIDEE-MICT, i.e., soil moisture, snowpack, water on the canopy and water stored for the routing reservoirs. In each grid cell, the corresponding 5-year average is removed from the 2003–2007 time series of TWS output from ORCHIDEE-MICT. The comparison of simulated TWS with GRACE data over seven basins in this study (Fig. 2b) is statistically satisfactory given large surface areas occupied by these basins (> 400,000 km<sup>2</sup> (see Table S4); Swenson et al., 2003).

### 5.2.2 Snow water mass

The GlobSnow v2.0 SWE dataset is based on a data-assimilation approach combining space-borne passive radiometer data (SMMR, SSM/I and SSMIS) with data from ground-based synoptic weather stations. The record provides SWE information on a daily, weekly and monthly basis, at a spatial resolution of approximately 25 km (in EASE-Grid format) and applies to non-mountainous regions of the northern hemisphere, excluding glaciers and Greenland. The product is based on the SWE retrieval methodology developed by Pulliainen (2006) complemented by a time-series melt-detection algorithm (Takala et al., 2009). A complete description of its methodology is given in Takala et al. (2011). SWE estimates are complemented with uncertainty data at the grid cell scale. This work uses monthly-averages of original daily data, interpolated to a 1° grid for appropriate comparison with ORCHIDEE-MICT output.

### 5.2.3 Evapotranspiration (split into components)

The Global Land Evaporation Amsterdam Model (GLEAM) (Miralles et al., 2011) estimates the daily terrestrial ET rate and its components (transpiration, bare-soil evaporation, open-water evaporation, interception loss and sublimation), as well as root-zone soil moisture, at a spatial resolution of 0.25°. It is driven by microwave remote sensing observations and uses satellite soil moisture to constrain potential evaporation rate. The latter is computed with the equation of Priestley and Taylor (1972), which is based on air temperature and net radiation. In the GLEAM v3 product (Martens et al., 2017), representation of evaporative stress has been improved by the use of microwave vegetation optical depth as a proxy for vegetation water content. Other improvements include the use of satellite-based SWE, reanalysis air temperature and radiation, and a multi-source precipitation product. GLEAM v3 contains three sub-versions: Version v3.0a is an assimilated product derived from satellite

observations and multiple reanalysis climate forcing, while v3.0b and v3.0c are purely satellite-based products. As the latter two do not provide full coverage of high-latitude regions (only 50°S — 50°N), the v3.0a, which covers the period 1980-2014, is used in our study. We use monthly averages of original daily data, interpolated to a 1° grid for appropriate comparison with ORCHIDEE-MICT output.

#### 5 5.2.4 River discharge

We use two river discharge databases, one global and another exclusively covering Siberia: The Global Runoff Data Centre (GRDC) product is a global database which collected river discharge data from nearly 7800 stations in 156 countries for the [maximum] period 1807-2017. We use one gauging station per river, the closest to the mouth of the river. For Siberia, we also use daily river discharge at the 3 gauges in three large Siberian rivers (the Ob, Yenisei and Lena) which have been reconstructed (naturalized discharge) to exclude the human impact from the data. This has been performed by using the Hydrograph Routing Model (HRM) developed at the University of New Hampshire (USA) in collaboration with the Arctic and Antarctic Research Institute (Russia) (Shiklomanov and Lammers, 2009). We use monthly averages of original daily data.

#### 5.2.5 Top-soil moisture

The global ESA CCI SM product (v2.2) provides daily soil moisture (in volumetric units  $\text{m}^3 \text{m}^{-3}$ ), at a spatial resolution of 25 km with quality flags specifying potential sources of errors linked to the presence of water bodies, dense vegetation, snow or frozen soils in the pixel area. Three datasets are provided based on different combinations of active and passive satellite sensors. In this work, we used the combined active and passive product which covers the whole period from November 1978 to December 2014. The product results from the merging of soil moisture estimations inversed from two types of instruments and two methodologies: passive microwave radiometers (SMMR, SSM/I, TMI, AMSR-E, AMSR2 and WindSat) using the methodology developed by Owe et al. (2008) and active microwave instruments (ERS-1&2 and ASCAT) using the algorithm developed by TU-Wien (Wagner et al., 1999; Bartalis et al., 2007). These two records were first rescaled together and merged with topsoil (10 cm) soil moisture simulations of the GLDAS-NOAH LSM using a CDF-matching approach. Data accuracy was estimated against in-situ measurements (Dorigo et al., 2015) to  $0.05 \text{m}^3 \text{m}^{-3}$ .

#### 5.2.6 Root-zone soil moisture

The GLEAM root-zone soil moisture v3.0a database (Martens et al., 2017) is based on both satellite observations and reanalysis data. It is a mixture of soil water content of three soil tiles: bare soil (0-10cm), herbaceous (0-100cm) and tall vegetation (0-250cm). The tile fractions are static and derived from MODerate resolution Imaging Spectroradiometer (MODIS) Global Vegetation Continuous Fields product (MOD44B, Hansen et al. 2005). Obviously, it is hard to compare the ORCHIDEE soil moisture with the GLEAM root-zone soil moisture directly not only due to the mismatch of defined soil tile depth but also due to the difference of soil tile fractions. Thus we implement the comparison after normalization of the relative soil moisture values with their respective standard deviation, following Koster et al. (2009).

### 5.3 Evaluation datasets for air-to-soil temperature continuum

#### 5.3.1 Snow depth

Realistically simulating snow depth is one of the prerequisites for accurately modeling soil thermal dynamics, particularly in winter. Daily snow depth in-situ data (1975-2005) from 524, 72, 528, 128 and 528 stations in Europe, Canada, USA, Russia and China were obtained from European Climate Assessment & Dataset (ECA&D), the National Climate Data and Information Archive of Environment Canada, the United States Historical Climatology Network (USHCN), the Russian Research Institute of Hydrometeorological Information—World Data Center (RHHMI-WDC) (Bulygina et al., 2011), and the National Meteorological Information Center of the China Meteorological Administration (Peng et al., 2010), respectively. Monthly averages were calculated when more than 2/3 of daily snow depth data were available.

#### 5.3.2 Surface soil temperature

The soil temperature product used here (André et al., 2015) resulted from the use of combined passive microwave and thermal infrared data to estimate land surface temperature (LST), during summer snow-free periods (snow-covered pixels are masked) in the northern high latitudes. The product is based on the use of SSM/I-SSMIS 37GHz measurements at both vertical and horizontal polarizations and is provided on a 25 km-resolution EASE grid at a one-hour time step for the period 2000-2011. LST retrievals are based on the assumption of a relationship between surface emissivities at both polarizations (Royer and Poirier, 2010) which was calibrated at pixel scale using cloud-free independent LST data from MODIS instruments. The SSM/I-SSMIS and MODIS data were synchronized by fitting a diurnal cycle model built on skin temperature with reanalysis provided by the European Centre for Medium-Range Weather Forecasts (ECMWF). This product was evaluated at local and circumpolar scales against MODIS LST, and the results show a mean Root Mean Square Error (RMSE) of the order of 2.5 K.

#### 5.3.3 In-situ air and soil temperatures

Simultaneous measurements of snow depth, near-surface air temperature and top soil (at 20 cm depth) temperature from Russian meteorological stations (Bulygina et al., 2011; Sherstiukov, 2012) are used to evaluate the thermal insulating effect of snow during winter, by comparing the relationships between snow depth and the air-to-topsoil temperature gradient ( $\Delta T$ ) (Wang et al., 2016b). This dataset includes monthly mean values at 268 sites for the period 1980-2000.

#### 5.3.4 Active-layer thickness

The Circumpolar Active Layer Monitoring (CALM) network's in-situ data set was used to evaluate modeled soil active layer thickness. The CALM network aims to observe the response of near-surface permafrost and the active layer to climate change over long time scales. Thaw depth is measured at the end of the thawing season, so that it should be comparable to the maximum active layer thickness in the model. Before comparing to ORCHIDEE-MICT, the data from 221 sites were averaged from 1990 to 2015.

In addition, we used the active layer thickness map derived by Beer et al. (2013) over Yakutia at a spatial resolution of 0.5°. It is based on regional surveys of landscapes and permafrost conditions in Yakutia during the time period 1960–1987. The gridded datasets can be accessed at the PANGAEA repository. Values of active layer thickness (ALT) in this region range between 0.3 m north of 70°N to 3 m south of 65°N. Uncertainty increases with ALT and is highest (up to 1m) in the south because of the occurrence of discontinuous and sporadic permafrost landscapes.

## **5.4 Evaluation datasets for leaf area, carbon stocks and fluxes**

### **5.4.1 Leaf Area Index**

The Leaf Area Index (LAI) can be derived from satellite measurements, using reflectance measurements and land cover maps. Comparing the seasonal cycle of the satellite products with simulated LAI gives information on the appropriateness of the phenology modules. The amplitude of the cycle is also informative with respect to several photosynthetic parameters (e.g. maximum carboxylation rate  $V_{max}$  or specific leaf area).

#### **GIMMS**

The Global Inventory Modeling and Mapping Studies (GIMMS) LAI3g dataset is derived from Advanced Very High Resolution Radiometers (AVHRR) measurements. It is more specifically computed from the GIMMS Normalized Difference Vegetation Index (NDVI) 3g, using a neural network and the International Geosphere Biosphere Programme (IGBP) land cover classes (Zhu et al., 2013). The 1/12 degree files cover 30 years from July 1981 to December 2011, at a bi-weekly resolution. They were aggregated at a 1.0° spatial resolution on the model grid, to enable comparisons of the time-series and maps.

#### **GLASS**

The Global Land Surface Products (GLASS) LAI product (Liang and Xiao, 2012) is built from AVHRR reflectances over the period 1982–1999 and then from MODIS data over 2000–2012 (Collection 5). The processing is also based on a neural network and the 8 biomes of the MODIS land cover type 3 (Xiao et al., 2014). The files are available at a spatial resolution of 0.05° with an 8-day frequency and were aggregated at 1.0° spatial resolution.

### **5.4.2 NEE from atmospheric inversions**

We use estimates of NEE (here the net land-atmosphere CO<sub>2</sub> fluxes excluding fossil fuel CO<sub>2</sub> emissions) from two atmospheric inversions where NEE is optimized at the resolution of a global atmospheric transport model each month to match observed CO<sub>2</sub> concentration gradients from a global network of ground-based stations. The inversions are the Copernicus Atmosphere Monitoring Service (CAMS) inversion system (Chevallier et al., 2010) and the Jena CarboScope (Rödenbeck, 2005).

The inversion from CAMS uses atmospheric CO<sub>2</sub> concentration observations from a total of 132 sites covering the period 1979–2015 and includes all sites being added to the network through time, combined with the LMDZ-INCA atmospheric transport model and prior information about fossil fuel CO<sub>2</sub> emissions (from EDGAR3.2 FastTrack 2000 database) as well as



land/ocean-atmosphere fluxes, to minimize a cost function in order to calculate NEE. Here we use the latest version (15r4), in which monthly fluxes are calculated at  $1.875^{\circ} \times 3.75^{\circ}$  lat/lon resolution.

The Jena CarboScope atmospheric inversion uses the same set of atmospheric  $\text{CO}_2$  observation sites available during the temporal observation period (1996-2015) referred to as Jena s96. Jena CarboScope inversion uses fossil fuel emissions from EDGAR 4.2 and the TM3 global atmospheric transport models. Here we use monthly NEE from the latest version of Jena s96 (v3.8), which is provided at  $3.8^{\circ} \times 5.0^{\circ}$  lat-lon resolution.

### 5.4.3 Site-level GPP and NPP observations

#### In-situ GPP and NPP measurements

While there are several NPP datasets especially for forest (e.g. Cannel et al., 1992; Olson et al., 2001; Michaletz et al., 2014), few provide both GPP and NPP at the same locations. We used a recent database of in situ co-located GPP and NPP measurements (Carnioli et al., 2015), which is an extension of the Luyssaert et al. (2007) database for forests. The database consists of 131 sites, for which annual NPP obtained from biometric measurements and associated uncertainties are provided. The criteria employed by Carnioli et al. (2015) for selecting the sites was the availability of methodologically independent and site-specific estimates of NPP and GPP. Carbon Use Efficiency (CUE) was calculated at each site by taking the ratio of NPP to GPP. Site-level biomes are classified as one of tundra, boreal peatlands, marshes, forests, grasslands or croplands. To study the temperature sensitivity of GPP and NPP, we selected only sites northward of  $30^{\circ}\text{N}$ , where model grid cells were representative of both the vegetation type and meteorological conditions observed at each site. We thus restricted the selection to sites whose IGBP vegetation type had a corresponding fraction greater than 0.5 in the grid cell of ISLSCP II MODIS IGBP Land Cover product (Friedl et al., 2010). Site-years where the absolute difference between local and the Mean Annual Temperature (MAT) of the GSWP3 or the CRUNCEP climate forcing fields was higher than 4 degrees were discarded. As no information was given about the time-period of site NPP and GPP observations in Carnioli et al. (2015), we compared them to mean values of the model over the period 2000-2007. The subset of selected sites consisted then in 52 sites out of 131 in the full dataset.

### 5.4.4 Gridded GPP and NPP observation-based data

#### GPP

In addition to GPP from sites, we used the monthly gridded GPP observation-based product Model Ensemble Tree GPP (MTE-GPP) from Jung et al. (2009, 2011) from year 1982 to 2010. This product is the result of a statistical model that combines measurements at FLUXNET sites with geospatial information from satellite remote sensing (fAPAR) and meteorological data. MTE-GPP is obtained as the median from an ensemble of the 25 best model trees out of an initial set of 1000 and the Mean Absolute Deviation (MAD) of the ensemble is used to define the uncertainty of MTE-GPP, which is also provided as a monthly gridded product. This uncertainty estimate was mentioned by Jung et al. (2011) to be underestimated as compared to the true RMSE. That is why we consider the RMSE from the cross-validation sites as an estimate of uncertainty. The RMSE of mean

MTE-GPP at cross-validation sites is  $270 \text{ gC m}^{-2} \text{ yr}^{-1}$ . At global scale and over the period 1982–2008, the mean MTE-GPP is  $933 \pm 46 \text{ gC m}^{-2} \text{ yr}^{-1}$  and the total is  $119 \pm 6 \text{ GtC yr}^{-1}$ , the uncertainties being derived from the MAD of the MTE (Jung et al., 2011).

## NPP

- 5 We use the MOD17A3.005 global annual NPP model driven by satellite observations from the NASA Land Processes Distributed Active Archive Center, at 1 km resolution and aggregated to the  $1^\circ$  grid of the model over the period 2000–2010. Yearly MODIS NPP is computed from yearly and daily components. Among the daily components, MODIS GPP is based upon a LUE model:

$$\text{GPP} = E * f\text{APAR} * \text{PAR} \quad (15)$$

- where PAR is the Photosynthetically Active Radiation, fAPAR the fraction of absorbed PAR and E is the radiation conversion efficiency. E depends on vegetation type and on meteorological conditions (temperature, Vapor Pressure Deficit (VPD)). PAR, temperature and VPD used in MOD17A3.005-NPP are from the NASA Data Assimilation Office (DAO) and fAPAR is a MODIS product (Knyazikhin et al., 1998). A daily Maintenance Respiration (MR) was computed for leaves (l) and fine roots (fr), using the LAI MODIS product (Knyazikhin et al., 1998). On a yearly basis, MR was computed for livewood (lw) and Growth Respiration (GR) for leaves, fine roots, livewood and deadwood (dw). These respirations were computed using a biome-specific parameter look-up table, derived from the BIOME-BGC terrestrial biosphere model (Heinsch et al., 2003). This set of parameters was recalibrated for the MOD17A3.005-NPP collection 5 (Zhao et al., 2005). Yearly MODIS-NPP was thus computed as follows (Running et al., 2004):
- 10  
15

- Turner et al. (2006) evaluated MODIS NPP at nine sites representing various biomes, using local meteorological, LAI and above ground NPP measurements as inputs for the BIOME-BGC model to derive NPP over a  $25 \text{ km}^2$  area. They showed globally no bias but NPP was underestimated at the most productive sites (attributed to a negative bias of E) and overestimated at low productivity sites (attributed to a positive bias of MODIS fAPAR). Zhao et al. (2005) evaluated MODIS NPP against the Ecosystem Model Data Intercomparison (EMDI) global database of NPP measurements (Olson et al., 2001) and found a good agreement except for tropical forests (spatial  $R^2 = 0.77$ ). Zhao et al. (2006) estimated the sensitivity of the MODIS NPP product to the use of different meteorological reanalysis input data, showing differences at global scale up to more than
- 20  
25  $20 \text{ PgC yr}^{-1}$ , with the largest differences in the tropics.

### 5.4.5 Soil carbon inventories

#### NCSCD

- The Northern Circumpolar Soil Carbon Database quantifies storage of organic carbon in soils of the northern circumpolar permafrost region, comprising four depths up to 3m (0–30cm, 30–100cm, 100–200cm and 200–300 cm) (Hugelius et al., 2013).
- 30 Total SOC storage in the 0–2m and 0–3m depth ranges in northern permafrost soils are estimated to be  $827 \pm 108$  and  $1035 \pm 150 \text{ PgC}$  respectively.

## ISRIC

ISRIC is a global soil information database with spatial predictions for a selection of soil properties, at 1 km resolution and six depths, up to a maximum of 2m (0-5cm, 5-15cm, 15-30cm, 30-60cm, 60-100cm and 100-200cm). About 110,000 soil profiles were used to generate the product, using statistical models based on climatic and biomass indices, lithology, and taxonomic mapping units (Hengl et al., 2014). The gridded maps of soil organic carbon mass fraction, soil bulk density, and volumetric fraction of coarse fragments was used to calculate SOC density in the unit of  $\text{kgCm}^{-2}$  to be comparable with model output (see Eq. 6 in Hengl et al., 2014). However, we found in this product a systematic overestimation of bulk density for organic-rich soils, which is similar to the issue found in the HWSD database Köchy et al. (2015). Therefore, we followed the adjusting method in Köchy et al. (2015) to correct bulk density for histosols and other soils with organic carbon mass fraction larger than 3%. This adjustment decreases total SOC stock in northern permafrost regions (the same domain as NCSCD) in the 0-2m depth range from the original 1724 PgC to 1177 PgC, which we considered acceptable though higher than NCSCD.

### 5.4.6 Biomass carbon stocks

Two forest biomass datasets are used to evaluate simulated forest biomass. The first is that from Avitabile et al. (2016), which, at a spatial resolution of  $0.01^\circ$ , merges two tropical aboveground forest biomass (AGB) datasets from Saatchi et al. (2011) and Baccini et al. (2012) with northern hemisphere volumetric forest stock growth data from Santoro et al. (2015). The second forest biomass dataset from Thurner et al. (2014) is given at a spatial resolution of  $0.01^\circ$  for 2010 and confined to the northern hemisphere ( $30^\circ\text{N}$ – $80^\circ\text{N}$ ). This is derived from the Santoro et al. (2011) forest standing stock data. The two datasets were re-gridded to  $1^\circ$  resolution for comparison with model output. The Avitabile et al. (2016) AGB was converted into total biomass, assuming a constant AGB to total biomass ratio of 0.8 and divided by two to obtain total biomass carbon, under the assumption that the carbon content of dry biomass is 50%. Liu et al. (2015) synthesized ratios of aboveground to total biomass for different forest biomes in temperate and boreal regions and found that they range between 0.76 and 0.84 according to regional forest inventory assessments. Our uniform 0.8 factor for converting the Avitabile et al. (2016) data thus yields a potential error of ~5% in inferred total biomass carbon, which is within the reported data uncertainties given by Avitabile et al. (2016). Simulated equivalent total forest biomass carbon over the period 2000–2007 was compared with observations.

### 5.4.7 Burned area and fire emissions

We compared the simulated burned area and fire carbon emission with the Global Fire Emissions Database (GFED4s) data set, which is based on the GFED modeling framework (van der Werf et al., 2010). The GFED4s burned area data is derived from the MODIS sensor data that is then complemented by specific algorithms to retrieve “small fires” (Randerson et al., 2012). Emissions are computed from a combination of burned area with the revised Carnegie-Ames-Stanford Approach biogeochemical model (CASA-GFED, described in van der Werf et al., 2010), that provides an estimation of fuel loads and combustion completeness. GFED4s data is provided at a  $0.25^\circ$  spatial resolution with a monthly time-step for 1997–2015.

Cropland fires are not yet implemented in the ORCHIDEE-MICT model, but are included in the GFED4s dataset. In order to compare the simulation with GFED4s, it was necessary to correct the simulated burned area for the omission of cropland fires. To do so, we computed for each grid cell the fraction of existing natural PFTs (i.e. all non-cropland PFTs), and divided the simulated burned area in each grid cell by this value. For carbon emissions, the GFED4s dataset provides separate emissions data for cropland and natural fires. In this way, we were able to simply remove cropland fire emissions from GFED for comparison with ORCHIDEE-MICT output.

## 6 Evaluation of the atmosphere-snow-soil continuum

In the following, we analyze model performance in replicating the transfer of heat from atmosphere to deep soils. This is done by evaluation against measured temperature gradients, starting from snow depth controls on winter  $\Delta T$ , followed by evaluation of surface (skin) temperature in summer, and the temperature gradients between the near-surface and deeper soils.

### 6.1 Snow insulation controls on the temperature gradient between air and topsoil

#### 6.1.1 Snow depth

ORCHIDEE-MICT correctly captures the spatial distribution of maximum monthly average snow depth (Fig. 3a,b), and the seasonal decrease of snow depth from March to June (Fig. 3c), but modeled snow depth strongly depends on the atmospheric forcing used. GSWP3 climate forcing tends to produce a larger maximum snow depth than CRUNCEP, greater than those observed in all Northern regions, especially over boreal Europe (BOEU) (Fig. 3c). This shows that uncertainty from climate forcing data is as large as the model bias compared with observations, making it difficult to attribute a model bias to a particular component of the snow model. However, the rate of sublimation in winter (Pomeroy et al., 1998) and the prescribed albedo value of fresh snow have been shown to be critical in determining the peak value and phase of both snow depth and SWE (Wang et al., 2015b).

#### 6.1.2 Snow conductivity and snow density

Mean snow density and mean snow thermal conductivity are computed at the month of maximum snow depth over the 1981-2007 period as weighted averages over the three snow layers. Gouttevin et al. (2012b) report density values of  $200 \text{ kg m}^{-3}$  for taiga and  $330 \text{ kg m}^{-3}$  for tundra and conductivity values of  $70 \text{ mW m}^{-1} \text{ K}^{-1}$  for taiga and  $250 \text{ mW m}^{-1} \text{ K}^{-1}$  for tundra, from Sturm and Johnson (1992) and Domine et al. (2010). These higher values over tundra were attributed to snow compaction by wind. This process is not modeled in ORCHIDEE-MICT and we thus simulate similarly high values of conductivity for both biomes (Fig. S3): approximating tundra with C3 grass PFT between  $55$  and  $85^\circ\text{N}$ , and taiga with the boreal forests PFTs between  $45$  and  $70^\circ\text{N}$  and considering only grid cells with a fraction of the dominant biome above  $0.6$ . The model yields a mean snow conductivity of  $266 \pm 203$  (GSWP3) and  $219 \pm 197$  (CRUNCEP)  $\text{mW m}^{-1} \text{ K}^{-1}$  for tundra compared to  $221 \pm 113$  (GSWP3) and  $182 \pm 100$  (CRUNCEP)  $\text{mW m}^{-1} \text{ K}^{-1}$  for taiga and a mean density of  $269 \pm 102$  (GSWP3) and  $239 \pm 103$

(CRUNCEP)  $\text{kg m}^{-3}$  for tundra and of  $233 \pm 67$  (GSWP3) and  $207 \pm 63$  (CRUNCEP)  $\text{kg m}^{-3}$  for taiga. Note that a recent study (Domine et al., 2016) suggests for tundra a complex structure with depth-hoar developing at the base of snowpack during the course of the snow season, causing conductivities as low as  $20 \text{ mW m}^{-1} \text{ K}^{-1}$  in late winter, whereas snow-compacted upper layers have conductivities of 200 to  $300 \text{ mW m}^{-1} \text{ K}^{-1}$ , more comparable to ORCHIDEE-MICT.

## 5 6.2 Summer land surface temperature

ORCHIDEE-MICT overestimates summer (June-August) LST of about  $1.36^\circ\text{C}$  when forced by GSWP3 (Fig. 4b) and  $0.49^\circ\text{C}$  by CRUNCEP (Fig. 4c). The bias is larger in northern and southern Siberia, where it can reach  $4^\circ\text{C}$ . These differences may be linked to the overall underestimation of ET (see Fig. 11). This is further addressed in the Discussion (Section 10).

## 6.3 Soil temperature

- 10 The simulated spatial patterns of mean annual topsoil (0.2 m) temperature generally reproduce the observed gradient along a southwest-northeast transect in Siberia (Fig. 5a,b). However, the CRUNCEP-forced simulation results are colder than those from the GSWP3 ones as well as relative to observations in the permafrost region (Fig. 5a,b), mainly driven by a strong cold bias in CRUNCEP-based winter soil temperatures (Fig. 5c,d).

- During winter, the snowpack acts as an insulating layer above soil surface, reducing soil heat loss. Snow thus causes a large  
15 positive temperature gradient ( $\Delta T$ ), which is controlled by both snow depth and snow thermal properties, such as thermal conductivity, density and albedo. Generally, the model underestimates snow insulation in the early (Nov to Jan) and late (Feb to Apr) cold seasons for the same snow depth, as compared to observations (Fig. 6). This indicates that relatively congruent wintertime soil temperatures in the GSWP3-forced simulation in permafrost regions (Fig. 5c) may be due to a bias compensation from overestimated snow depth (Fig. 3) and underestimated snow thermal insulation.

- 20 A prominent component of the  $\Delta T$  / snow depth relationship observed at Russian stations (black in Fig. 6) is the significantly lower insulation during the late cold season, probably due to snow compaction and densification leading to higher snow conductivity (Decharme et al., 2016). This differential sensitivity of  $\Delta T$  to snow depth between the two periods is effectively captured by ORCHIDEE-MICT, despite small modeled negative  $\Delta T$  values (i.e. topsoil colder than air) compared to observations at the termination of the snow season, when snow depth is diminished ( $< 20 \text{ cm}$ ).

- 25 Summer soil temperatures are higher in the GSWP3-forced simulation relative to those of CRUNCEP, ~~and warmer than observations by  $1\text{--}2^\circ\text{C}$  at 0.8 and 1.6 m depth at thaw season (Fig. 5c,d). This is consistent with an overestimation of ALT~~ and warmer than observations from the Russian meteorological stations in continuous permafrost region by  $1\text{--}2^\circ\text{C}$  on average at 0.8 and 1.6 m depths (Fig. 5c,d). Spatially however, the bias in peak summer soil temperature varies for different regions, with a large warm bias for the Lena basin below 0.8 m, and some cold bias for the further eastern sites (Fig. S4). This is  
30 consistent with the overestimation of ALT for Yakutia (Fig. S5) compared to the empirically-derived map by Beer et al. (2013) (see Section 6.4). Differences between the two simulations in summertime soil temperatures may be partly driven by warmer GSWP3 land skin temperatures during summer (Fig. 4) than CRUNCEP. In addition, the cold bias in winter soil temperatures

in the CRUNCEP-forced simulation might be carried over to summer via soil thermal inertia. This would partly offset the warm bias in CRUNCEP land surface temperatures during summer (Fig. 4c).

#### 6.4 Active layer thickness and permafrost area

Figure 7a,b shows the simulated spatial pattern of ALT, as calculated from modeled soil temperatures using a linear interpolation between soil layers to locate the first depth that remains frozen (below 0°C) year-round. Compared with CALM observations, in which most of the sites have ALT <1m, the GSWP3-forced model generally overestimates ALT by more than 1 m (also see Fig. S35 which compares modeled ALT of Yakutia, East Siberia with the map of Beer et al., 2013), whereas CRUNCEP-forced output shows relatively better agreement with the observations. ~~This is consistent with our diagnosis that GSWP3-forced soil temperatures (Fig. 5) are overestimated in permafrost regions during summertime at depths of 0.8 m and 1.6 m. Similar results have been reported in several modeling studies (e.g. Ekici et al., 2014; Decharme et al., 2016), which show deeper modeled ALT compared to observations. Apart from the uncertainty induced by climate forcing differences, the model-data mismatch may also arise from scale differences, since models forced by 1°x1° resolution data cannot capture the sub-grid scale heterogeneity of land cover, soil texture, organic matter modulated soil physical properties and topographic microclimates.~~ Apart from the uncertainty induced by climate forcing, the model-data mismatch may also arise from scale differences for the organic carbon content that is used to calculate soil physical properties for each grid cell. As mentioned in Section 4.2, the empirical SOC map from NCSCD (Hugelius et al., 2013) is prescribed for permafrost regions in the soil thermal and hydrological modules, which is upscaled by the model to the target spatial resolution (1° by 1° in this study). These SOC values thus do not represent the site-level soil conditions, aside from the uncertainty of the NCSCD database itself. To further investigate this impact, we conducted additional simulations for the sites that provide explicit organic layer thicknesses (in total 69 sites), forced by CRUNCEP. In these runs, we assumed pure organic soil, i.e.  $f_{i,SOC}$  in Eq. 9 equaling to one, for the soil layers above the site-specific organic layer thickness, while kept the SOC concentration unchanged below this thickness, i.e. from NCSCD. Note that the moss layer, vegetation mat, and/or organic root zone as described in some sites were all summed to derive a total organic layer thickness. The other configurations including climate forcing and soil texture were the same as the regional simulation. The result is displayed in Fig. S6, showing significantly shallower ALTs simulated by these site runs which better match the observations (Fig. S6a), with different magnitudes of ALT reductions among the sites (Fig. S6b).

For simulated permafrost extent, two typical definitions of permafrost in LSMs, one defined as ALT less than 3 m to give “near surface” permafrost (e.g. Koven et al., 2013), the other defined if any of the soil layers stay frozen (e.g. Ekici et al., 2014; Burke et al., 2017b), produce quite different permafrost extents (Fig. 7c,d). This result highlights that the intercomparison of permafrost areas among different LSMs with differing soil vertical discretizations may include uncertainties brought by the arbitrarily chosen definition, whereas evaluation and comparison directly for soil temperatures and ALT should be more robust. A qualitative comparison against the empirical IPA (International Permafrost Association) permafrost map (Brown et al., 2002) shows better agreement for CRUNCEP compared to GSWP3-forced output, since CRUNCEP-forced simulation

general matches the distribution of continuous permafrost using the former definition, while GSWP3-forced simulation seems to underestimate permafrost extent (Fig. 7c,d). This is consistent with the deeper simulated ALT under GSWP3 climate forcing.

## 7 Evaluation of large-scale water storage and fluxes

Simulated water budget components are evaluated over selected northern basins (Fig. 2b), most of which are underlain by permafrost (e.g. Lena, Kolyma), with the exception of the warmer Volga. The Ob and Yenisei catchments have contrasting north-south precipitation and temperature regimes, with attendant impacts on Arctic Ocean discharge and sub-basin scale water budgets. Here, only basin-scale averages are discussed.

### 7.1 Total terrestrial water storage change

A realistic phase and amplitude of TWS is simulated with both forcing datasets, although peak-to-peak amplitude is slightly overestimated in the Volga, Yenisei and Kolyma basins under GSWP3 input and the seasonal amplitude underestimated in the Yukon with both forcings, and in the Mackenzie and Lena with CRUNCEP forcing (Fig. 8). The positive temporal trend of TWS in the Ob, Lena and Kolyma basins is captured well by ORCHIDEE-MICT, where it reflects upward precipitation trends (not shown). In general, the GRACE TWS is well captured by ORCHIDEE-MICT with both forcings, at the seasonal scale and for the 5-year trends, except in the Yukon basin where observed TWS decreases are not reproduced in our simulations, with no precipitation decrease in the GSWP3-forced model. The Yukon TWS decline is likely due to glacier melt in the northwest Cordillera (Wang et al., 2015a). As glaciers are not represented in ORCHIDEE-MICT, the model does not capture these TWS trends. Note that groundwater storage changes related to the development of closed and open taliks (Muskett and Romanovsky, 2009) which contribute to existing TWS trends – increasing storage in the Lena and Yenisei, decreasing it in the Mackenzie basin, and no change in the Ob – are not modeled either in ORCHIDEE-MICT. Despite this, the model reproduces observed trends in these basins.

### 7.2 Snow related processes controlling land water storage in the cold season

The modeled seasonal cycle of the SWE and length of the snowmelt period are in agreement with observations (Fig. 9), suggesting a good parameterization of the snowmelt and sublimation processes. Yet, results differ strongly according to forcing inputs. In basins with a large permafrost fraction (Yukon, Lena and Kolyma) and, to a lesser extent, in the Mackenzie, Ob and Yenisei basins, SWE is underestimated throughout the year compared to GlobSnow data when ORCHIDEE-MICT is forced by CRUNCEP, and it is significantly larger in the GSWP3-forced simulation (in the Volga basin, the SWE is overestimated in the two simulations). This is due to the low basin-specific snowfall rate in CRUNCEP forcing compared to GSWP3 (Fig. S103), which is probably the result of the criterion used to partition rainfall and snowfall in CRUNCEP, and strongly affects the simulation of snow depth and SWE (Loth et al., 1993; Wen et al., 2013). By contrast, the GSWP3-forced model captures the early winter SWE accumulation in these basins. In spring, the SWE is systematically overestimated except over the Lena, whose seasonal cycle is well reproduced by ORCHIDEE-MICT. This corresponds to an excessive persistence of the snow



cover, which may be explained by the absence of hysteresis in the snow depletion curve relating the snow cover extent and the SWE (e.g. Magand et al., 2014). In the Yenisei and the Mackenzie basins, the SWE in winter is closer to observations with the GSWP3 forcing, with the exception of springtime values, which are better under CRUNCEP forcing. In basins where permafrost area is near-zero (Ob and Volga), the SWE from CRUNCEP-forced simulation is closer to Globsnow than those from GSWP3, in which SWE is overestimated in winter and spring. This is related to the large difference in snowfall over these basins between the two forcings (Fig. S103).

### 7.3 Soil moisture

#### In the topsoil

Seasonal evolution of topsoil moisture (first 2 cm) is compared to the ESA-CCI-SM product over the seven basins (Fig. 10a). Liquid soil moisture values from the model are used for the comparison because ESA-CCI-SM product measures only the topsoil moisture when temperature is above 0°C. Because of scaling issues (the satellite product is rescaled to the 10 cm top layer of NOAH model, as already noted, but is more representative of a thinner soil layer of about 2 cm), the comparison was performed on relative liquid soil moisture values after normalization with their respective standard deviation. The observed seasonal moisture variations are well captured by ORCHIDEE-MICT with both forcings over the seven basins (Fig. 10a). The maximum values occur in summer, in contrast to lower latitudes, because of the thawing processes occurring in summer. The local minimum simulated in summer in the Volga and Ob basins is underestimated, suggesting a too slow infiltration front of the water in the top soil layers of the model. Thus, less water in the root zone is available for transpiration which is found to be underestimated when compared to GLEAM (see Fig. 11b). Compared to observations, a more rapid increase (decrease) of the modeled topsoil moisture in spring during snow melt (in fall) is found in the Yukon and Mackenzie basins, related to a more rapid thawing (freezing) of the top soil.

#### In the root zone

The soil water deficit is of primary importance during spring and summer in the high latitudes because of its potential impacts on the vegetation transpiration, leading to a surface temperature increase and a reduction in the productivity. Yet, Aa soil water comparison between GLEAM and ORCHIDEE-MICT is difficult because of differences in soil depth, which in GLEAM varies with vegetation cover, but is fixed at 2 m in ORCHIDEE-MICT. Moreover, the fraction of soil tile of short vegetation and forests used in the two products is not the same. We therefore normalize the relative root soil moisture (Fig. 10b) by its standard deviation to compare the dynamic of the soil moisture rather than the total amount of water in the soil. The intensity of water uptake by the roots of the vegetation in summer is generally well simulated by the model in both simulations. In basins underlain by permafrost, except in the Lena basin, water uptake is delayed by one month for both forcing sets, while the rate of decrease in root soil moisture is underestimated for GSWP3-forced output only. The similarity in output in this respect, despite very different SWE, highlights the low impact of the latter on the root soil moisture, and further underscores how the snowmelt differential is lost through runoff rather than being available for vegetation.



#### 7.4 Evapotranspiration and component fluxes

The amplitude of the ET seasonal cycle is generally well captured by the model in most basins, when compared to the GLEAM product, despite a systematic underestimation by ORCHIDEE-MICT, whichever the input forcing (Fig. 11a). The disparity between modeled peak ET and GLEAM data is reduced under GSWP3 forcing in the Yukon, and under CRUNCEP forcing in the Yenisei and Lena. ET increase is underestimated in spring and early summer for both forcings, except in the Volga basin. This is consistent with modeled LAI increasing too late in spring (see Fig. 13) which could be due, at least partially, to an excessive persistence of the snow cover in spring. In autumn, the timing of the decrease in ET is reproduced by both forcings. Model biases with respect to GLEAM data in the sublimation, soil evaporation and transpiration components of ET are shown in Fig. 11b. In all basins, sublimation bias in simulations forced by GSWP3 is ~0 in winter, in agreement with GLEAM. By contrast, CRUNCEP forced simulations slightly overestimate sublimation in early spring across basins with the exception of the Volga. These results are consistent with SWE underestimation (except in the Volga) (Fig. 9) and higher downward shortwave radiation (Fig. S125) that results when CRUNCEP forcing is used. The general underestimation of summer ET by ORCHIDEE-MICT in the Yukon, Mackenzie and Kolyma basins is explained mainly by a too low transpiration despite bare soil evaporation being slightly over-estimated (Fig. 11b). When forced by CRUNCEP data, ORCHIDEE-MICT overestimates interception loss in all basins but the Kolyma and Yukon, which is consistent with CRUNCEP LAI overestimation (see Fig. 13).

#### 7.5 River discharge

By comparing the two simulations, it is clear that the meteorological forcing exerts a significant influence on the simulated river discharge (Fig. 12): GSWP3 leads to systematically higher river discharge than CRUNCEP, which is perfectly consistent with the SWE biases (Fig. 9). In a majority of basins, GSWP3-forced simulations better capture the seasonal cycle of observed discharge than CRUNCEP-forced ones, especially in the Yukon and the eastern Siberian basins, where the discharge is strongly underestimated under CRUNCEP forcing (between 60% in the Yenisei and 83% in the Yukon).

A first feature of the nival regime characterizing the studied high-latitude basins is the occurrence of low flows in winter, when water is frozen in the snowpack, soils, and river ice. This is well simulated by ORCHIDEE-MICT, despite a small underestimation in the Yukon, Mackenzie and Yenisei. Naturalized river discharge is available in the latter, and lower in winter than GRDC values, which reflects the effect of reservoir operations. The winter discharge simulated by ORCHIDEE-MICT in the Yenisei is closer to the naturalized estimates, as the model does not account for artificial reservoirs. No natural lakes are simulated by ORCHIDEE-MICT, which may contribute to the winter discharge underestimation, especially for the Mackenzie which includes massive lakes (Great Slave, Great Bear, and Athabasca).

The nival regime of the studied basins is also characterized by peak flow in late spring, which is broadly captured by the ORCHIDEE-MICT simulations, even if the magnitude of peak flow is strongly biased in some cases, with a strong link to the forcing used and the SWE biases. As an example, the Volga is the only river where both peak flow and SWE are overestimated

with both forcing, and closer to river discharge observations under CRUNCEP forcing. In this human-altered basin, the absence of the simulation of water withdrawals for irrigation in the model can explain the peak flow overestimation.

An almost systematic weakness of the simulated hydrographs is the underestimation of river discharge during summer and fall, which is very strong with both forcing in the Yukon, Mackenzie, Lena and Kolyma, and to a lesser extent in the Yenisei. This makes discharge closer to observations during the second half of the year under GSWP3 than under CRUNCEP, particularly in the Ob. This summer-fall underestimation propagates to underestimated annual mean discharge in a majority of basins (reaching -25 and -30% under GSWP3 in the Lena and Kolyma rivers) despite the overestimation of peak flows. Eventually, the discharge underestimation found in summer and fall, and on annual means in many basins, is not coherent with the low simulated ET compared to GLEAM data. Yet, according to Wang et al. (2015a), biases in precipitation and ET datasets, that are used to evaluate the models, are source of errors for the water imbalances found in the northern high-latitude basins.

These river discharge results highlight deficiencies in the model representation of water infiltration in frozen soils, which appear to be too drastic in their prevention of snowmelt infiltration under conditions of frozen topsoils. Alternative parameterizations of these dynamics are underway. For example, infiltration of meltwater into frozen soil could be permitted when accounting for sub-grid scale variability of topsoil freezing and drying which would enhance infiltration (Gray et al., 2001). Other perspectives are the improvement of the floodplain parameterization in ORCHIDEE-MICT (Lauerwald et al., 2017) and the inclusion of natural lakes and artificial reservoirs.

## 8 Evaluation of the leaf area index, gross and net CO<sub>2</sub> fluxes

### 8.1 Leaf area index

According to the two evaluation products, the seasonal cycle of LAI (Fig. 13) is similar across the basins, with values near zero during winter, and a maximum in summer. There is a consistent phasing of seasonal LAI between the two evaluation datasets, however maximum LAI in GIMMS is systematically higher than in GLASS. In all the basins, the LAI simulated by ORCHIDEE-MICT has a phase delay of up to one month compared to both products. This is due to a delay in the start of the growing season, which may be related to excessive persistence of the snow cover (Fig. 9). The phenological models in ORCHIDEE (detailed in MacBean et al., 2015, Appendix A) do not explicitly take into account this influence, unlike what is done in Van Wijk et al. (2003), who model the link of the start of the tussock tundra growing season to the soil thaw at 10-cm depth. However there is a first indirect link between the snow cover and the vegetation phenology through air temperature, which influences both the start of the growing season, determined in ORCHIDEE using growing degree days (GDD)-based phenological models for deciduous species, and the start of the snowmelt season. There is a second indirect link through snowmelt. While there is still a large amount of snow, the soil surface temperature is kept at zero degree Celsius or below and the soil cannot thaw. Only when snowmelt occurs and when the snow fraction is small enough, will the soil start thawing, thus increasing soil liquid water content. This impacts the start of the growing season for grasses and crops, which use both a GDD and a soil moisture thresholds, and also reduces water stress, thus favoring photosynthesis for all PFTs. Note here that ORCHIDEE-MICT is prone to overestimate the timing of senescence (MacBean et al., 2015). This is true in particular for

conifers, for which the model lacks an explicit senescence inception model. Except in the Yukon and the Mackenzie basins, the maximum LAI simulated by ORCHIDEE-MICT lies between the two satellite product estimates. Winter LAI of ~1.0 are overestimated in the Yukon, Mackenzie, Ob and Yenisei basins, however observations show values around zero. Given that these basins are covered by a larger fraction of evergreen forests compared to the others, these simulated values look reasonable. The mismatch with the observations could be explained here by data errors, the assessment of solar reflectance from space in winter in the high latitudes being less reliable.

## 8.2 Gross (GPP) and net (NPP) primary productivity

### 8.2.1 Spatial distribution and seasonal cycle of GPP

GPP in the high latitudes is co-limited by cold temperatures, constraining the duration of the growing season, and by summer water stress (Schulze et al., 1999) in northern Canada and Siberian boreal forests, for which the water balance is usually negative at this time. The simulated spatial pattern of GPP in Eurasia and North America is close to the MTE-GPP dataset (Fig. 14). Values lower than MTE-GPP were simulated in Eastern Siberia under the GSWP3 forcing, mainly due to water stress (see also underestimated biomass for eastern Siberia in Figs. 20). The seasonal GPP cycle in Fig. 14 (lower panel) is generally accurate with respect to observations at the scale of large boreal regions; however peak GPP is strongly overestimated for boreal North America (BONA).

Interestingly, comparing GPP forced by the two climate datasets shows higher values by CRUNCEP than by GSWP3 (Fig. 14), despite a generally lower precipitation in CRUNCEP (Fig. S17). This could be explained by the higher specific air humidity during summer in CRUNCEP than in GSWP3 (Fig. S14). A low air humidity increases the atmospheric vapor pressure deficit (VPD) and the leaf to air vapour pressure difference; plants then partially close the stomata to constrain a potentially fast transpiration (Oren et al., 1999; McAdam and Brodribb, 2015), which leads to reduced photosynthetic rate. The photosynthesis module in ORCHIDEE largely follows Yin and Struik (2009) in which stomata conductance decreases with an increasing VPD, thus is able to simulate a lower GPP under dry air conditions. A recent study (Novick et al., 2016) showed that, between the two factors that impact plant water stress, i.e. soil moisture supply and atmospheric demand for water (reflected by VPD), the latter limits evapotranspiration to a greater extent than the former in relatively wet forested ecosystems. In spite of its importance, the effect of VPD on vegetation productivity has been far less studied than soil water availability (Konings et al., 2017), warranting further investigations in both observations and land surface models.

### 8.2.2 Spatial distribution of NPP and site level comparison

The distribution of NPP for both ORCHIDEE-MICT simulations is compared to forest site data from Campioli et al. (2015) (Fig. 15), in which NPP measurements are collocated with GPP (118 sites in total north of 30°N). Despite no sampling from Western Russia and Siberia, the model is able to reasonably capture NPP gradients in BONA and BOEU. In temperate forests of western Europe and eastern US, modeled NPP is too low (see also Fig. 16b for warm sites), possibly due to high water stress in ORCHIDEE-MICT (see below) ~~or due to a lack of N-deposition combined with soil fertility effects in modeled NPP~~.

Considering only the 52 sites that were selected (as described in [Section 5.4 Appendix C3](#)), we computed the distributions of GPP and NPP by 5 °C mean annual temperature bins for the sites, the MTE-GPP and MODIS-NPP products, and for the ORCHIDEE-MICT simulations. We plot the 95th percentiles of these distributions in Fig. 16, which arguably defines an upper envelope for temperature-limited GPP and NPP (see Fig. 3 of Luyssaert et al., 2007), and allows us to evaluate the spatial sensitivity of GPP and NPP to mean annual temperature. The behavior of all products is similar to the sensitivities derived from site observations, with a strong positive relationship between GPP, NPP and mean annual temperature over the range -10 to 10°C. ORCHIDEE-MICT then captures the decrease of GPP (NPP) at warmer sites but underestimate the values above 5°C. Global gridded data products generated independently from the Campioli et al. (2015) dataset exhibit sensitivities comparable to those derived from local sites, whereas modeled GPP and NPP saturate for temperatures above 10°C, indicating water stress dominant controls.

The local measurements and the global observations products give similar median CUE values (51% and 49% respectively) and first and third quartiles whereas the model gives a narrower distribution range, with higher median values (57% for GSWP3 and 53% for CRUNCEP) (Fig. 17). ~~This high CUE bias can be expected, as ORCHIDEE-MICT omits the effects of low nutrient availability on CUE (Vicea et al., 2012).~~

### 8.3 Spatial distribution of burned area and fire emissions

The spatial distribution of burned area is largely reproduced by ORCHIDEE-MICT, with a higher fraction of burned area in central Eurasia, and a decrease in burned area toward higher latitudes (Fig. 18). For the GSWP3 simulation, modeled total burned area over the period 1997-2007 (in the unit of  $\text{Mha yr}^{-1}$ ) is smaller than GFED4s for BONA (1.7 vs. 2.2 in GFED4s), and higher for BOEU (8.1 vs. 5.0) and BOAS (16.1 vs. 10.4). Modelled spatial distribution of natural fire emissions has greater discrepancies with respect to evaluation data than burned area, bearing in mind that GFED4s data is based on a biosphere model (CASA), not observations. Higher emissions in eastern Eurasia are reproduced by ORCHIDEE-MICT. However, modelled emissions are overestimated for central Eurasia and underestimated in boreal America, with respect to GFED4s. As a result, regional carbon emissions (in the unit of  $\text{TgC yr}^{-1}$ ) in ORCHIDEE-MICT are lower than in GFED4s for BONA (20 vs. 48), much higher for BOEU (45 vs. 6), and in good agreement for BOAS (104 vs. 111). One possible reason for the discrepancy in BOEU is the lack of forest management and fire suppression measures in ORCHIDEE-MICT, leading to higher simulated burned area and carbon emissions than in GFED4s. Changing the climate forcing from GSWP3 to CRUNCEP yields a smaller burned area (Fig. 18 b and c). Because the reduction of burned area mainly occurs in grassland, the impact on carbon emission is small.

Carbon emissions peak in summer for both GFED4s and ORCHIDEE-MICT for BONA and BOEU (Fig. S69d,e). However, for BONA the fire season starts one month later in ORCHIDEE-MICT (Fig. S69a). For BOAS and BOEU, there are stronger discrepancies in seasonal carbon emission (Fig. S69e,f). In particular, ORCHIDEE-MICT fails to account for the April peak in carbon emissions. A possible explanation for the missing April emissions in ORCHIDEE-MICT may be the late timing of snowmelt (Fig. 9) and the delayed spring increase in LAI (Fig. 13). This would cause an unavailability of fuel in spring time.

Changing the climate forcing from GSWP3 to CRUNCEP has only a very small effect on burned area and carbon emission seasonality.

#### 8.4 Seasonal cycle of NEE

5 Monthly NEE from inversions, originally provided at the spatial resolution of each transport model, was aggregated at the scale of the 3 high-latitude sub-regions (Fig. 2a). Spatially averaged NEE is expected to be more consistent between different inversions at coarser spatial scale, given the sparseness of atmospheric CO<sub>2</sub> stations and differences in transport models. Nevertheless, the seasonal cycle of NEE is generally consistently estimated by the two inversions in each sub-region, although Jena CarboScope estimates generally higher seasonal NEE amplitude, and in BOEU NEE from CAMS peaks one month earlier  
10 than Jena CarboScope (Fig. 19). Simulated seasonal NEE (defined as GPP minus autotrophic and heterotrophic respiration, fire emissions and emissions from LUC) magnitude and evolution is very similar in both simulations and, in general, ORCHIDEE-MICT was able to reproduce the timing and magnitude of the transition between winter release and spring NEE uptake, despite a later onset of spring uptake for all three boreal sub-regions, and a smaller peak summer NEE for BOAS.

Since the timing of NEE uptake in Spring and release in Autumn is well constrained in the inversions from the observed  
15 periodical drawdown and buildup of CO<sub>2</sub> at atmospheric stations, the modeled delayed onset of Spring uptake is consistent with the ~1-month lag between ORCHIDEE-simulated and satellite-observed LAI in Fig. 13 (and ET in Fig. 11a) in the basins of the three sub-regions. Even though GPP is overestimated in BONA during spring and summer (Fig. 14), the resulting NEE gives a slightly weaker uptake.

Modeled autumn and winter emissions (positive NEE) from soil respiration are well reproduced in the BOEU sub-region,  
20 even though the release is overestimated in autumn and underestimated in winter. The underestimation of NEE from ~~November~~October to March in BONA and BOAS suggests that ~~wintertime~~cold season respiration in soils is underestimated in the model. This may be because: (i) decomposition in the model is cut off at T = -1°C, whereas observations suggest it can be sustained well below the freezing point in liquid films of soil pores (Schaefer and Jafarov, 2016); and (ii) insufficient snow insulation of soils (see Fig. 6); and (iii) ~~the lack of the carbon cycle of mosses and lichens which could have respiration under winter low~~  
25 ~~temperatures~~ (Atanasiu, 1971). This suggests that we should either allow decomposition below the freezing point, perhaps to account for heat produced in the soil by microbial decomposition (Zimov et al., 1993; Hollesen et al., 2015), improve the snow insulating, ~~or incorporate an additional organic layer of insulating topsoil (e.g. mosses, O-horizons observed in boreal forests, see O'Donnell et al., 2014) into the soil module.~~prescribe an organic layer of insulating topsoil (e.g. mosses, O-horizons observed in boreal forests, see O'Donnell et al., 2011) into the thermal module, or explicitly represent the moss/lichen plants  
30 including their carbon cycle and physical effects (Porada et al., 2016; Druel et al., 2017).

NEE discrepancies may be related to the different seasonal contributions of evergreen and deciduous trees to GPP and LAI (Fig. S710), as well as to the relative sensitivity of heterotrophic respiration, autotrophic respiration and disturbances to the warming (cooling) at the beginning of the growing season (winter).

## 9 Evaluation of carbon stocks

### 9.1 Biomass

Consistent with the LAI and GPP results, the simulation forced by CRUNCEP shows higher forest biomass carbon densities than the GSWP3 simulation (Fig. 20). The two simulations, however, exhibit similar patterns, which reproduce the general spatial pattern of observed biomass, being higher in western and eastern regions of North America, with a declining gradient across Eurasia from the west to the east. However, in general, the model tends to overestimate carbon density in regions of observed high carbon density (e.g., Northwestern Europe and European Russia, eastern North America, Korean Peninsula and Japan). The model also misplaced the region of the highest biomass density in Northwestern Europe and European Russia rather than Central Europe as shown by the observation data. Likewise, the model estimates extremely large biomass in eastern North America, especially by CRUNCEP forcing, which almost doubles the observed amount. For the rest of the study region, simulated carbon density is slightly lower than observations, by around 5–25 MgCha<sup>-1</sup>. The two observation data sets show considerable similarities.

Both model output and observation data are subject to the spatial uncertainties introduced by the use of satellite-derived land cover maps. We thus used the forest cover map as prescribed in the model for both data sets, and calculated latitudinal averages to compare with model results (Fig. S8). GSWP3-forced model output agrees well with observations averaged over the whole study region, while CRUNCEP-forced output overestimates biomass at all latitudes (Fig. S8a). For the sub-regions, the overestimation of biomass in BONA is consistent with that of GPP (Fig. 14), while biomass is more overestimated in BOEU compared with GPP, probably because of the lack of forest management and forest age structure for Europe. Comparing the two model results, CRUNCEP-forced biomass is much higher than GSWP3-forced biomass, which cannot solely be explained by the higher GPP by CRUNCEP (Fig. 14), indicating a bias in the allocation scheme in ORCHIDEE. As detailed in Krinner et al. (2005), the photosynthates are partitioned into leaves, roots, sapwood, and carbohydrate reserve, dependent on soil moisture etc. If LAI is above a PFT-specific maximum value, carbon will not be allocated to leaves but to the sapwood, which slowly converts to heartwood. Therefore, a higher LAI forced by CRUNCEP leads to more carbon allocation to the wood for some areas, the turnover time of which is much longer than leaves. A new allocation scheme based on the pipe model was implemented in another branch of ORCHIDEE (Naudts et al., 2015), which provides a physiologically meaningful relationship between foliage, roots, and wood. This would be incorporated in ORCHIDEE-MICT in the future developments. When latitudinal distributions of total forest biomass are examined by using forest area distribution as prescribed in the model input for all data sets, strong agreement is found between observed data and GSWP3-forced model output (Fig. S5a) for the whole study region. CRUNCEP-forced model output shows much higher biomass at all latitudes. For BOEU, the model estimated higher biomass observations for regions above 55°N (Fig. S5e), whereas for BOAS, model output fell between the observation datasets at latitudes of >50°N (Fig. S5b). The overestimation of biomass in BONA is consistent with that of GPP (Fig. 14), while biomass is more overestimated in BOEU compared with GPP, probably because of simulated longer biomass turnover and lack of forest management and forest age structure. Over the whole study region, ORCHIDEE-MICT-simulated a total forest biomass carbon stock for the whole domain of 95 PgC under GSWP3 forcing (165 PgC under CRUNCEP), close

to estimates from forest inventory data in Pan et al. (2011) of 92.1 PgC. Somewhat lower estimates are derived by Avitabile et al. (2016) and Thurner et al. (2014), of 73 and 84 PgC, respectively.

## 9.2 Soil carbon

5 SOC stocks simulated by the model fit [to some extent](#) the spatial pattern from observed inventory data, including that of NCSCD's permafrost region near-surface SOC density (Fig. 21), but generate lower values than those from ISRIC in BONA and BOAS. ORCHIDEE-MICT tends to underestimate SOC density in deep soils below ~1m. This underestimation is maybe because the model does not include the sedimentation of soils that characterized peat formation during the Holocene, nor simulation of loess carbon or yedoma deposits during the last glacial period (Zimov et al., 2009; Zhu et al., 2016). Further, spin  
10 up was performed with the coarse approximating assumption that Holocene climate is equal to recent climate. Nevertheless, it is important to note that ISRIC and NCSCD products also have important dissimilarities in overlapping regions suggesting that high systematic uncertainties exist in boreal and Arctic SOC inventories. We also observe that modeled SOC stocks are rather dependent on climate forcing inputs, particularly for BONA and BOAS (Fig. 21). Higher stocks are produced under CRUNCEP forcing, mainly due to the higher primary productivity and therefore higher inputs to SOC associated with this  
15 forcing data.

## 10 Discussion

The model performances against datasets being documented in previous sections, we focus here on key mechanisms expected from a “high-latitude” model: 1) the conversion of winter snow and ice storage into a peak river discharge in spring, 2) the ability to capture the rectification of the seasonal amplitude of air temperature through the insulating snowpack and its  
20 attenuation at depth in the soil profile, and 3) the ability to reproduce the large scale gradients of carbon input to ecosystems from GPP and NPP and its further partitioning into live biomass and SOC pools driving soil heterotrophic respiration.

### 10.1 Conversion of winter water storage to spring and summer river discharge

We have shown that the model simulated very different river discharge values according to the atmospheric forcing used. The spatial distribution estimate of winter snowfall in high latitudes in climate datasets is very difficult. The first reason is  
25 the low density of meteorological stations, in particular in the CRU data (Burke et al., 2013), which leads to lower total precipitation in winter and spring (November to April) in CRUNCEP compared to GSWP3 forcing (Fig. S14). Another well known reason is the difficulty to catch snowfall in a gauge, because of its lower density; the wind prevents the snowfall from being vertical, leading to systematic undercatch (Yang et al., 2005). Then, the way the total precipitation is partitioned into rainfall and snowfall in the atmospheric forcing can also change the SWE in winter and spring. The threshold of 0°C in 2-m  
30 temperature, used to partition the precipitation in the CRUNCEP forcing, can lead to very different results compared to the physical partitioning performed within the dynamical downscaling for the GSWP3 forcing. In the Yukon and Kolyma basins, the proportion of snowfall to total precipitation in winter and spring being 100% in both forcings, the precipitation partitioning



has no influence in these basins. By contrast, the proportion of snowfall to total precipitation in winter and spring largely differs between the two forcings in the other basins, in particular in the Volga (77% in GSWP3 and 53% in CRUNCEP), Ob (84% and 72%), Yenisei (87% and 80%) and Lena (96% and 89%) basins.

5 We can assume that most of the forcing-related bias in SWE is converted into a river discharge bias in spring. Regressing the spring (April to June) river discharge bias for each forcing against the bias of SWE during the pre-melt season (February to March), over the same period of spring discharge and pre-melt season than Rawlins et al. (2007), shows that across the seven basins, 69% of the spatial variance of the bias of the spring flow is explained by the SWE bias during the pre-melt season (Fig. 22). With GSWP3 forcing, the positive ratio of pre-melt SWE bias to spring flow bias ranges between 7 and 97% in  
10 the different basins (Table S5). With the CRUNCEP forcing, this range is from 9 to 34%. In some basins such as the Ob or the Mackenzie with CRUNCEP or the Lena with GSWP3, the errors in spring discharge cannot be evidently related to the SWE bias (negative ratio). This can be explained by the periods chosen for spring and pre-melt which are not necessarily the same for all basins, in particular in the Ob, Mackenzie and Lena, the rivers for which the advance of simulated peak flow is the largest. The higher sublimation in the end of winter with CRUNCEP compared to GSWP3 (Fig. 11b), probably related  
15 to higher downward shortwave radiation and lower specific humidity in CRUNCEP (Figs. S125 and S147 respectively), also contributes to the reduced spring discharge between the two simulations (Fig. 12). Over the Lena basin, higher soil evaporation together with higher interception loss (Fig. 11b), which is coherent with higher LAI (Fig. 13), lead to higher ET in summer with CRUNCEP (Fig. 11a), which reduces the spring discharge compared to GSWP3 (Fig. 12). The excessive persistence of the snow cover can explain the delay of the LAI increase occurrence in the simulations, which may in turn contribute to too early  
20 simulated peak flow, also coupled to floodplain buffering. The Yukon, Mackenzie, Ob and Lena rivers have large floodplain areas and wetlands (and, in the Yukon and Mackenzie basins, lakes that enhance the floodplain buffer) that retain snowmelt water (Ringeval et al., 2012), and should attenuate the peak discharge in spring and sustain a significant summer discharge. The ORCHIDEE-MICT model uses a predefined floodplain map from GLWD (Lehner and Döll, 2004), but flooded area data products (e.g. Prigent et al., 2016) show that wider areas are flooded in these catchments in spring, suggesting that better  
25 discharge could be obtained by connecting these larger floodplains and wetlands to the river routing scheme. This highlights complex potential interactions between the snow, vegetation, and river discharge dynamics, the overall results of which remain highly uncertain.

## 10.2 Seasonal rectification of soil temperature in the atmosphere-snow-soil continuum

A realistic modeling of soil temperature in cold regions requires not only a good soil thermal scheme, but also a good parameterization of snow insulation processes which strongly modulate soil temperatures during winter, as well as a realistic soil-plant hydrology which affects heat transfer in the air-soil interface during summer through ET, and in spring and autumn through latent heat uptake and release. Therefore, a better understanding and evaluation of model behaviors call for an integrated examination of all these elements.

-In this study, we showed that biases in modeled winter soil temperatures (Fig. 5) are connected with biases in snow depth (Fig. 3), while error compensations seem to exist in terms of snow depth and snow thermal conductivity. Model forced by



the GSWP3 climate forcing, which has higher snowfalls than CRUNCEP forcing (Fig. S103), produces significantly larger snow depth and warmer winter soil temperatures than by CRUNCEP. However, the diagnosis from the relationship between  $\Delta T$  and snow depth, which reveals the model's intrinsic parameterization of snow insulation irrespective of snow depth, shows similar patterns with both GSWP3 and CRUNCEP forcings. The model can generally capture the broad characters in the  $\Delta T$ -snow depth relationship, including the different  $\Delta T$  regimes for different periods of the snow season, yet underestimates the insulation given the same snow depth, compared to observations. This indicates an overestimation of effective thermal conductivity of snow. In the snow module of ORCHIDEE-MICT, snow compaction is parameterized such that snow density increases due to the weight of the overlying snow, while the top layer of snow keeps relatively low-density due to fresh snowfall; thermal conductivity of snow then evolves with time and depth, calculated as a non-linear function of snow density (Wang et al., 2013). Recent field measurements for Arctic snow, however, show that the formation of a thin layer of soft depth hoar at the base of snowpack, with its low thermal conductivity around  $0.025 \text{ W m}^{-1} \text{ K}^{-1}$  which is ten times smaller than the intermediate snow layers, significantly re-shapes the vertical profile of snow thermal conductivity (Domine et al., 2016). It is therefore important to account for such complex metamorphic conditions of snowpack, especially in Arctic/sub-Arctic regions, in order to better model the soil thermal regime in permafrost regions.

Soil temperature during the thawing season is a key driver for microbial decomposition of soil organic carbon in the high latitudes; hence its realistic representation is crucial in modeling carbon cycle in permafrost regions. ORCHIDEE-MICT produces reasonable summer-time soil temperature and ALT in general, but significant discrepancies exist between the results of the model forced by the two climate forcings (Figs. 5 and 7). As the monthly mean gridded air temperature of GSWP3 and CRUNCEP forcings are almost identical (Fig. S912), both coming from the CRU data, the differences in the two results are mainly induced by other meteorological fields. A model intercomparison study (Peng et al., 2016) has reported that the surface incident longwave radiation is one of the dominant drivers of soil temperature trends in LSMs. Indeed, GSWP3 forcing has a systematically higher downward longwave radiation than CRUNCEP (Fig. S136), partly explaining the higher land skin and soil temperatures and deeper ALT with GSWP3 than with CRUNCEP forcing (Figs. 4, 5 and 7). What's more, the interactions with hydrology and vegetation also play an important role. The higher ET in Siberian permafrost regions with CRUNCEP than with GSWP3 forcing (Fig. 11a), which is probably driven by the higher LAI in the same regions (Fig. 13), leads to more cooling for the land surface through latent heat release. Besides, the strong cold bias in winter soil temperatures with CRUNCEP forcing affects summer temperature through thermal inertia, especially for the deeper soils below topsoil. All these elements are interconnected, again highlighting the importance of synthetic evaluation of different yet related model aspects to improve our understanding of the system.

Previous land surface modeling studies have shown the critical role of organic matter in soil thermodynamics in permafrost regions (e.g. Lawrence et al., 2008; Chadburn et al., 2015b), while different parameterizations of such effects are implemented in different models. Most of the recent models, like CLM (Lawrence et al., 2008), JULES (Chadburn et al., 2015a), ISBA (Decharme et al., 2016), and ORCHIDEE-MICT in this study, assume weighted combinations of organic soil and mineral soil in the calculation of soil physical parameters for each soil layer in the model. This structure is more flexible than a fixed thickness of organic layer or moss layer as the implementation in JSBACH (Ekici et al., 2014), since the former could

approximate the latter by assuming 100% organic soil above the prescribed thickness. Note that for the insulating effect of moss/lichen layer, the same values of thermal properties to that of the organic soil are usually used in recent models (Chadburn et al., 2015a; Porada et al., 2016). In this study, however, we did not apply a fixed moss layer in the thermal module for the regional simulations, due to the lack of a gridded map for moss/lichen ground covers especially on the boreal forest floor, and to the lack of a representation for dynamic moss/lichen coverage as in JULES (Chadburn et al., 2015a) and JSBACH (Porada et al., 2016). This could partly explain the regionally overestimated ALT compared to the empirical map for Yakutia (Fig. S5). An explicit representation for non-vascular plants in ORCHIDEE (Druel et al., 2017) has been worked in parallel with this study at the moment, but would be incorporated in ORCHIDEE-MICT in the future developments.

### 10.3 Ability of ORCHIDEE-MICT to simulate northern land carbon fluxes and pools

Figure 23 shows the land carbon fluxes and pools modeled and derived from observations (datasets used in the text) in the model domain  $> 30^{\circ}\text{N}$ . Modeled estimates encompass the range of observations excepted for the deep SOC pool where the model underestimates SOC. The same figures for each northern region are given in Fig. S811. At the regional level, model-observed agreement still holds.

Arguably, the main variables that should be well simulated for a high-latitude land ecosystem model regarding future carbon-cycle climate feedbacks (Koven et al., 2013; Burke et al., 2017a) are the carbon stocks in biomass, litter and soils. In this respect, ORCHIDEE-MICT performs generally well for biomass with GSWP3 forcing, including the latitudinal profile, showing a peak between  $50$  and  $60^{\circ}\text{N}$ , despite the simple constant background mortality used for forest (Krinner et al., 2005) in a region where harsh climate is known to induce mortality events. Note that the DGVM version of ORCHIDEE-MICT has a climate and PFT dependent mortality (Zhu et al., 2015) which was adjusted to reproduce the distribution of vegetation types, but it was not activated in this study aiming to reproduce observed stocks from an observed vegetation map. The inclusion of climate dependent fires adds another factor of mortality in this study. From simulated burned area and fire induced biomass suppression, we calculated that fires add a mortality of  $0.4\% \text{ yr}^{-1}$  of biomass in BONA,  $0.3\% \text{ yr}^{-1}$  in BONEU and  $0.6\% \text{ yr}^{-1}$  in BOAS, compared to the fixed background mortality rates of  $1.25\%$ . Thus fire is a non negligible component of biomass mortality, controlling the modeled turnover of forest biomass (Thurner et al., 2016). Interestingly, ORCHIDEE-MICT can reproduce the latitudinal gradient of biomass stocks well, even in warmer regions where NPP is lower than observed (compare Figs. 20 and 15b). This suggests a possible error compensation with too small mortality and NPP in the southern boreal forests. Lack of representation of forest management, and under-representation of other natural disturbances (insect, wind, etc.) could collectively contribute to an overall underestimation of biomass turnover. Carvalhais et al. (2014) reported an average turnover of 53.5 years for boreal forests, using extrapolated global soil database and satellite-derived biomass and GPP estimations. Their estimation of the turnover time integrates both biomass and soil C. Given a longer soil carbon turnover than biomass, the biomass turnover of boreal forests should be smaller than 53.5 years. While ORCHIDEE-derived turnover is between 54 (adding together fire and background mortality) and 80 years (assuming only a  $1.25\%$  background mortality) with a lower boundary close to Carvalhais et al. (2014), given that forest fires impact only a small fraction of forests each year ( $0.1\text{--}3\%$  on annual basis), the ultimate turnover time in the model could be much longer than Carvalhais et al. (2014). The last factor for

overestimation of biomass is that ORCHIDEE represents a mature forest state everywhere due to lack of age structure in the model, whereas in reality disturbances and management have led to forest age younger than a mature one.

Regarding SOC, it is critical for carbon climate feedbacks that models can simulate the large current SOC stocks of the high latitudes. This is a particularly difficult problem since frozen high-latitude SOC was formed during the Pleistocene (Yedoma) or the Holocene (peat and tundra soils) through processes that are not incorporated as routine in current models, despite efforts in this direction, e.g. Zhu et al. (2016) for Yedoma and Kleinen et al. (2012) and Spahni et al. (2011) for Holocene peat deposits. The incorporation of slow forming high-latitude carbon deposits also requires climate history for deeper past periods. In order to reproduce the burial of SOC below the active layer, the strategy followed in ORCHIDEE-MICT is inspired from Koven et al. (2009) who used a diffusion equation to move carbon in the permafrost where no decomposition occurs. Other studies (Koven et al., 2013; Burke et al., 2017a) further limited the rate of decomposition of SOC at depth, to reproduce the lack of oxygen inhibiting decomposition. It should be noted also that the decomposition scheme of SOC is still based on Parton et al. (1988) as classically done in land surface models (Friedlingstein et al., 2006). Different approaches were proposed in models focusing only on SOC decomposition (Wutzler and Reichstein, 2008; Manzoni and Porporato, 2009) based on different assumptions (substrate driven, decomposer driven, etc.), but no clear consensus emerged up to date to revise the SOC decomposition scheme in land surface models (Luo et al., 2016). Here we argue that key to the simulation of a high and realistic SOC stock is the correct representation of soil thermics, illustrated in Fig. 5. A detailed study of the sensitivity of modeled SOC to soil thermics during the spin-up phase will be presented in a follow up study but we already found comparing SOC with CRUNCEP and GSWP3 climate forcings, the latter giving warmer soil temperature profiles, that SOC is indeed lower in GSWP3. Overall, the simulated circumboreal SOC stocks are comparable with NCSD observations (Fig. 21). This suggests that simulating the formation of frozen carbon with a diffusion-related burial process gives a good performance for the model, although the too stiff profile of SOC (Fig. 21) indicates that a further inhibition of respiration in deep horizons could improve the modeled vertical distribution. Other processes that would tend to make the vertical profile of SOC more uniform with depth is the omission of interactions between SOM and the physical soil environment (Doetterl et al., 2015; Tipping et al., 2016). Modeled SOC is too low in regions covered by peat (lowland Hudson Bay, Ob northern and central basin) and Yedoma (Northeastern Siberia and Alaska) (Fig. 21), suggesting that adding these two SOC formation processes should further improve the modeled SOC patterns in longitude, and that in non-peat and non-Yedoma regions, there is no significant model bias for SOC.

Second, for research related to biophysical vegetation-climate feedbacks and understanding the current variability of the carbon cycle, having a correct seasonal representation of GPP, NPP and of the related plant transpiration is a critical requirement for a LSM. The phase of simulated LAI in spring lags satellite observations, in particular for BONA and BOAS sub-regions (Fig. 13). We argued that this lag is related to the late persistence of the snow cover. However a recent work (Chadburn et al., 2017) shows a late onset even in the absence of snow persistence on site simulations by ORCHIDEE. This calls for a revisit of the phenology-related thresholds in the high-latitudes, perhaps by introducing new PFTs (arctic C3 grass and shrub, and non-vascular plants), with their separate set of parameters calibrated, to better represent Arctic vegetation and their phenology (Druel et al., 2017). There is also a lag of GPP (Fig. 14), and of ET (Fig. 11a), but of smaller length (~15 days) compared to

the lag of LAI. In order to further analyze why the lag of GPP (and ET) in spring is shorter than the one of LAI, we represent separately monthly GPP and LAI for deciduous and evergreen PFTs (Fig. S710). The phase of GPP coincides with that of LAI in deciduous, whereas GPP increases in spring as soil thaw faster and earlier than LAI for evergreens. Thus, it is the modeled spring decoupling between GPP and LAI in evergreen forest that explains the phase bias of GPP being smaller than that of LAI. Note that evergreen trees GPP is modeled to increase rapidly in early spring ORCHIDEE-MICT, even if the model has an inhibition of  $V_{cmax}$  when previous monthly air temperature remains below  $-4^{\circ}\text{C}$  (Krinner et al., 2005) to reproduce impaired photosynthesis observed in conifers, a physiological measure to avoid frost damage of photosystems (Leinonen, 1996; Tanja et al., 2003).

Peak summer GPP is overestimated in BONA and BOEU sub-regions compared to MTE-GPP (Fig. 14) and mean GPP is underestimated at southern sites in W. Europe and US where  $T > 7^{\circ}\text{C}$  (Fig. 16a). We checked that underestimated GPP at warmer sites and in Eastern Siberia for the deciduous needle-leaved forest PFT (Larch) is due to an overestimation of water stress in summer by the model, since ET is also lower than observed at all the warmer sites. Without the dynamic root allocation to optimize water use in the root zone (see Section 4.3), GPP in Eastern Siberia would be even lower compared to MTE-GPP (not shown). Last, regarding the timing of GPP decrease in autumn, ORCHIDEE-MICT shows good performances (Fig. 14) despite the lack of an explicit senescence model for conifers.

Interestingly the reasonable seasonal phase of simulated GPP (Fig. 14) can be contrasted with the larger lag of spring NEE uptake compared to inversions results (Fig. 19). ~~This suggests that in the model, ecosystem respiration may peak too early in spring when topsoil thaws, possibly because it lacks a representation of  $\text{CO}_2$  trapping in the soil column before complete thawing of the soil, as observed in a tundra site (Elberling and Brandt, 2003).~~ This may be partly explained by the underestimated GPP at the beginning of the growing season (Fig. 14), and also by a possible too big soil respiration in spring. The moss/lichen surface coverage could be over 70% under the vast boreal forests (Porada et al., 2016), but we did not prescribe an additional moss layer in the regional simulations, which could lead to a too early thawing of the soil in spring.

The connection between modeled GPP and carbon pools takes place through NPP and its allocation to interconnected pools of variable turnovers. Of critical importance is thus the ability of the model to reproduce CUE which defines the ratio of NPP to GPP, i.e. carbon available for ecosystem pools. Piao et al. (2010) showed that CUE of forest decrease from 0.5 to 0.3 between  $10^{\circ}\text{C}$  and  $-10^{\circ}\text{C}$ , due to higher maintenance needs of trees e.g. to recover from cavitation in spring or maintain tissues during the cold-season period. While mean CUE is correctly simulated (Fig. 17), ORCHIDEE-MICT does not capture its observed decrease towards colder temperatures (data not shown), possibly because allocation and NPP are not driven by sink considerations (Körner, 2003; Fatichi et al., 2014) and thus too much GPP is used to make NPP at colder temperatures.

## 11 Conclusions

This study has described the inclusion of parameterizations which link soil carbon content and its decomposition rates with permafrost physics and hydrology in the ORCHIDEE-MICT land surface model. ~~The effects of soil organic matter on soil thermal and hydraulic properties are incorporated.~~ The model was evaluated against temperature gradients between the at-

mosphere and deep soils, and reasonable captured active layer thickness, northern permafrost extent, and soil carbon stocks and profiles. We have shown that the simulated water balance components and their seasonal transition between cold season storage, mostly in solid form, and warm season loss are comparable to observations. Naturally, there remains significant room for improvement. The model appears to underestimate evapotranspiration and overestimate surface temperatures, particularly in the southern portion of the ~~northern~~-boreal zone, ~~particularly over southeastern Siberia~~. Simulated phenology shows generally a delay in the onset of growing season. And the snow module underestimates the thermal insulation of snow. Through the use of two different climate forcing datasets across all diagnostic variables, we found that for a large number of these, the variation in output arising from choice of a forcing was as large as the discrepancy between model and observations. This raises a caution flag against ‘over-calibrating’ the parameters of a LSM to match measured quantities in the presence of large biases in climate input datasets. One critical aspect of forcing data in this respect is the partitioning of precipitation between snowfall and rainfall, which appears to produce a large difference in modeled snow mass and depth. Future improvements of the ORCHIDEE-MICT model include the need for a better floodplain parameterization (Lauerwald et al., 2017) and a better representation of floodplain areas and wetland effects on river discharge. Including an explicit senescence inception model in ORCHIDEE-MICT may contribute to correct the excessive persistence of LAI during the fall, or even winter in the Yukon and Mackenzie basins. The subgrid-scale representation of permafrost hydrology in discontinuous permafrost areas may be necessary, as might a more realistic description of the slow processes that accumulate carbon in the soil (peat, yedoma, erosion) on very long time scales.

## 12 Appendix

### 20 Appendix A: Evaluation datasets for the water budget

#### A1 Total terrestrial water storage

The change of total terrestrial water storage (TWS) is an integrated measure of the ability of a LSM to partition the cold season storage of water as snow and ice, and its decrease from losses to downstream river discharge, plant and soil evapotranspiration, and soil moisture increase. Alkama et al. (2010) showed with the ISBA-TRIP model that seasonal variations in TWS over high-latitude basins resulted primarily from snow accumulation in the cold season, despite a wintertime underestimation of TWS values. Decreasing soil moisture contributed to a small decrease in springtime TWS, while exports with riverflow increased dramatically during the snowmelt period. The GRACE (Gravity Recovery And Climate Experiment) satellite mission permits estimation of monthly TWS variations through measurements of the Earth’s gravitational field. Three solutions, at 1° resolution, based on spherical harmonic coefficients (Release 05), are obtained by different processing centers: CSR (Center for Space Research), JPL (Jet Propulsion Laboratory) and GFZ (GeoForschungsZentrum Potsdam) and provided in the GRACE Tellus dataset (Swenson and Wahr, 2006; Swenson, 2012; Landerer and Swenson, 2012). We use the ensemble mean of these solutions to reduce uncertainty in GRACE data (Sakumura et al., 2014). To compare TWS simulated by ORCHIDEE-MICT to GRACE data over the common time period between the two meteorological forcing data sets and GRACE data (July 2003-December

2007), we summed the water stocks simulated by ORCHIDEE-MICT, i.e., soil moisture, snowpack, water on the canopy and water stored for the routing reservoirs. In each grid cell, the corresponding 5-year average is removed from the 2003–2007 time series of TWS output from ORCHIDEE-MICT. The comparison of simulated TWS with GRACE data over seven basins in this study (Fig. 2b) is statistically satisfactory given large surface areas occupied by these basins ( $> 400,000 \text{ km}^2$  (see Table S4), Swenson et al., 2003).

## **A2 Snow water mass**

The GlobSnow v2.0 SWE dataset is based on a data-assimilation approach combining space-borne passive radiometer data (SMMR, SSM/I and SSMIS) with data from ground-based synoptic weather stations. The record provides SWE information on a daily, weekly and monthly basis, at a spatial resolution of approximately 25 km (in EASE-Grid format) and applies to non-mountainous regions of the northern hemisphere, excluding glaciers and Greenland. The product is based on the SWE retrieval methodology developed by Pulliainen (2006) complemented by a time-series melt-detection algorithm (Takala et al., 2009). A complete description of its methodology is given in Takala et al. (2011). SWE estimates are complemented with uncertainty data at the grid cell scale. This work uses monthly-averages of original daily data, interpolated to a  $1^\circ$  grid for appropriate comparison with ORCHIDEE-MICT output.

## **A3 Evapotranspiration (split into components)**

The Global Land Evaporation Amsterdam Model (GLEAM) (Miralles et al., 2011) estimates the daily terrestrial ET rate and its components (transpiration, bare-soil evaporation, open-water evaporation, interception loss and sublimation), as well as root-zone soil moisture, at a spatial resolution of  $0.25^\circ$ . It is driven by microwave remote sensing observations and uses satellite soil moisture to constrain potential evaporation rate. The latter is computed with the equation of Priestley and Taylor (1972), which is based on air temperature and net radiation. In the GLEAM v3 product (Martens et al., 2017), representation of evaporative stress has been improved by the use of microwave vegetation optical depth as a proxy for vegetation water content. Other improvements include the use of satellite based SWE, reanalysis air temperature and radiation, and a multi-source precipitation product. GLEAM v3 contains three sub-versions: Version v3.0a is an assimilated product derived from satellite observations and multiple reanalysis climate forcing, while v3.0b and v3.0c are purely satellite-based products. As the latter two do not provide full coverage of high-latitude regions (only  $50^\circ\text{S} - 50^\circ\text{N}$ ), the v3.0a, which covers the period 1980–2014, is used in our study. We use monthly-averages of original daily data, interpolated to a  $1^\circ$  grid for appropriate comparison with ORCHIDEE-MICT output.

## **A4 River discharge**

We use two river discharge databases, one global and another exclusively covering Siberia: The Global Runoff Data Centre (GRDC) product is a global database which collected river discharge data from nearly 7800 stations in 156 countries for the [maximum] period 1807–2017. We use one gauging station per river, the closest to the mouth of the river. For Siberia, we also

use daily river discharge at the 3 gauges in three large Siberian rivers (the Ob, Yenisei and Lena) which have been reconstructed (naturalized discharge) to exclude the human impact from the data. This has been performed by using the Hydrograph Routing Model (HRM) developed at the University of New Hampshire (USA) in collaboration with the Arctic and Antarctic Research Institute (Russia) (Shiklomanov and Lammers, 2009). We use monthly-averages of original daily data.

#### **A5 Top-soil moisture**

The global ESA CCI SM product (v2.2) provides daily soil moisture (in volumetric units  $\text{m}^3 \text{m}^{-3}$ ), at a spatial resolution of 25 km with quality flags specifying potential sources of errors linked to the presence of water bodies, dense vegetation, snow or frozen soils in the pixel area. Three datasets are provided based on different combinations of active and passive satellite sensors. In this work, we used the combined active and passive product which covers the whole period from November 1978 to December 2014. The product results from the merging of soil moisture estimations inversed from two types of instruments and two methodologies: passive microwave radiometers (SMMR, SSM/I, TMI, AMSR-E, AMSR2 and WindSat) using the methodology developed by Owe et al. (2008) and active microwave instruments (ERS-1&2 and ASCAT) using the algorithm developed by TU-Wien (Wagner et al., 1999; Bartalis et al., 2007). These two records were first rescaled together and merged with topsoil (10 cm) soil moisture simulations of the GLDAS-NOAH LSM using a CDF-matching approach. Data accuracy was estimated against in-situ measurements (Dorigo et al., 2015) to  $0.05 \text{m}^3 \text{m}^{-3}$ .

#### **A6 Root-zone soil moisture**

The GLEAM root-zone soil moisture v3.0a database (Martens et al., 2017) is based on both satellite observations and reanalysis data. It is a mixture of soil water content of three soil tiles: bare soil (0-10cm), herbaceous (0-100cm) and tall vegetation (0-250cm). The tile fractions are static and derived from MODerate resolution Imaging Spectroradiometer (MODIS) Global Vegetation Continuous Fields product (MOD44B, Hansen et al. 2005). Obviously, it is hard to compare the ORCHIDEE soil moisture with the GLEAM root-zone soil moisture directly not only due to the mismatch of defined soil tile depth but also due to the difference of soil tile fractions. Thus we implement the comparison after normalization of the relative soil moisture values with their respective standard deviation, following Koster et al. (2009).

### **Appendix B: Evaluation datasets for air-to-soil temperature continuum**

#### **B1 Snow depth**

Realistically simulating snow depth is one of the prerequisites for accurately modeling soil thermal dynamics, particularly in winter. Daily snow depth in-situ data (1975-2005) from 524, 72, 528, 128 and 528 stations in Europe, Canada, USA, Russia and China were obtained from European Climate Assessment & Dataset (ECA&D), the National Climate Data and Information Archive of Environment Canada, the United States Historical Climatology Network (USHCN), the Russian Research Institute of Hydrometeorological Information—World Data Center (RIHMI-WDC) (Bulygina et al., 2011), and the National Meteoro-



rological Information Center of the China Meteorological Administration (Peng et al., 2010), respectively. Monthly averages were calculated when more than 2/3 of daily snow depth data were available.

## **B2 Surface soil temperature**

- 5 The soil temperature product used here (André et al., 2015) resulted from the use of combined passive microwave and thermal infrared data to estimate land surface temperature (LST), during summer snow-free periods (snow-covered pixels are masked) in the northern high latitudes. The product is based on the use of SSM/I-SSMIS 37GHz measurements at both vertical and horizontal polarizations and is provided on a 25 km resolution EASE grid at a one-hour time step for the period 2000-2011. LST retrievals are based on the assumption of a relationship between surface emissivities at both polarizations (Royer and
- 10 Poirier, 2010) which was calibrated at pixel scale using cloud-free independent LST data from MODIS instruments. The SSM/I-SSMIS and MODIS data were synchronized by fitting a diurnal cycle model built on skin temperature with reanalysis provided by the European Centre for Medium-Range Weather Forecasts (ECMWF). This product was evaluated at local and circumpolar scales against MODIS LST, and the results show a mean Root Mean Square Error (RMSE) of the order of 2.5 K.

## **B3 In-situ air and soil temperatures**

- 15 Simultaneous measurements of snow depth, near-surface air temperature and top soil (at 20 cm depth) temperature from Russian meteorological stations (Bulygina et al., 2011; Sherstiukov, 2012) are used to evaluate the thermal insulating effect of snow during winter, by comparing the relationships between snow depth and the air-to-topsoil temperature gradient ( $\Delta T$ ) (Wang et al., 2016b). This dataset includes monthly mean values at 268 sites for the period 1980-2000.

## **B4 Active-layer thickness**

- 20 The Circumpolar Active Layer Monitoring (CALM) network's in-situ data set was used to evaluate modeled soil active layer thickness. The CALM network aims to observe the response of near-surface permafrost and the active layer to climate change over long time scales. Thaw depth is measured at the end of the thawing season, so that it should be comparable to the maximum active layer thickness in the model. Before comparing to ORCHIDEE-MICT, the data from 221 sites were averaged from 1990 to 2015.
- 25 In addition, we used the active layer thickness map derived by Beer et al. (2013) over Yakutia at a spatial resolution of 0.5°. It is based on regional surveys of landscapes and permafrost conditions in Yakutia during the time period 1960–1987. The gridded datasets can be accessed at the PANGAEA repository. Values of active layer thickness (ALT) in this region range between 0.3 m north of 70°N to 3 m south of 65°N. Uncertainty increases with ALT and is highest (up to 1m) in the south because of the occurrence of discontinuous and sporadic permafrost landscapes.

## **30 Appendix C: Evaluation datasets for leaf area, carbon stocks and fluxes**

### **C1: Leaf Area Index**



The Leaf Area Index (LAI) can be derived from satellite measurements, using reflectance measurements and land cover maps. Comparing the seasonal cycle of the satellite products with simulated LAI gives information on the appropriateness of the phenology modules. The amplitude of the cycle is also informative with respect to several photosynthetic parameters (e.g. maximum carboxylation rate  $V_{\text{cmax}}$  or specific leaf area).

## **GIMMS**

The Global Inventory Modeling and Mapping Studies (GIMMS) LAI3g dataset is derived from Advanced Very High Resolution Radiometers (AVHRR) measurements. It is more specifically computed from the GIMMS Normalized Difference Vegetation Index (NDVI) 3g, using a neural network and the International Geosphere Biosphere Programme (IGBP) land cover classes (Zhu et al., 2013). The 1/12 degree files cover 30 years from July 1981 to December 2011, at a bi-weekly resolution. They were aggregated at a 1.0° spatial resolution on the model grid, to enable comparisons of the time-series and maps.

## **GLASS**

The Global Land Surface Products (GLASS) LAI product (Liang and Xiao, 2012) is built from AVHRR reflectances over the period 1982-1999 and then from MODIS data over 2000-2012 (Collection 5). The processing is also based on a neural network and the 8 biomes of the MODIS land cover type 3 (Xiao et al., 2014). The files are available at a spatial resolution of 0.05° with an 8-day frequency and were aggregated at 1.0° spatial resolution.

## **C2: NEE from atmospheric inversions**

We use estimates of NEE (here the net land-atmosphere  $\text{CO}_2$  fluxes excluding fossil fuel  $\text{CO}_2$  emissions) from two atmospheric inversions where NEE is optimized at the resolution of a global atmospheric transport model each month to match observed  $\text{CO}_2$  concentration gradients from a global network of ground based stations. The inversions are the Copernicus Atmosphere Monitoring Service (CAMS) inversion system (Chevallier et al., 2010) and the Jena CarboScope (Rödenbeck, 2005).

The inversion from CAMS uses atmospheric  $\text{CO}_2$  concentration observations from a total of 132 sites covering the period 1979-2015 and includes all sites being added to the network through time, combined with the LMDZ INCA atmospheric transport model and prior information about fossil-fuel  $\text{CO}_2$  emissions (from EDGAR3.2 FastTrack 2000 database) as well as land/ocean-atmosphere fluxes, to minimize a cost function in order to calculate NEE. Here we use the latest version (15r4), in which monthly fluxes are calculated at 1.875° x 3.75° lat/lon resolution.

The Jena CarboScope atmospheric inversion uses the same set of atmospheric  $\text{CO}_2$  observation sites available during the temporal observation period (1996-2015) referred to as Jena s96. Jena CarboScope inversion uses fossil fuel emissions from EDGAR 4.2 and the TM3 global atmospheric transport models. Here we use monthly NEE from the latest version of Jena s96 (v3.8), which is provided at 3.8° x 5.0° lat-lon resolution.

### C3: Site-level GPP and NPP observations

#### In-situ GPP and NPP measurements

While there are several NPP datasets especially for forest (e.g. Cannel et al., 1992; Olson et al., 2001; Michaletz et al., 2014), few provide both GPP and NPP at the same locations. We used a recent database of in situ co-located GPP and NPP measurements (Capioli et al., 2015), which is an extension of the Luyssaert et al. (2007) database for forests. The database consists of 131 sites, for which annual NPP obtained from biometric measurements and associated uncertainties are provided. The criteria employed by Capioli et al. (2015) for selecting the sites was the availability of methodologically independent and site-specific estimates of NPP and GPP. Carbon Use Efficiency (CUE) was calculated at each site by taking the ratio of NPP to GPP. Site-level biomes are classified as one of tundra, boreal peatlands, marshes, forests, grasslands or croplands. To study the temperature sensitivity of GPP and NPP, we selected only sites northward of 30°N, where model grid cells were representative of both the vegetation type and meteorological conditions observed at each site. We thus restricted the selection to sites whose IGBP vegetation type had a corresponding fraction greater than 0.5 in the grid cell of ISLSCP II MODIS IGBP Land Cover product (Friedl et al., 2010). Site-years where the absolute difference between local and the Mean Annual Temperature (MAT) of the GSWP3 or the CRUNCEP climate forcing fields was higher than 4 degrees were discarded. As no information was given about the time-period of site NPP and GPP observations in Capioli et al. (2015), we compared them to mean values of the model over the period 2000-2007. The subset of selected sites consisted then in 52 sites out of 131 in the full dataset.

### C4: Gridded GPP and NPP observation-based data

#### GPP

In addition to GPP from sites, we used the monthly gridded GPP observation-based product Model Ensemble Tree GPP (MTE-GPP) from Jung et al. (2009, 2011) from year 1982 to 2010. This product is the result of a statistical model that combines measurements at FLUXNET sites with geospatial information from satellite remote sensing (fAPAR) and meteorological data. MTE-GPP is obtained as the median from an ensemble of the 25 best model trees out of an initial set of 1000 and the Mean Absolute Deviation (MAD) of the ensemble is used to define the uncertainty of MTE-GPP, which is also provided as a monthly gridded product. This uncertainty estimate was mentioned by Jung et al. (2011) to be underestimated as compared to the true RMSE. That is why we consider the RMSE from the cross-validation sites as an estimate of uncertainty. The RMSE of mean MTE-GPP at cross validation sites is  $270 \text{ gC m}^{-2} \text{ yr}^{-1}$ . At global scale and over the period 1982-2008, the mean MTE-GPP is  $933 \pm 46 \text{ gC m}^{-2} \text{ yr}^{-1}$  and the total is  $119 \pm 6 \text{ GtC yr}^{-1}$ , the uncertainties being derived from the MAD of the MTE (Jung et al., 2011).

#### NPP

We use the MOD17A3.005 global annual NPP model driven by satellite observations from the NASA Land Processes Distributed Active Archive Center, at 1 km resolution and aggregated to the 1° grid of the model over the period 2000-2010. Yearly

MODIS NPP is computed from yearly and daily components. Among the daily components, MODIS GPP is based upon a LUE model:

$$GPP = E * fAPAR * PAR \quad (15)$$

5

where PAR is the Photosynthetically Active Radiation, fAPAR the fraction of absorbed PAR and E is the radiation conversion efficiency. E depends on vegetation type and on meteorological conditions (temperature, Vapor Pressure Deficit (VPD)). PAR, temperature and VPD used in MOD17A3.005-NPP are from the NASA Data Assimilation Office (DAO) and fAPAR is a MODIS product (Knyazikhin et al., 1998). A daily Maintenance Respiration (MR) was computed for leaves (l) and fine roots (fr), using the LAI MODIS product (Knyazikhin et al., 1998). On a yearly basis, MR was computed for livewood (lw) and Growth Respiration (GR) for leaves, fine roots, livewood and deadwood (dw). These respirations were computed using a biome-specific parameter look-up table, derived from the BIOME-BGC terrestrial biosphere model (Heinsch et al., 2003). This set of parameters was recalibrated for the MOD17A3.005-NPP collection 5 (Zhao et al., 2005). Yearly MODIS-NPP was thus computed as follows (Running et al., 2004):

10

$$NPP = \left\{ \sum_{\text{days}} [GPP - MR(l) - MR(fr)] \right\} - MR(lw) - GR(l) - GR(fr) - GR(lw) - GR(dw) \quad (16)$$

Turner et al. (2006) evaluated MODIS NPP at nine sites representing various biomes, using local meteorological, LAI and above ground NPP measurements as inputs for the BIOME-BGC model to derive NPP over a 25 km<sup>2</sup> area. They showed globally no bias but NPP was underestimated at the most productive sites (attributed to a negative bias of E) and overestimated at low productivity sites (attributed to a positive bias of MODIS fAPAR). Zhao et al. (2005) evaluated MODIS NPP against the Ecosystem Model-Data Intercomparison (EMDI) global database of NPP measurements (Olson et al., 2001) and found a good agreement except for tropical forests (spatial R<sup>2</sup> = 0.77). Zhao et al. (2006) estimated the sensitivity of the MODIS NPP product to the use of different meteorological reanalysis input data, showing differences at global scale up to more than 20PgCyr<sup>-1</sup>, with the largest differences in the tropics.

20

## 25 C5: Soil carbon inventories

### NCSCD

The Northern Circumpolar Soil Carbon Database quantifies storage of organic carbon in soils of the northern circumpolar permafrost region, comprising four depths up to 3m (0-30cm, 30-100cm, 100-200cm and 200-300 cm) (Hugelius et al., 2013).

Total SOC storage in the 0-2m and 0-3m depth ranges in northern permafrost soils are estimated to be  $827 \pm 108$  and  $1035 \pm 150$  PgC respectively.

## ISRIC

- 5 ISRIC is a global soil information database with spatial predictions for a selection of soil properties, at 1 km resolution and six depths, up to a maximum of 2m (0-5cm, 5-15cm, 15-30cm, 30-60cm, 60-100cm and 100-200cm). About 110,000 soil profiles were used to generate the product, using statistical models based on climatic and biomass indices, lithology, and taxonomic mapping units (Hengl et al., 2014). The gridded maps of soil organic carbon mass fraction, soil bulk density, and volumetric fraction of coarse fragments was used to calculate SOC density in the unit of  $\text{kgC m}^{-2}$  to be comparable with model output (see
- 10 Eq. 6 in Hengl et al., 2014). However, we found in this product a systematic overestimation of bulk density for organic-rich soils, which is similar to the issue found in the HWSD database (Köchy et al., 2015). Therefore, we followed the adjusting method in Köchy et al. (2015) to correct bulk density for histosols and other soils with organic carbon mass fraction larger than 3%. This adjustment decreases total SOC stock in northern permafrost regions (the same domain as NCSCD) in the 0-2m depth range from the original 1724 PgC to 1177 PgC, which we considered acceptable though higher than NCSCD.

## 15 C6: Biomass carbon stocks

- Two forest biomass datasets are used to evaluate simulated forest biomass. The first is that from Avitabile et al. (2016), which, at a spatial resolution of  $0.01^\circ$ , merges two tropical aboveground forest biomass (AGB) datasets from Saatchi et al. (2011) and Baccini et al. (2012) with northern hemisphere volumetric forest stock growth data from Santoro et al. (2015). The second forest biomass dataset from Thurner et al. (2014) is given at a spatial resolution of  $0.01^\circ$  for 2010 and confined to the northern
- 20 hemisphere ( $30^\circ\text{N}$ – $80^\circ\text{N}$ ). This is derived from the Santoro et al. (2011) forest standing stock data. The two datasets were re-gridded to  $1^\circ$  resolution for comparison with model output. The Avitabile et al. (2016) AGB was converted into total biomass, assuming a constant AGB to total biomass ratio of 0.8 and divided by two to obtain total biomass carbon, under the assumption that the carbon content of dry biomass is 50%. Liu et al. (2015) synthesized ratios of aboveground to total biomass for different forest biomes in temperate and boreal regions and found that they range between 0.76 and 0.84 according to regional forest
- 25 inventory assessments. Our uniform 0.8 factor for converting the Avitabile et al. (2016) data thus yields a potential error of  $\sim 5\%$  in inferred total biomass carbon, which is within the reported data uncertainties given by Avitabile et al. (2016). Simulated equivalent total forest biomass carbon over the period 2000–2007 was compared with observations.

## C7: Burned area and fire emissions

- We compared the simulated burned area and fire carbon emission with the Global Fire Emissions Database (GFED4s) data set,
- 30 which is based on the GFED modeling framework (van der Werf et al., 2010). The GFED4s burned area data is derived from the MODIS sensor data that is then complemented by specific algorithms to retrieve “small fires” (Randerson et al., 2012). Emissions are computed from a combination of burned area with the revised Carnegie-Ames-Stanford-Approach biogeochem-

ical model (CASA-GFED, described in van der Werf et al., 2010), that provides an estimation of fuel loads and combustion completeness. GFED4s data is provided at a 0.25° spatial resolution with a monthly time-step for 1997–2015.

Cropland fires are not yet implemented in the ORCHIDEE-MICT model, but are included in the GFED4s dataset. In order to compare the simulation with GFED4s, it was necessary to correct the simulated burned area for the omission of cropland fires. To do so, we computed for each grid cell the fraction of existing natural PFTs (i.e. all non-cropland PFTs), and divided the simulated burned area in each grid cell by this value. For carbon emissions, the GFED4s dataset provides separate emissions data for cropland and natural fires. In this way, we were able to simply remove cropland fire emissions from GFED for comparison with ORCHIDEE-MICT output.

## 10 Author contributions

M. Guimberteau and D. Zhu led the study. D. Zhu performed the simulations. F. Maignan and A. Jornet-Puig performed the code versioning of ORCHIDEE-MICT; F. Maignan managed the scientific aspects of the updates from the main branch (TRUNK); A. Jornet-Puig optimized the code; they both assisted with technical aspects of the model and its run environment. H. Kim built the GSWP3 dataset and J. Chang adapted its format to be compatible with ORCHIDEE-MICT. D. Zhu developed SOC-dependant soil physical properties in the model and introduced a reformulation of the hydric stress above the permafrost table. She conducted the evaluation of the soil temperature and permafrost area simulated by ORCHIDEE-MICT. Y. Huang provided the comparison of the snow depth with dataset. C. Ottlé and S. Dantec-Nédélec processed the satellite datasets and interpreted the soil moisture, land surface temperature and albedo results. A. Ducharne contributed to the improvement of the hydrological modeling in the initial ORCHIDEE version and M. Guimberteau led the water budget evaluation. Z. Yin contributed to the comparison of soil moisture with GLEAM. M. Guimberteau, R. Lauerwald and S. Bowring supervised the routing scheme. C. Yue built an updated version of SPITFIRE in ORCHIDEE-MICT and, with P. Laurent, analyzed the results in biomass carbon stocks, burned area and fire emissions. A. Bastos provided the comparison of NEE with atmospheric inversions. F. Maignan performed the NPP and GPP evaluations over the sites database. B. Guenet, M. Tifafi and D. Zhu conducted the soil carbon evaluation. Y. Wang synthesized the carbon budget evaluation. C. Ottlé, T. Wang, S. Peng and G. Krinner supervised the representation of the high-latitude processes in ORCHIDEE-MICT in coherence with the initial version of ORCHIDEE. D. Goll, X. Wang, E. Joetzjer, C. Qiu and F. Wang assisted in ensuring consistency between the carbon cycle, water cycle and various other processes in ORCHIDEE. P. Ciais contributed to the theoretical concept of the modeling of high-latitude processes. All authors contributed to the writing of the manuscript.

## Competing interests

30 The authors declare that they have no conflict of interest.

## Code availability

~~The SVN version of the code branch is <https://forge.ipsl.jussieu.fr/orchidee/browser/branches/ORCHIDEE-MICT> revision 4126 from the 28th February 2017. Please contact the corresponding author for the code of the ORCHIDEE-MICT if you plan an application of the model and envisage longer-term scientific collaboration.~~The source code for ORCHIDEE-MICT

- 5 version 8.4.1 is available online but its access is restricted. Consequently, it is required to communicate to the corresponding author for a username and password. The source code can be found at the following address: [https://forge.ipsl.jussieu.fr/orchidee/browser/branches/ORCHIDEE-MICT/tags/ORCHIDEE\\_MICT\\_8.4.1](https://forge.ipsl.jussieu.fr/orchidee/browser/branches/ORCHIDEE-MICT/tags/ORCHIDEE_MICT_8.4.1)

## Data license

This software is governed by the CeCILL license under French law and abiding by the rules of distribution of free software.

- 10 You can use, modify and/ or redistribute the software under the terms of the CeCILL license as circulated by CEA, CNRS and INRIA at the following URL: <http://www.cecill.info>.

## Data availability

Primary data and scripts used in the analysis and other supplementary information that may be useful in reproducing the author's work can be obtained by contacting the corresponding author.

- 15 *Acknowledgements.* This work was financially supported by the European Research Council Synergy grant ERC-2013-SyG-610028 IMBALANCE-P. We acknowledge the ORCHIDEE-PROJET group, which developed and tested the ORCHIDEE model revision 3976. We thank Natasha MacBean who contributed to the production of the land cover map. We thank Arsène Druel who processed the GLASS LAI data. Simulations with ORCHIDEE-MICT were performed using computational facilities of the Institut du Développement et des Ressources en Informatique Scientifique (IDRIS, CNRS, France) and of the CCRT (CEA). S. Dantec-Nédélec, G. Krinner, P. Ciais and C. Ottlé were supported by
- 20 the CLASSIQUE national project (ANR Grant No. ANR 2010-CEPL-012-02) and the PAGE 21 project (grant agreement number 282700, funded by the EC seventh Framework Programme theme FP7-ENV-2011). A. Ducharne was supported by the IGEM project 'Impact of Groundwater in Earth system Models' (ANR Grant No. ANR-14-CE01-0018-01). Discharge station data is kindly provided by the Global Runoff Data Centre, 56068 Koblenz, Germany. GRACE land are available at <http://grace.jpl.nasa.gov>, supported by the NASA MEaSUREs Program. The MOD17A3.005 data product was retrieved from the online Data Pool, courtesy of the NASA EOSDIS Land Processes Distributed Active Archive Center (LP DAAC), USGS/Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota
- 25 ([https://lpdaac.usgs.gov/data\\_access/data\\_pool](https://lpdaac.usgs.gov/data_access/data_pool)). Jena CarboScope inversion data were provided by Christian Rödenbeck and are available at [http://www.bgc-jena.mpg.de/CarboScope/?ID=s96\\_v3.8](http://www.bgc-jena.mpg.de/CarboScope/?ID=s96_v3.8).

## References

- Alkama, R., Decharme, B., Douville, H., Becker, M., Cazenave, A., Sheffield, J., Voldoire, A., Tyteca, S., and Le Moigne, P.: Global evaluation of the ISBA-TRIP continental hydrological system. Part I: Comparison to GRACE terrestrial water storage estimates and in situ river discharges, *J. Hydrometeorol.*, 11, 583–600, doi:10.1175/2010jhm1211.1, 2010.
- André, C., Ottlé, C., Royer, A., and Maignan, F.: Land surface temperature retrieval over circumpolar Arctic using SSM/I–SSMIS and MODIS data, *Remote Sens. Environ.*, 162, 1–10, doi:10.1016/j.rse.2015.01.028, 2015.
- Atanasiu, L.: Photosynthesis and respiration of three mosses at winter low temperatures, *Bryologist*, 74, 23–27, doi:10.2307/3241751, 1971.
- Avitabile, V., Herold, M., Heuvelink, G. B. M., Lewis, S. L., Phillips, O. L., Asner, G. P., Armston, J., Ashton, P. S., Banin, L., Bayol, N., Berry, N. J., Boeckx, P., de Jong, B. H. J., DeVries, B., Girardin, C. A. J., Kearsley, E., Lindsell, J. A., Lopez-Gonzalez, G., Lucas, R., Malhi, Y., Morel, A., Mitchard, E. T. A., Nagy, L., Qie, L., Quinones, M. J., Ryan, C. M., Ferry, S. J. W., Sunderland, T., Laurin, G. V., Gatti, R. C., Valentini, R., Verbeeck, H., Wijaya, A., and Willcock, S.: An integrated pan-tropical biomass map using multiple reference datasets, *Glob. Change Biol.*, 22, 1406–1420, doi:10.1111/gcb.13139, 2016.
- Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nature Clim. Change*, 2, 182–185, doi:10.1038/nclimate1354, <http://dx.doi.org/10.1038/nclimate1354>, 2012.
- Baldocchi, D. D., Vogel, C. A., and Hall, B.: Seasonal variation of energy and water vapor exchange rates above and below a boreal jack pine forest canopy, *J. Geophys. Res.-Atmos.*, 102, 28 939–28 951, doi:10.1029/96jd03325, 1997.
- Bartalis, Z., Wagner, W., Naeimi, V., Hasenauer, S., Scipal, K., Bonekamp, H., Figa, J., and Anderson, C.: Initial soil moisture retrievals from the METOP-A Advanced Scatterometer (ASCAT), *Geophys. Res. Lett.*, 34, doi:10.1029/2007GL031088, <http://dx.doi.org/10.1029/2007GL031088>, 120401, 2007.
- Beer, C., Lucht, W., Gerten, D., Thonicke, K., and Schmullius, C.: Effects of soil freezing and thawing on vegetation carbon density in Siberia: A modeling analysis with the Lund-Potsdam-Jena Dynamic Global Vegetation Model (LPJ-DGVM), *Global Biogeochem. Cycles*, 21, doi:10.1029/2006gb002760, 2007.
- Beer, C., Fedorov, A. N., and Torgovkin, Y.: Permafrost temperature and active-layer thickness of Yakutia with 0.5-degree spatial resolution for model evaluation, *Earth Syst. Sci. Data*, 5, 305–310, doi:10.5194/essd-5-305-2013, <http://www.earth-syst-sci-data.net/5/305/2013/>, 2013.
- Beven, K.: Infiltration into a class of vertically non-uniform soils, *Hydrolog. Sci. J.*, 29, 425–434, doi:10.1080/02626668409490960, 1984.
- Beven, K. and Germann, P.: Macropores and water flow in soils, *Water Resour. Res.*, 18, 1311–1325, doi:10.1029/WR018i005p01311, 1982.
- Biancamaria, S., Bates, P. D., Boone, A., and Mognard, N. M.: Large-scale coupled hydrologic and hydraulic modelling of the Ob river in Siberia, *J. Hydrol.*, 379, 136–150, doi:10.1016/j.jhydrol.2009.09.054, 2009.
- Bontemps, S., Defourny, P., Radoux, J., Van Bogaert, E., Lamarche, C., Achard, F., Mayaux, P., Boettcher, M., Brockmann, C., Kirches, G., Zulkhe, M., Kalogirou, V., Seifert, F., and Arino, O.: Consistent global land cover maps for climate modelling communities: current achievements of the ESA's land cover CCI, in: *Proceedings of the ESA Living Planet Symposium*, Edinburgh, pp. 9–13, 2013.
- Boone, A. and Etchevers, P.: An Intercomparison of Three Snow Schemes of Varying Complexity Coupled to the Same Land Surface Model: Local-Scale Evaluation at an Alpine Site, *Journal of Hydrometeorology*, 2, 374–394, [http://dx.doi.org/10.1175/1525-7541\(2001\)002<0374:AIOTSS>2.0.CO;2](http://dx.doi.org/10.1175/1525-7541(2001)002<0374:AIOTSS>2.0.CO;2), 2001.

- Botta, A., Viovy, N., Ciais, P., Friedlingstein, P., and Monfray, P.: A global prognostic scheme of leaf onset using satellite data, *Glob. Change Biol.*, 6, 709–725, doi:10.1046/j.1365-2486.2000.00362.x, 2000.
- Brown, J., Ferrians Jr, O., Heginbottom, J., and Melnikov, E.: Circum-Arctic map of permafrost and ground-ice conditions (Version 2), 45, 2002.
- Bulygina, O., Groisman, P. Y., Razuvaev, V., and Korshunova, N.: Changes in snow cover characteristics over Northern Eurasia since 1966, *Environ. Res. Lett.*, 6, 045 204, doi:10.1088/1748-9326/6/4/045204, 2011.
- Burke, E. J., Dankers, R., Jones, C. D., and Wiltshire, A. J.: A retrospective analysis of pan Arctic permafrost using the JULES land surface model, *Clim. Dyn.*, 41, 1025–1038, doi:10.1007/s00382-012-1648-x, 2013.
- Burke, E. J., Chadburn, S. E., and Ekici, A.: A vertical representation of soil carbon in the JULES land surface scheme (vn4.3\_permafrost) with a focus on permafrost regions, *Geosci. Model Dev.*, 10, 959–975, doi:10.5194/gmd-10-959-2017, <http://www.geosci-model-dev.net/10/959/2017/>, 2017a.
- Burke, E. J., Ekici, A., Huang, Y., Chadburn, S. E., Huntingford, C., Ciais, P., Friedlingstein, P., Peng, S., and Krinner, G.: Quantifying uncertainties of permafrost carbon-climate feedbacks, *Biogeosciences Discuss.*, 2017, 1–42, doi:10.5194/bg-2016-544, <http://www.biogeosciences-discuss.net/bg-2016-544/>, 2017b.
- Campioli, M., Vicca, S., Luyssaert, S., Bilcke, J., Ceschia, E., III, F. S. C., Ciais, P., Fernández-Martínez, M., Malhi, Y., Obersteiner, M., Olefeldt, D., Papale, D., Piao, S. L., Peñuelas, J., Sullivan, P. F., Wang, X., Zenone, T., and Janssens, I. A.: Biomass production efficiency controlled by management in temperate and boreal ecosystems, *Nat. Geosci.*, 8, 843–846, doi:10.1038/ngeo2553, 2015.
- Campoy, A., Ducharne, A., Cheruy, F., Hourdin, F., Polcher, J., and Dupont, J.: Response of land surface fluxes and precipitation to different soil bottom hydrological conditions in a general circulation model, *J. Geophys. Res.-Atmos.*, 118, 10,725–10,739, doi:10.1002/jgrd.50627, 10.1002/jgrd.50627, 2013.
- Cannell, M., Malcolm, D., and Robertson, P.: *The Ecology of Mixed-Species Stands of Trees: Forest Dynamics*, Oxford University Press, Oxford, 1992.
- Carsel, R. and Parrish, R.: Developing joint probability distributions of soil water retention characteristics, *Water Resour. Res.*, 24, 755–769, doi:10.1029/WR024i005p00755, 1988.
- Carvalhais, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., Mu, M., Saatchi, S., Santoro, M., Thurner, M., et al.: Global covariation of carbon turnover times with climate in terrestrial ecosystems, *Nature*, 514, 213–217, doi:10.1038/nature13731, 2014.
- Chadburn, S., Burke, E., Essery, R., Boike, J., Langer, M., Heikenfeld, M., Cox, P., and Friedlingstein, P.: An improved representation of physical permafrost dynamics in the JULES land-surface model, *Geosci. Model Dev.*, 8, 1493–1508, doi:10.5194/gmd-8-1493-2015, 2015a.
- Chadburn, S., Burke, E., Essery, R., Boike, J., Langer, M., Heikenfeld, M., Cox, P. M., and Friedlingstein, P.: Impact of model developments on present and future simulations of permafrost in a global land-surface model, *The Cryosphere*, 9, 1505–1521, doi:10.5194/tc-9-1505-2015, 2015b.
- Chadburn, S., Burke, E., Ekici, A., Beer, C., Porada, P., Krinner, G., Wang, T., Brutel-Vuilmet, C., Beletti Marchesini, L., van Huissteden, K., et al.: Carbon stocks and fluxes in the high-latitudes: Using site-level data to evaluate earth system models, *Biogeosciences Discuss.*, doi:10.5194/bg-2017-197, 2017.
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Fröhlich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morgui, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen,



- A. T., Wofsy, S., and Worthy, D.: CO<sub>2</sub> surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *J. Geophys. Res.-Atmos.*, 115, doi:10.1029/2010JD013887, <http://dx.doi.org/10.1029/2010JD013887>, d21307, 2010.
- Chini, L., Hurtt, G., and Frolking, S.: Harmonized Global Land Use for Years 1500–2100, V1. Data set, Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, 2014.
- Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Brönnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., Maugeri, M., Mok, H. Y., Nordli, Ø., Ross, T. F., Trigo, R. M., Wang, X. L., Woodruff, S. D., and Worley, S. J.: The twentieth century reanalysis project, *Q.J.Roy. Meteor. Soc.*, 137, 1–28, doi:10.1002/qj.776, <http://dx.doi.org/10.1002/qj.776>, 2011.
- de Rosnay, P., Bruen, M., and Polcher, J.: Sensitivity of surface fluxes to the number of layers in the soil model used in GCMs, *Geophys. Res. Lett.*, 27, 3329–3332, doi:10.1029/2000gl011574, 10.1029/2000GL011574, 2000.
- de Rosnay, P., Polcher, J., Bruen, M., and Laval, K.: Impact of a physically based soil water flow and soil-plant interaction representation for modeling large-scale land surface processes, *J. Geophys. Res.-Atmos.*, 107, 4118, doi:10.1029/2001JD000634, 2002.
- Decharme, B., Brun, E., Boone, A., Delire, C., Le Moigne, P., and Morin, S.: Impacts of snow and organic soils parameterization on northern Eurasian soil temperature profiles simulated by the ISBA land surface model, *The Cryosphere*, 10, 853–877, doi:10.5194/tc-10-853-2016, 2016.
- Doetterl, S., Stevens, A., Six, J., Merckx, R., Oost, K. V., Pinto, M. C., Casanova-Katny, A., Muñoz, C., Boudin, M., Venegas, E. Z., and Boeckx, P.: Soil carbon storage controlled by interactions between geochemistry and climate, *Nat. Geosci.*, 8, 780–783, doi:10.1038/ngeo2516, 2015.
- Domine, F., Krinner, G., Menegoz, M., Bock, J., Gallet, J.-C., and Anttila, K.: Snow-climate feedbacks driven by changes in snow thermal conductivity, in: IPY Oslo science conference, 2010.
- Domine, F., Barrere, M., and Sarrazin, D.: Seasonal evolution of the effective thermal conductivity of the snow and the soil in high Arctic herb tundra at Bylot Island, Canada, *The Cryosphere*, 10, 2573–2588, doi:10.5194/tc-10-2573-2016, <http://www.the-cryosphere.net/10/2573/2016/>, 2016.
- d’Orgeval, T., Polcher, J., and de Rosnay, P.: Sensitivity of the West African hydrological cycle in ORCHIDEE to infiltration processes, *Hydrol. Earth Syst. Sc.*, 12, 1387–1401, doi:10.5194/hessd-5-2251-2008, 2008.
- Dorigo, W., Gruber, A., De Jeu, R., Wagner, W., Stacke, T., Loew, A., Albergel, C., Brocca, L., Chung, D., Parinussa, R., et al.: Evaluation of the ESA CCI soil moisture product using ground-based observations, *Remote Sens. Environ.*, 162, 380–395, doi:10.1016/j.rse.2014.07.023, 2015.
- Druel, A., Peylin, P., Krinner, G., Ciais, P., Viovy, N., Peregon, A., Bastrikov, V., Kosykh, N., and Mironycheva-Tokareva, N.: Towards a more detailed representation of high-latitude vegetation in the global land surface model ORCHIDEE (ORC-HL-VEGv1.0), *Geosci. Model Dev. Discuss.*, doi:10.5194/gmd-2017-65, 2017, 2017.
- Ekici, A., Beer, C., Hagemann, S., Boike, J., Langer, M., and Hauck, C.: Simulating high-latitude permafrost regions by the JSBACH terrestrial ecosystem model, *Geosci. Model Dev.*, 7, 631–647, doi:10.5194/gmd-7-631-2014, <http://dx.doi.org/10.5194/gmd-7-631-2014>, 2014.
- FAO: FAO, IIASA, ISRIC, ISSCAS and JRC: Harmonized World Soil Database (version 1.2), Tech. rep., FAO, Rome, Italy and IIASA, Laxenburg, Austria, 2012.
- Fatichi, S., Leuzinger, S., and Körner, C.: Moving beyond photosynthesis: from carbon source to sink-driven vegetation modeling, *New Phytologist*, 201, 1086–1095, doi:10.1111/nph.12614, <http://dx.doi.org/10.1111/nph.12614>, 2014.

- Fekete, B. M., Vörösmarty, C. J., Roads, J. O., and Willmott, C. J.: Uncertainties in precipitation and their impacts on runoff estimates, *J. Climate*, 17, 294–304, doi:10.1175/1520-0442(2004)017<0294:uipati>2.0.co;2, 2004.
- Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X.: MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sens. Environ.*, 114, 168–182, doi:10.1016/j.rse.2009.08.016, 2010.
- 5 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., et al.: Climate–carbon cycle feedback analysis: results from the C4MIP model intercomparison, *J. Climate*, 19, 3337–3353, doi:10.1175/jcli3800.1, 2006.
- Gouttevin, I., Krinner, G., Ciais, P., Polcher, J., and Legout, C.: Multi-scale validation of a new soil freezing scheme for a land-surface model with physically-based hydrology, *The Cryosphere*, 6, 407–430, doi:10.5194/tc-6-407-2012, 2012a.
- 10 Gouttevin, I., Menegoz, M., Domine, F., Krinner, G., Koven, C., Ciais, P., Tarnocai, C., and Boike, J.: How the insulating properties of snow affect soil carbon distribution in the continental pan-Arctic area, *J. Geophys. Res.*, 117, G02020, doi:10.1029/2011jg001916, 2012b.
- Gray, D., Toth, B., Zhao, L., Pomeroy, J., and Granger, R.: Estimating areal snowmelt infiltration into frozen soils, *Hydrol. Process.*, 15, 3095–3111, doi:10.1002/hyp.320, 2001.
- Green, W. H. and Ampt, G.: Studies on soil physics, 1. The flow of air and water through soils, *J. Agr. Sci.*, 4, 1–24, 1911.
- 15 Guimberteau, M., Drapeau, G., Ronchail, J., Sultan, B., Polcher, J., Martinez, J. M., Prigent, C., Guyot, J. L., Cochonneau, G., Espinoza, J. C., Filizola, N., Fraizy, P., Lavado, W., De Oliveira, E., Pombosa, R., Noriega, L., and Vauchel, P.: Discharge simulation in the sub-basins of the Amazon using ORCHIDEE forced by new datasets, *Hydrol. Earth Syst. Sci.*, 16, 911–935, 10.5194/hess-16-911-2012, 2012.
- Haddeland, I., Clark, D. B., Franssen, W., Ludwig, F., Voß, F., Arnell, N. W., Bertrand, N., Best, M., Folwell, S., Gerten, D., Gomes, S., Gosling, S. N., Hagemann, S., Hanasaki, N., Harding, R., Heinke, J., Kabat, P., Koirala, S., Oki, T., Polcher, J., Stacke, T., Viterbo, P.,
- 20 Weedon, G. P., and Yeh, P.: Multimodel Estimate of the Global Terrestrial Water Balance: Setup and First Results., *J. Hydrometeorol.*, 12, 869–884, doi:10.1175/2011jhm1324.1, <http://dx.doi.org/10.1175/2011jhm1324.1>, 2011.
- Hansen, M. C., Townshend, J. R. G., DeFries, R. S., and Carroll, M.: Estimation of tree cover using MODIS data at global, continental and regional/local scales, *Int. J. Remote Sens.*, 26, 4359–4380, doi:10.1080/01431160500113435, <http://dx.doi.org/10.1080/01431160500113435>, 2005.
- 25 Heinsch, F. A., Reeves, M., Votava, P., Kang, S., Milesi, C., Zhao, M., Glassy, J., Jolly, W. M., Loehman, R., Bowker, C. F., et al.: GPP and NPP (MOD17A2/A3) Products NASA MODIS Land Algorithm, Tech. rep., 2003.
- Hengl, T., de Jesus, J. M., MacMillan, R. A., Batjes, N. H., Heuvelink, G. B. M., Ribeiro, E., Samuel-Rosa, A., Kempen, B., Leenaars, J. G. B., Walsh, M. G., and Gonzalez, M. R.: SoilGrids1km – Global Soil Information Based on Automated Mapping, *PLOS ONE*, 9, 1–17, doi:10.1371/journal.pone.0105992, <https://doi.org/10.1371/journal.pone.0105992>, 2014.
- 30 Hollesen, J., Matthiesen, H., Møller, A. B., and Elberling, B.: Permafrost thawing in organic Arctic soils accelerated by ground heat production, *Nature Clim. Change*, 5, 574–578, doi:10.1038/nclimate2590, 2015.
- Hudson, B. D.: Soil organic matter and available water capacity, *J. Soil Water Conserv.*, 49, 189–194, 1994.
- Hugelius, G., Tarnocai, C., Broll, G., Canadell, J., Kuhry, P., and Swanson, D.: The Northern Circumpolar Soil Carbon Database: spatially distributed datasets of soil coverage and soil carbon storage in the northern permafrost regions, *Earth Syst. Sci. Data*, 5, 3, doi:10.5194/essdd-
- 35 5-707-2012, 2013.
- Jarvis, P., Massheder, J., Hale, S., Moncrieff, J., Rayment, M., and Scott, S.: Seasonal variation of carbon dioxide, water vapor, and energy exchanges of a boreal black spruce forest, *J. Geophys. Res.-Atmos.*, 102, 28 953–28 966, doi:10.1029/97jd01176, 1997.
- Jung, M., Reichstein, M., and Bondeau, A.: Towards global empirical upscaling of FLUXNET eddy covariance observations: validation of a model tree ensemble approach using a biosphere model, *Biogeosciences*, 6, 2001–2013, doi:10.5194/bg-6-2001-2009, 2009.

- Jung, M., Reichstein, M., Margolis, H., Cescatti, A., Richardson, A., Arain, M., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., Giffonne, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, *J. Geophys. Res.-Biogeo.*, 116, G00J07, doi:10.1029/2010JG001566, 2011.
- Kajimoto, T., Matsuura, Y., Osawa, A., Prokushkin, A. S., Sofronov, M. A., and Abaimov, A. P.: Root system development of *Larix gmelinii* trees affected by micro-scale conditions of permafrost soils in central Siberia, *Plant Soil*, 255, 281–292, doi:10.1007/978-94-017-2923-9\_27, 2003.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, doi:10.1175/1520-0477(1996)077<0437:tnyrp>2.0.co;2, [http://dx.doi.org/10.1175/1520-0477\(1996\)077<0437:tnyrp>2.0.co;2](http://dx.doi.org/10.1175/1520-0477(1996)077<0437:tnyrp>2.0.co;2), 1996.
- Kleinen, T., Brovkin, V., and Schuldt, R. J.: A dynamic model of wetland extent and peat accumulation: results for the Holocene, *Biogeosciences*, 9, 235–248, doi:10.5194/bg-9-235-2012, <http://www.biogeosciences.net/9/235/2012/>, 2012.
- Knyazikhin, Y., Martonchik, J., Diner, D., Myneni, R., Verstraete, M., Pinty, B., and Gobron, N.: Estimation of vegetation canopy leaf area index and fraction of absorbed photosynthetically active radiation from atmosphere-corrected MISR data, *J. Geophys. Res.-Atmos.*, 103, 32 239–32 256, doi:10.1029/98jd02461, 1998.
- Köchy, M., Hiederer, R., and Freibauer, A.: Global distribution of soil organic carbon–Part 1: Masses and frequency distributions of SOC stocks for the tropics, permafrost regions, wetlands, and the world, *Soil*, 1, 351–365, doi:10.5194/soil-1-351-2015, 2015.
- Konings, A., Williams, A., and Gentine, P.: Sensitivity of grassland productivity to aridity controlled by stomatal and xylem regulation, *Nat. Geosci.*, 10, 284–288, doi:10.1038/ngeo2903, 2017.
- Körner, C.: *Alpine plant life: functional plant ecology of high mountain ecosystems; with 47 tables*, Springer Science & Business Media, 2003.
- Koster, R. D., Guo, Z., Yang, R., Dirmeyer, P. A., Mitchell, K., and Puma, M. J.: On the nature of soil moisture in land surface models, *J. Climate*, 22, 4322–4335, doi:10.1175/2009jcli2832.1, 2009.
- Koven, C., Friedlingstein, P., Ciais, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.: On the formation of high-latitude soil carbon stocks: Effects of cryoturbation and insulation by organic matter in a land surface model, *Geophys. Res. Lett.*, 36, doi:10.1029/2009GL040150, <http://dx.doi.org/10.1029/2009GL040150>, 121501, 2009.
- Koven, C., Riley, W., Subin, Z., Tang, J., Torn, M., Collins, W., Bonan, G., Lawrence, D., and Swenson, S.: The effect of vertically resolved soil biogeochemistry and alternate soil C and N models on C dynamics of CLM4, *Biogeosciences*, 10, 7109–7131, doi:10.5194/bgd-10-7201-2013, 2013.
- Krinner, G., Viovy, N., de Noblet-Ducoudre, N., Ogee, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochem. Cy.*, 19, 1–33, doi:10.1029/2003gb002199, doi:10.1029/2003GB002199, 2005.
- Krull, E. S., Skjemstad, J. O., and Baldock, J. A.: *Functions of soil organic matter and the effect on soil properties*, Cooperative Research Centre for Greenhouse Accounting, 2004.
- Landerer, F. W. and Swenson, S.: Accuracy of scaled GRACE terrestrial water storage estimates, *Water Resour. Res.*, 48, doi:10.1029/2011wr011453, 2012.

- Lauerwald, R., Regnier, P., Camino-Serrano, M., Guenet, B., Guimberteau, M., Ducharne, A., Polcher, J., and Ciais, P.: ORCHILEAK: A new model branch to simulate carbon transfers along the terrestrial-aquatic continuum of the Amazon basin, *Geosci. Model Dev. Discuss.*, 2017, 1–58, doi:10.5194/gmd-2017-79, <http://www.geosci-model-dev-discuss.net/gmd-2017-79/>, 2017.
- 5 Lawrence, D. M. and Slater, A. G.: Incorporating organic soil into a global climate model, *Clim. Dyn.*, 30, 145–160, doi:10.1007/s00382-007-0278-1, <http://dx.doi.org/10.1007/s00382-007-0278-1>, 2008.
- Lawrence, D. M., Slater, A. G., Romanovsky, V. E., and Nicolsky, D. J.: Sensitivity of a model projection of near-surface permafrost degradation to soil column depth and representation of soil organic matter, *J. Geophys. Res.-Earth*, 113, doi:10.1029/2007jf000883, 2008.
- Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296, 1–22, doi:10.1016/j.jhydrol.2004.03.028, 2004.
- 10 Leinonen, I.: A simulation model for the annual frost hardiness and freeze damage of Scots pine, *Annals of Botany*, 78, 687–693, doi:10.1006/anbo.1996.0178, 1996.
- Liang, S. and Xiao, Z.: Global land surface products: Leaf area index product data collection (1985–2010), Tech. rep., 2012.
- Liu, Y. Y., Van Dijk, A. I., De Jeu, R. A., Canadell, J. G., McCabe, M. F., Evans, J. P., and Wang, G.: Recent reversal in loss of global terrestrial biomass, *Nature Clim. Change*, 5, 470–474, doi:10.1038/nclimate2581, 2015.
- 15 Loth, B., Graf, H.-F., and Oberhuber, J. M.: Snow cover model for global climate simulations, *J. Geophys. Res.*, 98, 451–10, doi:10.1029/93jd00324, 1993.
- Luo, Y., Ahlström, A., Allison, S. D., Batjes, N. H., Brovkin, V., Carvalhais, N., Chappell, A., Ciais, P., Davidson, E. A., Finzi, A., et al.: Toward more realistic projections of soil carbon dynamics by Earth system models, *Global Biogeochem. Cycles*, 30, 40–56, doi:10.1002/2015gb005239, 2016.
- 20 Luysaert, S., Inglisma, I., Jung, M., Richardson, A. D., Reichstein, M., Papale, D., Piao, S., SCHULZE, E.-D., Wingate, L., Matteucci, G., et al.: CO<sub>2</sub> balance of boreal, temperate, and tropical forests derived from a global database, *Glob. Change Biol.*, 13, 2509–2537, doi:10.1111/j.1365-2486.2007.01439.x, 2007.
- MacBean, N., Maignan, F., Peylin, P., Bacour, C., Bréon, F.-M., and Ciais, P.: Using satellite data to improve the leaf phenology of a global terrestrial biosphere model, *Biogeosciences*, 12, 7185–7208, doi:10.5194/bgd-12-13311-2015, 2015.
- 25 Magand, C., Ducharne, A., Le Moine, N., and Gascoin, S.: Introducing hysteresis in snow depletion curves to improve the water budget of a land surface model in an Alpine catchment, *J. Hydrometeorol.*, 15, 631–649, doi:10.1175/jhm-d-13-091.1, 2014.
- Manzoni, S. and Porporato, A.: Soil carbon and nitrogen mineralization: theory and models across scales, *Soil Biol. Biochem.*, 41, 1355–1379, doi:10.1016/j.soilbio.2009.02.031, 2009.
- 30 Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A Reconstruction of Regional and Global Temperature for the Past 11,300 Years, *Science*, 339, 1198–1201, doi:10.1126/science.1228026, <http://science.sciencemag.org/content/339/6124/1198>, 2013.
- Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., Beck, H. E., Dorigo, W. A., and Verhoest, N. E. C.: GLEAM v3: satellite-based land evaporation and root-zone soil moisture, *Geosci. Model Dev.*, 10, 1903–1925, doi:10.5194/gmd-10-1903-2017, <http://dx.doi.org/10.5194/gmd-2016-162>, 2017.
- 35 McAdam, S. A. and Brodribb, T. J.: The evolution of mechanisms driving the stomatal response to vapor pressure deficit, *Plant Physiol.*, 167, 833–843, doi:10.1104/pp.114.252940, 2015.
- McGuire, A. D., Koven, C., Lawrence, D. M., Clein, J. S., Xia, J., Beer, C., Burke, E., Chen, G., Chen, X., Delire, C., Jafarov, E., MacDougall, A. H., Marchenko, S., Nicolsky, D., Peng, S., Rinke, A., Saito, K., Zhang, W., Alkama, R., Bohn, T. J., Ciais, P., Decharme, B., Ekici, A., Gouttevin, I., Hajima, T., Hayes, D. J., Ji, D., Krinner, G., Lettenmaier, D. P., Luo, Y., Miller, P. A., Moore, J. C., Romanovsky, V.,

- Schädel, C., Schaefer, K., Schuur, E. A., Smith, B., Sueyoshi, T., and Zhuang, Q.: Variability in the sensitivity among model simulations of permafrost and carbon dynamics in the permafrost region between 1960 and 2009, *Global Biogeochem. Cycles*, 30, 1015–1037, doi:10.1002/2016gb005405, <http://dx.doi.org/10.1002/2016GB005405>, 2016.
- 5 Medlyn, B. E., Zaehle, S., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hanson, P. J., Hickler, T., Jain, A. K., Luo, Y., Parton, W., et al.: Using ecosystem experiments to improve vegetation models, *Nature Clim. Change*, 5, 528–534, doi:10.1038/nclimate2621, 2015.
- Michaletz, S. T., Cheng, D., Kerkhoff, A. J., and Enquist, B. J.: Convergence of terrestrial plant production across global climate gradients, *Nature*, 512, 39–43, doi:10.1038/nature13470, 2014.
- Miralles, D., Holmes, T., De Jeu, R., Gash, J., Meesters, A., and Dolman, A.: Global land-surface evaporation estimated from satellite-based  
10 observations, *Hydrol. Earth Syst. Sc.*, 15, 453–469, doi:10.5194/hess-15-453-2011, 2011.
- Morris, P. J., Baird, A. J., and Belyea, L. R.: Bridging the gap between models and measurements of peat hydraulic conductivity, *Water Resour. Res.*, 51, 5353–5364, doi:10.1002/2015wr017264, <http://dx.doi.org/10.1002/2015WR017264>, 2015.
- Motoya, K., Masuda, K., Takata, K., and Oki, T.: Sensitivity of precipitation gauge correction for the estimation of the global water balance, *Eos Transactions, American Geophysical Union*, 83, 2002.
- 15 Mualem, Y.: A new model for predicting the hydraulic conductivity of unsaturated porous media, *Water Resour. Res.*, 12, 513–522, doi:10.1029/WR012i003p00513, 1976.
- Muskett, R. R. and Romanovsky, V. E.: Groundwater storage changes in arctic permafrost watersheds from GRACE and in situ measurements, *Environ. Res. Lett.*, 4, 045 009, doi:10.1088/1748-9326/4/4/045009, <http://dx.doi.org/10.1088/1748-9326/4/4/045009>, 2009.
- Naudts, K., Ryder, J., McGrath, M. J., Otto, J., Chen, Y., Valade, A., Bellasen, V., Berhongaray, G., Bönisch, G., Campioli, M., Ghattas,  
20 J., Groote, T. D., Haverd, V., Kattge, J., MacBean, N., Maignan, F., Merilä, P., Penuelas, J., Peylin, P., Pinty, B., Pretzsch, H., Schulze, E. D., Solyga, D., Vuichard, N., Yan, Y., and Luyssaert, S.: A vertically discretised canopy description for ORCHIDEE (SVN r2290) and the modifications to the energy, water and carbon fluxes, *Geosci. Model Dev.*, 7, 8565–8647, doi:10.5194/gmd-8-2035-2015, <http://dx.doi.org/10.5194/gmd-8-2035-2015>, 2015.
- New, M., Hulme, M., and Jones, P.: Representing twentieth-century space-time climate variability. Part II: Development of 1901–96 monthly  
25 grids of terrestrial surface climate, *J. Climate*, 13, 2217–2238, doi:10.1175/1520-0442(2000)013<2217:rtcstc>2.0.co;2, 2000.
- Ngo-Duc, T., Polcher, J., and Laval, K.: A 53-year forcing data set for land surface models, *J. Geophys. Res.-Atmos.*, 110, D06 116, doi:10.1029/2004JD005434, 2005.
- Novick, K. A., Ficklin, D. L., Stoy, P. C., Williams, C. A., Bohrer, G., Oishi, A. C., Papuga, S. A., Blanken, P. D., Noormets, A., Sulman, B. N., et al.: The increasing importance of atmospheric demand for ecosystem water and carbon fluxes, *Nature Clim. Change*, 6, 1023–  
30 1027, doi:10.1038/nclimate3114, 2016.
- O'Donnell, J. A., Harden, J. W., McGuire, A. D., and Romanovsky, V. E.: Exploring the sensitivity of soil carbon dynamics to climate change, fire disturbance and permafrost thaw in a black spruce ecosystem, *Biogeosciences*, 8, 1367–1382, doi:10.5194/bg-8-1367-2011, <http://www.biogeosciences.net/8/1367/2011/>, 2011.
- Oki, T., Nishimura, T., and Dirmeyer, P.: Assessment of annual runoff from land surface models using Total Runoff Integrating Pathways  
35 (TRIP), *J. Meteorol. Society of Japan*, 77, 235–255, doi:10.2151/jmsj1965.77.1b\_235, 1999.
- Olson, R., Johnson, K., Zheng, D., and Scurlock, J.: Global and regional ecosystem modeling: Databases of model drivers and validation measurements, ORNL Technical Memorandum TM-2001/196. Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA, doi:10.2172/814241, 2001.

- Oren, R., Sperry, J., Katul, G., Pataki, D., Ewers, B., Phillips, N., and Schäfer, K.: Survey and synthesis of intra-and interspecific variation in stomatal sensitivity to vapour pressure deficit, *Plant, Cell & Environment*, 22, 1515–1526, doi:10.1046/j.1365-3040.1999.00513.x, 1999.
- Owe, M., de Jeu, R., and Holmes, T.: Multisensor historical climatology of satellite-derived global land surface moisture, *J. Geophys. Res.-Earth*, 113, doi:10.1029/2007jf000769, 2008.
- Pan, Y., Chen, J. M., Birdsey, R., McCullough, K., He, L., and Deng, F.: Age structure and disturbance legacy of North American forests, *Biogeosciences*, 8, 715, doi:10.5194/bg-8-715-2011, 2011.
- Papa, F., Prigent, C., and Rossow, W.: Monitoring flood and discharge variations in the large Siberian rivers from a multi-satellite technique, *Surv. Geophys.*, 29, 297–317, doi:10.1007/s10712-008-9036-0, 2008.
- Parton, W., Schimel, D. S., Cole, C., and Ojima, D.: Analysis of factors controlling soil organic matter levels in Great Plains grasslands, *Soil Science Society of America Journal*, 51, 1173–1179, doi:10.2136/sssaj1987.03615995005100050015x, 1987.
- Parton, W. J., Stewart, J. W., and Cole, C. V.: Dynamics of C, N, P and S in grassland soils: a model, *Biogeochemistry*, 5, 109–131, doi:10.1007/bf02180320, 1988.
- Peng, S., Piao, S., Ciais, P., Fang, J., and Wang, X.: Change in winter snow depth and its impacts on vegetation in China, *Glob. Change Biol.*, 16, 3004–3013, doi:10.1111/j.1365-2486.2010.02210.x, <http://dx.doi.org/10.1111/j.1365-2486.2010.02210.x>, 2010.
- Peng, S., Ciais, P., Krinner, G., Wang, T., Gouttevin, I., McGuire, A. D., Lawrence, D., Burke, E., Chen, X., Decharme, B., Koven, C., MacDougall, A., Rinke, A., Saito, K., Zhang, W., Alkama, R., Bohn, T. J., Delire, C., Hajima, T., Ji, D., Lettenmaier, D. P., Miller, P. A., Moore, J. C., Smith, B., and Sueyoshi, T.: Simulated high-latitude soil thermal dynamics during the past 4 decades, *The Cryosphere*, 10, 179–192, doi:10.5194/tc-10-179-2016, <http://dx.doi.org/10.5194/tc-10-179-2016>, 2016.
- Peng, S., Ciais, P., Maignan, F., Li, W., Chang, J., Wang, T., and Yue, C.: Sensitivity of land use change emission estimates to historical land use and land cover mapping, *Global Biogeochem. Cycles*, 31, 626–643, doi:10.1002/2015GB005360, <http://dx.doi.org/10.1002/2015GB005360>, 2015GB005360, 2017.
- Piao, S., Luysaert, S., Ciais, P., Janssens, I. A., Chen, A., Cao, C., Fang, J., Friedlingstein, P., Luo, Y., and Wang, S.: Forest annual carbon cost: a global-scale analysis of autotrophic respiration, *Ecology*, 91, 652–661, doi:10.1890/08-2176.1, <http://dx.doi.org/10.1890/08-2176.1>, 2010.
- Polcher, J.: Les processus de surface à l'échelle globale et leurs interactions avec l'atmosphère, *Habilitation à diriger des recherches*, 2003.
- Pomeroy, J. W., Parviainen, J., Hedstrom, N., and Gray, D. M.: Coupled modelling of forest snow interception and sublimation, *Hydrol. Process.*, 12, 2317–2337, doi:10.1002/(SICI)1099-1085(199812)12:15<2317::AID-HYP799>3.0.CO;2-X, [http://dx.doi.org/10.1002/\(SICI\)1099-1085\(199812\)12:15<2317::AID-HYP799>3.0.CO;2-X](http://dx.doi.org/10.1002/(SICI)1099-1085(199812)12:15<2317::AID-HYP799>3.0.CO;2-X), 1998.
- Porada, P., Ekici, A., and Beer, C.: Effects of bryophyte and lichen cover on permafrost soil temperature at large scale, *The Cryosphere*, 10, 2291, doi:10.5194/tc-10-2291-2016, 2016.
- Poulter, B., Ciais, P., Hodson, E., Lischke, H., Maignan, F., Plummer, S., and Zimmermann, N.: Plant functional type mapping for earth system models, *Geosci. Model Dev.*, 4, 993–1010, doi:10.5194/gmd-4-993-2011, 2011.
- Poulter, B., MacBean, N., Hartley, A., Khlystova, I., Arino, O., Betts, R., Bontemps, S., Boettcher, M., Brockmann, C., Defourny, P., Hagemann, S., Herold, M., Kirches, G., Lamarche, C., Lederer, D., Ottlé, C., Peters, M., and Peylin, P.: Plant functional type classification for earth system models: results from the European Space Agency's Land Cover Climate Change Initiative, *Geosci. Model Dev.*, 8, 2315–2328, doi:10.5194/gmd-8-2315-2015, <http://www.geosci-model-dev.net/8/2315/2015/>, 2015.
- Priestley, C. and Taylor, R.: On the assessment of surface heat flux and evaporation using large-scale parameters, *Mon. Weather Rev.*, 100, 81–92, doi:10.1175/1520-0493(1972)100<0081:otaosh>2.3.co;2, 1972.

- Prigent, C., Lettenmaier, D. P., Aires, F., and Papa, F.: Toward a High-Resolution Monitoring of Continental Surface Water Extent and Dynamics, at Global Scale: from GIEMS (Global Inundation Extent from Multi-Satellites) to SWOT (Surface Water Ocean Topography), *Surv. Geophys.*, 37, 339–355, doi:10.1007/978-3-319-32449-4\_7, 2016.
- 5 Pulliainen, J.: Mapping of snow water equivalent and snow depth in boreal and sub-arctic zones by assimilating space-borne microwave radiometer data and ground-based observations, *Remote Sens. Environ.*, 101, 257 – 269, doi:https://doi.org/10.1016/j.rse.2006.01.002, http://www.sciencedirect.com/science/article/pii/S0034425706000241, 2006.
- Randerson, J. T., Chen, Y., van der Werf, G. R., Rogers, B. M., and Morton, D. C.: Global burned area and biomass burning emissions from small fires, *J. Geophys. Res.-Biogeo.*, 117, doi:10.1029/2012JG002128, http://dx.doi.org/10.1029/2012JG002128, g04012, 2012.
- 10 Rawlins, M. A., Fahnestock, M., Frolking, S., and Vörösmarty, C. J.: On the evaluation of snow water equivalent estimates over the terrestrial Arctic drainage basin, *Hydrol. Process.*, 21, 1616–1623, doi:10.1002/hyp.6724, 2007.
- Rawls, W., Pachepsky, Y., Ritchie, J., Sobecki, T., and Bloodworth, H.: Effect of soil organic carbon on soil water retention, *Geoderma*, 116, 61–76, doi:https://doi.org/10.1016/S0016-7061(03)00094-6, http://www.sciencedirect.com/science/article/pii/S0016706103000946, quantifying agricultural management effects on soil properties and processes, 2003.
- 15 Reynolds, C., Jackson, T., and Rawls, W.: Estimating soil water-holding capacities by linking the Food and Agriculture Organization soil map of the world with global pedon databases and continuous pedotransfer functions, *Water Resour. Res.*, 36, 3653–3662, doi:10.1029/2000wr900130, 2000.
- Ringeval, B., Decharme, B., Piao, S. L., Ciais, P., Papa, F., de Noblet-Ducoudré, N., Prigent, C., Friedlingstein, P., Gouttevin, I., Koven, C., and Ducharne, A.: Modelling sub-grid wetland in the ORCHIDEE global land surface model: evaluation against river discharges and remotely sensed data, *Geosci. Model Dev.*, 5, 941–962, doi:10.5194/gmdd-5-683-2012, http://dx.doi.org/10.5194/gmdd-5-683-2012, 2012.
- 20 Rödenbeck, C.: Estimating CO<sub>2</sub> sources and sinks from atmospheric mixing ratio measurements using a global inversion of atmospheric transport, Technical Reports, Max Planck Institute for Biogeochemistry, 2005.
- Royer, A. and Poirier, S.: Surface temperature spatial and temporal variations in North America from homogenized satellite SMMR-SSM/I microwave measurements and reanalysis for 1979–2008, *J. Geophys. Res.-Atmos.*, 115, doi:10.1029/2009JD012760, http://dx.doi.org/10.1029/2009JD012760, d08110, 2010.
- 25 Running, S. W., Nemani, R. R., Heinsch, F. A., Zhao, M., Reeves, M., and Hashimoto, H.: A continuous satellite-derived measure of global terrestrial primary production, *BioScience*, 54, 547–560, doi:10.1641/0006-3568(2004)054[0547:acsmog]2.0.co;2, 2004.
- Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *P. Natl. Acad. Sci. U.S.A.*, 108, 9899–9904, doi:10.1073/pnas.1019576108, http://dx.doi.org/10.1073/pnas.1019576108, 2011.
- 30 Sakumura, C., Bettadpur, S., and Bruinsma, S.: Ensemble prediction and intercomparison analysis of GRACE time-variable gravity field models, *Geophys. Res. Lett.*, 41, 1389–1397, doi:10.1002/2013GL058632, http://dx.doi.org/10.1002/2013GL058632, 2014.
- Santoro, M., Beer, C., Cartus, O., Schmullius, C., Shvidenko, A., McCallum, I., Wegmüller, U., and Wiesmann, A.: Retrieval of growing stock volume in boreal forest using hyper-temporal series of Envisat ASAR ScanSAR backscatter measurements, *Remote Sens. Environ.*, 115, 490–507, doi:https://doi.org/10.1016/j.rse.2010.09.018, http://www.sciencedirect.com/science/article/pii/S0034425710002919, 2011.
- 35 Santoro, M., Beaudoin, A., Beer, C., Cartus, O., Fransson, J. E., Hall, R. J., Pathe, C., Schmullius, C., Schepaschenko, D., Shvidenko, A., Thurner, M., and Wegmüller, U.: Forest growing stock volume of the northern hemisphere: Spatially explicit estimates for 2010 derived

- from Envisat ASAR, *Remote Sens. Environ.*, 168, 316 – 334, doi:<https://doi.org/10.1016/j.rse.2015.07.005>, <http://www.sciencedirect.com/science/article/pii/S003442571530064X>, 2015.
- Schaefer, K. and Jafarov, E.: A parameterization of respiration in frozen soils based on substrate availability, *Biogeosciences*, 13, 1991–2001, doi:10.5194/bg-13-1991-2016, 2016.
- 5 Schulze, E.-D., Lloyd, J., Kelliher, F., Wirth, C., Rebmann, C., Lühker, B., Mund, M., Knohl, A., Milyukova, I., Schulze, W., et al.: Productivity of forests in the Euro Siberian boreal region and their potential to act as a carbon sink—a synthesis, *Glob. Change Biol.*, 5, 703–722, doi:10.1046/j.1365-2486.1999.00266.x, 1999.
- Sherstiukov, A.: Statistical quality control of soil temperature dataset, *RIHMI-WDC*, 176, 224–232, 2012.
- 10 Shiklomanov, A. and Lammers, R.: Record Russian river discharge in 2007 and the limits of analysis, *Environ. Res. Lett.*, 4, 045 015, doi:10.1088/1748-9326/4/4/045015, 2009.
- Spahni, R., Wania, R., Neef, L., van Weele, M., Pison, I., Bousquet, P., Frankenberg, C., Foster, P. N., Joos, F., Prentice, I. C., and van Velthoven, P.: Constraining global methane emissions and uptake by ecosystems, *Biogeosciences*, 8, 1643–1665, doi:10.5194/bg-8-1643-2011, <http://www.biogeosciences.net/8/1643/2011/>, 2011.
- 15 Sturm, M. and Johnson, J. B.: Thermal conductivity measurements of depth hoar, *J. Geophys. Res.-Sol Ea.*, 97, 2129–2139, doi:10.1029/91JB02685, <http://dx.doi.org/10.1029/91JB02685>, 1992.
- Swenson, S.: GRACE monthly land water mass grids NETCDF RELEASE 5.0. Ver. 5.0 PO.DAAC, CA, USA. Dataset accessed [2016-01-25] at <http://dx.doi.org/10.5067/TELND-NC005>, Tech. rep., 2012.
- Swenson, S. and Wahr, J.: Post-processing removal of correlated errors in GRACE data, *Geophys. Res. Lett.*, 33, L08 402, doi:10.1029/2005GL025285, 2006.
- 20 Swenson, S., Wahr, J., and Milly, P.: Estimated accuracies of regional water storage variations inferred from the Gravity Recovery and Climate Experiment (GRACE), *Water Resour. Res.*, 39, 1223, doi:10.1029/2002WR001808, 2003.
- Takala, M., Pulliainen, J., Metsamäki, S. J., and Koskinen, J. T.: Detection of snowmelt using spaceborne microwave radiometer data in Eurasia from 1979 to 2007, *IEEE Trans. Geosci. Remote Sensing*, 47, 2996–3007, doi:10.1109/tgrs.2009.2018442, 2009.
- 25 Takala, M., Luojus, K., Pulliainen, J., Derksen, C., Lemmetyinen, J., Kärnä, J.-P., Koskinen, J., and Bojkov, B.: Estimating northern hemisphere snow water equivalent for climate research through assimilation of space-borne radiometer data and ground-based measurements, *Remote Sens. Environ.*, 115, 3517 – 3529, doi:<https://doi.org/10.1016/j.rse.2011.08.014>, <http://www.sciencedirect.com/science/article/pii/S0034425711003166>, 2011.
- Tanja, S., Berninger, F., Vesala, T., Markkanen, T., Hari, P., Mäkelä, A., Ilvesniemi, H., Hänninen, H., Nikinmaa, E., Huttula, T., et al.: Air temperature triggers the recovery of evergreen boreal forest photosynthesis in spring, *Glob. Change Biol.*, 9, 1410–1426, doi:10.1046/j.1365-2486.2003.00597.x, 2003.
- 30 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochem. Cycles*, 23, doi:10.1029/2008GB003327, <http://dx.doi.org/10.1029/2008GB003327>, gB2023, 2009.
- 35 Thonicke, K., Spessa, A., Prentice, I., Harrison, S., Dong, L., and Carmona-Moreno, C.: The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model., *Biogeosciences*, 7, 1991–2011, doi:10.5194/bg-7-1991-2010, 2010.




- Thurner, M., Beer, C., Santoro, M., Carvalhais, N., Wutzler, T., Schepaschenko, D., Shvidenko, A., Kompter, E., Ahrens, B., Levick, S. R., and Schmulilius, C.: Carbon stock and density of northern boreal and temperate forests, *Global Ecol. Biogeogr.*, 23, 297–310, doi:10.1111/geb.12125, <http://dx.doi.org/10.1111/geb.12125>, 2014.
- 5 Thurner, M., Beer, C., Carvalhais, N., Forkel, M., Santoro, M., Tum, M., and Schmulilius, C.: Large-scale variation in boreal and temperate forest carbon turnover rate related to climate, *Geophys. Res. Lett.*, 43, 4576–4585, doi:10.1002/2016GL068794, <http://dx.doi.org/10.1002/2016GL068794>, 2016GL068794, 2016.
- Tipping, E., Somerville, C. J., and Luster, J.: The C:N:P:S stoichiometry of soil organic matter, *Biogeochemistry*, 130, 117–131, doi:10.1007/s10533-016-0247-z, <http://dx.doi.org/10.1007/s10533-016-0247-z>, 2016.
- 10 Turner, D. P., Ritts, W. D., Cohen, W. B., Gower, S. T., Running, S. W., Zhao, M., Costa, M. H., Kirschbaum, A. A., Ham, J. M., Saleska, S. R., and Ahl, D. E.: Evaluation of MODIS NPP and GPP products across multiple biomes, *Remote Sensing of Environment*, 102, 282 – 292, doi:10.1016/j.rse.2006.02.017, <http://www.sciencedirect.com/science/article/pii/S0034425706000873>, 2006.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009),  
15 *Atmos. Chem. Phys.*, 10, 11 707–11 735, doi:10.5194/acp-10-11707-2010, <http://www.atmos-chem-phys.net/10/11707/2010/>, 2010.
- Van Genuchten, M. T.: A closed-form equation for predicting the hydraulic conductivity of unsaturated soils, *Soil Sci. Soc. Am. J.*, 44, 892–898, doi:10.2136/sssaj1980.03615995004400050002x, 1980.
- Van Wijk, M., Williams, M., Laundre, J., and Shaver, G.: Interannual variability of plant phenology in tussock tundra: modelling interactions of plant productivity, plant phenology, snowmelt and soil thaw, *Glob. Change Biol.*, 9, 743–758, doi:10.1046/j.1365-2486.2003.00625.x,  
20 2003.
- Vörösmarty, C. J., Fekete, B., Meybeck, M., and Lammers, R.: Global system of rivers: Its role in organizing continental land mass and defining land-to-ocean linkages, *Global Biogeochem. Cy.*, 14, 599–621, doi:10.1029/1999gb900092, 2000.
- Wagner, W., Lemoine, G., and Rott, H.: A Method for Estimating Soil Moisture from ERS Scatterometer and Soil Data, *Remote Sens. Environ.*, 70, 191 – 207, doi:[https://doi.org/10.1016/S0034-4257\(99\)00036-X](https://doi.org/10.1016/S0034-4257(99)00036-X), <http://www.sciencedirect.com/science/article/pii/S003442579900036X>,  
25 S003442579900036X, 1999.
- Wang, F., Cheruy, F., and Dufresne, J.-L.: The improvement of soil thermodynamics and its effects on land surface meteorology in the IPSL climate model, *Geosci. Model Dev.*, 9, 363–381, doi:10.5194/gmdd-8-8411-2015, <http://www.geosci-model-dev-discuss.net/8/8411/2015/>, 2016a.
- Wang, S., Huang, J., Yang, D., Pavlic, G., and Li, J.: Long-term water budget imbalances and error sources for cold region drainage basins,  
30 *Hydrol. Process.*, 29, 2125–2136, doi:10.1002/hyp.10343, 2015a.
- Wang, T., Ottlé, C., Boone, A., Ciais, P., Brun, E., Morin, S., Krinner, G., Piao, S., and Peng, S.: Evaluation of an improved intermediate complexity snow scheme in the ORCHIDEE land surface model, *J. Geophys. Res.-Atmos.*, 118, 6064–6079, doi:10.1002/jgrd.50395, 2013.
- Wang, T., Peng, S., Krinner, G., Ryder, J., Li, Y., Dantec-Nédélec, S., and Ottlé, C.: Impacts of Satellite-Based Snow Albedo Assimilation on Offline and Coupled Land Surface Model Simulations, *PLoS One*, 10, e0137 275, doi:10.1371/journal.pone.0137275, 2015b.
- 35 Wang, W., Rinke, A., Moore, J. C., Ji, D., Cui, X., Peng, S., Lawrence, D. M., McGuire, A. D., Burke, E. J., Chen, X., Decharme, B., Koven, C., MacDougall, A., Saito, K., Zhang, W., Alkama, R., Bohn, T. J., Ciais, P., Delire, C., Gouttevin, I., Hajima, T., Krinner, G., Lettenmaier, D. P., Miller, P. A., Smith, B., Sueyoshi, T., and Sherstiukov, A. B.: Evaluation of air–soil temperature relationships

- simulated by land surface models during winter across the permafrost region, *The Cryosphere*, 10, 1721–1737, doi:10.5194/tc-10-1721-2016, <http://dx.doi.org/10.5194/tc-10-1721-2016>, 2016b.
- Wen, L., Nagabhatla, N., Lü, S., and Wang, S.-Y.: Impact of rain snow threshold temperature on snow depth simulation in land surface and regional atmospheric models, *Adv. Atmos. Sci.*, 30, 1449–1460, doi:10.1007/s00376-012-2192-7, <http://dx.doi.org/10.1007/s00376-012-2192-7>, 2013.
- Wutzler, T. and Reichstein, M.: Colimitation of decomposition by substrate and decomposers? a comparison of model formulations, *Biogeo-sciences*, 5, 749–759, 2008.
- Xiao, Z., Liang, S., Wang, J., Chen, P., Yin, X., Zhang, L., and Song, J.: Use of general regression neural networks for generating the GLASS leaf area index product from time-series MODIS surface reflectance, *IEEE Trans. Geosci. Remote Sensing*, 52, 209–223, doi:10.1109/tgrs.2013.2237780, 2014.
- Yang, D., Robinson, D., Zhao, Y., Estilow, T., and Ye, B.: Streamflow response to seasonal snow cover extent changes in large Siberian watersheds, *J. Geophys. Res.-Atmos.*, 108, doi:10.1029/2002jd003149, 2003.
- Yang, D., Kane, D., Zhang, Z., Legates, D., and Goodison, B.: Bias corrections of long-term (1973–2004) daily precipitation data over the northern regions, *Geophys. Res. Lett.*, 32, doi:10.1029/2005gl024057, 2005.
- Yin, X. and Struik, P.: C<sub>3</sub> and C<sub>4</sub> photosynthesis models: An overview from the perspective of crop modelling, *NJAS-Wagen. J. Life Sc.*, 57, 27–38, 2009.
- Yoshimura, K. and Kanamitsu, M.: Dynamical global downscaling of global reanalysis, *Mon. Weather Rev.*, 136, 2983–2998, doi:10.1175/2008mwr2281.1, 2008.
- Yue, C., Ciais, P., Cadule, P., Thonicke, K., Archibald, S., Poulter, B., Hao, W. M., Hantson, S., Mouillot, F., Friedlingstein, P., Maignan, F., and Viovy, N.: Modelling the role of fires in the terrestrial carbon balance by incorporating SPITFIRE into the global vegetation model ORCHIDEE–Part 1: simulating historical global burned area and fire regimes, *Geosci. Model Dev.*, 7, 2747–2767, doi:10.5194/gmd-7-2747-2014, <http://dx.doi.org/10.5194/gmd-7-2747-2014>, 2014.
- Yue, C., Ciais, P., Cadule, P., Thonicke, K., and Van Leeuwen, T.: Modelling the role of fires in the terrestrial carbon balance by incorporating SPITFIRE into the global vegetation model ORCHIDEE–Part 2: Carbon emissions and the role of fires in the global carbon balance, *Geosci. Model Dev.*, 8, 1321–1338, doi:10.5194/gmd-8-1321-2015, 2015.
- Zhao, M., Heinsch, F. A., Nemani, R. R., and Running, S. W.: Improvements of the MODIS terrestrial gross and net primary production global data set, *Remote Sens. Environ.*, 95, 164 – 176, doi:<http://dx.doi.org/10.1016/j.rse.2004.12.011>, <http://www.sciencedirect.com/science/article/pii/S0034425705000106>, 2005.
- Zhao, M., Running, S. W., and Nemani, R. R.: Sensitivity of Moderate Resolution Imaging Spectroradiometer (MODIS) terrestrial primary production to the accuracy of meteorological reanalyses, *J. Geophys. Res.-Biogeo.*, 111, doi:10.1029/2004JG000004, <http://dx.doi.org/10.1029/2004JG000004>, g01002, 2006.
- Zhao, Y., Ciais, P., Peylin, P., Viovy, N., Longdoz, B., Bonnefond, J. M., Rambal, S., Klumpp, K., Olioso, A., Cellier, P., Maignan, F., Eglin, T., and Calvet, J. C.: How errors on meteorological variables impact simulated ecosystem fluxes: a case study for six French sites, *Biogeosciences*, 9, 2537–2564, doi:10.5194/bg-9-2537-2012, <http://dx.doi.org/10.5194/bg-9-2537-2012>, 2012.
- Zhou, S., Medlyn, B., Sabaté, S., Sperlich, D., Prentice, I. C., et al.: Short-term water stress impacts on stomatal, mesophyll and biochemical limitations to photosynthesis differ consistently among tree species from contrasting climates, *Tree Physiol.*, 34, 1035–46, doi:10.1093/treephys/tpu072, 2014.


- Zhu, D., Peng, S. S., Ciais, P., Viovy, N., Druel, A., Kageyama, M., Krinner, G., Peylin, P., Ottlé, C., Piao, S. L., Poulter, B., Schepaschenko, D., and Shvidenko, A.: Improving the dynamics of northern vegetation in the ORCHIDEE ecosystem model, *Geosci. Model Dev.*, 8, 2263–2283, doi:10.5194/gmd-8-2263-2015, <http://dx.doi.org/10.5194/gmd-8-2263-2015>, 2015.
- 5 Zhu, D., Peng, S., Ciais, P., Zech, R., Krinner, G., Zimov, S., and Grosse, G.: Simulating soil organic carbon in yedoma deposits during the Last Glacial Maximum in a land surface model, *Geophys. Res. Lett.*, 43, 5133–5142, doi:10.1002/2016gl068874, 2016.
- Zhu, Z., Bi, J., Pan, Y., Ganguly, S., Anav, A., Xu, L., Samanta, A., Piao, S., Nemani, R. R., and Myneni, R. B.: Global data sets of vegetation leaf area index (LAI) 3g and Fraction of Photosynthetically Active Radiation (FPAR) 3g derived from Global Inventory Modeling and Mapping Studies (GIMMS) Normalized Difference Vegetation Index (NDVI3g) for the period 1981 to 2011, *Remote Sens.*, 5, 927–948, doi:10.3390/rs5020927, 2013.
- 10 Zimov, N. S., Zimov, S. A., Zimova, A. E., Zimova, G. M., Chuprynin, V. I., and Chapin, F. S.: Carbon storage in permafrost and soils of the mammoth tundra-steppe biome: Role in the global carbon budget, *Geophys. Res. Lett.*, 36, doi:10.1029/2008gl036332, <http://dx.doi.org/10.1029/2008GL036332>, 102502, 2009.
- Zimov, S. A., Zimova, G. M., Daviodov, S. P., Daviodova, A. I., Voropaev, Y. V., Voropaeva, Z. V., Prosiannikov, S. F., Prosiannikova, O. V., Semiletova, I. V., and Semiletov, I. P.: Winter biotic activity and production of CO<sub>2</sub> in Siberian soils: A factor in the greenhouse effect, *J. Geophys. Res.-Atmos.*, 98, 5017–5023, doi:10.1029/92JD02473, <http://dx.doi.org/10.1029/92JD02473>, 1993.
- 15

Dataset	Variable	Resolution	Period	URL	References
<b>Evaluation datasets for water budget</b>					
GRACE	TWS	1°	Jul. 2003-Dec. 2007	<a href="http://grace.jpl.nasa.gov">http://grace.jpl.nasa.gov</a>	Swenson and Wahr, 2006; Swenson, 2012 Landerer and Swenson, 2012
GlobSnow	Snow water mass	25 km	1979-2013	<a href="http://www.globsnow.info">www.globsnow.info</a>	Takala et al. (2009)
GLEAM v3.0a	Evapotranspiration	0.25°	1980-2014	<a href="http://www.gleam.eu">http://www.gleam.eu</a>	Miralles et al. (2011)
GRDC	River discharge	-	1981-2007	<a href="http://www.bafg.de/GRDC/EN/Home/homepage_node.html">http://www.bafg.de/GRDC/EN/Home/homepage_node.html</a>	-
Naturalized discharge	River discharge	-	1981-2007	<a href="http://www.r-arcticnet.sr.unh.edu/ObservedAndNaturalizedDischarge-Website">http://www.r-arcticnet.sr.unh.edu/ObservedAndNaturalizedDischarge-Website</a>	Shiklomanov and Lammers (2009)
ESA CCI SM v2.2	Top-soil moisture	25 km	Nov. 1978-Dec. 2014	<a href="http://esa-soilmoisture-cci.org">http://esa-soilmoisture-cci.org</a>	-
GLEAM v3.0a	Root-zone soil moisture	0.25°	1980-2014	<a href="http://www.gleam.eu">http://www.gleam.eu</a>	Martens et al. (2017)
<b>Evaluation datasets for air-to-soil temperature continuum</b>					
ECA&D	Snow depth	-	1975-2005	<a href="http://ecad.knmi.nl/dailydata/predefinedseries.php">http://ecad.knmi.nl/dailydata/predefinedseries.php</a>	-
National Climate Data and Info. Archive of Env. Canada	Snow depth	-	1975-2005	<a href="http://climate.weather.gc.ca/historical_data/search_historic_data_e.html">http://climate.weather.gc.ca/historical_data/search_historic_data_e.html</a>	-
USHCN	Snow depth	-	1975-2005	<a href="http://cdiac.ornl.gov/epubs/ndp/ushcn/ushcn.html">http://cdiac.ornl.gov/epubs/ndp/ushcn/ushcn.html</a>	-
RIHMI-WDC	Snow depth	-	1975-2005	-	Bulygina et al. (2011)
National Meteo. Info. Center of the China Meteo. Admin.	Snow depth	-	1975-2005	-	Peng et al. (2010)
-	Surface soil temperature	25km	2000-2011	<a href="http://doi.pangaea.de/10.1594/PANGAEA.833409">http://doi.pangaea.de/10.1594/PANGAEA.833409</a>	André et al. (2015)
-	In-situ air and soil temperatures	-	1980-2000	-	Sherstiukov (2012)
CALM	Active-layer thickness	-	1990-2015	-	-
For Yakutia	Active-layer thickness	-	1960-1987	<a href="https://doi.org/10.1594/PANGAEA.808240">doi:10.1594/PANGAEA.808240</a>	Beer et al. (2013)
<b>Evaluation datasets for leaf area, carbon stocks and fluxes</b>					
GIMMS	Leaf Area Index	0.08°	Jul. 1981 to Dec. 2011	<a href="http://nasanex.s3.amazonaws.com/AVHRR/GIMMS/LAI3G">http://nasanex.s3.amazonaws.com/AVHRR/GIMMS/LAI3G</a>	Zhu et al. (2013)
GLASS	Leaf Area Index	0.05°	1982-2012	<a href="http://glcf.umd.edu/data/lai">http://glcf.umd.edu/data/lai</a>	Liang and Xiao (2012)
CAMS	NEE	1.875° x 3.75°	1979-2015	<a href="https://apps-test.ecmwf.int/datasets/data/cams-ghg-inversions">https://apps-test.ecmwf.int/datasets/data/cams-ghg-inversions</a>	Chevallier et al. (2010)
Jena s96 v3.8	NEE	3.8° x 5.0°	1996-2015	<a href="http://www.bgc-jena.mpg.de/CarboScope/?ID=s96_v3.8">http://www.bgc-jena.mpg.de/CarboScope/?ID=s96_v3.8</a>	Rödenbeck (2005)
-	GPP	-	Not known	-	Campioli et al. (2015)
MTE-GPP	GPP	0.5°	1982-2010	-	Jung et al. (2009, 2011)
-	NPP	-	Not known	-	Campioli et al. (2015)
MOD17A3.005	NPP	1 km	2000-2010	-	-
NCSCD	Soil carbon inventories	0.01°	-	<a href="http://bolin.su.se/data/ncscd/netcdf.php">http://bolin.su.se/data/ncscd/netcdf.php</a>	Hugelius et al. (2013)
ISRIC	Soil carbon inventories	1 km	-	<a href="https://doi.org/10.5879/ecds/00000001">doi:10.5879/ecds/00000001</a>	Hengl et al. (2014)
-	Biomass carbon stocks	0.01°	-	-	Avitabile et al. (2016)
-	Biomass carbon stocks	0.01°	-	<a href="http://www.biomasar.orghttp://www.bgc-jena.mpg.de/geodb/projects/Home.php">http://www.biomasar.orghttp://www.bgc-jena.mpg.de/geodb/projects/Home.php</a>	Thurner et al. (2014)
GFED4s	Burned area and fire emissions	0.25°	1997-2015	<a href="http://www.globalfiredata.org/data.html">http://www.globalfiredata.org/data.html</a>	van der Werf et al. (2010)


**Table 1.** List of the data sets used for the ORCHIDEE-MICT evaluation, with their references, the original spatial resolution, and period of availability



**Figure 1.** Temperature, water and carbon interactions in the initial version of ORCHIDEE (black), and processes included in ORCHIDEE-MICT in this study (red)




**Figure 2.** (a) The three high-latitude sub-regions used in this study, including boreal North America (BONA), boreal Europe (BOEU) and boreal Asia (BOAS), following McGuire et al. (2016). Blue and red lines indicate the extent of continuous permafrost and all permafrost categories respectively, according to the IPA permafrost map (Brown et al., 2002). (b) The seven high-latitude basins selected for this study with the gauge stations (red circles on the map, more information in Table S4).




**Figure 3.** Maximum monthly snow depth (m) simulated (background maps) with (a) GSWP3 and (b) CRUNCEP forcings compared to observations (color filled circles), averaged over the period 1975-2005. (c) Monthly mean seasonal snow depth (m) from observation and the two simulations, averaged over the observation sites in the three high-latitude sub-regions (shown in Fig. 2a).


**Figure 4 (a,b)** Mean snow-density ( $\text{kg m}^{-3}$ ) and (c,d) snow-conductivity ( $\text{mW m}^{-1} \text{K}^{-1}$ ) computed over the three snow layers, averaged over the period 1981-2007. Results from the (a,c) GSWP3 and (b,d) CRUNCEP-forced simulation.




**Figure 4.** (a) Observed mean summer (JJA) land surface temperature (°C) and bias in the (b) GSWP3 and (c) CRUNCEP-forced simulation, averaged over the period 1996-2007.



**Figure 5.** Mean annual soil temperature at 0.2 m depth (°C) in the (a) GSWP3 and (b) CRUNCEP-forced simulation (background maps), compared to the site observations (color filled circles), averaged over the period 1981-2000. Monthly mean seasonal soil temperatures at different depths (°C) in the (c) GSWP3 and (d) CRUNCEP-forced simulation, compared with the observation, averaged over the 51 sites in continuous permafrost region (according to the IPA map) and over the period 1981-2000. The spatial patterns of maximum monthly soil temperature are also shown in Fig. S4.




**Figure 6.** Relationship between  $\Delta T$  (soil temperature at 20 cm depth minus air temperature) and snow depth (cm) over the period 1981-2000, for site-level observations (black), and for model results (red) (9612 site-month values in total), forced by (left) GSWP3 and (right) CRUNCEP. Circles and squares are medians of 5 cm snow depth bins, representing early (Nov, Dec, Jan) and late (Feb, Mar, Apr) snow season respectively. Upper and lower bars indicate the 25th and 75th percentiles of each bin. The size of circles/squares indicates frequency of occurrence in each bin.




**Figure 7.** Active layer thickness (ALT in m) from the (a) GSWP3 and (b) CRUNCEP-forced simulation (background maps), compared to the observed ALT from the CALM network (color filled circles), averaged over the period 1990-2007. Permafrost extent from the (c) GSWP3 and (d) CRUNCEP-forced simulation according to two different definitions (yellow and red lines) on top of the IPA permafrost map (Brown et al., 2002).







**Figure 8.** Interannual monthly variation and trend (line) of TWS (mm) simulated with the two forcings compared to GRACE data over the seven basins (see Fig. 2b), for the period July 2003-December 2007.




**Figure 9.** Monthly mean seasonal SWE (mm) simulated with the two forcings compared to GlobSnow over the seven basins (see Fig. 2b), averaged over the period 1981-2007.




**Figure 10.** Monthly mean seasonal relative (a) topsoil (-) and (b) root soil moisture (-), both normalized by their multi-year standard deviation, simulated with the GSWP3 and CRUNCEP forcings over the seven basins (see Fig. 2b), averaged over the period 1981-2007. The results are compared with (a) satellite-derived observations from ESA CCI and (b) the GLEAM data driven model assimilated against satellite data.




**Figure 11.** (a) Monthly mean seasonal evapotranspiration ( $\text{mm d}^{-1}$ ) simulated with the two forcings compared to GLEAM data-driven model over the seven basins (see Fig. 2b), averaged over the period 1981-2007. (b) Seasonal bias of ET components ( $\text{mm d}^{-1}$ ) averaged over the same period, with the GSWP3 (solid line) and CRUNCEP (dashed line) forcings.




**Figure 12.** Monthly mean seasonal river discharge ( $\text{m}^3 \text{s}^{-1}$ ) simulated with the two forcings compared to observed non-naturalized (GRDC) and the Siberian naturalized river discharge dataset at the gauge stations of the seven basins (see Fig. 2b), averaged over the period 1981-2007.




**Figure 13.** Monthly mean seasonal LAI (-) simulated with the two forcings compared to GIMMS and GLASS products over the seven basins (see Fig. 2b), averaged over the period 1981-2007.




**Figure 14. (Upper panel)** Simulated annual GPP ( $\text{gC m}^{-2} \text{ day}^{-1}$ ) from the two climate forcings, compared to the data-driven MTE-GPP, averaged over the period 2000-2007. **(Lower panel)** Monthly mean seasonal GPP ( $\text{gC m}^{-2} \text{ day}^{-1}$ ) simulated with the two forcings compared to MTE-GPP over the three high-latitude sub-regions (shown in Fig. 2a), averaged over the same period.



**Figure 15.** Mean annual NPP ( $\text{gC m}^{-2} \text{yr}^{-1}$ ) simulated with (a) GSWP3 and (b) CRUNCEP forcings (background maps) compared to the site observations (color filled circles), on averaged over the period 2000-2007.




**Figure 16.** 95th percentiles of mean annual (a) GPP ( $\text{gC m}^{-2} \text{yr}^{-1}$ ) and (b) NPP ( $\text{gC m}^{-2} \text{yr}^{-1}$ ) distributions per temperature bins of  $5^\circ\text{C}$  for in situ measurements, the gridded MTE-GPP (MODIS-NPP) product sampled at the sites location and the two simulation results, averaged over the period 2000-2007.




**Figure 17.** Mean annual CUE (%) over the 52 Campioli sites, averaged over the period 2000-2007. The first black boxplot is computed using the local estimations of the Campioli et al. (2015) database and the second one (global observations) using MODIS-NPP and MTE-GPP. The red boxplots uses the values of the two simulations. For each boxplot the median value is the short horizontal bar within the rectangle, whose bottom and top sides illustrate the 25th and 75th percentiles of the distribution, while the vertical segments link those sides to the points representing respectively the minimum and maximum values.






**Figure 18.** (a-c) Mean annual fractional burned area (%) from (a) satellite observation in GFED4s and (b) GSWP3 and (c) CRUNCEP-forced simulations. (d-f) Mean annual carbon emissions from natural fires ( $\text{gC} \cdot \text{m}^{-2} \cdot \text{yr}^{-1}$ ) from (d) satellite observation in GFED4s and (e) GSWP3 and (f) CRUNCEP-forced simulations, averaged over the period 1997-2007. The burned area fraction simulated from ORCHIDEE-MICT is corrected for the omission of cropland fires in the simulation.







**Figure 19.** Monthly mean seasonal net land-atmosphere CO<sub>2</sub> fluxes (NEE in PgCmonth<sup>-1</sup>) from the GSWP3 and CRUNCEP-forced simulations compared to atmospheric inversions, over the three high-latitude sub-regions (shown in Fig. 2a), averaged over the period 2000-2007. The grey shaded areas corresponds to the standard deviation of the monthly values for each inversion. Note that a negative sign in NEE corresponds to CO<sub>2</sub> uptake from the atmosphere, and a positive sign to release to the atmosphere.




**Figure 20.** Total forest biomass carbon density (MgCha<sup>-1</sup>) from the (a) GSWP3 and (b) CRUNCEP-forced simulations compared with satellite-derived observation products from (c) Avitabile et al. (2016) and (d) Thurner et al. (2014), averaged over the period 2000-2007.



**Figure 21.** Soil organic carbon from the GSWP3 and CRUNCEP-forced simulations compared with the two inventory datasets NCSCD (Hugelius et al., 2013) and ISRIC, averaged over the period 2000-2007. (**Upper panel**) Spatial distribution ( $\text{kgC m}^{-2}$ ). (**Lower panel**) Vertical profiles ( $\text{kgC m}^{-3}$ ) averaged over the three high-latitude sub-regions (shown in Fig. 2a). Since NCSCD does not encompass the whole domain, only grid cells with available data in NCSCD are averaged for BONA and BOAS so that the four vertical profiles are comparable; while for BOEU, NCSCD is not shown as it has few data in this sub-region.



**Figure 22.** Scatter plot of the pre-melt SWE bias ( $\text{m}^3 \text{s}^{-1}$ ) defined as the difference between simulated and observed values during the pre-melt season (between February and March) vs. the spring river discharge bias (total discharge at the mouth of the river between April and June) ( $\text{m}^3 \text{s}^{-1}$ ), averaged over the period 1981-2007. Each number corresponds to one basin: 1: Yukon, 2: Mackenzie, 3: Volga, 4: Ob, 5: Yenisei, 6: Lena and 7: Kolyma (see Fig. 2b). One color represents the result with one atmospheric forcing.



**Figure 23.** Annual mean land carbon fluxes ( $\text{PgCyr}^{-1}$ ) and pool sizes ( $\text{PgC}$ ), averaged over the terrestrial domain higher than  $30^\circ\text{N}$ , for the period 2000-2007. The red numbers are the results of the model forced by (left) GSWP3 and (right) CRUNCEP. The black numbers are from observation-based estimates used in the text : NPP from MODIS (NTSG), fire emissions from (left) GFED4s and (right) GFAS datasets, harvest fluxes include crop harvest and wood product decay in the model (for simplicity the change in wood product pools is not represented), NEE from the two atmospheric inversions (left) Jena CarboScope and (right) CAMS, forest biomass from (left) Avitabile et al. (2016) and (right) Thurner et al. (2014), soil carbon from NCSCD (Hugelius et al., 2013) in permafrost regions and HWSD in non-permafrost regions.

# ORCHIDEE-MICT (~~revision 4126~~)(v8.4.1), a land surface model for the high-latitudes: model description and validation

Guimberteau et al.

## Soil thermal parameters

	$\theta_{\text{sat}}$ (or porosity) ( $\text{m}^3 \text{m}^{-3}$ )	$\lambda_{\text{dry}}$ ( $\text{W m}^{-1} \text{K}^{-1}$ )	$\lambda_{\text{solid}}$ ( $\text{W m}^{-1} \text{K}^{-1}$ )	$c_{\text{dry}}$ ( $10^6 \text{J K}^{-1} \text{m}^{-3}$ )
sand	0.43	0.22	6.91	1.47
loamy sand	0.41	0.23	6.04	1.41
sandy loam	0.41	0.23	4.49	1.34
silt loam	0.45	0.20	2.80	1.27
silt	0.46	0.20	3.30	1.21
loam	0.43	0.22	3.43	1.21
sandy clay loam	0.39	0.25	4.49	1.18
silty clay loam	0.43	0.22	3.30	1.32
clay loam	0.41	0.23	3.21	1.23
sandy clay	0.38	0.26	4.03	1.18
silty clay	0.36	0.28	3.30	1.15
clay	0.38	0.26	2.80	1.09
pure organic soil	0.92	0.05	0.25	2.50

$\theta_{\text{sat}}$ : volumetric moisture content at saturation (porosity);  $\lambda_{\text{dry}}$  : dry soil thermal conductivity, calculated by the empirical equation:

$$\lambda_{\text{dry}} = \frac{0.135\rho+64.7}{2700-0.947\rho} \text{ with } \rho = 2700 (1 - \theta_{\text{sat}})$$

$\lambda_{\text{solid}}$  : thermal conductivity of soil solids;  $c_{\text{dry}}$  : dry soil heat capacity.

**Table S1.** Soil thermal parameters for mineral soil (including 12 USDA soil textures) and organic soil used in Eq. 4.

## Soil hydraulic parameters

	$\theta_r$ ( $\text{m}^3 \text{m}^{-3}$ )	$\alpha$	n	$\theta_{fc}$ ( $\text{m}^3 \text{m}^{-3}$ )	$\theta_{wp}$ ( $\text{m}^3 \text{m}^{-3}$ )
sand <sup>a</sup>	0.045	1.45	2.68	0.049	0.045
loamy sand <sup>a</sup>	0.057	1.24	2.28	0.071	0.057
sandy loam <sup>a</sup>	0.065	0.75	1.89	0.122	0.066
silt loam	0.067	0.20	1.41	0.240	0.104
silt	0.034	0.16	1.37	0.258	0.090
loam	0.078	0.36	1.56	0.165	0.088
sandy clay loam	0.100	0.59	1.48	0.169	0.111
silty clay loam	0.089	0.10	1.23	0.338	0.197
clay loam	0.095	0.19	1.31	0.270	0.150
sandy clay	0.100	0.27	1.23	0.267	0.170
silty clay	0.070	0.05	1.09	0.337	0.266
clay	0.068	0.08	1.09	0.347	0.271

**Table S2.** Coefficients in the van Genuchten equation (Eq. 11) and derived field capacity ( $\theta_{fc}$ ) and permanent wilting point ( $\theta_{wp}$ ) for the 12 USDA soil textures. <sup>a</sup>: for the three sandy soils, field capacity corresponds to  $\psi = -10$  kPa instead of -33 kPa.

## Number of ignitions in the SPITFIRE module in ORCHIDEE-MICT

Region	BONA	TENA	CEAM	NHSA	SHSA	EURO	MIDE	NHAF	SHAF	BOAS	CEAS	SEAS	EQAS	AUST
Ratio	2.09	0.04	1.02	0.62	0.67	1.20	2.85	56.24	31.67	3.39	1.68	12.64	99.80	3.56

**Table S3.** Region-specific values used to scale the simulated number of ignitions in the SPITFIRE module in ORCHIDEE-MICT. Refer to <http://www.globalfiredata.org/data.html> for the acronyms of different regions.

### Selected northern watersheds for discharge and water balance evaluation

Station name	Basin name	Longitude (°E)	Latitude (°N)	Basin area (km <sup>2</sup> )	$Q_{\text{mean}}$ GRDC (m <sup>3</sup> s <sup>-1</sup> )	Naturalized $Q_{\text{mean}}$ (m <sup>3</sup> s <sup>-1</sup> )	Permafrost area (%)
Pilot station	Yukon	-162.88	61.93	831390	6553	-	87
Arctic Red River	Mackenzie	-133.75	67.46	1660000	9044	-	42
Volgograd Power Plant	Volga	44.59	48.81	1360000	8246	-	-
Salekhard	Ob	66.53	66.57	2949998	12960	13005	4
Igarka	Yenisei	86.50	67.48	2440000	19400	19234	42
Kusur	Lena	127.65	70.70	2430000	17550	17426	90
Kolymaskaya	Kolyma	158.72	68.73	526000	3252	-	100


**Table S4.** List of the gauge stations of the seven major Arctic rivers studied.  $Q_{\text{mean}}$  is the mean annual discharge from the two databases, averaged over the period 1981-2007. Permafrost coverage (continuous + discontinuous permafrost) is calculated using data from Brown et al. (1998) over the entire basin considered (Holmes et al., 2012).

### Relationships between river discharge bias in Spring and pre-melt SWE bias

Basin name	Pre-melt SWE bias (m <sup>3</sup> s <sup>-1</sup> )		Spring flow bias (m <sup>3</sup> s <sup>-1</sup> )		Spring flow bias explained by the pre-melt SWE bias (%)	
	[average Feb.-Mar.]		[sum Apr.-Jun.]			
	GSWP3	CRUNCEP	GSWP3	CRUNCEP	GSWP3	CRUNCEP
1: Yukon	+1825	-6420	+13180	-18900	14	34
2: Mackenzie	+2700	-3200	+38350	+1880	7	-170
3: Volga	+9140	+2615	+57360	+30030	16	9
4: Ob	+11825	-3510	+64960	+30730	18	-11
5: Yenisei	+6235	-4730	+6430	-46470	97	10
6: Lena	-530	-9515	+18160	-39550	-3	24
7: Kolyma	+560	-2580	+4430	-11175	13	23


**Table S5.** SWE bias during the pre-melt season (average between February and March) and the bias of discharge in spring (total discharge between April and June) (both in m<sup>3</sup> s<sup>-1</sup>) with the two forcings in each basin, over the period 1981-2007. The two last columns are the contribution of the pre-melt SWE bias to the spring discharge bias (in %). Bias is defined here exactly like in Fig. 232.

## Soil vertical discretization in ORCHIDEE-MICT




**Figure S1.** Soil vertical discretization in ORCHIDEE-MICT. Solid lines and black dots represent the interfaces and nodes of each layer respectively. The soil hydrological module has 11 layers to 2 meters; the soil thermal and soil carbon modules have 32 layers to 38 meters, in which the first 11 layers are identical to that in hydrology, except that the nodes of the 1st and 11th layer in hydrology lie at the interface.

## Soil moisture content as a function of soil organic carbon



**Figure S2.** Parameterization of soil moisture content at field capacity ( $\theta_{fc}$ ) and permanent wilting point ( $\theta_{wp}$ ) as a function of soil organic carbon, for three representative soil textures in the northern high latitudes. The difference between  $\theta_{fc}$  and  $\theta_{wp}$  represents the water holding capacity of the soil as a function of SOC.


## Snow density and conductivity



**Figure S3.** (a,b) Mean snow density ( $\text{kg m}^{-3}$ ) and (c,d) snow conductivity ( $\text{mW m}^{-1} \text{K}^{-1}$ ) computed over the three snow layers, averaged over the period 1981-2007. Results from the (a,c) GSWP3 and (b,d) CRUNCEP-forced simulation.




## Maximum monthly soil temperature




**Figure S4.** Maximum monthly soil temperature at different depths in **(a)** GSWP3 and **(b)** CRUNCEP-forced simulation (background maps), compared to the site observations (color filled circles), averaged over the period 1981-2000.

# Active layer thickness in North Eastern Siberia




**Figure S35.** (a) Active layer thickness (ALT) of Yakutia, East Siberia, upscaled based on the map of landscapes and permafrost conditions in Yakutia by Beer et al. (2013). (b,c) Modeled ALT for the same region with GSWP3 and CRUNCEP, averaged over the period 1960-1987 to be consistent with Beer et al. (2013).



**Figure S6.** (a) Scatter plot of modeled ALT forced by CRUNCEP compared with observed ALT from the CALM network, averaged over the period 1990-2007. The black circles represent the grid cells taken from the regional simulation (shown in Fig. 8b). The blue circles represent a subset of the CALM sites for which we performed additional runs using site-specific organic layer thicknesses, with the result shown by the red circles. (b) Illustration for the difference of the additional site simulations. Each grey arrow connects the same site, pointing from the blue circles using soil carbon content from NCSCD (upscaled at  $1^\circ$  by  $1^\circ$  resolution) to calculate soil physical properties, to the red circles using the organic layer thickness provided by the sites. See text for further information.


## Albedo



**Figure S47.** Albedo (left) observed (ESA-GlobAlbedo) for the months of (**Upper panel**) February and (**Lower panel**) August, averaged over the period 1998-2011. Albedo bias with (**middle**) GSWP3 and (**right**) CRUNCEP climate forcing.


In February, when the snow cover is maximum, the plots show an overall underestimation of the modeled albedo up to 0.25 in the snow-covered regions particularly with CRUNCEP forcing and a slight overestimation elsewhere (Fig. S47). These differences may be partly explained by the SWE underestimation already highlighted and also by weaknesses in the parametrization of the albedo, especially over high vegetation (forested areas in North America, central Europe and Siberia). In these forested regions, the albedo seems to be overestimated in presence of snow. In August, the comparison shows very similar values in the modeled product whatever the atmospheric forcing used and the differences with the observations are strongly related to the land cover map. Grasslands in the northern regions presents underestimated albedo values up to 0.1, whereas the albedo is overestimated slightly elsewhere with a bias of about +0.02. These discrepancies may be explained by the LAI summer lag already discussed which leads to an overestimation of the vegetation cover in August and consequently an underestimation of the albedo (since the albedo of bare soils is generally larger than those of vegetation).

## Latitudinal distribution of fire carbon emissions




**Figure S58.** Latitudinal distribution of total annual forest biomass carbon stocks ( $\text{PgC degree}^{-1}$ ) in the GSWP3 and CRUNCEP-forced simulations, compared to observations by Avitabile et al. (2016) and Thurner et al. (2014) for **(a)** the whole model domain. High-latitude sub-regions of **(b)** BONA, **(c)** BOEU and **(d)** BOAS have been extended to  $30^\circ\text{N}$  here compared to their shapes in Fig. 2a.

## Seasonal cycle of burned area and fire carbon emissions




**Figure S69.** Monthly mean seasonal **(a-c)** fire burned area ( $\text{Mha month}^{-1}$ ) and **(d-f)** carbon emissions to the atmosphere ( $\text{TgC month}^{-1}$ ) from the GFED4s dataset and the two simulations, averaged over the period 1997–2007. For carbon emissions, cropland and natural fires are represented separately.

## LAI and GPP seasonality between deciduous and evergreen forests



**Figure S710.** Monthly mean seasonal (first and third panels) GPP (gCm<sup>-2</sup> day<sup>-1</sup>) and (second and fourth panels) LAI (-) for deciduous (light green) and conifer forests (dark green), over the three high-latitude sub-regions (shown in Fig. 2a), averaged over the period 2000-2007. The lower panel shows absolute values and the upper one values scaled for a better comparison.


## Land carbon fluxes and pools modeled and derived from observations



**Figure S811.** Annual mean land carbon fluxes (PgCyr<sup>-1</sup>) and pool sizes (PgC) over the three high-latitude sub-regions (shown in Fig. 2a) **(a) BONA**, **(b) BOEU** and **(c) BOAS**, averaged over the period 2000-2007. The red numbers are the model results forced by (left) GSWP3 and (right) CRUNCEP. The black numbers are from observation-based estimates used in the text : NPP from MODIS (NTSG), fire emissions from (left) GFED4s and (right) GFAS datasets, harvest fluxes include crop harvest and wood product decay in the model (for simplicity the change in wood product pools is not represented), NEE from the two atmospheric inversions (left) Jena CarboScope and (right) CAMS, biomass from (left) Avitabile et al. (2016) and (right) Thurner et al. (2014), litter from Pan et al. (2011), soil carbon from NCSCD (Hugelius et al., 2013) in permafrost regions and HWSD in non-permafrost regions.




Differences of atmospheric variables between GSWP3 and CRUNCEP forcings (GSWP3 minus CRUNCEP)




**Figure S912.** Difference in monthly air temperature (°C) between GSWP3 and CRUNCEP (GSWP3 minus CRUNCEP), averaged over the period 1981-2007.






**Figure S103.** Same as Fig. S912 but for snowfall ( $\text{mm yr}^{-1}$ )




**Figure S114.** Same as Fig. S912 but for total precipitation ( $\text{mm yr}^{-1}$ )




**Figure S125.** Same as Fig. S912 but for downward shortwave radiation ( $\text{W m}^{-2}$ )







**Figure S136.** Same as Fig. S912 but for downward longwave radiation ( $\text{W m}^{-2}$ )



**Figure S147.** Same as Fig. S912 but for specific humidity ( $\text{g kg}^{-1}$ )



**Figure S158.** Same as Fig. S912 but for wind speed at 10 m ( $\text{ms}^{-1}$ )



**Figure S169.** Same as Fig. S912 but for wind speed at 2 m in ORCHIDEE outputs ( $\text{m s}^{-1}$ )



## References

- Avitabile, V., Herold, M., Heuvelink, G. B. M., Lewis, S. L., Phillips, O. L., Asner, G. P., Armston, J., Ashton, P. S., Banin, L., Bayol, N., Berry, N. J., Boeckx, P., de Jong, B. H. J., DeVries, B., Girardin, C. A. J., Kearsley, E., Lindsell, J. A., Lopez-Gonzalez, G., Lucas, R., Malhi, Y., Morel, A., Mitchard, E. T. A., Nagy, L., Qie, L., Quinones, M. J., Ryan, C. M., Ferry, S. J. W., Sunderland, T., Laurin, G. V.,  
5 Gatti, R. C., Valentini, R., Verbeeck, H., Wijaya, A., and Willcock, S.: An integrated pan-tropical biomass map using multiple reference datasets, *Glob. Change Biol.*, 22, 1406–1420, doi:10.1111/gcb.13139, 2016.
- Beer, C., Fedorov, A. N., and Torgovkin, Y.: Permafrost temperature and active-layer thickness of Yakutia with 0.5-degree spatial resolution for model evaluation, *Earth Syst. Sci. Data*, 5, 305–310, doi:10.5194/essd-5-305-2013, <http://www.earth-syst-sci-data.net/5/305/2013/>, 2013.
- 10 Brown, J., Ferrians Jr, O., Heginbottom, J., and Melnikov, E.: Revised February 2001. Circum-Arctic map of permafrost and ground-ice conditions. Boulder, CO: National Snow and Ice Data Center/World Data Center for Glaciology, Digital media, 1998.
- Holmes, R. M., McClelland, J. W., Peterson, B. J., Tank, S. E., Bulygina, E., Eglinton, T. I., Gordeev, V. V., Gurtovaya, T. Y., Raymond, P. A., Repeta, D. J., Staples, R., Striegl, R. G., Zhulidov, A. V., and Zimov, S. A.: Seasonal and Annual Fluxes of Nutrients and Organic Matter from Large Rivers to the Arctic Ocean and Surrounding Seas, *Estuaries Coasts*, 35, 369–382, doi:10.1007/s12237-011-9386-6, 2012.
- 15 Hugelius, G., Tarnocai, C., Broll, G., Canadell, J., Kuhry, P., and Swanson, D.: The Northern Circumpolar Soil Carbon Database: spatially distributed datasets of soil coverage and soil carbon storage in the northern permafrost regions, *Earth Syst. Sci. Data*, 5, 3, doi:10.5194/essdd-5-707-2012, 2013.
- Pan, Y., Chen, J. M., Birdsey, R., McCullough, K., He, L., and Deng, F.: Age structure and disturbance legacy of North American forests, *Biogeosciences*, 8, 715, doi:10.5194/bg-8-715-2011, 2011.
- 20 Thurner, M., Beer, C., Santoro, M., Carvalhais, N., Wutzler, T., Schepaschenko, D., Shvidenko, A., Kompter, E., Ahrens, B., Levick, S. R., and Schmullius, C.: Carbon stock and density of northern boreal and temperate forests, *Global Ecol. Biogeogr.*, 23, 297–310, doi:10.1111/geb.12125, <http://dx.doi.org/10.1111/geb.12125>, 2014.