

Interactive comment on “3D-Var versus Optimal Interpolation for Aerosol Assimilation: a Case Study over the Contiguous United States” by Youhua Tang et al.

Y. Tang

Youhua.tang@noaa.gov

Received and published: 12 July 2017

Thank you for your prompt comments. In this study, the observations and observation errors (constant 0.1) used in these two assimilation tools are same. However, their model errors are not the same, which is partly due to their different engineering structures. As we mentioned that GSI is 3D-var system, and its error covariance and length scales have vertical variation as shown in Figure 1. The OI assimilation is made in horizontally in 11×11 grid box (12km resolution each grid), so its horizontal length scale is approximately $11/2 \times 12 = 66$ km. Vertically, OI applies its adjustment ratio up to PBL top for surface PM_{2.5} assimilation, or whole column for AOD assimilation over

C1

each grid location. OI's model errors for surface PM_{2.5} assimilation have diurnal variation and are varied from location to location based on the raw run's statistics, which is carried out in this study (Figure 2 of Tang et al., 2015). The GSI's model errors have no temporal variation or horizontal changes, but have vertical variations. Their biggest behaviors difference can be seen from the Figures 4a,b: the OI assimilation mainly affect local or nearby grids while GSI 3D-var increment expands more broadly.

Interactive comment on Geosci. Model Dev. Discuss., <https://doi.org/10.5194/gmd-2017-147>, 2017.

C2

Model Relative Uncertainties use in OI's surface PM_{2.5} assimila




Figure 2 of Tang, Y., T. Chai, L. Pan, P. Lee, Daniel Tong, H. Kim, and W. Chen, Using Interpolation to Assimilate Surface Measurements and Satellite AOD for Ozone and for July 2011. Journal of the Air & Waste Management Association, DOI:10.1080/10962247.2015.1062439, 2015