

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14 **The Carbon Dioxide Removal Model Intercomparison Project**  
15 **(CDR-MIP): Rationale and experimental design**

16  
17  
18 David P. Keller<sup>1,\*</sup>, Andrew Lenton<sup>2,3</sup>, Vivian Scott<sup>4</sup>, Naomi E. Vaughan<sup>5</sup>, Nico  
19 Bauer<sup>6</sup>, Duoying Ji<sup>7</sup>, Chris D. Jones<sup>8</sup>, Ben Kravitz<sup>9</sup>, Helene Muri<sup>10</sup>, Kirsten  
20 Zickfeld<sup>11</sup>

21  
22  
23  
24  
25  
26  
27  
28  
29  
30 \*Corresponding author email: dkeller@geomar.de

31 <sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany

32 <sup>2</sup>CSIRO Oceans and Atmospheres, Hobart, Australia

33 <sup>3</sup>Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Australia

34 <sup>4</sup>School of GeoSciences, University of Edinburgh

35 <sup>5</sup>Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of  
36 East Anglia, Norwich, UK.

37 <sup>6</sup>Potsdam Institute for Climate Impact Research, Research Domain Sustainable Solutions, 14473  
38 Potsdam, Germany

39 <sup>7</sup>College of Global Change and Earth System Science, Beijing Normal University, Beijing, China

40 <sup>8</sup> Met Office Hadley Centre, Exeter, UK,

41 <sup>9</sup>Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory,  
42 Richland, WA, USA.

43 <sup>10</sup>Department of Geosciences, University of Oslo, Oslo, Norway.

44 <sup>11</sup>Department of Geography, Simon Fraser University, Burnaby, Canada

45  
46

47 **Abstract**

48  
49 The recent IPCC reports state that continued anthropogenic greenhouse gas  
50 emissions are changing the climate, threatening "severe, pervasive and  
51 irreversible" impacts. Slow progress in emissions reduction to mitigate climate  
52 change is resulting in increased attention on what is called *Geoengineering*,  
53 *Climate Engineering*, or *Climate Intervention* – deliberate interventions to counter  
54 climate change that seek to either modify the Earth's radiation budget or remove  
55 greenhouse gases such as CO<sub>2</sub> from the atmosphere. When focused on CO<sub>2</sub>, the  
56 latter of these categories is called Carbon Dioxide Removal (CDR). Future  
57 emission scenarios that stay well below 2°C, and all emission scenarios that do  
58 not exceed 1.5°C warming by the year 2100, require some form of CDR. At  
59 present, there is little consensus on the climate impacts and atmospheric CO<sub>2</sub>  
60 reduction efficacy of the different types of proposed CDR. To address this need  
61 the Carbon Dioxide Removal Model Intercomparison Project (or CDR-MIP) was  
62 initiated. This project brings together models of the Earth system in a common  
63 framework to explore the potential, impacts, and challenges of CDR. Here, we  
64 describe the first set of CDR-MIP experiments that are designed to address  
65 questions concerning CDR-induced climate "reversibility", the response of the  
66 Earth system to direct atmospheric CO<sub>2</sub> removal (direct air capture and storage),  
67 and the CDR potential and impacts of afforestation/reforestation, as well as  
68 ocean alkalinization.

69

70

71 **1. Introduction**

72

73 The Earth system is sensitive to the concentration of atmospheric  
74 greenhouse gases (GHG) because they have a direct impact on the planetary  
75 energy balance (Hansen, 2005), and in many cases also on biogeochemical  
76 cycling (IPCC, 2013). The concentration of one particularly important GHG,  
77 carbon dioxide (CO<sub>2</sub>), has increased from approximately 277 ppm in the year  
78 1750 to over 400 ppm today as a result of anthropogenic activities (Dlugokencky  
79 and Tans, 2016; Le Quéré et al., 2015). This CO<sub>2</sub> increase, along with other GHG  
80 increases and anthropogenic activities (e.g. land use change), has perturbed the  
81 Earth's energy balance leading to an observed global mean surface air  
82 temperature increase of around 0.8 °C above preindustrial (year 1850) levels in  
83 the year 2015 [updated from Morice et al. (2012)]. Biogeochemistry on land and  
84 in the ocean has also been affected by the increase in CO<sub>2</sub>, with a well-observed  
85 decrease in ocean pH being one of the most notable results (Gruber, 2011;  
86 Hofmann and Schellnhuber, 2010). Many of the changes attributed to this rapid  
87 temperature increase and perturbation of the carbon cycle have been  
88 detrimental for natural and human systems (IPCC, 2014a).

89 While recent trends suggest that the atmospheric CO<sub>2</sub> concentration is  
90 likely to continue to increase (Peters et al., 2013; Riahi et al., 2017), the Paris  
91 Agreement of the 21st session of the Conference of Parties (COP21) on climate  
92 change (UNFCCC, 2016) has set the goal of limiting anthropogenic warming to  
93 well below 2°C (ideally no more than 1.5°C) relative to the global mean  
94 preindustrial temperature. To do this a massive climate change mitigation effort  
95 to reduce the sources or enhance the sinks of greenhouse gases (IPCC, 2014b)  
96 must be undertaken. Even if significant efforts are made to reduce CO<sub>2</sub>  
97 emissions, it will likely take decades before net emissions approach zero (Bauer  
98 et al., 2017; Riahi et al., 2017; Rogelj et al., 2015a), a level that is likely required  
99 to reach and maintain such temperature targets (Rogelj et al., 2015b). Changes  
100 in the climate will therefore continue for some time, with future warming  
101 strongly dependent on cumulative CO<sub>2</sub> emissions (Allen et al., 2009; IPCC, 2013;  
102 Matthews et al., 2009), and there is the possibility that “severe, pervasive and  
103 irreversible” impacts will occur if too much CO<sub>2</sub> is emitted (IPCC, 2013, 2014a).

104 The lack of agreement on how to sufficiently reduce CO<sub>2</sub> emissions in a timely  
105 manner, and the magnitude of the task required to transition to a low carbon  
106 world has led to increased attention on what is called *Geoengineering*, *Climate*  
107 *Engineering*, or *Climate Intervention*. These terms are all used to define actions  
108 that deliberately manipulate of the climate system in an attempt to ameliorate or  
109 reduce the impact of climate change by either modifying the Earth's radiation  
110 budget (Solar Radiation Management, or SRM), or by removing the primary  
111 greenhouse gas, CO<sub>2</sub>, from the atmosphere (Carbon Dioxide Removal, or CDR)  
112 (National Research Council, 2015). In particular, there is an increasing focus and  
113 study on the potential of carbon dioxide removal (CDR) methods to offset  
114 emissions and eventually to enable "net negative emissions", whereby more CO<sub>2</sub>  
115 is removed via CDR than is emitted by anthropogenic activities, to complement  
116 emissions reduction efforts. CDR has also been proposed as a means of  
117 "reversing" climate change if too much CO<sub>2</sub> is emitted, i.e., CDR may be able to  
118 reduce atmospheric CO<sub>2</sub> to return radiative forcing to some target level.

119 All Integrated Assessment Model (IAM) scenarios of the future state that  
120 some form of CDR will be needed to prevent the mean global surface  
121 temperature from exceeding 2°C (Bauer et al., 2017; Fuss et al., 2014; Kriegler et  
122 al., 2016; Rogelj et al., 2015a). Most of these limited warming scenarios feature  
123 overshoots in radiative forcing around mid-century, which is closely related to  
124 the amount of cumulative CDR up until the year 2100 (Kriegler et al., 2013).  
125 Despite the prevalence of CDR in these scenarios, and its increasing utilization in  
126 political and economic discussions, many of the methods by which this would be  
127 achieved at this point rely on immature technologies (National Research Council,  
128 2015; Schäfer et al., 2015). Large scale CDR methods are not yet a commercial  
129 product, and hence questions remain about their feasibility, realizable potential  
130 and risks (Smith et al., 2015; Vaughan and Gough, 2016).

131 Overall, knowledge about the potential climatic, biogeochemical,  
132 biogeophysical, and other impacts in response to CDR is still quite limited, and  
133 large uncertainties remain, making it difficult to comprehensively evaluate the  
134 potential and risks of any particular CDR method and make comparisons  
135 between methods. This information is urgently needed to allow us to assess:

137       i. The degree to which CDR could help mitigate or perhaps reverse climate  
138       change;

140       ii. The potential risks/benefits of different CDR proposals; and

142       iii. To inform how climate and carbon cycle responses to CDR could be  
143       included when calculating and accounting for the contribution of CDR in  
144       mitigation scenarios, i.e., so that CDR is better constrained when it is  
145       included in IAM generated scenarios.

146  
147       To date, modelling studies of CDR focusing on the carbon cycle and  
148       climatic responses have been undertaken with only a few Earth system models  
149       (Arora and Boer, 2014; Boucher et al., 2012; Cao and Caldeira, 2010; Gasser et al.,  
150       2015; Jones et al., 2016a; Keller et al., 2014; MacDougall, 2013; Mathesius et al.,  
151       2015; Tokarska and Zickfeld, 2015; Zickfeld et al., 2016). However, as these  
152       studies all use different experimental designs, their results are not directly  
153       comparable, consequently building a consensus on responses is challenging. A  
154       model intercomparison study with Earth System Models of Intermediate  
155       Complexity (EMICS) that addresses climate reversibility, among other things, has  
156       recently been published (Zickfeld et al., 2013), but the focus was on the very  
157       distant future rather than this century. Moreover, in many of these studies,  
158       atmospheric CO<sub>2</sub> concentrations were prescribed rather than being driven by  
159       CO<sub>2</sub> emissions and thus, the projected changes were independent of the strength  
160       of feedbacks associated with the carbon cycle.

161       Given that Earth system models are one of the few tools available for  
162       making quantifications at these scales, as well as for making projections into the  
163       future, CDR assessments must include emissions-driven modeling studies to  
164       capture the carbon-cycle feedbacks. However, such an assessment cannot be  
165       done with one or two models alone, since this will not address uncertainties due  
166       to model structure and internal variability. Below we describe the scientific foci  
167       and several experiments (Table 1) that comprise the initial phase of the Carbon  
168       Dioxide Removal Model Intercomparison Project (CDR-MIP).

170 **1.2 CDR-MIP Scientific Foci**

171

172        There are three principal science motivations behind CDR-MIP. First and  
173 foremost, CDR-MIP will provide information that can be used to help assess the  
174 potential and risks of using CDR to address climate change. A thorough  
175 assessment will need to look at both the impacts of CDR upon the Earth system  
176 and human society. CDR-MIP will focus primarily on Earth system impacts, with  
177 the anticipation that this information will also be useful for understanding  
178 potential impacts upon society. The scientific outcomes will lead to more  
179 informed decisions about the role CDR may play in climate change mitigation  
180 (defined here as a human intervention to reduce the sources or enhance the  
181 sinks of greenhouse gases). CDR-MIP experiments will also provide an  
182 opportunity to better understand how the Earth system responds to  
183 perturbations, which is relevant to many of the Grand Science Challenges posed  
184 by the World Climate Research Program (WCRP; [https://www.wcrp-](https://www.wcrp-climate.org/grand-challenges/grand-challenges-overview)  
185 [climate.org/grand-challenges/grand-challenges-overview](https://www.wcrp-climate.org/grand-challenges/grand-challenges-overview)). CDR-MIP  
186 experiments provide a unique opportunity because the perturbations are often  
187 opposite in sign to previous CMIP perturbation experiments (CO<sub>2</sub> is removed  
188 instead of added). Second, CDR-MIP results may also be able to provide  
189 information that helps to understand how model resolution and complexity  
190 cause systematic model bias. In this instance, CDR-MIP experiments may be  
191 especially useful for gaining a better understanding of the similarities and  
192 differences between global carbon cycle models because we invite a diverse  
193 group of models to participate in CDR-MIP. Finally, CDR-MIP results can help to  
194 quantify uncertainties in future climate change scenarios, especially those that  
195 include CDR. In this case CDR-MIP results may be useful for calibrating CDR  
196 inclusion in IAMs during the scenario development process.

197        The initial foci that are addressed by CDR-MIP include (but are not limited  
198 to):

199

200        (i) Climate “reversibility”: assessing the efficacy of using CDR to return high  
201 future atmospheric CO<sub>2</sub> concentrations to lower levels. This topic is highly  
202 idealized, as the technical ability of CDR methods to remove such enormous

203 quantities of CO<sub>2</sub> on relatively short timescales (i.e., this century) is doubtful.  
204 However, the results will provide information on the degree to which a changing  
205 and changed climate could be returned to a previous state. This knowledge is  
206 especially important since socio-economic scenarios that limit global warming to  
207 well below 2° C often feature radiative forcing overshoots that must be  
208 "reversed" using CDR. Specific questions on reversibility will address:

209

210 1) What components of the Earth's climate system exhibit "reversibility"  
211 when CO<sub>2</sub> increases and then decreases? On what timescales do these  
212 "reversals" occur? And if reversible, is this complete reversibility or  
213 just on average (are there spatial and temporal aspects)?  
214 2) Which, if any, changes are irreversible?  
215 3) What role does hysteresis play in these responses?

216

217 (ii) The potential efficacy, feedbacks, and side effects of specific CDR methods.  
218 Efficacy is defined here as CO<sub>2</sub> removed from the atmosphere, over a specific  
219 time horizon, as a result of a specific unit of CDR action. This topic will help to  
220 better constrain the carbon sequestration potential and risks and/or benefits of  
221 selected methods. Together, a rigorous analysis of the nature, sign, and  
222 timescales of these CDR-related topics will provide important information for the  
223 inclusion of CDR in climate mitigation scenarios, and in resulting mitigation and  
224 adaptation policy strategies. Specific questions on individual CDR methods will  
225 address:

226

227 1) How much CO<sub>2</sub> would have to be removed to return to a specified  
228 concentration level e.g. present day or pre-industrial?  
229 2) What are the short-term carbon cycle feedbacks (e.g. rebound)  
230 associated with the method?  
231 3) What are the short- and longer-term physical/chemical/biological  
232 impacts and feedbacks, and potential side effects of the method?  
233 4) For methods that enhance natural carbon uptake, e.g., afforestation  
234 or ocean alkalization, where is the carbon stored (land and

235 ocean) and for how long (i.e. issues of permanence; at least as  
236 much as this can be calculated with these models)?

237

238 **1.3 Structure of this document**

239

240 Our motivation for preparing this document is to lay out in detail the  
241 CDR-MIP experimental protocol, which we request all modelling groups to follow  
242 as closely as possible. Firstly, in Section 2, we review the scientific background  
243 and motivation for CDR in more detail than covered in this introduction. Section  
244 3 describes some requirements and recommendations for participating in CDR-  
245 MIP and describes links to other CMIP6 activities. Section 4 describes each CDR-  
246 MIP simulation in detail. Section 5 describes the model output and data policy.  
247 Section 6 presents an outlook of potential future CDR-MIP activities and a  
248 conclusion. Section 7 describes how to obtain the model code and data used  
249 during the production of this document.

250

251 **2. Background and motivation**

252

253 At present, there are two main proposed CDR approaches, which we  
254 briefly introduce here. The first category encompasses methods that are  
255 primarily designed to enhance the Earth's natural carbon sequestration  
256 mechanisms. Enhancing natural oceanic and terrestrial carbon sinks is suggested  
257 because these sinks have already *each* taken up over a quarter of the carbon  
258 emitted as a result of anthropogenic activities (Le Quéré et al., 2016) and have  
259 the capacity to store additional carbon, although this is subject to environmental  
260 limitations. Some prominent proposed sink enhancement methods include  
261 afforestation or reforestation, enhanced terrestrial weathering, biochar, land  
262 management to enhance soil carbon storage, ocean fertilization, ocean  
263 alkalinization, and coastal management of blue carbon sinks.

264 The second general CDR category includes methods that rely primarily on  
265 technological means to directly remove carbon from the atmosphere, ocean, or  
266 land and isolate it from the climate system, e.g., storage in a geological reservoir  
267 (Scott et al., 2015). Methods that are primarily technological are suggested

268 because they may not be as limited by environmental constraints. Some  
269 prominent proposed technological methods include direct CO<sub>2</sub> air capture with  
270 storage and seawater carbon capture (and storage). One other proposed CDR  
271 method, bioenergy with carbon capture and storage (BECCS), relies on both  
272 natural processes and technology. BECCS is thus, constrained by some  
273 environmental limitations (e.g., suitable land area), but because the carbon is  
274 removed and ultimately stored elsewhere, it may have a higher CDR potential  
275 than if the same deployment area were used for a sink-enhancing CDR method  
276 like afforestation that stores carbon permanently above ground and reaches a  
277 saturation level for a given area (Smith et al., 2015).

278 From an Earth system perspective, the potential and impacts of proposed  
279 CDR methods have only been investigated in a few individual studies - see recent  
280 climate intervention assessments for a broad overview of the state of CDR  
281 research (National Research Council, 2015; Rickels et al., 2011; The Royal  
282 Society, 2009; Vaughan and Lenton, 2011) and references therein. These studies  
283 agree that CDR application at a large scale ( $\geq 1\text{Gt CO}_2\text{ yr}^{-1}$ ) would likely have a  
284 substantial impact on the climate, biogeochemistry and the ecosystem services  
285 that the Earth provides (i.e., the benefits humans obtain from ecosystems)  
286 (Millennium Ecosystem Assesment, 2005). Idealized Earth system model  
287 simulations suggest that CDR does appear to be able to limit or even reverse  
288 warming and changes in many other key climate variables (Boucher et al., 2012;  
289 Tokarska and Zickfeld, 2015; Wu et al., 2014; Zickfeld et al., 2016). However,  
290 less idealized studies, e.g., when some environmental limitations are accounted  
291 for, suggest that many methods have only a limited individual mitigation  
292 potential (Boysen et al., 2016, 2017; Keller et al., 2014; Sonntag et al., 2016).

293 Studies have also focused on the carbon cycle response to the deliberate  
294 redistribution of carbon between dynamic carbon reservoirs or permanent  
295 (geological) carbon removal. Understanding and accounting for the feedbacks  
296 between these reservoirs in response to CDR is particularly important for  
297 understanding the efficacy of any method (Keller et al., 2014). For example,  
298 when CO<sub>2</sub> is removed from the atmosphere in simulations, the rate of oceanic  
299 CO<sub>2</sub> uptake, which has historically increased in response to increasing emissions,  
300 is reduced and might eventually reverse (i.e., net outgassing), because of a

301 reduction in the air-sea flux disequilibrium (Cao and Caldeira, 2010; Jones et al.,  
302 Tokarska and Zickfeld, 2015; Vichi et al., 2013). Equally, the terrestrial  
303 carbon sink also weakens in response to atmospheric CO<sub>2</sub> removal, and can also  
304 become a source of CO<sub>2</sub> to the atmosphere (Cao and Caldeira, 2010; Jones et al.,  
305 Tokarska and Zickfeld, 2015). This 'rebound' carbon flux response that  
306 weakens or reverses carbon uptake by natural carbon sinks would oppose CDR  
307 and needs to be accounted for if the goal is to limit or reduce atmospheric CO<sub>2</sub>  
308 concentrations to some specified level (IPCC, 2013).

309 In addition to the climatic and carbon cycle effects of CDR, most methods  
310 appear to have side effects (Keller et al., 2014). The impacts of these side effects  
311 tend to be method specific and may amplify or reduce the climate change  
312 mitigation potential of the method. Some significant side effects are caused by  
313 the spatial scale (e.g., millions of km<sup>2</sup>) at which many methods would have to be  
314 deployed to have a significant impact upon CO<sub>2</sub> and global temperatures (Boysen  
315 et al., 2016; Heck et al., 2016; Keller et al., 2014). Side effects can also potentially  
316 alter the natural environment by disrupting biogeochemical and hydrological  
317 cycles, ecosystems, and biodiversity (Keller et al., 2014). For example, large-  
318 scale afforestation could change regional albedo and evapotranspiration and so  
319 have a biogeophysical impact on the Earth's energy budget and climate (Betts,  
320 2000; Keller et al., 2014). Additionally, if afforestation were done with non-  
321 native plants or monocultures to increase carbon removal rates this could impact  
322 local biodiversity. For human societies, this means that CDR-related side effects  
323 could potentially impact the ecosystem services provided by the land and ocean  
324 (e.g., food production), with the information so far suggesting that there could be  
325 both positive and negative impacts on these services. Such effects could change  
326 societal responses and strategies for climate change adaptation if large-scale  
327 CDR were to be deployed.

328 CDR deployment scenarios have focused on both preventing climate  
329 change and reversing it. While there is some understanding of how the Earth  
330 system may respond to CDR, as described above, another dynamic comes into  
331 play if CDR were to be applied to "reverse" climate change. This is because if  
332 CDR were deployed for this purpose, it would deliberately change the climate,  
333 i.e., drive it in another direction, rather than just prevent it from changing by

334 limiting CO<sub>2</sub> emissions. Few studies have investigated how the Earth system may  
335 respond if CDR is applied in this manner. The link between cumulative CO<sub>2</sub>  
336 emissions and global mean surface air temperature change has been extensively  
337 studied (IPCC, 2013). Can this change simply be reversed by removing the CO<sub>2</sub>  
338 that has been emitted since the preindustrial era? Little is known about how  
339 reversible this relationship is, or whether it applies to other Earth system  
340 properties (e.g., net primary productivity, sea level, etc.). Investigations of CDR-  
341 induced climate reversibility have suggested that many Earth system properties  
342 are "reversible", but often with non-linear responses (Armour et al., 2011;  
343 Boucher et al., 2012; MacDougall, 2013; Tokarska and Zickfeld, 2015; Wang et al.,  
344 2014; Wu et al., 2014; Zickfeld et al., 2016). However, these analyses were  
345 generally limited to global annual mean values, and most models did not include  
346 potentially important components such as permafrost or terrestrial ice sheets.  
347 Thus, there are many unknowns and much uncertainty about whether it is  
348 possible to "reverse" climate change. Obtaining knowledge about climate  
349 "reversibility" is especially important as it could be used to direct or change  
350 societal responses and strategies for adaptation and mitigation.

351

## 352 **2.1 Why a model intercomparison study on CDR?**

353

354       Although ideas for controlling atmospheric CO<sub>2</sub> concentrations were  
355 proposed in the middle of the last century, it is only recently that CDR methods  
356 have received widespread attention as climate intervention strategies (National  
357 Research Council, 2015; Schäfer et al., 2015; The Royal Society, 2009; Vaughan  
358 and Lenton, 2011). While some proposed CDR methods do build upon  
359 substantial knowledge bases (e.g., soil and forest carbon, and ocean  
360 biogeochemistry), little research into large scale CDR has been conducted and  
361 limited research resources applied (National Research Council, 2015; Oschlies  
362 and Klepper, 2017). The small number of existing laboratory studies and small-  
363 scale field trials of CDR methods were not designed to evaluate climate or carbon  
364 cycle responses to CDR. At the same time it is difficult to conceive how such an  
365 investigation could be carried out without scaling a method up to the point  
366 where it would essentially be "deployment". The few natural analogues that exist

367 for some methods (e.g., weathering or reforestation) only provide limited insight  
368 into the effectiveness of deliberate large scale CDR. As such, beyond syntheses of  
369 resource requirements and availabilities, e.g., Smith, (2016), there is a lack of  
370 observational constraints that can be applied to the assessment of the  
371 effectiveness of CDR methods. Lastly, many proposed CDR methods are pre-  
372 mature at this point and technology deployment strategies would be required to  
373 overcome this barrier (Schäfer et al., 2015), which means that they can only be  
374 studied in an idealized manner, i.e., through model simulations.

375 Understanding the response of the Earth system to CDR is urgently  
376 needed because CDR is increasingly being utilized to inform policy and economic  
377 discussions. Examples of this include scenarios that are being developed with  
378 GHG emission forcing that exceeds (or overshoots) what is required to limit  
379 global mean temperatures to 2° C or 1.5 °C, with the assumption that  
380 reversibility is possible with the future deployment of CDR. These scenarios are  
381 generated using Integrated Assessment Models, which compute the emissions of  
382 GHGs, short-lived climate forcers, and land-cover change associated with  
383 economic, technological and policy drivers to achieve climate targets. Most  
384 integrated assessment models represent BECCS as the only CDR option, with  
385 only a few also including afforestation (IPCC, 2014b). During scenario  
386 development and calibration the output from the IAMs is fed into climate models  
387 of reduced complexity, e.g., MAGICC (Model for the Assessment of Greenhouse-  
388 gas Induced Climate Change) (Meinshausen et al., 2011), to calculate the global  
389 mean temperature achieved through the scenario choices, e.g., those in the  
390 Shared Socioeconomic Pathways (SSPs) (Riahi et al., 2017). These climate  
391 models are calibrated to Earth system models or based on modelling  
392 intercomparison exercises like the Coupled Model Intercomparison Phase 5  
393 (CMIP5), where much of the climate-carbon cycle information comes from the  
394 Coupled Climate-Carbon Cycle Model Intercomparison Project (C4MIP).  
395 However, since the carbon cycle feedbacks of large-scale negative CO<sub>2</sub> emissions  
396 have not been explicitly analyzed in projects like CMIP5, with the exception of  
397 Jones et al. (2016a), many assumptions have been made about the effects of CDR  
398 on the carbon cycle and climate. Knowledge of these short-term carbon cycle

399 feedbacks is needed to better constrain the effectiveness of the CDR technologies  
400 assumed in the IAM generated scenarios.

401 This relates to the policy relevant question of whether in a regulatory  
402 framework, CO<sub>2</sub> removals from the atmosphere should be treated like emissions  
403 except for the opposite (negative) sign or if specific methods, which may or may  
404 not have long-term consequences (e.g., afforestation/reforestation vs. direct CO<sub>2</sub>  
405 air capture with geological carbon storage), should be treated differently. The  
406 lack of this kind of analyses is a knowledge gap in current climate modeling  
407 (Jones et al., 2016a) and relevant for IAM models and political decisions. There is  
408 an urgent need to close this gap since additional CDR options like the enhanced  
409 weathering of rocks on land or direct air capture continue to be included in IAMs,  
410 e.g., Chen and Tavoni (2013). For the policy relevant questions it is also  
411 important to analyze the carbon cycle effects given realistic policy scenarios  
412 rather than idealized perturbations.

413

### 414 **3. Requirements and recommendations for participation in CDR-MIP**

415

416 The CDR-MIP initiative is designed to bring together a suite of Earth  
417 System Models, Earth System Models of Intermediate Complexity (EMICs), and  
418 potentially even box models in a common framework. Models of differing  
419 complexities are invited to participate because the questions posed above cannot  
420 be answered with any single class of models. For example, ESMs are primarily  
421 suited for investigations spanning only the next century because of the  
422 computational expense, while EMICs and box models are well suited to  
423 investigate the long-term questions surrounding CDR, but are often highly  
424 parameterized and may not include important processes, e.g., cloud feedbacks.  
425 The use of differing models will also provide insight into how model resolution  
426 and complexity controls modeled short- and long-term climate and carbon cycle  
427 responses to CDR.

428 All groups that are running models with an interactive carbon cycle are  
429 encouraged to participate in CDR-MIP. We desire diversity and encourage groups  
430 to use older models, with well-known characteristics, biases and established  
431 responses (e.g. previous CMIP model versions), as well as state-of-the-art CMIP6

432 models. For longer model simulations, we would encourage modellers when  
433 possible to include additional carbon reservoirs, such as ocean sediments or  
434 permafrost, as these are not always implemented for short simulations. Models  
435 that only include atmospheric and oceanic carbon reservoirs are welcome, and  
436 will be able to participate in some experiments. All models wishing to participate  
437 in CDR-MIP must provide clear documentation that details the model version,  
438 components, and key run-time and initialization information (model time  
439 stepping, spin-up state at initialization, etc.). Furthermore, all model output must  
440 be standardized to facilitate analyses and public distribution (see Sections 4 and  
441 5).

442

### 443 **3.1 Relations to other MIPs**

444

445 There are no existing MIPs with experiments focused on climate  
446 "reversibility", direct CO<sub>2</sub> air capture (with storage), or ocean alkalinization.  
447 However, this does not mean that there are no links between CDR-MIP and other  
448 MIPs. CMIP6 and CMIP5 experiments, analyses, and assessments both provide a  
449 valuable baseline and model sensitivities that can be used to better understand  
450 CDR-MIP results and we highly recommend that participants in CDR-MIP also  
451 conduct other MIP experiments. Further, to maximize the use of computing  
452 resources CDR-MIP uses experiments from other MIPs as a control run for a  
453 CDR-MIP experiment or to provide a pathway from which a CDR-MIP experiment  
454 branches (Sections 3.2 and 4, Tables 2- 7). Principle among these is the CMIP  
455 Diagnostic, Evaluation, and Characterization of Klima (DECK) and historical  
456 experiments as detailed in Eyring et al. (2016) for CMIP6, since they provide the  
457 basis for many experiments with almost all MIPs leveraging these in some way.

458 Here, we additionally describe links to ongoing MIPs that are endorsed by  
459 CMIP6, noting that earlier versions of many of these MIPs were part of CMIP5  
460 and so provide a similar synergy for any CMIP5 models participating in CDR-MIP.

461 Given the emphasis on carbon cycle perturbations in CDR-MIP, there is a  
462 strong synergy with C4MIP which provides a baseline, standard protocols, and  
463 diagnostics for better understanding the relationship between the carbon cycle  
464 and the climate in CMIP6 (Jones et al., 2016b). For example, the C4MIP

465 emissions-driven SSP5-8.5 scenario (a high CO<sub>2</sub> emission scenario with a  
466 radiative forcing of 8.5 Wm<sup>-2</sup> in year 2100) simulation, *esm-ssp585*, is a control  
467 run and branching pathway for several CDR-MIP experiments. CDR-MIP  
468 experiments may equally be valuable for understanding model responses during  
469 related C4MIP experiments. For example, the C4MIP experiment *ssp534-over-bgc*  
470 is a concentration driven "overshoot" scenario simulation that is run in a  
471 partially coupled mode. The simulation required to analyze this experiment is a  
472 fully coupled CO<sub>2</sub> concentration driven simulation of this scenario, *ssp534-over*,  
473 from the Scenario Model Intercomparison Project (ScenarioMIP). The novel CDR-  
474 MIP experiment, *C2\_overshoot*, which is a fully coupled CO<sub>2</sub> emission driven  
475 version of this scenario, will provide additional information that can be used to  
476 extend the analyses to better understand climate-carbon cycle feedbacks.

477 The Land Use Model Intercomparison Project (LUMIP) is designed to  
478 better understand the impacts of land-use and land-cover change on the climate  
479 (Lawrence et al., 2016). The three main LUMIP foci overlap with some of the  
480 CDR-MIP foci, especially in regards to land management as a CDR method (e.g.,  
481 afforestation/reforestation). To facilitate land-use and land-cover change  
482 investigations LUMIP provides standard protocols and diagnostics for the  
483 terrestrial components of CMIP6 Earth system models. The inclusion of these  
484 diagnostics will be important for all CDR-MIP experiments performed with  
485 CMIP6 models. The CDR-MIP experiment on afforestation/reforestation, C3  
486 (*esm-ssp585-ssp126Lu-ext*), is also an extension of the LUMIP *esm-ssp585-*  
487 *ssp126Lu* simulation beyond 2100 to investigate the long-term consequences of  
488 afforestation/reforestation in a high-CO<sub>2</sub> world (Section 4.3).

489 ScenarioMIP is designed to provide multi-model climate projections for  
490 several scenarios of future anthropogenic emissions and land use changes  
491 (O'Neill et al., 2016), and provides baselines or branching for many MIP  
492 experiments. The ScenarioMIP SSP5-3.4-OS experiments, *ssp534-over* and  
493 *ssp534-over-ext*, which prescribe atmospheric CO<sub>2</sub> to follow an emission  
494 overshoot pathway that is followed by aggressive mitigation to reduce emissions  
495 to zero by about 2070, with substantial negative global emissions thereafter, are  
496 used as control runs for the CDR-MIP CO<sub>2</sub> emission driven version of this  
497 scenario. Along with the partially coupled C4MIP version of this experiment,

498 these experiments will allow for qualitative comparative analyses to better  
499 understand climate-carbon cycle feedbacks in an "overshoot" scenario with  
500 negative emissions (CDR). If it is found that the carbon cycle effects of CDR are  
501 improperly accounted for in the scenarios, then this information can be used to  
502 recalibrate older CDR-including IAM scenarios and be used to better constrain  
503 CDR when it is included in new scenarios.

504 The Ocean Model Intercomparison Project (OMIP), which primarily  
505 investigates the ocean-related origins and consequences of systematic model  
506 biases, will help to provide an understanding of ocean component functioning for  
507 models participating in CMIP6 (Griffies et al., 2016). OMIP will also establish  
508 standard protocols and output diagnostics for ocean model components. The  
509 biogeochemical protocols and diagnostics of OMIP (Orr et al., 2016) are  
510 particularly relevant for CMIP6 models participating in CDR-MIP. While the  
511 inclusion of these diagnostics will be important for all CDR-MIP experiments,  
512 these standards will be particularly important for facilitating the analysis of our  
513 marine CDR experiment, *C4* (Section 4.4).

514

### 515 **3.2 Prerequisite and recommended CMIP simulations**

516

517 The following CMIP experiments are considered prerequisites for  
518 specified CDR-MIP experiments (Tables 2- 7) and analyses:

519

520 • The CMIP prescribed atmospheric CO<sub>2</sub> pre-industrial control simulation,  
521 *piControl*. This is required for all CDR-MIP experiments (many control  
522 runs and experiment prerequisites branch from this) and is usually done  
523 as part of the spin-up process.

524

525 • The CMIP6 pre-industrial control simulation with interactively simulated  
526 atmospheric CO<sub>2</sub> (i.e., the CO<sub>2</sub> concentration is internally calculated, but  
527 emissions are zero), *esm-piControl*. This is required for CDR-MIP  
528 experiments *C2\_pi-pulse*, *C2\_overshoot*, *C3*, and *C4*.

529

530     • The CMIP 1 % per year increasing CO<sub>2</sub> simulation, *1pctCO<sub>2</sub>*, that is  
531       initialized from a pre-industrial CO<sub>2</sub> concentration with CO<sub>2</sub> then  
532       increasing by 1% per year until the CO<sub>2</sub> concentration has quadrupled  
533       (approximately 139 years). This is required for CDR-MIP experiment *C1*.

534  
535     • The CMIP6 historical simulation, *historical*, where historical atmospheric  
536       CO<sub>2</sub> forcing is prescribed along with land use, aerosols, and non-CO<sub>2</sub>  
537       greenhouse gases forcing. This is required for CDR-MIP experiment  
538       *C2\_yr2010-pulse*.

539  
540     • The CMIP6 emissions driven historical simulation, *esm-hist*, where the  
541       atmospheric CO<sub>2</sub> concentration is internally calculated in response to  
542       historical anthropogenic CO<sub>2</sub> emissions forcing. Other forcing such as land  
543       use, aerosols, and non-CO<sub>2</sub> greenhouse gases are prescribed. This is  
544       required for CDR-MIP experiments *C2\_overshoot*, *C3*, and *C4*.

545  
546     • The LUMIP *esm-ssp585-ssp126Lu* simulation, which simulates  
547       afforestation in a high CO<sub>2</sub> emission scenario, is the basis for CDR-MIP  
548       experiment *esm-ssp585-ssp126Lu-ext*.

549  
550     • The C4MIP *esm-ssp585* simulation, which is a high emission scenario and  
551       serves as a control run and branching pathway for CDR-MIP *C4*  
552       experiment.

553  
554     We also highly recommend that groups run these additional C4MIP and  
555       ScenarioMIP simulations:

556  
557     • The ScenarioMIP *ssp534-over* and *ssp534-over-ext* simulations, which  
558       prescribe the atmospheric CO<sub>2</sub> concentration to follow an emission  
559       overshoot pathway that is followed by aggressive mitigation to reduce  
560       emissions to zero by about 2070, with substantial negative global  
561       emissions thereafter. These results can be qualitatively compared to CDR-

562 MIP experiment *C2\_overshoot*, which is the same scenario, but driven by  
563 CO<sub>2</sub> emissions.

564

- 565 • The C4MIP *ssp534-over-bgc* and *ssp534-over-bgcExt* simulations, which  
566 are biogeochemically-coupled versions of the *ssp534-over* and *ssp534-*  
567 *over-ext* simulations, i.e., only the carbon cycle components (land and  
568 ocean) see the prescribed increase in the atmospheric CO<sub>2</sub> concentration;  
569 the model's radiation scheme sees a fixed preindustrial CO<sub>2</sub>  
570 concentration. These results can be qualitatively compared to CDR-MIP  
571 experiment *C2\_overshoot*, which is a fully coupled version of this scenario.

572

### 573 3.3 Simulation ensembles

574

575 We encourage participants whose models have internal variability to  
576 conduct multiple realizations, i.e. ensembles, for all experiments. While these are  
577 highly desirable, they are neither mandatory, nor a prerequisite for participation  
578 in CDR-MIP. Therefore, the number of ensemble members is at the discretion of  
579 each modeling group. However, we strongly encourage groups to submit at least  
580 three ensemble members if possible.

581

### 582 3.4 Climate sensitivity calculation

583

584 Knowing the climate sensitivity of each model participating in CDR-MIP is  
585 important for interpreting the results. For modelling groups that have not  
586 already calculated their model's climate sensitivity, the required CMIP *1pctCO<sub>2</sub>*  
587 simulation can be used to calculate both the transient and equilibrium climate  
588 sensitivities. The transient climate sensitivity can be calculated as the difference  
589 in the global annual mean surface temperature between the start of the  
590 experiment and a 20-year period centered on the time of CO<sub>2</sub> doubling. The  
591 equilibrium response can be diagnosed following Gregory et al. (2004), Frölicher  
592 et al. (2013), or if possible (desirable) by running the model to an equilibrium  
593 state at 2×CO<sub>2</sub> or 4×CO<sub>2</sub>.

594

595 **3.5 Model drift**

596

597 Model drift (Gupta et al., 2013; Séférian et al., 2015) is a concern for all  
598 CDR-MIP experiments because if a model is not at an equilibrium state when the  
599 experiment or prerequisite CMIP experiment begins, then the response to any  
600 experimental perturbations could be confused by drift. Thus, before beginning  
601 any of the experiments a model must be spun-up to eliminate long-term drift in  
602 carbon reservoirs or fluxes. Groups participating in CMIP6 should follow the  
603 C4MIP protocols described in Jones et al. (2016b), to ensure that drift is  
604 acceptably small. This means that land, ocean and atmosphere carbon stores  
605 should each vary by less than 10 GtC per century (long-term average  $\leq 0.1$  Gt C  
606  $\text{yr}^{-1}$ ). We leave it to individual groups to determine the length of the run required  
607 to reach such a state. If older model versions, e.g., CMIP5, are used for any  
608 experiments, any known drift should be documented.

609

610 **4. Experimental Design and Protocols**

611

612 To facilitate multiple model needs, the experiments described below have  
613 been designed to be relatively simple to implement. In most cases, they were also  
614 designed to have high signal-to-noise ratios to better understand how the  
615 simulated Earth system responds to significant CDR perturbations. While there  
616 are many ways in which such experiments could be designed to address the  
617 questions surrounding climate reversibility and each proposed CDR method, the  
618 CDR-MIP like all MIPs, must be limited to a small number of practical  
619 experiments. Therefore, after careful consideration, one experiment was chosen  
620 specifically to address climate reversibility and several more were chosen to  
621 investigate CDR by idealized direct air capture of CO<sub>2</sub> (DAC),  
622 afforestation/reforestation, and ocean alkalization (Table 1). Experiments are  
623 prioritized based on a tiered system, although, we encourage modelling groups  
624 to complete the full suite of experiments. Unfortunately, limiting the number of  
625 experiments means that a number of potentially promising or widely utilized  
626 CDR methods or combinations of methods must wait until a later time, i.e., a 2<sup>nd</sup>

627 phase, to be investigated in a multi-model context. In particular, the exclusion of  
628 Biomass Energy with Carbon Capture and Storage (BECCS) is unfortunate, as this  
629 is the primary CDR method in the Representative Concentration Pathways (RCP)  
630 and Shared Socio-economic Pathways (SSP) scenarios used in CMIP5 and 6,  
631 respectively. However, there was no practical way to design a less idealized  
632 BECCS experiment as most state-of-the-art models are either incapable of  
633 simulating a biomass harvest with permanent removal or would require a  
634 substantial amount of reformulating to do so in a manner that allows comparable  
635 multi-model analyses.

636 In some of the experiments described below we ask that non-CO<sub>2</sub> forcing  
637 (e.g., land use change, radiative forcing from other greenhouse gases, etc.) be  
638 held constant, e.g. at that of a specific year, so that only changes in other forcing,  
639 like CO<sub>2</sub> emissions, drive the main model response. For some forcing, e.g. aerosol  
640 emissions, this may mean that monthly changes in forcing are repeated  
641 throughout the rest of the simulation as if it was always one particular year.  
642 However, we recognize that models apply forcing in different ways and leave it  
643 to individual modelling groups to determine the best way hold forcing constant.  
644 We request that the methodology for holding forcing constant be documented  
645 for each model.

646

#### 647 **4.1. Climate and carbon cycle reversibility experiment (C1)**

648

649 If CO<sub>2</sub> emissions are not reduced quickly enough, and more warming  
650 occurs than is desirable or tolerable, then it is important to understand if CDR  
651 has the potential to "reverse" climate change. Here we propose an idealized Tier  
652 1 experiment that is designed to investigate CDR-induced climate "reversibility"  
653 (Fig. 1, Table 2). This experiment investigates the "reversibility" of the climate  
654 system by leveraging the prescribed 1% yr<sup>-1</sup> CO<sub>2</sub> concentration increase  
655 experiment that was done for prior CMIPs, and is a key run for CMIP6 (Eyring et  
656 al., 2016; Meehl et al., 2014). The CDR-MIP experiment starts from the 1% yr<sup>-1</sup>  
657 CO<sub>2</sub> concentration increase experiment, *1pctCO<sub>2</sub>*, and then at the 4×CO<sub>2</sub>  
658 concentration level prescribes a -1% yr<sup>-1</sup> removal of CO<sub>2</sub> from the atmosphere to  
659 pre-industrial levels [Fig. 1; this is also similar to experiments in Boucher et al.,

660 (2012) and Zickfeld et al., (2016)]. This approach is analogous to an unspecified  
661 CDR application or DAC, where CO<sub>2</sub> is removed to permanent storage to return  
662 atmospheric CO<sub>2</sub> to a prescribed level, i.e., a preindustrial concentration. To do  
663 this, CDR would have to counter emissions (unless they have ceased) as well as  
664 changes in atmospheric CO<sub>2</sub> due to the response of the ocean and terrestrial  
665 biosphere. We realize that the technical ability of CDR methods to remove such  
666 enormous quantities of CO<sub>2</sub> on such a relatively short timescale (i.e. in a few  
667 centuries) is unrealistic. However, branching from the existing CMIP *1pctCO2*  
668 experiment provides a relatively straightforward opportunity, with a high signal-  
669 to-noise ratio, to explore the effect of large-scale removal of CO<sub>2</sub> from the  
670 atmosphere and issues involving reversibility (Fig. 2 shows exemplary *C1* results  
671 from two models).

672

#### 673 **4.1.1 Protocol for *C1***

674

675 *Prerequisite simulations*: Perform the CMIP *piControl* and the *1pctCO2*  
676 experiments. The *1pctCO2* experiment branches from the DECK *piControl*  
677 experiment, which should ideally represent a near-equilibrium state of the  
678 climate system under imposed year 1850 conditions. Starting from year 1850  
679 conditions (*piControl* global mean atmospheric CO<sub>2</sub> should be 284.7 ppm) the  
680 *1pctCO2* simulation prescribes a CO<sub>2</sub> concentration increase at a rate of 1% yr<sup>-1</sup>  
681 (i.e., exponentially). The only externally imposed difference from the *piControl*  
682 experiment is the change in CO<sub>2</sub>, i.e., all other forcing is kept at that of year 1850.  
683 A restart must be generated when atmospheric CO<sub>2</sub> concentrations are four  
684 times that of the *piControl* simulation (1138.8 ppm; this should be 140 years into  
685 the run). Groups that have already performed the *piControl* and *1pctCO2*  
686 simulations for CMIP5 or CMIP6 may provide a link to them if they are already  
687 on the Earth System Grid Federation (ESGF) that host CMIP data.

688

689 *1pctCO2-cdr* simulation: Use the 4×CO<sub>2</sub> restart from *1pctCO2* and prescribe a 1%  
690 yr<sup>-1</sup> removal of CO<sub>2</sub> from the atmosphere (start removal at the beginning of the  
691 140<sup>th</sup> year: January 1<sup>st</sup>.) until the CO<sub>2</sub> concentration reaches 284.7 ppm (140  
692 years of removal). As in *1pctCO2* the only externally imposed forcing should be

693 the change in CO<sub>2</sub> (all other forcing is kept at that of year 1850). The CO<sub>2</sub>  
694 concentration should then be held at 284.7 ppm for as long as possible (a  
695 minimum of 60 years is required), with no change in other forcing. EMICs and  
696 box models are encouraged to extend runs for at least 1000 years (and up to  
697 5000 years) at 284.7 ppm CO<sub>2</sub> to investigate long-term climate system and  
698 carbon cycle reversibility (see Fig. 2 b and d for examples of why it is important  
699 to understand the long-term response).

700

## 701 **4.2 Direct CO<sub>2</sub> air capture with permanent storage experiments (C2)**

702

703 The idea of directly removing excess CO<sub>2</sub> from the atmosphere (i.e.,  
704 concentrations above pre-industrial levels) and permanently storing it in some  
705 reservoir, such as a geological formation, is appealing because such an action  
706 would theoretically address the main cause of climate change, anthropogenically  
707 emitted CO<sub>2</sub> that remains in the atmosphere. Laboratory studies and small-scale  
708 pilot plants have demonstrated that atmospheric CO<sub>2</sub> can be captured by several  
709 different methods that are often collectively referred to as Direct Air Capture  
710 (DAC) technology (Holmes and Keith, 2012; Lackner et al., 2012; Sanz-Pérez et  
711 al., 2016). Technology has also been developed that can place captured carbon in  
712 permanent reservoirs, i.e., Carbon Capture and Storage (CCS) methods (Matter et  
713 al., 2016; Scott et al., 2013, 2015) . DAC technology is currently prohibitively  
714 expensive to deploy at large scales and may be technically difficult to scale up  
715 (National Research Council, 2015), but does appear to be a potentially viable  
716 CDR option. However, aside from the technical questions involved in developing  
717 and deploying such technology, there remain questions about how the Earth  
718 system would respond if CO<sub>2</sub> were removed from the atmosphere.

719 Here we propose a set of experiments that are designed to investigate and  
720 quantify the response of the Earth system to idealized large-scale DAC. In all  
721 experiments, atmospheric CO<sub>2</sub> is allowed to freely evolve to investigate carbon  
722 cycle and climate feedbacks in response to DAC. The first two idealized  
723 experiments described below use an instantaneous (*pulse*) CO<sub>2</sub> removal from the  
724 atmosphere - approach for this investigation. Instantaneous CO<sub>2</sub> removal  
725 perturbations were chosen since *pulsed* CO<sub>2</sub> addition experiments have already

726 been proven useful for diagnosing carbon cycle and climate feedbacks in  
727 response to CO<sub>2</sub> perturbations. For example, previous positive CO<sub>2</sub> pulse  
728 experiments have been used to calculate Global Warming Potential (GWP) and  
729 Global Temperature change Potential (GTP) metrics (Joos et al., 2013). The  
730 experiments described below build upon the previous positive CO<sub>2</sub> pulse  
731 experiments, i.e., the PD100 and PI100 impulse experiments described in Joos et.  
732 al. (2013) where 100 Gt C is instantly added to preindustrial and near present  
733 day simulated climates. However, our experiments also prescribe a negative CDR  
734 pulse as opposed to just adding CO<sub>2</sub> to the atmosphere. Two experiments are  
735 desirable because the Earth system response to CO<sub>2</sub> removal will be different  
736 when starting from an equilibrium state versus starting from a perturbed state  
737 (Zickfeld et al., 2016). One particular goal of these experiments is to estimate a  
738 Global Cooling Potential (GCP) metric based on a CDR Impulse Response  
739 Function (IRF<sub>CDR</sub>). Such a metric will be useful for calculating how much CO<sub>2</sub> is  
740 removed by DAC and how much DAC is needed to achieve a particular climate  
741 target.

742 The third experiment, which focuses on "negative emissions", is based on  
743 the Shared Socio-economic Pathway (SSP) 5-3.4-overshoot scenario and its long-  
744 term extension (Kriegler et al., 2016; O'Neill et al., 2016). This scenario is of  
745 interest to CDR-MIP because after an initially high level of emissions, which  
746 follows the SSP5-8.5 unmitigated baseline scenario until 2040, CO<sub>2</sub> emissions are  
747 rapidly reduced with net CO<sub>2</sub> emissions becoming negative after the year 2070  
748 and continuing to be so until the year 2190 when they reach zero. In the original  
749 SSP5-3.4-OS scenario, the negative emissions are achieved using BECCS.  
750 However, as stated earlier there is currently no practical way to design a good  
751 multi-model BECCS experiment. Therefore, in our experiments negative  
752 emissions are achieved by simply removing CO<sub>2</sub> from the atmosphere and  
753 assuming that it is permanently stored in a geological reservoir. While this may  
754 violate the economic assumptions underlying the scenario, it still provides an  
755 opportunity to explore the response of the climate and carbon cycle to  
756 potentially achievable levels of negative emissions.

757 According to calculations done with a simple climate model, MAGICC  
758 version 6.8.01 BETA (Meinshausen et al., 2011; O'Neill et al., 2016), the SSP5-3.4-

759 OS scenario considerably overshoots the  $3.4 \text{ W m}^{-2}$  forcing level, with a peak  
760 global mean temperature of about  $2.4^\circ \text{ C}$ , before returning to  $3.4 \text{ W m}^{-2}$  at the  
761 end of the century. Eventually in the long-term extension of this scenario, the  
762 forcing stabilizes just above  $2 \text{ W m}^{-2}$ , with a global mean temperature that should  
763 equilibrate at about  $1.25^\circ \text{ C}$  above pre-industrial temperatures. Thus, in addition  
764 to allowing an investigation into the response of the climate and carbon cycle to  
765 negative emissions, this scenario also provides the opportunity to investigate  
766 issues of reversibility, albeit on a shorter timescale and with less of an  
767 "overshoot" than in experiment *C1*.

768

#### 769 **4.2.1 Instantaneous CO<sub>2</sub> removal / addition from an unperturbed climate 770 experimental protocol (*C2\_pi-pulse*)**

771

772 This idealized Tier 1 experiment is designed to investigate how the Earth  
773 system responds to DAC when perturbed from an equilibrium state (Fig. 3, Table  
774 3). The idea is to provide a baseline system response that can later be compared  
775 to the response of a perturbed system, i.e., experiment *C2\_yr2010-pulse* (Section  
776 4.2.3). By also performing another simulation where the same amount of CO<sub>2</sub> is  
777 added to the system, it will be possible to diagnose if the system responds in an  
778 inverse manner when the CO<sub>2</sub> pulse is positive. Many modelling groups will have  
779 already conducted the prerequisite simulation for this experiment in preparation  
780 for other modelling research, e.g., during model spin-up or for CMIP, which  
781 should minimize the effort needed to perform the complete experiment. The  
782 protocol is as follows:

783

784 *Prerequisite simulation* - Control simulation under preindustrial conditions with  
785 freely evolving CO<sub>2</sub>. All boundary conditions (solar forcing, land use, etc.) are  
786 expected to remain constant. This is also the CMIP5 *esmControl* simulation  
787 (Taylor et al., 2012) and the CMIP6 *esm-piControl* simulation (Eyring et al.,  
788 2016). Note that this is exactly the same as PI100 run 4 in Joos et. al. (2013).

789

790 *esm-pi-cdr-pulse* simulation - As in *esm-Control* or *esm-piControl*, but with 100 Gt  
791 C instantaneously (within 1 time step) removed from the atmosphere in year 10.

792 If models have CO<sub>2</sub> spatially distributed throughout the atmosphere, we suggest  
793 removing this amount in a uniform manner. After the negative pulse ESMs  
794 should continue the run for at least 100 years, while EMICs and box models are  
795 encouraged to continue the run for at least 1000 years (and up to 5000 years if  
796 possible). Figure 4 shows example *esm-pi-cdr-pulse* model responses.

797

798 *esm-pi-co2pulse* simulation - The same as *esm-pi-cdr-pulse*, but add a positive 100  
799 Gt C pulse (within 1 time step) as in Joos et. al. (2013), instead of a negative one.  
800 If models have CO<sub>2</sub> spatially distributed throughout the atmosphere, we suggest  
801 adding CO<sub>2</sub> in a uniform manner. Note that this would be exactly the same as the  
802 PI100 run 5 in Joos et. al. (2013) and can thus, be compared to this earlier study.

803

804 **4.2.3 Instantaneous CO<sub>2</sub> removal from a perturbed climate experimental  
805 protocol (C2\_yr2010-pulse)**

806

807 This Tier 3 experiment is designed to investigate how the Earth system  
808 responds when CO<sub>2</sub> is removed from an anthropogenically-altered climate not in  
809 equilibrium (Fig. 5, Table 4). Many modelling groups will have already conducted  
810 part of the first run of this experiment in preparation for other modelling  
811 research, e.g., CMIP, and may be able to use a "restart" file to initialize the first  
812 run, which should reduce the effort needed to perform the complete experiment.

813

814 *Prerequisite simulation* - Prescribed CO<sub>2</sub> run. Historical atmospheric CO<sub>2</sub> is  
815 prescribed until a concentration of 389ppm is reached (~year 2010; Fig. 5 top  
816 panel). Other historical forcing, i.e., from CMIP, should also be applied. An  
817 existing run or setup from CMIP5 or CMIP6 may also be used to reach a CO<sub>2</sub>  
818 concentration of 389ppm, e.g., the RCP 8.5 CMIP5 simulation or the CMIP6  
819 *historical* experiment. During this run, compatible emissions should be  
820 frequently diagnosed (at least annually).

821

822 *yr2010co2* simulation - Atmospheric CO<sub>2</sub> should be held constant at 389 ppm  
823 with other forcing, like land use and aerosol emissions, also held constant (Fig. 5  
824 top panel). ESMs should continue the run at 389ppm for at least 105 years, while

825 EMICs and box models are encouraged to continue the run for as long as needed  
826 for the subsequent simulations (e.g., 1000+ years). During this run, compatible  
827 emissions should be frequently diagnosed (at least annually). Note that when  
828 combined with the prerequisite simulation described above this is exactly the  
829 same as the PD100 run 1 in Joos et. al. (2013).

830

831 *esm-hist-yr2010co2-control* simulation - Diagnosed emissions control run. The  
832 model is initialized from the pre-industrial period (i.e., using a restart from  
833 either *piControl* or *esm-piControl*) with the emissions diagnosed in the *historical*  
834 and *yr2010co2* simulations, i.e., year 1850 to approximately year 2115 for ESMs  
835 and longer for EMICs and box models (up to 5000 years). All other forcing should  
836 be as in the *historical* and *yr2010co2* simulations. Atmospheric CO<sub>2</sub> must be  
837 allowed to freely evolve. The results should be quite close to those in the  
838 *historical* and *yr2010co2* simulations. If there are significant differences, e.g., due  
839 to climate-carbon cycle feedbacks that become evident when atmospheric CO<sub>2</sub> is  
840 allowed to freely evolve, then they must be diagnosed and used to adjust the CO<sub>2</sub>  
841 emission forcing. In some cases it may be necessary to perform an ensemble of  
842 simulations to diagnose compatible emissions. Note that this is exactly the same  
843 as the PD100 run 2 in Joos et. al. (2013). As in Joos et al. (2013), if computational  
844 time is an issue and if a group is sure that CO<sub>2</sub> remains at a nearly constant value  
845 with the emissions diagnosed in *yr2010co2*, the *esm-hist-yr2010co2-control*  
846 simulation may be skipped. This may only apply to ESMs and it is strongly  
847 recommended to perform the *esm-hist-yr2010co2-control* simulation to avoid  
848 model drift.

849

850 *esm-yr2010co2-cdr-pulse* simulation - CO<sub>2</sub> removal simulation. Setup is initially  
851 as in the *esm-hist-yr2010co2-control* simulation. However, a "negative" emissions  
852 pulse of 100 GtC is subtracted instantaneously (within 1 time step) from the  
853 atmosphere 5 years after the time at which CO<sub>2</sub> was held constant in the *esm-*  
854 *hist-yr2010co2-control* simulation (this should be at the beginning of the year  
855 2015), with the run continuing thereafter for at least 100 years (up to 5000  
856 years, if possible). If models have CO<sub>2</sub> spatially distributed throughout the  
857 atmosphere, we suggest removing this amount in a uniform manner. It is crucial

858 that the negative pulse be subtracted from a constant background concentration  
859 of  $\sim 389$  ppm. All forcing, including CO<sub>2</sub> emissions, must be exactly as in the *esm-*  
860 *hist-yr2010co2-control* simulation so that the only difference between these runs  
861 is that this one has had CO<sub>2</sub> instantaneously removed from the atmosphere.

862  
863 *esm-yr2010co2-noemit* - A zero CO<sub>2</sub> emissions control run. Setup is initially as in  
864 the *esm-yr2010co2-cdr-pulse* simulation. However, at the time of the "negative"  
865 emissions pulse in the *esm-yr2010co2-cdr-pulse* simulation, emissions are set to  
866 zero with the run continuing thereafter for at least 100 years. If possible extend  
867 the runs for at least 1000 years (and up to 5000 years). All other forcing must be  
868 exactly as in the *esm-yr2010co2-control* simulation. This experiment will be used  
869 to isolate the Earth system response to the negative emissions pulse in the *esm-*  
870 *yr2010co2-cdr-pulse* simulation, which convolves the response to the negative  
871 emissions pulse with the lagged response to the preceding positive CO<sub>2</sub>  
872 emissions (diagnosed with the zero emissions simulation). The response to the  
873 negative emissions pulse will be calculated as the difference between *esm-*  
874 *yr2010co2-cdr-pulse* and *esm-yr2010co2-noemit* simulations.

875  
876 *esm-yr2010co2-co2pulse* simulation - CO<sub>2</sub> addition simulation. Setup is initially as  
877 in the *esm-yr2010co2-cdr-pulse* simulation. However, a "positive" emissions  
878 pulse of 100 GtC is added instantaneously (within 1 time step), with the run  
879 continuing thereafter for a minimum of 100 years. If models have CO<sub>2</sub> spatially  
880 distributed throughout the atmosphere, we suggest adding CO<sub>2</sub> in a uniform  
881 manner. If possible extend the runs for at least 1000 years (and up to 5000  
882 years). It is crucial that the positive pulse be added to a constant background  
883 concentration of  $\sim 389$  ppm. All forcing, including CO<sub>2</sub> emissions, must be exactly  
884 as in the *esm-hist-yr2010co2-control* simulation so that the only difference  
885 between these runs is that this one has had CO<sub>2</sub> instantaneously added to the  
886 atmosphere. Note that this would be exactly the same as PD100 run in Joos et. al.  
887 (2013). This will be used to investigate if, after positive and negative pulses,  
888 carbon cycle and climate feedback responses, which are expected to be opposite  
889 in sign, differ in magnitude and temporal scale. The results can also be compared  
890 to Joos et. al. (2013).

891

892   **4.2.5 Emission driven SSP5-3.4-OS experimental protocol (C2\_overshoot)**

893

894       This Tier 2 experiment explores CDR in an "overshoot" climate change  
895 scenario, the SSP5-3.4-OS scenario (Fig. 6, Table 5). To start, groups must  
896 perform the CMIP6 emission driven historical simulation, *esm-hist*. Then using  
897 this as a starting point, conduct an emissions-driven SSP5-3.4-OS scenario  
898 simulation, *esm-ssp534-over*, (starting on January 1, 2015) that includes the long-  
899 term extension to the year 2300, *esm-ssp534-over-ext*. All non-CO<sub>2</sub> forcing should  
900 be identical to that in the ScenarioMIP *ssp534-over* and *ssp534-over-ext*  
901 simulations. If computational resources are sufficient, we recommend that the  
902 *esm-ssp534-over-ext* simulation be continued for at least another 1000 years with  
903 year 2300 forcing, i.e., the forcing is held constant at year 2300 levels as the  
904 simulation continues for as long as possible; up to 5000 years, to better  
905 understand processes that are slow to equilibrate, e.g., ocean carbon and heat  
906 exchange or permafrost dynamics.

907

908   **4.3 Afforestation/reforestation experiment (C3)**

909

910       Enhancing the terrestrial carbon sink by restoring or extending forest  
911 cover, i.e., reforestation and afforestation, has often been suggested as a potential  
912 CDR option (National Research Council, 2015; The Royal Society, 2009).  
913 Enhancing this sink is appealing because terrestrial ecosystems have  
914 cumulatively absorbed over a quarter of all fossil fuel emissions (Le Quéré et al.,  
915 2016) and could potentially sequester much more. Most of the key questions  
916 concerning land use change are being addressed by LUMIP (Lawrence et al.,  
917 2016). These include investigations into the potential and side effects of  
918 afforestation/reforestation to mitigate climate change, for which they have  
919 designed four experiments (LUMIP Phase 2 experiments). However, three of  
920 these experiments are CO<sub>2</sub> concentration driven, and thus are unable to fully  
921 investigate the climate-carbon cycle feedbacks that are important for CDR-MIP.  
922 The LUMIP experiment where CO<sub>2</sub> emissions force the simulation, *esm-ssp585-*  
923 *ssp126Lu*, will allow for climate-carbon cycle feedbacks to be investigated.

924 Unfortunately, since this experiment ends in the year 2100 it is too short to  
925 answer some of the key CDR-MIP questions (Section 1.2). We have therefore  
926 decided to extend this LUMIP experiment within the CDR-MIP framework as a  
927 Tier 2 experiment (Table 6) to better investigate the longer-term CDR potential  
928 and risks of afforestation/reforestation.

929 The LUMIP experiment, *esm-ssp585-ssp126Lu*, simulates  
930 afforestation/reforestation by combining a high SSP CO<sub>2</sub> emission scenario,  
931 SSP5-8.5, with a future land use change scenario from an alternative SSP  
932 scenario, SSP1-2.6, which has much greater afforestation/reforestation (Kriegler  
933 et al., 2016; Lawrence et al., 2016). By comparing this combination to the SSP5-  
934 8.5 baseline scenario, it will be possible to determine the CDR potential of this  
935 particular afforestation/reforestation scenario in a high CO<sub>2</sub> world. This is  
936 similar to the approach of Sonntag et al. (2016) using RCP 8.5 emissions  
937 combined with prescribed RCP 4.5 land use.

938

#### 939 **4.3.1 C3 Afforestation/reforestation experimental protocol**

940

941 *Prerequisite simulations* - Conduct the C4MIP emission-driven *esm-ssp585*  
942 simulation, which is a control run, and the LUMIP Phase 2 experiment *esm-*  
943 *ssp585-ssp126Lu* (Lawrence et al., 2016). Generate restart files in the year 2100.

944

945 *esm-ssp585-ssp126Lu-ext* simulation - Using the year 2100 restart from the *esm-*  
946 *ssp585-ssp126Lu* experiment, continue the run with the same LUMIP protocol  
947 (i.e., an emission driven SSP5-8.5 simulation with SSP1-2.6 land use instead of  
948 SSP5-8.5 land use) until the year 2300 using the SSP5-8.5 and SSP1-2.6 long-  
949 term extension data (O'Neill et al., 2016). If computational resources are  
950 sufficient, we recommend that the simulation be continued for at least another  
951 1000 years with year 2300 forcing (i.e., forcing is held at year 2300 levels as the  
952 simulation continues for as long as possible; up to 5000 years). This is to better  
953 understand processes that are slow to equilibrate, e.g., ocean carbon and heat  
954 exchange or permafrost dynamics, and the issue of permanence.

955

956 *esm-ssp585ext* simulation - The emission-driven esmSSP5-8.5 simulation must be  
957 extended beyond the year 2100 to serve as a control run for the *esm-ssp585-*  
958 *ssp126Lu-ext* simulation. This will require using the ScenarioMIP *ssp585-ext*  
959 forcing, but driving the model with CO<sub>2</sub> emissions instead of prescribing the CO<sub>2</sub>  
960 concentration. If computational resources are sufficient, the simulation should be  
961 extended even further than in the official SSP scenario, which ends in year 2300,  
962 by keeping forcing constant after this time (i.e., forcing is held at year 2300 levels  
963 as the simulation continues for as long as possible; up to 5000 years).

964

#### 965 **4.4. Ocean alkalinization experiment (C4)**

966

967 Enhancing the natural process of weathering, which is one of the key  
968 negative climate-carbon cycle feedbacks that removes CO<sub>2</sub> from the atmosphere  
969 on long time scales (Colbourn et al., 2015; Walker et al., 1981), has been  
970 proposed as a potential CDR method (National Research Council, 2015; The  
971 Royal Society, 2009). Enhanced weathering ideas have been proposed for both  
972 the terrestrial environment (Hartmann et al., 2013) and the ocean (Köhler et al.,  
973 2010; Schuiling and Krijgsman, 2006). We focus on the alkalinization of the  
974 ocean given its capacity to take up vast quantities of carbon over relatively short  
975 time periods and its potential to reduce the rate and impacts of ocean  
976 acidification (Kroeker et al., 2013). The idea is to dissolve silicate or carbonate  
977 minerals in seawater to increase total alkalinity. Total alkalinity, which can  
978 chemically be defined as the excess of proton acceptors over proton donors with  
979 respect to a certain zero level of protons, is a measurable quantity that is related  
980 to the concentrations of species of the marine carbonate system (Wolf-Gladrow  
981 et al., 2007). It plays a key role determining the air-sea gas exchange of CO<sub>2</sub>  
982 (Egleston et al., 2010). When total alkalinity is artificially increased in surface  
983 waters, it basically allows more CO<sub>2</sub> to dissolve in the seawater and be stored as  
984 ions such as bicarbonate or carbonate, i.e., the general methodology increases  
985 the carbon storage capacity of seawater.

986 Theoretical work and idealized modelling studies have suggested that  
987 ocean alkalinization may be an effective CDR method that is more limited by  
988 logistic constraints (e.g., mining, transport, and mineral processing) rather than

natural ones, such as available ocean area, although chemical constraints and side effects do exist (González and Ilyina, 2016; Ilyina et al., 2013; Keller et al., 2014; Köhler et al., 2010, 2013). One general side effect of ocean alkalinization, is that it increases the buffering capacity and pH of the seawater. While such a side effect could be beneficial or even an intended effect to counter ocean acidification (Feng et al., 2016), high levels of alkalinity may also be detrimental to some organisms (Cripps et al., 2013). Ocean alkalinization likely also has method specific side effects. Many of these side effects are related to the composition of the alkalinizing agent, e.g., olivine may contain nutrients or toxic heavy metals, which could affect marine organisms and ecosystems (Hauck et al., 2016; Köhler et al., 2013). Other side effects could be caused by the mining, processing, and transport of the alkalinizing agent, which in some cases may offset the CO<sub>2</sub> sequestration potential of specific ocean alkalinization methods (e.g., through CO<sub>2</sub> release by fossil fuel use or during the calcination of CaCO<sub>3</sub>) (Kheshgi, 1995; Renforth et al., 2013).

Although previous modelling studies have suggested that ocean alkalinization may be a viable CDR method, these studies are not comparable due to different experimental designs. Here we propose an idealized Tier 2 experiment (Table 7) that is designed to investigate the response of the climate system and carbon cycle to ocean alkalinization. The amount of any particular alkalinizing agent that could be mined, processed, transported, and delivered to the ocean in a form that would easily dissolve and enhance alkalinity is poorly constrained (Köhler et al., 2013; Renforth et al., 2013). Therefore, the amount of alkalinity that is to be added in our experiment is set (based on exploratory simulations conducted with the CSIRO-Mk3L-COAL model) to have a cumulative effect on atmospheric CO<sub>2</sub> by the year 2100 that is comparable to the amount removed in the CDR-MIP instantaneous DAC simulations, i.e., an atmospheric reduction of ~100 Gt C; experiments *C2\_pi-pulse* and *C2\_yr2010-pulse*. The idea here is not to test the maximum potential of such a method, which would be difficult given the still relatively coarse resolution of many models and the way in which ocean carbonate chemistry is simulated, but rather to compare the response of models to a significant alkalinity perturbation. We have also

1021 included an additional "termination" simulation that can be used to investigate  
1022 an abrupt stop in ocean alkalinization deployment.

1023

1024 **4.4.1 C4 Ocean alkalinization experimental protocol**

1025

1026 Prerequisite simulation - Conduct the C4MIP emission-driven *esm-ssp585*  
1027 simulation as described by Jones et al., (2016b). This is the SSP5-8.5 high CO<sub>2</sub>  
1028 emission scenario, and it serves as the control run and branching point for the  
1029 ocean alkalinization experiment. A restart must be generated at the end of the  
1030 year 2019.

1031

1032 *esm-ssp585-ocean-alk* simulation - Begin an 80 year run using the *esm-ssp585*  
1033 year 2020 restart (starting on Jan. 1, 2020) and add 0.14 Pmol Total Alkalinity  
1034 (TA) yr<sup>-1</sup> to the upper grid boxes of each model's ocean component, i.e., branch  
1035 from the C4MIP *esm-ssp585* simulation in 2020 until 2100. The alkalinity  
1036 additions should be limited to mostly ice free, year-round ship accessible waters,  
1037 which for simplicity should set to be between 70°N and 60°S (note that this  
1038 ignores the presence of seasonal sea-ice in some small regions). For many  
1039 models, this will in practice result in an artificial TA flux at the air-sea interface  
1040 with realized units that might, for example, be something like  $\mu\text{mol TA s}^{-1} \text{cm}^{-2}$ .  
1041 Adding 0.14 Pmol TA yr<sup>-1</sup> is equivalent to adding 5.19 Pg yr<sup>-1</sup> of an alkalinizing  
1042 agent like Ca(OH)<sub>2</sub> or 4.92 Pg yr<sup>-1</sup> of forsterite (Mg<sub>2</sub>SiO<sub>4</sub>), a form of olivine  
1043 [assuming theoretical net instant dissolution reactions which for every mole of  
1044 Ca(OH)<sub>2</sub> or Mg<sub>2</sub>SiO<sub>4</sub> added sequesters 2 or 4 moles, respectively, of CO<sub>2</sub> (Ilyina et  
1045 al., 2013; Köhler et al., 2013)]. As not all models include marine iron or silicate  
1046 cycles, the addition of these nutrients, which could occur if some form of olivine  
1047 were used as the alkalinizing agent, is not considered here. All other forcing is as in  
1048 the *esm-ssp585* control simulation. If the ocean alkalinization termination  
1049 simulation (below) is to be conducted, generate a restart at the beginning of the  
1050 year 2070.

1051

1052 Optional (Tier 3) *esm-ssp585-ocean-alk-stop* simulation - Use the year 2070  
1053 restart from the *esm-ssp585-ocean-alk* simulation and start a simulation

1054 (beginning on Jan. 1, 2070) with the SPP5-8.5 forcing, but without adding any  
1055 additional alkalinity. Continue this run until the year 2100, or beyond, if  
1056 conducting the *esm-ssp585-ocean-alk-ext* simulation (below).

1057

1058 Optional (Tier 3) ocean alkalinization extension simulations:

1059

1060 *esm-ssp585ext* simulation - If groups desire to extend the ocean alkalinization  
1061 experiment beyond the year 2100, an optional simulation may be conducted to  
1062 extend the control run using forcing data from the ScenarioMIP *ssp585ext*  
1063 simulation, i.e., conduct a longer emission-driven control run, *esm-ssp585ext*.  
1064 This extension is also a control run for those conducting the CDR-MIP C3  
1065 afforestation/reforestation simulation (Section 4.3). If computational resources  
1066 are sufficient, the simulation should be extended even further than in the official  
1067 SSP scenario, which ends in year 2300, by keeping the forcing constant after this  
1068 time (i.e., forcing is held at year 2300 levels as the simulation continues for as  
1069 long as possible; up to 5000 years).

1070

1071 *esm-ssp585-ocean-alk-ext* simulation - Continue the ocean alkalinization  
1072 experiment described above (i.e., adding 0.14 Pmol Total Alkalinity (TA)  $\text{yr}^{-1}$  to  
1073 the upper grid boxes of each model's ocean component) beyond the year 2100  
1074 (up to 5000 years) using forcing from the *esm-ssp585-ext* simulation.

1075

1076 **5. Model output, data availability, and data use policy**

1077 **5.1 Gridded model output**

1078

1079 Models capable of generating gridded data must use a NetCDF format. The  
1080 output (see Appendix A web link for the list of requested variables) follows the  
1081 CMIP6 output requirements in frequency and structure. This allows groups to  
1082 use CMOR software (Climate Model Rewriter Software, available at  
1083 <http://cmor.llnl.gov/>) to generate the files that will be available for public  
1084 download (Section 5.5). CMOR3 tables for CDR-MIP are available at [www.kiel-earth-institute.de/files/media/downloads/CDRmon.json](http://www.kiel-earth-institute.de/files/media/downloads/CDRmon.json) (table for monthly  
1085 output) and [www.kiel-earth-institute.de/files/media/downloads/CDRga.json](http://www.kiel-earth-institute.de/files/media/downloads/CDRga.json)  
1086 (table for annual output).

1087 (table for global annual mean output). The resolution of the data should be as  
1088 close to native resolution as possible, but on a regular grid. Please note as  
1089 different models have different formulations, only applicable outputs need be  
1090 provided. However, groups are encouraged to generate additional output, i.e.,  
1091 whatever their standard output variables are, and can also make this data  
1092 available (preferably following the CMIP6 CMOR standardized naming  
1093 structure).

1094

## 1095 **5.2 Conversion factor Gt C to ppm**

1096

1097 For experiments where carbon must be converted between GtC (or Pg)  
1098 and ppm CO<sub>2</sub>, please use a conversion factor of 2.12 GtC per ppm CO<sub>2</sub> to be  
1099 consistent with Global Carbon Budget (Le Quere et al., 2015) conversion factors.

1100

## 1101 **5.3 Box model output**

1102

1103 For models that are incapable of producing gridded NetCDF data (i.e., box  
1104 models), output is expected to be in an ASCII format (Appendix B). All ASCII files  
1105 are expected to contain tabulated values (at a minimum global mean values),  
1106 with at least two significant digits for each run. Models must be able to calculate  
1107 key carbon cycle variables (Appendix C) to participate in CDR-MIP experiments  
1108 C1 and C2. Please submit these files directly to the corresponding author who  
1109 will make them available for registered users to download from the CDR-MIP  
1110 website.

1111

## 1112 **5.4 Model output frequency**

1113

1114 The model output frequency is listed in Table 8. In all experiments box  
1115 models and EMICs without seasonality are expected to generate annual mean  
1116 output for the duration of the experiment, while models with seasonality are  
1117 expected to generate higher spatial resolution data, i.e., monthly, for most  
1118 simulations.

1119 In experiment C1 for the control run, *piControl*, we request that 100 years  
1120 of 3-D model output be written monthly (this should be the last 100 years if  
1121 conducting a 500+ year run for CMIP6). For the *1pctCO2* and *1pctCO2-cdr*  
1122 simulations 3-D model output should also be written monthly, i.e., as the  
1123 atmospheric CO<sub>2</sub> concentration is changing. We suggest that groups that have  
1124 already performed the *piControl* and *1pctCO2* simulations for CMIP5 or CMIP6  
1125 with an even higher output resolution (e.g., daily) continue to use this resolution  
1126 for the *1pctCO2-cdr* simulation, as this will facilitate the analysis. For groups  
1127 continuing the simulations for up to 5000 years after CO<sub>2</sub> has returned to 284.7  
1128 ppm, at a minimum, annual global mean values (non-gridded output) should be  
1129 generated after the initial minimum 60 years of higher resolution output.

1130 For experiment *C2\_pi\_pulse* if possible, 3-D model output should be  
1131 written monthly for 10 years before the negative pulse and for 100 years  
1132 following the pulse. For groups that can perform longer simulations, e.g.,  
1133 thousands of years, at a minimum, annual global mean values (non-gridded  
1134 output) should be generated. Data for the control run, i.e., the equilibrium  
1135 simulation *esm-piControl*, must also be available for analytical purposes. CMIP  
1136 participants may provide a link to the *esm-Control* or *esm-piControl* data on the  
1137 ESGF.

1138 For experiment *C2\_yr2010-pulse* the *historical* and *yr2010co2* simulations  
1139 output is only needed to diagnose annual CO<sub>2</sub> emissions and will not be archived  
1140 on the ESGF. Gridded 3-D monthly mean output for the *esm-hist-yr2010co2-*  
1141 *control* (starting in the year 2010), *esm-yr2010co2-cdr-pulse*, *esm-yr2010co2-*  
1142 *noemit*, and *esm-yr2010co2-co2pulse* simulations should be written for the initial  
1143 100 years of the simulation. Thereafter, for groups that can perform longer  
1144 simulations (up to 5000 years), at a minimum annual global mean values (non-  
1145 gridded output) should be generated. CMIP participants are requested to provide  
1146 a link to the *historical* simulation data on the ESGF.

1147 For experiment *C2\_overshoot*, if possible, 3-D model output should be  
1148 written monthly until the year 2300. We suggest that groups that have already  
1149 performed the ScenarioMIP *ssp534-over* and *ssp534-over-ext* and C4MIP *ssp534-*  
1150 *over-bgc* and *ssp534-over-bgcExt* CMIP6 simulations with an even higher output  
1151 resolution (e.g., daily) continue to use this resolution as this will facilitate

1152 analyses. For groups that can perform longer simulations, e.g., thousands of  
1153 years, at a minimum annual global mean values (non-gridded output) should be  
1154 generated for every year beyond 2300. We recommend that CMIP participants  
1155 provide a link to the *esm-hist* data on the ESGF. For analytical purposes, we also  
1156 request that ScenarioMIP and C4MIP participants provide links to any completed  
1157 *ssp534-over*, *ssp534-over-ext*, *ssp534-over-bgc* and *ssp534-over-bgcExt* simulation  
1158 data on the ESGF.

1159 For experiment C3 if possible, 3-D model output should be written  
1160 monthly until the year 2300. LUMIP participants may provide a link to the *esm-*  
1161 *hist* and *esm-ssp585-ssp126Lu* data on the ESGF for the first portions of this run  
1162 (until the year 2100). For groups that can perform longer simulations, e.g.,  
1163 thousands of years, at a minimum annual global mean values (non-gridded  
1164 output) should be generated for every year beyond 2300.

1165 For experiment C4 if possible, 3-D gridded model output should be  
1166 written monthly for all simulations. For groups that can perform longer  
1167 simulations, e.g., thousands of years, at a minimum annual global mean values  
1168 (non-gridded output) should be generated for every year beyond 2300  
1169

## 1170 **5.5 Data availability and use policy**

1171

1172 The model output from the CDR-MIP experiments described in this paper  
1173 will be publically available. All gridded model output will, to the extent possible,  
1174 be distributed through the Earth System Grid Federation (ESGF). Box model  
1175 output will be available via the CDR-MIP website (<http://www.kiel-earth->  
1176 [institute.de/cdr-mip-data.html](http://institute.de/cdr-mip-data.html)). The CDR-MIP policy for data use is that if you  
1177 use output from a particular model, you should contact the modeling group and  
1178 offer them the opportunity to contribute as authors. Modeling groups will  
1179 possess detailed understanding of their models and the intricacies of performing  
1180 the CDR-MIP experiments, so their perspectives will undoubtedly be useful. At  
1181 minimum, if the offer of author contribution is not taken up, CDR-MIP and the  
1182 model groups should be credited in acknowledgments with for example a  
1183 statement like: "*We acknowledge the Carbon Dioxide Removal Model*  
1184 *Intercomparison Project leaders and steering committee who are responsible for*

1185 *CDR-MIP and we thank the climate modelling groups (listed in Table XX of this*  
1186 *paper) for producing and making their model output available."*

1187 The natural and anthropogenic forcing data that are required for some  
1188 simulations are described in several papers in the Geoscientific Model  
1189 Development CMIP6 special issue. These data will be available on the ESGF.  
1190 Links to all forcing data can also be found on the CMIP6 Panel website  
1191 (<https://www.wcrp-climate.org/wgcm-cmip/wgcm-cmip6>). CMIP6 and CMIP5  
1192 data should be acknowledged in the standard way.

1193

## 1194 **6. CDR-MIP outlook and conclusion**

1195

1196 It is anticipated that this will be the first stage of an ongoing project  
1197 exploring CDR. CDR-MIP welcomes input on the development of other (future)  
1198 experiments and scenarios. Potential future experiments could include Biomass  
1199 Energy with Carbon Capture and Storage (BECCS) or ocean fertilization. Future  
1200 experiments could also include the removal of non-CO<sub>2</sub> greenhouse gases, e.g.,  
1201 methane, as these in many cases have a much higher global warming potential  
1202 (de\_Richter et al., 2017; Ming et al., 2016). We also envision that it will be  
1203 necessary to investigate the simultaneous deployment of several CDR or other  
1204 greenhouse gas removal methods since early studies suggest that there is likely  
1205 not an individually capable method (Keller et al., 2014). It is also anticipated that  
1206 scenarios will be developed that might combine Solar Radiation Management  
1207 (SRM) and CDR in the future, such as a joint GeoMIP (Geoengineering Model  
1208 Intercomparison Project) CDR-MIP experiment.

1209 In addition to reductions in anthropogenic CO<sub>2</sub> emissions, it is very likely  
1210 that CDR will be needed to achieve the climate change mitigation goals laid out in  
1211 the Paris Agreement. The potential and risks of large scale CDR are poorly  
1212 quantified, raising important questions about the extent to which large scale CDR  
1213 can be depended upon to meet Paris Agreement goals. This project, CDR-MIP, is  
1214 designed to help us better understand how the Earth system might respond to  
1215 CDR. Over the past two years the CDR-MIP team has developed a set of numerical  
1216 experiments to be performed with Earth system models of varying complexity.

1217 The aim of these experiments is to provide coordinated simulations and analyses  
1218 that addresses several key CDR uncertainties including:

1219

1220 • The degree to which CDR could help mitigate climate change or even  
1221 reverse it.

1222

1223 • The potential effectiveness and risks/benefits of different CDR proposals  
1224 with a focus on direct CO<sub>2</sub> air capture, afforestation/reforestation, and  
1225 ocean alkalization.

1226

1227 • To inform how CDR might be appropriately accounted for within an Earth  
1228 system framework and during scenario development.

1229

1230 We anticipate that there will be numerous forthcoming studies that utilize  
1231 CDR-MIP data. The model output from the CDR-MIP experiments will be  
1232 publically available and we welcome and encourage interested parties to  
1233 download this data and utilize it to further investigate CDR.

1234

1235

1236 **7. Code and/or data availability**

1237

1238 As described in Section 5.5, the output from models participating in CDR-  
1239 MIP will be made publically available. This will include data used in exemplary  
1240 Figs. 2 and 4. All gridded model output will be distributed through the Earth  
1241 System Grid Federation (ESGF). Box model output will be available via the CDR-  
1242 MIP website (<http://www.kiel-earth-institute.de/cdr-mip-data.html>). The code  
1243 from the models used to generate the exemplary figures in this document (Figs. 2  
1244 and 4, Appendix D) will be made available here via a web link when this  
1245 manuscript is accepted for publication. To obtain code from modelling groups  
1246 who are participating in CDR-MIP please contact the modelling group using the  
1247 contact information that accompanies their data.

1248

1249 *Acknowledgements.* D. P. Keller and N. Bauer acknowledge funding received from  
1250 the German Research Foundation's Priority Program 1689 "Climate Engineering"  
1251 (project CDR-MIA; KE 2149/2-1). K. Zickfeld acknowledges support from the  
1252 Natural Sciences and Engineering Research Council of Canada (NSERC)  
1253 Discovery grant program. The Pacific Northwest National Laboratory is  
1254 operated for the U.S. Department of Energy by Battelle Memorial Institute under  
1255 contract DE-AC05-76RL01830. D. Ji acknowledges support from the National  
1256 Basic Research Program of China under grant number 2015CB953600. CDJ was  
1257 supported by the Joint UK BEIS/ Defra Met Office Hadley Centre Climate  
1258 Programme ( GA01101) and by the European Union's Horizon 2020 research  
1259 and innovation programme under grant agreement No 641816 ( CRESCENDO).  
1260 H. Muri was supported by Norwegian Research Council grant 261862/E10.

1261

1262

1263

1264 **Appendix A. Requested model output variables**

1265

1266 A spreadsheet of the requested model output variables and their format can be  
1267 found at: [www.kiel-earth-institute.de/files/media/downloads/CDR-MIP\\_model\\_output\\_requirements.pdf](http://www.kiel-earth-institute.de/files/media/downloads/CDR-MIP_model_output_requirements.pdf). Please note as different models have  
1268 different formulations, only applicable outputs need be provided. However,  
1269 groups are encouraged to generate additional output, i.e., whatever their  
1270 standard output variables are, and can also make this data available.  
1271

1272

1273 **Appendix B. Box model output formatting**

1274

1275 Box model ASCII formatting example:

1276

1277 File name format: RUNNAME\_MODELNAME\_Modelversion.dat

1278 C1\_MYBOXMODEL\_V1.0\_.dat

1279 Headers and formats:

1280 *Example:*

1281     • Start each header comment line with a #

- *Line 1:* Indicate run name, e.g., "# *esm-pi-cdr-pulse*"
- *Line 2:* Provide contact address, e.g., "# B. Box, Uni of Box Models, CO2 Str., BoxCity 110110, BoxCountry"
- *Line 3:* Provide a contact email address, e.g., "# *bbox@unibox.bx*"
- *Line 4:* Indicate model name, version, e.g., "# *MyBoxModel Version 2.2*"
- *Line 5:* Concisely indicate main components, e.g., "# two ocean boxes (upper and lower), terrestrial biosphere, and one atmospheric box"
- *Line 6:* Indicate climate sensitivity of model, the abbreviation TCS may be used for transient climate sensitivity and ECS for equilibrium climate sensitivity, e.g., "# TCS=3.2 [deg C], ECS=8.1 [deg C]"
- *Line 7:* Description of non-CO<sub>2</sub> forcing applied, e.g., "# Forcing: solar"
- *Line 8:* Indicate the output frequency and averaging, e.g., "# Output: global mean values"
- *Line 9:* List tabulated output column headers with their units in brackets (see table below), e.g., "# *year tas[K]*"

1298 Complete Header Example:

```

1299 # esm-pi-cdr-pulse
1300 # B. Box, Uni. of Box Models, CO2 Str., BoxCity 110110, BoxCountry
1301 # bbox@unibox.bx
1302 # MyBoxModel Version 2.2
1303 # two ocean boxes (upper and lower), terrestrial biosphere, and one
1304 atmospheric box
1305 # TCS=3.2 deg C, ECS=8.1 deg C
1306 # Forcing: solar
1307 # Output: global mean values
1308 # year tas[K] co2[Gt C] nep[Gt C yr-1] fgco2[Gt C yr-1]
1309

```

1310 **Appendix C. Requested box model output variables**

1311

1312 Table of requested box model output (at a minimum as global mean values). To
1313 participate in CDR-MIP at a minimum the variables *tas*, *xco2*, and *fgco2* must be
1314 provided.

1315

Long name	Column Header Name*	Units	Comments
Relative year	year	year	
Near-surface Air Temperature	tas	K	
Atmospheric CO <sub>2</sub>	xco2	ppm	
Surface Downward CO <sub>2</sub> flux into the ocean	fgco2	kg m <sup>-2</sup>	This is the net air-to-ocean carbon flux (positive flux is into the ocean)
Total Atmospheric Mass of CO <sub>2</sub>	co2mass	kg	
Net Carbon Mass Flux out of Atmosphere due to Net Ecosystem Productivity on Land.	nep	kg m <sup>-2</sup>	This is the net air-to-land carbon flux (positive flux is into the land)
Total ocean carbon	cOcean	Gt C	If the ocean contains multiple boxes this output can also be provided, e.g., as cOcean_up and cOcean_low for upper and lower ocean boxes
Total land carbon	cLand	Gt C	This is the sum of all C pools
Ocean Potential Temperature	thetao	K	Please report a mean value if there are multiple ocean boxes
Upper ocean pH	pH	1	Negative log of hydrogen ion concentration with the concentration expressed as mol H kg <sup>-1</sup> .
Carbon Mass Flux out of Atmosphere due to Net Primary Production on Land	npp	kg m <sup>-2</sup>	This is calculated as gross primary production – autotrophic respiration (gpp-ra)
Carbon Mass Flux into Atmosphere due to Heterotrophic Respiration on Land	rh	kg m <sup>-2</sup>	
Ocean Net Primary Production by Phytoplankton	intpp	kg m <sup>-2</sup>	

1316

1317 \*Column header names follow the CMIP CMOR notation when possible

1318

1319 **Appendix D. Model descriptions**

1320  
1321        The two models used to develop and test CDR-MIP experimental  
1322        protocols and provide example results (Figs. 2 and 4) are described below.

1323        The University of Victoria Earth System Climate model (UVic), version 2.9  
1324        consists of three dynamically coupled components: a three-dimensional general  
1325        circulation model of the ocean that includes a dynamic-thermodynamic sea ice  
1326        model, a terrestrial model, and a simple one-layer atmospheric energy-moisture  
1327        balance model (Eby et al., 2013). All components have a common horizontal  
1328        resolution of 3.6° longitude x 1.8° latitude. The oceanic component, which is in  
1329        the configuration described by Keller et al. (2012), has 19 levels in the vertical  
1330        with thicknesses ranging from 50 m near the surface to 500 m in the deep ocean.  
1331        The terrestrial model of vegetation and carbon cycles (Meissner et al., 2003) is  
1332        based on the Hadley Center model TROLL (Top-down Representation of  
1333        Interactive Foliage and Flora Including Dynamics). The atmospheric energy-  
1334        moisture balance model interactively calculates heat and water fluxes to the  
1335        ocean, land, and sea ice. Wind velocities, which are used to calculate the  
1336        momentum transfer to the ocean and sea ice model, surface heat and water  
1337        fluxes, and the advection of water vapor in the atmosphere, are determined by  
1338        adding wind and wind stress anomalies. These are determined from surface  
1339        pressure anomalies that are calculated from deviations in pre-industrial surface  
1340        air temperature to prescribed NCAR/NCEP monthly climatological wind data  
1341        (Weaver et al., 2001). The model has been extensively used in climate change  
1342        studies and is also well validated under pre-industrial to present day conditions  
1343        (Eby et al., 2009, 2013; Keller et al., 2012).

1344        The CSIRO-Mk3L-COAL Earth system model consists of a climate model,  
1345        Mk3L (Phipps et al., 2011), coupled to a biogeochemical model of carbon,  
1346        nitrogen and phosphorus cycles on land (CASA-CNP) in the Australian  
1347        community land surface model, CABLE (Mao et al., 2011; Wang et al., 2010), and  
1348        an ocean biogeochemical cycle model (Duteil et al., 2012; Matear and Hirst,  
1349        2003). The atmospheric model has a horizontal resolution of 5.6° longitude by  
1350        3.2° latitude, and 18 vertical layers. The land carbon model has the same  
1351        horizontal resolution as the atmosphere. The ocean model has a resolution of

1352 2.8° longitude by 1.6° latitude, and 21 vertical levels. Mk3L simulates the  
1353 historical climate well, as compared to the models used for earlier IPCC  
1354 assessments (Phipps et al., 2011). Furthermore, the simulated response of the  
1355 land carbon cycle to increasing atmospheric CO<sub>2</sub> and warming are consistent  
1356 with those from the Coupled Model Intercomparison Project Phase 5 (CMIP5)  
1357 (Zhang et al., 2014). The ocean biogeochemical model was also shown to  
1358 realistically simulate the global ocean carbon cycle (Duteil et al., 2012; Matear  
1359 and Lenton, 2014).

1360 **References**

1361

1362 Allen, M. R., Frame, D. J., Huntingford, C., Jones, C. D., Lowe, J. A., Meinshausen, M.  
1363 and Meinshausen, N.: Warming caused by cumulative carbon emissions towards  
1364 the trillionth tonne, *Nature*, 458(7242), 1163–1166, doi:10.1038/nature08019,  
1365 2009.

1366 Armour, K. C., Eisenman, I., Blanchard-Wrigglesworth, E., McCusker, K. E. and  
1367 Bitz, C. M.: The reversibility of sea ice loss in a state-of-the-art climate model,  
1368 *Geophys. Res. Lett.*, 38(16), 1–5, doi:10.1029/2011GL048739, 2011.

1369 Arneth, A., Sitch, S., Pongratz, J., Stocker, B. D., Ciais, P., Poulter, B., Bayer, A.,  
1370 Bondeau, A., Calle, L., Chini, L., Gasser, T., Fader, M., Friedlingstein, P., Kato, E., Li,  
1371 W., Lindeskog, M., Nabel, J. E. M. S., Pugh, T. A. M., Robertson, E., Viovy, N., Yue, C.  
1372 and Zaehle, S.: Historical carbon dioxide emissions due to land use changes  
1373 possibly larger than assumed, *Nat. Geosci.*, (January), doi:10.1038/ngeo2882,  
1374 2017.

1375 Arora, V. K. and Boer, G. J.: Terrestrial ecosystems response to future changes in  
1376 climate and atmospheric CO<sub>2</sub> concentration, *Biogeosciences*, 11(15), 4157–4171,  
1377 doi:10.5194/bg-11-4157-2014, 2014.

1378 Bauer, N., Calvin, K., Emmerling, J., Fricko, O., Fujimori, S., Hilaire, J., Eom, J., Krey,  
1379 V., Kriegler, E., Mouratiadou, I., Sytze de Boer, H., van den Berg, M., Carrara, S.,  
1380 Daioglou, V., Drouet, L., Edmonds, J. E., Gernaat, D., Havlik, P., Johnson, N., Klein,  
1381 D., Kyle, P., Marangoni, G., Masui, T., Pietzcker, R. C., Strubegger, M., Wise, M.,  
1382 Riahi, K. and van Vuuren, D. P.: Shared Socio-Economic Pathways of the Energy  
1383 Sector – Quantifying the Narratives, *Glob. Environ. Chang.*, 42, 316–330,  
1384 doi:10.1016/j.gloenvcha.2016.07.006, 2017.

1385 Betts, R. A.: Offset of the potential carbon sink from boreal forestation by  
1386 decreases in surface albedo, *Nature*, 409, 187–190, 2000.

1387 Betts, R. A., Boucher, O., Collins, M., Cox, P. M., Falloon, P. D., Gedney, N.,  
1388 Hemming, D. L., Huntingford, C., Jones, C. D., Sexton, D. M. H. and Webb, M. J.:  
1389 Projected increase in continental runoff due to plant responses to increasing  
1390 carbon dioxide, *Nature*, 448(7157), 1037–1041,  
1391 doi:[http://www.nature.com/nature/journal/v448/n7157/supplinfo/nature06045\\_S1.html](http://www.nature.com/nature/journal/v448/n7157/supplinfo/nature06045_S1.html), 2007.

1393 Boucher, O., Halloran, P. R., Burke, E. J., Doutriaux-Boucher, M., Jones, C. D., Lowe,  
1394 J., Ringer, M. a, Robertson, E. and Wu, P.: Reversibility in an Earth System model  
1395 in response to CO<sub>2</sub> concentration changes, *Environ. Res. Lett.*, 7(2), 24013,  
1396 doi:10.1088/1748-9326/7/2/024013, 2012.

1397 Boysen, L. R., Lucht, W., Gerten, D. and Heck, V.: Impacts devalue the potential of  
1398 large-scale terrestrial CO<sub>2</sub> removal through biomass plantations, *Environ. Res.*  
1399 *Lett.*, 11(9), 95010, doi:10.1088/1748-9326/11/9/095010, 2016.

1400 Boysen, L. R., Lucht, W., Gerten, D., Heck, V., Lenton, T. M. and Schellnhuber, H. J.:  
1401 The limits to global-warming mitigation by terrestrial carbon removal, *Earth's*  
1402 *Futur.*, 1–12, doi:10.1002/2016EF000469, 2017.

1403 Cao, L. and Caldeira, K.: Atmospheric carbon dioxide removal: long-term  
1404 consequences and commitment, *Environ. Res. Lett.*, 5(2), 24011,  
1405 doi:10.1088/1748-9326/5/2/024011, 2010.

1406 Chen, C. and Tavoni, M.: Direct air capture of CO<sub>2</sub> and climate stabilization: A  
1407 model based assessment, *Clim. Change*, 118(1), 59–72, doi:10.1007/s10584-  
1408 013-0714-7, 2013.

1409 Claussen, M., Brovkin, V. and Ganopolski, A.: Biogeophysical versus  
1410 biogeochemical feedbacks of large-scale land cover change, *Geophys. Res. Lett.*,  
1411 28(6), 1011–1014, doi:10.1029/2000GL012471, 2001.

1412 Colbourn, G., Ridgwell, A. and Lenton, T. M.: The time scale of the silicate  
1413 weathering negative feedback on atmospheric CO<sub>2</sub>, , 1–14,  
1414 doi:10.1002/2014GB005054. Received, 2015.

1415 Cripps, G., Widdicombe, S., Spicer, J. I. and Findlay, H. S.: Biological impacts of  
1416 enhanced alkalinity in *Carcinus maenas*, *Mar. Pollut. Bull.*, 71(1–2), 190–198,  
1417 doi:10.1016/j.marpolbul.2013.03.015, 2013.

1418 de\_Richter, R., Ming, T., Davies, P., Liu, W. and Caillol, S.: Removal of non-CO<sub>2</sub>  
1419 greenhouse gases by large-scale atmospheric solar photocatalysis, *Prog. Energy*  
1420 *Combust. Sci.*, 60, 68–96, doi:10.1016/j.pecs.2017.01.001, 2017.

1421 Dlugokencky, E. (NOAA/ESRL) and Tans, P. (NOAA/ESRL): NOAA/ESRL, 2016.

1422 Duteil, O., Koeve, W., Oschlies, a., Aumont, O., Bianchi, D., Bopp, L., Galbraith, E.,  
1423 Matear, R., Moore, J. K., Sarmiento, J. L. and Segschneider, J.: Preformed and  
1424 regenerated phosphate in ocean general circulation models: can right total  
1425 concentrations be wrong?, *Biogeosciences*, 9(5), 1797–1807, doi:10.5194/bg-9-

1426 1797-2012, 2012.

1427 Eby, M., Zickfeld, K., Montenegro, a., Archer, D., Meissner, K. J. and Weaver, a. J.:  
1428 Lifetime of Anthropogenic Climate Change: Millennial Time Scales of Potential CO  
1429 2 and Surface Temperature Perturbations, *J. Clim.*, 22(10), 2501–2511,  
1430 doi:10.1175/2008JCLI2554.1, 2009.

1431 Eby, M., Weaver, a. J., Alexander, K., Zickfeld, K., Abe-Ouchi, A., Cimatoribus, a. a.,  
1432 Crespin, E., Drijfhout, S. S., Edwards, N. R., Eliseev, a. V., Feulner, G., Fichefet, T.,  
1433 Forest, C. E., Goosse, H., Holden, P. B., Joos, F., Kawamiya, M., Kicklighter, D.,  
1434 Kienert, H., Matsumoto, K., Mokhov, I. I., Monier, E., Olsen, S. M., Pedersen, J. O. P.,  
1435 Perrette, M., Philippon-Berthier, G., Ridgwell, A., Schlosser, A., Schneider von  
1436 Deimling, T., Shaffer, G., Smith, R. S., Spahni, R., Sokolov, a. P., Steinacher, M.,  
1437 Tachiiri, K., Tokos, K., Yoshimori, M., Zeng, N. and Zhao, F.: Historical and  
1438 idealized climate model experiments: an intercomparison of Earth system  
1439 models of intermediate complexity, *Clim. Past*, 9(3), 1111–1140, doi:10.5194/cp-  
1440 9-1111-2013, 2013.

1441 Egleston, E. S., Sabine, C. L. and Morel, F. M. M.: Revelle revisited: Buffer factors  
1442 that quantify the response of ocean chemistry to changes in DIC and alkalinity,  
1443 *Global Biogeochem. Cycles*, 24(1), n/a-n/a, doi:10.1029/2008GB003407, 2010.

1444 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J. and Taylor,  
1445 K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6)  
1446 experimental design and organization, *Geosci. Model Dev.*, 9(5), 1937–1958,  
1447 doi:10.5194/gmd-9-1937-2016, 2016.

1448 Feng, E. Y., Keller, D. P., Koeve, W. and Oschlies, A.: Could artificial ocean  
1449 alkalinization protect tropical coral ecosystems from ocean acidification?,  
1450 *Environ. Res. Lett.*, 11(7), 74008, doi:10.1088/1748-9326/11/7/074008, 2016.

1451 Friedlingstein, P. and Prentice, I. C.: Carbon-climate feedbacks: a review of model  
1452 and observation based estimates, *Curr. Opin. Environ. Sustain.*, 2, 251–257, 2010.

1453 Frölicher, T. L., Winton, M. and Sarmiento, J. L.: Continued global warming after  
1454 CO<sub>2</sub> emissions stoppage, *Nat. Clim. Chang.*, 4(1), 40–44,  
1455 doi:10.1038/nclimate2060, 2013.

1456 Fuss, S., Canadell, J. G., Peters, G. P., Tavoni, M., Andrew, R. M., Ciais, P., Jackson, R.  
1457 B., Jones, C. D., Kraxner, F., Nakicenovic, N., Quéré, C. Le, Raupach, M. R., Sharifi,  
1458 A., Smith, P. and Yamagata, Y.: Betting on negative emissions, *Nat. Publ. Gr.*,

1459 4(10), 850–853, doi:10.1038/nclimate2392, 2014.

1460 Gasser, T., Guivarc'h, C., Tachiiri, K., Jones, C. D. and Ciais, P.: Negative emissions  
1461 physically needed to keep global warming below 2 °C, *Nat. Commun.*, 6, 7958,  
1462 doi:10.1038/ncomms8958, 2015.

1463 González, M. F. and Ilyina, T.: Impacts of artificial ocean alkalinization on the  
1464 carbon cycle and climate in Earth system simulations, *Geophys. Res. Lett.*,  
1465 43(12), 6493–6502, doi:10.1002/2016GL068576, 2016.

1466 Gregory, J. M.: A new method for diagnosing radiative forcing and climate  
1467 sensitivity, *Geophys. Res. Lett.*, 31(October 2003), 2–5,  
1468 doi:10.1029/2003GL018747, 2004.

1469 Griffies, S. M., Danabasoglu, G., Durack, P. J., Adcroft, A. J., Balaji, V., B??ning, C. W.,  
1470 Chassignet, E. P., Curchitser, E., Deshayes, J., Drange, H., Fox-Kemper, B., Gleckler,  
1471 P. J., Gregory, J. M., Haak, H., Hallberg, R. W., Heimbach, P., Hewitt, H. T., Holland,  
1472 D. M., Ilyina, T., Jungclaus, J. H., Komuro, Y., Krasting, J. P., Large, W. G., Marsland,  
1473 S. J., Masina, S., McDougall, T. J., George Nurser, A. J., Orr, J. C., Pirani, A., Qiao, F.,  
1474 Stouffer, R. J., Taylor, K. E., Treguier, A. M., Tsujino, H., Uotila, P., Valdivieso, M.,  
1475 Wang, Q., Winton, M. and Yeager, S. G.: OMIP contribution to CMIP6:  
1476 Experimental and diagnostic protocol for the physical component of the Ocean  
1477 Model Intercomparison Project, *Geosci. Model Dev.*, 9(9), 3231–3296,  
1478 doi:10.5194/gmd-9-3231-2016, 2016.

1479 Gruber, N.: Warming up, turning sour, losing breath: ocean biogeochemistry  
1480 under global change., *Philos. Trans. A. Math. Phys. Eng. Sci.*, 369(1943), 1980–96,  
1481 doi:10.1098/rsta.2011.0003, 2011.

1482 Gupta, A. Sen, Jourdain, N. C., Brown, J. N. and Monselesan, D.: Climate Drift in the  
1483 CMIP5 Models\*, *J. Clim.*, 26(21), 8597–8615, doi:10.1175/JCLI-D-12-00521.1,  
1484 2013.

1485 Hansen, J.: Efficacy of climate forcings, *J. Geophys. Res.*, 110(D18), D18104,  
1486 doi:10.1029/2005JD005776, 2005.

1487 Hartmann, J., West, J., Renforth, P., Köhler, P., De La Rocha, C. L., Wolf-Gladrow, D.,  
1488 Dürr, H. and Scheffran, J.: Enhanced Chemical Weathering as a Geoengineering  
1489 Strategy to Reduce Atmospheric Carbon Dioxide, a Nutrient Source and to  
1490 Mitigate Ocean Acidification, *Rev. Geophys.*, in press, 2013.

1491 Hauck, J., Köhler, P., Wolf-Gladrow, D. and Völker, C.: Iron fertilisation and

1492 century-scale effects of open ocean dissolution of olivine in a simulated CO<sub>2</sub>  
1493 removal experiment, *Environ. Res. Lett.*, 11(2), 24007, doi:10.1088/1748-  
1494 9326/11/2/024007, 2016.

1495 Heck, V., Gerten, D., Lucht, W. and Boysen, L. R.: Is extensive terrestrial carbon  
1496 dioxide removal a “green” form of geoengineering? A global modelling study,  
1497 *Glob. Planet. Change*, 137, 123–130, doi:10.1016/j.gloplacha.2015.12.008, 2016.

1498 Hofmann, M. and Schellnhuber, H. J.: Ocean acidification: a millennial challenge,  
1499 *Energy Environ. Sci.*, 3, 1883–1896, 2010.

1500 Holmes, G. and Keith, D. W.: An air-liquid contactor for large-scale capture of CO<sub>2</sub>  
1501 from air, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 370(1974), 4380–4403,  
1502 doi:10.1098/rsta.2012.0137, 2012.

1503 Ilyina, T., Wolf-Gladrow, D., Munhoven, G. and Heinze, C.: Assessing the potential  
1504 of calcium-based artificial ocean alkalinization to mitigate rising atmospheric CO<sub>2</sub>  
1505 and ocean acidification, *Geophys. Res. Lett.*, 40(22), 5909–5914,  
1506 doi:10.1002/2013GL057981, 2013.

1507 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working  
1508 Group I to the Fifth Assessment Report of the Intergovernmental Panel on  
1509 Climate Change., 2013.

1510 IPCC: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A:  
1511 Global and Sectoral Aspects. Contribution of Working Group II to the Fifth  
1512 Assessment Report of the Intergovernmental Panel on Climate Change [Field,  
1513 C.B., V.R. Barros, D.J. Dokken, K.J., Cambridge University Press, Cambridge, United  
1514 Kingdom and New York, NY, USA., 2014a.

1515 IPCC: Climate Change 2014: Mitigation of Climate Change, Cambridge University  
1516 Press., 2014b.

1517 Jones, C. D., Ciais, P., Davis, S. J., Friedlingstein, P., Gasser, T., Peters, G. P., Rogelj,  
1518 J., van Vuuren, D. P., Canadell, J. G., Cowie, A., Jackson, R. B., Jonas, M., Kriegler, E.,  
1519 Littleton, E., Lowe, J. A., Milne, J., Shrestha, G., Smith, P., Torvanger, A. and  
1520 Wiltshire, A.: Simulating the Earth system response to negative emissions,  
1521 *Environ. Res. Lett.*, 11(9), 95012, doi:10.1088/1748-9326/11/9/095012, 2016a.

1522 Jones, C. D., Arora, V., Friedlingstein, P., Bopp, L., Brovkin, V., Dunne, J., Graven, H.,  
1523 Hoffman, F., Ilyina, T., John, J. G., Jung, M., Kawamiya, M., Koven, C., Pongratz, J.,  
1524 Raddatz, T., Randerson, J. and Zaehle, S.: The C4MIP experimental protocol for

1525 CMIP6, *Geosci. Model Dev.*, (1), 1–52, doi:10.5194/gmd-2016-36, 2016b.

1526 Joos, F., Roth, R., Fuglestvedt, J. S., Peters, G. P., Enting, I. G., von Bloh, W., Brovkin, V., Burke, E. J., Eby, M., Edwards, N. R., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A. and Weaver, A. J.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmos. Chem. Phys.*, 13(5), 2793–2825, doi:10.5194/acp-13-2793-2013, 2013.

1534 Keller, D. P., Oschlies, A. and Eby, M.: A new marine ecosystem model for the University of Victoria Earth System Climate Model, *Geosci. Model Dev.*, 5(5), 1195–1220, doi:10.5194/gmd-5-1195-2012, 2012.

1537 Keller, D. P., Feng, E. Y. and Oschlies, A.: Potential climate engineering effectiveness and side effects during a high carbon dioxide-emission scenario, *Nat. Commun.*, 5, 1–11, doi:10.1038/ncomms4304, 2014.

1540 Kheshgi, H. S.: Sequestering atmospheric carbon dioxide by increasing ocean alkalinity, *Energy*, 20(9), 915–922, doi:[http://dx.doi.org/10.1016/0360-5442\(95\)00035-F](http://dx.doi.org/10.1016/0360-5442(95)00035-F), 1995.

1543 Köhler, P., Hartmann, J. and Wolf-Gladrow, D. a: Geoengineering potential of artificially enhanced silicate weathering of olivine., *Proc. Natl. Acad. Sci. U. S. A.*, 107(47), 20228–20233, doi:10.1073/pnas.1000545107, 2010.

1546 Köhler, P., Abrams, J. F., Völker, C., Hauck, J. and Wolf-Gladrow, D. A.: Geoengineering impact of open ocean dissolution of olivine on atmospheric CO<sub>2</sub>, surface ocean pH and marine biology, *Environ. Res. Lett.*, 8(1), 14009 [online] Available from: <http://stacks.iop.org/1748-9326/8/i=1/a=014009>, 2013.

1550 Kriegler, E., Tavoni, M., ABOUMAHBOUB, T., LUDERER, G., CALVIN, K., DEMAERE, G., KREY, V., RIAHI, K., R?SLER, H., SCHAEFFER, M. and VAN VUUREN, D. P.: WHAT DOES THE 2°C TARGET IMPLY FOR A GLOBAL CLIMATE AGREEMENT IN 2020? THE LIMITS STUDY ON DURBAN PLATFORM SCENARIOS, *Clim. Chang. Econ.*, 4(4), 1340008, doi:10.1142/S2010007813400083, 2013.

1555 Kriegler, E., Bauer, N., Popp, A., Humpenöder, F., Leimbach, M., Strefler, J., Baumstark, L., Bodirsky, B. L., Hilaire, J., Klein, D., Mouratiadou, I., Weindl, I., Bertram, C., Dietrich, J.-P., Luderer, G., Pehl, M., Pietzcker, R., Piontek, F., Lotze-

1558 Campen, H., Biewald, A., Bonsch, M., Giannousakis, A., Kreidenweis, U., Müller, C.,  
1559 Rolinski, S., Schultes, A., Schwanitz, J., Stevanovic, M., Calvin, K., Emmerling, J.,  
1560 Fujimori, S. and Edenhofer, O.: Fossil-fueled development (SSP5): An energy and  
1561 resource intensive scenario for the 21st century, *Glob. Environ. Chang.*,  
1562 doi:10.1016/j.gloenvcha.2016.05.015, 2016.

1563 Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte,  
1564 C. M. and Gattuso, J.-P.: Impacts of ocean acidification on marine organisms:  
1565 quantifying sensitivities and interaction with warming, *Glob. Chang. Biol.*, 19(6),  
1566 1884–1896, doi:10.1111/gcb.12179, 2013.

1567 Lackner, K. S., Brennan, S., Matter, J. M., Park, a.-H. a., Wright, a. and van der  
1568 Zwaan, B.: The urgency of the development of CO<sub>2</sub> capture from ambient air,  
1569 *Proc. Natl. Acad. Sci.*, 109(33), 13156–13162, doi:10.1073/pnas.1108765109,  
1570 2012.

1571 Lawrence, D. M., Hurt, G. C., Arneth, A., Brovkin, V., Calvin, K. V., Jones, A. D.,  
1572 Jones, C. D., Lawrence, P. J., de Noblet-Ducoudré, N., Pongratz, J., Seneviratne, S. I.  
1573 and Shevliakova, E.: The Land Use Model Intercomparison Project (LUMIP):  
1574 Rationale and experimental design, *Geosci. Model Dev. Discuss.*, 0, 1–42,  
1575 doi:10.5194/gmd-2016-76, 2016.

1576 MacDougall, A. H.: Reversing climate warming by artificial atmospheric carbon-  
1577 dioxide removal: Can a Holocene-like climate be restored?, *Geophys. Res. Lett.*,  
1578 40(20), 5480–5485, doi:10.1002/2013GL057467, 2013.

1579 Mao, J., Phipps, S. J., Pitman, A. J., Wang, Y. P., Abramowitz, G. and Pak, B.: The  
1580 CSIRO Mk3L climate system model v1.0 coupled to the CABLE land surface  
1581 scheme v1.4b: evaluation of the control climatology, *Geosci. Model Dev.*, 4(4),  
1582 1115–1131, doi:10.5194/gmd-4-1115-2011, 2011.

1583 Matear, R. J. and Hirst, A. C.: Long-term changes in dissolved oxygen  
1584 concentrations in the ocean caused by protracted global warming, *Global  
1585 Biogeochem. Cycles*, 17(4), n/a-n/a, doi:10.1029/2002GB001997, 2003.

1586 Matear, R. J. and Lenton, a.: Quantifying the impact of ocean acidification on our  
1587 future climate, *Biogeosciences*, 11(14), 3965–3983, doi:10.5194/bg-11-3965-  
1588 2014, 2014.

1589 Mathesius, S., Hofmann, M., Caldeira, K. and Schellnhuber, H. J.: Long-term  
1590 response of oceans to CO<sub>2</sub> removal from the atmosphere, *Nat. Clim. Chang.*,

1591 (August), doi:10.1038/nclimate2729, 2015.

1592 Matter, J. M., Stute, M., Snaebjornsdottir, S. O., Oelkers, E. H., Gislason, S. R.,

1593 Aradottir, E. S., Sigfusson, B., Gunnarsson, I., Sigurdardottir, H., Gunnlaugsson, E.,

1594 Axelsson, G., Alfredsson, H. A., Wolff-Boenisch, D., Mesfin, K., Taya, D. F. d. l. R.,

1595 Hall, J., Dideriksen, K. and Broecker, W. S.: Rapid carbon mineralization for

1596 permanent disposal of anthropogenic carbon dioxide emissions, *Science* (80-. ),

1597 352(6291), 1312–1314, doi:10.1126/science.aad8132, 2016.

1598 Matthews, H. D., Gillett, N. P., Stott, P. a and Zickfeld, K.: The proportionality of

1599 global warming to cumulative carbon emissions., *Nature*, 459(7248), 829–32,

1600 doi:10.1038/nature08047, 2009.

1601 Meehl, G. a., Moss, R., Taylor, K. E., Eyring, V., Stouffer, R. J., Bony, S. and Stevens,

1602 B.: Climate Model Intercomparisons: Preparing for the Next Phase, *Eos, Trans.*

1603 *Am. Geophys. Union*, 95(9), 77–78, doi:10.1002/2014EO090001, 2014.

1604 Meinshausen, M., Raper, S. C. B. and Wigley, T. M. L.: Emulating coupled

1605 atmosphere-ocean and carbon cycle models with a simpler model, *MAGICC6 -*

1606 *Part 1: Model description and calibration*, *Atmos. Chem. Phys.*, 11(4), 1417–

1607 1456, doi:10.5194/acp-11-1417-2011, 2011.

1608 Meissner, K. J., Weaver, A. J., Matthews, H. D. and Cox, P. M.: The role of land

1609 surface dynamics in glacial inception: A study with the UVic Earth System Model,

1610 *Clim. Dyn.*, 21, 515–537, 2003.

1611 Millennium Ecosystem Assesment: *Ecosystems and Human Well-Being: Synthesis*, Island Press, Washington, D.C., 2005.

1612 Ming, T., de\_Richter, R., Shen, S. and Caillol, S.: Fighting global warming by

1613 greenhouse gas removal: destroying atmospheric nitrous oxide thanks to

1614 synergies between two breakthrough technologies, *Environ. Sci. Pollut. Res.*,

1615 doi:10.1007/s11356-016-6103-9, 2016.

1616 Morice, C. P., Kennedy, J. J., Rayner, N. a. and Jones, P. D.: Quantifying

1617 uncertainties in global and regional temperature change using an ensemble of

1618 observational estimates: The HadCRUT4 data set, *J. Geophys. Res. Atmos.*, 117(8),

1619 1–22, doi:10.1029/2011JD017187, 2012.

1620 National Research Council: *Climate Intervention*, National Academies Press,

1621 Washington, D.C., 2015.

1622 O'Neill, B. C., Tebaldi, C., Van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G.,

1624 Knutti, R., Kriegler, E., Lamarque, J. F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K. and  
1625 Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for  
1626 CMIP6, *Geosci. Model Dev.*, 9(9), 3461–3482, doi:10.5194/gmd-9-3461-2016,  
1627 2016.

1628 Orr, J. C., Najjar, R. G., Aumont, O., Bopp, L., Bullister, J. L., Danabasoglu, G.,  
1629 Doney, S. C., Dunne, J. P., Dutay, J.-C., Graven, H., Griffies, S. M., John, J. G., Joos, F.,  
1630 Levin, I., Lindsay, K., Matear, R. J., McKinley, G. A., Mouchet, A., Oschlies, A.,  
1631 Romanou, A., Schlitzer, R., Tagliabue, A., Tanhua, T. and Yool, A.: Biogeochemical  
1632 protocols and diagnostics for the CMIP6 Ocean Model Intercomparison Project  
1633 (OMIP), *Geosci. Model Dev. Discuss.*, 0(July), 1–45, doi:10.5194/gmd-2016-155,  
1634 2016.

1635 Oschlies, A. and Klepper, G.: Research for assessment, not deployment, of Climate  
1636 Engineering: The German Research Foundation's Priority Program SPP 1689,  
1637 *Earth's Futur.*, 5(1), 128–134, doi:10.1002/2016EF000446, 2017.

1638 Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré, C.,  
1639 Marland, G., Raupach, M. R. and Wilson, C.: The challenge to keep global warming  
1640 below 2 °C, *Nat. Clim. Chang.*, 3(1), 4–6, doi:10.1038/nclimate1783, 2013.

1641 Phipps, S. J., Rotstayn, L. D., Gordon, H. B., Roberts, J. L., Hirst, a. C. and Budd, W.  
1642 F.: The CSIRO Mk3L climate system model version 1.0 – Part 1: Description and  
1643 evaluation, *Geosci. Model Dev.*, 4(2), 483–509, doi:10.5194/gmd-4-483-2011,  
1644 2011.

1645 Pongratz, J., Reick, C. H., Raddatz, T., Caldeira, K. and Claussen, M.: Past land use  
1646 decisions have increased mitigation potential of reforestation, *Geophys. Res.*  
1647 *Lett.*, 38(15), 1–5, doi:10.1029/2011GL047848, 2011.

1648 Le Quéré, C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I.,  
1649 Friedlingstein, P., Peters, G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J.  
1650 I., Keeling, R. F., Tans, P., Arneth, A., Bakker, D. C. E., Barbero, L., Bopp, L., Chang,  
1651 J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M., Feely, R. A., Gkritzalis, T., Harris, I.,  
1652 Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Klein Goldewijk, K., Koven, C.,  
1653 Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N.,  
1654 Millero, F., Munro, D. R., Murata, A., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y.,  
1655 O'Brien, K., Olsen, A., Ono, T., Pérez, F. F., Pfeil, B., Pierrot, D., Poulter, B., Rehder,  
1656 G., Rödenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R., Steinhoff, T.,

1657 Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Laan-Luijkx, I. T.,  
1658 van der Werf, G. R., van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle,  
1659 S. and Zeng, N.: Global Carbon Budget 2015, *Earth Syst. Sci. Data*, 7(2), 349–396,  
1660 doi:10.5194/essd-7-349-2015, 2015.

1661 Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P.,  
1662 Manning, A. C., Boden, T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S.,  
1663 Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L. P., Ciais,  
1664 P., Currie, K., Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I.,  
1665 Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K., Kato, E.,  
1666 Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi,  
1667 D., Melton, J. R., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M.,  
1668 S., Nakaoka, S., O&apos;Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D.,  
1669 Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R.,  
1670 Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der  
1671 Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J. and  
1672 Zaehle, S.: Global Carbon Budget 2016, *Earth Syst. Sci. Data Discuss.*, 1–3,  
1673 doi:10.5194/essd-2016-51, 2016.

1674 Renforth, P., Jenkins, B. G. and Kruger, T.: Engineering challenges of ocean liming,  
1675 *Energy*, 60, 442–452, doi:10.1016/j.energy.2013.08.006, 2013.

1676 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S.,  
1677 Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC,  
1678 S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T.,  
1679 Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J.,  
1680 Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G.,  
1681 Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont,  
1682 Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A. and Tavoni, M.:  
1683 The Shared Socioeconomic Pathways and their energy, land use, and greenhouse  
1684 gas emissions implications: An overview, *Glob. Environ. Chang.*, 42, 153–168,  
1685 doi:10.1016/j.gloenvcha.2016.05.009, 2017.

1686 Rickels, W., Klepper, G., Dovern, J., Betz, G., Brachatzek, N., Cacean, S., Güssow, K.,  
1687 Heintzenberg, J., Hiller, S., Hoose, C., Leisner, T., Oschlies, A., Platt, U., Proelß, A.,  
1688 Renn, O., Schäfer, S. and Zürn, M.: Large-Scale Intentional Interventions into the  
1689 Climate System? Assessing the Climate Engineering Debate., 2011.

1690 Ridgwell, A., Maslin, M. A. and Watson, A. J.: Reduced effectiveness of terrestrial  
1691 carbon sequestration due to an antagonistic response of ocean productivity,  
1692 *Geophys. Res. Lett.*, 29(6), doi:10.1029/2001GL014304, 2002.

1693 Rogelj, J., Luderer, G., Pietzcker, R. C., Kriegler, E., Schaeffer, M., Krey, V. and Riahi,  
1694 K.: Energy system transformations for limiting end-of-century warming to below  
1695 1.5 °C, *Nat. Clim. Chang.*, 5(6), 519–527, doi:10.1038/nclimate2572, 2015a.

1696 Rogelj, J., Schaeffer, M., Meinshausen, M., Knutti, R., Alcamo, J., Riahi, K. and Hare,  
1697 W.: Zero emission targets as long-term global goals for climate protection,  
1698 *Environ. Res. Lett.*, 10(10), 105007, doi:10.1088/1748-9326/10/10/105007,  
1699 2015b.

1700 Sanz-Pérez, E. S., Murdock, C. R., Didas, S. A. and Jones, C. W.: Direct Capture of CO  
1701 <sub>2</sub> from Ambient Air, *Chem. Rev.*, acs.chemrev.6b00173,  
1702 doi:10.1021/acs.chemrev.6b00173, 2016.

1703 Schäfer, S., Lawrence, M., Stelzer, H., Born, W., Low, S., Aaheim, A., Adriázola, P.,  
1704 Betz, G., Boucher, O., Carius, A., Devine-Right, P., Gullberg, A. T., Haszeldine, S.,  
1705 Haywood, J., Houghton, K., Ibarrola, R., Irvine, P., Kristjansson, J.-E., Lenton, T.,  
1706 Link, J. S. A., Maas, A., Meyer, L., Muri, H., Oschlies, A., Proelß, A., Rayner, T.,  
1707 Rickels, W., Ruthner, L., Scheffran, J., Schmidt, H., Schulz, M., Scott, V., Shackley, S.,  
1708 Tänzler, D., Watson, M. and Vaughan, N.: The European Transdisciplinary  
1709 Assessment of Climate Engineering (EuTRACE): Removing Greenhouse Gases  
1710 from the Atmosphere and Reflecting Sunlight away from Earth., 2015.

1711 Schuiling, R. D. and Krijgsman, P.: Enhanced weathering: An effective and cheap  
1712 tool to sequester CO <sub>2</sub>, *Clim. Change*, 74(1–3), 349–354, doi:10.1007/s10584-  
1713 005-3485-y, 2006.

1714 Scott, V., Gilfillan, S., Markusson, N., Chalmers, H. and Haszeldine, R. S.: Last  
1715 chance for carbon capture and storage, *Nat. Clim. Chang.*, 3(2), 105–111, doi:10.1038/Nclimate1695, 2013.

1717 Scott, V., Haszeldine, R. S., Tett, S. F. B. and Oschlies, A.: Fossil fuels in a trillion  
1718 tonne world, *Nat. Clim. Chang.*, 5(5), 419–423, doi:10.1038/nclimate2578, 2015.

1719 Séférian, R., Gehlen, M., Bopp, L., Resplandy, L., Orr, J. C., Marti, O., Dunne, J. P.,  
1720 Christian, J. R., Doney, S. C., Ilyina, T., Lindsay, K., Halloran, P., Heinze, C.,  
1721 Segschneider, J. and Tjiputra, J.: Inconsistent strategies to spin up models in  
1722 CMIP5: implications for ocean biogeochemical model performance assessment,

1723 Geosci. Model Dev. Discuss., 8(10), 8751–8808, doi:10.5194/gmdd-8-8751-2015,  
1724 2015.

1725 Smith, P.: Soil carbon sequestration and biochar as negative emission  
1726 technologies, *Glob. Chang. Biol.*, n/a-n/a, doi:10.1111/gcb.13178, 2016.

1727 Smith, P., Davis, S. J., Creutzig, F., Fuss, S., Minx, J., Gabrielle, B., Kato, E., Jackson, R.  
1728 B., Cowie, A., Kriegler, E., van Vuuren, D. P., Rogelj, J., Ciais, P., Milne, J., Canadell, J.  
1729 G., McCollum, D., Peters, G., Andrew, R., Krey, V., Shrestha, G., Friedlingstein, P.,  
1730 Gasser, T., Grübler, A., Heidug, W. K., Jonas, M., Jones, C. D., Kraxner, F., Littleton,  
1731 E., Lowe, J., Moreira, J. R., Nakicenovic, N., Obersteiner, M., Patwardhan, A.,  
1732 Rogner, M., Rubin, E., Sharifi, A., Torvanger, A., Yamagata, Y., Edmonds, J. and  
1733 Yongsung, C.: Biophysical and economic limits to negative CO<sub>2</sub> emissions, *Nat. Clim. Chang.*, 6(1), 42–50, doi:10.1038/nclimate2870, 2015.

1735 Sonntag, S., Pongratz, J., Reick, C. H. and Schmidt, H.: Reforestation in a high-CO<sub>2</sub>  
1736 world-Higher mitigation potential than expected, lower adaptation potential  
1737 than hoped for, *Geophys. Res. Lett.*, (May), 1–8, doi:10.1002/2016GL068824,  
1738 2016.

1739 Taylor, K. E., Stouffer, R. J. and Meehl, G. a.: An Overview of CMIP5 and the  
1740 Experiment Design, *Bull. Am. Meteorol. Soc.*, 93(4), 485–498, doi:10.1175/BAMS-  
1741 D-11-00094.1, 2012.

1742 The Royal Society: *Geoengineering the climate.*, 2009.

1743 Tokarska, K. B. and Zickfeld, K.: The effectiveness of net negative carbon dioxide  
1744 emissions in reversing anthropogenic climate change, *Environ. Res. Lett.*, 10(9),  
1745 94013, doi:10.1088/1748-9326/10/9/094013, 2015.

1746 UNFCCC: Paris Agreement of the 21st session of the Conference of Parties on  
1747 climate change., 2016.

1748 Unger, N.: Human land-use-driven reduction of forest volatiles cools global  
1749 climate, *Nat. Clim. Chang.*, 4(October), 907–910, doi:10.1038/nclimate2347,  
1750 2014.

1751 Vaughan, N. E. and Gough, C.: Expert assessment concludes negative emissions  
1752 scenarios may not deliver, *Environ. Res. Lett.*, 11(9), 95003, doi:10.1088/1748-  
1753 9326/11/9/095003, 2016.

1754 Vaughan, N. E. and Lenton, T. M.: A review of climate geoengineering proposals,  
1755 *Clim. Change*, 109, 745–790, doi:10.1007/s10584-011-0027-7, 2011.

1756 Vichi, M., Navarra, A. and Fogli, P. G.: Adjustment of the natural ocean carbon  
1757 cycle to negative emission rates, *Clim. Change*, 1–14, doi:10.1007/s10584-012-  
1758 0677-0, 2013.

1759 Walker, J. C. G., Hays, P. B. and Kasting, J. F.: A negative feedback mechanism for  
1760 the long-term stabilization of Earth's surface temperature, *J. Geophys. Res.*,  
1761 86(C10), 9776, doi:10.1029/JC086iC10p09776, 1981.

1762 Wang, X., Heald, C. L., Ridley, D. a., Schwarz, J. P., Spackman, J. R., Perring, a. E.,  
1763 Coe, H., Liu, D. and Clarke, a. D.: Exploiting simultaneous observational  
1764 constraints on mass and absorption to estimate the global direct radiative  
1765 forcing of black carbon and brown carbon, *Atmos. Chem. Phys.*, 14(20), 10989–  
1766 11010, doi:10.5194/acp-14-10989-2014, 2014.

1767 Wang, Y. P., Law, R. M. and Pak, B.: A global model of carbon, nitrogen and  
1768 phosphorus cycles for the terrestrial biosphere, *Biogeosciences*, 7(7), 2261–  
1769 2282, doi:10.5194/bg-7-2261-2010, 2010.

1770 Weaver, A. J., Eby, M., Wiebe, E. C., Bitz, C. M., Duffy, P. B., Ewen, T. L., Fanning, A.  
1771 F., Holland, M. M., MacFadyen, A., Matthews, H. D., Meissner, K. J., Saenko, O.,  
1772 Schmittner, A., Wang, H. and Yoshimori, M.: The UVic earth system climate  
1773 model: Model description, climatology, and applications to past, present and  
1774 future climates, *Atmosphere-Ocean*, 39(4), 361–428,  
1775 doi:10.1080/07055900.2001.9649686, 2001.

1776 Wolf-Gladrow, D. a., Zeebe, R. E., Klaas, C., Kötzinger, A. and Dickson, A. G.: Total  
1777 alkalinity: The explicit conservative expression and its application to  
1778 biogeochemical processes, *Mar. Chem.*, 106(1–2), 287–300,  
1779 doi:10.1016/j.marchem.2007.01.006, 2007.

1780 Wu, P., Ridley, J., Pardaens, A., Levine, R. and Lowe, J.: The reversibility of CO<sub>2</sub>  
1781 induced climate change, *Clim. Dyn.*, doi:10.1007/s00382-014-2302-6, 2014.

1782 Zhang, Q., Wang, Y. P., Matear, R. J., Pitman, A. J. and Dai, Y. J.: Nitrogen and  
1783 phosphorous limitations significantly reduce future allowable CO<sub>2</sub> emissions,  
1784 *Geophys. Res. Lett.*, 41(2), 632–637, doi:10.1002/2013GL058352, 2014.

1785 Zickfeld, K., Eby, M., Weaver, A. J., Alexander, K., Crespin, E., Edwards, N. R.,  
1786 Eliseev, A. V., Feulner, G., Fichefet, T., Forest, C. E., Friedlingstein, P., Goosse, H.,  
1787 Holden, P. B., Joos, F., Kawamiya, M., Kicklighter, D., Kienert, H., Matsumoto, K.,  
1788 Mokhov, I. I., Monier, E., Olsen, S. M., Pedersen, J. O. P., Perrette, M., Philippon-

1789 Berthier, G. G., Ridgwell, A., Schlosser, A., Schneider Von Deimling, T., Shaffer, G.,  
1790 Sokolov, A., Spahni, R., Steinacher, M., Tachiiri, K., Tokos, K. S., Yoshimori, M.,  
1791 Zeng, N. and Zhao, F.: Long-Term Climate Change Commitment and Reversibility:  
1792 An EMIC Intercomparison, *J. Clim.*, 26(16), 5782–5809, doi:10.1175/jcli-d-12-  
1793 00584.1, 2013.  
1794 Zickfeld, K., MacDougall, A. H. and Matthews, H. D.: On the proportionality  
1795 between global temperature change and cumulative CO<sub>2</sub> emissions during  
1796 periods of net negative CO<sub>2</sub> emissions, *Environ. Res. Lett.*, 11(5), 55006,  
1797 doi:10.1088/1748-9326/11/5/055006, 2016.  
1798  
1799

## CDR-MIP GMDD manuscript tables

Table 1. Overview of CDR-MIP experiments. In the "Forcing methods" column, "All" means "all anthropogenic, solar, and volcanic forcing". Anthropogenic forcing includes aerosol emissions, non-CO<sub>2</sub> greenhouse gas emissions, and land use changes.

Short Name	Long Name	Tier	Experiment Description	Forcing methods	Major purpose
<i>C1</i>	Climate and carbon cycle reversibility experiment	1	CO <sub>2</sub> prescribed to increase at 1% yr <sup>-1</sup> to 4x pre-industrial CO <sub>2</sub> and then decrease at 1% yr <sup>-1</sup> until again at a pre-industrial level, after which the simulation continues for as long as possible	CO <sub>2</sub> concentration prescribed	Evaluate climate reversibility
<i>C2_pi-pulse</i>	Instantaneous CO <sub>2</sub> removal / addition from an unperturbed climate experiment	1	100 Gt C is instantly removed (negative pulse) from a steady-state pre-industrial atmosphere; 100 Gt C is instantly added (positive pulse) to a steady-state pre-industrial atmosphere	CO <sub>2</sub> concentration calculated (i.e., freely evolving)	Evaluate climate and C-cycle response of an unperturbed system to atmospheric CO <sub>2</sub> removal; comparison with the positive pulse response
<i>C2_yr2010-pulse</i>	Instantaneous CO <sub>2</sub> removal / addition from a perturbed climate experiment	3	100 Gt C is instantly removed (negative pulse) from a near present-day atmosphere; 100 Gt C is instantly added (positive pulse) to a near present-day atmosphere	All; CO <sub>2</sub> concentration calculated (i.e., emission driven)*	Evaluate climate and C-cycle response of a perturbed system to atmospheric CO <sub>2</sub> removal; comparison with the positive pulse response
<i>C2_overshoot</i>	Emission driven SSP5-3.4-OS scenario experiment	2	SSP5-3.4-overshoot scenario where CO <sub>2</sub> emissions are initially high and then rapidly reduced, becoming negative	All; CO <sub>2</sub> concentration calculated (i.e., emission driven)	Evaluate the Earth system response to CDR in an overshoot climate change scenario
<i>C3</i>	Afforestation/ reforestation experiment	2	Long-term extension of an experiment with forcing from a high CO <sub>2</sub> emission scenario (SSP5-8.5), but with land use prescribed from a scenario with high levels of afforestation and reforestation (SSP1-2.6)	All; CO <sub>2</sub> concentration calculated (i.e., emission driven)	Evaluate the long-term Earth system response to afforestation/ reforestation during a high CO <sub>2</sub> emission climate change scenario
<i>C4</i>	Ocean alkalization experiment	2	A high CO <sub>2</sub> emission scenario (SSP5-8.5) with 0.14 Pmol yr <sup>-1</sup> alkalinity added to ice-free ocean surface waters from the year 2020 onward	All; CO <sub>2</sub> concentration calculated (i.e., emission driven)	Evaluate the Earth system response to ocean alkalization during a high CO <sub>2</sub> emission climate change scenario

\*In this experiment CO<sub>2</sub> is first prescribed to diagnose emissions, however, the key simulations calculate the CO<sub>2</sub> concentration.

Table 2. Climate and carbon cycle reversibility experiment (C1) simulations. All simulations are required to complete the experiment.

Simulation ID	Simulation description	Owning MIP	Run length (years)	Initialized using a restart from
<i>piControl</i>	Pre-industrial prescribed CO <sub>2</sub> control simulation	CMIP6 DECK	100*	The model spin-up
<i>1pctCO2</i>	Prescribed 1% yr <sup>-1</sup> CO <sub>2</sub> increase to 4× the pre-industrial level	CMIP6 DECK	140**	<i>piControl</i>
<i>1pctCO2-cdr</i>	1% yr <sup>-1</sup> CO <sub>2</sub> decrease from 4× the pre-industrial level until the pre-industrial CO <sub>2</sub> level is reached and held for as long as possible	CDR-MIP	200 min. 5000 max.	<i>1pctCO2</i>

\*This CMIP6 DECK should have been run for at least 500 years. Only the last 100 years are needed as a control for C1.

\*\*This CMIP6 DECK experiment is 150 years long. A restart for C1 should be generated after 139 years when CO<sub>2</sub> is 4 times that of *piControl*.

Table 3. Instantaneous CO<sub>2</sub> removal from an unperturbed climate experiment (C2\_pi-pulse) simulations.

Simulation ID	Simulation description	Owning MIP	Run length (years)	Initialized using a restart from
<i>esm-piControl</i>	Pre-industrial freely evolving CO <sub>2</sub> control simulation	CMIP6 DECK	100*	The model spin up
<i>esm-pi-cdr-pulse</i>	100 Gt C is instantly removed (negative pulse) from a pre-industrial atmosphere	CDR-MIP	100 min. 5000 max.	<i>esm-piControl</i>
<i>esm-pi-co2pulse</i>	100 Gt C is instantly added to (positive pulse) a pre-industrial atmosphere	CDR-MIP	100 min. 5000 max.	<i>esm-piControl</i>

\*This CMIP6 DECK should have been run for at least 500 years. Only the last 100 years are needed as a control for C2.1.

Table 4. Instantaneous CO<sub>2</sub> removal from a perturbed climate experiment (C2\_yr2010-pulse) simulations.

Simulation ID	Simulation description	Owning MIP	Run length (years)	Initialized using a restart from
<i>historical</i>	Historical atmospheric CO <sub>2</sub> (and other forcing) is prescribed until a concentration of 389ppm CO <sub>2</sub> is reached	CMIP6 DECK	160*	<i>piControl</i>
<i>yr2010co2</i>	Branching from <i>historical</i> , atmospheric CO <sub>2</sub> is held constant (prescribed) at 389ppm; other forcing is also held constant at the 2010 level	CDR-MIP	105 min. 5000 max.	<i>historical</i>
<i>esm-hist-yr2010co2-control</i>	Control run forced using CO <sub>2</sub> emissions diagnosed from <i>historical</i> and <i>yr2010co2</i> simulations; other forcing as in <i>historical</i> until 2010 after which it is constant	CDR-MIP	265 min. 5160 max.	<i>esm-hist- piControl</i> or <i>piControl</i>
<i>esm-yr2010co2-noemit</i>	Control run that branches from <i>esm-hist-yr2010co2-control</i> in year 2010 with CO <sub>2</sub> emissions set to zero 5 years after the start of the simulation	CDR-MIP	105 min. 5000 max.	<i>esm-hist- yr2010co2-control</i>
<i>esm-yr2010co2-cdr-pulse</i>	Branches from <i>esm-hist-yr2010co2-control</i> in year 2010 with 100 Gt C instantly removed (negative pulse) from the atmosphere 5 years after the start of the simulation	CDR-MIP	105 min. 5000 max.	<i>esm-hist- yr2010co2-control</i>
<i>esm-yr2010co2-co2pulse</i>	Branches from <i>esm-hist-yr2010co2-control</i> in year 2010 with 100 Gt C instantly added to (positive pulse) the atmosphere 5 years after the start of the simulation	CDR-MIP	105 min. 5000 max.	<i>esm-hist- yr2010co2-control</i>

\*This CMIP6 DECK continues until the year 2015 but only the first 160 years are need for C2\_yr2010-pulse.

Table 5. Emission driven SSP5-3.5-OS scenario experiment (*C2\_overshoot*) simulations. All simulations are required to complete the experiment.

Simulation ID	Simulation description	Owning MIP	Run length (years)	Initialized using a restart from
<i>esm-hist</i>	Historical simulation forced with CO <sub>2</sub> emissions	CMIP6 DECK	265	<i>esm-piControl</i> or <i>piControl</i>
<i>esm-ssp534-over</i>	CO <sub>2</sub> emission-driven SSP5-3.4 overshoot scenario simulation	CDR-MIP	85	<i>esm-hist</i>
<i>esm-ssp534-over-ext</i>	Long-term extension of the CO <sub>2</sub> emission-driven SSP5-3.4 overshoot scenario	CDR-MIP	200 min. 5000 max.	<i>esm-ssp534-over</i>

Table 6. Afforestation/ reforestation experiment (*C3*) simulations. All simulations are required to complete the experiment.

Simulation ID	Simulation description	Owning MIP	Run length (years)	Initialized using a restart from
<i>esm-ssp585</i>	CO <sub>2</sub> emission driven SSP5-8.5 scenario	C4MIP	85	<i>esm-hist</i>
<i>esm-ssp585-ssp126Lu</i>	CO <sub>2</sub> emission driven SSP5-8.5 scenario with SSP1-2.6 land use forcing	LUMIP	85	<i>esm-hist</i>
<i>esm-ssp585-ssp126Lu-ext</i>	CO <sub>2</sub> emission-driven SSP5-3.4 overshoot scenario simulation	CDR-MIP	200 min. 5000 max.	<i>esm-ssp585-ssp126Lu</i>
<i>esm-ssp585ext</i>	Long-term extension of the CO <sub>2</sub> emission-driven SSP5-8.5 scenario	CDR-MIP	200 min. 5000 max.	<i>esm-ssp585</i>

Table 7. Ocean alkalinization (C4) experiment simulations. "Pr" in the Tier column indicates a prerequisite experiment.

Simulation ID	Tier	Simulation description	Owning MIP	Run length (years)	Initialized using a restart from
<i>esm-ssp585</i>	Pr	CO <sub>2</sub> emission driven SSP5-8.5 scenario	C4MIP	85	<i>esm-hist</i>
<i>esm-ssp585-ocean-alk</i>	2	SSP5-8.5 scenario with 0.14 Pmol yr <sup>-1</sup> alkalinity added to ice-free ocean surface waters from the year 2020 onward	CDR-MIP	65	<i>esm-ssp585</i>
<i>esm-ssp585-ocean-alk-stop</i>	3	Termination simulation to investigate an abrupt stop in ocean alkalinization in the year 2070	CDR-MIP	30*	<i>esm-ssp585-ocean-alk</i>
<i>esm-ssp585ext</i>	3	Long-term extension of the CO <sub>2</sub> emission-driven SSP5-8.5 scenario	CDR-MIP	200 min. 5000 max.	<i>esm-ssp585</i>
<i>esm-ssp585-ocean-alk-ext</i>	3	Long-term extension of the <i>esm-ssp585-ocean-alk</i> simulation	CDR-MIP	200 min. 5000 max.	<i>esm-ssp585-ocean-alk</i>

\*If the *esm-ssp585ext* simulation is being conducted this may be extended for more than 200 more years (up to 5000 years).

Table 8. Model output frequency for 3-D models with seasonality. Box models and EMICs without seasonality are expected to generate annual global mean output for the duration of all experiments. For longer simulations (right column) if possible 3-D monthly data should be written out for one year every 100 years. For models with interannual variability, e.g., ESMs, monthly data should be written out for a 10-year period every 100 years so that a climatology may be developed. The years referred to in the table indicate simulations years, e.g. years from the start of the run, not that of any particular scenario.

Experiment Short Name	Individual simulation output frequency	
	Monthly gridded 3-D output	Annual global mean output + climatological output at 100 year intervals
<i>C1</i>	<i>piControl</i> (last 100 years) <i>1pctCO2</i> <i>1pctCO2-cdr</i> (initial 200 years)	<i>1pctCO2-cdr</i> (from year 200 onward)
<i>C2_pi-pulse</i>	<i>esm-piControl</i> <i>esm-pi-cdr-pulse</i> (initial 100 years) <i>esm-pi-co2pulse</i> (initial 100 years)	<i>esm-pi-cdr-pulse</i> (from year 100 onward) <i>esm-pi-co2pulse</i> (from year 100 onward)
<i>C2_yr2010-pulse</i>	<i>esm-hist-yr2010co2-control</i> (initial 105 years) <i>esm-yr2010co2-noemit</i> <i>esm-yr2010co2-cdr-pulse</i> <i>esm-yr2010co2-co2pulse</i>	<i>esm-hist-yr2010co2-control</i> <i>esm-yr2010co2-noemit</i> <i>esm-yr2010co2-cdr-pulse</i> <i>esm-yr2010co2-co2pulse</i>
<i>C2_overshoot</i>	<i>esm-hist</i> <i>esm-ssp534-over</i> <i>esm-ssp534-over-ext</i> (initial 200 years)	<i>esm-ssp534-over-ext</i> (from year 200 onward)**
<i>C3</i>	<i>esm-ssp585ext</i> (initial 200 years) <i>esm-ssp585-ssp126Lu</i> <i>esm-ssp585-ssp126Lu-ext</i> (initial 200 years)	<i>esm-ssp585ext</i> (from year 200 onward)** <i>esm-ssp585-ssp126Lu-ext</i> (from year 200 onward)**
<i>C4</i>	<i>esm-ssp585</i> <i>esm-ssp585-ocean-alk</i> <i>esm-ssp585-ocean-alk-stop</i> (initial 200 years) <i>esm-ssp585ext</i> (initial 200 years) <i>esm-ssp585-ocean-alk-ext</i> (initial 200 years)	<i>esm-ssp585-ocean-alk-stop</i> (from year 200 onward)** <i>esm-ssp585ext</i> (from year 200 onward)** <i>esm-ssp585-ocean-alk-ext</i> (from year 200 onward)**

\*In the *historical* and *yr2010co2* simulations output is needed only to diagnose (at least annually) CO<sub>2</sub> emissions.

\*\*This is from scenario year 2300 onward.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

1 Figure 1. Schematic of the CDR-MIP climate and carbon cycle reversibility  
2 experimental protocol (*C1*). From a preindustrial run at steady state atmospheric  
3 CO<sub>2</sub> is prescribed to increase and then decrease over a ~280 year period, after  
4 which it is held constant for as long as computationally possible.

5

6 Figure 2. Exemplary climate and carbon cycle reversibility experiment (*C1*)  
7 results with the CSIRO-Mk3L-COAL Earth system model and the University of  
8 Victoria (UVic) Earth system model of intermediate complexity (models are  
9 described in Appendix D). The left panels show annual global mean (a)  
10 temperature anomalies (°C; relative to pre-industrial temperatures) and (c) the  
11 atmosphere to ocean carbon fluxes (Pg C yr<sup>-1</sup>) versus the atmospheric CO<sub>2</sub> (ppm)  
12 during the first 280 years of the experiment (i.e., when CO<sub>2</sub> is increasing and  
13 decreasing). The right panels show the same (b) temperature anomalies and (d)  
14 the atmosphere to ocean carbon fluxes versus time. Note that the CSIRO-Mk3L-  
15 COAL simulation was only 400 years long.

16

17 Figure 3. Schematic of the CDR-MIP instantaneous CO<sub>2</sub> removal / addition from  
18 an unperturbed climate experimental protocol (*C2\_pi-pulse*). Models are spun-up  
19 for as long as possible with a prescribed preindustrial atmospheric CO<sub>2</sub>  
20 concentration. Then atmospheric CO<sub>2</sub> is allowed to freely evolve for at least 100  
21 years as a control run. The negative / positive pulse experiments are conducted  
22 by instantly removing or adding 100 Gt C to the atmosphere of a simulation  
23 where the atmosphere is at steady state and CO<sub>2</sub> can freely evolve. These runs  
24 continue for as long as computationally possible.

25

26 Figure 4. Exemplary instantaneous CO<sub>2</sub> removal from a preindustrial climate  
27 experiment (*C2\_pi-pulse*) results from the *esm-pi-cdr-pulse* simulation with the  
28 CSIRO-Mk3L-COAL Earth system model and the University of Victoria (UVic)  
29 Earth system model of intermediate complexity (models are described in  
30 Appendix D). (a) shows atmospheric CO<sub>2</sub> vs. time, (b) the land to atmosphere  
31 carbon flux vs. time, and (c) the ocean to atmosphere carbon flux vs. time. Note  
32 that the Mk3L-COAL simulation was only 184 years long.

33

34 Figure 5. Schematic of the CDR-MIP instantaneous CO<sub>2</sub> removal / addition from a  
35 perturbed climate experimental protocol (*C2\_yr2010-pulse*). Top panel: Initially  
36 historical CO<sub>2</sub> forcing is prescribed and then held constant at 389 ppm (~ year  
37 2010) while CO<sub>2</sub> emissions are diagnosed. Bottom panel: A control simulation is  
38 conducted using the diagnosed emissions. The negative / positive pulse  
39 experiments are conducted by instantly removing or adding 100 Gt C to the  
40 atmosphere of the CO<sub>2</sub> emission-driven simulation 5 years after CO<sub>2</sub> reaches 389  
41 ppm. Another control simulation is also conducted that sets emissions to zero at  
42 the time of the negative pulse. The emission-driven simulations continue for as  
43 long as computationally possible.

44

45

46 Figure 6. Schematic of the CDR-MIP emission-driven SSP5-3.4-OS scenario  
47 experimental protocol (*C2\_overshoot*). A CO<sub>2</sub> emission-driven historical  
48 simulation is conducted until the year 2015. Then an emission-driven simulation  
49 with SSP5-3.4-OS scenario forcing is conducted. This simulation is extended until  
50 the year 2300 using SSP5-3.4-OS scenario long-term extension forcing.  
51 Thereafter, runs may continue for as long as computationally possible with  
52 constant forcing after the year 2300.