

Original comments are in black text. Our responses are in blue text.

**Associate Editor (Remarks to Author):**

In my role as Executive editor of GMD, I would like to bring to your attention our Editorial version 1.1:

<http://www.geosci-model-dev.net/8/3487/2015/gmd-8-3487-2015.html>

This highlights some requirements of papers published in GMD, which is also available on the GMD website in the ‘Manuscript Types’ section:

[http://www.geoscientific-model-development.net/submission/manuscript\\_types.html](http://www.geoscientific-model-development.net/submission/manuscript_types.html)

In particular, please note that for your paper, the following requirements have not been met in the Discussions paper:

- "The main paper must give the model name and version number (or other unique identifier) in the title."

You applied this rule only for the first half of the title. Please include also the acronym and version number of the GHG forecast model, e.g. GEM-MACH-GHG vY.z .

Thank you for your comment. We modified the title and the relevant text in the revised version of the manuscript. The new title is “*Coupling the Canadian Terrestrial Ecosystem Model (CTEM v. 2.0) to Environment and Climate Change Canada’s greenhouse gas forecast model (v.107-glb)*”.

**Anonymous Referee #1**

**General comments**

This paper describes the coupling between the Canadian Terrestrial Ecosystem Model (CTEM v2.0) and Environment and Climate Change Canada’s greenhouse gas forecast model (GEM-MACH-GHG). The radiation, surface temperature, and precipitation fields etc. calculated from GEM-MACH-GHG feed to CTEMv2.0 every 30 minutes, and the net biosphere fluxes from CTEM v2.0 are used as surface boundary conditions to drive CO<sub>2</sub> simulations by GEM-MACH-GHG. The ultimate goal of this coupling is to do carbon cycle data assimilation to constrain biosphere model parameters and surface carbon fluxes. The authors tested the performance of such coupling by evaluating the meteorology fields, and carbon fluxes including gross primary production and net terrestrial biosphere fluxes. At last, they tested the impact of using net biosphere fluxes from the coupled system as priors on atmospheric flux inversion with GEOS-Chem 4D-Var system. The paper is well structured. However, the rationale of such coupling is not well described, the evaluation of the model performance can be further improved, and some conclusions in the paper are not supported by the results. My detailed comments are below.

We thank the referee for the careful review of our paper, and for the helpful comments. In the revised version of the paper, we significantly improve the rationale for coupling CTEM and GEM-MACH-GHG (see the response to the detailed comment #1). We believe that the revised

manuscript has greatly benefitted from the comments of both Reviewers. Below are detailed responses to each comment.

### Detailed comments

1. It is not clear to me why it is necessary to couple CTEM v2.0 with GEM-MACH-GHG for the purpose of carbon cycle data assimilation. CTEM v2.0 and GEM-MACH-GHG can run in parallel, and the GEM-MACH-GHG read in the fluxes from CTEM-v2.0 every 30 minutes. In that case, CTEM v2.0 can use the best possible meteorology fields it can get. As shown in this paper, the CTEM v2.0 forced by CRUNCEP performs better. I don't see the benefits of having consistent meteorology between CTEM v2.0 and GEM-MACH-GHG. The errors in meteorology affect CTEM v2.0 and GEM-MACH-GHG in quite different ways. The rationale discussed in the introduction is not convincing. I would recommend adding more discussions about the benefits of coupling these two models together. If the authors can give a specific example, it would be clearer.

We agree with the reviewer that much can be accomplished without coupling CTEM v2.0 with the GEM-MACH-GHG meteorology. Specifically, prescribed fluxes can be used for flux inversions. However, there are two important reasons why eventually this coupling must be done. First, we know that theoretically it is the same meteorology that drives both the weather and ecosystem models. Thus, climate models that simulate the carbon cycle include interactive ocean and terrestrial ecosystem models. Weather forecast models are also increasingly coupled atmosphere-land-ocean models. The carbon cycle involves coupled systems as well, and eventually, coupled models may be used for greenhouse gas simulations. Second and more importantly, we are developing an ensemble Kalman Filter (EnKF) for greenhouse gas state and flux estimation. The EnKF requires realizations of prior flux errors (among other things) to directly simulate sources of forecast uncertainty. By coupling CTEM 2.0 online with the atmospheric model, we can directly perturb uncertain parameters to get an ensemble of prior fluxes. In turn, the assimilation process will be able to provide more direct feedback to the CTEM developers. Without the data assimilation prospect, CTEM is usually evaluated against flux towers at selected sites. It is because of this constant feedback on CTEM that becomes possible with data assimilation that ECCC has chosen the approach of online coupling CTEM and GEM for eventual implementation. Thus, this work helps to pave the way for the coupled meteorological and ecosystem model within the EnKF. This rationale was not explained in the original manuscript so we have added it to the introduction in the revised version.

We have also added more discussion in the introduction about the importance of coupling bottom-up and weather models. For example, previous studies Lin et al., (2011) and Garnaud et al., (2014) (and many other, including the current paper) have shown that uncertainties in meteorological forcings (i.e. temperature, specific humidity, shortwave radiation) of ecosystem models contribute significantly to uncertainties in the simulated biospheric fluxes. At the same time, Miller et al., (2015) have shown that many of these meteorological variables (i.e. temperature, specific humidity, zonal wind, and planetary boundary layer) are also correlated with and contribute to biases in modeled atmospheric

transport. Therefore, coupling between TEMs and atmospheric transport models that account for meteorological uncertainty is considered an important step toward understanding how meteorological uncertainties impact both terrestrial CO<sub>2</sub> flux and transport errors.

Finally, we disagree that CTEM forced by CRU-NCEP performs better. A new figure (Taylor diagram in Figure 11) more clearly reveals that overall, CTEM forced by GEM generally better matches observations compared to CTEM forced by CRU-NCEP. However, the latter is indeed better in the tropics but that is where all prior (and posterior flux estimates) disagree and where observations are also inconsistent. These results can be explained by the fact that

1. The Northern Hemisphere, which is well observed, dominates the global seasonal cycle of carbon fluxes (GPP, Reco, and NEE) because it has the largest land areas that are mainly dominated by forest ecosystems. At the same time, the net contributions from the tropical and the Southern Hemisphere regions are close to zero due to their small and opposite seasonal cycles (Figures 8 and 9).
2. Figure 9 shows that even though there is a large difference in the amplitude of the seasonal cycle of GPP from CTEM-GEM compared to CTEM-CRUNCEP, the difference is much smaller in NEE, which is what is used in the inversion. Even though GPP have large biases compared to observation-based estimates, the biases in NEE are much smaller because NEE is the difference between two large terms (GPP and Reco).

So, given that CLASS-CTEM will provide only a prior estimate of NEE (at least in the first stage) for flux inversions in EC-CAS, we feel that CTEM forced by GEM is acceptable as a source of a priori fluxes for flux inversions.

2. It is not clear whether the energy fluxes (e.g., latent heat flux) and water fluxes (e.g., evaporation) required running GEM-MACH-GHG is from the CTEM v2.0 or from somewhere else. I would recommend adding descriptions whether the coupling is one-way or two-way.

This implementation is an offline coupling in that GEM is run first and the required meteorological variables are written to an auxiliary file at half-hourly time interval. These meteorological variables are then used as atmospheric forcing for CLASS-CTEM to simulate the land carbon fluxes which then used as a priori fluxes in GEM model. This is now stated explicitly (page 9, line 2-6) as well as in the introduction (page 3, lines 33-34).

3. The left panels and right panels in Figure 1 are basically the same. I would recommend either plotting only one year or averaging over two years. The same applies for Figures 3, 5, and 7.

In the revised version of the paper, we limited the plotting to the average of 2009-2010 for Figures 1, 3, 5 and 7 as suggested by the reviewer.

4. As discussed in the paper, precipitation from reanalysis product is not the best product available. I would recommend using Global Precipitation Climatology Project (GPCP) or CPC Merged Analysis of Precipitation (CMAP) precipitation as validation data set.

First, the main focus of this comparison is to assess how the GEM meteorology differs from CRU-NCEP since that is the default forcing that is used in CLASS-CTEM. This was needed to interpret the CTEM fluxes produced with the two different sources of meteorology. To assess these differences we compared to observation-based data from CRU. We agree that GPCP and CMAP could also have been used instead of CRU but any precipitation product has its own limitations (e.g. GPCP and CMAP are based on a combination of satellite and ground observations which are affected by issues of coverage, accuracy, etc.). In particular, the different observational datasets show the largest differences in the tropics (as noted in the original manuscript (revised manuscript page 12 and 13, lines 33-35, and 1-5, respectively)). It is therefore very challenging to justify which global precipitation product is better (particularly over the tropics) to use for comparison. Second, the point of comparing to ERAI is only to get a sense of how well a reputable reanalysis product compares to independent observations. Since our meteorological products are only analyses they are not expected to perform as well as reanalysis products like ERAI or MERRA or JRA-55. Thus if reanalyses also have difficulty in matching observations, this provides context or bounds for the kind of agreement we can expect from our analyses. However, we do not condone the use of reanalyses in place of observational datasets for the purpose of validation. This, we believe, is the Reviewer's concern and we share this concern. In the revised manuscript, we have clarified this point in section 2.5 (page 8, lines 5-8).

5. In Figure 7, the authors compared model simulated GPP to an up-scaled GPP product based on flux towers (B10). But the B10 data are over different periods. The FLUXCOM that is from the same research group as B10 has GPP product over 2009 and 2010. I would recommend comparing the model simulated GPP to the FLUXCOM GPP over the same time period.

As suggested by the reviewer, we replaced B10 by FLUXCOM GPP data and changed the text accordingly. However, we first compared GPP from FLUXCOM (mean of 2009-2010) and the climatological mean GPP from B10. The comparison showed that B10 and FLUXCOM are very similar in terms of the zonal mean (Figure R1) and the spatial distribution (Figure R2). Thus, the conclusions did not change by using FLUXCOM data instead of B10.




Figure R1: The zonal mean of GPP from FLUXCOM and B10.




Figure R2: The spatial difference between FLUXCOM and B10.

6. Figures 10 and 11 are not very informative. The differences shown in Figure 10 are a convolution of transport errors and the errors in the surface fluxes. Figure 11 uses the same transport model, so the differences are only due to surface fluxes, which have been discussed in Figure 9.

Actually, Figure 10 does not involve any convolution of transport errors since the same model (GEM-MACH-GHG) was used to integrate two different sets of prior fluxes (CTEM-GEM and CTEM-CRUNCEP). The CT2013B curve shown for comparison was directly taken from the mole fractions downloaded from NOAA's website. However, we agree that Figure 10 could be much more informative. Therefore, following the suggestion from the second reviewer to "consider using more sites for the comparison in

Figure 10”, we changed Figure 10 to compare the time series of the modeled and observed monthly CO<sub>2</sub> concentration at multiple sites. Also, we added a Taylor diagram (which will become figure 11) for a quantitative comparison between the modeled and observed CO<sub>2</sub> at the selected sites. We removed the figure that shows the zonal mean CO<sub>2</sub> (old Figure 11) as suggested by the Reviewer.

7. Tables 4 and 5 list the RMS and bias statistics of posterior CO<sub>2</sub> relative to independent CO<sub>2</sub> observations from TCCON and HIPPO campaigns. I would recommend adding one plot showing time series comparison between model simulated CO<sub>2</sub> and TCCON, and one latitudinal plot showing the comparison between model simulated CO<sub>2</sub> and HIPPO data, which may be more informative than the final statistics. I would also recommend adding a figure showing the comparison between posterior CO<sub>2</sub> and the CO<sub>2</sub> flask data assimilated in the flux inversion system, which will show how well the inversion system fits the assimilated data.

To assess the performance or tuning of the assimilation system, we normally look at the chi-squared diagnostic. This is now mentioned in the revised manuscript (section 4.3.3, page 20, lines 7-8). The chi-squared values are 0.913662, 0.892746, 0.756677 after assimilating 5365, 5393, and 5601 observations using CTEM-CRUNCEP, CTEM-GEM and BEPS, respectively. We also added histograms of modelled minus observed values over all assimilated observations (revised Figure 15). The fit of the three inversions to the assimilated observations is comparable. The standard deviations are similar and the biases are all small relative to the standard deviations. This is now mentioned in the revised text (page 20, lines 12-15).

Since we now have 15 figures in the manuscript, we are reluctant to also add the time series plots. However, we have included the time series comparisons to TCCON here for the benefit of the Reviewers and to assist with the review process. Sites at 3 different latitudes were chosen for the plots and only the middle of the 2-year study period was shown for clarity and to avoid error due to spin-up and spin-down effects. The magnitude and temporal variation of the errors for the 3 experiments are quite similar at the three sites. The largest differences are apparent at Bialystok with BEPS being more dissimilar to the two CTEM-based priors.





Figure R3: The temporal variation of the errors (model - obs) for 3 sites at different latitudes.



Reviewer 2 also suggested a latitudinal stratification of the HIPPO comparison. In particular, a stratification of Table 5 by latitude was requested. This was done in the revised Table 5. The text referring to Table 5 was modified accordingly (page 20, lines 19-24). We also include latitudinal plots of the HIPPO comparisons below, for the benefit of the review process, as requested.



**Figure R4:** Comparison of modelled and HIPPO observations as a function of latitude for observations below 5 km (top panel) and for those from 5-10 km (bottom panel).

8. The inversions use the 3D CO<sub>2</sub> fields from CarbonTracker as initial CO<sub>2</sub> fields in the inversion. This is risky since the transport models between TM5 used in the



CarbonTracker and GEOS-Chem is different. The initial fields works for the CarbonTracker does not necessarily the best for GEOS-Chem.

The Reviewer is correct to be concerned about the choice of initial state. However, there are two different issues associated with the initial conditions. The first is the issue of the global mean bias in the initial conditions. Any mean bias in the initial state will persist throughout the 2-year simulation length. In that regard, using the optimized 3D fields provides a means of mitigating this bias. The second issue is discrepancies in the spatial structure in the initial CO<sub>2</sub> field due to differences in transport between GEOS-Chem and TM5. On time scales of about three months, any intrahemispheric discrepancies in the initial state will be removed by large-scale mixing in the model, so on the time scale of the assimilation conducted here (2 years), these differences will not be important. Interhemispheric discrepancies, however, will matter. In our case, CarbonTracker was the best choice. Compared to the GEOS-Chem initial state, the CarbonTracker initial state actually produced less bias. This may suggest that the interhemispheric gradient in TM5 is better than in GEOS-Chem. The revised manuscript now includes this rationale for the choice of initial state (page 10, lines 31-33).

9. Some descriptions in the paper are not well justified by the results. For example, in the last paragraph in section 4.2.1: “CTEM-GEM flux estimates are within the range of the other estimates from TEMs used as a priori estimates in flux inversions (i.e., BEPS) or measurement-constrained fluxes (i.e., CT2013 B).”. This is not well justified since CTEM-GEM apparently has large differences with other fluxes over the tropics due to the bias in precipitation. This further reinforces my first point that it is not necessary to have consistent meteorology between CTEM and GEM to do carbon cycle data assimilation. It is important to have best meteorology.


While we agree with the referee that we should use the best meteorological forcings, it is challenging to identify the “best meteorology” data (precipitation, in particular) for all regions since measurements are incomplete or entirely lacking in some regions (tropics for example) [see references cited in section 4.2]. It is therefore challenging to identify the best flux estimates particularly over the tropics, where all models and observation-based estimates show large differences. Not surprisingly, CTEM has issues in the tropics, just as most other models do. This study gives us more specific insights about model deficiencies that need to be considered in the EC-CAS development processes. Moreover, even though CTEM-GEM shows large differences over the tropics, the Taylor diagram [revised Figure 11] shows that the modeled CO<sub>2</sub> using CTEM-GEM is generally relatively closer to the observed CO<sub>2</sub> compared to the one from CTEM-CRUNCEP (for both the forward and inversion simulations (revised Figure 11 and Figures R5, R6 below). Also Table A2 below shows that CTEM-GEM-based CO<sub>2</sub> fields have relatively smaller RMSEs at most sites than CTEM-CRUNCEP. This further supports our view that CTEM-GEM provides reasonable a priori-fluxes that are consistent with the atmospheric CO<sub>2</sub> signal.

**Table A1:** Locations of selected tropical stations measuring atmospheric CO<sub>2</sub> concentrations used in the inversion simulations.


Code	Name	Country	Latitude	Longitude
ASC	Ascension Island	United Kingdom	-7.967	-14.400
ASK	Assekrem	Algeria	23.262	5.632
BKT	Bukit Kototabang	Indonesia	-0.202	100.318
EIC	Easter Island	Chile	-27.160	-109.428
IZO	Izana, Tenerife, Canary Islands	Spain	28.309	-16.499
NMB	Gobabeb	Namibia	-23.580	15.030
RPB	Ragged Point	Barbados	13.165	-59.432
SEY	Mahe Island	Seychelles	-4.682	55.532

**Table A2:** The mean differences and RMSEs (in ppm) of the posterior CO<sub>2</sub> fields, based on CTEM-CRUNCEP, CTEM-GEM, and BEPS fluxes, with respect to surface observations at selected tropical sites (Table A1).

	Mean (mod – obs)			RMSE (mod – obs)		
	CTEM-CRUNCEP	CTEM-GEM	BEPS	CTEM-CRUNCEP	CTEM-GEM	BEPS
BKT	-0.08	0.86	0.15	2.70	2.96	3.14
NMB	0.34	0.19	0.30	1.67	1.24	1.24
ACS	0.05	0.12	0.09	0.97	0.83	0.85
ASK	-0.10	-0.07	-0.03	0.91	0.72	0.94
RPB	-0.28	-0.23	-0.18	0.91	0.68	0.91
IZO	0.41	0.34	0.39	1.67	1.41	1.67
SEY	-0.27	-0.16	-0.24	0.91	0.80	0.84
EIC2	0.42	0.49	0.47	0.93	0.94	0.95



**Figure R5.** Comparison of the posteriori modelled CO<sub>2</sub> using land the optimized fluxes from CTEM-CRUNCEP (blue circle), CTEM-GEM (blue diamond), and BEPS (green) with surface observations (red) at selected sites (Table A1). The modelled CO<sub>2</sub> produced by GEOS-Chem model.



**Figure R6:** Taylor diagram of modeled CO<sub>2</sub> produced by GEOS-Chem model using land fluxes from BEPS (Orange), GEM (Blue), and CRUNCEP (Green) and observed CO<sub>2</sub> (identified as "1.0") at selected sites (shapes) (listed in Table A1).

## Anonymous Referee #2

### General comments

This paper addresses the challenge of coupling a terrestrial ecosystem model to an NWP model that has been adapted to forecast CO<sub>2</sub> for the purpose of estimating CO<sub>2</sub> fluxes using a flux inversion analysis system. This coupling is required in order to provide boundary conditions to the forward CO<sub>2</sub> model and prior estimates for the flux inversion system. The main challenge stems from the differences in timescales required for the TEM to spin up and the short timescales used in NWP. The paper C1

presents an alternative configuration for the spin up suitable for the NWP model and compares the results with the standard spin up procedure. The results emphasize the large impact of meteorological biases on the biogenic CO<sub>2</sub> flux biases, in particular over the tropics. Although there is a small impact on the estimation of CO<sub>2</sub> growth rate when these fluxes are used as priors in flux inversion systems, there is a significant impact on the spatial distribution of the optimized fluxes, particularly in the tropics. All these aspects addressed are relevant scientific modelling questions within the scope of GMD, which are important to advance the use of earth system models to monitor the climate change.

The paper is well written and well structured. The methods used are valid and clearly outlined and the results are based on sound simulations with independent validation based on observations. However, the validation could be expanded to make better use of observations in the identification of regional biases both for the forward model and the optimized fluxes (see specific comments below). Test of statistical significance would also be highly recommended in order to strengthen the conclusions drawn from the results.

We are grateful to the referee for the careful review of our paper, and for the useful comments. We believe that the revised manuscript has greatly benefitted from the comments of both Reviewers. A point-by-point response to each comment is provided below.

### Specific Comments

1. Page 2, line 31: The sentence "Theoretically, in CCDAS, parameters of a TEM can also be optimized..." is a bit confusing. Isn't that what CCDAS aims to do?

In retrospect, we agree with the reviewer that the statement is confusing. Our original intention was to contrast the CCDAS approach from inversion systems (such as CarbonTracker or those based on GEOS-Chem) which optimize surface fluxes but not parameters of the ecosystem models. However, the intention was not realized. Therefore, we have changed this statement to the following "In CCDAS, key parameters of a TEM can also be optimized..."

2. Page 3, line 2: The limitations of CCDAS should also be mentioned (e.g. inability to correct for model structure and missing processes).

This is a good point. We added one sentence (section 1, Page 3, line 4-6) regarding the limitation of CCDAS including references.

3. Page 3, line 9: Shouldn't "reduced CO<sub>2</sub>" be "increased CO<sub>2</sub>"

Yes. We changed "reduced CO<sub>2</sub>" to "reduced CO<sub>2</sub> uptake" to be more consistent with the previous sentence.

4. Page 6, line 15: Please remove sentence "(e.g. Agusti-Panareda et al. (2016) also had similar issues)". The TEM used in Agusti-Panareda et al. (2016) doesn't have carbon stocks, so it does not require a spin up period.

The original intention was not to point out issues with the spin up period but rather the challenges of dealing with a constantly changing system. Agusti-Panareda et al. (2016) encountered this issue when trying to construct a consistent, long-term model climatology. However, the reviewer is correct that the statement as it stands implies something else. Since it is not worth trying to explain this reference, we removed the statement as suggested by the Reviewer.

5. Page 7, line 29: Note that the TCCON observations and HIPPO data in free troposphere are not ideal to assess impact of fluxes because their sensitivity to the surface fluxes is small compared to in situ surface stations. Why weren't surface stations used for validation?

Flask observations from surface stations were assimilated by GEOS-Chem in the inversion so they only provide a check on the consistency or set-up of the assimilation system. We still need to compare to independent observations that were not assimilated for validation. That is why we chose to focus on the TCCON and HIPPO comparisons and to not show the comparisons to surface data. The reason for focusing on TCCON and HIPPO is now added to the manuscript (page 20, lines 16-18).

Continuous surface observations are other sources of independent observations, but they provide limited insight (based on many years of experience) because of the coarse 4°×5° resolution used by GEOS-Chem inversions. Specifically, comparisons to continuous measurements reveal the ability to capture the synoptic, seasonal and longer timescales well. But this is also evident from comparisons to TCCON observations. It is not possible to capture the finer time scale variations with the coarse resolution we use. Again, this is already well known from our experience with the system. Thus, this observation set does not much help our specific goal here of studying the different prior fluxes in the context of our flux inversions.

Nevertheless, comparisons to the surface flask data that were assimilated are now better described. Chi-squared diagnostics are mentioned (page 20, lines 7-12) and histograms of the residuals between modelled and observed values are included in the revised manuscript (revised Figure 15). These comparisons serve to assess the setup of the inversion system and all 3 experiments fit the assimilated data about equally well.

6. Page 12, line 19: "the range of the other \*model\* estimat (Melton and Arora, 2014, Table 2)".

Changed

7. Page 15, line 2: Why are only 2 sites used in the evaluation of the forward model? This evaluation is not enough to draw conclusions on the impact of the coupled fluxes on the atmospheric CO<sub>2</sub> spatial/temporal variability at global scale. Please consider using more sites, if possible one site per TransCom region.

As suggested, we have replaced Figure 10 with a multi-panel plot of time series comparisons and added a Taylor diagram comparing the modeled and observed CO<sub>2</sub> at 9 sites as a revised Figure 11 that are representative of various global regions. Also, we removed the original Figure 11 to reduce the number of figures.

8. Page 15, line 22: I do not agree with this statement. The differences between red and blue lines can be substantial in summer and autumn as shown in Figure 10. Please also consider the use of more sites for the validation to make the results more robust in terms of spatial distribution (see previous comment).

In response to the previous point, we have replaced the original Figure 11 with a Taylor diagram comparing the modeled and observed CO<sub>2</sub> at multiple sites, and the text has been changed accordingly.

9. Page 15, line 23-24: Again, I don't think one can say that the forward runs with the CTEM and the posterior fluxes are similar in Figure 10 when the different are around 5 ppm both at both Alert and Mauna Loa during spring, summer and autumn.

We changed the whole text in section 4.2.2 to discuss the new figures (10 and 11) showing the monthly time series and the Taylor diagram of the modeled and observed CO<sub>2</sub> at multiple sites, in response to comment #8.

Nevertheless, we agree with the reviewer regarding the original Figure. However the new figures show that the general pattern of the seasonal cycle from CTEM simulations is fairly good in comparison to CT2013B, given that CTEM fluxes are not yet optimized. Interestingly, during the NH growing season, in particularly in 2009, CTEM-GEM is closer to the observation than both CT2013B and CTEM-CRUNCEP at ALT, BRW, IZO, MLO, and ZEP. Figure 9 shows that both CTEM-GEM and CTEM-CRUNCEP simulate higher CO<sub>2</sub> fluxes during winter compared to CT2013B, which can explain the enhanced CO<sub>2</sub> concentration during the winter season at the selected sites from the CTEM-based simulations. This also suggests that the model structure has some



deficiencies in simulating carbon fluxes during the winter season. This point has been added to the revised version (page 17, line 10-15; page 19, line 6-7; and page 22, lines 1-7).

10. Page 16, lines 10,11: There are significant differences in CTEM-based posterior estimates both in phase and amplitudes for N America, Europe and Asia.

We totally agree with the reviewer that there are significant differences in CTEM-based posterior estimates both in phase and amplitudes in NH regions. But, what was meant here is that the spread in the NH regions is much smaller than that in the tropical regions. We had already mentioned the large difference in Eurasia boreal. We have clarified this point in the revised version. Also, we bear in mind that CT2013 is not the truth and it is not well constrained over many regions of the globe, in particular the tropics, where measurements are incomplete or entirely lacking.

11. Page 16, line 24: A small flux increment does not necessarily mean a more accurate posterior estimate. In order to conclude that it is necessary to compare the resulting posterior estimate with independent observations of fluxes or compare posterior atmospheric CO<sub>2</sub> concentrations with independent observations at sites that are sensitive to the biogenic fluxes.

We meant here that the a posteriori from CTEM-CRUNCEP has been shifted to be almost close to the a priori CTEM-GEM which didn't change much. The comparison of the modeled and observed CO<sub>2</sub> for the forward simulation showed that CTEM-GEM is much closer to observation than CTEM-CRUNCEP, which might explain why the optimized fluxes from CTEM-GEM didn't change much from its a priori value. This point has been clarified in the revised version (see page 18, lines 28-33). We also compared the optimized CO<sub>2</sub> fields with independent data from TCCON and HIPPO and the results show that CTEM-GEM have a good agreement with these observations. In addition, in a subsequent analysis (Byrne et al., 2017) we conducted a more detailed evaluation of the CTEM-based a priori fluxes for NH ecosystems and found that CTEM-GEM does indeed provide a better simulation of atmospheric CO<sub>2</sub> data than CTEM-CRUNCEP. The Byrne et al. (2017) manuscript is in review in the Journal of Geophysical Research, but we now cite it in the revised manuscript in sections 4.2.1 and 4.3.1.

12. Page 17, line 26: Are these small differences in RMSE statistically significant?  
We thank the reviewer for highlighting this point. The differences in the RMSEs between CTEM-CRUNCEP and CTEM-GEM are not statistically significant (added in page 20, line 28-29). Our subsequent analysis in Byrne et al. (submitted, 2017) does indicate that the a priori CTEM-GEM provide a better simulation of variability in atmospheric CO<sub>2</sub>, but the statistics in this manuscript does not support the argument that the a posteriori fields based on CTEM-GEM are better. Furthermore, for the revised manuscript we restricted the TCCON comparison period to July 2009 to June 2010 and we found that the RMSEs are the same for CTEM-CRUNCEP and CTEM-GEM (see revised Table 5).

13. Page 18, lines 1,2 : These statement should be supported with a significance test of the error differences.

As noted above, it is not supported by the significance test so the statement was changed.

14. Page 18, line 26: ".. datasets of analyses are simply not possible to obtain" unless re-analysis datasets are used.

We changed it as suggested.

15. Page 19, lines 28-30: It is still not clear if the small differences in the error resulting from using CTEM-GEM or CTEM-CRUNCEP are statistically significant. Unless this is shown, this statement should not be used. Also, there is not explanation as to why one would expect the CTEM-GEM to provide a better constraint than CTEM-CRUNCEP for the inversion system.

Actually, based on the results of Byrne et al. (submitted, 2017) we would expect CTEM-GEM to be better. However, in the context of the analyses conducted here, we do not necessarily expect CTEM-GEM to provide a better constraint than CTEM-CRUNCEP for the inversion, given the far-from-ideal spinup process used for the former. The key points for our analysis here are (1) whether deficiencies in the CTEM-GEM prior fluxes would be evident in the context of a flux inversion when observations can correct for prior flux errors, and (2) how the deficiencies in CTEM-GEM prior fluxes compare to other prior fluxes. If they are no worse than other sources of prior fluxes, then they are a potential starting point for our flux estimation system. The differences, as noted by the reviewer, are small and are not statistically significant. We have, therefore, removed the statement. Instead, we acknowledge that CTEM-GEM can provide useful prior fluxes for flux inversions.

16. Page 19, line 31: This study provides insight into the deficiencies attributed to errors in the meteorological forcing (e.g. dry bias in precipitation over the tropics). It is not clear where is the insight into the deficiencies in the model.

Here we were referring to the deficiencies in simulating the seasonal cycle from CTEM noted in Page 19- lines 7-13. These deficiencies were uncovered in the course of our study and the authors of CTEM (Arora, Melton) ended up revising CTEM's formulation to address this issue. We tried to clarify this point in the revised version (page 17, line 10-15; page 19, line 6-7; and page 22, lines 1-7). We also mention the deficiencies in the model regarding the high CO<sub>2</sub> fluxes during the winter season (see the response to the comment #9).

17. Page 19, line 33: It is not clear how can the approach in this paper help improve the performance of CLASS-CTEM. Please provide an example.

As mentioned in the previous comment, the analysis shows some deficiencies in simulating the phase of the seasonal cycle as well as the winter fluxes which will form the basis for future development of CLASS-CTEM. The developers of CLASS-CTEM (Arora, Melton) are co-authors in this paper so they are aware about this issue and are

presently working on solving it. However, we agree that this detail was not evident in the original manuscript. This has now been clarified in the revised version (page 17, line 10-15; page 19, line 6-7; and page 22, lines 1-7), and we revised the relevant conclusion statement.

18. Page 21, line 6: Please update the reference.

Done.

19. Figure 8 shows a larger impact from meteorological forcing than from model formulation in NH summer, tropics and SH. This message should be emphasized in the paper.

This is a very good point and we are grateful to the Reviewer for this insight. We now explicitly mention that the impact of meteorological forcings can exceed that due to model formulation. We also note that we demonstrate that using different meteorological forcings result in the same biases in the winter season, indicating there is a problem in the model phenology as well. We now mention this in different places in the paper and have added it to the conclusions as well.

20. Figure 13 shows a large difference of NEE budget in Europe between CTEM-GEM and CTEM-CRUNCEP. This is not mentioned in the paper.

We will mention to this in the revised paper (page 19, lines 19-20). But, basically there are large differences in all regions, which highlight the sensitivity of the inversion to the a priori fluxes (as noted in the paper).

21. Table 4: Please show station to station error (e.g. std of station bias) in order to evaluate the spatial variability of posterior CO<sub>2</sub>. This is commonly done in evaluation of CO<sub>2</sub> satellite data. The ability of reproducing the global mean does not reflect the impact that the prior has on the posterior regional patterns.

Table 4 has now been revised to include station to station error and the text had been modified to mention this.

22. Table 5: The error with respect to the HIPPO data could be stratified in latitude bands in order to evaluate the interhemispheric gradient.

Table 5 was modified as suggested and the revised manuscript now discusses the interhemispheric gradient.

# Coupling the Canadian Terrestrial Ecosystem Model (CTEM v. 2.0) to Environment and Climate Change Canada's greenhouse gas forecast model (v.107-glb)

Bakr Badawy<sup>1,\*</sup>, Saroja Polavarapu<sup>1</sup>, Dylan B. A. Jones<sup>2</sup>, Feng Deng<sup>2</sup>, Michael Neish<sup>1</sup>, Joe R. Melton<sup>3</sup>, Ray Nassar<sup>1</sup>, and Vivek K. Arora<sup>3</sup>

<sup>1</sup>Climate Research Division, Environment and Climate Change Canada, Toronto, Canada

<sup>2</sup>Department of Physics, University of Toronto, Toronto, Canada

<sup>3</sup>Climate Research Division, Environment and Climate Change Canada, Victoria, Canada

\*Now at Faculty of Environment, University of Waterloo, Canada

Correspondence to: Bakr Badawy(bbadawy@uwaterloo.ca)

**Abstract.** The Canadian Land Surface Scheme and the Canadian Terrestrial Ecosystem Model (CLASS-CTEM) together form the land surface component in the family of Canadian Earth System Models (CanESM). Here, CLASS-CTEM is coupled to Environment and Climate Change Canada (ECCC)'s weather and greenhouse gas forecast model (GEM-MACH-GHG) to consistently model atmosphere-land exchange of CO<sub>2</sub>. The coupling between the land and the atmospheric transport model ensures consistency between meteorological forcing of CO<sub>2</sub> fluxes and CO<sub>2</sub> transport. The procedure used to spin up carbon pools for CLASS-CTEM for multi-decadal simulations needed to be significantly altered to deal with the limited availability of consistent meteorological information from a constantly changing operational environment in the GEM-MACH-GHG model. Despite the limitations in the spin up procedure, the simulated fluxes obtained by driving the CLASS-CTEM model with meteorological forcing from GEM-MACH-GHG were comparable to those obtained from CLASS-CTEM when it is driven with standard meteorological forcing (CRU-NCEP). This is due to the similarity of the two meteorological datasets in terms of temperature and radiation. However notable discrepancies in the seasonal variation and spatial patterns of precipitation estimates, especially in the tropics, were reflected in the estimated carbon fluxes, as they significantly affected the magnitude of the vegetation productivity and, to a lesser extent, the seasonal variations in carbon fluxes. Nevertheless, the simulated fluxes based on the meteorological forcing from the GEM-MACH-GHG model are ~~within the range of~~ consistent to some extent with other estimates from bottom-up or top-down approaches. Indeed, when simulated fluxes obtained by driving the CLASS-CTEM model with meteorological data from the GEM-MACH-GHG model are used as prior estimates for an atmospheric CO<sub>2</sub> inversion analysis using the adjoint of the GEOS-Chem model, the retrieved CO<sub>2</sub> flux estimates are comparable to those obtained from other systems in terms of the global budget and the total flux estimates for the northern extratropical regions, which have good observational coverage. In data poor regions, as expected, differences in the retrieved fluxes due to the prior fluxes become apparent, ~~but fall within the uncertainty bounds based on multi-inversion analyses. The coupling of~~ Coupling CLASS-CTEM ~~to an atmospheric transport model with carbon assimilation capabilities also provides insights into the limitations of CLASS-CTEM simulated~~ into EC-CAS is considered an important step toward understanding how

~~meteorological uncertainties affect both CO<sub>2</sub> fluxes through comparisons of simulated atmospheric CO<sub>2</sub> with observations at selected flask stations. This capability can be used to continually assess and improve flux estimates and modelled atmospheric transport. Ultimately, such an approach will provide more direct feedback to the terrestrial ecosystem modules of the CLASS-CTEM model developers, and thus help to improve the performance of CLASS-CTEM by identifying the model limitations based on atmospheric constraints.~~

*Copyright statement.* TEXT

## 1 Introduction

Terrestrial ecosystems play a crucial role in the global climate-carbon system. Therefore, there is a need to better understand terrestrial biospheric processes related to the carbon cycle in order to obtain more reliable projections of their behavior under a changing climate. Given the great heterogeneity of vegetation and soils, the coverage and accuracy of the flux measurements are not sufficient for obtaining large-scale flux estimates with high confidence (Jung et al., 2009; Beer et al., 2010). As a result, considerable efforts have been made to develop terrestrial ecosystem models (TEMs) (whether simple regression or process-oriented) in order to quantify the magnitude, geographical distribution, and evolution of sources and sinks of carbon at regional and global scales (Potter et al., 1993; McGuire et al., 2001; Sitch et al., 2003; Thornton et al., 2005; Krinner et al., 2005; Reichstein et al., 2005; Badawy et al., 2013; Arora and Boer, 2005; Melton and Arora, 2016). However, systematic errors and uncertainties in the models can result from driving or forcing data (Jung et al., 2007; Clein et al., 2007; Zhao et al., 2006; Garnaudo et al., 2014; Dalmonech et al., 2015; Anav et al., 2015; Wei et al., 2014), process formulation (also called model structure) (Sitch et al., 2015), model parameter specification, and initial conditions (Carvalhais et al., 2008, 2010; Melton et al., 2015; Zhu and Zhuang, 2015), leading to differing estimates of CO<sub>2</sub> fluxes from different models (McGuire et al., 2001; Piao et al., 2013; Sitch et al., 2015). Such differences in TEMs are among the main sources of uncertainty in future projections from coupled carbon-climate models (Anav et al., 2013; Friedlingstein et al., 2006; Arora et al., 2013; Friedlingstein et al., 2014). Therefore, there is a need to evaluate the performance of TEMs in order to identify and diagnose their weaknesses and strengths and ultimately reduce model uncertainties. Indeed, this is the motivation behind TEM multimodel intercomparisons efforts such as the Multi-scale Synthesis and Terrestrial Ecosystem Model Intercomparison Project (MsTMIP) (Huntzinger et al., 2013).

Inverse models (which relate observed concentrations to fluxes using an atmospheric transport model) are powerful tools to quantify carbon fluxes over large regions (Rödenbeck et al., 2003; Peters et al., 2007; Peylin et al., 2013) and can be used to evaluate the TEM results. However, inverse models suffer from deficiencies and uncertainties (Peylin et al., 2013) arising from transport errors, choice of observation network, observation uncertainties, and prior flux errors. Inverse models also provide little or no information about the underlying processes responsible for the estimated fluxes. Hence, they cannot be used to understand or predict the future behavior of the carbon cycle. Alternatively, there are carbon cycle data assimilation

systems (CCDAS), which couple the strengths of the top-down (inversion) and bottom-up (i.e. TEM) approaches by embedding a TEM within a comprehensive climate model and using measurements from multiple streams to constrain the TEM (Scholze et al., 2003; Rayner et al., 2005; Koffi et al., 2013). The benefit is that biospheric models can then be validated on the global scale using atmospheric measurements of CO<sub>2</sub> that integrate the CO<sub>2</sub> signal at various spatial and temporal scales. ~~Theoretically, in CCDAS, In CCDAS, key~~ parameters of a TEM can also be optimized to improve its fit to atmospheric CO<sub>2</sub> observations (Scholze et al., 2003; Rayner et al., 2005; Koffi et al., 2013; Kaminski et al., 2013), which can potentially yield greater understanding about underlying processes, and thus can help to improve the model performance. Nevertheless, CCDAS is sometimes challenging and has its limitations. For example, the optimized fluxes (and parameters) are sensitive to CCDAS configurations (Kaminski et al., 2013) (i.e. atmospheric transport, background fluxes, observational network, processes representations, missing process, etc).

Comprehensive Earth System Models need to include TEMs because the ecosystem responds to a changing climate. However, weather and carbon fluxes are also interconnected so that coupled weather and CO<sub>2</sub> prediction models operating on weather or seasonal timescales can also benefit from online TEMs. ~~Specifically, if such a coupled model uses a prior flux estimate of terrestrial biosphere activity from an offline TEM,~~ Previous studies (Lin et al., 2011; Garnaud et al., 2014) have shown that uncertainties in meteorological forcings (i.e. temperature, specific humidity, shortwave radiation) contribute significantly to uncertainties in the simulated fluxes. In addition, Miller et al. (2015) have shown that several meteorological variables (i.e. temperature, specific humidity, zonal wind, and planetary boundary layer) are correlated and contribute to biases in modelled atmospheric transport. Therefore, inconsistencies may arise when the TEM's meteorological driving data differs from that in the weather model. For example, at a given point in time, a TEM's grid cell might have experienced sunny weather and thus produced large CO<sub>2</sub> uptake whereas the weather model may indicate cloudy conditions and reduced CO<sub>2</sub> uptake. An online TEM constrained by the model's weather would have predicted this reduced CO<sub>2</sub> uptake. If such inconsistent CO<sub>2</sub> predictions are used to constrain inverse models there is a risk of misattributing some of the model-data mismatch to the flux estimate. Therefore, coupling between TEMs and atmospheric transport models (i.e. using the same meteorological variables) is considered an important step toward understanding how meteorological uncertainties impact both CO<sub>2</sub> flux estimates and modelled atmospheric transport. Forecasting systems that integrate land and ocean CO<sub>2</sub> fluxes within numerical weather prediction (NWP) models have recently been developed (Agusti-Panareda et al., 2014; Ott et al., 2015) to produce short term predictions of atmospheric CO<sub>2</sub>.

At Environment and Climate Change Canada (ECCC), a Carbon Assimilation System (EC-CAS) (Polavarapu et al., 2016) is being developed ~~to~~ by implementing an ensemble Kalman Filter (EnKF) for greenhouse gas state and flux estimation. EC-CAS will assimilate satellite and in situ data to generate hindcasts of atmospheric CO<sub>2</sub> and estimates of regional fluxes of CO<sub>2</sub>. ~~A-EC-CAS uses an ensemble of coupled meteorological and constituent forecasts to directly compute the complex transport error covariances needed for proper flux estimation (Miller et al., 2015). The meteorological model is an adaptation of the operational weather prediction model GEM-MACH (Global Environmental Multi-scale - Modelling Air quality and CHemistry) (Moran et al., 2010; Robichaud and Ménard, 2014; Makar et al., 2015) for greenhouse gasses (called GEM-MACH-GHG hereafter, see Section 2.2).~~ In addition to an ensemble of meteorological fields, EC-CAS also requires an ensemble of prior

fluxes to simulate prior flux uncertainty. With an online ecosystem model, we can directly perturb parameters to get this ensemble. In turn, the assimilation process would provide continual feedback on the ecosystem model, just as in CCDAS. For this reason, we envision an online ecosystem model eventually within EC-CAS. However, we first consider the simpler approach of offline coupling wherein the atmospheric model's meteorology is used to drive the ecosystem model. Thus, a key objective of the work here is to assess the viability of land surface fluxes of CO<sub>2</sub> from the Canadian Terrestrial Ecosystem Model (CTEM) (Melton and Arora, 2014), coupled to the Canadian Land Surface Scheme (CLASS) (Verseghy, 2012), as a source of a priori biospheric fluxes of CO<sub>2</sub> for EC-CAS and other CO<sub>2</sub> flux inversion systems. ~~EC-CAS is being created by adapting the operational weather prediction model GEM-MACH (Global Environmental Multi-scale—Modelling Air quality and CHemistry) (Moran et al., 2010; Robichaud and Ménard, 2014; Makar et al., 2015) for greenhouse gasses (called GEM-MACH-GHG hereafter, see Section 2.2).~~ CLASS-CTEM is a process-based TEM which simulates the exchange of carbon, water and energy fluxes between the land surface and the atmosphere. It is similar in level of complexity to other TEMs (such as CASA (Potter et al., 1993) or SiB (Sellers et al., 1996)) which have been used for flux inversions and which have participated in multimodel intercomparisons such as that of Huntzinger et al. (2012). In recent studies (Melton and Arora, 2014; Melton et al., 2015; Melton and Arora, 2016; Badawy et al., 2016), CLASS-CTEM was calibrated based on observation-based climate data from the Climate Research Unit (CRU) (Harris et al., 2014) combined with reanalysis fields from the National Centers for Environmental Prediction (NCEP) (Kalnay et al., 1996). By coupling CLASS-CTEM with the atmospheric model, this work helps to pave the way for the coupled meteorological and ecosystem model within the EnKF (e.g. see conclusions of Miller et al. (2015)).

Although incorporating CLASS-CTEM within EC-CAS is potentially mutually beneficial, the incorporation of a TEM designed for Earth System Modelling (decadal timescales) into a data assimilation system designed for short timescales (i.e. months to a few years) is not without its challenges. For example, the spin-up of carbon pools to present climate needs to be merged with the switch in climate data from reanalyses to that from the weather forecasting model (e.g. EC-CAS). The challenge is that operational weather forecasting systems are, by definition, constantly changing so that long archives of consistent analyses (i.e. with the same horizontal or vertical resolution or model coordinates or variable, etc.) are not available (see also ~~Agusti-Panareda et al. (2016)~~ Agusti-Panareda et al. (2017) for example), contrary to the case of reanalyses (e.g. ERA-Interim (Dee et al., 2011) or MERRA (Rienecker et al., 2011)). Given that the spin-up procedure is known to impact TEM predictions (Wutzler and Reichstein, 2007; Carvalhais et al., 2008, 2010), how will this affect the use of CLASS-CTEM in the flux estimation context? In addition, the environmental drivers of TEMs also impact their results, so will the change in the climate forcing of CLASS-CTEM negatively impact its predictions on these short "climate timescales"? Garnaud et al. (2014) show that carbon pools and fluxes from CLASS-CTEM are sensitive to climate datasets for the case of a limited area domain (North America). On the other hand, large changes in fluxes are not necessarily detectable by observing systems such as Greenhouse Gases Observing SATellite (GOSAT) (Ott et al., 2015) so such differences may not be perceptible in flux inversion results. Finally, given that there is already a well-documented sensitivity to prior flux estimates in data sparse regions (Gurney et al., 2004; Peylin et al., 2013), do such deficiencies in spin-up procedure and environmental drivers matter? In other words, despite the unavoidable imperfections in coupling a TEM from an Earth System Model to a Carbon Assimilation System focussed on



short climate timescales, will the flux inversion results obtained using CTEM fall within the range of uncertainty encompassed by an ensemble of recognized flux inversion systems? The goal of this work is to answer these questions.

We begin in Section 2 with a description of the various models and datasets involved in this study, followed by the experimental design (Section 3). In order to interpret differences in fluxes resulting from the change in meteorological forcing, we first compare the quality of the meteorological inputs from GEM-MACH-GHG against the standard climate forcing (CRU-NCEP) that was used to drive CLASS-CTEM, as well as against independent sources of data (Section 4.1). Then, we examine the sensitivity of the simulated carbon fluxes to the change in meteorological forcing to determine whether biases in the simulated carbon fluxes can be attributed to biases in the meteorological variables (Section 4.2). The simulated fluxes are assessed both directly as well as indirectly through their impact on CO<sub>2</sub> concentrations. Finally, in Section 4.3, the a priori fluxes from CTEM are used in a flux inversion system and the results are analyzed in terms of the seasonal cycle and annual totals of the optimized fluxes and the a posteriori CO<sub>2</sub> concentrations. The conclusions are presented in Section 5.

## 2 Models and Data

Before presenting the experimental design, we first introduce the TEM and the coupled meteorological and tracer transport model to which the TEM will be coupled. Then, ~~CRU-NCEP and other~~ the validation datasets used to assess the various sources of climate forcing are described, followed by the experimental methodology.

### 2.1 CLASS-CTEM

The coupled CLASS-CTEM model used here is based on CLASS v3.6 (Versegny, 2012) and an updated version of CTEM v1.2 (Melton and Arora, 2014) and runs globally on a Gaussian 128×64 grid that corresponds to  $\sim 2.8^\circ \times 2.8^\circ$  grid spacing. CLASS calculates the biophysical exchange of energy and water fluxes between the land surface (soil, snow, and vegetation canopy) and the atmosphere. The model includes three soil layers, which extend to a total depth of 4.1 m, and one vegetation canopy and one snow layer. The model solves for the energy and hydrological balances at each grid cell using a half-hourly time step. The land surface of each grid cell is divided into four subareas: bare soil, vegetation, snow over bare soil and snow with vegetation. The vegetation within a grid cell, in CLASS, can be composed of 4 PFTs (Plant Functional Types): needleleaf trees, broadleaf trees, crops and grasses. For each PFT, prescribed physiological characteristics, such as albedo, annual maximum and minimum leaf area index (LAI), vegetation height, canopy mass, and rooting depth have to be specified. When coupled to CTEM, these structural vegetation attributes are dynamically simulated by CTEM with a daily time step and then passed to CLASS.

CTEM is a process-based terrestrial biosphere model that grows vegetation from bare ground and simulates the main processes governing carbon fluxes between the land biosphere and atmosphere. The model is parametrized and designed to simulate land-atmosphere exchanges of carbon through photosynthesis, ecosystem respiration (sum of autotrophic and heterotrophic respiration), phenology, turnover, mortality, allocation, fire and land use change (Arora, 2003; Arora and Boer, 2005; Melton and Arora, 2016). The model is represented by three living vegetation pools (leaves, stems, and roots) and two dead carbon

pools (soil organic matter and litter). The terrestrial ecosystem processes are calculated for nine PFTs: Needleleaf evergreen, Needleleaf deciduous, broadleaf evergreen, broadleaf cold deciduous, broadleaf drought/dry deciduous, crops ( $C_3$  and  $C_4$ ) and grasses ( $C_3$  and  $C_4$ ). When coupled, CTEM provides time-varying vegetation structure attributes to CLASS and the calculated variables for the nine PFTs are averaged (weighted by the fractional coverage of each PFT) to obtain the four PFTs in CLASS that share similar functionality.

Within CTEM, photosynthesis and leaf respiration sub-modules operate on a half-hourly time step as in CLASS in order to model the effect of the  $CO_2$  concentration on stomatal conductance. Other terrestrial ecosystem processes, including stem, root, and heterotrophic respiration are modelled at a daily time step. Recently, Badawy et al. (2016) modified CTEM to add the capability to simulate all respiratory fluxes at the same time step as CLASS (i.e. half-hourly) in order to model their diurnal variation caused by subdiurnal signals in the driving climate data. The current version of CTEM does not include the nitrogen cycle and its interactions with carbon cycle. Nevertheless, the model constrains the response of terrestrial photosynthesis to elevated  $CO_2$  via an empirical formulation based on experimental plant growth studies (Arora et al., 2009). The model structure and its parametrizations are documented in Arora (2003), Arora and Boer (2005), and Melton and Arora (2016), in which a comprehensive description of model subroutines is provided.

Besides the meteorological inputs (shortwave and longwave downward radiation, air temperature, precipitation, specific humidity, surface pressure, wind speed (see Section 2.4)), the model requires data on soil texture (i.e. percentage of sand and clay for the three soil layers), fractional vegetation coverage for each PFT, organic matter content, permeable soil depth, and atmospheric  $CO_2$ . The soil texture information is based on Zobler (1986). The vegetation fractional coverage for the nine PFTs in CTEM are adapted from Arora and Boer (2010) but using the HYDE v3.1 data set for crop area (Hurt et al., 2011) to reconstruct the historical land cover. The model uses inputs of annual mean atmospheric  $CO_2$  concentrations, which are based on phase 5 of the Coupled model Intercomparison Project (CMIP5) (Meinshausen et al., 2011).

## 2.2 GEM-MACH-GHG

GEM-MACH is based on the dynamics and physics of the Global Environmental Multiscale (GEM) model (Côté et al., 1998a; Girard et al., 2013) at the Canadian Meteorological Centre (CMC). GEM is used for operational weather forecasting in both global and regional (North America) domains, whereas GEM-MACH includes an online chemical model that is fully integrated into the meteorological model to provide air quality forecasts over North America. GEM-MACH-GHG (v.107-glb) (Polavarapu et al., 2016) is a variant of GEM-MACH that removes the reactive chemistry and replaces it with climate-chemistry (e.g. OH climatology). In addition, a number of modifications to GEM-MACH were made, including the implementation of a mass conservation scheme, and modifying the vertical mixing in the boundary layer. A horizontal resolution of  $0.9^\circ$  ( $400 \times 200$  grid points), and a time step of 30 minutes are used.

In this study, the meteorological fields required to drive CLASS-CTEM are produced from GEM-MACH-GHG (v.107-glb) following the same approach as in Polavarapu et al. (2016) for the 2009-2010 period. Prior to 22 June 2009, the operational analyses were produced using a model with a lid at 10 hPa. Since that date, the operational model has used a much higher lid of 0.1 hPa and since the period of interest for greenhouse gas simulations commences with the launch of the Greenhouse

Gases Observing Satellite (GOSAT) (Kuze et al., 2009; Yokota et al., 2009) in 2009, GEM-MACH-GHG uses the more recent model configuration. As a result, it is difficult to make use of CMC analyses prior to 22 June 2009. Thus, early in 2009, these analyses were supplemented by CMC archives of the "parallel run" (the system during its testing phase) and a preliminary run. Given that GEM-MACH-GHG was under development during this study, only a few years were simulated (2009-2010). There will always be an unsatisfactory length of analyses available for a TEM spin up period whenever operational weather forecast system is involved (e.g. Agusti-Panareda et al. (2016) also had similar issues). Moreover, greenhouse gas assimilation systems are constrained (by time, computational expense and the observing system) and thus often focus on a few years of study at one time (e.g Deng et al. (2014, 2016)). Thus, the challenge is to merge this small dataset into the spin-up procedure used for the TEM. As we shall see, despite this considerable challenge, the resulting impact on net fluxes are still within the bounds of uncertainty provided by an ensemble of TEMs comparable to other observation-based estimates and model simulations. The meteorological fields are initialized at the start of each 24h cycle with archived analyses from the CMC which were produced by the previously operational four-dimensional variational (4D-Var) data assimilation system (Charron et al., 2012), interpolated to GEM-MACH-GHG's 0.9° resolution. The 24-hour forecasts of shortwave and longwave radiation, surface temperature, wind speed, surface pressure, total precipitation, and specific humidity, were generated every 30 minutes, and then interpolated to the CLASS-CTEM grid.

### 2.3 GEOS-Chem

Previous inversion studies show that optimized fluxes are sensitive to prior fluxes particularly for regions that are poorly constrained by atmospheric observations such as the tropics (Peylin et al., 2013). In order to assess the quality of NEE from CTEM-GEM in comparison to other flux estimates, it is necessary to perform some inversion studies. Ideally, such inversions would be conducted with GEM-MACH-GHG but since the assimilation capability of EC-CAS is still under development, an alternative inversion system based on the GEOS-Chem model (<http://geos-chem.org>) is used. The GEOS-Chem model has often been used to simulate atmospheric CO<sub>2</sub> (e.g. Suntharalingam et al. (2004); Nassar et al. (2010)). This model is a global 3-D chemical transport model driven by assimilated meteorology from the Goddard Earth Observing System (GEOS-5) of the NASA Global Modelling and Assimilation Office (GMAO). Nassar et al. (2010) described an update of the atmospheric CO<sub>2</sub> simulation in GEOS-Chem. In this study, the model has a horizontal resolution of 4° × 5°, with 47 vertical layers from the surface to 0.01 hPa. The assimilation system is a 4D-Var data assimilation system in which a set of scaling factors is optimized to adjust the fluxes in each model grid box to better reproduce the observations over a given time period. In the 4D-Var system, the adjoint of the GEOS-Chem model is used to optimize the fluxes. Details of the GEOS-Chem adjoint model are given in Henze et al. (2007) and a description of its application for inverse modeling-modelling of atmospheric CO<sub>2</sub> is provided in Deng et al. (2014, 2016).

### 2.4 CRU-NCEP

The observation-based 0.5° monthly climatology from the Climate Research Unit (CRU, version TS3.2) (Harris et al., 2014) and the ~ 2.5°, 6-hourly reanalysis fields from the National Centers for Environmental Prediction (NCEP) (Kalnay et al., 1996)

were combined to produce the CRU-NCEP global climate data set (Viovy, 2016) that has been described in Wei et al. (2014). The CRU-NCEP dataset provides globally gridded ( $0.5^\circ \times 0.5^\circ$ ) 6-hourly time-varying climatology products that covers the period 1901-2014. The input data from CRU-NCEP includes shortwave and longwave radiation, surface temperature, wind speed, surface pressure, total precipitation, and specific humidity. These climate data were interpolated to the CLASS-CTEM's  
5 grid and disaggregated to a half-hourly time step as described in Arora and Boer (2005), and Melton and Arora (2014).

## 2.5 Other Datasets

To evaluate the quality of the GEM driving data, the forecasted fields of shortwave radiation, temperature, and precipitation for 2009 and 2010 are compared with CRU-NCEP, and both are evaluated against the CRU dataset and the ERA-Interim reanalysis (hereafter called ERAI) of the European Centre for Medium-Range Weather Forecasts (ECMWF) (Berrisford et al., 2011; Dee  
10 et al., 2011). The  $2.5^\circ$  monthly ERAI data is available at the ECMWF data server. Comparing to ERAI is done only to get a sense of how well a reputable reanalysis product compares to independent observations. Since the GEM products are only analyses they are not expected to perform as well as reanalysis products like ERAI. Thus, if reanalyses also have difficulty in matching observations, this provides context or bounds for the kind of agreement we can expect from our analyses.

To assess the impact of using alternative driving data on the simulated fluxes, the CLASS-CTEM fluxes obtained with  
15 GEM and CRU-NCEP meteorology are compared and evaluated against independent observation-based flux estimates and other model results. For example, the simulated GPP was compared with the ~~observation-based estimates of gross primary productivity (GPP) produced by Beer et al. (2010) (called B10 hereafter). They are based on eddy covariance flux data and several data-driven models, and averaged for the period 1998 to 2005.~~ upscaled GPP from FLUXCOM (Jung et al., 2017). The FLUXCOM GPP data is based on machine learning methods that integrate in situ flux measurements, satellite-based  
20 vegetation indices, and meteorological data (Tramontana et al., 2016; Jung et al., 2017). In this study, we used the ensemble mean of GPP estimates (for 2009-2010) generated using multivariate regression splines (MARS), random forests (RF), and artificial neural networks (ANN) available through the Data Portal of the Max Planck Institute for Biogeochemistry (https://www.bgc-jena.mpg.de). The model results are also compared with the multi-year average 3-hourly GPP and ecosystem respiration ( $R_{eco}$ ) from the Boreal Ecosystem Productivity Simulator (BEPS) (Chen et al., 2012), ~~in which~~ which is normally  
25 used by GEOS-Chem in flux inversions (Deng et al., 2014, 2016). In BEPS, the annual terrestrial ecosystem exchange imposed in each grid box ( $4^\circ \times 5^\circ$ ) is neutral (Deng and Chen, 2011) (i.e.  $GPP = R_{eco}$ ). BEPS is driven by NCEP reanalysis dataset. Finally, the model results are evaluated using the a posteriori  $CO_2$  fluxes from the CarbonTracker data assimilation system (Peters et al., 2007) (version CT2013B) available at <http://carbontracker.noaa.gov>. All datasets used in evaluating the model's results are re-gridded to the CLASS-CTEM grid.

30 We evaluate the results of the inversion analyses (described in section 3.3) using the GEOS-Chem model by comparing the a posteriori  $CO_2$  fields to atmospheric  $CO_2$  observations from the Total Carbon Column Observing Network (TCCON) from which the column-averaged dry-air mole fractions of  $CO_2$  ( $XCO_2$ ) are retrieved (Wunch et al., 2011). TCCON data were obtained from the TCCON Data Archive, hosted by the Carbon Dioxide Information Analysis Center (CDIAC) (<http://tcccon.ornl.gov/>). For the comparisons, we use observations from the current TCCON GGG2014 data set from 13 different

sites (Table 2) (see also Deng et al. (2014)) in 2009 and 2010. We also evaluate the inversion analyses using aircraft data from the HIAPER Pole-to-Pole Observations (HIPPO) project (<http://hippo.ornl.gov/>). We use the 10-second averaged data from the HIPPO-1, HIPPO-2, HIPPO-3 campaigns (Wofsy, 2011; Wofsy et al., 2012), for 9 to 21 January 2009, 31 October to 22 November 2009, and 24 March to 16 April 2010, respectively.

### 5 3 Experimental Design

In this study, we performed an offline coupling between CLASS-CTEM and GEM-MACH-GHG. We first run GEM-MACH-GHG to produce the necessary meteorological variables required to drive CLASS-CTEM. Then, the simulated fluxes from CLASS-CTEM are used in GEM-MACH-GHG to simulate the CO<sub>2</sub> concentrations. Given the complexity of online coupling (which is harder to implement), the offline simulations provide an affordable means to better isolate and assess the sensitivity of the model to different climate forcings. Offline coupling is also a step toward online coupling.

When coupling CLASS-CTEM to ~~EC-CAS~~GEM-MACH-GHG, we first identify a necessarily-imperfect spin-up procedure that transitions from climate data forcing from a standard dataset such as CRU-NCEP to a short sequence of operational meteorological analyses (Section 3.1). Once fluxes are available from CLASS-CTEM for CRU-NCEP meteorology with the standard spin-up procedure and from GEM-MACH-GHG with the modified spin-up procedure, the simulations of CO<sub>2</sub> that are performed with GEM-MACH-GHG are described in Section 3.2. Finally, the a priori fluxes from CLASS-CTEM are tested in a flux inversion experiment which is described in Section 3.3.

These different simulations will assess whether the deficiencies of CTEM-GEM prior fluxes would be evident in the context of flux inversion when observations can correct for prior flux errors, and how the deficiencies in CTEM-GEM prior fluxes compare to other prior fluxes. If they are consistent or no worse than other sources of prior fluxes, then they are a potential starting point for our flux estimation system.

#### 3.1 CLASS-CTEM Runs

To test the sensitivity of the simulated carbon fluxes to the meteorological forcing, we performed a series of experiments with CLASS-CTEM using two different meteorological inputs from (1) CRU-NCEP (hereafter called CTEM-CRUNCEP), which has been used to drive CLASS-CTEM simulations in previous studies (Melton and Arora, 2014; Melton et al., 2015; Badawy et al., 2016), and (2) GEM-MACH-GHG (hereafter called CTEM-GEM). For the CTEM-CRUNCEP run, the model was first initialized (to represent the pre-industrial period 1861-1900) by running it to equilibrium using repeated 1901-1940 CRU-NCEP climate, a constant globally uniform CO<sub>2</sub> of 286.37 ppm, and a fixed vegetation fractional coverage corresponding to the year 1861 until carbon pools and fluxes were in steady state (zero mean annual net ecosystem exchange (NEE)). The model was then run from 1901-2010 using varying CO<sub>2</sub> concentrations and CRU-NCEP meteorology.

For the CTEM-GEM run, the meteorological inputs from GEM-MACH-GHG were only available for 2009-2010 at the time of this study, and hence no global climate data available for the pre-industrial run. In general, reanalysis output begins around 1949 (e.g. NCEP-NCAR reanalyses) when the observing system had sufficient coverage, and as noted earlier, analyses from

operational systems are restricted to much shorter and recent periods because of the constant change in model, observations and assimilation schemes. Therefore, the spin-up simulation was performed with a constant uniform CO<sub>2</sub> concentration of 387.4 ppm (corresponding to 2009) and a fixed vegetation fractional coverage corresponding to the same year. The spin-up simulations were driven with repeated meteorological data for the 2009-2010 period until model pools reached equilibrium.

- 5 The transient simulation for the 2009-2010 period was then initialized from the spin-up simulations using varying CO<sub>2</sub> concentrations and GEM meteorology.

To assess the impact of using present climate to spin up the model on the simulated carbon pools and fluxes, we also performed a special run that used repeated meteorological data for 2009-2010 from CRU-NCEP, and constant uniform CO<sub>2</sub> of 387.4 ppm, and a fixed vegetation fractional coverage corresponding to the year 2009 until the model pools reach equilibrium

- 10 (hereafter CTEM-CRUNCEP2yr).

Note that fire and land use change are not taken into account in the current model's simulations due to the large uncertainty in the global land use history (Houghton et al., 2012) that may yield significant biases in the simulated CO<sub>2</sub> fluxes. Also, the standard model parameters were not changed or tuned to improve model performance when using alternative meteorological inputs. Hence the main differences between the CLASS-CTEM runs are the meteorological inputs, and the set-up of the spin-up

- 15 simulations.

### 3.2 Forward simulation using GEM-MACH-GHG model

Forward simulations are performed using the GEM-MACH-GHG model to evaluate how well CLASS-CTEM, using meteorological inputs from GEM-MACH-GHG, is able to reproduce temporal variations in atmospheric CO<sub>2</sub> at monitoring stations. The estimated NEE from CTEM-CRUNCEP and CTEM-GEM were used as a surface boundary condition in GEM-MACH-GHG, which transports the signal from the surface fluxes throughout the atmosphere, to validate the resulting modelled concentrations against observations of atmospheric CO<sub>2</sub>. The other fluxes are kept the same as in Polavarapu et al. (2016). Specifically, the anthropogenic emissions from fossil fuel burning and cement manufacturing, biomass burning, ocean-atmosphere carbon exchange and initial atmospheric concentration (Jan 1, 2009) are based on CT2013B (Peters et al., 2007).

### 3.3 Inversion Analysis Configuration in the GEOS-Chem Model

- 25 Because flux inversions have been performed for over a decade with the in situ measurements, there is a considerable body of literature of such inversion results (e.g. Rödenbeck et al. (2003); Peters et al. (2007), and Peylin et al. (2013)). Consequently, for our experiments, we use this observing network as opposed to a combined one that includes the more recent satellite missions. Thus, the GEOS-Chem flux inversions use the flask observations of atmospheric CO<sub>2</sub> collected by NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network sites (Dlugokencky et al., 2015) and ECCC sampling sites (Worthy et al., 2009).
- 30 We use the same set of observation sites as described in Deng et al. (2014) (see their Section 2.1.2).

In this study, we use the similar a priori CO<sub>2</sub> fluxes of the anthropogenic emissions from fossil fuel burning and cement manufacturing, biomass burning, and ocean-atmosphere carbon exchange described in Deng et al. (2014) in order to maximize comparability with the those results. However, for the biospheric flux of CO<sub>2</sub>, we conducted three runs using three different

NEE priors from CTEM-GEM, CTEM-CRUNCEP, and BEPS. The optimized 3-D CO<sub>2</sub> mixing ratio field from CarbonTracker was used as the initial CO<sub>2</sub> field in the inversion runs. This choice was made because the CarbonTracker initial state produced a smaller global mean bias than the GEOS-Chem initial state. This may indicate that the interhemispheric gradient in TM5 is better than that in GEOS-Chem. When assimilating the flask data, we use a threshold of 8 ppm for rejecting observations in the GEOS-Chem assimilations, and this means when the difference of observed and the modeled mixing ratio is larger than 8 ppm, the observation is not ingested in the model and does not provide any information on the fluxes. We assimilated 5365, 5393, and 5601 observations into our model in inversions using CTEM-CRUNCEP, CTEM-GEM, and BEPS prior fluxes, respectively.

## 4 Results and Discussion

For the meteorological data, we compare temperature, shortwave radiation, and precipitation, which are considered to be the most important variables controlling land carbon dynamics (Piao et al., 2013). We also compare the component fluxes of GPP,  $R_{eco}$ , and net ecosystem exchange ( $NEE = R_{eco} - GPP$ ) in order to identify the potential drivers of differences between model simulations. To examine regional differences, data and model output are also spatially aggregated to the 11 land regions of the TransCom inverse model inter-comparison project (Gurney et al., 2003).

### 4.1 Differences in Meteorological Forcing

Here, we evaluate the meteorological data from GEM by comparing it against CRU-NCEP, CRU and ERAI datasets where possible. Figure 1 shows the spatial patterns of the differences in mean annual temperature (averaged over the period 2009-2010) between GEM, CRU-NCEP, and CRU ~~for 2009 and 2010.~~. The differences between CRU-NCEP and CRU show cold biases in middle and high Northern latitudes and warm biases in Africa and South America. CRU-NCEP retains the monthly climatology of CRU but adds the daily and diurnal variations of NCEP reanalyses (Wei et al., 2014). Thus, differences in annual mean temperature of CRU and CRU-NCEP should be small by design. In contrast, GEM is warmer than CRU over the North high latitudes and generally cooler elsewhere. The comparison also shows that CRU-NCEP is cooler than GEM in Northeastern North America, Eastern Europe, and Eastern Asia, and warmer in Africa, Southwestern Asia, South America, and the west coastline of North America. The differences in ~~Figs. 1e and 1f~~ Fig. 1c are much larger than those seen in ~~Figs. 1a and 1b~~ Fig. 1a and 1b because GEM analyses are completely independent of CRU. NCEP reanalyses are constrained by the global meteorological observing system and the datasets used in 2009-2010 are likely broadly similar to that used by operational centers such as ECCC. Indeed, Zhao et al. (2006) compared meteorological fields from NCEP, the Data Assimilation Office (DAO) (currently called the GMAO), and ECMWF for the 2000-2003 period and found that the NCEP fields had a cold bias at all latitudes and that the bias was largest in the tropics, which is similar to the bias in the GEM fields. Zhao et al. (2006) also found that the ECMWF ERA-40 (the precursor to ERAI) and DAO fields had smaller zonal mean biases compared to NCEP, but the ERA-40 fields were similar to those from GEM in that they had a high bias at high latitudes.



To better illustrate the differences between the datasets, we have plotted in Fig. 2 the monthly mean temperature averaged for the 11 TransCom land regions. All the data show the same seasonal variations, with opposite phases of temperature between hemispheres. The largest differences are found in the tropics and the Southern hemisphere. GEM tends to be biased low compared to the other data in Northern Africa, Southern Africa and temperate South America. In these regions, ERAI is closer to CRU observations, but in tropical South America, GEM is closer to observations. Since ERAI is a reanalysis product, it is expected to be much better than an operational analysis. Our GEM analyses are also further degraded (in terms of resolution) from ECCC operational analyses, so any reasonable comparability to ERAI results are considered promising. CRU-NCEP overall is in better agreement with the observations (CRU). This is not surprising given that CRU-NCEP was produced by combining CRU and NCEP/NCAR Reanalysis products.

Figure 3 shows the spatial distribution of the differences of mean annual shortwave radiation (averaged over the period 2009-2010) between GEM and CRU-NCEP, and ERAI datasets~~for 2009 and 2010.~~. Shortwave radiation estimates are not available in the CRU data set. The comparison indicates that CRU-NCEP is approximately  $15\text{--}70\text{ W m}^{-2}$  higher (sunny bias) than ERAI in the high latitudes and in the tropical land regions. In arid areas (i.e. Australia, Sahara, South Africa, Southern North America, Tibetan Plateau, and West Asia), CRU-NCEP is approximately  $15\text{--}50\text{ W m}^{-2}$  lower than ERAI. In contrast, GEM is approximately  $10\text{--}60\text{ W m}^{-2}$  higher than ERAI over all land regions, with the highest values ( $40\text{--}60\text{ W m}^{-2}$ ) over tropical lands. The shortwave radiation estimates from GEM is approximately  $10\text{--}80\text{ W m}^{-2}$  higher than those from CRU-NCEP over nearly all land regions, with the exception of Europe, Eastern North America and in a few grid cells in the tropical regions, where CRU-NCEP is higher ( $10\text{--}80\text{ W m}^{-2}$ ).

Figure 4 shows the monthly mean shortwave radiation averaged for the TransCom land regions. GEM and ERAI have more similar seasonal variability compared to CRU-NCEP in most of the land regions, especially in the tropics. However, ERAI shows slightly lower monthly mean values in Eurasia regions, and in tropical South America. Zhao et al. (2006) also found that ERA-40 (the precursor to ERAI) underestimated shortwave radiation in the tropics. Differences between the datasets in the tropics may be due to cloudiness biases over the Intertropical Convergence Zone (ITCZ) region, which have a large impact on radiative forcing (Dee et al., 2011).

The comparisons of the differences in annual total precipitation between GEM, CRU-NCEP, and CRU datasets are shown in Fig. 5. The smallest differences are between CRU-NCEP and CRU. The largest differences in magnitude between CRU-NCEP and CRU are mainly in the tropics, particularly tropical Asia, and along the west coast of South America. Also, the largest differences between GEM and CRU are in the tropics. The comparison also indicates that CRU-NCEP is wetter than GEM in the tropics and sub-tropics, and in the temperate regions, but is drier than GEM in some areas of the boreal regions, and over a few grid cells in central Africa, and China. In general, the tropics exhibit the largest differences between the GEM and CRU-NCEP datasets.

The comparisons between the monthly total precipitation integrated over the TransCom land regions are shown in Fig. 6. Unlike temperature and shortwave radiation (well represented by global models), there is a very clear difference in monthly total precipitation among the datasets, except between CRU-NCEP and CRU, which agree very well with some differences in the tropics. It is clear that the largest differences occur mainly during summer in each hemisphere, which is associated with high

precipitation. GEM tends to be drier mainly during summer. Despite the differences in the seasonal amplitude, GEM shows a quite similar seasonal variability compared to other datasets. We should keep in mind that precipitation estimates from the reanalysis/forecast systems are normally associated with large errors (Harris et al., 2014), particularly over land. These errors are due to problems with the convective parametrization in the models, and the fact that ground-based precipitation observations are not yet used in the data assimilation systems. ~~On the other hand~~Also, CRU monthly precipitation suffers large biases in areas where observations are sparse (i.e. tropics and southern Hemisphere) (Harris et al., 2014). In fact, the observation-based datasets are not based only on measurements, but are also sometimes model-dependent (i.e. filling gaps, interpolation, etc) (Harris et al., 2014). These deficiencies as well as the different spatial/temporal resolutions among models and observations can explain some of the differences between the datasets. Deficiencies in ERAI and CRU have been investigated in previous studies (Simmons et al., 2010; Balsamo et al., 2010; Szczypka et al., 2011).

In summary, the meteorological fields from GEM are similar in quality to those from reanalyses (ERAI) and observation-based (CRU and CRU-NCEP) datasets. However, there are some notable discrepancies in seasonal variations and spatial distribution patterns between GEM and CRU-NCEP, particularly in precipitation estimates in the tropics, which will be reflected in the estimated carbon fluxes. Biases in precipitation may indicate that the convective scheme used in GEM system needs to be improved, in particular, over the tropics. CLASS-CTEM driven by GEM precipitation will be impacted by these biases.

## 4.2 Impact of Meteorological Forcing on Carbon Fluxes

Here, we assess the impact of changing meteorological inputs on the simulated carbon fluxes to determine whether biases in fluxes can be attributed to biases in the meteorological variables.

### 4.2.1 Differences in Simulated Carbon Fluxes

To evaluate the spin-up procedure, the simulated global values of primary carbon pools and fluxes are summarized in Table 3 for the spin-up simulations from CTEM-CRUNCEP (which used the 1901-1940 climate data for the spin-up), and CTEM-GEM (which used the 2009-2010 climate data for the spin-up). CTEM-GEM produces smaller values of carbon pools and fluxes compared to CTEM-CRUNCEP. One possible explanation for this is the use of the present climate to spin up the model in the case of CTEM-GEM. Table 3 shows also the global values of carbon pools and fluxes simulated by the CTEM-CRUNCEP2yr experiment, which also uses just the 2009-2010 climate to spin up the model. Rather than reducing the size of carbon pools, CTEM-CRUNCEP2yr produces much higher values compared to both CTEM-CRUNCEP and CTEM-GEM. Table 3 also compares the mean areal land precipitation globally as well as for the tropical land band (30°N-30°S) averaged over the 1901-1940 (for CTEM-CRUNCEP), and 2009-2010 (for CTEM-CRUNCEP2yr and CTEM-GEM). For the CRU-NCEP runs, 2009-2010 is wetter than the 1901-1940 period at global and tropical scales, which can explain the higher productivity in CTEM-CRUNCEP2yr compared to CTEM-CRUNCEP. ~~On the other hand,~~GEM precipitation (2009-2010) is slightly higher than CRU-NCEP (1901-1940) at the global scale, but lower over the tropical band for the same periods. The drier tropical band is reflected in the estimated tropical GPP from CTEM-GEM (Table 3), which dominates the global total GPP (Beer et al. (2010), Anav et al. (2015), and many others). This may explain the low carbon pools simulated by CTEM-GEM. This comparison

suggests that precipitation plays a significant role in plant productivity in the tropics, and thus accurate precipitation patterns are necessary to establish realistic initial values for carbon pools and fluxes during the spin-up runs. Despite the differences in model inputs and spin-up configuration, the initial global carbon pools and fluxes from CTEM-GEM, however, are still within the range of the ~~other estimates (Melton and Arora, 2014, Table 2)~~preindustrial values from other modelling studies listed in Melton and Arora (2014, Table 2).

The low GPP values from CTEM-GEM warrant further discussion given that the initial estimates of carbon pools and fluxes are critical to obtain an accurate estimate of historical CO<sub>2</sub> fluxes (Exbrayat et al., 2014; Tian et al., 2015). Carbon stocks are often not well ~~modeled-modelled~~ in TEMs (Houghton et al., 2012; Tian et al., 2015). The ~~modeled-modelled~~ pool sizes can be adjusted by tuning the model parameters in order to match observation-based estimates of carbon stocks. For example, Carvalhais et al. (2008, 2010) have reported the limitation of the carbon cycle steady state assumption in TEMs. Carvalhais et al. (2010), therefore, introduced a new parameter in the CASA model that forced the adjustment of both vegetation and soil carbon pools from equilibrium (after spin-up) allowing for model runs to be initialized either as net sinks or sources. They found that including this new parameter yielded better model performance in simulating carbon fluxes in comparison to observations. Moreover, their ~~modeled-modelled~~ soil carbon stocks became closer to observations. However, large uncertainties and errors in measurements can produce biased parameters, and hence poor model performance. Thus, forcing agreement to a given global mean value of GPP (e.g. 120 ~~Pg C/yr~~PgC year<sup>-1</sup>) by tuning model parameters may lead to worse model performance and is not justifiable given the observational uncertainty in this value. Given that CLASS-CTEM will provide only a prior estimate of NEE (~~not GPP and R<sub>eco</sub> separately~~, at least in the first stage) for flux inversions in EC-CAS, adjusting the initial carbon pools ~~modeled-modelled~~ by CLASS-CTEM is not necessary and would not likely change the major conclusions derived here. Moreover, tuning of CLASS-CTEM specifically for the far-from-ideal spin-up process that we employed for the GEM fields would be dubious and would make comparison with CTEM-CRUNCEP results difficult. Beyond the global budget, which is well constrained by atmospheric data (Peylin et al., 2013), the main focus is to assess the ability of the model to simulate the spatial and temporal flux variations ~~in~~ (mainly the NEE that is used in flux inversions) in response to changes in environmental conditions and its ability to match the atmospheric signal.

For transient simulations, the simulated terrestrial carbon fluxes ~~for 2009-2010~~ from the two simulations (CTEM-CRUNCEP and CTEM-GEM) are compared to each other as well as to observation-based estimates (where possible) or independent model results. Figure 7 shows the annual spatial difference of GPP simulated by CLASS-CTEM (averaged over the period 2009-2010) and the observation-based GPP estimates from ~~B10 (averaged over the period 1998 to 2005)~~FLUXCOM. The figure also shows the spatial difference between the ~~modeled-modelled~~ GPP from BEPS and ~~B10-FLUXCOM~~ and the zonal distribution of GPP from all datasets. There are significant differences in the annual GPP between the two simulations (CTEM-CRUNCEP and CTEM-GEM) and the evaluation data, particularly in the most highly vegetated areas (i.e. the tropics, and the boreal and temperate regions). CTEM-CRUNCEP and CTEM-GEM have similar spatial differences over Western Europe and boreal Asia, and to a lesser extent over North America but they show poor agreement in the tropics. Tropical GPP from CTEM-CRUNCEP is overestimated compared to ~~B10-FLUXCOM~~ (see also Fig. 7b). In contrast, it is underestimated in the Amazonian region, western Africa, and tropical Asia with CTEM-GEM. In comparison to other model results, the spatial distribution of

the difference between BEPS and ~~B40-FLUXCOM~~ (Fig. 7a) reveals ~~different patterns almost everywhere~~ relatively smaller different patterns compared to CTEM. The zonally averaged GPP in Fig. 7b indicates that CTEM-CRUNCEP and BEPS agrees very well with ~~B40-FLUXCOM~~ compared to CTEM-GEM, which ~~underestimate GPP over~~ underestimates GPP in the tropics.

5 Since the formulation of most models, including CLASS-CTEM and BEPS, links respiration to photosynthesis (Melton and Arora, 2016),  $R_{eco}$  estimates from both simulations and BEPS show a similar pattern to GPP (spatially and zonally), with significant differences in the most productive ecosystems (not shown here). The large discrepancies in seasonal variations and spatial distribution patterns between GEM and CRU-NCEP are due to the precipitation differences (temperature and shortwave radiation have much better agreement), particularly in the tropics. Figures 5 and 7 suggest that the differences in the spatial  
10 pattern of GPP are more closely associated with precipitation than temperature or shortwave radiation differences over the tropics. This is consistent with previous findings (Nemani et al., 2003; Jung et al., 2007; Beer et al., 2010; Piao et al., 2013; Anav et al., 2015) that interannual variation of productivity is primarily correlated with the precipitation over the tropics.

To examine regional differences, the seasonal variation and the annual mean of GPP from both simulations, and BEPS are also spatially aggregated to the 11 TransCom land regions and compared to FLUXCOM in Fig. 8. For  $R_{eco}$  (not shown  
15 here), the same conclusions can be drawn as from the GPP figures. Figure 8 shows that the seasonal variations of GPP from CTEM-CRUNCEP are consistently higher than those from CTEM-GEM. In the northern hemisphere regions, flux estimates have large seasonal variations, i.e., small values in winter and high values in summer, reflecting the seasonal change in carbon uptake by the land vegetation. The largest differences between CTEM-CRUNCEP and CTEM-GEM, in terms of the amplitude of the seasonal cycle, are found in the tropics, with CTEM-GEM having smaller amplitudes. However, in tropical Asia, the  
20 seasonal cycle from CTEM-GEM agrees well with BEPS and FLUXCOM compared to CTEM-CRUNCEP, which has larger annual GPP and  $R_{eco}$  (not shown here). Also, CTEM-GEM agrees very well with FLUXCOM in Europe and North America boreal compared to CTEM-CRUNCEP, especially during the growing season. In general, CTEM-GEM have some differences compared to CTEM-CRUNCEP over all regions mainly in terms of the amplitude, and to a lesser extent in the phase of the seasonal cycle. This is consistent with the findings of Dalmonech et al. (2015) who tested the impact of coupled and  
25 uncoupled configurations of JSBACH land surface component of the Max Planck Institute Earth System Model (MPI-ESM) on the simulated land carbon fluxes. They found that biases in the meteorological forcing to a large extent control the magnitude of GPP rather than the phenology and seasonal cycle of productivity, which could be more related to the model formulations (i.e. the timing and length of the growing season). In summary, these results indicate that the simulated fluxes from CLASS-CTEM are sensitive to the meteorological forcings over all land regions, which is consistent with the finding of Garnaud et al. (2014).

30 The annual GPP,  $R_{eco}$ , and the net flux are given in Table 4. Annual GPP values from CTEM-GEM for 2009 and 2010 are smaller than the multi-year average of GPP from BEPS (119.5 ~~Pg-C~~ PgC) (Deng et al., 2014), and from ~~B40-Beer et al. (2010)~~ (123  $\pm 8$  ~~Pg-C~~). On the other hand, annual PgC. Annual GPP values from CTEM-CRUNCEP for 2009 and 2010 are higher than those from BEPS and ~~B40~~ the calculated GPP from FLUXCOM (119.83 and 120 PgC for 2009, 2010 respectively). This leads to a stronger land carbon sink from CTEM-CRUNCEP compared to CTEM-GEM (Table 4). The weaker sink in CTEM-GEM  
35 is due to the lower precipitation estimates in the tropics (the region that mainly controls interannual variability in the carbon

cycle), and hence lower global GPP (Piao et al., 2013; Beer et al., 2010). Previous studies have reported a wide range of the global terrestrial GPP. For example, Piao et al. (2013) showed that GPP averaged across 10 land models is  $133 \pm 15$  (ranging from  $111 \pm 4$  PgC to  $151 \pm 4$  PgC). Other studies that are based on carbon cycle data assimilation suggest a GPP around 150 PgC (Koffi et al., 2012; Peylin et al., 2016), similar to Welp et al. (2011).

5 To assess the impact of the differences in GPP and  $R_{eco}$  from CTEM-CRUNCEP and CTEM-GEM on the seasonal cycle of NEE (the difference between GPP and  $R_{eco}$ ), Fig. 9 compares the NEE seasonal cycle from both simulations with the simulated prior NEE from BEPS (multi-year average) and the optimized NEE from CT2013B for 2009 and 2010 over the TransCom land regions. BEPS produces the smallest amplitude of the seasonal cycle of NEE while CTEM-CRUNCEP has the largest amplitude in northern land regions, except boreal Eurasia where the optimized NEE from CT2013B exhibits the largest  
10 amplitude (Fig. 9). For the South American tropical region, all models show considerable disagreement in the seasonal cycle, sometimes with opposite phases. In the northern hemisphere, CTEM-GEM and CTEM-CRUNCEP have better agreement with each other during winter than in summer. CTEM-GEM also tends to have the peak of the growing season one month earlier than CTEM-CRUNCEP (i.e. Eurasian boreal and North America temperate) due to the differences in GPP seasonal cycle (see Fig. 8). Even though there is large difference in the amplitude of the seasonal cycle of GPP (the same for  $R_{eco}$  - not shown  
15 here) from CTEM-GEM compared to CTEM-CRUNCEP (Fig. 8), the difference is much smaller in NEE. This is due to the fact that NEE is the difference between two large terms (GPP and  $R_{eco}$ ). That means, even though GPP and  $R_{eco}$  have large biases compared to observation-based estimates over the tropics, the biases in NEE are much smaller (4).

Despite the significant differences in model inputs, A recent study by Byrne et al. (JGR, under review, personal communication) evaluated the fluxes from CTEM-GEM and CTEM-CRUNCEP (the same fluxes used in the current study) in northern mid-latitude  
20 ecosystems by comparing GPP against Global Ozone Monitoring Experiment-2 (GOME-2) solar induced fluorescence (SIF) (Joiner et al., 2013; Khler et al., 2015) and NEE against total column  $CO_2$  ( $XCO_2$ ) from TCCON (Wunch et al., 2011). They found that while GPP from CTEM-CRUNCEP showed closer agreement with SIF than CTEM-GEM, CTEM-GEM showed closer agreement with TCCON  $XCO_2$  than CTEM-CRUNCEP. This implies that the biases in GPP might be compensated by biases in  $R_{eco}$ , which result in improved NEE fluxes from CTEM-GEM.

25 The NEE values in Table 4 show that CTEM-CRUNCEP tends to have a higher carbon sink compared to CTEM-GEM. CTEM-GEM simulates a land carbon sink of -1.2 and ~~differences in model structure and methods, the -2.2 PgC year<sup>-1</sup>~~ for 2009 and 2010, respectively, which is close to the optimized land sink from the CCDAS study of Peylin et al. (2016) (around 2.2 PgC year<sup>-1</sup>, for the 2000-2009 period). NEE from CTEM-GEM ~~flux estimates are within the range of the other estimates from TEMs used as a priori estimates in flux inversions (i.e. BEPS) or measurement-constrained fluxes (i.e. CT2013B)~~ is also  
30 more compatible with the Global Carbon Budget (GCP) (Quéré et al., 2015) ( $2.4 \pm 0.8$  PgC year<sup>-1</sup> for the 2000-2009 period).

In summary, CTEM-GEM has some issue in the tropics but that is where all prior (and posterior flux estimates) disagree and where observations are also inconsistent. However, Figs. 8 and 9 show that the Northern Hemisphere (where CTEM-GEM is fairly consistent with the other estimates) dominate the global seasonal cycle of carbon fluxes because it has the largest land  
35 areas that mainly dominated by forest ecosystems. At the same time, the net contribution from the tropical and the Sothern

Hemisphere regions are close to zero due to their opposite (and relatively small) seasonal cycles. Accordingly, the prior NEE information from CTEM-GEM is considered to be suitable for testing in the data assimilation context. However, flux estimates in the tropics from CTEM-GEM should be treated with caution.

#### 4.2.2 Modelled CO<sub>2</sub> Concentration

- 5 To assess the quality of the CO<sub>2</sub> fluxes from CLASS-CTEM simulations, terrestrial NEE fluxes from CTEM-CRUNCEP and CTEM-GEM are used as a priori land fluxes in the GEM-MACH-GHG global atmospheric CO<sub>2</sub> transport model. For comparisons, GEM-MACH-GHG was also run using the posterior NEE fluxes from CT2013B as described in Polavarapu et al. (2016). In these forward simulations, the anthropogenic emissions from fossil fuel burning and cement manufacturing, biomass burning, ocean-atmosphere carbon exchange are based on CT2013B (Peters et al., 2007) so that the only difference between  
10 the three runs is the terrestrial NEE fluxes.

Figure 10 compares the ~~modeled monthly times series of the modelled~~ CO<sub>2</sub> ~~time-series~~ from the two CLASS-CTEM simulations with that based on the CT2013B posterior fluxes, and ~~with~~ observed CO<sub>2</sub> at ~~Alert Bay, Canada (82.45°N, 62.52°W) and Mauna Loa, Hawaii (19.53°N, 155.58°W)~~ selected sites that are representative of various global regions (listed in Table 1) for continuous CO<sub>2</sub> measurements (Worthy et al., 2009; Dlugokencky et al., 2015). At ~~both all~~ observation sites, the simulations  
15 forced with CTEM-CRUNCEP and CTEM-GEM NEE fluxes (~~red-green~~ and blue curves) have a similar overestimation of the observed atmospheric CO<sub>2</sub> from December to ~~June~~ May, but the simulation forced with CT2013B fluxes (~~green-red~~ curves) has a much better match to observations. This makes sense because CT2013B fluxes have been informed by atmospheric observations whereas the other two fluxes have not. ~~The differences between the modeled and observed~~ Nevertheless, the overestimation of the CO<sub>2</sub> ~~might also indicate deficiencies in the seasonal cycle of flux estimates from CTEM-CRUNCEP and CTEM-GEM~~ concentrations can be attributed to the smaller net uptake simulated by CLASS-CTEM in the Northern Hemisphere (NH) regions during the winter season (see Fig. 9). At ALT and BRW, the two CLASS-CTEM simulations, in particularly CTEM-CRUNCEP, show a 1 month shift of the peak of the growing season compared to the observations. These differences in the seasonal cycle might indicate some limitations in the phenology of CLASS-CTEM (i.e. the larger source of carbon fluxes in wintertime and the seasonal cycle shift) and should be further investigated in future studies. In-  
20 ~~terestingly, for Alert, the simulation forced with~~ at ALT, BRW, IZO, MLO, and ZEP, during the NH growing season, in particularly in 2009, CTEM-GEM has a better match to ~~observations during the autumn of both years compared to those driven by CTEM-CRUNCEP and the observation than~~ CT2013B ~~which underestimate atmospheric CO<sub>2</sub>. As explained in Polavarapu et al. (2016), the autumn underestimation at Alert and CTEM-CRUNCEP.~~ This is promising as it indicates that the CTEM-GEM fluxes may be suitable to be used in the global inversion system (EC-CAS).

- 30 The large differences in the modelled CO<sub>2</sub> concentrations from the two CLASS-CTEM simulations highlight the sensitivity of the transport model to land fluxes. This might contradict the finding of (Ott et al., 2015) that large differences in prior flux estimates result in only small differences in CO<sub>2</sub> concentrations. The autumn underestimation with CT2013B fluxes (e.g. at ALT, BRW, and IZO) is likely due to a mismatch in seasonal-scale meridional transport between GEM-MACH-GHG and TM5 (the model used to produce CT2013B) (Polavarapu et al., 2016). We should also keep in mind that the performance of



the forward simulations also depends on the site specific conditions (e.g. topographic features, local sources of emissions, its location from urban areas, accuracy of the measurements, etc.) (Peters et al., 2005; Niwa et al., 2012; Shirai et al., 2017) and also on the performance of the transport model (Gurney et al., 2003, 2004). For Mauna Loa, CTEM-GEM has worse agreement with the observations, especially during winter. The overestimation of atmospheric CO<sub>2</sub> at Mauna Loa with CTEM-GEM is likely due to the less net uptake over the tropics as discussed before.

Figure ?? shows the zonal mean To better assess the quality of the modelled CO<sub>2</sub> for the 3 simulations (concentrations obtained with the two CLASS-CTEM simulations, Fig. 11 shows the Taylor diagram (Taylor, 2001) that compares the modelled CO<sub>2</sub> from CTEM-CRUNCEP, CTEM-GEM, and CT2013B ) for selected winter and summer dates. In general, the zonal mean fields from the three simulations have good agreement in the upper levels with the greatest differences near the surface. In winter, the CTEM-GEM simulation (center panels) produces high net emission (less carbon uptake) over the northern hemisphere, while CTEM-CRUNCEP (left panels) and CT2013B (right panels) show quite similar zonal means. In summer the CTEM-CRUNCEP simulation shows higher with the observed CO<sub>2</sub> over the northern hemisphere compared to CTEM-GEM and CT2013B, which have better agreement but to a lesser extent in summer 2010.

Overall, the comparison reveals that the modelled CO<sub>2</sub>, at Alert and Mauna Loa, from both CTEM-CRUNCEP and CTEM-GEM are generally in agreement, despite the differences in the meteorological inputs and the spin-up approach. Thus, despite the imperfect TEM spin-up procedure used for CTEM-GEM and the change in meteorological forcing at the selected sites (listed in Table 1). The position of each dot appearing on the plot quantifies how closely the modelled concentrations match the observations. The centered root mean square (RMS) difference between the simulated and observed patterns is proportional to the distance to the point on the x-axis identified as "1.0", which is the observations. The normalized standard deviation (dividing the standard deviation of simulated by the standard deviation of the observed) of the simulated pattern is proportional to the radial distance from the origin and it represents the agreement in the amplitude of the variability between the modelled and observed concentrations. As mentioned before, the simulation driven by retrieved fluxes (forced with the optimized fluxes from CT2013B ) is similar to that achieved with CLASS-CTEM prior fluxes. This result is consistent with the finding of (Ott et al., 2015) that large differences in prior flux estimates result in only small differences in CO<sub>2</sub> concentrations. The reason is that once the fluxes enter the atmosphere, the gradients they create are slowly smoothed out by various modes of atmospheric variability. Thus, concentrations reflect the time integrated history of past fluxes smoothed by atmospheric mixing, has a much better match to observations. Also, Fig. 11 shows that CTEM-GEM agrees better with the observations at all selected sites compared to CTEM-CRUNCEP in terms of correlations and the amplitude of the variability, which indicates an overall reasonable performance of the CTEM-GEM, especially during the growing season as mentioned before (see Fig. 10).

### 4.3 Inversion Analyses

The results in the previous two sections revealed that the GEM-MACH-GHG simulation of atmospheric CO<sub>2</sub> using CTEM-GEM and CTEM-CRUNCEP fluxes are comparable. They also indicate that the model driven with CTEM-GEM fluxes is able to reproduce temporal variations in atmospheric CO<sub>2</sub> at the selected sites. Since CTEM-GEM will be used as the land



component of EC-CAS, which is presently under development and thus not yet available, here we use the GEOS-Chem data assimilation system to examine the impact on regional flux estimates of using CTEM-GEM and CTEM-CRUNCEP as prior fluxes in the context of an atmospheric CO<sub>2</sub> inversion analysis. To determine how the retrieved fluxes obtained with the two CTEM-based priors compare to other documented inverse modelling results, we also perform an inversion analysis using BEPS prior fluxes, which is the ecosystem model used in the GEOS-Chem inversions of Deng et al. (2014, 2016), and we compare our results to the retrieved fluxes from CT2013.

### 4.3.1 Seasonal Cycle of the Flux Estimates

Figure 12 shows the seasonal cycle of the a posteriori NEE from the GEOS-Chem inversion analyses using the three different a priori estimates of NEE (CTEM-GEM, CTEM-CRUNCEP, and BEPS), together with the optimized NEE from CT2013B. The a priori NEE from CTEM-GEM, CTEM-CRUNCEP and BEPS are also shown. For northern land, there ~~is good agreement~~ are some differences between the optimized NEE in terms of the amplitude and the growing season, ~~except in especially in temperate region and in boreal Eurasia where the peak carbon uptake is greatest for CT2013B. Since the range of various posteriors can be taken as an estimate of the uncertainty of flux retrievals (e.g., Gurney et al., 2004; Peylin et al., 2013), we conclude that the CTEM-based~~ However, the spread in the posterior estimates for most northern land regions ~~are comparable to those from CT2013B and BEPS. This occurs~~ is smaller (compared to the tropical regions) because the surface observation network can reasonably constrain the northern extratropical latitudes (Peylin et al., 2013). ~~However, the~~ The spread between the fluxes is larger in the tropical regions, where NEE seasonal cycles show less agreement in both phase and magnitude. There are large differences in the seasonal cycle between the CTEM-based fluxes (CTEM-CRUNCEP and CTEM-GEM) and the BEPS-based fluxes in northern Africa and tropical South America that are present in both the prior and posterior fluxes. As a result of the limited observational coverage in the tropics, the posterior fluxes are strongly influenced by the prior fluxes. Consequently, the differences in the prior fluxes across the inversions are reflected in the posterior fluxes. Similarly, in the southern extratropics the posterior fluxes primarily reflect the prior flux distributions due to the sparsity of observations. This result is consistent with that of Peylin et al. (2013) who also found more disagreement of various inversion results in the tropics and southern hemisphere.

The amplitude of seasonal cycle of the optimized NEE from CTEM-CRUNCEP is significantly reduced compared to the a priori seasonal cycle in almost all land regions. The changes in the amplitude of the seasonal cycle for CTEM-GEM are smaller than those for CTEM-CRUNCEP. ~~This might indicate that Figure 9 shows that CTEM-CRUNCEP tends to have a larger amplitude of NEE compared to CTEM-GEM has the ability to simulate the seasonal cycle of CO<sub>2</sub> fluxes that is and the evaluation data in all regions. Also, the comparison in Figs. 10 and 11 revealed that modelled CO<sub>2</sub> concentrations from CTEM-GEM has a much better match to observations compared to CTEM-CRUNCEP, especially during the growing season. This might explain the significant shift in the optimized NEE from CTEM-CRUNCEP compared to CTEM-GEM. This also might indicate that the simulated fluxes from CTEM-GEM are more consistent with the atmospheric CO<sub>2</sub> signal - than CTEM-CRUNCEP. As noted above, the recent study by Byrne et al. (JGR, under review, personal communication) evaluated the a priori fluxes from CTEM-GEM and CTEM-CRUNCEP and found that compared to TCCON data, CTEM-GEM~~

NEE provided a better simulation of the atmospheric CO<sub>2</sub> seasonal cycle. They found that CTEM-CRUNCEP NEE produced a seasonal cycle in atmospheric CO<sub>2</sub> with a larger amplitude and with the onset of the springtime drawdown delayed by 10 days relative to CTEM-GEM NEE. However, both CLASS-CTEM simulations tend to have a large carbon source in wintertime, which is subsequently reduced in the optimized NEE (Fig. 12), especially in Europe, and NH temperate regions.

5 This demonstrates the capacity of inversion systems to constrain the phenological cycle in CLASS-CTEM, and ultimately can be used to optimize model parameters.

#### 4.3.2 Annual Mean Flux Estimates

The total annual a priori and a posteriori NEE from the GEOS-Chem inversion analyses for 2009-2010 are shown in Fig. 13 for the 11 land TransCom regions, along with the optimized values from CT2013B. Note that the tropical Asia panel has a different scale. All models estimate a sink (both for the a priori and the a posteriori) for the North America temperate, South American tropical, and Eurasian regions (except for temperate Eurasia, which has a source for CTEM-GEM prior). The largest difference between the a priori and a posteriori NEE in terms of the sign and magnitude were obtained for the Southern American temperate and Northern and Southern African regions. This is due to the fact that the tropics and Southern Hemisphere are poorly constrained by the current CO<sub>2</sub> network. Although the CT2013B fluxes tend to have stronger uptake in boreal Eurasia and the two African regions, BEPS has the largest uptake in the extra-tropical regions (North America temperate, Europe, Eurasia temperate) and tropical Asia. The two CLASS-CTEM simulations show large differences, from each other, in the optimized NEE in all land regions. Some of the difference between CT2013B and all of the GEOS-Chem estimates can be attributed to different transport models and configurations of data assimilation (Peylin et al., 2013) used in GEOS-Chem and CT2013B (uses Tracer Transport Model - version 5 (TM5) (Peters et al., 2007)). Figure 13 also indicates that, for 2009, GEOS-Chem allocates the strongest sink in Tropical Asia for CTEM-CRUNCEP (2 Pg-C-PgC year<sup>-1</sup>). Given that the inversion with CTEM-CRUNCEP fluxes suggest much weaker sinks (or larger sources) for the South American Tropical and Northern African regions compared to CTEM-GEM and BEPS, we suspect that stronger uptake in tropical Asia could reflect the inversion compensating for the larger sources in tropical south America and northern Africa. The hypothesis is that CTEM-CRUNCEP start with a larger a priori sink in tropical Asia and it gets enhanced in the inversion to compensate for the other tropical regional biases.

10

15

20

25

Figure 14 shows the global annual totals of NEE (a priori and a posteriori) for 2009-2010, as well as annual totals, aggregated into three latitudinal bands: Southern Hemisphere (SH): 90° S - 30° S, Tropics (TR): 30° S - 30° N, and Northern Hemisphere (NH): 30° N - 90° N). At the global scale, there is a good agreement between the optimized NEE in terms of magnitude. This indicates that the observations sufficiently constrain the global carbon budget so that the choice of prior fluxes is not critical. This is in agreement with Bruhwiler et al. (2011) who examine the impact of changing observation networks on flux estimates. However, optimized NEE for the three latitudinal bands show some differences, mainly in the tropics, where the observational coverage is poor.

30

These results are consistent with previous findings (e.g., Peters et al., 2007; Miller et al., 2015), which showed that optimized CO<sub>2</sub> fluxes in inversion analyses are heavily influenced by the spatial patterns in the a priori CO<sub>2</sub> fluxes, particularly in regions where observations are sparse (i.e. tropics and southern Hemisphere).

### 4.3.3 Evaluation of the Inversions

5 As described in section 3.3, we assimilated 5365, 5393, and 5601 observations into the inversion analysis using CTEM-CRUNCEP, CTEM-GEM, and BEPS prior fluxes. Based on these varied numbers of assimilated observations, we obtained  $\chi^2$  (chi-squared, which is a measure of the consistency of actual model-data mismatch with prescribed error covariances of model-data mismatch and where values close to 1.0 indicate consistency) for the three inversions are 0.91, 0.89, 0.76, and the correlation coefficients between observations and modeled root mean square (RMS) are 0.8398, 0.8436, and 0.8529, respectively, for CRUNCEP, 10 CTEM-GEM, and BEPS prior fluxes, respectively. The  $\chi^2$  and the correlation coefficient values suggests that the performance of CTEM-GEM and CTEM-CRUNCEP are quite similar and reliable in terms of fitting the assimilated observations. Figure 15 shows the histogram of the residuals between modelled (using CTEM-CRUNCEP, CTEM-GEM, and BEPS prior fluxes) and observed CO<sub>2</sub> concentrations. It is clear that there is no significant differences in the shape or width of the frequency distributions. Also, the standard deviations ( $\sigma$ ) are similar and the biases ( $\mu$ ) are all small relative to the standard deviations 15 biases for all simulations, indicating that all 3 experiments fit the assimilated data about equally well.

The flask observations from surface stations, which were assimilated by GEOS-Chem in the inversion, only provide a check on the consistency or set-up of the assimilation system. We still need to compare to independent observations that were not assimilated for validation. To more effectively evaluate the assimilation results, we compare the a posteriori CO<sub>2</sub> fields with independent data that were not ingested in the assimilation. Listed in Table 5 are the mean differences and root-mean-square 20 errors (RMSEs) of the a posteriori CO<sub>2</sub> relative to TCCON data in-between July 2009 and June 2010. The middle of the 2-year study period was chosen to avoid error due to spin-up and spin-down effects. All three fluxes reproduce the TCCON data well, but the BEPS-based CO<sub>2</sub> fields have the smallest RMSEs of 1.22 ppm and 1.18 ppm with respect to the data in 2009 and 2010, mean difference, RMSE, and inter-station bias of 0.49 ppm, 1.23 ppm, and 0.49 ppm, respectively. We find that the RMSEs RMSE for the CO<sub>2</sub> fields based on the CTEM-GEM fluxes are smaller than those from the CTEM-CRUNCEP 25 fluxes, with values of 1.24 ppm and 1.39 ppm in 2009 and 2010, compared to 1.42 ppm in both years for the fields from CTEM-CRUNCEP fluxes are identical. We also find that the fields based on CTEM-GEM have a smaller inter-station bias of 0.53 ppm, whereas those based on CTEM-CRUNCEP fluxes have a smaller mean difference of 0.53 ppm.

We also compare the a posteriori CO<sub>2</sub> fields with HIPPO aircraft data in the lower troposphere (see Table 6). As with the comparison to TCCON data, we find that the a posteriori fields based on BEPS fluxes produce the smallest RMSEs relative to 30 mean difference and RMSE relative to all of the aircraft data (between 0 - 10 km and 60°S - 60°N), with the fields from CTEM-GEM fluxes producing smaller RMSEs than those obtained from the CTEM-CRUNCEP fluxes. For the HIPPO-1 campaign, the a posteriori fields based on CTEM-CRUNCEP fluxes produce the smallest mean difference (-0.01 ppm), whereas for HIPPO-2 and HIPPO-3 the a posteriori CO<sub>2</sub> fields based on the BEPS and However, the differences in the RMSE between CTEM-CRUNCEP and CTEM-GEM fluxes produce the smallest mean differences (-0.41 ppm and 0.16 ppm), respectively are

~~not statistically significant. Overall, the comparisons show that that of the a posteriori results to the independent data indicate that CTEM-GEM provides relatively better agreement (in terms of the RMSEs) with the independent observations compared to CTEM-CRUNCEP. This implies that the spatial pattern of the can provide useful a priori fluxes from CTEM-GEM provides a better constraint than CTEM-CRUNCEP for the inversion system for CO<sub>2</sub> inversion analyses.~~

## 5 5 Conclusions

CLASS-CTEM will be used to provide first-guess (a priori) terrestrial fluxes for the Environment Canada Carbon Assimilation System (EC-CAS) (Polavarapu et al., 2016). The transport model of EC-CAS that relates surface fluxes to atmospheric CO<sub>2</sub> concentrations is based on the GEM-MACH-GHG model (Polavarapu et al., 2016). To ensure consistency between the land and transport model, CLASS-CTEM will be driven by the standard meteorological forcing simulated (24 h forecast) by GEM-MACH-GHG. Therefore, the main focus of this study was to assess the impact of using the meteorological inputs from GEM-MACH-GHG in simulating both regional and global carbon fluxes by CLASS-CTEM.

We first evaluated the quality of the meteorological inputs from GEM-MACH-GHG against the standard meteorological forcing (CRU-NCEP) that is used to drive the latest versions of CLASS-CTEM, as well as against observation-based or reanalysis datasets. The comparison ~~between the datasets indicates that the meteorological fields from GEM-MACH-GHG used in this study are similar in quality to those from observations-based datasets (CRU and CRU-NCEP) and reanalysis (ERA-Interim). The comparison also~~ shows that radiation and temperature data from GEM-MACH-GHG and CRU-NCEP are in good agreement. However, there are some notable discrepancies between GEM and CRU-NCEP in terms of seasonal variations and spatial patterns of precipitation estimates, especially in the tropics, with GEM being drier than CRU-NCEP, ERA-Interim, and CRU. That might indicate that the convective scheme used in GEM-MACH-GHG system needs to be improved in particular over the tropics.

~~Fluxes produced with GEM meteorology were obtained using a modified spin-up procedure based on current climate only. While it is clearly unsatisfactory to use a short climatology to spin-up carbon pools, it is an inevitable problem when coupling a TEM to an assimilation system since the latter focus on only a few years at a time. Moreover, operational weather assimilation systems are constantly changing so long datasets of analyses are simply not possible to obtain, unless reanalysis datasets are used.~~ The differences in the precipitation fields between GEM-MACH-GHG and CRU-NCEP was reflected in the estimated carbon fluxes (GPP and  $R_{\text{eco}}$ ). The amplitude and, to a lesser extent, the phase of the seasonal cycle are different between the two simulations, especially in the tropics. This is consistent with the findings of Dalmonech et al. (2015), who found that meteorological biases significantly control the magnitude of the productivity rather than the phenology and the seasonal cycle of carbon fluxes. ~~Fluxes produced with GEM meteorology were obtained using a modified spin-up procedure based on current climate only. While it is clearly unsatisfactory to use a short climatology to spin-up carbon pools, it is an inevitable problem when coupling a TEM to an assimilation system since the latter focus on only a few years at a time. Moreover, operational weather assimilation systems are constantly changing so long datasets of analyses are simply not possible to obtain.~~ Overall, the differences in the simulated fluxes between the two CLASS-CTEM simulations indicate that the model is sensitive to

the meteorological forcings over all land regions, and agree with the finding of Garnaud et al. (2014). Despite the deficiencies in the spin-up procedure, ~~the fluxes produced from CTEM-GEM were comparable in quality to those produced from CTEM-CRUNCEP. Moreover, the global constraint from observations is sufficient to determine the global budget irrespective of the choice of a priori flux. In fact, some inverse models use a neutral annual a priori flux to better assess the ability of the~~ observations to constrain the flux estimates (Deng et al., 2014). However, regional flux estimates are affected by the choice of a priori flux, particularly in data poor regions such as the tropics and southern hemisphere. Overall, the CTEM-GEM prior flux estimates are within the range of the other estimates from BEPS or top-down approaches (i.e. CT2013B), despite the significant differences in the model structure/approaches simulates a land carbon sink of -1.2 and -2.2 PgC year<sup>-1</sup> for 2009 and 2010, respectively, which is close to the optimized land sink from the CCDAS study of Peylin et al. (2016) (around 2.2 PgC year<sup>-1</sup>, for the 2000-2009 period), and also compatible with the Global Carbon Budget (GCP) (Quéré et al., 2015) (2.4 ±0.8 PgC year<sup>-1</sup> for the 2000-2009 period). However, flux estimates over the tropics from CTEM-GEM should be treated with caution due to the negative biases in the precipitation fields compared to all other datasets (i.e. ERA-Interim, CRU, and CRU-NCEP).

To assess their ability to model CO<sub>2</sub> at monitoring stations, NEE fluxes from CTEM-CRUNCEP and CTEM-GEM were used as a priori land fluxes in the GEM-MACH-GHG global atmospheric CO<sub>2</sub> transport model. The comparison ~~indicated that the simulated shows that the modelled CO<sub>2</sub> based on concentrations forced with CTEM-GEM compared reasonably well with observed have a better match to the observations of CO<sub>2</sub> in terms of temporal variations. However, the time series of the modelled concentration during the NH growing season compared to CTEM-CRUNCEP. The differences in the modelled CO<sub>2</sub> at the two selected sites indicated concentrations from the two CLASS-CTEM simulations highlight the sensitivity of~~ the transport model to land fluxes. The results also provided insights into the deficiencies in CLASS-CTEM. For example, ~~the comparison indicated that CTEM-CRUNCEP and CTEM-GEM NEE are overestimated compared to observations at all the selected sites during the NH winter season. The overestimation of the CO<sub>2</sub> concentrations can be attributed to the larger carbon source simulated by CLASS-CTEM in the Northern Hemisphere (NH) regions in wintertime. The results also show that the two CLASS-CTEM simulations have~~ some difficulties in capturing the phase of seasonal cycle of the observations, which ~~can be attributed to the deficiencies in simulating the seasonal cycle of NEE from indicates deficiencies in the phenology scheme of CLASS-CTEM in terms of phase as well as the magnitude, especially for the fluxes based on the CRU-NCEP meteorological data and this should be further investigated in future studies.~~ The deficiencies in simulating the seasonal cycle was also noticed in the study by Arora et al. (2009), who compared simulated monthly CO<sub>2</sub> from CanESM1 (CTEM used as the land component of that model) against observations at selected sites and found that there was a shift in the seasonal cycle (about a month later) at Barrow, Niwot Ridge, and Mauna Loa (see their Figure 11). The study by Anav et al. (2013), which compared 18 Earth system models, also showed that CanESM2 has some limitations reproducing the net uptake of carbon during spring and summer months.

To examine the impact of using fluxes from CTEM-GEM and CTEM-CRUNCEP as a priori flux estimates in atmospheric inversion analyses, we used the GEOS-Chem data assimilation system since EC-CAS is still under development. We assimilated in situ atmospheric CO<sub>2</sub> observations from the surface network to estimate optimized monthly mean NEE fluxes for

2009-2010. The time series of the estimated fluxes, integrated over different land regions, revealed that the optimized NEE is shifted from its a priori pattern in order to fit the data. For comparison with the CTEM-based fluxes we also used ~~BEP~~ BEPS fluxes (Deng et al., 2014) as an a priori in the inversion analyses. We found that the CTEM-based optimized fluxes produced atmospheric CO<sub>2</sub> concentrations that were consistent with those based on BEPS. ~~For example, the mean differences between independent TCCON data in 2010 and the modeled CO<sub>2</sub> based the CTEM-GEM, CTEM-CRUNCEP, and BEPS optimized fluxes were 0.80, 0.78, and 0.54 ppm, with RMSEs of 1.39, 1.42, and 1.18 ppm, respectively. For observations from the HIPPO-3 aircraft campaign, the mean differences between the observations and the CO<sub>2</sub> simulated from the optimized CTEM-GEM, CTEM-CRUNCEP, and BEPS fluxes were 0.16, 0.26, -0.28 ppm, with RMSEs of 0.94, 1.04, and 0.87 ppm, respectively, and they fit the assimilated data about equally well.~~ The results are promising for the EC-CAS project as they

demonstrate that the CTEM-GEM fluxes can provide useful a priori fluxes for the global inversion system. ~~When incorporated into the GEOS-Chem assimilation system, the inversion is capable of correcting the CTEM-GEM fluxes over annual and seasonal time scales in order to match the variability in the atmospheric data. In addition, the results of the optimized fluxes as well as the comparison of the a posteriori CO<sub>2</sub> to observations suggests that the spatial pattern of the a priori fluxes from CTEM-GEM provide a better constraint than CTEM-CRUNCEP for the inversion system.~~

Finally, this study provided insights into the deficiencies in the model, and data constraints (both meteorological and atmospheric CO<sub>2</sub> data). By coupling CLASS-CTEM into EC-CAS, a CCDAS approach becomes possible where observational constraints give feedback on a TEM. CLASS-CTEM into EC-CAS, this study helps to pave the way for the coupled meteorological and ecosystem model within the EnKF (e.g. see conclusions of Miller et al. (2015)) and is considered an important step toward understanding how meteorological uncertainties affect both CO<sub>2</sub> flux estimates and modelled atmospheric transport. Ultimately, such an approach ~~can help to improve the performance of~~ will provide more direct feedback to the CLASS-CTEM developers, and thus ~~improve the help to improve the performance of CLASS-CTEM by identifying the model limitations based on atmospheric constraints. This can also lead to improvements in the~~ CanESM which is used to address the question of the feedback between climate change and the carbon cycle.

*Code and data availability.* Fortran code for CLASS-CTEM modelling frame-work is available on request and upon agreeing to ECCC's licensing agreement available at: <http://collaboration.cmc.ec.gc.ca/science/rpn.comm>. Please contact the coauthor, Joe Melton (joe.melton@canada.ca), to obtain model code. The GEM and GEM-MACH source codes are integrated into the unique operational computing environments of ECCC. These source codes are copyrighted but are available upon request subject to the GNU Lesser General Public License (LGPL v2.1) agreement (contact the coauthor, Michael Neish (Michael.Neish@canada.ca)). Some documentation on GEM is available at: <http://collaboration.cmc.ec.gc.ca/science/rpn/gem/gemdm/gemdm.html> and [http://collaboration.cmc.ec.gc.ca/science/rpn/gef\\_html\\_public/](http://collaboration.cmc.ec.gc.ca/science/rpn/gef_html_public/). ECCC's model output data are available at: [https://weather.gc.ca/grib/index\\_e.html](https://weather.gc.ca/grib/index_e.html). The GEOS-Chem model, including the adjoint code, is freely available to the public and is distributed through GitLab. Instructions for obtaining and running the model are available on the GEOS-Chem wiki: (<http://wiki.seas.harvard.edu/geos-chem/>). The NOAA in situ CO<sub>2</sub> observations used in the inversion analysis are available from [ftp://aftp.cmdl.noaa.gov/data/trace\\_gases/co2/flask/surface/](ftp://aftp.cmdl.noaa.gov/data/trace_gases/co2/flask/surface/). All data generated by CLASS-CTEM is available from ECCC upon completion of a licensing agreement.

*Author contributions.* Bakr Badawy and Saroja Polavarapu designed the experiments. Bakr Badawy generated the required meteorological fields for CLASS-CTEM from GEM-MACH-GHG and performed the simulations for CLASS-CTEM and GEM-MACH-GHG and analysed the experiments and created the figures. Feng Deng performed the simulations for GEOS-Chem in coordination with Dylan Jones who contributed to the analysis and the interpretation of the assimilation results. Michael Neish developed and contributed to the GEM-MACH-GHG model diagnostics. Joe Melton and Vivek Arora provided the code of CLASS-CTEM used in this study and provided input on the implementation of the model. Ray Nassar provided helpful discussions. Bakr Badawy prepared the manuscript with contributions from all co-authors.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* Bakr Badawy was supported by Environment and Climate Change Canada (ECCC). We gratefully acknowledge all the data providers including the NOAA CarbonTracker assimilation system team for making their model products publicly available at <http://carbontracker.noaa.gov>. The NOAA in situ CO<sub>2</sub> observations used in the inversion analysis are available from <https://www.esrl.noaa.gov/gmd/dv/data/>. We [thank all the data providers who made their data available at the Global Atmosphere Watch \(GAW\) observing network.](#) We also would like to thank Doug Worthy for developing and maintaining ECCCs greenhouse gas measurement network and for providing the CO<sub>2</sub> concentration measurements. We gratefully acknowledge TCCON PIs (cited in Table 2) for making their data available. TCCON data were obtained from the TCCON Data Archive, hosted by the Carbon Dioxide Information Analysis Center (CDIAC) - [tccon.onrl.gov](http://tccon.onrl.gov). We gratefully thank the HIPPO (HIAPER Pole-to-Pole Observations, National Science Foundation, NSF, NSF/NCAR Gulfstream-V (GV)) science team for making their data freely available at <http://hippo.ornl.gov>. The collection of the original HIPPO data were supported by NSF and NOAA. ERA-Interim data used in this study have been obtained from the ECMWF data server: <http://data.ecmwf.int/data>. We thank [M. Jung for providing access to the GPP FLUXCOM data.](#) We thank Douglas Chan who evaluated the manuscript for the internal review process at ECCC and provided helpful comments, which improved the manuscript. [We would like to thank the two anonymous referees for insightful comments that helped to improve the paper.](#)



## References

- Agusti-Panareda, A., Massart, S., Chevallier, F., Boussetta, S., Balsamo, G., Beljaars, A., Ciais, P., Deutscher, N. M., Engelen, R., Jones, L., Kivi, R., Paris, J.-D., Peuch, V.-H., Sherlock, V., Vermeulen, A. T., Wennberg, P. O., and Wunch, D.: Forecasting global atmospheric CO<sub>2</sub>, *Atmospheric Chemistry and Physics*, 14, 11959–11983, <https://doi.org/10.5194/acp-14-11959-2014>, <http://www.atmos-chem-phys.net/14/11959/2014/>, 2014.
- Agusti-Panareda, A., Diamantakis, M., Bayona, V., Klappenbach, F., and Butz, A.: Improving the inter-hemispheric gradient of total column atmospheric CO<sub>2</sub> and CH<sub>4</sub> in simulations with the ECMWF semi-Lagrangian atmospheric global model, *Geoscientific Model Development Discussions*, 2016, 1–31, <https://doi.org/10.5194/gmd-2016-143>, <http://www.geosci-model-dev-discuss.net/gmd-2016-143/>, 2016.
- Agusti-Panareda, A., Diamantakis, M., Bayona, V., Klappenbach, F., and Butz, A.: Improving the inter-hemispheric gradient of total column atmospheric CO<sub>2</sub> and CH<sub>4</sub> in simulations with the ECMWF semi-Lagrangian atmospheric global model, *Geoscientific Model Development*, 10, 1–18, <https://doi.org/https://doi.org/10.5194/gmd-10-1-2017>, 2017.
- Anav, A., Friedlingstein, P., Kidston, M., Bopp, L., Ciais, P., Cox, P., Jones, C., Jung, M., Myneni, R., and Zhu, Z.: Evaluating the Land and Ocean Components of the Global Carbon Cycle in the CMIP5 Earth System Models, *Journal of Climate*, 26, 6801–6843, <https://doi.org/10.1175/JCLI-D-12-00417.1>, <http://dx.doi.org/10.1175/JCLI-D-12-00417.1>, 2013.
- Anav, A., Friedlingstein, P., Beer, C., Ciais, P., Harper, A., Jones, C., Murray-Tortarolo, G., Papale, D., Parazoo, N. C., Peylin, P., Piao, S., Sitch, S., Viovy, N., Wiltshire, A., and Zhao, M.: Spatiotemporal patterns of terrestrial gross primary production: A review, *Reviews of Geophysics*, 53, 785–818, <https://doi.org/10.1002/2015RG000483>, <http://dx.doi.org/10.1002/2015RG000483>, 2015RG000483, 2015.
- Arora, V. K.: Simulating energy and carbon fluxes over winter wheat using coupled land surface and terrestrial ecosystem models, *Agricultural and Forest Meteorology*, 118, 21–47, 2003.
- Arora, V. K. and Boer, G. J.: Fire as an interactive component of dynamic vegetation models, *Journal of Geophysical Research-Biogeosciences*, 110, G02008, doi:10.1029/2005JG000042, 2005.
- Arora, V. K. and Boer, G. J.: Uncertainties in the 20th century carbon budget associated with land use change, *Global Change Biology*, 16, 3327–3348, <https://doi.org/10.1111/j.1365-2486.2010.02202.x>, 2010.
- Arora, V. K., Boer, G. J., Christian, J. R., Curry, C. L., Denman, K. L., Zahariev, K., Flato, G. M., Scinocca, J. F., Merryfield, W. J., and Lee, W. G.: The Effect of Terrestrial Photosynthesis Down Regulation on the Twentieth-Century Carbon Budget Simulated with the CCCma Earth System Model, *J. Climate*, 22, 6066–6088, <https://doi.org/10.1175/2009JCLI3037.1>, 2009.
- Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C. D., Christian, J. R., Bonan, G., Bopp, L., Brovkin, V., Cadule, P., Hajima, T., Ilyina, T., Lindsay, K., Tjiputra, J. F., and Wu, T.: Carbon-Concentration and Carbon-Climate Feedbacks in CMIP5 Earth System Models, *Journal of Climate*, 26, 5289–5314, <https://doi.org/10.1175/JCLI-D-12-00494.1>, <http://dx.doi.org/10.1175/JCLI-D-12-00494.1>, 2013.
- Badawy, B., Rödenbeck, C., Reichstein, M., Carvalhais, N., and Heimann, M.: Technical Note: The Simple Diagnostic Photosynthesis and Respiration Model (SDPRM), *Biogeosciences*, 10, 6485–6508, <https://doi.org/10.5194/bg-10-6485-2013>, 2013.
- Badawy, B., Arora, V. K., Melton, J. R., and Nassar, R.: Modeling the diurnal variability of respiratory fluxes in the Canadian Terrestrial Ecosystem Model (CTEM), *J. Adv. Model. Earth Syst.*, 8, 614–633, <http://dx.doi.org/10.1002/2015MS000540>, 2016.
- Balsamo, G., Boussetta, S., Lopez, P., and Ferranti, L.: Evaluation of ERA-Interim and ERA-Interim GPCP-rescaled precipitation over the U.S.A., *ERA Report Series 5*, pp. 10., ECMWF, Reading, England, 2010.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rodenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luyssaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal,

- E., Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial Gross Carbon Dioxide Uptake: Global Distribution and Covariation with Climate, *Science*, 329, 834, <https://doi.org/10.1126/science.1184984>, 2010.
- Berrisford, P., Dee, D., Poli, P., Brugge, R., Fielding, K., Fuentes, M., Källberg, P., Kobayashi, S., Uppala, S., and Simmons, A.: The ERA-Interim archive Version 2.0, Shinfield Park, Reading, 2011.
- 5 Blumenstock, T., Hase, F., Schneider, M., Garcia, O., and Sepulveda, E.: TCCON data from Izana, Tenerife, Spain, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.izana01.R0/1149295>, <https://doi.org/10.14291/tccon.ggg2014.izana01.R0/1149295>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- Bruhwyler, L. M. P., Michalak, A. M., and Tans, P. P.: Spatial and temporal resolution of carbon flux estimates for 1983–2002, *Biogeosciences*, 8, 1309–1331, <https://doi.org/10.5194/bg-8-1309-2011>, <http://www.biogeosciences.net/8/1309/2011/>, 2011.
- 10 Carvalhais, N., Reichstein, M., Seixas, J., Collatz, G. J., Pereira, J. S., Berbigier, P., Carrara, A., Granier, A., Montagnani, L., Papale, D., Rambal, S., Sanz, M. J., and Valentini, R.: Implications of the carbon cycle steady state assumption for biogeochemical modeling performance and inverse parameter retrieval, *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2007GB003033>, <http://dx.doi.org/10.1029/2007GB003033>, gB2007, 2008.
- 15 Carvalhais, n., reichstein, m., ciais, p., collatz, g. j., mahecha, m. d., montagnani, l., papale, d., rambal, s., and seixas, j.: Identification of vegetation and soil carbon pools out of equilibrium in a process model via eddy covariance and biometric constraints, *Global Change Biology*, 16, 2813–2829, <https://doi.org/10.1111/j.1365-2486.2010.02173.x>, <http://dx.doi.org/10.1111/j.1365-2486.2010.02173.x>, 2010.
- Charron, M., Polavarapu, S., Buehner, M., Vaillancourt, P. A., Charette, C., Roch, M., Morneau, J., Garand, L., Aparicio, J. M., MacPherson, S., Pellerin, S., St-James, J., and Heilliette, S.: The Stratospheric Extension of the Canadian Global Deterministic Medium-Range Weather
- 20 Forecasting System and Its Impact on Tropospheric Forecasts, *Monthly Weather Review*, 140, 1924–1944, <https://doi.org/10.1175/MWR-D-11-00097.1>, <http://dx.doi.org/10.1175/MWR-D-11-00097.1>, 2012.
- Chen, J. M., Mo, G., Pisek, J., Liu, J., Deng, F., Ishizawa, M., and Chan, D.: Effects of foliage clumping on the estimation of global terrestrial gross primary productivity, *Global Biogeochemical Cycles*, 26, <https://doi.org/10.1029/2010GB003996>, <http://dx.doi.org/10.1029/2010GB003996>, gB1019, 2012.
- 25 Clein, J., McGuire, A. D., Euskirchen, E. S., and Calef, M.: The Effects of Different Climate Input Datasets on Simulated Carbon Dynamics in the Western Arctic, *Earth Interactions*, 11, 1–24, <https://doi.org/10.1175/EI229.1>, <http://dx.doi.org/10.1175/EI229.1>, 2007.
- Côté, J., Gravel, S., Méthot, A., Patoine, A., Roch, M., and Staniforth, A.: The Operational CMC-MRB Global Environmental Multi-scale (GEM) Model. Part I: Design Considerations and Formulation, *Mon. Wea. Rev.*, 126, 1373–1395, [https://doi.org/10.1175/1520-0493\(1998\)126<1373:TOCMGE>2.0.CO;2](https://doi.org/10.1175/1520-0493(1998)126<1373:TOCMGE>2.0.CO;2), [http://dx.doi.org/10.1175/1520-0493\(1998\)126<1373:TOCMGE>2.0.CO;2](http://dx.doi.org/10.1175/1520-0493(1998)126<1373:TOCMGE>2.0.CO;2), 1998a.
- 30 Dalmonech, D., Zaehle, S., Schuermann, G. J., Brovkin, V., Reick, C., and Schnur, R.: Separation of the Effects of Land and Climate Model Errors on Simulated Contemporary Land Carbon Cycle Trends in the MPI Earth System Model version 1, *Journal of Climate*, 28, 272–291, <https://doi.org/10.1175/JCLI-D-13-00593.1>, <http://dx.doi.org/10.1175/JCLI-D-13-00593.1>, 2015.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kallberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, <http://dx.doi.org/10.1002/qj.828>, 2011.

- Deng, F. and Chen, J. M.: Recent global CO<sub>2</sub> flux inferred from atmospheric CO<sub>2</sub> observations and its regional analyses, *Biogeosciences*, 8, 3263–3281, <https://doi.org/10.5194/bg-8-3263-2011>, <http://www.biogeosciences.net/8/3263/2011/>, 2011.
- Deng, F., Jones, D. B. A., Henze, D. K., Bousserrez, N., Bowman, K. W., Fisher, J. B., Nassar, R., O'Dell, C., Wunch, D., Wennberg, P. O., Kort, E. A., Wofsy, S. C., Blumenstock, T., Deutscher, N. M., Griffith, D. W. T., Hase, F., Heikkinen, P., Sherlock, V., Strong, K., Sussmann, R., and Warneke, T.: Inferring regional sources and sinks of atmospheric CO<sub>2</sub> from GOSAT XCO<sub>2</sub> data, *Atmospheric Chemistry and Physics*, 14, 3703–3727, <https://doi.org/10.5194/acp-14-3703-2014>, <http://www.atmos-chem-phys.net/14/3703/2014/>, 2014.
- Deng, F., Jones, D. B. A., O'Dell, C. W., Nassar, R., and Parazoo, N. C.: Combining GOSAT X CO<sub>2</sub> observations over land and ocean to improve regional CO<sub>2</sub> flux estimates, *Journal of Geophysical Research: Atmospheres*, 121, 1896–1913, <https://doi.org/10.1002/2015JD024157>, <http://dx.doi.org/10.1002/2015JD024157>, 2015JD024157, 2016.
- 10 Deutscher, N., Notholt, J., Messerschmidt, J., Weinzierl, C., Warneke, T., Petri, C., Grupe, P., and Katrynski, K.: TC-CON data from Bialystok, Poland, Release GGG2014R1, <http://dx.doi.org/10.14291/tccon.ggg2014.bialystok01.R1/1183984>, <https://doi.org/10.14291/tccon.ggg2014.bialystok01.R1/1183984>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- 15 Dlugokencky, E., Lang, P., Masarie, K., Crotwell, A., and Crotwell, M.: Atmospheric Carbon Dioxide Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1968–2014, Version: 2015-08-03, [ftp://afpt.cmdl.noaa.gov/data/trace\\_gases/co2/flask/surface/](ftp://afpt.cmdl.noaa.gov/data/trace_gases/co2/flask/surface/), 2015.
- Exbrayat, J.-F., Pitman, A. J., and Abramowitz, G.: Response of microbial decomposition to spin-up explains CMIP5 soil carbon range until 2100, *Geoscientific Model Development*, 7, 2683–2692, <https://doi.org/10.5194/gmd-7-2683-2014>, <http://www.geosci-model-dev.net/7/2683/2014/>, 2014.
- 20 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-Carbon Cycle Feedback Analysis: Results from the C4MIP Model Intercomparison, *J. Climate*, 19, 3337–3353, <https://doi.org/10.1175/JCLI3800.1>, 2006.
- Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K., and Knutti, R.: Uncertainties in CMIP5 Climate Projections due to Carbon Cycle Feedbacks, *Journal of Climate*, 27, 511–526, <https://doi.org/10.1175/JCLI-D-12-00579.1>, <http://dx.doi.org/10.1175/JCLI-D-12-00579.1>, 2014.
- 25 Garnaud, C., Sushama, L., and Arora, V. K.: The effect of driving climate data on the simulated terrestrial carbon pools and fluxes over North America, *International Journal of Climatology*, 34, 1098–1110, <https://doi.org/10.1002/joc.3748>, 2014.
- Girard, C., Plante, A. and Desgagné, M., McTaggart-Cowan, R., Côté, J., Charron, M., Gravel, S., Lee, V., Patoine, A., Qaddouri, A., Roch, M., Spacek, L., Tanguay, M., Vaillancourt, P. A., and Zadra, A.: Staggered Vertical Discretization of the Canadian Environmental Multiscale (GEM) Model Using a Coordinate of the Log-Hydrostatic-Pressure Type, *Mon. Wea. Rev.*, 142, 1183–1196, <https://doi.org/10.1175/MWR-D-13-00255.1>, <http://dx.doi.org/10.1175/MWR-D-13-00255.1>, 2013.
- 30 Griffith, D. W. T., Deutscher, N., Velazco, V. A., Wennberg, P. O., Yavin, Y., Aleks, G. K., Washenfelder, R., Toon, G. C., Blavier, J.-F., Murphy, C., Jones, N., Kettlewell, G., Connor, B., Macatangay, R., Roehl, C., Ryzcek, M., Glowacki, J., Culgan, T., and Bryant, G.: TCCON data from Darwin, Australia, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.darwin01.R0/1149290>, <https://doi.org/10.14291/tccon.ggg2014.darwin01.R0/1149290>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014a.
- 35

- Griffith, D. W. T., Velazco, V. A., Deutscher, N., Murphy, C., Jones, N., Wilson, S., Macatangay, R., Kettlewell, G., Buchholz, R. R., and Riggensbach, M.: TCCON data from Wollongong, Australia, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.wollongong01.R0/1149291>, <https://doi.org/10.14291/tccon.ggg2014.wollongong01.R0/1149291>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014b.
- 5 Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S. M., Fung, I. Y., Gloor, M., Heimann, M., Higuchi, K., John, J., Kowalczyk, E., Maki, T., Maksyutov, S., Peylin, P., Prather, M., Pak, B. C., Sarmiento, J., Taguchi, S., Takahashi, T., and Yuen, C.-W.: TransCom 3 CO<sub>2</sub> inversion intercomparison: 1. Annual mean control results and sensitivity to transport and prior flux information, *Tellus Series B-Chemical and Physical Meteorology*, 55, 555–579, 2003.
- 10 Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Pak, B. C., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fung, I. Y., Heimann, M., John, J., Maki, T., Maksyutov, S., Peylin, P., Prather, M., and Taguchi, S.: Transcom 3 inversion intercomparison: Model mean results for the estimation of seasonal carbon sources and sinks, *Global Biogeochemical Cycles*, 18, GB1010, doi:10.1029/2003GB002111, 2004.
- Harris, I., Jones, P., Osborn, T., and Lister, D.: Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset, *Int. J. Climatol.*, 34, 623–642, <http://dx.doi.org/10.1002/joc.3711>, 2014.
- 15 Hase, F., Blumenstock, T., Dohe, S., Gross, J., and Kiel, M.: TCCON data from Karlsruhe, Germany, Release GGG2014R1, <http://dx.doi.org/10.14291/tccon.ggg2014.karlsruhe01.R1/1182416>, <https://doi.org/10.14291/tccon.ggg2014.karlsruhe01.R1/1182416>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- 20 Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS-Chem, *Atmospheric Chemistry and Physics*, 7, 2413–2433, <https://doi.org/10.5194/acp-7-2413-2007>, <http://www.atmos-chem-phys.net/7/2413/2007/>, 2007.
- Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quéré, C., and Ramankutty, N.: Carbon emissions from land use and land-cover change, *Biogeosciences*, 9, 5125–5142, <https://doi.org/10.5194/bg-9-5125-2012>, <http://www.biogeosciences.net/9/5125/2012/>, 2012.
- 25 Huntzinger, D., Post, W., Wei, Y., Michalak, A., West, T., Jacobson, A., Baker, I., Chen, J., Davis, K., Hayes, D., Hoffman, F., Jain, A., Liu, S., McGuire, A., Neilson, R., Potter, C., Poulter, B., Price, D., Raczká, B., Tian, H., Thornton, P., Tomelleri, E., Viovy, N., Xiao, J., Yuan, W., Zeng, N., Zhao, M., and Cook, R.: North American Carbon Program (NACP) regional interim synthesis: Terrestrial biospheric model intercomparison, *Ecological Modelling*, 232, 144 – 157, <https://doi.org/http://dx.doi.org/10.1016/j.ecolmodel.2012.02.004>, <http://www.sciencedirect.com/science/article/pii/S0304380012000725>, 2012.
- 30 Huntzinger, D. N., Schwalm, C., Michalak, A. M., Schaefer, K., King, A. W., Wei, Y., Jacobson, A., Liu, S., Cook, R. B., Post, W. M., Berthier, G., Hayes, D., Huang, M., Ito, A., Lei, H., Lu, C., Mao, J., Peng, C. H., Peng, S., Poulter, B., Ricciuto, D., Shi, X., Tian, H., Wang, W., Zeng, N., Zhao, F., and Zhu, Q.: The North American Carbon Program Multi-Scale Synthesis and Terrestrial Model Intercomparison Project - Part 1: Overview and experimental design, *Geoscientific Model Development*, 6, 2121–2133, <https://doi.org/10.5194/gmd-6-2121-2013>, <http://www.geosci-model-dev.net/6/2121/2013/>, 2013.
- 35 Hurtt, G., Chini, L., Frolking, S., Betts, R., Feddema, J., Fischer, G., Fisk, J., Hibbard, K., Houghton, R., Janetos, A., Jones, C., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D., and Wang, Y.: Harmonization of land-use scenarios for the period 1500-2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117–161–, 2011.

- Joiner, J., Guanter, L., Lindstrot, R., Voigt, M., Vasilkov, A. P., Middleton, E. M., Huemmrich, K. F., Yoshida, Y., and Frankenberg, C.: Global monitoring of terrestrial chlorophyll fluorescence from moderate-spectral-resolution near-infrared satellite measurements: methodology, simulations, and application to GOME-2, *Atmospheric Measurement Techniques*, 6, 2803–2823, <https://doi.org/10.5194/amt-6-2803-2013>, 2013.
- 5 Jung, M., Vetter, M., Herold, M., Churkina, G., Reichstein, M., Zaehle, S., Ciais, P., Viovy, N., Bondeau, A., Chen, Y., Trusilova, K., Feser, F., and Heimann, M.: Uncertainties of modeling gross primary productivity over Europe: A systematic study on the effects of using different drivers and terrestrial biosphere models, *Global Biogeochemical Cycles*, 21, <https://doi.org/10.1029/2006GB002915>, <http://dx.doi.org/10.1029/2006GB002915>, gB4021, 2007.
- Jung, M., Reichstein, M., and Bondeau, A.: Towards global empirical upscaling of FLUXNET eddy covariance observations: validation of a  
10 model tree ensemble approach using a biosphere model, *Biogeosciences*, 6, 2013, 2009.
- Jung, M., Reichstein, M., Schwalm, C. R., Huntingford, C., Sitch, S., Ahlström, A., Arneth, A., Camps-Valls, G., Ciais, P., Friedlingstein, P., Gans, F., Ichii, K., Jain, A. K., Kato, E., Papale, D., Poulter, B., Raduly, B., Rdenbeck, C., Tramontana, G., Viovy, N., Wang, Y.-P., Weber, U., Zaehle, S., and Zeng, N.: Compensatory water effects link yearly global land CO<sub>2</sub> sink changes to temperature, *Nature*, 541, 516–520, <https://doi.org/10.1038/nature20780>, 2017.
- 15 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Met. Soc.*, 77, 437–471, 1996.
- Kaminski, T., Knorr, W., Schürmann, G., Scholze, M., Rayner, P. J., Zaehle, S., Blessing, S., Dorigo, W., Gayler, V., Giering, R., Gobron, N., Grant, J. P., Heimann, M., Hooker-Stroud, A., Houweling, S., Kato, T., Kattge, J., Kelley, D., Kemp, S., Koffi, E. N., Köstler, C., Mathieu,  
20 P.-P., Pinty, B., Reick, C. H., Rödenbeck, C., Schnur, R., Scipal, K., Sebald, C., Stacke, T., van Scheltinga, A. T., Vossbeck, M., Widmann, H., and Ziehn, T.: The BETHY/JSBACH Carbon Cycle Data Assimilation System: experiences and challenges, *Journal of Geophysical Research: Biogeosciences*, 118, 1414–1426, <https://doi.org/10.1002/jgrg.20118>, 2013.
- Kivi, R., Heikkinen, P., and Kyro, E.: TCCON data from Sodankyla, Finland, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.sodankyla01.R0/1149280>, <https://doi.org/10.14291/tccon.ggg2014.sodankyla01.R0/1149280>,  
25 TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- Koffi, E. N., Rayner, P. J., Scholze, M., and Beer, C.: Atmospheric constraints on gross primary productivity and net ecosystem productivity: Results from a carbon-cycle data assimilation system, *Global Biogeochemical Cycles*, 26, n/a–n/a, <https://doi.org/10.1029/2010gb003900>, 2012.
- 30 Koffi, E. N., Rayner, P. J., Scholze, M., Chevallier, F., and Kaminski, T.: Quantifying the constraint of biospheric process parameters by CO<sub>2</sub> concentration and flux measurement networks through a carbon cycle data assimilation system, *Atmospheric Chemistry and Physics*, 13, 10 555–10 572, <https://doi.org/10.5194/acp-13-10555-2013>, 2013.
- Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochemical Cycles*, 19, GB1015, [doi:10.1029/2003GB002199](https://doi.org/10.1029/2003GB002199), 2005.  
35
- Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Initial Onboard Performance of TANSO-FTS on GOSAT, in: *Advances in Imaging*, p. FTuC2, Optical Society of America, <https://doi.org/10.1364/FTS.2009.FTuC2>, <http://www.osapublishing.org/abstract.cfm?URI=FTS-2009-FTuC2>, 2009.

- Khler, P., Guanter, L., and Joiner, J.: A linear method for the retrieval of sun-induced chlorophyll fluorescence from GOME-2 and SCIAMACHY data, *Atmospheric Measurement Techniques*, 8, 2589–2608, <https://doi.org/10.5194/amt-8-2589-2015>, 2015.
- Lin, J. C., Pejam, M. R., Chan, E., Wofsy, S. C., Gottlieb, E. W., Margolis, H. A., and McCaughey, J. H.: Attributing uncertainties in simulated biospheric carbon fluxes to different error sources, *Global Biogeochemical Cycles*, 25, n/a–n/a, <https://doi.org/10.1029/2010gb003884>, 2011.
- Makar, P., Gong, W., Milbrandt, J., Hogrefe, C., Zhang, Y., Curci, G., Žabkar, R., Im, U., Balzarini, A., Baró, R., Bianconi, R., Cheung, P., Forkel, R., Gravel, S., Hirtl, M., Honzak, L., Hou, A., Jiménez-Guerrero, P., Langer, M., Moran, M., Pabla, B., Pérez, J., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., Zhang, J., and Galmarini, S.: Feedbacks between air pollution and weather, Part 1: Effects on weather, *Atmospheric Environment*, 115, 442 – 469, <https://doi.org/http://dx.doi.org/10.1016/j.atmosenv.2014.12.003>, <http://www.sciencedirect.com/science/article/pii/S1352231014009510>, 2015.
- McGuire, A. D., Sitch, S., Clein, J. S., Dargaville, R., Esser, G., Foley, J., Heimann, M., Joos, F., Kaplan, J., Kicklighter, D. W., Meier, R. A., Melillo, J. M., Moore III, B., Prentice, I. C., Ramankutty, N., Reichenau, T., Schloss, A., Tian, H., Williams, L. J., and Wittenberg, U.: Carbon balance of the terrestrial biosphere in the twentieth century: Analyses of CO<sub>2</sub>, climate and land use effects with four process-based ecosystem models, *Global Biogeochemical Cycles*, 15, 183–206, 2001.
- Meinshausen, M., Smith, S., Calvin, K., Daniel, J., Kainuma, M., Lamarque, J.-F., Matsumoto, K., Montzka, S., Raper, S., Riahi, K., Thomson, A., Velders, G., and van Vuuren, D.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Climatic Change*, 109, 213–241–, 2011.
- Melton, J. R. and Arora, V. K.: Sub-grid scale representation of vegetation in global land surface schemes: implications for estimation of the terrestrial carbon sink, *Biogeosciences*, 11, 1021–1036, <https://doi.org/10.5194/bg-11-1021-2014>, 2014.
- Melton, J. R. and Arora, V. K.: Competition between plant functional types in the Canadian Terrestrial Ecosystem Model (CTEM) v. 2.0, *Geoscientific Model Development*, 9, 323–361, <https://doi.org/doi:10.5194/gmd-9-323-2016>, 2016.
- Melton, J. R., Shrestha, R. K., and Arora, V. K.: The influence of soils on heterotrophic respiration exerts a strong control on net ecosystem productivity in seasonally dry Amazonian forests, *Biogeosciences*, 12, 1151–1168, <https://doi.org/10.5194/bg-12-1151-2015>, 2015.
- Miller, S. M., Hayek, M. N., Andrews, A. E., Fung, I., and Liu, J.: Biases in atmospheric CO<sub>2</sub> estimates from correlated meteorology modeling errors, *Atmospheric Chemistry and Physics*, 15, 2903–2914, <https://doi.org/10.5194/acp-15-2903-2015>, <http://www.atmos-chem-phys.net/15/2903/2015/>, 2015.
- Moran, M., Meñard, S., Talbot, D., Huang, P., Makar, P., Gong, W., Landry, H., Gravel, S., Gong, S., Crevier, L., Kallaur, A., and Sassi, M.: Particulate-matter forecasting with GEM-MACH15, a new Canadian air-quality forecast model, in: Steyn, D.G. and Rao, S.T (Eds.)s), *Air Pollution Modelling and Its Application XX*, Springer, Dordrecht, 2010.
- Nassar, R., Jones, D. B. A., Suntharalingam, P., Chen, J. M., Andres, R. J., Wecht, K. J., Yantosca, R. M., Kulawik, S. S., Bowman, K. W., Worden, J. R., Machida, T., and Matsueda, H.: Modeling global atmospheric CO<sub>2</sub> with improved emission inventories and CO<sub>2</sub> production from the oxidation of other carbon species, *Geoscientific Model Development*, 3, 689–716, <https://doi.org/10.5194/gmd-3-689-2010>, <http://www.geosci-model-dev.net/3/689/2010/>, 2010.
- Nemani, R. R., Keeling, C. D., Hashimoto, H., Jolly, W. M., Piper, S. C., Tucker, C. J., Myneni, R. B., and Running, S. W.: Climate-driven increases in global terrestrial net primary production from 1982 to 1999, *Science*, 300, 1560–1563, 2003.
- Niwa, Y., Machida, T., Sawa, Y., Matsueda, H., Schuck, T. J., Brenninkmeijer, C. A. M., Imasu, R., and Satoh, M.: Imposing strong constraints on tropical terrestrial CO<sub>2</sub> fluxes using passenger aircraft based measurements, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2012jd017474>, 2012.

- Notholt, J., Petri, C., Warneke, T., Deutscher, N., Buschmann, M., Weinzierl, C., Macatangay, R., and Grupe, P.: TC-CON data from Bremen, Germany, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.bremen01.R0/1149275>, <https://doi.org/10.14291/tccon.ggg2014.bremen01.R0/1149275>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- 5 Ott, L. E., Pawson, S., Collatz, G. J., Gregg, W. W., Menemenlis, D., Brix, H., Rousseaux, C. S., Bowman, K. W., Liu, J., Eldering, A., Gunson, M. R., and Kawa, S. R.: Assessing the magnitude of CO<sub>2</sub> flux uncertainty in atmospheric CO<sub>2</sub> records using products from NASA's Carbon Monitoring Flux Pilot Project, *Journal of Geophysical Research: Atmospheres*, 120, 734–765, <https://doi.org/10.1002/2014JD022411>, <http://dx.doi.org/10.1002/2014JD022411>, 2014JD022411, 2015.
- Peters, W., Miller, J. B., Whitaker, J., Denning, A. S., Hirsch, A., Krol, M. C., Zupanski, D., Bruhwiler, L., and Tans, P. P.: An ensemble  
10 data assimilation system to estimate CO<sub>2</sub> surface fluxes from atmospheric trace gas observations, *Journal of Geophysical Research-Atmospheres*, 110, 2005.
- Peters, W., Jacobson, A. R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., Miller, J. B., Bruhwiler, L. M. P., Pétron, G., Hirsch, A. I., Worthy, D. E. J., van der Werf, G. R., Randerson, J. T., Wennberg, P. O., Krol, M. C., and Tans, P. P.: An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker, *PNAS*, 104, 18 925–18 930, 2007.
- 15 Peylin, P., Law, R. M., Gurney, K. R., Chevallier, F., Jacobson, A. R., Maki, T., Niwa, Y., Patra, P. K., Peters, W., Rayner, P. J., Rödenbeck, C., van der Laan-Luijkx, I. T., and Zhang, X.: Global atmospheric carbon budget: results from an ensemble of atmospheric CO<sub>2</sub> inversions, *Biogeosciences*, 10, 6699–6720, <https://doi.org/10.5194/bg-10-6699-2013>, <http://www.biogeosciences.net/10/6699/2013/>, 2013.
- Peylin, P., Bacour, C., MacBean, N., Leonard, S., Rayner, P., Kuppel, S., Koffi, E., Kane, A., Maignan, F., Chevallier, F., Ciais, P., and Prunet, P.: A new stepwise carbon cycle data assimilation system using multiple data streams to constrain the simulated land surface carbon cycle,  
20 *Geoscientific Model Development*, 9, 3321–3346, <https://doi.org/10.5194/gmd-9-3321-2016>, 2016.
- Piao, S., Sitch, S., Ciais, P., Friedlingstein, P., Peylin, P., Wang, X., Ahlstrom, A., Anav, A., Canadell, J. G., Cong, N., Huntingford, C., Jung, M., Levis, S., Levy, P. E., Li, J., Lin, X., Lomas, M. R., Lu, M., Luo, Y., Ma, Y., Myneni, R. B., Poulter, B., Sun, Z., Wang, T., Viovy, N., Zaehle, S., and Zeng, N.: Evaluation of terrestrial carbon cycle models for their response to climate variability and to CO<sub>2</sub> trends, *Global Change Biology*, 19, 2117–2132, <https://doi.org/10.1111/gcb.12187>, <http://dx.doi.org/10.1111/gcb.12187>, 2013.
- 25 Polavarapu, S. M., Neish, M., Tanguay, M., Girard, C., de Grandpré, J., Semeniuk, K., Gravel, S., Ren, S., Roche, S., Chan, D., and Strong, K.: Greenhouse gas simulations with a coupled meteorological and transport model: the predictability of CO<sub>2</sub>, *Atmospheric Chemistry and Physics*, 16, 12 005–12 038, <https://doi.org/10.5194/acp-16-12005-2016>, <http://www.atmos-chem-phys.net/16/12005/2016/>, 2016.
- Potter, C. S., Randerson, J. T., Field, C. B., Matson, P. A., Vitousek, P. M., Mooney, H. A., and Klooster, S. A.: Terrestrial ecosystem production: a process model based on global satellite and surface data, *Global Biogeochemical Cycles*, 7, 811–841, 1993.
- 30 Quéré, C. L., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P., Peters, G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J. I., Keeling, R. F., Tans, P., Arneeth, A., Bakker, D. C. E., Barbero, L., Bopp, L., Chang, J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M., Feely, R. A., Gkritzalis, T., Harris, I., Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Goldewijk, K. K., Koven, C., Landschutzer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N., Millero, F., Munro, D. R., Murata, A., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T., Pérez, F. F., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G.,  
35 Rdenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle, S., and Zeng, N.: Global Carbon Budget 2015, *Earth System Science Data*, 7, 349–396, <https://doi.org/10.5194/essd-7-349-2015>, 2015.




- Rayner, P. J., Scholze, M., Knorr, W., Kaminski, T., Giering, R., and Widmann, H.: Two decades of terrestrial carbon fluxes from a carbon cycle data assimilation system (CCDAS), *Global Biogeochemical Cycles*, 19, 20 PP., doi:10.1029/2004GB002254, 2005.
- Reichstein, M., Kätterer, T., Andrén, O., Ciais, P., Schulze, E.-D., Cramer, W., Papale, D., and Valentini, R.: Temperature sensitivity of decomposition in relation to soil organic matter pools: critique and outlook, *Biogeosciences*, 2, 317–321, 2005.
- 5 Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648, <https://doi.org/10.1175/JCLI-D-11-00015.1>, <http://dx.doi.org/10.1175/JCLI-D-11-00015.1>, 2011.
- 10 Robichaud, A. and Ménard, R.: Multi-year objective analyses of warm season ground-level ozone and PM<sub>2.5</sub> over North America using real-time observations and Canadian operational air quality models, *Atmospheric Chemistry and Physics*, 14, 1769–1800, <https://doi.org/10.5194/acp-14-1769-2014>, <http://www.atmos-chem-phys.net/14/1769/2014/>, 2014.
- Rödenbeck, C., Houweling, S., Gloor, M., and Heimann, M.: CO<sub>2</sub> flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport, *Atmospheric Chemistry and Physics*, 3, 1919–1964, 2003.
- 15 Scholze, M., Kaplan, J. O., Knorr, W., and Heimann, M.: Climate and interannual variability of the atmosphere-biosphere 13CO<sub>2</sub> flux, *Geophysical Research Letters*, 30, 4 PP., doi:10.1029/2002GL015631, 2003.
- Sellers, P. J., Randall, D. A., Collatz, G. J., Berry, J. A., Field, C. B., Dazlich, D. A., Zhang, C., Collelo, C. D., and Bounoua, L.: A revised land surface parameterization (SiB2) for atmospheric GCMs: Part 1. Model formulation, *Journal of Climate*, 9, 676–705, 1996.
- Sherlock, V., Connor, B., Robinson, J., Shiona, H., Smale, D., and Pollard, D.: TCCON data from Lauder, New Zealand, 125HR, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.lauder02.R0/1149298>, <https://doi.org/10.14291/tccon.ggg2014.lauder02.R0/1149298>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- 20 Shirai, T., Ishizawa, M., Zhuravlev, R., Ganshin, A., Belikov, D., Saito, M., Oda, T., Valsala, V., Gomez-Pelaez, A., Langenfelds, R., and Maksyutov, S.: A decadal inversion of CO<sub>2</sub> using the Global Eulerian–Lagrangian Coupled Atmospheric model (GELCA): sensitivity to the ground-based observation network, *Tellus B: Chemical and Physical Meteorology*, 69, 1291–158, <https://doi.org/10.1080/16000889.2017.1291158>, 2017.
- 25 Simmons, A. J., Willett, K. M., Jones, P. D., Thorne, P. W., and Dee, D. P.: Low-frequency variations in surface atmospheric humidity, temperature, and precipitation: Inferences from reanalyses and monthly gridded observational data sets, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012442>, <http://dx.doi.org/10.1029/2009JD012442>, d01110, 2010.
- 30 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Global Change Biology*, 9, 161–185, 2003.
- Sitch, S., Friedlingstein, P., Gruber, N., Jones, S. D., Murray-Tortarolo, G., Ahlström, A., Doney, S. C., Graven, H., Heinze, C., Huntingford, C., Levis, S., Levy, P. E., Lomas, M., Poulter, B., Viovy, N., Zaehle, S., Zeng, N., Arneth, A., Bonan, G., Bopp, L., Canadell, J. G., Chevallier, F., Ciais, P., Ellis, R., Gloor, M., Peylin, P., Piao, S. L., Le Quéré, C., Smith, B., Zhu, Z., and Myneni, R.: Recent trends and drivers of regional sources and sinks of carbon dioxide, *Biogeosciences*, 12, 653–679, <https://doi.org/10.5194/bg-12-653-2015>, <http://www.biogeosciences.net/12/653/2015/>, 2015.
- 35


- Strong, K., Mendonca, J., Weaver, D., Fogal, P., Drummond, J., Batchelor, R., and Lindenmaier, R.: TCCON data from Eureka, Canada, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.eureka01.R0/1149271>, <https://doi.org/10.14291/tccon.ggg2014.eureka01.R0/1149271>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- 5 Suntharalingam, P., Jacob, D. J., Palmer, P. I., Logan, J. A., Yantosca, R. M., Xiao, Y. P., Evans, M. J., Streets, D. G., Vay, S. L., and Sachse, G. W.: Improved quantification of Chinese carbon fluxes using CO<sub>2</sub>/CO correlations in Asian outflow, *Journal of Geophysical Research-Atmospheres*, 109, D18S18, doi:10.1029/2003JD004362, 2004.
- Sussmann, R. and Rettinger, M.: TCCON data from Garmisch, Germany, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.garmisch01.R0/1149299>, <https://doi.org/10.14291/tccon.ggg2014.garmisch01.R0/1149299>,
- 10 TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- Szczypka, C., Calvet, J.-C., Albergel, C., Balsamo, G., Boussetta, S., Carrer, D., Lafont, S., and Meurey, C.: Verification of the new ECMWF ERA-Interim reanalysis over France, *Hydrology and Earth System Sciences*, 15, 647–666, <https://doi.org/10.5194/hess-15-647-2011>, <http://www.hydrol-earth-syst-sci.net/15/647/2011/>, 2011.
- 15 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *Journal of Geophysical Research: Atmospheres*, 106, 7183–7192, <https://doi.org/10.1029/2000JD900719>, <http://dx.doi.org/10.1029/2000JD900719>, 2001.
- Thornton, P. E., Cook, R. B., Braswell, B. H., Law, B. E., Post, W. M., Shugart, H. H., Rhyne, B. T., and Hook, L. A.: Archiving numerical models of biogeochemical dynamics, *EOS Transactions, American Geophysical Union*, 86, p. 431, 2005.
- Tian, H., Lu, C., Yang, J., Banger, K., Huntzinger, D. N., Schwalm, C. R., Michalak, A. M., Cook, R., Ciais, P., Hayes, D., Huang, M., Ito,
- 20 A., Jain, A. K., Lei, H., Mao, J., Pan, S., Post, W. M., Peng, S., Poulter, B., Ren, W., Ricciuto, D., Schaefer, K., Shi, X., Tao, B., Wang, W., Wei, Y., Yang, Q., Zhang, B., and Zeng, N.: Global patterns and controls of soil organic carbon dynamics as simulated by multiple terrestrial biosphere models: Current status and future directions, *Global Biogeochem. Cycles*, 29, 775–792, <http://dx.doi.org/10.1002/2014GB005021>, 2015.
- Tramontana, G., Jung, M., Schwalm, C. R., Ichii, K., Camps-Valls, G., Ráduly, B., Reichstein, M., Arain, M. A., Cescatti, A., Kiely, G.,
- 25 Merbold, L., Serrano-Ortiz, P., Sickert, S., Wolf, S., and Papale, D.: Predicting carbon dioxide and energy fluxes across global FLUXNET sites with regression algorithms, *Biogeosciences*, 13, 4291–4313, <https://doi.org/10.5194/bg-13-4291-2016>, 2016.
- Versegny, D.: CLASS - the Canadian Land Surface Scheme (Version 3.6), Technical Documentation, Tech. Rep. pp 179, Science and Technology Branch, Environment and Climate Change Canada, Toronto, 2012.
- Viovy, N.: CRUNCEP data set version 6, available at: [https://vesg.ipsl.upmc.fr/thredds/catalog/store/p529viov/cruncep/V6\\_1901\\_2014/](https://vesg.ipsl.upmc.fr/thredds/catalog/store/p529viov/cruncep/V6_1901_2014/catalog.html)
- 30 [catalog.html](https://vesg.ipsl.upmc.fr/thredds/catalog/store/p529viov/cruncep/V6_1901_2014/catalog.html) [last access: June 2016], 2016.
- Warneke, T., Messerschmidt, J., Notholt, J., Weinzierl, C., Deutscher, N., Petri, C., Grupe, P., Vuillemin, C., Truong, F., Schmidt, M., Ramonet, M., and Parmentier, E.: TCCON data from Orleans, France, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.orleans01.R0/1149276>, <https://doi.org/10.14291/tccon.ggg2014.orleans01.R0/1149276>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge,
- 35 Tennessee, U.S.A., 2014.
- Wei, Y., Liu, S., Huntzinger, D. N., Michalak, A. M., Viovy, N., Post, W. M., Schwalm, C. R., Schaefer, K., Jacobson, A. R., Lu, C., Tian, H., Ricciuto, D. M., Cook, R. B., Mao, J., and Shi, X.: The North American Carbon Program Multi-scale Synthesis and Terrestrial Model In-

- tercomparison Project - Part 2: Environmental driver data, Geoscientific Model Development, 7, 2875–2893, <https://doi.org/10.5194/gmd-7-2875-2014>, <http://www.geosci-model-dev.net/7/2875/2014/>, 2014.
- Welp, L. R., Keeling, R. F., Meijer, H. A. J., Bollenbacher, A. F., Piper, S. C., Yoshimura, K., Francey, R. J., Allison, C. E., and Wahlen, M.: Interannual variability in the oxygen isotopes of atmospheric CO<sub>2</sub> driven by El Niño, *Nature*, 477, 579–582, <https://doi.org/10.1038/nature10421>, 2011.
- Wennberg, P. O., Roehl, C., Wunch, D., Toon, G. C., Blavier, J.-F., Washenfelder, R., Keppel-Aleks, G., Allen, N., and Ayers, J.: TC-CON data from Park Falls, Wisconsin, USA, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.parkfalls01.R0/1149161>, <https://doi.org/10.14291/tccon.ggg2014.parkfalls01.R0/1149161>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014a.
- Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., Allen, N., Dowell, P., Teske, K., Martin, C., and Martin, J.: TCCON data from Lamont, Oklahoma, USA, Release GGG2014R0, <http://dx.doi.org/10.14291/tccon.ggg2014.lamont01.R0/1149159>, <https://doi.org/10.14291/tccon.ggg2014.lamont01.R0/1149159>, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014b.
- Wofsy, S., Daube, B., Jimenez, R., Kort, E., Pittman, J., Park, S., Commane, R., Xiang, B., Santoni, G., Jacob, D., Fisher, J., Pickett-Heaps, C., Wang, H., Wecht, K., Wang, Q.-Q., Stephens, B., Shertz, S., Watt, A., Romashkin, P., Campos, T., Haggerty, J., Cooper, W., Rogers, D., Beaton, S., Hendershot, R., Elkins, J., Fahey, D., Gao, R., Moore, F., Montzka, S., Schwarz, J., Perring, A., Hurst, D., Miller, B., Sweeney, C., Oltmans, S., Nance, D., Hints, E., Dutton, G., Watts, L., Spackman, J., Rosenlof, K., Ray, E., Hall, B., Zondlo, M., Diao, M., Keeling, R., Bent, J., Atlas, E., Lueb, R., and Mahoney, M.: HIPPO Merged 10-second Meteorology, Atmospheric Chemistry, and Aerosol Data (R20121129), [https://doi.org/10.3334/cdiac/hippo\\_010](https://doi.org/10.3334/cdiac/hippo_010), 2012.
- Wofsy, S. C.: HIPPER Pole-to-Pole Observations (HIPPO): fine-grained, global-scale measurements of climatically important atmospheric gases and aerosols, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 369, 2073–2086, 2011.
- Worthy, D. E. J., Chan, E., Ishizawa, M., Chan, D., Poss, C., Dlugokencky, E. J., Maksyutov, S., and Levin, I.: Decreasing anthropogenic methane emissions in Europe and Siberia inferred from continuous carbon dioxide and methane observations at Alert, Canada, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD011239>, <http://dx.doi.org/10.1029/2008JD011239>, d10301, 2009.
- Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W., Sherlock, V., and Wennberg, P. O.: The total carbon column observing network, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 369, 2087–2112, 2011.
- Wutzler, T. and Reichstein, M.: Soils apart from equilibrium & consequences for soil carbon balance modelling, *Biogeosciences*, 4, 125–136, <https://doi.org/10.5194/bg-4-125-2007>, <http://www.biogeosciences.net/4/125/2007/>, 2007.
- Yokota, T., Yoshida, Y., Eguchi, N., Ota, Y., Tanaka, T., Watanabe, H., and Maksyutov, S.: Global Concentrations of CO<sub>2</sub> and CH<sub>4</sub> Retrieved from GOSAT: First Preliminary Results, *SOLA*, 5, 160–163, <https://doi.org/10.2151/sola.2009-041>, 2009.
- Zhao, M., Running, S. W., and Nemani, R. R.: Sensitivity of Moderate Resolution Imaging Spectroradiometer (MODIS) terrestrial primary production to the accuracy of meteorological reanalyses, *J. Geophys. Res.*, 111, <http://dx.doi.org/10.1029/2004JG000004>, 2006.
- Zhu, Q. and Zhuang, Q.: Ecosystem biogeochemistry model parameterization: Do more flux data result in a better model in predicting carbon flux?, *Ecosphere*, 6, 1–20, <https://doi.org/10.1890/ES15-00259.1>, <http://dx.doi.org/10.1890/ES15-00259.1>, 2015.


Zobler, L.: A World Soil File for Global Climate Modelling. NASA Technical Memorandum 87802, <http://daac.ornl.gov/SOILS/guides/ZoblerSoil1.html>, 1986.



**Figure 1.** Comparison of spatial distribution patterns of annual mean temperature ( $^{\circ}\text{C}$ ) ~~for 2009 and 2010~~ (averaged over the period 2009-2010): (~~a and b~~) CRU-NCEP minus CRU, (~~e and d~~) CRU minus GEM, and (~~e and f~~) CRU-NCEP minus GEM.




**Figure 2.** Monthly mean temperature (°C) averaged for the TransCom land regions.




**Figure 3.** Comparison of spatial distribution patterns of annual mean shortwave radiation ( $\text{W m}^{-2}$ ) ~~for 2009 and 2010~~ (averaged over the period 2009-2010): (a and b) CRU-NCEP minus ERAI, (c and d) GEM minus ERAI, and (e and f) CRU-NCEP minus GEM.







**Figure 4.** Monthly mean shortwave radiation ( $\text{W m}^{-2}$ ) averaged for the 11 TransCom land regions.




**Figure 5.** Comparison of spatial distribution patterns of annual total precipitation ( $\text{mm year}^{-1}$ ) for 2009 and 2010 (averaged over the period 2009-2010): (a) and (b) CRU-NCEP minus CRU, (c) and (d) CRU minus GEM, and (e) and (f) CRU-NCEP minus GEM.




**Figure 6.** Monthly total precipitation (mm month<sup>-1</sup>) for the 11 TransCom land regions.




**Figure 7.** The annual spatial difference of GPP ( $\text{gC m}^{-2} \text{ year}^{-1}$ ) for CTEM-GEM, CTEM-CRUNCEP, and BEPS ~~for 2009 and 2010~~ against the observation-based GPP estimates from ~~B10~~ FLUXCOM (averaged over the period ~~1998 to 2005~~ 2009-2010). The zonal distributions of GPP from all datasets are shown (top-right).



**Figure 8.** The seasonal cycle of GPP from CTEM-GEM, CTEM-CRUNCEP, ~~and~~-BEPS, and FLUXCOM integrated over the 11 TransCom land regions.




**Figure 9.** The seasonal cycle of NEE from CTEM-GEM, CTEM-CRUNCEP, and BEPS in comparison to the optimized NEE from CT2013B integrated over the 11 TransCom land regions.




**Figure 10.** Comparison of the monthly time series of modelled  $\text{CO}_2$  concentrations using land prior fluxes from CTEM-CRUNCEP (redgreen) and CTEM-GEM (blue) with surface observations (black) at Alert-selected sites (toplisted in Table 1), and Mauna-Loa (bottom); and-modelled  $\text{CO}_2$  using posterior fluxes from CT2013B (greenred). The modelled  $\text{CO}_2$  was produced by a forward run of GEM-MACH-GHG.







**Figure 11.** Zonal-mean Taylor diagram of modelled monthly  $\text{CO}_2$  on 1 July 2009 and 31 Dec. 2010 concentrations produced by a forward run of GEM-MACH-GHG using land fluxes from CT2013B (top two rows red), and 1 July 2009 and 31 Dec. 2010 CTEM-GEM (bottom two rows blue) for, and CTEM-CRUNCEP (left column green), CTEM-GEM and observed  $\text{CO}_2$  (middle column identified as "1.0"), and CT2013B at selected sites (right column shapes) (listed in Table 1). The modelled  $\text{CO}_2$  was produced by a forward run of GEM-MACH-GHG.




**Figure 12.** The seasonal cycle of the optimized NEE from GEOS-Chem using three different prior estimates of NEE from CTEM-GEM, CTEM-CRUNCEP, and BEPS (indicated as well) in comparison to the optimized NEE from CT2013B integrated over the 11 TransCom land regions.



**Figure 13.** The annual total of the optimized NEE from GEOS-Chem using three different prior flux estimates of NEE from CTEM-GEM, CTEM-CRUNCEP, and BEPS (indicated as well) in comparison to the ~~the~~ optimized NEE from CT2013B integrated over the 11 TransCom land regions.



**Figure 14.** The monthly (left) and annual total (right) of the optimized NEE from GEOS-Chem using three different prior flux estimates of NEE from CTEM-GEM, CTEM-CRUNCEP, and BEPS (indicated as well) in comparison to the **red**-optimized NEE from CT2013B integrated over three latitudinal bands.



**Figure 15.** Frequency distributions of the residuals between modelled (using A) CTEM-CRUNCEP, B) CTEM-GEM, and C) BEPS prior fluxes, respectively) and observed CO<sub>2</sub> concentrations, where  $n$  is the number of the assimilated observations,  $\mu$  is the mean bias, and  $\sigma$  is the mean standard deviation.

**Table 1.** Locations of selected stations measuring atmospheric CO<sub>2</sub> concentrations used for comparison in the forward simulations.

<u>Code</u>	<u>Site name</u>	<u>Country</u>	<u>Longitude</u>	<u>Latitude</u>	<u>Elevation (m)</u>
<u>ALT</u>	<u>Alert</u>	<u>Canada</u>	<u>-62.5</u>	<u>82.45</u>	<u>210.</u>
<u>BRW</u>	<u>Barrow</u>	<u>USA</u>	<u>-156.6</u>	<u>71.32</u>	<u>11.</u>
<u>EGB</u>	<u>Egbert</u>	<u>Canada</u>	<u>-79.8</u>	<u>44.23</u>	<u>253.</u>
<u>IZO</u>	<u>Tenerife</u>	<u>Spain</u>	<u>-16.5</u>	<u>28.30</u>	<u>2367.</u>
<u>MLO</u>	<u>Mauna Loa</u>	<u>USA</u>	<u>-155.6</u>	<u>19.54</u>	<u>3397.</u>
<u>PAL</u>	<u>Pallas-Sammaltunturi</u>	<u>Finland</u>	<u>24.1</u>	<u>67.97</u>	<u>560.</u>
<u>YON</u>	<u>Yonagunijima</u>	<u>Japan</u>	<u>123.0</u>	<u>24.47</u>	<u>30.</u>
<u>ZEP</u>	<u>Zeppelinfjelle</u>	<u>Norway</u>	<u>11.9</u>	<u>78.90</u>	<u>475.</u>
<u>ZSF</u>	<u>Zugspitze</u>	<u>Germany</u>	<u>11.0</u>	<u>47.42</u>	<u>2656.</u>

**Table 2.** TCCON sites used in this study.

Site Name	Lat	Lon	Reference
Eureka, Canada	80.05 N	86.42 W	Strong et al. (2014)
Sodankyla, Finland	67.37 N	26.63 E	Kivi et al. (2014)
Bialystok, Poland	53.23 N	23.03 E	Deutscher et al. (2014)
Bremen, Germany	53.10 N	8.85 E	Notholt et al. (2014)
Karlsruhe, Germany	49.10 N	8.44 E	Hase et al. (2014)
Orleans, France	47.97 N	2.11 E	Warneke et al. (2014)
Garmisch, Germany	47.48 N	11.06 E	Sussmann and Rettinger (2014)
Park Falls, USA	45.95 N	90.27 W	Wennberg et al. (2014a)
Lamont, USA	36.60 N	97.49 W	Wennberg et al. (2014b)
Izana, Tenerife, Spain	28.3 N	16.5 W	Blumenstock et al. (2014)
Darwin, Australia	12.42 S	130.90 E	Griffith et al. (2014a)
Wollongong, Australia	34.41 S	150.88 E	Griffith et al. (2014b)
Lauder, New Zealand	45.04 S	169.68 E	Sherlock et al. (2014)

**Table 3.** Simulated global values of primary carbon pools and fluxes for the spin-up simulations using CTEM-CRUNCEP, CTEM-GEM and CTEM-CRUNCEP2yr. Values are a 20 year average at the end of model simulations. Mean areal precipitation (global land and for the 30°N-30°S land band) averaged for the 1901-1940 period used to spin-up CTEM-CRUNCEP, and for the 2009-2010 period used to spin-up CTEM-GEM and CTEM-CRUNCEP2yr, and the correspondence GPP estimates.

Variable	CTEM-CRUNCEP	CTEM-GEM	CTEM-CRUNCEP2yr
Gross primary productivity (Pg C yr <sup>-1</sup> )	118.0	97.0	139.8
Net primary productivity (Pg C yr <sup>-1</sup> )	58.0	47.0	70.0
Autotrophic respiration (Pg C yr <sup>-1</sup> )	60.5	49.6	69.8
Heterotrophic respiration (Pg C yr <sup>-1</sup> )	57.5	47.4	70.0
Litter carbon respiration (Pg C yr <sup>-1</sup> )	40.8	33.4	49.4
Soil carbon respiration (Pg C yr <sup>-1</sup> )	16.7	13.7	20.5
Vegetation biomass (Pg C)	674.0	544.0	829.2
Litter mass (Pg C)	97.0	79.0	108.9
Soil carbon mass (Pg C)	1410.0	1162.0	1843.0
Mean areal precipitation (mm yr <sup>-1</sup> ) (global)	760.0	762.0	828.0
Mean areal precipitation (mm yr <sup>-1</sup> ) (30°N-30°S)	1047.0	984.0	1139.0
Gross primary productivity (Pg C yr <sup>-1</sup> ) (30°N-30°S)	80.7	60.9	95.5



**Table 4.** Annual GPP,  $R_{\text{eco}}$ , and NEE ( $\text{Pg-C-PgC}\cdot\text{year}^{-1}$ ) from CTEM-CRUNCEP and CTEM-GEM for the transient simulations. The transient simulation was initialized from the spin-up simulations using varying  $\text{CO}_2$  concentrations and meteorology.

	CTEM-CRUNCEP		CTEM-GEM		other estimates
	2009	2010	2009	2010	multi-year average
GPP	133.6	137.4	99.3	100.9	119.5 (Deng et al., 2014) 123 $\pm$ 8 (Beer et al., 2010)
$R_{\text{eco}}$	130.4	132.2	98.1	98.7	
NEE	-3.2	-5.2	-1.2	-2.2	

**Table 5.** The mean differences and RMSEs (in ppm) of the a posteriori  $\text{CO}_2$  fields, based on CTEM-CRUNCEP, CTEM-GEM, and BEPS fluxes, with respect to TCCON data ~~in for July 2009 and to June 2010.~~ Station to station error is also shown.

	CTEM-CRUNCEP	CTEM-GEM	BEPS	<del>CTEM-CRUNCEP</del>	<del>CTEM-GEM</del>	<del>BEPS</del>
<del>2009</del> <u>Mean (mod – obs)</u>	<del>0.13</del> <u>0.53</u>	<del>0.27</del> <u>0.64</u>			<del>0.50</del> <u>0.49</u>	
<u>RMSE (mod – obs)</u>	<del>1.42</del> <u>1.36</u>	<del>1.24</del> <u>1.36</u>			<del>1.22</del> <u>1.23</u>	
<del>2010</del> <u>heightinter-station bias</u>	<del>0.78</del> <u>0.61</u>	<del>0.80</del> <u>0.53</u>			<del>0.54</del> <u>0.49</u>	<del>1.42</del> <u>1.39</u> <u>1.18</u>

**Table 6.** The mean differences and RMSEs (in ppm) of the a posteriori  $\text{CO}_2$  fields, based on CTEM-CRUNCEP, CTEM-GEM, and BEPS fluxes, with respect to aircraft data from the HIPPO-1, HIPPO-2, and HIPPO-3 campaigns.

<u>Altitude</u>	<u>Latitude</u>	Mean (mod – obs)			RMSE (mod – obs)		
		CTEM-CRUNCEP	CTEM-GEM	BEPS	CTEM-CRUNCEP	CTEM-GEM	BEPS
<del>HIPPO-1</del> <u>0 - 5 Km</u>	<del>-0.01</del> <u>60°S - 30°S</u>	<del>-0.11</del> <u>0.07</u>	<del>-0.20</del> <u>0.00</u>	<del>2.85</del> <u>0.02</u>	<del>2.65</del> <u>0.70</u>	<del>1.90</del> <u>0.70</u>	<u>0.74</u>
<del>HIPPO-2</del>	<del>-0.69</del> <u>30°S - 0°</u>	<u>-0.61</u>	<u>-0.58</u>	<u>-0.33</u>	<u>0.87</u>	<u>0.83</u>	<u>0.52</u>
	<u>0° - 30°N</u>	<u>-0.54</u>	<u>-0.35</u>	<u>-0.26</u>	<u>1.03</u>	<u>0.95</u>	<u>0.75</u>
	<u>30°N - 60°N</u>	<u>-0.48</u>	<u>-0.34</u>	<u>-0.68</u>	<u>1.72</u>	<u>1.72</u>	<u>1.58</u>
<u>5 - 10 Km</u>	<u>60°S - 30°S</u>	<u>-0.45</u>	<u>-0.46</u>	<u>-0.39</u>	<u>0.78</u>	<u>0.80</u>	<u>0.78</u>
	<u>30°S - 0°</u>	<u>-0.56</u>	<u>-0.44</u>	<u>-0.16</u>	<u>0.80</u>	<u>0.75</u>	<u>0.69</u>
	<u>0° - 30°N</u>	<u>-0.91</u>	<u>-0.67</u>	<del>-0.41</del> <u>-0.34</u>	<del>1.84</del> <u>1.12</u>	<del>1.77</del> <u>0.95</u>	<del>1.55</del> <u>0.74</u>
<del>HIPPO-3</del>	<del>0.26</del> <u>30°N - 60°N</u>	<del>0.16</del> <u>-0.86</u>	<del>-0.28</del> <u>-0.56</u>	<del>1.04</del> <u>-0.49</u>	<del>0.94</del> <u>1.46</u>	<del>0.87</del> <u>1.30</u>	<u>1.20</u>
<u>0 - 10 Km</u>	<u>60°S - 60°N</u>	<u>-0.56</u>	<u>-0.43</u>	<u>-0.36</u>	<u>1.17</u>	<u>1.10</u>	<u>0.99</u>