

Response to reviewers for the manuscript: Constraining DALEC v2 using multiple data streams and ecological constraints.

Sylvain Delahaies¹, Ian Roulstone¹, and Nancy Nichols²

¹Department of Mathematics, University of Surrey, Guildford, UK.

²Department of Mathematics, University of Reading, Reading, UK.

Correspondence to: S. Delahaies (s.b.delahaies@surrey.ac.uk)

We thank both referees for their thorough reading of the manuscript and for useful comments. We have addressed all comments and tried our best to clarify the manuscript. Our responses to both referees are described in the following sections. Together with this response we submit a revision of the manuscript which accounts for the changes described below.

5 **1 Response to referee 1**

1.1 Tables references

All tables references have been checked and corrected.

1.2 Range of parameters

The range of parameters in Table 1 have been corrected, they correspond to what is used in Bloom
10 and Williams (2015).

1.3 Ecological constraints

Our implementation of the EDCs leads to a set of 29 inequalities denoted EDC_1 to EDC_{29} , we omitted to take into account the carbon pools growth constraints in the original manuscript.

In the original manuscript we chose to provide only a heuristic description of the inequalities, the
15 complete description of which can be found in Bloom and Williams (2015). We thought that although justified for the sake of self consistency, detailing the EDCs did not bring any insight into the question addressed here and increased significantly unnecessary mathematical notation. Nonetheless we acknowledge the comment made by referee 1 and we have made substantial changes to section

2.2 to incorporate a complete description of the EDCs, running from line 123 to line 180 in the
20 revised manuscript.

1.4 Other comments

A discussion section has been added to the manuscript to address the remaining comments.

Global scale experiments are discussed in reference to Rayner et al. (2005) and CCDAS. The work
of Kemp et al. (2014), which directly relates to what we discuss in our paper, is also cited. Although
25 this preliminary work is not reported in our manuscript the three approaches presented in Kemp et al.
(2014) were evaluated for the preparation of the manuscript, and our conclusion was different: in our
case incorporating the EDCs by adding a penalty term to the cost function was the most successful
approach to constrain unresolved parameters and, most importantly in our case, to allow for a better
uncertainty quantification.

30 We also refer to Ziehn et al. (2012) for their comparison between MCMC method and 4DVAR in
CCDAS. A MCMC method is used in Bloom and Williams (2015), and in Safta et al. (2015), added
in reference to the revised manuscript, a detailed analysis of MCMC for DALEC is performed. A
comparison between MCMC and 4DVAR was beyond the scope of this paper, our intention was
rather to establish 4DVAR as a suitable method for DALEC and the EDCs. Nonetheless a compari-
35 son between 4DVAR and fully non-linear methods is necessary, it is one of the aspects of our current
work.

Finally we mention our current work on a hybrid ensemble-variational method. This approach pro-
vides an adjoint-free formulation of the variational problem and show promising results in the con-
text discussed in the manuscript. This work is part of a paper in preparation.

40 2 Response to referee 2

The version of DALEC used for this study, referred to as DALECV2 and described in details in
Bloom and Williams (2015), is a nonlinear dynamical system, it is not a linear autonomous system.
The trajectories of the carbon pools \mathbf{C} are computed using the recursion formula

$$\mathbf{C}^{t+1} = \mathbf{C}^t + \mathbf{f}(\mathbf{C}^t, \mathbf{p}, \phi(t))\Delta t, \quad (1)$$

45 where \mathbf{f} is a nonlinear vector valued function of the carbon pools, the parameters \mathbf{p} and the meteo-
rological drivers $\phi(t)$, Δt denoting the step time in month in our case. The nonlinear nature of the
model is stressed out in the revised manuscript at lines 77 and 83. Moreover, the definition of \mathbf{h} is
clarified at line 197.

The main focus of the paper is on the vector $\mathbf{x} = \log([\mathbf{p}, \mathbf{C}_0])^T$. In section 2.3 first where we investi-
50 gate the sensitivity of different outputs with respect to \mathbf{x} and its components, and then in subsequent
sections where \mathbf{x} is estimated using inverse methods. The vector \mathbf{x} , denoting fixed quantities as ini-
tial conditions and parameters for the dynamical system DALECV2, is seen as the variable from the

point of view of sensitivity analysis and inverse modelling and therefore its components are referred to as state variables, input variables or parameters interchangeably throughout the manuscript. This
55 choice of terminology, stressed out at line 228, have been reinforced in the revised manuscript by adding the present paragraph at line 103.

References

- Bloom, A. A. and Williams, M.: Constraining ecosystem carbon dynamics in a data-limited world: integrating ecological "common sense" in a model–data fusion framework, *Biogeosciences*, 12, 1299–1315, doi:10.5194/bg-12-1299-2015, 2015.
- 60 Kemp, S., Scholze, M., Ziehn, T., and Kaminski, T.: Limiting the parameter space in the Carbon Cycle Data Assimilation System (CCDAS), *Geoscientific Model Development*, 7, 1609–1619, doi:10.5194/gmd-7-1609-2014, <http://www.geosci-model-dev.net/7/1609/2014/>, 2014.
- Rayner, P. J., Scholze, M., Knorr, W., Kaminski, T., Giering, R., and Widmann, H.: Two decades of terrestrial carbon fluxes from a carbon cycle data assimilation system (CCDAS), *Global Biogeochemical Cycles*, 19, n/a–n/a, doi:10.1029/2004GB002254, <http://dx.doi.org/10.1029/2004GB002254>, gB2026, 2005.
- 65 Safta, C., Ricciuto, D. M., Sargsyan, K., Debusschere, B., Najm, H. N., Williams, M., and Thornton, P. E.: Global sensitivity analysis, probabilistic calibration, and predictive assessment for the data assimilation linked ecosystem carbon model, *Geoscientific Model Development*, 8, 1899–1918, doi:10.5194/gmd-8-1899-2015, <http://www.geosci-model-dev.net/8/1899/2015/>, 2015.
- 70 Ziehn, T., Scholze, M., and Knorr, W.: On the capability of Monte Carlo and adjoint inversion techniques to derive posterior parameter uncertainties in terrestrial ecosystem models, *Global Biogeochemical Cycles*, 26, n/a–n/a, doi:10.1029/2011GB004185, <http://dx.doi.org/10.1029/2011GB004185>, gB3025, 2012.

Constraining DALEC v2 using multiple data streams and ecological constraints: analysis and application.

Sylvain Delahaies¹, Ian Roulstone¹, and Nancy Nichols²

¹Department of Mathematics, University of Surrey, Guildford, UK.

²Department of Mathematics, University of Reading, Reading, UK.

Correspondence to: S. Delahaies (s.b.delahaies@surrey.ac.uk)

Abstract. We use a variational method to assimilate multiple data streams into the terrestrial ecosystem carbon cycle model DALECv2. Ecological and dynamical constraints have recently been introduced to constrain unresolved components of this otherwise ill-posed problem. Here we recast these constraints as a multivariate ~~gaussian~~ Gaussian distribution to incorporate them into the variational framework and we demonstrate their benefit through a linear analysis. Using an adjoint method we study a linear approximation of the inverse problem: firstly we perform a sensitivity analysis of the different outputs under consideration, and secondly we use the concept of resolution matrices to diagnose the nature of the ill-posedness and evaluate regularisation strategies. We then study the non linear problem with an application to real data. Finally, we propose a modification to the model: introducing a spin-up period provides us with a built-in formulation of some ecological constraints which facilitates the variational approach.

1 Introduction

Carbon is a fundamental constituent of life and understanding its global cycle is a key challenge for the modelling of the Earth system. Through the processes of photosynthesis and respiration, ecosystems play a major role in the carbon cycle and thus in the dynamics of the global climate system. Our knowledge of the biogeochemical processes of ecosystems and an ever-growing amount of Earth observation systems can be combined using inverse modelling strategies to improve model predictions and uncertainty quantification.

The data assimilation linked ecosystem model (DALEC) is a simple box model for terrestrial ecosystems simulating a large range of processes occurring at different time scales from days to millennia. The work of Williams et al. (2005) established the benefit of using DALEC together with net ecosystem exchange of CO_2 (NEE) measurements in a Bayesian framework to estimate initial carbon stocks and model parameters, to improve flux predictions for ecosystem models, and to quantify un-

certainties. ~~Intereomparison~~ Inter-comparison experiments (Fox et al. (2009); Hill et al. (2012)) have
25 then demonstrated the relative merit of various inverse modelling strategies using NEE and MODIS
leaf area index observations: most results agreed on the fact that parameters and initial stocks directly
related to fast processes were best estimated with narrow confidence intervals, whereas those related
to slow processes were poorly estimated with very large uncertainties. Other studies have tried to
overcome this difficulty by adding complementary data streams, see Richardson et al. (2010), or
30 by considering longer observation windows, see Hill et al. (2012). Recently Bloom and Williams
(2015) defined a set of ecological and dynamical constraints (EDCs) to reject unrealistic parameter
combinations in the absence of additional data. However, to date ~~no~~ very few systematic analysis
has been carried out to explain the large differences among results.

As with many inverse problems, assimilating Earth observations into DALEC is an ill-posed prob-
35 lem: the model-observation operator which relates parameters and initial carbon stocks to the obser-
vations is rank deficient and not all variables can be estimated, or the model-observation operator is
ill-conditioned and small observational noise may lead to a solution we can have little confidence
in. Solving the problem amounts first to transforming it into a tractable problem in order to ensure
a robust, meaningful and stable solution. This can be achieved by using regularisation techniques;
40 the most popular one involves combining the observations and prior information, assuming it exists,
through Bayesian inference. The choice of regularisation method depends on the nature of the prob-
lem and on the inverse modelling approach adopted.

So far, off-the-shelf methods such as ensemble Kalman filter (EnKF) and Monte Carlo Markov Chain
(MCMC) were adopted to perform model-data fusion with DALEC. For its ability to ~~accomodate~~
45 accommodate non-linearity and any kind of probability distributions, the MCMC method, in the limit
of a large number of samples, may be considered as the gold standard. However, ~~despite~~ despite be-
ing well suited for this type of small scale problem, the computational complexity of MCMC method
makes it ~~untractable~~ intractable for more complex situations. Here we adopt a variational approach
(4DVAR) where a cost function measuring the mismatch between the model and observations is
50 ~~minimized~~ minimised using a gradient method based on the adjoint of the model. At Ameriflux sites
(see <http://ameriflux.lbl.gov/>), we use MODIS monthly mean leaf area index (LAI) observations
over a 13 year time window together with flux tower measurements of NEE and gross primary pro-
duction (GPP). 4DVAR facilitates the diagnosis of the ill-posedness of the inverse problem: using
~~resolution matrices, successfully adapted from the field of tomography,~~ model resolution matrices
55 we can assess the resolution and stability properties of the observation operators and of the regu-
larisation terms. We transcribe the EDCs into a novel variational framework and use some of this
additional knowledge to estimate the otherwise undetermined variables. We consider a modification
of the DALEC model by adding a spin-up period where carbon stocks are brought to equilibrium,
this offers an alternative to including all the EDCs and helps reducing the confidence intervals for
60 the predicted fluxes.

The paper is organised as follows. In section 2 we present DALECV2 and the observation streams used in this study; we ~~briefly~~ review the EDCs introduced in Bloom and Williams (2015), and we perform a sensitivity analysis of the different outputs of DALECV2 of interest for our experiments. In section 3 we recall basic principles of inverse theory from a Bayesian perspective, we introduce
65 the variational formulation and we show how to incorporate the EDCs into this framework. Section 4 is devoted to a résumé of the linearised problem, using the tangent linear model, where the challenges of ill-posed problems and their regularisation can be explored in detail using simple linear algebra. Using a singular value decomposition we illustrate the effect of observational noise on ~~ill-conditioned~~ ill-conditioned systems, and we investigate solution strategies from the point of
70 view of resolution matrices. In section 5 we conduct a series of nonlinear inverse modelling experiments using multiple data streams and EDCs. In section 6 we modify DALECV2 to include a spin-up period which offers a built-in formulation of some EDCs, and then we reproduce the nonlinear experiments. ~~Finally in~~ In section 7 we discuss several extension to our manuscript and finally in section 8 we draw conclusions.

75 2 Model, constraints and observations

2.1 DALECV2

DALECV2 depicts a terrestrial ecosystem as a set of six carbon pools (labile C_{lab} , foliar C_f , wood C_w , root C_r , litterfall C_l and soil organic matter C_s) linked via allocation fluxes. At a monthly time step the gross primary production (GPP) is calculated using the Aggregated canopy model (Williams et al. (1997)) as a nonlinear function of meteorological drivers (temperature, radiation, atmospheric
80 ~~CO₂~~ CO₂ concentration), foliar carbon and foliar nitrogen. Following a mass conservation principle GPP is then allocated to the different carbon pools or released in the atmosphere via respiration. The schematic for DALECV2 is represented in Figure 1 and a complete description of the model can be found in Bloom and Williams (2015). DALECV2 combines the two previous DALEC-evergreen and
85 DALEC-deciduous into a single model where the non-differentiable phenology process of DALEC-deciduous has been replaced with a differentiable process. DALECV2 is a nonlinear dynamical system and the carbon pools are dynamical variables ~~parametrized~~ parametrised by their initial values C_0 and by 17 parameters p whose range and description can be found in Table 1. The magnitudes and ranges of the parameters and the initial values vary drastically; therefore to avoid the compu-
90 tational problems caused by these different scales the variational methods will be formulated and implemented in terms of the log transformed ~~variables~~ variable $x = \log([p, C_0])^T$. However in order to limit unnecessary notation and definition, in the remainder of this paper p and C_0 will stand for their log transform.

The meteorological drivers are extracted from 0.125°x0.125°ERA-interim reanalysis datasets.
95 For the purpose of our inverse modelling experiments we use four different observation streams:




Figure 1. DALECV2 links the carbon pools (C) via allocation fluxes (green), litterfall fluxes (red), **decomposition** **decomposition** (black). Respiration is represented by the blue arrows. The orange arrow represents the feedback of foliar carbon to gross primary production (GPP).

Table 1. DALECV2 dynamical variables and parameters with their respective range. The units of the non dimensionless quantities are given in brackets.

Label	Variable	description	range
$C_0(1)$	C_{lab}	initial labile C pool (gCm^{-2})	20 - 2000
$C_0(2)$	C_f	initial foliar C pool (gCm^{-2})	20 - 2000
$C_0(3)$	C_r	initial fine root C pool (gCm^{-2})	20 - 2000
$C_0(4)$	C_w	initial above and below ground woody C pool (gCm^{-2})	100 - 10^5
$C_0(5)$	C_l	initial litter C pool (gCm^{-2})	20 - 2000
$C_0(6)$	C_s	initial soil organic matter C pool (gCm^{-2})	100 - 2×10^5
p_1	θ_{min}	Litter mineralisation rate (day^{-1})	10^{-5} - 10^{-2}
p_2	f_a	autotrophic respiration fraction	0.3 - 0.7
p_3	f_f	fraction of GPP allocated to C_f	0.01 - 0.5
p_4	f_r	fraction of GPP allocated to C_r	0.01 - 0.5
p_5	c_{lf}	Annual Leaf Loss Fraction (season)	1 - 8
p_6	θ_w	C_w turnover rate (day^{-1})	2.5×10^{-5} - 10^{-3}
p_7	θ_r	C_r turnover rate (day^{-1})	10^{-4} - 10^{-2}
p_8	θ_l	C_l turnover rate (day^{-1})	10^{-4} - 10^{-2}
p_9	θ_s	C_s turnover rate (day^{-1})	10^{-7} - 10^{-3}
p_{10}	Θ	temperature dependence exponent factor	0.018 - 0.08
p_{11}	c_{eff}	Canopy Efficiency Parameter	10 - 100
p_{12}	d_{onset}	Leaf Onset Day (day)	1 - 365
p_{13}	f_l	fraction of GPP allocated to C_{lab}	0.01 - 0.5
p_{14}	c_{ronset}	C_{lab} release period (day)	10 - 100
p_{15}	d_{fall}	Leaf Fall Day (day)	1 - 365
p_{16}	c_{rfall}	Leaffall period (day)	20 - 150
p_{17}	c_{lma}	Leaf Mass per area (gCm^{-2})	10 - 400

LAI, NEE, GPP and RESP. LAI monthly mean observations for Ameriflux sites are extracted from MOD15A2 LAI 8-day version 005 1km-resolution product. These observations together with the meteorological drivers are provided by A. Bloom and J. Exbrayat: details about their construction can be found in Bloom and Williams (2015). At Ameriflux sites we use the level 4 data product
100 (available at <http://cdiac.ornl.gov/ftp/ameriflux/data/Level4/>), which provides monthly means for NEE and GPP. NEE and GPP are then used to define total respiration (RESP) as $RESP=NEE+GPP$. The meteorological drivers span over a period of twelve years from 2001 to 2013. LAI observations are available during the full period but for NEE and GPP, and thus RESP, shorter ~~reecords~~-records are available depending on the Ameriflux site. In this study we consider the Morgan Monroe state
105 forest located in Indiana, US (39.3,-86.4). This Ameriflux site is composed in majority of mixed hardwood broadleaf deciduous trees and classifies as a humid subtropical climate.

In the remainder of the paper the main focus is on the vector $x = \log([p, C_0])^T$: in section 2.3 first where we investigate the sensitivity of different outputs with respect to x and its components, and then in subsequent sections where x is estimated using inverse methods. The vector x , denoting
110 fixed quantities as initial conditions and parameters for the dynamical system DALECV2, is seen as the variable from the point of view of sensitivity analysis and inverse modelling and therefore its components will be referred to as state variables, input variables or parameters interchangeably throughout the manuscript.

2.2 Ecological constraints

115 Over the last decade many inverse ~~modeling~~-modelling studies have used NEE measurements from the fluxnet network, together with other types of observations when available, to provide information about processes controlled by parameters with respect to which NEE is weakly sensitive. Though it contains an ~~ever-ineresing~~-ever-increasing amount of information, the flux tower network only provides sparse coverage of terrestrial ecosystems. On the other hand, despite a good spatial and temporal coverage, MODIS LAI monthly mean observations only constrain a limited set of DALECV2
120 state variables, and additional information is required in order to ~~regularize~~-regularise the ill-posed problem and obtain a meaningful solution.

Additional information can be obtained by imposing priors on the variables or by adding other observation streams (biomass, soil organic matter, ...). As an alternative, Bloom and Williams introduced
125 a set of constraints, referred to as ecological and dynamical constraints (EDCs). These constraints, detailed in Bloom and Williams (2015) can be divided into two groups: static and ~~dynamieie~~-dynamic constraints. The static constraints which directly impose conditions on the parameters are:

- turnover rates constraints which ensure that turnover rates ratios are consistent with knowledge of the carbon pools residence times.

$$130 \quad \underline{EDC_1 : p_9 < p_8}, \quad (1)$$

$$\underline{EDC_2 : p_9 < p_1}, \quad (2)$$

$$\underline{EDC_3 : p_6 < 1/(p_5 \times 365.25)}, \quad (3)$$

$$\underline{EDC_4 : p_7 > p_9 \exp p_{10} \bar{T}}, \quad (4)$$

$$135 \quad \underline{EDC_5 : p_{12} + 45 < p_{15}}, \quad (5)$$

where \bar{T} denotes the mean temperature within the drivers time window. EDC_4 is a modification to the constraint proposed in Bloom and Williams (2015), it is currently used in the CARDAMON framework (<http://www.geos.ed.ac.uk/homes/mwilliam/CARDAMOM.html>).

- Root-foliar allocation which allows for a strong correlation between parameters controlling allocation to foliage and roots.

$$140 \quad \underline{EDC_6 : f_{\text{root}} < 5(f_{\text{fol}} + f_{\text{lab}})}, \quad (6)$$

$$\underline{EDC_7 : f_{\text{fol}} + f_{\text{lab}} < 5f_{\text{root}}}, \quad (7)$$

where the allocation fractions f_{fol} , f_{lab} and f_{root} are defined by

$$145 \quad \underline{f_{\text{auto}} = p_2}, \quad (8)$$

$$\underline{f_{\text{fol}} = (1 - f_{\text{auto}})p_3}, \quad (9)$$

$$\underline{f_{\text{lab}} = (1 - f_{\text{auto}} - f_{\text{fol}})p_{13}}, \quad (10)$$

$$\underline{f_{\text{root}} = (1 - f_{\text{auto}} - f_{\text{fol}} - f_{\text{lab}})p_4}. \quad (11)$$

150 The dynamic constraints, for which a model run is performed to define attractors, limit the application of the model to ecosystems with no major recent disturbance. They are defined by:

- Root-foliar mean dynamics

$$\underline{EDC_8 : \bar{C}_r < 5\bar{C}_f}, \quad (12)$$

$$155 \quad \underline{EDC_9 : \bar{C}_f < 5\bar{C}_r}, \quad (13)$$

where \bar{C}_f and \bar{C}_r denote the mean of C_f and C_r over the simulation period.

- Yearly carbon pools growth rate is limited to 10%.

$$\underline{EDC_{10-15} : \bar{C}^n / \bar{C}^1 < 1 + \zeta(n-1)/10}, \quad (14)$$

160 where for each pool \bar{C}^i denotes the mean carbon pool size over year i and the growth factor ζ
is set to 0.1.

- Carbon pools are not expected to show rapid exponential decay; therefore parameter sets are required to satisfy the condition that the half-life period of carbon pools is more than three years.

165 EDC_{16–21} : $\gamma < 3 \times 365 / \log 2$. (15)

The trajectory of each carbon pool is approximated using an exponential decay curve $a + b \exp \gamma t$
where a, b and γ are the fitted exponential decay parameters and t the time variable, in days in
this case.

- 170 – Carbon pools are expected to be within an order of magnitude of a steady state attractor.
~~Our implementation of the EDCs leads to a set of 23 inequalities, denoted EDC₁ to EDC₂₃,~~
~~constraining all state variables: 7 for the static EDCs constraining only p_1 to p_{10} , p_{12} , p_{13} and~~
 ~~p_{15} .~~

175 EDC_{22–29} : $C_0/10 < C^\infty < 10C_0$, (16)

where for each of the carbon pools C_s, C_l, C_w and C_r, C_0 denotes the initial state and C^∞
denotes the steady state attractor defined by

$$C_{\text{som}}^\infty = \frac{(f_{\text{wood}} + (f_{\text{fol}} + f_{\text{root}} + f_{\text{lab}})p_1)\bar{G}}{(p_1 + p_9)p_8 \exp \bar{T}p_{10}},$$
 (17)

$$C_{\text{lit}}^\infty = \frac{(f_{\text{fol}} + f_{\text{root}} + f_{\text{lab}})\bar{G}}{p_9 \exp \bar{T}p_{10}},$$
 (18)

180
$$C_{\text{wood}}^\infty = \frac{f_{\text{wood}}\bar{G}}{p_6},$$
 (19)

$$C_{\text{root}}^\infty = \frac{f_{\text{root}}\bar{G}}{p_7},$$
 (20)

where \bar{G} denotes the mean gross primary production and $f_{\text{wood}}, f_{\text{som}}$ and ~~16 for the dynamic~~
~~EDCs constraining all state variables.~~ f_{lit} are given by

185
$$f_{\text{wood}} = 1 - f_{\text{auto}} - f_{\text{fol}} - f_{\text{lab}} - f_{\text{root}},$$
 (21)

$$f_{\text{som}} = f_{\text{wood}} + (f_{\text{root}} + f_{\text{lab}} + f_{\text{fol}})p_1 / (p_1 + p_8),$$
 (22)

$$f_{\text{lit}} = (f_{\text{root}} + f_{\text{lab}} + f_{\text{fol}}).$$
 (23)

To the original EDCs, we found useful to add the three following constraints:

190 $\text{EDC}_{24} : \text{LAI}(\text{summer}) < \alpha, \alpha > 0,$

$\text{EDC}_{25} : \text{LAI}(\text{final day}) > 0,$

$\text{EDC}_{26,27} : -\beta < \text{E}[\text{NEE}] < \beta, \beta > 0,$

$\text{EDC}_{30} : \text{LAI}(\text{summer}) < \alpha, \alpha > 0,$ (24)

195 $\text{EDC}_{31} : \text{LAI}(\text{final day}) > 0,$ (25)

$\text{EDC}_{32,33} : -\beta < \text{E}[\text{NEE}] < \beta, \beta > 0,$ (26)

where α and β are real constants that need to be adjusted, $\text{LAI}(\text{summer})$ denotes the modelled LAI during summer and $\text{LAI}(\text{final day})$ denotes the modelled LAI at the end of the model run. These new constraints guarantee that LAI and the mean NEE remain between realistic bounds.

Bloom and Williams demonstrated the efficiency of incorporating EDCs using a Monte Carlo method to improve parameter estimates and NEE predictions. We propose an approach to apply these extra constraints within a variational framework.

2.3 Sensitivity analysis

205 Sensitivity analysis studies how the variations of the output \mathbf{h} of a model can be attributed to ~~variations~~ variations of the input variables x_i . Such information is crucial for model design, inverse modelling and reduction of complex nonlinear models. A global sensitivity analysis for DALEC was recently performed in Safta et al. (2015), here we consider a local approach where first order derivatives are used to build sensitivity indices that help understanding the influence of input variables
 210 on the output.

~~Following Zhu and Zhuang (2014), we consider the mean normalized sensitivity (MNS) defined by~~

$$s_i = \text{E} \left(\frac{\frac{\partial \mathbf{h}}{\partial x_i} \left| \frac{\sigma_i}{\sigma_h} \right|}{\sum_j \left| \frac{\partial \mathbf{h}}{\partial x_j} \frac{\sigma_j}{\sigma_h} \right|} \right),$$

~~where \mathbf{h} denotes~~ We denote by h_t the function that maps $\mathbf{x} = \log(\mathbf{p}, \mathbf{C}^0)$ to the value of an output of the model (here LAI, NEE, GPP and RESP) at time t and we denote by $\mathbf{h} = (h_{t_1}, \dots, h_{t_n})$

215 the time series of the model output(LAI and NEE in the following example), and. Following Zhu and Zhuang (2014), we consider the mean normalised sensitivity (MNS) defined by

$$s_i = \text{E} \left(\frac{\frac{\partial \mathbf{h}}{\partial x_i} \left| \frac{\sigma_i}{\sigma_h} \right|}{\sum_j \left| \frac{\partial \mathbf{h}}{\partial x_j} \frac{\sigma_j}{\sigma_h} \right|} \right), \quad (27)$$

where $\text{E}(\cdot)$ denotes the average of the time series. The scalars σ_i and σ_h denote the parameter variance, set as 40% of the parameter range, and the variance of the output respectively. The partial




Figure 2. Mean ~~normalized~~ normalised sensitivities : 100 parameter sets satisfying EDCs are sampled at the Morgan Monroe State Forest. Parameters are ranked in decreasing order according to their sensitivity, the blue dots represent the mean of the MNS (dimensionless quantity) and the intervals represent 1-sigma error bars; the red dots correspond to null sensitivity.

220 derivatives are computed using the adjoint derived using the method described in Giering and Kaminski (1998). The MNS s_i is a dimensionless number that allows us to compare between parameters. We consider the Morgan Monroe State Forest over a thirteen year period. We sample 100 parameter sets satisfying the ecological constraints. For each parameter set we compute the MNS for DALEC simulated mean fluxes LAI, NEE. In Figure 2 parameters are ranked with respect to their mean

225 MNS. We see that for LAI only 12 out of the 23 variables are sensitive, namely $p_5, p_{17}, p_2, p_{13}, p_{11}, p_{15}, p_{16}, C_f, p_{12}, p_3, C_{lab}$ and p_{14} . Therefore using LAI only in an inverse ~~modeling~~ modelling experiment provides, at best, information about those twelve sensitive variables. For NEE we see that all variables are sensitive. Sensitivity analysis for GPP shows similar characteristics with LAI and so does RESP with NEE. For the four outputs under consideration (LAI, NEE, GPP and RESP)

230 the most sensitive variables are the autotrophic respiration, p_2 , the annual leaf loss fraction, p_5 , the leaf mass per area, p_{17} , the fraction of GPP ~~allocated~~ allocated to labile pool, p_{13} , the nitrogen use efficiency, p_{11} , and the leaf fall day p_{15} .

Here our focus is on the mean of the time series of DALEC fluxes (LAI, NEE) over a thirteen year period. Finer analysis could be carried out by looking at seasonal aspects of the carbon cycle, identifying what variables are the most sensitive ~~in a growing or in a decaying season for example at~~ certain time of the year for example as studied in Safta et al. (2015).

3 Data assimilation

In this section we introduce concepts and methods that allow for a close mathematical scrutiny of inverse problems and we present the variational method that will be used for applications.

240 3.1 Ill posed problem

A generic inverse problem consists in finding a n -dimensional state vector \mathbf{x} such that

$$\mathbf{h}(\mathbf{x}) = \mathbf{y}, \quad (28)$$

for a given N -dimensional observation vector \mathbf{y} , including random noise, and a given model \mathbf{h} . In the remainder of the paper the terms state vector, state variable, input variable or parameters will
245 be used interchangeably to denote the vector \mathbf{x} to be estimated using inverse methods and defined in the previous section as $\mathbf{x} = \log([\mathbf{p}, \mathbf{C}])^T$. The problem is well posed in the sense of Hadamard (1923) if the three following conditions hold: 1) there exists a solution, 2) the solution is unique and 3) the solution depends continuously on the input data. If at least one of these conditions is violated the problem is said to be ill-posed. The inverse problem (28) is often ill-posed, and a ~~regularization~~
250 ~~regularisation~~ method is required to replace the original problem with a well-posed-posed problem. Solving (28) amounts to 1) constructing a solution \mathbf{x} , 2) assessing the validity of the solution, 3) ~~characterizing~~-~~characterising~~ its uncertainty. Each inverse problem has its own features which need to be understood in order to ~~characterize~~-~~characterise~~ properly the solution and its uncertainty.

255 3.2 Bayesian inference: 4DVAR

Inverse problems are generally presented in a probabilistic framework where most methods can be expressed through a Bayesian formulation. The Bayesian approach provides a full characterisation of all possible solutions, their relative probabilities and uncertainties.

From Bayes' theorem the probability density function (pdf) of the model state \mathbf{x} given the set of
260 observations \mathbf{y} , $p(\mathbf{x}|\mathbf{y})$, is given by

$$p(\mathbf{x}|\mathbf{y}) \propto p(\mathbf{y}|\mathbf{x})p(\mathbf{x}), \quad (29)$$

where $p(\mathbf{y}|\mathbf{x})$ is the pdf of the observations given \mathbf{x} and $p(\mathbf{x})$ is the prior pdf of \mathbf{x} . A special case is given when $p(\mathbf{y}|\mathbf{x})$ and $p(\mathbf{x})$ are ~~gaussian~~-~~Gaussian~~ pdf given by

$$p(\mathbf{x}) = \exp \left[-\frac{1}{2}(\mathbf{x} - \mathbf{x}_0)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_0) \right], \quad (30)$$

265 and

$$p(\mathbf{y}|\mathbf{x}) = \exp \left[-\frac{1}{2}(\mathbf{h}(\mathbf{x}) - \mathbf{y})^T \mathbf{R}^{-1}(\mathbf{h}(\mathbf{x}) - \mathbf{y}) \right], \quad (31)$$

where \mathbf{B} is the covariance matrix of the prior term \mathbf{x}_0 , and \mathbf{R} is the covariance matrix of the observation error. When the operator \mathbf{h} is linear then the posterior pdf $p(\mathbf{x}|\mathbf{y})$ is ~~gaussian~~-~~Gaussian~~ and thus fully ~~characterized~~-~~characterised~~ by its mean and covariance matrix. The mean is obtained by
270 ~~minimizing~~-~~minimising~~ the modulus of the log of the joint probability distribution, that is the cost

function J given by

$$J(\mathbf{x}) = J_0(\mathbf{x}) + J_y(\mathbf{x}) = \frac{1}{2}\|\mathbf{x} - \mathbf{x}_0\|_{\mathbf{B}}^2 + \frac{1}{2}\|\mathbf{h}(\mathbf{x}) - \mathbf{y}\|_{\mathbf{R}}^2. \quad (32)$$

Many methods can be considered to ~~minimize~~ minimise this cost function. A Monte Carlo method is employed in Bloom and Williams (2015). Here we use a variational approach which applies a
 275 gradient based method where the gradient is given by

$$\nabla J = \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_0) + \mathbf{H}^T \mathbf{R}^{-1}(\mathbf{h}(\mathbf{x}) - \mathbf{y}), \quad (33)$$

with \mathbf{H}^T denoting the adjoint operator. The covariance matrix of the solution, \mathbf{C} , is given by the inverse of the hessian of the cost function

$$\mathbf{C} = [\text{Hess}(J)]^{-1} = [\mathbf{B}^{-1} + \mathbf{H}^T \mathbf{R}^{-1} \mathbf{H}]^{-1}. \quad (34)$$

280 When the observation operator \mathbf{h} is non-linear, the cost function J can have multiple local minima and the posterior pdf may no longer be a ~~gaussian~~ Gaussian pdf. However, locally, the pdf $N(\tilde{\mathbf{x}}, \mathbf{C})$, where \mathbf{C} is given by equation (34) evaluated at a minimum $\tilde{\mathbf{x}}$, provides a ~~gaussian~~ Gaussian approximation of the posterior pdf $p(\mathbf{x}|\mathbf{y})$.

The first term in the cost function (32) is a regularisation term encoding the ~~gaussian~~ Gaussian
 285 prior $p(\mathbf{x})$. As we will show in the next sections the problem of assimilating EO observations (LAI,GPP,NEE,RESP) into DALEC is a highly ill-posed problem and regularisation is required. The sensitivity analysis of section 2.3 showed that LAI and GPP are not sensitive to all variables. Moreover all observations streams show very small sensitivities to some variables. Therefore, as will be illustrated in section 4.1, the solution (if any) is likely to be subject to large uncertainties. Apart
 290 from a couple of extensively studied sites, our prior knowledge about the variables is so far limited to their upper and lower bounds given in Table 1. As performed in Zhu and Zhuang (2014), it is a common practice to use this information to define a ~~gaussian~~ Gaussian prior $p(\mathbf{x}) \sim N(\mathbf{x}_0, \mathbf{B})$, where \mathbf{x}_0 is given by the ~~center~~ centre of the variables ranges and \mathbf{B} is the diagonal matrix whose diagonal elements are the squares of 40% of the variables ranges. While using this kind of regulari-
 295 sation is necessary to ensure any solution at all when no better source of information is available, this introduces some ~~biaises~~ biases in the solution. The EDCs introduced by Bloom and Williams (2015) provide new prior information about the variables. One of the purposes of this paper is to incorporate the EDCs as a regularisation term within 4DVAR. In the next section we propose a strategy to achieve this goal.

300 3.3 EDCs and 4DVAR

Incorporating the EDCs from an optimisation point of view can be easily performed by considering an inequality constraint optimisation problem where we aim at solving

$$\min_{\mathbf{x}} J_y(\mathbf{x}) \quad \text{subject to} \quad \mathbf{l} < \mathbf{x} < \mathbf{u} \quad \text{and} \quad \mathbf{g}(\mathbf{x}) < \mathbf{0},$$




Figure 3. distribution and ~~gaussian~~-Gaussian fit for EDC₄ and EDC₆.

where g is the nonlinear operator defining the EDCs described in Section 2.2, and l and u are the
 305 lower and upper bounds defined in Table 1. This approach provides an efficient, robust and quick
 strategy to find an acceptable solution; however, stability properties are not easily determined, see
 Roesse-Koerner et al. (2012).

We are seeking for a multivariate ~~gaussian~~-Gaussian distribution that would encode the EDCs. At a
 forest site, we start by sampling the parameter space to obtain an ensemble of 1000 parameter sets
 310 satisfying the EDCs; each parameter set \mathbf{x} is randomly created and required to satisfy $g(\mathbf{x}) < \mathbf{0}$.
 We denote this ensemble by $\mathcal{X}_{\text{EDCs}}$. For most parameters, the sampling gives rise to undetermined
 pdfs which can certainly not be represented by ~~gaussian~~-Gaussian pdfs. However inspecting the
 distribution $g(\mathbf{x})$, for all \mathbf{x} in $\mathcal{X}_{\text{EDCs}}$, we see that the distribution $\log(g(\mathbf{x}))$ can be fairly accurately
 approximated by a multivariate ~~gaussian~~-Gaussian pdfs $N(\mathbf{c}, \Sigma)$ where \mathbf{c} denotes the mean of the
 315 distribution $\log(g(\mathbf{x}))$ and Σ denotes its covariance matrix. As an example Figure 3 shows the
 marginals ~~log(g₂(x)) and log(g₃(x))~~ log(g₄(x)) and log(g₆(x)), corresponding to the EDCs 4 and
 6 respectively, together with a ~~gaussian~~-Gaussian fit.

Using Bayes' Theorem we can then write

$$p(\mathbf{x}|\mathbf{y}, \mathbf{c}) \propto p(\mathbf{y}|\mathbf{x})p(\mathbf{c}|\mathbf{x})p(\mathbf{x}). \quad (35)$$

320 Finding a ~~gaussian~~-Gaussian approximation for $p(\mathbf{x}|\mathbf{y}, \mathbf{c})$ amounts then to minimising the cost func-
 tion

$$J(\mathbf{x}) = \frac{1}{2} \|\mathbf{h}(\mathbf{x}) - \mathbf{y}\|_{\mathbf{R}}^2 + \frac{1}{2} \|\log(g(\mathbf{x})) - \mathbf{c}\|_{\Sigma}^2 + \frac{1}{2} \|\mathbf{x} - \mathbf{x}_0\|_{\mathbf{B}}^2, \quad (36)$$

The gradient of J is given by

$$\begin{aligned} \nabla J(\mathbf{x}) = & \mathbf{H}^T \mathbf{R}^{-1} (\mathbf{h}(\mathbf{x}) - \mathbf{y}) \\ & + \frac{1}{g(\mathbf{x})} \mathbf{G}^T \Sigma^{-1} (\log(g(\mathbf{x})) - \mathbf{c}) + \mathbf{B}^{-1} (\mathbf{x} - \mathbf{x}_0), \end{aligned} \quad (37)$$

and the hessian of the cost function can be approximated by

$$\mathcal{H} = \mathbf{H}^T \mathbf{R}^{-1} \mathbf{H} + \frac{1}{(g(\mathbf{x}))^2} \mathbf{G}^T \Sigma^{-1} \mathbf{G} + \mathbf{B}^{-1}, \quad (37)$$

evaluated at the minimiser $\tilde{\mathbf{x}}$. The operator \mathbf{G}^T denotes the adjoint of the tangent linear model \mathbf{G} whose key ingredient is given by the adjoint of DALECV2. The approximation of $p(\mathbf{x}|\mathbf{y}, \mathbf{c})$ is then given by the ~~gaussian~~-Gaussian distribution $N(\tilde{\mathbf{x}}, \mathcal{H}^{-1})$. In Section 5 we will perform experiments using real data to validate this approach.

4 Linear analysis

~~Consirable~~-Considerable theoretical insights into the nature of the inverse problem, and the ill-posedness, can be obtained by studying a linearisation of the operator \mathbf{h} . A first approximation to the inverse problem consists in finding a perturbation \mathbf{z} which best ~~satisfies~~-satisfies the linear equation

$$\mathbf{H}\mathbf{z} = \mathbf{d}, \quad (38)$$

where \mathbf{H} is the tangent linear operator for \mathbf{h} and \mathbf{d} is a perturbation of the observations. The linear operator \mathbf{H} is commonly referred to as the *observability matrix* (see Johnson et al. (2005)). The least squares formulation of this problem is to solve the ~~optimization~~-optimisation problem

$$\min_{\mathbf{z}} J(\mathbf{z}) = \min_{\mathbf{z}} \frac{1}{2} \|\mathbf{H}\mathbf{z} - \mathbf{d}\|^2. \quad (39)$$

The minimisation can be performed using an iterative method such as the conjugate gradient method, where the gradient is given by

$$\nabla J = \mathbf{H}^T(\mathbf{H}\mathbf{z} - \mathbf{d}). \quad (40)$$

The inverse hessian of the cost function, $(\mathbf{H}^T\mathbf{H})^{-1}$, gives the covariance matrix of the least squares solution. In the next section we consider a direct solution method based on the singular value decomposition of the operator \mathbf{H} , which allows us to investigate the nature of the ill-posedness of the problem. We illustrate regularisation using a truncated singular value decomposition.

4.1 Singular value decomposition

We consider a singular value decomposition of \mathbf{H} of the form

$$\mathbf{H} = \mathbf{U}\mathbf{S}\mathbf{V}^T, \quad (41)$$

where \mathbf{U} is a $N \times N$ unitary matrix, \mathbf{V} is a $n \times n$ unitary matrix and \mathbf{S} is the $N \times n$ diagonal matrix whose diagonal elements are the singular values $s_1 \geq \dots \geq s_n \geq 0$. Using this decomposition, the solution \mathbf{z}_{LS} to (39) can be written as

$$\mathbf{z}_{\text{LS}} = \mathbf{V}\mathbf{S}^\dagger\mathbf{U}^T\mathbf{y} = \mathbf{H}^\dagger\mathbf{y}. \quad (42)$$




Figure 4. Singular values of the observability matrix for NEE (log scale)

The matrix $\mathbf{H}^\dagger = \mathbf{V}\mathbf{S}^\dagger\mathbf{U}^T$ is the *pseudo-inverse* of \mathbf{H} where \mathbf{S}^\dagger is the diagonal matrix obtained by transposing \mathbf{S} and replacing the non zero elements by their inverse s_i^{-1} . The covariance of the solution is given by

$$\text{Cov}(\mathbf{z}_{\text{LS}}) = \mathbf{H}^{\dagger T} \mathbf{H}^\dagger. \quad (43)$$

Much can be learned about the stability of the solution (42) by inspecting the singular values of \mathbf{H} . Assuming that \mathbf{H} is full rank, it can be shown, (see Golub and Van Loan (1996)), that the relative error in the solution, defined as the left hand side of the above inequality, is bounded by

$$\frac{\|\mathbf{z}_{\text{LS}} - \mathbf{z}_0\|}{\|\mathbf{z}_0\|} \leq \kappa(\mathbf{H}) \frac{\|\boldsymbol{\epsilon}\|}{\|\mathbf{d}\|}, \quad (44)$$

where $\kappa(\mathbf{H})$ is the condition number of \mathbf{H} defined by $\kappa(\mathbf{H}) = s_1/s_n$, \mathbf{z}_0 denotes the truth (possibly unknown) and $\boldsymbol{\epsilon}$ represents observational noise. When the condition number is large the matrix is said to be ill-conditioned, the problem is ill-posed and the solution (42) is unstable: small perturbations to the system can lead to very large perturbations in the solution.

4.2 Stability for NEE operator

As an example we consider the problem of assimilating NEE observations into DALECV2 to estimate model parameters and initial conditions at Morgan Monroe State forest. We linearise equation (28) about a point \mathbf{x}^* satisfying the EDCs, we form the observability matrix \mathbf{H} and compute its singular value decomposition. The singular values, shown in figure 4, reveal a condition number of order 10^5 . For a signal-to-noise ratio, namely $\|\boldsymbol{\epsilon}\|/\|\mathbf{d}\|$, of magnitude 0.1, inequality (44) gives an upper bound for the relative error in the solution of order 10^4 , which does not give much credit to the least square solution. How sharp is this bound? Are we overestimating the error? To answer these questions we create a set of noisy observations with noise variance $\sigma = 0.1$ and we compute the solution (42). The relative error for each component of the solution, η_i , and the variance ν_i , are given in Table 2. Despite a relatively good match between the modelled NEE ~~perturbations~~ perturbations and the observations, as shown in figure 5, the results of Table 2 show very large relative errors and variances for most variables. Moreover these results are in agreement with the results of REFLEX: parameters directly




Figure 5. Solution of the ~~linearized-linearised~~ inverse problem for NEE. The red points represent the observations, the red curve is the true trajectory, the green curve is the trajectory obtained using the unstable solution and the blue curve is obtained using the TSVD solution.

linked to foliage and GPP are better estimated than parameters related to allocation to and turnover of
 385 fine root/wood. The results of Table 2 reflect the sensitivity analysis shown on figure 2. The variables
 with respect to which NEE is the most (resp. least) sensitive are the less (resp. more) affected by the
 noise.

To reduce the impact of observational noise on the solution, regularisation is required. The *truncated
 singular value decomposition* (TSVD) is a simple and popular method for regularisation. TSVD
 390 consists in truncating the pseudo-inverse in equation (42) in order to remove the smallest singular
 values, the most affected by the noise. The solution $z^{(k)}$ is then given by

$$z^{(k)} = \mathbf{V}_k \mathbf{S}_k^\dagger \mathbf{U}_k^T \mathbf{y} = \mathbf{H}_k^\dagger \mathbf{y}, \quad (45)$$

where k is the truncation rank, \mathbf{S}_k , \mathbf{U}_k and \mathbf{V}_k are the rectangular matrices formed by the first k
 columns of \mathbf{S} , \mathbf{U} and \mathbf{V} . The covariance of the solution is given by

$$395 \text{Cov}(z^{(k)}) = \mathbf{V}_k (\mathbf{S}_k^\dagger)^{-2} \mathbf{V}_k^T. \quad (46)$$

The truncation rank k can be chosen using the L-curve method. The L-curve is a log-log plot of the
 norm of the solution $\|z^{(k)}\|$ against the norm of the residual $\|\mathbf{H}z^{(k)} - \mathbf{d}\|$ ~~parametrized-parametrised~~
 by the regularisation parameter k . The optimal parameter corresponds to the point of maximum
 curvature of the L-curve. Further details on the L-curve method can be found in Hansen and O'Leary
 400 (1993).

In our example with NEE we use Hansen's ~~regularization-regularisation~~ tools, see Hansen (2007),
 to perform the TSVD method. The truncation rank obtained using the L-curve method is $k = 7$. The
 last three columns of Table 2, presenting the TSVD solution, the relative error of each components
 and the variances, can be compared with the results for the unstable solution. Whereas the relative
 405 errors in the unstable solution range from 5.3×10^{-2} to 3.8×10^4 the relative errors in the ~~regularized~~
~~regularised~~ solution range from 5.3×10^{-2} to 5.1. We see that TSVD has the effect of keeping small
 the variables that cannot be estimated correctly. As previously stated the results of the regularisation
 can be related to the sensitivity analysis depicted on Figure 2: TSVD prevents the variables with

Table 2. Results of the linear inverse problem showing: 1- the solution components for the least square solution \mathbf{z}_{LS} together with their relative error η_i (dimensionless quantity) and variance ν_i ; 2- the solution components for the TSVD solution $\mathbf{z}^{(k)}$ together with their relative error $\eta_i^{(k)}$ and variance $\nu_i^{(k)}$.

	\mathbf{x}^*	\mathbf{z}^*	\mathbf{z}_{LS}	η_i	ν_i	$\mathbf{z}^{(k)}$	$\eta_i^{(k)}$	$\nu_i^{(k)}$
p_1	-6.984	-0.070	3.715	54.190	19182.5715	0.004	1.052	0.0001
p_2	-1.114	-0.011	0.342	31.715	23.2871	0.003	1.242	0.0005
p_3	-3.480	-0.035	-2.690	76.285	7384.9940	0.001	1.022	0.0000
p_4	-2.745	-0.027	3.389	124.470	82380.0339	-0.000	1.000	0.0000
p_5	0.086	0.001	0.048	54.082	0.6575	-0.004	5.139	0.0009
p_6	-8.776	-0.088	67.445	769.477	1581516.7404	0.000	1.001	0.0000
p_7	-5.265	-0.053	-0.999	17.970	57.1478	-0.005	0.900	0.0001
p_8	-6.640	-0.066	-0.344	4.176	7981.3944	-0.016	0.757	0.0013
p_9	-10.292	-0.103	133.504	1298.187	494620.7529	-0.006	0.946	0.0002
p_{10}	-3.035	-0.030	-0.003	0.889	2.6980	-0.011	0.632	0.0008
p_{11}	3.539	0.035	1.075	29.370	79.5237	-0.003	1.083	0.0003
p_{12}	4.736	0.047	0.045	0.053	0.0256	0.044	0.080	0.0003
p_{13}	-0.772	-0.008	1.042	135.879	8676.9499	-0.028	2.616	0.0033
p_{14}	3.261	0.033	0.072	1.196	4.3971	0.004	0.866	0.0001
p_{15}	5.533	0.055	0.084	0.515	0.0548	0.058	0.053	0.0003
p_{16}	4.082	0.041	0.092	1.265	1.2521	0.004	0.904	0.0001
p_{17}	5.178	0.052	1.631	30.497	8160.2559	0.000	0.997	0.0005
C_{lab}	6.237	0.062	1.002	15.073	8237.5716	0.019	0.697	0.0020
C_f	4.073	0.041	1.348	32.090	8070.9246	0.005	0.888	0.0001
C_r	6.858	0.069	1.788	25.067	8300.6094	-0.012	1.170	0.0008
C_w	8.341	0.083	-318.175	3815.484	7436253.3370	0.000	0.999	0.0000
C_l	5.961	0.060	0.568	8.532	8550.3479	-0.006	1.097	0.0002
C_s	8.956	0.090	-134.334	1500.869	483025.4281	-0.006	1.064	0.0002

respect to which NEE is the least sensitive from growing ~~unboundedly~~unbounded.

410 In the next section we consider the concept of a resolution matrix, which allows for a finer analysis of the solution of the linear problem.

4.3 Resolution matrix

As suggested by equations (42) and (45), finding a solution \mathbf{z} amounts to constructing a *generalized generalised* inverse \mathbf{H}^g such that formally

415
$$\mathbf{z} = \mathbf{H}^g \mathbf{d}. \tag{47}$$

The *generalized-generalised* inverse is the operator representing any method, direct or iterative, used to solve the linear inverse problem, including or not any kind of regularisation. In the previous section we considered two examples of generalised inverse, the pseudo inverse and the truncated inverse obtained using TSVD. The *generalized-generalised* inverse can be used to define operators

420 which directly address the conditions for well posedness for the linearised problem. Assuming a true state z^* exists, possibly unknown, then using equation (38) and (47) we can define an operator \mathbf{N} called the *model resolution matrix* which relates the solution z to the true state

$$z = \mathbf{H}^g \mathbf{H} z^* = \mathbf{N} z^*. \quad (48)$$

This matrix gives a ~~pratical~~practical tool to analyse the resolution power of an inverse method, that is its ability to retrieve the true state, including or not any ~~regularization~~regularisation method: the closer \mathbf{N} is to the identity the better the resolution. Moreover the trace of the matrix defines a natural notion of *information content* (IC). Similarly a *data resolution matrix* can be defined to study how well data can be reconstructed and its diagonal elements naturally define a notion of *data importance*. For the two examples of ~~generalized~~generalised inverse presented in the previous section we obtain
430 the following resolution matrices:

$$\mathbf{N} = \mathbf{H}^\dagger \mathbf{H}, \quad (49)$$

for the pseudo-inverse and

$$\mathbf{N} = \mathbf{V}_k \mathbf{V}_k^T, \quad (50)$$

for the truncated pseudo inverse. In the first case the IC equals the number of non zero singular values, in the second case the IC equals the truncation rank k . An in-depth theoretical and ~~pratical~~practical analysis of these concepts and those introduced in the remainder of this section can be found in Menke (1984).
435

While the model resolution matrix allows us to see how a solution strategy maps the true state variables to the solution of the inverse problem, and to see how well and how independently the state variables can be recovered, one also needs to assess the uncertainty of the solution. This can
440 be studied using the so called *unit covariance matrix*, \mathbf{C} , defined using the ~~generalized~~generalised inverse as

$$\mathbf{C} = \mathbf{H}^{gT} \mathbf{H}^g. \quad (51)$$

By characterising the degree of error amplification that occurs in the mapping from the true state to the solution of the inverse problem, the unit covariance matrix is a crucial object to study the
445 stability of the solution with respect to observational noise. The unit covariance matrix defined by (51) agrees with the covariance matrices given in the previous section by (43) for the pseudo-inverse, and by (46) when TSVD is applied.

4.4 Resolution for LAI operator

450 We now study the model resolution matrix for the LAI observation operator at Morgan Monroe State Forest. In the first instance we will demonstrate the resolution power of the LAI signal without regularisation using the pseudo-inverse as ~~generalized~~generalised inverse first, and then apply TSVD to

show how using the truncated pseudo-inverse affects resolution. In a second case we will study the added value of the EDCs in terms of resolution.

As previously, we linearise equation (28) about the point x^* given in Table 2. The trace of the reso-




Figure 6. Model resolution matrix for the LAI operator.

455

lution matrix obtained using the pseudo-inverse as generalised inverse is 10, and this means that 10 independent variables can be estimated using LAI. These independent-independent variables are not the variables in which the system is expressed but a linear transformation can be found to express the system in terms of the independent variables. Figure 6 shows the model resolution matrix for LAI.

460

As shown in section 2.3 with the sensitivity analysis, 11 out of the 23 variables are not sensitive to LAI, and this can be seen in the resolution matrix by the diagonal terms which are zero, represented by the blue color-colour. In contrast the diagonal elements corresponding to sensitive variables have positive values, represented by colors-colours ranging from light blue to red. Figure 6 also shows that whereas p_5 , p_{11} , p_{12} , p_{14} , p_{15} and p_{16} are perfectly resolved (the corresponding elements are

465

colored-coloured in brown or dark red), there exist linear combinations between the remaining sensitive variables and this explains why only 10 independent-independent variables can be estimated from the 12 sensitive variables.

For the study of the unit covariance matrix we restrict ourselves to the sensitive variables, this amounts to removing the columns corresponding to the non sensivite-sensitive variables, containing

470

only null elements, from the observability matrix. The dependency of the solution on observational noise can be studied by looking at Figure 7 where the diagonal elements of the unit covariance matrix, corresponding to the variance of each element of the solution obtained using the pseudo-inverse, are represented in log scale. Except for p_5 , p_{12} , p_{15} and p_{16} all variances are shown to be large.

As previously, we illustrate a simple regularisation strategy, TSVD, and show its effects on both

475

resolution and stability. Figure 8 shows the resolution matrix for LAI with optimal truncation rank




Figure 7. Diagonal elements (log scale) of the unit covariance matrix for the LAI operator using the pseudo inverse in green, using TSVD in yellow.




Figure 8. Model resolution matrix for the LAI operator using TSVD with truncation rank $k = 5$.

$k = 6$. The IC decreases to 6. We see that whereas p_5 , p_{12} , p_{15} and p_{16} remain almost perfectly resolved, p_{13} , p_{17} and C_{lab} are only partially resolved and the remaining variables are not resolved properly. Figure 7 shows the corresponding diagonal elements of the unit covariance matrix, we see that the variances have been drastically reduced. This example shows how regularisation ensures stability at the price of losing resolution.

480

We now consider the effect of incorporating the static EDCs into the variational framework in terms of resolution. The static EDCs are given by the seven first EDCs, the linear problem is then given by

$$\begin{bmatrix} \mathbf{H} \\ \tilde{\mathbf{G}} \end{bmatrix} z = \begin{pmatrix} d \\ f \end{pmatrix}, \quad (52)$$

485 where

$$\tilde{\mathbf{G}} = g(\mathbf{x}^*)^{-1} \Sigma^{-1/2} \mathbf{G} \quad (53)$$




Figure 9. Model resolution matrix for LAI and static EDCs as defined by equation 52.

with $\Sigma^{-1/2}$ the inverse of the symmetric square root of the covariance matrix Σ , defined in Section 3.3, restricted to the seven first components. The static EDCs depend only on 13 out of the 23 variables, namely p_1 to p_{10} , p_{12} , p_{13} and p_{15} , this can be seen on the matrix \mathbf{G} where the columns corresponding to the remaining variables are null. Together with LAI observations, whose sensitive variables are represented in Figure 2, we therefore have 19 sensitive variables. The model resolution matrix corresponding to the operator on the left hand side of equation (52), obtained using the pseudo inverse, is depicted in Figure 9. The trace of the model resolution matrix gives an IC of 16, 13 variables are perfectly resolved, 4 variables show linear dependencies (p_2 , p_3 , p_4 and p_{13}). However, although p_9 and p_{10} are sensitive variables they do not appear to be resolved at all: inspecting the linear operator \mathbf{G} shows that the non zero components corresponding to p_9 and p_{10} are several order of magnitude smaller than the other components.

This example shows clearly the benefit of introducing the static EDCs to help estimating poorly constrained or otherwise undetermined components.

500 5 Experiments at Ameriflux sites

We now consider a real experiment at the Morgan Monroe state forest. At this Ameriflux site, 13 years of MODIS LAI monthly mean observations from 2001 to 2013, NEE, GPP and thus RESP observations from 2001 to 2005 are available. Our goal is to study two different aspects; the first one is the impact of using multiple data streams: how does it affect uncertainty of the predicted fluxes? how well do we predict non-observed fluxes? And the second one is to use the static EDCs and to assess their utility in constraining poorly sensitive variables.

Table 3. Experiment set up summary: in Exp 1 we use LAI and bounds constraints (BDS), in Exp 2 we use LAI, NEE and BDS and so on.

	LAI	NEE	GPP	RESP	BDS	EDCs
Exp 1	•				•	
Exp 2	•	•			•	
Exp 3	•	•	•	•	•	
Exp 4	•				•	•
Exp 5	•	•			•	•
Exp 6	•	•	•	•	•	•

When including all terms the cost function, J_{TOT} , becomes

$$\begin{aligned}
 J_{TOT}(\mathbf{x}) &= \frac{\lambda_L}{2} \|\mathbf{h}_L(\mathbf{x}) - \mathbf{y}_L\|^2 + \frac{\lambda_N}{2} \|\mathbf{h}_N(\mathbf{x}) - \mathbf{y}_N\|^2 \\
 &\quad + \frac{\lambda_G}{2} \|\mathbf{h}_G(\mathbf{x}) - \mathbf{y}_G\|^2 + \frac{\lambda_R}{2} \|\mathbf{h}_R(\mathbf{x}) - \mathbf{y}_R\|^2 \\
 510 \quad &\quad + \frac{\lambda_c}{2} \|\log(\mathbf{g}(\mathbf{x})) - \mathbf{c}\|_{\Sigma}^2 + \frac{\lambda_0}{2} \|\mathbf{x} - \mathbf{x}_0\|_{\mathbf{B}}^2 \\
 &= J_L + J_N + J_G + J_R + J_c + J_0,
 \end{aligned}$$

where subscripts L, N, G and R stand for LAI, NEE, GPP and RESP respectively. The vectors \mathbf{y}_L , \mathbf{y}_N , \mathbf{y}_G and \mathbf{y}_R represent the observation vectors for LAI, NEE, GPP and RESP respectively. The
 515 scalars λ_L , λ_N , λ_G and λ_R take the value 0 or 1 depending on whether or not the corresponding data stream is included in the experiment. The scalar λ_c takes the value 0 or 1 depending on whether we include the EDCs and λ_0 takes the value 1.

We perform six experiments ~~summarized~~ summarised in Table 3. In experiment 1, Exp 1, we use only LAI observations and bounds constraints so that in the cost function J_{TOT} we set $\lambda_L = 1$ and
 520 $\lambda_0 = 1$, and the other λ s are set to zero. For Exp 2, we use LAI and NEE observations, that is we set $\lambda_L = 1$, $\lambda_N = 1$ and $\lambda_0 = 1$; the other λ s are set to zero. We proceed similarly for the remaining experiments. Here we assimilate all data streams simultaneously; it is not our intention to question what method best ~~accommodates~~ accommodates multiple data streams, MacBean et al. (2016) addresses this question using a simple C-cycle model. Moreover we choose to assume the same sta-
 525 tistical error for all data streams and set their error covariance matrix equal to the identity. To avoid being trapped at meaningless local minima, the experiments are performed multiple times using different initialisation parameter sets and results for the best candidate only are reported.

The results of the experiments are presented in Table 4 where each element of J_{TOT} is given for all experiments, and in Table 5 where the solution components and their variance are presented for
 530 all experiments. Results of Table 4 show that J_L is the smallest in Exp 1 when LAI only is used. In Exp 2, when adding NEE we see that J_N decreases from 109.012 in Exp 1 to 15.263, and J_G slightly decreases as compared to Exp 1, but on the contrary J_R increases. In Exp 3 we see that all costs drastically decrease compared to their initial values. Going from Exp 1 to Exp 3 J_0 slightly

increases, adding more data streams constrains more parameters and the parameters shift from their prior value which may cause J_0 to increase. Similar observations can be made for Exp 4 to Exp 6, moreover we see that including the EDCs only slightly affects the costs, a reason for this might be that EDCs help constraining the less sensitive parameters for which the costs are the less sensitive as suggested by the sensitivity analysis depicted in Figure 2. To see the effect of the EDCs we need to look at Table 5, which details the solution components together with their relative variance defined by the ratio of the variance by the parameter range. In Exp 1 we see that the variables with the smallest relative variance are the most sensitive parameters as illustrated in Figure 2; they are p_2 , p_5 , p_{10} , p_{12} , p_{14} , p_{15} , p_{16} and p_{17} . We recall that the sensitivity analysis of section 2.3 was performed by averaging sensitivities for an ensemble of initial parameter sets, therefore the ranking shown in figure 2 may not be reflected in the relative variances. As we include NEE in Exp2 we see that most relative variances decrease, especially for p_8 , p_9 , p_{10} , p_{13} , C_{lab} , C_f and C_1 . The only variable whose relative variance increases is p_{14} , but as shown in figure 2 p_{14} has very small sensitivity. In Exp 3 most relative variances decrease. The values are still large though for p_1 , p_3 , p_4 , p_6 , p_9 , C_r and C_1 . Again similar features can be observed for Exp 4 to Exp 6 but a clear improvement can be seen for most variables except for C_r which is not constrained by the seven first EDCs. Finally, the last column of Table 4 shows the computation time for each experiments. As expected we see that the more observation streams we consider the longer the ~~experiemnt~~ experiment takes to run, and incorporating the EDCs increases the computation time. However we stress that these figures are several orders of magnitude less than the time required to perform the same experiments using the current gold standard MCMC approach used in Bloom and Williams (2015).

Figures 10 and 11 show the predicted fluxes for LAI, NEE, GPP and RESP for the result of Exp 6. We can see a good agreement between modelled fluxes and observations. The uncertainty of the predicted fluxes is evaluated by ~~modeling~~ modelling an ensemble of trajectories from a 95% ellipsoid of the posterior truncated ~~gaussian~~ Gaussian distribution: these trajectories are represented as ~~gray~~ grey curves on Figure 10 and 11. Figure 12 shows the posterior parameter distribution marginals for p_1 , p_7 , p_{16} , and C_f for Exp 6, illustrating the four different cases where: as for p_{16} most of the marginal is contained in the parameter range; the marginal is truncated on the left or the right as for p_7 or C_f and truncated on both sides for p_1 .

6 DALEC-SP

In the previous section we used EDCs within 4DVAR and showed their benefit in reducing drastically the uncertainty of otherwise undetermined variables. However we only included the static EDCs which do not require a model run: including more EDCs often leads to convergence issues, the solution and its uncertainty become subject to caution.

As shown in Chuter et al. (2015) for the previous DALEC evergreen and deciduous models, the

Table 4. Costs for the results of the inverse modelling experiments. The last column reports the computation time in seconds for the experiment.

	J_L	J_N	J_G	J_R	J_0	J_c	time (s)
x_{init}	179.525	353.229	1265.556	419.696	0.003	7.157	0.000
Exp 1	14.083	109.012	153.475	45.415	0.017	2.498	2.722
Exp 2	19.188	15.263	145.349	131.963	0.018	3.704	7.541
Exp 3	25.089	16.737	36.155	17.842	0.020	4.643	5.886
Exp 4	14.083	107.420	152.908	45.480	0.016	2.498	5.012
Exp 5	19.193	15.262	145.254	131.878	0.018	3.701	9.045
Exp 6	25.059	16.699	36.143	17.826	0.019	4.642	8.215

Table 5. Results of the inverse modelling experiments. The solution components together with their relative variance, in bracket, are given for each experiment. The vector x_{init} is the randomly chosen parameter set satisfying the EDCs that initialises the minimisation routine.

	x_{init}	Exp 1	Exp 2	Exp 3	Exp 4	Exp 5	Exp 6
p_1	-5.172	-8.059 (1.727)	-8.248 (1.471)	-8.282 (1.021)	-6.323 (0.125)	-6.486 (0.095)	-6.415 (0.087)
p_2	-0.947	-0.885 (0.207)	-1.106 (0.120)	-1.085 (0.030)	-0.770 (0.126)	-0.960 (0.093)	-0.984 (0.019)
p_3	-4.318	-2.673 (0.955)	-2.944 (0.849)	-3.376 (0.894)	-3.001 (0.950)	-4.419 (0.965)	-3.461 (0.880)
p_4	-1.493	-2.649 (0.978)	-2.813 (0.961)	-2.692 (0.936)	-1.308 (0.111)	-1.703 (0.142)	-1.354 (0.097)
p_5	1.123	0.117 (0.003)	0.153 (0.002)	0.085 (0.000)	0.149 (0.002)	0.172 (0.001)	0.091 (0.000)
p_6	-7.959	-8.752 (0.922)	-8.870 (0.911)	-8.707 (0.919)	-7.151 (0.170)	-7.039 (0.108)	-7.681 (0.458)
p_7	-7.432	-6.908 (1.151)	-6.373 (0.941)	-5.064 (0.224)	-7.236 (0.221)	-5.961 (0.109)	-6.144 (0.108)
p_8	-5.281	-6.908 (1.151)	-6.906 (0.316)	-6.522 (0.078)	-5.565 (0.113)	-5.753 (0.052)	-6.160 (0.039)
p_9	-16.012	-11.513 (2.303)	-10.075 (0.995)	-11.411 (1.514)	-14.592 (2.303)	-8.384 (0.036)	-11.266 (0.143)
p_{10}	-3.041	-3.272 (0.373)	-3.296 (0.085)	-3.036 (0.055)	-3.244 (0.373)	-3.045 (0.072)	-3.079 (0.055)
p_{11}	2.792	3.829 (0.540)	4.026 (0.163)	3.542 (0.003)	3.630 (0.286)	4.198 (0.232)	3.544 (0.003)
p_{12}	3.549	4.626 (0.002)	4.739 (0.000)	4.735 (0.000)	4.627 (0.002)	4.733 (0.000)	4.735 (0.000)
p_{13}	-1.768	-0.693 (0.130)	-0.996 (0.067)	-0.795 (0.046)	-0.779 (0.039)	-1.193 (0.037)	-0.801 (0.029)
p_{14}	3.343	4.013 (0.034)	3.291 (0.123)	3.292 (0.052)	3.940 (0.029)	3.526 (0.048)	3.247 (0.079)
p_{15}	5.656	5.512 (0.001)	5.528 (0.001)	5.531 (0.000)	5.521 (0.001)	5.540 (0.000)	5.534 (0.000)
p_{16}	4.529	4.115 (0.068)	3.993 (0.025)	4.095 (0.011)	4.003 (0.061)	4.071 (0.019)	4.082 (0.010)
p_{17}	5.351	5.289 (0.213)	5.278 (0.198)	5.138 (0.165)	5.142 (0.135)	5.032 (0.110)	5.098 (0.098)
C_{lab}	3.979	5.950 (0.115)	6.031 (0.059)	6.187 (0.040)	5.884 (0.078)	5.841 (0.039)	6.151 (0.025)
C_f	5.389	4.677 (0.282)	4.868 (0.068)	4.038 (0.066)	4.421 (0.302)	4.342 (0.093)	4.018 (0.051)
C_w	7.045	5.298 (1.151)	5.829 (0.900)	6.520 (0.096)	5.309 (1.022)	6.910 (0.226)	7.061 (0.135)
C_r	9.753	8.406 (1.554)	8.188 (1.533)	8.318 (1.544)	8.687 (1.554)	6.771 (1.030)	8.538 (1.415)
C_1	3.992	5.298 (1.151)	7.307 (0.300)	6.226 (0.161)	5.963 (0.990)	5.670 (0.263)	5.985 (0.088)
C_s	9.721	8.406 (1.900)	9.546 (1.188)	8.603 (1.633)	8.895 (1.900)	7.140 (0.561)	9.003 (0.718)




Figure 10. DALECV2 monthly estimates for LAI and NEE at Morgan Monroe State forest. The red dots are the observations, the blue trajectories are obtained using the 4DVAR analysis, the ~~gray~~-grey trajectories are ensemble runs obtained from a 95% confidence sample of the posterior pdf.




Figure 11. DALECV2 monthly estimates for GPP and RESP at Morgan Monroe State forest. The red dots are the observations, the blue trajectories are obtained using the 4DVAR analysis, the ~~gray~~-grey trajectories are ensemble runs obtained from a 95% confidence sample of the posterior pdf.




Figure 12. Posterior parameter distributions for parameters p_1 , p_7 , p_{16} and C_f for Exp 6. For each plot the limits of the abscissa correspond to the parameter range. The red curve is the gaussian-Gaussian posterior distribution and the blue bars represent-represent the sample used to produce the gray-grey trajectories in figures 10 and 11.




Figure 13. Pseudo periodical seasonal cycle for DALECV2. Using a given set of parameters and initial values for C_w and C_s , 100 DALECV2 runs are performed using random initial values for C_{lab} , C_f , C_r , C_l . The plot shows the 100 trajectories for C_f .

evolution of the carbon pools for DALECV2 show a tipping point which depends on the parameters p_1 to p_{17} . Given a set of parameters \mathbf{p} the fast carbon pools C_{lab} , C_f , C_r and C_l grow or decay rapidly to an equilibrium state. This equilibrium-equilibrium is either zero, the forest dies out, or a pseudo-periodical seasonal cycle as shown on Figure 13 for C_f . Moreover there exists a limit value below which any initial condition leads to the zero equilibrium and above which the equilibrium is a strictly positive pseudo-periodical seasonal cycle.

Here we consider ecosystems with no recent major disturbance, where the fast carbon pools are expected to be close to their pseudo-periodical cycle. To model these ecosystems, one can either restrict the parameter space by using the dynamic EDCs, or we can introduce a spin-up period during which the carbon pools reach their attractor. Given parameters p_1 to p_{17} and initial values for C_w and C_s , a first run of DALECV2 is performed to obtain a state which is closer to a pseudo-periodical

Table 6. Results of the inverse modelling experiments with DALEC-SP model showing the log of the initial carbon pools (gCm^{-2}). The rows labelled (SP) correspond to the solution components for DALEC-SP, the other rows reproduce the results for DALEC as reported in Table 5. As previously the relative variances are given in bracket.

	x_{init}	Exp 1	Exp 2	Exp 3	Exp 4	Exp 5	Exp 6
C_{lab}	3.979	5.950 (0.115)	6.031 (0.059)	6.187 (0.040)	5.884 (0.078)	5.841 (0.039)	6.151 (0.025)
C_{lab} (SP)	NA	5.461 (0.106)	5.961 (0.015)	6.064 (0.008)	5.097 (1.572)	5.914 (0.074)	6.048 (0.011)
C_{f}	5.389	4.677 (0.282)	4.868 (0.068)	4.038 (0.066)	4.421 (0.302)	4.342 (0.093)	4.018 (0.051)
C_{f} (SP)	NA	4.358 (0.112)	4.602 (0.034)	3.966 (0.029)	4.102 (0.633)	4.586 (0.079)	4.024 (0.043)
C_{r}	7.045	5.298 (1.151)	5.829 (0.900)	6.520 (0.096)	5.309 (1.022)	6.910 (0.226)	7.061 (0.135)
C_{r} (SP)	NA	4.702 (0.765)	5.069 (0.575)	5.072 (0.665)	6.078 (0.260)	5.323 (0.316)	6.398 (0.116)
C_{w}	9.753	8.406 (1.554)	8.188 (1.533)	8.318 (1.544)	8.687 (1.554)	6.771 (1.030)	8.538 (1.415)
C_{w} (SP)	7.201	5.298 (0.853)	5.289 (0.852)	5.285 (0.853)	5.298 (0.853)	5.320 (0.852)	5.203 (0.853)
C_{l}	3.992	5.298 (1.151)	7.307 (0.300)	6.226 (0.161)	5.963 (0.990)	5.670 (0.263)	5.985 (0.088)
C_{l} (SP)	NA	6.361 (0.276)	6.590 (0.215)	6.791 (0.181)	5.412 (0.226)	5.928 (0.074)	6.056 (0.033)
C_{s}	9.721	8.406 (1.900)	9.546 (1.188)	8.603 (1.633)	8.895 (1.900)	7.140 (0.561)	9.003 (0.718)
C_{s} (SP)	9.793	8.406 (1.900)	8.550 (0.799)	9.324 (1.359)	8.406 (1.900)	8.940 (0.222)	10.453 (1.090)

580 cycle for the fast carbon pools. The steady state trajectories are then used to initialise the fast carbon pools. For this DALECV2-"spin-up" model, DALEC-SP, the state variable is therefore formed of the seventeen parameters p_1, \dots, p_{17} and the initial conditions for C_{w} and C_{s} .

DALEC-SP offers several advantages: some of the EDCs such as those controlling the growth and the half life period of carbon pools are almost automatically satisfied; this reduces largely the time
585 required to generate the pdf $p(c|\mathbf{x})$. Moreover as the sensitivity analysis and the resolution matrices showed, the fast carbon pools are variables that are not highly sensitive to the signals that we observe, and therefore reducing the number of variables by removing the fast carbon pools is likely to improve the overall [conditioning-conditioning](#) of the inverse problem. To investigate this assertion we perform Exp 1 to 6 using DALEC-SP. The solution components and their variance for the carbon
590 pools are presented in Table 6; the results for the parameters p_1 to p_{17} are not reported as they do not significantly differ from what was observed and reported in Table 5 for DALECV2. For the fast carbon pools, which are not directly estimated during the assimilation process, we start by taking a sample of the posterior pdf and then we run DALEC-SP for this sample. The values presented in Table 6 represent the means and variances of the fast carbon pools after the spin up period. Except
595 for two anomalies in Exp 4, where the uncertainty for C_{lab} and C_{f} is larger with DALEC-SP as compared to DALECV2, almost all uncertainties for all experiments are smaller with DALEC-SP. Despite some improvement in Exp 2, 4 and 5, the uncertainty for C_{s} is still large.

7 Discussion

600 To our knowledge, this paper presents the first application of variational methods for an inverse
modelling experiment using DALEC. Over the last fifteen years many studies have validated the
use of DALEC together with various types of data streams to infer ecological parameters at site
level but ensemble Kalman filter first and then Monte Carlo methods were privileged. At the same
time 4DVAR was successfully used at global scale to constrain ecosystem parameters in carbon
cycle data assimilation system (CCDAS). In Rayner et al. (2005) the Biosphere Energy Transfer
605 Hydrology model (BETHY) is coupled with the transport model TM2, and satellite observations
of photo-synthetically active radiation and atmospheric CO₂ concentration observations are used
to optimise model parameters. In this context Kemp et al. (2014) investigated how to constrain the
4DVAR problem in CCDAS through a number of different methods: using constrained optimisation,
adding a penalty term and applying parameter transformations. They concluded that using parameter
610 transformations give the best results. In our context the three methods were investigated: Gaussian
anamorphosis where priors based on the distribution of parameters satisfying the EDCs were considered,
constrained optimisation as stated in section 3.3, and adding a penalty term to account for the EDCs.
The latter solution which is the main interest of this publication was found to be the most successful
in our case.

615 The complexity of global scale experiments still limit the application of fully nonlinear methods such
as MCMC. In Ziehn et al. (2012) a comparison between the MCMC Metropolis-Hastings approach
and 4DVAR for the BETHY-TM2 CCDAS framework is performed. This study reports a computation
time of less than one hour for the variational method and about eight months for the overall MCMC
computation. For our setting, DALECV2 site based experiment, the complexity is relatively small
620 and a MCMC approach is affordable. Used in Bloom and Williams (2015), the MCMC approach for
DALEC is studied in detail in Safta et al. (2015), the resulting parameter distributions suggest that
4DVAR and the inherent Gaussian approximation provides a reasonable posterior distribution.

As most variational methods, the analysis and application presented in this paper rely heavily on the
possibility to derive the tangent linear model and its adjoint. DALECV2 was designed to take into
625 account this requirement, in particular replacing the phenology process of the DALEC deciduous
model was suggested in order to obtain differentiable processes. The model resolution matrix and
the gradient of the cost function, including the additional term encoding the EDCs, are computed
using adjoint techniques. Despite the increasing capacities offered by automatic differentiation tools,
deriving and maintaining an adjoint code can be a complicated task, and, besides its limiting hypothesis,
630 this is certainly one of the main reason for choosing alternatives to 4DVAR. In a paper in preparation
we use ensemble methods to approximate the gradient of the cost function and to derive approximate
resolution matrices, and the experiments presented in this paper are reproduced. The approach,
which no longer requires the adjoint, shows very promising results: firstly in terms of estimating
parameters, and secondly in terms of computation time by using graphic processing units (GPU) to

635 perform massive parallel computations.

Designing a global scale experiment involving a coupling between DALEC and a transport model has been considered but is still at an early stage. As presented in Bloom and Williams (2015), the EDCs were originally introduced to constrain unresolved parameters at site level where, in the absence of any other information, only MODIS LAI observations were available. In theory there is no restriction
640 to readily apply the same constraints at global scale however their efficiency highly depends on the nature of the coupling between the ecosystem model and the transport model and on the observation streams considered. Nonetheless in this context 4DVAR remains the only reasonable method to consider in terms of computer resources, and our study demonstrate that the current research efforts to develop regularisation strategies fit well into the variational framework.

645 **8 Conclusions**

We used DALECV2 and combined multiple data streams - MODIS monthly LAI and monthly NEE, GPP and RESP at an Ameriflux site - together with ecological constraints to estimate model parameters and initial conditions and to provide uncertainty characterisation for predicted fluxes. DALECV2 is a simple model; it represents the basic processes at the heart of more sophisticated models of the
650 carbon cycle and, besides its large modelling skills, its simplicity allows for close mathematical scrutiny. Here we adopted a variational approach where the tangent linear model and its adjoint play a major role in 1) facilitating a linear analysis which allows to understand the nature of the ill-posed problem and to evaluate strategies to ~~regularize~~ regularise it; 2) finding a posterior distribution for the state variables.

655 We performed a sensitivity analysis using a direct method that consists in studying the first order derivatives of the output computed using an adjoint method. The sensitivity analysis is a prerequisite to any work with a model, but there is a paucity of literature on this topic in connection with DALEC. Our analysis reveals generic issues that will be encountered in many inverse modelling ~~stategies~~ strategies. Studying the first order inverse problem, we discussed how noise affects the stability of the solution and we illustrated a simple regularisation method. We then introduced the notion of ~~resolution matrix, which we adapted from the field of seismic tomography,~~ model resolution matrix and showed how this can be used to diagnose the ill-posedness of an inverse problem and evaluate the result of regularisation strategies. While some of our findings may be anticipated in the framework of a simple model, it is important to describe these tools and their interpretation as
665 similar analyses can be readily applied to a wide range of more complex models.

Bloom and Williams (2015) proved the benefit of the EDCs in constraining poorly resolved components of the carbon cycle and ~~recommanded~~ recommended their use for inverse ~~modeling~~ modelling problems. We successfully incorporated the EDCs within the context of variational data assimilation. Our results confirm that the EDCs regularise an otherwise ill-posed problem and efficiently reduce

670 the uncertainty of predicted fluxes, and thus ~~confort the recommandation~~ comfort the recommendation
of Bloom and Williams (2015). Moreover, our ~~modification~~ modification to DALECV2, DALEC-SP
which includes a spin-up period, offers an alternative to some EDCs that facilitates the variational
approach.

This study did not aim at providing an exhaustive account for the capability of variational tools, nor
675 at exploring all aspects of the EDCs for the inverse problem for DALEC. The objectives were to use
4DVAR and show that it offers a suitable framework to solve efficiently, robustly and quickly the
inverse problem for DALEC, and to present some methodology to analyse some issues that affect
most methods based on Bayesian inference.

9 Code availability

680 The model and inversion code, together with the drivers, observational data and experiment results
are available at: <https://zenodo.org/record/269937>.

Acknowledgements. The authors would like to thank the anonymous referees for their valuable comments
which helped to improve the manuscript. This project was funded by the NERC National Centre for Earth Obser-
vation, NCEO, UK. We acknowledge US-MMS AmeriFlux site for its data records. Research at the MMSF site
685 was supported by the Office of Science (BER), U.S. Department of Energy, Grant No. DE-FG02-07ER64371.
AmeriFlux is funded by the United States Department of Energy (DOE – TES), Department of Commerce
(DOC – NOAA), the Department of Agriculture (USDA – Forest Service), the National Aeronautics and Space
Administration (NASA), the National Science Foundation (NSF). We are grateful to J. Exbrayat and A. Bloom
for providing us with meteorological drivers, MODIS LAI observations and DALECV2 code. Finally we are
690 grateful to M. Williams, J. Exbrayat, A. Bloom and T. Hill for ~~feedbacks~~ comments and useful discussions.

References

- Bloom, A. A. and Williams, M.: Constraining ecosystem carbon dynamics in a data-limited world: integrating ecological "common sense" in a model–data fusion framework, *Biogeosciences*, 12, 1299–1315, doi:10.5194/bg-12-1299-2015, 2015.
- 695 Chuter, A. M., Aston, P. J., Skeldon, A. C., and Roulstone, I.: A dynamical systems analysis of the data assimilation linked ecosystem carbon (DALEC) models, *Chaos*, 25, 036401, doi:10.1063/1.4897912, 2015.
- Fox, A., Williams, M., Richardson, A. D., Cameron, D., Gove, J. H., Quaife, T., Ricciuto, D., Reichstein, M., Tomelleri, E., Trudinger, C. M., and Van Wijk, M. T.: The REFLEX project: Comparing different algorithms and implementations for the inversion of a terrestrial ecosystem model against eddy covariance data, *Agricultural and Forest Meteorology*, 149, 1597 – 1615, doi:10.1016/j.agrformet.2009.05.002, 2009.
- 700 Giering, R. and Kaminski, T.: Recipes for Adjoint Code Construction, *ACM Trans. Math. Softw.*, 24, 437–474, doi:10.1145/293686.293695, 1998.
- Golub, G. H. and Van Loan, C. F.: *Matrix Computations* (3rd Ed.), Johns Hopkins University Press, Baltimore, MD, USA, 1996.
- 705 Hadamard, J.: *Lectures on Cauchy's problem in linear partial differential equations*, Yale University Press, 1923.
- Hansen, P. C.: Regularization Tools version 4.0 for Matlab 7.3, *Numerical Algorithms*, 46, 189–194, doi:10.1007/s11075-007-9136-9, 2007.
- Hansen, P. C. and O'Leary, D. P.: The Use of the L-Curve in the Regularization of Discrete Ill-Posed Problems, *SIAM Journal on Scientific Computing*, 14, 1487–1503, doi:10.1137/0914086, 1993.
- 710 Hill, T. C., Ryan, E., and Williams, M.: The use of CO₂ flux time series for parameter and carbon stock estimation in carbon cycle research, *Global Change Biology*, 18, 179–193, doi:10.1111/j.1365-2486.2011.02511.x, 2012.
- Johnson, C., Hoskins, B. J., and Nichols, N. K.: A singular vector perspective of 4D-Var: Filtering and interpolation, *Quarterly Journal of the Royal Meteorological Society*, 131, 1–19, doi:10.1256/qj.03.231, 2005.
- 715 Kemp, S., Scholze, M., Ziehn, T., and Kaminski, T.: Limiting the parameter space in the Carbon Cycle Data Assimilation System (CCDAS), *Geoscientific Model Development*, 7, 1609–1619, doi:10.5194/gmd-7-1609-2014, <http://www.geosci-model-dev.net/7/1609/2014/>, 2014.
- MacBean, N., Peylin, P., Chevallier, F., Scholze, M., and Schürmann, G.: Consistent assimilation of multiple data streams in a carbon cycle data assimilation system, *Geoscientific Model Development*, 9, 3569–3588, doi:10.5194/gmd-9-3569-2016, 2016.
- 720 Menke, W.: *Geophysical Data Analysis: Discrete Inverse Theory*, Academic Press, doi:10.1016/B978-0-12-490920-5.50001-6, 1984.
- Rayner, P. J., Scholze, M., Knorr, W., Kaminski, T., Giering, R., and Widmann, H.: Two decades of terrestrial carbon fluxes from a carbon cycle data assimilation system (CCDAS), *Global Biogeochemical Cycles*, 19, n/a–n/a, doi:10.1029/2004GB002254, <http://dx.doi.org/10.1029/2004GB002254>, gB2026, 2005.
- 725 Richardson, A. D., Williams, M., Hollinger, D. Y., Moore, D. J. P., Dail, D. B., Davidson, E. A., Scott, N. A., Evans, R. S., Hughes, H., Lee, J. T., Rodrigues, C., and Savage, K.: Estimating parameters of a forest ecosystem C model with measurements of stocks and fluxes as joint constraints, *Oecologia*, 164, 25–40, doi:10.1007/s00442-010-1628-y, 2010.
- 730

- Roose-Koerner, L., Devaraju, B., Sneeuw, N., and Schuh, W.-D.: A stochastic framework for inequality constrained estimation, *Journal of Geodesy*, 86, 1005–1018, doi:10.1007/s00190-012-0560-9, 2012.
- 735 Safta, C., Ricciuto, D. M., Sargsyan, K., Debusschere, B., Najm, H. N., Williams, M., and Thornton, P. E.: Global sensitivity analysis, probabilistic calibration, and predictive assessment for the data assimilation linked ecosystem carbon model, *Geoscientific Model Development*, 8, 1899–1918, doi:10.5194/gmd-8-1899-2015, <http://www.geosci-model-dev.net/8/1899/2015/>, 2015.
- Williams, M., Rastetter, E. B., Fernandes, D. N., Goulden, M. L., Shaver, G. R., and Johnson, L. C.: Predicting gross primary productivity in terrestrial ecosystems, *Ecological Applications*, 7, 882–894, doi:10.1890/1051-0761(1997)007[0882:PGPPIT]2.0.CO;2, 1997.
- 740 Williams, M., Schwarz, P., Law, B., Irvine, J., and Kurpius, M.: An improved analysis of forest carbon dynamics using data assimilation, *Global Change Biology*, 11, 89–105, doi:10.1111/j.1365-2486.2004.00891.x, 2005.
- Zhu, Q. and Zhuang, Q.: Parameterization and sensitivity analysis of a process-based terrestrial ecosystem model using adjoint method, *Journal of Advances in Modeling Earth Systems*, 6, 315–331, doi:10.1002/2013MS000241, 2014.
- 745 Ziehn, T., Scholze, M., and Knorr, W.: On the capability of Monte Carlo and adjoint inversion techniques to derive posterior parameter uncertainties in terrestrial ecosystem models, *Global Biogeochemical Cycles*, 26, n/a–n/a, doi:10.1029/2011GB004185, <http://dx.doi.org/10.1029/2011GB004185>, gB3025, 2012.