

An advanced method of contributing emissions to short-lived chemical species (OH and HO₂): The TAGGING 1.1 submodel based on the Modular Earth Submodel System (MESSy 2.53)

Vanessa S. Rieger^{1,2}, Mariano Mertens¹, and Volker Grewe^{1,2}

¹Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

²also at: Delft University of Technology, Aerospace Engineering, Section Aircraft Noise and Climate Effects, Delft, Netherlands

Correspondence to: Vanessa S. Rieger (vanessa.rieger@dlr.de)

Abstract. To mitigate human impact on climate change, it is essential to determine the contribution of emissions to the concentration of trace gases. In particular, source attribution of short-lived species such as OH and HO₂ is important as they play a crucial role for atmospheric chemistry. This study presents an advanced version of a tagging method for OH and HO₂ (HO_x) which attributes HO_x concentrations to emissions. While the former version V1.0 only considered 12 reactions in the troposphere, the new version V1.1, presented here, takes 19 reactions in the troposphere into account. For the first time, also the main chemical reactions for the HO_x chemistry in the stratosphere are regarded (in total 27 reactions). To fully take into account the main HO₂ source by the reaction of H and O₂, the tagging of H radical is introduced. In order to ensure the steady-state assumption, we introduce rest terms which balance the deviation of HO_x production and loss. This closes the budget between the sum of all contributions and the total concentration. The contributions to OH and HO₂ obtained by the advanced tagging method V1.1 deviate from V1.0 in certain source categories. For OH, major changes are found in the categories biomass burning, biogenic emissions and methane decomposition. For HO₂, the contributions differs strongly in the categories biogenic emissions and methane decomposition. As HO_x reacts with ozone (O₃), carbon monoxide (CO), reactive nitrogen compounds (NO_y), non-methane hydrocarbons (NMHC) and peroxyacyl nitrates (PAN), the contributions to these species are also modified by the advanced HO_x tagging method V1.1. The contributions to NO_y, NMHC and PAN show only little changes, whereas O₃ from biogenic emissions and methane decomposition increases in the tropical troposphere. Variations for CO from biogenic emissions and biomass burning are only found in the Southern Hemisphere.

1 Introduction

The radicals hydroxyl (OH) and hydroperoxyl (HO₂) are crucial for atmospheric chemistry. Both radicals are very reactive and have a lifetime of only a few seconds. OH is frequently converted to HO₂ and vice versa. Thus, OH and HO₂ radicals are closely linked and often referred together as chemical family HO_x. The ratio of OH to HO₂ in an air parcel strongly depends on the chemical background, in particular on the composition of nitrogen oxides NO_x (= NO + NO₂) and non-methane hydrocarbons (NMHC) (Heard and Pilling, 2003).

HO_x impacts global warming and local air quality in various ways: Reacting with greenhouse gases such as methane (CH₄) and ozone (O₃), OH reduces their atmospheric residence time (e.g. Stevenson et al. (2006); Voulgarakis et al. (2013); Righi et al. (2015)). Hence, HO_x controls the impact of CH₄ and O₃ on global warming. Moreover, being the main oxidizer in the troposphere, OH is involved in the decomposition of pollutants as well as in the production of ground-level ozone, photochemical smog and secondary organic aerosols (e.g. Lawrence et al. (2001); Heard and Pilling (2003)). Consequently, to quantify human impact on climate and air quality, it is essential to understand the distribution and variability of OH and HO₂ in the atmosphere.

However, the determination of OH and HO₂ concentrations in the atmosphere is still challenging due to their short lifetimes. In field campaigns HO_x concentrations are measured on a local scale which is generally difficult to compare with global models (e.g. Ren et al., 2003; Olson et al., 2006). For certain environments, such as marine boundary layer, model studies compare well with measurements. Other regions, such as unpolluted forest areas, show large discrepancies (Heard and Pilling, 2003; Stone et al., 2012). On regional and global scale, no direct HO_x measurement is available. So far, OH can only be estimated indirectly by measurements and emission rates of methyl chloroform (CH₃CCl₃) (Prinn et al., 2005). As emissions of CH₃CCl₃ steadily decline, Liang et al. (2017) suggest an alternative method: They combine several trace gases such as CH₂F₂, CH₂FCF₃, CH₃CHF₂ and CHClF₂ in a gradient-trend based two-box model approach to derive a global OH concentration of $11.2 \cdot 10^5$ molec cm⁻³. Overall, global chemistry climate models estimate a tropospheric OH concentration of around $11 \cdot 10^5$ molec cm⁻³ (Naik et al., 2013), which compares well with the observation-based results from Prinn et al. (2005) and Liang et al. (2017).

To mitigate human impact on climate change or pollution in general, it is crucial to determine the contribution of an emission sector to the concentration of certain chemical species (Grewe et al., 2012; Clappier et al., 2017). To do so, we use a "tagging" method: the theoretical framework of this tagging method is given in Grewe et al. (2010) and Grewe (2013), whereas the implementation is described in (Grewe et al., 2017). This method splits up all chemical species which are important for O₃ production and destruction into ten source categories: emissions from anthropogenic non-traffic (e.g. industry and households), road traffic, shipping, aviation, biogenic sources, biomass burning, lightning, methane (CH₄) and nitrous oxide (N₂O) decompositions and stratospheric ozone production. Subsequently, the contributions of these sources to the concentrations of O₃, CO, OH, HO₂, peroxyacyl nitrates (PAN), reactive nitrogen compounds (NO_y, e.g. NO, NO₂, HNO₄, ...) and non-methane hydrocarbons (NMHC) are diagnosed. The contribution calculations are based on chemical reaction rates, online emissions (e.g. lightning), offline emissions (e.g. road traffic) and deposition rates. It considers the competition of NO_y, CO and NMHC in producing and destroying O₃.

The tagging method of the long-lived species O₃, CO, PAN, NO_y and NMHC and of the short-lived species OH and HO₂ base on the same principles of apportioning the contributions. (In this study, O₃, CO, PAN, NO_y and NMHC are denoted as long-lived species because their atmospheric lifetime is significantly longer than the lifetime of OH and HO₂.) However, the implementation for long-lived and short-lived species differs. For the long-lived species, each source tracer is transported, receives the corresponding online or offline emissions, is deposited and reacts with other species. Based on these processes, the





Figure 1. Sketch of chemistry used in advanced tagging mechanism V1.1. Blue boxes indicate tagged long-lived species, orange boxes display tagged short-lived species.

tagging method determines the concentration of the source tracers. A detailed description of the implementation of the tagging method for long-lived species is given in Grewe et al. (2017).

However, the short-lived species HO_x are not transported and experience neither emissions nor deposition. Thus, the same implementation of the tagging method as for long-lived species is not possible. Tsati (2014) and Grewe et al. (2017) introduced a modified approach for tagging HO_x : since the lifetime of OH and HO_2 is very short, a steady-state between the production and destruction of OH and HO_2 is assumed. Using the main chemical reactions of HO_x chemistry, the contributions of each source category to OH and HO_2 are determined.

The contributions to long-lived and short-lived species are closely linked (see Fig. 1). For example, the reaction



involves the long-lived species O_3 and the short-lived species OH and HO_2 . Hence, this reaction is considered in the implementation of the tagging method for long-lived and short-lived species. The contribution of, for example, shipping emissions to O_3 influences the contribution of shipping emissions to HO_2 : the higher the contribution to O_3 is the more HO_2 is attributed to shipping emissions. Furthermore, OH from shipping emissions destroys O_3 and thus reduces the contribution of shipping emissions to O_3 .

The implementation of the tagging method for the short-lived species HO_x , presented by Grewe et al. (2017), is referred to HO_x tagging method V1.0. It did not consider all relevant reactions for the production and loss of HO_x . Especially, the reactions which are important in the stratosphere were not taken into account. Moreover, the steady-state assumption between

HO_x production and loss was not fulfilled. In this study, we present a revised version V1.1 of the HO_x tagging method, largely improving these shortcomings. It includes the main chemical reactions of HO_x chemistry in the troposphere and stratosphere. This is enabled by introducing the tagging of the hydrogen radical (H). Special care is taken for the steady-state assumption.

The paper is structured as follows: After introducing the model setup in Section 2, we present the advanced HO_x tagging method V1.1 in Section 3. In Section 4, the results are compared with the tagging method V1.0 by Grewe et al. (2017). Finally, Section 5 concludes the method and the results of this study.

2 Model description of EMAC and MECO(n)

To evaluate the further developed HO_x tagging method we use the same model setup as Grewe et al. (2017). A global climate simulation is performed with the ECHAM/MESSy Atmospheric Chemistry (EMAC) chemistry climate model. EMAC is a numerical chemistry and climate simulation system that includes submodels describing tropospheric and middle atmosphere processes and their interaction with oceans, land and human influences (Jöckel et al., 2010). It uses the second version of the Modular Earth Submodel System (MESSy2.53) to link multi-institutional computer codes. The core atmospheric model is the 5th generation European Centre Hamburg general circulation model (ECHAM5, Roeckner et al. (2006)). For the present study we apply EMAC in the T42L90MA-resolution, i.e. with a spherical truncation of T42 (corresponding to a quadratic Gaussian grid of approx. 2.8° by 2.8° in latitude and longitude) with 90 vertical hybrid pressure levels up to 0.01 hPa. For the simulation presented in this study, the time span of July 2007 to December 2008 is considered: half a year as a spin-up and one year for the analysis.

For the chemical scheme, we use the submodel MECCA (Module Efficiently Calculating the Chemistry of the Atmosphere) which is based on Sander et al. (2011) and Jöckel et al. (2010). The chemical mechanism includes 218 gas phase, 12 heterogeneous and 68 photolysis reactions. In total 188 species are considered. It regards the basic chemistry of OH, HO₂, O₃, CH₄, nitrogen oxides, alkanes, alkenes, chlorine and bromine. Alkynes, aromatics and mercury are not considered.

Total global emissions of lightning NO_x are scaled to approximately 4 Tg(N) a⁻¹ (parametrized according to Grewe et al. (2001)). The submodel ONEMIS (Kerkweg et al., 2006) calculates NO_x emissions from soil (parametrized according to Yienger and Levy (1995)) and biogenic C₅H₈ emissions (parametrized according to Guenther et al. (1995)). Direct CH₄ emissions are not considered, instead pseudo-emissions are calculated using the submodel TNUDGE (Kerkweg et al., 2006). This submodel relaxes the mixing ratios in the lowest model layer towards observations by Newtonian relaxation (more details are given by Jöckel et al. (2016)).

To show the effect of the HO_x tagging method on a regional scale, a further simulation with the coupled model system MESSyified ECHAM and COSMO models nested n-times (MECO(n)) is performed. The nested system couples the global chemistry climate model EMAC online with the regional chemistry climate model COSMO/MESSy (Kerkweg and Jöckel, 2012a, b). To test the HO_x tagging in MECO(n), we conduct a simulation using one COSMO/MESSy nest over Europe with a resolution of 0.44°. EMAC is applied in a horizontal resolution of T42 with 31 vertical levels. The period from July 2007 to

December 2008 is simulated. The setup of the simulation is identical to the one described in Grewe et al. (2017). A detailed chemical evaluation of the setup is given in Mertens et al. (2016).


Both model simulations are based on the quasi chemistry-transport model (QCTM) mode in which the chemistry is decoupled from the dynamics (Deckert et al., 2011). The anthropogenic emissions are taken from the MACCity emission inventory (Granier et al., 2011). The TAGGING submodel (as described by Grewe et al. (2017)) is coupled to the detailed chemical solver MECCA from which it obtains information about tracer concentrations and reaction rates. Based on this information, it calculates the contributions of source categories to O₃, CO, NO_y, PAN and NMHC concentration. The contributions of OH and HO₂ are calculated with the advanced method V1.1 presented in the next section. The implementation is based on MESSy2.53 and will be available in MESSy2.54.

10 3 Tagging method of short-lived species

3.1 Tagging method V1.0

The tagging method V1.0 described by Grewe et al. (2017) determines the contribution of source categories to O₃, NO_y, CO, NMHC, PAN, OH and HO₂ concentrations. Ten source categories are considered and every species included in the tagging method is decomposed into these categories: For example, the concentration of O₃ is split up into O₃ from anthropogenic non-traffic (e.g. industry) emissions (O₃^{ant}), road traffic emissions (O₃^{tra}), ship emissions (O₃^{shp}), air traffic emissions (O₃^{air}), biogenic emissions (O₃^{bio}), biomass burning (O₃^{bb}), lightning (O₃^{lig}), methane decomposition (O₃^{CH₄}), nitrous oxide decomposition (O₃^{N₂O}) and stratospheric ozone production (O₃^{str}). These tagged species go through the same chemical reactions and the same deposition loss processes as O₃. The tagging method uses a combinatoric approach to determine the contributions: It redistributes the production and loss rates of each species to the ten source categories according to the concentrations of the tagged species. Details on the tagging theory and implementation in EMAC and MECO(n) are found in Grewe (2013) and Grewe et al. (2017), respectively.

For the first time, V1.0 determined the contribution of source categories to OH and HO₂ concentrations. The tagging method V1.0 was based on 12 reactions for the HO_x chemistry (reactions marked with "o" in last column of Table 1). It included the main production and loss reactions of HO_x with O₃, NO_y, NMHC, CO and CH₄. V1.0 only regarded reactions which are important in the troposphere. Reactions which mainly occur in the stratosphere were not taken into account. However, the main HO₂ production by the reaction (1) H + O₂ → HO₂ (see Table 1) was not regarded. It was combined with reaction (11) CO + OH → H + CO₂ (see Table 1) to



But not all H radicals in the troposphere are produced by the reaction of CO + OH. Also the reactions (7) OH + O(³P), (10) H₂ + OH and (28) HCHO + *hν* produce H (Table 2). These reactions were neglected in V1.0. Thus, only 80 % of the H production

and therefore only 80 % of the HO₂ production by reaction (1) was considered in the troposphere. In the stratosphere, the reaction of CO + OH becomes less important and most of H is produced by reactions (7) and (28). Consequently, only 6 % of the H and also of HO₂ production by reaction (1) was regarded by this approach. (Numbers are derived from a EMAC simulation as described in Sect. 2.)

5 In the troposphere, the most important reactions not covered in V1.0 are reaction (1) H + O₂ as well as reaction (15) NO₂ + HO₂ and (18) decomposition of HNO₄. In the stratosphere, reactions (1) H + O₂, (5) HO₂ + O(³P) and (7) OH + O(³P) play a leading role and were not included in V1.0.

Most reaction rates used in the tagging method corresponds to the production and loss rates directly provided by the chemical scheme MECCA of EMAC. However for reactions with NMHC, the reaction rates were obtained indirectly. The reaction rate
10 of OH with NMHC (reaction 21, Table 1) was determined via the production rates of CO by assuming that each reaction of OH with NMHC produces one CO molecule. This method neglects all intermediate oxidation reactions of NMHC and considers only these reactions when NMHC is finally oxidized to CO. For the reaction rates of NO_y and HO₂ with NMHC (reaction 22 and 23), only the reaction of HO₂ with methylperoxy radical (CH₃O₂) was considered.

To derive the contributions to OH and HO₂, steady-state between HO_x production and loss was assumed. However, the
15 steady-state assumption was not completely fulfilled for V1.0 (see Sect. 3.4). Moreover, the sum over the contributions of the ten source categories to the OH and HO₂ concentrations did not equal the total OH and HO₂ concentrations. It deviated by about 70 %.

3.2 Reduced HO_x reaction system V1.1

OH and HO₂ react with many chemical species. To reduce the calculation time of a simulation, we reduce the HO_x chemistry
20 used in chemical scheme MECCA to the most important reactions which occur in the troposphere and stratosphere. We consider only reactions with a tropospheric or stratospheric annual mean reaction rate larger than 10⁻¹⁵ mol mol⁻¹s⁻¹ (see Table 1). Hence, we increase the number of reactions from 12 (V1.0) to 27 (V1.1), which still constitutes a reduced set of reactions compared to the full chemical scheme MECCA used in EMAC. In the following, we call this set *reduced HO_x reaction system V1.1*.

25 The reactions which are important in the troposphere are indicated in Table 1. As stated above, reaction (1) of H and O₂ dominates the HO₂ production in the troposphere. It produces 49 % of tropospheric HO₂. In V1.0, only part of this HO₂ source was regarded (see Sect. 3.1). The most important HO₂ loss is the reaction with NO (reaction 14) followed by the reaction with itself producing H₂O₂ (reaction 3) which accounts for 32 % and 12 % of tropospheric HO₂ loss. The production via H₂O and O(¹D) produces about 21 % of tropospheric OH (reaction 2). The excited oxygen radical (O(¹D)) originates from
30 the photolysis of O₃. Also reaction (14) of NO and HO₂ produces 32 % of tropospheric OH. OH is mostly destroyed by CO (reaction 11, 38 %) followed by NMHC (reaction 21, 25 %).

In the stratosphere different chemical reactions become important. Here, OH is mainly destroyed by O₃, producing 40 % of stratospheric HO₂. The reaction is partly counteracted by the reaction (14) which produces 21 % of OH and destroys 24 % of

	reaction		rates	tropos.	stratos.	V1.1
1	$\text{H} + \text{O}_2 \longrightarrow \text{HO}_2$		R_1	x	x	(x)
2	$\text{H}_2\text{O} + \text{O}(^1\text{D}) \longrightarrow 2 \text{OH}$		R_2	x	x	o
3	$\text{HO}_2 + \text{HO}_2 \longrightarrow \text{H}_2\text{O}_2 + \text{O}_2$		R_3	x		o
4	$\text{HO}_2 + \text{O}_3 \longrightarrow \text{OH} + 2 \text{O}_2$		R_4	x	x	o
5	$\text{HO}_2 + \text{O}(^3\text{P}) \longrightarrow \text{OH} + \text{O}_2$		R_5		x	x
6	$\text{OH} + \text{O}_3 \longrightarrow \text{HO}_2 + \text{O}_2$		R_6	x	x	o
7	$\text{OH} + \text{O}(^3\text{P}) \longrightarrow \text{H} + \text{O}_2$		R_7		x	x
8	$\text{HO}_2 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{O}_2$		R_8	x	x	o
9	$\text{H}_2\text{O}_2 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{HO}_2$		R_9	x		x
10	$\text{H}_2 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{H}$		R_{10}	x		x
11	$\text{CO} + \text{OH} \longrightarrow \text{H} + \text{CO}_2$		R_{11}	x	x	o
12	$\text{CH}_4 + \text{OH} \longrightarrow \text{CH}_3 + \text{H}_2\text{O}$		R_{12}	x	x	o
13	$\text{ClO} + \text{OH} \longrightarrow 0.94 \text{Cl} + 0.94 \text{HO}_2 + 0.06 \text{HCl} + 0.06 \text{O}_2$		R_{13}		x	x
14	$\text{NO} + \text{HO}_2 \longrightarrow \text{NO}_2 + \text{OH}$		R_{14}	x	x	o
15	$\text{NO}_2 + \text{HO}_2 \longrightarrow \text{HNO}_4$		R_{15}	x	x	x
16	$\text{NO} + \text{OH} \longrightarrow \text{HONO}$		R_{16}		x	x
17	$\text{NO}_2 + \text{OH} \longrightarrow \text{HNO}_3$		R_{17}		x	o
18	$\text{HNO}_4 \longrightarrow \text{NO}_2 + \text{HO}_2$		R_{18}	x		x
19	$\text{HONO} + h\nu \longrightarrow \text{NO} + \text{OH}$		R_{19}		x	x
20	$\text{HNO}_3 + h\nu \longrightarrow \text{NO}_2 + \text{OH}$		R_{20}		x	x
21	$\text{NMHC} + \text{OH} \longrightarrow \text{NMHC}$		R_{21}	x		o
22	$\text{NMHC} + \text{HO}_2 \longrightarrow \text{NMHC}$		R_{22}	x		o
23	$\text{NMHC} + \text{NO}_y \longrightarrow \text{HO}_2 + \text{NMHC} + \text{NO}_y$		R_{23}	x	x	o
24	$\text{NMHC} + \text{OH} \longrightarrow \text{NMHC} + \text{HO}_2$		R_{24}	x		x
25	$\text{NMHC} + h\nu \longrightarrow \text{NMHC} + \text{HO}_2$		R_{25}	x		x
26	$\text{ClO} + \text{HO}_2 \longrightarrow \text{HOCl} + \text{O}_2$		R_{26}		x	x
27	$\text{BrO} + \text{HO}_2 \longrightarrow \text{HOBr} + \text{O}_2$		R_{27}		x	x

Table 1. Reduced HO_x reaction system V1.1 describes the main reactions of HO_x chemistry in troposphere and stratosphere. These 27 reactions are used for the tagging method V1.1. In the column "tropos." ("stratos."), reactions which are important in the troposphere (stratosphere) are marked. In the column "V1.1", reactions marked with "o" were already included in V1.0. Reactions marked with "x" are added in V1.1. Reactions marked with "(x)" were only partly taken into account in V1.0. The numbers of reactions are referenced in the text.

HO₂. Since large quantities of O₃ are found in the stratosphere, O₃ or the excited oxygen radical (O(³P)) destroys about 62 % of HO₂. Reactions with NMHC, CO and CH₄ play only a minor role in the stratosphere.

Reactions of OH and HO₂ with chlorine and bromide were not considered in V1.0. We add these reactions, which take place only in the stratosphere, to the tagging method V1.1. Reactions (21) to (25) involve the chemical family NMHC which contains several species such as formaldehyde (HCHO), ethylene (C₂H₄) and propane (C₃H₈). The rate for reaction (21) is determined by adding up the rates of all reactions of OH with each single species of the family NMHC. Reaction rate (23) contains all rates of the reactions between the species of the chemical families NO_y and NMHC. All reaction rates are directly derived by MECCA mechanism of EMAC.

Table 1 does not consider all reactions with annual reaction rates larger than 10⁻¹⁵ mol mol⁻¹s⁻¹. The photolysis of hydrogen peroxide (H₂O₂), hypochlorous acid (HOCl) and hypobromous acid (HOBr) are excluded from the reduced HO_x reaction system V1.1 as the tagging method can not be applied. The specific reasons are explained in Appendix A.

3.3 Deductions of tagged species

To derive how much OH and HO₂ is produced and destroyed by a source category *i*, the tagging approach described in Grewe et al. (2010, 2017) is used. In general, bimolecular reactions with two chemical species A + B → C are tagged as follows:

Each tagged species is split up into its contribution from *n* source categories $A = \sum_{i=1}^n A^i$, $B = \sum_{i=1}^n B^i$ and $C = \sum_{i=1}^n C^i$. These contributions (A^i, B^i, C^i) go through the same reactions as their main species (A, B, C). If A from category *i* reacts with B from category *j*, then the resulting species C belongs half to the category *i* and half to the category *j*:



Consequently, the production *P* and loss *L* of a species from the category *i* (here $LossA^i$, $LossB^i$ and $ProdC^i$) are determined by regarding all possible combinations of the reaction between A^i and B^j :

$$LossA^i = LossB^i = ProdC^i = k \left(A^i B^i + \sum_{j=1, j \neq i}^n \frac{1}{2} A^i B^j + \sum_{j=1, j \neq i}^n \frac{1}{2} A^j B^i \right) = \frac{1}{2} R \left(\frac{A^i}{A} + \frac{B^i}{B} \right) \quad (2)$$

with *k* being the reaction rate coefficient and $R = kAB$ being the respective reaction rate. For unimolecular reactions A → B + C, the distribution of categories from the educts is completely passed to the products:

$$LossA^i = ProdB^i = ProdC^i = R \frac{A^i}{A} \quad (3)$$

with the reaction rate $R = kA$.

As described above, the long-lived species O₃, CO, NO_y and NMHC are tagged according to tagging method described in Grewe et al. (2017). Due to limited computational resources, other species such as H₂, H₂O₂, CH₄, ClO and BrO are not tagged (as in V1.0). Here, different approaches are derived to retain the ratio of contribution to total concentration $\frac{A^i}{A}$:

1. If a tagged species reacts with a non-tagged species, the non-tagged species does not contribute and the tagging method for a unimolecular reaction is applied (see equation 3). Examples are reactions (9), (10) and (14).

	reaction		rates	tropos.	stratos.	
1	H + O ₂	→	HO ₂	R ₁	x	x
7	OH + O(³ P)	→	H + O ₂	R ₇		x
10	H ₂ + OH	→	H ₂ O + H	R ₁₀	x	
11	CO + OH	→	H + CO ₂	R ₁₁	x	x
28	HCHO + O ₂ + hv	→	H + CO + HO ₂	R ₂₈	x	

Table 2. Reduced H reaction system describes the main reactions of H. In the column "tropos." ("stratos."), reactions which are important in the troposphere (stratosphere) are marked. The numbers of the reactions correspond to the numbers in Table 1.

2. Using the family concept as described in Grewe et al. (2017) allows the assumption that all tags are distributed equally among the species within the same chemical family. It follows:

$$\frac{NO^i}{NO} = \frac{NO_2^i}{NO_2} = \frac{HNO_4^i}{HNO_4} = \frac{NO_y^i}{NO_y} \quad (4)$$

- 5 As mentioned in Grewe et al. (2017), all species which are frequently converted back and forth to ozone are considered as an "ozone storage" (Crutzen and Schmailzl, 1983). These species together with O₃ are lumped into one chemical family "ozone". Both O(¹D) and O(³P) belong to this chemical family. Hence, as in Grewe et al. (2017), we apply the family concept and set:

$$\frac{O(^1D)^i}{O(^1D)} = \frac{O(^3P)^i}{O(^3P)} = \frac{O_3^i}{O_3} \quad (5)$$

- 10 3. In reaction (1), neither H nor O₂ are tagged. To obtain the ratio $\frac{HO_2^i}{HO_2}$, we set up an extra tagging of H itself. As the H radical is very reactive, we assume that H production balances H loss (see Sect. 3.4). Table 2 presents the main reactions for H which still constitute a subset of full H chemistry implemented in MECCA. Based on Table 2, we set up the H production $ProdH^i$ and H loss $LossH^i$ for the contribution of a source category i :

$$ProdH^i = \frac{1}{2}R_7 \left(\frac{OH^i}{OH} + \frac{O_3^i}{O_3} \right) + R_{10} \frac{OH^i}{OH} + \frac{1}{2}R_{11} \left(\frac{CO^i}{CO} + \frac{OH^i}{OH} \right) + R_{28} \frac{NMHC^i}{NMHC} \quad (6)$$

$$LossH^i = R_1 \frac{H^i}{H} \quad (7)$$

- 15 As mentioned above, the family concept also sets $\frac{HCHO^i}{HCHO} = \frac{NMHC^i}{NMHC}$. Since the steady-state assumption applies for H (see Sect. 3.4), the H production per source category i $ProdH^i$ equals the loss $LossH^i$. After setting eq. (6) and (7) equal to each other, we obtain:

$$\frac{H^i}{H} = \frac{1}{2} \frac{R_7}{R_1} \left(\frac{OH^i}{OH} + \frac{O_3^i}{O_3} \right) + \frac{R_{10}}{R_1} \frac{OH^i}{OH} + \frac{1}{2} \frac{R_{11}}{R_1} \left(\frac{CO^i}{CO} + \frac{OH^i}{OH} \right) + \frac{R_{28}}{R_1} \frac{NMHC^i}{NMHC} \quad (8)$$

- 20 These different approaches are applied to the reduced HO_x reaction system V1.1 (Table 1) to derive the contributions of source categories to OH and HO₂ in Sect. 3.5.

		OH		HO ₂		H	
		prod.	loss	prod.	loss	prod.	loss
total - MECCA	tropos.	0.49	0.49	0.49	0.49	0.24	0.24
	<i>stratos.</i>	<i>2.78</i>	<i>2.78</i>	<i>2.48</i>	<i>2.48</i>	<i>7.09</i>	<i>7.09</i>
reduced - V1.1	tropos.	0.43	0.48	0.47	0.49	0.24	0.24
	<i>stratos.</i>	<i>2.49</i>	<i>2.76</i>	<i>2.47</i>	<i>2.48</i>	<i>7.06</i>	<i>5.99</i>
reduced - V1.0	tropos.	0.43	0.47	0.29	0.42	-	-
	<i>stratos.</i>	<i>0.86</i>	<i>1.30</i>	<i>1.19</i>	<i>0.84</i>	-	-

Table 3. Annual mean of OH, HO₂ and H production and loss rates (air mass weighted) in 10⁻¹³ mol mol⁻¹ s⁻¹ for the total rates (derived from the complete chemical scheme MECCA in EMAC) and for the rates of the reduced reaction system of the tagging method V1.0 and V1.1. The first row gives the rates for the troposphere, the second row for the stratosphere (written in italic).

3.4 Steady-state assumption

The steady-state assumption of the HO_x chemistry is the basic principle of the tagging method for short-lived species (Tsati, 2014; Grewe et al., 2017). In steady-state, the production and loss of OH and HO₂ balance each other. Table 3 shows annual means of HO_x and H production and loss rates of the reduced reaction system for the tagging methods V1.0 and V1.1 as well as the total production and loss rates derived from the complete chemical scheme MECCA in EMAC. The production and loss rates are obtained from an EMAC simulation following the setup described in Sect. 2. Note that for V1.0 no values for the H production and loss are available since the tagging of H was not considered in V1.0.

In general, total OH production (derived by MECCA) equals total OH loss in the troposphere and stratosphere. The same holds for HO₂ and H. In the troposphere, the OH loss of V1.1 and V1.0 represents well the total OH loss in the troposphere. However, the OH production for V1.1 and V1.0 differs by 12 % from the total OH production. Considering HO₂ in the troposphere, the total production and loss rates are well reflected by V1.1. In contrast, the HO₂ production and loss of V1.0 differs by 14 % and 41 % from the total rates.

In the stratosphere, V1.1 represents the total rates very well. However, the OH production of V1.1 misses 10 % of the total OH production. Since V1.0 was only developed for the troposphere, not all reactions which are important in the stratosphere were considered. Thus, the OH and HO₂ production and loss rates of V1.0 considerably underestimated the total production and loss rates.

The reduced H reaction system in V1.1 (Table 2) represents the total H production and loss in the troposphere very well. However in the stratosphere, H loss in V1.1 deviates by 17 % from the total H loss.

Summing up, the reduced HO_x reaction system V1.1 represents well the total HO_x production and loss in the troposphere and stratosphere. V1.1 reproduces the HO_x chemistry better than V1.0. However, OH production in troposphere and stratosphere as

well as H loss in the stratosphere of V1.1 deviates from the total rates derived by MECCA. Thus, the state-state for the reduced HO_x and H reaction system (Tables 1 and 2) is not completely fulfilled.

But steady-state between production and loss is crucial for the tagging method for short-lived species. To re-establish steady-state, it would be necessary to include the complete HO_x and H chemistry in the tagging method. However, this is not possible as the tagging method does not apply to all reactions of the HO_x and H chemistry (for examples see Appendix A). Consequently, we introduce rest terms *resOH*, *resHO₂* and *resH* for OH, HO₂ and H to compensate for the deviations from steady-state. Each rest term is calculated by subtracting the production rate of the reduced reaction system from the loss rate (Tables 1 and 2). The resulting rest terms are shown in the Supplement (Fig. S1).

Considering the rest terms *resOH*, *resHO₂* and *resH* leads to the closure of the budget. In V1.0, the sum over the contributions from all source categories did not balance the total concentration. The averaged deviations for OH and HO₂ in troposphere were about 70 % of the total concentrations. Since the stratosphere was not considered in V1.0, the deviations were even larger (104 % for OH and 89 % for HO₂). In V1.1, the sum of OH and HO₂ now perfectly balances the total OH and HO₂ concentrations. The deviations are negligible (below 10⁻⁴ % for OH and below 10⁻³ % for HO₂). Consequently, including the rest terms to the tagging method is mandatory for the steady-state assumption and also closes the budget.

3.5 Determination of HO_x contributions

Taking the above considerations into account, we finally derive the OH and HO₂ production and loss terms per source category *i*. In the reduced HO_x reaction system V1.1 (Table 1), OH is produced by the reactions (2) H₂O + O(¹D), (4) HO₂ + O₃, (5) HO₂ + O(³P), (14) NO + HO₂, (19) HONO + *hν* and (20) HNO₃ + *hν*. Applying the partitioning described in Sect. 3.3, the OH production for a source category *i* *ProdOHⁱ* is determined as follows:

$$\begin{aligned}
 ProdOH^i = & 2 \cdot R_2 \frac{O_3^i}{O_3} + \frac{1}{2} R_4 \left(\frac{HO_2^i}{HO_2} + \frac{O_3^i}{O_3} \right) + \frac{1}{2} R_5 \left(\frac{HO_2^i}{HO_2} + \frac{O_3^i}{O_3} \right) + \frac{1}{2} R_{14} \left(\frac{NO_y^i}{NO_y} + \frac{HO_2^i}{HO_2} \right) \\
 & + R_{19} \frac{NO_y^i}{NO_y} + R_{20} \frac{NO_y^i}{NO_y}
 \end{aligned} \tag{9}$$

OH is destroyed by the reactions (6) OH + O₃, (7) OH + O(³P), (8) HO₂ + OH, (9) H₂O₂ + OH, (10) H₂ + OH, (11) CO + OH, (12) CH₄ + OH, (13) ClO + OH, (16) NO + OH, (17) NO₂ + OH, (21) NMHC + OH and (24) NMHC + OH. The OH loss per source category *i* *LossOHⁱ* is:

$$\begin{aligned}
 LossOH^i = & \frac{1}{2} R_6 \left(\frac{OH^i}{OH} + \frac{O_3^i}{O_3} \right) + \frac{1}{2} R_7 \left(\frac{OH^i}{OH} + \frac{O_3^i}{O_3} \right) + \frac{1}{2} R_8 \left(\frac{HO_2^i}{HO_2} + \frac{OH^i}{OH} \right) + \frac{1}{2} R_9 \left(\frac{HO_2^i}{HO_2} + \frac{OH^i}{OH} \right) \\
 & + R_{10} \frac{OH^i}{OH} + \frac{1}{2} R_{11} \left(\frac{CO^i}{CO} + \frac{OH^i}{OH} \right) + R_{12} \frac{OH^i}{OH} + R_{13} \frac{OH^i}{OH} + \frac{1}{2} R_{16} \left(\frac{NO_y^i}{NO_y} + \frac{OH^i}{OH} \right) \\
 & + \frac{1}{2} R_{17} \left(\frac{NO_y^i}{NO_y} + \frac{OH^i}{OH} \right) + \frac{1}{2} R_{21} \left(\frac{NMHC^i}{NMHC} + \frac{OH^i}{OH} \right) + \frac{1}{2} R_{24} \left(\frac{NMHC^i}{NMHC} + \frac{OH^i}{OH} \right)
 \end{aligned} \tag{10}$$

HO₂ is produced by reactions (1) H + O₂, (6) OH + O₃, (9) H₂O₂ + OH, (13) ClO + OH, (18) HNO₄, (23) NMHC + NO_y, (24) NMHC + OH and (25) NMHC + *hν*. However, H in reaction (1) is not tagged. To be able to determine the HO₂ production by

reaction (1) $R_1 \frac{H^i}{H}$, we apply the introduced H tagging (see Sect. 3.3) and replace $\frac{H^i}{H}$ with equation (8). Besides, reaction (13) constitutes a simplified reaction producing $0.94 \cdot \text{HO}_2$. Consequently, the HO_2 production per source category i $ProdHO_2^i$ is:

$$\begin{aligned}
ProdHO_2^i = & \frac{1}{2}R_6 \left(\frac{OH^i}{OH} + \frac{O_3^i}{O_3} \right) + \frac{1}{2}R_7 \left(\frac{OH^i}{OH} + \frac{O_3^i}{O_3} \right) + \frac{1}{2}R_9 \left(\frac{HO_2^i}{HO_2} + \frac{OH^i}{OH} \right) + R_{10} \frac{OH^i}{OH} \\
& + \frac{1}{2}R_{11} \left(\frac{CO^i}{CO} + \frac{OH^i}{OH} \right) + 0.94 \cdot R_{13} \frac{OH^i}{OH} + R_{18} \frac{NO_y^i}{NO_y} + \frac{1}{2}R_{23} \left(\frac{NMHC^i}{NMHC} + \frac{NO_y^i}{NO_y} \right) \\
& + \frac{1}{2}R_{24} \left(\frac{NMHC^i}{NMHC} + \frac{OH^i}{OH} \right) + R_{25} \frac{NMHC^i}{NMHC} + R_{28} \frac{NMHC^i}{NMHC}
\end{aligned} \tag{11}$$

5 The HO_2 loss is determined by reactions (3) $\text{HO}_2 + \text{HO}_2$, (4) $\text{HO}_2 + \text{O}_3$, (5) $\text{HO}_2 + \text{O}(\text{}^3\text{P})$, (8) $\text{HO}_2 + \text{OH}$, (14) $\text{NO} + \text{HO}_2$, (15) $\text{NO}_2 + \text{HO}_2$, (22) $\text{NMHC} + \text{HO}_2$, (26) $\text{ClO} + \text{HO}_2$ and (27) $\text{BrO} + \text{HO}_2$. Hence, the HO_2 loss per source category i $LossHO_2^i$ is:

$$\begin{aligned}
LossHO_2^i = & R_3 \frac{HO_2^i}{HO_2} + \frac{1}{2}R_4 \left(\frac{HO_2^i}{HO_2} + \frac{O_3^i}{O_3} \right) + \frac{1}{2}R_5 \left(\frac{HO_2^i}{HO_2} + \frac{O_3^i}{O_3} \right) + \frac{1}{2}R_8 \left(\frac{HO_2^i}{HO_2} + \frac{OH^i}{OH} \right) \\
& + \frac{1}{2}R_{14} \left(\frac{NO_y^i}{NO_y} + \frac{HO_2^i}{HO_2} \right) + \frac{1}{2}R_{15} \left(\frac{NO_y^i}{NO_y} + \frac{HO_2^i}{HO_2} \right) + \frac{1}{2}R_{22} \left(\frac{NMHC^i}{NMHC} + \frac{HO_2^i}{HO_2} \right) \\
& + R_{26} \frac{HO_2^i}{HO_2} + R_{27} \frac{HO_2^i}{HO_2}
\end{aligned} \tag{12}$$

Sect. 3.4 shows that the steady-state assumption for OH and HO_2 is justified when the rest terms $resOH$, $resHO_2$ and $resH$ are regarded. Therefore, the rest terms are divided by the number of source categories n to add them to the contributions of a category i . In steady-state, production of OH^i and HO_2^i equals the loss:

$$ProdOH^i - LossOH^i + resOH/n = 0 \tag{13}$$

$$ProdHO_2^i - LossHO_2^i + resHO_2/n + resH/n = 0 \tag{14}$$

The equations (13) and (14) are rewritten as follows:

$$15 \quad 0 = A^i - L^{OH} \frac{OH^i}{OH} + P^{OH} \frac{HO_2^i}{HO_2} + \frac{resOH}{n} \tag{15}$$

$$0 = B^i + P^{HO_2} \frac{OH^i}{OH} - L^{HO_2} \frac{HO_2^i}{HO_2} + \frac{resHO_2}{n} + \frac{resH}{n} \tag{16}$$

with the variables P^{OH} , L^{OH} , P^{HO_2} , L^{HO_2} , A^i and B^i (compare to Grewe et al. (2017) equations (25) to (28)):

$$P^{OH} = \frac{1}{2}R_4 + \frac{1}{2}R_5 + \frac{1}{2}R_{14} - \frac{1}{2}R_8 \tag{17}$$

$$L^{OH} = \frac{1}{2}R_6 + \frac{1}{2}R_7 + \frac{1}{2}R_8 + R_9 + R_{10} + \frac{1}{2}R_{11} + R_{12} + R_{13} + \frac{1}{2}R_{16} + \frac{1}{2}R_{17} + \frac{1}{2}R_{21} + \frac{1}{2}R_{24} \tag{18}$$

$$20 \quad P^{HO_2} = \frac{1}{2}R_6 + \frac{1}{2}R_7 + R_9 + R_{10} + \frac{1}{2}R_{11} + 0.94 \cdot R_{13} + \frac{1}{2}R_{24} - \frac{1}{2}R_8 \tag{19}$$

$$L^{HO_2} = 2 \cdot R_3 + \frac{1}{2}R_4 + \frac{1}{2}R_5 + \frac{1}{2}R_8 + \frac{1}{2}R_{14} + \frac{1}{2}R_{15} + \frac{1}{2}R_{22} + R_{26} + R_{27} \tag{20}$$

$$\begin{aligned}
A^i = & 2 \cdot R_2 \frac{O_3^i}{O_3} + \frac{1}{2} R_4 \frac{O_3^i}{O_3} + \frac{1}{2} R_5 \frac{O_3^i}{O_3} + \frac{1}{2} R_{14} \frac{NO_y^i}{NO_y} + R_{19} \frac{NO_y^i}{NO_y} + R_{20} \frac{NO_y^i}{NO_y} \\
& - \frac{1}{2} R_6 \frac{O_3^i}{O_3} - \frac{1}{2} R_7 \frac{O_3^i}{O_3} - \frac{1}{2} R_{11} \frac{CO^i}{CO} - \frac{1}{2} R_{16} \frac{NO_y^i}{NO_y} - \frac{1}{2} R_{17} \frac{NO_y^i}{NO_y} - \frac{1}{2} R_{21} \frac{NMHC^i}{NMHC} - \frac{1}{2} R_{24} \frac{NMHC^i}{NMHC}
\end{aligned} \quad (21)$$

$$\begin{aligned}
B^i = & \frac{1}{2} R_6 \frac{O_3^i}{O_3} + \frac{1}{2} R_7 \frac{O_3^i}{O_3} + \frac{1}{2} R_{11} \frac{CO^i}{CO} + R_{18} \frac{NO_y^i}{NO_y} + \frac{1}{2} R_{23} \left(\frac{NMHC^i}{NMHC} + \frac{NO_y^i}{NO_y} \right) + \frac{1}{2} R_{24} \frac{NMHC^i}{NMHC} \\
& + R_{25} \frac{NMHC^i}{NMHC} + R_{28} \frac{NMHC^i}{NMHC} - \frac{1}{2} R_4 \frac{O_3^i}{O_3} - \frac{1}{2} R_5 \frac{O_3^i}{O_3} - \frac{1}{2} R_{14} \frac{NO_y^i}{NO_y} - \frac{1}{2} R_{15} \frac{NO_y^i}{NO_y} - \frac{1}{2} R_{22} \frac{NMHC^i}{NMHC}
\end{aligned} \quad (22)$$

5 Solving equations (15) and (16), we finally obtain the contributions of a source category i to the OH and HO₂ concentration (same equations as equations (29) and (30) in Grewe et al. (2017), but with differently defined coefficients):

$$\frac{OH^i}{OH} = \frac{A^i L^{HO_2} + B^i P^{OH}}{L^{OH} L^{HO_2} - P^{OH} P^{HO_2}} \quad (23)$$

$$\frac{HO_2^i}{HO_2} = \frac{A^i P^{HO_2} + B^i L^{OH}}{L^{OH} L^{HO_2} - P^{OH} P^{HO_2}} \quad (24)$$

10 These equations are implemented in the TAGGING submodel and EMAC and MECO(n) simulations according to Sect. 2 are performed. The results for the OH and HO₂ contributions are analysed and compared with V1.0 in the following Section.

4 Results of model simulations

4.1 Contribution of short-lived species (HO_x)

15 Figures 2 and 3 show the zonal mean of OH and HO₂ contributions up to 200 hPa for the ten source categories derived by V1.1 (first and third columns) and V1.0 (second and forth columns). The zonal mean of OH and HO₂ contributions from 1 to 200 hPa are shown in Appendix B (Figs. B1, B2). First, the OH and HO₂ contributions of V1.1 are described in the following. For the categories which are determined by anthropogenic emissions, such as "shipping", "road traffic" and "anthropogenic non-traffic", the maximum values of OH and HO₂ contributions occur in the lower troposphere in the Northern Hemisphere. This clearly shows that for anthropogenic dominated categories the OH and HO₂ contributions are caused by anthropogenic emissions. The contributions vary among these categories of surface emissions as not only the amount but also the composition of the emissions differs. For the category "aviation", maximum OH contribution are found in the Northern Hemisphere between 200 and 250 hPa. However, HO₂ contribution has a minimum in this region and a maximum in the lower troposphere. The OH values for the categories "CH₄ decomposition", "N₂O decomposition", "lightning" and "biogenic emissions" are largest in the upper troposphere. Most OH contributions of "biomass burning" are found in lower tropical troposphere. In contrast, negative values occur in the upper tropical troposphere. Concerning the HO₂ contribution, the residual categories show a maximum in the tropical lower troposphere. In addition, the category "lightning" shows a strong HO₂ loss in the upper tropical troposphere which is caused by reaction (14).

OH




Figure 2. Contribution of ten source categories to OH in $10^{-15} \text{ mol mol}^{-1}$. Zonal means of the year 2008 are shown. First and third columns show the tagging method V1.1. Second and fourth columns show the tagging method V1.0. Simulation is performed with EMAC.

The results obtained by V1.1 are compared to the OH and HO₂ zonal profiles of V1.0 only in the troposphere (Figs. 2 and 3). The HO_x tagging method V1.0 was only developed for the troposphere. Hence, a comparison in the stratosphere is not

HO₂




Figure 3. Contribution of ten source categories to HO₂ in 10⁻¹⁵ mol mol⁻¹. Zonal means of the year 2008 are shown. First and third columns show the tagging method V1.1. Second and fourth columns show the tagging method V1.0. Simulation is performed with EMAC.

reasonable. In general, contributions to OH and HO₂ concentrations of V1.1 are larger in the troposphere compared to V1.0. This overall shift towards larger values is explained by the re-establishment of the steady-state and thus the closure of the budget in V1.1. In V1.0 the budget was not closed and thus the contributions were underestimated.

For OH, the categories "lightning" and "aviation" show no large changes in the general pattern of the zonal means between V1.0 and V1.1. Considering the HO₂ contributions, no large changes are found for the categories "biomass burning", "anthropogenic non-traffic", "road traffic" and "shipping".

The contribution of the category "aviation" to HO₂ in V1.1 shows roughly the same pattern compared to V1.0. However, the HO₂ destruction along the flight path is not as pronounced any more which is caused by the inclusion of reaction (15) and (18) to V1.1. Reaction (15) adds the term $\frac{1}{2}R_{15}\frac{NO_y^i}{NO_y}$ to the HO₂ loss (eq. 12) and reaction (18) adds the term $R_{18}\frac{NO_y^i}{NO_y}$ to the HO₂ production (eq. 11). As reaction rate R_{15} equals the rate R_{18} , this leads to a larger HO₂ production than HO₂ loss ($R_{18}\frac{NO_y^i}{NO_y} > \frac{1}{2}R_{15}\frac{NO_y^i}{NO_y}$). Consequently, the addition of reaction (15) and (18) to the reduced HO_x reaction system V1.1 constitutes an extra HO₂ source.

Larger values of the categories "N₂O decomposition" and "lightning" to HO₂ in the upper troposphere are explained by a larger HO₂ production in V1.1 compared to V1.0. The H tagging in V1.1 considers all relevant HO₂ sources (reaction (7), (10), (11) and (28)) leading to a larger HO₂ production. Also the addition of reactions (15) and (18) (explanation see above) as well as the addition of reaction (23) which considers more reactions than in V1.0 increase the HO₂ contribution of the categories "N₂O decomposition" and "lightning".

Large changes in pattern are observed for the contributions of "biogenic emissions" and "CH₄ decomposition" to OH and HO₂ as well as for the contributions of "biomass burning" and "anthropogenic non-traffic" to OH. In V1.1, these categories mainly constitute a source of OH and HO₂ in the troposphere. The addition of reaction (24) and (25) to the reduced HO_x reaction system V1.1 presents a HO₂ source increasing OH and HO₂ contributions. Furthermore, reactions of NMHC with OH, HO₂ and NO_y (reaction 21, 22 and 23) are important throughout the whole troposphere. In contrast to V1.0, V1.1 considers all reactions of NMHC with OH, HO₂ and NO_y (see Sect. 3.2) significantly changing the pattern of "biogenic emissions", "CH₄ decomposition", "biomass burning" and "anthropogenic non-traffic".

To quantify the impact of the advanced HO_x tagging method on regional scale, Fig. 4 shows the contributions of ship emissions to OH and HO₂ in the boundary layer simulated with the high resolution model MECO(n) (see Sect. 2). The ship paths in the Atlantic, Mediterranean and Red Sea are clearly visible and lead to OH and HO₂ production along these paths. In the polluted area at the coast of Marseille the OH and HO₂ contributions are reduced, although NO_y from shipping emission is larger than in the Mediterranean Sea. However, O₃ from shipping is also larger in this region. This causes a larger HO₂ and OH loss via reaction (4) and (6) than in the Mediterranean Sea.

The tagging method V1.0 (Grewe et al. (2017) their Fig. 6) showed negative HO₂ shipping contribution along the ship paths. This was explained by reaction (14): NO destroys HO₂ and leads to negative contributions. However, in V1.1 HO₂ shipping contributions are positive. The change of sign is caused by the addition of reaction (15) and (18) to the reduced HO_x reaction system V1.1 which constitutes a net HO₂ production leading to positive HO₂ contributions (explanation see above). The comparison shows that HO₂ contributions in V1.0 were systematically and erroneously underestimated.




Figure 4. Contribution of shipping emissions to OH and HO₂ in 10⁻¹⁵ mol mol⁻¹. Monthly means of ground level values in August 2007 are shown. Simulation is performed with MECO(n).

To summarize, the contributions to OH and HO₂ concentrations show larger values in V1.1 compared to V1.0. This is explained by the re-establishment of the steady-state. For OH, no large changes are found in the categories "lightning" and "aviation". However, large changes are found for "biomass burning", "CH₄ decomposition" and "biogenic emissions". For HO₂, no large differences occur in the categories "biomass burning", "anthropogenic non-traffic", "road traffic" and "shipping".

5 In comparison, the categories "biogenic emissions" and "CH₄ decomposition" differ strongly. The differences between the contributions of V1.1 and V1.0 are traced back to the addition of certain reactions to the reduced reaction system considered in the HO_x tagging method.

4.2 Effects on long-lived species

The tagging of short-lived and long-lived species closely intertwine (see Fig. 1). Changes in the contributions to OH and HO₂ influence the contributions to the long-lived tracers O₃, NO_y, CO, NMHC and PAN. For example, Figure 5 shows the zonal mean of the contributions of the ten source categories to O₃. Grewe et al. (2017) presents the same figure for the HO_x tagging method V1.0 (their Fig. 4). For consistency, we compare our results with the results of Grewe et al. (2017) only for the year 2008.

In general, no large differences between V1.1 and V1.0 for long-lived species are found. The category "biogenic emissions" and "CH₄ decomposition" show an O₃ increase in the tropical troposphere. "Stratospheric O₃ production" slightly increases in the Southern Hemisphere. Small O₃ changes are found for the categories "lightning" and "N₂O decomposition". Regarding the remaining long-lived species (see Figures S3 – S6 in the Supplement), the contribution of "biomass burning" to CO decreases while the contributions of "biogenic emissions" to CO increases in the Southern Hemisphere. The remaining sectors stay rather unchanged. NO_y, NMHC and PAN show only minor changes. Even though, major differences in OH and HO₂ occur between V1.0 and V1.1, these do not have a large effect on the long-lived species.




Figure 5. Annual mean contributions of ten source categories to O_3 concentration in %.

5 Discussion and Conclusion

We present an extension of the HO_x tagging method described by Grewe et al. (2017). 15 new reactions producing and destroying HO_x are added to tagging mechanism. In Grewe et al. (2017), the HO_x tagging method V1.0 was restricted to the troposphere only. We further include the reactions which are essential for HO_x production and loss in the stratosphere. Moreover, we introduce an equivalent tagging method to obtain the contributions to the H radical. This step is mandatory to fully account for the main HO_2 source: the reaction of H with O_2 .

In V1.0, the steady-state assumption was not completely fulfilled resulting in an unclosed budget: the sum of the HO_x contributions and the total HO_x concentration deviated by about 70 %. To re-establish steady-state, we add more reactions to the reduced HO_x reaction system and introduce rest terms to balance the deviation of HO_x production and loss. This leads to the closure of the budget. Thus, the tagging mechanism introduced by Grewe et al. (2010) operates not only for long-lived but also for short-lived species.

The advanced HO_x tagging method V1.1 was implemented in the global chemistry climate model EMAC and in the regional model MECO(n). A 1-year simulation was performed in both model systems and compared to V1.0. For most categories, the general zonal pattern of the contributions to OH and HO_2 show minor differences. In contrast, large changes are observed in

the category "biogenic emissions" and "CH₄ decomposition" which are traced back to the addition of certain reactions to V1.1. Although the contributions of long-lived and short-lived species influence each other, no large changes are found for long-lived species.

The mechanism presented in this study (and introduced by Tsati (2014) and Grewe et al. (2017)) is the first method for tagging short-lived species. Other studies quantify the source attributions of chemical species with a significant longer lifetime. The idea of source attribution is applied to attribute CO to different emission types and regions (e.g. Granier et al., 1999; Pfister et al., 2004, 2011), to attribute NO_x concentrations to emission sources (Horowitz and Jacob, 1999) or to trace stable isotopic compositions (Gromov et al., 2010). Also for the source attribution of tropospheric O₃, several tagging approaches exist attributing tropospheric O₃ only to NO_x sources (Lelieveld and Dentener, 2000; Grewe, 2004; Grewe et al., 2012; Emmons et al., 2012), only to NMHC sources (Butler et al., 2011; Coates and Butler, 2015) or to NO_y, CO and NMHC emissions simultaneously (Grewe et al., 2017).

A common technique to quantify the impact of emissions to OH is the so called perturbation method which compares two simulations: one simulation with all emissions and one simulation with reduced emissions (e.g. Niemeier et al., 2006; Hoor et al., 2009). However, if the underlying chemical processes are non-linear (as it is the case for OH), the perturbation method largely underestimates the contribution (Grewe et al., 2012; Emmons et al., 2012; Mertens et al., 2017). Consequently, the tagging approach presented in this study delivers the actual contribution of the emission source while the perturbation method displays the impact of the emission reduction.

To conclude, the further developed HO_x tagging method can be used to identify the contribution of anthropogenic emissions on the atmospheric composition. In particular, the contribution of emission sectors on the concentrations of OH and HO₂ in the troposphere and stratosphere can be achieved. This method will be applied for re-evaluating the impact of the traffic sector on climate.

Code availability. The Modular Earth Submodel System (MESSy) is continuously further developed and applied by a consortium of institutions. The usage of MESSy and access to the source code is licensed to all affiliates of institutions, which are members of the MESSy Consortium. Institutions can become a member of the MESSy Consortium by signing the MESSy Memorandum of Understanding. More information can be found on the MESSy Consortium Web-site (<http://www.messy-interface.org>). The submodel TAGGING 1.1 will be included in MESSy version 2.54. The code being used to obtain the presented results is available upon personal request.

Appendix A: Exclusion of reactions from reduced HO_x reaction system V1.1

The annual mean reaction rates of the following three reactions are also greater than 10^{-15} mol mol⁻¹s⁻¹ and thus would usually be accounted to the reduced HO_x reaction system V1.1:



However, the tagging method can not be applied for these three reactions.

To include the OH production by the photolysis of H₂O₂ (reaction A1), we would need to tag H₂O₂. Since the production and the loss of H₂O₂ are not balanced, we can not assume a steady-state. Thus, a similar tagging approach as for HO_x and H is not valid for H₂O₂. Consequently, we exclude the reaction (A1) from the HO_x tagging method. This reaction contributes about 8 % to the total OH production in the troposphere.

Hypochlorous acid (HOCl) and hypobromous acid (HOBr) are photolysed in the stratosphere and produce OH (reaction A2 and A3), but HOCl and HOBr are not tagged. Although the steady-state assumption is globally valid, locally the production and loss of HOCl and HOBr are not balanced everywhere. In the stratosphere, for about 65 % of the model grid boxes the production deviates by more than 10 % from the loss of HOCl and HOBr. In particular, in the transition area between day and night in the polar region, the production deviates strongly from the loss. Also at night where the reactions mostly occur, the steady-state is not fulfilled everywhere. Moreover, since both species are not radicals, their lifetimes can not be assumed to be short. Hence, we can not apply the tagging method, so we have to omit the reactions (A2) and (A3) from the reduced HO_x reaction system V1.1.

20 Considering reactions (A1), (A2) and (A3) to the reduced HO_x reaction system V1.1 would lead to a significantly larger OH production in the troposphere representing about 98 % of the total OH production rate derived by MECCA. In the stratosphere, 91 % of the total OH production would be regarded. Hence, excluding these reactions from the reduced HO_x reaction system V1.1 worsen the steady-state assumption between OH production and loss. The rest term *resOH* introduced in Sect. 3.4 compensates this deviation from production and loss rate.

25 Appendix B: HO_x contributions in the stratosphere

Figs. B1 and B2 show the zonal mean of OH and HO₂ from 1 to 200 hPa. As OH concentration strongly rises with increasing height, so do the contributions to OH. The category "biomass burning" shows negative OH values in the tropopause region. In this region, also large CO values from "biomass burning" occur. CO effectively destroys OH by reaction (11) which causes this OH loss. The large OH loss in the lower stratosphere of the category "stratospheric O₃ production" is mainly caused by the destruction of OH by O₃ (reaction 6).




Figure B1. Contributions of ten source categories to OH in the stratosphere. Zonal means of the year 2010 are shown. Black line indicates the tropopause. Simulation is performed with EMAC. Note the logarithmic scale of the contour levels.

The contributions to HO_2 in the stratosphere increases with height as well. The categories "biogenic emissions", "lightning", "biomass burning", "anthropogenic non-traffic", "road traffic", "shipping" and "aviation" show a local maximum at around 5 hPa which is caused by omitting the photolysis of HOCl (see Appendix A).

For the category "lightning", HO_2 is destroyed by reaction (14) in the tropopause region. The category "N₂O decomposition" shows negative values in the lower stratosphere and a strong negative minimum at around 10 hPa which is also caused by reaction (14). The local maximum with positive HO_2 contributions indicates that in this region the HO_2 production via reaction (1) and (6) dominates the HO_2 loss via reaction (14).

Competing interests. There are no competing interests.




Figure B2. Contributions of ten source categories to HO_2 in the stratosphere. Zonal means of the year 2010 are shown. Black line indicates the tropopause. Simulation is performed with EMAC. Note the logarithmic scale of the contour levels.

Acknowledgements. This study has been carried out in the framework of the project VEU2 funded by DLR. We used the NCAR Command Language (NCL) for data analysis and to create the figures of this study. NCL is developed by UCAR/NCAR/CISL/TDD and available on-line (doi: 10.5065/D6WD3XH5). We gratefully acknowledge the computer systems provided by the Deutsches Klimarechenzentrum (DKRZ) which we used for our simulations. We thank M. Righi from DLR for helpful comments.

References

- Butler, T., Lawrence, M., Taraborrelli, D., and Lelieveld, J.: Multi-day ozone production potential of volatile organic compounds calculated with a tagging approach, *Atmospheric Environment*, 45, 4082 – 4090, doi:<https://doi.org/10.1016/j.atmosenv.2011.03.040>, <http://www.sciencedirect.com/science/article/pii/S1352231011003001>, 2011.
- 5 Clappier, A., Belis, C. A., Pernigotti, D., and Thunis, P.: Source apportionment and sensitivity analysis: two methodologies with two different purposes, *Geoscientific Model Development*, 10, 4245–4256, doi:10.5194/gmd-10-4245-2017, <https://www.geosci-model-dev.net/10/4245/2017/>, 2017.
- Coates, J. and Butler, T. M.: A comparison of chemical mechanisms using tagged ozone production potential (TOPP) analysis, *Atmospheric Chemistry and Physics*, 15, 8795–8808, doi:10.5194/acp-15-8795-2015, <https://www.atmos-chem-phys.net/15/8795/2015/>, 2015.
- 10 Crutzen, P. J. and Schmailzl, U.: Chemical budgets of the stratosphere, *Planetary and Space Science*, 31, 1009 – 1032, doi:[https://doi.org/10.1016/0032-0633\(83\)90092-2](https://doi.org/10.1016/0032-0633(83)90092-2), <http://www.sciencedirect.com/science/article/pii/0032063383900922>, 1983.
- Deckert, R., Jöckel, P., Grewe, V., Gottschaldt, K.-D., and Hoor, P.: A quasi chemistry-transport model mode for EMAC, *Geoscientific Model Development*, 4, 195–206, doi:10.5194/gmd-4-195-2011, <http://www.geosci-model-dev.net/4/195/2011/>, 2011.
- Emmons, L. K., Hess, P. G., Lamarque, J.-F., and Pfister, G. G.: Tagged ozone mechanism for MOZART-4, CAM-chem and other chemical transport models, *Geoscientific Model Development*, 5, 1531–1542, doi:10.5194/gmd-5-1531-2012, <https://www.geosci-model-dev.net/5/1531/2012/>, 2012.
- Granier, C., Müller, J., Pétron, G., and Brasseur, G.: A three-dimensional study of the global CO budget, *Chemosphere - Global Change Science*, 1, 255 – 261, doi:[https://doi.org/10.1016/S1465-9972\(99\)00007-0](https://doi.org/10.1016/S1465-9972(99)00007-0), <http://www.sciencedirect.com/science/article/pii/S1465997299000070>, 1999.
- 20 Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., van Aardenne, J., van der Werf, G. R., and van Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109, 163, doi:10.1007/s10584-011-0154-1, <https://doi.org/10.1007/s10584-011-0154-1>, 2011.
- 25 Grewe, V.: Technical Note: A diagnostic for ozone contributions of various NO_x emissions in multi-decadal chemistry-climate model simulations, *Atmospheric Chemistry and Physics*, 4, 729–736, doi:10.5194/acp-4-729-2004, <https://www.atmos-chem-phys.net/4/729/2004/>, 2004.
- Grewe, V.: A generalized tagging method, *Geoscientific Model Development*, 6, 247–253, doi:10.5194/gmd-6-247-2013, <https://www.geosci-model-dev.net/6/247/2013/>, 2013.
- 30 Grewe, V., Brunner, D., Dameris, M., Grenfell, J., Hein, R., Shindell, D., and Staehelin, J.: Origin and variability of upper tropospheric nitrogen oxides and ozone at northern mid-latitudes, *Atmospheric Environment*, 35, 3421 – 3433, doi:[http://dx.doi.org/10.1016/S1352-2310\(01\)00134-0](http://dx.doi.org/10.1016/S1352-2310(01)00134-0), <http://www.sciencedirect.com/science/article/pii/S1352231001001340>, 2001.
- Grewe, V., Tsati, E., and Hoor, P.: On the attribution of contributions of atmospheric trace gases to emissions in atmospheric model applications, *Geoscientific Model Development*, 3, 487–499, doi:10.5194/gmd-3-487-2010, <http://www.geosci-model-dev.net/3/487/2010/>, 2010.
- 35

- Grewe, V., Dahlmann, K., Matthes, S., and Steinbrecht, W.: Attributing ozone to NO_x emissions: Implications for climate mitigation measures, *Atmospheric Environment*, 59, 102 – 107, doi:<http://dx.doi.org/10.1016/j.atmosenv.2012.05.002>, <http://www.sciencedirect.com/science/article/pii/S1352231012004335>, 2012.
- Grewe, V., Tsati, E., Mertens, M., Frömming, C., and Jöckel, P.: Contribution of emissions to concentrations: The TAGGING 1.0 sub-model based on the Modular Earth Submodel System (MESSy 2.52), *Geoscientific Model Development Discussions*, 2017, 1–33, doi:10.5194/gmd-2016-298, <http://www.geosci-model-dev-discuss.net/gmd-2016-298/>, 2017.
- Gromov, S., Jöckel, P., Sander, R., and Brenninkmeijer, C. A. M.: A kinetic chemistry tagging technique and its application to modelling the stable isotopic composition of atmospheric trace gases, *Geoscientific Model Development*, 3, 337–364, doi:10.5194/gmd-3-337-2010, <https://www.geosci-model-dev.net/3/337/2010/>, 2010.
- 10 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., Mckay, W. A., Pierce, T., Sc-holes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *Journal of Geophysical Research: Atmospheres*, 100, 8873–8892, doi:10.1029/94JD02950, <http://dx.doi.org/10.1029/94JD02950>, 1995.
- Heard, D. E. and Pilling, M. J.: Measurement of OH and HO_2 in the Troposphere, *Chemical Reviews*, 103, 5163–5198, doi:10.1021/cr020522s, <http://dx.doi.org/10.1021/cr020522s>, PMID: 14664647, 2003.
- 15 Hoor, P., Borken-Kleefeld, J., Caro, D., Dessens, O., Endresen, O., Gauss, M., Grewe, V., Hauglustaine, D., Isaksen, I. S. A., Jöckel, P., Lelieveld, J., Myhre, G., Meijer, E., Olivier, D., Prather, M., Poberaj, C. S., Shine, K., Staehelin, J., Tang, Q., van Aardenne, J., van Velthoven, P., and Sausen, R.: The impact of traffic emissions on atmospheric ozone and OH: results from QUANTIFY, *Atmospheric Chemistry and Physics*, 9, 3113–3136, <http://centaur.reading.ac.uk/17096/>, 2009.
- Horowitz, L. W. and Jacob, D. J.: Global impact of fossil fuel combustion on atmospheric NO_x , *Journal of Geophysical Research: Atmo-spheres*, 104, 23 823–23 840, doi:10.1029/1999JD900205, <http://dx.doi.org/10.1029/1999JD900205>, 1999.
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geoscientific Model Development*, 3, 717–752, doi:10.5194/gmd-3-717-2010, <http://www.geosci-model-dev.net/3/717/2010/>, 2010.
- Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., 25 Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Mod-elling (ESCiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geoscientific Model Development*, 9, 1153–1200, doi:10.5194/gmd-9-1153-2016, <https://www.geosci-model-dev.net/9/1153/2016/>, 2016.
- Kerkweg, A. and Jöckel, P.: The 1-way on-line coupled atmospheric chemistry model system MECO(n) – Part 1: Description of the limited- 30 area atmospheric chemistry model COSMO/MESSy, *Geoscientific Model Development*, 5, 87–110, doi:10.5194/gmd-5-87-2012, <https://www.geosci-model-dev.net/5/87/2012/>, 2012a.
- Kerkweg, A. and Jöckel, P.: The 1-way on-line coupled atmospheric chemistry model system MECO(n) – Part 2: On-line coupling with the Multi-Model-Driver (MMD), *Geoscientific Model Development*, 5, 111–128, doi:10.5194/gmd-5-111-2012, <https://www.geosci-model-dev.net/5/111/2012/>, 2012b.
- 35 Kerkweg, A., Sander, R., Tost, H., and Jöckel, P.: Technical note: Implementation of prescribed (OFFLEM), calculated (ONLEM), and pseudo-emissions (TNUDGE) of chemical species in the Modular Earth Submodel System (MESSy), *Atmospheric Chemistry and Physics*, 6, 3603–3609, doi:10.5194/acp-6-3603-2006, <http://www.atmos-chem-phys.net/6/3603/2006/>, 2006.

- Lawrence, M. G., Jöckel, P., and von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmospheric Chemistry and Physics*, 1, 37–49, doi:10.5194/acp-1-37-2001, <https://www.atmos-chem-phys.net/1/37/2001/>, 2001.
- Lelieveld, J. and Dentener, F. J.: What controls tropospheric ozone?, *Journal of Geophysical Research: Atmospheres*, 105, 3531–3551, doi:10.1029/1999JD901011, <http://dx.doi.org/10.1029/1999JD901011>, 2000.
- 5 Liang, Q., Chipperfield, M. P., Fleming, E. L., Abraham, N. L., Braesicke, P., Burkholder, J. B., Daniel, J. S., Dhomse, S., Fraser, P. J., Hardiman, S. C., Jackman, C. H., Kinnison, D. E., Krummel, P. B., Montzka, S. A., Morgenstern, O., McCulloch, A., Mühle, J., Newman, P. A., Orkin, V. L., Pitari, G., Prinn, R. G., Rigby, M., Rozanov, E., Stenke, A., Tummon, F., Velders, G. J. M., Visionsi, D., and Weiss, R. F.: Deriving Global OH Abundance and Atmospheric Lifetimes for Long-Lived Gases: A Search for CH₃CCl₃ Alternatives, *Journal of Geophysical Research: Atmospheres*, 122, 11,914–11,933, doi:10.1002/2017JD026926, <http://dx.doi.org/10.1002/2017JD026926>, 2017.
- 10 Mertens, M., Kerkweg, A., Jöckel, P., Tost, H., and Hofmann, C.: The 1-way on-line coupled model system MECO(n) – Part 4: Chemical evaluation (based on MESSy v2.52), *Geoscientific Model Development*, 9, 3545–3567, doi:10.5194/gmd-9-3545-2016, <http://www.geosci-model-dev.net/9/3545/2016/>, 2016.
- Mertens, M., Grewe, V., Rieger, V. S., and Jöckel, P.: Revisiting the contribution of land transport and shipping emissions to tropospheric ozone, *Atmospheric Chemistry and Physics Discussions*, 2017, 1–35, doi:10.5194/acp-2017-747, <https://www.atmos-chem-phys-discuss.net/acp-2017-747/>, 2017.
- 15 Naik, V., Voulgarakis, A., Fiore, A. M., Horowitz, L. W., Lamarque, J.-F., Lin, M., Prather, M. J., Young, P. J., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R., Eyring, V., Faluvegi, G., Folberth, G. A., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., van Noije, T. P. C., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R., Shindell, D. T., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., and Zeng, G.: Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmospheric Chemistry and Physics*, 13, 5277–5298, doi:10.5194/acp-13-5277-2013, <https://www.atmos-chem-phys.net/13/5277/2013/>, 2013.
- 20 Niemeier, U., Granier, C., Kornblüeh, L., Walters, S., and Brasseur, G. P.: Global impact of road traffic on atmospheric chemical composition and on ozone climate forcing, *Journal of Geophysical Research: Atmospheres*, 111, n/a–n/a, doi:10.1029/2005JD006407, <http://dx.doi.org/10.1029/2005JD006407>, d09301, 2006.
- 25 Olson, J. R., Crawford, J. H., Chen, G., Brune, W. H., Faloona, I. C., Tan, D., Harder, H., and Martinez, M.: A reevaluation of airborne HO_x observations from NASA field campaigns, *Journal of Geophysical Research: Atmospheres*, 111, n/a–n/a, doi:10.1029/2005JD006617, <http://dx.doi.org/10.1029/2005JD006617>, d10301, 2006.
- Pfister, G., Pétron, G., Emmons, L. K., Gille, J. C., Edwards, D. P., Lamarque, J.-F., Attie, J.-L., Granier, C., and Novelli, P. C.: Evaluation of CO simulations and the analysis of the CO budget for Europe, *Journal of Geophysical Research: Atmospheres*, 109, n/a–n/a, doi:10.1029/2004JD004691, <http://dx.doi.org/10.1029/2004JD004691>, d19304, 2004.
- 30 Pfister, G. G., Avise, J., Wiedinmyer, C., Edwards, D. P., Emmons, L. K., Diskin, G. D., Podolske, J., and Wisthaler, A.: CO source contribution analysis for California during ARCTAS-CARB, *Atmospheric Chemistry and Physics*, 11, 7515–7532, doi:10.5194/acp-11-7515-2011, <https://www.atmos-chem-phys.net/11/7515/2011/>, 2011.
- 35 Prinn, R. G., Huang, J., Weiss, R. F., Cunnold, D. M., Fraser, P. J., Simmonds, P. G., McCulloch, A., Harth, C., Reimann, S., Salameh, P., O'Doherty, S., Wang, R. H. J., Porter, L. W., Miller, B. R., and Krummel, P. B.: Evidence for variability of atmospheric hydroxyl radicals over the past quarter century, *Geophysical Research Letters*, 32, n/a–n/a, doi:10.1029/2004GL022228, <http://dx.doi.org/10.1029/2004GL022228>, 107809, 2005.

- Ren, X., Harder, H., Martinez, M., Leshner, R. L., Oliger, A., Simpas, J. B., Brune, W. H., Schwab, J. J., Demerjian, K. L., He, Y., Zhou, X., and Gao, H.: OH and HO₂ Chemistry in the urban atmosphere of New York City, *Atmospheric Environment*, 37, 3639 – 3651, doi:[http://dx.doi.org/10.1016/S1352-2310\(03\)00459-X](http://dx.doi.org/10.1016/S1352-2310(03)00459-X), <http://www.sciencedirect.com/science/article/pii/S135223100300459X>, 2003.
- 5 Righi, M., Eyring, V., Gottschaldt, K.-D., Klinger, C., Frank, F., Jöckel, P., and Cionni, I.: Quantitative evaluation of ozone and selected climate parameters in a set of EMAC simulations, *Geoscientific Model Development*, 8, 733–768, doi:10.5194/gmd-8-733-2015, <https://www.geosci-model-dev.net/8/733/2015/>, 2015.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornbluh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of Simulated Climate to Horizontal and Vertical Resolution in the ECHAM5 Atmosphere Model, *Journal of Climate*, 19, 3771–3791, doi:10.1175/JCLI3824.1, <http://dx.doi.org/10.1175/JCLI3824.1>, 2006.
- 10 Sander, R., Baumgaertner, A., Gromov, S., Harder, H., Jöckel, P., Kerkweg, A., Kubistin, D., Regelin, E., Riede, H., Sandu, A., Taraborrelli, D., Tost, H., and Xie, Z.-Q.: The atmospheric chemistry box model CAABA/MECCA-3.0, *Geoscientific Model Development*, 4, 373–380, doi:10.5194/gmd-4-373-2011, <https://www.geosci-model-dev.net/4/373/2011/>, 2011.
- Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, 15 M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F., Lawrence, M. G., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *Journal of Geophysical Research: Atmospheres*, 111, n/a–n/a, doi:10.1029/2005JD006338, <http://dx.doi.org/10.1029/2005JD006338>, d08301, 2006.
- Stone, D., Whalley, L. K., and Heard, D. E.: Tropospheric OH and HO₂ radicals: field measurements and model comparisons, *Chem. Soc. Rev.*, 41, 6348–6404, doi:10.1039/C2CS35140D, <http://dx.doi.org/10.1039/C2CS35140D>, 2012.
- 20 Tsati, E. E.: Investigation of the impacts of emissions on the trace gas budgets in the troposphere by using global climate chemistry model simulations, Ph.D. thesis, <https://edoc.ub.uni-muenchen.de/17524/>, 2014.
- Voulgarakis, A., Naik, V., Lamarque, J.-F., Shindell, D. T., Young, P. J., Prather, M. J., Wild, O., Field, R. D., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Folberth, G. A., Horowitz, L. W., Josse, B., 25 MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Stevenson, D. S., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Analysis of present day and future OH and methane lifetime in the ACCMIP simulations, *Atmospheric Chemistry and Physics*, 13, 2563–2587, doi:10.5194/acp-13-2563-2013, <https://www.atmos-chem-phys.net/13/2563/2013/>, 2013.
- Yienger, J. J. and Levy, H.: Empirical model of global soil-biogenic NO_x emissions, *Journal of Geophysical Research: Atmospheres*, 100, 11 447–11 464, doi:10.1029/95JD00370, <http://dx.doi.org/10.1029/95JD00370>, 1995.