

Dear Brian,

Once again thank you for the time spent in reviewing this paper and your useful suggestions. Our responses are below.

The authors have generally done a good job in responding to reviewer comments, and I only have a couple of minor suggestions remaining.

The revisions to address the issue of the representation of biospherical effects of land use are adequate, although I note that the argument was not that the FAIR model should include latitude dependence, but rather that it is unclear whether it is a good idea to represent only part of the biospherical effects of land use. However the revised text acknowledges the other factors at play, and that the sign of the effect on forcing remains uncertain. And I would agree that a dependence on cumulative emissions from land use remains a reasonable simple relationship to rely on.

Thank you for the understanding of our argument, which we trust is satisfactory.

Land use forcing remains an uncertainty that is difficult to relate to projections where limited information is available. As stated in the response to the other reviewer, we have now allowed the option of specifying a land-use forcing directly. The scaling with cumulative land-use CO₂ emissions remains available and is the default option.

Following your original suggestion in your first review, we intend to allow a gridded land-use transition dataset to be convoluted with the contributions to surface albedo change with deforestation from Jones et al. (2015) in a future model version, once this data becomes available.

In section 3 a particular approach to constraining model parameters using the observed record of global average temperature is described. However, it needs to be made clear that while this is one approach, there is a sub-field of work on this topic, aiming to estimate model parameters (including ECS) from simple models and historical data, and there are quite a number of methodological issues to contend with, including the influence of the length of the historical record, the means of distinguishing the forced response from natural variability, the method of specifying and updating prior distributions, the accounting for individual components of historical forcing, etc.

You are quite correct. We update the paragraph at the end of section 3.1 as follows:

“It should be stressed that there are several issues to consider when attempting to derive plausible parameter sets from observational data. These include the type of observational constraints to employ (Meinshausen et al., 2009), the length of the historical record (e.g.

Otto et al. (2013)), the separation of forced response from natural variability (Haustein et al., 2017), and assumptions surrounding prior distributions (Frame et al., 2005).”

p. 18 line 12 the revised sentence is somewhat awkwardly stated, or perhaps there is a typo. Maybe clearer to say ”The RCP8.5 temperature change is lower despite higher 21st century ERF..”

Thank you for this suggestion. We have clarified this sentence.

References

- Frame, D. J., Booth, B. B. B., Kettleborough, J. A., Stainforth, D. A., Gregory, J. M., Collins, M., and Allen, M. R. (2005). Constraining climate forecasts: The role of prior assumptions. *Geophys. Res. Lett.*, 32(9). L09702.
- Haustein, K., Allen, M., Forster, P., Otto, F., Mitchell, D., Matthews, H., and Frame, D. (2017). A real-time global warming index. *Scientific Reports*, 7(1):15417.
- Jones, A. D., Calvin, K. V., Collins, W. D., and Edmonds, J. (2015). Accounting for radiative forcing from albedo change in future global land-use scenarios. *Climatic Change*, 131(4):691–703.
- Meinshausen, M., Meinshausen, N., Hare, W., Raper, S. C., Frieler, K., Knutti, R., Frame, D. J., and Allen, M. R. (2009). Greenhouse-gas emission targets for limiting global warming to 2 C. *Nature*, 458(7242):1158–1162.
- Otto, A., Otto, F. E. L., Boucher, O., Church, J., Hegerl, G., Forster, P. M., Gillett, N. P., Gregory, J., Johnson, G. C., Knutti, R., Lewis, N., Lohmann, U., Marotzke, J., Myhre, G., Shindell, D., Stevens, B., and Allen, M. R. (2013). Energy budget constraints on climate response. *Nat. Geosci.*, 6:415–416.

Dear William,

Once again thank you for your time spent in reviewing this manuscript and the useful comments you provided. Original comments are given in bold, which are responded to point-by-point in regular font.

Implementing the aerosol changes you suggest have changed model results slightly, so to reflect this new default behaviour we have incremented the model version to 1.3.

Second review of Smith et al.

The authors have given detailed and thoughtful responses to my first review. I just have two main outstanding points.

As I stated before it is dangerous to hard-wire conversions from land CO₂ emissions to albedo forcing, and from aircraft NO_x to contrails. While these (by design) give the correct response to the RCP8.5 scenario, this model is likely to be taken up and (the authors should hope) used by many diverse groups for different purposes and different scenarios. It is very likely that this hard-wiring will persist out in the community even when the authors have updated their own version. As an example the RCP6.0 scenario has strong negative land CO₂ emissions which presumably will give incorrect land albedo forcing. The SSP2.6 and 1.9 also have negative land emissions. If the input data is land use area then at least any future user is forced to think about where this data is coming from.

Thank you for this comment.

We use this model for carbon budget calculations for the 600+ scenarios that are used in the IPCC Special Report on 1.5 Degrees. For these scenarios all that are provided are annual global mean emissions of the 39 species mentioned in table 1 in the manuscript. We therefore do not have any better data on which to calculate forcing due to land use change or aviation contrails in these scenarios. However, we have expanded the options available for both forcing categories to allow the user to select a more appropriate option.

Land use

We have now added an option to allow the user to specify their own time series of land use ERF externally, thereby not calculating it through a scaling with CO₂ land use emissions. The scaling with cumulative land-use CO₂ emissions remains the default option.

A further option we will develop is to use an external dataset of annual land-use transitions to and from forest/shrubland from the LUH datasets, and multiply these by the effect on albedo change in each grid cell calculated from the GCAM model in Jones et al. (2015),

figure 1. The Jones et al. (2015) data is currently not available to us, but this feature will be implemented when it is.

Contrails

We understand your original concern and therefore provide two alternative methods.

As for land use, a user can specify a time series of contrail forcing directly, bypassing any internally-calculated routines.

Alternatively, we provide a relationship based on a user-supplied time series of kerosene jet-fuel consumption. For RCP scenarios this data does not appear to be available so cannot be used for our demonstration of future projections, but a user could implement this if they did have the data.

To respond to the author's responses on methane lifetime:

- 1. If using the Holmes et al. parameters gives a slightly different result to MAGICC (see also their Fig 7) that doesn't seem an issue as there is no reason to assume MAGICC is the "truth". I agree with RCP8.5 there are two opposing effects – methane increases the lifetime (+29%), climate decreases the lifetime (-17%), but that might not be the same in all scenarios e.g. where climate increases but methane doesn't or vice-versa.**
- 2. The relationship has been shown (in theory, and in models) to hold for methane doubling in many papers, particularly by Michael Prather and Chris Holmes.**
- 3. Applying a simple methane feedback won't work so it is not surprising it doesn't reproduce the historical or future without extreme natural emissions. Over the historical the methane self-feedback effect will be offset by increases in NOx emissions and changes in climate, Naik et al. 2013 ACP find these effects more or less cancel. In the future, as above Holmes et al. find climate change offsets more than half of the methane self-feedback in RCP8.5.**

Not using a state-varying methane lifetime remains a frustration and is an avenue for future development.

Specific comments:

Page 2, line 13: Do you really mean different responses to "forcings" here (i.e. different efficacies) or do you mean different responses to emissions?

We meant in response to emissions, and have updated this sentence.

Page 2, line 15: Gasser et al. 2017 “Accounting for the climate–carbon feedback in emission metrics” Earth Syst. Dynam. have quantified the effect of non-CO₂ agents on the carbon cycle.

This reference has been added, and a recent paper on the effects of non-CO₂ forcing on carbon budgets (Tokarska et al., 2018) has also been cited.

Page 2, line 23: I should have been clearer in my comment on this the first time round, the Joos parameter certainly do include feedbacks for both biospheric carbon uptake and temperature (see AR5 section 8.7.1.4 and above Gasser paper).

Thank you for this clarification. We have tried to simplify further to indicate that the issue is that the Joos et al. (2013) relationship, which is designed for a particular experiment (100 GtC pulse against a background of 389ppm and present-day conditions) does not represent C4MIP carbon cycles well for a more general background state. This was the carbon-cycle relationship introduced in Millar et al. (2017). As Joos et al. (2013) models the multi-model carbon cycle response, it does include carbon-cycle-climate feedbacks.

“A simple emulation of the carbon cycle of full- and intermediate-complexity earth system models was developed by Joos et al. (2013) and used in the IPCC Fifth Assessment Report (AR5) for the purposes of calculating global warming potentials. The model was developed for a 100 GtC pulse against a background CO₂ concentration of 389 ppm. Millar et al. (2017) showed that this model does not sufficiently capture the time-evolving dependency of carbon sinks against different background conditions.”

Page 3, line 16: It is not obvious whether the methane is already accounted for in the carbon budgets (Boucher et al. 2009). See my further comment on page 6.

We agree it is not obvious. From Gillenwater (2008) and Boucher et al. (2009) it appears that our treatment of including oxidised methane as CO₂ is correct, and MAGICC also includes a proportion of methane oxidised to CO₂ (eq. A1 of (Meinshausen et al., 2011)). The assumption that methane oxidised to CO₂ is not included in national emissions inventories has been moved to this section.

Gillenwater (2008) suggests that this treatment adds between 0.5 and 0.7% to global emissions on a GWP basis. So, neglecting this effect (or erroneously including it) does not introduce large errors.

The fraction of fossil methane oxidised to CO₂ is a user-specified input which defaults to 0.61 following Boucher. The time series of total methane emissions attributable to fossil sources is scenario-dependent, and RCP-calculated values are included in FAIR for the user to import. By setting either of these numbers to zero, the user can switch off methane oxidation to CO₂. This has been highlighted in section 2.1.3.

We have tested this in RCP4.5, and found that the difference in CO₂ concentrations with the methane conversion switched off is 0.3 ppm in 2015 and about 1.2 ppm in 2500.

Page 3, line 28 & page 4, line 23: Either use “molar mixing ratio” instead of “concentration”, or specifically state that the units are mol mol⁻¹. While it is generally fine to use “concentration” colloquially, here specific equations are being presented and the reader needs to know whether the results are in kg m⁻³ or mol mol⁻¹.

Thank you for this suggestion which has been implemented.

Page 5, lines 19-25. If you are going to mention the perturbation lifetime, I think you should explicitly say why it is not appropriate (changes in emissions of NO_x & VOCs, changes in climate). Naik et al. 2013 ACP find the methane lifetime is essentially unchanged over the historical period due to compensating terms (models range from -15% to +12%).

Thank you for this suggestion. The following commentary has been added:

“As emissions of OH-affecting species (NO_x, NMVOCs, CO) and temperature have varied substantially over the historical period, the perturbation lifetime definition is not appropriate as the background state is not constant, and furthermore there is no robust evidence of a change in methane lifetime between pre-industrial and present-day from the ACCMIP inter-model comparison (Naik et al., 2013).”

Page 6, lines 1-2. The exponential decay of methane has no affect at all on the double counting as this decay is to CO₂. Even if the decay was instantaneous each mole of CH₄ would end up as a mole of CO₂.

We have deleted this sentence to prevent confusion, and have mentioned the fact that methane oxidation fraction and proportion of methane from fossil origin are user-specifiable.

Page 8, line 29-30: The semi-direct effect applies (almost) entirely to BC in IPCC. No significant semi-direct effect has been quantified for the scattering aerosols. Therefore your BC coefficient should be scaled down (and certainly not up by 1.37) until you get a forcing of -0.45, and all other coefficients unchanged.

Thank you for this suggestion. As with many numbers supplied in this paper, the aerosol radiative efficiencies for each species are inputs to the model that can be changed by the user, and we report the defaults. Therefore we have updated these default coefficients.

The new relationship uses the 1750-2010 direct radiative forcings from Myhre et al. (2013) for non-BC aerosols and scales the direct forcing from BC down by 0.1 W m⁻² to account for the semi-direct effect to match the best estimate ERF_{ari}. Table 5 and the aerosol section in the text have been updated.

Because this change in default aerosol treatment also changes the results, the headline results reported in the paper have also changed slightly. To highlight the fact that this is a different (default) model setup (although scientifically the same model since the last paper revision), we decided it was cleaner to increment the model version to 1.3.

Section 2.2.9: It is very dangerous to have a conversion from CO₂ emissions to land-use hard-wired into FAIR – for instance in RCP6.0 land use CO₂ emissions are strongly negative (I don't know why). It would be much safer to have land use area as an input. RCP data can then be pre-processed to create a land use area input. As the authors note the latitudinal dependence is very large. The new SSPs provide land use area change for different cover types.

As noted earlier in our response, we have now provided an alternative method to directly specify land use forcing in FAIR. We have requested the gridded deforestation-to-forcing dataset from Andy Jones (but we do not yet have this), which would allow gridded land-use datasets to be imported directly.

The SSPs do provide land use area change, but they do not appear to be gridded (at least from the SSP database at <https://tntcat.iiasa.ac.at/SspDb>). Land use (presumably MAGICC-derived) forcing data does not appear to be available from these SSP scenarios either, but even if it was, this would only be tuning responses to MAGICC.

References

- Boucher, O., Friedlingstein, P., Collins, B., and Shine, K. P. (2009). The indirect global warming potential and global temperature change potential due to methane oxidation. *Environ. Res. Lett.*, 4(4):044007.
- Gillenwater, M. (2008). Forgotten carbon: indirect CO₂ in greenhouse gas emission inventories. *Environ. Sci. Policy*, 11(3):195–203.
- Jones, A. D., Calvin, K. V., Collins, W. D., and Edmonds, J. (2015). Accounting for radiative forcing from albedo change in future global land-use scenarios. *Climatic Change*, 131(4):691–703.
- Joos, F., Roth, R., Fuglestedt, J., Peters, G., Enting, I., von Bloh, W., Brovkin, V., Burke, E., Eby, M., Edwards, N., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J. (2013). Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis. *Atmos. Chem. Phys.*, 13(5):2793–2825.

- Lee, D. S., Fahey, D. W., Forster, P. M., Newton, P. J., Wit, R. C., Lim, L. L., Owen, B., and Sausen, R. (2009). Aviation and global climate change in the 21st century. *Atmos. Environ.*, 43(22):3520–3537.
- Meinshausen, M., Raper, S., and Wigley, T. (2011). Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration. *Atmos. Chem. Phys.*, 11:1417–1456.
- Millar, R. J., Nicholls, Z. R., Friedlingstein, P., and Allen, M. R. (2017). A modified impulse-response representation of the global near-surface air temperature and atmospheric concentration response to carbon dioxide emissions. *Atmos. Chem. Phys.*, 2017:7213–7228.
- Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C. (2013). Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations. *Atmos. Chem. Phys.*, 13(4):1853–1877.
- Naik, V., Voulgarakis, A., Fiore, A. M., Horowitz, L. W., Lamarque, J.-F., Lin, M., Prather, M. J., Young, P. J., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R., Eyring, V., Faluvegi, G., Folberth, G. A., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., van Noije, T. P. C., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R., Shindell, D. T., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., and Zeng, G. (2013). Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP). *Atmos. Chem. Phys.*, 13(10):5277–5298.
- Tokarska, K. B., Gillett, N. P., Arora, V. K., Lee, W. G., and Zickfeld, K. (2018). The influence of non-co₂ forcings on cumulative carbon emissions budgets. *Environmental Research Letters*, 13(3):034039.

FAIR v1.32: A simple emissions-based impulse response and carbon cycle model

Christopher J. Smith¹, Piers M. Forster¹, Myles Allen², Nicholas Leach², Richard J. Millar^{3,4}, Giovanni A. Passerello¹, and Leighton A. Regayre¹

¹School of Earth and Environment, University of Leeds, Leeds, UK

²Atmospheric Physics Department, University of Oxford, Oxford, UK

³College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK

⁴Environmental Change Institute, University of Oxford, Oxford, UK

Correspondence to: C.J.Smith (C.J.Smith1@leeds.ac.uk)

Abstract. Simple climate models can be valuable if they are able to replicate aspects of complex fully coupled earth system models. Larger ensembles can be produced, enabling a probabilistic view of future climate change. A simple emissions-based climate model, FAIR, is presented which calculates atmospheric concentrations of greenhouse gases and effective radiative forcing (ERF) from greenhouse gases, aerosols, ozone and other agents. Model runs are constrained to observed temperature change from 1880 to 2016 and produce a range of future projections under the Representative Concentration Pathway (RCP) scenarios. The constrained estimates of equilibrium climate sensitivity (ECS), transient climate response (TCR) and transient climate response to cumulative CO₂ emissions (TCRE) are 2.86 (2.01 to 4.22) ~~2.93 (2.04 to 4.32)~~ K, 1.53 (1.05 to 2.41) ~~1.59 (1.07 to 2.50)~~ K and 1.40 (0.96 to 2.23) ~~1.44 (0.97 to 2.31)~~ K (1000 GtC)⁻¹ (median and 5–95% credible intervals). These are in good agreement, with tighter uncertainty bounds, than the AR5 likely range, noting that AR5 estimates were derived from a combination of climate models, observations and expert judgement. The ranges of future projections of temperature and ranges of estimates of ECS, TCR and TCRC are somewhat sensitive to the prior distributions of ECS/TCR parameters, but less sensitive to the ERF from a doubling of CO₂ or the observational temperature dataset used to constrain the ensemble. Taking these sensitivities into account, there is no evidence to suggest that the median and credible range of observationally constrained TCR or ECS differ from climate model-derived estimates. The range of temperature projections under RCP8.5 for 2081–2100 in the constrained FAIR model ensemble is lower than the emissions-based estimate reported in AR5 by half a degree, owing to differences in forcing assumptions and ECS/TCR distributions.

Copyright statement. ©Author(s) 2018. This work is distributed under the Creative Commons Attribution 4.0 License.

1 Introduction

Most multi-model studies, such as the Coupled Model Intercomparison Project (CMIP) which produces headline climate projections for the Intergovernmental Panel on Climate Change (IPCC) assessment reports, compare atmosphere-ocean general

circulation models that are run with prescribed concentrations of greenhouse gases. Greenhouse gas and aerosol emissions time series are provided by integrated assessment modelling groups based on socio-economic narratives (Moss et al., 2010; Meinshausen et al., 2011b), which are then converted to atmospheric concentrations by simple climate-carbon-cycle models such as MAGICC6 (Meinshausen et al., 2011a). Earth system models can be run in emissions mode, where emissions of carbon dioxide are used as a starting point and the atmospheric CO₂ concentrations are calculated interactively in the model, with atmospheric concentration changes being the residual of emissions minus absorption by land and ocean sinks. While many models include the functionality to be run in CO₂ emissions-driven mode, these integrations were not the main focus of CMIP5 (the fifth phase of CMIP; Taylor et al. (2012)).

Earth system models in CMIP5 all show a positive carbon-cycle feedback, meaning that as surface temperature increases, land and ocean carbon sinks become less effective at absorbing CO₂ and a larger proportion of any further emitted carbon will remain in the atmosphere (Friedlingstein, 2015). The various feedback strengths are nevertheless model dependent (Friedlingstein et al., 2006). While CO₂ is the most important climate forcer, individual models may also differ in their responses to non-CO₂ emissions forcings. These emissions forcings can also introduce uncertainty that is not captured in concentration-driven or CO₂-only driven model experiments (Matthews and Zickfeld, 2012; Tachiiri et al., 2015; Gasser et al., 2017). As non-CO₂ forcing impacts temperature which affects the efficiency of carbon sinks, non-CO₂ forcing agents themselves influence the carbon cycle (MacDougall et al., 2015; Tokarska et al., 2018).

Simple models can be used to emulate radiative forcing and temperature responses to emissions and atmospheric concentrations, and can be tuned to replicate the behaviour of individual climate and earth system models (Meinshausen et al., 2011a; Good et al., 2011, 2013; Geoffroy et al., 2013). A simple emulation of the carbon cycle of full- and intermediate-complexity earth system models was developed by Joos et al. (2013) and used in the IPCC Fifth Assessment Report (AR5) for the purposes of calculating global warming potentials. The model was developed for a 100 GtC pulse against a background CO₂ concentration of 389 ppm. In the IPCC Fifth Assessment Report (AR5), a simple carbon-cycle model was suggested, calibrated to present-day conditions (Supplementary Material to Myhre et al., 2013b). This model used fixed time constants for the decay of atmospheric CO₂ where time constants were taken from a multi-model intercomparison of full- and intermediate-complexity earth system models (Joos et al., 2013) with no feedbacks assumed for biospheric carbon uptake or temperature. It was introduced for the purposes of calculating global warming potentials, built on the impulse response model of Boucher and Reddy (2008). Millar et al. (2017) showed that this AR5 model does not sufficiently capture the time-evolving dependency of carbon sinks against different background conditions with cumulative carbon emissions and temperatures. They introduced the Finite Amplitude Impulse Response (FAIR) model (version 1.0) that tracks the time-integrated airborne fraction of carbon and uses this to determine the efficiency of carbon sinks, in turn calculating atmospheric CO₂ concentrations, radiative forcing, and temperature change.

FAIR v1.0 is well-calibrated to the temperature and carbon cycle response of earth system models. FAIR v1.3v1.2 is extended to calculate non-CO₂ greenhouse gas concentrations from emissions, aerosol forcing from aerosol precursor emissions, tropospheric and stratospheric ozone forcing from the emissions of precursors, and forcings from black carbon on snow, stratospheric methane oxidation to water vapour, contrails and land use change. Forcings from volcanic eruptions and solar irradiance

fluctuations are supplied externally. These forcings are then converted to a temperature change, taking into account the different thermal responses of the ocean mixed layer and deep ocean.

The model philosophy in FAIR is to represent these processes as simply as possible, and to be able to emulate the [historical](#) ERF time series in AR5 given input emissions. FAIR is written in Python and open source. The extension to non-CO₂ emissions makes FAIR [v1.3v1.2](#) applicable for assessing scenarios with a broad range of emissions pathways.

This paper introduces the FAIR model in section 2, including the key changes from versions 1.0 to [1.3v1.2](#). Section 3 then discusses the generation of a large ensemble of input parameters to the FAIR model which is run and results described in section 4. A sensitivity analysis to some of the key inputs to the large ensemble is given in section 5. Section 6 provides a summary.

2 Development of FAIR [v1.3v1.2](#) and differences to v1.0

FAIR [v1.3v1.2](#) takes emissions of greenhouse gases and short-lived climate forcers as its main input. This is an array of size (number of years \times 40) (see table 1), and is based on the order provided in the RCP emissions datasets (Meinshausen et al., 2011b). Additional [optionsinputs](#) that can be specified by the user include the treatment of aviation contrail and land use forcing, ~~there are the fraction of total nitrogen oxides (NO_x) emitted by the aviation sector,~~ fraction of total methane attributable to fossil fuels, natural emissions of CH₄ and N₂O, and natural forcing from solar variability and volcanoes. The atmospheric concentrations of greenhouse gases are calculated from new emissions minus the decay of the current atmospheric burden, which is determined by the atmospheric lifetime of each gas, and output as a (years \times 31) array (table 2). For CO₂, atmospheric concentrations are calculated from a simple representation of the carbon cycle which includes temperature and saturation dependency of land and ocean sinks. ~~The calculated CO₂ concentrations at each timestep also and~~ includes a proportion of methane oxidised to CO₂. ~~This is on the assumption of a mole of oxidised methane from fossil sources is not also counted as a mole of CO₂ when reported in national emissions inventories (Gillenwater, 2008; Boucher et al., 2009).~~

The effective radiative forcing (ERF) from 13 different forcing agents (table 3) is determined from the concentrations of each greenhouse gas, plus emissions of short-lived climate forcers and natural forcing and is output from the model as a (years \times 13) array. From the ERF, temperature change is calculated. The change in temperature feeds back into the carbon cycle, which impacts the atmospheric lifetime of carbon dioxide. A flow diagram outlining the key processes is provided in fig. 1. It is also possible to run FAIR using only CO₂ emissions as inputs [or purely in forcing-only mode](#), where a time series of non-CO₂ [or total](#) forcing can be optionally specified rather than calculated from emissions.

2.1 Greenhouse gases: emissions to concentrations

2.1.1 Carbon dioxide and carbon cycle

The carbon cycle component in FAIR-~~v1.0~~ is described in detail by Millar et al. (2017) and an overview is provided here. The FAIR model uses anthropogenic fossil and land-use CO₂ emissions as input and partitions them into four boxes R_i (with

partition fraction a_i and $\sum_0^3 a_i = 1$) representing the differing time scales of carbon uptake by geological processes (τ_0), the deep ocean (τ_1), biosphere (τ_2) and ocean mixed layer (τ_3). The atmospheric **molar mixing ratios** of CO₂ and its relationship to each box is

$$C_{\text{CO}_2} = C_{\text{CO}_2,\text{pi}} + \sum_{i=0}^3 \frac{R_i}{M_a} \frac{w_{\text{CO}_2}}{w_a} \quad (1)$$

with $C_{\text{CO}_2,\text{pi}}$ equal to 278 ppm and pi representing pre-industrial. $M_a = 5.1352 \times 10^{18}$ kg is the dry mass of the atmosphere and w_{CO_2} and w_a are the molecular weights of CO₂ and dry air. R_i is in kilograms. The governing equations for the four boxes are

$$15 \quad \frac{dR_i}{dt} = a_i E_{\text{CO}_2} - \frac{R_i}{\alpha \tau_i}; \quad i = 0, \dots, 3 \quad (2)$$

with E_{CO_2} the emissions of CO₂.

The four time constants τ_i are scaled by a factor α depending on the 100-year integrated impulse response function (iIRF₁₀₀), which represents the 100-year average airborne fraction of a pulse of CO₂ (Joos et al., 2013). α is found by equating two different expressions for iIRF₁₀₀

$$15 \quad \sum_{i=0}^3 \alpha a_i \tau_i \left[1 - \exp\left(\frac{-100}{\alpha \tau_i}\right) \right] = r_0 + r_C C_{\text{acc}} + r_T T \quad (3)$$

and finding the unique root α (Millar et al., 2017). The right hand side of eq. (3) proposed by Millar et al. (2017) is a simplified expression for iIRF₁₀₀ that depends on the total accumulated carbon in land and ocean sinks $C_{\text{acc}} = (\sum_t E_{\text{CO}_2,t}) - (C_{\text{CO}_2} - C_{\text{CO}_2,\text{pi}})$ and temperature change T since the pre-industrial era that simulates the behaviour of earth system models well. This increase in iIRF₁₀₀ and scaling of the time constants by α accounts for the land and ocean carbon sinks changing absorption efficacy as more carbon is added to them (r_C parameter). In earth system models it is also observed that the efficiency of carbon sinks decreases with increasing temperature (r_T parameter; Fung et al. (2005); Friedlingstein et al. (2006)). Following Millar et al. (2017) we use $r_C = 0.019$ yr GtC⁻¹ and $r_T = 4.165$ yr K⁻¹, but in contrast to Millar et al. (2017) a pre-industrial $r_0 = 35$ years is used rather than their 32.4 years. This facilitates better agreement with present-day CO₂ atmospheric concentrations when spun up from 1765 with historical CO₂ and non-CO₂ emissions. This parameter combination is consistent with a present-day iIRF₁₀₀ diagnosed from more complex carbon-cycle models (Joos et al., 2013) with a fixed background CO₂ concentration of 389 ppm.

2.1.2 Other greenhouse gases

A one-box decay model is assumed for other greenhouse gases where the sink is an exponential decay of the existing gas concentration anomaly. New emissions are converted to **the equivalent increase in molar mixing ratios** δC_t in year t by

$$\delta C_t = \frac{E_t}{M_a} \frac{w_a}{w_f} \delta_t \quad (4)$$

where E_t is the emissions of gas in year t , $M_a = 5.1352 \times 10^{18}$ kg is the mass of the atmosphere and w_f is the molecular mass of the greenhouse gas ($\delta_t = 1$ for annual emissions data). The model updates the atmospheric [molar mixing ratios](#) [concentrations](#)

5 at year t based on new emissions and the natural sink by

$$C_t = C_{t-1} + \frac{1}{2}(\delta C_{t-1} + \delta C_t) - C_{t-1}(1 - \exp(-1/\tau)) \quad (5)$$

where τ is the atmospheric lifetime of each gas (table 2).

For CH_4 and N_2O , time-varying natural emissions are included in E_t (fig. 2) in order to match the observed atmospheric concentrations of these gases in Meinshausen et al. (2011b), including the 1765 concentrations when the 1765 natural emissions
10 are run to steady state. Beyond 2005, natural emissions of CH_4 and N_2O are fixed at $191 \text{ Mt CH}_4 \text{ yr}^{-1}$ and $8.99 \text{ Mt N}_2\text{-eq yr}^{-1}$ which are close to the best-estimate present-day emissions of $202 \text{ Mt CH}_4 \text{ yr}^{-1}$ and $9.1 \text{ Mt N}_2\text{-eq yr}^{-1}$ (Prather et al., 2012). We prefer to use varying natural emissions with a fixed atmospheric lifetime of CH_4 and N_2O , firstly because this provides a good match to observed and projected concentrations and secondly because this is consistent with the simple model philosophy. Other methods of calculating concentrations of these gases are possible, for example using a fixed natural background emission
15 and relating any differences between observed and calculated historical concentrations as an error term (either in the natural or anthropogenic time series or missing processes), or by adjusting the atmospheric lifetime of each gas over the historical period in order to match the observed concentrations at each time step.

Natural emissions of CO_2 are not included as the carbon cycle model is more complex than the single box used for other gases and it is assumed that natural sources and natural sinks are in balance. For other greenhouse gases, natural emissions
20 are assumed to be zero except for CF_4 , CH_3Br and CH_3Cl . Pre-industrial concentrations of these three minor compounds are estimated by running the 1765 emissions from Meinshausen et al. (2011b) to steady state using the lifetimes in table 2. Unlike CH_4 and N_2O , natural emissions of CF_4 , CH_3Br and CH_3Cl are included in the anthropogenic emissions data. In total, 31 greenhouse gas species are used (table 2). Other than CO_2 , CH_4 and N_2O , the remaining gases can be subdivided into those covered by the Kyoto Protocol (HFCs, PFCs, SF_6), and the ozone depleting substances (ODSs) covered by the Montreal
25 Protocol (CFCs, HCFCs and other chlorinated and brominated compounds).

The best estimate of τ for each gas except methane is used from AR5 (Myhre et al., 2013b, table 8.A.1), consistent with using AR5 estimates for parameters where possible. We find that using a constant methane lifetime of 9.3 years results in reasonable levels of historical natural emissions and also agrees well with the [MAGICC6-projected RCP concentration](#) scenarios in the future. The [global](#) methane lifetime of 9.3 years used is significantly lower than the [perturbation lifetime](#) of 12.4 years in AR5.
30 This latter figure includes the feedback of methane emissions on its own lifetime [due to the depletion of the OH radical which is the main tropospheric sink for methane](#) (a factor of 1.34; Holmes et al. (2013), also used in AR5) and is used for perturbation calculations against a constant background concentration. [As emissions of OH-affecting species \(NOx, NMVOCs, CO\) and temperature have varied substantially over the historical period, the background state is not constant, so the perturbation lifetime is not appropriate](#) ~~Dividing the perturbation lifetime by 1.34 gives the atmospheric burden lifetime of 9.3 years.~~

2.1.3 Methane oxidation to CO₂

The oxidation of CH₄ produces additional CO₂ if it is of fossil origin, which is accounted for in the model. Methane is assumed to be from fossil sources if it arises from the transport, energy or industry sectors. ~~We use the breakdown of emissions by sector from the RCP Database (2009).~~ A best estimate of 61% of the methane lost through reaction with the hydroxyl radical in the troposphere (the dominant loss pathway) is converted to CO₂ (Boucher et al., 2009). This is treated as additional emissions of CO₂:

$$E_{\text{CH}_4 \rightarrow \text{CO}_2} = 0.61 f_{\text{CH}_4 \text{ fos}} (C_{\text{CH}_4} - C_{\text{CH}_4, \text{ pi}}) (1 - \exp(-1/\tau_{\text{CH}_4})) \quad (6)$$

where $f_{\text{CH}_4 \text{ fos}}$ is the fraction of anthropogenic methane attributable to fossil sources and τ_{CH_4} is 9.3 years. ~~Both the fraction of methane converted (61%) and the timeseries of the fraction of methane that is of fossil origin $f_{\text{CH}_4 \text{ fos}}$ are user-specifiable, and RCP-derived values available from the RCP Database (2009) can be imported. The user can therefore switch off methane oxidation by setting either of these values to zero. As atmospheric methane concentrations are reduced by exponential decay in eq. (5), this prevents against (approximately) double-counting an emitted fossil-fuel CH₄ molecule as both CH₄ and CO₂ after it has been oxidised.~~

Oxidation of CO and non-methane volatile organic compounds (NMVOCs) to CO₂ is not included as to not double count the carbon that is included in national CO₂ emissions inventories (Daniel and Solomon, 1998; Gillenwater, 2008).

2.2 Effective radiative forcing

The ERF from 13 different forcing agent groups are considered: CO₂, CH₄, N₂O, other greenhouse gases, tropospheric O₃, stratospheric O₃, stratospheric water vapour, contrails, aerosols, black carbon on snow, land use change, solar irradiance and volcanoes (table 3). ERF, which accounts for all (stratospheric plus tropospheric) rapid adjustments, corresponds better to temperature change than “traditional” stratospherically adjusted radiative forcing (RF) (Myhre et al., 2013b; Forster et al., 2016). Therefore, we use relationships for ERF where they exist.

2.2.1 Carbon dioxide, methane and nitrous oxide

We use the updated Etminan et al. (2016) RF relationships for CO₂, CH₄ and N₂O, which for the first time includes band overlaps between CO₂ and N₂O. It also includes a significant upward revision of the CH₄ RF due to inclusion of previously neglected shortwave absorption, compared to the previous relationships of Myhre et al. (1998) used in AR5. Although Etminan et al. (2016) calculate RF, Myhre et al. (2013b) concluded that over the industrial era there was not sufficient evidence to state that ERF was significantly different from RF for these three gases, and ERF is taken to equal RF, although with a doubled uncertainty range. The Etminan et al. (2016) relationships are reproduced in eqs. (7) to (9), where C (ppm), M and N (ppb) have been used to represent concentrations of CO₂, CH₄ and N₂O, and the subscript pi representing pre-industrial

concentrations.

$$F_{\text{CO}_2} = [(-2.4 \times 10^{-7})(C - C_{\text{pi}})^2 + (7.2 \times 10^{-4})|C - C_{\text{pi}}| - (1.05 \times 10^{-4})(N + N_{\text{pi}}) + 5.36] \times \log\left(\frac{C}{C_{\text{pi}}}\right) \quad (7)$$

$$F_{\text{N}_2\text{O}} = [(-4.0 \times 10^{-6})(C + C_{\text{pi}}) + (2.1 \times 10^{-6})(N + N_{\text{pi}}) - (2.45 \times 10^{-6})(M + M_{\text{pi}}) + 0.117] \times (\sqrt{N} - \sqrt{N_{\text{pi}}}) \quad (8)$$

$$F_{\text{CH}_4} = [-(6.5 \times 10^{-7})(M + M_{\text{pi}}) - (4.1 \times 10^{-6})(N + N_{\text{pi}}) + 0.043] \times (\sqrt{M} - \sqrt{M_{\text{pi}}}). \quad (9)$$

- 5 Finally, a scaling to F_{CO_2} is made to ensure that a doubling of CO_2 using eq. (7) along with pre-industrial N_2O concentrations equals the user-specified value of $F_{2\times}$, which defaults to 3.71 W m^{-2} .

2.2.2 Other well-mixed greenhouse gases

For all well-mixed greenhouse gases in table 2 except CO_2 , CH_4 and N_2O the ERF is assumed to be a linear relationship of the change in gas concentration C_i since the pre-industrial era by its radiative efficiency η_i [W m^{-2} ppb]:

$$10 \quad F_i = \eta_i(C_i - C_{i,\text{pi}}); \quad i \in \{\text{gas indices } 3, 4, \dots, 30\} \quad (10)$$

where radiative efficiencies are given in table 2, i refers to index numbers in table 2 and C_i are converted to ppb. This is an established method for small greenhouse gas forcings, also used in MAGICC.

2.2.3 Tropospheric ozone

Tropospheric ozone is formed from a complex chemical reaction chain from emissions of CH_4 , NO_x , CO and NMVOC .

- 15 Furthermore its concentration is more variable in space and time than for the well-mixed greenhouse gases. Therefore, we do not calculate a globally averaged concentration. We use coefficients from Stevenson et al. (2013) to estimate tropospheric ozone ERF from emissions of NO_x , CO and NMVOC , and concentrations of methane, assuming linearity between atmospheric burden and ozone forcing:

$$F_{\text{O}_3\text{tr}} = \beta_{\text{CH}_4}(C_{\text{CH}_4} - C_{\text{CH}_4,\text{pi}}) + \beta_{\text{NO}_x}(E_{\text{NO}_x} - E_{\text{NO}_x,\text{pi}}) + \beta_{\text{CO}}(E_{\text{CO}} - E_{\text{CO},\text{pi}}) + \beta_{\text{NMVOC}}(E_{\text{NMVOC}} - E_{\text{NMVOC},\text{pi}}) + f(T) \quad (11)$$

- 20 and

$$f(T) = \min\{0, 0.032 \exp(-1.35T) - 0.032\} \quad (12)$$

The β coefficients in eq. (11) are provided in table 4 and eq. (12) is a small negative climate feedback, estimated using a curve fit to year 2000, 2030 and 2100 temperature changes under RCP8.5 in Stevenson et al. (2013). As Stevenson et al. (2013) used 1850 as their baseline for forcing calculations based on emissions data from Lamarque et al. (2010), in RCP scenarios adjusted

25 coefficients can optionally be specified for times prior to 1850 where “pre-industrial” anthropogenic emissions are taken from Skeie et al. (2011). This ensures that ERF is both equal to zero in 1765 and equal to the best estimates in Stevenson et al. (2013) for 2005. Both the differing treatment prior to 1850 and the climate feedback can optionally be switched off by the user.

2.2.4 Stratospheric ozone

30 The stratospheric ozone ERF is calculated using the functional relationship borrowed from Meinshausen et al. (2011a), namely

$$F_{O_3st} = a(bs)^c. \quad (13)$$

$a = -1.46 \times 10^{-5}$, $b = 2.05 \times 10^{-3}$ and $c = 1.03$ in eq. (13) are fitting parameters that are found by a least-squares curve fit between eq. (13) and the stratospheric ozone ERF timeseries from AR5; due to this data fitting approach, our parameters differ from MAGICC. s is the equivalent effective stratospheric chlorine (EESC) from all ozone depleting substances, calculated as
5 (Newman et al., 2007)

$$s = r_{CFC11} \sum_{i \in ODS} \left(n_{Cl}(i) C_i \frac{r_i}{r_{CFC11}} + 45 n_{Br}(i) C_i \frac{r_i}{r_{CFC11}} \right). \quad (14)$$

r_i represents fractional release values for each ODS compound taken from Daniel and Velders (2011) and reproduced in table 2. n_{Cl} and n_{Br} represent the number of chlorine and bromine atoms in compound i with the factor of 45 in eq. (14) indicating that bromine is 45 times more effective at stratospheric ozone depletion than chlorine (Daniel et al., 1999). The concentrations C_i
10 are expressed in ppb.

2.2.5 Stratospheric water vapour from methane oxidation

In AR5, the ERF from the stratospheric water vapour oxidation of methane was assumed to be 15% of the methane ERF. This was based on the methane forcing relationship of Myhre et al. (1998), which is about 20% lower than the Etminan et al. (2016) methane forcing used in FAIR. As there has been no substantial revision to the stratospheric water vapour forcing we define
15 stratospheric water vapour ERF as 12% of the methane ERF.

2.2.6 Contrails

Meinshausen et al. (2011b) did not include a forcing timeseries for contrails or contrail-induced cirrus, which contribute a small positive ERF (Boucher et al., 2013). [Three different methods to supply contrail ERF are available in FAIR: \(1\) scaling with aviation-based NOx emissions; \(2\) scaling with global supply of jet kerosene fuel, or \(3\) supplying an external forcing time series.](#) In method 1, ~~it is assumed that contrail ERF is proportional to the level of air traffic, which is in turn is proportional to aircraft emissions. We use aviation NOx emissions for this purpose, which are obtained from the RCP database. The ERF from contrails F_{con} is scaled by the ratio of aircraft NOx emissions in a given year $E_{NOx,avi}$ compared to 20052014 and multiplied by the 20052014 ERF from Lee et al. (2009) of 0.0448 W m⁻²;~~
20 F_{con} is scaled by the ratio of aircraft NOx emissions in a given year $E_{NOx,avi}$ compared to 20052014 and multiplied by the 20052014 ERF from Lee et al. (2009) of 0.0448 W m⁻²;

$$F_{con} = \frac{E_{NOx,avi}}{E_{NOx,avi,2005}} F_{con,2005}. \quad (15)$$

25 This gives a coefficient of $F_{con,2005}/E_{NOx,avi,2005} = 0.0152 \text{ W m}^{-2} (\text{Mt-aviNOx yr}^{-1})^{-1}$.

Method 2 is similar, based on kerosene fuel supplied S_{kerosene} , as a proxy for activity data. We take a 2005 global kerosene supply of 236 Gt from International Energy Agency (2018) to anchor the forcing time series calculation:

$$F_{\text{con}} = \frac{S_{\text{kerosene}}}{S_{\text{kerosene},2005}} F_{\text{con},2005}. \quad (16)$$

For method 1, the past and future aviation NOx emissions from the RCP scenarios are available in FAIR, and is introduced as the fraction of total NOx emissions attributable to aviation.

2.2.7 Aerosols

Aerosols have a lifetime of the order of days (Kristiansen et al., 2016), and the emissions are converted to forcing without an intermediate concentration step.

5 The aerosol ERF contains contributions from aerosol-radiation interactions (ERFari) and from aerosol-cloud interactions (ERFaci). ERFari includes the direct radiative effect of aerosols, in addition to rapid adjustments due to changes in the atmospheric temperature, humidity and cloud profile (formerly the semi-direct effect; Boucher et al. (2013)).

We use the multi-model results from AeroCom (Myhre et al., 2013a) ~~and to provide coefficients of how emissions affect ERFari. Again we~~ assume a linear relationship between emissions and forcing:

$$10 F_{\text{ari}} = \gamma_{\text{BC}} E_{\text{BC}} + \gamma_{\text{OC}} E_{\text{OC}} + \gamma_{\text{SOx}} E_{\text{SOx}} + \gamma_{\text{NOx}} E_{\text{NOx}} + \gamma_{\text{NH}_3} E_{\text{NH}_3} + \gamma_{\text{SOA}} E_{\text{NMVOC}} \quad (17)$$

where the default coefficients for each γ are provided in table 5 and calculated from the difference in anthropogenic emissions between 17651850 and 20102000 (Lamarque et al., 2010). Users are free to specify their own species-dependent γ for each aerosol precursor ~~and scaled for rapid adjustments (see below)~~. We assume emitted black carbon (BC) and organic carbon (OC) correspond directly to BC and OC forcing, and that emissions of sulfur compounds (SOx) correspond directly to sulfate forcing. Following Shindell et al. (2009) we assume a 60% contribution to nitrate aerosol forcing from NH₃ and 40% from NOx. We allow formation of secondary organic aerosol (SOA) to scale with emissions of anthropogenic NMVOC. Biomass burning aerosol has a net zero forcing in 2011 and is ignored, and mineral dust, which does not scale directly with an emitted component, is also disregarded. The sum of the direct effects of each component is ~~-0.35-0.328~~ W m⁻² in 2010+ assuming RCP4.5. The difference between this and the best estimate ERFari of -0.45 W m⁻² in AR5 is assumed to be due to rapid adjustments (semi-direct effects). ~~There is evidence that scattering aerosols do not exhibit significant semi-direct effects (Boucher et al., 2013), so the overall difference of -0.1 W m⁻² is assumed to be due to BC semi-direct effects. The net impact is to reduce the radiative efficiency coefficient of BC, and therefore the radiative efficiency of each species is scaled by a factor of 1.37.~~

ERFaci describes how aerosols affect clouds in the radiation budget; the two main mechanisms are changes in cloud droplet size, which changes cloud albedo (Twomey, 1977) and changes in cloud lifetime and precipitation efficiency which affects cloud fraction (Albrecht, 1989; Boucher et al., 2013). There is evidence that the ERFaci is not linear with emissions (Carslaw et al., 2013) and as such a simple linear scaling as for ERFari may not be appropriate.

In FAIR we use an emulation of the global aerosol model of Ghan et al. (2013) to estimate ERFaci from precursor emissions. The Ghan et al. (2013) method contains a series of non-linear equations that require iterative solutions and currently has not

been optimised for use in FAIR. We therefore emulate the ERF_{aci} by varying the emissions of SO_x and primary organic aerosol (the sum of BC and OC). Secondary organic aerosol is also an input to the model, but it is found that ERF_{aci} is only weakly dependent on NMVOC emissions and a simple functional form could not be found, so was eliminated as a predictor.

Informed by the simple aerosol model of Stevens (2015), we use a logarithmic dependence of ERF_{aci} on emissions that can vary both as a function of SO_x and BC+OC, which represents increasing saturation of the cloud-albedo effect with increasing emissions. We find a relationship of the form

$$G(E_{\text{SO}_x}, E_{\text{BC+OC}}) = -1.95 \log(1 + 0.0111 E_{\text{SO}_x} + 0.0139 E_{\text{BC+OC}}) \quad (18)$$

where the coefficients in eq. (18) are found with a least-squares optimisation routine ($r^2 = 0.938$). The modelled and simulated outputs are compared in figure S1.

Equation (18) was derived from a climate model, and produces a present-day ERF_{aci} that is stronger than the central estimate of -0.45 W m^{-2} from AR5. We therefore scale eq. (18) in order to obtain a forcing of -0.45 W m^{-2} in 2011 under RCP4.5 emissions:

$$F_{\text{aci}} = -0.45 \frac{G(E) - G(E_{1765})}{G(E_{2011}) - G(E_{1765})} \quad (19)$$

where $E = (E_{\text{SO}_x}, E_{\text{BC+OC}})$ refers to emissions, and a numerical subscript refers to a particular year. The emissions for 1750 from Skeie et al. (2011) are used for year 1765, and a linear interpolation between 1765 and 1850 applied.

2.2.8 Black carbon on snow

The best estimate ERF of 0.04 W m^{-2} in AR5 for 2011 is compared to the BC emissions in 2011 from Meinshausen et al. (2011b), with this scaling factor assumed to hold for all years. The relationship is given by

$$F_{\text{BCsnow}} = 0.00494 E_{\text{BC}}, \quad (20)$$

where E_{BC} is BC emissions in Mt yr^{-1} .

2.2.9 Land use change

Land use forcing is a result of surface albedo change (Andrews et al., 2017) and changes in evapotranspiration patterns (Jones et al., 2015), which is often due to deforestation for agriculture (Myhre and Myhre, 2003). Cropland has a higher albedo than the forest that it replaces, reflecting more incident solar radiation and therefore resulting in a negative ERF; additionally, deforestation in boreal regions may unmask snow-covered ground, again increasing albedo.

Deforestation produces land-use related CO₂ emissions. The total amount of deforestation since pre-industrial times could therefore be expected to scale with cumulative land-use related CO₂ emissions. This is the default approach taken in FAIR. A regression of non-fossil CO₂ emissions against land use ERF in AR5 gives

$$F_{\text{landuse}} = -1.14 \times 10^{-3} \sum_{j=0}^t E_{\text{CO}_2\text{land},j}, \quad (21)$$

where the coefficient has units $\text{W m}^{-2} (\text{Gt C})^{-1}$.

The simple relationship in eq. (21) does not take into account latitude dependence of surface albedo or any evapotranspiration changes. As a zero-dimensional model, FAIR does not include geographical dependence of individual forcing effects, which may differ significantly between forcing pathways (for example the scenarios typically used to drive integrated assessment models). Inclusion of evapotranspiration effects, again which differ between tropical and boreal regions (Jones et al., 2015), is
5 challenging as they do not directly relate to an emitted species or a change in radiative forcing (Pielke et al., 2002). Nevertheless, we conclude that this simple treatment is acceptable, firstly as the range of land use forcing uncertainty is relatively large (so much that the sign of the forcing is not known with confidence (Myhre et al., 2013b)), secondly because at least in the best estimate the forcing is a small fraction of the present-day total, and thirdly because the future trajectory of the land use forcing in the RCP datasets is very similar to that predicted by FAIR, suggesting that a dependence on cumulative land-use CO_2
10 emissions is an important component of the land use forcing in MAGICC.

Noting that this simple relationship may not be suitable in all cases, the user is free to supply their own time series of ERF from land use change. If gridded land-use data is available, the transitions to and from forested land each year can be convoluted with the marginal contribution to land use forcing per square kilometre deforestation (e.g. from Jones et al. (2015)).

2.2.10 Solar variability

15 The SOLARIS-HEPPA v3.2 solar irradiance dataset prepared for CMIP6 is used to generate the solar ERF, which includes projections of the variation in future solar cycles from 1850 to 2300 (Matthes et al., 2017). ERF from solar forcing is calculated as the change in solar constant since 1850 divided by 4 (average insolation) and multiplied by 0.7 (representing planetary co-albedo). This approach is also used in Meinshausen et al. (2011b). Prior to 1850, we revert to the solar forcing from AR5.

2.2.11 Volcanic aerosol

20 Historical volcanic forcing is punctuated by several large eruptions that cause large but short-lived negative forcing episodes, with several smaller eruptions that cause year-to-year changes in the volcanic forcing. In order to generate a historical volcanic ERF time series we first start with gridded volcanic optical depths taken from the Easy Volcanic Aerosol generator over the 1850–2014 period (Toohey et al., 2016) which will be used to drive CMIP6 models. A number of time slice experiments with various scalings of the historical mean volcanic optical depth are run in the HadGEM3-GA7.1 climate model (Walters et al.,
25 2017), where it was found that aerosol ERF scales as $-18\tau_{\text{vol}}$ (where τ_{vol} is globally averaged volcanic aerosol optical depth at 550 nm). This scaling factor is consistent with other HadGEM models (Gregory et al., 2016), although weaker than the value of $-26\tau_{\text{vol}}$ adopted in AR5. The discrepancy is claimed to be due to rapid adjustments, in which case our adoption of the less negative value is consistent with the ERF definition.

In the context of measuring forcing since the pre-industrial, we have to assume an “average” level of volcanic background
30 aerosol. We therefore define the 1850–2014 period to have a mean volcanic forcing of zero. To achieve this we subtract the mean (negative) forcing from the historical period, resulting in a quiescent year ERF of around $+0.1 \text{ W m}^{-2}$. A similar approach was taken in Meinshausen et al. (2011b), with a higher quiescent year forcing of about $+0.2 \text{ W m}^{-2}$. Prior to 1850 we use the

AR5 dataset, scaled by 18/26 to match the differences in optical depth/forcing relationships, and from 2015 onwards volcanic forcing is defined to be zero. For solar and volcanic forcing, users are free to provide a custom forcing time series or to use the AR5 or RCP datasets which are both available in FAIR.

5 2.3 Temperature change

In simple impulse-response models, forcing is related to total temperature change in year t , T_t , by a two-time constant model (Boucher and Reddy, 2008; Myhre et al., 2013b; Millar et al., 2015, 2017). FAIR v1.3v1.2 takes this approach with a small modification compared to FAIR v1.0 to allow for forcing-specific efficacies ϵ_j such that

$$T_{t,i} = T_{t-1,i} \exp(1/d_i) + \sum_{j=0}^{12} (q_i \epsilon_j F_j (1 - \exp(1/d_i))); \quad i = 1, 2. \quad (22)$$

- 10 Owing to the use of ERF rather than RF in FAIR v1.3v1.2 and its better correspondence with temperature, efficacies are assumed to be unity for all forcing agents except black carbon on snow ($j = 9$), where an efficacy of 3 is used following Bond et al. (2013). The coefficients d_1 and d_2 govern the slow ($i = 1$) and fast ($i = 2$) temperature changes from a response to forcing from the upper ocean and the deep ocean respectively (Millar et al., 2015). The total temperature change in year t is the sum of the slow and fast components, i.e. $T_t = T_{t,1} + T_{t,2}$. F_j represents the 13 individual forcing agents in year t calculated
- 15 in section 2.2 (see also table 3). d_1 and d_2 default to 239 and 4.1 years which are fit to match the mean of CMIP5 models (Geoffroy et al., 2013). The coefficients q_1 and q_2 (units $\text{K W}^{-1} \text{m}^2$) are determined by solving a matrix equation given TCR, ECS, d_1 , d_2 and the ERF from a doubling of CO_2 , $F_{2\times} = 3.71 \text{ W m}^{-2}$ (Myhre et al., 2013b):

$$T_{\text{ECS}} = F_{2\times} (q_1 + q_2); \quad (23)$$

$$T_{\text{TCR}} = F_{2\times} \left(q_1 \left(1 - \frac{d_1}{D} \left(1 - \exp\left(-\frac{D}{d_1}\right) \right) \right) + q_2 \left(1 - \frac{d_2}{D} \left(1 - \exp\left(\frac{D}{d_2}\right) \right) \right) \right) \quad (24)$$

- 20 giving the relative contributions to the fast and slow components of the warming. $D = \log(2)/\log(1.01) \approx 69.7$ years is the time to a doubling of CO_2 with a compound 1% per year increase in CO_2 concentrations.

There is some evidence that ECS and TCR have not been constant values over the historical period (Gregory and Andrews, 2016; Gregory et al., 2015), and that ECS does not necessarily assume a constant value in CMIP5 modelling experiments (Armour, 2017). FAIR has the capability to model time-evolving ECS and TCR by updating the q_1 and q_2 values in each time

- 25 step.

3 Projections using a large ensemble

To test the model response to a range of forcing pathways, we perform a 100,000-member Monte Carlo simulation using emissions from the RCP datasets (Meinshausen et al., 2011b). Emissions themselves are not altered from the RCP timeseries, but the TCR, ECS, carbon cycle response to increasing temperature (r_T) and cumulative emissions (r_C) along with the pre-industrial value of iIRF₁₀₀ (r_0), plus the ERF scale factors for each of the 13 forcing agents, are drawn from distributions. FAIR is run from 1765 (the start of the RCP emissions datasets) to 2100.

3.1 Constraint to historical temperature observations

As a wide range of forcing, and thus temperature, scenarios can be generated, there are a proportion of ensemble members generated that fall outside the range of plausibility. We constrain the full 100,000 member ensemble (hereafter FULL) to the observed temperature change from the Cowtan and Way (2014) dataset (hereafter C&W) to assess plausibility; ensemble members that satisfy the temperature constraint are designated as Not Ruled Out Yet (NROY) and the majority of the discussion of the results in section 4 focuses on this dataset. We rebase all of the temperatures to the 1861–1880 mean following Richardson et al. (2016), to represent a “pre-industrial” state that is relatively free from volcanic eruptions but with a reasonable global coverage of temperature observations. An ordinary least-squares regression of temperature change versus time from 1880–2016 is used to calculate the linear warming trend in each ensemble member. The regression is also performed for the C&W “observational” dataset to estimate the observed warming rate. The confidence interval around the C&W warming rate is inflated by a factor that represents the lag-1 autocorrelation of residuals (i.e. the trend-line estimate from the regression minus the C&W “observations”) which accounts for internal climate variability (Santer et al., 2008; Thompson et al., 2015) and is the same method used in AR5 to estimate linear temperature trends. The constraint is satisfied for an ensemble member if the modelled trend falls within the 5–95% range of trend from C&W of 0.95 ± 0.17 K.

The C&W observed warming from 1880–2016 higher than the HadCRUT4 estimate of 0.91 ± 0.18 K for the same timeframe. The infilling of grid boxes where no or limited data are available accounts for these differences, as sparse observations are typically in polar regions which warm faster than the global mean (Cowtan and Way, 2014). Under this constraint approximately 26% of the FULL ensemble is retained in NROY.

It should be stressed that there are several issues to consider when attempting to derive plausible parameter sets from observational data. These include the type of observational constraints to employ (Meinshausen et al., 2009), the length of the historical record (e.g. Otto et al. (2013)), the separation of forced response from natural variability (Haustein et al., 2017), and assumptions surrounding prior distributions (Frame et al., 2005).

3.2 Sampling ECS and TCR

The ECS and TCR from CMIP5 models (Forster et al., 2013) are used to generate a joint lognormal distribution. Random variables are sampled using the R package MethylCapSig¹ using the mean, standard deviation and correlation coefficient ($r = 0.81$) between ECS and TCR in CMIP5. A lognormal distribution is representative of distributions of ECS and TCR in the literature (Meinshausen et al., 2009; Rogelj et al., 2012; Flato et al., 2013; Millar et al., 2017). The sampled joint and marginal distributions are shown as black contours and curves in fig. 3. We sample 100,000 ECS/TCR pairs; sampled pairs where $ECS < TCR$ are rejected and redrawn. A joint distribution is used because ECS and TCR are highly correlated and low values of the realised warming fraction (TCR divided by ECS) are inconsistent with models and observations (Millar et al., 2015). For other sampled quantities in this section a 100,000 member ensemble is also generated.

¹<https://cran.r-project.org/package=MethylCapSig>

3.3 Sampling thermal response and carbon cycle parameters

We allow $F_{2\times}$, the ERF due to a doubling of CO_2 , to assume a Gaussian distribution with 5–95% confidence interval of 20% around the best estimate ERF of 3.71 W m^{-2} (Myhre et al., 2013b). d_1 (mean 239 years, standard deviation 63 years) and d_2 (mean 4.1 years, standard deviation 1.0 years) in eq. (22) are also varied based on truncated Gaussian distributions (no values outside $\pm 3\sigma$ allowed, primarily to prevent unrealistically small or negative values of the slow response d_1). Although FAIR is able to model the response to time-varying ECS and TCR, we use time-invariant values in our ensemble.

Some uncertainty in the carbon cycle parameters is assumed with samples of r_0 , r_C and r_T taken from Gaussian distributions. r_0 , r_C and r_T are given 5–95% confidence intervals of 13% of the default parameter value following Millar et al. (2017).

3.4 Sampling ERF uncertainties

The uncertainty in each of the 13 forcing components is modelled following the 5–95% confidence intervals for each forcing from AR5 (Myhre et al., 2013b, Table 8.6) in 2011 (table 3). This is achieved by scaling the ERF values calculated in section 2.2; the uncertainty ranges are given in table 3. The scaling factor is applied to the whole time series. [A time-varying scale factor for forcing can be used, and an option is provided to exactly replicate the AR5 historical time series for each component, but we do not apply it here.](#) Most uncertainties are assumed to be Gaussian, the exceptions being contrails and BC on snow which are lognormally distributed (with geometric standard deviations 1.92 and 1.65 respectively, following AR5), and aerosols which are modelled as two half-Gaussian distributions, treating values above and below the best estimate separately. These ERF uncertainties are assumed to be uncorrelated with each other.

20 4 Results from the NROY ensemble for the RCP scenarios

4.1 ECS and TCR

The FULL and NROY joint and marginal distributions of ECS and TCR are shown in fig. 3.

The temperature constraint in NROY results in distributions of ECS and TCR that are lower than in FULL. Some of the prior sample space in which ECS and TCR are larger than the AR5 likely ranges has been rejected in the NROY distribution. While the possibility that $\text{ECS} > 5 \text{ K}$ cannot be ruled out, it appears less likely than would be inferred from CMIP5 models, although it should be stressed that time-varying feedbacks are not accounted for in this large ensemble which would allow ECS to increase over time (Armour, 2017). From the marginal distributions, we estimate that ECS and TCR are [2.86 \(2.01 to 4.22\)](#)~~2.93 (2.04 to 4.32)~~ K and [1.53 \(1.05 to 2.41\)](#)~~1.59 (1.07 to 2.50)~~ K (median; (5–95% range)) respectively in the NROY ensemble, similar to but a little more tightly constrained than the AR5 *likely* (>66% probability) ranges of 1.5 to 4.5 K and 1.0 to 2.5 K, noting that the AR5 ranges are estimated from a combination of models, observations and expert judgement. The ratio of TCR to ECS, the realised warming fraction (RWF), is approximately independent of TCR in CMIP5 models (Millar et al., 2015) and the prior distribution could alternatively be defined in terms of the TCR and RWF joint distribution, which is

explored in section 5. The FULL and NROY median and 5 to 95% ranges of RWF are 0.56 (0.41 to 0.76) and 0.54 (0.40 to 0.72) respectively, which is close to the range of CMIP5 models (0.45 to 0.75, Millar et al. (2017)).

4.2 Historical and future greenhouse gas concentrations

5 The historical (1765–2005) greenhouse gas concentrations from the RCP scenarios in Meinshausen et al. (2011b) were assimilated from observations of in-situ and ice core records and represent a best estimate of the actual concentrations over this period. We therefore assume that the RCP data represents the best estimate of the historical concentrations and compare our estimates from the NROY ensemble using the emissions-driven model.

The FAIR model reproduces the historical concentrations of greenhouse gases (fig. 4). The atmospheric concentrations of CO₂ estimated from FAIR are up to 9 ppm lower than MAGICC6 in the period 1880–1950 (fig. 4a). A simple carbon cycle model cannot reproduce the kinks in the observational CO₂ trend without large changes in the input emissions. However, between 1950 and 2005, the differences between the two curves are small. The post-2005 atmospheric CO₂ concentrations are slightly higher than those estimated by MAGICC6 for the RCP scenarios, but the MAGICC6 concentrations are within the uncertainty range from the NROY ensemble. FAIR projects best estimate CO₂ concentrations of 427 (413 to 443), 552 (527 to 578), 695 (661 to 731) and 979 (926 to 1040) ppm for RCP2.6, RCP4.5, RCP6 and RCP8.5 in 2100. Here, the uncertainty in CO₂ concentrations relates to the range of carbon cycle parameters and the temperature dependence on carbon uptake sampled in the large ensemble.

The historical CH₄ and N₂O concentrations in FAIR have been tuned to agree with Meinshausen et al. (2011b) by adjusting natural emissions as described previously (fig. 4b,c). There are some small differences in the future CH₄ and N₂O concentrations in the RCPs using fixed atmospheric lifetimes and constant (present-day) natural emissions.

Kyoto Protocol gases have been grouped as HFC134a-eq based on their radiative efficiency, and ODSs grouped as CFC12-eq similarly (fig. 4d,e). Small differences between the models in future scenarios may be a result of the assumption of a change in the rate of the Brewer-Dobson circulation in MAGICC6 (Meinshausen et al., 2011a), which increases the efficiency of the stratospheric sink for these gases. This temperature-dependent effect is not included in FAIR. Over the historical period, the differences are a result of the natural emissions of CF₄ (contributing to HFC134a-eq), and CH₃Br and CH₃Cl (contributing to CFC12-eq) providing a non-zero background state of these greenhouse gas equivalents in FAIR. In the RCP historical dataset these background concentrations have not been added to the HFC134a-eq and CFC12-eq timeseries.

4.3 Historical, present and future radiative forcing

Figure 5 shows the comparison between FAIR and MAGICC6 for the 13 forcing agents considered in FAIR for the NROY ensemble. The ERF time series for the historical period in AR5 is also shown (IPCC, 2013). The updated radiative forcing relationship for CH₄ increases radiative forcing substantially (fig. 5b). The new relationship for N₂O results in a slightly lower ERF estimate in FAIR than RF in MAGICC6 (fig. 5c) which is offset by the higher concentrations of N₂O in FAIR. The FAIR estimate of CO₂ forcing is also higher than MAGICC for the RCPs, but the ERFs from FAIR and RFs from MAGICC for the

minor greenhouse gases are similar (fig. 5a,d). For non-CO₂ gases, AR5 did not provide a breakdown of ERF by individual gas, so the total ERF has been scaled by the ratios in the MAGICC6 timeseries.

For tropospheric ozone, the Stevenson et al. (2013) relationship agrees well with AR5 until around 1970, from which point it is larger than AR5. There is also a large relative difference between this relationship and the MAGICC estimate in AR5 out to 2100 (fig. 5e). The shape of the stratospheric ozone ERF curve between AR5 and MAGICC6 differs, but it can be seen that the AR5 historical ERF is well emulated as it uses the same functional relationship as AR5 (fig. 5f). Stratospheric water vapour from methane oxidation depends on the underlying methane forcing and is similar to the AR5 timeseries (fig. 5g). Contrail ERF shows a similar time evolution over the historical period to AR5 (fig. 5h). Historically, ERF from aviation contrails has been small, but may become more substantial in the future. The median aerosol ERF in FAIR is slightly more negative than in AR5 from around 1900 to 2011 (fig. 5i) but is less negative than the RCP projections. We reiterate here that the RCPs report RF from MAGICC rather than ERF.

BC on snow has a smaller ERF in FAIR than the corresponding RF in MAGICC6, although the efficacy factor of 3 used in FAIR results in a similar effect on temperature between the models (fig. 5j). Estimates of future land use forcing in FAIR follow a similar shape to the Meinshausen et al. (2011b) dataset with slightly less negative best estimates to the AR5 ERF 2011 best estimate; agreement in the historical period to either MAGICC6 or AR5 is less good, but the general trajectory of forcing is correct (fig. 5k). There are substantial differences between the volcanic forcing datasets in FAIR, AR5 and the RCPs that are not easy to discern at the resolution of the plot (fig. 5l): generally, the AR5 dataset gives more negative forcing than the RCPs during volcanically active years, and also defines the absence of volcanoes as zero forcing whereas the RCP and FAIR datasets define zero to be the average of the historical period. Solar forcing is used from the new CMIP6 dataset which is reasonably similar to the RCP time series for historical forcing but exhibits some differences in the future owing to the assumed inter-cycle variability that was not present in CMIP5 (fig. 5m).

Figure 5n shows the sum of the forcing components. The best estimate sum of ERF follows AR5 closely over the historical period, which is intentional. In the RCP future scenarios, the FAIR best estimates of ERF are higher than the corresponding RF estimates in MAGICC. This is in part due to the increased CH₄, tropospheric ozone and contrail forcing in FAIR, and less negative total aerosol ERF for RCP2.6. The FAIR model projects 2100 ERFs (median and (5–95% credible intervals)) of 2.62 (1.79 to 3.64), 4.62 (3.30 to 6.22), 5.84 (4.07 to 8.00) and 9.34 (6.84 to 12.44) for RCP2.6, RCP4.5, RCP6.0 and RCP8.5 respectively.

4.4 Relationship between forcing components, ECS and TCR

The distribution of ERF in 2017 for aerosols, greenhouse gases and the anthropogenic total in both the FULL and the NROY ensembles assuming the RCP8.5 forcing pathway is shown in fig. 6 and table 6. The temperature constraint in NROY permits a wider range of greenhouse gas ERF than shifts the distribution of ERF for greenhouse gases slightly to the left but is similar to the FULL ensemble. For aerosols, the distribution of ERF in NROY is again slightly wider than in FULL, with a reduction in the probability of a strong present-day aerosol forcing. The median estimate of net anthropogenic ERF of

2.632-70 W m^{-2} in 2017 for NROY is a little lower than similar to the unconstrained FULL estimate of 2.73 W m^{-2} with a wider uncertainty range.

There are negative correlations between aerosol radiative forcing and ECS/TCR (fig. 7). A large negative aerosol forcing requires a high ECS to balance and recreate realistic observed temperatures (Forest et al., 2006). Millar et al. (2015) highlighted the necessity of anti-correlation between TCR and aerosol forcing in observational constraints. The aerosol forcing on TCR constraint is tighter than that on ECS, evidenced by the narrower mass of points in the TCR plot (fig. 7b) compared to the ECS plot (fig. 7a). A high value for TCR (greater than 2.5 K) or ECS (greater than 5 K) is only possible with a strong negative present-day aerosol forcing (more negative than about -1.0 W m^{-2}).

4.5 Observed and future temperature changes

Figure 8a shows the transient historical and RCP-projected temperature change for 1850–2100 along with the 2081–2100 median, 1σ (16–84% range) and 5–95% credible range for the NROY ensemble (fig. 8b), (c.f. Rogelj et al. (2012), Collins et al. (2013, fig. 12.8b)). For the RCP scenarios, the median NROY estimates of temperature change for 2081–2100 are 1.49, 2.29, 2.66 and 3.914 , 53 , 2.37 , 2.74 and 4.03 K above pre-industrial for RCPs 2.6, 4.5, 6.0 and 8.5 respectively. The median, 1σ and 5–95% ranges of total temperature change predicted from FAIR are a little lower for RCPs 2.6, 4.5 and 6.0 than about the same for RCP2.6 and RCP4.5 as those predicted by the emissions-driven MAGICC6 experiments which are reported in AR5 (Rogelj et al., 2012; Meinshausen et al., 2009), a little lower for RCP6, and substantially lower for RCP8.5. The RCP8.5 temperature change is lower ease is despite higher 21st century ERF profiles in FAIR compared to RF in MAGICC. The difference of 0.60 , 5 K in the median end-of-century warming in RCP8.5 could be particularly important in policy assessments.

Differences between the models can arise from many sources. The results of Rogelj et al. (2012) are based on best estimates of the ECS/TCR and radiative forcing from the IPCC Fourth Assessment Report (AR4), whereas we guide FAIR using AR5 forcings. Differences between this study and Rogelj et al. (2012) could be due to differences in the historical radiative forcing time series. The RF over the 1861–1880 to 2005 period, which forms the bulk of the period used to constrain the ensemble to observed temperatures, in Meinshausen et al. (2011b) is 1.72 W m^{-2} whereas the ERF differences are 1.98 W m^{-2} in AR5 and 1.97 , 1.96 W m^{-2} in FAIR over the same period. Therefore, the same observed temperature change would be recreated with a smaller RF in Rogelj et al. (2012) than the corresponding ERF in FAIR, and the same future forcing in MAGICC6 would lead to a higher temperature change than in FAIR. Other differences between the studies include a different selection of ECS and TCR priors in Meinshausen et al. (2009) and Rogelj et al. (2012) (based on AR4, but not substantially different from the CMIP5 models used in this study), a different method of constraining to observed temperatures, and different assumptions regarding the strength of future aerosol and ozone forcing. The sensitivity to some of these assumptions is tested in section 5.

4.6 Transient Climate Response to Emissions

There is an approximately linear relationship between cumulative CO_2 emissions and temperature, independent of the actual emissions pathway taken, providing temperature is still increasing (Allen et al., 2009; Collins et al., 2013). Using this linearity

we can diagnose the transient climate response to emissions (TCRE), defined as the change in temperature for a 1000 Gt cumulative emission of carbon.

We show both the TCRE assuming CO₂ forcing alone and the temperature change due to all forcing agents but measured against cumulative carbon emissions (fig. 9). When including the effect of non-CO₂ forcing on the total temperature change, the temperature response is substantially larger than for CO₂ forcing alone. This indicates that a smaller cumulative CO₂ emission is required to reach the same temperature change, and is a result of the total non-CO₂ forcing being positive. This same conclusion was reached in Collins et al. (2013) when assessing a suite of earth system models.

To determine the TCRE we run FAIR in CO₂-only mode. We measure cumulative CO₂ emissions and temperature change since 1870, as this is the date from which reliable estimates of carbon emissions start (Le Quéré et al., 2016) and is also at the centre of the 1861–1880 period used to evaluate temperature changes.

The NROY ensemble in FAIR shows a TCRE of 0.95 to 2.220.97 to 2.31 K for a cumulative carbon emission of 1000 Gt with a best estimate of 1.394.44 K. We diagnose TCRE based on the RCP8.5 simulation. The TCRE range from FAIR is within the range of estimates from AR5 (0.8 to 2.5 K, Collins et al. (2013)). Towards higher cumulative CO₂ emissions in RCP8.5 the temperature response has a slightly concave shape. The slight (rather than moderate) downward curvature is also present in CMIP5 earth system models, as the increase in airborne fraction of CO₂ with emissions almost cancels out the logarithmic relationship between CO₂ concentration and temperature (Millar et al., 2016).

~~The TCRE curve can be inverted to consider the remaining carbon budget until either 1.5 K or 2 K total warming is reached (the limits imposed in the Paris Agreement). In RCP8.5, a total of 555 GtC has been emitted over the period 1870–2016, which is close to the observational best estimate of 565 GtC (Le Quere et al., 2016). Considering all forcing agents (the red curve in fig. 9), the cumulative emissions limit required to ensure peak warming remains under 1.5 K is 725 (590 to 943) GtC. To remain under 2 K of total warming, the allowable cumulative emissions are 968 (734 to 1323) GtC.~~

4.7 Top of atmosphere energy imbalance

The top of atmosphere energy imbalance N can be diagnosed from (Forster et al., 2013)

$$N = F - \lambda T \tag{25}$$

where λ is the climate feedback parameter and $\lambda = F_{2\times}/ECS$. In fig. 10 we compare FAIR model outputs from the NROY ensemble under RCP4.5 to observations of the earth’s energy imbalance from satellites (Clouds and the Earth’s Radiant Energy System; CERES) and from the array of Argo floats (Argo, 2000), which measure ocean temperature which is the largest component of the change in earth’s energy budget. Both datasets are taken from Johnson et al. (2016).

For most years from 2001 to 2015 the net energy balance from CERES is within the uncertainty range estimated from the FAIR NROY ensemble. The Argo estimate of N is more variable prior to 2005, after which coverage of the Argo floats saw a large increase (Johnson et al., 2016). From 2005 onwards, all Argo estimates but one fall inside the credible range of FAIR estimates. Both CERES and Argo observations from 2005 onwards are clustered towards the lower half of the credible range

from FAIR, which may indicate that ECS over the 2005–2015 period could be towards the lower end of the credible range estimated from the NROY ensemble.

5 5 Sensitivity to prior distributions and constraints

To determine the robustness of the results of the NROY ensemble, the input assumptions were varied or the ensemble members subjected to a different constraint as described in this section. The results are summarised in tables 7 to 9.

5.1 Prior distributions of ECS and TCR

The prior distributions of ECS and TCR have a large influence on the posterior distributions attained (Pueyo, 2012). Here we test the dependence of the shape of the posterior distributions of ECS and TCR in the constrained samples on the choice of prior distributions.

As the RWF is approximately independent of TCR we use an alternative prior starting with the distributions of TCR and RWF. Noting the analysis of Collins et al. (2013), the AR5 likely range of TCR of 1.0 to 2.5 K is taken to be most probable, with values between 0.5–1.0 K and 2.5–3.5 K possible but unlikely. A trapezoidal distribution in TCR with these limits is constructed, therefore not expressing any prior judgement about the most likely value of TCR within the AR5 likely range. The RWF is sampled from a Gaussian distribution with mean 0.6 and 5–95% range of 0.45–0.75 following Millar et al. (2017), truncated to fall within the 0.2–1.0 range. These ranges are subjective choices based on evidence from CMIP5 models. The posterior distribution of ECS in particular can be sensitive to the choice of prior distribution (Frame et al., 2005; Pueyo, 2012). Figure S2 shows the alternative prior distributions and the posteriors obtained as a result of constraining to the C&W observed temperatures.

The best estimate and credible range of ERF is very similar to NROY with the alternative prior distributions (table 8). However the future temperature projections under the RCPs span a wider range than in NROY (table 9). This is due to the wider range of ECS and TCR admitted in the posterior distributions (table 7) using this alternative prior.

5.2 ERF from a doubling of CO₂

The canonical RF value of $F_{2\times} = 3.71 \text{ W m}^{-2}$ may not be applicable when considering all land surface and tropospheric rapid adjustments in the definition of ERF. For CO₂ forcing rapid adjustments include cloud changes that are not driven by temperature change (Gregory and Webb, 2008) and land surface adjustments consequential to plant stomatal conductance (Doutriaux-Boucher et al., 2009). The mean ERF for a doubling of CO₂ in CMIP5 models was found to be 3.44 W m^{-2} (Forster et al., 2013). The simulation is repeated with this new lower ERF value for a doubling of CO₂, with the same uncertainty of 20%.

It is found that this lower value of $F_{2\times}$ slightly lowers the best estimate and credible range of ECS, TCR and ERF, but the temperature change under the RCP scenarios are higher than in NROY due to non-CO₂ forcings. This behaviour can be analysed with the help of eq. (25). In equilibrium states, the TOA energy imbalance $N = 0$ and eq. (25) is rearranged to yield

$T = F_{2\times}/\lambda$. If $F_{2\times}$ is found to take a lower value, and ensemble members are constrained to the same observed temperature, then the climate sensitivity $1/\lambda$ must be higher to compensate. Therefore, the same positive future non-CO₂ forcing time series will produce higher temperatures in the future.

5.3 Historical temperature constraint

Historical temperatures were also constrained using the HadCRUT4 dataset without infilling (Morice et al., 2012), along with the GISTEMP (Hansen et al., 2010), Berkeley Earth (Berkeley Earth, 2017) and NOAA (Zhang et al., 2017) observational datasets. The linear 1880–2016 trends are 0.91 ± 0.18 K, 0.99 ± 0.22 K, 1.07 ± 0.16 K and 0.93 ± 0.24 K respectively. All datasets, including C&W (0.95 ± 0.17 K), were accessed on 17 October 2017.

We also perform analysis on the FULL dataset, where the input assumptions are guided by CMIP5 models and AR5 uncertainty ranges but no constraint to historical temperature is performed. We show in tables 7 to 9 that ECS, TCR, TCRE, ERF and temperature change depend slightly on the dataset of constraint, with datasets showing more warming over the historical period also projecting warmer 2100 temperatures under the RCP scenarios. Using the FULL ensemble however leads to wide uncertainty bounds and higher median estimates of these diagnosed parameters than using any of the constrained ensembles. Therefore, using a historical temperature constraint rejects parameter combinations that produce larger future temperature changes.

6 Conclusions

We present a simple model, FAIR v1.3v1.2, that calculates global temperature change, effective radiative forcing from a variety of drivers, and concentrations of greenhouse gases. The emissions-based model is based on the FAIR v1.0 carbon cycle-climate model with an extension for emissions of non-CO₂ greenhouse gases, ozone precursors and aerosols. This version of FAIR, which is tuned to the effective radiative forcing timeseries in AR5 over the historical period, provides ERFs that are close to the target radiative forcings from the RCP scenarios in 2100. FAIR was not tuned to emulate the radiative forcing in the MAGICC6 model, however it closely matches the concentrations of greenhouse gases projected in that model.

Within FAIR, the response of the carbon cycle model can be adjusted via the rate of uptake of carbon by land and ocean processes parameterised as a function of total temperature change and cumulative carbon emissions (iIRF₁₀₀). Emissions and concentrations are converted to effective radiative forcing and the relationship of ERF to temperature change is governed by the TCR, ECS, and the efficacy of each of the 13 separate forcing categories considered in the model. The emulation of specific earth system models is therefore possible as discussed by Millar et al. (2017).

Using a correlated joint lognormal prior distribution of ECS and TCR based on CMIP5 models, running a 100,000 member ensemble in FAIR and keeping only those ensemble members that match the rate of temperature change from 1880–2016 in from Cowtan & Way (the not ruled out yet or NROY ensemble), we find that the median and 5–95% credible range of ECS and TCR to be 2.86 (2.01 to 4.22) K and 1.53 (1.05 to 2.41) 2.93 (2.04 to 4.32) K and 1.58 (1.07 to 2.50) K respectively. The transient climate response to CO₂ emissions (TCRE) is diagnosed from a CO₂-only ensemble and found to be 1.40 (0.96 to

35 ~~2.23~~~~1.44~~~~(0.97 to 2.31)~~ K (1000 GtC)⁻¹. These ranges are similar to the IPCC AR5 likely ranges for ECS, TCR and TCRE, albeit with tighter credible bounds. The NROY best estimates and ranges are not very sensitive to a lower estimate of the ERF from a doubling of CO₂ or a different observational temperature datasets to constrain the historical temperature change rather than the Cowtan and Way (2014) dataset. They are more sensitive to the prior distribution of ECS and TCR, particularly for the constraint on ECS. All methods of constraint lead to lower median and credible range estimates of ECS, TCR and TCRE than
5 not constraining to temperature at all (the FULL ensemble, with input parameters estimated from the distribution of CMIP5 models and ERF uncertainties based on AR5 estimates).

Our estimate of TCR is not as low as the range derived by Otto et al. (2013) from observational constraints (0.9 to 2.0 K). Similarly our best estimate of ECS is higher than the estimate provided by Gregory and Andrews (2016) of around 2 K using observed sea-surface temperatures and sea-ice in two atmosphere-only GCMs. While we cannot absolutely rule out values of
10 ECS greater than 5 K or TCR greater than 2.5 K, this would require a strong present-day aerosol forcing (at least as negative as -1.0 W m⁻², ~~but probably more so~~) to balance. Progress towards tightening these upper bounds could therefore be achieved with a better understanding of the present-day aerosol forcing (Stevens et al., 2016).

Temperature changes projected in the NROY ensemble in 2100 are ~~a little lower than~~~~similar to~~ those from Rogelj et al. (2012) for the RCP scenarios, except for RCP8.5 where ~~the~~FAIR is ~~0.60~~~~5~~ K lower in the median response. This is due to the lower ensemble estimates of ECS and TCR in NROY, and the differences in present-day minus 1850 radiative forcing between AR5/NROY and the RCP radiative forcing in Meinshausen et al. (2011b). Nevertheless, under RCP8.5 the median
5 year 2100 temperature projection is ~~4.324~~~~45~~ K above the pre-industrial in our NROY ensemble, which would have very severe global consequences. Conversely the median estimate for RCP2.6 is ~~1.48~~~~52~~ K, suggesting about a 50% chance of limiting end-of-century warming to 1.5 K under this pathway.

FAIR is useful for creating large ensembles of future temperature change based on input uncertainties in the carbon cycle parameters and effective radiative forcing strengths. This can be used for instance to assess the impacts of emissions commitment scenarios or committed warming (Ehlert and Zickfeld, 2017), or if a certain category of emissions such as aerosols are
10 increased or decreased in the future. FAIR can be used with integrated assessment models to calculate the social cost of carbon in the presence of non-CO₂ forcing agents. Following the 2015 Paris Agreement and in anticipation of the 2018 IPCC Special Report, the FAIR model can be used to investigate emissions pathways consistent with 1.5 K and 2 K total warming limits, including remaining carbon budgets, and give probabilistic indications of the likelihood of these limits being breached.

15 *Code availability.* The source code can be obtained at <https://github.com/OMS-NetZero/FAIR> and can also be installed from the Python Package Index (<https://pypi.org/project/fair/>). A user guide is included in the supplementary material.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. CJS and PMF acknowledge financial support from the Natural Environment Research Council under grant NE/N006038/1. The authors are grateful to Zebedee Nicholls (University of Melbourne) and Robert Gieseke (PIK Potsdam) for assistance with repository management and code streamlining. We also thank Steve Ghan (Pacific Northwest University) for providing his original code for calculating global aerosol indirect effects.

References

- Albrecht, B. A.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245, 1227–1231, 1989.
- Allen, M. R., Frame, D. J., Huntingford, C., Jones, C. D., Lowe, J. A., Meinshausen, M., and Meinshausen, N.: Warming caused by cumulative carbon emissions towards the trillionth tonne, *Nature*, 458, 1163–1166, <https://doi.org/10.1038/nature08019>, 2009.
- Andrews, T., Betts, R. A., Booth, B. B. B., Jones, C. D., and Jones, G. S.: Effective radiative forcing from historical land use change, *Clim. Dynam.*, 48, 3489–3505, <https://doi.org/10.1007/s00382-016-3280-7>, 2017.
- Argo: Argo float data and metadata from Global Data Assembly Centre (Argo GDAC). SEANOE, <https://doi.org/10.17882/42182>, 2000.
- Armour, K.: Energy budget constraints on climate sensitivity in light of inconstant climate feedbacks, *Nat. Clim. Change*, 7, 331–335, <https://doi.org/10.1038/nclimate3278>, 2017.
- Berkeley Earth: Land + Ocean (1850 - Recent) Monthly Global Average Temperature, http://berkeleyearth.lbl.gov/auto/Global/Land_and_Ocean_complete.txt, accessed 17 October 2017, 2017.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- Boucher, O. and Reddy, M.: Climate trade-off between black carbon and carbon dioxide emissions, *Energ. Policy*, 36, 193–200, <https://doi.org/10.1016/j.enpol.2007.08.039>, 2008.
- Boucher, O., Friedlingstein, P., Collins, B., and Shine, K. P.: The indirect global warming potential and global temperature change potential due to methane oxidation, *Environ. Res. Lett.*, 4, 044 007, <https://doi.org/10.1088/1748-9326/4/4/044007>, 2009.
- 5 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.: Clouds and Aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 571–658, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 10 Carslaw, K., Lee, L., Reddington, C., Pringle, K., Rap, A., Forster, P., Mann, G., Spracklen, D., Woodhouse, M., Regayre, L., and Pierce, J.: Large contribution of natural aerosols to uncertainty in indirect forcing, *Nature*, 503, 67, <https://doi.org/10.1038/nature12674>, 2013.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A., and Wehner, M.: Long-term Climate Change: Projections, Commitments and Irreversibility, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 1029–1136, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.024>, 2013.
- 15 Cowtan, K. and Way, R. G.: Coverage bias in the HadCRUT4 temperature series and its impact on recent temperature trends, *Q. J. Roy. Meteor. Soc.*, 140, 1935–1944, <https://doi.org/10.1002/qj.2297>, 2014.
- 20 Daniel, J. and Velders, G.: A focus on information and options for policymakers, in: *Scientific Assessment of Ozone Depletion*, edited by Ennis, C. A., p. 516, World Meteorological Organization, Geneva, Switzerland, 2011.

- Daniel, J. S. and Solomon, S.: On the climate forcing of carbon monoxide, *J. Geophys. Res.-Atmos.*, 103, 13 249–13 260, <https://doi.org/10.1029/98JD00822>, 1998.
- Daniel, J. S., Solomon, S., Portmann, R. W., and Garcia, R. R.: Stratospheric ozone destruction: The importance of bromine relative to chlorine, *J. Geophys. Res.-Atmos.*, 104, 23 871–23 880, <https://doi.org/10.1029/1999JD900381>, 1999.
- 25 Doutriaux-Boucher, M., Webb, M. J., Gregory, J. M., and Boucher, O.: Carbon dioxide induced stomatal closure increases radiative forcing via a rapid reduction in low cloud, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2008GL036273>, 102703, 2009.
- Ehlert, D. and Zickfeld, K.: What determines the warming commitment after cessation of CO₂ emissions?, *Environ. Res. Lett.*, 12, 015 002, <https://doi.org/10.1088/1748-9326/aa564a>, 2017.
- 30 Etminan, M., Myhre, G., Highwood, E. J., and Shine, K. P.: Radiative forcing of carbon dioxide, methane, and nitrous oxide: A significant revision of the methane radiative forcing, *Geophys. Res. Lett.*, 43, 12,614–12,623, <https://doi.org/10.1002/2016GL071930>, 2016GL071930, 2016.
- Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., and Rummukainen, M.: Evaluation of Climate Models, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 741–866, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 35 Forest, C. E., Stone, P. H., and Sokolov, A. P.: Estimated PDFs of climate system properties including natural and anthropogenic forcings, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2005GL023977>, 101705, 2006.
- Forster, P., Richardson, T., Maycock, A., Smith, C., Samset, B., Myhre, G., Andrews, T., Pincus, R., and Schulz, M.: Recommendations for diagnosing effective radiative forcing from climate models from CMIP6, *J. Geophys. Res.*, 121, 12 460–12 475, <https://doi.org/10.1002/2016JD025320>, 2016.
- 5 Forster, P. M., Andrews, T., Good, P., Gregory, J. M., Jackson, L. S., and Zelinka, M.: Evaluating adjusted forcing and model spread for historical and future scenarios in the CMIP5 generation of climate models, *J. Geophys. Res.-Atmos.*, 118, 1139–1150, <https://doi.org/10.1002/jgrd.50174>, 2013.
- 10 Frame, D. J., Booth, B. B. B., Kettleborough, J. A., Stainforth, D. A., Gregory, J. M., Collins, M., and Allen, M. R.: Constraining climate forecasts: The role of prior assumptions, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2004GL022241>, 109702, 2005.
- Friedlingstein, P.: Carbon cycle feedbacks and future climate change, *Philos. T. R. Soc. A.*, 373, <https://doi.org/10.1098/rsta.2014.0421>, 2015.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate–Carbon Cycle Feedback Analysis: Results from the C4MIP Model Intercomparison, *J. Climate*, 19, 3337–3353, <https://doi.org/10.1175/JCLI3800.1>, 2006.
- 15 Fung, I. Y., Doney, S. C., Lindsay, K., and John, J.: Evolution of carbon sinks in a changing climate, *P. Natl. Acad. Sci. USA*, 102, 11 201–11 206, <https://doi.org/10.1073/pnas.0504949102>, 2005.
- 20 Gasser, T., Peters, G., Fuglestedt, J., Collins, W., Shindell, D., and Ciais, P.: Accounting for the climate–carbon feedback in emission metrics, *Earth System Dynamics*, 8, 235–253, <https://doi.org/10.5194/esd-8-235-2017>, <https://www.earth-syst-dynam.net/8/235/2017/>, 2017.

- Geoffroy, O., Saint-Martin, D., Olivié, D. J. L., Voldoire, A., Bellon, G., and Tytéca, S.: Transient Climate Response in a Two-Layer Energy-Balance Model. Part I: Analytical Solution and Parameter Calibration Using CMIP5 AOGCM Experiments, *J. Climate*, 26, 1841–1857, <https://doi.org/10.1175/JCLI-D-12-00195.1>, 2013.
- 25 Ghan, S. J., Smith, S. J., Wang, M., Zhang, K., Pringle, K., Carslaw, K., Pierce, J., Bauer, S., and Adams, P.: A simple model of global aerosol indirect effects, *J. Geophys. Res.-Atmos.*, 118, 6688–6707, <https://doi.org/10.1002/jgrd.50567>, 2013.
- Gillenwater, M.: Forgotten carbon: indirect CO₂ in greenhouse gas emission inventories, *Environ. Sci. Policy*, 11, 195–203, <https://doi.org/10.1016/j.envsci.2007.09.001>, 2008.
- Good, P., Gregory, J. M., and Lowe, J. A.: A step-response simple climate model to reconstruct and interpret AOGCM projections, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2010GL045208>, 101703, 2011.
- 30 Good, P., Gregory, J. M., Lowe, J. A., and Andrews, T.: Abrupt CO₂ experiments as tools for predicting and understanding CMIP5 representative concentration pathway projections, *Clim. Dynam.*, 40, 1041–1053, <https://doi.org/10.1007/s00382-012-1410-4>, 2013.
- Gregory, J. and Webb, M.: Tropospheric adjustment induces a cloud component in CO₂ forcing, *J. Climate*, 21, 58–71, <https://doi.org/10.1175/2007JCLI1834.1>, 2008.
- 35 Gregory, J. M. and Andrews, T.: Variation in climate sensitivity and feedback parameters during the historical period, *Geophys. Res. Lett.*, 43, 3911–3920, <https://doi.org/10.1002/2016GL068406>, 2016GL068406, 2016.
- Gregory, J. M., Andrews, T., and Good, P.: The inconstancy of the transient climate response parameter under increasing CO₂, *Philos. T. R. Soc. A*, 373, <https://doi.org/10.1098/rsta.2014.0417>, 2015.
- Gregory, J. M., Andrews, T., Good, P., Mauritsen, T., and Forster, P. M.: Small global-mean cooling due to volcanic radiative forcing, *Clim. Dynam.*, 47, 3979–3991, <https://doi.org/10.1007/s00382-016-3055-1>, 2016.
- Hansen, J., Ruedy, R., Sato, M., and Lo, K.: Global surface temperature change, *Rev. Geophys.*, 48, RG4004, <https://doi.org/10.1029/2010RG000345>, accessed 17 October 2017, 2010.
- 5 Haustein, K., Allen, M., Forster, P., Otto, F., Mitchell, D., Matthews, H., and Frame, D.: A real-time Global Warming Index, *Scientific Reports*, 7, 15417, 2017.
- Holmes, C. D., Prather, M. J., Søvde, O. A., and Myhre, G.: Future methane, hydroxyl, and their uncertainties: key climate and emission parameters for future predictions, *Atmos. Chem. Phys.*, 13, 285–302, <https://doi.org/10.5194/acp-13-285-2013>, 2013.
- International Energy Agency: IEA Statistics, [http://www.iea.org/statistics/statisticssearch/report/?country=WORLD&product=oil&year=](http://www.iea.org/statistics/statisticssearch/report/?country=WORLD&product=oil&year=2005)
- 10 2005, 2018.
- IPCC: Annex II: Climate System Scenario Tables, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)], edited by Prather, M., Flato, G., Friedlingstein, P., Jones, C., Lamarque, J.-F., Liao, H., and Rasch, P., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 15 Johnson, G., Lyman, J., and Loeb, N.: Improving estimates of Earth’s energy imbalance, *Nat. Clim. Change*, 6, 639–640, <https://doi.org/10.1038/nclimate3043>, 2016.
- Jones, A. D., Calvin, K. V., Collins, W. D., and Edmonds, J.: Accounting for radiative forcing from albedo change in future global land-use scenarios, *Climatic Change*, 131, 691–703, <https://doi.org/10.1007/s10584-015-1411-5>, 2015.
- Joos, F., Roth, R., Fuglestedt, J., Peters, G., Enting, I., von Bloh, W., Brovkin, V., Burke, E., Eby, M., Edwards, N., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J.: Carbon dioxide and

- climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmos. Chem. Phys.*, 13, 2793–2825, 2013.
- Kristiansen, N. I., Stohl, A., Oliv  , D. J. L., Croft, B., S  vde, O. A., Klein, H., Christoudias, T., Kunkel, D., Leadbetter, S. J., Lee, Y. H.,
25 Zhang, K., Tsigaridis, K., Bergman, T., Evangeliou, N., Wang, H., Ma, P.-L., Easter, R. C., Rasch, P. J., Liu, X., Pitari, G., Di Genova, G.,
Zhao, S. Y., Balkanski, Y., Bauer, S. E., Faluvegi, G. S., Kokkola, H., Martin, R. V., Pierce, J. R., Schulz, M., Shindell, D., Tost, H., and
Zhang, H.: Evaluation of observed and modelled aerosol lifetimes using radioactive tracers of opportunity and an ensemble of 19 global
models, *Atmos. Chem. Phys.*, 16, 3525–3561, <https://doi.org/10.5194/acp-16-3525-2016>, 2016.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G.,
30 Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V.,
Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and
aerosols: methodology and application, *Atmos. Chem. and Phys.*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, [https://www.
atmos-chem-phys.net/10/7017/2010/](https://www.atmos-chem-phys.net/10/7017/2010/), 2010.
- Le Qu  r  , C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P., Manning, A. C., Boden, T. A., Tans, P. P., Houghton,
35 R. A., Keeling, R. F., Alin, S., Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Currie, K.,
Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K.,
Kato, E., K  rtzinger, A., Landsch  tzer, P., Lef  vre, N., Lenton, A., Lienert, S., Lombardozzi, D., Melton, J. R., Metz, N., Millero, F.,
Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., O’Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D., Poulter,
B., R  denbeck, C., Salisbury, J., Schuster, U., Schwinger, J., S  f  rian, R., Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian,
H., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., and Zaehle, S.: Global Carbon
Budget 2016, *Earth Syst. Sci. Data*, 8, 605–649, <https://doi.org/10.5194/essd-8-605-2016>, 2016.
- 5 Lee, D. S., Fahey, D. W., Forster, P. M., Newton, P. J., Wit, R. C., Lim, L. L., Owen, B., and Sausen, R.: Aviation and global climate change
in the 21st century, *Atmos. Environ.*, 43, 3520–3537, <https://doi.org/10.1016/j.atmosenv.2009.04.024>, 2009.
- MacDougall, A. H., Zickfeld, K., Knutti, R., and Matthews, H. D.: Sensitivity of carbon budgets to permafrost carbon feedbacks and non-CO
2 forcings, *Environ. Res. Lett.*, 10, 125 003, <https://doi.org/10.1088/1748-9326/11/1/019501>, 2015.
- Matthes, K., Funke, B., Andersson, M. E., Barnard, L., Beer, J., Charbonneau, P., Clilverd, M. A., Dudok de Wit, T., Haberreiter, M., Hendry,
10 A., Jackman, C. H., Kretzschmar, M., Kruschke, T., Kunze, M., Langematz, U., Marsh, D. R., Maycock, A. C., Misios, S., Rodger, C. J.,
Scaife, A. A., Sepp  l  , A., Shangguan, M., Sinnhuber, M., Tourpali, K., Usoskin, I., van de Kamp, M., Verronen, P. T., and Versick, S.:
Solar forcing for CMIP6 (v3.2), *Geosci. Model Dev.*, 10, 2247–2302, <https://doi.org/10.5194/gmd-10-2247-2017>, 2017.
- Matthews, H. and Zickfeld, K.: Climate response to zeroed emissions of greenhouse gases and aerosols, *Nat. Clim. Change*, 2, 338–341,
<https://doi.org/10.1038/nclimate1424>, 2012.
- 15 Meinshausen, M., Meinshausen, N., Hare, W., Raper, S. C., Frieler, K., Knutti, R., Frame, D. J., and Allen, M. R.: Greenhouse-gas emission
targets for limiting global warming to 2 C, *Nature*, 458, 1158–1162, 2009.
- Meinshausen, M., Raper, S., and Wigley, T.: Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6
– Part 1: Model description and calibration, *Atmos. Chem. Phys.*, 11, 1417–1456, <https://doi.org/10.5194/acp-11-1417-2011>, 2011a.
- Meinshausen, M., Smith, S., Calvin, K., Daniel, J., Kainuma, M., Lamarque, J.-F., Matsumoto, K., Montzka, S., Raper, S., Riahi, K., Thom-
20 son, A., Velders, G., and van Vuuren, D.: The RCP Greenhouse Gas Concentrations and their Extension from 1765 to 2300, *Climatic
Change*, <https://doi.org/10.1007/s10584-011-0156-z>, 2011b.

- Millar, R., Allen, M., Rogelj, J., and Friedlingstein, P.: The cumulative carbon budget and its implications, *Oxford Rev. Econ. Pol.*, 32, 323–342, <https://doi.org/10.1093/oxrep/grw009>, 2016.
- 25 Millar, R. J., Otto, A., Forster, P. M., Lowe, J. A., Ingram, W. J., and Allen, M. R.: Model structure in observational constraints on transient climate response, *Climatic Change*, 131, 199–211, <https://doi.org/10.1007/s10584-015-1384-4>, 2015.
- Millar, R. J., Nicholls, Z. R., Friedlingstein, P., and Allen, M. R.: A modified impulse-response representation of the global near-surface air temperature and atmospheric concentration response to carbon dioxide emissions, *Atmos. Chem. Phys.*, 2017, 7213–7228, <https://doi.org/10.5194/acp-17-7213-2017>, 2017.
- 30 Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D.: Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 dataset, *J. Geophys. Res.*, 117, D08 101, <https://doi.org/10.1029/2011JD017187>, accessed 17 October 2017, 2012.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Vuuren, D. P. v., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, <https://doi.org/10.1038/nature08823>, 35 2010.
- Myhre, G. and Myhre, A.: Uncertainties in Radiative Forcing due to Surface Albedo Changes Caused by Land-Use Changes, *J. Climate*, 16, 1511–1524, <https://doi.org/10.1175/1520-0442-16.10.1511>, 2003.
- Myhre, G., Highwood, E. J., Shine, K. P., and Stordal, F.: New estimates of radiative forcing due to well mixed greenhouse gases, *Geophys. Res. Lett.*, 25, 2715–2718, <https://doi.org/10.1029/98GL01908>, 1998.
- 5 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmos. Chem. Phys.*, 13, 1853–1877, <https://doi.org/10.5194/acp-13-1853-2013>, <http://www.atmos-chem-phys.net/13/1853/2013/>, 2013a.
- 10 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 659–740, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013b.
- 15 Newman, P. A., Daniel, J. S., Waugh, D. W., and Nash, E. R.: A new formulation of equivalent effective stratospheric chlorine (EESC), *Atmos. Chem. Phys.*, 7, 4537–4552, <https://doi.org/10.5194/acp-7-4537-2007>, <https://www.atmos-chem-phys.net/7/4537/2007/>, 2007.
- Otto, A., Otto, F. E. L., Boucher, O., Church, J., Hegerl, G., Forster, P. M., Gillett, N. P., Gregory, J., Johnson, G. C., Knutti, R., Lewis, N., Lohmann, U., Marotzke, J., Myhre, G., Shindell, D., Stevens, B., and Allen, M. R.: Energy budget constraints on climate response, *Nat. Geosci.*, 6, 415–416, <https://doi.org/10.1038/ngeo1836>, 2013.
- 20 Pielke, R. A., Marland, G., Betts, R. A., Chase, T. N., Eastman, J. L., Niles, J. O., Niyogi, D. D. S., and Running, S. W.: The Influence of Land-Use Change and Landscape Dynamics on the Climate System: Relevance to Climate-Change Policy beyond the Radiative Effect of Greenhouse Gases, *Philos. T. R. Soc. A*, 360, 1705–1719, 2002.

- Prather, M. J., Holmes, C. D., and Hsu, J.: Reactive greenhouse gas scenarios: Systematic exploration of uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL051440>, 109803, 2012.
- Pueyo, S.: Solution to the paradox of climate sensitivity, *Climatic change*, 113, 163–179, <https://doi.org/10.1007/s10584-011-0328-x>, 2012.
- 25 RCP Database: RCP Database version 2.0 hosted at IIASA, <http://www.iiasa.ac.at/web-apps/tnt/RcpDb>. Accessed 27/07/2017, 2009.
- Richardson, M., Cowtan, K., Hawkins, E., and Stolpe, M. B.: Reconciled climate response estimates from climate models and the energy budget of Earth, *Nat. Clim. Change*, 2016.
- Rogelj, J., Meinshausen, M., and Knutti, R.: Global warming under old and new scenarios using IPCC climate sensitivity range estimates, *Nat. Clim. Change*, 2, 248–253, <https://doi.org/10.1038/nclimate1385>, 2012.
- 30 Santer, B. D., Thorne, P. W., Haimberger, L., Taylor, K. E., Wigley, T. M. L., Lanzante, J. R., Solomon, S., Free, M., Gleckler, P. J., Jones, P. D., Karl, T. R., Klein, S. A., Mears, C., Nychka, D., Schmidt, G. A., Sherwood, S. C., and Wentz, F. J.: Consistency of modelled and observed temperature trends in the tropical troposphere, *Int. J. Climatol.*, 28, 1703–1722, <https://doi.org/10.1002/joc.1756>, <http://dx.doi.org/10.1002/joc.1756>, 2008.
- Shindell, D. T., Faluvegi, G., Koch, D. M., Schmidt, G. A., Unger, N., and Bauer, S. E.: Improved Attribution of Climate Forcing to Emissions, *Science*, 326, 716–718, <https://doi.org/10.1126/science.1174760>, <http://science.sciencemag.org/content/326/5953/716>, 2009.
- 35 Skeie, R., Berntsen, T., Myhre, G., Tanaka, K., Kvalevåg, M., and Hoyle, C.: Anthropogenic radiative forcing time series from pre-industrial times until 2010, *Atmos. Chem. Phys.*, 11, 11 827–11 857, <https://doi.org/10.5194/acp-11-11827-2011>, <https://www.atmos-chem-phys.net/11/11827/2011/>, 2011.
- Stevens, B.: Rethinking the Lower Bound on Aerosol Radiative Forcing, *J. Climate*, 28, 4794–4819, <https://doi.org/10.1175/JCLI-D-14-00656.1>, 2015.
- Stevens, B., Sherwood, S. C., Bony, S., and Webb, M. J.: Prospects for narrowing bounds on Earth’s equilibrium climate sensitivity, *Earth’s Future*, 4, 512–522, <https://doi.org/10.1002/2016EF000376>, 2016EF000376, 2016.
- 5 Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsoren, S. B., Myhre, G., Berntsen, T. K., Folberth, G. A., Rumbold, S. T., Collins, W. J., MacKenzie, I. A., Doherty, R. M., Zeng, G., van Noije, T. P. C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D. A., Strode, S. A., Horowitz, L., Lee, Y. H., Szopa, S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K. W., Wild, O., and Archibald, A.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*,
- 10 13, 3063–3085, <https://doi.org/10.5194/acp-13-3063-2013>, 2013.
- 855 Tachiiri, K., Hajima, T., and Kawamiya, M.: Increase of uncertainty in transient climate response to cumulative carbon emissions after stabilization of atmospheric CO₂ concentration, *Environ. Res. Lett.*, 10, 125 018, 2015.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *B. Am. Meteorol. Soc.*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- Thompson, D. W. J., Barnes, E. A., Deser, C., Foust, W. E., and Phillips, A. S.: Quantifying the Role of Internal Climate Variability in Future
- 860 Climate Trends, *J. Climate*, 28, 6443–6456, <https://doi.org/10.1175/JCLI-D-14-00830.1>, 2015.
- Tokarska, K. B., Gillett, N. P., Arora, V. K., Lee, W. G., and Zickfeld, K.: The influence of non-CO₂ forcings on cumulative carbon emissions budgets, *Environmental Research Letters*, 13, 034 039, <https://doi.org/10.1088/1748-9326/aaafdd>, 2018.
- Toohey, M., Stevens, B., Schmidt, H., and Timmreck, C.: Easy Volcanic Aerosol (EVA v1.0): an idealized forcing generator for climate simulations, *Geosci. Model Dev.*, 9, 4049–4070, <https://doi.org/10.5194/gmd-9-4049-2016>, <https://www.geosci-model-dev.net/9/4049/>
- 865 2016/, 2016.

Twomey, S.: The influence of pollution on the shortwave albedo of clouds, *J. Atmos. Sci.*, 34, 1149–1152, 1977.

870 Walters, D., Baran, A., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J., Furtado, K., Hill, P., Lock, A., Manners, J., Morcrette, C., Mulcahy, J., Sanchez, C., Smith, C., Stratton, R., Tennant, W., Tomassini, L., Van Weverberg, K., Vosper, S., Willett, M., Browse, J., Bushell, A., Dalvi, M., Essery, R., Gedney, N., Hardiman, S., Johnson, B., Johnson, C., Jones, A., Mann, G., Milton, S., Rumbold, H., Sellar, A., Ujiie, M., Whittall, M., Williams, K., and Zerroukat, M.: The Met Office Unified Model Global Atmosphere 7.0/7.1 and JULES Global Land 7.0 configurations, *Geosci. Model Dev. Discuss.*, 2017, 1–78, <https://doi.org/10.5194/gmd-2017-291>, <https://www.geosci-model-dev-discuss.net/gmd-2017-291/>, 2017.

875 Zhang, H.-M., Huang, B., Lawrimore, J., Menne, M., and Smith, T.: NOAA Global Surface Temperature Dataset (NOAAGlobalTemp), Version 4.0. NOAA National Centers for Environmental Information, <https://doi.org/10.7289/V5FN144H>, accessed 17 October 2017, 2017.




Figure 1. Simplified overview of the FAIR v1.3v1.2 model.




Figure 2. Natural emissions of methane and nitrous oxide used in the FAIR model. Future emissions are fixed at their 2011 values. Also shown are the present-day best estimates of Prather et al. (2012).




Figure 3. (a) Joint distributions (FULL and NROY) of ECS and TCR. (b) Marginal distributions of ECS. (c) Marginal distributions of TCR. In (a), the FULL distribution is shown with open black contours and the NROY distribution with coloured squares. Probability density is plotted on a log scale. In (b) and (c), the FULL distribution is shown as a black curve and the NROY distribution with yellow histogram bars, both plotted on a linear scale. The NROY distributions contains only those ensemble members which agree with the C&W observed historical temperatures. CMIP5 models are depicted with red circles.

Historical and RCP WMGHG concentrations in FAIR and MAGICC6




Figure 4. Comparison of the historical and RCP greenhouse gas concentrations in FAIR (heavy solid lines) with 5-95% confidence intervals (shading) for CO₂. Dashed lines show the concentrations from MAGICC6 (Meinshausen et al., 2011b). For CFC12-eq, the RCP4.5 and RCP6.0 lines lie underneath the RCP8.5 line.




Figure 5. Comparison of the radiative forcing from RCP2.6, RCP4.5, RCP6.0 and RCP8.5 derived from 13 separate components (subplots a–m), along with the total radiative forcing (subplot n). ERF from FAIR (solid lines) with 5-95% confidence intervals (shading), RF from MAGICC6 (dashed lines, Meinshausen et al. (2011b)) and RF from AR5 Annex II for 1850–2011 (green solid lines, IPCC (2013)).




Figure 6. ERF from aerosols (blue), greenhouse gases (red) and total anthropogenic (black) for present-day (2017, based on RCP8.5) runs from FAIR constrained to observed temperature change (NROY; histograms) and from prior distributions (FULL; curves); compare Myhre et al. (2013b, Figure 8.16). Greenhouse gas forcing includes contributions from ozone and stratospheric water vapour from methane. Anthropogenic total is the sum of greenhouse gas, aerosol, contrails, BC on snow and land use change. The latter three distributions are not shown separately.




Figure 7. Relationship between (a) ECS and aerosol ERF; (b) TCR and aerosol ERF for the NROY ensemble. Aerosol ERF is shown for 2017 under the RCP8.5 scenario.




Figure 8. (a) Historical constrained modelled temperature and future probabilistic temperature scenarios from FAIR driven by emissions and forcing from the RCPs for 1850–2100 expressed as temperature change since 1861–1880. Also shown is observed temperature from C&W. (b) Comparison of 2081–2100 mean temperature from FAIR compared to the emission-driven ensemble from MAGICC6 (Rogelj et al., 2012).




Figure 9. Transient climate response to CO₂ emissions (TCRE) for FAIR based on RCP8.5 for all forcing (red) and for CO₂-only forcing (black).




Figure 10. Comparison of earth's energy imbalance N to observations from Argo and CERES, from Johnson et al. (2016).

Table 1. Emissions time series input used in FAIR, based on the RCP emissions datasets in Meinshausen et al. (2011b).

Index	Species	Unit	Remark
0	Year	Year	Important for running RCP scenarios as some treatments differ before 1850
1	CO ₂ fossil	Gt C yr ⁻¹	
2	CO ₂ land-use	Gt C yr ⁻¹	
3	CH ₄	Mt yr ⁻¹	Only anthropogenic emissions
4	N ₂ O	Mt N ₂ yr ⁻¹	Only anthropogenic emissions, expressed as N ₂ equivalent mass
5	SO _x	Mt S yr ⁻¹	Only anthropogenic emissions
6	CO	Mt yr ⁻¹	Only anthropogenic emissions
7	NMVOC	Mt yr ⁻¹	Only anthropogenic emissions
8	NO _x	Mt N yr ⁻¹	Only anthropogenic emissions
9	BC	Mt yr ⁻¹	Only anthropogenic emissions
10	OC	Mt yr ⁻¹	Only anthropogenic emissions
11	NH ₃	Mt yr ⁻¹	Only anthropogenic emissions
12	CF ₄	kt yr ⁻¹	Natural emissions should be included
13	C ₂ F ₆	kt yr ⁻¹	
14	C ₆ F ₁₄	kt yr ⁻¹	
15	HFC23	kt yr ⁻¹	
16	HFC32	kt yr ⁻¹	
17	HFC43-10	kt yr ⁻¹	
18	HFC125	kt yr ⁻¹	
19	HFC134a	kt yr ⁻¹	
20	HFC143a	kt yr ⁻¹	
21	HFC227ea	kt yr ⁻¹	
22	HFC245fa	kt yr ⁻¹	
23	SF ₆	kt yr ⁻¹	
24	CFC11	kt yr ⁻¹	
25	CFC12	kt yr ⁻¹	
26	CFC113	kt yr ⁻¹	
27	CFC114	kt yr ⁻¹	
28	CFC115	kt yr ⁻¹	
29	CCl ₄	kt yr ⁻¹	
30	Methyl chloroform	kt yr ⁻¹	
31	HCFC22	kt yr ⁻¹	
32	HCFC141b	kt yr ⁻¹	
33	HCFC142b	kt yr ⁻¹	
34	Halon 1211	kt yr ⁻¹	
35	Halon 1202	kt yr ⁻¹	
36	Halon 1301	kt yr ⁻¹	
37	Halon 2401	kt yr ⁻¹	
38	CH ₃ Br	kt yr ⁻¹	Natural emissions should be included
39	CH ₃ Cl	kt yr ⁻¹	Natural emissions should be included

Table 2. The set of greenhouse gases used in FAIR. With the exception of methane lifetime, radiative efficiencies and lifetimes are from AR5 (Myhre et al., 2013b, table 8.A.1). For ozone-depleting substances, the fractional release coefficients r_i (Daniel and Velders, 2011) and the number of chlorine and bromine atoms are also given, for calculation of equivalent effective stratospheric chlorine (eq. (14)).

Index	Gas	Molecular weight w_f (g mol ⁻¹)	Radiative efficiency η (W m ⁻² ppb ⁻¹)	Lifetime τ (yr)	r_i	n_{Cl}	n_{Br}
Major gases							
0	CO ₂	44.01	N/A	Variable			
1	CH ₄	16.04	N/A	9.3			
2	N ₂ O	44.01	N/A	121			
Kyoto Protocol gases							
3	CF ₄	88.00	0.09	50000			
4	C ₂ F ₆	138.01	0.25	10000			
5	C ₆ F ₁₄	338.04	0.44	3100			
6	HFC23	70.01	0.18	222			
7	HFC32	52.02	0.11	5.2			
8	HFC43-10	252.06	0.42	16.1			
9	HFC125	120.02	0.23	28.2			
10	HFC134a	102.03	0.16	13.4			
11	HFC143a	84.04	0.16	47.1			
12	HFC227ea	170.03	0.26	38.9			
13	HFC245fa	134.05	0.24	7.7			
14	SF ₆	146.06	0.57	3200			
Ozone depleting substances							
15	CFC11	137.37	0.26	45	0.47	3	0
16	CFC12	120.91	0.32	100	0.23	2	0
17	CFC113	187.38	0.30	85	0.29	3	0
18	CFC114	170.92	0.31	190	0.12	2	0
19	CFC115	154.47	0.20	1020	0.04	1	0
20	CCl ₄	153.81	0.17	26	0.56	4	0
21	Methyl chloroform	133.40	0.07	5	0.67	3	0
22	HCFC22	86.47	0.21	11.9	0.13	1	0
23	HCFC141b	116.94	0.16	9.2	0.34	2	0
24	HCFC142b	100.49	0.19	17.2	0.17	1	0
25	Halon 1211	165.36	0.29	16.0	0.62	1	1
26	Halon 1202	209.82	0.27	2.9	0.62	0	2
27	Halon 1301	148.91	0.30	65	0.28	0	1
28	Halon 2401	259.82	0.30	20	0.65	0	2
29	CH ₃ Br	94.94	0.004	0.8	0.60	0	1
30	CH ₃ Cl	50.49	0.01	1	0.44	1	0

Table 3. The 13 separate forcing groups considered in FAIR v1.3v1.2 in the calculation of effective radiative forcing. The ERF uncertainty represents the 5–95% range and is used in the generation of the large ensemble (section 3). ERF uncertainties from Myhre et al. (2013b) are used except for CH₄ where we use the Myhre et al. (2013b) estimate inflated by the additional uncertainty in the new methane forcing relationship in Etminan et al. (2016), which also affects the uncertainty in stratospheric water vapour oxidation from methane.

Index	Forcing agent	Depends on	ERF uncertainty
0	CO ₂	CO ₂ emissions; CH ₄ fossil fraction; cumulative C emissions; total temperature change	±20%
1	CH ₄	CH ₄ emissions	±28%
2	N ₂ O	N ₂ O emissions	±20%
3	Other greenhouse gases	Emissions of other greenhouse gases	±20%
4	Tropospheric ozone	Emissions of CH ₄ and short-lived climate forcers	±50%
5	Stratospheric ozone	Concentrations of ozone depleting substances (subset of minor greenhouse gases)	±200%
6	Stratospheric water vapour	CH ₄ ERF	±72%
7	Contrails	Aviation NOx fraction; total NOx emitted	−66 to +191%
8	Aerosols	Emissions of short-lived climate forcers	−89 to +111%
9	Black carbon on snow	Emissions of black carbon	−56 to +128%
10	Land use change	Cumulative emissions of land-use related CO ₂	±167%
11	Volcanic	Externally supplied forcing from volcanoes	±50%
12	Solar	Externally supplied forcing from solar variability	±0.05 W m ^{−2}

Table 4. Contribution to tropospheric ozone ERF from each precursor. Pre-industrial emissions from Skeie et al. (2011), pre-industrial CH₄ from Meinshausen et al. (2011b), 1850 and 2000 emissions from Lamarque et al. (2010) and 2000 minus 1850 ERF from Stevenson et al. (2013).

Species	ERF in 2000 (W m ⁻²)	Ozone forcing efficiency β_i (eq. (11))		Pre-industrial value
		Pre-1850	Post-1850	
CH ₄	0.178	1.73×10^{-4}	$1.73 \times 10^{-4} \text{ W m}^{-2} \text{ ppb}^{-1}$	722 ppb
CO	0.076	4.76×10^{-5}	$8.51 \times 10^{-5} \text{ W m}^{-2} (\text{Mt yr}^{-1})^{-1}$	170 Mt CO yr ⁻¹
NMVOC	0.044	1.88×10^{-4}	$2.25 \times 10^{-4} \text{ W m}^{-2} (\text{Mt yr}^{-1})^{-1}$	5 Mt NMVOC yr ⁻¹
NO _x	0.125	5.72×10^{-4}	$9.08 \times 10^{-4} \text{ W m}^{-2} (\text{Mt yr}^{-1})^{-1}$	2 Mt N yr ⁻¹

Table 5. Contribution to ERFari from each aerosol precursor species and contribution to 2011 ERFari.

Species	ERFari in 2011 (W m^{-2})	Radiative efficiency ($10^{-3} \text{ W m}^{-2} (\text{Mt yr}^{-1})^{-1}$)
SOx	-0.34	-6.22 (Mt S)
BC	+0.13	+16.0
OC	-0.05	-1.45
NH ₃	-0.066	-1.56
NOx	-0.044	-1.17 (Mt N)
SOA	-0.08	-0.38

Table 6. Median and 5–95% credible intervals for effective radiative forcing from greenhouse gases, aerosols and anthropogenic total from the FULL and NROY FAIR ensembles in 2017. Anthropogenic total contains contributions from contrails, BC on snow and land use change and therefore is not equal to the sum of greenhouse gas and aerosol forcing. Compare fig. 6.

Forcing type	Effective radiative forcing (W m^{-2})	
	Before temperature constraint (FULL)	After temperature constraint (NROY)
Greenhouse gases	3.69 (3.18 to 4.21)	3.68 (2.90 to 4.61)
Aerosols	−0.91 (−1.63 to −0.37)	−0.96 (−1.65 to −0.27)
Anthropogenic total	2.73 (1.85 to 3.50)	2.63 (1.74 to 3.73)

Table 7. Sensitivity in the ECS, TCR, TCRC to variations in the underlying assumptions in the FAIR large ensemble. For the sensitivity experiments the section number in the manuscript describing the change is given. The Accepted column details the proportion of the 100,000 member FULL ensemble that satisfied the specified temperature constraint.

Variation (section)	Accepted	ECS (K)			TCR (K)			TCRE (K (Eg C) ⁻¹)		
		5%	50%	95%	5%	50%	95%	5%	50%	95%
C&W temperature constraint (NROY)	26.1%	2.01	2.86	4.22	1.05	1.53	2.41	0.96	1.40	2.23
C&W with alternative ECS/TCR prior (5.1)	21.6%	1.57	2.62	4.76	0.99	1.55	2.68	0.89	1.35	2.55
C&W with $F_{2\times} = 3.44 \text{ W m}^{-2}$ (5.2)	24.9%	1.99	2.84	4.22	1.03	1.52	2.41	0.93	1.36	2.15
HadCRUT4 temperature constraint (5.3)	26.4%	1.97	2.82	4.19	1.02	1.51	2.38	0.93	1.37	2.21
GISTEMP temperature constraint (5.3)	33.2%	2.03	2.89	4.26	1.06	1.56	2.44	0.97	1.42	2.27
Berkeley Earth temperature constraint (5.3)	23.4%	2.11	2.97	4.32	1.13	1.61	2.48	1.02	1.47	2.30
NOAA temperature constraint (5.3)	35.2%	1.98	2.85	4.22	1.03	1.52	2.41	0.93	1.39	2.23
No temperature constraint (FULL)	100%	2.00	3.11	4.86	1.01	1.73	2.96	0.91	1.58	2.78

Table 8. Sensitivity in the effective radiative forcing to variations in the underlying assumptions in the FAIR large ensemble.

Variation	Effective radiative forcing in 2100, W m^{-2}											
	RCP 2.6			RCP 4.5			RCP 6.0			RCP 8.5		
	5%	50%	95%	5%	50%	95%	5%	50%	95%	5%	50%	95%
C&W temperature constraint (NROY)	1.79	2.62	3.64	3.30	4.62	6.22	4.07	5.84	8.00	6.84	9.34	12.40
C&W with alternative ECS/TCR prior	1.76	2.61	3.68	3.27	4.61	6.25	4.04	5.84	8.04	6.81	9.34	12.42
C&W with $F_{2\times} = 3.44 \text{ W m}^{-2}$	1.67	2.45	3.40	3.12	4.34	5.85	3.83	5.47	7.48	6.51	8.83	11.69
HadCRUT4 temperature constraint	1.76	2.59	3.59	3.27	4.57	6.17	4.03	5.78	7.93	6.78	9.26	12.28
GISTEMP temperature constraint	1.80	2.64	3.67	3.31	4.63	6.27	4.09	5.87	8.05	6.86	9.37	12.45
Berkeley Earth temperature constraint	1.85	2.70	3.73	3.36	4.71	6.36	4.16	5.97	8.17	6.94	9.50	12.58
NOAA temperature constraint	1.77	2.60	3.62	3.27	4.59	6.21	4.04	5.80	7.98	6.79	9.29	12.36
No temperature constraint (FULL)	1.65	2.67	3.90	3.17	4.67	6.55	3.91	5.91	8.40	6.66	9.42	12.85

Table 9. Sensitivity in the 2100 temperature change in RCP scenarios to variations in the underlying assumptions in the FAIR large ensemble.

Variation	Temperature change in 2100 from 1861–1880 mean, K											
	RCP 2.6			RCP 4.5			RCP 6.0			RCP 8.5		
	5%	50%	95%	5%	50%	95%	5%	50%	95%	5%	50%	95%
C&W temperature constraint (NROY)	1.10	1.48	2.09	1.73	2.37	3.45	2.08	2.84	4.14	3.11	4.33	6.50
C&W with alternative ECS/TCR prior	0.95	1.42	2.26	1.52	2.31	3.83	1.86	2.80	4.59	2.82	4.29	7.24
C&W with $F_{2\times} = 3.44 \text{ W m}^{-2}$	1.10	1.49	2.11	1.73	2.39	3.51	2.07	2.85	4.19	3.13	4.38	6.63
HadCRUT4 temperature constraint	1.06	1.44	2.05	1.67	2.31	3.39	2.00	2.77	4.07	3.01	4.22	6.40
GISTEMP temperature constraint	1.11	1.51	2.14	1.74	2.42	3.53	2.09	2.90	4.25	3.14	4.42	6.62
Berkeley Earth temperature constraint	1.19	1.59	2.20	1.88	2.52	3.62	2.27	3.03	4.36	3.39	4.60	6.79
NOAA temperature constraint	1.06	1.47	2.09	1.67	2.35	3.46	2.00	2.81	4.15	3.01	4.29	6.51
No temperature constraint (FULL)	0.92	1.66	2.97	1.52	2.66	4.73	1.81	3.20	5.75	2.80	4.90	8.72