

A representation of the collisional ice break-up process in the two-moment microphysics scheme LIMA v1.0 of Meso-NH

Thomas Hoarau¹, Jean-Pierre Pinty², and Christelle Barthe¹

¹Laboratoire de l'Atmosphère et Cyclones, UMR 8105, CNRS/Météo-France/Université de La Réunion, St Denis, La Réunion, France

²Laboratoire d'Aérodynamique, University of Toulouse/CNRS/UPS, 14 avenue Edouard Belin, F-31400 Toulouse, France

Correspondence to: jean-pierre.pinty@aero.obs-mip.fr

Abstract. The paper describes a switchable parameterization of CIBU (Collisional Ice Break-Up), an ice multiplication process that fits in with the two-moment microphysical scheme LIMA (Liquid Ice Multiple Aerosols). The LIMA scheme with three ice types (pristine cloud ice crystals, snow-aggregates and graupel-hail) was developed in the cloud-resolving mesoscale model Meso-NH. Here the CIBU process assumes that collisional break-up is mostly efficient for the small snow-aggregate class of particles with a fragile structure when hit by large dense graupel particles. The increase of cloud ice number concentration depends on a prescribed number (or a random number) of fragments being produced per collision. This point is discussed and analytical expressions of the newly contributing CIBU terms in LIMA are given.

5 The scheme is run in the cloud resolving mesoscale model Meso-NH to simulate a first case of a three-dimensional deep convective event with a heavy production of graupel. The consequence of dramatically changing the number of fragments produced per collision is investigated by examining the rainfall rates and the changes in small ice concentrations and mass mixing ratios. Many budgets of the ice phase are shown and the sensitivity of CIBU to the **initial IFN concentration** is explored.

15 Then the scheme is tested for another deep convective case **but** with a varying CAPE (Convective Available Potential Energy). The results confirm the strong impact of CIBU with up to a one thousand fold increase in small ice concentrations, a reduction of the rainfall or precipitating area and an invigoration of the convection with higher cloud tops.

Finally it is concluded that the efficiency of the ice crystal fragmentation needs to be tuned carefully. The proposed parameterization of CIBU is easy to implement in any two-moment microphysics

schemes. It could be used in this form to simulate deep tropical cloud systems where preferential occurrence of anomalously high concentrations of small ice crystals is suspected.

1 Introduction

In a series of papers, Yano and Phillips (2011, 2016) and Yano et al. (2016) brought the Collisional
25 Ice Break-Up (hereafter CIBU) process to the fore again as a possible secondary ice production
mechanism in clouds. Using an analytical model, they showed that the CIBU could lead to an ex-
plosive growth of small ice crystal concentrations. Afterwards Sullivan et al. (2017) tried to include
CIBU in a parcel model of six species, assumed to be monodispersed here, in an attempt to make
this finding specific. However intriguingly and in contrast to the Hallett-Mossop (hereafter H-M) ice
30 multiplication mechanism¹ (Hallett and Mossop, 1974), the vast majority of microphysics schemes
do not include the CIBU process. Yet, the CIBU process is very likely to be active in the case of in-
homogeneous cloud regions where ice crystals of different sizes and types are locally mixed (Hobbs
and Rangno, 1985; Rangno and Hobbs, 2001). For instance, collisions between large dense graupel
growing by riming, and plane vapour-grown dendrites or irregular weakly rimed assemblages
35 are the most conceivable scenario for generating multiple ice debris as envisioned by Hobbs and
Farber (1972) and by Griggs and Choulaton (1986). So a legitimate quest for a two-moment mixed-
phase microphysics scheme, where number concentrations and mixing ratios of the ice crystals are
predicted, is to find ways to include an ice-ice break-up **effect** and to characterize its importance, rel-
atively to other ice generating processes such as ice heterogeneous nucleation. Our wish to introduce
40 CIBU in a microphysics scheme was initially motivated essentially by the detection of unexplained
high ice water contents that sometimes largely exceed the concentration of ice nucleating particles
(Leroy et al., 2015; Field et al., 2017; Ladino et al., 2017).

As recalled by Yano and Phillips (2011), the first few laboratory experiments dedicated to the
study of ice collisions were conducted in the 1970s following investigations concerning the promis-
45 ing H-M process. The pioneering work of Vardiman (1978) was a rare experimental reference show-
ing evidence for the mechanical fracturing of natural ice crystals. An interesting outcome of the
study was the finding that the number of fragments was dependent on the shape of the initial collid-
ing crystal and on the momentum change following the collision. According to a concluding remark
by Vardiman (1978), this 'secondary' production of ice could lead to concentrations as high as 100
50 to 1000 times the natural concentrations of ice crystals in clouds expected from heterogeneous nu-
cleation on ice freezing nuclei. Another laboratory study by Takahashi et al. (1995) also revealed a
huge production of splinters after collisions between rimed and deposition-grown graupels. However
the experimental set-up used there was more appropriate to very big, artificially grown crystals and
to large impact velocities because as many as 400 fragments could be obtained.

¹H-M is based on the explosive riming of "big" droplets on graupel particles in a narrow range of temperature

55 For clarity, this study does not focus on cloud conditions leading to an explosive ice multiplication
by mechanical break-up in ice-ice collisions (Yano and Phillips, 2011). Neither does it attempt to
reformulate this process on the basis of collisional kinetic energy with many empirical parameters as
proposed by Phillips et al. (2017), or earlier by Hobbs and Farber (1972) with the breaking energy,
mostly applicable to "bin" microphysics schemes. Here, the goal is rather to implement an empirical
60 but realistic parameterization of CIBU in the LIMA (an acronym for Liquid, Ice, Multiple Aerosols)
scheme (Vié et al., 2016) to cooperate with other microphysical processes (heterogeneous ice nucle-
ation, droplet freezing, H-M process, etc.) to determine the concentration of small ice crystals. Our
idealization of CIBU is the formation of cloud ice crystals as the result of collisions between big
graupel particles and small aggregates after which the graupel particles **lose the mass of the** aggre-
65 gates. The parameterization of CIBU relies on the laboratory observations by Vardiman (1978) to
set limits on the number of fragments per collision. However, the large uncertainties attached to this
parameter encourage us to run exploratory experiments with several fixed values and also to model
the number of fragments by means of a random process with a span of two decades.

The LIMA scheme was inserted in Meso-NH (Lafore et al., 1998). Several sensitivity experiments
70 were performed to evaluate the importance of the CIBU process and the impact of the tuning, i.e. the
number of fragments produced per collision. The efficiency of CIBU in dramatically increasing the
concentration of small ice crystals can be scaled by the ice number concentration from nucleation.
The case of a three-dimensional continental deep convective storm, the well-known STERAO case
simulated by Skamarock et al. (2000), provided a framework for several adjustments of the number
75 of ice fragments. A series of experiments was then performed for the same case to see how much
the CIBU process altered the precipitation and the persistence of convective plumes. The question
of the number of ice nuclei necessary to initiate CIBU (Field et al., 2017; Sullivan et al., 2018)
was also tackled. A second case of deep convective cloud (Weisman and Klemp, 1984) was run to
confirm the impact of CIBU in a series of **varying** CAPE environments. The simulations showed
80 the invigoration of the convection when the CIBU efficiency was strong, so leading to larger cloud
covers and an increase of the mean cloud top height. Finally, a conclusion is drawn on the importance
of calibrating the parameterization of CIBU and the need to systematically include CIBU and other
ice multiplication processes in microphysics bulk schemes.

2 Introduction of CIBU into the LIMA scheme

85 2.1 General considerations

In contrast to the work of Yano and Phillips (2011), where large and small graupel particles fuelled
the CIBU process, we consider collisions involving two types of precipitating ice here: small ice
particles growing by deposition and aggregation (aggregates including dendritic pristine ice crystals
with a size larger than $\sim 150 \mu\text{m}$ and big, massive graupel particles growing by riming. Collisions

90 between graupel particles of different sizes are not considered because, according to Griggs and
 Choulaton (1986), the fragmentation of rime is very unlikely to occur in natural clouds. **For the
 sake of simplicity and because the impact velocity of the graupel particles should be well above 1
 m s⁻¹ to remain in the break-up regime of the aggregates, the particle sizes are selected to enable a
 substantial occurrence of CIBU.**

95 A general form of the equation describing the CIBU process can be written

$$\frac{\partial n_i}{\partial t} = \alpha n_s n_g \quad (1)$$

where n is the particle size distribution of the cloud ice (subscript "i"), the snow-aggregates ("s") and
 the graupel particles ("g"). α is the snow-aggregate-graupel collision kernel times \mathcal{N}_{sg} , the number
 of ice fragments produced by collision. An expression for α , which does not include thermal and
 100 mechanical energy effects, is

$$\alpha = \mathcal{N}_{sg} V_{sg} \frac{\pi}{4} D_g^2 \quad (2)$$

where V_{sg} is the impact velocity of a graupel particle of size D_g at the surface of the aggregate.

In Eq. 2, it is assumed that the size of the aggregate is negligible compared to D_g . V_{sg} is ex-
 pressed as the difference of fall speed between the colliding graupel and the aggregate target so
 105 $V_{sg} = (\rho_{00}/\rho_a)^{0.4} \times (c_g D_g^{d_g} - c_s D_s^{d_s})$ using the generic formula of the particle fall speeds $V_x =$
 $(\rho_{00}/\rho_a)^{0.4} \times c_x D_x^{d_x}$ with the air density correction of Foote and du Toit (1969) due to the drag
 force exerted by the particles during their fall. ρ_{00} is the reference air density ρ_a at normal pressure.

As introduced above and suggested in Yano and Phillips (2011), **the impact velocity V_{sg} should
 be such that a minimum value is guaranteed to enable CIBU.** An easy way to do this is to restrict the
 110 size of the aggregates to the range [$D_{smin}=0.2$ mm, $D_{smax}=1$ mm] and to introduce a minimum size
 of $D_{gmin}=2$ mm for the graupel particles. The reasons for these choices are discussed below. The
 lower bound value, D_{smin} , is an estimate **that results in the collision efficiency with a graupel particle
 approaching unity.** For $D_s < D_{smin}$, big crystals or aggregates stay outside the path of capture which
 explains the observation of bimodal ice spectra. Field (2000) reported minimum values of 150-200
 115 μ m for D_{trough} , a critical size separating cloud ice and aggregate regimes. The D_{smin} value is also
 consistent with an upper bound of the cloud ice crystal size distribution that results from the critical
 diameter of 125 μ m to convert cloud ice to snow by deposition (see Harrington et al. (1995) for
 the original and analytical developments and Vié et al. (2016) for the implementation in LIMA).
 The choice of round numbers for D_{smax} and D_{gmin} is above all dictated by the empirical rule that
 120 $V_{sg} > 1$ m s⁻¹. With the setup in LIMA which is [c_x, d_x] = [5.1, 0.27] for "x = s" and [124, 0.66]
 for "x = g" in MKS units, we obtain $V_{sg} > 1.26$ m s⁻¹ at ground level.

The number of fragments, \mathcal{N}_{sg} , is the critical parameter for ice multiplication. From scaling argu-
 ments Yano and Phillips (2011) recommended taking $\mathcal{N}_{sg} = 50$. Recently Yano and Phillips (2016)
 introduced a notion of random fluctuations into the production of fragments leading to a stochastic

125 equation of the ice crystal concentration due to the realization of a noise process for α (Eq. 2). The
parameterization of \mathcal{N}_{sg} as a function of collisional kinetic energy (Phillips et al., 2017) enables a
differentiated treatment of the fragmentation of a variety of ice crystals. All these results stem from
Fig. 6 in Vardiman (1978) which suggests that \mathcal{N}_{sg} is a function of momentum change, ΔM_g , af-
ter the collision. As $\Delta M_g \sim 0.1 \text{ g cms}^{-1}$ for $D_g=2 \text{ mm}$, the corresponding \mathcal{N}_{sg} lies between 10
130 (for collision with plane dendrites) and 40 (for rimed spatial crystals). These values are consistent
with those found by Yano and Phillips (2011) for rimed assemblages. In conclusion, it is tempting
to run both deterministic and stochastic simulations to test the sensitivity to \mathcal{N}_{sg} ~~but~~ in the range
suggested by laboratory experiments. In the following \mathcal{N}_{sg} was set successively to 0.1 (weak ef-
fect) **or alternatively** one fragment per ten collisions, to 1.0 (moderate effect) and to 10.0 or even 50
135 (strong effect) fragments per collision. Additional experiments were performed by first generating a
random variable X uniformly distributed over $[0, 1.0]$ and then by applying an empirical formula,
 $\mathcal{N}_{sg} = 10^{2.0 \times X - 1.0}$, to generate numbers over two decades $[0.1, 10.0]$ of \mathcal{N}_{sg} . The randomization
of \mathcal{N}_{sg} reflects the fact that the number of fragments depends on the positioning of the impact, on
the tip or on the body of the fragile particle, and also on the energy lost by the possible rotation of
140 the residual particle.

2.2 Characteristics of the LIMA microphysics scheme

The LIMA microphysics scheme (Vié et al., 2016) includes a representation of the aerosols as a mix-
ture of Cloud Condensation Nuclei (CCN) and Ice Freezing Nuclei (IFN) with an accurate budget
equation (transport, activation or nucleation, scavenging by rain) for each aerosol type. The CCN
145 are selectively activated to produce the cloud droplets which grow by condensation and coalescence
to produce the rain drops (Cohard and Pinty, 2000). The ice phase is more complex as we consider
nucleation by deposition on insoluble IFN (black carbon and dust) and nucleation by immersion
(glaciation of tagged droplets because they are formed on partially soluble CCN, containing an in-
soluble core). Homogeneous freezing of the droplets is possible when the temperature drops below
150 -35° C . The Hallett-Mossop mechanism generates ice crystals during the riming of the graupel and
the snow-aggregates. The H-M efficiency depends strongly on the temperature and on the size distri-
bution of the droplets (Beheng, 1987). The initiation of the snow-aggregates category is the result of
depositional growth of large pristine crystals beyond a critical size (Harrington et al., 1995). Aggre-
gation and riming are computed explicitly. Heavily rimed particles (graupel) can experience a dry or
155 wet growth mode. The freezing of raindrops by contact with small ice crystals leads to frozen drops
which are merged with the graupel category. The melting of snow-aggregates leads to graupel and
shed raindrops while the graupel particles melt directly into rain. **The sedimentation of all particle
types is considered.** The snow-aggregates and graupel particles are characterized by their mixing ra-
tios only. The LIMA scheme assumes a strict saturation of the water vapour over the cloud droplets

160 while the small ice crystals are subject to super- or under-saturated conditions (no instantaneous equilibrium).

2.3 Representation of CIBU in the LIMA scheme

In a 2-moment bulk scheme, the zeroth order (total number concentration) and "bth" order (mixing ratio)² moments of the size distributions are computed. From Eqs.1 and 2 with expansion, the CIBU
165 tendency of the number concentration of the cloud ice N_i (here in # kg⁻¹) can be written as:

$$\frac{\partial N_i}{\partial t} = \frac{N_{sg}}{\rho_{dref}} \frac{\pi}{4} \left(\frac{\rho_{00}}{\rho_{dref}} \right)^{0.4} \int_{D_{smin}}^{D_{smax}} n_s(D_s) \left\{ \int_{D_{gmin}}^{\infty} D_g^2 (c_g D_g^{d_g} - c_s D_s^{d_s}) n_g(D_g) dD_g \right\} dD_s \quad (3)$$

where $\rho_{dref}(z)$ is a reference density profile of dry air (Meso-NH is anelastic) and with a further approximation $\rho_a = \rho_{dref}$.

In LIMA, the size distributions follow a generalized gamma law:

$$170 \quad n(D)dD = N \frac{\alpha}{\Gamma(\nu)} \lambda^{\alpha\nu} D^{\alpha\nu-1} e^{-(\lambda D)^\alpha} dD$$

where α and ν are fixed shape parameters, N is the total number concentration and λ is the slope parameter. With the definition of the moments $M_x^{INC}(p; X)$ of the incomplete gamma law given in Appendix A, integration of Eq. 3 leads to:

$$175 \quad \frac{\partial N_i}{\partial t} = \frac{N_{sg}}{\rho_{dref}} \frac{\pi}{4} \left(\frac{\rho_{00}}{\rho_{dref}} \right)^{0.4} N_s N_g \times \left\{ c_g \left(M_s^{INC}(0; D_{smin}) - M_s^{INC}(0; D_{smax}) \right) \left(M_g(2 + d_g) - M_g^{INC}(2 + d_g; D_{gmin}) \right) - c_s \left(M_s^{INC}(d_s; D_{smin}) - M_s^{INC}(d_s; D_{smax}) \right) \left(M_g(2) - M_g^{INC}(2; D_{gmin}) \right) \right\} \quad (4)$$

with $N_s = C_s \lambda_s^{x_s}$ and $N_g = C_g \lambda_g^{x_g}$. The set of **flexible** parameters used in LIMA is $C_s = 5$, $C_g =$
180 $5. \times 10^5$, $x_s = 1$, $x_g = -0.5$. These values were chosen to generalize the classical Marshall-Palmer law, $n(D) = N_0 \exp(-\lambda D)$, a degenerate form of the generalized gamma law when $\alpha = \nu = 1$, leading to a total concentration $N = N_0 \lambda^{-1}$ with a fixed intercept parameter N_0 .

Concerning the mixing ratios, the mass of the newly formed cloud ice fragments is simply taken as the product of the mean mass of the pristine ice crystals by the N_i tendency (Eq. 3). The mass loss
185 of the aggregates after collisional break-up is equal to the mass of the ice fragments. The mass of the graupel is unchanged. The mass transfer from aggregates to small ice crystals is constrained by the mass of individual aggregates that may break up completely. This limiting mixing ratio tendency is

²Ice mixing ratios are computed by integration over the size distribution of the mass of individual particles given by a mass-size relationship $m(D) = aD^b$, a power law with a non-integer exponent "b"

given by:

$$\frac{\partial r_i}{\partial t} = -\frac{\partial r_s}{\partial t} = \frac{a_s}{\rho_{dref}} \frac{\pi}{4} \left(\frac{\rho_{00}}{\rho_{dref}} \right)^{0.4} \int_{D_{smin}}^{D_{smax}} D_s^{b_s} n_s(D_s) \left\{ \int_{D_{gmin}}^{\infty} D_g^2 (c_g D_g^{d_g} - c_s D_s^{d_s}) n_g(D_g) dD_g \right\} dD_s. \quad (5)$$

190 In the above expression the mass of an aggregate of size D_s is given by $a_s D_s^{b_s}$ with $a_s=0.02$ and $b_s=1.9$ in LIMA, meaning that aggregates are quasi two-dimensional particles. After integration the mixing ratio tendencies are expressed as:

$$195 \frac{\partial r_i}{\partial t} = -\frac{\partial r_s}{\partial t} = \frac{a_s}{\rho_{dref}} \frac{\pi}{4} \left(\frac{\rho_{00}}{\rho_{dref}} \right)^{0.4} N_s N_g \times \left\{ c_g \left(M_s^{INC}(b_s; D_{smin}) - M_s^{INC}(b_s; D_{smax}) \right) \left(M_g(2 + d_g) - M_g^{INC}(2 + d_g; D_{gmin}) \right) - c_s \left(M_s^{INC}(b_s + d_s; D_{smin}) - M_s^{INC}(b_s + d_s; D_{smax}) \right) \left(M_g(2) - M_g^{INC}(2; D_{gmin}) \right) \right\} \quad (6)$$

This expression is independent of the number of fragments \mathcal{N}_{sg} .

3 Simulation of a 3-dimensional deep convective case

200 The test case is illustrated by idealized numerical simulations of the 10 July 1996 thunderstorm in the Stratospheric-Tropospheric Experiment: Radiation, Aerosols, and Ozone (STRAO) experiment (Dye et al., 2000). This case is characterized by a multicellular storm which becomes supercellular after 2 hours. The simulations were initialized with the sounding of north eastern Colorado given in Skamarock et al. (2000) and convection was triggered by three 3K-buoyant bubbles aligned along
 205 the main diagonal of the X,Y **plan** in the wind axis. Meso-NH was run for 5 hours over a domain of **320 × 320** with 1 km-horizontal grid spacing. There were 50 unevenly spaced vertical levels up to 23 km height. With the exception of the wind components advected with a fourth-order scheme, all the fields including microphysics, were transported by an accurate, conservative, positive-definite PPM (Piecewise Parabolic Method) scheme (Colella and Woodward, 1984). There were no surface fluxes
 210 **but** the 3D turbulence scheme of Meso-NH was activated. Open lateral boundary conditions were imposed. The upper level damping layer of the upward moving gravity waves started above 12500 m.

The aerosols were initialized as for the simulated squall-line case in Vié et al. (2016). A summary is given in Table 1 for the soluble Cloud Condensation Nuclei (CCN) and for the insoluble Ice
 215 Freezing Nuclei (IFN). Homogeneous vertical profiles are assumed for the aerosols. Although the LIMA scheme incorporates size distribution parameters and differentiates between the chemical compositions of the CCN and the IFN, the characteristics of the five aerosol modes are standard for

the simulations shown here, except for the sensitivity of CIBU to the initial concentration of the IFN which is explored in Section 3.5.

220 3.1 Impact on precipitation

Figure 1 shows the accumulated precipitation at ground level after 4 hours of simulation for the four experiments corresponding to $\mathcal{N}_{sg}=0.0, 0.1, 1.0$ and 10.0 . The highest amount of rainfall is obtained when the CIBU process is ignored ($\mathcal{N}_{sg}=0.0$) in Fig. 1a. Then, stepping up the CIBU efficiency by decades from $\mathcal{N}_{sg}=0.1$, Fig. 1b-d clearly shows a steady reduction of precipitation and a fine scale modification of the precipitation pattern. Furthermore, Fig. 1d reveals that the spread of the precipitation field, caused by the motion of the multicellular storm, is significantly reduced when $\mathcal{N}_{sg}=10.0$. The results of Fig. 1 suggest empirically that a plausible range for \mathcal{N}_{sg} is between 0.1 and 10.0 fragments per collision. A value lower than 0.1 leads to a negligible effect of CIBU in the simulation, while taking $\mathcal{N}_{sg}>10.0$ has an excessive impact on the storm rainfall (the " $\mathcal{N}_{sg}=50.0$ " case is not shown). In complement, Fig 2 shows the results of a simulation, called "RANDOM" hereafter, where \mathcal{N}_{sg} is generated by a random process as explained above but providing $0.1 < \mathcal{N}_{sg} < 10.0$. The perturbation caused by CIBU is noticeable in this case too but it remains weak for the precipitation field. These first 3D numerical experiments show that inclusion of CIBU can strongly modify surface precipitation when $\mathcal{N}_{sg} > 10.0$ fragments per aggregate-graupel collision. Taking $0.1 < \mathcal{N}_{sg} < 10.0$ and also considering \mathcal{N}_{sg} as the realization of a random process seems to be a more satisfactory approach. Admittedly, the limit $\mathcal{N}_{sg} \sim 10$ is more an order of magnitude but our conclusion is to recommend an upper bound value of \mathcal{N}_{sg} much lower than the former $N=50$, used by Yano and Phillips (2011) with their notation in the box model.

3.2 Changes in the microphysics

240 Basically, intensifying the CIBU process by increasing \mathcal{N}_{sg} leads to higher cloud ice crystal concentrations which deplete the supersaturation of water vapour that would otherwise contribute to the deposition growth of the snow-aggregates. However, a further effect is possible because the partial mass sink of the snow-aggregate particles also slows down the flux of graupel particles, which form essentially by heavy riming and conversion of the snow-aggregates. This point is now examined by looking at the ice in the high levels of the STERAO cells. Figures 3-5 reproduce the 10 minute average of the mixing ratios r_i, r_s and r_g at 12 km height from the 4 experiments $\mathcal{N}_{sg}=0.0, 0.1, 1.0$ and 10.0 after 4 hours. The increase of the cloud ice mixing ratio with \mathcal{N}_{sg} is clear in the area covered by the 0.2 g kg^{-1} isocontour in Fig. 3. Simultaneously, a slight decrease of r_s , indicating a slow erosion of the mass of the aggregates, is visible in Fig. 4. The effect on the graupel (Fig. 5) is even smaller but appears clearly for the case $\mathcal{N}_{sg}=10.0$, where less graupel is found. A last illustration is provided by Fig. 6, showing the number concentration of cloud ice N_i at a higher altitude of 15 km. Again, the increase of N_i follows \mathcal{N}_{sg} with an explosive multiplication of N_i when $\mathcal{N}_{sg}=10.0$ (N_i is well above

1000 crystals kg^{-1} of dry air in this case). Figure 7 summarizes the behaviour of r_i , r_s , r_g at 12 km height, and of N_i at 15 km height, for the "RANDOM" simulation. The results are those expected but, when comparing these results with Figs 3-6, it is not possible to find microphysics anomalies equivalent to the case where CIBU is not accounted for, so "RANDOM" is a **full** simulation scenario that is intermediate between $\mathcal{N}_{sg}=1$ and $\mathcal{N}_{sg}=10$.

The analysis of the STERAO simulations continues by looking at the vertical profiles of microphysics budgets. The profiles are 10 minute averages of all cloudy columns that contain at least 10^{-3} g kg^{-1} of condensate at any level. The column selection is updated at each time step because of the evolution and motion of the storm. Figure 8 shows the mixing ratio profiles in three cases: $\mathcal{N}_{sg} = 0.0$, "RANDOM" and $\mathcal{N}_{sg} = 10.0$. A key feature that shows up in Fig. 8a-c is the increase of the r_i peak value at 11 km altitude. This change is accompanied by a reduction of r_s (more visible between cases b) and c)) and by a reduction of r_g , which stands out at $z=8,000$ m. The decrease of r_g , even if graupels are passive colliders for CIBU, is the result of the decrease of r_s in the growth chain of the precipitating ice. The low value of the mean r_r profiles, compared to the mixing ratios of the ice phase above, is explained by the fact that rain is spread over fewer grid points than the ice in the anvil (the mixing ratio profiles are averaged over the same number of columns).

3.3 Budget of ice mixing ratios

The next step is dedicated to the microphysics tendencies (10 minute average again with the nomenclature of the processes provided in Table 3) of the ice mixing ratios in Fig. 9-11 to assess the impact of the CIBU process. We do not discuss the case of the liquid phase here because the tendencies (not shown) are not very much affected by CIBU.

As expected, **the tendencies of r_i (Fig. 9a-c) are the most affected by the CIBU process.** The main processes standing out in Fig. 9a, when CIBU is not activated, are CEDS (Deposition-Sublimation), essentially a gain term, and AGGS (Aggregation), the main loss of r_i by aggregation with a rate of $0.5 \times 10^{-3} \text{ g kg}^{-1} \text{ s}^{-1}$. The loss of r_i by CFRZ (Drop Freezing by Contact) makes a moderate contribution as some raindrops are present in the glaciated part of the storm. Above $z=10,000$ m, the net loss of r_i (AGGS and SEDI, the Cloud Ice Sedimentation) is balanced by the convective vertical transport (not shown). When $\mathcal{N}_{sg}=\text{RANDOM}$, the r_i tendencies are amplified, even with a modest contribution of $\sim 0.2 \times 10^{-3} \text{ g kg}^{-1} \text{ s}^{-1}$ for CIBU itself. The growth of AGGS, which doubles at 10 km height, is caused by CIBU and by an increase in the convection because SEDI (a loss there) is amplified in response to an increase of r_i in the upper levels. The CFRZ contribution is also increased. The last case, with $\mathcal{N}_{sg}=10$ (Fig. 9c) confirms a further increase of the rates except for CFRZ, interpreted here as a lack of raindrops.

The budget of the snow/aggregate mixing ratio in Fig. 10 contains many processes of equivalent importance in the range $\pm 0.05 \times 10^{-3} \text{ g kg}^{-1} \text{ s}^{-1}$ but SEDS (Sedimentation of Snow-aggregates) dominates negatively at $z=11,000$ m and positively at $z = 7,000$ m. The inclusion of CIBU (Fig. 10b-

c) mostly leads to an increase of AGGS, the other processes remaining almost the same. Finally many
 290 processes contribute to the evolution of the graupel mixing ratio profiles (Fig. 11). The strongest
 loss is in the GMLT term (Melting of graupel) that converts graupel into rain (down to -0.3×10^{-3}
 $\text{g kg}^{-1} \text{s}^{-1}$) while the contact freezing of the raindrops (CFRZ) reaches $0.15 \times 10^{-3} \text{g kg}^{-1} \text{s}^{-1}$.
 The sedimentation term SEDG (Sedimentation of Graupel) lies between $-0.3 \times 10^{-3} \text{g kg}^{-1} \text{s}^{-1}$ at
 $z = 10,000 \text{ m}$ and $0.15 \times 10^{-3} \text{g kg}^{-1} \text{s}^{-1}$ at $5,000 \text{ m}$. Another noticeable effect is the sign change
 295 of DEPG (Growth of Graupel by Deposition, $\pm 0.07 \times 10^{-3} \text{g kg}^{-1} \text{s}^{-1}$) showing that the water
 vapour is super(under)saturated above(below) $z=7,000 \text{ m}$ on average. The relative importance of
 these processes does not change very much when CIBU is increased but tendencies weaken. In
 summary, the impact of CIBU is modest for the microphysics mixing ratios. The increase of ice
 fragments in r_i is approximately compensated by an increase of AGGS (see Fig. 9 and 10).

300 3.4 Budget of cloud ice concentration

The next point examines the behaviour of the cloud ice number concentration as a function of the
 strength of the CIBU process after 4 hours of simulation. Figure 12 shows that the altitude of the N_i
 peak value decreases when \mathcal{N}_{sg} increases. In the absence of CIBU ($\mathcal{N}_{sg} = 0$), the source of N_i is the
 heterogeneous nucleation processes on insoluble IFN and on coated IFN (nucleation by immersion)
 305 which are more efficient at low temperature. Nucleation on IFN provides a mean peak value $N_i =$
 400 kg^{-1} at $z = 11,500 \text{ m}$. In contrast, the $\mathcal{N}_{sg} = 10$ case (here scaled by a factor 0.1 for plotting
 reasons) keeps the trace of an explosive production of cloud ice concentration, $N_i = 7,250 \text{ kg}^{-1}$,
 due to CIBU. The altitude of the maximum of N_i in this case ($z = 10,000 \text{ m}$) is consistent with
 the location of the maximum value of the $r_s \times r_g$ product (see Fig. 8). The "RANDOM" simulation
 310 produces $N_i = 1100 \text{ kg}^{-1}$ at $z = 11,000 \text{ m}$, a number concentration which is similar to that found
 for the $\mathcal{N}_{sg} = 2$ case. Table 2 reports the peak amplitude of the N_i profiles as a function of \mathcal{N}_{sg} but
 after 3 hours of simulation, when the CIBU rate is strongly dominant. Additional cases were run to
 cover $0.1 < \mathcal{N}_{sg} < 50$ with a logarithmic progression above $\mathcal{N}_{sg} = 1.0$. The CIBU enhancement factor,
 CIBU_{ef} , is computed as $N_i(\mathcal{N}_{sg})/N_i(\mathcal{N}_{sg} = 0) - 1$ as $N_i(\mathcal{N}_{sg} = 0)$ stands as a baseline not affected
 315 by CIBU. The results show that the growth of N_i is fast when \mathcal{N}_{sg} reaches ~ 5 (CIBU_{ef} switches
 from 135% to 913% when \mathcal{N}_{sg} moves from 2 to 5). Taking $\mathcal{N}_{sg} = 50$ leads to an extremely high
 peak value of N_i .

The N_i tendencies are the subject of Fig. 13. Many processes are involved during the temporal
 integration of N_i . The $\mathcal{N}_{sg} = 0$ case confirms the importance of the heterogeneous nucleation process
 320 by deposition, HIND, (refer to Table 3) and, to a lesser degree, by immersion (HINC) at 8 km
 height. HIND peaks at three altitudes with two sources of IFN (Table 1). This case also reveals the
 importance of the HMG (Hallett-Mossop on Graupel, $1.3 \text{ kg}^{-1} \text{s}^{-1}$) and HMS (Hallett-Mossop on
 Snow, $0.85 \text{ kg}^{-1} \text{s}^{-1}$) processes. Here, we consider that H-M also operates for the snow-aggregates
 because this category of ice includes lightly rimed particles that can rime further to form graupel

325 particles. These processes are first compensated by AGGS (capture of cloud ice by the aggregates).
 There is also a loss of cloud ice due to CFRZ and CEDS with the full sublimation of individual cloud
 ice crystals that replenish the IFN reservoir. The sedimentation profile transports ice from cloud top
 (SEDI<0) to mid-level cloud (SEDI>0). Then, taking $\mathcal{N}_{sg} = \text{RANDOM}$ shows the domination of
 the CIBU process, which reaches $2.5 \text{ kg}^{-1}\text{s}^{-1}$ at 5 km height. The enhancement of HIND at cloud
 330 top can also be noted. The CIBU source of ice crystals is balanced by an increase of AGGS and,
 above all, of CEDS (here CEDS represents the sublimation of the ice crystal concentration when the
 crystals are detrained in the low level of the cloud vicinity, below the anvil for instance). Finally, the
 $\mathcal{N}_{sg} = 10$ case demonstrates the reality of the exponential-like growth of N_i because the three main
 driving terms CIBU, CEDS and AGGS are growing at a similar rate, which is multiplied by a factor
 335 of approximately 5.

3.5 Sensitivity to the initial concentration of freezing nuclei

The purpose of the last series of experiments was to look more closely at the sensitivity of the cloud
 ice concentration to N_{IFN} , the initial concentration of the IFN. Numerical simulations were run
 with N_{IFN} decreasing by decades from 100 dm^{-3} to 0.001 dm^{-3} for each IFN mode (see Table 1).
 340 Two different cases were considered. In the first one, CIBU was activated with the RANDOM set-up
 while, in the second, CIBU effects were ignored. All the results are summarized in the plots of Fig.
 14.

Figure 14a shows that N_i concentrations did not change very much for a wide range of N_{IFN}
 concentrations, which were scanned by decades. This clearly illustrates the predominance of the
 345 CIBU effect for current IFN concentrations, which disconnects N_i concentrations from the underlying
 abundance of IFN particles. In this vein, the small hump superimposed on all profiles at 5,000
 m height reveals a residual effect of the Hallett-Mossop process. Another remarkable feature is that
 a fairly low IFN concentration ($N_{IFN} = 0.001 \text{ dm}^{-3}$) suffices to initiate the CIBU process and to
 reach $N_i \sim 500 \text{ kg}^{-1}$. In contrast, and in the absence of CIBU (Fig. 14b), the N_i profiles show a
 350 sensitivity to IFN nucleation that is, indeed, difficult to interpret because of the non-monotonic trend
 of the N_i profiles with respect to N_{IFN} . Some insight can be gained by checking the concentra-
 tion of the nucleated IFN of the first IFN mode (dust particles). In Fig. 14c, the IFN profiles are
 rescaled (multiplication by an appropriate numbers of powers of ten) to be comparable. Here, this is
 equivalent to computing an IFN nucleation efficiency. The important result here is that the number
 355 of nucleated IFN evolves in close proportion to the initially available IFN concentrations, meaning
 that the nucleating properties of the IFN do not depend on the IFN concentration as expected. The
 last plot (Fig. 14d) reproduces the normalized differences of N_i profiles between twin simulations
 performed with CIBU and without CIBU. Even if simulations made with the same initial concentra-
 tion N_{IFN} ; diverge because of additional non-linear effects (vertical transport, enhanced or reduced
 360 cloud ice sink processes), the figure gives a flavour of the bulk sensitivity of CIBU to the IFN. The

enhancement ratio due to CIBU remains low (less than 1 for $N_{IFN} \sim 100 \text{ dm}^{-3}$) but can reach a factor of 20 at 9,000 m height in the case of moderate IFN concentration i.e. $N_{IFN} \sim 1 \text{ dm}^{-3}$. The behaviour of LIMA can be explained in the sense that increasing N_{IFN} too much leads to smaller pristine crystals that need a longer time to grow because the conversion to the next category of snow-aggregates is size-dependent (see Harrington et al. (1995) and Vié et al. (2016)). On the other hand, a low concentration of N_{IFN} initiates fewer snow-aggregate and thus fewer graupel particles, so the whole CIBU efficiency is also reduced. Consequently, this study confirms the essential role of CIBU in compensating for IFN deficit when cloud ice concentrations are building up.

4 Simulation of a 3-dimensional idealized supercell storm with varying atmospheric stability

The test case (referred as WK) was suggested by the idealized sounding of Weisman and Klemp (1982, 1984) where the intensity of the CAPE can be easily modified by changing a reference water vapour mixing ratio. The simulation conditions were close to those of the STERAO case with the same set-up for the physics and the aerosol characteristics. The domain simulation was 180×180 at 1 km resolution and 70 levels with a mean vertical grid spacing of 350 m. Convection was triggered by a domain-centered single 2K-buoyant air parcel of 10 km radius and 3 km height. The base of the upper level Rayleigh damper was set at 15 km above the ground.

Meso-NH was initialized with the analytic sounding of Weisman and Klemp (1984) with a low 2-dimensional shear. The hodograph in Fig. 15 features a three-quarter-cycle with a constant wind of 6.4 m s^{-1} (in modulus) above the height of 5 km. When running Meso-NH a constant translation speed ($U_{trans}=5 \text{ m s}^{-1}$ and $V_{trans}=1 \text{ m s}^{-1}$) was added to the wind to keep the convection well centered in the domain of simulation. As explained in Weisman and Klemp (1982), buoyancy is varied by altering the magnitude of the surface water vapour mixing ratio q_{v0} in Weisman and Klemp notation. So three water vapour profiles were defined taking $q_{v0} = 13.5 \text{ g kg}^{-1}$, hereafter the "Low" CAPE case of 1970 J kg^{-1} ; $q_{v0} = 14.5 \text{ g kg}^{-1}$ as the "Mid" CAPE case of 2400 J kg^{-1} , and $q_{v0} = 15.5 \text{ g kg}^{-1}$, the "High" CAPE case with 2740 J kg^{-1} . Four 4h-experiments were performed for each CAPE case by changing the magnitude of \mathcal{N}_{sg} .

4.1 Sensitivity to the mean ice concentrations

The mean concentrations of the small ice crystals between 9.5 and 10.5 km levels are plotted on a log scale in Fig. 16 after 4 hours of simulation. In addition, two CTH (Cloud Top Height) isocontours delineate the 11 km (dotted line) and 13 km (solid line) levels. The $\mathcal{N}_{sg} = 0, \text{RANDOM}, 10$ and 50 cases, are explored for each sounding ("Low", "Mid" and "High" CAPE). In the absence of CIBU (first row in Fig. 16), the cloud ice concentrations N_i are in the range of what was simulated for the STERAO case (see Figs. 6 and 7d). The N_i peak values do not increase with the initial CAPE (Figs 16a-b) but the area of $\text{CTH} > 11 \text{ km}$ is larger in the "Mid CAPE" case. The "High CAPE" case is a

395 little more difficult to analyse because of an earlier development of the convection, spreading out
ahead of the main system, showing up in the "Low" and "Mid" CAPE cases. However, the N_i peak
values of the "High" CAPE case are in the range of the "Low" CAPE case, meaning that a higher
environmental instability is not decisive in fixing the N_i peak values. Jumping now to the $\mathcal{N}_{sg} = 10$
and 50 cases, we retrieve the dramatic increase of N_i due to an increasing CIBU efficiency. The
400 enhancement is locally as high as one thousand fold in the strongest case ($\mathcal{N}_{sg} = 50$). There are also
other noteworthy features: an increase of the N_i area coverage with \mathcal{N}_{sg} (less visible in the "Low"
CAPE case) and a higher CTH which exceeds 13 km for the "Mid" and "High" CAPE cases. All
these observations strongly suggest that convection is invigorated when the CIBU effect is increased.
In contrast, the simulations run with $\mathcal{N}_{sg} = \text{RANDOM}$ with values taken in the 0.1-10 range (see
405 Section 2.1), show a moderate effect of CIBU. Locally, N_i values reach $1 \times 10^4 \text{ kg}^{-1}$, which is one
hundred less than N_i peak values in the $\mathcal{N}_{sg} = 50$ cases but approximately, ten times more than in
the "no CIBU" case ($\mathcal{N}_{sg} = 0$). Finally the simulation results suggest that the \mathcal{N}_{sg} parameter could
be constrained by satellite data because of the sensitivity of CIBU to the cloud ice coverage and the
cloud top height.

410 4.2 Sensitivity to the precipitation

The 4-hour accumulated precipitation maps are presented in Fig. 17. On each row, precipitation
increase from the "Low" to "High" CAPE cases. This is because the CAPE is enhanced by the
addition of more and more water vapour to the atmosphere. Looking now at the sensitivity of the
accumulated precipitation to \mathcal{N}_{sg} , it is not easy to draw a general conclusion on the decrease of
415 the precipitation peak with \mathcal{N}_{sg} as for the STERAO case (see section 3.1). The reason is the highly
concentrated precipitation field, which leads to a sharp gradient around the location of the peak
value. However, the decrease of the precipitation with \mathcal{N}_{sg} is observed in the "Low" and "High"
CAPE cases. In the "Mid" case, the precipitation peak value remains high when $\mathcal{N}_{sg} = 50$ but the
area where the precipitation is less than 10 mm shrinks continuously. The reduction of the area where
420 the precipitation amount is greater than 10 mm when \mathcal{N}_{sg} is increased, operates in all CAPE cases
(not shown).

In conclusion, the simulations illustrate the fact that the precipitation patterns are affected by
the value of the \mathcal{N}_{sg} parameter. When \mathcal{N}_{sg} is increased from zero up to 50, the precipitation is
reduced, either for the peak value or at least for the precipitating area. This is consistent with our
425 previous results concerning the STERAO case. The conversion efficiency of the small ice crystals to
precipitating ice particles is lower when the cloud ice concentration is high because the deposition
growth of individual small crystals is limited by the amount of supersaturated water vapour available.

4.3 Sensitivity to the ice thickness

The last analysis is concerned with the **ice thicknesses** computed as the integral along the vertical
430 of $\rho_{dref} r_x$ where r_x refers to the mixing ratio with $x \in i, s, g$ standing for the cloud ice, the snow-
aggregates and the graupel-hail, respectively. Fig. 18 displays the total ice thickness, a sum of three
terms, in mm (coloured area) with the superimposed cloud ice thickness (THIC), contoured at 1 mm.
A remarkable feature is that the total ice thickness seems almost insensitive to the CIBU process for
a given CAPE case as there is no great modification in the plots when moving from $\mathcal{N}_{sg} = 0$ to
435 $\mathcal{N}_{sg} = 50$. **This is in contrast with the cloud ice thickness, for which the area increases with \mathcal{N}_{sg} .**
The rise in the maximum value of THIC was also expected for growing values of \mathcal{N}_{sg} . However, the
increase of THIC_{\max} with the CAPE is much more moderate between the "Low" and "High" cases
because a higher CAPE regime with higher humidity tends to favour the horizontal spread of the
cloud ice mass more.

440 5 Summary and perspectives

The aim of this work was to study a **comprehensive** parameterization of the Collisional Ice Break-
Up for a bulk 2-moment microphysics scheme LIMA running in a cloud resolving mesoscale model
(Meso-NH in our case). While the process is suspected to occur in real clouds, it is not included in
current bulk microphysics schemes. Because of uncertainties, the present parameterization has been
445 kept as simple as possible. It considers only collisions between small aggregates and large dense
graupel particles. The number of ice fragments that results from a single collision, \mathcal{N}_{sg} , is a key
parameter, which is estimated from only very few past experiments (Vardiman, 1978). A merit of
this study is to suggest an upper bound to the value of \mathcal{N}_{sg} because of the sensitivity of \mathcal{N}_{sg} to the
simulated precipitation. **We found that taking $\mathcal{N}_{sg} > 10$ significantly reduces the precipitation at the**
450 **ground. This is problematic since most of the cloud schemes (running without the CIBU process) are**
tuned for quantitative precipitation forecasts. Going further, we suggest that \mathcal{N}_{sg} could be considered
as the realization of a random process because delicate radiating crystals undergoing fragmentation
lead to a variety of crystals with a missing arm or to many irregular fragments as illustrated and
discussed by Hobbs and Farber (1972). As a result, it has been shown, that running LIMA with
455 $\mathcal{N}_{sg} > 10$ for the **STERAO and WK** deep convection cases, alters the precipitation at the ground
because the conversion of cloud ice crystals into precipitating ice is slowed down. In any case, the
increase of the number concentration of the small ice crystals due to the application of CIBU is
clearly significant (up to one thousand fold in the WK simulations with $\mathcal{N}_{sg} = 50$).

The microphysics perturbation due to the activation of CIBU has been studied in detail for the
460 STERAO case by looking at the profiles of the mixing ratios, ice concentrations and corresponding
budget terms. In particular, the CIBU effect on the pristine ice and aggregate mixing ratios is com-
pensated by an enhancement of the capture of the small crystals by the aggregates. The sensitivity

of the ice concentration to \mathcal{N}_{sg} is demonstrated with a mean multiplication factor as high as 25 for $\mathcal{N}_{sg} = 10$. The last study on the sensitivity of the simulations to the initial IFN concentration showed that CIBU was mostly efficient for current IFN concentrations of $\sim 1 \text{ dm}^{-3}$. Furthermore, the CIBU process was still active for very low IFN concentrations, down to 0.001 dm^{-3} , which were sufficient to initiate the ice phase.

The effects of CIBU have been confirmed by the **additional** WK simulations. The enhancement of the cloud ice concentration is very high when $\mathcal{N}_{sg} > 10$ and a loss of surface precipitation (peak value and reduction of the precipitating areas) is found. Higher ice concentrations lead to a larger coverage of ice clouds and higher cloud tops for the most vigorous convective cells. In contrast, the total ice thickness is almost insensitive to CIBU. An increase of cloud ice mass with \mathcal{N}_{sg} is balanced by a slight decrease of the precipitating ice (aggregates and graupels).

The proposed parameterization is very easy to implement. It would be useful to evaluate it in other microphysics schemes where the conversion of the cloud ice and the growth of precipitating ice (aggregates and rimed particles) are treated differently. The basic adjustment of the scheme can be revised as soon as laboratory experiments are available for more precise fixing of the sizes and the shapes of the crystals that break following collisions, to examine any possible thermal effect and to estimate the variety of fragment numbers more accurately. Another way to determine the acceptable range of values for \mathcal{N}_{sg} is to work with satellite data, as the WK experiments demonstrated an enhancement of the cloud top ice cover (and possibly the cloud top height) with \mathcal{N}_{sg} .

With new imagers, counters and improvements in data analysis (Ladino et al., 2017), more and more evidence is being presented that ice multiplication is an essential process in natural deep convective clouds. However, the explanation of anomalously high ice crystal concentrations is still difficult to link to a precise process (Rangno and Hobbs, 2001; Field et al., 2017). So the next step in the LIMA scheme is to introduce the shattering of the raindrops during freezing as proposed by Lawson et al. (2015) in order to complete the LIMA scheme, since the different ingredients, raindrops and small ice crystals, offer another pathway for ice multiplication. A task is then to study whether all the known sources of small ice crystals, nucleation and secondary ice production, are able **to cooperate** in microphysics schemes to reproduce the very high values (units of cm^{-3}) of ice concentrations sometimes observed. Quantitative cloud data gathered in the tropics during HAIC/HIWC (High Altitude Ice Crystals/ High Ice water Content) field project (Leroy et al., 2015; Ladino et al., 2017) could be a starting point to evaluate the capability of high resolution cloud simulations to reproduce events where high cloud ice contents were recorded.

6 Code availability

The Meso-NH code is publicly available at <http://mesonh.aero.obs-mip.fr/mesonh51>. Here the model development and the simulations were carried with version "MASDEV5-1 BUG2". The modifica-

tions made to the LIMA scheme (v1.0) are available upon request from Jean-Pierre Pinty and in the Supplement related to this article, available at <http://doi.org/10.5281/zenodo.1078527>.

500 Appendix A: Moments of the gamma and incomplete gamma functions

The p^{th} moment of the generalized gamma function (see definition in the text) is

$$M(p) = \int_0^{\infty} D^p n(D) dD = \frac{\Gamma(\nu + p/\alpha)}{\Gamma(\nu)} \frac{1}{\lambda^p} \quad (A1)$$

where the gamma function is defined as:

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt. \quad (A2)$$

505 The p^{th} moment of the incomplete gamma function is written

$$M^{INC}(p; X) = \int_0^X D^p n(D) dD. \quad (A3)$$

The algorithm of the "GAMMA_INC($p; X$)" function (Press et al., 1992) is useful to tabulate $M^{INC}(p; X) \times \Gamma(p)$ in addition to the "GAMMA" function algorithm of Press et al. (1992). A change of variable is necessary to take the generalized form of the gamma size distributions into account.

510 As a result, $M^{INC}(p; X)$ is written:

$$M^{INC}(p; X) = M(p) \times \text{GAMMA_INC}(\nu + p/\alpha; (\lambda X)^\alpha) \quad (A4)$$

with $M(p)$ given by Eq. A1.

Acknowledgements. J.-P. Pinty wishes to thank V. Phillips for discussions about his original work on the topic. This work was done during the PhD of T. Hoarau who is financially supported by Reunion Island Regional
 515 Council and the European Union Council. T. Hoarau thanks the University of La Réunion for supporting a short stay at the Laboratoire d'Aérodologie. Mrs Susan Becker corrected the English of the manuscript. Preliminary computations were performed on the 36 node home-made cluster of Lab. Aérodologie. J.-P. Pinty acknowledges CALMIP (CALcul Midi-Pyrénées) of the University of Toulouse for access to the "eos" supercomputer where useful additional simulations were performed. T. Hoarau and C. Barthe acknowledge the GENCI
 520 ressources for access to the "Occigen" supercomputer. This work was supported by the French national programme LEFE/INSU through the LIMA-TROPIC project. The authors thank the reviewers for their pertinent comments which helped to greatly improve the first version of the manuscript.

References

- 525 Colella, P. and Woodward, P.: The piecewise parabolic method (PPM) for gas-dynamical simulations, *J. Comput. Phys.*, 54, 174–201, 1984.
- Dye, J. E., Ridley, B. A., Skamarock, W., Barth, M., Venticinque, M., Defer, E., Blanchet, P., Théry, C., Laroche, P., Baumann, K., Hubler, G., Parrish, D. D., Ryerson, T., Trainer, M., Frost, G., Holloway, J. S., Matejka, T., Bartels, D., Fehsenfeld, F. C., Tuck, A., Rutledge, S. A., Lang, T., Stith, J., and Zerr, R.: An overview of the Stratospheric-Tropospheric Experiment: Radiation, Aerosols, and Ozone (STERAO)-Deep
530 Convection experiment with results for the July 10, 1996 storm, *J. Geophys. Res.*, 105, 10,023–10,045, doi:10.1029/1999JD901116, 2000.
- Field, P. R.: Bimodal ice spectra in frontal clouds, *Quart. J. Roy. Meteor. Soc.*, 126, 379–392, doi:10.1002/qj.49712656302, 2000.
- Field, P. R., Lawson, R. P., Brown, P. R. A., Lloyd, G., Westbrook, C., Moisseev, D., Miltenberger, A.,
535 Nenes, A., Blyth, A., Choulaton, T., Connolly, P., Buehl, J., Crosier, J., Cui, Z., Dearden, C., DeMott, P., Flossmann, A., Heymsfield, A., Huang, Y., Kalesse, H., Kanji, Z. A., Korolev, A., Kirchgaessner, A., Lasher-Trapp, S., Leisner, T., McFarquhar, G., Phillips, V., Stith, J., and Sullivan, S.: Secondary ice production: current state of the science and recommendations for the future, *Meteor. Monogr.*, 58, 7.1–7.20, doi:10.1029/AMSMONOGRAPHS-D-16-0014.1, 2017.
- 540 Foote, G. B. and du Toit, P. S.: Terminal velocity of raindrops, *J. Appl. Meteor.*, 8, 249–253, 1969.
- Griggs, D. G. and Choulaton, T. W.: A laboratory study of secondary ice article production by the fragmentation of rime and vapour-grown ice crystals, *Quart. J. Roy. Meteor. Soc.*, 112, 149–163, doi:10.1002/qj.49711247109, 1986.
- Hallett, J. and Mossop, S. C.: Production of secondary ice particles during the riming process, *Nature*, 249,
545 26–28, doi:10.1038/249026a0, 1974.
- Harrington, J. Y., Meyers, M. P., Walko, R. L., and Cotton, W. R.: Parameterization of Ice Crystal Conversion Processes Due to Vapor Deposition for Mesoscale Models Using Double-Moment Basis Functions. Part I: Basic Formulation and Parcel Model Results, *J. Atmos. Sci.*, 52, 4344–4366, doi:10.1175/1520-0469(1995)052<4344:POICCP>2.0.CO;2, 1995.
- 550 Hobbs, P. V. and Farber, R. J.: Fragmentation of ice particles in clouds, *J. Rech. Atmos.*, 6, 245–258, 1972.
- Hobbs, P. V. and Rangno, A. L.: Ice particle concentrations in clouds, *J. Atmos. Sci.*, 42, 2523–2549, 1985.
- Ladino, L. A., Korolev, A., Heckman, I., Wolde, M., Fridlind, A. M., and Ackerman, A. S.: On the role of ice-nucleating aerosol in the formation of ice particles in tropical mesoscale convective systems, *Geophys. Res. Lett.*, 44, doi:10.1002/2016GL072455, 2017.
- 555 Lafore, J. P., Stein, J., Asencio, N., Bougeault, P., Ducrocq, V., Duron, J., Fischer, C., Héreil, P., Mascart, P., Masson, V., Pinty, J.-P., Redelsperger, J.-L., Richard, E., and Vila-Guerau de Arellano, J.: The Meso-NH atmospheric simulation system. Part I: adiabatic formulation and control simulations, *Annales Geophysicae*, 16, 90–109, doi:10.1007/s00585-997-0090-6, 1998.
- Lawson, R. P., Woods, S., and Morrison, H.: The microphysics of ice and precipitation development in tropical
560 cumulus clouds, *J. Atmos. Sci.*, 72, 2429–2445, doi:10.1175/JAS-D-14-0274.1, 2015.
- Leroy, D., Fontaine, E., Schwarzenboeck, A., Strapp, J. W., Lilie, L., Delanoe, J., Protat, A., Dezitter, F., and Grandin, A.: HAIC/HIWC Field Campaign - Specific findings on PSD microphysics in high IWC regions

- from in situ measurements: median mass diameters, particle size distribution characteristics and ice crystal shapes, SAE Technical Paper, doi:10.4271/2015-01-2087, 2015.
- 565 Phillips, V. T., Yano, J.-I., and Khain, A.: Ice multiplication by break-up in ice-ice collisions. Part I: Theoretical formulation, *J. Atmos. Sci.*, 74, 1705–1719, doi:10.1175/JAS-D-16-0224.1, 2017.
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., and Flannery, B. P.: *Numerical Recipes in FORTRAN: The Art of Scientific Computing*, Cambridge University Press, New-York, 1992.
- Rangno, A. L. and Hobbs, P. V.: Ice particles in stratiform clouds in the Arctic and possible mechanisms for the
570 production of high ice concentrations, *J. Geophys. Res.*, 106, 15 065–15 075, 2001.
- Skamarock, W. C., Powers, J. G., Barth, M., Dye, J. E., Matejka, T., Bartels, D., Baumann, K., Stith, J., Parrish, D. D., and Hübner, G.: Numerical simulations of the July 10 Stratospheric-Tropospheric Experiment: Radiation, Aerosol, and Ozone/Deep Convection Experiment convective system: Kinematics and transport, *J. Geophys. Res.*, 105, 19,973–19,990, 2000.
- 575 Sullivan, S. C., Hoose, C., and Nenes, A.: Investigating the contribution of secondary ice production to in-cloud ice crystal numbers, *J. Geophys. Res.*, 122, doi:10.1002/2017JD026 546, 2017.
- Sullivan, S. C., Hoose, C., Kiselev, A., Leisner, T., and Nenes, A.: Initiation of secondary ice production in clouds, *Atmos. Chem. Phys.*, 18, doi:10.5194/acp-18-1593-2018, 2018.
- Takahashi, T., Nagao, Y., and Kushiyama, Y.: Possible high ice particle production during graupel-graupel col-
580 lisions, *J. Atmos. Sci.*, 52, 4523–4527, 1995.
- Vardiman, L.: The generation of secondary ice particles in clouds by crystal–crystal collision, *Journal of the Atmospheric Sciences*, 35, 2168–2180, doi:10.1029/1978-0352168:JGOSIP>2.0.CO;2, 1978.
- Vié, B., Pinty, J.-P., Berthet, S., and Leriche, M.: LIMA (v1.0): A quasi two-moment microphysical scheme driven by a multimodal population of cloud condensation and ice freezing nuclei, *Geoscientific Model Development*, 9, 567–586, doi:10.5194/gmd-9-567-2016, 2016.
585
- Weisman, M. L. and Klemp, J. B.: The dependence of numerically simulated convective storms on vertical wind shear and buoyancy, *Mon. Wea. Rev.*, 110, 504–520, 1982.
- Weisman, M. L. and Klemp, J. B.: The structure and classification of numerically simulated convective storms in directionally varying wind shear, *Mon. Wea. Rev.*, 112, 2479–2498, 1984.
- 590 Yano, J.-I. and Phillips, V. T.: Ice–Ice Collisions: An Ice Multiplication Process in Atmospheric Clouds, *J. Atmos. Sci.*, 68, 322–333, doi:10.1175/2010JAS3607.1, 2011.
- Yano, J.-I. and Phillips, V. T.: Explosive Ice Multiplication Induced by multiplicative-Noise fluctuation of Mechanical Break-up in Ice-Ice Collisions, *J. Atmos. Sci.*, 73, 4685–4697, doi:10.1175/JAS-D-16-0051.1, 2016.
- Yano, J.-I., Phillips, V. T., and Kanawade, V.: Explosive ice multiplication by mechanical break-up in ice–ice
595 collisions: a dynamical system-based study, *Quart. J. Roy. Meteor. Soc.*, 142, 867–879, doi:10.1002/qj.2687, 2016.

CCN	Aitken mode	Accumulation mode	Coarse mode
N (cm^{-3})	300	140	50
d_X (μm)	0.23	0.8	2.0
σ_X	2.0	1.5	1.6

IFN	Dust mode	BC+Organics mode
N (dm^{-3})	10	10
d_X (μm)	0.8	0.2
σ_X	2.0	1.6

Table 1. Background CCN and IFN configuration for the STERAO idealized case simulations.

\mathcal{N}_{sg} (no unit)	0.0	0.1	1.0	2.0	5.0	10.0	20.0	50
N_i ($\#kg^{-1}$)	790	940	1,160	1,860	8,000	25,670	62,010	112,740
CIBU _{ef} (%)	0	19	47	135	913	3149	7749	14171

Table 2. After 3 hours of simulation, maximum value of the cloud ice number concentration $N_{i,max}$ as a function of the number of fragments produced per snow/aggregate-graupel collision \mathcal{N}_{sg} . The last row is the CIBU enhancement factor CIBU_{ef} in percent (see text).

Process Acronym	Description
ACC	Raindrop accretion on snow to produce graupel
AGGS	Snow growth by capture of cloud ice
BERFI	Growth of cloud ice by Bergeron-Findeisen process
CEDS	Deposition/sublimation of water vapour on cloud ice
CFRZ	Raindrop Freezing by contact with cloud ice
CIBU	Snow break-up by collision with graupel
CMEL	Conversion Melting of snow into graupel
CNVI	Decreasing snow converted back to cloud ice
CNVS	Growing cloud ice converted into snow
DEPG	Water vapour deposition on graupel
DEPS	Water vapour deposition on snow
DRYG	Graupel dry growth (water can freeze fully)
HINC	Heterogeneous nucleation by immersion
HIND	Heterogeneous nucleation by deposition
HONC	Homogeneous freezing of the cloud droplets
HONH	Haze homogeneous freezing
HMG	Droplet riming and Hallett-Mossop process on graupel
HMS	Droplet riming and Hallett-Mossop process on snow
IMLT	Melting of cloud ice
RIM	Riming of cloud droplets on snow to produce graupel
SEDI	Sedimentation of cloud ice, snow or graupel
WETG	Graupel wet growth (water is partially frozen)

Table 3. Nomenclature of the microphysics processes of the budget profiles.




Figure 1. 4-h accumulated precipitation of the STERAO simulations where a) to d) refers to cases with $N_{sg}=0.0, 0.1, 1.0$ and 10.0 ice fragments per collision, respectively. The plots are for a fraction of the computational domain.




Figure 2. Same as Fig. 1, but for the "RANDOM" simulation.




Figure 3. Mixing ratios of the cloud ice (r_i in log scale) of the STERAO simulations at 12 km height, where a) to d) refer to cases with $N_{sg}=0.0, 0.1, 1.0$ and 10.0 ice fragments per collision, respectively. The plots are for a fraction of the computational domain.




Figure 4. Same as Fig. 3 but for the mixing ratios of snow-aggregates (r_s).




Figure 5. Same as Fig. 3 but for the mixing ratios of graupel (r_g).




Figure 6. Number concentration of the cloud ice (N_i in log scale) of the STERAO simulations at 15 km height, where a) to d) refer to cases with $\mathcal{N}_{sg}=0.0, 0.1, 1.0$ and 10.0 ice fragments per collision, respectively. The plots are for a fraction of the computational domain.




Figure 7. "RANDOM" case of the STERAO simulations showing the mixing ratios of a) the cloud ice (r_i), b) the snow-aggregates (r_s), and c) the graupel (r_g) at 12 km height. Plot d) refers to the number concentration of the cloud ice crystals (N_i) at 15 km height. The plots are for a fraction of the computational domain.




Figure 8. Mean profiles of condensate mixing ratios r_c, r_r, r_i, r_s and r_g ; in g kg^{-1}) of the STERAO simulations corresponding to a) the $\mathcal{N}_{sg}=0.0$ case, b) the "RANDOM" case and c) the case with $\mathcal{N}_{sg} = 10.0$.




Figure 9. Mean microphysics profiles of cloud ice mixing ratio tendencies of the STERAO simulations corresponding to a) the $\mathcal{N}_{sg} = 0.0$ (no CIBU) case, b) the "RANDOM" case and c) the case with $\mathcal{N}_{sg} = 10.0$. The dashed lines are associated with processes having no significant impact on these budgets.




Figure 10. Same as Fig. 9 but for snow-aggregates.




Figure 11. Same as Fig. 9 but for graupel.




Figure 12. Mean profiles of the cloud ice crystal concentrations N_i (g kg^{-1}) of the STERAO simulations corresponding to different values of N_{sg} (see the legend for details). The profiles drawn with a dashed line have been divided by 10 to fit into the plot.




Figure 13. Mean microphysics profiles of the cloud ice crystal concentration tendencies of the STERAO simulations corresponding to a) the $\mathcal{N}_{sg} = 0.0$ (no CIBU) case, b) the "RANDOM" case and c) the case with $\mathcal{N}_{sg} = 10.0$ (Note that the horizontal scale increases from a) to c)). The dashed lines of the list box are associated with processes having no significant impact on these budgets.




Figure 14. Mean profiles of cloud ice crystal concentration for 6 decades of initial IFN concentrations from 100 dm^{-3} to 0.001 dm^{-3} of the STERAO simulations corresponding to a) the CIBU simulation and "RANDOM" case and b) the non-CIBU simulation. The mean profiles of the nucleated IFN concentrations are plotted in c) after rescaling to fit the [0.0-1.0] range. The rough estimate of CIBU enhancement factor of N_i is plotted in d) as a function of the initial IFN concentrations.




Figure 15. Vertical profile of the horizontal wind components of the WK84 simulations. The solid line with a constant shear ($2.5 \times 10^{-2} \text{ s}^{-1}$) refers to U , the x -component of the wind and the dashed line with a jet-like structure, refers to V , the y -component of the wind. U and V are constant above 5 km height.




Figure 16. Small ice concentration N_i average between 9.5 and 10.5 km height after 4 hours of the WK84 simulations, where a) to c) refer to no CIBU cases ($N_{sg}=0.0$), d) to f) to cases with random CIBU ($0.1 < N_{sg} < 10$) and g) to i) to cases with a high CIBU effect ($N_{sg}=10.0$), and j) to l) to cases with an intense CIBU effect ($N_{sg}=50.0$). The isocontours are the cloud top heights with dotted lines for 11 km and solid lines for 13 km.




Figure 17. As in fig. 16, but for the 4-h accumulated precipitation of the WK84 simulations. The peak value (max in mm) corresponds to the peak value of precipitation of the main convective clouds in the centre of the simulation domain.




Figure 18. As in fig. 16, but for the total ice thickness in mm after 4 hours of the WK84 simulations. The additional isocontours are the small ice thickness component (THIC) taken at 1 mm. The peak value of THIC (THIC_{max} is given in mm).