

# TAMSAT-ALERT v1: A new framework for agricultural decision support (Supplementary information)

Dagmawi Asfaw<sup>1</sup>, Emily Black<sup>1</sup>, Matt Brown<sup>2</sup>, Kathryn Jane Nicklin<sup>3</sup>, Frederick Out-Larbi<sup>4</sup>, Ewan Pinnington<sup>1</sup>, Andrew Challinor<sup>3</sup>, Ross Maidment<sup>1</sup>, Tristan Quaife<sup>1</sup>

<sup>1</sup>Department of Meteorology, University of Reading, Reading, UK

<sup>2</sup>Department of Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, UK

<sup>3</sup>School of Earth and Environment, University of Leeds, Leeds, UK

<sup>4</sup>Ghana Meteorological Agency, Accra, Ghana

**Table S1. GLAM–maize parameter values used for the simulation of maize yield in Ghana**

Parameter	Explanation	Value	Reference
<b>Leaves</b>			
DLDTMX	Maximum daily increase in LAI	0.1	(Watson et al., 2015)
MASPA	Minimum daily increase in senesced LAI during leaf senescence.	0.03	(Bergamaschi et al., 2013)
SWF_THRESH	Critical value of soil water stress factor for leaves	0.7	(Bergamaschi et al., 2013)
NDSLAI	Specific Leaf Area control	5 days	(Challinor and Wheeler, 2008)
SLA_INI	Specific leaf area control (see Challinor and Wheeler 2008)	350 cm <sup>2</sup> g <sup>-1</sup>	(Ashraf and Hafeez , 2004)
MAX_ISTG_SLA	Max development stage for SLA control	2 (until flowering)	
<b>Evaporation and transpiration</b>			
ALBEDO	Albedo	0.25	(Oguntunde and van de Giesen, 2004)
CRIT_LAI_T	LAI below which transpiration is physiologically limited	2.7	(Al-Kaisi et al., 1989; Bergamaschi et al., 2013)
P_TRANS_MAX	Max value of potential transpiration	0.73 cm day <sup>-1</sup>	(Al-Kaisi et al., 1989)
VPD_CTE	Used to calculate vapour pressure deficit	0.7	(Tanner and Sinclair, 1983)
VPD_REF	Used to calculate Priestly–Taylor coefficient	1kPa	(Steiner et al., 1991)
SHF_CTE	Used to calculate soil heat flux	0.4	(Choudhury et al., 1987)
EXTC	Extinction coefficient	0.5	(Bergamaschi et al., 2013)

<b>Soil and roots</b>			
RLL	Lower limit of volumetric soil water ( $\text{m}^3 \text{m}^{-3}$ )	$0.083 \text{ m}^3 \text{m}^{-3}$	
DUL	Drained upper limit of volumetric soil water ( $\text{m}^3 \text{m}^{-3}$ )	$0.195 \text{ m}^3 \text{m}^{-3}$	
SAT	Saturated volumetric soil water ( $\text{m}^3 \text{m}^{-3}$ )	$0.409 \text{ m}^3 \text{m}^{-3}$	
ASWS	Initial soil water (fraction of water holding capacity)	0	
NWBDAYS	Number of days water balance is run before start of planting window	30 days	
NSL	Number of soil layers	25	(Challinor et al., 2004)
ZSMAX	Maximum rooting depth	150 cm	
RKCTE	Used to calculate saturated hydraulic conductivity of the soil	$75 \text{ cm day}^{-1}$	(Suleiman and Ritchie, 2001)
UPDIFC	Uptake diffusion coefficient	$0.25 \text{ cm}^2 \text{day}^{-1}$	(Jamieson and Ewert, 1999; Robertson and Fukai, 1994).
EFV	Extraction front velocity	$2 \text{ cm day}^{-1}$	(Dardanelli et al., 1997; Bergamashi et al., 2013)
DLDLAI	Increase in root length density at surface with LAI	$1 \text{ cm cm}^{-3}$	(Bergamaschi et al., 2013)
RLVEF	Root length density at the extraction front	$0.3 \text{ cm cm}^{-3}$	(Watson et al., 2015)
<b>Biomass and yield</b>			
TE	Transpiration efficiency	7 Pa (1994–2006) 8 Pa (2007–2014)	(Walker, 1986; Adamtey et al., 2010)
TEN_MAX	Max transpiration efficiency	$10 \text{ g kg}^{-1}$	(Walker, 1986; Adamtey et al., 2010)
RUE	Radiation Use Efficiency	$3.4 \text{ g MJ}^{-1}$	(Kiniry et al., 1989; Lindquist et al. , 2005)
DHDT	Maximum daily increase in harvest index	$0.011 \text{ day}^{-1}$	(Birch et al., 1999)
MAX_HI_MAIZE	Maximum harvest index for maize	0.6	(Hay and Gilbert, 2001)
<b>Development</b>			
FSWSOW	Fractional soil moisture for intelligent planting and emergence	0.5	(Challinor et al., 2004)
IEMDAY	Number of days from planting to emergence	6 days	(Ashagre et al., 2014)

I_TTCALC	Shape of function to calculate thermal time	1 (flat top)	(Keating et al., 1992)
TBMAI	Base temperature for maize	10 °C	(Sanchez et al., 2014)
TOMAI	Optimum temperature for maize	30 °C	(Sanchez et al., 2014)
TMMAI	Maximum temperature for maize	40 °C	(Sanchez et al., 2014)
I_PHEN	Complex (I_PHEN=0) or simple (I_PHEN=1) simulation of phenology	1	
TLIMFLW	Thermal time from emergence to silking	680 °C days	
TLIMPFL	Thermal time from silking to start of grain filling.	145 °C days	(Tumbo et al., 2010)
TLIMGFP	Thermal time from start of grain filling to maturity	750 °C days	
<b>Optional processes</b>			
TETRS	Switch for temperature dependence of TE and RUE	1(on)	
TETR1	Temperature above which TE and RUE begin linear reduction	35 °C	(Yang et al., 2004; Carberry et al., 1989)
TETR2	Temperature above which TE and RUE are zero	47 °C	(Yang et al., 2004; Carberry et al., 1989)
TETR3	Temperature below which TE and RUE are zero	7 °C	(Yang et al., 2004; Carberry et al., 1989)
TETR4	Temperature below which TE and RUE begin linear reduction	18 °C	(Yang et al., 2004; Carberry et al., 1989)
HTS	Switch for high temperature stress around flowering	0(off)	(Gourdji et al., 2013)
TDS	Switch for terminal drought stress (TDS)	1(on)	
HIMIN	Minimum harvest index for TDS to occur	0.25	(Challinor et al., 2009)
SWC_FAC	Fraction of water holding capacity below which TDS occurs	0.05	(Challinor et al., 2009)
MIN_ISTG_TDS	Minimum ISTG for TDS to occur.	4 (start of grain filling)	
NMAXTDS	Number of days of TDS needed for the crop to be harvested.	8 days	(Jones and Thornton, 2003)
IEMER	Switch for intelligent emergence	-1 (off)	

TRKILL	Switch for lethal temperature parameterisation	-1 (off)	
WS	Switch for water stress around flowering	0 (off)	
TRLAI	Switch for temperature dependence of LAI growth	1 (on)	
TRLAIB	Base temperature below which leaf growth does not occur	7.3 °C	(Sanchez et al., 2014)
TRLAIO	Optimum temperature for leaf growth	31.1 °C	(Sanchez et al., 2014)
TRLAIM	Maximum temperature for leaf growth	41.3 °C	(Sanchez et al., 2014)

## References

- Adamtey, N., Cofie, O., Ofosu-Budu, K. G., Ofosu-Anim, J., Laryea, K. B., and Forster, D.: Effect of N-enriched co-compost on transpiration efficiency and water-use efficiency of maize (*Zea mays* L.) under controlled irrigation, *Agricultural Water Management*, 97, 995–1005, doi:10.1016/j.agwat.2010.02.004, 2010.
- Al-Kaisi, M., Brun, L. J., and Enz, J. W.: Transpiration and evapotranspiration from maize as related to leaf area index, *Agricultural and Forest Meteorology*, 48, 111–116, doi:10.1016/0168-1923(89)90010-5, 1989.
- Ashagre, H., Hamza, I. A., Tesfaye, B., Derebachew, D., and Ayana, B.: Emergence and Seedling Growth of Corn (*Zea mays* L.) as Influenced by Irrigation Schedules on Vertisol, *International Research Journal of Plant Science*, 5, 17–22, doi:http://dx.doi.org/10.14303/irjps.2013.061, 2014.
- Ashraf, M. and Hafeez, M.: Thermotolerance of pearl millet and maize at early growth stages: Growth and nutrient relations, *Biologia Plantarum*, 48, 81–86, doi:10.1023/B:BIOP.0000024279.44013.61, 2004.
- Bergamaschi, H., da Costa, S. M. S., Wheeler, T. R., and Challinor, A. J.: Simulating maize yield in sub-tropical conditions of southern Brazil using Glam model, *Pesquisa Agropecuaria Brasileira*, 48, 132–140, doi:10.1590/S0100-204X2013000200002, 2013.
- Birch, C. J., Hammer, G. L., and Rickert, K. G.: Dry matter accumulation and distribution in five cultivars of maize (*Zea mays*): Relationships and procedures for use in crop modelling, *Australian Journal of Agricultural Research*, 50, 513–527, doi:10.17700/jai.2015.6.1, 1999.
- Carberry, P. S., Muchow, R. C., and McCown, R. L.: Testing the CERES-Maize simulation model in a semi-arid tropical environment, *Field Crops Research*, 20, 297–315, doi:10.1016/0378-4290(89)90072-5, 1989.
- Challinor, A. J. and Wheeler, T. R.: Use of a crop model ensemble to quantify CO<sub>2</sub> stimulation of water-stressed and well watered crops, *Agricultural and Forest Meteorology*, 148, 1062–1077, doi:10.1016/j.agrformet.2008.02.006, 2008.
- Challinor, A. J., Wheeler, T. R., Craufurd, P. Q., Slingo, J. M., and Grimes, D. I. F.: Design and optimisation of a large-area process-based model for annual crops, *Agricultural and Forest Meteorology*, 124, 99–120, doi:10.1016/j.agrformet.2004.01.002, 2004.
- Challinor, A. J., Wheeler, T., Hemming, D., and Upadhyaya, H. D.: Ensemble yield simulations: Crop and climate uncertainties, sensitivity to temperature and genotypic adaptation to climate change, *Climate Research*, 38, 117–127, doi:10.3354/cr00779, 2009.
- Choudhury, B. J., Idso, S. B., and Reginato, R. J.: Analysis of an empirical model for soil heat flux under a growing wheat crop for estimating evaporation by an infrared-temperature based energy balance equation, *Agricultural and Forest Meteorology*, 39, 283–297, doi:10.1016/0168-1923(87)90021-9, 1987.
- Dardanelli, J. L., Bachmeier, O. A., Sereno, R., and Gil, R.: Rooting depth and soil water extraction patterns of different crops in a silty loam haplustoll, *Field Crops Research*, 54, 29–38, doi:10.1016/S0378-4290(97)00017-8, 1997.
- Gourdji, S. M., Sibley, A. M., and Lobell, D. B.: Global crop exposure to critical high temperatures in the reproductive period: Historical trends and future projections, *Environmental Research Letters*, 8, 0–10, doi:10.1088/1748-9326/8/2/024041, 2013.
- Hay, R. and Gilbert, R.: Variation in the harvest index of tropical maize: evaluation of recent evidence from Mexico and Malawi, *Annals of Applied Biology*, 138, 103–109, doi:10.1111/j.1744-7348.2001.tb00090.x, 2001.
- Jamieson, P. D. and Ewert, F.: The role of roots in controlling soil water extraction during drought: An analysis by simulation, *Field Crops Research*, 60, 267–280, doi:10.1016/S0378-4290(98)00144-0, 1999.
- Jones, P. G. and Thornton, P. K.: The potential impacts of climate change on maize production in Africa and Latin America in 2055, *Global Environmental Change*, 13, 51–59, doi:10.1016/S0959-3780(02)00090-0, 2003.
- Keating, B. A., Wafula, B. M., and Watiki, J. M.: Development of a modelling capability for maize in semi-arid eastern Kenya, in: *A Search for Strategies for Sustainable Dryland Cropping in Semi-arid Eastern Kenya*, edited by Probert, M. E., 41, pp. 26–32, ACIAR Proceedings No 41, Nairobi, Kenya, URL [http://aciarc.gov.au/files/node/2274/sustainable{\\\_}dryland{\\\_}cropping{\\\_}in{\\\_}semi{\\\_}arid{\\\_}eastern{\\\_}{\\\_}{\\\_}1109.pdf{\#}page=11](http://aciarc.gov.au/files/node/2274/sustainable{\_}dryland{\_}cropping{\_}in{\_}semi{\_}arid{\_}eastern{\_}{\_}{\_}1109.pdf{\#}page=11), 1992.
- Kiniry, J. R., Jones, C. A., O'toole, J. C., Blanchet, R., Cabelguenne, M., and Spanel, D. A.: Radiation-use efficiency in biomass accumulation prior to grain-filling for five grain-crop species, *Field Crops Research*, 20, 51–64, doi:10.1016/0378-4290(89)90023-3, 1989.

- Lindquist, J. L., Arkebauer, T. J., Walters, D. T., Cassman, K. G., and Dobermann, A.: Maize Radiation Use Efficiency under Optimal Growth Conditions, *Agronomy Journal*, 97, 72–78, 2005.
- Oguntunde, P. G. and Van De Giesen, N.: Crop growth and development effects on surface albedo for maize and cowpea fields in Ghana, West Africa, *International Journal of Biometeorology*, 49, 106–112, doi:10.1007/s00484-004-0216-4, 2004.
- Robertson, M. J. and Fukai, S.: Comparison of water extraction models for grain sorghum under continuous soil drying, *Field Crops Research*, 36, 145–160, doi:10.1016/0378-4290(94)90063-9, 1994.
- Sanchez, B., Rasmussen, A., and Porter, J. R.: Temperatures and the growth and development of maize and rice: A review, *Global Change Biology*, 20, 408–417, doi:10.1111/gcb.12389, 2014.
- Steiner, J., Howell, T., and Schneider, A.: Lysimetric evaluation of daily potential evapotranspiration models for grain sorghum, *Agronomy Journal*, 83, 240–247, doi:10.2134/agronj1991.00021962008300010055x, 1991.
- Suleiman, a. and Ritchie, J. T.: Estimating saturated hydraulic conductivity from soil porosity, *Transactions of the ASAE*, 44, 235–239, doi:10.13031/2013.4683, 2001.
- Tanner, C. B. and Sinclair, T. R.: Efficient Water Use in Crop Production: Research or Re-Search?, in: *Limitations to Efficient Water Use in Crop Production*, edited by Tanner, C. and Sinclair, T. R., pp. 1–27, American Society of Agronomy, Crop Science Society of America and Soil Science Society of America, Madison, Wisconsin, 1983.
- Tumbo, S. D., Rwehumbiza, F. B., Kahimba, F. C., Enfors, E., and Henry, F.: Agronomic Management Strategies for Adaptation to the Current Climate Variability: The Case of North-Eastern Tanzania, <http://hdl.handle.net/10625/46075>, 2010.
- Walker, G. K.: Transpiration efficiency of field-grown maize, *Field Crops Research*, 14, 29–38, doi:10.1016/0378-4290(86)90044-4, 1986.
- Watson, J., Challinor, A. J., Fricker, T. E., and Ferro, C. A.: Comparing the effects of calibration and climate errors on a statistical crop model and a process-based crop model, *Climatic Change*, 132, 93–109, doi:10.1007/s10584-014-1264-3, 2015.
- Yang, H. S., Dobermann, A., Lindquist, J. L., Walters, D. T., Arkebauer, T. J., and Cassman, K. G.: Hybrid-maize - A maize simulation model that combines two crop modeling approaches, *Field Crops Research*, 87, 131–154, doi:10.1016/j.fcr.2003.10.003, 2004.