Explicit representation and parametrised impacts of under ice shelf seas in the z* coordinate ocean model NEMO 3.6

Pierre Mathiot^{1,2}; Adrian Jenkins¹; Christopher Harris²; Gurvan Madec³

¹ British Antarctic Survey, NERC, Cambridge, UK

5 ²Met Office, Exeter, UK

³Sorbonne Universités (University Pierre et Marie Curie Paris 6)-CNRS-IRD-MNHN, LOCEAN Laboratory, Paris, France Correspondence to: Pierre Mathiot (pierre.mathiot@metoffice.gov.uk)

Abstract. Ice shelf/ocean interactions are a major source of freshwater on the Antarctic continental shelf and have a strong impact on ocean properties, ocean circulation and sea ice. However, climate models based on the ocean/sea ice model
NEMO (Nucleus for European Modelling of the Ocean) currently do not include these interactions in any detail. The capability of explicitly simulating the circulation beneath ice shelves is introduced in the non-linear free surface model NEMO. Its implementation into the NEMO framework and its assessment in an idealised and realistic circum-Antarctic configuration is described in this study.

Compared with the current prescription of ice shelf melting (i.e. at the surface), inclusion of open sub-ice-shelf cavities leads

- 15 to a decrease in sea ice thickness along the coast, a weakening of the ocean stratification on the shelf, a decrease in salinity of High Salinity Shelf Water on the Ross and Weddell Sea shelves and an increase in the strength of the gyres that circulate within the over-deepened basins on the West Antarctic continental shelf. Mimicking the overturning circulation under the ice shelves by introducing a prescribed meltwater flux over the depth range of the ice shelf base, rather than at the surface, is also assessed. It yields similar improvements in the simulated ocean properties and circulation over the Antarctic continental
- 20 shelf to those from the explicit ice shelf cavity representation. With the ice shelf cavities opened, the widely used "three equation" ice shelf melting formulation, which enables an interactive computation of melting, is tested. Comparison with observational estimates of ice shelf melting indicates realistic results for most ice shelves. However, melting rates for Amery, Getz and George VI ice shelves are considerably overestimated.

1 Introduction

- 25 Ice shelf melting, which accounts for 55% of the ice mass loss from Antarctica, is one of the main sources of freshwater input to the Antarctic coastal ocean. The net basal meltwater flux released to the Southern Ocean is estimated to be 1,500±237 Gt y⁻¹ (or 48±8 mSv), compared with 1,265±141Gt y⁻¹ (or 39±4 mSv) from iceberg calving (Rignot, et al., 2013). The total Antarctic mass discharge is thus similar to the 76 mSv due to surface atmospheric forcing (P-E) south of 63°S (Silva et al., 2006). The ice shelf melting contribution to the Southern Ocean freshwater forcing is different from the iceberg
- 30 melting and precipitation. Ice shelf melting is injected into the ocean at depth whereas precipitation is input at the surface

and icebergs inject melt water at a range of depths, but primarily in the top ~ 100 m. Therefore, the effect of ice shelf melting on coastal ocean stratification and circulation is very different from that of iceberg melt and precipitation.

The net ice shelf discharge (melting and calving) does not directly contribute to eustatic sea level change, because ice shelves are already floating, but does make a small steric contribution, because of the associated freshening (Jenkins and

- 35 Holland, 2007). However, the strong mechanical coupling between ice sheet and ice shelf controls the ice flux across the grounding line from the ice sheet. Modifications to the ice shelf geometry associated with changes in ice thickness or extent lead to changes in buttressing at the grounding line. A reduction in buttressing can trigger a speed up of the discharge from the ice sheet, a process that has been implicated in widespread mass loss from the Antarctic Ice Sheet (Scambos et al., 2004, Rignot et al., 2004, Favier et al., 2014). Therefore, understanding of ice shelf/ocean interaction is a key factor in advancing our understanding of the ice sheet contribution to sea level rise.
- Basal melting of ice shelves is driven by the properties of the water masses that are present over the continental shelves, enter the ocean cavities and reach the grounding line where they initiate melting. The associated input of buoyancy triggers an overturning circulation with inflow at depth and outflow along the ice shelf base that carries meltwater upward. The process is referred to as an ice pump when the ascending waters cause refreezing (Lewis and Perkin, 1986). Jacobs et al.
- 45 (1992) identified three modes of overturning, depending on the inflowing water mass, which could be either High Salinity Shelf Water (HSSW, Mode 1), modified forms of Circumpolar Deep Water (CDW, Mode 2), or less saline water masses that could collectively be referred to as Antarctic Surface Water (AASW, Mode 3). Mode 1 melt is low, because HSSW has a temperature close to the surface freezing point and can melt ice at depth only because of the lowering of its freezing point with increasing pressure. Mode 2 melt can be high if almost unmodified CDW has access to the sub-ice-shelf cavities. Mode
- 50 3 melt is intermediate and variable, depending on whether only the near-freezing core of ASSW, often designated Winter Water (WW), or the seasonally warmer upper layers can access the cavities. When the inflow has a temperature at or close to the surface freezing point (HSSW or WW), melting at depth is accompanied by partial refreezing at higher levels, as the falling pressure results in a rising freezing point temperature. In this case, the out flowing water mass produced is designated as Ice Shelf Water (ISW), and has a temperature below the surface freezing point. At the edge of the broad continental
- 55 shelves of the southern Weddell and Ross seas and along the Adelie Land coast, ISW mixes with CDW and HSSW to form Antarctic Bottom Water (Foldvik, et al., 1985, Williams et al., 2008) that contributes to the global overturning circulation. A modelling study (Hellmer, 2004) further suggests that 20 cm of the total sea ice thickness in the Ross and Weddell seas results from the cooling and freshening of shelf water by ice shelf melting.

To improve the representation of the Antarctic coastal ocean and global sea level rise in the coupled Ocean/Sea-ice model NEMO (Nucleus for European Modelling of the Ocean), ice-shelf/ocean interactions need to be properly included. In previous NEMO simulations, ice-shelf melt was uniformly distributed around the coast of Antarctica and input at the surface. Global conservation is an important issue, as the ocean/sea-ice model is also used as a component within Earth System Models. To tackle this issue, a z* vertical coordinate has been included within the NEMO framework (Madec, 2012), and the ice shelf module as well as the ice shelf parametrisation are developed using this vertical coordinates and considering

65 ice shelf melting as a mass flux.

This study is based on that of Losch (2008) (hereafter L08), describing the development of an ice shelf module within MITgcm. We follow a similar strategy to introduce ice shelf-ocean interactions into the NEMO framework (Madec, 2012). The work is a first step towards adding an ice sheet component and its interaction within NEMO, and including these interactions within climate models such as IPSL (Dufresne, et al., 2013), the Hadley Centre models (Hewitt, et al., 2011 and

70 Hewitt, et al., 2016), EC earth (Hazeleger, et al., 2010), CNRM (Voldoire, et al., 2013) and CMCC (Scoccimarro, et al., 2011).

Ice shelves range in size from the giant Ross ice shelf $(500,000 \text{ km}^2)$ to the tiny Ferrigno ice shelf (117 km^2) . This means that current global ocean model configurations are not able to resolve explicitly all the ice shelf cavities. For this reason, a simple way to include unresolved ice shelf melting in the ocean model that mimics the circulation driven by ice shelf melting at

75 depth is also presented here.

The paper is structured as follows: firstly, the NEMO model (Sec. 2.1), as well as the ice shelf module (Sec. 2.2 and Sec. 2.3), are described, then idealized experiments are presented to validate the ice shelf module (Sec. 3) and ice shelf parametrisation (Sec. 4), followed by its application to a realistic circum-Antarctic configuration at 0.25° resolution (Sec. 5). The sensitivity of the ocean and sea ice properties to the inclusion of the ice shelf cavity (Sec. 5.3 and Sec 5.4), the effect of

the ice shelf cavity parametrisation under prescribed ice shelf melting (Sec. 5.5) and the resulting meltwater flux (Sec. 5.6) are then discussed. Finally, in a summary section (Sec. 6), the major results as well as the remaining issues are highlighted, and we conclude with details of code availability.

2 Model description

2.1 Ocean model

NEMO is a primitive equation ocean model, and this study uses version 3.6 of the code. The variables are distributed on an Arakawa C-grid; i.e. the scalar point (temperature, salinity) is defined on the centre of the cell and the vector points (zonal, meridional, vertical velocity) are defined on the centre of each face (Arakawa, 1966). We also make use of the time varying z* vertical coordinate; i.e. the variation of the water column thickness due to sea-surface undulations is not concentrated in the surface level, as in the z-coordinate formulation, but is distributed over the full water column (Adcroft and Campin, 2004).

A complete description of the schemes and options available in NEMO is available in the documentation (Madec, 2012). A full description of the configurations used in this study is presented in Sect. 3.1 for the idealised configuration and in Sect. 4.1 for the realistic configuration.

2.2 Ice shelf/ocean interaction description

95 2.2.1 Ocean dynamics

The z* vertical coordinate can be used with a sea ice model (Campin et al., 2008) in NEMO (Madec et al., 2012). However, modelling the ocean circulation within an ice shelf cavity in z* coordinates requires some modification of the existing code. Beneath sea ice, the number of ocean levels is kept constant, and the levels are squeezed between the bottom surface of the ice and the seabed. The resulting pressure gradient error term is small because the ratio of sea ice thickness to total water

100 column thickness is small and almost spatially constant. Within an ice shelf cavity, a z* coordinate used as a surface following coordinate will face the same limitation as terrain following coordinates, especially along the ice shelf front. The pressure gradient error will be large, particularly at the vertical ice front, and the tiny vertical cell thickness where the water column is thin will limit the stable time-step that is achievable.

To avoid these issues, we follow the idea of Grosfeld et al. (1997) for an s-coordinate model. All cells between the surface (z=0) and the ice shelf base are masked at the model initialisation stage. By masking the ice shelf cells, the z* iso-surfaces are close to horizontal and the associated slopes are small, even at the ice front. Outside the ice shelf cavity, the definition of the cell thickness and the computation of the pressure gradient are not changed compared with the original code that follows Adcroft and Campin (2004). Within the cavities, the ice shelf thickness and the associated masked cells are constant over time, so the z* iso-surfaces are defined as:

110
$$Z_w(1) = 0$$
 (1)

$$if \ k < k_{isf}, \ Z_w(k) = \sum_{kz=1}^{k-1} dz_{0,T}(kz)$$
(2)

$$if \ k \ge k_{isf}, \ Z_w(k) = \sum_{kz=1}^{k_{isf}-1} dz_{0,T}(kz) + \sum_{kz=k_{isf}}^{k-1} dz_{t,T}(kz)$$
(3)

$$dz_{t,T}(kz) = dz_{0,T}(kz) \left(1 + \frac{\eta}{H}\right)$$
(4)

with Z_w the depth of the w interface (interface between 2 cells along the z axis of the Arakawa C-grid, positive down), $dz_{t,T}$ 115 the vertical level thickness at time t, $dz_{0,T}$ the vertical level thickness at time 0, k the model level (k=1 is the first level), η the sea surface height (positive up), H the total water column thickness (sum of all the wet cell vertical thicknesses at time 0) and k_{isf} the first wet level.

The pressure p at a depth z is computed in a standard way (Beckmann et al., 1999; L08). We assume the ice shelf to be in hydrostatic equilibrium in water at the reference density ρ_{isf} , taken to be the density of water at a temperature of -1.9° C

120 (freezing point) and a salinity of 34.4 PSU (mean salinity over the Antarctic continental shelves). The total pressure at any depth is computed from the sum of the ice shelf load and the pressure due to the water column above that depth. The pressure gradient is formulated as suggested by Campin et al. (2004) for z* coordinate models:

$$p(z) = \int_{z_{isf}}^{0} \rho_{isf} g dz + \int_{z}^{z_{isf}} \rho g dz$$
(5)

$$\nabla_z p(z) = \nabla_{z*} p(z) + \rho g \nabla_{z*} z \tag{6}$$

125 where p(z) is the pressure at depth z, ρ is the water density at depth z, and z_{isf} is the ice shelf draft. The hydrostatic pressure gradient at a given level, k, (first term in Eq. 6) is computed by adding the pressure gradient due to the ice shelf load (defined as the first term of Eq. 5) to the vertical integral of the in-situ density gradient along the model level from the surface to that level.

In this study, we assume the ice shelf to be in an equilibrium state (i.e. the ice shelf draft is temporally constant), so that all

130 the ice melted by the ocean is assumed to be replaced by the seaward advection of new ice. The pressure of the ice shelf on the ocean therefore stays constant, but the ocean volume increases due to ice shelf melting. Dealing with an evolving ice shelf thickness is beyond the scope of this paper.

Representation of the bottom topography is difficult in z coordinate models. The partial cell scheme allows a more accurate representation of bottom topography through the use of partially wet cells (Adcroft et al., 1997). Solutions obtained with this

- 135 scheme compare favourably with those obtained with sigma coordinate models (Adcroft et al., 1997) and also with more realistic solutions (Barnier et al., 2006). Following L08, we apply the partial cell scheme developed for the bottom topography to the top cells beneath the ice shelf base. For stability reasons, the minimum thickness of the bottom and top cells is set to the smaller of 25 m or 20% of a full cell. However, representation of density driven flow in a z coordinate model (even with partial cells), like the overflow, is challenging (Legg et al., 2006). Thus the representation of the buoyancy driven flow along an ice shelf base is expected to present analogous problems.
- Where the water column is thinner than two cells, vertical circulation cannot be represented. In order to simulate the overturning circulation generated by ice shelf melting in such regions, we modify the bathymetry or the ice shelf draft sufficiently to open a new cell in the water column. In places where the cavity is thin and the slopes of the bathymetry and ice shelf draft are steep, it would sometimes be necessary to create more than one new cell in order to open a minimum of
- 145 two cells at the velocity points (at the centre of the cell faces on the Arakawa C-grid). Rather than making such extensive modifications to the topography, we regard the combination of vertical and horizontal resolution as too coarse to represent the sub-ice cavity geometry in these places, and instead we ground the ice shelf. Consequently some ice shelves have a reduced area.

For regional configurations with open boundaries, the normal barotropic velocity around the boundary at each time step is

150 corrected to force the total volume to be constant. The correction ensures that the net inflow (the combination of inflow at the open boundary, runoff, ice shelf melting and precipitation) and net outflow (the combination of outflow at the open boundary, ice shelf freezing and evaporation) are balanced.

2.2.2 Thermodynamics

Two formulations of the ice shelf melt rate are available: a simple one used in the idealised cases, for consistency with earlier studies and the Ice Shelf-Ocean Model Intercomparison Project (ISOMIP), and a more sophisticated one used in the realistic configuration. For the idealized study, the heat flux and the freshwater flux (negative for melting) resulting from ice shelf melting/freezing are parameterized following Grosfeld et al. (1997). This formulation is based on a balance between the vertical diffusive heat flux across the ocean top boundary layer and the latent heat due to melting/freezing:

$$Q_{h} = \rho c_{p} \gamma (T_{w} - T_{f})$$

$$q = \frac{-Q_{h}}{L_{f}}$$
(8)

where $Q_h(W m^{-2})$ is the heat flux, q (kg s⁻¹ m⁻²) the freshwater flux, L_f the specific latent heat, T_w the temperature averaged over a boundary layer below the ice shelf (explained below), T_f the freezing point computed from Millero (1978) using the pressure at the ice shelf base and the salinity of the water in the boundary layer, and γ the thermal exchange coefficient. Hereafter, Eq. (7) and (8) are referred to as the ISOMIP formulation.

For realistic studies, the heat and freshwater fluxes are parameterized following Jenkins et al. (2001, Eq. 24). This formulation is based on three equations: a balance between the vertical diffusive heat flux across the boundary layer and the latent heat due to melting/freezing of ice plus the vertical diffusive heat flux into the ice shelf (Eq. 9); a balance between the vertical diffusive salt flux across the boundary layer and the salt source or sink represented by the melting/freezing (Eq. 10); and a linear equation for the freezing temperature of sea water (Eq. 11, Jenkins, 1991):

$$c_{p}\rho\gamma_{T}(T_{w} - T_{b}) = -L_{f}q - \rho_{i}c_{p,i}\kappa\frac{T_{s} - T_{b}}{h_{isf}}$$

$$\rho\gamma_{S}(S_{w} - S_{b}) = (S_{i} - S_{b})q$$

$$T_{b} = \lambda_{1}S_{b} + \lambda_{2} + \lambda_{3}z_{isf}$$

$$(11)$$

160

165

170

T_b is the temperature at the interface, S_b the salinity at the interface, γ_T and γ_S the exchange coefficients for temperature and 175 salt, respectively, S_i the salinity of the ice (assumed to be 0), h_{isf} the ice shelf thickness, ρ_i the density of the ice shelf, c_{p,i} the specific heat capacity of the ice, κ the thermal diffusivity of the ice and T_s the atmospheric surface temperature (at the ice/air interface, assumed to be -20°C). The linear system formed by Eq. 9, Eq. 10 and the linearised equation for the freezing temperature of sea water (Eq. 11) can be solved for S_b or T_b. Afterward, the freshwater flux (q) and the heat flux (Q_h) can be computed. γ_T and γ_S are velocity dependent (Jenkins et al., 2010) and can be written as:

180
$$\gamma_T = \sqrt{C_d} u_w \Gamma_T$$
 (12)
 $\gamma_S = \sqrt{C_d} u_w \Gamma_S$ (13)

with u_w the ocean velocity in the top boundary layer, C_d the drag coefficient and $\Gamma_{T/S}$ a constant. The choices of the thermal Stanton number ($\sqrt{C_d}\Gamma_T = 0.0011$) and the diffusion Stanton number ($\sqrt{C_d}\Gamma_S = 3.1 \times 10^{-5}$) are based on the recommendation of Jenkins et al. (2010). The drag coefficient is chosen to be 1.0×10^{-3} . This value lies within the range

185

used in the literature. However, there are no direct measurements of the drag coefficient beneath an ice shelf. Dansereau et al. (2014) highlight that the range of values used for the top drag coefficient is large (from 1.0×10^{-3} to 9.7×10^{-3}). Furthermore, uncertainties in the Stanton numbers are also large, as the study used to determine their values (Jenkins, et al., 2010) is based on data from a single borehole. Parameter values used in Eqs. 7-12 are defined in Table 1. Hereafter, Eqs. 10-

12 are referred to as the "three equation" ice shelf melting formulation. Unlike in more sophisticated models of the freezing

190 process (Galton-fenzi et al., 2012), the parameters used in the "three equation" formulation are not dependent of the surface state (freezing or melting) and the freezing only occurs at the ice/ocean interface.

Following L08, in the idealized experiments, the ice shelf forcing is applied as an effective heat flux and a virtual salt flux (no ocean volume change). For realistic configurations, the velocity divergence at the ice shelf base is adjusted in order to apply the ice shelf melting as a volume flux of freshwater at the freezing point temperature.

- 195 L08 shows that z coordinate models with partial cells generate a noisy melt rate pattern due to the variation of the top cell thickness. The melt rate is proportional to the difference between the in-situ basal temperature and in-situ temperature in the first wet cell. Because the largest cells cool down more slowly than the thinnest cells, for a given initial basal temperature, the melt rate in the thickest cells is larger than in the smallest cells. Following L08, the noise due to the spatially varying size of the top cells is suppressed by computing T_w and S_w in Eq. 7, 9 and 10 as the mean value over a constant thickness,
- 200 assumed to represent the top boundary layer thickness (H_{TBL} , i.e. properties are averaged over the cells entirely included in the top boundary layer and a fraction of the deepest wet cell partly included in the top boundary required to make up the constant H_{TBL}). The top ocean velocity u_w is defined as the velocity magnitude derived from the mean zonal/meridional velocity at U/V points within the top boundary layer averaged at T points. The heat and freshwater fluxes are distributed over the same constant thickness. If the first wet cell is thicker than the specified top boundary layer thickness, H_{TBL} is set to
- 205 the top cell thickness. A complete description of this parametrisation is available in L08. Using z^* instead of pure z coordinates does not alter the noise seen in the melt rate. Therefore, the parametrisation proposed by L08 is applied in each simulation used in this study. H_{TBL} is set to a default value of 30 m, but different values are used for the simulations with various vertical resolutions, as presented in Table 2.

2.3 Simplified representation of ice shelf melting

whatever the grounding line depth is.

210 Global ocean model configurations are typically unable to resolve all the ice shelves around Antarctica. Despite their limited extent, the smaller ice shelves nevertheless make a significant contribution to the total meltwater flux from the ice sheet. We therefore need a way to mimic the impact of unresolved cavities on the ocean.

Beckmann and Goosse (2003, hereafter BG03) suggest a simple parametrisation for the melting beneath an ice shelf and prescribe the input of melt water at the ocean level corresponding to the base of the ice shelf (Fig. 1c). One of the mainissues with this parametrisation is that, for the same ice shelf melting, the effect on the ocean dynamics will be the same

The idea tested in this paper is to spread the freshwater due to ice shelf melting evenly between the grounding line depth and the depth of the calving front. In this case, the model creates its own plume along the vertical wall (Fig. 1d, no cavity in this case) and thus an overturning between the grounding line depth and the equilibrium depth (the depth where the density of the

220 plume is equal to the density of the ambient water). Fig. 1a and 1b are discussed in Sec. 5.2.

In this part of the study we focus on how to inject the observed ice shelf meltwater flux into the ocean model. Therefore, the ice shelf melting is prescribed and the heat flux is derived from the freshwater flux using Eq. 8. The computation of the melt rate from the off-shore ocean properties and ice shelf geometry could be included using the BG03 parametrisation or some adaptation of the Jenkins (2011) plume model. The parametrisation tested in this study is kept as simple as possible for ease

225 of use in a wide range of applications. Further testing of other interactive melt parametrisations or fresh water distributions that are functions of the ice shelf geometry or melt rate is beyond the scope of this study.

3 Academic case

In order to compare the sub-ice shelf cavity capability in NEMO with that in other models, the idealized configuration used in this study is the one described in the Ice Shelf-Ocean Model Inter comparison Project (ISOMIP). ISOMIP is an open, international effort to identify systematic errors in sub-ice-shelf cavity ocean models and the reference configuration is based

230

) international effort to identify systematic errors in sub-ice-shelf cavity ocean models and the reference configuration is based on a very simple setup, briefly described below.

3.1 ISOMIP setup

The ISOMIP setup follows the recommendations of the inter comparison project for experiment 1.01 (Hunter, 2006). The geometry is based on a closed domain with a flat seabed fixed at 900 m. The grid extends over 15° in longitude, from 0 to

235 15°E with a resolution of 0.3°, and 10° in latitude, from 80°S to 70°S with a resolution of 0.1°. The spatial resolution ranges from 6 km at the southern boundary to 11 km at the northern boundary. The whole domain is covered with an ice shelf, and includes no open ocean region. The ice shelf draft is uniform in the east-west direction, is set at 200 m between the northern boundary and 76°S and deepens linearly south of 76°S down to 700 m at the southern boundary. The water is initially at rest and has a potential temperature of -1.9° C and a salinity of 34.4 PSU. No restoring is applied to either the temperature and

240 salinity.

245

The vertical resolution is uniform and fixed at 30 m, allowing a direct comparison with the results of L08. The density is computed using the polyEOS80-bsq function. It takes the same polynomial form as the polyTEOS10 function (Roquet et al., 2015), but the coefficients have been optimized to accurately fit EOS-80 (Roquet, personnal communication). The melt formulation is the "ISOMIP" one. All the results presented are taken from day 10 000 at which time the system is in steady state.

3.2 Model comparison

The ISOMIP experiment has been carried out with many models using different vertical coordinates during the last 10 years, including ROMS¹, OzPOM², MITgcm (Losch, 2008) and POP (Asays-Davis, 2012). All these models agree on a common

¹ http://www.ccpo.odu.edu/~msd/ISOMIP/

² http://staff.acecrc.org.au/~bkgalton/ISOMIP/

circulation and melt pattern. The melting and freezing along the base of the ice shelf drives an overturning circulation of

- 250 about 0.1 Sv. Associated with the meridional overturning circulation, all the models generate a cyclonic gyre with a western boundary current beneath the sloping ice shelf of about 0.3 Sv. This horizontal circulation drives water that is warmer than the freezing point into the south-eastern part of the cavity. The inflow of warm water causes melting at the ice shelf base that is concentrated along the eastern and southern boundaries. On the western side of the ice shelf cavity, the boundary current advects colder water towards the ice front. Shoaling of the ice shelf base causes super-cooling of the water in contact with
- 255 the ice and thus drives freezing. A detailed discussion of this circulation can be found in Grosfeld et al. (1997). The maximum melting/freezing rates are model dependent. The range is 0.7 1.8 m y⁻¹ for the maximum freezing rate and 0.7-2.4 m y⁻¹ for the maximum melting rate.

The NEMO response to the ISOMIP setup (simulation I_30M) is shown in Fig. 2. It is similar to that previously simulated with a z coordinate model (L08). The strength of the overturning circulation is 0.11 Sv. The transport of the western

260 boundary current generated by the cyclonic gyre beneath the sloping ice shelf is 0.32 Sv. The pattern of melting and freezing is similar to that in L08. The melting occurs, as expected, in the south-eastern corner with a maximum of 2.7 m y⁻¹ and the freezing takes place beneath the western boundary current with a maximum of 1.9 m y⁻¹. The low noise is the result of the L08 parametrisation (Fig. 2). In simulations without this parametrisation (not shown) the noise in the melt pattern is as shown in L08.

265 **3.3 Sensitivity of ocean circulation to the vertical resolution**

Depending on the scientific question to be addressed, the ocean models commonly used have very different vertical resolutions, ranging from 1m to 100m. The representation of the top boundary layer is strongly affected by the choice of vertical resolution. To evaluate the impact of this choice on the ocean circulation beneath the ice shelf, nine simulations with vertical resolution ranging from 5m (I_5M) to 150m (I_150M) have been carried out (Table 2).

- 270 The choice of vertical resolution and Losh H_{TBL} strongly affects the ice shelf melting. When H_{TBL} is tied to the vertical resolution, finer resolution gives lower melting. Under melting conditions, a thin, fresh and cold top boundary layer appears in the top metres of the ocean next to the ice shelf base. With finer vertical resolution, a thinner and colder top boundary layer can be resolved, resulting in weaker melting (Fig. 3a). Our sensitivity experiments show a maximum melt rate 4 times higher in the I_150M simulation (4.3 m y⁻¹) and 3 times higher in the I_60M simulation (3.1 m y⁻¹) than in the I_5M
- 275 simulation (0.9 m y⁻¹). In analogous experiments, L08 found a similar sensitivity, with maximum melting 3 times larger at 45 m resolution than at 10m resolution. However, when H_{TBL} is kept constant (I_5M30M, I_10M30M and I_30M), the total melt is insensitive to the vertical resolution. The total melt at high vertical resolution (5 m or 10 m) with a 30 m Losh top boundary layer thickness (respectively I_5M30M and I_10M30M) is converging toward I_30M (Fig. 3a). This suggests that a more physical definition of H_{TBL} (based on stratification, melt rate, etc ...), rather than a constant H_{TBL}, could significantly
 280 shares the melt rate in a high resolution models (although investigation of this is housed the second of the paper).
- 280 change the melt rate in a high resolution models (although investigation of this is beyond the scope of the paper).

With very coarse resolution (I_100M/I_150M), the model is unable to represent a top boundary layer at all and the total melting saturates. Total melting is 20% smaller in the I_5M simulation than in both the I_100M and I_150M simulations, which have the same total melt (Fig. 3a). With variable vertical resolution (I_31L, I_46L and I_75L), such as is typically used in global configurations of NEMO (Timmermann et al, 2005, Drakkar group, 2007 and Megann et al., 2014), the

- 285 coarsest resolution in the cavity seems to determine the total melt. This is because more than 50% of the melting occurs between 500 m and 700 m depth where the resolution is coarsest (not shown). This could be an issue for modelling ice shelf melting with the standard configuration used for climate applications because Dutrieux et al. (2013) show that, for some ice shelves with high melt rates, most of the melt may occur over a small area close to the grounding line, where the resolution is coarsest.
- 290 The vertical resolution also has a major impact on the noise pattern (Fig. 4). As the noise in the melt pattern is closely linked with variations in the thickness of the first wet cell, the finer the vertical resolution, the weaker the noise. In contrast, the barotropic stream function and the overturning circulation in the cavity are not altered by any choice of vertical resolution between 5 and 150 m (Fig. 3b). One of the reasons could be that with the bulk formulation of melting used in the ISOMIP simulations, there is no direct link between the ocean current velocity at the ice-shelf/ocean interface and the
- 295 melt rate, because the thermal exchange coefficient is defined to be a constant.

4 Ice shelf cavity parametrisation

300

While the ice shelf module as described so far works well in idealised cases, for a wider range of applications (including ice shelves of varying extent at all likely horizontal resolutions) we also need the capability of representing the impact of circulation and melting within unresolved cavities. In this section, we investigate the ability of our ice shelf cavity parametrisation to mimic the circulation and water mass properties produced by the full cavity simulation, and compare the results with those produced by the parametrisation of BG03. Both parametrisations are evaluated in an idealised

configuration derived from the ISOMIP setup.

The geometry is the one for ISOMIP experiment 2.01, which is the same as that for ISOMIP experiment 1.01 except in the top 200 m, where the flat ice shelf is replaced by open water (Fig. 5a). The simulations are initialised with a warm linear

305 profile typical of conditions on the continental shelves of the Amundsen and Bellingshausen Seas (Fig. 6 in Asay Davis et al., 2016 with constant value between 720 m and 900 m). Radiative open boundary conditions are applied at the northern boundary (Treguier et al., 2001). The vertical eddy viscosity and diffusivity, in unstable conditions, are increased from 0.1 m².s⁻¹ to 10 m².s⁻¹ to reduce the noise generated along the ice shelf front.

Three experiments are run for 30 years: one with the ice shelf cavity open (A_ISF, Fig. 1b), but with a steady pattern of basal

310 melt/freeze imposed; another with the open ocean circulation driven by the cavity parametrisation of BG03 (A_BG03, Fig. 1c); and a third with the cavity parametrised as outlined in Sect. 2.3 (A_PAR, Fig. 1d). In all these experiments the same

heat and freshwater fluxes are applied, derived from the basal melt/freeze pattern obtained in the last month of a dedicated 30-year run with explicit ice shelf melting calculated using the "ISOMIP" formulation.

- A_ISF drives a deep inflow toward the ice shelf, and corresponding outflow in the top 400 m toward the open ocean, of 0.9 315 Sv at the northern boundary (Fig. 5a). In a stratified ocean, this circulation has a crucial effect on the total amount of heat advected toward the ice shelf, on the properties of the water drawn into the overturning circulation and on the overall stratification in the basin. In A_BG03 the overturning is too weak (0.6 Sv compared with 0.9 Sv in A_ISF) and too shallow (200 m compared with 400 m in A_ISF). Consequently, the water masses drawn into the overturning come from a different depth and have different T/S properties, and the resulting stratification is too strong, with colder surface waters and warmer
- 320 deep waters (Fig. 5c). In A_PAR, because the freshwater flux is distributed over the same depth range as in A_ISF (between 200 m and 700 m), the vertical extent of the overturning and the water masses drawn in are the same in both A_PAR and A_ISF. The result is a circulation on the shelf that is similar in depth and magnitude and a stratification that is similar in strength to those simulated in A_ISF (Fig. 5b).

With far-field conditions typical of the cold, salty continental shelves of the Ross and Weddell seas, where the water column

325 is well mixed by brine rejection from growing sea ice in winter and less heat is available at depth, the differences in the stratification resulting from the two parametrisations and the simulation with the open ice shelf cavity should be smaller.

5 Real ocean application

In the ISOMIP test cases, the ocean circulation in the cavity compares well with that simulated by other models. Furthermore, the suggested parametrisation of ice shelf melting mimics well the circulation and water properties generated 330 by the presence of an open ice shelf cavity. Nevertheless, the bathymetry and ice shelf draft are smooth in these idealised cases and the heat transfer coefficient is constant, so the favourable comparison with other models in the idealised ISOMIP setup between models as well as the good match between the idealised A_ISF and A_PAR experiment might not be reproduced in a realistic configuration. In the next section, we assess both the explicit ocean cavity representation and the cavity parametrisation in a realistic circumpolar configuration.

335 **5.1 Antarctic configuration setup**

ePERIANT025 is a circum-Antarctic configuration based on the PERIANT025 configuration (Dufour et al., 2012) covering the ocean from 86.5°S to 30°S, using a ¹/₄° isotropic Mercator grid. A feature of the Mercator grid is that the mesh spacing reduces with decreasing distance from the South Pole, so that the farthest south grid boxes strongly constrain the model time step. To maintain a model time step equal to that used in current global ¹/₄° configurations, the Mercator grid is replaced

340 south of 67°S with 2 quasi-isotropic bipolar grids, one for the Bellingshausen, Amundsen and Ross sea sector and one for the Weddell sea sector (Fig. 6). Each sector is built following the semi-analytical method used to create the tripolar ORCA grid north of 22°N (Madec and Imbard, 1996). The effective resolution is 13.8 km at 60°S, increasing to 3.8 km at 86.5°S, where a pure Mercator grid would have a resolution of 2.2 km. The model uses 75 vertical levels with thicknesses varying from 1m at the surface to 200 m at 6000 m depth, giving a vertical resolution ranging from 10 to 150 m beneath the ice shelves. See

The bathymetry used for the model domain north of the Antarctic continental shelf is that described by Megann et al., (2014). Over the Antarctic continental shelves the IBCSO data set (Arndt et al., 2013) is used. The two bathymetry data sets are merged between the 1000 m and 2000 m isobath along the Antarctic continental slope. Under the ice shelves, bathymetry (included in the IBCSO data set) and ice draft are taken from BEDMAP 2 (Fretwell et al., 2013). The resulting model

Sec. 3.3 for the effect of this resolution on ice shelf melting in an idealised case.

345

- bathymetry is shown in Fig. 6. Note that for some ice shelves, Fretwell et al. (2013) enforced floatation by lowering the sea bed. In addition, we impose a minimum of two vertical grid cells within the ocean cavities so that an overturning cell can develop. Where necessary, either the bathymetry or the ice shelf draft, depending on the local configuration, is modified to fit the criterion. If more than one cell has to be modified to fit the water column criterion, the entire water column is masked. Using this procedure, Totten and Dalton (Moscow University in Rignot et al., 2013) ice shelves and the deepest part of Amery Ice Shelf are almost fully masked.
- Other choices (the momentum advection, tracer advection, diffusion, viscosity, vertical mixing, double diffusion, bottom friction, bottom boundary layer and tidal mixing parametrisations) are as used in Megann et al., (2014). For the sea ice we use the Louvain-la-Neuve sea-ice model LIM2 (Fichefet and Morales, 1997) with ice rheology based on an elasto-visco-plastic law as described in Bouillon et al. (2013).
- 360 The geothermal heat flux is assumed to be constant and set to 86 mW/m² (Emile-Geay and Madec, 2010), while the internal wave energy used in the tidal mixing parametrisation (0 under the ice shelf for simplicity) is derived from the tide model FES 2012 (Carrère et al., 2012). Sea surface salinity restoring is applied north of 55°S, river runoff comes from Dai and Trenberth (2002), and iceberg melting based on Rignot et al. (2013) is evenly distributed at the surface along the Antarctica coast. Ice shelf melt is applied either into the open cavities, at depth following our parameterisation, or as surface runoff. The
- 365 total ice shelf melt in each individual cavity is either interactively computed using the "three equation" formulation or prescribed following the Rignot et al. (2013) estimates.

Radiative boundary conditions are applied at the northern open boundary (Treguier et al., 2001) using velocity, temperature and salinity data from a global NEMO ORCA025 simulation (Barnier et al., 2012) forced by the DFS5.2 atmospheric forcing developed by the DRAKKAR project. To minimize inconsistency, the model is also driven by the same DFS5.2 atmospheric

370 forcing. The methodology applied to build the DFS forcing series is described in Brodeau et al. (2010), and the details of the DFS5.2 are given in a report by Dussin et al. (2016). Initial conditions come from the World Ocean Atlas 2013 (Locarnini et al., 2013 and Zweng et al., 2013). The model is run for 10 years starting in 1979 and ending in 1988, and the first order response is investigated using output from the last year of the simulation.

5.2 Experiment description

- 375 In order to evaluate both the explicit ice shelf module (Sect. 2.2) and the improved parametrisation (Sect. 2.3) in this realistic case, 4 simulations are run:
 - R_noISF: a simulation without ice shelf cavities. Both the ice shelf freshwater flux and the latent heat flux associated with melting of the ice are prescribed at the surface (Fig. 1a).
 - R_ISF: a simulation with explicit ice shelf cavities (Fig. 1b), but where both the melt rate of the ice shelves and the latent heat flux at the ice shelf/ocean interface are specified.
 - R_PAR: a simulation without ice shelf cavities (Fig. 1d). Both freshwater and latent heat fluxes from the ice shelves are uniformly distributed along the calving front from its base down to the grounding line depth, or the seabed if it is shallower.
 - R_MLT: a simulation with explicit ice shelf cavities and interactive melt rates computed by the "three equation" formulation (Fig. 1b).

For R_ISF, R_noISF and R_PAR the same total inputs of freshwater and latent heat are prescribed for each ice shelf and the fluxes are constant over time; only the location of the input changes. The melting pattern used in R_ISF is provided by the simulation R_MLT, while the magnitude is scaled so that the total for each ice shelf matches that from Rignot et al. (2013). The associated latent heat flux is derived from the melt rate using Eq. 8.

390 Initially, results from R_noISF and R_ISF are used to evaluate the sensitivity of the ocean and sea ice properties to the presence of ice shelf cavities in a control setup with prescribed melting. Next, results from R_PAR are compared with those from R_noISF and R_ISF in order to evaluate and validate the ice shelf parameterisation in a realistic case. Finally, results from R_MLT are used to evaluate the modelled ice shelf melting in our circum-Antarctic configuration using the "three equation" ice shelf melting formulation.

395 5.3 Sensitivity of ocean properties to the ice shelf cavities

In both R_noISF and R_ISF, large-scale open ocean features are well represented. Simulated ACC transport (135 Sv) and Weddell gyre transport (56 Sv) are similar and compare well with the observations of 137 Sv for the ACC transport (Cunningham et al., 2003) and 56 Sv for the Weddell gyre transport (Klatt et al., 2005). Temperature and salinity properties north of the continental shelves are also similar in both simulations and compare reasonably with WOA2013 (Figs. 7-8). In

400 contrast, the presence of ice shelf cavities in R_ISF substantially affects the ocean properties and dynamics in the coastal Antarctic seas (Figs. 7, 8 and 10).

Over the Bellingshausen and Amundsen seas, the input of freshwater at the surface in R_noISF leads to strong stratification in the upper 250 m, weak stratification below (Fig. 9), a weak and shallow vertical circulation (maximum overturning is 0.01 Sv at 70 m depth) and a weak barotropic circulation over the continental shelf (Fig. 10). In R_ISF, the input of buoyancy at

405 the ice shelf base activates the buoyancy-forced overturning, driving upwelling along the ice shelf/ocean interface. The

385

380

overturning circulation entrains 0.23 Sv of a mix of ambient water (CDW) and meltwater along the ice shelf base. This upwelling generates a barotropic circulation that follows the f/h contours over the Amundsen and Bellingshausen sea continental shelf (Figs. 10a,c) as explained in Grosfeld et al. (1997). The resulting mixture of CDW and meltwater stabilises at an equilibrium depth between 400 m and 60 m (not shown). The upwelling of CDW into the surface mixed layer weakens

410 the thermohaline stratification and warms and salinifies the surface layer. These changes in ocean dynamics on the shelf lead to a more realistic continental shelf temperature and salinity distribution (Figs. 7-8) and stratification (Fig. 9) in R_ISF compared with R_noISF.

In Pine Island Bay and elsewhere on the Amundsen and Bellingshausen Sea shelves, the bottom water properties in the overdeepened basins are determined by the properties in the open ocean at the sill depth (Walker et al., 2007) close to the shelf

- 415 break. So the bottom temperature bias present in R_ISF could be strongly affected by the model bias in the ACC, the possible sources of which are beyond the scope of this paper. In R_noISF, as the overturning is not activated, there is no process to flush the bottom water trapped in the over-deepened basins, so the waters there are not affected by external forcing, and the bottom properties still match the initial conditions after ten years of the run (Fig. 9).
- Over the Ross and Weddell Sea continental shelves, the cold, salty HSSW in R_noISF matches the observations and spreads northward across the shelf break toward the open ocean. In R_ISF, the HSSW produced is too fresh (-0.2 PSU, Fig. 8). Weak winds in the atmospheric forcing (Dinniman et al., 2015), in addition to a fresher coastal current (Nakayama et al., 2014), the opening of a new pathway for HSSW circulation beneath the ice shelves (Budillon et al., 2003; Nicholls et al., 2009), mixing of HSSW with light surface waters all year long, and a deficiency of the sea-ice model in representing coastal polynyas could all help to explain the absence of HSSW in R_ISF.

425 **5.4 Sensitivity of sea ice properties to the ice shelf cavities**

Winter sea ice extent compares well with the 18.3 million km^2 estimated from satellite observations (Comiso, 2000) in both R_ISF (18.2 million km^2) and R_noISF (18.4 million km^2). The position of the sea-ice edge, being too far south in the Amundsen Sea and too far north in the Weddell Sea and around East Antarctica in both simulations, is not changed significantly by the presence of ice shelf cavities (Fig. 11).

- 430 Over the warm continental shelves of the Amundsen and Bellingshausen seas, sea ice is thicker in the R_noISF than in the R_ISF simulation (+1 metre, Fig. 11a). In R_noISF, because the freshwater and the latent heat sink from the melting of land ice are prescribed at the surface, the consequent freshening and cooling of the surface waters considerably enhances the formation of sea ice. This leads to very thick sea ice in R_noISF, greater than 3 m locally (Fig. 11c). In R_ISF, the overturning circulation driven by melting at the ice shelf ocean interface entrains warm CDW and mixes it into the surface
- 435 layer. This upward heat flux decreases the sea ice formation and has a direct effect on sea ice thickness (Fig. 11a). Over the cold continental shelves of the Ross and Weddell seas and around the coast of East Antarctica, sea ice thickness differences between R ISF and R noISF are much smaller, typically about 20 cm (Fig. 11). The ocean is well mixed and the

shelf water temperature is close to the freezing point (Fig. 7). So the amount of heat entrained into the buoyant overturning along the ice shelf base is smaller, as is the impact on sea ice.

440 Comparison with spring sea ice thickness estimates derived from sea-ice freeboard and snow thickness measurements (Fig. 11d, Kurtz and Markus, 2012) shows that sea ice thickness in R_ISF is closer to observation by about 1 m over the warm shelves of West Antarctica. Over the cold shelves, the modelled sea-ice thicknesses are similar in both simulations (less than 20 cm differences) and comparable with the observations, which are subject to +/- 40 cm uncertainties.

5.5 Assessment of the simplified ice shelf representation

- The implementation of the ice shelf cavities in a realistic configuration showed a great improvement in the circulation on the Antarctic continental shelves, especially in the Amundsen and Bellingshausen seas. However, many climate models lack the horizontal and vertical resolution needed to represent all these cavities. Our parametrisation described in Sect 2.3 has been developed to address this issue. The evaluation of our parametrisation in a simple idealised case showed very encouraging results. Here, by comparing R_ISF and R_noISF with R_PAR, we evaluate the parametrisation for all ice shelves of the Southern Ocean.
 - Over the warm shelves of West Antarctica, R_PAR reproduces well the R_ISF shelf properties and circulation (Fig. 12a-b and Fig. 10). Critically, the prescription of the ice shelf meltwater flux at depth drives an overturning circulation and spins up the associated gyres within the over-deepened basins. The magnitudes of the gyres are similar between the R_ISF and the R PAR simulations (Fig. 10b,c). Shelf water properties generated by R ISF are better reproduced by R PAR than by
- 455 R_noISF over all the West and East Antarctic shelves (Fig. 12a-d). Over the Amundsen shelf, R_PAR also decreases the stratification and improves the mean temperature and salinity profiles compared with R_noISF (Fig. 9). Over the Ross and Weddell sea shelves, HSSW produced in R_PAR is saltier than in R_ISF (+0.1 PSU). The salinity gradient between the salty western side and the fresher eastern side of the shelves is larger than in R_ISF (Fig. 12c) and
- 460 Ross (Budillon et al., 2003) and Filchner-Ronne (Nicholls et al., 2009) ice shelves that in reality carries HSSW formed in the west over to the central or eastern shelf. Instead of this sub-ice shelf circulation that is captured in R_ISF (Fig. 10), R_PAR drives individual gyre circulations within each of the over-deepened basins, similar in structure to, but stronger than, those in R noISF.

Sea ice extent and thickness in R_PAR match well the R_ISF sea ice characteristics (Fig. 11). Thickness is smaller by more

larger than in the observations (Fig. 8). In R PAR, this is due to the lack of a HSSW circulation pathway beneath the giant

465 than 1m in West Antarctica compared with the R_noISF simulation. Around East Antarctica, and over the Ross and Weddell sea shelves, despite the deficiency in representing the ocean circulation beneath the giant ice shelves, sea ice thickness in R_PAR is similar to that in R_ISF.

These comparisons between R_ISF/R_PAR and R_noISF suggest that not only the presence and the amount of melt water are important, but also the depth at which it is input to the model. The parametrisation directly addresses this latter feature of

470 the sub-ice-shelf ocean circulation and so is able to represent the ocean dynamics associated with the overturning circulation

within the cavity. However, the parametrisation is not fully adapted to mimic the large-scale horizontal gyre circulation that is spun up under the giant ice shelves. This may not be a significant problem because, current coarse resolution ocean models have a nominal resolution of $1^{\circ} \cos(\theta)$, where θ is the latitude, which is sufficient to explicitly represent the two giant ice shelves (L08, Hellmer et al., 2004, Hellmer et al., 2012).

475 5.6 Ice shelf melting

In the previous section we showed that specifying a realistic melting pattern at the ice shelf/ocean interface gives convincing results with major improvements in the properties and circulation of the ocean beyond the ice shelves, especially in the Amundsen and Bellingshausen seas. However, prescribing the freshwater flux represents a strong constraint on the range of applications, since the specified fluxes will only be valid for the present oceanic state. To compute melt rates for other

480 oceanic states interactively, and eventually to couple the ocean model to an evolving ice sheet model, requires the "three equation" formulation for ice shelf melting. Next, we evaluate the ability of the described circum-Antarctic configuration with the "three equation" ice shelf melting formulation to simulated realistic ice shelf melting.

The total ice shelf melting simulated in R_MLT (1864 Gt y⁻¹) is slightly above the range of the observational estimate of Rignot et al. (2013) (Table 3). In R_MLT, as in the observations, we can separate the ice shelves into two different regimes based on the temperature of the water masses on the continental shelves (Fig. 7d) and the average melt rate: the cold water

(Fig. 13b-d) and the warm water (Fig. 13a) ice shelves. As the ice shelf cavity geometry is based on recent estimates (Fretwell et al., 2013) and the ice shelf regime modelled in R_MLT are similar to those in recent observations, the modelled ice shelf melt rate are compared with the Rignot et al. (2013) estimates.

5.6.1 Cold water ice shelves

490 For the Ross, Weddell and East Antarctic continental shelves, the agreement between computed and observed ice shelf melt rates varies. The total melt in R_MLT for these ice shelves (722 Gt y⁻¹) lies within the range of the observations (475-867 Gt y⁻¹) (Table 3). These ice shelves all experience low melt rates (Fig. 13b-d) due to the presence of cold water on the shelves (Fig. 8).

For Filchner-Ronne Ice Shelf (FRIS) the total melt in R_MLT is in agreement with the observation based estimates (Table

- 3), while the spatial pattern of melting and freezing is also similar to other simulations without tidal forcing (Makinson, et al., 2011). FRIS experiences strong melt close to the grounding line, along the ice front and along the paths of the main inflows. Large freezing rates occur along the paths of the main outflows that follow the eastern coasts of the Antarctic Peninsula, Berkner Island and Henry Ice Rise. The latter generates a particularly large area of intense freezing in the central part of the ice shelf, north of the ice rises, in agreement with the observation based distributions of Joughin and Padman
- 500 (2003) and Moholdt et al. (2014).

For Ross Ice Shelf, R_MLT generates a total melt of 111 Gt y^{-1} , with high melt rates concentrated along the ice front, and lower freezing rates in the central part of the ice shelf (Fig. 13). The total melt is within the range of previous model based

estimates (51-260 Gt y⁻¹) and the melting/freezing pattern is in good agreement with earlier modelling studies (Timmermann et al., 2012, Holland et al., 2003 and Dinniman et al., 2007). However, the total melt simulated in R_MLT is 30 Gt y⁻¹ above

505 the observational range, because melt rates along the ice front and on the western side of the ice shelf are larger than those inferred from observation (Rignot, et al.,2013; Moholdt, et al.,2014).

Total melt of Amery Ice Shelf is overestimated by at least a factor of 5 (Table 3), because the waters on the continental shelf in front of the cavity are warmer than observed by more than 1.2°C (Fig. 14). As a consequence, the freezing within the cavity, evaluated from remote sensing and in situ data (Wen, et al.,2010) and simulated by Galton-Fenzi et al. (2012), is

510 absent in R_MLT.

5.6.2 Warm water ice shelves

The ice shelves along the West Antarctic coastline between the Ross and Weddell seas experience a large total melt rate in R_MLT (1142 Gt y⁻¹) (Fig. 12a), due to the presence of CDW on the continental shelf. This total melt is about twice the recent observation based estimate (541, Gt y⁻¹) (Table 3).

- 515 The melt rates in R_MLT are realistic for Abbot Ice Shelf (52 Gt y⁻¹) (Table 3), but slightly underestimated for Thwaites (74 Gt y⁻¹) and Pine Island Glacier (PIG, 87 Gt y⁻¹) compared with observation (Table 3). By comparison with previous modelling studies, R_MLT results for Abbot and PIG ice shelves are in the range of earlier work (Timmermann, et al., 2012; Nakayama, et al., 2014; Shodlock et al., 2016) while for Thwaites the results are above those obtained previously.
- Most of the warm ice shelf melting overestimate in R_MLT comes from Getz (337 Gt y⁻¹) and George VI (298 Gt y⁻¹) ice shelves (+178 Gt y⁻¹ and +181 Gt y⁻¹ respectively, table 3). R_MLT estimates are also well above earlier estimates obtained with FESOM by Timmermann et al. (2012) and Nakayama, et al. (2014) with RTOPO1 bathymetry (Timmerman et al., 2010), respectively, 164 and 127 Gt y⁻¹ for Getz Ice shelf, and 86 and 88 Gt y⁻¹ for George VI Ice Shelf. However, Schodlok et al. (2016) obtained similar melt rates using MITgcm with IBCSO bathymetry (respectively 303.9 and 373.1 Gt y⁻¹).
- These large inter-model differences could have three causes. Firstly, the bathymetry and ice shelf draft data used in 525 Timmermann, et al. (2012) and Nakayama, et al. (2014) come from RTOPO1, whereas Schodlok et al. (2016) and the present study use bathymetry data from IBCSO and ice shelf draft data from BEDMAP2. Differences in ice shelf geometry and bathymetry, particularly the height of seabed sills, can strongly affect ice-shelf melting (Rydt, et al., 2014). Secondly, the ability of off-shelf CDW to cross the shelf break and spread across the continental shelf is a key control on the
- water mass structure within the ice shelf cavities. In R_MLT (Fig. 14) and MITgcm (Shodlock et al., 2016), CDW flow onto
 the shelf is well established. However, in the FESOM simulations of Nakayama et al. (2014), the shelf water is colder than
 the observations by 0.5°C to 3°C, depending of the horizontal resolution used. Analysis of why CDW can cross the
 continental shelf break in some models and not in others is beyond of the scope of this paper.

Finally, NEMO and MITgcm both use z coordinates, while FESOM use a sigma-coordinate around the Antarctic margin. In sigma coordinate model the vertical resolution within the cavity is higher due to the concentration of level beneath the ice

535 shelf. In R_MLT, the number of wet levels in the cavities varies from ~10 levels near the ice fronts to two levels at the

grounding line, while in FESOM there are 21 levels everywhere. This allows better resolution near the grounding line and in the top boundary layer. Shodlok et al. (2016) and the sensitivity experiments performed in Sect. 3.3 show that some ice shelves (West, Dalton, Totten, George VI, Larsen C and FRIS for example) are highly sensitive to the vertical resolution, which affects the ocean properties on the continental shelf, the representation of the top boundary layer beneath the ice shelf, and the ability to resolve details of the cavity geometry.

5.6.3 Limitations

540

In addition to the inter-model differences described above, ice shelf/ocean models in general are still subject to several limitations. Most of them are specific to our model setup as well as the large uncertainties in geometry and forcing data, and critical gaps in our knowledge of dynamics at the ice/ocean interface.

- 545 The most recent bathymetry and ice shelf draft reconstruction of the Amundsen Sea (Millan et al., 2017) shows features that are missing in the BEDMAP2 data-set. In BEDMAP2, for many ice shelves, there are only indirect observations of ice draft, based on satellite surface elevation data, while the sub-ice bathymetry data are often poorly constrained. For some ice shelves (Getz, Venable, Stange, Nivlisen, Shackleton, Totten and Dalton ice shelves, some of the thickest areas of the Filchner, Ronne, Ross and Amery ice shelves and for the ice shelves of Dronning Maud Land), the flotation condition had
- to be enforced by lowering the sea bed arbitrarily from a level that itself was based on nothing more than extrapolation of cavity thickness from surrounding regions of grounded ice and 100 m thick cavity. Consequently, more data are needed for effective modelling (Fretwell, et al., 2013), because cavity geometry has a major impact on the simulated melting by controlling the water mass structure and circulation within the cavity (Rydt, et al., 2014).

Tides have a strong impact on high frequency variability in melting as well as the magnitude and spatial pattern of the temporal mean melt rate (Makinson et al., 2011), but they are not taken into account in the present study.

Subglacial runoff can enhance melting at the ice/ocean interface, especially near the grounding line (Jenkins, 2011). However, the location, magnitude and variability of subglacial outflows from beneath the Antarctic Ice Sheet are poorly known (Dierssen et al, 2002, Fricker et al., 2007).

The drag coefficient, as well as the friction law, affect the top velocity and hence the turbulent exchange coefficients (Eq. 12

- and 13). The appropriate drag coefficient for the base of an ice shelf of unknown roughness is highly speculative, and the range of values discussed in the literature is wide, ranging from 1.5×10^{-3} (Holland and Jenkins, 1999) to 9.7×10^{-3} (Jenkins, et al., 2010), while the basal melting simulated in models is sensitive to the value chosen (Dansereau et al. 2014; Gwyther, et al., 2015, Jourdain et al., 2016). Furthermore, the friction law commonly used to compute the drag is overly simplistic. The same drag coefficient and friction law are used to compute the stress whatever the dynamic regime
- 565 appropriate for the grid point location beneath the ice shelf (i.e. whether it lies within the boundary layer or the free stream flow beyond).

Recent observations beneath George VI ice shelf exhibit thermohaline staircases in the top 20 m below the melting ice shelf base, due to double-diffusive convection (Kimura et al., 2015). These observations raise a doubt about the applicability of

the widely used three-equation model to predict the melt rate in regions where the flow beneath the ice shelf is weak. More

570 experiments, observations, and numerical simulations are needed to fully understand the role of turbulence and thermohaline staircases controlling the heat flux to melting ice shelves.

In addition, Dutrieux et al. (2013) suggest that melting can be concentrated around kilometre-scale heterogeneities in ice thickness, such as keels and channels, especially near the grounding line. Furthermore, Stanton et al. (2013), from density measurements in the top 30m of the ocean beneath Pine Island Glacier, suggest that the top boundary layer can be less than 5

575 m thick. This means either very high horizontal and vertical resolution or a better melt formulation, or both, are needed to improve the representation of processes near the grounding line and the ice shelf base.

6 Conclusions

580

An ice shelf capability has been implemented and evaluated in the NEMO model framework following Losch et al. (2008). The work represents the first step toward a couple ice sheet/ocean model. The working hypothesis used here is that the ice shelf is in equilibrium, with the mass removed by melting being replenished by the flow of the ice shelf, so the shape of the

sub-ice-shelf cavity remains constant over time. In an idealised case (ISOMIP setup) the simulated ocean circulation and ice shelf melting are similar to those described by Losch et al. (2008) using the MITgcm model. Ice shelf melting appears to be sensitive to vertical resolution and the top boundary layer definition. When the Losch top boundary layer thickness is fixed, results are independent of vertical

585 resolution, converge toward those obtained with a vertical resolution equal to that of the top boundary layer. When top boundary layer thickness changes with the vertical resolution under melting conditions, models simulated a cold, fresh, top boundary layer that tends to decrease the thermal forcing and thus the simulated melt rate. At coarse resolution, the cold, top boundary layer is absent, leading to much larger melt rates..

To apply this work to a realistic case, a southward-extended global ORCA grid (eORCA) has been set up using two quasi-

- 590 isotropic bipolar grids south of 67°S. The impact of including the ice shelf cavities has been evaluated in a circum-Antarctic version of the eORCA grid, by comparison with a control simulation without ice shelf cavities. The freshwater and heat flux resulting from ice shelf melting is specified at the ice shelf/ocean interface for the simulation with cavities and at the ocean surface for the control run.
- For warm water shelves, prescribing the ice shelf melting at the surface (R_noISF) leads to a stratification that is too strong compared with the observations. With ice shelf cavities included (R_ISF), melting into the cavity drives a buoyant overturning circulation and entrains warm and salty CDW into the upwelling branch that subsequently mixes into the cold, fresh surface layers outside of the cavity. The entrainment of CDW thus weakens the thermocline by warming and increasing the salinity of the upper ocean layers, resulting in a decrease of the ocean stratification. The activation of the overturning circulation also creates a barotropic circulation that follows f/h contours on the continental shelf.

- 600 For cold water shelves, High Salinity Shelf Water (HSSW) simulated in R_noISF is slightly less dense than observations, but when ice shelf cavities are present, the model is unable to maintain HSSW on the shelf at all. Compared with the simulation without ice shelf cavities, two extra processes consume the HSSW. The vertical overturning circulation driven by melting acts to mix the HSSW with the upper layers all year long, and the presence of new pathways beneath Ross and Filchner-Ronne ice shelves increases the export of HSSW from its formation location on the western continental shelf. The loss of
- 605 HSSW with the ice shelf cavity opened is not balanced by increased dense water formation at the surface. This could be a result of deficiencies in any or all of the atmospheric forcing, the sea-ice model used in this study, or the representation of coastal polynyas.

The effects on sea ice are very dependent on the amount of ocean heat available at depth. Over warm water shelves, the CDW entrained into the cavity overturning circulation warms the surface layer all year long and thus restricts the sea ice

610 formation. This warming of the surface layer leads to thinning of the sea ice by more than 1m in coastal regions of the Bellingshausen and Amundsen seas (2 m locally). Over cold water shelves, including the sub-ice-shelf cavities has a smaller effect on sea ice thickness (less than 20 cm).

Hence, the inclusion of the ice shelf capability in NEMO has a major impact on ocean and sea ice properties. However, the ice shelves vary greatly in area, from $O(100 \text{ km}^2)$ to $O(100 000 \text{ km}^2)$, so depending on the application, more or fewer ice shelves will remain unresolved. In our ¹/₄° configuration the unresolved ice shelves contribute 25% of the total ice shelf

meltwater flux from Antarctica, and at coarser resolutions the majority of the freshwater source could be missing. To mimic the circulation driven by these unresolved ice shelves, the ice shelf melting is uniformly distributed over the depth and width of the unresolved cavity opening, from the mean ice front draft down to the seabed, or the grounding line depth if it is shallower. This simple representation of the ice shelf melting drives a buoyant overturning circulation along the coast

615

620 similar to that that would be present within the ice shelf cavity. Idealised and realistic circum-Antarctic experiments show that this parametrisation mimics the effect of the overturning circulation within small ice shelf cavities and its impact on water mass properties and circulation on the continental shelf. However, for large ice shelves, such as Ross and Filchner-Ronne, the parameterisation is unable to mimic the effect of the large-scale horizontal ocean circulation beneath the ice shelf. Thus, the redistribution of melt water and High Salinity Shelf Water between the different troughs on the continental shelf

625 via their connections under the ice shelf is missing. The specification of ice shelf melting, either over the area of the ice shelf base for resolved cavities or over the area of the cavity opening for unresolved cavities, leads to major improvements in the water mass properties, ocean circulation and sea ice state on the Antarctic continental shelf. However, a model that interactively computes ice shelf melting is crucial for simulating the ocean and ice sheet response to perturbations as well as for developing coupled ice-sheet-ocean models. With

630 the parameterised version of the ice shelf presented here, we only explain how to distribute the meltwater fluxes in an ocean model without ice shelf cavities in a physically sensible way. We do not describe a way to compute the melt rate itself. To tackle this issue, this work needs to be combined with a parameterisation of ice shelf melting (for example: Beckmann and Goosse, 2003; Jenkins et al., 2011).

With the ice shelf cavities opened, the widely-used "three equation" ice shelf melting formulation enables an interactive

- 635 computation of melting. The ability of the circum-Antarctic configuration with the "three equation" ice shelf melting formulation to simulated realistic ice shelf melting has been assessed. Comparison with observational estimates of ice shelf melting reported by Rignot et al. (2013) indicates realistic results for most ice shelves. However, melting rates for Amery, Getz and George VI ice shelves are considerably overestimated and some key ice shelves, such as Totten and Dalton, are missing because of inadequate horizontal and vertical resolution. Possible causes of the overestimated melt rates include
- poor representation of shelf water properties, inaccurate or poorly resolved cavity shape, unknown ice shelf ocean drag coefficient and poor representation of boundary layer processes.
 Despite some deficiencies in the simulation of ice shelf melting and the parametrisation of ocean processes in unresolved ice shelf cavities, this work is a step forward toward a better representation of ice-shelf-ocean interaction in the NEMO
 - framework for all model resolutions. In practice, for horizontal resolutions finer than 2° , some of the ice shelf cavities can be
- resolved (Ross ice shelf for example), while at almost any useable resolution some cavities will have to be parametrized. The most suitable choice of which can be explicitly resolved and which must be parameterised will depend on the combination of horizontal and vertical resolution used.

To apply this work to a global coupled ice sheet/ocean model, we will need some further developments. First, a better knowledge of sub-ice-shelf cavity geometries and key processes that contribute to melting (drag, tides, boundary layer, etc

650 ...) could lead to improvements in the ice shelf representation. Secondly, parameterisations need to be developed to represent the processes (melt and circulation) where the resolution is not fine enough to represent the ice shelf cavity geometry correctly as at the grounding line for example.. Finally, a conservative wetting and drying scheme needs to be developed to allow the grounding line (and calving front) to move back and forth.

Code availability

The model code for NEMO 3.6 is available from the NEMO website (www.nemo-ocean.eu). On registering, individuals can access the FORTRAN code using the open source subversion software (http://subversion.apache.org/). The branch used for both configurations used in this study is the 2015 development branch named dev_r5151_UKMO_ISF at revision 5204. The ice shelf module is now included in the public NEMO distribution.

Data availability

660 The ISOMIP configuration is distributed in NEMO version 3.6 as an unsupported configuration. No file is required to run ISOMIP configuration. For the Circum-Antarctic configuration, the input files (cpp keys, namelist, bathymetry, ice shelf draft, iceberg runoff, initial condition, river runoff, tidal mixing, and weights for the surface forcings) could be requested

from the authors. The surface forcing and the open boundary were provided by the DRAKKAR consortium (http://www.drakkar-ocean.eu).

665 Acknowledgements

The authors acknowledge financial support from the National Environmental Research Council and the UK Met Office. Computational resources were provided by the supercomputing facilities of the British Antarctic Survey and the ARCHER UK National Supercomputing Service. The DFS5.2 forcing fields and the open boundary conditions from the ORCA025-GRD100 simulation were provided by the DRAKKAR coordination (CNRS GDRI No 810). We thank Martin Losch, Julien

670 Le Sommer, Nicolas Jourdain and Jean-Marc Molines for their useful comments and suggestions. We thank, Xylar Asay-Davis and one anonymous reviewer for their detailed and very constructive comments on this work.

References

Adcroft, A. and Campin, J.-M.: Rescaled height coordinates for accurate representation of free-surface flows in ocean circulation models, Ocean Modelling, 7, 269-284, 2004

675 Adcroft, A., Hill, C. and Marshall, J.: Representation of Topography by Shaved Cells in a Height Coordinate Ocean Model, Mon. Wea. Rev., 125(9), 2293-2315, 1997.

Amante, C. and Eakins, B.: ETOPO1: 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis, NOAA Technical Memorandum NESDIS NGDC-24, 2009.

Arakawa, A.: Computational design for long-term numerical integration of the equations of fluid motion: Two-dimensional incompressible flow, Part I. Journal of Computational Physics, 1(1), 119-143, 1966.

- Arndt, J. E., Schenke, H. W., Jakobsson, M., Nitsche, F. O., Buys, G., Goleby, B., Rebesco, M., Bohoyo, F., Hong, J., Black,
 J., Greku, R., Udintsev, G., Barrios, F., Reynoso-Peralta, W., Taisei, M. and Wigley, R.: The International Bathymetric
 Chart of the Southern Ocean (IBCSO) Version 1.0/A new bathymetric compilation covering circum-Antarctic waters,
 Geophys. Res. Lett., 40(12), 3111-3117, 2013.
- Asay-Davis, X. S., Cornford, S. L., Durand, G., Galton-Fenzi, B. K., Gladstone, R. M., Gudmundsson, G. H., Hattermann, T., Holland, D. M., Holland, D., Holland, P. R., Martin, D. F., Mathiot, P., Pattyn, F. and Seroussi, H.: Experimental design for three interrelated marine ice sheet and ocean model intercomparison projects: MISMIP v. 3 (MISMIP+), ISOMIP v. 2 (ISOMIP+) and MISOMIP v. 1 (MISOMIP1), Geoscientific Model Development, 9(7), 2471--2497, 2016.
- Asay-Davis, X. S.: Simulations of ice-shelf cavities in POP, 17th Annual CESM Workshop, NCAR, 690 <u>http://www.cesm.ucar.edu/events/workshops/ws.2012/presentations/liwg/asaydavis.pdf</u>, 18 21 June, 2012.
 - Assmann, K., Hellmer, H. and Beckmann, A.: Seasonal variation in circulation and water mass distribution on the Ross Sea continental shelf. Ant, Sci., mar, 15(1), 3-11, 2003.

Assmann, K. M. and Timmermann, R.: Variability of dense water formation in the Ross Sea, Ocean Dynamics, 55(2), 68-87, 2005.

695 Barnier, B., Madec, G., Penduff, T., Molines, J.,-M., Treguier, A.-M., Le Sommer, J., Beckmann, A., Biastoch, A., Boening, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Remy, E. Talandier, C., Theetten, S., Maltrud, M., McClean, J. and De Cuevas, B.:Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy-permitting resolution, Ocean Dynamics, 56(5-6), 543-567, 2006.

Barnier, B., Dussin, R., Molines, J.-M.: Definition of the interannual experiment ORCA025.L75-GRD100, 1958-2010,

700 MEOM - LEGI – CNRS, LEGI-DRA-12-04-2012, <u>https://www.drakkar-ocean.eu/publications/reports/orca025-grd100-report-dussin</u>, 2012.

Beckmann, A. and Goosse, H.: A parameterization of ice shelf–ocean interaction for climate models. Ocean Modelling, 5(2), 157-170, 2003.

- Beckmann, A., Hellmer, H. H. and Timmermann, R.: A numerical model of the Weddell Sea: Large-scale circulation and water mass distribution, J. Geophys. Res., 104(C10), 23375-23391, 1999.
- Bouillon, S., Morales, M. a., Legat, V. and Fichefet, T. : An elastic-viscous-plastic sea ice model formulated on Arakawa B and C grids. Ocean Modelling, 27, 174-184, 2009. Bouillon, S., Fichefet, T., Legat, V., Madec, G.: The elastic–viscous–plastic method revisited, Ocean Modelling, 71, 2-12,

Bouillon, S., Fichefet, T., Legat, V., Madec, G.: The elastic–viscous–plastic method revisited, Ocean Modelling, 71, 2-12, http://dx.doi.org/10.1016/j.ocemod.2013.05.013, 2013.

- Brodeau, L., Barnier, B., Treguier, A.-M., Penduff, T. and Gulev, S.: An ERA40-based atmospheric forcing for global ocean circulation models, Ocean Modelling, 31 (3–4), 88-104, http://dx.doi.org/10.1016/j.ocemod.2009.10.005, 2010 Budillon, G., Pacciaroni, M., Cozzi, S., Rivaro, P., Catalano, G., Ianni, C., and Cantoni, C.: An optimum multiparameter mixing analysis of the shelf waters in the Ross Sea, Antarctic Science, 15, 105–118, 2003. Brodeau, L., Barnier, B., Treguier, A.-M., Penduff, T. and Gulev, G.: An ERA40-based atmospheric forcing for global ocean
- circulation models, Ocean Modelling, 31(3–4), 88-104, http://dx.doi.org/10.1016/j.ocemod.2009.10.005, 2010.
 Campin, J.-M., Marshall, J. and Ferreira, D.: Sea ice/ocean coupling using a rescaled vertical coordinate z*. Ocean Modelling, 24, 1-14, 2008.
 Carrère L. Lyard F. Cancet M. Guillet A and Poblen L : EES2012: A new global tidal model taking taking advantage of

Carrère L., Lyard, F., Cancet, M., Guillot, A. and Roblou, L.: FES2012: A new global tidal model taking taking advantage of nearly 20 years of altimetry, Proceedings of meeting "20 Years of Altimetry", 2012.

Cunningham, S. A., Alderson, S. G., King, B. A. and Brandon, M. A.: Transport and variability of the Antarctic Circumpolar Current in Drake Passage, Journal of Geophysical Research: Oceans, 108(C5), 2003.
 Dai, A. and Trenberth, K. E.: Estimates of Freshwater Discharge from Continents: Latitudinal and Seasonal Variations. J. Hydrometeor, 3(6), 660-687, 2002.

Dansereau, V., Heimbach, P. and Losch, M.: Simulation of subice shelf melt rates in a general circulation model: Velocitydependent transfer and the role of friction, Journal of Geophysical Research: Oceans, 119(3), 1765-1790, 2014. Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N.

730 and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Q.J.R. Meteorol. Soc., 137(656), 553-597, 2011.

Dinniman, M. S., Klinck, J.M., Bai, L.-S., Bromwich, D. H. and Hines, K. M.: The Effect of Atmospheric Forcing Resolution on Delivery of Ocean Heat to the Antarctic Floating Ice Shelves, Journal of climate, 28, 6067-6085, 2015.

Dinniman, M. S., Klinck, J. M. and Smith, W. O.: Influence of sea ice cover and icebergs on circulation and water mass formation in a numerical circulation model of the Ross Sea, Antarctica, J. Geophys. Res., 112(C11), 2007.

- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H.,
 Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de
 Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichefet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L.,
 Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G.,
- Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N. and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, Climate Dynamics, 40(9-10), 2123-2165, 2013.
 DRAKKAR Group: Eddy-permitting ocean circulation hindcasts of past decades. CLIVAR Exchanges 12, 3, 2007.
- 745 Dussin, R., Barnier, B. Brodeau, L. and Molines, J.-M.: The Making of the DRAKKAR FORCING SET DFS5, <u>https://www.drakkar-ocean.eu/publications/reports/report_DFS5v3_April2016.pdf</u>, DRAKKAR/MyOcean, Report01-04-16, 2016.

Dutrieux, P., Vaughan, D. G., Corr, H. F. J., Jenkins, A., Holland, P. R., Joughin, I., and Fleming, A. H.: Pine Island glacier ice shelf melt distributed at kilometre scales, The Cryosphere, 7(5), 1543-1555, 2013.

750 Emile-Geay, J. and Madec, G.: Geothermal heating, diapycnal mixing and the abyssal circulation, Ocean Science, 5(2), 203-217, 2009.

Favier, L., Durand, G., Cornford, S. L., Gudmundsson, G. H., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A. J. and Le Brocq, A. M.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, Nature Climate Change, 4, 117–121, doi:10.1038/nclimate2094, 2014

Fichefet, T. and Morales, M. A.: Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics,J. Geophys. Res., 102, 12609-12646, 1997.

Foldvik, A., Gammelsrod, T. and Torresen, T.: Circulation and water masses on the southern Weddell Sea shelf, Antarct. Res. Ser., 43, 1985.

Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G.,

- 760 Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M.,
- Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, The Cryosphere, 7, 375-393, doi:10.5194/tc-7-375-2013, 2013.
 Galton-Fenzi, B. K., Hunter, J.R., Coleman, R., S. J. Marsland and Warner, R. C.: Modeling the basal melting and marine ice accretion of the Amery Ice Shelf, J. Geophys. Res., sep, 117(C9), 2012.Grosfeld, K., Gerdes, R. & Determann, J.: Thermohaline circulation and interaction between ice shelf cavities and the adjacent open ocean, J. Geophys. Res., 102(C7),
- 15595-15610, 1997.
 Gwyther, D. E., Galton-Fenzi, B. K., Dinniman, M. S., Roberts, J. L. and Hunter, J. R.: The effect of basal friction on melting and freezing in ice shelf/ocean models, Ocean Modelling, 95, 38-52, 2015.
 Hazeleger, W., C. Severijns, T. Semmler, S. Ştefănescu, S. Yang, X. Wang, K. Wyser, E. Dutra, J. Baldasano, R. Bintanja, P. Bougeault, R. Caballero, A. Ekman, J. Christensen, B. van den Hurk, P. Jimenez, C. Jones, P. Kållberg, T. Koenigk, R.
- 775 McGrath, P. Miranda, T. Van Noije, T. Palmer, J. Parodi, T. Schmith, F. Selten, T. Storelvmo, A. Sterl, H. Tapamo, M. Vancoppenolle, P. Viterbo, and U. Willén: EC-Earth: A Seamless Earth-System Prediction Approach in Action, Bull. Amer. Meteor. Soc., 91(10), 1357-1363, 2010.

Hellmer, H. H.: Impact of Antarctic ice shelf basal melting on sea ice and deep ocean properties. Geophysical Research Letters, 31(10), 2004.

- Hellmer, H. H., Kauker F., Timmermann, R., Determann, J. and Rae, J.:Twenty-first-century warming of a large Antarctic ice-shelf cavity by a redirected coastal current, Nature, 485(7397), 225-228, 2012.
 Hewitt, H. T., Copsey, D., Culverwell, I. D., Harris, C. M., Hill, R. S. R., Keen, A. B., McLaren, A. J., and Hunke, E. C.: Design and implementation of the infrastructure of HadGEM3: the next-generation Met Office climate modelling system, Geosci. Model Dev., 4, 223-253, doi:10.5194/gmd-4-223-2011, 2011.
- 785 Hewitt, H. T., Roberts, M. J., Hyder, P., Graham, T., Rae, J., Belcher, S. E., Bourdallé-Badie, R., Copsey, D., Coward, A., Guiavarch, C., Harris, C., Hill, R., Hirschi, J. J.-M., Madec, G., Mizielinski, M. S., Neininger, E., New, A. L., Rioual, J.-C., Sinha, B., Storkey, D., Shelly, A., Thorpe, L., and Wood, R. A.: The impact of resolving the Rossby radius at mid-latitudes in the ocean: results from a high-resolution version of the Met Office GC2 coupled model, Geosci. Model Dev., 9, 3655-3670, doi:10.5194/gmd-9-3655-2016, 2016.
- 790 Hunter, J. R.: Specification for test models of ice shelf cavities, Tech. Rep., Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Australia, available at: http://staff.acecrc.org.au/~bkgalton/Hunter_2006.pdf, 2006.

Jackett, D. R. and Mcdougall, T. J.: Minimal Adjustment of Hydrographic Profiles to Achieve Static Stability, J. Atmos. Oceanic Technol., 12(2), 381-389, 1995.

Jacobs, S.S., Helmer, H.H., Doake, C.S.M., Jenkins, A. and Frolich, R.M.: 'Melting of ice shelves and the mass balance of Antarctica', Journal of Glaciology, 38(130), 375–387, doi: 10.1017/S0022143000002252, 1992.

Jacobs, S. S., Jenkins, A., Giulivi, C. F. and Dutrieux, P.: Stronger ocean circulation and increased melting under Pine Island Glacier ice shelf, Nature Geosci, 4(8), 519-523, 2011. Jenkins, A.: A one-dimensional model of ice shelf-ocean interaction, Journal of Geophysical Research: Oceans, 96(C11),

20671-20677, 1991.

- Jenkins, A.: The Impact of Melting Ice on Ocean Waters, J. Phys. Oceanogr., 29(9), 2370-2381, 1999.
 Jenkins, A., Hellmer, H. H. and Holland, D. M.: The Role of Meltwater Advection in the Formulation of Conservative Boundary Conditions at an Ice/Ocean Interface, J. Phys. Oceanogr., 31(1), 285-296, 2001.
 Jenkins, A., and Holland, D.: Melting of floating ice and sea level rise, Geophys. Res. Lett., 34, doi:10.1029/2007GL030784, 2007.
- Jenkins, A., Dutrieux, P., Jacobs, S. J., McPhail, S. D., Perrett, J. R., Webb, A. T. and White, D.: Observations beneath Pine Island Glacier in West Antarctica and implications for its retreat. Nature Geosci, 3(7), 468-472, 2010.
 Jenkins, A., Nicholls, K. W. and Corr, H. F. J.: Observation and Parameterization of Ablation at the Base of Ronne Ice Shelf, Antarctica, J. Phys. Oceanogr., 40(10), 2298-2312, 2010.
 Jenkins, A.: Convection-Driven Melting near the Grounding Lines of Ice Shelves and Tidewater Glaciers, J. Phys.
- Oceanogr., 41(12), 2279-2294, 2011.
 Jourdain, N., Mathiot, P., Merino, N., Durand, G., Le Sommer, J., Spence, P., Dutrieux, P. and Madec, G: Ocean circulation and sea-ice thinning induced by melting ice shelves in the Amundsen Sea, JGR-Ocean, 2017.
 Klatt, O., Fahrbach, E., Hoppema, M. and Rohardt, G.: The transport of the Weddell Gyre across the Prime Meridian, Deep Sea Research Part II: Topical Studies in Oceanography, 52, 513-528, 2005.
- 815 Kurtz, N. T. and Markus, T.: Satellite observations of Antarctic sea ice thickness and volume. Journal of Geophysical Research, 117(C8), 2012.

Lambrecht, A., Sandhäger, H., Vaughan, D. and Mayer, C.: New ice thickness maps of Filchner\textendashRonne Ice Shelf, Antarctica, with specific focus on grounding lines and marine ice, Antarctic Science, 19(04), 2007.

Leclair, M. and Madec, G.: A conservative leapfrog time stepping method. Ocean Modelling, 30, 88-94, 2009.

Legg, S., Hallberg, R. and Girton, J.: Comparison of entrainment in overflows simulated by z-coordinate, isopycnal and non-hydrostatic models. Ocean Modelling, 11, 69–97, 2006.
Lewis, E. L., and Perkin R. G. : Ice pumps and their rates, J. Geophys. Res., 91, 11756–11762, doi:10.1029/JC091iC10p11756, 1986.

Little, C. M., Gnanadesikan, A. and Hallberg, R.: Large-Scale Oceanographic Constraints on the Distribution of Melting and Freezing under Ice Shelves, Journal of Physical Oceanography, 38(10), 2242-2255, 2008. Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: World Ocean Atlas 2013, Volume 1: Temperature. S. Levitus, Ed., A. Mishonov Technical Ed.; NOAA Atlas NESDIS 73, 40 pp, 2013.

Losch, M.: Modeling ice shelf cavities in a z coordinate ocean general circulation model. Journal of Geophysical Research: 830 Oceans, 113(C8), 2008.

Madec, G. and Imbard, M.: A global ocean mesh to overcome the North Pole singularity, Climate Dynamics, 12, 381-388, doi:10.1007/BF00211684, 1996.

Madec, G. and the NEMO team: "NEMO ocean engine", NEMO reference manual 3_6_STABLE, Note du Pôle de modélisation, Institut Pierre-Simon Laplace (IPSL), France, No 27 ISSN No 1288-1619, 2016.

- Makinson, K., Holland, P. R., Jenkins, A., Nicholls, K. W., and Holland, D. M.: Influence of tides on melting and freezing beneath Filchner-Ronne Ice Shelf, Antarctica, Geophys. Res. Lett., 38, L06601, doi:10.1029/2010GL046462, 2011.
 Marchesiello, P., McWilliams, J.C. and Shchepetkin, A.: Open boundary conditions for long-term integration of regional ocean models, Ocean Modelling, 3,1-20, 2001.
 Mathiot, P., Barnier, B., Gallée, H., Molines, J.-M., Le Sommer, J., Juza, M. and Penduff, T.: Introducing katabatic winds in
- 840 global ERA40 fields to simulate their impacts on the Southern Ocean and sea-ice, Ocean Modelling, 35(3), 146-160, 2010. Mathiot, P., Goosse, H., Fichefet, T., Barnier, B., and Gallée, H.: Modelling the seasonal variability of the Antarctic Slope Current, Ocean Sci., 7, 455-470, doi:10.5194/os-7-455-2011, 2011.

Mathiot, P., Jourdain, N. C., Barnier, B., Gallée, H., Molines, J.-M., Le Sommer, J. and Penduff, T.: Sensitivity of coastal polynyas and high-salinity shelf water production in the Ross Sea, Antarctica, to the atmospheric forcing. Ocean Dynamics, 62(5), doi:10.1007/s10236-012-0531-v, 2012.

Megann, A., Storkey, D., Aksenov, Y., Alderson, S., Calvert, D., Graham, T., Hyder, P., Siddorn, J., and Sinha, B.: GO5.0: the joint NERC–Met Office NEMO global ocean model for use in coupled and forced applications, Geosci. Model Dev., 7, 1069-1092, doi:10.5194/gmd-7-1069-2014, 2014.

Millan, R., Rignot, E., Bernier, V., Morlighem, M. and Dutrieux, P.: Bathymetry of the Amundsen Sea Embayment sector of

850 West Antarctica from Operation IceBridge gravity and other data, Geo. phys. Res. Lett., 44, 1360–1368, doi:10.1002/2016GL072071, 2017.

Millero, F. J.: Freezing point of sea water. Eighth Report the joint of Panel of Oceanographic Tables and Standards, Appendix UNESCO Tech. 6, 28, 29-31, 1978

Moholdt, G., Padman, L. and Fricker, H. A.: Basal mass budget of Ross and Filchner-Ronne ice shelves, Antarctica, derived from Lagrangian analysis of ICESat altimetry, Journal of Geophysical Research: Earth Surface, 119(11), 2361-2380, 2014.

Morales Maqueda, M. A., Willmott, A. J. and Biggs, N. R. T.: Polynya Dynamics: a Review of Observations and Modeling, Reviews of Geophysics, 42(1), 2004.

855

Nakayama, Y., Timmermann, R., Rodehacke, C. B., Schröder, M. and Hellmer, H. H.: Modeling the spreading of glacial meltwater from the Amundsen and Bellingshausen Seas. Geophys. Res. Lett., 41(22), 7942-7949, 2014.

Nicholls, K. W., Osterhus, S., Makinson, K., Gammelsrod, T. and Fahrbach, E.: Ice-ocean processes over the continental shelf of the southern Weddell Sea, Antarctica: A review. Rev. Geophys., 47(3), doi: 10.1029/2007RG000250, 2009.
Orlanski, I.: A simple boundary condition for unbounded hyperbolic flows: Journal of Computational Physics, 21(3), 251-269, 1976.

Padmann, L., Erofeeva, S. and Joughin, I.: Tides of the Ross Sea and Ross Ice Shelf cavity, Ant. Sci., 15(1), 31-40, 2003.

865 Penduff, T., Juza, M., Brodeau, L., Smith, G. C., Barnier, B., Molines, J.-M., Treguier, A.-M., and Madec, G.: Impact of global ocean model resolution on sea-level variability with emphasis on interannual time scales, Ocean Sci., 6, 269-284, doi:10.5194/os-6-269-2010, 2010.

Rignot, E., Casassa, G., Gogineni, P., Krabill, W., Rivera, A. and Thomas, R.: Accelerated ice discharge from the Antarctic Peninsula following the collapse of Larsen B ice shelf. Geophys. Res. Lett., 31, L18401, doi:10.1029/2004GL020697, 2004.

- Rignot, E., Velicogna, I., van den Broeke, M. R., Monaghan, A. and Lenaerts, J. T. M.: Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise. Geophys. Res. Lett., 38, L05503, doi:10.1029/2011GL046583, 2011.
 Rignot, E., Jacobs, S., Mouginot, J. and Scheuchl, B.: Ice-Shelf Melting Around Antarctica. Science, 341, 266-270, 2013.
 Rydt, J. D., Holland, P. R., Dutrieux, P. and Jenkins, A. : Geometric and oceanographic controls on melting beneath Pine Island Glacier. Journal of Geophysical Research: Oceans, 119(4), 2420-2438, 2014.
- 875 Schmidt, G. A., Bitz, C. M., Mikolajewicz, U. and Tremblay, L.: Ice-ocean boundary conditions for coupled models, Ocean Modelling, 7, 59-74, 2004.

Scambos, T., Bohlander, J. A., Shuman, C. A. and Skvarca, P. : Glacier acceleration and thinning after ice shelf collapse in the Larsen B embayment, Antarctica, Geophys. Res. Lett., 31, L18402, 2004.

Schoof, C. Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, J. Geophys. Res., 112(F3), 2007.

880 Scoccimarro, E., Gualdi, S., Bellucci, A., Sanna, A., Giuseppe Fogli, P., Manzini, E., Vichi, M.,Oddo, P. and Navarra, A.: Effects of Tropical Cyclones on Ocean Heat Transport in a High-Resolution Coupled General Circulation Model, J. Climate, 24(16), pp. 4368-4384, 2011.

Schodlok, M. P., Menemenlis, D. and Rignot, E. J.: Ice shelf basal melt rates around Antarctica from simulations and observations, J. Geophys. Res. Oceans, 121, 1085–1109, doi:10.1002/2015JC011117, 2016

- Silva, T. A. M., Bigg, G. R. and Nicholls K. W. (2006): Contribution of giant icebergs to the Southern Ocean freshwater flux, J. Geophys. Res.,111, C03004, doi:10.1029/2004JC002843, 2006.
 Simmons, H. L., Jayne, S. R., Laurent, L. C. and Weaver, A. J. Tidally driven mixing in a numerical model of the ocean general circulation, Ocean Modelling, 6, 245-263, 2004.
 Stanton, T. P., Shaw, W.J., Truffer, M., Corr, H. F. J., Peters, L. E., Riverman, K. L., Bindschadler, R., Holland, D. M.,
- 890 Anandakrishnan, S.: Channelized Ice Melting in the Ocean Boundary Layer Beneath Pine Island Glacier, Antarctica. Science, 341(6151), pp. 1236-1239, 2013.

Timmermann, R., Goosse, H., Madec, G., Fichefet, T., Ethe, C., & Dulière, V.: On the representation of high latitude processes in the ORCA-LIM global coupled sea ice-ocean model. Ocean Modelling, 8(1–2), 175–201. https://doi.org/10.1016/j.ocemod.2003.12.009, 2005.

895 Timmermann, R., Brocq, A.L., Deen, T., Domack, E., Dutrieux, P., Galton-Fenzi, B., Hellmer, H., Humbert, A., Jansen, D., Jenkins, A., et al.: A consistent data set of Antarctic ice sheet topography, cavity geometry, and global bathymetry. Earth Syst. Sci. Data 2, 261–273, 2010.

Timmermann, R., Wang, Q. and Hellmer, H.: Ice-shelf basal melting in a global finite-element sea-ice/ice-shelf/ocean model, Annals of Glaciology, 53(60), 303-314, 2012.

Treguier, A. M., Barnier, B., de Miranda, A. P., Molines, J. M., Grima, N., Imbard, M., Madec, G., Messager, C., Reynaud, T. and Michel S.: An eddy-permitting model of the Atlantic circulation: Evaluating open boundary conditions, J. Geophys. Res., 106(C10), 22115–22129, doi:10.1029/2000JC000376, 2001.

Voldoire, A., Sanchez-Gomez, E., Salas y Mélia, D., Decharme, B., Cassou, C., Sénési, S., Beau, I., Alias, A., Chevallier,
M., Déqué, M., Deshayes, J., Douville, H., Fernandew, E. and Madec, G.:The CNRM-CM5.1 global climate model: description and basic evaluation. 40(9-10), pp. 2091-2121, 2013.

Wang, C. and Beckmann, A.: Investigation of the impact of Antarctic ice-shelf melting in a global ice-ocean model (ORCA2-LIM), Annals of Glaciology, 46(1), 78-82, 2007.

905

Wen, J. et al. Basal melting and freezing under the Amery Ice Shelf, East Antarctica, Journal of Glaciology, 56(195), 81-90, 2010.

910 Zweng, M.M, Reagan, J.R., Antonov, J.I., Locarnini, R.A., Mishonov, A.V., Boyer, T.P., Garcia, H.E, Baranova, O.K., Johnson, D.R., Seidov, D. and Biddle, M.M.: World Ocean Atlas 2013, Volume 2: Salinity. S. Levitus, Ed., A. Mishonov Technical Ed.; NOAA Atlas NESDIS 74, 39 pp., 2013.



- Masked area (land, ice sheet or closed ice shelf cavity)
- 915 Figure 1: Freshwater and associated latent heat introduced a) at the surface (R_ noISF), b) beneath the ice shelf (A_ISF, R_ISF and R_MLT), c) at the ice shelf base level (A_BG03) and d) over the depth range of the ice shelf base (A_PAR and R_PAR).



Figure 2: Near steady state (after 10000 days) solution of the I_30M ISOMIP experiment. a) Horizontal stream function (Psi) in Sv with a contour interval of 0.02 Sv, b) Meridional overturning circulation (moc) in Sv with a contour interval of 0.01. c) Melt rate in $m y^{-1}$ (negative for melting and positive for freezing) with a contour interval of 0.4 $m y^{-1}$.



925 Figure 3: a) Total melting rate versus total freezing rate, and b) meridional overturning circulation versus barotropic stream function (bsf) for all the ISOMIP sensitivity experiments (I_5M, I_10M, I_30M, I_60M, I_100M, I_150M, I_31L, I_46L and I_75L). The simulations I_XXM are with constant vertical resolutions of XX m and a H_{TBL} of 30 m, the simulations I_XXM yer with variable vertical resolution. Details are given in Table 2.



Figure 4: Melt rate in a) the 5M simulation, and b) the 100M simulation in m y^{-1} (negative for melting and positive for freezing) with a contour interval of 0.4 m y^{-1} .



935 Figure 5: (a) Zonal mean temperature (°C) after 30 years of the run. In contour, the meridional overturning stream function (MOC) in the A_ISF experiment. b) Mean temperature difference (°C) with respect to A_ISF experiment (A_PAR-A_ISF). In contour, the MOC in the A_PAR experiment. c) as b) but for A_BG03.



940 Figure 6: Bathymetry (m) over the Antarctic continental shelf and beneath the ice shelves. Black lines are the cell edges (plotted every 25 cells). The thick grey line is the limit of the Weddell sector of the grid and the thick dashed grey line is the limit of the Ross, Amudsen and Bellingshausen sector.



945 Figure 7: Temperature (°C) averaged between 300 and 1000m from a) R_ISF, b) R_PAR, c) R_ noISF, d) World Ocean Atlas 2013 (Locarnini et al., 2013 and Zweng et al., 2013).



Figure 8: Salinity (PSU) averaged between 300 and 1000m from a) R_ISF, b) R_PAR, c) R_ noISF, d) World Ocean Atlas 2013 (Locarnini et al., 2013 and Zweng et al., 2013).



Figure 9: Profiles (year 10, 1988) in Pine Island Bay in R_noISF (blue), R_ISF (red) and R_PAR (green) of a) salinity and b) temperature. Climatology from 1994 to 2012 (Dutrieux et al, 2014) is in black.



Figure 10: Barotropic stream function (Sv) on the Ross, Amundsen, Bellingshausen and Weddell continental shelves in a) R_ISF, b) R_PAR and c) R_ noISF. Stream function isolines out of the ± 2 Sv range are not plotted.



960 Figure 11: Mean sea ice thickness (m) from September to November (SON) in colour. Lines represent the sea ice extent (threshold set at 15% ice concentration) in the observations of Comiso (2000) (grey) and the corresponding simulation (black). a) R_ISF, b) R_PAR, c) R_ noISF, d) Kurtz and Markus (2012) data. The observational uncertainty is ±40 cm.



965 Figure 12: Map of temperature in °C (a,b) and salinity in PSU (c,d) differences between R_PAR and R_ISF (a,c) and R_noISF and R_ISF (b,d) averaged between 300 m and 1000 m.



Figure 13: Ice shelf melting (m y⁻¹, positive values mean melting) in the R_MLT simulation for a) the West Antarctic ice shelves, b) Ross Ice Shelf, c) Filchner-Ronne Ice Shelf and d) the East Antarctic ice shelves. Note that panels (a) and (b,c and d) have different colorbars.



Figure 14: 300-1000m mean temperature differences between R_MLT and observations from World Ocean Atlas 2013 (Locarnini et al., 2013 and Zweng et al., 2013). Grey area represents ice sheet, ice shelves or ocean shallower than 300m. The hatched area limited by the green line represents where the observational dataset is obtained by extrapolation.

980 Table 1: Parameters used in the ice shelf/ocean interaction formulation.

Symbol	Description	Value	Unit
Cp	Ocean specific heat	3992	J kg K ⁻¹
Lf	Ice latent heat of fusion	3.34×10^{5}	J kg ⁻¹
C _{p,i}	Ice specific heat	2000	J kg ⁻¹ K ⁻¹
ρί	Ice density	920	Kg m ⁻³
К	Heat diffusivity	1.54×10^{-6}	m ² s ⁻¹
Cd	Top drag coefficient	10 ⁻³	
$\sqrt{C_d}\Gamma_T$	Thermal Stanton number	1.1×10^{-3}	
$\sqrt{C_d}\Gamma_S$	Diffusion Stanton number	3.1×10^{-5}	
λ_1	Liquidus slope	-0.0575	°C PSU ⁻¹
λ_2	Liquidus intercept	0.0901	°C
λ_3	Liquidus pressure coefficient	-7.61×10^{-4}	°C m ⁻¹

Name	Vertical resolution in the	Losch	top	Melt rate	
	cavity	boundary layer			
1.514	5	5		E1	
1_3M	5 m	5 m		Expl.	
I_5M30M	5 m	30 m		Expl.	
I_10M	10 m	10 m		Expl.	
I_10M30M	10 m	30 m		Expl.	
I_30M	30 m	30 m		Expl.	
I_60M	60 m	30 m		Expl.	
I_100M	100 m	30 m		Expl.	
I_150M	150 m	30 m		Expl.	
I_31L	40-240 m	30 m		Expl.	
I_46L	40-110 m	30 m		Expl.	
I_75L	20-80 m	30 m		Expl.	
A_ISF	30 m (Fig. 1b)	30 m		Presc.	
A_PAR	No cavity (Fig. 1d)	N/A		Presc.	
A_BG03	No cavity (Fig. 1c)	N/A		Presc.	
R_ISF	20-80 m (Fig. 1b)	30 m		Presc.	
R_PAR	No cavity (Fig. 1d)	N/A		Presc.	
R_noISF	No cavity (Fig. 1a)	N/A		Presc.	
R_MLT	20-80 m (Fig. 1b)	30 m		Expl.	

Table 2: List of model runs. Expl. means the ice shelf melt rate is explicitly calculated. Presc. means the ice shelf melt rate is prescribed (i.e. independent of ocean temperature and salinity and constant in time).

Table 3: Basal melt in Gt y⁻¹ for the last year of simulation in R_MLT. Observations come from Rignot et al. (2013). Geometry column indicates the main modification to the BEDMAP2 bathymetry/ice shelf draft as follows: GL means the GL is moved seaward, "shallow" means the ice shelf is too shallow away from the grounding line and "narrow" means the narrowest passage

into the cavity is one cell wide. $++/+/0/-/-$ is a summary of the ocean temperature condition at the closest nor the WOA2013 observational dataset (Fig. 14). $++$ for ocean temperature differences wrt WOA2013 of differences in the range 0.5 and 1°C 0 differences in the range 0.5 and -0.5 °C - differences in the range 0.5	t non-extrapola 8 of more that	ted cell in
the WOA2013 observational dataset (Fig. 14). ++ for ocean temperature differences wrt WOA2013 of differences in the range 0.5 and ${}_{2}$ C of differences in the range 0.5 and	8 of more than	n 1°C. +
differences in the range 0.5 and 1° C. 0 differences in the range 0.5, and -0.5 °C differences in the range -0.5	, or more that	, .
and the function of the functi	e -0.5 and -1 °C	and for
ocean temperature differences greater than -1 °C.		

Ice shelf	Model	Obs (Rignot 2013)	Temperature error at the ice shelf edge (observation: WOA2013)	Geometry
Amery	207	13 – 59	++	GL
West	26	17 – 37	0	
Shackleton	14	58 - 88		GL
Ross	111	14 - 81	0	GL, shallow
Larsen C	46	-46 - 87	0	
FRIS	123	111 – 210	0	GL
Brunt + Riiser	39	-6 - 26	-	shallow
Fimbul	42	13 – 43	-	GL
Cold ice shelves	722	531 - 1033		
Getz	337	131 – 159	+ (east) (west)	shallow
Thwaites	74	91 – 105	+	
Pine Island	87	93 - 109	+	
Abbot	52	32 - 72	+	
George VI	298	72 – 106	+	narrow
Warm ice shelves	1142	452 - 630		
Others	408	214 - 425		
Total	1864	1263 – 1737		