

# ***Interactive comment on “Description and Validation of the Simple, Efficient, Dynamic, Global, Ecological Simulator (SEDGES v.1.0)” by Pablo Paiewonsky and Oliver Elison Timm***

**Pablo Paiewonsky and Oliver Elison Timm**

ppaiewonsky@albany.edu

Received and published: 15 July 2017

Printer-friendly version

Discussion paper



## Reply to Anonymous Referee #3

July 15, 2017

To begin with, we thank the reviewer for raising numerous important issues in his or her critique of our paper. In particular, we hope that in our response, our discussion on the constant  $c_i/c_a$  in our model is useful to others in the land surface modeling community, too.

My major concern is that the improvement of model performances between SimBA and SEDGES is not clearly demonstrated. Since SEDGES builds on SimBA, to emphasize the value of this work, the advance of SEDGES needs to be well manifested.

We agree with the reviewer that the value of the development of SEDGES would be highlighted by demonstrating its improved performance as compared to SimBA. It is important to remember that there are basically two main versions of SimBA. The older version (Kleidon et al., 2006) has been used in several studies (which are cited in the main paper) and is included in version 15 of the Planet Simulator (Lunkeit et al., 2007). The newer version of SimBA was developed by Pablo Paiewonsky from this older version and is included in version 16 of Planet Simulator (Lunkeit et al., 2011). Pablo Paiewonsky prefers to not evaluate the newer version of SimBA that he developed, because that version had no formal evaluation and, to his knowledge, has not been

[Printer-friendly version](#)

[Discussion paper](#)



used in any published study. His personal opinion is that SEDGES is a much better model. However, if the reviewer insists, we would be willing to compare SEDGES with that newer version of SimBA.

For example, the authors show particularly well-simulated GPP, but it is not clear how much SEDGES improves the representations and simulations of GPP in SimBA. In other words, whether this well-simulated GPP is due to the incorporated processes in SEDGES, or due to the original framework set up in SimBA model? The authors need to prove that SEDGES indeed improves the GPP simulation compared to that from SimBA. Maybe adding the SimBA simulations in those relevant figures is the easiest way to illustrate. It will be even better if the authors could provide a short summary of how GPP is modeled in SimBA.

The older version of SimBA has a serious problem of multiple steady states, when used in a coupled model (Dekker et al., 2010). When we recently forced the old SimBA offline using the ERA-Interim reanalysis forcing as described in our paper, we also found multiple equilibria (at the end of the 280ppm CO<sub>2</sub> spin-up period). One simulation started from bare soil, and the other one was initialized with a "tropical forest" level of biomass. The forest-initialized simulation is only slightly more realistic than the bare ground-initialized simulation.

Results (GPP and ET) from only the forest-initialized simulation are shown below in the two figures for the years indicated (i.e. near the end of the increasing CO<sub>2</sub> phase). The simulated GPP is very unrealistic (fig. 1 below); it is much further away from the observational datasets than is the GPP from SEDGES (figure 2 in the main text). The simulated ET (fig. 2 below) is generally too large due to the Manabe bucket "beta" formulation, which gives the potential rate of ET for soil wetness fractions  $\geq 0.25$ . This figure can be compared with the SEDGES simulation of ET, given in figure 12 of the

main paper, which is much more realistic. The GPP figure, especially, shows that this older version of SimBA has very severe deficiencies when forced offline with the ERA-Interim reanalysis data.

Similarly, the authors state that SEDGES improves most of the parameterization of SimBA, but again it is not clear whether or not those modifications of parameterization indeed improve SimBA simulation. The lesson we learnt from current LSMs tells us that increasing the complexity not necessarily guarantees a better model performance. Therefore, I am also concerned about the trade-off between realism and simplicity. To balance this trade-off with the purpose of improving the reliability and robustness of models, the added processes or modified parameterizations need to be proved as necessary for improving the reliability of the model.

First, we disagree that the modifications need to be formally shown to be necessary for improving model reliability. Informal evaluations occurred as the model was developed, and changes were made along the way as were deemed necessary. Second, for the majority of cases in which the SimBA parameterizations were changed, the resulting increase in model complexity was negligible. As is partially stated in the introduction, there are four main increases in complexity in SEDGES relative to SimBA: separation of ET into soil and vegetative components, inclusion of aerodynamic conductance in the formulation for carbon uptake by vegetation, full coupling of photosynthesis and transpiration through interactive canopy conductance, and soil organic carbon-dependent soil albedo. Of these three, we justify the need for only the fourth (in section 2.3.1). In the revised manuscript, we will list all four increases in complexity and explain what advantages are gained by incorporating them into SEDGES.

Below, we include a comparison between the old version of SimBA and SEDGES with respect to evapotranspiration and GPP.

Moreover, the simplifications also need to prove as reasonable. For example, the ratio of  $c_i/c_a$  in Equation (6) is considered as a constant, but has been shown that the optimal stomatal behavior allows  $c_i/c_a$  decreases with VPD, increase with temperature (e.g. Prentice et al. 2014, Medlyn et al. 2011, Lin et al. 2015). The variation in  $c_i/c_a$  seems quite important in terms of capturing the spatial pattern of GPP (Wang et al. 2014). This simplification needs a justification.

We agree with the reviewer that it would be preferable to have variable  $c_i/c_a$  in our model. In a future version of SEDGES, we hope to include this feature. Its incorporation into SEDGES is actually quite difficult because the simple relationships between  $c_i/c_a$  and VPD that are derived by Medlyn et al. (2011) and Prentice et al. (2014) are incompatible with the SEDGES framework. Medlyn et al. (2011) and Prentice et al. (2014) assume a Fick's Law of Diffusion transmission between the leaf and the outside air. In order to have consistency in moisture and  $\text{CO}_2$  fluxes between the land surface and atmosphere, the land surface scheme also needs to use a diffusive exchange between outside air and leaves, or at the least, reasonably approximate such diffusive exchange. Diffusive exchange of moisture and  $\text{CO}_2$  is not used in the SEDGES framework. Instead, transfer occurs from surface to atmosphere through a bulk aerodynamic formulation. This formulation only approximates the diffusive scheme when canopy resistance ( $r_c$ ) greatly exceeds aerodynamic resistance ( $r_a$ ). As is said in the main text, in early versions of SEDGES (coupled to Planet Simulator), it was found that this condition ( $r_c \gg r_a$ ) was frequently violated.

It should be mentioned that Pablo Paiewonsky has been extremely interested in incorporating variable  $c_i/c_a$  into SEDGES and did in fact include its dependency on VPD in an early, unpublished version of SEDGES, but later realized that it was not theoretically justifiable and thus removed it.

[Printer-friendly version](#)[Discussion paper](#)

It should also be noted that even though  $c_i/c_a$  decreases with VPD (Morison and Gifford, 1983), it is nevertheless conserved in the real world under many conditions, including a wide range of different light levels,  $\text{CO}_2$  levels, and nutrient levels (Wong et al., 1979).

While we view this constant  $c_i/c_a$  as a limitation of the model, we think that the reviewer overstates its importance with regards to capturing the spatial pattern of GPP. Our GPP results with SEDGES suggest that a fixed  $c_i/c_a$  is adequate for this purpose. Also, the paper that the reviewer cites to justify the importance of variable  $c_i/c_a$  on GPP, Wang et al. (2014), shows that the effect of including the RuBP-regeneration limited rate (which has the  $c_i$  dependency) has a much weaker impact on the spatial pattern of annual GPP than do the two key variables: incoming light levels and foliage cover, whose variation are indeed incorporated into SEDGES.

Moreover, LUE (light-use efficiency) models using remotely sensed data have been successfully used for many years, but typically lack explicit  $c_i$  dependence (e.g., Yuan et al., 2007). (Granted, though,  $c_i$  dependence may be implicit in productivity's dependence on VPD in some of those models). More specifically, with regards to the VPD dependency of  $c_i/c_a$  and its subsequent effects on GPP, as we describe below, dependency of GPP on VPD in SEDGES does occur by way of the water-limited rate, GPPW, i.e. by hydraulic transport limitation. In this sense, productivity limitation by VPD still occurs in SEDGES, albeit through a different mechanism than by  $c_i/c_a$  reduction. With regards to the temperature dependency of  $c_i/c_a$  and its subsequent effects on GPP, colder temperatures do result in lower  $c_i$  (Prentice et al., 2014; Lin et al., 2015), which, in isolation, reduces GPP in the Farquhar model of photosynthesis (Farquhar et al., 1980). However, this effect is substantially offset by the decrease in the  $\text{CO}_2$  compensation point with lower temperatures (Prentice et al., 2017). On the other hand, we think it would be beneficial to discuss in the manuscript what the ramifications are of using a fixed  $c_i/c_a$  on GPP, for the situations in which it is significant, and also the ramifications for simulating transpiration.

[Printer-friendly version](#)[Discussion paper](#)

Having a constant  $c_i/c_a$  is likely to play a more important role in transpiration (by way of affecting the water use efficiency) than (at least directly) in GPP. This is because, when  $r_c \gg r_a$  (i.e. using the diffusive approximation for heuristic purposes), transpiration is (unless it is occurring at the maximum rate in our model) proportional to  $VPD \cdot GPPL / (1 - c_i/c_a)$ , where GPPL is the light-limited rate of GPP. As such, changes in  $c_i/c_a$  (whose values are typically closer to 1 than 0 for C3 plants) have greater relative impact on transpiration than they do in the equations for either RuBP regeneration-limited or Rubisco-limited photosynthesis (at least for non-extreme conditions) in the Farquhar model (Farquhar et al., 1980).

I think it is  $c_i$  (the intercellular CO<sub>2</sub> concentration) that really matters in CO<sub>2</sub> fertilization. If you consider  $c_i$  for water-limited GPP in Equation (6), why not here for light-limited GPP?"

As stated in the paper, the parameterization for CO<sub>2</sub> fertilization (Equation 3) comes from Franks et al. (2013). Pablo Paiewonsky has had a similar concern as the reviewer and agrees that a dependency on  $c_i$  would make more sense, from a theoretical standpoint. The above fertilization parameterization is for the RuBP regeneration-limited rate of photosynthesis, but it makes the approximation that  $c_i \approx c_a$ . In Pablo Paiewonsky's current interpretation, the Franks et al. (2013) parameterization is with respect to  $c_a$  and not  $c_i$  because it seems that  $c_a$  and not always  $c_i$  was measured in the CO<sub>2</sub> sensitivity experiments that they used to validate their parameterization. So, strictly speaking, the empirical support of that fertilization parameterization is with respect to  $c_a$ , rather than  $c_i$ . Regardless, the impact of using  $c_a$  instead of  $c_i$  in that equation is not large. For instance, if one uses  $c_i$  in Equation 3 and  $c_i/c_a = 0.8$ , then it can be shown that there is only an  $\approx 6\%$  increase in the light-limited rate of GPP at CO<sub>2</sub> of 2000ppm as compared to using  $c_a$  in Equation 3. In fact, a potentially greater source of inaccuracy in the fertilization parameterization than the above is the assumption of a constant CO<sub>2</sub> compensation point.

[Printer-friendly version](#)[Discussion paper](#)

Typically, modeled LUE (not only fAPAR) is represented with a dependency on vapor pressure deficit, why the equation here does not include such an effect? Is it implicitly considered via the coupling with water-limited GPP?

Yes, it is correct that the effect of VPD is implicit within the water-limited GPP. Canopy conductance is limited by a maximum rate of transpiration, which, in turn, depends on the specific humidity deficit ( $\Delta q$ ), which, in turn, is approximately proportional to VPD. Note that the effect of VPD has nothing to do with optimality, though.

### References

Dekker, S., De Boer, H., Brovkin, V., Fraedrich, K., Wassen, M., and Rietkerk, M.: Biogeophysical feedbacks trigger shifts in the modelled vegetation-atmosphere system at multiple scales, *Biogeosciences*, 7, 1237-1245, 2010.

Dewar, R. C.: A simple model of light and water use evaluated for *Pinus radiata*, *Tree Physiology*, 17, 259-265, 1997.

Farquhar, G., von Caemmerer, S., and Berry, J.: A Biochemical Model of Photosynthetic CO<sub>2</sub> Assimilation in Leaves of C<sub>3</sub> Species, *Planta*, 149, 78-90, 1980.

Franks, P. J., Adams, M. A., Amthor, J. S., Barbour, M. M., Berry, J. A., Ellsworth, D. S., Farquhar, G. D., Ghannoum, O., Lloyd, J., McDowell, N., Norby, R. J., Tissue, D. T., and von Caemmerer, S.: Sensitivity of plants to changing atmospheric CO<sub>2</sub> concentration: from the geological past to the next century, *New Phytologist*, 197, 1077-1094, 2013.

Kleidon, A.: The climate sensitivity to human appropriation of vegetation productivity and its thermodynamic characterization, *Global and Planetary Change*, 54, 109-127, 2006.

Lunkeit, F., Bottinger, M., Fraedrich, K., Jansen, H., Kirk, E., Kleidon, A., and Luksch, U.: Planet Simulator-reference manual, version 15.0, Tech. rep., Meteorologisches Institut, Universität Hamburg, 2007.

[Printer-friendly version](#)[Discussion paper](#)



Lunkeit, F., Borth, H., Bottinger, M., Fraedrich, K., Jansen, H., Kirk, E., Kleidon, A., Luksch, U., Paiewonsky, P., S, S., Sielmann, S., and Wan, H.: Planet Simulator-reference manual, version 16, Tech. rep., Meteorologisches Institut, Universität Hamburg, 2011.

Medlyn, B. E., Duursma, R. A., Eamus, D., Ellsworth, D. S., Prentice, I. C., Barton, C. V., Crous, K. Y., de Angelis, P., Freeman, M., and Wingate, L.: Reconciling the optimal and empirical approaches to modelling stomatal conductance, *Global Change Biology*, 17, 2134-2144, 2011.

Morison, J.I. and Gifford, R.M.: Stomatal sensitivity to carbon dioxide and humidity, *Plant physiology*, 71, 789-796, 1983.

Mueller, B., Hirschi, M., Jimenez, C., Ciais, P., Dirmeyer, P., Dolman, A., Fisher, J., Jung, M., Ludwig, F., Maignan, F., et al.: Benchmark products for land evapotranspiration: LandFlux-EVAL multi-data set synthesis, *Hydrology and Earth System Sciences*, 17, 3707-3720, 2013.

Prentice, I. C., Dong, N., Gleason, S. M., Maire, V., and Wright, I. J.: Balancing the costs of carbon gain and water transport: testing a new theoretical framework for plant functional ecology, *Ecology letters*, 17, 82-91, 2014.

Prentice, I.C., Cleator, S.F., Huang, Y.H., Harrison, S.P., and Roulstone, I.: Reconstructing ice-age palaeoclimates: Quantifying low-CO<sub>2</sub> effects on plants, *Global and Planetary Change*, 149, 166-176, 2017.

Wang, H., Prentice, I. C., and Davis, T. W.: Biophysical constraints on gross primary production by the terrestrial biosphere, *Biogeosciences*, 11, 5987-6001, 2014.

Wong, S. C., Cowan, I. R., and Farquhar, G. D.: Stomatal conductance correlates with photosynthetic capacity, *Nature*, 282, 424-426, 1979.

Yuan, W., Liu, S., Zhou, G., Zhou, G., Tieszen, L. L., Baldocchi, D., Bernhofer, C., Gholz, H., Goldstein, A. H., Goulden, M. L., et al.: Deriving a light use efficiency model

[Printer-friendly version](#)[Discussion paper](#)

from eddy covariance flux data for predicting daily gross primary production across biomes, *Agricultural and Forest Meteorology*, 143, 189-207, 2007.

---

Interactive comment on *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2017-75>, 2017.

GMDD


---

Interactive  
comment

Printer-friendly version


Discussion paper





**Fig. 1.** Multi-year annual mean of gross primary productivity (GPP) for "old" SimBA (version 15 of Planet Simulator) and two reference datasets, MTE and CARBONES, for 1990-2009 over non-glaciated land.

[Printer-friendly version](#)[Discussion paper](#)



**Fig. 2.** Zonal monthly mean and annual mean climatologies of evapotranspiration for "old" SimBA (version 15 of Planet Simulator) and Mueller et al. (2013) reference dataset for 1989–2005 (non-glaciated land).

[Printer-friendly version](#)[Discussion paper](#)