

Answer to comments from the reviewer #1.

We thank reviewer for the constructive evaluation of the manuscript. Please find below our answers to questions/comments. Comments from the reviewer were left intentionally in this document and written in roman font. Our answers are written in italics.

Anonymous Referee #1

Received and published: 14 June 2018

Improvement of the soil modules in global carbon cycle models is a recurrent need claimed by the scientific community. The land surface model Orchidee is one of the important tools to analyse and predict future changes of the Earth's climate and biosphere. A recent study highlighted that current Earth system models predict a too young age for soil organic carbon. The present work introduces the radiocarbon isotope in the model to better constrain Orchidee. Based on the use of radiocarbon, the study furthermore improved the model itself, and model prediction, through a better representation of carbon movement within the soil profile. The article is fully relevant, clearly written and illustrated, and worth publication in GMD.

Thank you very much for the positive comments

But one point requires a significant change. Once this point fixed, the paper could be acceptable with only minor corrections.

Important point. (parameterization of the 'Model_Test_He') Line 233 authors state "... multiply by 14 the turnover rate and by 0.07 the flux...". And later Line 236 and in Table 2: "decrease the flux from 0.07 to 0.049. Consistency would either decrease the flux from 0.07 to 0.0049, or multiply it by a factor 0.7. I suppose that the initial intention of authors was to multiply by 0.07 the flux, so that the steady state stock of passive would be kept similar (multiplied by $14 \times 0.07 = 0.98$), but with a F14 much lower.

Here it seems from the results that the stock of the passive pool was multiplied by a factor almost 10 (less than 10 because of the duration of the spin-up), as expected by a factor $\times 0.7$ for the flux. The over estimation of both carbon content and age is obviously expected with such a parameterization. In the present state, I further recommend not to use the name of a person in the surname of the model.

Finally, I recommend that the authors either (i) remove this model_test from the paper, which would then be accepted with minor revision, or (ii) recalculate using a flux to passive 0.0049 instead of 0.049. Option (ii) is preferred, but is not mandatory, since the other parts bring significant results; note that option (i) would not affect the summary nor the conclusion.

Actually this was only a typo mistakes in the manuscript but we carefully checked the code and it was correct. We therefore corrected the manuscript.

Minor and typographical points.

Table 1. Be clear in the legend on what was averaged. Do "over the profiles" means a calculated mean for (0- 2.0 m)"?

We did not have data up to 2m so we calculated a mean for the different available layers that we applied to the entire profile of the model (0-2.0 m). We clarified the legend: "Table 1. General description of the studied sites. The mean bulk density, pH and clay fraction values calculated from the different soil layers depths available in the data were used as input for each site. For Mons and Feucherolles sites, min and max values of pH and clay fraction are provided between brackets"

Table 3 Data are in kg C m⁻³, which is a unit for local concentration, not for carbon stock. Is it: (a) kg C m⁻², i.e., the carbon stock per unit area; or (b) the average concentration over the 0-2.0 m profile (then the Stock would be 2 times the mean concentration value)? Option (a) would be preferred.

The table 3 was modified and all the results are now presented in kg C m⁻².

Line 786 Legend fig.6: indicate the variable in object (= F14C).

This is now added in the legend

Line 134. A brief statement of the formalism and parameterization of the priming would be welcome.

We modified the text in the revised version as following: "Briefly, priming is described following equation 1.

$$\frac{\partial SOC_{i,z}}{\partial t} = DOC_{Recycled,i,j}(t) - k_{SOC,i} \times (1 - e^{-c \times LOC_z(t)}) \times SOC(t)_{i,z} \times \theta(t) \times \tau(t) \quad (1)$$

with $DOC_{recycled}$ being the unrespired DOC that is redistributed into the pool i considered for each soil layer z in $g\ C\ m^{-2}\ days^{-1}$, k_{SOC} being a SOC decomposition rate constant ($days^{-1}$), and LOC being the stock of labile organic C defined as the sum of the C pools with a higher decomposition rate than the pool considered within each soil layer z . We therefore considered that for the active carbon pool LOC is the litter and DOC , but for the slow carbon pool LOC is the sum of the litter, DOC and so on. Finally, c is a parameter controlling the impact of the LOC pool on the SOC mineralization rate, i.e., the priming effect. The equation was parameterized based on soil incubations data and evaluated over litter manipulation experiments (Guenet et al. 2016)."

Line 179. In Eq (6), STRUC was excluded of total 14C. Was it? In the Century model, STRUC usually accounts for 10- 20% of C in 0-20 cm layer, and is therefore non negligible (your figure 7). It is considered as retrieved as material < 2mm (for a large part) and therefore often included in the "measured" total carbon. This exclusion may affect the comparison between observed and modelled values of F14.

In the model, structural litter may come from leaves or root litter production of the ongoing year. Soil scientists, before measurements, generally remove it. We agreed with the reviewer that a part can still be present after few years but we were not able to clearly define a time

step when structural litter is less than 2mm and therefore integrated in the measurements. To avoid overestimation of modern C we decided to not integrate the structural litter in the final calculation. If needed we can perform a sensitivity analysis adding or not the structural litter in the final calculation to estimate the impacts on our results.

Line 218. "OCC (wt/wt)" would be better than "OCC (wt %)"

Done

Line 232. "turnover rate" is an unclear term (might be the reciprocal of turnover time). Here turnover time?

Correction is done in the revised version of the manuscript.

Lines 314-326 and throughout: MSD values aren't in kg C m⁻³, but in kg² C m⁻⁶ (variance not standard deviation); or use squareroot(MSC)

We corrected the MSD units in the revised version of the manuscript.

Line 325. Arenosols are not very specific and are broadly represented on the planet.

Remove "for such specific conditions". Replace by "Probably due to an overestimation of decay rates by ORCHIDEE in sandy soils?"

This is now corrected in the revised version of the manuscript.

Lines 367-401. See Major comment.

As explained above, the error was in the table but not in the model.

Do is the boundary condition at depth 2.0 for constant diffusion affect the base of the profiles?

It is difficult to answer this question because changing the soil depth of the model would not only affect the carbon but also the hydrology, the plant uptake and in fine the carbon inputs.

Typography

All the typo mistakes are corrected in the revised version.

Line 71. "this"

Line 158. verify in the final edition the greek symbol delta (not ok in my pdf)

Line 186. "1.5_m" (= separate the units throughout)

Line 227. "(2016)" (=no square brackets)

Line 242. "et al. (2014)" (spaces)

Line 255. Point missing; also lines 326, 337 ... check.

Line 315: spaces before and after "=" (throughout the text)

Line 447 'processes

Answer to comments from the reviewer #2.

We thank reviewer for the constructive evaluation of the manuscript. Please find below our answers to questions/comments. Comments from the reviewer were left intentionally in this document and written in roman font. Our answers are written in italics.

Anonymous Referee #2

Received and published: 18 August 2018

This paper presents ORCHIDEE-SOM-¹⁴C, a new version of the IPSL-Land Surface Model, and tests it against data from four different sites. It makes an important contribution by implementing the isotopic tracer ¹⁴C in the model. This is a valuable addition to the ORCHIDEE-SOM model, which simulates depth-resolved soil carbon dynamics from 0-2m below the surface. The authors also demonstrate how the new model can be used to constrain SOC turnover times and internal model processes. In particular, they implement two variations on the model (“Model_Test_He” and “Model_Test_Diffusion”).

They follow the suggestions of He et al (2016) to slow turnover in the passive pool and reduce the flux from the slow to passive pool (pending comment by reviewer #1).

They also implement a version of the model with depth-dependent bioturbation rate following Jagercikova et al (2014). Conceptually, this paper is a nice demonstration of how F¹⁴C data could be used for comparison against different model implementations.

However, there are significant issues which should be addressed both with the implementation (see Reviewer #1 comments) and interpretation of the results (see below) prior to publication.

Thanks for the positive comments please see our answer to reviewer #1.

In its current form, this paper does not convincingly demonstrate that there are meaningful differences in the modeled profiles across sites, or that any differences reflect the modeled differences in climate, vegetation or soil properties. Figures 3 and 4 demonstrate that the model can broadly fit a generic soil profile. However, it is unclear if the model can reliably capture differences between sites (for example, in Fig 3, the model reasonably fits only two of the four profiles). Comparison to a somewhat larger number of published soil F¹⁴C profiles is needed to support current statements that the model can “reproduce soil organic carbon stocks and radiocarbon profiles” (for example, line 29). This additional analysis would significantly strengthen the paper. It would also be particularly interesting to see if the model is able to capture the wide differences in bulk soil ¹⁴C seen across soil taxa (for example as explored in Mathieu et al (2015)). Alternatively, if the authors feel that comparison to a wider suite of soil profiles is beyond the scope of the current work, the current model-data comparison should be rephrased as a proof-of-concept contribution. In either case, the discussion should address potential controls on the soil F¹⁴C profiles (for both data and model). For example, despite the important role of mineralogy and clay content in controlling the age of soil C, these topics are not mentioned in the current discussion. Relatedly, more discussion and exploration of the model processes and parameters that control the ¹⁴C profiles would be an important addition to this paper. Although I acknowledge that comparison to a wider suite of soil profiles may be beyond the scope of the current work, I would like to see more exploration and discussion of these issues prior to publication.

We agree that such an evaluation would be a good step forwards but one the difficulty to run the model over such a large database is that very often some boundaries conditions of the model are missing and we have to estimate them with large scale database that may not be accurate for a given site. In this study we decided to carefully choose some sites which have enough data to feed the model and which are also representative of different situations. We aimed to go for such large-scale evaluation but we thought that it would have been more useful to have first a model description papers evaluated on well-chosen site. We changed several parts of the document in the revised version of the manuscript to explore more this weakness of our study see for example:

“Nevertheless, the model evaluation performed here on only four sites should be considered as proof of concept and more in depth evaluation are needed, in particular using a large ^{14}C database available at global scale (Balesdent et al., 2018; Mathieu et al., 2015). Indeed, the $F^{14}\text{C}$ is largely controlled by pedo-climatic conditions such as clay content, climate and mineralogy (Mathieu et al., 2015) and the range of situations we covered here is relatively limited.”

or “Furthermore, here we used only one averaged value over the soil profile for soil boundary conditions (texture, pH, bulk density) but those variables are known to impact the $F^{14}\text{C}$ (Mathieu et al., 2015) and change with depth (Barré et al., 2009) and depth-varying boundary conditions may also help to improve the model.””

The authors make a good case for the addition of depth-varying parameters, both conceptually (eg line 69) and in the results, by making the important contribution of implementing He et al's suggested parameters in a depth-dependent context and updating the diffusion formulation. However, although the updated diffusion formulation is a key contribution of the paper, the impact of this model improvement should not be overstated, as the difference between the two different model profiles relative to the data is not large (fig 3 &4). The modest gains suggest that adding other depth-varying processes in the future could be valuable. Although implementation of depth-varying parameters is clearly important, diffusion alone is not a singular model fix, and the discussion and conclusion should be broadened where possible to reflect this (for example, “mainly for diffusion” in line 40 and 468 is misleading/overstated).

We agree with this statement and we add a paragraph in the conclusion to detail what should be the next step in the implementation of depth-varying parameters: “Here we presented the effect of a depth-varying diffusion constant but other parameters are depth dependent and should be represented in the next version of the model. For instance, belowground litter production in the model is simply represented by an exponential law without any representation of the effect of resource distribution on root profile (e.g. water or nutrients). This is a complex task in a land surface model aiming at running at large scale with a classical resolution of 0.5° but the soil modules of land surface models are quite sensitive to the NPP (Camino-Serrano et al., 2018; Todd-Brown et al., 2013) and a better constraint on the profile of the below ground litter production would probably improve the model performance.”

I agree with Reviewer #1 on the major technical issue presented. This should be corrected prior to publication. The contribution of implementing the He et al (2016) suggested parameters is a good idea, and a nice contribution to the paper, so I would suggest retaining this model fit after updating the values as suggested by reviewer #1.

In general, figures could be made more professional, and a careful reading for grammatical errors is needed prior to publication.

The error was a typo mistake in the manuscript but we carefully checked the code and it was correct.

In summary, this manuscript should be considered for publication after major revisions, including the technical fix presented by reviewer #1, model comparison to additional soil profiles, and/or an updated discussion of the results. Minor comments are listed below.

Specific comments:

Line 40 & 468: “mainly for diffusion” is misleading as discussed above

This is removed in the revised version.

Lines 71-84: In introduction, cite other work using radiocarbon profiles to constrain soil models (e.g. Braakhekke et al, 2014; Ahrens et al, 2015)

We added the citation the papers suggested by the reviewer: “Different authors have already successfully implemented radiocarbon in soil models and were able to clearly show that the introduction of pools with turnover time of thousands of year were unnecessary to fit radiocarbon data (Ahrens et al., 2015) whereas Braakhekke et al., (2014) showed that after a reparameterization of the models based on radiocarbon data the prediction of their model was quite different with more carbon in top soil and less in deep soil compared to the model without radiocarbon.”

Line 136-137: Please clarify, as this seems contradictory: “SOC diffusion is actually a representation of bioturbation processes (animal (and plant) activity), whereas DOC diffuses through concentration gradients.” This text suggests that implementation of SOC diffusion would not be based on a concentration gradient, while the Fick’s law formulation provided (138-140) relies on a concentration gradient. Also, what do you mean by “the amount of carbon in the pool subject to transport”?

Both are based on a concentration gradient but the mechanisms we aimed to represent are different since it is bioturbation for the SOC whereas it is “real” diffusion for the DOC. We clarified the sentence: “SOC diffusion is actually a representation of bioturbation processes (animal (and plant) activity), whereas DOC relies more on a non-biological diffusion. Both diffuse through concentration gradients.”

Line 181...: 14C data collection:

-Please clarify: was new data collected for this paper or is this published elsewhere?
More details are now given see answer below.

-Please include a table of ^{14}C data values, including sampling depth increments
We added the table 5 to present those data in the revised version of the manuscript.

-Please provide more methods details on soil collection and processing or reference to appropriate publication.

For the French sites information can be found in Jagercikova, M., S. Cornu, D. Bourlès, O. Evrard, C. Hatté, and J. Balesdent (2017), Quantification of vertical solid matter transfers in soils during pedogenesis by a multi-tracer approach, J. Soils Sediments, 17(2), 408–422, doi:10.1007/s11368-016-1560-9. The information is now added.

For the two other sites, data are not published yet so we added more details. See for instance for the Misiones sites: “Details on measurements and sampling can be found in Tifafi et al., in prep. Briefly, the soil was sampled in May 2015 at different depth: 0-5cm, 5-10cm, 10-15cm, 15-20cm, 20-30cm, 30-40cm, 40-50cm, 50-60cm, 60-80cm, 80-100cm. All sampled were crushed and air-dried. Once in the laboratory, they were homogenized, crushed, randomly subsampled and sieved at 200 μm . Then ^{14}C measurements were made using a new Compact Radiocarbon System called ECHoMICADAS (Environment, Climate, Human, Mini Carbon Dating System) following the recommendation of Tisnérat-Laborde et al., (2015).”

-How were litter and roots handled? Included/excluded? How does that correspond to model results?

Roots were removed when visible. In the model we used only the active, slow and passive pools to calculate the $F^{14}\text{C}$ but as mentioned by reviewer #1 structural litter might have been included in the calculation. Nevertheless, structural litter in the model can be part of the litter produced during the on-going year but can also be few years old. Fix a threshold to determine which part of the structural litter would have been included needs underlying assumptions difficult to test. We therefore considered that only the soil carbon pools must be included in the calculation.

Line 245-255: How are soil $F^{14}\text{C}$ values handled in the spinup? What is the potential influence on initial soil ^{14}C values? Spinup is only ~ 2 half-lives of ^{14}C and doesn't consider atmospheric variation prior to 1700.

$F^{14}\text{C}$ were considered as stable before 1700. We considered this is a reasonable assumption since the variations observed from 1700 are mainly anthropogenic. The initialization procedure may indeed impact the results. If needed, we can perform a sensitivity analysis to the initial $F^{14}\text{C}$.

Line 301: Please mention somewhere how comparisons are made between data and model, given differences in depths

We added this information: “The intervals of soil depth of the model outputs and the measurements were homogenized by interpolating linearly the data to common depth intervals defined for each site. The simulations and data were then compared for each depth interval.”

Line 309-313 & Table 3: Visually, and discussed in the text, the sites Misiones and Feucherolles appear to have quite good fits for total soil carbon, while the fit is the worst for Mons, and also poor for Kissoko. However, the correlation coefficients are highest for Mons, but lowest for Kissoko. Is this a meaningful metric?

The good correlation coefficient for Mons is due to the relative good representation of the shape of the profile even though the mean bias is quite important as it is shown in Fig. 4. To clarify this point we added few words on this aspect at: “The correlation coefficient for Mons is relatively high compared to other site (Table 3) whereas Fig. 3 shows that the model performance was not very good for this site. This is mainly due to a large SB whereas other MSD components were rather low.”

Table 3&4: Is there a reason all values have been rounded to end in .05 or .00?

It was pure random and following the recommendation of reviewer #1 we change the units of the total carbon from kg C m^{-3} in kg C m^{-2} and the values from Table 3 do not all finished by .00 or .05

Line 320-326/Fig 3: Any comments on why the model does so well in one French Luvisol (Feucherolles) and so poorly on the other (Mons) for total soil carbon? From the site description the sites sound very similar.

This model like all the models following a similar structure are quite sensitive to the litter production. For Mons a net primary production (NPP) of $6.7 \text{ t ha}^{-1} \text{ yr}^{-1}$ was estimated by the technical institute for pasture in this region of France based on the annual yields, whereas the model predicts a NPP of $7.5 \text{ t ha}^{-1} \text{ yr}^{-1}$. The large over estimation might be a consequence of a bias in NPP. As far as we know no NPP estimation is available for Feucherolles. We added this information: “For Mons a net primary production (NPP) of $6.7 \text{ t ha}^{-1} \text{ yr}^{-1}$ was estimated by the technical institute for pasture in this region of France based on the annual yields, whereas the model predicts a NPP of $7.5 \text{ t ha}^{-1} \text{ yr}^{-1}$. The large overestimation of the SOC stocks may therefore be due to an overestimation of the NPP.”

Line 334: “The vertical profile of the SOC stock simulated was thereby globally not very far from that of the data”. This seems like an overstatement based on results in Table 3. For example, although reported model total soil carbon is 1.7 and 2.1 overestimated at two sites with better fits, it is overestimated by a factor of 8.5 and 4.6 at the other two sites.

We rephrase to avoid overstatement. See line: “The vertical profiles of the SOC stock were fairly represented by the model”

Fig 3: Relatedly, what depth ranges are used for comparison between data and model? How does this influence the results? For example, model and data look quite similar in Fig 3 for Misiones and Feucherolles, but the mean total soil carbon is reported to be overestimated by nearly a factor of 2.

This information is now added in the method section “The intervals of soil depth of the model outputs and the measurements were homogenized by interpolating linearly the data to common depth intervals defined for each site. The simulations and data were then compared for each depth interval.”

Lines 364-366: Interesting, and nice to build on He et al (2016) using a depth-resolved approach

Thanks for the positive comments.

Line 392: More explanation of the results/implications of the priming effect mentioned here would be interesting, but not required

Since we did not run our model without priming we prefer to not increase the discussion section as it is to avoid over-interpretation.

Lines 407-408: “Using a fixed diffusion constant implicitly suggests that soil fauna activity is uniform over the entire soil profile”. Please add more explanation of the link between fauna activity and the diffusion term formulation for the reader. This diffusion term will vary with depth and across sites, because the Fick’s law formulation also relies on the concentration gradient with depth. For example, in Kissoko, for much of the profile there is almost no change in total soil carbon with depth, so the diffusion term here would be zero. Does that imply that there is no soil fauna activity? Or simply that soil fauna activity does not result in a change in the soil carbon profile?

Here we were wanted to talk about the diffusion rate and not the entire diffusion fluxes. We clarified the sentence: “Fick’s law of diffusion is classically used in models to represent bioturbation assuming that soil fauna activity may be represented following the Fick’s law of diffusion (Elzein and Balesdent, 1995; Guenet et al., 2013; Koven et al., 2013; O’Brien and Stout, 1978; Wynn et al., 2005). Using a fixed diffusion constant (D in eq. 2) implicitly suggests that soil fauna activity is uniform over the entire soil profile. This is generally the case of several models of diffusion especially used at the level of an ecosystem (Bruun et al., 2007; Guimberteau et al., 2017; O’Brien and Stout, 1978). However, soil faunal activity vary naturally with depth and the diffusion constant should be depth-dependent (Jagercikova et al., 2014).”

Lines 449-454: Well-stated summary of model contributions

Thanks.

Line 457: Please mention and cite any other land surface models that incorporate soil 14C either here or in introduction.

We added a paper by Koven et al., 2013 in Biogeosciences.

Lines 466-468: “This suggests that, from now on, model improvements should mainly focus on a depth dependent parameterization, mainly for diffusion.” Although diffusion did improve model results, the change was not dramatic. Please make sure the language used here reflects the results.

This was rephrased in the revised version

-Broadly, figure aesthetics should be updated to look more professional throughout prior to publication. For example:

-Fig 7. Please label x & y axis. Please write depth increments for each bar on y-axis instead of 1-11. Also, in some of the panels numbers 11 and 12 are cutoff (eg 1..)

-Fig 3-7: Use more professional titles and punctuation on figures (eg. rather than “Model_Control” , “Model_Test He”, etc.)

-Fig 7: It appears there are stray line numbers throughout the figures which will presumably be removed once the line numbers have been removed (eg fig 4,6,7)

-Update “litter structural below” and “litter metabolic below” to more clear and professional names

All the figures have redo to more professional aspects.

-Fig 7 is instructive and interesting. However, what is the reason for the “litter structural below” to decrease then increase again at the deepest depths in some of the profiles?

The question might be that the diffusion constant D in deep layers has very low values in deep soil because of the depth-varying equations we used. Therefore the diffusion fluxes are quite limited in deep layers. Furthermore, in deep soil the temperature is rather stable and those layers don't face important temperature increase in summer leading to high decomposition rates. Then, in deep soil the decomposition is limited and diffusion is not strong enough to homogenize the profile.

Language Comments: A careful and significant reading for grammatical errors and typos is needed prior to publication. A large number of very small changes are required.

All the grammatical errors were corrected and a native English speaker read the revised manuscript.

Here are a few examples (not comprehensive):

Line 59: “simulate” should be “simulates”

Line 71: typo “thIS”

Lines 74-77: very confusingly worded sentence

Line 81: “have” should be “has”

Line 84: “because of the conceptual description by pools non measurable” – fix grammar

Line 92: “yielded for the abrupt increase of atmospheric ^{14}C concentration that doubles in 2-3 years.” -clarify language

Line 198: “Congo Republic” should be “Republic of Congo”

Line 337: Missing period at end of sentence

Lines 659-660: “over the profile according to total soil carbon” - Meaning is unclear

Additional references:

Ahrens et al (2015). Contribution of sorption, DOC transport and microbial interactions to the ^{14}C age of a soil organic carbon profile: Insights from a calibrated process model. *Soil Biology and Biochemistry*, 88. pp. 390-402.

Braakhekke et al (2014). The use of radiocarbon to constrain current and future soil organic matter turnover and transport in a temperate forest. *Journal of Geophysical Research: Biogeosciences*, 119(3).

Mathieu et al (2015). Deep soil carbon dynamics are driven more by soil type than by climate: a worldwide meta-analysis of radiocarbon profiles. *Global Change Biology*, 21. pp. 4278-4292.

Answer to comments from the reviewer #3.

We thank reviewer for the constructive evaluation of the manuscript. Please find below our answers to questions/comments. Comments from the reviewer were left intentionally in this document and written in roman font. Our answers are written in italics.

Anonymous Referee #3

Received and published: 23 August 2018

The cycling of organic matter through soil ecosystems is highly simplified in land surface models. This is a major source of uncertainty in projections of the terrestrial carbon sink under global climate change. Measurements of the radioactive carbon isotope ^{14}C provides a powerful constraint for soil carbon models which include a radiocarbon tracer component. This manuscript documents the addition of a radiocarbon tracer component into the ORCHIDEE land model in order to enable radiocarbon constraints in it and in the IPSL Earth System Model it is coupled with. This study then demonstrated applying this constraint to the model based on several vertically-resolved soil radiocarbon profiles.

General comments:

The paper represents a substantial advance in the ORCHIDEE/IPSL model, which is an important tool in climate science, and has broader implications for other models. As such, it is well within the scope of GMD, and would represent a meaningful contribution to the field. However, there are several issues that would need to be addressed before I could recommend it for publication. I have detailed these issues below, and I hope that by addressing them, the authors will return with an improved presentation of this worthwhile research.

Thanks for the positive comments.

Major issue 1:

There are a couple of major issues with the Model_Test_He experiment. He et al (2006) suggested scaling the passive pool turnover time in IPSL/ORCHIDEE by 14, while scaling the slow-to-passive transfer coefficient by 0.07. I applaud the authors' effort to test this suggestion. However, the manuscript lacks a detailed explanation of exactly which quantities

were scaled, and which of the arrows in Figure 1 corresponds with the first column of Table 2. The reduced complexity models of He et al consisted of three pools in series, whereas Figure 1 implies that ORCHIDEE has three soil pools that each independently exchange with a single pool of free DOC. Therefore, it seems that ORCHIDEE does not have a single transfer coefficient between slow and passive pools.

To avoid making the manuscript too long we did not give all the details of the model construction (we mainly refer to Camino-Serrano et al. 2018) and mainly information on the ¹⁴C-related part. Nevertheless, in the model when the decomposed SOC goes to DOC, we keep track of the pool where it came from and the redistribution of the DOC once decomposed into SOC follows the same parameterization than the ORCHIDEE version incorporated to the IPSL-ESM used by He et al., (2016). We therefore considered that using those parameter values still makes sense.

Furthermore, as pointed out in RC1, there seems to be an arithmetic error in the scaling of this transfer coefficient. The first and third rows of Table 2 imply that ORCHIDEE has some parameter with a value of 0.07 (this parameter being what needs improved explanation). Multiplying this by the scaling factor suggested in He et al would yield 0.0049, but it seems that 0.049 was used instead. The result is that the passive pool turnover time is increased by an order of magnitude without an equivalent adjustment to the inputs to this pool, leading to a large accumulation of radiocarbon-depleted SOM. This explains why the Model_Test_He experiment is so far off in Figures 3 and 5, and why the standard bias is so high in Figures 4 and 6.

I would encourage the authors to re-run this experiment with the correct values and keep it in the manuscript (and, unlike RC1, I have no problem with the name). I understand that the recommended values were for a previous version of IPSL/ORCHIDEE, and that some of the changes since then (yielding ORCHIDEE-SOM, detailed in Camino-Serano et al , 2017) make the recommended changes superfluous by accounting for priming. Nevertheless, I think that testing these recommendations is a worthwhile exercise, even with this updated model version, and I would be interested in seeing it done correctly.

The error was a typo mistake in the manuscript but we carefully checked the code and we used the good value for the simulations.

Major issue 2:

There is insufficient explanation of the depths at which the observational (field) data were sampled, and how that was compared with the model output. Figure 1 explains sufficiently the depth of the soil layers in the ORCHIDEE model (though an explanation in the main text would be welcome as well). The depth of the field measurements can be seen in Figures 3 and 5, but not with enough resolution to really understand. Was each field profile sampled at the exact same depths as the layers in ORCHIDEE, or is there some interpolation going on between one or the other?

More information is now given in the method section: “The intervals of soil depth of the model outputs and the measurements were homogenized by interpolating linearly the data to common depth intervals defined for each site. The simulations and data were then compared for each depth interval.”

The statistics in Section 2.6 are all over a dimension i , which I assume to represent the layers over depth, but this is not clearly stated. Given the importance of this i , we need more detail as to what it is.

We added this information: “ x refers to the model outputs and y to the measurements, while i refers to soil depth.”

I would prefer to see an additional table or additional information in Table 1 to indicate how many samples were taken at each site and at what depths. And, most importantly, some explanation in the methods of how layer depths were harmonized between the model and observations, including an indication of the size of I (i.e., the n in the equations of Section 2.6).

We added a table given the different layers and we add information on the interpolation method we used to compare data and model outputs (see previous answer).

Moreover, the specific depths at which the observed and modeled layers are compared should be clearly visible in Figures 3 and 5. The field observations are shown as points, with a single depth. Were measurements taken just at those single depths? Or were entire layers sampled with an upper and lower boundary depth? The model is presumably providing an average concentration of carbon (and radiocarbon) for entire layers, but the lines in Figures 3 and 5 make it seem like the data are continuous rather than discrete.

Since in the revised version of the manuscript we give both the interpolation method and the layers depth we hope that enough information is given to avoid misunderstanding.

Finally, the absence of explicit field data hinders the reproducibility of the study. The methods are described sufficiently to reproduce the study, and the model source code is available (though the web link has a problem, see below). But the study cannot be truly replicated without having access to the field data that were used. Including the field data in tabular format (perhaps as supplementary material) would go a long way toward making the methods more understandable and facilitating reproducibility.

Some of the data are already published and we refer now to the proper citation but others will be presented in an in prep. manuscript by the same authors. More details are now given. For instance: “Details on measurements and sampling can be found in Tifafi et al., In prep. Briefly, the soil was sampled in May 2015 at different depth: 0-5cm, 5-10cm, 10-15cm, 15-20cm, 20-30cm, 30-40cm, 40-50cm, 50-60cm, 60-80cm, 80-100cm. All sampled were crushed and air-dried. Once in the laboratory, they were homogenized, crushed, randomly subsampled and sieved at 200 μ m. Then ^{14}C measurements were made using a new Compact Radiocarbon System called ECHoMICADAS (Environment, Climate, Human, Mini Carbon Dating System) following the recommendations of Tisnérat-Laborde et al., (2015).”

Major issue 3:

The authors provide some interpretation of each of the individual results in Section 3, but the manuscript lacks an overall discussion of the big-picture implications of these results and how they serve to advance scientific knowledge. The introduction section provides a compelling motivation for the study, but the manuscript lacks a sufficient discussion of how the current study informs these issues, what can be learned about SOM processes and soil-climate interactions, and what the implications are for the use of ESMs to project future climate change. I would like to see an expanded discussion of how these results fit in with the larger body of literature. The authors neglect to acknowledge that radiocarbon has already been implemented in a well known ESM (the Community Earth System Mode, CESM), and therefore do not discuss how their results relate to the existing work. The authors do cite the paper that would be relevant for this (Koven et al., 2013) in the context of diffusion representing bioturbation (line 406), but I would like to see an expanded discussion of how the results from the two papers potentially inform each other.

The paper by Koven et al., 2013 is now properly cited to its contribution “ORCHIDEE-SOM- ^{14}C , is one of the first land surface models that incorporates the ^{14}C dynamic in the soil (Koven et al., 2013).”

And more discussion has been added. For instance: “We limited our work here to depth-varying diffusion, but other parameters are also depth dependent and should be represented as such in the next version of the model. For instance, belowground litter production in the model is simply represented by an exponential law without any representation of the effect of resource distribution on root profile (e.g. water or nutrients). This is a complex task in a land surface model running at large scale with a classical resolution of 0.5° , but the soil modules of land surface models are quite sensitive to the NPP (Camino-Serrano et al., 2018; Todd-Brown et al., 2013) and a better constraint on the profile of the below ground litter production would likely improve the model performance. Furthermore, here we used only one averaged value over the soil profile for soil boundary conditions (texture, pH, bulk density) but those variables are known to impact the $F^{14}\text{C}$ (Mathieu et al., 2015) and change with depth (Barré et al., 2009) and depth-varying boundary conditions may also help to improve the model.”

Or “Nevertheless, the model evaluation performed here on only four sites should be considered as proof of concept and more in depth evaluation are needed, in particular using a large ^{14}C database available at global scale (Balesdent et al., 2018; Mathieu et al., 2015). Indeed, the $F^{14}\text{C}$ is largely controlled by pedo-climatic conditions such as clay content, climate and mineralogy (Mathieu et al., 2015) and the range of situations we covered here is relatively limited.”

Minor issues and technical corrections:

All the minor issues and technical corrections were taken into account.

Abbreviations: there are some abbreviations that are used without an explicit definition. In some cases, they are defined later, but they should be defined in the first instance of use. I

would avoid abbreviating SOC and SOM in the abstract, since neither one is used again in the abstract and just use the full text instead (but then define the abbreviation and begin using it when it first appears in the main body of the text). The abbreviation "F14C" for fraction modern is used in the abstract, but not explicitly defined. "IPSL" is used several times before it is defined on line 105, and ORCHIDEE is never defined.

Line 71: spurious capitalization in the word "this"

Line 74: The sentence that begins on this line is too long, and should be broken up into at least two sentences to be understandable.

Line 75: "implementing" should be "to implement"

Lines 91-92: The decades should not have apostrophes (e.g., 1950s, not 1950's)

Line 93: Remove the word "since"

Line 94: Should be "As WITH any other carbon isotopes"

Lines 106–113: I am not sure how useful it is to list the names of the sub-components of ORCHIDEE without any further indication of how these components fit in to the present study. Instead, I would prefer to see a description of how ORCHIDEE fits into the larger ESM (e.g., which fluxes and state variables coupled it with the atmospheric model).

Line 158: There is some rendering issue with the δ (delta) symbol in $\delta^{13}\text{C}$; please double check.

Line 162: The abbreviations A_{sample} and A_{ref} should be explicitly defined for the sake of the reader who may be new to the concepts of radiocarbon.

Lines 167–179: There is some inconsistency between the main text and the equations regarding abbreviations. The text uses " ^{14}C " while the equations use "carbon14". I believe these are supposed to represent the same thing, and should therefore have the same abbreviations for clarity.

Lines 184–212: Some measurements include a space between the quantity and the units (e.g., "680 mm" on line 185) while others do not (e.g., "1.5m" on line 186)

Line 192: Define the abbreviation LSCE

Line 194: Define the abbreviation LMC14

Line 197: Define the abbreviation SOERE F-ORE-T

Line 232: The term "turnover rate" is ambiguous. I assume the authors mean "turnover time" since this is what He et al suggest should be scaled by 14, which would be the inverse of the decay "rate".

Line 252: What assumptions were made about the atmospheric ^{14}C content during spinup?

Line 256: Were simulations actually run at a yearly time step? Section 2.1 indicates that some model components have a much shorter time step. Also, for comparison with the field data, was the final (2011) time step used?

Lines 339–340: Something is wrong with this sentence grammatically, which makes it difficult to interpret.

Lines 392–393: The 50

Line 408: Remove the word "fact" or add the word "in" before it.

Line 465-466: Please revise this sentence for grammatical accuracy.

Line 477: The provided website address links to a page that has issues with the SSL certificate, and will not load in any web browser without having to make a security exception. Providing the link as http rather than https would fix this issue, though the preferred solution would be maintain the https link and insure that the website has a valid SSL certificate.

The use of radiocarbon ^{14}C to constrain carbon dynamics in the soil module of the land surface model ORCHIDEE (SVN r5165)

Marwa Tifafi¹, Marta Camino-Serrano^{2,3}, Christine Hatté¹, Hector Morras⁴, Lucas Moretti⁵, Sebastián Barbaro⁵, Sophie Cornu⁶, Bertrand Guenet¹

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France.

²CREAF, Cerdanyola del Vallès, 08193, Catalonia, Spain

³CSIC, Global Ecology Unit CREAF-CSIC-UAB, Bellaterra 08193, Catalonia, Spain

⁴INTA-CIRN, Instituto de Suelos, 1712 Castelar, Buenos Aires, Argentina

⁵INTA-EEA Cerro Azul, 3313 Cerro Azul, Misiones, Argentina

⁶Aix Marseille Univ, CNRS, IRD, INRA, Coll France, CEREGE, Aix-en-Provence, France

Corresponding authors: Marwa Tifafi (marwa.tifafi@lsce.ipsl.fr)

Abstract. Despite the importance of soil as a large component of the terrestrial ecosystems, the soil compartments are not well represented in the Land Surface Models (LSMs). Indeed, soils in current LSMs are generally represented based on a very simplified schema that can induce a misrepresentation of the deep dynamics of soil carbon. Here, we present a new version of the Institut Pierre Simon Laplace (IPSL) Land Surface Model called ORCHIDEE-SOM (ORganizing Carbon and Hydrology in Dynamic EcosystEms-Soil Organic Matter), incorporating the ^{14}C dynamic in the soil. ORCHIDEE-SOM first simulates soil carbon dynamics for different layers, down to 2 m depth. Second, concentration of dissolved organic carbon and its transport are modeled. Finally, soil organic carbon decomposition is considered taking into account the priming effect.

After implementing ^{14}C in the soil module of the model, we evaluated model outputs against observations of soil organic carbon and modern ^{14}C fraction ($F^{14}\text{C}$) for different sites with different characteristics. The model managed to reproduce the soil organic carbon stocks and the $F^{14}\text{C}$ along the vertical profiles for the sites examined. However, an overestimation of the total carbon stock was noted, primarily on the surface layer. Due to ^{14}C , it is possible to probe carbon age in the soil, which was found to underestimated. Thereafter, two different tests on this new version have been established. The first was to increase carbon residence time of the passive pool and decrease the flux from the slow pool to the passive pool. The second was to establish an equation of diffusion, initially constant throughout the profile, making it vary exponentially as a function of depth. The first modifications did not improve the capacity of the model to reproduce observations whereas the second test improved both estimation of surface soil carbon stock as well as soil carbon age. This demonstrates that we should focus more on vertical variation of soil parameters as a function of depth, in order to upgrade the representation of global carbon cycle in LSMs, thereby helping to improve predictions of the of soil organic carbon to environmental changes.

Bertrand Guenet 30/10/y 10:10

Supprimé: -

Bertrand Guenet 30/10/y 15:58

Supprimé: -

Auteur inconnu 31/10/y 07:58

Supprimé: ,

Auteur inconnu 31/10/y 07:58

Supprimé: ,

Bertrand Guenet 30/10/y 15:57

Supprimé: (DOC)

Bertrand Guenet 30/10/y 15:57

Supprimé: (SOC)

Auteur inconnu 31/10/y 07:59

Supprimé: the

Bertrand Guenet 30/10/y 15:58

Supprimé: activity

Auteur inconnu 31/10/y 07:59

Supprimé: chosen here

Auteur inconnu 31/10/y 08:00

Supprimé: but was mostly marked

Auteur inconnu 31/10/y 08:01

Supprimé: Then, thanks to the introduction of

Auteur inconnu 31/10/y 08:01

Supprimé: has

Auteur inconnu 31/10/y 08:02

Supprimé: been possible to highlight an underestimation of the age of carbon in the soil

Auteur inconnu 31/10/y 08:05

Supprimé: showed a decrease of the soil carbon stock overestimation, especially at the surface and an improvement of the estimates of the carbon age

Auteur inconnu 31/10/y 08:06

Supprimé: assumes

Bertrand Guenet 31/10/y 19:17

Supprimé: (?)

Bertrand Guenet 30/10/y 11:23

Supprimé: mainly for diffusion,

Bertrand Guenet 31/10/y 19:17

Supprimé: future response

Bertrand Guenet 31/10/y 19:17

Supprimé: global warming

Bertrand Guenet 31/10/y 19:18

Supprimé: -

... [1]

68 1 Introduction

69 The complexity of the mechanisms involved in controlling soil activity (Jastrow et al., 2007)
70 and therefore the carbon flux from the soil to the atmosphere makes predicting the response of
71 these systems to climate change extremely complex. Thus our ability to predict future changes
72 in carbon stocks in soils using global climate models is currently heavily criticized (Todd-
73 Brown et al., 2013; Wieder et al., 2013). Indeed, Earth System Models (ESMs) are
74 increasingly used today in order to predict the future evolution of the climate. For instance,
75 results of a set of ESMs are taken into account within the Intergovernmental Panel on Climate
76 Change (IPCC) (Taylor et al., 2012) for assessment of the impacts of climate change and
77 design of mitigation strategies. Hence, their predictions need to be as accurate as possible.
78 These models represent the physical, chemical and biological processes within and between
79 the atmosphere, ocean and terrestrial biosphere. They allow us to follow and understand both
80 the effect of the climate on carbon storage and vice versa. However, ESMs are continuously
81 under development and some key processes in the global carbon cycle are still missing or not
82 represented with the necessary details. One of the components of an ESM is the land surface
83 model (LSM). This component primarily manages the carbon cycle, energy and water on land
84 and simulates the carbon exchange between the land surface and the atmosphere, namely the
85 gross primary production (GPP), the autotrophic and heterotrophic respiration.

86 Despite the importance of soils as a large component of the global carbon storage, soil
87 compartments are not well represented in LSMs (Todd-Brown et al., 2013). Indeed, carbon
88 dynamics in soil described in LSMs are based on the “Century” (Parton et al., 1987) or
89 Roth-C models (Coleman et al., 1997) where soil carbon is represented as several pools with
90 different turnover rates for each pool. Carbon is decomposed in each pool, one part of which
91 is then transferred from one pool to another and the other part is lost through heterotrophic
92 respiration. In addition, soils are generally represented as a single-layer box in LSMs that do
93 not take into account the evolution and variation of soil organic processes as a function of
94 depth (Todd-Brown et al., 2013).

95 One way to reconcile this simplified representation of carbon dynamics of the models with the
96 complexity of the data collected in the field is to integrate isotopic tracers into the models
97 themselves and thus facilitate the comparison between model outputs and data (He et al.,
98 2016). Moreover, thanks to an additive constraints on the model structure, this may improve
99 the model performances. For instance, radiocarbon is an important tool for studying the
100 dynamics of soil organic matter (Trumbore, 2000). Indeed, ^{14}C data acquired from soil
101 organic matter provide complementary information on the dynamics (temporal dimension) of
102 soil organic matter. This tracer has the major advantage of being integrator of carbon
103 dynamics on long time scales (a few decades to several centuries). It is therefore a very
104 powerful tool to constrain conceptual schemes that may not be directly compared to variables
105 measured in the field (Elliott et al., 1996). Different authors have already successfully
106 implemented radiocarbon in soil models and were able to clearly show that the introduction of
107 pools with turnover time of thousands of year were unnecessary to fit radiocarbon data
108 (Ahrens et al., 2015) whereas Braakhekke et al., (2014) showed that after a reparameterization
109 of the models based on radiocarbon data the prediction of their model was quite different with
110 more carbon in top soil and less in deep soil compared to the model without radiocarbon.

Auteur inconnu 31/10/y 08:08

Supprimé: of the processes governing storage and destocking at variable time and space scales ...s currently heavily criticized ([2])

Bertrand Guenet 31/10/y 19:21

Supprimé: ...nd heterotrophic soil ([3])

Auteur inconnu 31/10/y 08:57

Supprimé: the...soil compartments ar ([4])

Bertrand Guenet 30/10/y 10:10

Supprimé: thIS

Auteur inconnu 31/10/y 08:59

Supprimé: to

Bertrand Guenet 31/10/y 19:21

Supprimé: Indeed, in order to be more pertinentrelevant to in evaluating the parameters and the equations of the newly implemented processes, it is of interest implementing carbon isotope dynamics in the model itself, this to facilitate the comparison between model outputs and available observations...Moreover,... but also... ([5])

Auteur inconnu 31/10/y 09:01

Supprimé: on...soil organic matter, ([6])

Bertrand Guenet 30/10/y 13:36

Supprimé: ve...the major advantage o ([7])

Auteur inconnu 31/10/y 09:03

Supprimé: succesfully...radiocarbon ([8])

Bertrand Guenet 30/10/y 11:30

Supprimé: (

Auteur inconnu 31/10/y 09:03

Supprimé: were

173 Radiocarbon is produced naturally at a constant rate in the upper atmosphere through
 174 bombardment of cosmic rays. It thus provides information on the dynamics of organic matter
 175 that has been stabilized by interaction with mineral surfaces and stored long enough for
 176 significant radioactive decay (Trumbore, 2000), as the half-life of ^{14}C is about 5730 years. We
 177 must also take into account radiocarbon produced during atmospheric tests of thermonuclear
 178 weapons in the early sixties (Delibrias et al., 1964; Hua et al., 2013). Atmospheric bomb
 179 testing in the late 1950s and early 1960s lead to an abrupt doubling of atmospheric ^{14}C
 180 concentration in a span of 2-3 years. Through exchange with ocean and terrestrial reservoirs,
 181 it has decreased but still remains above the natural background. As with any other carbon
 182 isotopes, this ^{14}C was metabolized by the vegetation and transferred to soil. By measuring ^{14}C
 183 activity of a soil sample, it is possible to evaluate the amount of carbon introduced into the
 184 soil since the 1960s (Balesdent and Guillet, 1982; Scharpenseel and Schiffmann, 1977).

185 In this study, we present a new version of the IPSL-Land Surface Model called ORCHIDEE-
 186 SOM incorporating ^{14}C dynamics in the soil. Thanks to this tracer, we can evaluate the SOC
 187 dynamics, in particular by looking at the ^{14}C peak produced by atmospheric weapons testing
 188 and observed in the soils at four different sites having different biomes.

189

190 2 Materials and methods

191 2.1 ORCHIDEE-SOM overview

192 ORCHIDEE is the Land Surface Model of the IPSL Earth System Model (Krinner et al.,
 193 2005). It is composed of three different modules. First, SECHIBA (Ducoudré et al., 1993;
 194 Rosnay and Polcher, 1998), the surface-vegetation-atmosphere transfer scheme, describes the
 195 soil water budget and energy and water exchanges. The time step of this module is 30 min.
 196 Second, the module of the vegetation dynamics has been taken from the dynamic global
 197 vegetation model LPJ (Sitch et al., 2003). The time step of this module is one year. Finally,
 198 the STOMATE (Saclay Toulouse Orsay Model for the Analysis of Terrestrial Ecosystems)
 199 module simulates vegetation phenology and carbon dynamics with a time step of one day.

200 ORCHIDEE can be run coupled to a global circulation model where the boundary conditions
 201 of the model are provided by the atmospheric modules (temperature, precipitation,
 202 atmospheric CO_2 concentration, etc.). In return ORCHIDEE provides the land surface carbon,
 203 energy and water fluxes. However, since our study focuses on changes in the land surface
 204 rather than on the interaction with climate, we ran ORCHIDEE in the off-line configuration.

205 In this case, atmospheric conditions such as temperature, humidity and wind are read from a
 206 meteorological dataset. The climate data CRUNCEP used for our study (6-hourly climate data
 207 over several years) were obtained from the combination of two existing datasets: the Climate
 208 Research Unit (CRU) (Mitchell et al., 2004) and the National Centers for Environmental
 209 Prediction (NCEP) (Kalnay et al., 1996).

210 Our starting point is a ORCHIDEE-SOM version based on the SVN r3340 (Krinner et al.,
 211 2005), which is presented in detail in Camino-Serrano et al. (2017). Figure 1 represents how
 212 the soil is described in this new version. Indeed, the major particularity of ORCHIDEE-SOM
 213 is that it simulates the dynamics of soil carbon for eleven layers from the surface to two

Auteur inconnu 31/10/y 09:04

Supprimé: First, there is the natural ... [9]

Bertrand Guenet 30/10/y 16:02

Supprimé: '... and early 1960'... yiel ... [10]

Auteur inconnu 31/10/y 09:07

Supprimé: increase... of atmospheric ... [11]

Bertrand Guenet 30/10/y 16:03

Supprimé: since ...ut still remains ab ... [12]

Auteur inconnu 31/10/y 09:11

Supprimé: the

Bertrand Guenet 30/10/y 15:58

Supprimé: (Institut Pierre Simon Laplace)

Auteur inconnu 31/10/y 09:12

Supprimé: S...rface-vegetation-atmo ... [13]

Auteur inconnu 31/10/y 09:14

Supprimé: ies...conditions of the mo ... [14]

Auteur inconnu 31/10/y 09:16

Supprimé: the... ORCHIDEE-SOM ... [15]

266 meters depth. First, litter is divided into four pools: metabolic or structural litter pools which
 267 can be found below or aboveground. Only the belowground litter is modeled on eleven levels,
 268 from surface to 2 m depth, as the aboveground litter layer has a fixed thickness of 10 mm.
 269 Second, SOC is divided into three pools (active, passive and slow), following Parton et al.
 270 (1988), which differ in their turnover rates and which are discretized into 11 layers up to a
 271 depth of two meters. Then, dissolved organic carbon (DOC) is represented as two pools and
 272 also discretized over 11 layers up to a depth of two meters: labile DOC has a high
 273 decomposition rate and recalcitrant DOC has a low decomposition rate (Camino-Serrano et
 274 al., 2018). Finally, another particularity of this version of ORCHIDEE-SOM is that the SOC
 275 decomposition is modified to account for the priming effect following Guenet et al. (2016).
 276 Briefly, priming is described following equation 1.

$$277 \frac{\partial SOC_{i,z}}{\partial t} = DOC_{Recycled,i,j}(t) - k_{SOC,i} \times (1 - e^{-c \times LOC_z(t)}) \times SOC(t)_{i,z} \times \theta(t) \times \tau(t) \quad (1)$$

278 with $DOC_{recycled}$ being the unrespired DOC that is redistributed into the pool i considered for
 279 each soil layer z in $g\ C\ m^{-2}\ days^{-1}$, k_{SOC} being a SOC decomposition rate constant ($days^{-1}$), and
 280 LOC being the stock of labile organic C defined as the sum of the C pools with a higher
 281 decomposition rate than the pool considered within each soil layer z . We therefore considered
 282 that for the active carbon pool LOC is the litter and DOC, but for the slow carbon pool LOC
 283 is the sum of the litter, DOC and so on. Finally, c is a parameter controlling the impact of the
 284 LOC pool on the SOC mineralization rate, i.e., the priming effect. The equation was
 285 parameterized based on soil incubations data and evaluated over litter manipulation
 286 experiments (Guenet et al. 2016).

287 Since the soil profile is divided into 11 layers, SOC and DOC transport following the
 288 diffusion must also be described. SOC diffusion is actually a representation of bioturbation
 289 processes (animal and plant activity), whereas DOC relies more on non-biological diffusion.
 290 Both diffuse through concentration gradients.

291 This is represented using the Fick's law (Braakhekke et al., 2011; Elzein and Balesdent, 1995;
 292 O'Brien and Stout, 1978; Wynn et al., 2005):

$$293 F_D = -D * \frac{\partial^2 C}{\partial z^2} \quad (2)$$

294 Where F_D is the flux of carbon transported by diffusion in $g\ C\ m^{-3}\ day^{-1}$, D is the diffusion
 295 coefficient ($m^2\ day^{-1}$) and C is the amount of carbon in the pool (DOC or SOC) subject to
 296 transport ($g\ C\ m^{-3}$). The diffusion coefficient is assumed to be constant across the soil profile
 297 in ORCHIDEE-SOM but the diffusion parameters (D) used in the equations for SOC and
 298 DOC can differ.

299 2.2 ORCHIDEE-SOM-¹⁴C

300 In ORCHIDEE-SOM, the different compartments (soil carbon input, litter, SOC, DOC and
 301 heterotrophic respiration) are presented as a matrix with a single dimension referring to the
 302 total carbon. In order to introduce the ¹⁴C, a new dimension has been added to all the
 303 variables cited above. Thus, all processes that apply to the total soil carbon are now also
 304 represented for ¹⁴C. We label this new version including ¹⁴C as ORCHIDEE-SOM-¹⁴C.

Auteur inconnu 31/10/y 09:19
Supprimé: 11
Auteur inconnu 31/10/y 09:19
Supprimé: horizons
Auteur inconnu 31/10/y 09:19
Supprimé: however
Auteur inconnu 31/10/y 09:19
Supprimé: ,
Auteur inconnu 31/10/y 09:19
Supprimé: ,
Auteur inconnu 31/10/y 09:20
Supprimé: into
Auteur inconnu 31/10/y 09:20
Supprimé: the
Auteur inconnu 31/10/y 09:20
Supprimé: with
Auteur inconnu 31/10/y 09:20
Supprimé: the
Auteur inconnu 31/10/y 09:20
Supprimé: with
Bertrand Guenet 31/10/y 19:28
Mis en forme: Default, Gauche, Espace
Après : 6 pt
Auteur inconnu 31/10/y 09:22
Supprimé: not
Auteur inconnu 31/10/y 09:22
Supprimé: -
Bertrand Guenet 31/10/y 19:27
Mis en forme: Non Exposant/ Indice
Auteur inconnu 31/10/y 09:50
Supprimé: is
Auteur inconnu 31/10/y 09:50
Supprimé: (
Auteur inconnu 31/10/y 09:50
Supprimé:)
Auteur inconnu 31/10/y 09:51
Supprimé: a
Bertrand Guenet 30/10/y 11:38
Supprimé: s
Bertrand Guenet 31/10/y 19:30
Supprimé: . . .
Bertrand Guenet 30/10/y 10:43
Supprimé: 1
Auteur inconnu 31/10/y 09:51
Supprimé: are
Auteur inconnu 31/10/y 09:51
Supprimé: different
Auteur inconnu 31/10/y 09:52
Supprimé: T
Auteur inconnu 31/10/y 09:52
Supprimé: the
Auteur inconnu 31/10/y 09:52
Supprimé: will be called

329 Several ways of reporting ^{14}C activity levels are available. We chose to use the *fraction*
 330 *modern*, with the $F^{14}\text{C}$ symbol as advocated by Reimer et al. (2004) rather than absolute
 331 concentration of ^{14}C (reported as Bq).

$$332 \quad F^{14}\text{C} = \left(\frac{A_s}{0.95 A_{\text{OX1}}} \right) * \left(\frac{0.975}{0.981} \right)^2 * \left[\left(\frac{1 + \delta^{13}\text{C}_{\text{OX1}}/1000}{1 + \delta^{13}\text{C}_s/1000} \right) \right]^2 \quad (3)$$

333 with $A_s = ^{14}\text{C}/^{12}\text{C}$, S for sample, OX1 for Oxalic Acid 1, the ^{14}C international standard.
 334 $F^{14}\text{C}$ is twice normalized: i) it takes into account isotopic fractionation by being normalized to
 335 a $\delta^{13}\text{C}_s = -25\text{‰}$ and ii) it corresponds to a deviation towards an international standard (i.e.
 336 95% of OX1 as measured in 1950 – (Stuiver and Polach, 1977)). By propagating $F^{14}\text{C}$ from
 337 atmosphere at the origin of vegetal photosynthesis to soil respired CO_2 , there is no need to
 338 focus on ^{13}C isotopic fractionation all along the organic matter mineralization with $F^{14}\text{C}$.

339 To make the reading of the paper easier, we will further express $F^{14}\text{C}$ as $F^{14}\text{C} = A_{\text{sample}}/A_{\text{ref}}$
 340 with A_{sample} being the A of the measured (or modeled) data and A_{ref} an international reference.
 341 Normalizations are included in A_{ref} and $F^{14}\text{C}$ will be written as F^{14} to simplify notation
 342 involving superscripts and subscripts.

343 Since we focus on SOC dynamics, we did not include the ^{14}C in the plants but did include ^{14}C
 344 in the litter. The ^{14}C -litter is obtained by multiplying the atmospheric value by the total carbon
 345 in the litter:

$$346 \quad \text{Litter } (^{14}\text{C}) = F_{\text{atm}}^{14} * \text{Litter } (\text{C}) \quad (4)$$

347 where F_{atm}^{14} is the $F^{14}\text{C}$ of atmosphere at the time of leaf growth (figure 2).

348 Thus, from the litter, all processes defined in section 2.1 that apply to total soil carbon are also
 349 represented for ^{14}C .

350 We also take into account the radioactive decay of ^{14}C . For that, we calculate the amount of
 351 ^{14}C as follow:

$$352 \quad ^{14}\text{C} = ^{14}\text{C} - K_{\text{decrease}} * ^{14}\text{C} \quad (5)$$

353 Where k_{decrease} is the radioactive decay constant ($= \text{Ln}2/5730$) (Godwin, 1962)

354 The $F^{14}\text{C}$ of the soil is then calculated back for carbon, per pool:

$$355 \quad F_{\text{Pool},z}^{14} = \frac{{}^{14}\text{C}_{\text{Pool},z}}{\text{C}_{\text{Pool},z}} \quad (6)$$

356 with pool representing the active, slow or passive pool.

357 Finally, we calculate a mean $F^{14}\text{C}$ value per soil layer, according to the depth:

$$358 \quad F_{\text{Mean},z}^{14} = \frac{F_{\text{active},z}^{14} * {}^{14}\text{C}_{\text{active},z} + F_{\text{slow},z}^{14} * {}^{14}\text{C}_{\text{slow},z} + F_{\text{passive},z}^{14} * {}^{14}\text{C}_{\text{passive},z}}{{}^{14}\text{C}_{\text{active},z} + {}^{14}\text{C}_{\text{slow},z} + {}^{14}\text{C}_{\text{passive},z}} \quad (7)$$

360 2.3 Site descriptions

361 2.3.1 French sites

Auteur inconnu 31/10/y 09:53

Supprimé: that should be

Bertrand Guenet 30/10/y 09:59

Commentaire [1]: Delta symbol to checked in the pdf (comments from ref#1)

Bertrand Guenet 31/10/y 19:31

Supprimé: ... 2 ... [16]

Bertrand Guenet 30/10/y 10:43

Supprimé:)

Auteur inconnu 31/10/y 09:53

Supprimé: -

Bertrand Guenet 31/10/y 22:53

Supprimé: d ^{13}C =- ... [17]

Auteur inconnu 31/10/y 09:53

Supprimé: -

Auteur inconnu 31/10/y 09:53

Supprimé: easier ... the reading of the ... [18]

Bertrand Guenet 30/10/y 16:08

Supprimé: normalizations

Auteur inconnu 31/10/y 09:54

Supprimé: to

Bertrand Guenet 31/10/y 19:31

Supprimé: ... [19]

Auteur inconnu 31/10/y 09:55

Supprimé: the ... notation with ... invol ... [20]

Auteur inconnu 31/10/y 09:55

Supprimé: directly ... n the litter. The ... [21]

Bertrand Guenet 30/10/y 10:43

Supprimé: 3

Bertrand Guenet 31/10/y 19:31

Supprimé: = - $K_{\text{decrease}} * \dots$ [22]

Bertrand Guenet 30/10/y 10:10

Supprimé: (=

Auteur inconnu 31/10/y 09:59

Supprimé: Then ... he, ... [23]

Bertrand Guenet 30/10/y 10:43

Supprimé: 5

Auteur inconnu 31/10/y 10:00

Mis en forme: Police : (Par défaut) Times New Roman, 12 pt, Italique, Anglais (E.U.), Motif : Transparente (Blanc)

Auteur inconnu 31/10/y 10:00

Supprimé: So f

Bertrand Guenet 30/10/y 10:43

Supprimé: 6

Auteur inconnu 31/10/y 10:01

Supprimé: s

Two Luvisol (WRB, 2006) profiles located in the northern France were selected: the Feucherolles and Mons sites. In Mons (49.87°N, 3.03°E), Luvisol, the soils sit under grassland, and are developed from several meters of loess and therefore, well drained. The mean annual air temperature is 11°C and the annual precipitation is about 680 mm (Keyvanshokouhi et al., 2016). In Feucherolles (48.9°N, 1.97°E), the soil sits under oak forest and clay and gritstone deposits are found at approximately 1.5 m depth. The mean annual air temperature is 11.2°C and the annual precipitation is about 660 mm (Keyvanshokouhi et al., 2016). Both soils are neutral to slightly acidic and are characterized by the presence of a clay accumulation Bt horizon with clay content reaching 30 % for Feucherolles and 27 % for Mons, while the upper horizons are poorer in clay (17 % for Feucherolles and 20% for Mons).

The ¹⁴C data from the soils of both sites were obtained after chemical treatment done at Laboratoire des Sciences du Climat et de l'Environnement (LSCE) using a protocol adapted to achieve carbonate leaching without any loss of organic carbon. ¹⁴C activity was measured by AMS at the French Laboratoire de mesure du ¹⁴C (LMC14) facility (Cottéreau et al., 2007). Details on measurements and sampling can be found in Jagercikova et al., (2017)

2.3.2 Congo site

The studied site is located in Kissoko (4.35°S, 11.75°E). It belongs to the SOERE F-ORE-T (Site de l'Observatoire de Recherche en Environnement sur le Fonctionnement des écosystèmes fOReTiers) field observation sites of Pointe Noire, Republic of Congo. The mean annual air temperature is about 25°C with low seasonal variation (± 5°C), and average annual precipitation of 1400mm, and a dry season between June and September. The deep acidic sandy soil is a ferralic Arenosol (WRB, 2006). The soil is characterized by a sand content larger than 90% (Laclau et al., 2000). A soil profile was taken under native savanna vegetation dominated by C4 plants (Epron et al., 2009). The soil was sampled in May 2014 at different depths: 0-5cm, 5-10cm, 10-15cm, 15-20cm, 20-30cm, 30-40cm, 40-50cm, 50-60cm, 60-80cm, 80-100cm, 100-120cm. All samples were crushed and air-dried. Once in the laboratory, they were homogenized, crushed, randomly subsampled and sieved at 200µm. Then ¹⁴C measurements were made the same way as the two French sites, using the LSCE chemical treatment and the French LMC14 facility following recommendations by Cottéreau et al., (2007).

2.3.3 Argentina site

The Province of Misiones is located in northeastern Argentina. The climate is subtropical humid without a dry season, an annual mean temperature of 20°C, and 1850mm of mean annual rainfall (Morrás et al., 2009). The profile used in this study is located in the southern part of Misiones (27°S, 55°W). Native vegetation is a forest dominated by C3 plants. The soil selected is an Acrisol (WRB, 2006). It's a red clay soil, strongly to very strongly acid with a clay content varying from 40% at the surface to 60% at 1m depth. ¹⁴C measurements were made using a new Compact Radiocarbon System called ECHOMICADAS (Environment, Climate, Human, Mini Carbon Dating System) (Tisnérat-Laborde et al., 2015). Details on measurements and sampling can be found in Tifafi et al., In prep. Briefly, the soil was sampled in May 2015 at different depths: 0-5cm, 5-10cm, 10-15cm, 15-20cm, 20-30cm, 30-40cm, 40-50cm, 50-60cm, 60-80cm, 80-100cm. All samples were crushed and air-dried. Once

- Auteur inconnu 31/10/y 10:01
Supprimé: (
- Auteur inconnu 31/10/y 10:02
Supprimé: are
- Auteur inconnu 31/10/y 10:02
Supprimé: thus
- Auteur inconnu 31/10/y 10:02
Supprimé: s
- Auteur inconnu 31/10/y 10:03
Supprimé: , site (48.9°N, 1.97°E),
- Bertrand Guenet 30/10/y 10:10
Supprimé: 5m
- Auteur inconnu 31/10/y 10:05
Supprimé: ,
- Auteur inconnu 31/10/y 10:05
Supprimé: and ¹
- Auteur inconnu 31/10/y 10:05
Supprimé: measurement performed
- Bertrand Guenet 30/10/y 12:59
Supprimé: (
- Bertrand Guenet 30/10/y 13:38
Supprimé: Republic
- Auteur inconnu 31/10/y 10:05
Supprimé: of
- Auteur inconnu 31/10/y 10:05
Supprimé: s
- Auteur inconnu 31/10/y 10:06
Supprimé: and
- Auteur inconnu 31/10/y 10:06
Supprimé: averages
- Auteur inconnu 31/10/y 10:06
Supprimé: with
- Auteur inconnu 31/10/y 10:06
Supprimé: N
- Auteur inconnu 31/10/y 10:07
Supprimé: is a savanna
- Auteur inconnu 31/10/y 10:07
Supprimé: and the selected soil profile is under this native savanna vegetation
- Auteur inconnu 31/10/y 10:07
Supprimé: d
- Bertrand Guenet 30/10/y 13:21
Supprimé: ¹⁴C analyses were made in
- Bertrand Guenet 30/10/y 13:21
Supprimé: with the measurements for
- Bertrand Guenet 30/10/y 13:21
Supprimé: (
- Auteur inconnu 31/10/y 10:48
Supprimé: of mean annual temperature
- Auteur inconnu 31/10/y 10:49
Supprimé: d

482 in the laboratory, they were homogenized, crushed, randomly subsampled and sieved at
 483 200µm. Then ¹⁴C measurements were made using a new Compact Radiocarbon System called
 484 ECHoMICADAS (Environment, Climate, Human, Mini Carbon Dating System) following the
 485 recommendations of Tisnérat-Laborde et al., (2015).

486 For the four sites, the SOC (kg m⁻³), for each depth z , was calculated using carbon content and
 487 bulk density data using the following equation:

$$488 \text{SOC}_z = \text{OCC}_z * \text{BD}_z \quad (8)$$

489 Where OCC (wt/wt) is the carbon content and BD (kg m⁻³) is the bulk density.

490 2.4 Different model tests

491 After the implementation of radiocarbon in the model, different tests were carried out (Table
 492 2). Here we represent the outputs provided by three simulations:

- 493 i- Simulation using the initial version ORCHIDEE-SOM-¹⁴C (labelled, “Control” in
 494 figures and tables) in which no changes were made. The diffusion was kept constant
 495 throughout the profile ($D = 1.10^{-4} \text{ m}^2 \text{ year}^{-1}$) and the other parameters are those of the
 496 detailed version in Camino-Serrano et al., (2017).
- 497 ii- Simulation using the initial version ORCHIDEE-SOM-¹⁴C in which we modified
 498 some parameters following He et al. (2016), (“He et al., (2016) parameterization” in
 499 figures and tables). In brief, the authors used ¹⁴C data from 157 globally distributed
 500 soil profiles sampled to 1-meter depth to evaluate CMIP5 models. Their results show
 501 that ESMs underestimated the mean age of soil carbon by a factor of more than six and
 502 overestimated the carbon sequestration potential of soils by a factor of nearly two. So,
 503 the suggestion (that we apply in this simulation) for the IPSL model was to multiply
 504 the turnover time of the passive pool by 14 and the flux from slow pool to passive pool
 505 by 0.07 (Table 2). The diffusion was kept constant throughout the profile ($D = 1.10^{-4}$
 506 $\text{m}^2 \text{ year}^{-1}$) but the turnover time of the passive pool increased from 462 years to 6468
 507 years and the flux from the slow pool to the passive pool decreased from 0.07 to
 508 0.0049.
- 509 iii- Simulation using the initial version ORCHIDEE-SOM-¹⁴C in which we assume that
 510 the diffusion varies as a function of the depth (“Depth-varying diffusion constant” in
 511 figures and tables) according to the equation below:

$$512 D(z) = 5.42. 10^{-4} e^{(-0.04z)} \quad (9)$$

513 Where D is the diffusion ($\text{m}^2 \text{ year}^{-1}$) at a specific depth and z is the depth. This equation of
 514 diffusion varying as a function of depth following Jagercikova et al., (2014) and assumes that
 515 bioturbation is higher in the top soil than in deep soil.

516 2.5 Model simulations

517 In order to reach a steady state of the soil module, we ran the model over 12700 years
 518 (spinup). The state at the last time step of this spinup was used as the initial state for the
 519 simulations. For this, the CRUNCEP meteorological data for the period 1901-1910 were used.
 520 This has been applied for Misiones, Feucherolles and Mons. However, for Kissoko, a first
 521 spinup similar to the other sites was carried out but a second one (over approximately 4200

Auteur inconnu 31/10/y 10:49
Supprimé: using the following equation

Bertrand Guenet 30/10/y 10:43
Supprimé: 7

Bertrand Guenet 30/10/y 10:50
Supprimé: %

Auteur inconnu 31/10/y 10:50
Supprimé: made

Auteur inconnu 31/10/y 10:50
Supprimé: so-called

Bertrand Guenet 30/10/y 14:18
Supprimé: Model_Control

Auteur inconnu 31/10/y 10:50
Supprimé: has...as been [24]

Bertrand Guenet 30/10/y 10:10
Supprimé: [...[...016]...] [25]

Auteur inconnu 31/10/y 10:51
Supprimé: so-called

Bertrand Guenet 30/10/y 14:20
Supprimé: Model_Test_He

Auteur inconnu 31/10/y 10:51
Supprimé: they...he authors used [26]

Bertrand Guenet 30/10/y 10:52
Supprimé: rate

Auteur inconnu 31/10/y 10:52
Supprimé: by 0.07 ...he flux from [27]

Bertrand Guenet 30/10/y 10:10
Supprimé: 049

Auteur inconnu 31/10/y 10:52
Supprimé: , initially constant throughout the profile, ...aries as a function of [28]

Bertrand Guenet 30/10/y 14:23
Supprimé: Model_Test_Diffusion

Bertrand Guenet 30/10/y 10:43
Supprimé: 8

Bertrand Guenet 30/10/y 10:00
Supprimé:

Auteur inconnu 31/10/y 10:53
Supprimé: First of all, i... order to re [29]

years) was also done after the end of the first to take into account the change of the land cover from a tropical forest to a C4 savanna at this site (Schwartz et al., 1992). The atmospheric CO₂ concentration has been set at 296 ppm (year 1901, (Keeling and Whorf, depth-varying diffusion constant)) for the spinups and the F¹⁴C has been set to pre-industrial values. For each site, specific pH, clay content and bulk density values were used (Table 1). It should be noted that for these last data, only one value (the mean value on the profile) is provided as input for the model.

The simulations were outputted at a yearly time step, from 1900 to 2011. A yearly atmospheric CO₂ concentration value (Keeling and Whorf, depth-varying diffusion constant) is read for the sites. The same specific pH, clay content and bulk density values were used (Table 1).

Figure 2 shows the evolution of the F¹⁴C values in the atmosphere used in our model for Argentina, Congo and France (Figure 5 from Hua et al. (2013)). The values provided are classified into five zones, three in the Northern Hemisphere (NH) and two in the Southern Hemisphere (SH), corresponding to different levels of ¹⁴C. For France, the values correspond to the NH zone 2, for the Congo to the SH zone 3 and finally for Argentina to the SH zone 1-2. Thus, for our simulations, a yearly value is read for each site.

An F¹⁴C value of 1.8 represents a doubling of the amount of ¹⁴C in atmospheric CO₂. In figure 2, it can be noted that the values recorded in France (northern hemisphere) are higher than those in the Congo and Argentina (southern hemisphere). This is due to the preponderance of atmospheric tests in the northern hemisphere and the time required to mix air across the equator.

2.6 Statistical analysis

Simulating carbon processes in soil requires comparison between the model outputs and the measurements to test the model accuracy and possibly implement further improvement. Statistical analysis based on the statistics of deviation were done to evaluate the model-measurement discrepancy according to Kobayashi and Salam (2000) (where a detailed description of the method is provided). Here, we only reproduce the different equations used. x refers to the model outputs and y to the measurements, while i refers to soil depth. The intervals of soil depth of the model outputs and the measurements were homogenized by linearly interpolating the data to common depth intervals defined for each site. The simulations and data were then compared for each depth interval.

$$RMSD = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - y_i)^2} \quad (10)$$

RMSD is the Root Mean Squared Deviation, which represents the mean distance between simulation and measurement.

$$MSD = \frac{1}{n} \sum_{i=1}^n (x_i - y_i)^2 = (\bar{x} - \bar{y})^2 + \frac{1}{n} \sum_{i=1}^n [(x_i - \bar{x}) - (y_i - \bar{y})]^2 \quad (11)$$

MSD, the Mean Squared Deviation, is the square of RMSD. The lower the value of MSD, the closer the simulation results are to the measurements.

$$SB = (\bar{x} - \bar{y})^2 \quad (12)$$

Bertrand Guenet 30/10/y 16:51
Supprimé: 2006

Bertrand Guenet 30/10/y 16:15
Supprimé: then run

Bertrand Guenet 30/10/y 16:51
Supprimé: 2006

Auteur inconnu 31/10/y 10:55
Supprimé: Of course, t

Auteur inconnu 31/10/y 10:55
Supprimé: In fact, t

Auteur inconnu 31/10/y 10:55
Supprimé: 3

Auteur inconnu 31/10/y 10:55
Supprimé: 2

Auteur inconnu 31/10/y 10:56
Supprimé: O

Auteur inconnu 31/10/y 10:56
Supprimé: represent

Bertrand Guenet 30/10/y 15:52
Mis en forme: Police :Italique

Bertrand Guenet 30/10/y 15:52
Mis en forme: Police :Italique

Auteur inconnu 31/10/y 10:57
Supprimé: linearly

Bertrand Guenet 30/10/y 10:44
Supprimé: 9

Bertrand Guenet 30/10/y 10:44
Supprimé: 10

Auteur inconnu 31/10/y 10:57
Supprimé: is

Bertrand Guenet 30/10/y 10:44
Supprimé: 11

624 Where are the means of x_i (model outputs) and y_i (measurements) respectively.
 625 SB is a part of the MSD (Eq.14) and represents the bias of the simulation from the
 626 measurement.

$$627 \quad SD_s = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2} \quad (13)$$

628 SD_s is the Standard Deviation of the simulation.

$$629 \quad SD_m = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2} \quad (14)$$

630 SD_m is the Standard Deviation of the measurements.

$$631 \quad r = \frac{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{SD_m SD_s} \quad (15)$$

632 r is the correlation coefficient between the simulation and measurements.

$$633 \quad SDSD = (SD_s - SD_m)^2 \quad (16)$$

634 SDSD is the difference in the magnitude of fluctuation between the simulation and
 635 measurements.

$$636 \quad LCS = 2SD_s SD_m (1 - r) \quad (17)$$

637 LSC represents the lack of positive correlation weighted by the standard deviations.

638 The MSD can be therefore be rewritten as:

$$639 \quad MSD = SB + SDSD + LCS \quad (18)$$

640 For the different simulations, the MSD and its components were calculated according to the
 641 total soil carbon and to the $F^{14}C$.

642

643 3 Model results and evaluation

644 3.1 Outputs from simulation using the initial version of the model ORCHIDEE-SOM- 645 ^{14}C (Control)

646 3.1.1 Simulated total soil carbon

647 Results from the initial version of ORCHIDEE-SOM- ^{14}C show that in all the studied sites, the
 648 model succeeds in reproducing the trend of the total carbon profiles, with more carbon at the
 649 surface which then decreases according to the depth (Figure 3). Moreover, total soil carbon
 650 stock simulated down to 2m depth is in accordance with data in the case of Misiones and
 651 Feucherolles where the major difference mainly lies on the surface. This results in correlation
 652 coefficients of 0.44 and 0.2 respectively (Table 3). For the sites of Kissoko and Mons, an
 653 over-estimation of the total soil carbon is found to a depth of 50cm for Kissoko and up to a
 654 depth of 120cm for Mons. Correlation coefficients are 0.14 and 0.49 for Kissoko and Mons
 655 respectively (Table 3).

656 Metrics presented in Figure 4, showed that this version (ORCHIDEE-SOM- ^{14}C) represents
 657 relatively well the observation from Feucherolles ($MSD = 206 \text{ kg C m}^{-6}$), whereas the other

Bertrand Guenet 30/10/y 10:45

Supprimé: 3

Bertrand Guenet 30/10/y 10:44

Supprimé: 12

Bertrand Guenet 30/10/y 10:44

Supprimé: 13

Bertrand Guenet 30/10/y 10:44

Supprimé: 14

Auteur inconnu 31/10/y 10:58

Supprimé: Where

Auteur inconnu 31/10/y 10:58

Mis en forme: Police : (Par défaut) Times New Roman, 12 pt, Italique, Anglais (E.U.)

Bertrand Guenet 30/10/y 10:44

Supprimé: 15

Auteur inconnu 31/10/y 10:58

Supprimé: here,

Bertrand Guenet 30/10/y 10:44

Supprimé: 16

Auteur inconnu 31/10/y 10:58

Supprimé: Finally, with all the above terms combined, the

Bertrand Guenet 30/10/y 10:44

Supprimé: 17

Bertrand Guenet 30/10/y 14:18

Supprimé: Model_Control

Auteur inconnu 31/10/y 11:00

Supprimé: then

Bertrand Guenet 30/10/y 14:05

Supprimé: 55

Bertrand Guenet 30/10/y 14:05

Supprimé: 6

Auteur inconnu 31/10/y 11:00

Supprimé: marked

Auteur inconnu 31/10/y 11:00

Supprimé: deep

Auteur inconnu 31/10/y 11:01

Supprimé: (then it decreases)

Auteur inconnu 31/10/y 11:01

Supprimé: deep

Bertrand Guenet 30/10/y 14:06

Supprimé: 75

Bertrand Guenet 30/10/y 10:53

Supprimé: 3

are highly overestimated (Kissoko, MSD = 1343 kg C m⁻⁶; Misiones MSD = 2180 kg C m⁻⁶; Mons MSD = 3355 kg C m⁻⁶). By detailing the different components of the MSD (Figure 4), we note that for Mons and Kissoko, standard bias (SB) is the major component of the MSD with contributing 70% and 60% respectively. This reflects that the average of total soil carbon over the soil profile simulated by the model is primarily the origin of the deviation of the model outputs from data. The mean total soil carbon estimated by the model (Table 3) is almost three times higher than the mean total carbon measured for Mons (2.37 kg C m⁻² against 0.8 kg C m⁻² respectively) and it is more than five times that measured for Kissoko (2.44 kg C m⁻² against 0.42 kg C m⁻² respectively). For Mons a net primary production (NPP) of 6.7 t ha⁻¹ yr⁻¹ was estimated by the technical institute for pasture in this region of France based on the annual yields, whereas the model predicts a NPP of 7.5 t ha⁻¹ yr⁻¹. The large overestimation of the SOC stocks may therefore be due to an overestimation of the NPP. This significant gap recorded in the case of the Kissoko site, where the measured SOC is very low, is probably due to an overestimation of decay rates by ORCHIDEE in sandy soils. The correlation coefficient for Mons is relatively high compared to other site (Table 3) whereas Fig. 3 shows that the model performance was not very good for this site. This is mainly due to a large SB whereas other MSD components were rather low.

However, the main components of MSD for Feucherolles and Misiones are both SB (46% and 56% for Feucherolles and Misiones, respectively) and also LCS (53 and 31% for Feucherolles and Misiones, respectively). This means that for these two sites, the deviation between model outputs and measurements is mainly due to a variation of carbon stock estimation throughout the profile. The mean total soil carbon estimated in these both cases (Table 3) is only slightly higher than those measured (2.03 kg C m⁻² estimated against 2.14 kg C m⁻² measured for Misiones and 0.7 kg C m⁻² estimated against 0.68 kg C m⁻² measured for Feucherolles).

The vertical profiles of the SOC stock were fairly represented by the model. The overestimation, especially at the top, suggests that the distribution of the litter following the root profile and / or the vertical transport of SOC by diffusion are not correctly described in the model.

3.1.2 Simulated F¹⁴C

Regarding the ¹⁴C activity, bulk F¹⁴C profiles show a classical pattern with higher ¹⁴C activity on the top, slightly influenced by the peak bomb enriched years. Subsequently profiles show decreasing ¹⁴C activity with depth (Figure 5).

The estimated profiles (Model-Control) follow the same trend with a decrease from the surface to the depth. However, there is a significant difference between the estimated values and those measured throughout the profile. The statistical analyzes (Figure 6) provide MSD values: 0.02 for Mons and Misiones, 0.03 for Kissoko and 0.09 for Feucherolles. The major component of the MSD in the four sites is the LCS, with a proportion reaching 90% for Mons, 80% for Misiones and 70% for Congo, but only 55% for Feucherolles. The high proportions of LCS suggest that the model fails to reproduce the shape of the profile. The lower values estimated by the models reflect a more modern carbon age than in reality. This can be

Bertrand Guenet 30/10/y 14:03

Supprimé: ³...; Misiones MSD=...= 2 ... [30]

Auteur inconnu 31/10/y 11:01

Supprimé: Then, b

Bertrand Guenet 30/10/y 14:32

Supprimé: more ...lmost than four... [31]

Bertrand Guenet 30/10/y 11:03

Supprimé: ORCHIDEE is a global model that is not parameterized for such specific soil conditions

Bertrand Guenet 30/10/y 14:12

Supprimé: only 1.7 to 2...nly slightly... [32]

Bertrand Guenet 30/10/y 14:43

Supprimé: simulated was thereby globally not very far from that of the data

Auteur inconnu 31/10/y 11:03

Supprimé: ...on the top, slightly infl... [33]

Auteur inconnu 31/10/y 11:04

Supprimé: always greater than 50% and which...eaching is even...90% for M... [34]

771 explained, first, by the fact that the root profile puts too much fresh organic carbon in deep
 772 soil. Afterwards, in ORCHIDEE, root profile is assumed to follow an exponential [function](#)
 773 without modulation due to environmental conditions.

774 [SB's contribution to the MSD does](#), not exceed 7% for Misiones, Kissoko and Mons but
 775 reaches about 40% for Feucherolles. This reflects that the mean value of the $F^{14}C$ estimated
 776 by the model and that obtained after the measurements are not very different, except for
 777 Feucherolles site (Table 4). Indeed, the average value estimated for Misiones is 0.920, very
 778 close to that measured at 0.930, 0.995 for Kissoko against 0.985 measured and 0.860 for
 779 Mons against 0.815 measured. Yet, the difference is greater for the Feucherolles site, the
 780 estimated value being 0.915 while the measurement is 0.725. This difference might be caused
 781 by the low $F^{14}C$ value measured at 150cm (0.257), that the model is not able to capture. This
 782 suggests that modeled deep soil carbon is much younger than the observed total soil carbon,
 783 probably because ORCHIDEE-SOM simulates a relatively small proportion of passive pool in
 784 the lower soil horizons (Figure 7), while an increasing proportion of passive carbon with soil
 785 depth could be expected.

786 In brief, SOC stocks are generally overestimated and soil carbon age in deep soils (as shown
 787 by the $F^{14}C$) is underestimated, suggesting that the turnover rate of [the](#) passive pool is subject
 788 to improvements in ORCHIDEE-SOM.

789 3.2 Outputs from simulation using the initial version of the model ORCHIDEE-SOM- 790 ^{14}C including He's suggestion ([He et al., \(2016\) parameterization](#))

791 3.2.1 Simulated total soil carbon

792 Figure 3 shows profiles output after [He et al., \(2016\)](#)'s suggestion [was](#) implemented into
 793 ORCHIDEE-SOM- ^{14}C (green dotted curves). Resulting profiles follow the same trend than
 794 observations but in this case ([He et al., \(2016\) parameterization](#)), the overestimation is very
 795 high [across the whole profile](#). This is further confirmed by the metrics analysis (Figure 4).
 796 MSD values markedly increased, resulting in an even higher variance. Obviously, the major
 797 component of MSD in all cases is the SB (varying from 80% to 87%) reflecting an even more
 798 marked overestimation of the mean total carbon estimates: [7.38 kg C m⁻²](#) against [2.14 kg C m⁻²](#)
 799 [for Misiones](#), [2.44 kg C m⁻²](#) against [0.42 kg C m⁻²](#) for Kissoko, [2.33 kg C m⁻²](#) against [0.66 kg](#)
 800 [C m⁻²](#) for Feucherolles and [9.99 kg C m⁻²](#) against [0.8 kg C m⁻²](#) for Mons.

801 3.2.2 Simulated $F^{14}C$

802 [He et al., \(2016\) parameterization](#) outputs (Figure 5, green dotted curves) for $F^{14}C$ are once
 803 again even further away from observations and MSDs (Figure 6) are much higher, except for
 804 Feucherolles. The MSD components for [the](#) Feucherolles site show that the LCS increases
 805 from 0.05 to 0.06 whereas [the SB decreases](#) from 0.04 to 0.03, again reflecting a variation of
 806 the profile more than a difference from the means.

807 Improvement of the model-measurement fit for the $F^{14}C$ at 150 cm in Feucherolles confirms
 808 that the deep soil carbon simulated by the control version of ORCHIDEE-SOM- ^{14}C was
 809 excessively young, since the longer residence time of the passive pool reported by He et al.
 810 (2016) resulted in a higher proportion of passive pool across the soil profile (Figure 7), thus
 811 improving deep soil carbon age. Nevertheless, this test only improves the simulation of deep

Auteur inconnu 31/10/y 11:05

Supprimé: Then,

Auteur inconnu 31/10/y 11:05

Supprimé: es

Bertrand Guenet 30/10/y 14:20

Supprimé: Model_Test He

Bertrand Guenet 30/10/y 14:20

Supprimé: Model_Test He

Auteur inconnu 31/10/y 11:09

Supprimé: from the surface to the depth

Bertrand Guenet 30/10/y 14:13

Supprimé: 128

Bertrand Guenet 30/10/y 14:13

Supprimé: 3

Bertrand Guenet 30/10/y 14:13

Supprimé: 31

Bertrand Guenet 30/10/y 14:13

Supprimé: 3

Bertrand Guenet 30/10/y 14:14

Supprimé: 53

Bertrand Guenet 30/10/y 14:14

Supprimé: 3

Bertrand Guenet 30/10/y 14:14

Supprimé: 3

Bertrand Guenet 30/10/y 14:14

Supprimé: 3

Bertrand Guenet 30/10/y 14:14

Supprimé: 4

Bertrand Guenet 30/10/y 14:14

Supprimé: 3

Bertrand Guenet 30/10/y 14:14

Supprimé: 14

Bertrand Guenet 30/10/y 14:14

Supprimé: 3

Bertrand Guenet 30/10/y 14:14

Supprimé: 131

Bertrand Guenet 30/10/y 14:14

Supprimé: 3

Bertrand Guenet 30/10/y 14:14

Supprimé: 15

Bertrand Guenet 30/10/y 14:14

Supprimé: 3

Bertrand Guenet 30/10/y 14:20

Supprimé: Model_Test He

Auteur inconnu 31/10/y 11:10

Supprimé: , which MSD value in this case is lower

Auteur inconnu 31/10/y 11:10

Supprimé: it is

Auteur inconnu 31/10/y 11:10

Supprimé: which

Auteur inconnu 31/10/y 11:10

Supprimé: d

838 soil carbon in Feucherolles. On the contrary, this increase in carbon residence time increases
839 model deviation from observations, for all the other cases (Figure 5 and 6).

840 Indeed, taking the priming effect into account in this new version of ORCHIDEE has
841 contributed to a 50% of decrease in carbon storage over the historical period. He, et al.,
842 (2016)'s correction was also aimed at reducing this storage and is of the same order of
843 magnitude as the priming effect. Thus, applying He's correction to this version of the model,
844 which takes into account the priming effect, contributes to a double correction for the same
845 target, which then generates this important difference between model outputs and
846 measurements. Moreover, the work of He et al. (2016) is done under the standard
847 parameterization of ORCHIDEE based on Century, while ORCHIDEE-SOM was re-
848 parameterized after adding several different processes, the priming effect among them
849 (Camino-Serrano et al., 2017), which makes it difficult to compare results from between the
850 two studies.

851 3.3 Outputs from simulation using the initial version of the model ORCHIDEE-SOM- 852 ¹⁴C with diffusion varying according to the depth (Depth-varying diffusion constant)

853 3.3.1 Simulated total soil carbon

854 Fick's law of diffusion is classically used in models to represent bioturbation assuming that
855 soil fauna activity may be represented following the Fick's law of diffusion (Elzein and
856 Balesdent, 1995; Guenet et al., 2013; Koven et al., 2013; O'Brien and Stout, 1978; Wynn et
857 al., 2005). Using a fixed diffusion constant (D in eq. 2) implicitly suggests that soil fauna
858 activity is uniform over the entire soil profile. This is generally the case of several models of
859 diffusion, in particular, at the level of an ecosystem (Bruun et al., 2007; Guimberteau et al.,
860 2017; O'Brien and Stout, 1978). However soil faunal activity vary naturally with depth and
861 the diffusion constant, should therefore be depth-dependent (Jagercikova et al., 2014).

862 With Depth-varying diffusion constant, the carbon profiles (orange dashed curves) was
863 improved compared to the initial outputs (Control). The overestimation at the surface
864 decreases at the four sites (Figure 3). In particular, the Misiones outputs fit very well the
865 observed profiles. This is confirmed with lower MSDs for the four sites for this version
866 compared to Control, (Figure 4).

867 The total SOC stocks simulated according to this third simulation are closer to the measured
868 values and describing the vertical transport of SOC through diffusion varying according to the
869 depth improves significantly the model outputs.

870 3.3.2 Simulated F¹⁴C

871 Regarding the F¹⁴C outputs, the simulations using the initial version ORCHIDEE-SOM-¹⁴C in
872 which we assume that the diffusion varies as a function of the depth (Depth-varying diffusion
873 constant) results in an improvement of the F¹⁴C profiles (orange dashes curves), in particular,
874 for the sites Misiones, Mons and Kissoko (Figure 5). Statistical analyzes prove it with
875 significantly lower MSDs. In addition, the proportion of LCS is 98%, 92% and 88% for
876 Mons, Misiones and Kissoko, respectively, highlighting an estimated average very close to
877 the measurements with a clear disparity, less marked than with the first two simulations,
878 throughout the profile (Figure 6). Overall, the simulated F¹⁴C to 2 m of depth according to

Auteur inconnu 31/10/y 11:11
Supprimé: has even more deviated the outputs of the model

Bertrand Guenet 31/10/y 19:39
Supprimé: 's

Auteur inconnu 31/10/y 11:12
Supprimé: then

Auteur inconnu 31/10/y 11:13
Supprimé: what

Auteur inconnu 31/10/y 11:13
Supprimé: associate

Auteur inconnu 31/10/y 11:13
Supprimé: her and this study

Bertrand Guenet 30/10/y 14:23
Supprimé: Model_Test Diffusion

Bertrand Guenet 30/10/y 14:54
Supprimé: fact

Auteur inconnu 31/10/y 11:14
Supprimé: especially used

Bertrand Guenet 30/10/y 14:55
Supprimé: , in addition, the characteristics of a soil, i.e. its structure and pore distribution, may vary depending on the depth, so,

Bertrand Guenet 30/10/y 14:55
Supprimé: coefficient

Bertrand Guenet 30/10/y 14:23
Supprimé: Model_Test Diffusion

Bertrand Guenet 30/10/y 14:18
Supprimé: Model_Control

Bertrand Guenet 30/10/y 14:18
Supprimé: Model_Control

Auteur inconnu 31/10/y 11:15
Supprimé: showing a much smaller deviation from the measurements

Auteur inconnu 31/10/y 11:15
Supprimé: Anyway, t

Bertrand Guenet 30/10/y 14:23
Supprimé: Model_Test Diffusion

Auteur inconnu 31/10/y 11:15
Supprimé: especially

901 | this third simulation are in a better agreement with the measured values, and thus
902 | incorporating diffusion that varies with depth, significantly improves the model outputs.

903 | Using a diffusion coefficient that varies as a function of the depth, seems to correct the
904 | overestimation of the surface total soil carbon by increasing the proportion of labile soil
905 | carbon pools in the first soil layers.

906 | When we sum the total soil carbon at each soil layer and look at the relative proportion of
907 | each of the soil carbon pools, (Figure 7), we note that it is mainly the distribution of the litter
908 | according to the depth which varies. In fact, the structural litter proportion is multiplied by
909 | about 2 in all four cases, and this proportion remains relatively constant across the profile.
910 | This increase in litter proportion has also resulted in a decrease in the passive pool, more
911 | pronounced at the surface but also important at depth (except for Feucherolles where the
912 | decrease is only marked at the bottom). It suggests that the vertical carbon distribution, which
913 | is largely modified by the diffusion coefficient, greatly impacts the SOC and ^{14}C profiles,
914 | which is in line with Dwivedi et al. (2017) who found that the vertical carbon input profiles
915 | were important controls over the ^{14}C depth distribution.

916 | In this study, the vertical transport of SOC and litter through diffusion has been improved by
917 | varying diffusion according to the depth. Further model development should explore the
918 | impact of the other processes defining the soil carbon pools vertical distribution and
919 | especially the distribution of the litter according to the root profile.

920 | Overall, by using radiocarbon (^{14}C) measurements we have been able to diagnose internal
921 | model biases (underestimation of deep soil carbon age) and to propose further model
922 | improvements (depth-dependent diffusion). Therefore, the use of radiocarbon (^{14}C) tracers in
923 | global models emerges as a promising tool to constrain, not only SOC turnover times in the
924 | long-term (He et al., 2016), but also internal SOC processes and fluxes that have no direct
925 | comparison with field measurements. Nevertheless, the model evaluation performed here on
926 | only four sites should be considered as proof of concept and more in depth evaluation are
927 | needed, in particular using a large ^{14}C database available at global scale (Balesdent et al.,
928 | 2018; Mathieu et al., 2015). Indeed, the F^{14}C is largely controlled by pedo-climatic conditions
929 | such as clay content, climate and mineralogy (Mathieu et al., 2015) and the range of situations
930 | we covered here is relatively limited.

931

932 | 4 conclusion

933 | ORCHIDEE-SOM- ^{14}C , is one of the first land surface models that incorporates the ^{14}C
934 | dynamics in the soil (Koven et al., 2013). Its starting point is ORCHIDEE-SOM, a recently
935 | developed soil model. We evaluated the new model ORCHIDEE-SOM- ^{14}C for four sites in
936 | different biomes. The model almost managed to reproduce the soil organic carbon stocks and
937 | the ^{14}C content along the vertical profiles at all four sites. However, an overestimation of the
938 | total carbon stock throughout the profile was noted, with the greatest deviation at the surface.
939 | By using radiocarbon (^{14}C) measurements, we have been able to diagnose internal model
940 | biases (underestimation of deep soil carbon age) and to propose further model improvements
941 | (depth-dependent diffusion). These results demonstrate the importance of depth-dependent

Auteur inconnu 31/10/y 11:16

Supprimé: ,

Auteur inconnu 31/10/y 11:16

Supprimé: varying according to the depth

Auteur inconnu 31/10/y 11:16

Supprimé: significantly

Auteur inconnu 31/10/y 11:17

Supprimé: ,

Auteur inconnu 31/10/y 11:17

Supprimé: summing the total soil carbon at each soil layer

Auteur inconnu 31/10/y 11:17

Supprimé: d

Auteur inconnu 31/10/y 11:18

Supprimé: as large at the surface as at depth

Bertrand Guenet 30/10/y 10:10

Supprimé: process

Auteur inconnu 31/10/y 11:19

Supprimé: t

Auteur inconnu 31/10/y 11:19

Supprimé: are

Auteur inconnu 31/10/y 11:19

Supprimé: has

Auteur inconnu 31/10/y 11:20

Supprimé: the

Auteur inconnu 31/10/y 11:20

Supprimé: but

Auteur inconnu 31/10/y 11:20

Supprimé: was mostly marked on

Auteur inconnu 31/10/y 11:20

Supprimé: Then, b

diffusion to improving model outputs with regards to observations. This suggests that, from now on, model improvements should mainly focus on a depth dependent parameterization. We limited our work here to depth-varying diffusion, but other parameters are also depth dependent and should be represented as such in the next version of the model. For instance, belowground litter production in the model is simply represented by an exponential law without any representation of the effect of resource distribution on root profile (e.g. water or nutrients). This is a complex task in a land surface model running at large scale with a classical resolution of 0.5°, but the soil modules of land surface models are quite sensitive to the NPP (Camino-Serrano et al., 2018; Todd-Brown et al., 2013) and a better constraint on the profile of the below ground litter production would likely improve the model performance. Furthermore, here we used only one averaged value over the soil profile for soil boundary conditions (texture, pH, bulk density) but those variables are known to impact the F¹⁴C (Mathieu et al., 2015) and change with depth (Barré et al., 2009) and depth-varying boundary conditions may also help to improve the model. Finally, the next step will deal with the comparison of model outputs to data at larger scales to be able to run the new version ORCHIDEE-SOM-¹⁴C at both regional and global scales.

974

975

976

977

978 Code availability

979 The version of the code is freely available here:

980 http://forge.ipsl.jussieu.fr/orchidee/wiki/GroupActivities/CodeAvalaibilityPublication/ORCHIDEE_gmd-2018-14C

982

983 Acknowledgement

984 This study, part of the MT's PhD, financed by the University of Versailles Saint Quentin, is within the scope of the ANR-14-CE01-0004 DeDyCAS project. Marta Camino-Serrano acknowledges funding from the European Research Council Synergy grant ERC- 2013-SyG-610028 IMBALANCE-P. Part of the data were acquired in the frame of the AGRIPED project (ANR 2010 BLAN 605). We thank Matthew McGrath for his valuable comments on the manuscript.

990

991 References

- 992 Ahrens, B., Braakhekke, M. C., Guggenberger, G., Schrumpf, M. and Reichstein, M.: Contribution of sorption, DOC transport and microbial interactions to the 14C age of a soil organic carbon profile: Insights from a calibrated process model, Soil Biol. Biochem., 88, 390–402, doi:10.1016/j.soilbio.2015.06.008, 2015.
- 996 Balesdent, J. and Guillet, B.: Les datations par le 14C des matières organiques des sols.

Auteur inconnu 31/10/y 11:21

Supprimé: The importance of diffusion has also been highlighted as by making it

Bertrand Guenet 30/10/y 16:17

Supprimé: varies according to the depth

Auteur inconnu 31/10/y 11:21

Supprimé: depth-dependent, the model outputs have been improved

Bertrand Guenet 30/10/y 11:24

Supprimé: , mainly for the diffusion

Auteur inconnu 31/10/y 11:23

Supprimé: Here we presented the effect of a

Auteur inconnu 31/10/y 11:23

Supprimé: constant

Auteur inconnu 31/10/y 11:23

Supprimé: aiming at

Auteur inconnu 31/10/y 11:24

Supprimé: probably

Auteur inconnu 31/10/y 11:24

Supprimé: ies

Bertrand Guenet 30/10/y 15:18

Supprimé: The

Bertrand Guenet 30/10/y 16:18

Supprimé: s

Auteur inconnu 31/10/y 11:24

Supprimé: g

1011 Contribution à l'étude de l'humification et du renouvellement des substances
1012 humiqueétriques, *Sci. du sol*, 2, 93–111, 1982.

1013 Balesdent, J., Basile-Doelsch, I., Chadoeuf, J., Cornu, S., Derrien, D., Fekiacova, Z. and
1014 Hatté, C.: Atmosphere–soil carbon transfer as a function of soil depth, *Nature*, 23, 1,
1015 doi:10.1038/s41586-018-0328-3, 2018.

1016 Barré, P., Berger, G. and Velde, B.: How element translocation by plants may stabilize illitic
1017 clays in the surface of temperate soils, *Geoderma*, 151(1–2), 22–30,
1018 doi:10.1016/j.geoderma.2009.03.004, 2009.

1019 Braakhekke, M., Beer, C., Schrumpf, M., Ekici, A., Ahrens, B., Hoosbeek, M. R., Kruijt, B.,
1020 Kabat, P. and Reichstein, M.: The use of radiocarbon to constrain current and future soil
1021 organic matter turnover and transport in a temperate forest, *J. Geophys. Res. Biogeosciences*,
1022 372–391, doi:10.1002/2013JG002420.Received, 2014.

1023 Braakhekke, M. C., Beer, C., Hoosbeek, M. R., Reichstein, M., Kruijt, B., Schrumpf, M. and
1024 Kabat, P.: SOMPROF: A vertically explicit soil organic matter model, *Ecol. Modell.*,
1025 222(10), 1712–1730, doi:10.1016/j.ecolmodel.2011.02.015, 2011.

1026 Bruun, S., Christensen, B. T., Thomsen, I. K., Jensen, E. S. and Jensen, L. S.: Modeling
1027 vertical movement of organic matter in a soil incubated for 41 years with ¹⁴C labeled straw,
1028 *Soil Biol. Biochem.*, 39(1), 368–371, doi:10.1016/j.soilbio.2006.07.003, 2007.

1029 Camino-Serrano, M., Guenet, B., Luyssaert, S., Ciais, P., Bastrikov, V., De Vos, B., Gielen,
1030 B., Gleixner, G., Jornet-Puig, A., Kaiser, K., Kothawala, D., Lauerwald, R., Peñuelas, J.,
1031 Schrumpf, M., Vicca, S., Vuichard, N., Walmsley, D. and Janssens, I. A.: ORCHIDEE-SOM:
1032 Modeling soil organic carbon (SOC) and dissolved organic carbon (DOC) dynamics along
1033 vertical soil profiles in Europe, *Geosci. Model Dev.*, 11, 937–957, doi:10.5194/gmd-11-937-
1034 2018, 2018.

1035 Coleman, K., Jenkinson, D. S., Crocker, G. J., Grace, P. R., Klír, J., Körschens, M., Poulton,
1036 P. R. and Richter, D. D.: Simulating trends in soil organic carbon in long-term experiments
1037 using RothC-26.3, *Geoderma*, 81(1–2), 29–44, doi:10.1016/S0016-7061(97)00079-7, 1997.

1038 Cottreau, E., Arnold, M., Moreau, C., Baqué, D., Bavay, D., Caffy, I., Comby, C.,
1039 Dumoulin, J.-P., Hain, S., Perron, M., Salomon, J. and Setti, V.: Artemis, the New ¹⁴C AMS
1040 at LMC14 in Saclay, France, *Radiocarbon*, 49(2), 291–299, doi:10.2458/azu_js_rc.49.2928,
1041 2007.

1042 Delibrias, G., Guillier, M. T. and Labeyrie, J.: Saclay natural radiocarbon measurements i,
1043 *Radiocarbon*, 6, 233–250, 1964.

1044 Ducoudré, N. I., Laval, K. and Perrier, A.: SECHIBA, a New Set of Parameterizations of the
1045 Hydrologic Exchanges at the Land-Atmosphere Interface within the LMD Atmospheric
1046 General Circulation Model, *J. Clim.*, 6(2), 248–273, doi:10.1175/1520-
1047 0442(1993)006<0248:SANSOP>2.0.CO;2, 1993.

1048 Dwivedi, D., Riley, W., Torn, M., Spycher, N., Maggi, F. and Tang, J.: Mineral properties,
1049 microbes, transport, and plant-input profiles control vertical distribution and age of soil
1050 carbon stocks, *Soil Biol. Biochem.*, 107(2017), 244–259, doi:10.1016/j.soilbio.2016.12.019,
1051 2017.

1052 Elliott, E. T., Paustian, K. and Frey, S. D.: Modeling the Measurable or Measuring the
1053 Modelable: A Hierarchical Approach to Isolating Meaningful Soil Organic Matter
1054 Fractionations, in *Evaluation of Soil Organic Matter Models: Using Existing Long-Term*
1055 *Datasets*, vol. 1994, pp. 161–179., 1996.

1056 Elzein, A. and Balesdent, J.: Mechanistic Simulation of vertical distribution of carbon
1057 concentrations and residence times in soils, *Soil Sci. Soc. Am. J.*, 59, 1328–1335,
1058 doi:10.1017/CBO9781107415324.004, 1995.

1059 Epron, D., Marsden, C., M'Bou, A. T., Saint-André, L., d'Annunzio, R. and Nouvellon, Y.:
1060 Soil carbon dynamics following afforestation of a tropical savannah with Eucalyptus in
1061 Congo, *Plant Soil*, 323(1), 309–322, doi:10.1007/s11104-009-9939-7, 2009.

1062 Godwin, H.: Half-life of radiocarbon, *Nature*, 195(4845), 984, doi:10.1038/195984a0, 1962.

1063 Guenet, B., Eglin, T., Vasilyeva, N., Peylin, P., Ciais, P. and Chenu, C.: The relative
1064 importance of decomposition and transport mechanisms in accounting for soil organic carbon
1065 profiles, *Biogeosciences*, 10(4), 2379–2392, doi:10.5194/bg-10-2379-2013, 2013.

1066 Guenet, B., Moyano, F. E., Peylin, P., Ciais, P. and Janssens, I. A.: Towards a representation
1067 of priming on soil carbon decomposition in the global land biosphere model ORCHIDEE
1068 (version 1.9.5.2), *Geosci. Model Dev.*, 9(2), 841–855, doi:10.5194/gmd-9-841-2016, 2016.

1069 Guimberteau, M., Zhu, D., Maignan, F., Huang, Y., Yue, C., Dantec-Nédélec, S., Ottlé, C.,
1070 Jornet-Puig, A., Bastos, A., Laurent, P., Goll, D., Bowring, S., Chang, J., Guenet, B., Tifafi,
1071 M., Peng, S., Krinner, G., Ducharne, A., Wang, F., Wang, T., Wang, X., Wang, Y., Yin, Z.,
1072 Lauerwald, R., Joetzjer, E., Qiu, C., Kim, H. and Ciais, P.: ORCHIDEE-MICT (revision
1073 4126), a land surface model for the high-latitudes: model description and validation, *Geosci.*
1074 *Model Dev. Discuss.*, 1–65, doi:10.5194/gmd-2017-122, 2017.

1075 He, Y., Trumbore, S. E., Torn, M. S., Harden, J. W., Vaughn, L. J. S., Allison, S. D. and
1076 Randerson, J. T.: Radiocarbon constraints imply reduced carbon uptake by soils during the
1077 21st century, *Science* (80-.), 353(6306), 1419–1424, 2016.

1078 Hua, Q., Barbetti, M. and Rakowski, A. Z.: Atmospheric Radiocarbon for the Period 1950–
1079 2010, *Radiocarbon*, 55(4), 2059–2072, doi:10.2458/azu_js_rc.v55i2.16177, 2013.

1080 Jagercikova, M., Evrard, O., Balesdent, J., Lefèvre, I. and Cornu, S.: Modeling the migration
1081 of fallout radionuclides to quantify the contemporary transfer of fine particles in Luvisol
1082 profiles under different land uses and farming practices, *Soil Tillage Res.*, 140(2014), 82–97,
1083 doi:10.1016/j.still.2014.02.013, 2014.

1084 Jagercikova, M., Cornu, S., Bourlès, D., Evrard, O., Hatté, C. and Balesdent, J.:
1085 Quantification of vertical solid matter transfers in soils during pedogenesis by a multi-tracer
1086 approach, *J. Soils Sediments*, 17(2), 408–422, doi:10.1007/s11368-016-1560-9, 2017.

1087 Jastrow, J. D., Amonette, J. E. and Bailey, V. L.: Mechanisms controlling soil carbon turnover
1088 and their potential application for enhancing carbon sequestration, *Clim. Change*, 80, 5–23,
1089 doi:10.1007/s10584-006-9178-3, 2007.

1090 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha,
1091 S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J.,
1092 Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, a., Reynolds, R., Jenne, R. and Joseph, D.:
1093 The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Meteorol. Soc.*, 77(3), 437–471,
1094 doi:10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2, 1996.

1095 Keeling, C. D. and Whorf, T. P.: Atmospheric CO₂ records from sites in the SIO air sampling
1096 network, Oak Ridge Natl. Lab. U.S. Dept. of Energy, Oak Ridge, Tenn., 2006.

1097 Keyvanshokouhi, S., Cornu, S., Samouëlian, A. and Finke, P.: Evaluating SoilGen2 as a tool
1098 for projecting soil evolution induced by global change, *Sci. Total Environ.*, 571, 110–123,
1099 doi:10.1016/j.scitotenv.2016.07.119, 2016.

1100 Kobayashi, K. and Salam, M. : Comparing simulated and measured values using mean
1101 squared deviation and its components, *Agron. J.*, 92(2), 345–352 [online] Available from:
1102 [http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Comparing+Simulated+and](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Comparing+Simulated+and+Measured+Values+Using+Mean+Squared+Deviation+and+its+Components#0)
1103 [+Measured+Values+Using+Mean+Squared+Deviation+and+its+Components#0](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Comparing+Simulated+and+Measured+Values+Using+Mean+Squared+Deviation+and+its+Components#0) (Accessed 2
1104 February 2012), 2000.

1105 Koven, C. D., Riley, W. J., Subin, Z. M., Tang, J. Y., Torn, M. S., Collins, W. D., Bonan, G.
1106 B., Lawrence, D. M. and Swenson, S. C.: The effect of vertically resolved soil
1107 biogeochemistry and alternate soil C and N models on C dynamics of CLM4, *Biogeosciences*,
1108 10(11), 7109–7131, doi:10.5194/bg-10-7109-2013, 2013.

1109 Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P.,
1110 Ciais, P., Sitch, S. and Prentice, I. C.: A dynamic global vegetation model for studies of the
1111 coupled atmosphere-biosphere system, *Glob. Biogeochem. Cycles*, 19(1),
1112 doi:10.1029/2003GB002199, 2005.

1113 Laclau, J. P., Bouillet, J. P. and Ranger, J.: Dynamics of biomass and nutrient accumulation in
1114 a clonal plantation of Eucalyptus in Congo, *For. Ecol. Manage.*, 128(3), 181–196,
1115 doi:10.1016/S0378-1127(99)00146-2, 2000.

1116 Mathieu, J. a., Hatté, C., Balesdent, J. and Parent, É.: Deep soil carbon dynamics are driven
1117 more by soil type than by climate: a worldwide meta-analysis of radiocarbon profiles, *Glob.*
1118 *Chang. Biol.*, n/a-n/a, doi:10.1111/gcb.13012, 2015.

1119 Mitchell, T. D., Carter, T. R., Jones, P. D., Hulme, M. and New, M.: A comprehensive set of
1120 high-resolution grids of monthly climate for Europe and the globe: the observed record
1121 (1901–2000) and 16 scenarios (2001–2100), ... *Cent. Clim. ...*, (July), 1–30 [online]
1122 Available from: <http://www.tyndall.ac.uk/sites/default/files/wp55.pdf>, 2004.

1123 Morrás, H., Moretti, L., Piccolo, G. and Zech, W.: Genesis of subtropical soils with stony
1124 horizons in NE Argentina: Autochthony and polygenesis, *Quat. Int.*, 196(1), 137–159,
1125 doi:10.1016/j.quaint.2008.07.001, 2009.

1126 O'Brien, B. J. and Stout, J. D.: Movement and turnover of soil organic matter as indicated by
1127 carbon isotope measurements, *Soil Biol. Biochem.*, 10(4), 309–317, doi:10.1016/0038-
1128 0717(78)90028-7, 1978.

1129 Parton, W., Schimel, D. S., Cole, C. and Ojima, D.: Analysis of factors controlling soil
1130 organic matter levels in Great Plains grasslands, *Soil Sci. Soc. Am. J.*, 51, 1173–1179 [online]
1131 Available from: [http://agris.fao.org/agris-](http://agris.fao.org/agris-search/search/display.do?f=1990/US/US90315.xml;US9004388)
1132 [search/search/display.do?f=1990/US/US90315.xml;US9004388](http://agris.fao.org/agris-search/search/display.do?f=1990/US/US90315.xml;US9004388) (Accessed 27 July 2011),
1133 1987.

1134 Parton, W. J., Stewart, J. W. B. and Cole, C. V.: Dynamics of C, N, P and S in grassland soils:
1135 a model, *Biogeochemistry*, 5(1), 109–131, doi:10.1007/BF02180320, 1988.

1136 Reimer, P. J., Brown, T. A. and Reimer, R. W.: Discussion: reporting and calibration of post-
1137 bomb ¹⁴C data., *Radiocarbon*, 46(1), 1299–1304, doi:10.2458/azu_js_rc.46.4183, 2004.

1138 Rosnay, P. de and Polcher, J.: MOdelling root water uptake in a complex land surface scheme
1139 coupled to a GCM, *Hydrol. Earth Syst. ...*, 2, 239–255 [online] Available from:
1140 <http://adsabs.harvard.edu/abs/1998HESS....2..239D> (Accessed 26 March 2013), 1998.

1141 Scharpenseel, H. W. and Schiffmann, H.: Radiocarbon dating of soils, a review, *Zeitschrift*
1142 *für Pflanzenernährung und Bodenk.*, 140(2), 159–174, doi:10.1002/jpln.19771400205, 1977.

1143 Schwartz, D., Mariotti, A., Trouve, C., Van Den Borg, K. and Guillet, B.: Etude des profils
1144 isotopiques ¹³C et ¹⁴C d'un sol ferrallitique sableux du littoral congolais : implications sur la

1145 dynamique de la matière organique et l'histoire de la végétation, *Comptes Rendus l'Académie*
1146 *des Sci. 2 Mécanique...*, 1992, 315, p. 1411-1417. ISSN 0249-6305, 315, 1411-1417, 1992.

1147 Sitch, S., Smith, B., Prentice, I. C., Arneth, a., Bondeau, a., Cramer, W., Kaplan, J. O.,
1148 Levis, S., Lucht, W., Sykes, M. T., Thonicke, K. and Venevsky, S.: Evaluation of ecosystem
1149 dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global
1150 vegetation model, *Glob. Chang. Biol.*, 9(2), 161-185, doi:10.1046/j.1365-2486.2003.00569.x,
1151 2003.

1152 Stuiver, M. and Polach, H. A.: Reporting of ^{14}C data, *Radiocarbon*, 19(3), 355-363,
1153 doi:10.1016/j.forsciint.2010.11.013, 1977.

1154 Taylor, K. E., Stouffer, R. J. and Meehl, G. a.: An overview of CMIP5 and the experiment
1155 design, *Bull. Am. Meteorol. Soc.*, 93(4), 485-498, doi:10.1175/BAMS-D-11-00094.1, 2012.

1156 Tisnérat-Laborde, N., Thil, F., Synal, H.-A., Hatté, C., Cersoy, S., Gauthier, C., Kaltnecker,
1157 E., Massault, M., Michelot, J.-L., Noret, A., Noury, C., Siani, G., Tombret, O., Vigne, J.-D.,
1158 Wacker, L. and Zazzo, A.: A new compact AMS facility measuring ^{14}C dedicated to
1159 Environment, Climate and Human Sciences, in 22nd International Radiocarbon Conference,
1160 Dakar, Senegal. November 2015. Oral presentation, pp. 16-20., 2015.

1161 Todd-Brown, K. E. O., Randerson, J. T., Post, W. M., Hoffman, F. M., Tarnocai, C., Schuur,
1162 E. a. G. and Allison, S. D.: Causes of variation in soil carbon simulations from CMIP5 Earth
1163 system models and comparison with observations, *Biogeosciences*, 10(3), 1717-1736,
1164 doi:10.5194/bg-10-1717-2013, 2013.

1165 Trumbore, S.: Age of Soil Organic Matter and Soil Respiration : Radiocarbon Constraints on
1166 Belowground C Dynamics, *Ecol. Appl.*, 10(2), 399-411, doi:10.1890/1051-
1167 0761(2000)010[0399:AOSOMA]2.0.CO;2, 2000.

1168 Wieder, W. R., Bonan, G. B. and Allison, S. D.: Global soil carbon projections are improved
1169 by modelling microbial processes, *Nat. Clim. Chang.*, 3(8), 1-4, doi:10.1038/nclimate1951,
1170 2013.

1171 WRB: World reference base for soil resources 2006: a framework for international
1172 classification, correlation and communication., 2006.

1173 Wynn, J. G., Bird, M. I. and Wong, V. N. L.: Rayleigh distillation and the depth profile of
1174 $^{13}\text{C}/^{12}\text{C}$ ratios of soil organic carbon from soils of disparate texture in Iron Range National
1175 Park, Far North Queensland, Australia, *Geochim. Cosmochim. Acta*, 69(8), 1961-1973, 2005.

1176
1177
1178
1179
1180
1181 |

1182
1183
1184
1185

1186
1187

1188
1189
1190
1191

Table 1. General description of the studied sites. The mean bulk density, pH and clay fraction values calculated from the different soil layers depths available from the data were used as input for each site. For the Mons and Feucherolles sites, min and max values of pH and clay fraction are provided between brackets.

Site name	Feucherolles	Mons	Kissoko	Misiones
Sampling Date	April 2011	March 2011	May 2014	May 2015
Location	France	France	Congo	Argentina
Coordinates	48.90°N, 1.97°E	49.87°N, 3.03°E	4.35°S, 11.75°E	27.65°S, 55.42°W
Elevation (m)	120	88	100	NA
Mean Annual Rainfall (mm)	660	680	1400	1850
Mean Annual Temperate (°C)	11.2	11	25	20
Soil Type (WRB)	Luvisol	Luvisol	Arenosol	Acrisol
Land Use	Temperate broad-leaved summergreen forest	Grassland	Native savanna	Tropical broad-leaved evergreen forest
Mean Bulk Density (g cm ⁻³)	1.34	1.4	1.48	1.15
Mean pH	5.9 (5.12-8.55)	6.9 (6.70-7.56)	5.2	5.2
Mean Clay Fraction (%)	20 % (13-30 %)	23 % (19-27 %)	5 %	58 %

Table 2. The main differences between the three simulations

	Flux from slow pool to passive pool	Turnover time of the passive pool (year)	Diffusion (m ² year ⁻¹)
<u>Control</u>	0.07	462	$D(z) = 1.10^{-4}$
<u>He et al., (2016) parameterization</u>	<u>0.0049</u>	6468	$D(z) = 1.10^{-4}$
<u>Depth-varying diffusion constant</u>	0.07	462	$D(z) = 5.42. 10^{-4} e^{(-0.04z)}$

Bertrand Guenet 30/10/y 10:08
Supprimé: over the profiles
Auteur inconnu 31/10/y 11:25
Supprimé: in

Bertrand Guenet 30/10/y 10:34
Supprimé: .

Bertrand Guenet 30/10/y 14:18
Supprimé: Model_Control
Bertrand Guenet 30/10/y 14:20
Supprimé: Model_Test He
Bertrand Guenet 30/10/y 10:10
Supprimé: 049
Bertrand Guenet 30/10/y 14:23
Supprimé: Model_Test Diffusion

1199
1200
1201
1202
1203
1204

Table 3. The correlation coefficient (r) between model outputs and measurements for carbon stock (kg C m⁻²) over the soil profile, for the four sites. The results of the initial version of the model ORCHIDEE-SOM-¹⁴C (Control) as well as those from the version including the modification according to (He et al., 2016) (He et al., (2016) parameterization) and diffusion varying according to the depth (Depth-varying diffusion constant), are provided.

		r	Mean total soil carbon (kg C m ⁻²) Model	Mean total soil carbon (kg C m ⁻²) Measurements
Misiones	Control	0.44	2.03	
	He et al., (2016) parameterization	0.69	7.38	2.14±0.30
	Depth-varying diffusion constant	0.46	2.23	
Kissoko	Control	0.14	0.76	
	He et al., (2016) parameterization	0.55	2.44	0.42±0.38
	Depth-varying diffusion constant	0.13	0.88	
Feucherolles	Control	0.20	0.70	
	He et al., (2016) parameterization	0.11	2.33	0.66±0.08
	Depth-varying diffusion constant	0.22	0.77	
Mons	Control	0.49	2.37	
	He et al., (2016) parameterization	-0.14	9.99	0.8±0.10
	Depth-varying diffusion constant	0.48	2.42	

1205
1206
1207
1208
1209
1210
1211
1212

Table 4. The correlation coefficient (r) between model outputs and measurements and the mean values (provided by the model and the measurements) over the profile according to F¹⁴C, for the four sites. The results of the initial version of the model ORCHIDEE-SOM-¹⁴C (Control) as well as those from the version including the modification according to (He et al., 2016) (He et al., (2016) parameterization) and diffusion varying according to the depth (Depth-varying diffusion constant), are provided.

		r	Mean Model	Mean Measurements
Misiones	Control	0.55	0.920	
	He et al., (2016) parameterization	0.50	0.560	0.930±0.009
	Depth-varying diffusion constant	0.60	0.900	
Kissoko	Control	0.40	0.995	
	He et al., (2016) parameterization	0.30	0.620	0.985±0.004
	Depth-varying diffusion constant	0.55	0.995	
Feucherolles	Control	0.55	0.915	
	He et al., (2016) parameterization	0.55	0.550	0.725±0.005
	Depth-varying diffusion constant	0.60	0.890	
Mons	Control	0.75	0.860	
	He et al., (2016) parameterization	0.70	0.510	0.815±0.005
	Depth-varying diffusion constant	0.80	0.835	

1213

Bertrand Guenet 30/10/y 10:40
Supprimé: and the mean values (pro... [35]
Auteur inconnu 31/10/y 11:26
Supprimé: if the passive pool turnov... [36]
Bertrand Guenet 30/10/y 14:20
Supprimé: Model_Test He...e et al., ... [37]
Auteur inconnu 31/10/y 11:26
Supprimé: ,
Bertrand Guenet 30/10/y 10:10
Supprimé: 3
Bertrand Guenet 30/10/y 10:10
Supprimé: 3
Bertrand Guenet 30/10/y 14:18
Supprimé: Model_Control
Bertrand Guenet 30/10/y 10:10
Supprimé: 55
Bertrand Guenet 30/10/y 10:10
Supprimé: 65
Bertrand Guenet 30/10/y 14:20
Supprimé: Model_Test He
Bertrand Guenet 30/10/y 10:10
Mis en forme ... [38]
Bertrand Guenet 30/10/y 10:10
Supprimé: 50
Bertrand Guenet 30/10/y 10:10
Supprimé: 128
Bertrand Guenet 30/10/y 10:10
Supprimé: 31
Bertrand Guenet 30/10/y 14:23
Supprimé: Model_Test Diffusion
Bertrand Guenet 30/10/y 10:10
Mis en forme ... [39]
Bertrand Guenet 30/10/y 10:10
Supprimé: 60
Bertrand Guenet 30/10/y 10:10
Supprimé: 57
Bertrand Guenet 30/10/y 14:18
Supprimé: Model_Control
Bertrand Guenet 30/10/y 10:10
Mis en forme ... [40]
Bertrand Guenet 30/10/y 10:10
Supprimé: 40
Bertrand Guenet 30/10/y 10:10
Supprimé: 34
Bertrand Guenet 30/10/y 14:20
Supprimé: Model_Test He
Bertrand Guenet 30/10/y 10:10
Mis en forme ... [42]
Bertrand Guenet 30/10/y 10:10
Supprimé: 40
Bertrand Guenet 30/10/y 10:10
Supprimé: 53
Bertrand Guenet 30/10/y 10:10
Mis en forme ... [41]
Bertrand Guenet 30/10/y 10:10
Supprimé: 30
Bertrand Guenet 30/10/y 14:23
Supprimé: Model_Test Diffusion
Bertrand Guenet 30/10/y 10:10
Supprimé: 50
Bertrand Guenet 30/10/y 10:10
Supprimé: 31
Bertrand Guenet 30/10/y 14:18
Supprimé: Model_Control
Bertrand Guenet 30/10/y 10:10
Bertrand Guenet 30/10/y 10:10
Bertrand Guenet 30/10/y 14:20
Bertrand Guenet 30/10/y 10:10
Mis en forme ... [43]
Bertrand Guenet 30/10/y 10:10
Bertrand Guenet 30/10/y 10:10
Bertrand Guenet 30/10/y 10:10
Bertrand Guenet 30/10/y 14:23

1356

Table 5. F¹⁴C profile obtained for each site.

Sites	Soil depth (cm)	F ¹⁴ C
Misiones	0-5	1.08
	5-10	1.04
	10-15	1.05
	15-20	0.99
	20-30	0.99
	30-40	0.87
	40-50	0.91
	50-60	0.76
	60-80	0.79
	80-100	0.79
Kissoko	0-5	1.06
	5-10	1.07
	10-15	1.07
	15-20	1.08
	20-30	1.05
	30-40	1.04
	40-50	1.02
	50-60	0.97
	60-80	0.90
	80-100	0.81
	100-120	0.72
Feucherolles	0-2	1.08
	16-18	1.05
	40-45	0.92
	75-85	0.69
	105-115	0.54
	125-135	0.53
	147-157	0.26
Mons	0-2	1.02
	2-4	1.03
	18-20	1.03
	45-50	0.87
	60-65	0.71
	82-92	0.65
	102-112	0.64
	142-152	0.55

1357

Bertrand Guenet 30/10/y 11:43

Mis en forme: Gauche, Interligne : multiple 1,15 li


Bertrand Guenet 30/10/y 11:43

Mis en forme: Exposant

Auteur inconnu 31/10/y 11:27

Supprimé: s

1359



1360


1361

1362 **Figure 1.** Overview of the different fluxes and processes in soil as presented in the version of
1363 ORCHIDEE-SOM adapted from Camino-Serrano et al. (2017).
1364

Bertrand Guenet 30/10/y 10:33
Supprimé: -

1366

F¹⁴C



1367


1368

1369

1370

Figure 2. Evolution of the F¹⁴C of atmospheric CO₂ in Argentina, Congo and France (data from Hua et al. 2013).

Bertrand Guenet 31/10/y 19:42



Atmospheric M

France
Congo
Argentina

Supprimé:

Bertrand Guenet 30/10/y 10:33

Mis en forme: Justifié, Interligne : multiple 1,15 li




Figure 3. Total soil carbon (kg C m^{-3}) according to the depth for the four sites. The results of the initial version of the model ORCHIDEE-SOM- ^{14}C (**Control**) as well as those from the version including the modification according to (He et al., 2016) (**He et al., (2016) parameterization**) and diffusion varying according to the depth (**Depth-varying diffusion constant**), are shown.

Supprimé: [50]

Supprimé: ,

Unknown

Mis en forme: Police : (Par défaut) Times New Roman

Unknown

Mis en forme: Police : (Par défaut) Times New Roman

Supprimé: Model_Control

Supprimé: if the passive pool turnover rate and the slow-to-passive flux revised

Supprimé: Model_Test He

Supprimé: Model_Test Diffusion

Supprimé: ,

Supprimé: ,

Kisso

Total Soil C

Supprimé:




Figure 4. Mean Squared Deviation (MSD) and its components for total soil carbon (kg C m^{-2}): lack of correlation weighted by the standard deviation (LCS), squared difference between standard deviations (SDSD) and the squared bias (SB). For the four sites, the results of the initial version of the model ORCHIDEE-SOM- ^{14}C (Control) as well as those from the version including the modification according to (He et al., 2016) (He et al., (2016) parameterization) and diffusion varying according to the depth (Depth-varying diffusion constant), are shown.

Bertrand Guenet 31/10/y 19:43

Supprimé: [51]

Bertrand Guenet 31/10/y 19:43

Mis en forme: Gauche, Interligne : multiple 1,15 li

Bertrand Guenet 30/10/y 10:54

Supprimé: 3

Bertrand Guenet 30/10/y 14:18

Supprimé: Model_Control

Auteur inconnu 31/10/y 11:28

Supprimé: if the passive pool turnover rate and the slow-to-passive flux revised


Bertrand Guenet 30/10/y 14:20

Supprimé: Model_Test He

Bertrand Guenet 30/10/y 14:23

Supprimé: Model_Test Diffusion

Bertrand Guenet 31/10/y 19:43



Supprimé:




Figure 5. Modern fraction $F^{14}C$ according to the depth, for the four sites. The results of the initial version of the model ORCHIDEE-SOM- ^{14}C (**Control**) as well as those from the version including the modification according to He et al., (2016) (**He et al., (2016) parameterization**) and diffusion varying according to the depth (**Depth-varying diffusion constant**), are shown.

Bertrand Guenet 30/10/y 10:25

Misio

Supprimé:

Unknown

Mis en forme: Police : (Par défaut) Times New Roman

Bertrand Guenet 30/10/y 14:18

Supprimé: Model_Control

Auteur inconnu 31/10/y 11:28

Supprimé: if the passive pool turnover rate and the slow-to-passive flux revised

Bertrand Guenet 30/10/y 14:20

Supprimé: Model_Test He

Bertrand Guenet 30/10/y 14:23

Supprimé: Model_Test Diffusion

Auteur inconnu 31/10/y 11:28

Supprimé: ,




Figure 6. Mean Squared Deviation (MSD) and its components: lack of correlation weighted by the standard deviation (LCS), squared difference between standard deviations (SDSD) and the squared bias (SB) calculated for modern fraction $F^{14}C$. For the four sites, the results of the initial version of the model ORCHIDEE-SOM- ^{14}C (Control) as well as those from the version including the modification according to He et al., (2016) (He et al., (2016) parameterization) and diffusion varying according to the depth (Depth-varying diffusion constant) are shown.

Bertrand Guenet 31/10/y 19:44

Supprimé:

Unknown

Mis en forme: Police :(Par défaut) Times New Roman

Unknown

Mis en forme: Police :(Par défaut) Times New Roman

Bertrand Guenet 30/10/y 14:18

Supprimé: Model_Control

Auteur inconnu 31/10/y 11:28

Supprimé: if the passive pool turnover rate and the slow-to-passive flux revised

Bertrand Guenet 30/10/y 14:20

Supprimé: Model_Test He

Bertrand Guenet 30/10/y 14:23

Supprimé: Model_Test Diffusion

Auteur inconnu 31/10/y 11:28

Supprimé:

Bertrand Guenet 30/10/y 10:31

Supprimé:




Figure 7. Relative proportion of each of the soil carbon pools summing the total soil carbon at each soil layer. The results of the initial version of the model ORCHIDEE-SOM-¹⁴C (Control, left pattern) as well as those from the version including the modification according to (He et al., 2016) (He et al., (2016) parameterization, pattern in the middle) and diffusion varying according to the depth (Depth-varying diffusion constant, right pattern), are shown.

Bertrand Guenet 31/10/y 19:44

Supprimé: -

[53]

Unknown

Mis en forme: Police :(Par défaut) Times New Roman

Bertrand Guenet 30/10/y 14:18

Supprimé: Model_Control

Auteur inconnu 31/10/y 11:29

Supprimé: if the passive pool turnover rate and the slow-to-passive flux revised

Bertrand Guenet 30/10/y 14:20

Supprimé: Model_Test He

Bertrand Guenet 30/10/y 14:23

Supprimé: Model_Test Diffusion

Auteur inconnu 31/10/y 11:29

Supprimé: ,