

Comments to the Author:

Thank you for your revised manuscript. I am sorry for the delay, which was caused by the previous topical editor no longer being available.

I think that you have answered most of the reviewers concerns, but there are still some problems with the manuscript.

Thank you for your time and effort to improve our manuscript. Please find below our response and a revised version of the manuscript.

Language...

The response to the reviewers and the revised manuscript are both very difficult to understand. I think the main problem is incoherence of language rather than incoherence of thought, but I cannot accept this manuscript until I have seen a version with much improved English.

Some of the problems:

Words are spelled incorrectly;

Words are used inappropriately;

Tenses are wrong;

Singular and plural are mixed up;

Punctuation is missing;

It is often not clear what "it" refers to;

Other grammatical errors that render sentences incomprehensible;

Several typos.

I will send you my copy of your response in which I have highlighted all the parts that I had to read multiple times in order to understand your meaning. Doing this helped me to understand your work. I hope it will also help you to improve the manuscript.

Note that "similar" means "resembling without being identical". If you write that something is similar to something else then you also need to mention in which ways it is different. If you actually mean identical/the same, then say so.

For example,

"Amongst these parameters, along with the basal effective pressure, the large scale bedrock curvature and/or sub-grid roughness could be used, similarly to Briggs et al. (2013)." could become,

"Amongst these parameters, along with the basal effective pressure, the large scale bedrock curvature and/or sub-grid roughness could be used, as in Briggs et al. (2013).", should this be the meaning that you intend.

Note, also, RMSE is not yielded or presented.

"... have an RMSE of ...", is sufficient.

None of the authors are English native and even with multiple thorough readings it makes it difficult for us to provide an error-free manuscript. We thank you for your time highlighting our mistakes, we did our best to provide a corrected revision in this new version.

On to the science...

Section 3 "Calibration of the Antarctic ice sheet"

This section is unconvincing. I think an additional paragraph is needed here, to explain what you are doing and why. What I would expect to see for a model calibration is the development of a cost

function that specifies a PDF of what you believe to be plausible models. The final set of models is then sampled from this PDF. In order to develop this PDF, discussion of the observations, the observational error, and the model adequacy (or discrepancy) need to be included.

What you have actually done seems rather bizarre. You appear to have 3 observations (total volume, ice thickness, ice velocity), and yet you have used only one of these (thickness). You have included in your final ensemble only those models with the lowest RMSE compared to that single metric. Observational error is not mentioned. Discussion of model (or in this case experimental) adequacy suddenly appears in a paragraph of the Discussion (the one that starts "Although widely used...") where you explain that you do not anyway believe the result of your calibration. I conclude from this that you in no way believe that your 12 "best" ensemble members are in fact your "best"! Would you be planning to use this set of 12 in any future work?

I might be being stupid, but I cannot find figures showing the observed thickness, or uncertainty in that thickness. In addition the plots of RMSE are not very informative as they do not indicate whether the ice is too thin or too thick. I also cannot tell whether (and where) the whole ensemble is biased, which is surely important information.

I wonder whether what you are doing is more of a sensitivity test than a calibration. It seems to me that what you are investigating is the range of model behaviours after a strong constraint is placed on ice thickness. If so, then the section needs quite a bit of rewriting to make this clear.

On the choice of the metric:

We have added a new paragraph to the new version of the manuscript to justify our choices.

it is true that we discuss three possible ways to constrain the model: ice volume, ice thickness and ice velocity. However, these three are not equivalent for the purpose of calibrating ice sheet models.

Ice velocities are a widely used target for ice sheet models, in particular when using inverse methods to tune ice dynamics. In this case, ice geometry (ice thickness) is generally prescribed to its present-day value and a dynamical parameter (e.g. basal drag coefficient) is tuned to reproduce the ice velocity field (e.g. Price et al., 2010; Gillet-Chaulet et al., 2012).

When no inversion methodology is used, a useful metric based on ice velocities is more complicated to define:

- Ice velocities are spatially highly variable and present their maximum values at the ice sheet margins. This means that small errors in the simulated extent of the ice sheet lead to important discrepancy with the observations. As such, marginal regions, which represent a small fraction of the ice sheet, have more weight for metrics such as the RMSE. For example: having the shape of a simulated ice shelf slightly different from the observations is not crucial for ice dynamics but have important consequences on a RMSE using ice velocities.
- In addition, ice velocities show fine scale structure resulting from the complex bedrock below the ice sheet (together with geologic bed properties, hydrological and thermal conditions). Our coarse-resolution model (40 km) is not suited to reproduce the fine scale structures and can only reproduce the broad pattern.
- Finally, aggregation of fine scale (~100 m) ice streams with neighbouring slow ice is not meaningful at 40 km resolution.

For all these reasons, ice velocities are in general not used as a tuning target for large spatial scale ice sheet simulations at coarse resolution.

Conversely, ice sheet geometry (in our case, assessed by ice thickness RMSE) is usually a primary target for ice sheet modellers (e.g. Ritz et al., 1997; Stone et al., 2010; Applegate et al., 2012; Pollard et al., 2016). The advantage of ice geometry is that it is relatively well constrained for present-day but also in the past (ice margin migrations from geological evidence, e.g. DATED and

RAISED community effort). This means that ice sheet models can sometimes be validated with respect to our knowledge on palaeo-geometries.

Ice volume is the spatial integration of ice thickness and is generally a poor metric to assess model performance due to potential compensatory biases. In turns, ice thickness RMSE is a more robust metric.

For all these reasons, we follow the approach commonly adopted by the community building our main metric on ice geometry. The justification of the approach at the end of Sec. 3.1 reads:

“In the following, individual member performance is assessed with the root mean square error (RMSE) computed from simulated and observed ice thickness (Fretwell et al., 2013). This metric puts a strong constraints on ice sheet geometry and avoids potential compensatory biases that could appear when using total ice volume. Observed ice velocities are not used as a metric because of high spatial variability that results from small-scale bed properties. With a coarse 40-km resolution, the model is intrinsically unable to reproduce such variability but is expected to reproduce the large scale pattern of ice thickness. However, at the end of the long 100-kyrs simulation, the velocities in the model are the balance velocities corresponding to the simulated topography. Thus, the minimisation of the RMSE in ice thickness should also reduce the error in velocities with respect to observations at the global scale.”

On observational error:

Following your advice we added a figure showing the ice thickness observational dataset together with its uncertainty (Fretwell et al., 2013). Fretwell et al. estimate generally an uncertainty of about ± 150 m. However, there are some regions for which the lack of in-situ measurements forces the authors to use gravity-derived ice thickness estimates only. The estimated uncertainty then reaches ± 1000 m. We agree that the addition of this figure helps the reader to interpret our model results. We explicitly refer to this figure when commenting the model bias (new reference in Sec. 3.2).

On model biases:

The response of the model within our large ensemble presents both positive and negative volume anomalies, with members overestimating (about +38% within AN40S and +27% within AN40T) and members underestimating (about -27%) total ice volume. This is shown in Fig. 5 and Fig. 6. Our ensemble is thus not biased towards smaller/larger ice sheets. However, as explained in the manuscript (Sec. 3.2), we have persisting model biases in some regions (East Antarctica and Ronnie-Filchner basin), as shown in Fig. 7 and Fig. 8.

On the usefulness of the parameters yielded during the calibration:

In the discussion we mention that a calibration based on glacial-interglacial simulations is ideally preferred to the approach followed in Sec. 3.1 (perpetual modern climate). However, the determination of a realistic climate forcing for palaeo-simulations is a considerable challenge and mechanical parameters inferred with such an approach will necessarily reflect the assumptions made on past climate change. Even though the parameter values found in Sec. 3.1 minimising the RMSE do not take into account past climate change, they are nonetheless useful since, as shown in Sec. 3.2, they are able to provide glacial-interglacial ice geometry changes that roughly fit available palaeo-data constraints. To emphasize this point, we added the following in the discussion:

“[...] behaviour for long-term integrations. Using the parameters calibrated under perpetual modern climate, the model is nonetheless able to reproduce ice geometry changes compatible with palaeo-constraints. Further work [...]”

Numerical solution methods...

I have rarely seen a manuscript in which the reviewers clamour so loudly for the code to be made available. While some description of the numerical solution methods has been added to the manuscript, it is still rather generic (eg "an upwind scheme" !!) and I think this is insufficient for GMD's aim of "scientific reproducibility". One possibility would be writing the difference schemes in an appendix or in the supplement, but this is unlikely to fully satisfy all readers, and a simpler solution would be to be more specific in the manuscript while also making the code developed in this paper available in the supplement.

I do not fully understand the grounds under which the code is (or is not) being made available at present. You wrote in your response,

"The public distribution needs to be done with a specific license (e.g. GNU public licence). Such a license is currently lacking in GRISLI. The model has been developed from the 90ies and has benefited from the additions from numerous contributors. To put a license to the current model, we have to get the permissions to all the contributors from the past 20 years. To date, we do not have this permission from all the contributors. However, as stated in the manuscript, any potential users are encouraged to get in touch with C. Dumas, A. Quiquet or C. Ritz, to start a collaboration."

My understanding is that it is the employers, rather than the individuals, who need to give permission. How many of these employers have you been unable to track down (are some no longer active?), and which parts of the code are still restricted? If you upload all parts of the code that are not restricted, will this include the parts of the code that the reviewers want to see (ie the numerical solution to the equations outlined in this manuscript). If so then this may be satisfactory solution for the present manuscript. It seems strange that you can grant access upon request, but not release the code - what is the current license that users get when given access to the repository? I do understand that it can take a long time (years!) to achieve full code access for older models, but I would like to see some evidence in the manuscript that this is being actively pursued. It is already foreseeable that in the future all code developed in scientific publications will have to be made publicly available, so you are strongly encouraged to make this a reality for your own model sooner rather than later.

Encouraged by referee #3 comment and yours, we started to put parts of the code under the CeCILL public licence v 2.1 which is our employer's requirement for compatibility with GNU/GPL. Previously requested code sections are attached to the present revision of the model and are intended to be public. Sections of the code currently under the CeCILL licence include:

- Resolution of the elliptic equation
- Implementation of Schoof (2007) and Tsai et al. (2015) in the model
- Basal hydrology
- Passive tracer

We hope that these files cover most of the critical sections of the model and will be helpful for the readers.

We are currently working to release the whole model code under the CeCILL license. This however is work in progress and will require additional work. We will update you with the status of our efforts by the end of the review process in order to make a final version of the code availability section.

References

- Applegate, P. J., Kirchner, N., Stone, E. J., Keller, K., and Greve, R.: An assessment of key model parametric uncertainties in projections of Greenland Ice Sheet behavior, *The Cryosphere*, 6, 589–606, doi:10.5194/tc-6-589-2012, 2012.
- Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, doi:10.5194/tc-6-1561-2012, 2012.
- Pollard, D., Chang, W., Haran, M., Applegate, P., and DeConto, R.: Large ensemble modeling of the last deglacial retreat of the West Antarctic Ice Sheet: comparison of simple and advanced statistical techniques, *Geosci. Model Dev.*, 9, 1697–1723, <https://doi.org/10.5194/gmd-9-1697-2016>, 2016.
- Price, S.F., Payne, A.J., Howat, I.M. and Smith, B.E. Committed sea-level rise for the next century from Greenland ice sheet dynamics during the past decade. *PNAS* 108, 8978-8983, doi:10.1073/pnas.1017313108, 2011.
- Ritz, C., Fabre, A., and Letréguilly, A.: Sensitivity of a Greenland ice sheet model to ice flow and ablation parameters: consequences for the evolution through the last climatic cycle, *Climate Dynamics*, 13, 11–23, doi:10.1007/s003820050149, 1997.
- Stone, E. J., Lunt, D. J., Rutt, I. C., and Hanna, E.: Investigating the sensitivity of numerical model simulations of the modern state of the Greenland ice-sheet and its future response to climate change, *The Cryosphere*, 4, 397–417, doi:10.5194/tc-4-397-2010, 2010.
- Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375–393, doi:10.5194/tc-7-375-2013, 2013.

The GRISLI ice sheet model (version 2.0): calibration and validation for multi-millennial changes of the Antarctic ice sheet

Aurélien Quiquet¹, Christophe Dumas¹, Catherine Ritz², Vincent Peyaud², and Didier M. Roche^{1,3}

¹Laboratoire des Sciences du Climat et de l'Environnement (LSCE), UMR8212, CEA/CNRS-INSU/UVSQ, Gif-sur-Yvette Cedex, France

²Université Grenoble Alpes, CNRS, IRD, IGE, F-38000 Grenoble, France

³Earth and Climate Cluster, Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

Correspondence to: A. Quiquet (aurelien.quiquet@lsce.ipsl.fr)

Abstract.

In this paper we present the GRISLI (Grenoble Ice Sheet and Land Ice) model in its newest revision (version 2.0). Whilst GRISLI is applicable to any given geometry ice sheet, we focus here on the Antarctic ice sheet because it highlights the importance of grounding line dynamics. Important improvements have been implemented in the model since its original version (Ritz et al., 2001) ~~including notably~~. Notably, GRISLI now includes a basal hydrology model and an explicit flux computation at the grounding line based on the analytical formulations of Schoof (2007) or Tsai et al. (2015) ~~and a basal hydrology model. A~~. We perform a full calibration of the ~~mechanical parameters of the~~ model based on an ensemble of 300 ~~members sampled with simulations sampling mechanical parameter space using~~ a Latin Hypercube method ~~is used~~. ~~The ensemble members performance~~. Performance of individual members is assessed relative to the deviation from present-day observed Antarctic ice thickness. ~~The model being designed for multi-millennial long-term integrations, we~~ We also present glacial-interglacial ice sheet changes throughout the last 400 kyr using the best ensemble members taking advantage of the capacity of the model to perform multi-millennial long-term integrations. To achieve this goal, we construct a simple climatic perturbation of present-day climate forcing fields based on two climate proxies, both atmospheric and oceanic. The model is able to reproduce expected grounding line advances during ~~glacials~~ glacial periods and subsequent retreats during terminations with reasonable glacial-interglacial ice volume changes.

1 Introduction

Continental ice sheets are a major ~~component for the~~ climatic component for Earth system dynamics ~~operating~~. They operate on a variety of timescales, from diurnal to multi-millennial, through multiple feedbacks such as temperature - surface albedo, gravity waves and oceanic circulation changes related to freshwater flux release. Over the last decades, ~~modern observation techniques~~ observations of the Greenland and Antarctic ice sheets (e.g. altimetry, gravimetry, echo sounding) have shown important changes such as an increase in surface and sub-shelf melt, glacier speed-up, dynamical thinning and drastic calving events (e.g. Mougnot et al., 2015; Paolo et al., 2015; Hall et al., 2013; Pritchard et al., 2009). ~~If the two major~~ While the two current ice sheets have been mostly stable for at least the last 1000 years, ~~their contribution to global~~ they are expected

~~to contribute to future~~ sea level rise ~~in the future is largely uncertain~~, albeit with a largely uncertain magnitude. Conversely, in the past, there is evidence of sea level rise as fast as four metres per century (e.g. Fairbanks, 1989; Hanebuth et al., 2000; Deschamps et al., 2012). The study of well-recorded past events can help us to constrain the fate of the ice sheets in a warming Earth and to disentangle the role of the different processes leading to rapid ice sheet destabilisation. The surface mass balance-
5 height feedback has often been proposed as a major driver for rapid northern Hemisphere ice ~~sheets~~ sheet disintegration (e.g. Abe-Ouchi et al., 2013; Gregoire et al., 2012). Another source of instability, for marine ice sheets such as the palaeo Kara-Barents or present-day Antarctic ice sheets, is related to the fact that large parts of the bedrock presents a retrograde slope from the grounding line ~~and deepens~~. Isostatic adjustment can, for instance, deepen the bedrock well below sea level ~~due, for instance, to isostatic adjustment or within glacial troughs~~. Such. Additionally, glacial troughs may form a deepening bedrock
10 from the grounding line towards the ice-sheet centre. Such bed properties lead to the marine ice sheet instability (MISI, Weertman, 1974; Schoof, 2007) responsible for an irreversible retreat of the grounding line in response to an initial perturbation such as local sea level change and/or increase in basal melting rate below ice shelves. ~~Such~~ The latter two processes are expected to play a crucial role for the stability of the Antarctic ice sheet in the future (e.g. Favier et al., 2014). Additional instabilities may also occur on neutral/prograde bed slopes in relation ~~with the~~ to structural instabilities of tall ice cliffs (marine ice cliff
15 instability, MICI, Pollard et al., 2015).

~~Because~~ Continental ice sheets are difficult to model because they include processes operating on ~~variety in a variety of~~ temporal, from diurnal to multi-millennial, and spatial scales, from a few metres to thousand of kilometres, but also ~~due to the lack of~~ because we lack crucial observations (e.g. basal conditions and internal thermo-mechanics), ~~continental ice sheets are difficult to model~~. Most numerical models consider ice sheets as an incompressible fluid, where motion can be described with the Navier-Stokes equations. Even if some processes have generally to be parameterised (e.g. ice anisotropy), the complete set of equations can be solved explicitly and does not require the use of any approximation. The most comprehensive continental ice sheet models, namely the Full-Stokes models, solve explicitly all the terms in the stress tensor (Gillet-Chaulet et al., 2012; Larour et al., 2012). Recent applications are promising but, due to computational cost, continental scale applications are currently limited to a few centuries. As such, they are not yet the most suitable tool for palaeo-reconstructions or multi-millennial future projections.

In order to decrease the degree of complexity, simpler models were historically developed that make use of the small aspect ratio of ice sheets (vertical to horizontal scale ratio) to derive approximations for the Navier-Stokes equations (e.g. Hindmarsh, 2004). Such models ~~present the advantage to be much cheaper compared to~~ are computationally much cheaper than Full-Stokes models, allowing for multi-millennial integrations. They are well-suited to study slow feedbacks such as glacio-isostasy or the impact of temperature and surface mass balance perturbations. The Grenoble ice sheet and land ice (GRISLI) model ~~belongs to the latter category is one of these simpler models~~ (Ritz et al., 2001). GRISLI consists of the combination of the inland ice model of Ritz (1992) & Ritz et al. (1997) and the ice shelf model of Rommelaere and Ritz (1996), extended to the case of ice
35 streams treated as *dragging ice shelves*. GRISLI was in the late nineties the first large scale ice sheet model with an hybrid

shallow ice / shallow shelf system of equations. Whilst since Ritz et al. (2001) the fundamental equations for ice dynamics have not drastically changed, the model has nonetheless benefited from ~~the numerous works which used it~~ numerous contributions. To date, 30 papers published or in peer-review discuss GRISLI model simulation results. The range of applications have been very wide ranging from ice sheet reconstructions for deep-time palaeo-climate (e.g. Benn et al., 2015; Donnadieu et al., 2011; 5 Ladant et al., 2014) and Quaternary (e.g. Peyaud et al., 2007; Alvarez-Solas et al., 2013; Quiquet et al., 2013; Colleoni et al., 2016) to future sea level rise projections (e.g. Ritz et al., 2015; Peano et al., 2017). GRISLI has participated in several inter-comparison exercises (Calov et al., 2010; Edwards et al., 2014; Koenig et al., 2015; Goelzer et al., 2017) and has been coupled to climate models of various complexities (e.g. Philippon et al., 2006; Roche et al., 2014; Le clec'h et al., 2018).

10 The aim of our current study is to provide a technical description of the GRISLI model in its current version (GRISLI version 2.0, hereafter GRISLI), including ~~the~~ several additional features from Ritz et al. (2001). In particular, we have now included an explicit flux computation at the grounding line following the analytical formulation from Schoof (2007) and Tsai et al. (2015) in order to have a better representation of the grounding line migration. We also provide details on some components (sub-glacial hydrology and tracking particle scheme embedded in GRISLI) which are currently not documented in international scientific 15 journals. In addition, we present a simple calibration of the mechanical parameters suitable for multi-millenia integrations and we show an example of the model response to glacial-interglacial forcing.

In Sec. 2 we describe the fundamental equations of the GRISLI ice sheet model with a particular emphasis on the model developments departing from Ritz et al. (2001). In sec. 3, we present a simple calibration methodology which aims at repro- 20 ducing the observed present-day Antarctic ice sheet geometry. In Sec. 4, we discuss the ability of the model to simulate the Antarctic ice sheet changes over the last four glacial-interglacial cycles.

2 The GRISLI ice sheet model

GRISLI constitutive equations were presented in Ritz et al. (2001) and we aim here at giving a broad comprehensive description 25 of the current model version, with its latest functionalities. Major model parameters are listed in Tab. 1.

2.1 Ice thermo-mechanics

2.1.1 Ice deformation and mass conservation

Ice deformation and mass conservation in GRISLI version 2.0 is mostly treated as in Ritz et al. (2001) with the notable excep- tion of the use of a polynomial flow law with the introduction of a linear, Newtonian, viscosity.

30 GRISLI considers the ice sheet as solely formed of pure ice with a constant and homogeneous density (ρ). In this approxima-

tion, the ice being considered as an incompressible fluid, the mass conservation equation can be written as:

$$\nabla \cdot \mathbf{u} = \frac{\partial u_x}{\partial x} + \frac{\partial u_y}{\partial y} + \frac{\partial u_z}{\partial z} = 0 \quad (1)$$

with (u_x, u_y, u_z) the Cartesian components of the ice velocity field.

The vertically integrated expression of the mass conservation equation (Eq. 1) provides the equation for the ice thickness, H :

$$5 \quad \frac{\partial H}{\partial t} = -\frac{\partial H \bar{u}_x}{\partial x} - \frac{\partial H \bar{u}_y}{\partial y} + M - b_{melt} \quad (2)$$

with \bar{u}_x and \bar{u}_y the vertically integrated velocities in x and y -directions, M the surface mass balance and b_{melt} is the basal melting rate.

The quasi-static approximation is used for the velocity field, in which the inertial terms of the momentum conservation equation are ignored. The gravity force being the sole external force acting on an infinitesimal cube of ice, we have:

$$10 \quad \begin{cases} \frac{\partial \sigma_x}{\partial x} + \frac{\partial \tau_{xy}}{\partial y} + \frac{\partial \tau_{xz}}{\partial z} = 0 \\ \frac{\partial \tau_{xy}}{\partial x} + \frac{\partial \sigma_y}{\partial y} + \frac{\partial \tau_{yz}}{\partial z} = 0 \\ \frac{\partial \tau_{xz}}{\partial x} + \frac{\partial \tau_{yz}}{\partial y} + \frac{\partial \sigma_z}{\partial z} = \rho g \end{cases} \quad (3)$$

where $\tau_{ij=x,y,z}$ are the shearing stress tensor terms and $\sigma_{i=x,y,z}$ the longitudinal stress tensor terms, defined as ($i = x, y, z$):

$$\sigma_i = \tau_{ii} \quad (4)$$

The pressure is defined as the first invariant of the stress tensor:

$$-P = \frac{\sigma_x + \sigma_y + \sigma_z}{3} \quad (5)$$

15 The deviatoric stress tensor is defined as (for $i, j = x, y, z$):

$$\tau'_{ij} = \tau_{ij} + \delta_{ij} P \quad (6)$$

with δ_{ij} being 1 for $i = j$, 0 otherwise.

Assuming isotropy, the deviatoric stress and the deformation rate ϵ'_{ij} are related by:

$$\tau'_{ij} = 2\eta \epsilon'_{ij} \quad (7)$$

20 where η is the ice viscosity.

Like most ice sheet models, GRISLI considers the ice as a non-Newtonian viscous fluid following a Norton-Hoff constitutive law (commonly named Glen flow law):

$$\frac{1}{\eta} = B_{AT} \tau^{n-1} \quad (8)$$

where B_{AT} is a temperature-dependent coefficient and τ is the effective shear stress, defined as (for $i = x, y, z$):

$$\tau^2 = \frac{1}{2} \sum_{i,j} \tau_{ij}^2 \quad (9)$$

The ~~temperature-dependant~~ temperature-dependent coefficient B_{AT} is computed ~~with an Arrhenius~~ following an Arrhenius' law:

$$5 \quad B_{AT} = S_f B_{BAT0} \exp\left(\frac{E_a}{R} \left(\frac{1}{T_m} - \frac{1}{T}\right)\right) \quad (10)$$

where S_f is a flow enhancement factor, B_{BAT0} is a flow law coefficient, E_a is the activation energy, R the gas constant, T_m the ice pressure melting temperature and T the local temperature.

To account for the fact that the activation energy increases close to the melting point (Le Gac, 1980), in GRISLI, the pair (E_a , B_{BAT0}) can take two different values for local temperatures higher or lower than a temperature threshold T^{trans} (Tab. 1).

10 The Glen flow law is an empirical formulation, derived from laboratory experiments. However, laboratory experiments ~~fail at reproducing the~~ cannot cover the full range of deviatoric stress ~~values expected operating~~ in real ice sheets. The timescale over which this stress is applied in real ice sheets is also not reproducible in laboratories. ~~If most modelling studies agree to~~ Most modelling studies use $n = 3$ for the Glen flow law exponent (Eq. 8), ~~there is room for uncertainties as a~~ A few studies suggest the possibility for a smaller exponent for small stress regime (Duval and Lliboutry, 1985; Pimienta, 1987; 15 Pettit and Waddington, 2003). Since the work of Dumas (2002), one ~~particularity~~ specificity of GRISLI is the possibility to simultaneously use a Glen viscosity with $n = 3$ and a linear, Newtonian, viscosity with $n = 1$. In this case, the two contributions simply add up:

$$2 \dot{\epsilon}_{ij} = (B_{AT1} + \tau^2 B_{AT3}) \tau'_{ij} \quad (11)$$

with B_{AT1} and B_{AT3} computed with Eq. 10, only using different activation energy (E_{a1} and E_{a3}), flow law coefficient 20 (B_{BAT01} and B_{BAT3}) and temperature threshold (T_1^{trans} and T_3^{trans}). The chosen values (Tab. 1) are based on Lipenkov et al. (1997).

Like other large scale ice sheet models, GRISLI does not explicitly take into account ~~explicitly~~ anisotropy, which tends to facilitate deformation due to vertical shear, but reduces deformation due to longitudinal stress. The role of the flow enhancement factor S_f in Eq. 10 is to artificially account for the effect of anisotropy. S_f takes different values for the two components of 25 the velocities computed in GRISLI (due to vertical shearing or longitudinal deformation, presented in the next section). In particular, we use a fixed ratio, typically ranging from 5:1 to 10:1 (Ma et al., 2010), between the value of the enhancement factor S_f for vertical shear and the one for longitudinal deformation. In the following, this ratio will be set to 8:1 and the enhancement factor S_f for vertical shearing with a Glen viscosity will be part one of the calibrated parameters. In turn, the flow enhancement factor for the vertical shearing with a linear viscosity is set to 1.

30 2.1.2 Ice velocity

Differing from Ritz et al. (2001), the velocity in GRISLI is now computed for the entire domain as the superposition of the shallow ice approximation (SIA) and the shallow shelf approximation (SSA) components, without using a sliding law to esti-

mate basal velocities.

The SIA (Hutter, 1982) assumes that the ~~horizontal derivatives of velocity~~ velocity horizontal derivatives are much smaller than its vertical derivatives, which is generally true for the major part of the ice sheet where the gravity driven flow induces a slow motion of the ice. GRISLI uses ~~the zero order~~ a zero order approximation of the SIA ~~approximation~~ in which the stress tensor components ~~simplified~~ simplify to:

$$\begin{cases} \tau_{xz} = \rho g \frac{\partial S}{\partial x} (S - z) \\ \tau_{yz} = \rho g \frac{\partial S}{\partial y} (S - z) \end{cases} \quad (12)$$

with S the surface elevation. The vertical velocity profile is computed as an integral from the bedrock elevation, B , to a given vertical coordinate, z (for $i = x, y$):

$$u_i(z) = u_{ib} + \int_B^z 2 \dot{\epsilon}_{iz} dz \quad (13)$$

where u_{ib} is the i component of the basal velocity.

The basal velocity can be computed with a sliding law (e.g. Bindschadler, 1983). However, recent versions of GRISLI use the shallow shelf approximation (SSA) as a sliding law as suggested by Bueler and Brown (2009). In this case, similarly to Winkelmann et al. (2011) we simply add up the two contributions of the SIA and SSA for the whole domain which ensure a smooth transition from non-sliding frozen regions to sliding over a thawed bed.

For ~~the~~ fast flowing regions, the vertical stresses are much smaller than the longitudinal shear stresses. In this case, the velocity fields with the SSA (MacAyeal, 1989) ~~reduced~~ reduce to the following elliptic equations:

$$\begin{cases} \frac{\partial}{\partial x} \left(2\bar{\eta} H \left(2 \frac{\partial u_x}{\partial x} + \frac{\partial u_y}{\partial y} \right) \right) + \frac{\partial}{\partial y} \left(\bar{\eta} H \left(\frac{\partial u_x}{\partial y} + \frac{\partial u_y}{\partial x} \right) \right) = \rho g H \frac{\partial S}{\partial x} - \tau_{bx} \\ \frac{\partial}{\partial y} \left(2\bar{\eta} H \left(2 \frac{\partial u_y}{\partial y} + \frac{\partial u_x}{\partial x} \right) \right) + \frac{\partial}{\partial x} \left(\bar{\eta} H \left(\frac{\partial u_y}{\partial x} + \frac{\partial u_x}{\partial y} \right) \right) = \rho g H \frac{\partial S}{\partial y} - \tau_{by} \end{cases} \quad (14)$$

where τ_b is the basal drag. The velocities u_x and u_y are identical along the vertical dimension.

The condition at the front of the ice shelf is given by the balance between the water pressure and the horizontal longitudinal stress (see also Sec. 2.3 on the numerical features).

25 2.1.3 Basal drag

For floating ice shelves, the basal drag, τ_b , is ~~neglected~~ negligible. For cold-based grounded ice we impose a large enough basal drag (typically 10^5 Pa) to ensure virtually no-slip conditions on the bedrock and the basal velocity is set to zero in this case.

For temperate-based grounded ice, a power-law basal friction (Weertman, 1957) is assumed:

$$\begin{cases} \tau_{bx}^m = -\beta u_{bx} \\ \tau_{by}^m = -\beta u_{by} \end{cases} \quad (15)$$

where the basal drag coefficient β is positive. In the experiment presented here, we assume the presence of a sediment at the base of the ice sheet allowing for a viscous deformation ($m = 1$).

5

In some recent applications of GRISLI, the basal drag coefficient has been inferred with an inverse method in order to match present-day ice sheet geometry (Ritz et al., 2015; Le clec'h et al., 2018). This approach has been followed to participate in the first phase of the recent ice sheet model intercomparison project (ISMIP6, Nowicki et al., 2016) for both the Greenland (Goelzer et al., 2017) and Antarctic ice sheets. In this context, GRISLI ~~achieves to provide~~ computes sea level rise projections by the end of the century in line with ~~the~~ results from more complex model (Edwards et al., 2014; Goelzer et al., 2017).

10

Inverse methods are ~~particularly adapted~~ especially suited to produce an ice sheet state (e.g. geometry and/or velocity) close to observations. However, by construction, such methods do not provide information ~~for grid points that are not glaciated today~~ by construction where no ice is present in observations. As such they are difficult to apply for palaeo reconstructions of the American or Eurasian ice sheets. More generally, inverse methods are no longer appropriate for long-term integrations, either palaeo or future, when ice thickness is very different from its present state and especially ~~when ice boundary if the ice margin~~ migrates from its present-day position. This motivates the use of a basal drag coefficient depending on computed from GRISLI internal variables. We generally assume that its value is modulated by the effective water pressure, N , at the base of the ice sheet:

15

20

$$\beta = C_f N \quad (16)$$

with C_f an internal parameter that needs calibration.

In our approach, any temperate-based grounded point will have a non-zero sliding velocity, depending on the C_f factor used. Previous works have used additional criteria to limit the extension of these ice streams. For example, Alvarez-Solas et al. (2010) only compute Eq. 16 where the present-day sediment thickness exceeds a certain threshold whilst Quiquet et al. (2013) restrict this to large-scale valleys. However these approaches have flaws. For example, the sediment distribution is only poorly known below present-day ice sheets and its past evolution is largely uncertain. Also, the definition of a typical spatial scale for ice streams is ~~somehow~~ somewhat arbitrary. For these reasons, in the following we use the simplest approach and compute Eq. 16 for any temperate-based grid point. The C_f parameter will be part of the calibrated parameters in our large ensemble.

25

2.1.4 Flux at the grounding line

In Ritz et al. (2001) the position of the grounding line was computed from a simple floatation criterion with no specific flux computation at the grounding line. Such an approach is in theory only valid for a very high spatial resolution, within tenths of meters, at the vicinity of the grounding line (Durand et al., 2009). Because the model runs typically at a much coarser resolution, in GRISLI version 2.0 we have implemented an explicit flux computation at the grounding line based on the analytical formulations from Schoof (2007) and Tsai et al. (2015). The two formulations differ from the assumption made on the sediment rheology.

The flux at the grounding line following Schoof (2007) is:

$$q_{gl}^S = - \left(\frac{\bar{A}(\rho g)^{n+1}(1 - \rho/\rho_w)^n}{4^n \beta} \right)^{\frac{m}{m+1}} H_{gl}^{\frac{m(n+3)+1}{m+1}} \phi_{bf}^{\frac{m-n}{m+1}} \quad (17)$$

with n and m being the exponents in the Glen flow law (Eq. 8) and in the friction law (Eq. 15) respectively, \bar{A} the vertically integrated temperature dependent coefficient in the Glen flow law (B_{AT} , Eq. 8), H_{gl} the ice thickness at the grounding line, and ϕ_{bf} a back force coefficient to take into account the buttressing role of ice shelves. β is the basal drag coefficient presented in Eq. 15.

Conversely, Tsai et al. (2015) proposed:

$$q_{gl}^T = Q_0 \frac{8\bar{A}(\rho g)^n}{4^n f} (1 - \rho/\rho_w)^{n-1} H_{gl}^{n+2} \phi_{bf}^{n-1} \quad (18)$$

with $Q_0 = 0.61$. In this case, the basal drag is assumed to vanish at the grounding and as such the coefficient β is not used. Instead Tsai et al. (2015) suggests a constant and homogeneous basal friction coefficient f set to 0.6.

In GRISLI, from the last grounded point in the direction of the flow, we compute the subgrid position of the grounding line in the x and y directions linearly interpolating the floatation criterion (dark green dots in Fig. 2). From this position, the flux at the grounding line is calculated using Eq. 17 or Eq. 18 (red arrows in Fig. 2). Because the flux at the sub-grid position is perpendicular to the local grounding line, ideally we should project this flux onto the x and y -axis. However, in the model, we assume that the grounding line is always perpendicular to either the x or y -axis (dashed brown line in Fig. 2). **Similarly to** [As in Füst \(2013\)](#), the value of the flux q_{gl} is linearly interpolated to the two closest downstream and upstream velocity grid points (dark blue arrows in Fig. 2) using the two bounding velocity points (light blue arrows in Fig. 2).

To evaluate the back force coefficient ϕ_{bf} , we solve the velocity equation twice. The first iteration is computed on the simulated geometry with no specific flux adjustment at the grounding line (i.e. not using Eq. 17 nor Eq. 18). The second iteration is computed the same way, except for the fact that the ice shelves are assigned to a very low viscosity so that they cannot exert any back force. The buttressing ratio ϕ_{bf} is then computed as the velocity ratio between these two computed velocities. Once ϕ_{bf} is estimated, we solve the velocity equation again, this time imposing the grounding line flux computation using Eq. 17

or Eq. 18, in order to estimate the velocity used in the mass conservation for this time step. We acknowledge the fact that this approach is computationally expensive but it allows for more accurate estimate for the buttressing role of ice shelves in the model.

2.1.5 Calving

5 Iceberg calving is not modelled explicitly. Instead, we used a simple ice thickness threshold criterion. Because this simple scheme can prevent ice shelf extension, we also maintain downstream ice shelf grid-points neighbouring the last grid-points meeting the criterion. The cut-off threshold may vary in space (e.g. oceanic depth dependency) and time. In the following, we use a constant and homogeneous thickness criterion (set to 250 m, roughly corresponding to the observed present-day Antarctic ice shelves front).

10 2.1.6 Ice temperature calculation

The ice temperature calculation has remained identical to Ritz et al. (2001). ~~Similarly to As in~~ most large-scale ice sheet models (e.g. Winkelmann et al., 2011; Pollard and DeConto, 2012; Pattyn, 2017), the temperature in GRISLI is computed by solving the general advection-diffusion equation of temperature:

$$\begin{aligned}
 \frac{\partial T}{\partial t} = & \underbrace{\frac{1}{\rho c} \frac{\partial}{\partial x} \left(k_i \frac{\partial T}{\partial x} \right) + \frac{1}{\rho c} \frac{\partial}{\partial y} \left(k_i \frac{\partial T}{\partial y} \right)}_{\text{horizontal diffusion}} + \underbrace{\frac{1}{\rho c} \frac{\partial}{\partial z} \left(k_i \frac{\partial T}{\partial z} \right)}_{\text{vertical diffusion}} \\
 & \underbrace{-u_x \frac{\partial T}{\partial x} - u_y \frac{\partial T}{\partial y}}_{\text{horizontal advection}} \quad \underbrace{-u_z \frac{\partial T}{\partial z}}_{\text{vertical advection}} \quad + \underbrace{\frac{Q}{\rho c}}_{\text{heat production}}
 \end{aligned}
 \tag{19}$$

15 with k_i the thermal conductivity of the ice ~~and~~ c the heat capacity ~~is~~ ~~and~~ u_z the vertical velocity, computed as in Ritz et al. (1997) (~~with~~ w_i in Eq. 14 of the original paper).

Horizontal diffusion is assumed to be ~~much smaller than the vertical one and is therefore neglected~~ negligible relative to the vertical diffusion.

The heat production is given by Hutter (1983) (for $i, j = x, y, z$):

$$20 \quad Q = \sum_{i,j} \dot{\epsilon}_{ij} \tau_{ij}
 \tag{20}$$

At the ice sheet surface, due to the absence of an explicit snowpack model, ice temperature is assumed to be equal to the near-surface air annual temperature (but not greater than the melting point). Depending on the surface mass balance parametrisation, the latent heat release due to refreezing is transferred to the first ice layer.

25 A geothermal heat flux ϕ_0 is applied at the base of a 3 km thick bedrock layer with a Neumann boundary condition:

$$\phi_0 = -k_b \frac{\partial T}{\partial z} \Big|_{\text{bedrock}}
 \tag{21}$$

with k_b the bedrock thermal conductivity.

Similarly to what is done for ice temperature, the heat equation is solved in the bedrock with similarly to Eq. 19 but with no advection nor heat production. From the temperature gradient in the bedrock (computed on four vertical levels) we compute a heat flux ϕ'_0 at the ice-bedrock interface. In case of ice dragging-When ice dragging occurs over the bedrock, an additional term due to friction, ϕ_f , is added to ϕ'_0 :

$$\phi_f = |\mathbf{u}_b \tau_b| \quad (22)$$

The ice-bedrock interface heat flux is used differently for cold and temperate based points:

- For cold based points, the heat at the ice-bedrock interface is transferred to the ice via a Neumann boundary condition:

$$k_i \frac{\partial T}{\partial z} |_{ice} = -\phi'_0 - \phi_f \quad (23)$$

with the ice thermal conductivity k_i computed as:

$$k_i = 3.1014 \cdot 10^8 \exp(-0.0057 (T + 273.15)) \quad (24)$$

- For temperate points, a Dirichlet boundary condition is applied as the temperature is kept at the pressure melting point. The excess heat in this case is used to compute basal melting:

$$b_{melt} = \frac{-\phi'_0 - \phi_f - k_i \frac{\partial T}{\partial z} |_{ice}}{L_f \rho} \quad (25)$$

with L_f is the latent heat of fusion.

Basal melting for oceanic points is usually imposed. For specific applications we have different values for deep ocean and continental shelves, or a geographical distribution depending on the oceanic basin.

The viscosity for the velocity grid points is the horizontal average of the viscosity on the centred grid and not the viscosity computed from the horizontal average of the temperature. This is preferable for regions with mixed frozen and temperate basal conditions.

2.2 Additional features

2.2.1 Basal hydrology

~~Since the work of Peyaud (2006), GRISLI accounts for~~ [Peyaud \(2006\) added](#) a simple diffusive basal hydrology scheme [in GRISLI](#). In the following we provide a complete description of the hydrology model because it has only been described in a French Phd dissertation (Peyaud, 2006) but currently lacks a description in an international scientific journal.

- 5 Using a Darcy law, the water produced by melting at the base of the ice sheet is routed outside glaciated areas following the highest gradient in the total water potential.

Such a gradient can be written as in Shreve (1972):

$$\nabla\Phi = \rho_w g \nabla h_w + \rho_w g \nabla B + \rho g \nabla H \quad (26)$$

where h_w is the hydraulic head, B is the bedrock height and H the ice thickness.

- 10 In GRISLI, we assume that the basal water flows within a sub-glacial till following a Darcy-type flow law:

$$\mathbf{Q}_w = -\frac{KD}{\rho_w g} \nabla\Phi \quad (27)$$

where \mathbf{Q}_w is the water flux vector in x and y directions, K is the hydraulic conductivity of the till and D is the water thickness within the till.

- The till is assumed to be present everywhere below the ice sheet with a constant and homogeneous thickness ($h_{till} = 20\text{m}$) and porosity ($\phi_{till} = 0.5$). The water thickness in the till, D , equals the hydraulic head, h_w , only for thicknesses lower than the effective thickness:

$$D = \min(h_w, \phi_{till} h_{till}) \quad (28)$$

The hydraulic conductivity of the till, K , is modulated by the effective pressure to take into account sediment dilatation:

$$K = \begin{cases} K_0 & \text{if } N > N_0 \\ K_0 N_0/N & \text{if } N \leq N_0 \end{cases} \quad (29)$$

- 20 with K_0 the reference conductivity, N the effective pressure and N_0 a constant (10^8Pa). The conductivity K_0 is poorly constrained and strongly depends on the material.

In GRISLI, we assume that the flow of water within the till can be described with a diffusivity equation for the hydraulic head:

$$\frac{\partial h_w}{\partial t} + \nabla \cdot \mathbf{Q}_w = b_{melt} - I_{gr} \quad (30)$$

with I_{gr} being the infiltration rate in the bedrock (kept constant at 1 mm yr^{-1}).

- 25 Using Eq. 26 and Eq. 27, this diffusivity equation can be written as:

$$\frac{\partial h_w}{\partial t} = b_{melt} - I_{gr} + \nabla \cdot \left(KD \nabla \left(B + \frac{\rho}{\rho_w} H \right) \right) + \nabla \cdot (KD \nabla h_w) \quad (31)$$

~~Similarly to what is done~~ [Eq. 31 can be solved using a semi-implicit relaxation method as](#) for the mass conservation equation [a semi-implicit relaxation method is used to solve Eq. 31](#).

From the hydraulic head, h_w , we can compute the water pressure, $p_w = \rho_w g h_w$, and the effective pressure, $N = \rho g H - p_w$.

2.2.2 Isostasy

As in Ritz et al. (2001), GRISLI computes the bedrock response to ice load with an elastic lithosphere - relaxed asthenosphere (ELRA) model (Le Meur and Huybrechts, 1996). This simple model evaluates the bedrock deformation to a local unit mass, scaled to the whole ice sheet. The relaxation time of the asthenosphere is usually set to 3000 years and the deflection of the lithosphere is assumed to follow a zero-order Kelvin function. Such a simple model has been shown to perform well compared to more sophisticated glacio-isostatic models (Le Meur and Huybrechts, 1996).

2.2.3 Passive tracer

GRISLI includes a passive tracer model that allows for the computation of vertical ice stratigraphy, i.e. time and location of ice deposition for the vertical model grid points. The model is the one of Lhomme et al. (2005) re-implemented in GRISLI by Quiquet et al. (2013). We use a semi-Lagrangian scheme following Clark and Mix (2002) in order to avoid the numerical instabilities of Eulerian schemes and information dispersion of Lagrangian schemes (Rybak and Huybrechts, 2003). For each timestep, the back trajectories of each grid points are computed and tri-linearly interpolated onto the model grid. This allows for a continuous information within the ice sheet at a low computational cost. Time and location of ice deposition can be convoluted for example with isotopic composition of precipitation (e.g. $\delta^{18}O$) in order to construct synthetic ice cores comparable with actual ice cores (Lhomme et al., 2005).

2.3 Numerical features

The model uses finite differences computed on a staggered Arakawa C-grid in the horizontal plane (Fig. 2). In the vertical, the model defines σ -reduced coordinates, $\zeta = (S - z)/H$, for 21 evenly spaced vertical layers, with the z vertical axis pointing upward and ζ pointing downward (0 at the surface and 1 at the bottom). The coordinate triplet (i, j, k) (in x , y and ζ direction) is representative of the centre of the grid cell. The horizontal resolution depends on the application, i.e. the extension of the geographical domain and the duration of the simulated period. For century scale applications, the resolution varies from 5 km for Greenland to 15 km for Antarctica (Peano et al., 2017; Ritz et al., 2015). For multi-millennial applications the resolution reduces to 15 km for Greenland and 40 km for the whole Northern Hemisphere and Antarctica.

The mass balance equation is solved as an advection-only equation with an upwind scheme in space and a semi-implicit scheme in time (velocities at the previous time step are used). The numerical resolution is performed with a point-relaxation method with a variable time step. The value of this time step is chosen to ensure that the matrix becomes strongly diagonal dominant to achieve convergence of the point-relaxation method. The criteria is thus a threshold that is inversely proportional to the fastest velocity on the whole grid. Note that this smaller time step is solely used for the mass conservation equation (Eq. 2) and subsequent variables (e.g. surface slopes, SIA velocity) while the rest of the model uses a main time step, typically ranging from 0.5 to 5 years depending on the horizontal resolution.

To solve ~~it~~, the ice shelves/ice streams equation (~~Eq. 14~~) ~~is~~ needs to be linearised. The viscosity is computed using an iterative method starting from the viscosity calculated from strain rates from the previous time step. As this equation is the most expensive part of the model, the iteration mode is not always used depending on the type of experiment (for instance not crucial when the objective is to reach the steady state). In this case the viscosity of the previous time step is used. The linear system is
5 solved with a direct method (Gaussian elimination, sgbv in the Lapack library (www.netlib.org/lapack)).

The resolution of the elliptic system (Eq. 14) is the most expensive part of the model. This is further amplified by the way we prescribe boundary conditions. As in Ritz et al. (2001), the ice shelf region is artificially extended towards the edges of the geographical domain. This artificial extension does not have any consequence on ice shelf velocity since added grid points (that
10 we call “ghost” nodes) are prescribed with a negligible ice viscosity (1500 Pa s). The front is then parallel to either x or y (Ritz et al., 2001) and thus the boundary condition there is easy to implement (see also Fig. S1 in the supplement). The boundary condition at the real ice shelf front is ~~implicitly done by solving~~ (solved implicitly with Eq. 14). However this method increases substantially the size of the linear system solved in (Eq. 14). To ~~go around~~ circumvent this issue, a simple reduction method is implemented. (Eq. 14) can be written as $\tilde{A} \tilde{\mathbf{u}} = \tilde{B}$ where $\tilde{\mathbf{u}}$ is a vector alternating u_x and u_y components for all the velocity
15 grid points, \tilde{A} is a band matrix (very sparse) and \tilde{B} is a vector corresponding to the right hand terms in Eq. 14. Every line of \tilde{A} and \tilde{B} are scaled so that the diagonal terms of \tilde{A} are equal to 1. If, for a given velocity node, all the non diagonal terms of the column are very small compared to 1, this means that this node is actually in practice not used by any other velocity node and this line of the matrix can be removed. The threshold to neglect nodes is related to the value of the integrated viscosity of “ghost” nodes. In practice, given its size, the matrix \tilde{A} is not ~~actually fully constructed, only the non-zero sub/sur-diagonals~~
20 are fully populated, being a band matrix with a bandwidth of one. An illustration of the matrix is shown in Fig. S2 in the supplement.

For the temperature equation (Eq. 19), we solved a 1D advection-diffusion equation for each model grid point. The resolution is performed with an upwind semi-implicit scheme (vertical velocity and heat production at the previous time step is
25 used). The ice thermal conductivity is computed as the geometric mean of the two neighbouring conductivities (Patankar, 1980). Because the horizontal diffusion is neglected, the only horizontal terms concern horizontal advection and are computed with an upwind explicit scheme. The heat production is computed at the velocity (staggered) grid points and is then summed up to the temperature (centred) grid points.

30 The model has been recently partially parallelised with OpenMP (www.openmp.org), which considerably shortens the length of the simulations (gain of 40% for the Antarctic at 40 km on four threads of an Intel® Xeon® CPU @ 3.47 GHz).

3 Calibration for the Antarctic ice sheet

3.1 Methods

In the following, we present a simple calibration methodology for the Antarctic ice sheet based on a large ensemble of model simulations. Given its degree of complexity, GRISLI is mostly designed for multi-millennial integrations. Due to long-term
5 diffusive response to SMB and temperature changes, an accurate methodology to select unknown parameters of the model would be to run long transient simulations with a climate forcing as close as possible from past climate states, ideally with a synchronous coupling between the ice sheet and the atmosphere. However, climate models generally fail at reproducing the regional climate changes during the last glacial-interglacial cycle as recorded by proxy data (Braconnot et al., 2012). Further-
10 more, the phase III of the Paleoclimate Modelling Intercomparison Project (PMIP3) has highlighted the large disagreement between participating climate models in simulating the Last Glacial Maximum (LGM) in the vicinity of northern Hemisphere ice sheets (e.g. Harrison et al., 2014). Given these uncertainties amongst climate models and the large sensitivity of the ice sheet model to climate forcing fields (e.g. Charbit et al., 2007; Quiquet et al., 2012; Yan et al., 2013), it is difficult to calibrate the mechanical parameters independently from that of the SMB, in particular for northern Hemisphere ice sheets.

15 For these reasons, here we suggest a simple calibration methodology for the Antarctic ice sheet in which the model is run for 100 kyrs under a perpetual modern climate forcing in order to reach an ice sheet equilibrium. In the following, we use the 27 km-grid atmospheric outputs, namely annual mean temperature and SMB, from the regional climate model RACMO2.3 (Van Wessem et al., 2014), averaged over the 1976-2016 time span. The basal melting rates under ice shelves are prescribed for the 18 sectors of the Antarctic ice sheet as defined in ISMIP-Antarctica project (Nowicki et al., 2016) and are shown in
20 Fig. 3. Their values are based on the sectoral average of sub-shelf melt rates that ensured stable ice shelves (minimal Eulerian ice thickness derivative) in the recent intercomparison exercise InitMIP-Antarctica (Nowicki et al., 2016), with slight modifications due to change in resolution. They are in line with observations-based estimates (Rignot et al., 2013). We do not apply any correction related to geometry changes to the climatic forcings during the calibration.

25 We choose to restrict this study to a coarse horizontal resolution, namely 40 km, as it allows for large ensembles of multi-millennial simulations. Whilst 6.7 hours on one thread of an Intel® Xeon® CPU@3.47 GHz (4 hours on four threads) are needed to perform 100 000 years of simulation over Antarctica on a 40-km grid (19 881 horizontal grid points), this time goes up significantly on a 16-km grid (145 161 points) for which we need 25 hours to perform 2000 years (17 hours on four threads). In addition, the 40 km resolution corresponds to the one used in the coupled version within the *i*LOVECLIM earth
30 system model of intermediate complexity (Roche et al., 2014). Whilst with such a resolution we do not expect to have an accurate representation of the ice sheet fine scale structures such as ice streams, we expect to reproduce the large scale behaviour of ice flow.

From our experience with GRISLI, we identified four unknown independent parameters that have a crucial role for ice dynamics ~~and that are independent from each other~~:

- The SIA flow enhancement factor S_f of the Glen flow law (Eq. 10). This coefficient is expected to have a large influence on shear-stress driven velocities.
- 5 – The basal drag coefficient C_f in Eq. 16. This coefficient is used to modulate the basal drag coefficient for temperate-based grid points where sliding occurs.
- The till conductivity K_0 . This parameter changes the efficiency of basal water routing (Eq. 29 and Eq. 31) and thus, basal effective pressure N . As such, this parameter is also influencing the basal drag coefficient β for temperate-based regions (Eq. 16).
- 10 – An ice shelf basal melting rate coefficient ϕ_{shelf} . For a specific Antarctic ice shelf sector i :

$$BMB^i = \phi_{shelf} BMB_0^i \quad (32)$$

with BMB_0^i the sub-shelf basal melting rate reference values shown in Fig. 3.

The parametric ensemble is designed with a Latin Hypercube Sampling (LHS) methodology. The LHS is used here because it has better space-filling quality than a standard Monte-Carlo sampling which might not explore sufficiently the tails of parameter distributions. This methodology has been used for calibration purposes in the ice sheet modelling community (e.g. Stone et al., 2010; Applegate et al., 2012). The size of the LHS consists of ~~300 model realisations. We perform two times the 300 member ensemble~~ 600 model realisations: half of it with the flux at the grounding line of Schoof (2007) (hereafter AN40S) and the rest with Tsai et al. (2015) (hereafter AN40T) The range of explored parameters are listed in Tab. 2. We assume a uniform statistical distribution within this range.

20

The initial ice sheet geometry, bedrock and ice thickness, is taken from the Bedmap2 dataset (~~Fretwell et al., 2013~~) (Fretwell et al., 2013, Fig. 1) via a spatial bi-linear interpolation to generate this data on the 40-km grid. The geothermal heat flux is from Shapiro and Ritzwoller (2004). Sensitivity to uncertainties in the forcing data are not explored in the ensemble as we aim at quantifying the model sensitivity to parameter choice even though we acknowledge for the fact that these could be the source of important model error (e.g. Stone et al., 2010; Pollard and DeConto, 2012).

25

In the following, individual member performance is assessed with the root mean square error (RMSE) computed from simulated and observed ice thickness (Fretwell et al., 2013). This metric puts a strong constraints on ice sheet geometry and avoids potential compensatory biases that could appear when using total ice volume. Observed ice velocities are not used as a metric because of high spatial variability that results from small-scale bed properties. With a coarse 40-km resolution, the model is intrinsically unable to reproduce such variability but is expected to reproduce the large scale pattern of ice thickness. However, at the end of the long 100-kyrs simulation, the velocities in the model are the balance velocities corresponding to the simulated

30

topography. Thus, the minimisation of the RMSE in ice thickness should also reduce the error in velocities with respect to observations at the global scale.

3.2 Calibration results

Figure 5 presents the Antarctic ice sheet volumes at the end of the 100 kyr simulations for each ensemble ~~members~~ member as a function of parameter values using the flux at the grounding line computed from Schoof (2007) (AN40S). We can see that there is a strong positive (respectively negative) correlation of ice volume with the basal drag coefficient (resp. enhancement factor). There is also a weak negative correlation for the sub-shelf basal melt coefficient, and a weak positive correlation with the till conductivity. Since the global volume is an integrated metric that does not account for potential systematic compensation, in Fig. 5 we also show the root mean square error (RMSE) in ice thickness for each ensemble members with respect to observations (Fretwell et al., 2013). Amongst the 300 model realisations, 120 members ~~present~~ have a RMSE lower than 350 m. ~~They are widespread across the range of parameter values~~ These members are widely distributed within the member spectrum. The lowest RMSE is 294 m. The 12 best ensemble members are outlined in red in Fig. 5 and ~~yield~~ have a RMSE not higher than 304.

The general model response ~~amongst the ensemble members~~ is not fundamentally different when the flux at the grounding line is computed from Tsai et al. (2015) (Fig. 6). However, the grounding line position is much more unstable with a greater number of members showing lower ice sheet volume. Only 62 model realisations ~~present~~ have a RMSE lower than 350 m (lowest RMSE at 295 m and 12 best not higher than 304), compared to 120 within the AN40S ensemble members. This difference in ~~term of~~ grounding line stability for the two flux formulations has already been highlighted by Pattyn (2017). Since basal drag vanishes at the grounding line in the Tsai et al. (2015) formulation, its coefficient has a smaller impact than in Schoof (2007), amplifying the role of the ice flow enhancement factor. As a consequence, for the AN40T ensemble, the enhancement factor requires values between 1.5 and 3 in order to reach a good agreement with observed ice thickness, whilst values within 1.5 to 4 are acceptable for AN40S.

In Fig. 7 (respectively Fig. 8) we show the ice thickness difference from the observations for the 12 ensemble members showing the lowest RMSE within the AN40S (resp. AN40T) model realisations. The differences are generally below 500 m even if persisting model biases are present across the ensemble members and model formulations. On the one hand, ice thickness in large parts of the East Antarctic ice sheet is systematically underestimated. On the other hand, the West Antarctic ice sheet shows more contrasted responses. Whilst for some ensemble members, the Ross embayment upstream region can be well represented (e.g. AN40S004 or AN40T065), the region feeding the Filchner-Ronne ice shelf show a quasi-systematic ice thickness underestimation. These model deficiencies can be attributed to our coarse model resolution, providing a poor representation of the complex bedrock structure in the Filchner-Ronne area. The model differences from the observations are very similar to results from the PISM-PIK model shown in Martin et al. (2011) in term of amplitude but also in term of structure. They are also generally similar to Pollard and DeConto (2009). Consistent model biases amongst these models, which use different input data, suggest a common source of error related to the coarse model resolution (20 to 40 km) or uncertainties in

the bedrock dataset, particularly large in East Antarctica (Fretwell et al., 2013)(Fig. 4).

Figure 9 (resp. Fig. 10) presents the root mean square error with respect to observations of the ensemble members in the two-dimensional parametric space within AN40S (resp. AN40T). The 12 best ensemble members are outlined in red. In most cases there is no clear relationship. However, there is a relationship emerging with a large basal drag coefficient being compensated by a large enhancement factor when using the Schoof (2007) (Fig. 9). When using the Tsai et al. (2015) formulation (Fig. 10), this relationship disappears as the enhancement factor is mostly driving the model response. A few model parameter combinations (30 out of 300) are able to provide a good representation of the present day Antarctic ice sheet, i.e. RMSE lower than 350 m, independently from the grounding flux computation used (not shown).

10

Although our quality metric is based on the ice sheet thickness we show in Fig 11 and in Fig. 12 the capability of the model to reproduce observed ice sheet velocity (Mouginot et al., 2017) for the best ensemble members with the two formulations of the flux at the grounding line (parameter values shown in Tab. 3). The model reproduces the general distribution of the velocity although it underestimates ice flow for the very fast grid points (velocity larger than 100 m yr^{-1} , generally ice shelves). The model has also difficulties ~~to reproduce in reproducing~~ well-defined ice streams such as the Amery or Filchner ice shelves tributaries. ~~These Such difficulties~~ could be due to the coarse resolution used but also to the simple scheme used to estimate basal drag.

15

4 Antarctic ice sheet changes for the last 400 kyrs

4.1 Methods

20 The main objective of this section is to show the ability of the model to reproduce large ice sheet geometry changes in response to Quaternary climate change. As a consequence of our limited knowledge of past climatic conditions in the Antarctic ice sheet region over glacial-interglacial cycles, we use here an idealised reconstruction of SMB, near surface air temperature and oceanic basal melting rates based on a limited number of proxy records. Our approach is somewhat similar to previous works (e.g. Ritz et al., 2001; Huybrechts, 2002; Pollard and DeConto, 2009; Greve et al., 2011; Golledge et al., 2014).

25

The near-surface air temperature, used in the model as a surface boundary condition for the advection-diffusion temperature equation, is assumed to follow the EPICA-DOME C deuterium record δD :

$$T_{palaeo} = T_0 + (1/\alpha^i) \delta D \quad (33)$$

with T_0 the annual mean near-surface air temperature from RACMO2.3 used for the present-day calibration. The isotopic slope for temperature, α^i , is set to $0.18 \text{ ‰}^\circ\text{C}^{-1}$ as in (Jouzel et al., 2007).

30

We also account for the additional temperature perturbation due to topography changes using a fixed and homogeneous lapse

rate λ :

$$T_{palaeo}^* = T_{palaeo} + \lambda(S - S_0) \quad (34)$$

with $S - S_0$ the local topography change from Bedmap2. In the following λ is set to $-8 \text{ }^\circ\text{C km}^{-1}$.

- 5 For a given near-surface air temperature change T_{palaeo}^* relative to present-day T_0 , we modify the present-day SMB field, SMB_0 :

$$SMB = SMB_0 \exp(-\gamma(T_0 - T_{palaeo}^*)) \quad (35)$$

with the precipitation ratio to temperature change γ set to $0.07 \text{ }^\circ\text{C}^{-1}$. The use of an exponential form in Eq. 35 is motivated by the Clausius-Clapeyron saturation vapour pressure for an ideal gas. Such a simple expression implies that SMB is driven
10 only by accumulation, an assumption justified by the very little surface ablation experienced by the Antarctic ice sheet under present-day climatic conditions. However, we may underestimate the surface melt for warmer past interglacial periods.

In order to account for changes in basal melting rates below ice shelves, there is the need to define a continuous proxy covering several glacial-interglacial cycles for past sub-surface oceanic conditions around Antarctica. To this end, and due to the lack of
15 such a record in the Southern ocean, we used the temperature derived from a benthic foraminifer $\delta^{18}\text{O}$ record from the North Atlantic. This temperature signal is considered to depict the North Atlantic Deep Water (NADW) temperature (Waelbroeck et al., 2002). Here, we assume that changes in NADW temperature drive changes in the temperature of waters upwelled in the Southern Ocean. This upward flow separates into surface equatorward and poleward flows, and thus influences surface and sub-surface temperature around coastal Antarctica (e.g. Ferrari et al., 2014). The basal melting rate below a specific ice shelf
20 sector i , BMB_{palaeo}^i for past periods is computed from its present-day value, BMB_0^i , corrected to account for past oceanic conditions:

$$BMB_{palaeo}^i = \max(BMB_0^i (1 + \delta^{oc}), 0.01 \text{ m yr}^{-1}) \quad (36)$$

using the palaeo-oceanic index δ^{oc} defined as:

$$\delta^{oc} = \alpha^{oc} \Delta T_{NA} / T_{NA0} \quad (37)$$

- 25 with T_{NA0} the pre-industrial temperature deduced from North Atlantic benthic foraminifera (Waelbroeck et al., 2002) and ΔT_{NA} the deviation from this temperature in the past. α^{oc} is a conversion coefficient, set to 1 in the following.

The atmospheric and oceanic indexes, $T_{palaeo} - T_0$ and δ^{oc} , used to drive the model for the last 400 kyr are presented in Fig. 13. In addition to these climatic perturbations we also use the eustatic sea level reconstruction of Waelbroeck et al. (2002) to account for sea level variations over glacial-interglacial cycles.

30

In the following, we discuss the model behaviour as the result in response to the 400 kyr forcing. We performed simulations using the 12 parameter combinations from Sec. 3 that yield have the lowest RMSE for the two groups AN40S and AN40T,

differing by the treatment of the flux at the grounding line. We used the 100-kyr integration under perpetual modern climate in Sec. 3 as a spin-up for the transient simulations. We ~~artificially expand the simulations for further compute~~ 10 kyr into the future with no climatic perturbation from the reference climate used in Sec. 3 in order to discuss the stability of the simulated present-day ice sheet state.

5 4.2 Transient simulation results

In Fig. 14 the simulated ice sheet volume is shown over the last 400 kyr. Across this time scale, a large glacial-interglacial volume variation is observed, in particular for the last two cycles where it reaches up to about 10 millions of km³. In our simulations, the Antarctic ice sheet volume increase at the last glacial maximum (LGM, 21 kaBP) relative to pre-industrial corresponds to about -10 to -20 m of global eustatic sea level drop depending on the simulations. These numbers mostly fall in the range of previous ice sheet model reconstructions (e.g. Huybrechts, 2002; Philippon et al., 2006; Pollard and DeConto, 2009), Antarctic contributions inferred as the difference from far-field and Northern Hemisphere near-field estimates (Peltier, 2004) or near-field estimates (Ivins and James, 2005; Argus et al., 2014; Whitehouse et al., 2012; Briggs et al., 2014). Our reconstructions are nonetheless at the higher ~~hand-end~~ of recent studies. This could be related to the fact that we do not account for different geologic bed types between today ice-free (with extensive amount of deformable till) and glaciated (mostly hard bed) continental shelf. To account for this, some authors have chosen a two-value basal drag for these different regions (e.g. Pollard and DeConto, 2012). Because of the large uncertainties related to the bed properties we have decided to ignore these differences, keeping in mind that this can bias our results towards thicker ice sheet when the ice expands over the continental shelf. In our simulations, the last interglacial (120 kaBP) ice volume ~~does not present has no~~ substantial changes relative to the present-day ice volume, ~~as the Antarctic ice sheet is contributing to contributing~~ less than 6 centimetres to the global eustatic sea level rise in the simulations with the lowest RMSE at 0k. This is well below recent estimates, ranging from 3 to 7 m, inferred from the limited contribution of Greenland to the last interglacial highstand (Dutton et al., 2015). Our crude representation of the last interglacial climate in which no surface melt is possible may be the cause for such a discrepancy. In addition, our proxy-based basal melting rate does not allow for above than present basal melting rates during the last interglacial.

The uncertainty related to the choice of the internal parameters within our subset leads generally to up to 3 10⁶ km³ differences in our framework but ~~de does~~ not change the model response to the forcings. In turn, the choice of either Schoof (2007) (AN40S) or Tsai et al. (2015) (AN40T) to compute the flux at the grounding line leads to important differences amongst temporal model responses. AN40T systematically start to retreat before AN40S. It also produces a larger glacial to interglacial volume ~~changes change~~. This confirms the fact that the Tsai et al. (2015) formulation leads to ~~more sensitive grounding line~~ higher grounding line migration variability as already highlighted in Sec. 3 and by other authors (Pattyn, 2017). The additional 10 kyr into the future with no climatic perturbation shows that the AN40S ensemble members do not produce an ice sheet at equilibrium at 0 ka BP. This means that, in our model, the Schoof (2007) formulation produces unrealistically too slow post-LGM retreat which induces a model drift persisting till 10 kyr in the future. Conversely, the Tsai et al. (2015) formulation leads to more rapid retreat rates which provides a stabilisation of the ice sheet during the Holocene.

Simulated ice sheet surface elevations at selected snapshots for the two ensemble members with the lowest RMSE at 0 ka BP after the transient simulations are presented in Fig. 15. Ice sheet geometry during the last interglacial resembles the present-day one. This is particularly true for the eastern part whilst the West Antarctic ice sheet is only slightly thinner. At the last glacial maximum, the grounding line advances towards the edge of the continental shelf, in agreement with geological reconstructions (Bentley et al., 2014). The choice of the flux at the grounding line formulation has an impact on the maximum ice sheet extent, with a less extended ice sheet using the Schoof (2007) formulation. As for the last interglacial, the eastern part of the ice sheet presents only small variations in surface elevation compared to the present-day geometry. There is no decrease in surface elevation at the last glacial maximum due to reduction in precipitation at this time since the larger extent and the colder climate tend to reduce the ice flow. The largest topography changes are occurring in the Weddel and Ross sea. The West Antarctic ice sheet is thus particularly dynamic during glacial-interglacial cycles.

The RMSE computed at 0 ka BP for the 24 members used for the transient simulations ranges from 372 to 467 m within AN40S and 326 to 376 m within AN40T. These numbers are only slightly greater than the ones obtained using a constant forcing (Sec. 3). The differences in ice sheet thickness between the transient simulations at 0 ka BP and the observations for the two ensemble members with the lowest RMSE (AN40S097 and AN40T059) are shown in Fig. 16. The pattern is similar to the one obtained during the calibration step (Sec. 3) with some notable differences. On the one hand, the East Antarctic ice sheet thickness underestimation is partly corrected when performing a transient simulation. This could be the result of a better representation of the temperature vertical profile in this case. On the other hand, whilst in other regions the model biases remain generally the same between an equilibrium and a transient simulation, important model biases appear at the margins of the ice sheet when using the transient simulations. This is particularly visible when using Schoof (2007) for which the retreat rate during the deglaciation is underestimated. Part of the misrepresentation of present-day margins could also be due to the over-simplified climatic perturbation used for the transient simulations.

5 Discussion and outlook

We have presented results from the updated version of the GRISLI model. Whilst the model is able to reproduce present-day Greenland (Le clec'h et al., 2017) and Antarctic (Ritz et al., 2015) ice sheets when using an inverse method to estimate the basal drag, our simulations with an interactive basal drag coefficients computed from the effective pressure show some important disagreements relative to observations. In particular there are some persisting model biases in ice thickness. In East Antarctica, the ice thickness is underestimated towards the pole and the Transantarctic mountains while it is overestimated towards the margins, from Queen Maud land to Wilkes land. In West Antarctica, there is an underestimation of ice thickness in the Ronnie-Filchner basin and an overestimation in the Ross basin. These model biases are also present in models of similar complexity when using an interactive basal drag computation (Pollard and DeConto, 2009; Martin et al., 2011). This data-model mismatch is mostly due to a poor representation of the bedrock-ice interface. In particular, the coarse resolution does not allow for the

consideration of fine scale troughs and pinning points. The persisting model biases can ~~be also~~ also be the consequence of our simplified basal drag computation that does not take into account bedrock physical properties (e.g. sediments).

We used a basal drag coefficient computed from an internal model parameter, namely the basal effective pressure. For long-term multi-millennial integrations, this is preferred to deducing the basal drag coefficient ~~deduced~~ from inversion using present-day geometry since it is fully consistent with the model physics and, in principle, remains valid for large ice sheet geometry change. However, by design, the fit with observations is systematically poorer compared to model results that make use of an inverse basal drag coefficient. A step forward would be to use the basal drag computed from inversion in order to deduce a formulation based solely on internal parameters. Amongst these parameters, along with the basal effective pressure, the large scale bedrock curvature and/or sub-grid roughness could be used, ~~similarly to as in~~ Briggs et al. (2013). However, some key basal features, such as the geologic bed type and the deformable till distribution, remain today largely unknown below present-day ice sheets and will contribute to large uncertainties in the basal drag formulation.

Although widely used for ice sheet model spin-up or calibration, long-term integrations under present-day forcing induce a warm bias in the vertical temperature profile because they discard the diffusion of glacial-interglacial changes in surface temperature. Calibrated parameters obtained with such a methodology tend to compensate for the under-estimated viscosity and are in theory not suitable for palaeo-reconstructions. Whilst a parameter calibration based on glacial-interglacial simulations is ideally preferred, the determination of a realistic climate forcing is a considerable challenge given the many degrees of freedom. Here, we presented a very simplified climate reconstructions for the last 400 kyr based on a minimal parameter set (proxy for atmospheric temperatures and oceanic conditions) in order to illustrate the model possible behaviour for long-term integrations. Using the parameters calibrated under perpetual modern climate, the model is nonetheless able to reproduce ice geometry changes compatible with palaeo-constraints. Further work will consist in the determination of more realistic climate reconstruction using general circulation model snapshots. We also aim at expanding the work of Roche et al. (2014) and couple the Antarctic geometry of GRISLI version 2.0 with the *i*LOVECLIM model.

The implementation of an explicit flux computation at the grounding line following Schoof (2007) and Tsai et al. (2015) lead to a more dynamic grounding line position compared to previous version of the model. As such, GRISLI version 2.0 is now more sensitive to both sub-shelf melt rate changes and also sea level variations. However, the current version of the model only considers an eustatic sea level perturbation with a regional bedrock adjustment. The explicit computation of local relative sea level could potentially have an important impact on grounding line migration for glacial-interglacial cycles (e.g. Gomez et al., 2013).

In the current version of the model, some important processes are still largely simplified. In particular, further developments will consist in the implementation of a new basal hydrology model relying on an explicit routing scheme (e.g. Kavanagh and Tarasov, 2017) ~~would allow to avoid~~ avoiding relaxed numerical solutions based on effective pressure. This could introduce fast basal water changes that are currently ignored and, ultimately, could yield ice streams abrupt speed-up or slow-down. Also,

calving processes are suspected to be a major driver for ice sheet evolution due to the importance of buttressing on inland ice dynamics (e.g. Pollard et al., 2015). GRISLI version 2.0 includes a very simplified calving representation that might prevent to assess the role of this process for multi-millennial ice dynamics. The inclusion of a physically based calving scheme (e.g. Christmann et al., 2016) would be a significant model improvement for future model revisions.

5 6 Conclusions

We have presented the GRISLI (version 2.0) model along with the significant improvements from the previous version of Ritz et al. (2001). Such improvements include an explicit flux computation at the grounding line, an interactive basal hydrology module and a semi-lagrangian tracking particle scheme. Thanks to its low computational cost, the model is suitable for long-term multi-millennial integrations. We performed a large ensemble of simulations of the Antarctic ice sheet forced by present-day climate conditions to calibrate the crucial unknown parameters. We have shown that the model is able to reproduce reasonably well the present-day geometry although the grounding line position in the model is much more unstable when we use Tsai et al. (2015) formulation of the flux at the grounding line instead of Schoof (2007). The model mismatch with respect to observed ice thickness shows some systematic biases (e.g. the East Antarctic ice sheet is too thick in the vicinity of the Transantarctic mountains and too thin elsewhere), that are similar to models of comparable complexity. We used the best ensemble members to simulate the Antarctic evolution throughout the last 400 kyr using an idealised climatic perturbation of present-day conditions. With this simple framework we reproduced the expected ice sheet geometry changes over glacial-interglacial cycles. A significant volume increase is simulated during glacial periods with a grounding line advance towards the edge of the continental shelf. The retreat during terminations is gradual when using our forcing scenario and is able to produce a final present-day ice volume and extent similar to observations. The Tsai et al. (2015) formulation produces a faster ice sheet retreat and yields an ice sheet near equilibrium during the Holocene contrary to Schoof (2007) for which the model is still drifting at +10 kyr into the future. This suggests that, in our model and under the climate forcing scenario we use, the Tsai et al. (2015) formulation produces a more realistic grounding line retreat rate.

7 Code availability

The developments on the GRISLI source code are hosted at <https://forge.ipsl.jussieu.fr/grisli>, but are not publicly available due to copyright restrictions. Access can be granted on demand by request to Christophe Dumas (christophe.dumas@lsce.ipsl.fr), Aurélien Quiquet (aurelien.quiquet@lsce.ipsl.fr) or Catherine Ritz (catherine.ritz@univ-grenoble-alpes.fr) to those who conduct research in collaboration with the GRISLI users group. For this work we used the model at revision 188.

Acknowledgements. We thank Michiel van den Broeke (IMAU, Utrecht University) for providing the RACMO2.3 model outputs. We also warmly thank Claire Waelbroeck for fruitful discussions on the construction of the index for sub-shelf melting rates. This is a contribution

to ERC project ACCLIMATE; the research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement 339108.

References

- Abe-Ouchi, A., Saito, F., Kawamura, K., Raymo, M. E., Okuno, J., Takahashi, K., and Blatter, H.: Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume, *Nature*, 500, 190–193, doi:10.1038/nature12374, 2013.
- Alvarez-Solas, J., Charbit, S., Ritz, C., Paillard, D., Ramstein, G., and Dumas, C.: Links between ocean temperature and iceberg discharge during Heinrich events, *Nature Geosci*, 3, 122–126, doi:10.1038/ngeo752, 2010.
- Alvarez-Solas, J., Robinson, A., Montoya, M., and Ritz, C.: Iceberg discharges of the last glacial period driven by oceanic circulation changes, *Proceedings of the National Academy of Sciences*, 110, 16 350–16 354, doi:10.1073/pnas.1306622110, 2013.
- Applegate, P. J., Kirchner, N., Stone, E. J., Keller, K., and Greve, R.: An assessment of key model parametric uncertainties in projections of Greenland Ice Sheet behavior, *The Cryosphere*, 6, 589–606, doi:10.5194/tc-6-589-2012, 2012.
- Argus, D. F., Peltier, W. R., Drummond, R., and Moore, A. W.: The Antarctica component of postglacial rebound model ICE-6G_C (VM5a) based on GPS positioning, exposure age dating of ice thicknesses, and relative sea level histories, *Geophysical Journal International*, 198, 537–563, doi:10.1093/gji/ggu140, 2014.
- Benn, D. I., Le Hir, G., Bao, H., Donnadieu, Y., Dumas, C., Fleming, E. J., Hambrey, M. J., McMillan, E. A., Petronis, M. S., Ramstein, G., Stevenson, C. T. E., Wynn, P. M., and Fairchild, I. J.: Orbitally forced ice sheet fluctuations during the Marinoan Snowball Earth glaciation, *Nature Geoscience*, 8, 704–707, doi:10.1038/ngeo2502, 2015.
- Bentley, M. J., Ó Cofaigh, C., Anderson, J. B., Conway, H., Davies, B., Graham, A. G. C., Hillenbrand, C.-D., Hodgson, D. A., Jamieson, S. S. R., Larter, R. D., Mackintosh, A., Smith, J. A., Verleyen, E., Ackert, R. P., Bart, P. J., Berg, S., Brunstein, D., Canals, M., Colhoun, E. A., Crosta, X., Dickens, W. A., Domack, E., Dowdeswell, J. A., Dunbar, R., Ehrmann, W., Evans, J., Favier, V., Fink, D., Fogwill, C. J., Glasser, N. F., Gohl, K., Golledge, N. R., Goodwin, I., Gore, D. B., Greenwood, S. L., Hall, B. L., Hall, K., Hedding, D. W., Hein, A. S., Hocking, E. P., Jakobsson, M., Johnson, J. S., Jomelli, V., Jones, R. S., Klages, J. P., Kristoffersen, Y., Kuhn, G., Leventer, A., Licht, K., Lilly, K., Lindow, J., Livingstone, S. J., Massé, G., McGlone, M. S., McKay, R. M., Melles, M., Miura, H., Mulvaney, R., Nel, W., Nitsche, F. O., O’Brien, P. E., Post, A. L., Roberts, S. J., Saunders, K. M., Selkirk, P. M., Simms, A. R., Spiegel, C., Stollendorf, T. D., Sugden, D. E., van der Putten, N., van Ommen, T., Verfaillie, D., Vyverman, W., Wagner, B., White, D. A., Witus, A. E., and Zwart, D.: A community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last Glacial Maximum, *Quaternary Science Reviews*, 100, 1–9, doi:10.1016/j.quascirev.2014.06.025, 2014.
- Bindschadler, R.: The Importance of Pressurized Subglacial Water in Separation and Sliding at the Glacier Bed, *Journal of Glaciology*, 29, 3–19, doi:10.3189/S0022143000005104, 1983.
- Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-Ouchi, A., Otto-Bliesner, B., and Zhao, Y.: Evaluation of climate models using palaeoclimatic data, *Nature Climate Change*, 2, 417–424, doi:10.1038/nclimate1456, 2012.
- Briggs, R., Pollard, D., and Tarasov, L.: A glacial systems model configured for large ensemble analysis of Antarctic deglaciation, *The Cryosphere*, 7, 1949–1970, doi:10.5194/tc-7-1949-2013, 2013.
- Briggs, R. D., Pollard, D., and Tarasov, L.: A data-constrained large ensemble analysis of Antarctic evolution since the Eemian, *Quaternary Science Reviews*, 103, 91–115, doi:10.1016/j.quascirev.2014.09.003, 2014.
- Bueler, E. and Brown, J.: Shallow shelf approximation as a “sliding law” in a thermomechanically coupled ice sheet model, *Journal of Geophysical Research: Earth Surface*, 114, F03 008, doi:10.1029/2008JF001179, 2009.

- Calov, R., Greve, R., Abe-Ouchi, A., Bueller, E., Huybrechts, P., Johnson, J. V., Pattyn, F., Pollard, D., Ritz, C., Saito, F., and Tarasov, L.: Results from the Ice-Sheet Model Intercomparison Project–Heinrich Event Intercomparison (ISMIP HEINO), *Journal of Glaciology*, 56, 371–383, doi:10.3189/002214310792447789, 2010.
- Charbit, S., Ritz, C., Philippon, G., Peyaud, V., and Kageyama, M.: Numerical reconstructions of the Northern Hemisphere ice sheets through the last glacial-interglacial cycle, *Climate of the Past*, 3, 15–37, doi:10.5194/cp-3-15-2007, 2007.
- Christmann, J., Plate, C., Müller, R., and Humbert, A.: Viscous and viscoelastic stress states at the calving front of Antarctic ice shelves, *Annals of Glaciology*, 57, 10–18, doi:10.1017/aog.2016.18, 2016.
- Clark, P. U. and Mix, A. C.: Ice sheets and sea level of the Last Glacial Maximum, *Quaternary Science Reviews*, 21, 1–7, doi:10.1016/S0277-3791(01)00118-4, 2002.
- 10 Colleoni, F., Kirchner, N., Niessen, F., Quiquet, A., and Liakka, J.: An East Siberian ice shelf during the Late Pleistocene glaciations: Numerical reconstructions, *Quaternary Science Reviews*, 147, 148–163, doi:10.1016/j.quascirev.2015.12.023, 2016.
- Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson, G. M., Okuno, J., and Yokoyama, Y.: Ice-sheet collapse and sea-level rise at the Bølling warming 14,600 years ago, *Nature*, 483, 559, doi:10.1038/nature10902, 2012.
- Donnadieu, Y., Dromart, G., Goddérès, Y., Pucéat, E., Brigaud, B., Dera, G., Dumas, C., and Olivier, N.: A mechanism for brief glacial episodes in the Mesozoic greenhouse, *Paleoceanography*, 26, PA3212, doi:10.1029/2010PA002100, 2011.
- 15 Dumas, C.: Modélisation de l'évolution de l'Antarctique depuis le dernier cycle glaciaire-interglaciaire jusqu'au futur : importance relative des différents processus physiques et rôle des données d'entrée, Ph.D. thesis, Université Joseph-Fourier - Grenoble I, 2002.
- Durand, G., Gagliardini, O., de Fleurian, B., Zwinger, T., and Le Meur, E.: Marine ice sheet dynamics: Hysteresis and neutral equilibrium, *Journal of Geophysical Research: Earth Surface*, 114, F03 009, doi:10.1029/2008JF001170, 2009.
- 20 Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., Horton, B. P., Rahmstorf, S., and Raymo, M. E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, aaa4019, doi:10.1126/science.aaa4019, 2015.
- Duval, P. and Lliboutry, L.: Superplasticity owing to grain growth in polar ices, *Journal of Glaciology*, 31, 60–62, 1985.
- Edwards, T. L., Fettweis, X., Gagliardini, O., Gillet-Chaulet, F., Goelzer, H., Gregory, J. M., Hoffman, M., Huybrechts, P., Payne, A. J., Perego, M., Price, S., Quiquet, A., and Ritz, C.: Effect of uncertainty in surface mass balance–elevation feedback on projections of the future sea level contribution of the Greenland ice sheet, *The Cryosphere*, 8, 195–208, doi:10.5194/tc-8-195-2014, 2014.
- 25 Fairbanks, R. G.: A 17,000-year glacio-eustatic sea level record: influence of glacial melting rates on the Younger Dryas event and deep-ocean circulation, *Nature*, 342, 637, doi:10.1038/342637a0, 1989.
- Favier, L., Durand, G., Cornford, S. L., Gudmundsson, G. H., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A. J., and Brocq, A. M. L.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, *Nature Climate Change*, 4, 117, doi:10.1038/nclimate2094, 2014.
- 30 Ferrari, R., Jansen, M. F., Adkins, J. F., Burke, A., Stewart, A. L., and Thompson, A. F.: Antarctic sea ice control on ocean circulation in present and glacial climates, *Proceedings of the National Academy of Sciences*, 111, 8753–8758, doi:10.1073/pnas.1323922111, 2014.
- Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langlely, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti,

- A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375–393, doi:10.5194/tc-7-375-2013, 2013.
- Fürst, J.: Dynamic response of the Greenland ice sheet to future climatic warming, Ph.D. thesis, Vrije Universiteit Brussel, Brussel, 2013.
- Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, doi:10.5194/tc-6-1561-2012, 5 2012.
- Goelzer, H., Nowicki, S., Edwards, T., Beckley, M., Abe-Ouchi, A., Aschwanden, A., Calov, R., Gagliardini, O., Gillet-Chaulet, F., Gollledge, N. R., Gregory, J., Greve, R., Humbert, A., Huybrechts, P., Kennedy, J. H., Larour, E., Lipscomb, W. H., Le clec’h, S., Lee, V., Morlighem, M., Pattyn, F., Payne, A. J., Rodehacke, C., Rückamp, M., Saito, F., Schlegel, N., Seroussi, H., Shepherd, A., Sun, S., van de Wal, R., and 10 Ziemen, F. A.: Design and results of the ice sheet model initialisation experiments initMIP-Greenland: an ISMIP6 intercomparison, *The Cryosphere Discuss.*, 2017, 1–42, doi:10.5194/tc-2017-129, 2017.
- Gollledge, N. R., Menviel, L., Carter, L., Fogwill, C. J., England, M. H., Cortese, G., and Levy, R. H.: Antarctic contribution to meltwater pulse 1A from reduced Southern Ocean overturning, *Nature Communications*, 5, 5107, doi:10.1038/ncomms6107, 2014.
- Gomez, N., Pollard, D., and Mitrovica, J. X.: A 3-D coupled ice sheet – sea level model applied to Antarctica through the last 40 ky, *Earth and Planetary Science Letters*, 384, 88–99, doi:10.1016/j.epsl.2013.09.042, 2013.
- 15 Gregoire, L. J., Payne, A. J., and Valdes, P. J.: Deglacial rapid sea level rises caused by ice-sheet saddle collapses, *Nature*, 487, 219, doi:10.1038/nature11257, 2012.
- Greve, R., Saito, F., and Abe-Ouchi, A.: Initial results of the SeaRISE numerical experiments with the models SICOPOLIS and IcIES for the Greenland ice sheet, *Annals of Glaciology*, 52, 23–30, doi:10.3189/172756411797252068, 2011.
- 20 Hall, D. K., Comiso, J. C., DiGirolamo, N. E., Shuman, C. A., Box, J. E., and Koenig, L. S.: Variability in the surface temperature and melt extent of the Greenland ice sheet from MODIS, *Geophysical Research Letters*, 40, 2114–2120, doi:10.1002/grl.50240, 2013.
- Hanebuth, T., Stattegger, K., and Grootes, P. M.: Rapid Flooding of the Sunda Shelf: A Late-Glacial Sea-Level Record, *Science*, 288, 1033–1035, doi:10.1126/science.288.5468.1033, 2000.
- Harrison, S. P., Bartlein, P. J., Brewer, S., Prentice, I. C., Boyd, M., Hessler, I., Holmgren, K., Izumi, K., and Willis, K.: Climate model 25 benchmarking with glacial and mid-Holocene climates, *Climate Dynamics*, 43, 671–688, doi:10.1007/s00382-013-1922-6, 2014.
- Hindmarsh, R. C. A.: A numerical comparison of approximations to the Stokes equations used in ice sheet and glacier modeling, *Journal of Geophysical Research: Earth Surface*, 109, F01 012, doi:10.1029/2003JF000065, 2004.
- Hutter, K.: A mathematical model of polythermal glaciers and ice sheets, *Geophysical & Astrophysical Fluid Dynamics*, 21, 201–224, doi:10.1080/03091928208209013, 1982.
- 30 Hutter, K.: *Theoretical glaciology: material science of ice and the mechanics of glaciers and ice sheets*, Springer, 1983.
- Huybrechts, P.: Sea-level changes at the LGM from ice-dynamic reconstructions of the Greenland and Antarctic ice sheets during the glacial cycles, *Quaternary Science Reviews*, 21, 203–231, doi:10.1016/S0277-3791(01)00082-8, 2002.
- Ivins, E. R. and James, T. S.: Antarctic glacial isostatic adjustment: a new assessment, *Antarctic Science*, 17, 541–553, doi:10.1017/S0954102005002968, 2005.
- 35 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S., Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J. P., Stenni, B., Stocker, T. F., Tison, J. L.,

- Werner, M., and Wolff, E. W.: Orbital and Millennial Antarctic Climate Variability over the Past 800,000 Years, *Science*, 317, 793–796, doi:10.1126/science.1141038, 2007.
- Kavanagh, M. and Tarasov, L.: BrAHMs V1.0: A fast, physically-based subglacial hydrology model for continental-scale application, *Geosci. Model Dev. Discuss.*, 2017, 1–20, doi:10.5194/gmd-2017-275, 2017.
- 5 Koenig, S. J., Dolan, A. M., de Boer, B., Stone, E. J., Hill, D. J., DeConto, R. M., Abe-Ouchi, A., Lunt, D. J., Pollard, D., Quiquet, A., Saito, F., Savage, J., and van de Wal, R.: Ice sheet model dependency of the simulated Greenland Ice Sheet in the mid-Pliocene, *Climate of the Past*, 11, 369–381, doi:10.5194/cp-11-369-2015, 2015.
- Ladant, J.-B., Donnadieu, Y., Lefebvre, V., and Dumas, C.: The respective role of atmospheric carbon dioxide and orbital parameters on ice sheet evolution at the Eocene-Oligocene transition, *Paleoceanography*, 29, 2013PA002593, doi:10.1002/2013PA002593, 2014.
- 10 Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *Journal of Geophysical Research: Earth Surface*, 117, F01022, doi:10.1029/2011JF002140, 2012.
- Le clec’h, S., Fettweis, X., Quiquet, A., Dumas, C., Kageyama, M., Charbit, S., Wyard, C., and Ritz, C.: Assessment of the Greenland ice sheet – atmosphere feedbacks for the next century with a regional atmospheric model fully coupled to an ice sheet model, *The Cryosphere Discuss.*, 2017, 1–31, doi:10.5194/tc-2017-230, 2017.
- 15 Le clec’h, S., Quiquet, A., Charbit, S., Dumas, C., Kageyama, M., and Ritz, C.: A rapidly converging spin-up method for the present-day Greenland ice sheet using the GRISLI ice-sheet model, *Geosci. Model Dev. Discuss.*, 2018, 1–21, doi:10.5194/gmd-2017-322, 2018.
- Le Gac, H.: Contribution à la détermination des lois de comportement de la glace polycristalline (anélasticité et plasticité), Ph.D. thesis, Université Joseph-Fourier - Grenoble I, 1980.
- Le Meur, E. and Huybrechts, P.: A comparison of different ways of dealing with isostasy: examples from modeling the Antarctic ice sheet during the last glacial cycle, *Annals of Glaciology*, 23, 309–317, 1996.
- 20 Lhomme, N., Clarke, G. K., and Marshall, S. J.: Tracer transport in the Greenland Ice Sheet: constraints on ice cores and glacial history, *Quaternary Science Reviews*, 24, 173–194, doi:10.1016/j.quascirev.2004.08.020, 2005.
- Lipenkov, V. Y., Salamatin, A. N., and Duval, P.: Bubbly-ice densification in ice sheets: II. Applications, *Journal of Glaciology*, 43, 397–407, doi:10.3189/S0022143000034973, 1997.
- 25 Ma, Y., Gagliardini, O., Ritz, C., Gillet-Chaulet, F., Durand, G., and Montagnat, M.: Enhancement factors for grounded ice and ice shelves inferred from an anisotropic ice-flow model, *Journal of Glaciology*, 56, 805–812, doi:10.3189/002214310794457209, 2010.
- MacAyeal, D. R.: Large-Scale Ice Flow Over a Viscous Basal Sediment: Theory and Application to Ice Stream B, Antarctica, *Journal of Geophysical Research*, 94, pp. 4071–4087, doi:10.1029/JB094iB04p04071, 1989.
- Martin, M. A., Winkelmann, R., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 2: Dynamic equilibrium simulation of the Antarctic ice sheet, *The Cryosphere*, 5, 727–740, doi:10.5194/tc-5-727-2011, 2011.
- 30 Mouginit, J., Rignot, E., Scheuchl, B., Fenty, I., Khazendar, A., Morlighem, M., Buzzi, A., and Paden, J.: Fast retreat of Zachariæ Isstrøm, northeast Greenland, *Science*, 350, 1357–1361, doi:10.1126/science.aac7111, 2015.
- Mouginit, J., Rignot, E., Scheuchl, B., and Millan, R.: Comprehensive Annual Ice Sheet Velocity Mapping Using Landsat-8, Sentinel-1, and RADARSAT-2 Data, *Remote Sensing*, 9, 364, doi:10.3390/rs9040364, 2017.
- Nowicki, S. M. J., Payne, A., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd, A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geosci. Model Dev.*, 9, 4521–4545, doi:10.5194/gmd-9-4521-2016, 2016.

- Paolo, F. S., Fricker, H. A., and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348, 327–331, doi:10.1126/science.aaa0940, 2015.
- Patankar, S. V.: Numerical heat transfer and fluid flow, Taylor & Francis, Washington, DC., 1980.
- Pattyn, F.: Sea-level response to melting of Antarctic ice shelves on multi-centennial timescales with the fast Elementary Thermomechanical Ice Sheet model (f.ETISH v1.0), *The Cryosphere*, 11, 1851–1878, doi:10.5194/tc-11-1851-2017, 2017.
- 5 Peano, D., Colleoni, F., Quiquet, A., and Masina, S.: Ice flux evolution in fast flowing areas of the Greenland ice sheet over the 20th and 21st centuries, *Journal of Glaciology*, 63, 499–513, doi:10.1017/jog.2017.12, 2017.
- Peltier, W. R.: Global glacial isostasy and the surface of the Ice-Age Earth: The ICE-5G (VM2) Model and GRACE, *Annual Review of Earth and Planetary Sciences*, 32, 111–149, doi:10.1146/annurev.earth.32.082503.144359, 2004.
- 10 Pettit, E. C. and Waddington, E. D.: Ice flow at low deviatoric stress, *Journal of Glaciology*, 49, 359–369, doi:10.3189/172756503781830584, 2003.
- Peyaud, V.: Rôle de la dynamique des calottes glaciaires dans les grands changements climatiques des périodes glaciaires-interglaciaires., Ph.D. thesis, Université Joseph-Fourier - Grenoble I, 2006.
- Peyaud, V., Ritz, C., and Krinner, G.: Modelling the Early Weichselian Eurasian Ice Sheets: role of ice shelves and influence of ice-dammed lakes, *Climate of the Past*, 3, 375–386, doi:10.5194/cp-3-375-2007, 2007.
- 15 Philippon, G., Ramstein, G., Charbit, S., Kageyama, M., Ritz, C., and Dumas, C.: Evolution of the Antarctic ice sheet throughout the last deglaciation: A study with a new coupled climate–north and south hemisphere ice sheet model, *Earth and Planetary Science Letters*, 248, 750–758, doi:10.1016/j.epsl.2006.06.017, 2006.
- Pimienta, P.: Etude du comportement mécanique des glaces polychristallines aux faibles contraintes ; applications aux glaces des calottes polaires, thèse, Université Joseph-Fourier - Grenoble I, 1987.
- 20 Pollard, D. and DeConto, R. M.: Modelling West Antarctic ice sheet growth and collapse through the past five million years, *Nature*, 458, 329–332, doi:10.1038/nature07809, 2009.
- Pollard, D. and DeConto, R. M.: Description of a hybrid ice sheet-shelf model, and application to Antarctica, *Geosci. Model Dev.*, 5, 1273–1295, doi:10.5194/gmd-5-1273-2012, 2012.
- 25 Pollard, D., DeConto, R. M., and Alley, R. B.: Potential Antarctic Ice Sheet retreat driven by hydrofracturing and ice cliff failure, *Earth and Planetary Science Letters*, 412, 112–121, doi:10.1016/j.epsl.2014.12.035, 2015.
- Pritchard, H. D., Arthern, R. J., Vaughan, D. G., and Edwards, L. A.: Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets, *Nature*, 461, 971–975, 2009.
- Quiquet, A., Punge, H. J., Ritz, C., Fettweis, X., Gallée, H., Kageyama, M., Krinner, G., Salas y Méliá, D., and Sjolte, J.: Sensitivity of a Greenland ice sheet model to atmospheric forcing fields, *The Cryosphere*, 6, 999–1018, doi:10.5194/tc-6-999-2012, 2012.
- 30 Quiquet, A., Ritz, C., Punge, H. J., and Salas y Méliá, D.: Greenland ice sheet contribution to sea level rise during the last interglacial period: a modelling study driven and constrained by ice core data, *Clim. Past*, 9, 353–366, doi:10.5194/cp-9-353-2013, 2013.
- Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-Shelf Melting Around Antarctica, *Science*, 341, 266–270, doi:10.1126/science.1235798, 2013.
- 35 Ritz, C.: Un modèle thermo-mécanique d'évolution pour le bassin glaciaire antarctique Vostok-Glacier Byrd: Sensibilité aux valeurs des paramètres mal connus, thèse, Université Joseph-Fourier - Grenoble I, 1992.
- Ritz, C., Fabre, A., and Letrégouilly, A.: Sensitivity of a Greenland ice sheet model to ice flow and ablation parameters: consequences for the evolution through the last climatic cycle, *Climate Dynamics*, 13, 11–23, doi:10.1007/s003820050149, 1997.

- Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years: Implications for altitude changes in the Vostok region, *Journal of Geophysical Research*, 106, 31 943–31 964, doi:10.1029/2001JD900232, 2001.
- Ritz, C., Edwards, T. L., Durand, G., Payne, A. J., Peyaud, V., and Hindmarsh, R. C. A.: Potential sea-level rise from Antarctic ice-sheet instability constrained by observations, *Nature*, 528, 115–118, doi:10.1038/nature16147, 2015.
- 5 Roche, D. M., Dumas, C., Bügelmayer, M., Charbit, S., and Ritz, C.: Adding a dynamical cryosphere to iLOVECLIM (version 1.0): coupling with the GRISLI ice-sheet model, *Geosci. Model Dev.*, 7, 1377–1394, doi:10.5194/gmd-7-1377-2014, 2014.
- Rommelaere, V. and Ritz, C.: A thermomechanical model of ice-shelf flow, *Annals of Glaciology*, 23, 13–20, 1996.
- Rybak, O. and Huybrechts, P.: A comparison of Eulerian and Lagrangian methods for dating in numerical ice-sheet models, *Annals of Glaciology*, 37, 150–158, doi:10.3189/172756403781815393, 2003.
- 10 Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research (Earth Surface)*, 112, 2007.
- Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica, *Earth and Planetary Science Letters*, 223, 213–224, doi:10.1016/j.epsl.2004.04.011, 2004.
- Shreve, R. L.: Movement of Water in Glaciers, *Journal of Glaciology*, 11, 205–214, doi:10.3189/S002214300002219X, 1972.
- 15 Stone, E. J., Lunt, D. J., Rutt, I. C., and Hanna, E.: Investigating the sensitivity of numerical model simulations of the modern state of the Greenland ice-sheet and its future response to climate change, *The Cryosphere*, 4, 397–417, doi:10.5194/tc-4-397-2010, 2010.
- Tsai, V. C., Stewart, A. L., and Thompson, A. F.: Marine ice-sheet profiles and stability under Coulomb basal conditions, *Journal of Glaciology*, 61, 205–215, doi:10.3189/2015JoG14J221, 2015.
- Van Wessem, J., Reijmer, C., Morlighem, M., Mouginot, J., Rignot, E., Medley, B., Joughin, I., Wouters, B., Depoorter, M., Bamber, J., and
20 et al.: Improved representation of East Antarctic surface mass balance in a regional atmospheric climate model, *Journal of Glaciology*, 60, 761–770, doi:10.3189/2014JoG14J051, 2014.
- Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J. C., McManus, J. F., Lambeck, K., Balbon, E., and Labracherie, M.: Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records, *Quaternary Science Reviews*, 21, 295–305, doi:10.1016/S0277-3791(01)00101-9, 2002.
- 25 Weertman, J.: On the Sliding of Glaciers, *Journal of Glaciology*, 3, 33–38, doi:10.3189/S0022143000024709, 1957.
- Weertman, J.: Stability of the Junction of an Ice Sheet and an Ice Shelf, *Journal of Glaciology*, 13, 3–11, doi:10.3189/S0022143000023327, 1974.
- Whitehouse, P. L., Bentley, M. J., and Le Brocq, A. M.: A deglacial model for Antarctica: geological constraints and glaciological modelling as a basis for a new model of Antarctic glacial isostatic adjustment, *Quaternary Science Reviews*, 32, 1–24, doi:10.1016/j.quascirev.2011.11.016, 2012.
- 30 Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 1: Model description, *The Cryosphere*, 5, 715–726, doi:10.5194/tc-5-715-2011, 2011.
- Yan, Q., Zhang, Z., Gao, Y., Wang, H., and Johannessen, O. M.: Sensitivity of the modeled present-day Greenland Ice Sheet to climatic forcing and spin-up methods and its influence on future sea level projections, *Journal of Geophysical Research: Earth Surface*, 118, 2012JF002 709, doi:10.1002/jgrf.20156, 2013.
- 35




Figure 1. Schematic representation of the different types of flows in GRISLI and their associated velocity profiles. The red arrows stand for the sliding velocity which is non-zero for temperate-based grounded regions.




Figure 2. Horizontal staggered grids used in the model. The blue arrows stand for the staggered velocity grid while the green circles represent the standard centred grid (for, e.g., ice thickness, temperature, effective pressure). The plain brown line is an illustration of the grounding line position with an example of the flux (q'_{gl}) at one location where the grounding line crosses the x-axis in the centred grid. In the model, the norm of the flux at this location is reported on the u velocity only (q_{gl}), i.e. assuming a grounding line perpendicular to the x-axis (dashed brown line). The dark blue arrows are the velocity nodes on which q_{gl} is interpolated onto, using the velocity values of the two bounding light blue arrows. The dark green dots are used to infer the sub-grid position of the grounding, interpolating the flotation criterion ($\rho H + \rho_w (B - \text{sealevel})$).

Table 1. GRISLI model parameters used in this study.

Variable	Identifier name	Value
Global constants		
Gravitational acceleration	g	9.81 m s^{-2}
Water density, liquid	ρ_w	1000 kg m^{-3}
Water density, ice	ρ	918 kg m^{-3}
Water density, ocean	ρ_o	1028 kg m^{-3}
Isostasy		
Mantle density	ρ_m	3300 kg m^{-3}
Relaxation time in isostasy	τ_m	3000 yr
Lithosphere flexural rigidity	D_l	$9.87 \times 10^{24} \text{ N m}$
Radius of relative stiffness	R_l	131910 m
Radius of action of a unit mass	R_{iso}	400 km
Hydrology		
Thickness of the till layer	h_{till}	20 m
Porosity of the till layer	ϕ_{till}	0.5
Deformation		
Transition temperature of deformation, Glen	T_3^{trans}	-6.5°C
Activation energy below transition, Glen	E_{a3}^{cold}	$7.820 \times 10^4 \text{ J mol}^{-1}$
Activation energy above transition, Glen	E_{a3}^{warm}	$9.545 \times 10^4 \text{ J mol}^{-1}$
Flow law coefficient below transition, Glen	B_{AT03}^{cold}	$1.660 \times 10^{-16} \text{ Pa}^{-3} \text{ yr}^{-1}$
Flow law coefficient above transition, Glen	B_{AT03}^{warm}	$2.000 \times 10^{-16} \text{ Pa}^{-3} \text{ yr}^{-1}$
Transition temperature of deformation, linear	T_1^{trans}	-10°C
Activation energy below transition, linear	E_{a1}^{cold}	$4.0 \times 10^4 \text{ J mol}^{-1}$
Activation energy above transition, linear	E_{a1}^{warm}	$6.0 \times 10^4 \text{ J mol}^{-1}$
Flow law coefficient below transition, linear	B_{AT01}^{cold}	$8.373 \times 10^{-8} \text{ Pa}^{-3} \text{ yr}^{-1}$
Flow law coefficient above transition, linear	B_{AT01}^{warm}	$8.373 \times 10^{-8} \text{ Pa}^{-3} \text{ yr}^{-1}$
Temperature		
Gas constant	R	$8.314 \text{ J mol}^{-1} \text{ K}^{-1}$
Ice melting temperature, at depth z	T_m	$9.35 \times 10^{-5} \rho g (S - z) \text{ K kPa}^{-1}$
Latent heat of ice fusion	L_f	$335 \times 10^3 \text{ J kg}^{-1}$
Ice thermal conductivity, at temperature T	k_i	$3.1014 \times 10^8 \exp(-0.0057(T + 273.15)) \text{ J m}^{-1} \text{ K}^{-1} \text{ yr}^{-1}$
Mantle thermal conductivity	k_b	$1.04 \times 10^8 \text{ J m}^{-1} \text{ K}^{-1} \text{ yr}^{-1}$

Table 2. Selected parameters included in the latin hypercube sampling (LHS) ensemble with their associated ranges.

Parameter	Units	Minimum	Maximum
S_f	–	1.	5.
C_f	yr m ⁻¹	0.5 10 ⁻³	5. 10 ⁻³
K_0	m yr ⁻¹	20.10 ⁻⁶	200.10 ⁻⁶
b_{melt}	–	0.75	1.25

Table 3. Parameter values for the ensemble members that yield the lowest RMSE with respect to observations at the end of 100-kyr simulation under perpetual present-day climate forcing.

	ensemble member	S_f	C_f (yr m ⁻¹)	K_0 (m yr ⁻¹)	ϕ_{shelf}
Using Schoof (2007)	123	3.19	2.4 10 ⁻³	188 10 ⁻⁶	1.05
Using Tsai et al. (2015)	213	2.33	4.6 10 ⁻³	114 10 ⁻⁶	0.86




Figure 3. Antarctic ice shelves sectors (left) and associated prescribed present-day sub-shelf basal melting rates in m yr⁻¹ (right). The melting rates are different for the shelf and the associated grounding line to mimic the higher values observed close to the grounding line (Rignot et al., 2013). Sub-shelf melting rate for the deep ocean (depth greater than 2500 m) are assigned a value of 5 m yr⁻¹.




Figure 4. Simulated total Bedmap2 ice volume for each ensemble members as a function of parameter values when using thickness (left) and associated uncertainty (right) in metre (Fretwell et al., 2013), interpolated on the Schoof (2007) formulation-GRISLI grid of the flux Antarctica at the grounding line 40 km resolution. Despite considerable improvements from Bedmap1, large areas present an important uncertainty ($AN40S \pm 1000$ m). The thick horizontal line stands for the observations (Fretwell et al., 2013). The colour shading corresponds to the root mean square error in ice thickness relative due essentially to observations. The stars outlined in red are the 12 ensemble members that yield the lowest RMSE a poor in-situ data coverage.




Figure 5. Simulated total ice volume for each ensemble members as a function of parameter values when using the Schoof (2007) formulation of the flux at the grounding line (AN40S). The thick horizontal line shows the observations (Fretwell et al., 2013). The colour shading corresponds to the root mean square error in ice thickness relative to observations. The stars outlined in red are the 12 ensemble members having the lowest RMSE.




Figure 6. Same as Fig. 5 but with the Tsai et al. (2015) formulation of the flux at the grounding line (AN40T).




Figure 7. Ice thickness difference with the observations [in metres](#) (simulated minus observed) from the 12 ensemble members showing the lowest RMSE when using the Schoof (2007) formulation of the flux at the grounding line (AN40S).




Figure 8. Same as Fig. 7 but with the Tsai et al. (2015) formulation of the flux at the grounding line (AN40T).




Figure 9. Ice thickness root mean square error respective to observations in the parameter space for the 300 model members using the Schoof (2007) formulation of the flux at the grounding line (AN40S). The 12 experiments showing the lowest error are outlined in red.




Figure 10. Same as Fig. 9 but with the Tsai et al. (2015) formulation of the flux at the grounding line (AN40T).




Figure 11. Observed velocity (Mouginot et al., 2017) against modelled velocity on the 40 km grid. Only the best ensemble members with the lowest RMSE is shown here for Schoof (2007) (AN40S123, left) and Tsai et al. (2015) (AN40T213, right). The parameter values for these experiments are shown in Tab. 3.




Figure 12. Map of observed (Mouginot et al., 2017) and simulated velocities in m yr^{-1} for the ensemble members with the lowest RMSE using Schoof (2007) (AN40S123) and Tsai et al. (2015) (AN40T213).




Figure 13. Climatic perturbation used in the 400 kyr glacial-interglacial simulations for the near-surface air temperature, $\delta T = (1/\alpha^i) \delta D$, and for the sub-shelf basal melting rate modifier, $\delta^{oc} = \alpha^{oc} \Delta T_{NA}/T_{NA0}$.




Figure 14. Simulated total ice sheet volume evolution over the last 400 kyr for the twelve ensemble members showing the lowest RMSE in Sec. 3 when using the flux at the grounding line computed from Schoof (2007) (AN40S, shade of reds) and Tsai et al. (2015) (AN40T, shade of blues). The glacial to interglacial difference in ice volume for the last termination corresponds to about -10 to -20 m of global sea level rise equivalent depending on the simulations.




Figure 15. Simulated surface elevation at selected snapshots for the two ensemble members that produce the minimal RMSE at 0 kaBP in the transient simulations (AN40S252, top, and AN40T213, bottom). The ice volume contributing to sea level change from present is -9.3 m (resp. -15.1 m) at 21 kaBP for AN40S252 (resp. AN40T213), whilst it is limited at 120 kaBP (-0.3 m and +0.6 m). The grounding line is indicated by the thick red line.




Figure 16. Ice thickness difference with the observations (simulated minus observed) at 0 ka BP for the two ensemble members that produce the minimal RMSE at 0 kaBP in the transient simulations (AN40S252, left, and AN40T213, right). The RMSE is 350 m (respectively 313 m) for AN40S252 (resp. AN40T213).