

Dear Min-Hui Lo,

This is the response to the referee comments (RCs) on the manuscript “A Hydrological Cycle Model for the Globally Resolved Energy Balance Model (GREB) v1.0” submitted to GMD by Stassen et al. 2018 (gmd-2018-131). We like to thank the editor and the two anonymous referees for the time and effort spent on reviewing this manuscript and for the many helpful comments they provided. We think the referee comments have helped to substantially improve this manuscript. Please find a point-by-point response to all referee comments. All page and line numbers refer to the original manuscript and might not match up with the revised manuscript. We hope this settles all of the referee concerns and we would be happy to submit the revised manuscript.

Kind regards,
Christian Stassen

Referee 1

Comments

1. Can the authors elaborate the difference between the GREB model and CMIP5 CGCMs? As atmospheric and oceanic circulations are not simulated in the GREB model, it is probable that the GREB simulations give rise to results lacking of dynamical contribution. Does this lacking component play some roles in affecting the performance in the new model?

Response: CMIP models are earth system models containing several sub-models (i.e. atmosphere, ocean, vegetation, etc.). The atmospheric model would solve the dynamical equations of the atmosphere (i.e. Navier-Stokes). Those lead to internal variability (i.e. weather). GREB is an energy balance model. It does not contain weather or has any internal variability. Thus, reaching equilibrium (i.e. constant temperature) quickly and then remaining at equilibrium if not disturbed by external forcing.

Atmospheric circulation is not dynamically responding but prescribed as an external boundary condition. The sensitivity experiments ENSO & Climate Change in GREB take into account a change in circulation. This is done by adding an anomaly of horizontal winds and omega (and surface temperature) on top of the GREB climatological field. The anomaly is obtained from a composite of El Ninos (or La Ninas) for the ENSO case and the ensemble mean of the CMIP5 RCP8.5 response for the climate change case.

Oceanic circulations are not considered in GREB. The effect of ocean currents on the atmosphere would be reflected through a change in sea surface temperature (SST) which, is part of the GREB models external forcing. Thus, the GREB model would respond to a change in ocean currents through the change of SST.

We changed page 3 line 16-18 to: 'Thus, the GREB model is conceptually very different from the CGCM simulations in CMIP5, as atmospheric and the oceanic circulations are not simulated but prescribed as an external boundary condition in the model. The effect of ocean circulation on the atmosphere is represented only through the sea surface temperature.'

2. Due to the fact that CMIP5 CGCMs have biases in simulating circulations (e.g., Yang et al., 2018, Journal of Climate), the differences, at least for circulations, between the GREB model and CMIP5 models could be traced to the differences between prescribed wind fields in ERA-Interim reanalysis product and simulated wind fields. In other words, different background wind states may be part of the reason generating the discrepancy. What if comparing results using the CMIP5 simulated mean fields to force the old/new GREB model? Do authors have insights toward this point?

Response: We performed a set of experiments where the GREB model is forced with boundary conditions (i.e. horizontal winds and omega) from CMIP5 models. We compared the GREB precipitation anomaly (GREB-ERAInterim forced *minus* GREB-CMIP forced) to the CMIP precipitation anomaly (CMIP *minus* ERAInterim). This showed a high pattern correlation (~0.8). For the majority of the CMIP models we looked at, most of the correlation was caused by forcing

GREB with omega. However, this needs more research and will be done in future work before we can confidently address this.

We added the following to page 12 line 11: 'A very recent study by (Yang et al., 2018) links circulation biases in CMIP models to biases in precipitation and moisture. Forcing GREB with the circulation of CMIP models could shed more light how discrepancies in circulation between CMIP models effect the hydrological cycle in the GREB model.'

3. If no daily weather systems are simulated, does that mean the temporal integration is performed in the time step of one month or more in the GREB model? I am confused because the model time step is 12hrs as mentioned in Page 3 Line 14. In addition, having no weather system simulated does not mean no internal variability generated in the model. I suggest authors rephrasing this paragraph or providing further explanation.

Response: There are two different time steps in the GREB model:

- The physics and the tendency equations (i.e. tendencies for the hydrological cycle or surface temperature) are integrated on a 12 hours time step.
- Circulation (Advection and Diffusion) is integrated using a sub-stepping of 0.5 hours or 24 sub-steps. This is necessary for the model to be numerically stable.

We rephrased Page 3 Line 13-16 to: "The tendency equations of the model (i.e. tendency equation of specific humidity) are solved with a time step of 12 hours. For circulation, a shorter time step of 0.5 hours is used. This is necessary for the model to remain numerically stable. The daily cycle of incoming solar radiation is not resolved instead the 24hrs mean incoming solar radiation is used."

If the GREB model has no external forcings it will reach equilibrium quickly, depending on the magnitude of the forcing. After reaching equilibrium the model is stable and the tendencies of the model go towards zero. This means the GREB model is not oscillating around an equilibrium point.

We rephrased Page 3 Line 19-21 to: "Additionally, the GREB model has no internal variability. This means the model will converge to its equilibrium point and the tendency equations converge to zero."

4. Page 1 Line 8: Does "the hydrological cycle" refer to the hydrological model? Similar mixture appears throughout the abstract.

Response: Yes. The terms 'hydrological cycle' and hydrological model' are used interchangeably. We changed this in page 1 line 8 and through the manuscript to be more precise.

5. Page 1 Line 9: The authors should clarify the meaning of "zero order". Does the "order" means numerical convergence rate in time or space?

Response: With zero order we mean that is was a first guess or a rudimentary approach. The term is clarified in a bit more detail on page line 12: 'The hydrological cycle in the GREB model was

only needed as a zero order estimate to model the latent heat in the energy balance and the atmospheric water vapour levels.’ We changed the word on page 1 Line 9 to ‘rudimentary’ to not cause confusion with order of numerical convergence or order of accuracy.

6. Page 2 Line 6: Authors may consider mentioning the computational efficiency of idealized model here.

Response: We changed Page 2 Line 6-7 to: ‘Because of their simplicity, they help to develop hypotheses about the processes involved and they can be run fast. The GREB numerical code computes one model year in a few seconds and on a standard personal computer. It therefore is a relatively fast tool, which allows conducting sensitivity studies to external forcing within minutes to hours (Dommenget & Floter, 2011)’

7. Page 2 Line 23: I suggest re-organizing the paragraphs that describe the GREB model. For example, the modl layer configuration and resolution in Page 3 Line 10 can be introduced before the description of the NCEP climatological fields used in the original GREB model. This may make introduction of the GREB model framework smoother.


Response: We moved the paragraph describing the GREB model to the beginning of chapter 2

8. Page 2 Lines 25-26: Any specific reason generating topography from an atmospheric model? Why not using ETOPO dataset?



Response: We adopted the approach form the original GREB model.

9. Page 2 Line 29: I suggest providing brief explanation for the reason of changing dataset here and directing further details to section 3.4. Also, the NCEP reanalysis datasets are used during 1950-2008, whereas the ERA-Interim reanalysis data during 1979-2015. Therefore, long-term mean climatology values may be different. What is the results using the NCEP data during 1979-2015?

NCEP (1979-2015) minus NCEP (1950-2008)



ERA-Interim (1979-2015) minus NCEP (1950-2008)



Response: The difference between the mean climatologies between NCEP 1950-2008 vs. NCEP 1979-2015 is small compared to difference between NCEP and ERA-Interim (see plot above). We added the following to page 2 Line 28: 'ERA-Interim reanalysis has a higher accuracy than NCEP and a better agreement with observations (Liu et al., 2017)' and page 8 line 28: 'The effect of changing the mean climatology from the years 1950-2008 to 1979-2015 is small compared to the differences between NCEP and ERA-Interim.'

10. Page 4 Line 3: *How important is the diffusion term compared to advection? Could we ignore diffusion effect in large-scale circulation dynamics and/or thermodynamics?*

Response: The diffusion term is about 1/5 of the magnitude of the advection term (annually and



Figure 1: Annual mean advection (left) and diffusion (right) for the GREB model in kg/m²/s.

globally averaged, see Figure 1 below). Therefore, we did not ignore diffusion. We added: 'The diffusion term is one fifth of the magnitude of the advection term in global average but it is more important in some locations and therefore not ignored in the GREB model (not shown).'

11. Page 10 Line 6: *ENSO events include both El Niño and La Niña. Is the analysis for La Niña shown in the manuscript? Figure 10 only shows analysis for El Niño composite, but not La Niña. How does response to La Niña look like?*

Response: The improvement in the response of the GREB model to La Niña is similar to the improvement in GREB simulated El Niño. It is however, not shown in the manuscript.

We added to Page 10 Line 6: 'La Nina events are qualitatively the same, but with opposite signs (not shown).'

12. Page 10 Line 9: *Figure 10g shows the improvement in precipitation anomalies. It could be informative also discussing extratropical precipitation response because ENSO-mid-latitude linkages were also well documented. In Figure 10g, no precipitation increases in the Southern U.S. region. Does that indicate ENSO teleconnection is still not resolved well in the new model?*

Response: We focused here on a first order approximation of the response and do not want to discuss all details. The figure is in mm/day and therefore is focussed on the tropics where the absolute response in precipitation is strong. The extratropical response in GREB roughly matches

observations but is weaker than observations. We added the following to page 10 line 11: '... precipitation response is somewhat weaker than observed, especially in the extra tropics.'

13. Page 1 Line 15: *It is better to mention the full name of "CMIP" in the abstract.*

Response: Changed Page 1 Line 15 to: 'The new hydrological cycle is evaluated against the Coupled Model Inter-comparison Project phase 5 (CMIP5) model simulations, ...'

14. Page 1 Line 16: *El Nino -> El Niño (and through the manuscript)*

Response: Changed El Nino to El Niño throughout the manuscript. Changed La Nina to La Niña throughout the manuscript.

15. Page 1 Line 17: *Add the full name of "CGCM".*

Response: Added Coupled General Circulation Models (CGCMs) to Page 1 Line 17.

16. Page 1 Line 25: *What is "CGCMs" stands for? If it refers to "General Circulation Models" at the end of Line 24, it should be abbreviated as GCMs.*

Response: Changed in manuscript.

17. Page 1 Line 26: *"(AR4)" only appears once here, there's no need to provide abbreviation.*

Response: Removed AR4

18. Page 3 Line 5: *Figure 2c and 3c -> Figures 2c and 3c. I found this kind of error appears through the manuscript (e.g., Page 4 Line 4 and Page 5 Line 23). Please check carefully and revise them consistently.*

Response: Changed to Figures 2c and 3c on page 3 line 5 and through the manuscript.

19. Page 3 Line 7: *rcp85 -> RCP8.5 (to be consistent to that in the caption of Figure 11, Page 27).*

Response: Changed in manuscript.

20. Page 4 Line 10: *RHS: (dqair/dt)obs minus simulated terms?*

Response: That is correct! Changed the order in the manuscript

21. Page 7 Line 7: remove the parenthesis.

Response: Removed parenthesis

22. Page 9 Line 7: remove the parenthesis.

Response: Removed parenthesis and added 'and' between citations.

23. Page 17: what are the color shadings and streamlines in Figure 1d?

Response: Added '...850 hPa wind direction (streamline) and strength (shading)' to the caption.

Referee 2

Comments

1. I believe, however, that the authors need to explain better how this model can be used. While I realize that this is a technical paper describing the model, I think there needs to be some justification as to why we need this model in the first place.

Response: We plan to apply the model for studying biases in CMIP models. This could be done by replacing boundary conditions through CMIP boundary conditions. We revised the manuscript to better highlight the use of the GREB model. For example we added the following to page 12 line 11 (also to address RC1.2): 'A very recent study by Yang et al., 2018 links circulation biases in CMIP models to biases in precipitation and moisture. Forcing GREB with the circulation of CMIP models could shed more light how discrepancies in circulation between CMIP models effect the hydrological cycle in the GREB model.'

2. It is nice that the new version of GREB is more successful in reproducing certain aspects of the hydrological cycle. On the other hand, given that the new version has more fitting parameters, is this really surprising? Using more parameters gives you a better fit but also carries the risk of overfitting. In particular, the model might be too constrained by present day climate to be useful for climate change projection because basic features of the present climate, such as the width of the Hadley circulation or the position of the ITCZ, may change.

Response: We addressed the problem of overfitting by different approaches: first we tested the development of the model in step-wise building up the complexity (see section 3). Secondly, we did a number of response experiments that test the model's skill beyond the information used to fit the parameters. For this we did three tests: Seasonal cycle, El Nino and climate change. In all three the new model showed skills in simulating changes in the hydrological cycle that would not have been achieved by overfitting the model. We added some additional information in the introduction of Section 4 to better highlight this problem.

3. Another way the model could be used is for understanding the climate change response of more complex models. For this purpose, it would seem that GREB's mixture of basic principles (e.g. energy balance), ad-hoc parameterization (e.g. standard deviation of omega), and fitting to observations (e.g. mean omega) does not lend itself to interpretation any more than model output itself.

Response: We agree that the mixture of basic principles and ad-hoc parameterisations helps understanding the climate change response of more complex models (i.e. their biases). The boundary conditions of GREB could for example be replaced with climatologies of CMIP models (i.e. replacing horizontal winds from ERA-Interim with horizontal winds from one CMIP model). By replacing only one or all boundary conditions would help to gain insight where changes in RCP-scenarios come from or where biases in the hydrological cycle originate from. This is indeed what we think could be a useful application of this GREB model.

We added the following to page 12 line 11 (also to address RC1.2 & RC2.0): ‘A very recent study by (Yang et al., 2018) links circulation biases in CMIP models to biases in precipitation and moisture. Forcing GREB with the circulation of CMIP models could shed more light how discrepancies in circulation between CMIP models effect the hydrological cycle in the GREB model.’

3. section 3.3 What causes f to be 2.5 rather than 1.0? Could there be an error in the calculation? Is this mismatch horizontally uniform?

Response: There are several sources of uncertainties:

- The value of the scaling height we use in GREB is larger than literature values
- The fact that GREB is a single layer model
- The coarse resolution of the GREB horizontal grid
- A mismatch of the omega climatology
- Calculating circulation as residual

f is a constant fitting parameter it has no dimensions (see table 2 page 15). It could be fitted to different regions (i.e. tropics only or extra-tropics only) to get estimate if it is horizontally uniform.

We added to page 8 line 14: ‘... vertical velocities may not perfectly match because of the coarse resolution, GREB uses a scaling height of water vapour that is larger than literature values and calculating circulation as residual could contain other uncertainties.’

4. p. 1, ll. 25-26: I think the authors mean AR5 here (whose models are quite dated by now). Also, “the best possible” is certainly debatable. Perhaps it suffices to say that they are very complex models.

Response: Changed line 25-26 to: ‘CGCMs evaluated by the Intergovernmental Panel on Climate Change (IPCC) for the fifth assessment report (AR5), are among the most complex simulations of the climate system.’

5. Equation (12): Please explain all the variables directly after introducing the equation. In particular, what is u_{star} ? It seems that it should be near-surface wind speed, but then, on p.7 ll. 2-3 it says that wind speed is lower over land for a given u_{star} , suggesting that it is something else.

Response: U_{star} is the absolute wind climatology explained in Table 2. We introduced Table 2 before any equation is mentioned in section 2.

6. p. 8, l. 26: *Is the good match with observations for Ireland very meaningful if what you are really interested in is the global quality of the data?*

Response: Deleted: 'and Mooney et al. (2011) found a higher correlation of surface temperature in ERA-Interim to observations than NCEP in Ireland'

7. *The English could use a little editing (e.g. number agreement). Some examples: p. 2, l. 19: "parameterisations . . . is described" -> "are described" p. 3, l. 16: "wind and cloud cover field are" -> "fields" p. 3, l. 23: "a autoregressive" -> "an autoregressive" p. 5, l. 13: "precipitation and its seasonal cycle is shown" -> "are shown" p. 5, l. 3: "It, however, has. . ." -> "It has, however, . . ." p. 6, l. 3: "range of uncertainty CMIP5 modelled" -> "of CMIP5"*

Response: Thank you for pointing this out. We carefully read over the manuscript and changed all errors pointed out above plus those we additionally found throughout the manuscript.

A Hydrological Cycle Model for the Globally Resolved Energy Balance Model (GREB) v1.0

Christian Stassen¹, Dietmar Dommenges¹, Nicholas Loveday²

¹School of Earth Atmosphere and Environment, ARC Centre of Excellence for Climate System Science, Monash University, Melbourne, Australia

²Bureau of Meteorology, Darwin, Northern Territory, Australia

Correspondence to: Christian Stassen (christian.stassen@monash.edu)

Abstract. This study describes the development of the hydrological cycle model for the Globally Resolved Energy Balance (GREB) model. Starting from a rudimentary hydrological cycle model included in the GREB model, we develop three new models: precipitation, evaporation and horizontal transport of water vapour. Precipitation is modelled based on the actual simulated specific and relative humidity in GREB and the prescribed boundary condition of vertical velocity. The evaporation bulk formula is slightly refined by considering differences in the sensitivity to winds between land and oceans, and by improving the estimates of the wind magnitudes. Horizontal transport of water vapour is improved by approximating moisture convergence by vertical velocity. The new parameterisations are fitted against the Global Precipitation Climatology Project (GPCP) data set and reanalysis data sets (ERA-Interim). The new hydrological cycle model is evaluated against the Coupled Model Inter-comparison Project phase 5 (CMIP5) model simulations, reduction in correction terms and by three different sensitivity experiments (Annual Cycle, El Niño Southern Oscillation and Climate Change). The skill of the hydrological cycle model in the GREB model is now within the range of more complex CMIP5 Coupled General Circulation Models, and capable of simulating key features of the climate system within the range of uncertainty of CMIP5 model simulations. The results illustrate that the new GREB model's hydrological cycle is a useful model to study the climate's hydrological response to external forcings and also to study inter-model differences or biases.

1 Introduction

One topic in climate change that deserves urgent attention is the changing pattern of the hydrological cycle (Donat et al., 2016). Changes of rainfall have direct impact on the environment and on human health (Dai, 2011; Parry et al., 2004; Patz et al., 2005). The projections on how rainfall is changing are primarily based on Coupled General Circulation Models (CGCMs). CGCMs evaluated by the Intergovernmental Panel on Climate Change (IPCC) for the fifth assessment report are among the most complex simulations of the climate system. However, it is far from trivial to understand even simple aspects of the climate system as several processes interact with each other (Dommenges & Floter, 2011).

Rainfall is generated by a multitude of different systems (e.g. mid-latitude cyclones, tropical convection), which makes it one of the most complex processes in the climate system to model and thus to forecast. Yet many aspects of the hydrological cycle (i.e. high precipitation in the inner tropical convergence zone (ITCZ)) seen in complex CGCMs can be found in

Deleted: In this study, we

Deleted: very

Deleted: simple zero order

Deleted: observations

Formatted: Not Highlight

Deleted: n

Deleted: CGCMs

Deleted: and of rainfall

Deleted: ourth

Deleted: (AR4)

Formatted: Highlight

Deleted: ,

Deleted: best possible and

Deleted: But

Deleted: '

Deleted: because

models with intermediate complexity such as the ‘CLIMBER-2’ (Petoukhov et al., 1999), the ‘UVic earth system climate model’ (Weaver et al., 2001) or the simple atmosphere-ocean-sea-ice model developed by Wang and Myask (2000). Additionally, idealized models like the omega and humidity based model by Pendergrass and Gerber (2016) or the simple enhanced advection model by Chadwick et al. (2016) are capable of representing many aspects of the climate change response seen in complex CGCMs. Simplified climate models and energy balance considerations, are capable of explaining the large-scale features of the climate system and climate change (e.g. arctic amplification and land-sea contrast (Dommenges & Floter, 2011; Izumi et al., 2015). They provide a framework to conceptually understand the hydrological response to climate change. Because of their simplicity, they help to develop hypotheses about the processes involved. The GREB code computes about one model year per second on a standard personal computer. It therefore is a relatively fast tool, which allows conducting sensitivity studies to external forcing within minutes to hours (Dommenges & Floter, 2011). This paper will present a simple hydrological cycle model for the Globally Resolved Energy Balance (GREB) model (Dommenges & Floter, 2011). The aim of this hydrological cycle model is to present a simple and fast model for studies of the large-scale climate in precipitation, its response to climate variability (e.g. El Niño or climate change) and external forcings. The GREB model was originally developed to simulate the globally resolved surface temperature, and in particular its response to a CO₂ forcing. The hydrological cycle in the GREB model was only needed as a zero order estimate to model the latent heat in the energy balance and the atmospheric water vapour levels. We now aim for a representation of the hydrological cycle, that will allow the study the hydrological cycle in the GREB model by itself. We will improve three separate parameterisations in the model: precipitation, evaporation and the circulation of water vapour. The model will be based on the dynamical variables (surface temperature, atmospheric temperature and humidity) in the GREB model and on the boundary conditions of the GREB model (horizontal and vertical winds).

This paper is organised as follows, the following section will present the data sets used, the original GREB model and the methods. In Section 3 the new parameterisations of the hydrological cycle model in the GREB model are described. Section 4, will present three different sensitivity experiments to test the new hydrological cycle model. Finally, we give a discussion and summary of the results.

25 2 Data and Methods

The GREB model is a three layer (surface, atmosphere and deep ocean) global climate model on a 3.75 x 3.75 horizontal latitude-longitude grid. The GREB model simulates the thermal (long-wave) and solar (short-wave) radiation, heat transport in the atmosphere by isotropic diffusion and advection with the mean winds, the hydrological cycle (evaporation, precipitation and water vapour transport), a simple ice/snow albedo feedback and heat uptake in the sub-surface ocean. The tendency equations of the model (i.e. tendency equation of specific humidity) are solved with a time step of 12 hours. For the atmospheric transport equations, a shorter time step of 0.5 hours is used. This is necessary for the model to remain numerically stable. The daily cycle of incoming solar radiation is not resolved, instead the 24 hours mean incoming solar

- Deleted: .
- Deleted: numerical
- Deleted: in a few
- Deleted: s and
- Deleted: In this paper, we
- Deleted: El Nino
- Deleted: globally
- Deleted: ,
- Deleted: o
- Deleted: in the model
- Deleted: and
- Deleted: in
- Deleted: , we
- Deleted: is
- Deleted: In the Model Validation
- Deleted: e

radiation is used. The tendency equation of surface temperature, deep ocean temperature and specific humidity are flux corrected towards reanalysis data. The wind and cloud cover fields are seasonally prescribed boundary conditions. Thus, the GREB model is conceptually very different from the CGCM simulations in CMIP5, as atmospheric circulations and cloud cover are not simulated but prescribed as external boundary conditions in the model. The effect of ocean circulation on the atmosphere is represented only through the sea surface temperature, but is not explicitly simulated. Additionally, the GREB model has no internal variability, as atmospheric fluid dynamics (e.g. weather systems) are not explicitly simulated. Subsequently, the model will converge to its equilibrium points (all tendency equations converge to zero), if all boundary conditions are constant. The control climate or response to forcings can therefore be estimated from one single year.

Deleted: .
Deleted: This means the model
Deleted: Subsequently
Deleted: , t

The original GREB model used climatological fields from the NCEP reanalysis data from 1950 to 2008 (Kalnay et al., 1996) for surface temperature, T_{surf} , specific humidity and horizontal winds. The cloud climatology is taken from the International Satellite Cloud Climatology Project (Rossow & Schiffer, 1991). The ocean mixed layer depth is taken from Lorbacher et al. (2006). Topographic data is taken from the ECHAM5 atmosphere model (Roeckner et al., 2003). For more details refer to Dommenget and Floter (2011). For the development of the new GREB hydrological cycle model we replaced the NCEP reanalysis boundary conditions for T_{surf} , specific humidity and horizontal winds by using ERA-Interim reanalysis data from 1979 to 2015 (Dee et al., 2011). ERA-Interim reanalysis has a higher accuracy than NCEP and a better agreement with observations (Liu et al., 2017). The reasoning for the changed data sets is further explained in section 3.4. Precipitation observations are taken from the Global Precipitation Climatology Project (GPCP) (Adler et al., 2003). The climatological boundary conditions and constraints for the GREB model are summarised in Figure 1. In the following we will refer to these datasets as observations.

Deleted: SCCP project

The observed hydrological cycle in terms of the annual mean and its seasonal cycle for precipitation, evaporation and moisture circulation are shown in Figures 2 and 3. The global pattern of precipitation is marked by the ITCZ, its seasonal cycle and by the storm tracks of the midlatitudes. The evaporation is strongest over subtropical oceans and has a complex seasonal cycle with generally more evaporation in the warm season over land. The horizontal moisture transport (Figures 2c and 3c) is dominated by large scale convergence and divergence zones over the oceans and their seasonal shift.

Deleted: is
Deleted: gs.
Deleted: inner tropical convergence zone (
Deleted:)
Deleted: and
Deleted: ,
Deleted: in
Deleted: rcp
Formatted: Font:Bold, English (US)
Deleted: Table 1

Model simulations, pre-industrial (pi-Control) and representative concentration pathway 8.5 (RCP8.5), from the Coupled Model Inter-comparison Project phase 5 (CMIP5) database are used for comparison (Taylor et al., 2012). All datasets are re-gridded to a horizontal resolution of 3.75 x 3.75 to match the GREB model grid. See Table 1 for a complete list of models used.

Deleted: The GREB model is a three layer (surface, atmosphere and deep ocean) global climate model on a 3.75 x 3.75 horizontal latitude-longitude grid. GREB simulates the thermal (long-wave) and solar (short-wave) radiation, heat transport in the atmosphere by isotropic diffusion and advection with the mean winds, the hydrological cycle (evaporation, precipitation and water vapour transport), a simple ice/snow albedo feedback and heat uptake in the sub-surface ocean. The model main time step is 12hrs and the daily cycle of incoming solar radiation is not resolved (e.g. 24hrs mean incoming solar radiation). The tendency equation of surface temperature, deep ocean temperature and specific humidity are flux corrected towards reanalysis data. The wind and cloud cover field are seasonally prescribed boundary conditions. Thus, the GREB model is conceptually very different from the CGCM simulations in CMIP5, as atmospheric and the oceanic circulations are not simulated. Additionally, the GREB model has no internal variability, as daily weather systems are not simulated. Subsequently, the control climate or response to forcings can be estimated from one single year.

The original GREB hydrological cycle model, which is the starting point for this study, is shortly presented below. All variables and parameters are listed and explained in Table 2. The precipitation is proportional to the specific humidity

Formatted: Highlight
Formatted: Font:Bold
Deleted: Eq. (2)

$$\Delta q_{precip} = \tau_{precip} \cdot q_{air} \quad (1)$$

with Eq. (1), which corresponds to an autoregressive model with a decorrelation (recirculation) time of about 14 days (Dommenget & Floter, 2011). Evaporation, Δq_{evap} , in the original GREB model is calculated using an extended bulk formula:

$$\Delta q_{eva} = \frac{\rho_{air} \cdot c_w \cdot |u_* + c_{turb}| \cdot \vartheta_{soll} \cdot (q_{air} - q_{sat})}{r_{qvivv}} \quad (2)$$

The Bulk formula depends on the saturation deficit ($q_{air} - q_{sat}$), the wind speed u_* , with a turbulent wind factor c_{turb} , the density of air ρ_{air} , the transfer coefficient c_w , and a linear regression factor, r_{qvivv} , which links surface humidity to the vertically integrated water vapour column (Dommenget & Floter, 2011; Rapti, 2005).

5 The saturation water vapour pressure is calculated after (Dommenget & Floter, 2011; James, 1995):

$$q_{sat} = e^{z_{topo}} \cdot 3.75 \cdot 10^{-3} \cdot e^{17.08085 \frac{T_{surf}-273.15}{T_{surf}-38.975}} \quad (3)$$

Together, this leads to the complete tendency equation of specific humidity in GREB

$$\frac{dq_{air}}{dt} = \Delta q_{eva} + \Delta q_{precip} + \kappa \cdot \nabla^2 q_{air} - \vec{u} \cdot \nabla q_{air} + \Delta q_{correct} \quad (4)$$

10 with the diffusion term $\kappa \cdot \nabla^2 q_{air}$, the advection term $\vec{u} \cdot \nabla q_{air}$ and the flux correction term $\Delta q_{correct}$. The simulated annual mean and seasonal cycle for precipitation, evaporation and mean horizontal moisture transport are shown in Figures 2 and 3 for the original GREB model as discussed above. **The diffusion term is only one fifth of the magnitude of the advection term in global average (not shown) but is more important in some locations and therefore not ignored in the GREB model.** The original GREB model simulated some of the main features of the regional differences in the precipitation and evaporation, but many important details are missing (e.g. ITCZ, **subtropical dry regions or extra-tropical storm tracks**). However, horizontal moisture transport is not simulated well by the original GREB model.

Deleted: is

Deleted: ing

15 The seasonally varying **flux** correction term $\Delta q_{correct}$ is calculated as the residual between the tendencies without flux corrections and observed tendencies:

Deleted: um

$$\Delta q_{correct} = \left. \frac{dq_{air}}{dt} \right|_{obs} - \Delta q_{eva} + \Delta q_{precip} + \kappa \cdot \nabla^2 q_{air} - \vec{u} \cdot \nabla q_{air} \quad (5)$$

Deleted: $-\left. \frac{dq_{air}}{dt} \right|_{obs}$

20 This effectively corrects the GREB model to have a climatological specific humidity as observed. The flux correction term $\Delta q_{correct}$ can help to evaluate the improvements in the hydrological cycle model. The better the model the smaller the correction term should be in Eq. (4). We can therefore split the flux correction into three diagnostic terms

$$\Delta q_{correct} = \Delta q_{cor-precip} + \Delta q_{cor-eva} + \Delta q_{cor-circul} \quad (6)$$

25 With each term on the RHS representing the fraction of the flux corrections attributed to precipitation, evaporation and circulation biases, respectively. Each term is estimated as the difference between the observed and the GREB model tendencies of the humidity resulting from precipitation, evaporation and circulation biases:

$$\Delta q_{cor-precip} = \Delta q_{precip-OBS} - \Delta q_{precip-GREB} \quad (7)$$

$$\Delta q_{cor-eva} = \Delta q_{eva-OBS} - \Delta q_{eva-GREB} \quad (8)$$

$$\Delta q_{cor-circul} = \Delta q_{circul-OBS} - \Delta q_{circul-GREB} \quad (9)$$

30 with the GREB model tendencies of the humidity resulting from circulation, $\Delta q_{circul-GREB}$, defined as:

$$\Delta q_{circul} = \kappa \cdot \nabla^2 q_{air} - \bar{u} \cdot \nabla q_{air} \quad (10)$$

The observed humidity tendencies resulting from circulation, $\Delta q_{circul-GREB}$, are defined by the residual of the total humidity tendency minus the precipitation and evaporation tendencies. By construction, all three flux correction terms (evaporation, precipitation and circulation) sum up to the total flux correction term.

5 3 Hydrological Cycle Model Development

The development of the new hydrological cycle model of the GREB model is based on the existing zero-order hydrological cycle model of the GREB model. The following section will outline the development of each of the three models and discuss how the change in the reference climatologies from NCEP to ERA-interim has affected the model. All variables are summarised in Table 2.

Formatted: Normal, Don't keep with next

Deleted: We shortly

Formatted: English (US)

Formatted: Font:Bold, English (US)

Deleted: .

... [1]

3.1 Precipitation

The original GREB precipitation model captures some large-scale aspects of the mean and seasonal cycle of observed precipitation, such as more precipitation in the tropics and during warm seasons over land (Figures 2 and 3). It has, however, substantial differences from the observed precipitation, as it cannot capture the high rainfall in the ITCZ, the enhanced precipitation over the midlatitudes storm track regions and misses many aspects of the seasonal cycle. The root-mean-square error for the annual mean of the original GREB model precipitation parameterisation is 1.46 mm/day.

Deleted: ,

Deleted: has

Deleted:

Deleted:

Deleted:

The new parameterisation of precipitation in the GREB model is assumed to be proportional to q_{air} , as in the original GREB model. We further assume that relative humidity, r_q , and upward air motion, ω , increase rainfall. The latter is assumed to be a function of the mean and the standard deviation of the daily mean variation, ω_{mean} and ω_{std} , respectively. The new precipitation parameterisation is:

Deleted: parameterization

$$\Delta q_{precip} = r_{precip} \cdot q_{air} \cdot (c_{r_q} \cdot r_q + c_{\omega} \cdot \omega_{mean} + c_{\omega_{std}} \cdot \omega_{std}) \quad (11)$$

The model parameters, r_{precip} , c_{r_q} , c_{ω} and $c_{\omega_{std}}$ are fitted to minimise the RMSE between observations and GREB simulated precipitation. The resulting mean precipitation and its seasonal cycle are shown in Figures 2g and 3g. The model is evaluated in a Taylor diagram in Figures 4a and b against observations. The new GREB precipitation model is now very

Deleted: is

close to the observed precipitation patterns in both the mean and annual cycle. It is actually closer to the observed precipitation than any CMIP5 model (Figures 4a and 4d). We further test the different elements of the precipitation model by only considering a subset of the variables in Eq. (11), setting the other terms to zero and fitting the parameterisations for these reduced models. This allows us to estimate the effect of each term in the equation, see Figures 4a and 4b and Figure 5.

Deleted: parameterization

Relative humidity (r_q) is widely used in climate models as a predictor for precipitation (Petoukhov et al., 2005; Petoukhov et al., 1999; Wang & Myask, 2000; Weaver et al., 2001). In the GREB model it increases precipitation mainly over humid regions like the Amazons basin (Figure 5c and amplifies the seasonal cycle (Figure 5d). The overall pattern of rainfall with

high precipitation in the tropics and decreasing towards higher latitudes is not changed. Including r_q gives some moderate improvement relative to the original GREB model (Figure 4a comparing marker '0' to marker 'b').

The mean vertical air motion (ω_{mean}) provides a substantial improvement of the precipitation model (Figures 4a and d comparing marker '0' to 'c'). Ascending air masses in the ITCZ lead to increased precipitation, whereas descending air masses (i.e. in the subtropics) suppress precipitation. It creates a sharper and more realistic gradient in precipitation than the original GREB model (compare Figures 3d & 5e). With adding ω_{mean} , GREB is in the range of uncertainty of more complex CMIP5 models in the annual mean and the seasonal cycle (Figures 4a and d).

The GREB precipitation model without ω_{std} has still fairly weak mean precipitation in the midlatitudes storm track regions (compare Figures 5g and 2g) and has a weak seasonal cycle with the wrong sign in these regions as well (compare Figures 5h and 3g). The transient pressure systems in these regions lead to large vertical motions (ω) on shorter, daily time scales that result into large precipitation, but have a near zero ω_{mean} . Thus, to capture the precipitation in regions with strong variability in ω , but weak ω_{mean} , we include ω_{std} . This mainly enhances rainfall in the mid- and high latitudes (Figures 2g and 3g).

In summary, the new GREB precipitation model is significantly better than the original model. The RMSE is reduced by 0.65 mm/day to 0.81 mm/day in the annual mean and by 1 mm/day in the seasonal cycle. GREB precipitation now has a comparable skill to more complex CGCMs and lies within the range of uncertainty of CMIP5 modelled precipitation. Introducing the new precipitation parameterisation globally reduces the flux corrections of specific humidity caused by precipitation, see Figures 6c and 6d. The root-mean-square of the flux corrections caused by precipitation are reduced by more than 40%, indicating that the new parametrization has indeed improved the simulation of the hydrological cycle in the GREB model. Similar improvements are gained for the seasonal cycle (Figures 7c and 7d). The original GREB model showed large flux corrections, especially in the tropics where the ITCZ is moving with seasons and in the midlatitudes. The pattern of the flux corrections of the new model still looks similar to the original model, but is only half as large in amplitude (Figures 6c & d and 7c & d).

3.2 Evaporation

In the original GREB model evaporation is calculated using a widely used bulk formula approach (see Eq. (1) in Richter and Xie (2008)). This model does capture the main aspects of the regional differences in the annual mean evaporation in GREB, with enhanced evaporation over subtropical oceans and weaker evaporation over land (Figure 2e). The seasonal cycle (Figure 3e) is, however, very different from observed and the land-sea differences are too strong.

For the new evaporation model, we retained the original bulk formula approach and included a few minor changes by considering land-sea differences, revised wind (u_*) estimates, scaled effectivity and skin temperature. The new evaporation model is:

$$\Delta q_{eva} = \frac{\rho_{air} \cdot c_{eva} \cdot c_w \cdot |u_* + c_{turb}| \cdot v_{soil} \cdot (q_{air} - q_{sat-skin})}{r_{qvivw}} \quad (12)$$

Deleted: from

Deleted:

The constant c_{eva} modifies the evaporation efficiency for a given mean wind speed, u_* . $q_{sat-skin}$ is an estimate of saturated humidity considering skin temperature. It is calculated using:

$$q_{sat-skin} = e^{\frac{z_{topo}}{z_{atmos}}} \cdot 3.75 \cdot 10^{-3} \cdot e^{17.08085 \frac{T_{surf} + c_{eva-temp} - 273.15}{T_{surf} + c_{eva-temp} - 38.975}} \quad (13)$$

The parameter $c_{eva-temp}$ is a constant temperature offset to mimic skin temperature difference to T_{surf} . The parameters c_{eva} , $c_{eva-temp}$ and c_{turb} are fitted against observations for ocean and land points individually to minimise the RMSE. The values we estimated are:

$$c_{eva} = \begin{cases} 0.25 \text{ over land} \\ 0.58 \text{ over ocean} \end{cases} \quad (14)$$

$$c_{eva-temp} = \begin{cases} 5 \text{ K over land} \\ 1 \text{ K over ocean} \end{cases} \quad (15)$$

$$c_{turb} = \begin{cases} 11.5 \text{ over land} \\ 5.4 \text{ over ocean} \end{cases} \quad (16)$$

10 The scaled effectivity (c_{eva}) is lower over land than over oceans reflecting the fact that for a given u_* , more evaporation is simulated over oceans. This appears to be realistic considering that land has lower wind speeds near the surface for a given u_* due to the topography and vegetation. The value of $c_{eva} \cdot c_w$ closely match the observed values over oceans (Anderson & Smith, 1981; Merlivat, 1978).

15 The skin temperature difference approximated by $c_{eva-temp}$ is larger over land. It reflects that the GREB model does not simulate the daily cycle and the larger daily cycle over land leads to an effectively larger difference between the simulated T_{surf} and the skin temperature. The offset of 1°C over oceans is also found by Feng et al. (2018).

The wind magnitudes (u_*) in the original GREB model were estimated on the basis of the monthly mean climatologies of the zonal and meridional wind components. This, however, is not an accurate estimate of the monthly mean wind magnitudes, as it neglects the turbulent term due to high frequent variability. In the new GREB model we estimate the monthly mean u_* climatology based on the original 6 hourly ERA-Interim time steps.

20 We can estimate how much each of these changes improved the evaporation model by including only one of these changes and fitting the parameters of these models individually, see Figures 4 b & e and Figure 8.

Fitting the evaporation efficiency c_{eva} and the turbulent wind factor improves evaporation over land, especially in the seasonal cycle (Figure 8d) and reduces the strength of evaporation over the ocean. The increase in evaporation over land is caused by the increase in the turbulent wind factor. c_{eva} would decrease the evaporation in the annual mean and the seasonal cycle. By including the new estimate of monthly mean wind speed u_* , the pattern of evaporation is getting closer to observations, especially over the oceans (i.e. Figure 8f, North Atlantic) and by including the new estimate of skin temperature the seasonal cycle is improving slightly (Figure 4e).

30 The original GREB model was evaporating too much on the annual mean (see Figure 2e) especially over the equatorial pacific and Atlantic. The new hydrological cycle model parameterisation largely decreases evaporation over these regions

Deleted: -

Deleted: (Feng et al., 2018)

Deleted: was

Deleted: has

and the flux corrections are reduced over the globe in the annual mean (Figures 6e & f). The correlation of the annual mean experiences the largest changes from changing the reference climatology (Figure 4b).

In the seasonal cycle, each included variable improves the simulation of evaporation in the GREB model (Figure 4e). The seasonal cycle of flux corrections caused by evaporation in the original GREB model are large over land and large over oceans. There are positive flux corrections around the equator and negative flux corrections over the oceans north of the equator (Figure 7e). The improved evaporation seasonal cycle mainly removes this distinct pattern over the oceans and reduces flux corrections over most land areas. (Figures 7e & f). Overall, the new evaporation model is slightly better than in the original GREB model, but it still has substantial limitation in simulating the seasonal cycle correctly (Figures 2h & 3h).

3.3 Transport

The original GREB model transport of moisture was very weak and had little agreement with observations (Figs. 2f and 3f). Atmospheric transport of moisture in GREB (Eq. (4)) is controlled by diffusion and advection with mean winds. This model considered a divergence free 2-dimensional flow.

However, moisture convergence, as it occurs for example in the ITCZ, is important for the transport of moisture in these regions. The mean convergence by advection including the moisture convergence term is:

$$\overline{\nabla(\vec{u} \cdot q_{air})} = \overline{\vec{u} \cdot \nabla q_{air}} + q_{air} \cdot \overline{\nabla \vec{u}} \quad (17)$$

The second term on the RHS was not considered in the original GREB model, but is now considered in the new model. The moisture convergence term can be approximated by knowing the vertical air flow assuming continuity and hydrostatic balance:

$$q_{air} \cdot \overline{\nabla \vec{u}} \approx q_{air} \cdot f \cdot \frac{dt_{crl}}{z_{vapour} \cdot \rho_{air} \cdot g} \cdot (-\omega) \quad (18)$$

with the known parameters scaling height of water vapour, z_{vapour} , density of air, ρ_{air} , gravitational acceleration, g , and the circulation time step, dt_{crl} . The scaling factor, f , should theoretically be 1.0, but the mean large-scale horizontal winds and vertical velocities may not perfectly match because of the coarse horizontal resolution. Other factors that influence f could be, the single layer approximation, the GREB scaling height of water vapour that is larger than literature values or calculating the reference circulation as residual. A fit of Eq. (18) to observations finds that $f = 2.5$.

This new model has now a fairly realistic transport in the annual mean and the seasonal cycle (Figures 2i and 3i), with clear moisture transport out of regions with diverging flow (e.g. in the subtropics of coast of Peru) and into converging zones (e.g. ITCZ). The new parameterisation of convergence also reduces the flux corrections in the annual mean and the seasonal cycle (Figs. 6g & h and 7g & h).

Deleted: e
Deleted:

3.4 Boundary Conditions and Input Data

The original GREB model used the NCEP reanalysis as boundary conditions and as references for estimating the parameterisation of the model. New generations of reanalysis products have improved, because of the use of better models, better input data and better assimilation products (Dee et al., 2011). This is shown by Chen (2016) who investigated the variability and trends of the vertically integrated water vapour and found that ECMWF's ERA-Interim reanalysis has a higher accuracy than NCEP and a better agreement with observations over oceans and in the tropics. NCEP underestimates water vapour in troposphere (Kishore et al., 2011). We therefore changed the reference climatology of specific humidity in the GREB model from NCEP to ERA-Interim. To get a consistent model we also take surface temperature, horizontal winds the climatology of omega and standard deviation of omega from ERA-Interim. The effect of changing the mean climatology from the years 1950-2008 to 1979-2015 is small compared to the differences between NCEP and ERA-Interim. The parameters of our new GREB hydrological cycle model are then fitted against the new reference climatologies.

Deleted: parameterization

We estimate the effect that the change in reference climatologies have on the new GREB hydrological cycle model by fitting the parameters of the new model as described above to both the NCEP and ERA-interim reanalysis. The resulting hydrological cycle models are evaluated against observations (GPCP and ERA-interim) in Taylor diagrams for the annual mean. Changing the reference climatology doesn't lead to major improvements in the representation of the hydrological cycle in the GREB model but, it increases the correlation of precipitation, evaporation and circulation and reduces the RMSE (Figure S1 in the supplementary plots). The main improvement is in the tropics and might be related to the underestimated value of specific humidity in the tropics found by Chen (2016) and Kishore et al. (2011).

Deleted: and Mooney et al. (2011) found a higher correlation of surface temperature in ERA-Interim to observations than NCEP in Ireland

Deleted: C

Deleted: -

Deleted: This

Deleted: however

Deleted: (Chen & Liu, 2016; Kishore et al., 2011)

4 Model Verification

We now test the new hydrological model in a series of three different sensitivity experiments. The focus in the discussion will be on evaluating the new model. The three examples test the hydrological cycle model response to changes in the boundary conditions. These changes are beyond those used to fit the model parameterisation and can therefore be a test of the model's skill. We will leave more in-depth analysis of some of these experiments to future studies.

Deleted: z

Deleted: models

4.1 Seasonal Cycle

The response of the hydrological cycle to seasonal changes is a good test for evaluating the skill of the hydrological cycle model. The GREB model applies monthly flux correction terms to maintain a mean atmospheric humidity as observed. Thus, by construction the specific humidity in each calendar month in the GREB model will be identical to the observations, see Figure 9a.

To illustrate that the seasonal cycle is not a feature of the seasonally varying flux corrections we changed the flux corrections to an annual mean value for the original GREB model (middle column in Figure 9) and for the new GREB model (right

column in Figure 9). This annual mean flux correction value is added on every time step to the tendency equation of specific humidity (Eq. (4)).

With the new parameterisations for precipitation, evaporation and circulation the new GREB model resolves the seasonal cycle better than the original GREB model (Figure 9). The seasonal cycle of the original GREB model was too weak in the northern hemisphere when compared to observations and throughout the year the GREB model was too dry (Figure 9b). For the southern hemisphere, the original GREB model was too wet. The new GREB model captures the high humidity in northern hemispheric summer and the low values in winter (Figure 9c). This makes the seasonal cycle stronger in the new GREB model and it is closer to the reference climatology. In summary, the new GREB hydrological cycle model simulates the seasonal evolution of the atmospheric humidity very well and significantly better than the original GREB model.

4.2 El Niño Southern Oscillation

Strong El Niño and La Niña events lead to significant changes in the tropical precipitation and associated hydrological cycle changes. Since these natural modes of climate variability are well documented they present a good test case for the GREB model.

We therefore conducted a set of sensitivity experiments with the GREB model forced by the mean conditions for strong El Niño and La Niña events. The GREB model was forced with mean composites of T_{surf} , horizontal winds and omega from observations for four El Niño (1982/83, 87/88, 91/92, 97/98) and La Niña (1988/89, 99/00, 07/08, 10/11) events. The anomalies are calculated around El Niño/La Niña from May before the peak in December to April in the following year and against the climatological mean. In the GREB model simulation they are added on top of the reference climatology. The observed anomalies in the hydrological cycle during these El Niño events are shown in Figures 10a-c. La Niña events are qualitatively the same, but with opposite signs. We clearly note strong regional changes in the precipitation in the tropical Pacific that match changes in moisture transport (Figure 10c), illustrating that ENSO events mark strong regional changes in the hydrological cycle related to changes in the circulation.

The new GREB response in precipitation shows a strong similarity with the observed changes (Figure 10g). There is a shift of rainfall from the Maritime Continent towards the NINO3.4 region (5°N to 5°S & 170°W to 120°W) over the Pacific.

However, the overall amplitude in the precipitation response is in general weaker than observed. In contrast, the original GREB model has nearly no precipitation response to the ENSO forcings. This is consistent with the weak response in the circulation in the original GREB model (Figure 10f). The correlation between the GREB simulated El Niño response increases from 0.0 for the original GREB model to 0.9 with the new GREB model.

The observed evaporation response to ENSO events in the tropical Pacific is somewhat counteracting the precipitation response, as we observe mostly decreased evaporation over regions with enhanced precipitation and increased evaporation over regions with reduced precipitation (Figures 10a and b). These evaporation changes are mostly caused by changes in winds, with decreased evaporation over regions where the winds have weakened (e.g. NINO3.4 region). The new GREB model somewhat captures this pattern, but shows a stronger evaporation response, which partly explains the weaker

Deleted: El Nino

Deleted: El Nino

Deleted: La Nina

Deleted: El Nino

Deleted: La Nina

Deleted: El Nino

Deleted: La Nina

Deleted: El Nino

Deleted: La Nina

Deleted: ENSO

Deleted: n

Deleted: n

Deleted: somewhat

Deleted: El Nino

Deleted: The improvement in GREB simulated La Niña is similar to the improvement in GREB simulated El Niño (not shown).

precipitation response. However, both the original and the new GREB model evaporation have only a weak spatial correlation (0.3) with the observed evaporation changes overall.

The observed strong changes in the circulation of atmospheric humidity (Figure 10c) is mostly due to changes in the convergence of moisture (e.g. omega). Since, convergence of moisture was not considered in the original GREB model, the simulated changes in the circulation are very weak in the original GREB model (Figure 10f). The new GREB model does consider convergence of moisture and simulates the changes in the circulation of atmospheric humidity very similar to the observed (Figure 10i). The new circulation parameterisation in [the new GREB model](#) improves the correlation between the observed and the simulated circulation tendency from 0.3 (original GREB) to 0.95.

In summary, [the new GREB model](#) does simulate the precipitation and circulation response to ENSO conditions fairly well, whereas the original GREB model had very little skill, illustrating the significant improvement of the new GREB model over the original GREB model. However, the evaporation response in both models is not as well simulated as the precipitation and circulation response.

4.3 Global Warming

The response of the hydrological cycle to global warming is one of the potential applications of the GREB model and a comparison of the GREB model with the CMIP model simulations response to global warming provides a good test. The CMIP5 ensemble mean response of precipitation shows a distinct increase of rainfall in the equatorial pacific, decreases of mean rainfall in some subtropical regions (i.e. east pacific) and increases in some areas of the midlatitudes, see Figure 11a. This pattern is normally referred to as wet-get-wetter paradigm (Held & Soden, 2006). Although this approach has been questioned by more recent studies (Chadwick et al., 2013) it still gives a good first order approach of the changes in the global hydrological cycle, although changes over land might be muted or even reversed (He & Soden, 2016).

To evaluate the GREB hydrological cycle model independent of the other GREB model components, such as the T_{surf} tendencies, we force the original and new GREB model with RCP8.5 equivalent CO_2 concentrations and all other input variables for the hydrological cycle model taken from CMIP model simulations. That is, we [add \$T_{surf}\$, horizontal winds and vertical velocity, RCP8.5 CMIP5 ensemble mean anomalies from the models described in Table 1, on top of the GREB control reference climatologies](#). In the control run the reference boundary conditions of T_{surf} , horizontal winds and omega are taken.

The precipitation response in the original GREB model is positive in all locations and it closely follows the pattern of specific humidity in the control simulation (see Eq. (1) and Figure 11d). This is mainly due to an increase in the saturation water vapour pressure of about 7% per degree of warming (Clausius-Calpeyron). The original GREB precipitation response pattern is not correlated to the CMIP5 ensemble mean response pattern (Figure 12a), suggesting that local differences in the precipitation response are very different from those in the CMIP simulations.

The improved GREB model response pattern is similar to the CMIP models with enhanced and reduced response roughly at similar locations, [which leads to a much improved correlation \(Figures 12a and c\)](#). This is strongly related to the moisture

Deleted: prescribe
Deleted: in the scenario run taken from the
Deleted: of the
Formatted: Font:Bold, English (US)
Deleted: Table 1

Deleted: .

transport changes. However, the overall global mean precipitation response in the new GREB model is shifted upwards compared to the CMIP5 ensemble mean, which is related to the much stronger response in evaporation (compare Figures 11b and h). In CMIP5 models, we see a muted response of evaporation mainly due to changes in surface relative humidity and surface stability (Richter & Xie, 2008).

Deleted:

5 Summary and Discussion

In this study, we introduced the newly develop hydrological cycle model for the GREB model. It consists of three parts: precipitation, evaporation and transport. The development of these models started from the existing zero order hydrological cycle model of the GREB model and used physical reasoning and observations for fitting parameters.

The simulation of precipitation and transport of moisture in the new hydrological cycle model is now comparable in skill to CMIP models in terms of annual mean and the seasonal cycle of rainfall. The simulation of precipitation in the GREB model is closer to the observed precipitation pattern than any CMIP5 model in both, the annual mean and the seasonal cycle. This is directly related to the fact that the GREB mode has a prescribed atmospheric circulation, which is the main driver of the global precipitation pattern.

The evaporation has only improved slightly, but does simulate the annual mean values fairly well. However, it is still different from the observed seasonal cycle and the skill is much lower than that of the CMIP model. This suggests that the evaporation model is still a limiting factor in the GREB model.

We applied the new hydrological cycle model to a number of sensitivity studies, that illustrated that the new hydrological cycle model is much improved over the original GREB model. The annual cycle simulation without any correction terms is very realistic with the new model and the precipitation response to ENSO events is now very similar to the observed, owing to the much-improved transport of moisture. Finally, the response to global warming now shows a precipitation response pattern that is comparable to that of the CMIP models. Again, a limiting factor in this sensitivity experiment was the evaporation response of the GREB model in comparison to that of CMIP models.

Deleted: ,

An interesting aspect of the GREB model is that it has the atmospheric circulation (vertical and horizontal winds), humidity and surface temperatures as boundary conditions. This allows the GREB model to be used as a diagnostic tool to understand how different boundary conditions affect aspects of the climate system, such as the hydrological cycle's response to global warming. It may also help to study how biases in the hydrological cycle in CMIP models related to different boundary conditions from the atmosphere, such as biases in the vertical winds. A recent study by (Yang et al., 2018) links circulation biases in CMIP models to biases in precipitation and moisture. Forcing GREB with the circulation of CMIP models could shed light on how discrepancies in circulation between CMIP models effect the hydrological cycle.

Deleted: very

The new GREB hydrological cycle model is therefore a good tool in helping to conceptually understand the hydrological cycle and its response to global warming or other external forcings. It will further help in understanding CMIP model biases in the simulation of the hydrological cycle.

Deleted: more

Deleted: in the GREB model

Code availability

The GREB model source code used in this paper as well as the data used to run the model is available on GitHub: <https://github.com/christianstassen/greb-hydro-develop-gmd.git>. The GitHub repository contains detailed documentation on how to download the source code and installation instructions along with an example script on how to plot data obtained from GREB model simulations. The GREB source code is tested on recent-generation Mac platforms.

Acknowledgments

This study was supported by the Australian Research Council (ARC), with additional support coming via the ARC Centre of Excellence in Climate System Science and the ARC Centre of Excellence in Climate Extremes.

We like to thank the two anonymous referees for their constructive and helpful comments which lead to a substantial improvement in the manuscript.

Deleted: .

... [2]

References

- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P. P., Janowiak, J., . . . Nelkin, E.: The Version2 Global Precipitation Climatology Project (GPCP) Monthly Precipitation Analysis, *Journal of Hydrometeorology*, 4, 1147-1167, 2003.
- 15 Anderson, R. J., and Smith, S. D.: Evaporation coefficient for the sea surface from eddy flux measurements, *Journal of Geophysical Research*, 86, 449-456, 1981.
- Chadwick, R., Boutle, I., and Martin, G.: Spatial Patterns of Precipitation Change in CMIP5: Why the Rich Do Not Get Richer in the Tropics, *Journal of Climate*, 26(11), 3803-3822, doi:10.1175/jcli-d-12-00543.1, 2013.
- Chadwick, R., Good, P., and Willett, K.: A Simple Moisture Advection Model of Specific Humidity Change over Land in Response to SST Warming, *Journal of Climate*, 29(21), 7613-7632, doi:10.1175/Jcli-D-16-0241.1, 2016.
- 20 Chen, B. Y.: Global water vapor variability and trend from the latest 36year (1979 to 2014) data of ECMWF and NCEP reanalyses, radiosonde, GPS, and microwave satellite, *Journal of Geophysical Research-Atmospheres*, 121(19), 11442-11462, doi:10.1002/2016jd024917, 2016.
- Dai, A. G.: Drought under global warming: a review, *Wiley Interdisciplinary Reviews-Climate Change*, 2(1), 45-65, doi:10.1002/wcc.81, 2011.
- 25 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., . . . Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553-597, doi:10.1002/qj.828, 2011.
- Dommengat, D., and Floter, J.: Conceptual understanding of climate change with a globally resolved energy balance model, *Climate Dynamics*, 37(11-12), 2143-2165, doi:10.1007/s00382-011-1026-0, 2011.
- 30 Donat, M. G., Lowry, A. L., Alexander, L. V., O'Gorman, P. A., and Maher, N.: More extreme precipitation in the world's dry and wet regions, *Nature Climate Change*, 6(5), 508+, doi:10.1038/nclimate2941, 2016.
- Feng, X. B., Haines, K., and de Boissesson, E.: Coupling of surface air and sea surface temperatures in the CERA-20C reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 144(710), 195-207, doi:10.1002/qj.3194, 2018.
- 35 He, J., and Soden, B. J.: A re-examination of the projected subtropical precipitation decline, *Nature Climate Change*, 7(1), 53-57, doi:10.1038/nclimate3157, 2016.
- Held, I. M., and Soden, B. J.: Robust responses of the hydrological cycle to global warming, *Journal of Climate*, 19(21), 5686-5699, doi:10.1175/Jcli3990.1, 2006.

- Izumi, K., Bartlein, P. J., and Harrison, S. P.: Energy-balance mechanisms underlying consistent large-scale temperature responses in warm and cold climates, *Climate Dynamics*, *44*(11-12), 3111-3127, doi:10.1007/s00382-014-2189-2, 2015.
- James, I. N.: Introduction to circulating atmospheres, *Cambridge University Press*, 1995.
- 5 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., . . . Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bulletin of the American Meteorological Society*, *77*(3), 437-471, doi:10.1175/1520-0477(1996)077<0437:Tnyrp>2.0.Co;2, 1996.
- Kishore, P., Ratnam, M. V., Namboothiri, S. P., Velicogna, I., Basha, G., Jiang, J. H., . . . Sivakumar, V.: Global (50 degrees S-50 degrees N) distribution of water vapor observed by COSMIC GPS RO: Comparison with GPS radiosonde, NCEP, ERA-Interim, and JRA-25 reanalysis data sets, *Journal of Atmospheric and Solar-Terrestrial Physics*, *73*(13), 1849-1860, doi:10.1016/j.jastp.2011.04.017, 2011.
- 10 Liu, C. H., Ikeda, K., Rasmussen, R., Barlage, M., Newman, A. J., Prein, A. F., . . . Yates, D.: Continental-scale convection-permitting modeling of the current and future climate of North America, *Climate Dynamics*, *49*(1-2), 71-95, doi:10.1007/s00382-016-3327-9, 2017.
- 15 Lorzbacher, K., Dommenges, D., Niiler, P. P., and Kohl, A.: Ocean mixed layer depth: A subsurface proxy of ocean-atmosphere variability, *Journal of Geophysical Research-Oceans*, *111*(C7), doi:Artn C07010 10.1029/2003jc002157, 2006.
- Merlivat, L.: The dependence of bulk evaporation coefficients on air-water interfacial conditions as determined by the isotopic method, *Journal of Geophysical Research*, *83*, 2977-2980, 1978.
- 20 Parry, M. L., Rosenzweig, C., Iglesias, A., Livermore, M., and Fischer, G.: Effects of climate change on global food production under SRES emissions and socio-economic scenarios, *Global Environmental Change-Human and Policy Dimensions*, *14*(1), 53-67, doi:10.1016/j.gloenvcha.2003.10.008, 2004.
- Patz, J. A., Campbell-Lendrum, D., Holloway, T., and Foley, J. A.: Impact of regional climate change on human health, *Nature*, *438*(7066), 310-317, doi:10.1038/nature04188, 2005.
- 25 Pendergrass, A. G., and Gerber, E. P.: The Rain Is Askew: Two Idealized Models Relating Vertical Velocity and Precipitation Distributions in a Warming World, *Journal of Climate*, *29*(18), 6445-6462, doi:10.1175/Jcli-D-16-0097.1, 2016.
- Petoukhov, V., Claussen, M., Berger, A., Crucifix, M., Eby, M., Eliseev, A. V., . . . Weaver, A. J.: EMIC Intercomparison Project (EMIP-CO2): comparative analysis of EMIC simulations of climate, and of equilibrium and transient responses to atmospheric CO2 doubling, *Climate Dynamics*, *25*(4), 363-385, doi:10.1007/s00382-005-0042-3, 2005.
- 30 Petoukhov, V., Ganopolski, A., Brovkin, V., Claussen, M., Eliseev, A., Kubatzki, C., and Rahmstorf, S.: CLIMBER-2: a climate system model of intermediate complexity. Part I: model description and performance for present climate, *Climate Dynamics*, *16*, 1-17, 1999.
- 35 Rapti, A. S.: Spectral optical atmospheric thickness dependence on the specific humidity in the presence of continental and maritime air masses, *Atmospheric Research*, *78*(1-2), 13-32, doi:10.1016/j.atmosres.2005.02.004, 2005.
- Richter, I., and Xie, S.-P.: Muted precipitation increase in global warming simulations: A surface evaporation perspective, *Journal of Geophysical Research*, *113*(D24), doi:10.1029/2008jd010561, 2008.
- Roeckner, E., Bauml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., . . . Tompkins, A.: The Atmospheric General Circulation Model ECHAM5. Part I: Model Description, *349*, 2003.
- 40 Rossow, W., and Schiffer, R.: Isccp cloud data products, *Bulletin of the American Meteorological Society*, *72*, 2-20, 1991.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of Cmp5 and the Experiment Design, *Bulletin of the American Meteorological Society*, *93*(4), 485-498, doi:10.1175/Bams-D-11-00094.1, 2012.
- 45 Wang, Z., and Myask, L. A.: A Simple Coupled Atmosphere-Ocean-Sea Ice-Land Surface Model for Climate and Paleoclimate Studies*, *Journal of Climate*, *13*, 1150-1172, 2000.
- Weaver, A. J., Eby, M., Wiebe, E. C., Bitz, C. M., Duffy, P. B., Ewen, T. L., . . . Yoshimori, M.: The UVic earth system climate model: Model description, climatology, and applications to past, present and future climates, *Atmosphere-Ocean*, *39*, 361-428, 2001.

Tables

Table 1: List of CMIP 5 models.

<i>Models</i>		
ACCESS1-0	CSIRO-Mk3-6-0	HadGEM2-ES
ACCESS1-3	CanESM2	inmcm4
bcc-csm1-1	EC-EARTH	MIROC-ESM-CHEM
bcc-csm1-1-m	FGOALS-g2	MIROC-ESM
BNU-ESM	FGOALS-s2	MIROC4h
CCSM4	FIO-ESM	MIROC5
CESM1-BGC	GFDL-CM3	MPI-ESM-LR
CESM1-CAM5	GFDL-ESM2G	MPI-ESM-MR
CESM1-FASTCHEM	GFDL-ESM2M	MPI-ESM-P
CESM1-WACCM	GISS-E2-H-CC	MRI-CGCM3
CMCC-CM	GISS-E2-H-R	NorESM1-M
CMCC-CM5	HadGEM2-CC	NorESM1-ME

Table 2: Variables of the GREB model

Variable	Dimension	Description
c_{eva}	constant	Evaporation efficiency
$c_{eva-temp}$	constant	Temperature scaling of evaporation
c_{turb}	constant	Turbulent wind offset for evaporation
c_{rq}	constant	Precipitation parameter for rel. humidity
c_{ω}	constant	Precipitation parameter for omega
$c_{\omega std}$	constant	Precipitation parameter for standard deviation of omega
f	constant	convergence scaling parameter
g	constant	gravitational acceleration
q_{air}	x, y, t	atmospheric humidity
q_{sat}	x, y, t	saturation pressure
$q_{sat-skin}$	x, y, t	saturation pressure with temperature offset
r_{precip}	constant	mean lifetime of water vapour
r_{qivwv}	constant	regression between atm. humidity and vertically

		integrated water vapour
r_q	x, y, t	relative humidity
T_{surf}	x, y, t	surface temperature
$ u_s $	x, y, t	Absolute wind climatology
\vec{u}	x, y, t	Horizontal wind climatology
z_{atmos}	constant	Scaling height of atmosphere
z_{topo}	x, y, t	Topographic height
z_{vapour}	constant	Scaling height of water vapour
θ_{soil}	x, y, t	Surface wetness fraction
ρ_{air}	constant	Density of air
ω_{std}	x, y, t	Standard deviation of vertical wind climatology
Δq_{eva}	x, y, t	mass flux for the atmospheric humidity by evaporation
Δq_{precip}	x, y, t	mass flux for the atmospheric humidity by precipitation
$\Delta q_{correct}$	x, y, t	Mass flux correction of specific humidity
$\Delta q_{cor-circul}$	x, y, t	Mass flux correction due to circulation
$\Delta q_{cor-eva}$	x, y, t	Mass flux correction due to evaporation
$\Delta q_{cor-precip}$	x, y, t	Mass flux correction due to precipitation
$\Delta q_{precip-GREB}$	x, y, t	Precipitation change in GREB
$\Delta q_{precip-OBS}$	x, y, t	Precipitation change in observations
Δt	constant	Model integration time step
dt_{crl}	constant	Model integration time step for circulation
κ	constant	Isotropic diffusion coefficient
ω	x, y, t	Vertical velocity in pressure coordinates

Figures




Figure 1: GREB mean state boundary conditions and reference climatologies: topography (a), surface temperature (b), surface humidity (c), 850 hPa wind **direction (streamline) and strength (shading)** (d), vertical velocity omega (e) and the daily standard deviation of vertical velocity omega (f).

Deleted: s




Figure 2: The decomposition of the hydrological cycle into its parts precipitation in mm/day (left column), evaporation (middle column) and circulation in kg/m²/s (right column) in observations (upper row), the original GREB model (middle row) and the new GREB model (lower row) for the annual mean.

5




Figure 3: As Fig. 2 but for the seasonal cycle (DJF-JJA). The decomposition of the hydrological cycle into its parts precipitation in mm/day (left column), evaporation (middle column) and circulation in kg/m²/s (right column) in observations (upper row), the original GREB model (middle row) and the new GREB model (lower row) for the seasonal cycle.

5





Figure 4: Precipitation (left column), evaporation (middle column) and circulation (right column) in the annual mean (top row) and seasonal cycle (bottom row) in mm/day in a Taylor diagram against observations from GPCP and ERA-Interim. Red colours indicate different GREB parametrizations with 0 being the original and * the best parametrization. Blue dots are pi-Control CMIP5 models and the green cross indicates the ensemble mean of all CMIP5 models.

5



5 **Figure 5:** Annual mean precipitation for four development steps of the GREB precipitation parametrisation (a, c, e, g) and their corresponding seasonal cycles (b, d, f, h) in mm/day. The first step was changing the specific humidity boundary climatology (a) and (b). Then subsequently more variables have been added to the precipitation parametrisation: adding only relative humidity (c, d), adding only omega (e, f), adding relative humidity and omega (g, h).

Flux corrections - Annual mean




Figure 6: Annual mean flux corrections of specific humidity for the original GREB model (a) and the improved GREB model (b). The flux corrections are then split into their contributions of precipitation (c, d), evaporation (e, f) and circulation (g, h) for the original GREB model (left column) and the improved GREB model (right column) in kg/m²/s. The top right shows the global root-mean-square (RMS).

5

Flux corrections - Seasonal cycle




Figure 7: As Fig. 6 but for the seasonal cycle (DJF-JJA). Flux corrections of specific humidity for the original GREB model (a) and the improved GREB model (b). The flux corrections are then split into their contributions of precipitation (c, d), evaporation (e, f) and circulation (g, h) for the original GREB model (left column) and the improved GREB model (right column) in kg/m²/s. The top right shows the global root-mean-square (RMS).

Evaporation




Figure 8: Annual mean evaporation for three development steps of the GREB evaporation parametrisation (a, c, e) and their corresponding seasonal cycles (b, d, f) in $\text{kg/m}^2/\text{s}$. The first step was changing the boundary climatology (a) and (b). Then subsequently more variables have been added to the evaporation parametrisation: fitting the evaporation parameters separately for ocean and land (c, d) and fitting parameters and prescribing the wind speed (e, f).




Figure 9: Annual cycle of specific humidity with seasonal varying flux corrections (a, d) and annual mean flux corrections for original GREB (b, e) and improved GREB (c, f) in g/kg. The top row shows the northern (solid) and southern (dashed) hemispheric mean for observations (black) and GREB (blue). The bottom shows the respective seasonal cycle (DJF-JJA). For the seasonally varying flux corrections (a) GREB (blue) matches observations (black).





Figure 10: The **El Niño** response of the hydrological cycle in: observations for precipitation (a) in mm/day, evaporation (b) and circulation (c) in kg/m²/s (upper), original GREB model for precipitation (d), evaporation (e) and circulation (f) (middle) and the improved GREB model for precipitation (g), evaporation (h) and circulation (i) (lower). GREB uses prescribed anomalies from an **El Niño** composite mean of surface temperature, horizontal winds and vertical winds (omega).

Deleted: El Niño

Deleted: El Niño



5 **Figure 11:** Response of the hydrological cycle to an RCP8.5 forcing in the: CMIP5 ensemble mean for precipitation (a) in mm/day, evaporation (b) and circulation (c) in kg/m²/s (upper), original GREB model for precipitation (d), evaporation (e) and circulation (f) (middle) and the improved GREB model for precipitation (g), evaporation (h) and circulation (i) (lower). GREB uses prescribed anomalies from CMIP5 ensemble mean of surface temperature, horizontal winds and vertical winds (ω). All responses are shown per degree of warming.




Figure 12: RCP8.5 response of CMIP5 models (blue), original GREB (0) and improved GREB (*) per degree of global warming against the CMIP5 ensemble mean (black star). Precipitation is shown on the left, evaporation in the middle and circulation on the right column. GREB uses prescribed anomalies from the CMIP5 ensemble mean of surface temperature, horizontal winds and vertical winds (ω). The correlation of the original GREB model precipitation response with the ensemble mean is zero. The original and improved GREB model have zero correlation with the ensemble mean evaporation and the standard deviation is one for both.




Figure S1: Precipitation (left column), evaporation (middle column) and circulation (right column) in the annual mean (top row) and seasonal cycle (bottom row) in mm/day in a Taylor diagram against observations. Red colours indicate different GREB parametrizations with 0 being the original and * and diamond the best parametrization. Star (*) is the best model for the ERA-Interim boundary conditions and diamond uses the NCEP boundary conditions. Blue dots are CMIP5 models and the green cross indicates the ensemble mean of all CMIP5 models.

Table 2