

Author response to reviews for “Evaluating Simplified Chemical Mechanisms within Present-Day Simulations of CESM Version 1.2 CAM-chem (CAM4): MOZART-4 vs. Reduced Hydrocarbon vs. Super-Fast Chemistry”

By Benjamin Brown-Steiner et al.

We first want thank the reviewers for their valuable and insightful comments, and for taking the time to review our manuscript.

We now respond to the reviewer comments, which are reproduced in black text below. Our responses follow immediately in red text, and any additions to the manuscript are included in italic red text, along with Line references, which refer to their locations in the revised manuscript.

Editor, F. O'Connor:

Dear Benjamin and co-authors,

As the topical editor for your manuscript, may I remind you that I'm requesting that the data be made available on a suitable repository with a digital object identifier (DOI) before final publication of your manuscript in GMD. Reference to this repository can then be included in your final manuscript. I trust that this will be acceptable to you.

Regards, Fiona O'Connor

We have uploaded the relevant data to a repository hosted on the MIT domain (<http://dspace.mit.edu/handle/1721.1/114993>). The Data/Code availability sections have been updated to direct readers to this repository, as is discussed in the comments below.

Anonymous Referee #1

Received and published: 13 March 2018

Brown-Steiner et al present a study of how two reduced chemical mechanisms perform in a number of comparisons against the more comprehensive MOZART scheme (MO). The Reduced Hydrocarbon mechanism (RH) contains around half the number of MOZART reactions, whilst the Super-Fast mechanism (SF) is about 1/6 of MO. The work has been done to explore how much of a compromise it is by choosing one of these simple mechanisms over the more comprehensive scheme, when considering computational time gains versus accuracy of the chemical predictions.

The model runs have been conducted on a global scale for 25 years. This has enabled the authors to pull out modelled data to compare with each other, and for the time periods and locations of ozonesonde, aircraft and CASTNET observations. Given how small SF is, it performs unexpectedly well, particularly against the CASTNET data, and in some cases better than the

other two schemes. The RH scheme often tracks MO quite well, but with some exceptions, particularly for CO.

The Super-Fast mechanism could be used to explore chemical sensitivity studies in a fraction of the time it would take to run MOZART in locations of low biogenic activity.

I think the manuscript fits within the GMD journal remit and should be published. I have a few minor comments and queries.

General comments:

Please check throughout for the consistency about the length of the run. Page 6 line 1. ‘we use MERRA.. for 25 years (1990 – 2014)’. However figure 1 plots maps for the year 2015, and figure 4 shows vertical distributions for 2015? Figure 6 also looks like it starts at 1991, not 1990 and carries on beyond 2014. Same applies to the statement on page 9, end of line 13 about the run being 1990-2014. Also figure 7 ‘for the full 1990-2015 time series’, which is 26 years.

We ran the simulations for 26 years (1990 – 2015) and used the first year as spin up, so the analyses are for 25 years (1991 – 2015). We have corrected the descriptions and added a line indicating the 1990 year as spin up:

Page 6, Line 2: “...for 26 years (1990 – 2015)...”

Page 6, Lines 3-4: “*The year 1990 is dropped to allow for spin-up.*”

Page 9, Line 24-25: “...sites throughout the 1991 – 2015 period...”

Figure 7, Caption: “...the full 1991 – 2015...”

Section 2.2.1 MOZART-4. This section is very short and doesn’t give the reader much information about MOZART other than to go searching through the suggested literature. I think a bit more information on what the scheme includes (e.g. how many alkanes/aromatics/biogenic species are considered explicitly) and omits would be useful, particularly as it is being used as the benchmark scheme.

We have added additional details and point again to Emmons et al. (2010) for a complete description.

Section 2.2.1: “*As described in detail in Emmons et al. (2010), MOZART-4 mechanism is a tropospheric mechanism that contains 85 gas-phase species and 12 bulk aerosol species, with 39 photolysis and 157 gas-phase reactions. Large alkanes, alkenes, and aromatics are lumped together (BIGALK, BIGENE, and TOLUENE, respectively), and monoterpenes are lumped together as C10H16 and treated as α -pinene.*”

Page 9 line 1. Please describe what ‘BAM’ means

Page 9, Line 10: “...the *Bulk Aerosol Model* (BAM) (see Tilmes et al., 2015)...”

Page 10. Line 20. There is a single line describing figure 6 and I then didn’t fully understand the results drawn from it. My assumption is that the range of ozone at each time step in the model has been extracted for the region and the differences in the percentiles plotted here (although the

figure caption says the CASTNET observations are only for JJA?). The results say it takes 5-10 years for the models to stabilise, but the plots also show that the CASTNET observations themselves need 5-10 years to stabilize? If the models are behaving similarly to the observations, why would we need the spin up? There's a bit of a leap of understanding, so I think a bit more description is needed. I can see why you would expect the range in predicted ozone to settle as times goes on using constant emissions – but why also in the observations?

In a concurrent paper under review in ACPD (Brown-Steiner et al., in review) we expand on the implications of this figure. We add additional clarifications of Figure 6 and point to reader to the concurrent paper for additional discussion. We also add this paper to the references.

Page 11, Line 5: “Brown-Steiner et al. (in review, ACPD) examines these implications, and also concludes that it takes approximately 10-years for long-term signals to emerge from meteorological variability. These results demonstrate the challenge in examining chemical signals in highly variable data, particularly if there are trends or changes to the ozone distribution, as is seen in the CASTNET data for the Southeastern US.”

Page 14 line 16 spelling. ‘mechanism’, not ‘mechanisms’.

Corrected.

Figures:

Figure 5, panels g,h and i. what does the 1x1 refer to?

They are individual grid cells within each region. This has been added to the caption.

Figure 5, caption: “Plots g, h, and i are individual grid cells from within each region.”

Figure 7. Titles overlap with plots.

The figure has been updated to correct this.

Figure 8. I’m struggling to see the orange N2O5 line in any of these plots. It could be that the line is hidden under the PAN+N2O5 line, but given the variation between PAN+N2O5 and N2O5 in the global ozone plot, I expected to see it?

The figure has been updated so that each line is more easily discernable.

Figure 8. Which single year are these seasonal cycles for? Why was this particular year chosen? How much variability is there between the first year run (1990) and the last (2014)?

Year 2015 was selected as a representative year, and we find some variability year-to-year due to meteorology, but all models tend to demonstrate the same year-to-year variability. For the sensitivity tests with PAN and N2O5, we only ran 2 years and selected the 2nd year (1991). We

also discovered a bug in the plotting code in the sensitivity tests which has been corrected. Some of the seasonal cycles were offset by 2 months, and but does not impact our conclusions.

Figure 8, Caption: "...single year (2015), averaged..."

Figure 9, Caption: "...SF mechanism (which were ran only for 2 years, 1990 – 1991, with 1991 being plotted here)..."

Figure 9. Please give the location of the grid cell, (lon, lat).

This has been added to the caption of Figure 9.

Figure 9, Caption: "(100° west and 47° north)"

Figure 10. please add units.

Added.

Figures, general comment:

About half way through the figures the colour scheme changes. In figures 2,3,5 and 6 MO is red, RH blue and SF is green. Later in figures 8 and 9, MO is black, RH is still blue but SF is now red. For the quick skimming reader, the assumption is that red is the benchmark scheme. It's a bit confusing.

All figures have been updated to remain consistent with the color schemes (MO red, RH blue, SF green).

Anonymous Referee #2

Received and published: 21 March 2018

Brown-Steiner et al. have performed an evaluation of the performance of a model of atmospheric chemistry run with three different chemical mechanisms to understand how big an impact there is by choosing a different chemical mechanism (network). This is an interesting paper, an important bit of science, and one of only a few examples in the literature to do this sort of work in a 3D sense. Evaluating the performance of these different mechanisms enables sound conclusions to be drawn about their utility. The aim is to see if a very simple mechanism, which would enable much longer (or many more) integrations as solving the coupled ODEs in the chemical network is computationally very expensive, is suitable. This is a very well written paper and I could hardly spot any typos or grammatical errors above those spotted by reviewer #1. However, I would like to see further experiments performed before I would recommend that this be published. As it stands, I don't think the suitability of the Super Fast (SF – simple chemistry) scheme to be used beyond a present day set up has been demonstrated.

And I think this is key for the argument that the SF scheme is suitable. The present work focuses on fairly long integrations (25 years or so), where anthropogenic emissions are fixed but

interactive biogenic emissions can change as the meteorology in the model changes. Some very nice analysis is then performed against surface observations of O₃ which emphasises that for these conditions, the SF scheme performs well – in accord with the other more complex schemes.

But, to be convinced that the SF scheme is suitable for long simulations of transient forcing, I would like to see simulations that test the response of the chemical schemes to the sorts of changes that have happened over the Anthropocene and for which the SF scheme may well end up being used for within CEMS (i.e. CMIP/AerChemMIP type experiments). For example, I would like to see, as a minimum, a set of simulations using ACCMIP pre-industrial emissions (you could keep the meteorology fixed as it is if that makes things easier) so that we can see what happens between these different schemes when they are perturbed with significant changes in NO_x and VOC (CH₄).

The Super-Fast mechanism was included in ACCMIP studies in both historical and future conditions (as reviewed in the Supplemental Material), although a full description is not in the scientific literature. We hope this manuscript allows other researchers to more easily find and utilize the SF mechanism and that this manuscript can serve as a baseline for future simulations and testing.

As this study is intended as a demonstration of the Super-Fast mechanism's utility, rather than a comprehensive evaluation, and as we have only analyzed present-day, we have added "Present-Day" to the manuscript title "Evaluating Simplified Chemical Mechanisms within Present-Day Simulations of CESM ..."

Demonstrating the capabilities of the Super-Fast mechanism for long-term transient forcing was beyond our scope and capabilities, and we agree that more work needs to be done with the Super-Fast mechanism (as well as the Reduced Hydrocarbon mechanism), and that care should always be taken when any model component is utilized outside of previously demonstrated periods and conditions. As such, we have made sure that the mechanism files are available (see reproducibility comment below), and we are in discussion as how to best integrate these mechanisms into available forms at the NCAR/CESM website.

In light of this, we have added language that makes clear the limitations of this manuscript and future research directions which will be needed to further study the utility and capabilities of the Super-Fast and Reduced Hydrocarbon mechanisms, as well as made clear where the code can be accessed (see comment below).

Abstract: "Here we present and compare three 25-year present-day offline simulations..."

Page 14, Lines 27-28: "...we have compared three chemical mechanisms of different levels of complexity within the CESM CAM-chem framework for present-day chemical and climatological conditions."

We have also added language in the conclusions highlighting that this study is only for present-day conditions:

Page 14, Lines 36-40: *“We examine present-day chemistry with MO, RH, and SF. Both MO and SF have been compared in other model intercomparisons, including for preindustrial conditions (see the Supplemental Material for additional information). We hope that the analysis presented in this paper, and the availability of the mechanism files (Supplemental Material) will provide a baseline for continuing research of both the RH and SF mechanisms.”*

I also am a bit concerned with the reproducibility of these experiments outside of the team working on this. There are no mentions of compsets (is that the word used within the CESM model set ups? I’m going from here

http://www.cesm.ucar.edu/models/cesm1.2/casename_conventions_cesm.html)

that have been used. Citing some rather old papers as the sources of the rate constants and reactions used, for example in the Reduced Hydrocarbon scheme, makes it difficult for others to test the schemes without large potential for making translational errors (I know, I have made many myself!). I would propose that the mechanism data files be made available (perhaps in a simple scv format?) or at least the compsets for these experiments be made available so that others can perform their own tests. GMD is a journal dedicated to holding high standards with code and I think that the mechanisms should be treated as a bit of complex code that should be archived in order to be more easily tested. This would be desirable but I can appreciate that this may not be top priority.

We have added to the data uploaded to the archive (<http://dspace.mit.edu/handle/1721.1/114993>) the chemical mechanism input files (reduced_hydrocarbon.in and superfast.in) and add additional text to the methods section and code availability section to direct readers to various CESM/NCAR resources. We clarify that we use the FMOZSOA compset for the MO simulation and make modifications to the chemical mechanism input file and speciation of species, as described in the text.

Page 6, Lines 22-25: *“The chemical mechanism input files for MO is available in the standard CESM release (<http://www.cesm.ucar.edu/models/cesm1.2/>) and the chemical mechanism input files used for RH and SF are archived (see section on Code Availability)”*

Page 6, Lines 36-39: *“We use the FMOZSOA compset (see <http://www.cesm.ucar.edu/models/cesm1.2/cesm/doc/modelnl/compsets.html>) and make modifications to the chemical mechanism input files (see section on Code Availability) and emission files for the following mechanisms.”*

Code Availability: *“The chemical mechanism files for both RH (reduced_hydrocarbon.in) and SF (superfast.in) are included in the Supplemental Material.”*

Supplemental Material: *“The SF mechanism is in the CESM code archive as an unsupported chemical mechanism, which can be activated using the option ‘-chem super_fast_llnl’.”*

Minor corrections/comments:

Page 8, line 23: A key conclusion of Squire et al. was that sign of the response to changes in emissions of isoprene was different in SF compared to more complex schemes traceable to our best understanding of the chemistry of isoprene (ie. The MCM). I think this needs to be acknowledged here in addition to current acknowledgement that "there are biases in regions of high biogenic chemistry".

We have added language to this section to highlight this Squire et al. (2015) conclusion:

Page 8, Lines 37-40: *“Schnell et al. (2015) also conclude that the SF mechanism responds differently than other more complex mechanisms, particular under different O_x production regimes (e.g. SF shows a net increase in O_x production when isoprene emissions increase in NO_x-limited regions, which the other mechanisms show a net decrease, or little change).”*

I note from Table S2 and from the discussion in Squire et al., that the SF scheme does not include NO₃. Presumably the bias in isoprene at night (Figure 9) could be solved by simulating NO₃ in the SF scheme? Have the authors considered this? It was not clear from the manuscript if that was tested in addition to the nice tests looking at the impacts of adding in PAN and N₂O₅.

The authors did not test the addition of NO₃ to the Super-Fast scheme. The addition of the PAN and N₂O₅ sensitivity tests are intended primarily as a demonstration of the type of simulations and sensitivity studies that the Super-Fast mechanism allows for.

To this point, we have added language in the manuscript that speculates about the addition of NO₃ to the Super-Fast mechanism:

Page 15, Lines 16-18: *“The SF mechanism does not include NO₃, which may also explain some of the nighttime biases. Future simulations in which NO₃ chemistry is added to the SF mechanism may correct some of these biases.”*

Table S2, reaction 14: "idential" should be "identical".

Corrected.

Figure 8: Axis labels are way too small. Please make bigger. As above for Figure 9.

Figure 8 font sizes have been increased, and we will work with the editors to see if this plot can be included as a full page. If not, we will work with the editors to make sure they are readable. Figure 9 font sizes have been increased.

How are the VOC emissions dealt with between the different schemes? I presume that there are different amounts of VOC that go into the simulations? Please can you clarify the magnitude and distribution amongst molecules of the VOC emissions. Emissions are a key part of the chemical mechanism in my opinion.

Supplemental Table S1 includes the mapping of VOC species from MO to RH (which is discussed in section 2.2.2). For SF, we mapped only the MO ISOP species directly to the SF ISOP species. We have also added some text discussing this point:

Page 8, Line 9: “*We map the MO isoprene directly to the single SF isoprene species (ISOP).*”

Evaluating Simplified Chemical Mechanisms within Present-Day Simulations of CESM Version 1.2 CAM-chem (CAM4): MOZART-4 vs. Reduced Hydrocarbon vs. Super-Fast Chemistry

5 Benjamin Brown-Steiner^{1,2,3}, Noelle E. Selin^{3,4,5}, Ronald Prinn^{2,3,5}, Simone Tilmes⁶, Louisa Emmons⁶,
Jean-François Lamarque⁶, Philip Cameron-Smith⁷

¹Now at Atmospheric and Environmental Research, 131 Hartwell Avenue, Lexington, MA 02421-3126

²Center for Global Change Science, Massachusetts Institute of Technology, 77 Massachusetts Ave,
Cambridge, MA 02139

10 ³Joint Program on the Science and Policy of Global Change, Massachusetts Institute of Technology, 77
Massachusetts Ave, Cambridge, MA 02139

⁴Institute for Data, Systems, and Society, Massachusetts Institute of Technology, 77 Massachusetts Ave,
Cambridge, MA 02139

⁵Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77

15 Massachusetts Ave, Cambridge, MA 02139

⁶Atmospheric Chemistry Observations and Modeling Lab, National Center for Atmospheric Research,
3450 Mitchell Lane, Boulder, CO 80301

⁷Lawrence Livermore National Laboratory, 7000 East Ave, Livermore, CA 94550

Correspondence to: Benjamin Brown-Steiner (bbrownst@aer.com)

20

Abstract. While state-of-the-art complex chemical mechanisms expand our understanding of atmospheric chemistry, their sheer size and computational requirements often limit simulations to short lengths, or ensembles to only a few members. Here we present and compare three 25-year [present-day](#) offline simulations with chemical mechanisms of different levels of complexity using CESM Version 1.2 CAM-chem (CAM4): the MOZART-4 mechanism, the Reduced Hydrocarbon mechanism, and the Super-Fast mechanism. We show that, for most regions and time periods, differences in simulated ozone chemistry between these three mechanisms is smaller than the model-observation differences themselves. The MOZART-4 mechanism and the Reduced Hydrocarbon are in close agreement in their representation of ozone throughout the troposphere during all time periods (annual, seasonal and diurnal). While the Super-Fast mechanism tends to have higher simulated ozone variability and differs from the MOZART-4 mechanism over regions of high biogenic emissions, it is surprisingly capable of simulating ozone adequately given its simplicity. We explore the trade-offs between chemical mechanism complexity and computational cost by identifying regions where the simpler mechanisms are comparable to the MOZART-4 mechanism, and regions where they are not. The Super-Fast mechanism is three times as fast as the MOZART-4 mechanism, which allows for longer simulations, or ensembles with more members, that may not be feasible with the MOZART-4 mechanism given limited computational resources.

Copyright Statement

- Authors retain the copyright of the article. Regarding copyright transfers please see below.
- Authors grant Copernicus Publications an irrevocable non-exclusive license to publish the article electronically and in print format and to identify itself as the original publisher. Authors grant Copernicus Publications commercial rights to produce hardcopy volumes of the journal for sale to libraries and individuals.
- Authors grant any third party the right to use the article freely as long as its original authors and citation details are identified.
- The article is distributed under the Creative Commons Attribution 4.0 License. Unless otherwise stated, associated published material is distributed under the same license.

1 Introduction

The anthropogenic influence on atmospheric chemistry is apparent at all spatial and temporal scales: human emissions have impacted local and very short-lived species (e.g. OH, see Prinn et al., 2001), very long-lived greenhouse gases (e.g. Collins et al., 2006) and everything in between (e.g. Baker et al., 2015; Solomon et al., 2016). Over the past decades, all three branches of modern atmospheric chemistry research (Abbatt et al., 2014) – observations, laboratory analysis, and modeling – have increased in both their sophistication and their capability to explain the chemistry of our atmosphere. However, while observational networks have significant growth potential (e.g. Sofen et al., 2016), and laboratory analysis still has significant challenges to overcome (Bocquet et al., 2015; Burkholder et al., 2017), chemistry modeling efforts are finding their growth potential ~~is~~-limited by the level of chemical complexity that can be included in models due to the constraint of the computational capabilities of even state-of-the-art supercomputers (Stockwell et al., 2012). Simulations that attempt to include all known species and reactions, such as the National Center for Atmospheric Research (NCAR) Master Mechanism (Madronich and Calvert, 1989; Aumont et al., 2000) or the Leeds Master Chemical Mechanism (Jenkin et al., 1997; Saunders et al., 2003), and even some species and reactions that have not been tested in any laboratory (e.g. Aumont et al. 2005; Szopa et al., 2005), are often limited to box-model level analysis (e.g. Emmerson and Evans, 2009; Squire et al., 2015). Modeling efforts that simulate regional- and global-scale atmospheric chemistry are forced, out of practical necessity, to utilize simplified, reduced form, and parameterized chemistry in order to address the large spatial and long temporal scales needed for ~~much~~-policy-relevant research.

Historically, as computational capacity has increased, modeling efforts have tended to maximize model resolution and complexity. This limits the capability to perform multi-scenario or multi-model ensembles to institutions with access to significant computational capabilities and storage. One way to increase the number of scenarios, or members, in an ensemble is to reduce the complexity of the chemical mechanism. This selection of a reduced-form chemical mechanism for different applications, and the advantages of the increased computational efficiency of a simplified mechanism, is the main focus of this paper. While there is a long history of publications (see Dodge, 2000) that compare different photochemical mechanisms within box models (e.g. Milford et al., 1992; Jimenez et al., 2003; Emmerson and Evans, 2009; Knote et al., 2015), studies that compare multiple mechanisms within a single 3D global model are rare (e.g. Squire et al., 2015). This study examines three chemical mechanisms within the Community Earth System Model Community Atmosphere Model with Chemistry Version 1.2 (CESM1.2 CAM-chem; Lamarque et al., 2012) framework: the MOZART-4 mechanism, the Reduced Hydrocarbon mechanism, and the Super-Fast chemical mechanism (described in Section 2), which is one of the simplest representations of atmospheric chemistry in the published literature.

This study examines the trade-offs and possibilities that arise from selection of a chemical mechanism that is simple enough to be computationally efficient – and thus capable of long simulations, or large ensembles at the global scale – as well as sophisticated enough to simulate the major features of tropospheric chemistry at the local and regional scale. Many climate studies include little to no chemistry, or prescribed chemistry, even though chemistry-climate feedbacks are well established to impact global and regional climate (e.g. Marsh et al. 2013; Fiore et al., 2015). Indeed, coarse grid (2°x2.5°) chemistry-climate studies which conduct 1,000 or more years of simulations using complex

chemistry are notable in their rarity (~~for notable exceptions~~, see Barnes et al., 2016 and Garcia-Menendez et al., 2015, 2017). This paper focuses on three primary lines of inquiry focusing on tropospheric ozone. First, what is lost or gained with the selection of a simplified chemical mechanism within a global model? Second, what is the nature of the uncertainties that arise with the selection of a particular chemical mechanism? And third, what are the tradeoffs that researchers make, either intentionally or tacitly, when they apply ~~that a specific~~ mechanism within a particular modeling framework? We focus this study on the short-lived gaseous species, in particular ozone and its precursors, that influence both the daily exposure of humans to pollutants as well as the decadal-scale global climate system. We focus primarily on a computationally efficient simulation of tropospheric gaseous chemistry within a single modeling framework, and leave further analysis of other aspects of atmospheric chemistry to future studies.

In Section 2, we describe the modeling framework, and describe each of the three aforementioned chemical mechanisms, including a detailed description and history of the Super-Fast mechanism, as it is not reported elsewhere in the literature, and the simulations and observations we use for comparison. In Section 3 we present spatial and temporal results, as well as compare various metrics of chemical accuracy. In Section 4, we explore the nature and the morphology of the chemical uncertainties, and the particular tradeoffs that are made by the selection of a single mechanism when faced with limited computational resources. We draw conclusions in Section 5.

2 Methods

Our analysis focuses on characterizing the ozone chemical uncertainties within a global chemistry model. We examine the morphology of the chemistry system, focusing specifically on the means, standard deviations, and variability (defined here as the standard deviation divided by the mean). We also include characterizations of the correlation of the ozone time series with the observations and of the extreme values (in particular the 90th and 99th percentiles) of the ozone distribution.

2.1 CESM1.2 CAM4-chem Simulations

The CESM1.2 CAM4-chem model (Tilmes et al., 2015; 2016) is a chemistry-climate model developed at the National Center for Atmospheric Research (NCAR) with other collaborators, including the U.S. Department of Energy. It has been utilized extensively in the Atmospheric Chemistry and Climate Intercomparison Project (ACCMIP) (Lamarque et al., 2013 and references therein), the Chemistry Climate Model Initiative (CCMI) (Morgenstern et al., 2017) and for a wide range of atmospheric chemistry research. We conduct our simulations using CESM CAM4-chem version 1.2 with the MOZART-4 chemical mechanism based on Emmons et al. (2010) with updates described in Tilmes et al. (2015), the Reduced Hydrocarbon mechanism (Houweling et al., 1998) as adapted to the CESM CAM-chem framework by Lamarque et al. (2008, 2010), which has a reduced form representation of hydrocarbon chemistry, and the Super-Fast mechanism (Cameron-Smith et al., 2006; Lamarque et al. 2013). Hereafter we will refer to these three mechanisms as MO, RH, and SF, respectively.

For meteorology we used the Modern-Era Retrospective analysis for Research and Applications (MERRA) reanalysis product (Rienecker et al., 2011) for ~~25-26~~ years (1990 – ~~2014~~2015), with a 50-hour Newtonian relaxation timing (roughly 1% nudging every 30 minutes). The year 1990 is dropped to allow for spin-up. All simulations are at 1.9°x2.5° resolution. Aerosols were represented by the bulk aerosol model (BAM) in the MO and RH mechanisms and is optional for the SF mechanism. The results presented here are without BAM aerosols. We keep anthropogenic emissions constant at year-2000 from the CCMI database (Lamarque et al., 2012) and include linearized chemistry for ozone in the stratosphere (McLinden et al., 2000; Hsu and Prather, 2009), and prescribe the concentration of other tracers above 50 hPa. We use an online biogenic emissions model (MEGAN; Guenther et al., 2012), and prescribed sea ice and sea surface temperatures. With the exception of a remapping of the MOZART species to the Reduced Hydrocarbon species (Supplemental Table S1), all parameterizations other than the chemical mechanism are identical between the three simulations, and thus any differences are due to differences among the mechanisms themselves. Ozone dry deposition was done as described in Val Martin et al. (2015). Because we run with prescribed meteorology, we do not include internal chemical feedback to the weather and climate other than that incorporated into the MERRA meteorology itself. All of these mechanisms can also be run with meteorology calculated internally by the CESM model, but since such simulations utilize a different number of vertical levels than simulations with prescribed meteorology, comparing to simulated meteorology runs is not straightforward, and so is omitted from the present study.

2.2 Mechanisms

Table 1 summarizes the characteristics of the three chosen mechanisms. The chemical mechanism input files for MO is available in the standard CESM release (<http://www.cesm.ucar.edu/models/cesm1.2/>) and the chemical mechanism input files used for RH and SF are archived (see section on Code Availability).

2.2.1 MOZART-4 (MO)

The Model for Ozone and Related Chemical Tracers, version 4 (MOZART-4) mechanism (Emmons et al., 2010; Lamarque et al., 2012; Tilmes et al., 2015) is the standard tropospheric chemical mechanism used within the CESM CAM-chem framework (Tilmes et al., 2015; 2016). It has been used in many model inter-comparison projects (e.g. Lamarque et al., 2013; Emmons et al., 2015), and extended to tagged tracer chemistry (Emmons et al., 2012). As described in detail in Emmons et al. (2010), MOZART-4 mechanism is a tropospheric mechanism that contains 85 gas-phase species and 12 bulk aerosol species, with 39 photolysis and 157 gas-phase reactions. Large alkanes, alkene, and aromatics are lumped together (BIGALK, BIGENE, and TOLUENE, respectively), and monoterpenes are lumped together as C₁₀H₁₆ and treated as α -pinene. We use the FMOZSOA compset (see <http://www.cesm.ucar.edu/models/cesm1.2/cesm/doc/modelnl/compsets.html>) and make modifications to the chemical mechanism input files (see section on Code Availability) and emission files for the following mechanisms.

2.2.2 Reduced Hydrocarbon (RH)

The Reduced Hydrocarbon (RH) chemical mechanism (Houweling et al., 1998; Lamarque et al., 2010) is a reduced-form mechanism based on the Carbon Bond Mechanism 4 (CBM-4) (Gery et al., 1989). The CBM-4 was developed to simulate polluted regional chemistry, and the RH mechanism updated and expanded this mechanism to also be capable of simulating background low-NO_x conditions (Houweling et al., 1998). As described in Houweling et al. (1998), the original RH mechanism has 30 tracers and 68 total reactions. It has been used extensively in model inter-comparisons (e.g. Pöchl et al., 2000) and is generally considered a satisfactory reduced hydrocarbon mechanism (e.g. Hauglustaine et al., 1998; Wang and Prinn, 1999; Granier et al., 2000; Pfister et al., 2014). Lamarque et al. (2008) incorporated the RH mechanism into the CESM CAM-chem framework with a few updates, and Lamarque et al. (2010) expanded it to 89 (to include the bulk aerosol model species) tracers and 202 total reactions. As the lumping of alkanes and alkenes in RH differs from the MO mechanism, a mapping between the differently aggregated species is necessary (see Supplemental Table S1).

For this work, we modified the RH mechanism to remove many of the tracers and reactions that are pertinent primarily to stratospheric chemistry (as introduced in Lamarque et al., 2008) since these simulations include specified long-lived stratospheric species (O₃, NO_x, HNO₃, N₂O, N₂O₅) as in MOZART-4 (Emmons et al., 2010). However, the unmodified RH mechanism can be run with the more complex stratospheric chemistry, but at a significant additional cost. This is not considered in this paper to allow a better comparison between the tropospheric-only mechanisms. The modified RH mechanism, which shows only minor differences in the simulated surface ozone concentration from the complete mechanism (not shown), contains 65 tracers and 127 reactions. This RH mechanism runs approximately twice as fast than the MO mechanism under our current configuration (Table 1).

2.2.3 Super-Fast (SF)

The Super-Fast (SF) mechanism is a highly simplified chemical mechanism designed to efficiently simulate background tropospheric ozone chemistry (Cameron-Smith et al., 2006, and supplementary material of Lamarque et al., 2013). It includes 15 chemical tracers with 6 photolysis reactions and 24 gas phase reactions, making it the simplest chemical mechanism to be included as a member of the ACCMIP ensembles (Lamarque et al., 2013). It was developed by the Lawrence Livermore National Laboratory (LLNL) and has not been described as implemented within the CESM code, so we include a description here and in our Supplementary Material.

Supplemental Table S2 summarizes the SF mechanism photolysis and gas-phase reactions, which consist of a basic methane oxidation scheme (CH₄, CH₃O₂, CH₃OOH, CH₂O, and CO), with basic oxidant chemistry (OH and O₃), along with simple sulfur chemistry (dimethyl sulfide (DMS), SO₂, and SO₄) and a single biogenic hydrocarbon species, isoprene (ISOP), with two oxidant pathways: ISOP + OH and ISOP + O₃. For reactions iii, vi, 10, 11, and 15 (Table S2), it is assumed that their products O, H, and CH₃OH are instantaneously converted to their ultimate products, O₃, HO₂, and HO₂, respectively. Nitric acid chemistry is limited to two reactions, one of which requires a heterogeneous reaction parameterization. Sulfur chemistry is limited to four reactions. Isoprene chemistry is highly parameterized. The reaction of isoprene with OH is based on the net effect of the reaction in the University of California Irvine (UCI) model (Wild and Prather, 2000), namely: ISOP + 2.5*OH →

2*CH₃O₂O. This particular parameterized reaction, which when originally implemented used a negative coefficient among the products (ISOP + OH → 2*CH₃O₂ - 1.5*OH) during the original implementation required a negative coefficient for the OH product, is not standard within the CESM chemical modeling framework and cannot be handled by the solver, so the equivalent triple reaction formulation of 21a, 21b, and 21c is required. The oxidation of isoprene by ozone is a simple parameterization (resulting in fractional production of only the species that already exist in the mechanism as part of the methane oxidation scheme: CH₂O, CH₃O₂, HO₂, and CO) derived from the net effect of the isoprene/ozone oxidation pathways from the full LLNL-IMPACT model (Rotman et al., 2004) and was included specifically to improve the simulation of surface ozone chemistry (Cameron-Smith et al., 2009). We map the MO isoprene directly to the single SF isoprene species (ISOP).

Much of the simplicity within the SF mechanism comes from what it does not include. Carbon chemistry is limited to the five single-carbon species used in the simple methane oxidation scheme, plus isoprene. There is no PAN (peroxy acetyl nitrate) or ammonia, and hence no nitrogen aerosols, although HNO₃ is created in reaction 8 and 16. These all impact ozone chemistry, but the inclusion of additional hydrocarbon, aerosol, or heterogeneous chemistry would introduce significant additional computational costs (similar to the more complete mechanisms). There are no halogen species, since this would require the inclusion of a significant number of additional chemical tracers, and as such there is no capability to describe the polar ozone hole phenomenon within the mechanism (Cameron-Smith et al., 2006), so it is implemented within Linoz using the simple loss parameterization of Cariolle, et al., (1990). The greatest simplifications in the SF mechanism arise from compacting all of the non-methane hydrocarbon chemistry (NMHC) into two isoprene reactions, and thus there is none of the complex chemistry that is required to adequately represent ozone chemistry in highly polluted regions. The simplicity of the SF mechanism allowed us to perform two short simulations in which we added reduced-form PAN and N₂O₅ chemistry (individually, and in conjunction) from the MOZART-4 mechanism into the SF mechanism, which we use as a demonstration of the type of sensitivity tests that are possible with the SF mechanism. This type of quick sensitivity test would be significantly more difficult with the more complex mechanisms, given the complexity of PAN and N₂O₅ chemistry.

The SF mechanism has been included in several model inter-comparison projects. We include an expanded review in the Supplemental Material. Unfortunately, the SF mechanism only simulates sulfate aerosol, and so was unable to be compared to the aerosol simulations of the other ACCMIP members (Lamarque et al., 2013). The SF simulations within ACCMIP demonstrated lower rates of ozone chemistry and deposition resulting in a low ozone burden bias and a high ozone lifetime bias (Young et al., 2013), and while projected changes in ozone radiative forcing fell within the ACCMIP range, the historical changes did not (Stevenson et al., 2013). Human health analysis with the SF simulations fell within the range of the other ACCMIP members (Silva et al., 2013; 2016; 2017). Squire et al. (2015) compared SF to more complicated isoprene schemes, and concluded that including the SF mechanisms is preferable to neglecting chemistry entirely, although there are biases in regions of high biogenic chemistry. Schnell et al. (2015) also conclude that the SF mechanism responds differently than other more complex mechanisms, particular under different O_x production regimes (e.g. SF shows a net increase in O_x production when isoprene emissions increase in NO_x-limited regions, whereas the other mechanisms show a net decrease, or little change). Finally, Schnell et al. (2015) compare seasonal and diurnal cycles to other mechanisms, and the SF mechanism simulates high ozone events in the

Formatted: Subscript

Comment [NES1]: I think this ought to be rephrased – I know I commented on this earlier but OH is a reactant not a product in this reaction.....

springtime, and that the SF mechanism outperforms others when compared to the observed summertime diurnal cycle.

2.3 Computational Requirements

5 The computational requirements of MO, RH, and SF as simulated on the NCAR Cheyenne
supercomputer are summarized in Table 1. The computational cost results from both the chemical
solver and the advection of the chemical tracers within CAM-chem. No load balancing was conducted,
which could potentially increase the efficiency of the RH and SF mechanisms. The CESM1.2 CAM-
chem model run with the SF mechanism is roughly three times faster than a run with the MO
10 | mechanism when ~~the Bulk Aerosol Model (BAM)~~ (see Tilmes et al., 2015) ~~BAM~~ aerosols are included
(which we do not examine in this present study), and a gas-phase-only simulation with the SF
mechanisms increases the speeds to nearly 4 times as fast. The RH mechanism is roughly twice as fast
as the MO mechanism. At higher spatial resolutions, and the computational advantage of the SF
mechanism over the more complex MO and RH schemes is likely to increase, since advection of tracers
15 typically becomes a larger fraction of the total model run-time.

2.4 Observations

 The ozone observational databases are of two types: the global database is ozonesonde data
compiled from Tilmes et al. (2012) while the US database comes from the EPA Clean Air Status and
20 Trends Network (CASTNET), which has more than 90 surface observational sites within the United
States and has been collecting surface meteorological and chemical data since 1990 (CASTNET, 2016
and <https://www.epa.gov/castnet>). We used data from all sites that reported complete ozone data from
each year, after removing data that the CASTNET database marked as invalid. The number of sites that
matched these criteria varied from year to year, but generally we have between 55 and 94 sites
25 | throughout the ~~1990-1991~~ – ~~2014~~ 2015 period. The CASTNET observational network is located
primarily in rural sites, and thus is a reasonable comparison to coarse grid cell model output e.g. Brown-
Steiner et al., 2015; Phalitnonkiat et al., 2016. In order to compare to the CESM CAM-chem
simulations, which has no emissions trend, we have detrended the CASTNET data for each region using
a simple linear regression. Regional averaging is first done by averaging all observational sites within a
30 single 1.9°x2.5° grid cell, and then averaged to the larger regions as needed. We also compare to ozone
precursor species observations from Tilmes et al., (2015).

3 Results

3.1 Spatial Comparisons

35 The spatial distribution of ozone and related species between the three mechanisms are
compared in Figure 1. Taylor-like diagrams comparing results to ozonesondes over different global
regions are provided in Figure 2, and comparisons to aircraft observations in Figure 3. Globally
averaged surface Daily Maximum 8-Hour (~~DM8HMDA8~~) O₃ is consistent across all mechanisms

(Table 2) with the largest spatial differences (especially with the SF mechanism) noted over regions of intense biomass burning or biogenic emissions, such as equatorial Africa and South America, as well as over northern hemisphere oceans within SF (Figure 1). Surface CO mixing ratios show small regional differences between MO and RF, while NO_x mixing ratios show very small and highly localized differences (Figure 1). All three mechanisms tend to have low CO biases over much of the northern hemisphere, with SF showing the largest bias. This coincides with starkly higher NO_x mixing ratios in the northern hemisphere (Figure 1, Figure 3), especially in the winter and spring seasons. This is explored in more detail below.

Zonal profiles (Figure 4) show that ozone is similar among all mechanisms for all seasons, especially in the lower troposphere. Compared to the MO mechanism, the SF mechanism simulates higher northern hemisphere ozone in the winter, and lower in the summer. Both the RH and SF mechanisms simulate lower CO mixing ratios than the MO mechanism in both the summer and winter, with the SF mechanism diverging the most in the northern hemisphere in the summer. The SF mechanism also simulates higher NO_x in the northern hemisphere winter, which (as we explore below) may in part be due to the lack of PAN chemistry.

At the largest spatial scales, all three mechanisms predict similar levels of surface ozone (Figure 5, Table 2), with global surface ozone estimates of 32.6 ± 0.93 , 33.9 ± 0.98 , and 31.5 ± 1.12 ppb for MO, RH, and SF, respectively. Even at the Continental US scale, all three mechanisms estimate similar surface ~~DM8HMDA8~~ O₃ values (56.7 ± 3.08 , 57.7 ± 3.23 , and 53.4 ± 3.59 ppb for MO, RH, and SF, respectively), which are consistent with the CASTNET observations of 56.1 ± 5.65 ppb. However, within the Northeastern US, the well-known high bias is apparent (74.4 ± 11.4 , 76.0 ± 11.9 , 72.6 ± 14.5 ppb for MO, RH, and SF, respectively, while the CASTNET observations are 57.4 ± 7.42 ppb). The MO and RH mechanisms are nearly identical at all spatial scales, while the SF mechanism simulates larger ~~DM8HMDA8~~ O₃ variability, especially at individual grid cells within the Eastern US. Taking into account the model ozone biases, the SF is a better characterization of the ozone distribution (as compared to CASTNET) for almost every spatial scale examined within the US. Indeed, in the Southeastern US, where we expect SF to perform poorly due to the simplified biogenic species chemistry, we actually find that the SF estimates the shape of the high ozone tail better than either MO or RF: CASTNET estimates at an individual grid-cell, that the 99th percentile for ~~DM8HMDA8~~ O₃ is 18% higher than the 90th percentile (Table 2), and while MO and RH estimate only 14% higher and 14% higher, respectively, the SF estimates 29% higher. In Section 4, we explore some of the implications of these differences, and in particular whether the biases within the SF mechanism are of the same magnitude as some of the biases within the MO and RF.

Figure 6 explores this finding, which plots the percentage difference between the 99th and the 90th percentile ozone as the length of the time series included grows. This comparison allows for a comparison of the relative distribution among mechanisms, here for the higher end of ozone values, to compare the overall shape of each mechanism's distribution when biases in the magnitudes are normalized. We note that: (1) it takes between 5 and 10 years before a consistent and stable estimate emerges with each simulation, indicating that simulations less than 10 years may be inadequate for comparisons between chemical mechanisms; (2) the CASTNET observations have a transient estimate, most notably in the Southeastern US, which indicates a divergence of the 99th and the 90th percentiles (i.e. a lengthening of the upper tail) that is not seen in the simulations; and (3) the SF mechanism is

inconsistent with the MO and RH mechanisms, which are nearly identical, but the SF mechanism estimate is also closer to the CASTNET estimate in the Midwestern and Southeastern US. Whether this is the result of fortunate biases within the SF mechanism or an implication that the more complex chemistry within the MO and RH mechanisms are underestimating the length of the ozone tail requires further study. Brown-Steiner et al. (in review, ACPD) examines these implications, and also concludes that it takes approximately 10 years for long-term signals to emerge from meteorological variability. These results demonstrate the challenge in examining chemical signals in highly variable data, particularly if there are trends or changes to the ozone distribution, as is seen in the CASTNET data for the Southeastern US.

However, while the SF mechanism performs as well as, or better than, the MO and RH mechanisms in certain regions, there are many regions – especially in the northernmost latitudes over land, and over equatorial land masses – where the SF mechanism is far less capable at simulating surface ozone than either the MO or RH mechanisms. Figure 7 plots R^2 values for the ~~DMSHMDA8~~ O_3 JJA time series (1990 – 2015) at every grid cell between the MO mechanism and both RH and SF, and it is clear that the RH mechanism has very high R^2 values ($R^2 > 0.75$) over much of the globe. And while the SF mechanism has large R^2 values over many regions – in particular the extratropics – over the equatorial regions, and especially over land, R^2 values drop below 0.5 and even 0.25.

3.2 Seasonal and Diurnal Comparisons

The seasonality of surface ozone is similar among all three mechanisms at the regional-scales (Figure 8), although differences occur at both the largest and smallest scales: (1) the SF mechanism simulates a dual-peaked maximum in surface ozone averaged at the global scale, a phenomenon also noted by Schnell et al. (2015); (2) this dual-peaked maximum is still apparent at the regional scales, although to a much lesser degree; and (3) the RH mechanism has a dual-peaked maximum over portions of the Southeastern US. The seasonal patterns for CO and NO_x are consistent across all models, although CO is lower in both RH and SF than in MO for all seasons. RH and MO NO_x levels are nearly identical, but SF simulates higher values for NO_x in all seasons, and particularly in the winter and spring seasons, as already noted. HO_x and isoprene seasonality is consistent across all mechanisms at most scales.

Diurnal cycles are compared for a single grid cell within the Central US in Figure 9. With the exception of isoprene within the SF mechanism, which does not adequately represent nighttime isoprene chemistry, the diurnal cycles are comparable across all mechanisms for most species. The MO and RH mechanisms are nearly identical, with the exception of CO values, as already mentioned. The SF mechanism tends to show more extreme peaks in OH and NO_x , and lower levels of O_3 , CO, H_2O_2 , and SO_4^- (Figure 9). Surface levels of O_3 and CO within the SF mechanisms are sensitive to the addition of PAN and N_2O_5 chemistry (the dotted lines in Figure 9), described below, although the sensitivity tends to be in the simulated magnitude and not the shape of the diurnal cycle.

Figures 8 and 9 also include two-year simulations (1990 – 1991, with year 2000 emissions) in which we included into the SF mechanism PAN and N_2O_5 chemistry taken (and reduced) from the MOZART-4 mechanism. We examine these mainly to demonstrate the potential for the modification of the SF mechanism to meet particular research needs. Largely, the addition of PAN chemistry (purple

lines) results in more substantial changes to various species than the addition of N_2O_5 chemistry (orange lines), but their combined addition (green lines) slightly modifies the simulated large-scale values of O_3 , CO , HO_x , and isoprene. The addition of PAN chemistry brings the SF mechanism simulations closer to the MO mechanism for the NO_x and HO_x seasonal cycles (Figure 8), and the CO diurnal cycle (Figure 9), but at the expense of the global-scale capability to simulate ozone and isoprene. Additional tuning of the parameterized reactions 21 and 22 (Table S2) may be able to correct these errors. Sulfate aerosol in the SF mechanisms is notably lower than both the MO and RH mechanisms, which may result from the simple aerosol scheme within the SF mechanism.

3.3 Comparison to Observations

Figure 10 compares the model estimates of surface ozone to observations (ozonesondes and CASTNET observations) for different spatial regions, as well as to each other. Generally, all three mechanisms simulate less variability over continental to global scale regions than the ozonesonde observations (Figure 10c,d,e) and show a high bias over many sites within North America, Europe, and Asia. Within the US, all mechanisms show a high bias in the Eastern US, and especially in the Northeastern US, but the variability is well-captured when compared to CASTNET (with slopes ranging from 0.61 – 1.24 in Figures 10f, g, and h). When compared to each other (Figures 10a,b,i,j), the RH mechanism and MO mechanism are nearly identical. The SF mechanism, while comparable to the MO mechanism at many sites, shows greater divergence, overestimating values in many grid cells throughout the globe (Figure 10b) and both over- and underestimating within the US (Figure 10j). Taylor-like diagrams are plotted in [Supplemental Figure S1 Figure 2](#) and show the close clustering of the MO and RH mechanisms, and that the SF mechanism differs from the observations at a similar magnitude than the MO and RH mechanism for some regions, but performs poorly in other regions (especially in the tropics, where tropospheric ozone is underestimated with the SF mechanism).

4 Discussion

Our primary objective has been to determine what is lost (or gained) with the selection of a simplified chemical mechanism, which we summarize here. We mostly discuss the SF mechanism, as the tradeoffs with the RH mechanisms are straightforward: we lose very little (Figure 10a and 10i) and gain about a 100% increase in simulation speed (Table 1). Many of the things that are lost with the use of the SF mechanism are expected: we lose the capability to directly simulate small-scale features of ozone chemistry in regions that depend strongly on complex biogenic chemistry. In particular, the equatorial landmasses – especially equatorial Africa and South America – are not well simulated (Figure 7). We also lose the capability to simulate some of the short-term features that require additional chemistry, such as the night-time behavior of isoprene (Figure 9), or the cold season CO and NO_x behavior (Figure 1 and 4). The addition of PAN and N_2O_5 chemistry do not rectify the nighttime behavior of isoprene (Figure 9), but do bring the cold-season simulated CO and NO_x mixing ratios closer to the MO mechanism (Figure 8). These deficiencies may result from the highly parameterized biogenic chemistry within the SF mechanisms (Supplemental Table S2), although it may also result

from the treatment of isoprene emissions, and future simulations will need to consider the trade-off between additional complexity and computational efficiency.

More surprisingly, there are several desirable capabilities that are not lost with the selection of the SF mechanism. For most regions, the selection of the SF mechanism does not degrade the estimate of surface ozone (both the magnitude and the variability), nor do we lose features of the daily variability that results from the meteorology. In many regions, and at many scales, we find that the selection of the SF mechanism introduces uncertainties that are smaller than the difference between the simulated and observed surface ozone mixing ratios (Figure 5). Surface layer ozone is adequately represented over many regions in all seasons within the SF mechanism (Figure 8), despite the high CO and low NO_x levels in the winter and spring seasons (Figure 4). For these seasons, the adequate ozone representation may be the result of compensating errors, and Schnell et al. (2015) previously found comparable cases where the SF mechanism outperforms more complex models, perhaps due to various sets of compensating biases or errors.

We now turn to the main question of this research: what do we gain when we select a simplified chemical mechanism? The primary thing we gain is the capability to simulate longer periods of time, or to include more members in an ensemble, in proportion to the simplicity of the mechanism. Our results show that, without any optimization of the code, the RH mechanism is ~ 100% faster than the MO mechanism, and the SF mechanism is up to 200% faster than the MO mechanism (Table 1). We feel that the capability to run three SF simulations for the price of one MO simulation under different sets of initial conditions, for example, can extend the quantification of parametric uncertainties which is largely unavailable to the most complex and most computationally demanding mechanisms.

For instance, there are many research frameworks where the “three-for-one” advantage of the SF mechanism could be utilized with the MO mechanism to allow for an expanded exploration of parametric uncertainties that would not otherwise be available with the MO mechanism alone. One simulation of a 5- or 10-year time slice with the MO mechanisms could be combined with three simulations of the SF mechanism, one matching the parameters of the MO mechanism (in order to provide a consistent baseline), and the other two exploring other parameter spaces (e.g. different initial conditions, or different emission scenarios). The establishment of a baseline comparison is particularly important, since the SF mechanism is a simplified mechanism, and should not be blindly trusted to reproduce the behavior of more complex mechanisms. For example, if a research group is interested in precise estimates of ozone concentrations in regions where the biogenic influence is significant, the SF mechanism would prove insufficient. The RH mechanism may be sufficient, but the more modest increase in computational speed – a “two-for-one” advantage over the MO mechanism – may not be enough to justify the simulation. If, however, the phenomenon of interest can be shown to be within the SF mechanism capabilities (e.g. simulating regional-scale ozone, as shown in this paper), the “three-for-one” advantage of the SF mechanism is readily apparent. The SF mechanism may be particularly desirable with chemistry-climate simulations at higher spatial resolutions.

In addition, the selection of a simplified mechanism allows for the capability to easily and efficiently test new forms and new representations of chemistry without the need to painstakingly update and test all possible interactions of any addition within a complex mechanism. For example, in this study, we added a simplified PAN and N₂O₅ representation to the SF mechanism (Figure 8 and 9) to see how it improves the simulations. This exercise offered a significant capability to test, simulate, and

further learn about improving atmospheric chemistry computations. This demonstrates that a hybrid approach (or tiered approach, as recommended in Uusitalo et al., 2015) – in which complex and trusted chemical mechanisms are used to evaluate simplified mechanisms that can run for longer periods or with increased ensemble members – has the potential to maximize computational capabilities and to get the most out of atmospheric chemistry modeling.

Furthermore, the selection of a simple chemical mechanism – especially when used in conjunction with more complex mechanisms within a consistent modeling framework – allows for better quantification of the uncertainties, and the relative importance, of particular pieces of the chemistry. Here, for instance, the SF mechanism's representation of biogenic species chemistry is insufficient to adequately represent equatorial landmasses, but the reduced form RH mechanism is nearly as capable as the MO mechanism over most regions and most species. This begs the question: is there a representation of biogenic chemistry somewhere between the RH and the SF mechanisms that can approach the efficiency of the SF mechanism and the accuracy of the RH mechanism? We hope that future research will address this question, as well as others, such as more globally oriented research pertaining to ozone budgets and the interaction between OH and CH₄ lifetime. In addition, comparisons of chemical mechanisms of different complexities, and particularly where the simplified mechanisms fail, could potentially identify regional chemical regimes. For instance, the SF mechanism cannot adequately represent the chemistry of equatorial forests (Figure 7), and the spatial regions that fail to simulate ozone chemistry are similar to the spatial distribution of the tropical forest chemical regime identified in Figure 4 of Sofen et al. (2016), which utilized a statistical clustering technique to identify chemical regimes. Finally, the capability to examine atmospheric chemistry complexity in a step-wise fashion could also be utilized to bridge the gap between the most complex 3D chemical models and the more efficient models utilized by the Earth Models of Intermediate Complexity (EMIC) or Integrated Assessment Model (IAM) communities.

5 Conclusion

In this study, we have compared three chemical mechanisms of different levels of complexity within the CESM CAM-chem framework for present-day chemical and climatological conditions. We conducted 25-year cycled emission simulations nudged to MERRA meteorology with the standard tropospheric MOZART-4 (MO) mechanism of Emmons et al. (2010), the Reduced Hydrocarbon (RH) mechanism of Houweling et al. (1998), and the Super-Fast (SF) mechanism of Cameron-Smith et al. (2006). The RH mechanism is roughly twice as efficient as the MO mechanism, and the SF mechanism is roughly three times as efficient as the MO mechanism, without any code optimization. As much as possible, we kept the parameterizations consistent across all mechanisms, although we had to remap some of the MO mechanism species to match up with the RH mechanism species.

We examine present-day chemistry with MO, RH, and SF. Both MO and SF have been compared in other model intercomparisons, including for preindustrial conditions (see the Supplemental Material for additional information). We hope that the analysis presented in this paper, and the availability of the mechanism files (Supplemental Material) will provide a baseline for continuing research of both the RH and SF mechanisms.

We find that all three mechanisms successfully capture surface ozone values at the larger spatial scales, but at smaller spatial scales, and especially within the Northeastern US, all three mechanisms have surface ozone biases when compared to CASTNET observations, but that the mean values for all three mechanisms are consistent with each other at a variety of spatial scales. The SF mechanism simulations show larger ozone variability than the MO and RH simulations, although when normalizing the distributions to account for the known ozone biases, the SF mechanism represents the shape and spread of the ozone distributions better than the MO or RH mechanisms, when compared to the CASTNET observations (Figure 5).

The RH mechanism is in close agreement with the MO mechanism for nearly every metric we examined, and any differences tend to be minor (both in magnitude and in spatial extent). The SF mechanism simulates higher NO_x and lower CO than the MO mechanism, and the NO_x deviations are particularly large in the winter season. In addition, the SF mechanism deviates from the MO mechanism over regions of high biogenic emissions, such as equatorial Africa and South America. These large deviations within the SF mechanism are likely a result of the simplicity of the mechanism, and especially the lack of biogenic species chemistry beyond a single-species, two-reaction representation, as well as a lack of PAN and N_2O_5 chemistry (Figures 8 and 9). The SF mechanisms do not include NO_3 , which may also explain some of the nighttime biases. Future simulations in which NO_3 chemistry is added to the SF mechanism may correct some of these biases. We also find that although the SF mechanism differs in the magnitude of the estimated ozone from the other two mechanisms, the simulated ozone variability is similar in all three mechanisms (Figures 4 and 10).

We find that there are significant gains that can be realized by a research approach that utilizes simulations with both a complex and a simplified chemical mechanism where the complex mechanisms are used to provide a more-trusted chemical result (especially for the mean values) and the simple mechanism could be used to efficiently simulate longer time periods to better understand the roles of meteorological variability. The capability of the SF mechanism to simulate adequate chemistry with interactive meteorology is not examined here, nor the coupling of the SF mechanism with modal aerosols, which is left for future research. These results encourage revitalizing or creating simplified chemical mechanisms within individual modeling frameworks, and examining the structural uncertainties that exist between different models with regards to simplified chemical mechanisms.

Finally, we note that there are many inherent uncertainties associated with the use and comparison of chemical mechanisms and climate-chemistry simulations, many of which are inherited with the adoption of a particular model. The CESM CAM-chem model has been used extensively to examine a variety of climate and chemistry phenomena, and uncertainties that arise from the individual choices made during the historical development of this chemical model (see Brasseur et al., 1998; Hauglustaine et al., 1998; Horowitz et al., 2003; Kinnison et al., 2007; Emmons et al., 2010) are still present in the CESM CAM-chem modeling framework, such as which scheme or parameterization was to be included and the specific metric and methodology of tuning the climate model to historical data (see Hourdin et al., 2017 and references therein). Future simulations using different model versions, or different choices of parameterizations, schemes, emissions, and other input datasets will need to examine the impact of those choices on the simulated chemical uncertainty and compare these to the uncertainty that arises from the selection of the different chemical mechanisms presented here.

Code Availability


CESM CAM-Chem code is available through the National Center for Atmospheric Research /University Corporation for Atmospheric Research (NCAR/UCAR) website (<http://www.cesm.ucar.edu/models/cesm1.2/>), and this project made no code modifications from the released model version. **The chemical mechanism files for both RH (reduced hydrocarbon.in) and SF (superfast.in) are available on Massachusetts Institute of Technology servers at: <http://dspace.mit.edu/handle/1721.1/114993>.**

Data Availability

The raw model output is archived on the NCAR servers, and processed data ~~will be made~~is available ~~upon publication on a public recognized repository with a unique digital object identifier (doi)~~on Massachusetts Institute of Technology servers at: <http://dspace.mit.edu/handle/1721.1/114993>.

5

Supplemental Link




Author Contribution

BBS prepared and ran the simulations and prepared the manuscript under direction and advisement of NES and RP. LE and ST aided in the development, preparation, and analysis of the simulations as well as reviewed the manuscript. JFL advised and aided in the Reduced Hydrocarbon simulation. PCS
5 | advised and aided in the Super-Fast simulation. JFL and PCS also, ~~and both~~ reviewed the manuscript.

Competing Interests

The authors declare that they have no conflict of interest.



Acknowledgements

This model development work was supported by the U.S. Department of Energy (DOE) Grant DE-FG02-94ER61937 to the MIT Joint Program on the Science and Policy of Global Change. The work of PC was supported through the Scientific Discovery through Advanced Computing (SciDAC) program funded by the DOE Office of Science, Advanced Scientific Computing Research and Biological and Environmental Research, and was performed under the auspices of the DOE by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. Computational resources for this project were provided by DOE and a consortium of other government, industry, and foundation sponsors of the Joint Program. For a complete list of sponsors, see: <http://globalchange.mit.edu>. Additional computing resources were provided by the Climate Simulation Laboratory at NCAR's Computational and Information Systems Laboratory (CISL), sponsored by the National Science Foundation and other agencies. The National Center for Atmospheric Research is funded by the National Science Foundation. The authors would also like to thank Daniel Rothenberg for efficient processing of the ozone files.

References

- Abbatt, J., George, C., Melamed, M., Monks, P., Pandis, S., and Rudich, Y.: New Directions: Fundamentals of atmospheric chemistry: Keeping a three-legged stool balanced, *Atmos. Environ.*, 48, 390–391, <https://doi.org/10.1016/j.atmosenv.2013.10.025>, 2014.
- 5 Aumont, B., Madronich, S., Bey, I., and Tyndall, G. S.: Contribution of secondary VOC to the composition of aqueous atmospheric particles: A modeling approach, *J. Atmos. Chem.*, 35, 59–75, <https://doi.org/10.1023/A:1006243509840>, 2000.
- Aumont, B., Szopa, S., and Madronich, S.: Modelling the evolution of organic carbon during its gas-phase tropospheric oxidation: development of an explicit model based on a self generating approach, *Atmos. Chem. Phys.*, 5, 2497–2517, <https://doi.org/10.5194/acp-5-2497-2005>, 2005.
- 10 Baker, L. H., Collins, W. J., Olivić, D. J. L., Cherian, R., Hodnebrog, Myhre, G., and Quaas, J.: Climate responses to anthropogenic emissions of short-lived climate pollutants, *Atmos. Chem. Phys.*, 15, 8201–8216, <https://doi.org/10.5194/acp-15-8201-20>, 2015.
- Barnes, E. A., Fiore, A. M., and Horowitz, L. W.: Detection of trends in surface ozone in the presence of climate variability, *J. Geophys. Res. Atmos.*, 121, 6112–6129, <https://doi.org/10.1002/2015JD024397>, 2016.
- 15 Bocquet, M., Elbern, H., Eskes, H., Hirtl, M., Aabkar, R., Carmichael, G. R., Flemming, J., Inness, A., Pagowski, M., Pérez Camacho, J. L., Saide, P. E., San Jose, R., Sofiev, M., Vira, J., Baklanov, A., Carnevale, C., Grell, G., and Seigneur, C.: Data assimilation in atmospheric chemistry models: Current status and future prospects for coupled chemistry meteorology models, *Atmos. Chem. Phys.*, 15, 5325–5358, <https://doi.org/10.5194/acp-15-5325-2015>, 2015.
- 20 Brasseur, G. P., Hauglustaine, D. A., Walters, S., Rasch, P. J., Müller, J., Granter, C., and Tie, X. X.: MOZART, a global chemical transport model for ozone 1. Model description, *J. Geophys. Res.*, 103, D21, 28265–28289, <https://doi.org/10.1029/98JD02397>, 1998.
- Brown-Steiner, B., Hess, P. G., and Lin, M. Y.: On the capabilities and limitations of GCCM simulations of summertime regional air quality: A diagnostic analysis of ozone and temperature simulations in the US using CESM CAM-chem, *Atmos. Environ.*, 101, 134–148, 2015.
- 25 Brown-Steiner, B., Selin, N. E., Prinn, R. G., Monier, E., Tilmes, S., Emmons, L., and Garcia-Menendez, F.: Maximizing Ozone Signals Among Chemical, Meteorological, and Climatological Variability, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-954>, in review, 2017.
- Burkholder, J.B., Abbatt, J. P. D., Barnes, I., Roberts, J.M., Melamed, M.L., Ammann, M., Bertram, A.K., Cappa, C.D., Carlton, A.G., Carpenter, L.J., Crowley, J.N., Dubowski, Y., George, C., Heard, D.E., Herrmann, H., Keutsch, F.N., Kroll, J.H., McNeill, V.F., Ng, N.L., Nizkorodov, S.A., Orlando, J.J., Percival, C.J., Picquet-Varraut, B., Rudich, Y., Seakins, P.W., Surratt, J.D., Tanimoto, H., Thornton, J.A., Tong, Z., Tyndall, G.S., Wahner, A., Weschler, C.J., Wilson, K.R., and Ziemann P.J.: The Essential Role for Laboratory Studies in Atmospheric Chemistry, *Environ. Sci. Technol.*, <https://doi.org/10.1021/acs.est.6b04947>, 2017.
- 30

- Cameron-Smith, P., Lamarque, J.-F., Connell, P., Chuang, C., and Vitt, F.: Toward an Earth system model: atmospheric chemistry, coupling, and petascale computing, *J. Phys.: Conf. Ser.*, 46, 343–350, <https://doi.org/10.1088/1742-6596/46/1/048>, 2006.
- 5 Cameron-Smith, P., Prather, M. J., Lamarque, J., Hess, P. G., Connell, P. S., Bergmann, D. J., and Vitt, F. M.: The super-fast chemistry mechanism for IPCC AR5 simulations with CCSM, in: AGU, Fall Meeting, 18 December 2009, San Francisco, A54A-08, 2009.
- Cariolle, D., A. Lasserre-Bigorry, J.-F. Royer, and J.-F. Geleyn.: A General Circulation Model Simulation of the Springtime Antarctic Ozone Decrease and Its Impact on Mid-Latitudes, *J. Geophys. Res.*, 95(D2), 1883–1898, doi:10.1029/JD095iD02p01883, 1990.
- 10 CASTNET. (2016). CASTNET 2014 Annual Report, Prepared by Environmental Engineering and Measurement Services, Inc. for the U.S. Environmental Protection Agency.
- Collins, W. D., Ramaswamy, V., Schwarzkopf, M. D., Sun, Y., Portmann, R. W., Fu, Q., Casanova, S. E. B., Dufresne, J.-L., Fillmore, D. W., Forster, P. M. D., Galin, V. Y., Gohar, L. K., Ingram, W. J., Kratz, D. P., Lefebvre, M.-P., Li, J., Marquet, P., Oinas, V., Tsushima, Y., Uchiyama, T., and Zhong, W. Y.: Radiative forcing by well-mixed greenhouse gases: Estimates from climate models in the Intergovernmental Panel on Climate Change (IPCC) Fourth Assessment Report (AR4), *J. Geophys. Res.*, 111, D14, D14317, <http://doi.org/10.1029/2005JD006713>, 2006.
- 15 Dodge, M. C.: Chemical oxidant mechanisms for air quality modeling: Critical review, *Atmos. Environ.*, 34, 2103–2130, [https://doi.org/10.1016/S1352-2310\(99\)00461-6](https://doi.org/10.1016/S1352-2310(99)00461-6), 2000.
- Emmerson, K. M., and Evans, M. J.: Comparison of tropospheric gas-phase chemistry schemes for use within global models, *Atmos. Chem. Phys.*, 9, 1831–1845, <http://doi.org/10.5194/acpd-8-19957-2008>, 2009.
- 20 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- 25 Emmons, L. K., Hess, P. G., Lamarque, J.-F., and Pfister, G. G.: Tagged ozone mechanism for MOZART-4, CAM-chem and other chemical transport models, *Geosci. Mod. Dev.*, 5, 1531–1542, <http://doi.org/10.5194/gmd-5-1531-2012>, 2012.
- Emmons, L. K., Arnold, S. R., Monks, S. A., Huijnen, V., Tilmes, S., Law, K. S., Thomas, J. L., Raut, J.-C., Bouarar, I., Turquety, S., Long, Y., Duncan, B., Steenrod, S., Strode, S., Flemming, J., Mao, J., Langner, J., Thompson, A. M., Tarasick, D., Apel, E. C., Blake, D. R., Cohen, R. C., Dibb, J., Diskin, G. S., Fried, A., Hall, S. R., Huey, L. G., Weinheimer, A. J., Wisthaler, A., Mikoviny, T., Nowak, J., Peischl, J., Roberts, J. M., Ryerson, T., Warneke, C., and Helmig, D.: The POLARCAT Model Intercomparison Project (POLMIP): overview and evaluation with observations, *Atmos. Chem. Phys.*, 15, 6721–6744, <http://doi.org/10.5194/acp-15-6721-2015>, 2015.
- 30 Fiore, A. M., Naik, V., and Leibensperger, E. M.: Air Quality and Climate Connections, *J. Air Waste Manag. Assoc.*, 65, 645–685, <http://doi.org/10.1080/10962247.2015.1040526>, 2015.

- Garcia-Menendez, F., Saari, R. K., Monier, E., and Selin, N. E.: U.S. Air Quality and Health Benefits from Avoided Climate Change under Greenhouse Gas Mitigation, *Environ. Sci. Technol.*, 49, 7580–7588, <https://doi.org/10.1021/acs.est.5b01324>, 2015.
- 5 Garcia-Menendez, F., Monier, E., and Selin, N. E.: The role of natural variability in projections of climate change impacts on U.S. ozone pollution, *Geophys. Res. Lett.*, 44, 2911–2921, <https://doi.org/10.1002/2016GL071565>, 2017.
- Gery, M. W., Whitten, G. Z., Killus, J. P., and Dodge, M. C.: A photochemical kinetics mechanism for urban and regional scale computer modeling, *J. Geophys. Res.*, 94, D10, 12925–12956, <https://doi.org/10.1029/JD094iD10p12925>, 1989.
- Granier, C., Pétron, G., Müller, J.-F., and Brasseur, G.: The impact of natural and anthropogenic hydrocarbons on the tropospheric budget of carbon monoxide, *Atmos. Environ.*, 34, 5255–5270. [https://doi.org/10.1016/S1352-2310\(00\)00299-5](https://doi.org/10.1016/S1352-2310(00)00299-5), 2000.
- 10 Guenther, a. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Mod. Dev.*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Hauglustaine, D. A., Brasseur, G. P., Walters, S., Rasch, P. J., Mfiller, J., Ernrons, L. K., and Carroll, M. A.: MOZART, a global chemical transport model for ozone and related chemical tracers 2. Model results and evaluation, *J. Geophys. Res.*, 103, D21, 28291–28335, <https://doi.org/10.1029/98JD02398>, 1998.
- 15 Horowitz, L. W.: A global simulation of tropospheric ozone and related tracers: Description and evaluation of MOZART, version 2, *J. Geophys. Res.*, 108, D24, 4784, <https://doi.org/10.1029/2002JD002853>, 2003.
- Hourdin, F., Mauritsen, T., Gettelman, A., Golaz, J.-C., Balaji, V., Duan, Q., Folini, D., Ji, Duoying, Klocke, D., Qian, Y., Rauser, F., Rio, C., Tomassini, L., Watanabe, M., and Williamson, D.: The Art and Science of Climate Model Tuning, *Bull. Amer. Meteor. Soc.*, 98, 589–602, <http://doi.org/10.1175/BAMS-D-15-00135.1>, 2017.
- 20 Houweling, S., Dentener, F., and Lelieveld, J.: The impact of nonmethane hydrocarbon compounds on tropospheric photochemistry, *J. Geophys. Res. Atmos.*, 103, D9, 10673–10696, <https://doi.org/10.1029/97JD03582>, 1998.
- Hsu, J., and M. J. Prather.: Stratospheric variability and tropospheric ozone. *J. Geophys. Res.*, 114, D06102, doi:10.1029/2008JD010942, 2009
- 25 Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: A protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, [https://doi.org/10.1016/S1352-2310\(96\)00105-7](https://doi.org/10.1016/S1352-2310(96)00105-7), 1997.
- Jimenez, P., Baldasano, J. M., and Dabdub, D.: Comparison of photochemical mechanisms for air quality modeling, *Atmos. Environ.*, 37, 4179–4194, [https://doi.org/10.1016/S1352-2310\(03\)00567-3](https://doi.org/10.1016/S1352-2310(03)00567-3), 2003.
- 30 Kinnison, D. E., Brasseur, G. P., Walters, S., Garcia, R. R., Marsh, D. R., Sassi, F., Harvey, V. L., Randall, C. E., Emmons, L. K., Lamarque, J.-F., Hess, P., Orlando, J. J., Tie, X. X., Randel, W., Pan, L. L., Gettelman, A., Granier, C., Diehl, T., Niemeier, U., and Simmons, A. J.: Sensitivity of chemical tracers to meteorological parameters in the MOZART-3 chemical transport model, *J. Geophys. Res.*, 112, D20, 1–24, <https://doi.org/10.1029/2006JD007879>, 2007.

- Knote, C., Tuccella, P., Curci, G., Emmons, L., Orlando, J. J., Madronich, S., Baró, R., Jimenez-Guerrero, P., Luecken, D., Hogrefe, C., Forkel, R., Werhahn, J., Hirtl, M., Pérez, J. L., San Jose, R., Giordano, L., Brunner, D., Yahya, K., and Zhang, Y.: Influence of the choice of gas-phase mechanism on predictions of key gaseous pollutants during the AQMEII phase-2 intercomparison, *Atmos. Environ.*, 115, 553-568, <https://doi.org/10.1016/j.atmosenv.2014.11.066>, 2015.
- Lamarque, J.-F., Kinnison, D. E., Hess, P. G., and Vitt, F. M.: Simulated lower stratospheric trends between 1970 and 2005: Identifying the role of climate and composition changes, *J. Geophys. Res.*, 113, D12, 1–19, <https://doi.org/10.1029/2007JD009277>, 2008.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Lioussé, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: Description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, <https://doi.org/10.5194/gmd-5-369-2012>, 2012.
- Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The atmospheric chemistry and climate model intercomparison Project (ACCMIP): Overview and description of models, simulations and climate diagnostics, *Geosci. Mod. Dev.*, 6, 179–206, <https://doi.org/10.5194/gmd-6-179-2013>, 2013.
- Madronich, S., and Calvert, J. G.: The NCAR Master Mechanism of the gas phase chemistry – Version 2.0, NCAR Technical Note, TN-333+SRT, Boulder, Colorado, 1989.
- Marsh, D. R., Mills, M. J., Kinnison, D. E., Lamarque, J.-F., Calvo, N., and Polvani, L. M.: Climate change from 1850 to 2005 simulated in CESM1 (WACCM), *J. Climate*, 26, 7372–7391, <https://doi.org/10.1175/JCLI-D-12-00558.1>, 2013.
- McLinden, C. A., S. C. Olsen, B. Hannegan, O. Wild, M. J. Prather, and J. Sundet.: Stratospheric ozone in 3-D models: A simple chemistry and the cross-tropopause flux, *Journal of Geophysical Research-Atmospheres*, 105(D11), 14653–14665, doi:10.1029/2000jd900124, 2000.
- Milford, J. B., Gao, D., Russell, A. G., and McRae, G. J.: Use of Sensitivity Analysis to Compare Chemical Mechanisms for Air-Quality Modeling, *Environ. Sci. Technol.*, 26, 1179–1189, <https://doi.org/10.1021/es50002a606>, 1992.
- Morgenstern, O., Hegglin, M., Rozanov, E., O'Connor, F., Luke Abraham, N., Akiyoshi, H., Archibald, A. T., Bekki, S., Butchart, N., Chipperfield, M. P., Deushi, M., Dhomse, S. S., Garcia, R. R., Hardiman, S. C., Horowitz, L. W., Jöckel,

5

10

15

20

25

30

P., Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M. E., Marchand, M., Marécal, V., Michou, M., Oman, L. D., Pitari, G., Plummer, D. A., Revell, L. E., Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T. Y., Silmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within phase 1 of the Chemistry-Climate Model Initiative (CCMI), *Geosci. Mod. Dev.*, 10, 639–671, <https://doi.org/10.5194/gmd-10-639-2017>, 2017.

Pfister, G., Walters, S., Lamarque, J.-F., Fast, J., Barth, M. C., Wong, J., Done, J., Holland, G., and Bruyère, C. L.: Projections of future summertime ozone over the U.S., *J. Geophys. Res. Atmos.*, 119, 5559–5582, <https://doi.org/10.1002/2013JD020932>, 2014.

[Phalitnonkiat, P., Sun, W., Grigoriu, M. D., Hess, P., and Samorodnitsky, G. Extreme ozone events: Tail behavior of the surface ozone distribution over the U.S. *Atmos. Environ.*, 128, 134–146. 2016.](#)

Pöschl, U., Von Kuhlmann, R., Poisson, N., and Crutzen, P. J.: Development and intercomparison of condensed isoprene oxidation mechanisms for global atmospheric modeling, *J. Atmos. Chem.*, 37, 29–52, <https://doi.org/10.1023/A:1006391009798>, 2000.

Prinn, R. G., Huang, J., Weiss, R. F., Cunnold, D. M., Fraser, P. J., Simmonds, P. G., McCulloch, A., Harth, C., Salameh, P., O’Doherty, S., Wang, R. H. J., Porter, L., and Miller, B. R.: Evidence for Substantial Variations of Atmospheric Hydroxyl Radicals in the Past Two Decades, *Science*, 292, 1882–1888, <https://doi.org/10.1126/science.1058673>, 2001.

Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, R., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA’s modern-era retrospective analysis for research and applications, *J. Climate*, 24, 3624–3648, <https://doi.org/10.1175/JCLI-D-11-00015.1>, 2011.

Rotman, D. A., Atherton, C. S., Bergmann, D. J., Cameron-Smith, P. J., Chuang, C. C., Connell, P. S., Dignon, J. E., Franz, A., Grant, K. E., Kinnison, D. E., Molenkamp, C. R., Proctor, D. D., and Tannahill, J. R.: IMPACT, the LLNL 3-D global atmospheric chemical transport model for the combined troposphere and stratosphere: Model description and analysis of ozone and other trace gases, *J. Geophys. Res.*, 109, D04303, <https://doi.org/10.1029/2002JD003155>, 2004.

Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, <https://doi.org/10.5194/acp-3-161-2003>, 2003.

Schnell, J. L., Prather, M. J., Josse, B., Naik, V., Horowitz, L. W., Cameron-Smith, P., Bergmann, D., Zeng, G., Plummer, D. A., Sudo, K., Nagashima, T., Shindell, D. T., Faluvegi, G., and Strode, S. A.: Use of North American and European air quality networks to evaluate global chemistry-climate modeling of surface ozone, *Atmos. Chem. Phys.*, 15, 10581–10596, <https://doi.org/10.5194/acp-15-10581-2015>, 2015.

Silva, R. A., West, J. J., Zhang, Y., Anenberg, S. C., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S., Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S., Skeie, R., Sudo, K., Takemura, T.,

- Bergmann, D., Cameron-Smith, P., Cionni, I., Doherty, R. M., Eyring, V., Josse, B., MacKenzie, I. A., Plummer, D., Righi, M., Stevenson, D. S., Strode, S., Szopa, S., and Zeng, G.: Global premature mortality due to anthropogenic outdoor air pollution and the contribution of past climate change, *Environ. Res. Lett.*, 8, 034005, <https://doi.org/10.1088/1748-9326/8/3/034005>, 2013.
- 5 Silva, R. A., West, J. J., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S., Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S. T., Sudo, K., Takemura, T., Bergmann, D., Cameron-Smith, P., Cionni, I., Doherty, R. M., Eyring, V., Josse, B., MacKenzie, I. A., Plummer, D., Righi, M., Stevenson, D. S., Strode, S., Szopa, S., and Zengast, G.: The effect of future ambient air pollution on human premature mortality to 2100 using output from the ACCMIP model ensemble, *Atmos. Chem. Phys.*, 16, 9847-9862, doi:10.5194/acp-16-9847-2016, 2016.
- 10 Silva, R. A., West, J. J., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Faluvegi, G., Folberth, G. A., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S. T., Sudo, K., Takemura, T., Bergmann, D., Cameron-Smith, P., Doherty, R. M., Josse, B., MacKenzie, I. A., Stevenson, D. S., and Zeng, G.: Future global mortality from changes in air pollution attributable to climate change, *Nature Climate Change*, 7, 647-651, doi:10.1038/nclimate3354, 2017.
- Squire, O. J., Archibald, A. T., Griffiths, P. T., Jenkin, M. E., Smith, D., and Pyle, J. A.: Influence of isoprene chemical
15 mechanism on modelled changes in tropospheric ozone due to climate and land use over the 21st century, *Atmos. Chem. Phys.*, 15, 5123-5143, <https://doi.org/10.5194/acp-15-5123-2015>, 2015.
- Sofen, E. D., Bowdalo, D., and Evans, M. J.: How to most effectively expand the global surface ozone observing network, *Atmos. Chem. Phys.*, 16, 1445-1457, 2016.
- Solomon, S., Ivy, D. J., Kinnison, D., Mills, M. J., Iii, R. R. N., and Schmidt, A.: Antarctic ozone layer, *Science*, 353, 269-
20 274, <http://doi.org/10.1126/science.aae0061>, 2016.
- Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsøren, S. B., Myhre, G., Berntsen, T. K., Folberth, G. A., Rumbold, S. T., Collins, W. J., MacKenzie, I. A., Doherty, R. M., Zeng, G., van Noije, T. P. C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D. A., Strode, S. A., Horowitz, L., Lee, Y. H., Szopa, S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K.
25 W., and Wild, O.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 3063-3085, <https://doi.org/10.5194/acp-13-3063-2013>, 2013.
- Stockwell, W. R., Lawson, C. V., Saunders, E., and Goliff, W. S.: A Review of Tropospheric Atmospheric Chemistry and Gas-Phase Chemical Mechanisms for Air Quality Modeling, *Atmosphere*, 3, 1-32,
30 <https://doi.org/10.3390/atmos3010001>, 2011.
- Szopa, S., Aumont, B., and Madronich, S.: Assessment of the reduction methods used to develop chemical schemes: building of a new chemical scheme for VOC oxidation suited to three-dimensional multiscale HO_x-NO_x-VOC chemistry simulations, *Atmos. Chem. Phys.*, 5, 2519-2538, <https://doi.org/10.5194/acpd-5-755-2005>, 2005.

- Tilmes, S., Lamarque, J.-F., Emmons, L. K., Conley, A., Schultz, M. G., Saunio, M., Thouret, V., Thompson, A. M., Oltmans, S. J., Johnson, B., and Tarasick, D.: Technical Note: Ozone sonde climatology between 1995 and 2011 : description, evaluation and applications, *Atmos. Chem. Phys.*, 12, 7475–7497, 2012.
- 5 Tilmes, S., Lamarque, J.-F., Emmons, L. K., Kinnison, D. E., Ma, P. L., Liu, X., Ghan, S., Bardeen, C., Arnold, S., Deeter, M., Vitt, F., Ryerson, T., Elkins, J. W., Moore, F., Spackman, J. R., and Val Martin, M.: Description and evaluation of tropospheric chemistry and aerosols in the Community Earth System Model (CESM1.2), *Geosci. Model Dev.*, 8, 1395–1426, <https://doi.org/10.5194/gmd-8-1395-2015>, 2015.
- 10 Tilmes, S., Lamarque, J.-F., Emmons, L. K., Kinnison, D. E., Marsh, D., Garcia, R. R., Smith, A. K., Neely, R. R., Conley, A., Vitt, F., Val Martin, M., Tanimoto, h., Simpson, I., Blake, D. R., and Blake, N.: Representation of the Community Earth System Model (CESM1) CAM4-chem within the Chemistry-Climate Model Initiative (CCMI), *Geosci. Model Dev.*, 9, 1853–1890, <https://doi.org/10.5194/gmd-9-1853-2016>, 2016.
- Uusitalo, L., Lehtikoinen, A., Helle, I., and Myrberg, K.: An overview of methods to evaluate uncertainty of deterministic models in decision support, *Environ. Model. Softw.*, 63, 24–31, <https://doi.org/10.1016/j.envsoft.2014.09.017>, 2015.
- 15 Val Martin, M., Heald, C. L., and Arnold, S. R.: Coupling dry deposition to vegetation phenology in the Community Earth System Model: Implications for the simulations of surface O₃, *Geophys. Res. Lett.*, 41, doi:10.1002/2014GL059651, 2015.
- Wang, C., and Prinn, R. G.: Impact of emissions, chemistry and climate on atmospheric carbon monoxide: 100-yr predictions from a global chemistry–climate model, *Chemosphere Global Change Sci.*, 1, 73–81, [https://doi.org/10.1016/S1465-9972\(99\)00016-1](https://doi.org/10.1016/S1465-9972(99)00016-1), 1999.
- 20 Wild, O., and Prather, M. J.: Excitation of the primary tropospheric chemical mode in a global three-dimensional model, *J. Geophys. Res.*, 105, D20, 24647–24660, <https://doi.org/10.1029/2000JD900399>, 2000.
- Young, P. J., Archibald, A. T., Bowman, K. W., Lamarque, J.-F., Naik, V., Stevenson, D. S., Tilmes, S., Voulgarakis, A., Wild, O., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Horowitz, L. W., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R. B., Shindell, D. T., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Pre-industrial to end 21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 2063–2090, <https://doi.org/10.5194/acp-13-2063-2013>, 2013.
- 25

Abbreviation	MO	RH*	SF**
Full Name	MOZART-4	Reduced Hydrocarbom	Super-Fast
Primary Citation	Emmons et al. (2010)	Houwelling et al. (1998)	Cameron-Smith et al. (2006)
Total Tracers	103	65 (89)	15
Total Reactions	212	127 (202)	30
Reactions in NHMC Chemistry	108	28	2
core hours / simulated year	615	319	165 (204)
simulated years / day	2.5	4.8	9.3 (7.5)
Efficiency (compared to MO)	1.0	1.9	3.7 (3.0)

*: unmodified RH listed in the parenthesis

** : SF + Bulk Aerosol Model (BAM) included in parentheses

Abbreviation	MO	RH*	SF**
Full Name	MOZART-4	Reduced Hydrocarbom	Super-Fast
Primary Citation	Emmons et al. (2010)	Houwelling et al. (1998)	Cameron-Smith et al. (2006)
Total Tracers	103	65 (89)	15
Total Reactions	212	127 (202)	30
Reactions in NHMC Chemistry	108	28	2
core hours / simulated year	615	319	165 (204)
simulated years / day	2.5	4.8	9.3 (7.5)
Efficiency (compared to MO)	1.0	1.9	3.7 (3.0)

*: unmodified RH listed in the parenthesis

** : SF + Bulk Aerosol Model (BAM) included in parenthesis

Comment [NES2]: Should be "parentheses" not parenthesis (which is only one side)

Table 1: Summary and comparison of the MOZART-4 (MO), Reduced Hydrocarbon (RH), and Super-Fast (SF) mechanisms included in this paper. All runs were conducted on the NCAR Cheyenne system with 64 CPUs on 2 nodes without any load optimization, and the values in this table represent the cost of the entire CESM CAM-chem model, not just the chemistry component. In this study, we removed many stratospheric species (see text), so we include both the modified and unmodified (in parentheses) RH mechanisms. The MO and RH mechanism include BAM.

		[ppbv] Mean	[ppbv] Median	[ppbv] Standard Deviation	[%] Variability	[ppbv] 90th Percentile	[ppbv] 99th Percentile	[ppbv] 99th - 90th Percentile	[%]
Global	MO	32.6	32.6	0.93	2.86	33.8	35.3	1.52	105
	RH	33.9	33.9	0.98	2.90	35.2	36.4	1.25	104
	SF	31.5	31.4	1.12	3.57	33.0	34.1	1.07	103
CO US	CASTNET	56.1	55.8	5.65	10.1	63.4	71.0	7.60	112
	MO	56.7	56.6	3.08	5.43	60.6	64.4	3.82	106
	RH	57.7	57.6	3.23	5.60	61.8	65.9	4.10	107
	SF	53.4	53.3	3.59	6.72	57.9	62.8	4.87	108
E US	CASTNET	56.4	56.0	6.41	11.4	64.5	73.3	8.78	114
	MO	58.6	58.4	5.77	9.85	66.1	72.8	6.70	110
	RH	59.7	59.5	6.06	10.2	67.5	74.7	7.17	111
	SF	56.5	56.1	7.12	12.6	66.0	74.8	8.77	113
NE US	CASTNET	57.4	56.9	7.42	12.9	66.6	78.1	11.4	117
	MO	74.4	73.7	11.4	15.4	89.8	104	13.8	115
	RH	76.0	75.1	11.9	15.6	92.0	107	14.8	116
	SF	72.6	71.3	14.5	20.0	91.8	114	21.9	124
NE US single grid cell	CASTNET	59.7	59.3	11.1	18.6	73.9	86.9	13.0	118
	MO	84.9	85.4	12.8	15.1	101	115	13.7	114
	RH	86.1	86.2	13.2	15.3	103	117	14.4	114
	SF	99.6	97.3	25.6	25.7	133	171	38.2	129

Table 2: Summary Statistics for the Daily Maximum 8-Hour (DM8HMDA8) O₃ over the globe and over the indicated regions in the US. Additional regions can be found in Supplemental Table S3.

Comment [NES3]: Same comment as on ACPD paper – should you alter the acronym here?




Figure 1: Maps of DJF and JJA O_3 , CO , and NO_x for MO, the difference between RH and MO, and between SF and MO for the year 2015. The chemical units are in ppb. Please note the difference in the chemical scales for each panel. Cool colors for the difference panels indicate MO is higher, and warm colors indicate that RH or SF is higher.

Comparison to Ozonesondes




Figure 2: Taylor-like diagrams comparing the mean and correlation of the seasonal cycle between observations (present-day ozonesonde climatology (Tilmes et al., 2012) from 1995 to 2011 for different regions (Tropics, Mid-Latitudes, and High-Latitudes) and different pressure levels (900 hPa, 250 hPa, and 50 hPa), as in Figure 12 of Tilmes et al. (2015)) and simulations (red: MO, blue: RH, green: SF).




Figure 3: Relative differences between [available](#) aircraft observations (black) and the MO, RH, and SF model configurations (colors) over different regions and seasons, averaged over 2 – 7 km, for O_3 , NO_x , CO , HNO_3 , H_2O_2 , and SO_2 as in Figure 17 of Tilmes et al. (2015).





Figure 4: Zonal Plots of Seasonal O_3 , CO , and NO_x for MO, the difference between RH and MO, and between SF and MO for the year 2015. The vertical axis is the model level, and the chemical units are in ppb. Please note the different vertical axis in each row. Cool colors for the different panels indicate

5 MO is higher, and warm colors indicate that RH or SF is higher.






Figure 5: Surface JJA ~~DM8HMDA8~~ O₃ boxplots for the 1991 – 2014 data for CASTNET (grey), MO (red), RH (blue), and SF (green) averaged over the various regions. Plots g, h, and i are individual grid cells from within each region (38.8° N and 87.5° W for g, 38.8° N and 80.0° W for h, and 33.2° N and 85.0° W for i). Global boxplots are included along with the Continental US. The units are in ppb, and for each boxplot the box contains the Inter Quartile Range (IQR), the horizontal line within the box is the median, and the whiskers extend out to the farthest point which is within 1.5 times the IQR with circles indicating any outliers. Note the scale difference between the top row and the rest of the panels.




Figure 6: The relative difference (%) between the 99th percentile and the 90th percentile of JJA DM8HMDA8 O₃ for CASTNET and the three mechanisms over three regions as a function of increasing length of simulation, from 1 day up to the full 25 years simulated. The vertical bars indicate the year 2000, for which the emissions for all three simulations were cycled.





Figure 7: R^2 values calculated at every grid-cell (for the full ~~1990-1991~~ – 2015 ~~DM8HMDA8~~ O_3 JJA time series) for MO and RH (left) and MO and SF (right).







Figure 8: Seasonal time series for O₃, CO, NO_x, HO_x, and ISOP for MO (black/red), RH (blue), and SF (red/green) for a single year (2015), averaged over different regions. The units are in ppb. Note the different scales in each panel. Also included are three sensitivity tests conducted with the SF mechanism (which were ran only for 2 years, 1990 – 1991, with 1991 being plotted here): adding PAN chemistry (purple), adding N₂O₅ chemistry (orange), and adding both PAN and N₂O₅ chemistry (green/black).








Figure 9: Example diurnal time series for various species for MO (~~black-red~~ circles), RH (blue triangles), and SF (~~red-green~~ diamonds) averaged over a single grid cell in the central US (100° west and 47° north). The units are in ppb. Also included are three sensitivity tests conducted with the SF mechanism: adding PAN chemistry (purple), adding N₂O₅ chemistry (orange), and adding both PAN and N₂O₅ chemistry (~~greenblack~~).





5 **Figure 10:** Scatterplots comparing model results to observations (two center columns) and to each other (two outer columns). Global regions (left) compare model results to ozonesondes (JJA averages), while regions within the US (right) compare the model results to CASTNET surface observations (JJA ~~MDA8~~ ~~MDA8~~ O₃). For the model-to-model comparisons, grey symbols additionally compare every grid cell in the model output. The numbers indicate the slope (upper left) and R² values (right) for each region. Each panel is labeled with the following convention: “y-axis” vs “x-axis.”

Supplemental Material for “Evaluating Simplified Chemical Mechanisms within Present-Day CESM Version 1.2 CAM-chem (CAM4): MOZART-4 vs. Reduced Hydrocarbon vs. Super-Fast Chemistry”

5 Benjamin Brown-Steiner^{1,2,3}, Noelle E. Selin^{3,4,5}, Ronald Prinn^{2,3,5}, Simone Tilmes⁶, Louisa Emmons⁶, Jean-François Lamarque⁶, Philip Cameron-Smith⁷

¹Now at Atmospheric and Environmental Research, 131 Hartwell Avenue, Lexington, MA 02421-3126

²Center for Global Change Science, Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139

³Joint Program on the Science and Policy of Global Change, Massachusetts Institute of Technology, 77 Massachusetts Ave,

10 Cambridge, MA 02139

⁴Institute for Data, Systems, and Society, Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139

⁵Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139

15 ⁶Atmospheric Chemistry Observations and Modeling Lab, National Center for Atmospheric Research, 3450 Mitchell Lane, Boulder, CO 80301

⁷Lawrence Livermore National Laboratory, 7000 East Ave, Livermore, CA 94550

Correspondence to: Benjamin Brown-Steiner (bbrownst@aer.com)

	CO	NO	NO2	SO2	C2H4	CH2O	CH3CHO	TOLUENE	NH3	ISOP	C10H16	OLE	PAR
CO	1												
NO		1											
NO2			1										
SO2				1									
BIGALK													5
BIGENE												1	2
C2H4					1								
C2H5OH													
C2H6													0.4
C3H6												1	1
C3H8													1.5
CH2O						1							
CH3CHO							1						
CH3COCH3													2
CH3OH													1
MEK													3
TOLUENE								1					
NH3									1				
ISOP										1			
C10H16											1		

Supplemental Table S1: Mapping of MOZART species (rows) to the lumped Reduced Hydrocarbon mechanism species

5 (columns)

Photolysis Reactions

	Reactant	Products	Relation to Emmons et al. (2010)
(i)	$O_3 + h\nu$	$2\cdot OH$	simplified, includes reaction of $O(^1D)$ with H_2O
(ii)	$H_2O_2 + h\nu$	$2\cdot OH$	identical
(iii)	$NO_2 + h\nu$	$NO + O_3$	simplified, $O \rightarrow O_3$
(iv)	$CH_2O + h\nu$	$CO + 2\cdot HO_2$	identical
(v)	$CH_2O + h\nu$	CO	simplified, no H_2
(vi)	$CH_3OOH + h\nu$	$CH_2O + HO_2 + OH$	simplified, $H \rightarrow HO_2$

Gas-Phase Reactions

	Reactants	Products	Rate	Relation to Emmons et al. (2010)
(1)	$O_3 + OH$	$HO_2 + O_2$	$1.70E-12 \cdot \exp(-940/T)$	identical
(2)	$HO_2 + O_3$	$2\cdot O_2 + OH$	$1.00E-14 \cdot \exp(-490/T)$	identical
(3)	$HO_2 + OH$	$H_2O + O_2$	$4.80E-11 \cdot \exp(250/T)$	identical
(4)	$HO_2 + HO_2$	$H_2O_2 + O_2$	$(2.3E-13 \cdot \exp(600/T) + 1.7E-33 \cdot [M] \cdot \exp(1000/T)) \cdot (1 + 1.4E-21 \cdot [H_2O] \cdot \exp(2200/T))$	identical
(5)	$H_2O_2 + OH$	$H_2O + HO_2$	$1.80E-12$	identical
(6)	$NO + O_3$	$NO_2 + O_2$	$3.00E-12 \cdot \exp(-1500/T)$	identical
(7)	$HO_2 + NO$	$NO_2 + OH$	$3.50E-12 \cdot \exp(250/T)$	different rates
(8)	$NO_2 + OH + M$	HNO_3	$k_o = 1.80E-30 \cdot (300/T)^{3.00}$, $k_i = 2.80E-11$; $f = 0.60$	identical
(9)	$CH_4 + OH$	$CH_3O_2 + H_2O$	$2.45E-12 \cdot \exp(-1775/T)$	identical
(10)	$CO + OH$	HO_2	$k_o \cdot m / (1 + (k_o \cdot m / k_{e,m})) \cdot 0.6 \cdot \exp((1/(1 + \log(k_o \cdot m / k_{e,m})))^2)) + (k_o / (1 + (k_o / k_{e,m}))) \cdot 0.6 \cdot \exp((1/(1 + \log(k_o / k_{e,m})))^2))$	simplified: no CO_2 , $H \rightarrow HO_2$, see note A
(11)	$CH_2O + OH$	$CO + H_2O + HO_2$	$5.50E-12 \cdot \exp(125/T)$	rates identical, simplified: $H \rightarrow HO_2$
(12)	$CH_3O_2 + HO_2$	$CH_3OOH + O_2$	$4.10E-13 \cdot \exp(750/T)$	identical
(13a)	$CH_3OOH + OH$	$CH_3O_2 + H_2O$	$2.70E-12 \cdot \exp(200/T)$	in combination, equivalent
(13b)	$CH_3OOH + OH$	$CH_3O + H_2O + OH$	$1.10E-12 \cdot \exp(200/T)$	
(14)	$CH_3O_2 + NO$	$CH_2O + HO_2 + NO_2$	$2.80E-12 \cdot \exp(300/T)$	identical
(15)	$CH_3O_2 + CH_3O_2$	$2\cdot CH_2O + 0.80\cdot HO_2$	$9.50E-14 \cdot \exp(390/T)$	different rates, simplified: 1 reaction instead of 2
(16)	$H_2O + NO_2$	$0.50\cdot HNO_3$	$4.00E-24$	no equivalent in MOZART, see note B
(17a)	$DMS + OH$	SO_2	$1.100E-11 \cdot \exp(-240/T)$	different, see note C
(17b)	$DMS + OH$	$0.75\cdot SO_2$	$2.00E-10 \cdot \exp(5820 \cdot [M]) / ((2.00E29 \cdot [O_2]) + \exp(6280 \cdot [M]))$	
(18)	$OH + SO_2 + M$	SO_4	$k_o = 3.30E-31 \cdot (300/T)^{4.30}$, $k_i = 1.60E-12$; $f = 0.60$	different, see note C
(19)	$H_2O_2 + SO_2$	SO_4	aqueous chemistry (see note D)	no equivalent, see note C
(20)	$O_3 + SO_2$	SO_4	aqueous chemistry (see note D)	no equivalent, see note C
(21a)	$ISOP + OH$	$2\cdot CH_3O_2$	$2.70E-11 \cdot \exp(390/T)$	different, see note E
(21b)	$ISOP + OH$	$ISOP$	$2.70E-11 \cdot \exp(390/T)$	
(21c)	$ISOP + OH$	$ISOP + 0.5\cdot OH$	$2.70E-11 \cdot \exp(390/T)$	different, see note E
(22)	$ISOP + O_3$	$.87\cdot CH_2O + 1.86\cdot CH_3O_2 + 0.06\cdot HO_2 + 0.05\cdot CO$	$5.59E-15 \cdot \exp(-1814/T)$	

NOTES:

A: For rate: $k_o = 5.90E-33 \cdot (300/T)^{1.4}$; $k_{e,m} = 1.10E-12 \cdot (T/300)^{1.3}$; $k_o \cdot m = k_o \cdot [M]$; $k_0 = 1.50E-13 \cdot (T/300)^{0.6}$; $k_{e,m} = (2.10E9 \cdot (T/300)^{6.1}) / [M]$

B: HNO_3 chemistry included only as reaction 8 and 16, with reaction 16 involving heterogeneous chemistry parameterization

C: DMS chemistry limited only to reaction with OH (reaction 17), SO_4 production simplified to reactions 18 with OH and 19 and 20 with aqueous chemistry (with a fixed pH in the cloud droplets)

D: Rate equations are included within the aerosol routines adapted from the MOZART-4 mechanism implementation within CAM-chem

E: Isoprene chemistry parameterized from UCI for ISOP + OH and from LLNL-IMPACT for ISOP + O_3 , see text for full details

Table S2: Full description of Super-Fast chemical mechanism as compared to the MOZART-4 mechanism of Emmons

et al. (2010). Reaction rates are written out if they are of the Arrhenius form, or otherwise formulated. If the reaction rates are of the Troe form, they list the k_o and k_i parameters, as in Emmons et al. (2010). The simplifications made in

the SF are noted by indicating what species is missing or modified when compared to Emmons et al. (2010). Chemical species are the same as in Emmons et al. (2010).

		[ppbv] Mean	[ppbv] Median	[ppbv] Standard Deviation	[%] Variability	[ppbv] 90th Percentile	[ppbv] 99th Percentile	[ppbv] 99th - 90th Percentile	[%]
W US	CASTNET	49.9	49.6	4.82	9.66	56.4	61.3	4.87	109
	MO	54.9	54.9	3.37	6.14	59.4	62.4	2.99	105
	RH	55.9	55.8	3.58	6.41	60.6	64.0	3.47	106
	SF	50.5	50.5	3.22	6.38	54.7	57.8	3.15	106
MW US	CASTNET	57.3	57.0	7.57	13.2	67.1	76.9	9.77	115
	MO	71.0	70.7	11.03	15.5	85.5	96.0	10.6	112
	RH	72.6	72.1	11.26	15.5	87.4	98.8	11.4	113
	SF	70.6	69.8	13.44	19.0	88.7	104	15.1	117
MW US single grid cell	CASTNET	62.4	60.6	14.5	23.2	81.7	104	22.7	128
	MO	84.6	83.8	16.2	19.2	106	126	19.4	118
	RH	86.2	85.3	15.8	18.4	107	126	19.5	118
	SF	85.4	82.5	22.4	26.2	116	149	32.8	128
SE US	CASTNET	51.5	51.3	8.18	15.9	62.3	71.9	9.64	115
	MO	60.2	59.5	9.22	15.3	72.5	83.7	11.2	115
	RH	61.3	60.5	10.1	16.6	74.9	87.9	12.9	117
	SF	58.6	57.8	11.4	19.4	73.9	87.1	13.2	118
SE US single grid cell	CASTNET	53.8	53.1	12.8	23.8	68.7	93.5	24.8	136
	MO	83.4	84.7	19.1	22.9	107	124	17.3	116
	RH	85.2	86.1	20.4	24.0	111	131	20.1	118
	SF	83.6	82.1	27.0	32.3	120	153	33.1	128

Table S3: Summary Statistics for the Daily Maximum 8-Hour (MDA8) O₃ over the globe other regions, accompanying Table 2. The last two columns indicate the difference between the 99th percentile and the 90th percentile, expressed both in absolute values (ppb) and as a percent.

Supplemental Description of the Super-Fast Chemical Mechanism

The SF mechanism is in the CESM code archive as an unsupported chemical mechanism, which can be activated using the option ‘-chem super_fast_llnl’. The SF mechanism has been included in several model inter-comparison projects, including the ACCMIP (e.g. Lamarque et al., 2013), a comparison of stratospheric dynamics and ozone production (Hsu et al., 2013), a comparison of isoprene mechanisms and ozone changes (Squire et al., 2015), and a multi-model assessment of surface ozone and observations (Schnell et al., 2015). The SF mechanism was also used to examine the role of DMS within ENSO (Xu et al., 2016). Here we briefly review the findings of these four model inter-comparison projects.

The SF only simulates sulfate (SO₄) and not the other aerosols, so the SF mechanism was not included in many of the ACCMIP aerosol comparisons (Lamarque et al., 2013). While the inclusion of non-sulfate aerosols within the CESM can be easily accomplished, there are two aerosol modules (either bulk or modal) to which aerosols could be added, which was beyond the scope of this project, so aerosol model capabilities are not examined in the present study.

We now summarize the ACCMIP results as they pertain to the SF mechanism. Within the ACCMIP, the SF mechanism has lower rates of ozone production and loss compared to the ACCENT models (biases of -24% and -22% respectively), as well as low ozone deposition (bias of -38%) (Young et al., 2013). In this comparison, natural emissions were not prescribed and different treatments of meteorology were used, which may account for some of the noted differences. This results in a high bias for the ozone lifetime (+3 days, or +14%), as well as a low ozone burden bias (-34 Tg, or -10%) (Young et al., 2013). In addition, the models that showed similarly low ozone production and loss rates have lower emissions of VOCs. The SF mechanism falls within the ACCMIP range for human health results due to ozone exposure (Silva et al., 2013). The SF mechanism simulated the 1850-2000 changes in the tropospheric ozone column within the range of the ACCMIP models, and projected changes to the ozone radiative forcing for future RCP scenarios also fell within the ACCMIP range (Stevenson et al., 2013). However, the calculated historical change in ozone RF fell outside of the ACCMIP range (+20% bias). The SF mechanism also has a high bias for global-mean OH (+16% compared to the ACCMIP mean) and a low bias for the calculation of the methane lifetime due to OH oxidation (-14%) (Voulgarakis et al., 2013).

The SF mechanism was tested against MOZART by Hsu et al. (2013) who concluded that the selection of a chemical mechanism was only a secondary influence on the stratospheric chemistry since they used a linearized scheme. However, the SF mechanism did produce a less stratified tropopause and a warmer troposphere due largely to the impact of ozone forcings on the simulated dynamics and thermodynamics. Unfortunately, the Hsu et al. (2013) analysis had a bug with their SF simulations, which resulted in the aerosols not being communicated to the cloud nucleation routines, but this didn’t affect their conclusions on the sensitivity of the stratosphere to uncertainty in the O₂ photolysis cross-section.

Squire et al. (2015) compared the SF isoprene scheme with three other schemes of much greater complexity. They concluded that the “1-species, 2-reaction” isoprene scheme from the SF mechanism, as simple as it is, is preferable to neglecting biogenic chemistry entirely, although the SF mechanism shows the highest biases in regions where isoprene chemistry is important for simulating accurate ozone concentrations. They also explored some of the other biases within the

SF mechanism scheme, which include: (1) under high-isoprene conditions, the SF mechanism overestimates O₃; (2) under low-isoprene and low-NO_x conditions, the SF mechanism overestimates O₃; (3) due to the simplicity of SF mechanism, HO_x is sequestered into the organic hydroperoxides, and methyl hydroperoxide (CH₃OOH) has low reactivity, which results in high levels of the peroxy radicals, an enhanced rate of CH₃O₂ + NO, and therefore a high bias (up to +80%) for ozone; and (4) the NO_x lifetime is too short, except in high-NO_x emission regions. They conclude that the addition of a PAN formation scheme would significantly improve the O₃ distribution. Finally, they find that many of the errors described above largely cancel each other out, which results in the globally averaged O₃ bias for SF mechanism to be small (-2.6% compared to the Master Chemical Mechanism).

The SF mechanism has a known anomalous annual cycle (see Schnell et al., 2015), in which peak ozone occurs in March/April rather than May. In the main article we show that this anomaly exists at global scales, but not within all regions. In addition, the size and extent of ozone pollution episodes is anomalously high, and these large events occur mainly in the springtime (Schnell et al., 2015). Interestingly, the SF mechanism outperforms many of the more sophisticated mechanisms in simulating the observed summertime diurnal cycle for ozone (Schnell et al., 2015).

References for Supplemental Material:

- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43-67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- Hsu, J., Prather, M. J., Bergmann, D., and Cameron-Smith, P.: Sensitivity of stratospheric dynamics to uncertainty in O₃ production, *J. Geophys. Res. Atmos.*, 118, 8984-8999, <https://doi.org/10.1002/jgrd.50689>, 2013.
- Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The atmospheric chemistry and climate model intercomparison Project (ACCMIP): Overview and description of models, simulations and climate diagnostics, *Geosci. Mod. Dev.*, 6, 179-206, <https://doi.org/10.5194/gmd-6-179-2013>, 2013.
- Schnell, J. L., Prather, M. J., Josse, B., Naik, V., Horowitz, L. W., Cameron-Smith, P., Bergmann, D., Zeng, G., Plummer, D. A., Sudo, K., Nagashima, T., Shindell, D. T., Faluvegi, G., and Strode, S. A.: Use of North American and European air quality networks to evaluate global chemistry-climate modeling of surface ozone, *Atmos. Chem. Phys.*, 15, 10581-10596, <https://doi.org/10.5194/acp-15-10581-2015>, 2015.
- Silva, R. A., West, J. J., Zhang, Y., Anenberg, S. C., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S., Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S., Skeie, R., Sudo, K., Takemura, T., Bergmann, D., Cameron-Smith, P., Cionni, I., Doherty, R. M., Eyring, V., Josse, B., MacKenzie, I. A., Plummer, D.,

Righi, M., Stevenson, D. S., Strode, S., Szopa, S., and Zeng, G.: Global premature mortality due to anthropogenic outdoor air pollution and the contribution of past climate change, *Environ. Res. Lett.*, 8, 034005, <https://doi.org/10.1088/1748-9326/8/3/034005>, 2013.

- 5 Squire, O. J., Archibald, A. T., Griffiths, P. T., Jenkin, M. E., Smith, D., and Pyle, J. A.: Influence of isoprene chemical mechanism on modelled changes in tropospheric ozone due to climate and land use over the 21st century, *Atmos. Chem. Phys.*, 15, 5123–5143, <https://doi.org/10.5194/acp-15-5123-2015>, 2015.
- 10 Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsøren, S. B., Myhre, G., Berntsen, T. K., Folberth, G. A., Rumbold, S. T., Collins, W. J., MacKenzie, I. A., Doherty, R. M., Zeng, G., van Noije, T. P. C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D. A., Strode, S. A., Horowitz, L.,
25 Lee, Y. H., Szopa, S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K. W., and Wild, O.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 3063–3085, <https://doi.org/10.5194/acp-13-3063-2013>, 2013.
- 15 Voulgarakis, A., Naik, V., Lamarque, J.-F., Shindell, D. T., Young, P. J., Prather, M. J., Wild, O., Field, R. D., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Folberth, G. A., Horowitz, L. W., Josse, B., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Stevenson, D. S., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Analysis of present day and future OH and methane lifetime in the ACCMIP simulations, *Atmos. Chem. Phys.*, 13, 2563–2587
- 20 Xu, L., Cameron-Smith, P., Russell, L. M., Ghan, S. J., Liu, Y., Elliott, S., Yang, y., Lou, S., Lamjiri, M. A., and Manizza, M.: DMS role in ENSO cycle in the tropics, *J. Geophys. Res. Atmos.*, 121, 537–558, <https://doi.org/10.1002/2016JD025333>, 2016.
- 25 Young, P. J., Archibald, A. T., Bowman, K. W., Lamarque, J.-F., Naik, V., Stevenson, D. S., Tilmes, S., Voulgarakis, A., Wild, O., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Horowitz, L. W., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R. B., Shindell, D. T., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Pre-industrial to end 21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 2063–2090, <https://doi.org/10.5194/acp-13-2063-2013>, 2013.

