

Reply reviewer 1

On the example of the North and Baltic Sea, the authors present a very simple way to extend a lower trophic level model (marine biogeochemistry) to an end-to-end model. Two state variables have to be added, both of them two-dimensional (only horizontally resolved). These are the biomasses of fish and macrozoobenthos.

Since these two “upper trophic levels” influence especially the mortality of zooplankton, the result is a spatially heterogenous mortality which replaces the previously uniform closure term of the LTL model. A comparison to a study for the North Sea where fish predation mortality is estimated from observed distributions of planktivorous fish shows, as the authors state, that the correct spatial patterns emerge. In this way, the model outlines a rather simple way to include, to a first order, higher trophic level effects onto the lower trophic food web. Compared to the previous study, this approach has much lower data requirements, which is an advantage not pointed out well enough.

When it comes to overall model performance, the authors could show a slight benefit when adding the two state variables as they compare lower trophic level variables such as nutrient concentrations and oxygen. At least for the North Sea, the authors could also show that fish and macrozoobenthos spatial patterns were similar to observations.

From my point of view, the largest benefit of the model is the opportunity to gain more realistic zooplankton mortality values. I would like this point to be more highlighted. Especially I would like to see the spatial patterns of zooplankton mortality as they emerge from this model compared to those of Maar et al. (2014). Contrary to what the authors state in the caption of Fig. 9, fish production is not a suitable indicator for zooplankton predation mortality, but only the ratio between fish production and zooplankton biomass gives the predation mortality. The comparison should be done to Fig. 10C+D in Maar et al., not to Fig. 4, and it would be very useful if the comparison was shown in this article.

Daewel et al: We agree that this comparison is necessary. We have compiled biomass specific mortality based on our model results as an average from 1980-1989 and separated the year into 1st and 2nd half as done in Maar et al. (2014) (see suppl. Figure 1a). The results are around the same magnitude as calculated by Maar et al. (2014) and the structure resembles the structure they had calculated for the first half of the year 2001. The results from Maar et al (2014) showed a clear difference between half-year 1 and half-year 2 with decreased biomass specific mortality in the second half-year in the central North Sea. This difference between the 1st and 2nd half-year is not evident in our model results. However we found a clear difference in magnitude when comparing winter and summer season (not shown). The reasons for the discrepancies between our model results and Maar et al (2014) are presumably related to interannual variations in fish consumption, which are not considered in a 10 year average, the fact that migration is not considered in the model and thus restrict the spatial variation, and that our functional group cannot resolve species and stage specific spatial and temporal variations like e.g. the increase in larval biomass in spring and changes in species composition. On the other hand, the approach from Maar et al. (2014) reveals uncertainties due to the fact that only parts of the North Sea fish assemblage is considered and that the fish biomass is prescribed and not dynamically coupled to zooplankton biomass. Considering the uncertainties in both approaches the comparison is actually quite good in both magnitude and spatial structure.

Action: We replaced the old Figure 9 with Suppl. Figure 1 (now the new Figure 9), (which also shows biomass specific zooplankton mortality due to macrobenthos), and substantially modified the text and discussion related to

Figure 9.

The discussion and conclusions sound reasonable to me, I would support them. The only exception is that I would not present total fish biomass as a model result, since it might be quite sensitive to the feeding efficiency which is poorly confined for the “average fish” I guess? I would rather interpret fish biomass as a tuning parameter which you have to fit to ensure that the ratio between predation mortality and background mortality is reasonable. In this way, quite simple measured quantities like total fish biomass and fishing mortality can be used to confine the zooplankton mortality, which was an arbitrary closure term before your extension.

Daewel et al.: We agree that showing total fish biomass as a model result is a bit critical in this context. On the other hand, this is one of the few possibilities to confirm that the model is able to estimate reasonable results in terms of magnitude and distribution, otherwise it will certainly be difficult show that the approach actually improves the formulation of the predation mortality for zooplankton. So we would rather leave that part in the ms. We have added the scenarios simulations to the manuscript to identify the sensitivity of the results. However, we suggest clarifying the uncertainty of the variable more thoroughly in the ms, together with the response to general remark 1 below.

General remarks:

1.) You say you can interpret the fish functional group as a “fish production potential”. In principle it is clear what you mean: Where there is food, there will be fish who eat it. You are not interested in where they will migrate afterwards, but you keep the fish biomass locally as a kind of bookkeeping of what the fish consumed here. However, see a few issues with this interpretation:

· Zooplankton or macrobenthos will not be consumed if the fish just “potentially” grow. So, the loss terms for these functional groups contradict this interpretation.

· Consumption of food by fish is limited by local fish biomass in the model. A “fish production potential” in reality could be larger, since fish from remote locations could migrate towards a spot of high prey abundance, which might lead to more efficient food consumption. I would like a clearer discussion of why neither of the two interpretations (fish biomass / fish production potential) is entirely correct.

Daewel et al: Yes we agree with the reviewer, this needs to be clarified in the ms. We will add the following explanation to the method section: “In the following, we will thus refer to “fish” as a functional group that comprises the fish biomass that emerges based on the lower trophic production at each horizontal grid cell. For clarification it needs to be noted that, even when called “fish production potential”, the fish biomass is a state variable in the model that interacts dynamically with the lower trophic level components and that will be used in the following to confirm the models ability to simulate spatial and temporal pattern of carbon transfer to higher trophic levels. On the other hand by constraining the horizontal migration capabilities of the fish group to one grid cell we will likely underestimate the local fish production potential by confining it to the locally available fish biomass.”

2.) Your extension has a very low computational cost. Not only is it just two state variables that are added, but also these are 2-d only which saves their advection, and the advection of the state variables is the most time consuming step. So your extension is especially suitable for LTL models which are simple by purpose, e.g. because they are used in long-term climate simulations where computational load is critical. You could highlight that even more.

Daewel et al.: Yes thank you. We will add this more clearly to the conclusion section.

3.) You mix up British and American English, e.g. P1L20: “analysed”, P1L24: “summarizes”. Please be consistent. –changed

4.) Also, please be consistent with “3d” vs. “3-d” / “end-to-end” vs. “End-to-End”. –will be changed

5.) Please capitalise “Figure 4” etc. –changed

6.) Commas are often missing in sentences which do not start with the subject, please add them. –corrected where commas are missing

Specific comments:

P1L20: “the observed pattern” -> “the observed patterns”? -ok

P1L26: “pattern agree” -> “patterns agree”? -ok

P2L10: “The differentiation of trophic levels” -> hard to understand what exactly you mean, can you rephrase? – Right! Changed to : “the separation of trophic levels ...”

P2L11: Citation missing in reference list. - added

P2L26: Please explain “foodweb models” as opposed to end-2-end models

Daewel et al: Here, a food web model is a model that resolves a complex food web on the basis of species or very specific functional groups. In principle a food web model does not necessarily need to be End-2-End. However the model Atlantis is both a food web model and End-2-End. We made that more clear in the ms.

P2L31: “based on environmental condition” -> “based on environmental conditions”? – changed

P2L31: “excluded” -> “excludes”? - changed

P3L25: “relative low” -> “relatively low”? - changed

P4L16-17: This is not a complete sentence. –combined with previous sentence

P4L21-22: Please cite Neumann and Schernewski (2008) who invented this approach in the model world. –added to the ms

P5L2-3: “with additional restriction as” -> “with the additional restriction that”?- changed

P5L8,10: Please use different symbols for the velocity vector and its vertical component. – changed

P5L10: A_v needs italics here. Actually, italics are missing a few times in this section when variables appear in the text. Please check. –changed

P5L15: “RC” needs formatting. - changed

P5L25: “the MB menu” -> “the MB diet”? –changed

P6L2: I suggest the use of X' instead of X to make clear it has nothing to do with the X in line 1. – The X in line 2 is the same X as in line 1. However, X was not clarified in the text. We modified the text as follows:

“Grazing rates GMB on prey type X ($X \in [Z_1; Z_2; P_1; P_2; DET; DOM; SED1]$) are estimated using ...”

P6L29: To be precise, the vertical integral of equation 1 reduces to the equation given in the text. Equation 1 itself would still keep a vertical migration term ($C w_m(z)$), even if the vertical migration velocity $w_m(z)$ is only known implicitly.

Daewel et al: Yes right that is not explained correctly. We changed the sentence to: “Following those three principles implies that equation 1 is simplified to $\frac{\partial C_{Fi}}{\partial t} + w_m(z) \frac{\partial C_{Fi}}{\partial z} = R_{Fi}$, where $w_m(z)$ is the vertical migration speed, which is given implicitly by the dynamical vertical distribution of the fish biomass in dependence of the vertical prey distribution. ...”

P7L13: Does this consumption even occur in anoxic layers? – No, fish consumption is confined to oxic conditions. We added the following sentence to the ms: “Note that, since fish do not tolerate anoxic conditions, only grid cells featuring positive oxygen concentrations were considered for the estimate of fish consumption.”

P7L14: There is no respiration of fish in the model? You rather treat fish respiration as an excretion and subsequent detritus mineralisation? Could you state this explicitly?

Daewel et al: The term excretion loss is related to respiration of MB and fish and does not include fecal excretion. It is considered directly as a source of dissolved inorganic nutrient (PO_4/NH_4 in principle also DIC if that were a state variable of the model)(see table 4 in the ms). See also explanation to P13L30-31 below.

Action: the role of the excretion term has now been clarified in the model description. We also modified table 4 since the reaction term for oxygen was not given in the earlier version of the ms.

P7L20: In your formulation, TK is always equal to 1/273.15 because T/T cancels out. This is certainly not correct.

Daewel et al: You are right. Thanks for pointing this out. The equation given

here is not correct. Will be changed to: $\varepsilon_{Fi} = \mu_{Fi} e^{\left(\frac{\theta_{Fi}}{k} TK\right)}$; $TK = \frac{T-T_0}{T \cdot T_0}$ with T is given in °K and $T_0=273.15$ °K.

Action: Equ. 11 has been changed.

P7L28-29: Which food web did the study consider? I assume this value differs a lot between different seas/regions. – The food web of the North Sea is considered in the study. - information has been added to the ms

P7L31: “is considerable higher” -> “is considerably higher with”? –changed

P7L31: Please state more explicitly that intraguild predation in zooplankton does not need to be represented in the model since it is not stage-resolving. –added to the ms: “Since the zooplankton model is not stage resolving intraguild predation is not explicitly prescribed as a mortality term, but is implicitly included in the background mortality.”

P8L21-27: Could you add a reference to the hydrodynamic model you used and the atmospheric forcing dataset? – More detail was added to the setup paragraph.

P9L2: “at each of the location” -> “at each location” / “at each of the locations”? - changed

P9L9: “on” -> “onto” - changed

P10L6: “equals” -> “equals to” -changed

P13L9: “concrete” -> “actual”? -changed

P13L12: “relative small” -> “relatively small”? –changed

P13L11-13: Could you give relative values (in % of average mass)?

Daewel et al: Yes we will give these values. “Values for both MB and fish biomass vary in the course of the year (Fig. 4). While the modelled seasonal amplitude for fish biomass is relatively small (North Sea: $94.7 \text{ mgC m}^{-3} \cong 11\%$; Baltic Sea: $84.8 \text{ mgC m}^{-3} \cong 9.7\%$ of the mean biomass) when compared to the average values, MB seasonality is substantial (North Sea: $4.4 \text{ gC m}^{-3} \cong 222\%$; Baltic Sea: $1.86 \text{ gC m}^{-3} \cong 183\%$ of the mean biomass).“ Following a suggestion by reviewer 2 we also added a discussion on the MB seasonality to a revised version of the ms.

P13L15: A minimum never falls, does it? A function of time falls until it reaches the minimum. – Yes, right. changed to “reach”

P13L30-31: Macrozoobenthos and fish can feed on their own excretions, since the energy contained in the carbon is not considered, correct? Also, there is no MB or fish respiration in the model, which would convert the organic carbon to DIC. So, there can be an infinite loop in the model where MB feeds and excretes, feeds and excretes, with no need to add energy from primary production. In Section 4 where you discuss future model improvements, could you comment on how to prevent this? I see, besides the way of explicitly defining “nutritional value classes” in detritus, the possibility to limit detritus consumption to ensure that at least a specific percentage of the total diet is fresh (non-detrital) material.

Daewel et al: We understand your concern, but we cannot agree on this

conclusion, and fear that the model description might be a bit unclear in that respect. On the one hand only parts of what is consumed can be assimilated while the rest can be considered as fecal excretion (this is probably what the reviewer is referring to). On the other hand the additional loss term for both fish and MB consists of a mortality and an excretion term. The mortality loss and fecal excretion contributes to the detritus pool, while the excretion loss is related to respiration of MB and fish and is considered directly as a source of dissolved inorganic nutrient (PO_4/NH_4 in principle also DIC if that were a state variable of the model)(see table 4 in the ms). Thus the feeding cycle features two specific loss term; i) the excretion term and ii) the remineralization of the dead organic matter, which would prevent the occurrence of such a self-containing feeding chain.

Action: We clarified these details in the model description of the ms (see also response to comment above P7L14). The role of the excretion term has been clarified. We also found an error in the last equation of table 4, which will be corrected.:” While fecal matter is accounted for through the use of assimilation efficiency, the excretion term from both fish and MB contributes directly to the nutrient reaction terms (see Table 4 equation for phosphate and ammonia).“

P13L33: This is because the North Sea is a tidal sea, correct? I would consider it as helpful for the non-European readers to state this difference here.

Daewel et al: Right, although the differences have been explained in the introduction, it is probably good to make that clear at this point.

Action: we changed the sentence to:” ... presumably due to the fact that a higher percentage of detritus is re-suspended in the tidal influenced, highly turbulent areas of the North Sea.“

P13L33: “Zooplankton and Phytoplankton is included” -> “Zooplankton and phytoplankton are included”? –changed

P15L5: “zooplankton form” -> “zooplankton forms”? –changed

P16L6-9: I would consider this as the main benefit of your model: You can obtain reasonable spatial patterns in zooplankton mortality without requiring data on planktivorous fish abundance. I would point this out already in the abstract and see it as the main point why your approach should be used.

Daewel et al: Thanks for pointing that out. We will follow the suggestion and add this more prominent to the abstract of the revised ms:” By using this approach we addressed the above-mentioned closure term problem in lower trophic ecosystem modelling at very low computational costs and thus provide an efficient method that requires very few data to obtain spatially and temporally dynamic zooplankton mortality.“

P16L12-16: Couldn't you split your model fish into pelagic and demersal “feeding groups” based on their diet-dependent vertical distribution and then compare the spatial patterns?

Daewel et al: We have tried to do so by differentiating between biomass in the water column and in the bottom layer (Suppl. Figure 2). However, clearly this is not directly comparable to the observation, where demersal and pelagic biomass is separated by species not actually by vertical appearance. Additionally the fish in our model has the chance to feed on all food sources at each time step.

Nonetheless we agree that this comparison can be helpful and thus will include it in a revised version of the manuscript together with a discussion. One thing that becomes quite apparent is, as already suggested in the discussion, that the pelagic fish biomass is underrepresented by the model, since, in contrast to the observations, the bottom biomass exceeds that of the water column biomass. On the other hand, similar to the observation, both stocks contribute to the latitudinal gradient in biomass.

Note that there was a unit error in the original version of the Figure 10d that was corrected in the new figure (Suppl. Figure 2).

Action: We modified the original figure 10 by the Suppl. Figure 2 and changed the discussion accordingly.

P16L25: What is zero at which boundary? – Here we mean that no fish enters the area through the open boundaries to the North Atlantic by e.g. migration. Action: we entered the following half sentence: , meaning that no fish enters or leave the model area over the lateral boundaries,

P17L8: “fish and MB is resolved” -> “are resolved”? - changed

P17L12: “for e.g.” -> “e.g.”, P18L31: the same - changed

Reply reviewer 2

Overall, this is a well written and interesting study linking a classic NPZD model to a single fish compartment. As a fisheries modeller and ecosystem modeller familiar with OSMOSE and Atlantic frameworks, I found the approach useful, however as the authors acknowledge, simplistic in its treatment of fish and fisheries. There is no mention of the impact that fisheries have on the ecosystem until the results section. There should be some introductory material about this as they are the biggest impact on fish populations, as many fisheries reduce more than half of fish biomass as a goal.

Daewel et al: As our model is formulated, fisheries can only be targeted in a very simplistic way. In particular species specific fish stock information are not possible through our functional group type approach. That is the reason why fisheries has not been addressed more extensively in the ms. However, we agree with the reviewer that fishing indeed plays an important role for the fish biomass in both ecosystems, North Sea and Baltic Sea, and that given additional consideration to the topic might become an additional component of the model (see conclusion). That is why we will follow the reviewer's suggestion and add some relevant sentences to the introduction. "The two systems also differ substantially in terms of ecosystem dynamics. The North Sea is known as a highly productive area inhabited by more than 26 zooplankton taxa (Colebrook et al., 1984) and over 200 fish species (Daan et al., 1990), with highest biomasses distributed among demersal gadoids, flatfish, clupeids and sandeel (*Ammodytes marinus*) (Daan et al., 1990). Consequently the North Sea is economically highly relevant with nine nations fishing in the area with landings of currently about 2 million tons annually (ICES, 2018b). Compared to the North Sea, species composition in the Baltic Sea is primarily limited by the low salinities and encompasses only a few key players for zooplankton (Möllmann et al., 2000) and fish (Fennel, 2010). Thus, compared to the North Sea, commercial fishing in the Baltic Sea includes only a few stocks with total landings of over 0.6 million tons annually (ICES, 2018a). Both regions have in common that landings peaked in the 1970s and have substantially (ca 50%) decline since then. Thus fishing has a substantial impact on the overall fish biomass in the region."

I would also like to see a more quantitative approach to calibrating the model to observed biomass of fish and fisheries catches. This data is spatially available, and a difference plot or map showing how well predicted vs observed fish biomasses compare in a spatially explicit analysis would be useful.

Daewel et al: We have deliberately avoided an active calibration of the model. For the following reasons: i) The emerging pattern for fish from our model cannot fully represent the actual observed fish biomass due to the specific restriction applied to the functional group in our model (see new ms methods page 7, lines 1-20). Therefore a quantitative calibration to observed pattern would not necessarily improve the models parameterization. ii) Other than the reviewer implies, we do not think that the relevant data (spatial distribution of total fish biomass) actually are available, even if our model were to simulate the real overall fish distribution. However, we agree with the reviewer that a spatial comparison to fish distribution is desirable. Therefore we chose to estimate fish biomass from the IBTS surveys (Figure 10a-c), which is the only long-term, consistently sampled dataset on fish available in the region and compared those sampled distributions qualitatively to our estimated distribution (Fig. 10 & discussion in paragraph 3.3). Fisheries catch data are probably not applicable since fishery mortality is not explicitly parameterized in the model. It is quite possible that there are relevant datasets available that we are not aware of and we would be grateful if the reviewer would share this information with us.

I'm also a bit concerned about the huge seasonality in biomass of the macrobenthos, much of which I presume is dominated by macroinvertebrates which don't vary in biomass as much as plankton communities do seasonally. & (page 13, line 11 - Is it reasonable for macrobenthos to vary that much annually? I would have expected there to be a much more constant standing stock similarly to fish. I'm thinking about the macro invertebrate community which doesn't vary that much through time.)

Daewel et al: See also comment above. We think that the reviewer's concern is valid in the way that the one functional group for MB might not adequately address all processes responsible for MB variation. However a comparison to observed seasonality shows that the seasonality for MB can be quite high in both systems. We will thus add a discussion at the respective position in the manuscript.: "This is in line with observations on seasonality of benthic infauna at three different locations in the North Sea published by Reiss and Krönke (2005), who found maximum biomass in late summer. Although the observed seasonality showed the highest magnitude in the German Bight the seasonality was clear at all three locations. The authors concluded that of the potential relevant factors (food availability/quality, water temperature, predation, hydrodynamic stress) food

quality plays the major role for infauna seasonality, thus is strongly related to primary production. They also suggest food limitation and predation pressure to be the main processes for decrease in abundance during winter. The same authors also looked at seasonality in the epibenthic community (Reiss and Krönke, 2004) showing that the epifaunal biomass varies less seasonally, especially in the off shore region, and that the main processes causing seasonal variations are related to migratory behaviour, which is not covered by our model. For the Baltic Sea only very local studies in seasonality of MB are available of which some indicate locally strong seasonality (Anders and Möller, 1983), while in other regions no seasonal changes in biomass were observed due to the dominance of long-lived species (Persson, 1983). In general the comparison to observations indicates that on the one hand the model is able to represent the main seasonality in MB even though epi- and infauna are not separated. On the other hand in future study the consideration of an additional functional group encompassing longer-lived species will be required for addressing MB seasonality more correctly.“

It's not clear to me why the Baltic and North Seas were combined into one model, as they exhibit very different environmental and fish production regimes.

Daewel et al: two reasons: First: Other than species specific models like OSMOSE and ATLANTIS, the methods we propose is based on a generic approach and shall be able to describe in a general way the transfer of matter and energy to the higher trophic levels. The specific advantage is, that it does not depend on species-specific information and the application to two systemically different regimes is in principle possible and a good test to the method and assumptions made here. Second: Even though North Sea and Baltic Sea are very different systems both in physical as well as in biological characteristics, they are dynamically tightly coupled to each other. On the one hand the Baltic Sea dynamics strongly determine the conditions in the Norwegian trench and are thus relevant for simulating the northern North Sea dynamics. On the other hand, the coupling to the North Sea is essential to simulate timing and characteristic of the Major Baltic Inflow events and hence is relevant for the dynamics in the Baltic Sea. Since the model setup used here is computationally relatively cheap and runs quickly, there is no need to separate two systems that are closely interlinked.

I think the closure terms where a lot of fish migration could be happening would be more important to focus on than connecting the two domains.

Daewel et al: As we commented on in the conclusions (old version of the ms P19 L5 et sqq.) we are aware that the assumption of “no migration” is a shortcoming of the method and needs to be addressed in continuative studies. It is however beyond the scope of this ms.

I would also like to see a more detailed treatment of fisheries mortality in the model, as this data is readily available and will be a huge driver of fish biomass given the very long exploitation history of the North Sea.

Daewel et al: A better representation of fisheries mortality is desirable but requires developmental work for integrating a dynamical representation of fisheries mortality and is thus beyond the scope of this ms (also see conclusion old version of the ms P19 L20 et sqq.).

Intro - You don't describe how this component contributes to the model and how the macrobenthos communities vary in the North and Baltic Seas.

Daewel et al: We agree that this has not been sufficiently addressed in the introduction. Action: We add additional information on macrobenthos in the North Sea and Baltic Sea to the introduction section:” Studies on the food web dynamics of the North Sea and Baltic Sea highlight additionally the relevance of benthic fauna for fish consumption (Greenstreet et al., 1997; Tomczak et al., 2012). The term benthos refers generally to all organisms inhabiting the sea floor. A comprehensive review on the topic has been given by Kröncke and Bergfeld (2003). The faunal components encompass over 5000 species generally divided by size into microfauna, meiofauna, and macrofauna. Additional differentiation can be made under consideration of the vertical habitat structure, with infauna inhabiting the inner part of the sediment and epifauna living above the sediments. While in the North Sea the macrobenthos assemblages are basically structured based on the spatial distribution of sediment characteristics and depth, the Baltic sea community is additionally influenced by oxygen availability (Ekeröth et al., 2016) and salinity (Gogina et al., 2010). Besides its role as prey and predator in the marine foodweb, macrobenthos additionally influences nutrient effluxes from the sediments and thus can modify temporal and spatial patterns in nutrient concentrations (Ekeröth et al., 2016). „

Results and Discussion - I suggest separating out your results first and then comparing to other studies. The way the two sections are intertwined makes it difficult to follow.

Daewel et al: We understand the reviewer's opinion. However, it is quite common in complex modeling studies to combine the results and discussion section, to avoid a repetition of major results in a separate discussion section. We thought carefully about the structure and still think that a combined section is more appropriate for this study.

page 2, lines 6-8 - the most common ecosystem models are Ecopath with Ecosim models which organize fish based on a combination of functional groups, species groups, and age-structured of species groups (see Tittensor et al. 2018 GMD).

Daewel et al: Right! A short paragraph on EwE will be included in the introduction." One of the most commonly used food-web modelling tools is the Ecopath with Ecosim (EwE) modelling software (Christensen and Walters, 2004), which provides an instantaneous snapshot of the trophic mass balance in marine food webs. Together with the dynamical modelling capability (Ecosim) and a tool that replicates the model on a spatial grid (Ecospace) it allows 2d estimates of the systems response to e.g. policy measures. However the approach still falls short in simulating ecosystem dynamics on high temporal and 3d spatial resolution."

page 3, lines 16-18 - How can questions about food webs be tested when there is only one fish functional group? There are different trophic levels of fish that are harvested which exert different controls on the macro food web. For example, forage fish have been shown to be important prey for many higher trophic levels and their exploitation has different effects than harvesting top predators (see Smith et al. 2010 Science).

Daewel et al: We agree that these kind of food-web interactions cannot be addressed by the current formulation of the model. However, the model presented in the ms is a first approach that can be further developed to represent a more complex food web (by e.g. distributing the fish group into separate feeding guilds), which can then be used to address this kind of question. We will make this more clearly in a revised version of the manuscript. Added to the introduction: "The approach we use cannot address changes in ecosystem structure related to variations in the fish assemblage or selected fishing activities. It does, however, provide the potential for further developments towards a more complex food web (e.g. by distributing fish into separate feeding guilds), which will then allow us to address specific changes in food web structure. "

page 4, line 1 - Awkward start to the sentence, suggest restructuring.

Daewel et al: Will be changed to: "Here we present a functional-type, E2E modelling approach, which relates food availability to potential fish growth and biomass distributions. In this manuscript we introduce..."

page 4, lines 16-17 - Awkward sentence, suggest combining with previous sentence in parentheses. - changed.

page 6, line 25 - What are the constraints for vertical fish movement based on oxygen and temperature limitations? This is one constraint of using only one fish group as there is a lot of variability among fish species in sensitivity to environmental conditions.

Daewel et al: Fish consumption is constrained by oxygen availability such the fish would not migrate into low oxygen regions. This information was missing in the ms and will be added to the method section. Temperature only plays a role for metabolic rates. : "Note that, since fish do not tolerate anoxic conditions, only grid cells featuring positive oxygen concentrations were considered for the estimate of fish consumption. ,,

The advantage of the generic functional group approach used in the model is that we simplify a complex community structure (also for phytoplankton and zooplankton) and reduce the information to the basic common features thus avoiding a huge parameter set and data requirements. Still the model is able to simulate relevant ecosystem dynamics.

We will add an additional explanation to the manuscript for clarification.

Action: the following sentence has been added to the conclusion:" The advantage of the generic functional group approach used in the model for all trophic levels is that we can simplify a complex community structure and reduce the information to the basic common features thus avoiding a huge parameter set and data requirements. Still the model is able to simulate relevant ecosystem dynamics on high spatial and temporal resolutions with relatively low computational requirements."

page 7, lines 17-18 - This sentence can be combined with the previous sentence. –changed

page 8, lines 1-12 - What about fisheries mortality on the fish compartment? Was this not included in the model? For the North Sea, this would comprise a significant proportion of total mortality.

Daewel et al: Fisheries has not been included in the reference simulation where mortality accounts for natural mortality and predation losses, which allows us to estimate the undisturbed “fish production potential”. We will clarify this in a revised version of the manuscript. To understand and clarify the impact of additional fisheries mortality we have additionally included the “fisheries scenarios” in the ms.

Action: Added to the method description: “Fisheries mortality has not been considered for the standard simulation, but was explicitly addressed in additional scenario experiments as described in section 2.3.”

page 10, lines 11-16 - This is the first mention of fisheries, if they are a component of the model, then they should have been included much earlier in the manuscript. How was the loss rate calculated? Much greater detail about the fisheries data that was used and how this was applied need to be included.

Daewel et al: Since simulating and understanding fisheries impacts on the ecosystem is not (and cannot be, due to the model simplifications applied) the primary research question addressed by the model, fisheries has not been addressed in the reference simulation (see comment above).

Therefore we included the scenarios to understand the effect of fisheries on the results. From estimated biomass and reported fisheries landings (see below our response to *page 12, line 28 - page 13, line 9*) we see that biomass losses due to fisheries are in both regions in the range of 20%-50% of the fish biomass per year. For convenience we decided to receive the fisheries mortality rate by scaling the natural mortality rate ($=0.001\text{d}^{-1}=0.365\text{yr}^{-1}$) with 0.5, 1 & 2. Thus receiving mortality rates that are representative for the bandwidth of the observed fisheries loss rates.

We will add a related explanation to a revised version of the manuscript.

Action: the following sentence has been added to the scenario description: “Following the fisheries overview of the North Sea and Baltic Sea region as published by ICES (2018a, 2018b) biomass losses due to fisheries are in both regions in the range of 20%-50% of the total fish biomass per year. For convenience we therefore decided to receive the fisheries mortality rate by scaling the natural mortality rate ($=0.001\text{d}^{-1}=0.365\text{yr}^{-1}$) with 0.5, 1 & 2.”

page 10, lines 18-23 - How was this value of 20% less loss rate derived? Any empirical data that supports this value?

Daewel et al: See explanation in the methods section (P7 and paragraph below)

page 12, lines 24-26 - This is a reason why it is important to consider functional groups/species/age classes/size classes when including fish in ecosystem models.

Daewel et al: As in many other models this depends on the research question that should be addressed with the model. If the main aim is indeed to estimate species-specific stock biomass or fisheries landings, we agree that the model requires a much greater detail than what we presented here. On the other hand if the question is related to the overall productivity of the ecosystem, energy and matter transfer processes and long-term variation of the latter, which is aimed for here, a more simplified approach with less data requirements and low computational costs is a helpful method. Of course that causes difficulties when comparing to observations. However, all models (also those that actually resolve species/age/size) are simplifications in a way and a large number of groups comes with a high degree of uncertainty in parameter settings and large computational costs.

page 12, line 28 - page 13, line 9 - How do the fish biomass estimates compare to reported fisheries landings from the regions? Is there enough biomass to support known landings?

Daewel et al: Following ICES, fisheries during the 1980's were in the range of 0.7-1 million tons in the Baltic Sea and 2-3 million tons in North Sea. Despite that the model underestimates fish production due to the no horizontal migration assumption and no fish migration over the lateral boundaries, the models' estimates of fish biomass in the North Sea: 5.13-10.27 million tons and in the Baltic Sea: 3.47-6.93 million tons would support the fisheries landing during that time. We will add explanations in the revised manuscript.

page 14, line 13 - You haven't introduced this analysis in the methods section to describe what it is.

Daewel et al: We understand the need for a more detailed description of the method. Since the method has been in detail introduced in an earlier manuscript, we suggest adding the method description to the ms as follows: The EOF analysis is a statistical method to understand major

modes of variability in multidimensional data fields. A detailed description on how this method has been applied is given in Daewel et al. (2015): “The annual values of the spatially explicit variable field form a $N \times M$ matrix χ (N : number of years; M : number of wet grid points). The empirical modes are given by the K eigenvectors of the covariance matrix with non-zero eigenvalues. Those modes are temporally constant and have the spatially variable pattern $p_k(m=1, \dots, M)$ where $k=1, \dots, K$. The time evolution $A_k(t=1, \dots, N)$ of each mode can then be obtained by projecting $p_k(m)$ onto the original data field χ such that $\chi(t, m) = \sum_{k=1}^K p_k(m) A_k(t)$. In the following we will refer to $A_k(t)$ as the principal components (PC) and to $p_k(m)$ as empirical orthogonal function (EOF). The percentage of the variance of the field χ explained by mode k is determined by the respective eigenvalues and is referred to as the global explained variance $\eta_g(k)$. Before using the method to analyse the spatiotemporal dynamics of the field, the data were demeaned (to account for the variability only) and normalized (to allow an analysis of the variability independent of its amplitude). The identified modes are not necessarily equally significant in all grid points of the data field. Thus, the local explained variance $\eta_{local,k}(m)$ could provide additional information about the regional relevance of an EOF mode and the corresponding PC in percent:

$$\eta_{local}^k(m) = \left[1 - \frac{\text{Var}(\chi(m,t) - p^k(m)A^k(t))}{\text{Var}(\chi(m,t))} \right] \cdot 100, \quad (12)$$

where $\text{Var}(X) = \sum_{t=1}^N (\bar{X} - X(t))^2$ denotes the variance of the field $X(t)$.”

New references

- Anders, K. and Möller, H.: Seasonal fluctuations in macrobenthic fauna of the Fucus belt in Kiel Fjord (western Baltic Sea), *Helgoländer Meeresuntersuchungen*, 36, 277–283, 1983.
- ICES: Baltic Sea Ecoregion – Fisheries overview, (November), 1–23, doi:<https://doi.org/10.17895/ices.pub.4648>, 2018a.
- ICES: Greater North Sea Ecoregion – Fisheries overview, (November), 1–31, doi:<https://doi.org/10.17895/ices.pub.4647>, 2018b.
- Persson, L.-E.: Temporal and spatial variation in coastal macrobenthic community structure, Hanö bay (southern Baltic), *J. Exp. Mar. Biol. Ecol.*, 68, 277–293, 1983.
- Reiss, H. and Krönke, I.: Seasonal variability of epibenthic communities in different areas of the southern North Sea, *ICES J. Mar. Sci.*, 61, 882–905, doi:10.1016/j.icesjms.2004.06.020, 2004.
- Reiss, H. and Krönke, I.: Seasonal variability of infaunal community structures in three areas of the North Sea under different environmental conditions, *Estuar. Coast. Shelf Sci.*, 65, 253–274, doi:10.1016/j.ecss.2005.06.008, 2005.

Towards End-2-End modelling in a consistent NPZD-F modelling framework (ECOSMOE2E_vs1.0): Application to the North Sea and Baltic Sea

Ute Daewel¹, Corinna Schrum^{1,2}, Jed Macdonald^{3,4}

¹Helmholtz Centre Geesthacht, Institute of Coastal Research, Max-Planck-Str. 1, 21502 Geesthacht, Germany

²Geophysical Institute, University of Bergen, Allegaten 41, 5007 Bergen, Norway

³Faculty of Life and Environmental Sciences, University of Iceland, 101 Reykjavik, Iceland

⁴Oceanic Fisheries Programme, Pacific Community (SPC), Noumea BP D5 98848, New Caledonia

Correspondence to: Ute Daewel (ute.daewel@hzg.de)

Abstract. Coupled physical-biological models usually resolve only parts of the trophic food chain and hence, run the risk of neglecting relevant ecosystem processes. Additionally, this imposes a closure term problem at the respective “ends” of the considered trophic levels. Here we aim to understand how the implementation of higher trophic levels in a NPZD model affects the simulated response of the ecosystem, by using a consistent NPZD-Fish modelling approach (ECOSMO E2E) in the combined North Sea and Baltic Sea system. By using this approach we addressed the above-mentioned closure term problem in lower trophic ecosystem modelling at very low computational costs and thus provide an efficient method that requires very few data to obtain spatially and temporally dynamic zooplankton mortality.

On the basis of the coupled ecosystem model ECOSMO II we implemented one functional group that represents fish and one group representing macrobenthos in the 3d model formulation. Both groups are linked to the lower trophic levels and to each other via predator-prey relationships, thus allowing the investigation of both, bottom-up processes and top-down mechanisms in the trophic chain of the North Sea and Baltic Sea ecosystem. Model results for a ten-year long simulation period (1980-1989) were analysed and discussed with respect to the observed patterns. To understand the impact of the newly implemented functional groups for the simulated ecosystem response, we compare the performance of the ECOSMO E2E to that of a respective truncated NPZD model (ECOSMO II) applied to the same time period. Additionally, we performed scenario tests to analyse the new role of the zooplankton mortality closure term in the truncated NPZD and the fish mortality term in the End-to-End model, which summarizes pressure imposed on the system by fisheries and mortality imposed by apex predators.

We found that the model-simulated macrobenthos and fish spatial and seasonal patterns agree well with current system understanding. Considering a dynamic fish component in the ecosystem model resulted in slightly improved model performance with respect to representation of spatial and temporal variations in nutrients, changes in modelled plankton seasonality and nutrient profiles. Model sensitivity scenarios showed that changes in the zooplankton mortality parameter are

Ute Daewel 13.3.2019 15:19

Deleted: School of BioSciences, The University of Melbourne, Parkville, Victoria 3052, Australia

transferred up and down the trophic chain with little attenuation of the signal, while major changes in fish mortality and in fish biomass cascade down the food chain.

1 Introduction

5 The majority of spatially-resolved marine ecosystem models are dedicated to a specific part of the marine food web. These models can be differentiated into lower trophic level Nutrient-Phytoplankton-Zooplankton models (so called NPZ models or LTL models) (e.g. Blackford et al., 2004; Daewel and Schrum, 2013; Maar et al., 2011; Schrum et al., 2006; Skogen et al., 2004) and, on the other end of the trophic chain, higher trophic level models (HTL models). The latter mainly simulate fish on a species level including both single-species Individual-Based Models (IBMs, e.g. Daewel et al., 2008; Megrey et al., 10 2007; Politikos et al., 2018; Vikebø et al., 2007), and multi-species models. Although some of these models are complex and already include many food-web components such as OSMOSE (Shin and Cury, 2004, 2001) or ERSEM (Butenschön et al., 2016), the [separation](#) of trophic levels often constrains such models' ability to simulate and distinguish between major control mechanisms on marine ecosystems (Cury and Shannon, 2004). The difficulty to resolve trophic feedback mechanisms increases the uncertainties when modelling the impacts of external controls on the trophic food chain (e.g. 15 Daewel et al., 2014; Peck et al., 2015).

In the last 10 to 15 years, major efforts have been made to link the different trophic levels together to cover the marine ecosystem from the lowest to the uppermost "end" (E2E) (Christensen and Walters, 2004; Fennel, 2009; Fulton, 2010; Heath, 2012; Shin et al., 2010; Travers et al., 2007; Watson et al., 2014). Although, some models such as Atlantis (Fulton et al., 2005), StrathE2E (Heath, 2012) or Ecopath with Ecosim (EwE) (Christensen and Walters, 2004) consider more trophic 20 levels (from phytoplankton to marine mammals and birds and/or fisheries) consistently within the model formulation, the majority of approaches couple conceptually different model types, either 'one-way' with no feedback on the lower trophic levels (e.g. Daewel et al., 2008; Rose et al., 2015; Utne et al., 2012), or 'two-way' (e.g. Megrey et al. 2007, Oguz et al. 2008), when linking the trophic levels. All these approaches work reasonably well in serving a specific purpose or scientific question, but are accompanied by different uncertainties and conceptual limitations to model ecosystem structuring under 25 external forcing. While one-way coupled approaches neglect feedbacks and hence imply difficulties at the model interfaces, [comprehensive food-web models like Atlantis resolve food webs on the basis of species or specific groups and are difficult to parameterize especially in complex ecosystems. One of the most commonly used food-web modelling tools is the Ecopath with Ecosim \(EwE\) modelling software \(Christensen and Walters, 2004\), which provides an instantaneous snapshot of the trophic mass balance in marine food webs. Together with the dynamical modelling capability \(Ecosim\) and a tool that replicates the model on a spatial grid \(Ecospace\) it allows 2d estimates of the systems response to e.g. policy measures. 30 However the approach still falls short in simulating ecosystem dynamics on high temporal and 3d spatial resolution.](#) In Peck et al. (2015), a number of different ecosystem models used in the European project VECTORS ([| |
|---|
| Ute Daewel 12.2.2019 14:07 |
| Deleted: differentiation |
| Unknown |
| Field Code Changed |
| Ute Daewel 12.2.2019 14:17 |
| Deleted: \(Curry and Shannon, 2004\) |
| Unknown |
| Field Code Changed |
| Ute Daewel 6.3.2019 16:27 |
| Deleted: \(Fennel, 2009; Fulton, 2010; Heath, 2012; Shin et al., 2010; Travers et al., 2007; Watson et al., 2014\) |
| Ute Daewel 4.3.2019 16:54 |
| Formatted: English \(UK\) |
| Unknown |
| Field Code Changed |
| Ute Daewel 6.3.2019 16:27 |
| Deleted: or |
| Unknown |
| Field Code Changed |
| Unknown |
| Field Code Changed |
| Unknown |
| Field Code Changed |
| Ute Daewel 14.3.2019 10:49 |
| Deleted: comprehensive E2E models like Atlantis |
| Unknown |
| Field Code Changed |](http://www.marine-</p>
</div>
<div data-bbox=)

vectors.eu/) were reviewed. Besides discussing statistical and physiology-based life cycle models, Peck et al. (2015) identified strength and weaknesses in food web models like Atlantis. While the strength of these models is the explicit consideration of species-specific responses, which are often vital for informing advice for management, a clear weakness of those models is the huge amount of data needed for model parameterization (Peck et al., 2015) and the sensitivity to assumptions made on the food-web structure and functioning. Another drawback of the recent End-2-End models is the lack in spatial resolution. They are either solved in 2d as EwE or, as Atlantis is, resolved in predefined (based on environmental conditions) larger area polygons. This consequently excludes the dynamic resolution of ecologically highly-relevant hydrographical structures such as tidal fronts or the thermocline, and implies that future changes in relevant hydrodynamics and their impacts cannot be considered through these models. To our knowledge, the first approach that attempted to resolve the trophic food web more consistently in a functional group framework with the spatial and temporal resolution of a state of the art physical model is the food web model presented by Fennel (2010, 2008) and Radtke et al. (2013). They proposed a nutrient-to-fish model where fish is included consistently in a NPZD model framework using a Eulerian approach. In their study they chose a species-specific way to introduce fish using size-structured formulations for the three major fish species in the Baltic Sea. The model has been proven to work well in the Baltic Sea, which is characterized by a relative simple food web (Casini et al., 2009; Fennel, 2008), but is likely more difficult to parameterize in other, more complex structured food webs involving more key species such as the North Sea.

Here, we aim to address these conceptual limitations in End-to-End modelling and present a different approach based on the assumption that food availability and corresponding energy and mass fluxes are the major controls in higher trophic production and spatial and temporal distribution of the fish biomass. A physical-biological coupled 3d NPZD ecosystem model is extended to include fish and macrobenthos (MB). Although our idea is inspired by the model presented by Fennel (2010, 2008) and Radtke et al. (2013), it is substantially different to their concept based on three key species. We used instead a functional group approach, which represents the entire fish population, and aims to be consistent with the functional group approach used for phytoplankton and zooplankton. This enables estimation of the total fish production potential and allows for resolving structuring impacts on the ecosystem. The advantage of this generic approach is its broad applicability. It allows for general and comparative studies on changing ecosystem structure and is not limited by unknown changes in key species for the respective ecosystems. The approach we use cannot address changes in ecosystem structure related to variations in the fish assemblage or selected fishing activities. It does, however, provide the potential for further developments towards a more complex food web (e.g. by distributing fish into separate feeding guilds), which will then allow us to address specific changes in food web structure.

We will present a first application of the Eulerian End-to-End model for the shelf sea system of the North Sea and Baltic Sea. The coupled North Sea and Baltic Sea system is located adjacent to the North Atlantic Ocean. Despite their close proximity, they are very different in physical and biogeochemical characteristics. The North Sea features pronounced co-oscillating tides combined with a major inflow from the North Atlantic. The Baltic Sea in contrast, has only a narrow opening to the North Sea leading to an almost enclosed, brackish system with weak tidal forcing (Müller-Navarra and Lange,

Ute Daewel 6.3.2019 15:41
Deleted: h
Unknown
Field Code Changed

Ute Daewel 14.3.2019 10:54
Deleted: and End-2-End type
Ute Daewel 14.2.2019 20:05
Deleted: both
Ute Daewel 14.2.2019 20:05
Deleted: types
Unknown
Field Code Changed

Ute Daewel 13.2.2019 09:39
Deleted: d
Ute Daewel 13.3.2019 15:19
Deleted: of
Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Ute Daewel 4.3.2019 16:58
Deleted: e
Ute Daewel 4.3.2019 16:58
Deleted: e
Ute Daewel 4.3.2019 16:57
Deleted: -
Unknown
Field Code Changed

Ute Daewel 6.3.2019 16:57
Deleted: -
Unknown
Field Code Changed

2004). The restricted exchange capacities and fresh water excess to the Baltic Sea leads to an estuarine-type circulation with strong stratification and relatively low salinities and to a relatively long water residence time of about 30 years (Omstedt and Hansson, 2006; Rodhe et al., 2006). Due to its brackish waters, winter sea ice develops regularly in the Baltic Sea, which might, in severe winters cover almost the entire surface (Seinä and Palosuo, 1996).

5 The two systems also differ substantially in terms of ecosystem dynamics. The North Sea is known as a highly productive area inhabited by more than 26 zooplankton taxa (Colebrook et al., 1984) and over 200 fish species (Daan et al., 1990), with highest biomasses distributed among demersal gadoids, flatfish, clupeids and sandeel (*Ammodytes marinus*) (Daan et al., 1990). Consequently, the North Sea is economically highly relevant with nine nations fishing in the area with landings of currently about 2 million tons annually (ICES, 2018b). Compared to the North Sea, species composition in the Baltic Sea is

10 primarily limited by the low salinities and encompasses only a few key players for zooplankton (Möllmann et al., 2000) and fish (Fennel, 2010). Thus, compared to the North Sea, commercial fishing in the Baltic Sea includes only a few stocks with total landings of over 0.6 million tons annually (ICES, 2018a). In both regions, landings peaked in the 1970s and have substantially (ca 50%) declined since then. Thus, fishing has a substantial impact on the overall fish biomass in the region. Studies on the food web dynamics of the North Sea and Baltic Sea highlight additionally the relevance of benthic fauna for fish consumption (Greenstreet et al., 1997; Tomeczak et al., 2012). The term benthos refers generally to all organisms inhabiting the sea floor. A comprehensive review on the topic has been given by Kröncke and Bergfeld (2003). The faunal components encompass over 5000 species generally divided by size into microfauna, meiofauna, and macrofauna. Additional differentiation can be made under consideration of the vertical habitat structure, with infauna inhabiting the inner part of the sediment and epifauna living above the sediments. While in the North Sea the macrobenthos assemblages are

20 structured based on the spatial distribution of sediment characteristics and depth, the Baltic sea community is additionally influenced by oxygen availability (Ekeröth et al., 2016) and salinity (Gogina et al., 2010). Besides its role as prey and predator in the marine foodweb, macrobenthos additionally influences nutrient effluxes from the sediments and thus can modify temporal and spatial patterns in nutrient concentrations (Ekeröth et al., 2016).

25 Here we present a functional-type, E2E modelling approach, which relates food availability to potential fish growth and biomass distributions. In this manuscript we introduce the conceptual basis of the model, discuss its characteristics and explore its performance with respect to observed fish and MB distributions. Further, we analyse model performance at the lower trophic levels in comparison to the NPZD modelling approach, and discuss the potential of our model to understand and compare basic regional ecosystem characteristics.

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Ute Daewel 11.2.2019 20:32
Moved (Insertion) [1]

Unknown
Field Code Changed

Ute Daewel 11.2.2019 20:32
Deleted: Benthic fauna has been observed to exhibit little tolerance to hypoxic and anoxic conditions (Kröncke and Bergfeld, 2003), wherefore macrobenthos growth was estimated only for positive oxygen concentrations in the model framework.

Unknown
Field Code Changed

Ute Daewel 8.2.2019 12:48
Deleted: The here

Ute Daewel 8.2.2019 12:48
Deleted: ed

Ute Daewel 8.2.2019 12:50
Deleted: In this study, we

Ute Daewel 8.2.2019 12:50
Deleted: present

2 Methods

2.1 Model Description

The E2E model builds on the coupled hydrodynamic-lower trophic level ecosystem model ECOSMO II (Barthel et al., 2012; Daewel and Schrum, 2013; Schrum et al., 2006; Schrum and Backhaus, 1999), which is further expanded for the present study. The latter model has been shown to accurately reproduce lower trophic level ecosystem dynamics in the coupled North Sea and Baltic Sea system. The model equations and a model validation on the basis of nutrients were presented in detail by Daewel & Schrum (2013), who showed that the model is able to reasonably simulate ecosystem productivity in the North Sea and Baltic Sea on seasonal up to decadal time scales. The NPZD module was designed to simulate different macronutrient limitation processes in targeted ecosystems and comprises 16 state variables. Besides the three relevant nutrient cycles (nitrogen, phosphorus and silica), three functional groups of primary producers (diatoms, flagellates & cyanobacteria) and two zooplankton (herbivore & omnivore) groups were resolved. Additionally, oxygen, biogenic opal, detritus and dissolved organic matter were considered. Sediment is implemented in the model as an integrated surface sediment layer, which accounts for consideration of sedimentation as well as resuspension (Neumann and Schernewski, 2008). Biogeochemical remineralisation is considered in surface sediments leading to inorganic nutrient fluxes into the overlying water column. To allow for nutrient specific processes in the sediment, the organic silicate content of the sediment is estimated in as separate state variable. A third sediment compartment is considered for iron-bound phosphorus in the sediment. To estimate total fish production and biomass in a consistent manner compared to lower trophic level production, we expanded the NPZD-type model by implementation of a wider food web in the system.

Although zooplankton in ECOSMO II could in principle grow also at the bottom, its parameterization as a passive tracer and the choice of parameterization for the functional groups makes it unsuitable for representing benthic production. Hence, the parameterization of a specific functional group representing benthic (meio- and macro-) fauna remains necessary. The group was designed similar to the zooplankton groups, but with the additional restriction that benthos grows only at the bottom and is neither exposed to advection nor diffusion. Benthic fauna has been observed to exhibit little tolerance to hypoxic and anoxic conditions (Kröncke and Bergfeld, 2003), wherefore macrobenthos growth was estimated only for positive oxygen concentrations in the model framework. In contrast to the benthic compartment in ERSEM (Butenschön et al., 2016), where the benthic predators are distributed into three different functional types, we here neglect different functional traits of infauna and epifauna and consider only one functional group, which we, for convenience, will refer to as macrobenthos (MB).

Each state variable C in ECOSMO II is estimated following prognostic equations in the form

$$C_t + (\vec{v} \cdot \nabla)C + (w_d)C_z = (A_v C_z)_z + R_c \quad (1)$$

with $C_t = \frac{\partial C}{\partial t}$, $C_z = \frac{\partial C}{\partial z}$ where t is time and z is the vertical coordinate. The equation includes advective transport ($\vec{v} \cdot \nabla$) ($\vec{v} = (u, v, w)$ current velocity vector), vertical turbulent sub-scale diffusion ($(A_v C_z)_z$ (A_v : turbulent sub-scale diffusion coefficient), sinking rates $(w_d)C_z$ (w_d : sinking rate is non-zero only for detritus, opal and cyanobacteria)) and chemical and

Unknown
Field Code Changed

Unknown
Field Code Changed

Ute Daewel 8.2.2019 12:52
Deleted: Whereof the zooplankton was divided into a herbivorous and an omnivorous group.

Unknown
Field Code Changed

Ute Daewel 4.3.2019 16:54
Formatted: English (UK)

Ute Daewel 11.2.2019 20:32
Moved up [1]: Studies on the food web dynamics of the North Sea and Baltic Sea highlight the relevance of benthic fauna for fish consumption (Greenstreet et al., 1997; Tomczak et al., 2012). Benthic fauna has been observed to exhibit little tolerance to hypoxic and anoxic conditions (Kröncke and Bergfeld, 2003), wherefore macrobenthos growth was estimated only for positive oxygen concentrations in the model framework. The term benthos refers generally to all organisms inhabiting the sea floor. A comprehensive review on the topic has been given by Kröncke and Bergfeld (2003). The faunal components encompass over 5000 species generally divided by size into microfauna, meiofauna, and macrofauna. Additional differentiation can be made under consideration of the vertical habitat structure, with infauna inhabiting the inner part of the sediment and epifauna living above the sediments.

Ute Daewel 13.2.2019 10:02
Deleted: as

Unknown
Field Code Changed

Unknown
Field Code Changed

Ute Daewel 13.2.2019 10:04
Deleted: v

Ute Daewel 13.2.2019 10:06
Deleted: v

Ute Daewel 4.3.2019 16:54
Formatted: Font:Italic

biological reactions R_c . Since MB production occurs locally at the bottom and the group is not exposed to mechanical displacement, Eq. 1 is simplified for MB to

$$\frac{dC_{MB}}{dt} = [R_{MB}]_{z=z_{bottom}} \quad (2)$$

Concurrently chemical and biological interactions are employed in the biological reaction term R_c , which is different for each variable (C) based on the relevant biochemical processes. For MB it divides into production, which is a function of consumption (R_{MB_Cons}) and assimilation efficiency (γ_{MB}) and a MB loss term.

$$R_{MB} = \gamma_{MB} R_{MB_Cons} - R_{MB_Loss} \quad (3)$$

The production of zooplankton in the model depends on the available food resources, which include phytoplankton, detritus, and, for the omnivorous zooplankton group, also herbivorous zooplankton. For the macrobenthos functional group, we assume a much wider range of potential prey items. The benthic community can be divided into benthic suspension/filter feeders feeding mainly on phytoplankton, detritus and bacteria, benthic deposit feeders ingesting bottom sediments, and larger individuals exerting predation pressure (among others) on the available zooplankton (Kröncke and Bergfeld, 2003). Thus, the prey spectrum of the simulated MB functional group includes, besides phyto- and zooplankton, also detritus and organic sediments. Since we assume that benthic suspension/filter feeders would also indirectly ingest dissolved organic matter, we chose to add the latter to the MB [diet](#).

Consumption of the MB group is estimated as the sum of consumption rates for the single prey items (herbivorous zooplankton (Z_1), omnivorous zooplankton (Z_2), flagellates (P_1), diatoms (P_2), detritus (DET), dissolved organic matter (DOM), organic sediment (SED1))

$$R_{MB_Cons} = C_{MB} \left(\sum_{i=1}^2 G_{MB}(C_{Z_i}) + \sum_{l=1}^2 G_{MB}(C_{P_l}) + G_{MB}(C_{DET}) + G_{MB}(C_{DOM}) + G_{MB}(C_{SED1}) \right) \quad (4)$$

Grazing rates G_{MB} on prey type X ($X \in \{Z_1; Z_2; P_1; P_2; DET; DOM; SED1\}$) are estimated using the Michaelis-Menten equation (Michaelis and Menten, 1913; Monod, 1942):

$$G_{MB}(C_X) = \sigma_{MB,X} \frac{a_{MB,X} C_X}{r_{MB,X} + F_{MB}} \quad (5)$$

where $F_{MB} = \sum_X a_{MB,X} C_X$.

The half saturation constant r_{MB} and values for grazing rates ($\sigma_{MB,X}$) are given in [Table 1](#) and feeding preferences ($a_{MB,X}$) are given in [Table 2](#).

The MB loss term consists of excretion ($\epsilon_{MB} C_{MB}$), natural mortality ($m_{MB} C_{MB}$) and predation mortality from the fish functional group ($C_{Fi} G_{Fi}(C_{MB})$). Values for excretion (ϵ_{MB}) and mortality rate (m_{MB}) are given in table 1.

$$R_{MB_Loss} = C_{Fi} G_{Fi}(C_{MB}) + m_{MB} C_{MB} + \epsilon_{MB} C_{MB} \quad (6)$$

We assume that fish is a prognostic variable, which is, in contrast to the other prognostic state variables in the ecosystem model, not exposed to passive transport processes and not moving actively horizontally. This can be translated into the assumption that characteristic fish migration is restricted to a spatial scale below the model grid size [in the order of 10km](#).

Ute Daewel 4.3.2019 16:54

Formatted: Font:Italic

Ute Daewel 4.3.2019 16:54

Formatted: Font:Italic, Subscript

Ute Daewel 4.3.2019 16:54

Formatted: Font:Italic

Ute Daewel 4.3.2019 16:54

Formatted: Font:Italic

Ute Daewel 4.3.2019 16:54

Formatted: Font:Italic, Subscript

Unknown

Field Code Changed

Ute Daewel 13.2.2019 10:08

Deleted: menu

Ute Daewel 4.3.2019 17:10

Formatted: Font:Italic

Unknown

Field Code Changed

Ute Daewel 13.3.2019 15:24

Deleted: table

Ute Daewel 13.3.2019 15:24

Deleted: table

Ute Daewel 13.3.2019 15:25

Deleted: O(

Ute Daewel 13.3.2019 15:25

Deleted:)

10

- 20

25

30

30

30

Ute Daewel 13.3.2019 15:25
Deleted: to
Ute Daewel 15.2.2019 11:20
Deleted: diversity
Ute Daewel 15.2.2019 11:23
Deleted: potentially emerging
Ute Daewel 15.2.2019 12:53
Deleted: However, this
Ute Daewel 15.2.2019 10:35
Deleted: The first principle
Ute Daewel 15.2.2019 10:35
Deleted: reduced
Ute Daewel 15.2.2019 10:36
Deleted: $\frac{dC_{PI}}{dt}$
Ute Daewel 15.2.2019 10:51
Deleted: in
Ute Daewel 4.3.2019 17:11
Deleted: .
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 16:54
Formatted
Ute Daewel 4.3.2019 17:11
Formatted: Normal1, Justified, Line spacing: 1.5 lines, Tabs: 12.75 cm, Left

Where P_X (X is one of four prey types (herbivorous zooplankton Z_1 , omnivorous zooplankton Z_2 , detritus DET, macrobenthos MB) available for fish in the model) is defined as the integrated biomass of the prey type X ($P_X = \sum_{k=1}^{k_{\max}} (C_{Xk} * \Delta z_k)$) over all vertical levels ($k=1:k_{\max}$) at the respective horizontal location (m,n). P_{Fi} is the corresponding vertically integrated biomass of fish. Grazing rates G_{Fi} are estimated using the Michaelis-Menten equation $G_{Fi}(P_X) = \sigma_{Fi,X} \frac{a_{Fi,X} P_{Fi}}{r + P_X}$ with $F = \sum_X a_{Fi,X} P_X$ and $a_{Fi,X}$ are the feeding preferences of fish on prey type X (values in Table 2), in a similar manner as for the zooplankton and MB groups.

In a second step, fish consumption in each grid box (m,n,k) is estimated by weighting the prey specific components of the consumption in each vertical layer based on the vertical distribution of the prey biomass with $C_X(m,n,k)/P_X(m,n)$ such that

$$R_{FiCons}(m,n,k) = P_{Fi} \left(\sum_{j=1}^2 G_{Fi} \left(P_{Zj} \right) * \frac{C_{Zj}}{P_{Zj}} + G_{Fi}(P_{DET}) * \frac{C_{DET}}{P_{DET}} + [G_{Fi}(P_{MB})]_{k=bottom} \right) \quad (9)$$

Note that, since fish do not tolerate anoxic conditions, only grid cells featuring positive oxygen concentrations were considered for the estimate of fish consumption.

The loss term for fish includes mortality and excretion.

$$R_{FiLoss} = m_{Fi} C_{Fi} + \varepsilon_{Fi} C_{Fi} \quad (10)$$

Mortality is considered as a linear mortality rate including biomass losses due to natural mortality and predation. Fisheries mortality has not been considered for the standard simulation, but was explicitly addressed in additional scenario experiments as described in section 2.4. Excretion is considered to be related to fish metabolism and consequently to respiration (Table 4 equation for oxygen) and hence has been parameterized in dependence of temperature (Clarke and Johnston, 1999; Gillooly et al., 2001). Reaction kinetics vary with temperature according to the Boltzmann's factor k and we formulated the fish excretion as

$$\varepsilon_{Fi} = \mu_{Fi} e^{\left(\frac{\theta_{Fi} L_{TK}}{k} \right)}; TK = \frac{T - T_0}{T - T_0} \text{ with } T \text{ is given in } ^\circ K \text{ and } T_0 = 273.15 ^\circ K. \quad (11)$$

All rates are given in Table 3.

Fish and macrobenthos predation, excretion and mortality are considered in addition to the pelagic lower trophic level biological reaction terms (see Daewel and Schrum, 2013), for nutrients, phytoplankton, zooplankton, detritus, dissolved organic matter or sediment (Table 4). While fecal matter is accounted for through the use of assimilation efficiency, the excretion term from both fish and MB contributes directly to the nutrient reaction terms (see Table 4 equation for phosphate and ammonia). The new zooplankton mortality term consists of fish predation and additional background mortality, which is 80% of the background mortality term used in ECOSMO II. In-situ and laboratory studies indicate that predation mortality accounts for 67-75% of the total mortality (Hirst and Kiørboe, 2002). Other sources of mortality are parasitism, disease and starvation. Including fish and macrobenthos as predators in the model does not, however, account for the overall predation exerted on zooplankton. By analyzing the pelagic food web of the North Sea Heath (2005) identified fish consumption of

Ute Daewel 4.3.2019 17:12

Formatted

[1]

Ute Daewel 4.3.2019 17:12

Formatted: Font:Italic

Ute Daewel 13.3.2019 15:27

Deleted: table

Ute Daewel 4.3.2019 16:54

Formatted

[2]

Ute Daewel 15.2.2019 14:15

Deleted: While the m...ortality is consider

[3]

Unknown

Field Code Changed

Ute Daewel 8.2.2019 13:12

Deleted: Following Gillooly et al. (2001).

Ute Daewel 13.2.2019 11:02

[4]

Ute Daewel 4.3.2019 16:54

Formatted

[5]

Ute Daewel 13.2.2019 11:03

Deleted: given in °K...

[6]

Ute Daewel 13.3.2019 15:28

Deleted: table

Ute Daewel 13.3.2019 15:31

Deleted: ...ADDIN CSL_CITATION

[7]

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

omnivorous zooplankton being on average $6.7 \text{ gC m}^{-2} \text{ year}^{-1}$. That value was recalculated in Heath (2007), after more specifically considering the role of fish pre-recruits feeding on zooplankton, to amount to $\sim 7.6 \text{ gC m}^{-2} \text{ year}^{-1}$, while the average consumption by carnivorous zooplankton (euphausiids and macroplankton) is considerably higher with $11 \text{ gC m}^{-2} \text{ year}^{-1}$. Since the zooplankton groups in the model are not stage resolving, intraguild predation is not explicitly prescribed as a mortality term, but is implicitly included in the background mortality. Although our model results also suggest that a substantial amount of zooplankton is consumed by macrobenthos in the shallow regions of the North Sea, assuming that about 20-30% of zooplankton mortality stems from the combined fish and benthos group seems to be a good first guess. Please consider that the reduction of the background mortality rate to 80% of its initial value does not necessary imply that the background mortality is 80% of the total mortality. By including a spatially and temporally variable mortality term in the model, this term can locally play a much larger (or smaller) role for the overall mortality. To evaluate the model sensitivity to the choice of this parameter, we performed scenario experiments described in section 2.4. The degradation products from MB, fish mortality and food consumption contribute to particulate (POM) and dissolved organic matter (DOM). The latter is distributed between the two partitions POM and DOM with a ratio 60%/40% (for explanation see Daewel and Schrum 2013b). Since MB is living at the sea floor we assume that the generated POM is directly contributing to the sediment pool (SED1), which via re-suspension might contribute to the suspended particulate matter (DET) in dependence of bottom stress. The fish contribution to the particulate organic matter is added to the detritus pool.

2.2 Experimental Setup

To evaluate the model performance after including two new functional groups, we chose to analyse model results from a 10-year long simulation period (1980-1989) based on two key requirements. First, since the characteristic time scale of the Baltic Sea is in the range of three decades and also the model results indicate an adaptation period of about 20-30 years for fish and MB in the Baltic Sea (not shown), the simulation was started in 1948 to allow for a sufficiently long spinup period with a realistic forcing. Second, we wanted to analyse a relatively undisturbed period with respect to hydrodynamic and biogeochemical conditions. Therefore we chose a period prior to the observed regime shift at the end of the 1980's and prior to the major Baltic inflow in 1993.

The model setup is similar to the one described by Daewel and Schrum (2013) for the long term simulations, with a hydrodynamic core model based on HAMSOM (Hamburg Shelf Ocean Model) as described in (Schrum and Backhaus, 1999) with additional modification of the advection scheme (Barthel et al., 2012). The model is formulated on a staggered Arakawa-C grid with a horizontal resolution of $6' \times 10'$ ($\sim 10\text{km}$) and a 20-min time step. The vertical dimension was resolved with 20 vertical levels, whereof the upper 40 m have a layer thickness of 5 m and the resolution is coarsening below that. The model requires boundary conditions at the atmosphere-ocean boundary (NCEP/NCAR re-analysis (Kalnay et al., 1996)) and at the open boundaries to the North Atlantic. Transport of freshwater and nutrient loads from land is considered. Details on the utilized boundary and forcing data are given by Daewel and Schrum (2013), who also gave a detailed description of analysis methods and validation datasets.

Ute Daewel 13.2.2019 11:06

Deleted: e

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Ute Daewel 7.3.2019 12:59

Deleted: 3

Unknown

Field Code Changed

Unknown

Field Code Changed

Ute Daewel 13.2.2019 13:48

Deleted: .

Unknown

Field Code Changed

Unknown

Field Code Changed

Ute Daewel 13.3.2019 15:33

Deleted: 20

Unknown

Field Code Changed

Unknown

Field Code Changed

2.3 Datasets and statistical methods for model analysis

As described in Daewel and Schrum (2013), we used observational data on surface (depth <10m) nutrients (nitrate and phosphate) in the North Sea, which are made available by ICES (International Council for the Exploration of the Seas, www.ices.dk), for nutrient validation. Observations and modelled surface nutrients were averaged over the upper 10 m of the water column and co-located in space and in time and corresponding statistics were calculated for the subareas specified in Figure 1. The seasonal cycle has not been removed prior to the analysis. The reason for doing so is the sparse data situation, which does not allow estimating a reliable seasonal cycle at each location. Additionally, the seasonal cycle changes from year to year. Thus, removing an average seasonal cycle from the data would add a bias to the data and hence increase the level of uncertainty. In the Baltic Sea, we used vertical nutrient profiles (NO₃, PO₄, O₂) at two distinct locations in the Baltic proper (BY5 (Lat/Lon: ~55.1°N/15.59°E), BY15 (Lat/Lon: ~57.2°N/20.03°E) see Figure 1) from the Baltic Sea monitoring network (see e.g. www.helcom.fi) that were continuously sampled since 1970. The data are available for download at www.ices.dk (accessed 05/2012). To account for inconsistencies in sampling frequencies, we co-located model data and observations prior to estimating average vertical profiles and standard deviations. For this purpose the model values were linearly interpolated onto a 1m vertical grid to allow best local comparison to the observations, while the observations were considered on the actual sampling depth. The statistical measures chosen for model analysis are the Pearson's correlation coefficient, the standard deviation and the root mean square deviation presented in a Taylor diagram (Taylor, 2001), and Empirical Orthogonal Functions (EOFs) as described in Storch and Zwiers (1999). The EOF analysis is a statistical method to understand major modes of variability in multidimensional data fields. A detailed description on how this method has been applied is given in (Daewel et al., 2015): "The annual values of the spatially explicit variable field form a NxM matrix χ (N: number of years; M: number of wet grid points). The empirical modes are given by the K eigenvectors of the covariance matrix with non-zero eigenvalues. Those modes are temporally constant and have the spatially variable pattern $p_k(m=1,...,M)$ where $k=1,...,K$. The time evolution $A_k(t=1,...,N)$ of each mode can then be obtained by projecting $p_k(m)$ onto the original data field χ such that $\chi(t, m) = \sum_{k=1}^K p_k(m) A_k(t)$. In the following we will refer to $A_k(t)$ as the principal components (PC) and to $p_k(m)$ as empirical orthogonal function (EOF). The percentage of the variance of the field χ explained by mode k is determined by the respective eigenvalues and is referred to as the global explained variance $\eta_g(k)$. Before using the method to analyse the spatiotemporal dynamics of the field, the data were demeaned (to account for the variability only) and normalized (to allow an analysis of the variability independent of its amplitude). The identified modes are not necessarily equally significant in all grid points of the data field. Thus, the local explained variance $\eta_{local,k}(m)$ could provide additional information about the regional relevance of an EOF mode and the corresponding PC in percent:

$$\eta_{local}^k(m) = \left[1 - \frac{\text{Var}(\chi(m,t) - p^k(m) A^k(t))}{\text{Var}(\chi(m,t))} \right] \cdot 100, \quad (12)$$

where $\text{Var}(X) = \sum_{t=1}^N (\bar{X} - X(t))^2$ denotes the variance of the field $X(t)$.

Unknown
Field Code Changed

Unknown
Field Code Changed

Ute Daewel 12.2.2019 13:37
Deleted: figure

Ute Daewel 13.2.2019 13:52
Deleted: of the

Ute Daewel 13.3.2019 15:37
Deleted: Fig.

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Information on the North Sea fish community are collected in the North Sea international bottom trawl survey (NS-IBTS) (ICES, 2012) and are freely available at the International council for the exploration of the sea (www.ices.dk). The NS-IBTS dataset contains spatially-resolved, species-specific information on fish length (for some target species also age) and catch per unit effort (CPUE: in numbers captured per hour). Given that our model estimated state variables on the base of carbon biomass, we converted fish length and abundance data to fish biomass (in grams captured per hour) based on published length-weight relationships (LWRs) for each species sampled in the NS-IBTS survey between 1980-1989 inclusive.

LWRs were derived from Coull et al. (1989), Froese et al. (2014) and FishBase (Froese and Pauly, 2000) for teleost species, McCully et al. (2012) and Templeman (1987) for rays and skates, Klaoudatos et al. (2013) for crabs, and Pierce et al. (1994) and Guerra and Rocha (1994) for squids. Total body weight (W) is a function of total length (L) following the relationship $W = aL^b$ where b is a parameter indicating isometric growth in body proportions if $b \sim 3$, and a is a parameter describing body shape and condition if $b \sim 3$.

Calculations were made at the species level, where species names were available, with the a and b parameter estimates taken from published sources in the following order:-

1. Studies specific to the North Sea (i.e. Coull et al., 1989; Froese and Pauly, 2000). If not available then:-
2. Studies specific to the British Isles (i.e. Coull et al., 1989; Froese and Pauly, 2000; McCully et al., 2012). If not available then:-
3. Studies specific to the North Atlantic (i.e. Coull et al., 1989; Froese and Pauly, 2000; McCully et al., 2012; Templeman, 1987). If not available then:-
4. Posterior mean estimates of a and b from a Bayesian hierarchical analysis for the species across all credible LWR studies, regardless of location (see Froese et al., 2014; Froese and Pauly, 2000).

For some species, e.g. herring, sprat, mackerel, Coull et al. (1989) provided mean parameter estimates for each month, or at least some months throughout the year, in addition to annual estimates. In these cases, we used annual estimates only. If information on genus only, family only or order only was available, the a and b parameter estimates for all species within that genus, family or order captured during the surveys were averaged to give a genus-, family- or order-specific value. If no information was available at any species level within a genus, family or order, then the geometric mean of that genus, family or order was computed from the data file accompanying Froese et al. (2014). The data were assorted onto the horizontal model grid based on their sampling location. For the data-model comparison we will only consider data from the first quarter of the year, since in the considered time period (1980-1989) this was the only systematically surveyed quarter in the survey.

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

2.4 Scenario Definition

Two sets of scenarios were performed to evaluate the simulated food-web response to specific changes in the food web parameterizations. We study the structuring effects of the new model closure term (higher trophic levels/fisheries) and the effects of changes in background zooplankton mortality. The first set of scenario experiments addresses the carbon loss through apex predators and fisheries by adding another source of mortality in addition to the natural fish mortality term.

Following the fisheries overview of the North Sea and Baltic Sea region as published by ICES (2018a, 2018b), biomass losses due to fisheries are in both regions in the range of 20%-50% of the total fish biomass per year. For convenience we therefore decided to receive the fisheries mortality rate by scaling the natural mortality rate ($=0.001\text{d}^{-1}=0.365\text{yr}^{-1}$) with 0.5, 1 & 2. As such, three different scenarios were calculated with an average (0.1825 yr^{-1}), high (0.365 yr^{-1}) and extreme (0.5475 yr^{-1}) loss rate.

Note that the latter two loss rates were chosen to provoke extreme responses in the fish biomass, and do not represent realistic catch rates in the areas.

The second set of scenario experiments was designed to understand the ecosystem response to changes in the zooplankton natural mortality. This term previously formed the sole closure term of the system and the rate has been reduced by 20% to account for the additional mortality induced to the system by MB and fish predation. Here, we chose four scenarios for the experiment. First we defined the control run, which considers the unchanged zooplankton mortality rate from ECOSMO II. Then, control -20%, is the reference setup for ECOSMO E2E. Additionally, we discussed control -40% and control +20% for comparison.

3 Results & Discussion

In the following, we discuss the basic characteristics of the model and assess its performance based on 10-year averages of the model variables. Specifically, we i) present and discuss the spatial dynamics of the newly introduced functional groups; ii) we discuss the seasonality of the ecosystem components and introduce the MB and fish diet composition emerging from the model; iii) we present the comparison of the simulated fish biomass distribution to observed data and repeat the nutrient validation analysis as previously presented for ECOSMO II in Daewel and Schrum (2013); and iv) we discuss the model sensitivity with respect to ecosystem model closure.

3.1 Description of modelled spatial pattern

The mean spatial pattern of calculated MB and fish vertically integrated biomass for the period (1980-1989) are presented in Figure 3. On average, estimated MB biomass (Figure 3a) in the North Sea is 1.98 gC m^{-2} for the considered time period. As we will see later, this value is highly sensitive to the parametrization of zooplankton mortality and fisheries effort (c.f. 3.4 and Figure 12). Heip et al. (1992) proposed an average of $7\text{g ash-free dry weight (AFDW) m}^{-2}$ based on a synoptic sampling of North Sea benthos in April-May 1986. This equates to approximately 3.5 gC m^{-2} when assuming a carbon fraction of ash-

Ute Daewel 7.3.2019 13:05
Deleted: (
Ute Daewel 7.3.2019 13:04
Deleted: ,
Ute Daewel 7.3.2019 13:06
Deleted: Three
Ute Daewel 4.3.2019 16:54
Formatted: English (UK)
Ute Daewel 4.3.2019 16:54
Formatted: English (UK)
Ute Daewel 4.3.2019 16:54
Formatted: English (UK)

Ute Daewel 4.3.2019 16:54
Formatted: English (UK)

Unknown
Field Code Changed
Ute Daewel 12.2.2019 13:37
Deleted: figure
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 4.3.2019 16:54
Formatted: English (UK)
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 4.3.2019 16:54
Formatted: English (UK)
Unknown
Field Code Changed
Ute Daewel 13.3.2019 15:38
Deleted: equals

free dry weight of 0.5 (Ingrid Krönke, Senckenberg am Meer, Wilhelmshaven, Germany, pers. communication), which is somewhat higher than our model estimates, but also includes benthic carnivore biomass (Greenstreet, 1997). Greenstreet (1997) estimated the biomass of the benthic filter feeder and deposit-feeder guild to amount to $\sim 3 \text{ gC m}^{-2}$ when the same carbon fraction of 0.5 is applied. Note that the comparison can only be an approximation due to the high variability in carbon content among species (e.g. Timmermann et al., 2012). Observational estimates of North Sea MB biomass indicate a decrease in biomass with increasing latitude according to Heip et al. (1992), and similar results were obtained from a subsequent sampling project in 2000 (Rees et al., 2007). Particularly high values of MB biomass were found in the shallow areas of the southern North Sea, including the coastal areas and Dogger Bank (Heip et al. 1992 their Figure 1), and in the river mouth areas along the English coast. This is in clear agreement with what has been estimated by our model.

In the Baltic Sea, the MB biomass was modelled to be on average 1.01 gC m^{-2} . This is in the range of what has been published by Timmermann et al. (2012) based on HELCOM data, who reported spatially-resolved values between $5\text{-}100 \text{ gWWt m}^{-2}$ (approx. $0.25 - 5 \text{ gC m}^{-2}$), as well as by Tomczak et al. (2012) who estimated macrobenthos biomass of about 30 t km^{-2} , which equals 1.5 gC m^{-2} using an Ecopath with Ecosim Baltic Proper food web model. The spatial distribution of MB modelled with our simplified model is consistent with the spatial distribution of major MB species in the Baltic Sea as presented by Gogina and Zettler (2010) based on species specific model estimates and observations. This applies specifically to the high abundances in the southern Baltic Sea, the near coastal areas and the Gulf of Riga.

Our model estimates highest MB biomass in both North Sea and Baltic Sea in shallower areas, especially near the coast and in bank regions, such as Dogger Bank, Fisher Banks and Oyster Ground, with a maximum of around 5 gCm^{-2} to be found in the southern North Sea. MB production in the model is constrained by the availability of oxygen; therefore, large areas of the central Baltic Sea are not inhabited by macrobenthos. In the North Sea, minimum MB biomass is estimated slightly offshore of the British coast and in the deeper parts of the Norwegian Trench region. Those minima in the North Sea were not caused by anoxic conditions, but by a lack of prey in the respective areas. The transition zone between North Sea and Baltic Sea, including the Skagerrak, Kattegat, Danish straits and the Fehmarn Belt, exhibits, in contrast, generally high values of MB biomass.

Simulated spatial variability in vertically integrated fish biomass shows a structured pattern both in the North Sea and the Baltic Sea. In the Baltic Sea, maxima of fish biomass are simulated in the coastal areas, the Gulf of Riga, in the southern Baltic Sea including Arkona Basin, Bornholm Basin and Bay of Gdansk and in the Åland Sea at the entrance to the Bothnian Sea. The deeper parts of the Eastern Gotland Basin, the Gulf of Finland and the Gulf of Bothnia in contrast feature very low fish biomass due to low prey biomass and oxygen depletion near the bottom. The modelled spatial distribution compares well with findings from the nutrient-to-fish model from Radtke et al. (2013). They integrated the model over a 4-year period 1980-1983. In their model approach fish follows specific rules for horizontal migration (food availability, spawning). However, their simulated spatial distribution of the combined biomass for the 3 different simulated fish species is very similar to our estimates. Differences between the simulated fish distributions occur specifically in time periods when predefined spawning areas determine the distribution. Other spatial differences were simulated for the Gulf of Finland where

Unknown
Field Code Changed
Unknown
Field Code Changed
Unknown
Field Code Changed
Unknown
Field Code Changed
Unknown
Field Code Changed
Ute Daewel 12.2.2019 13:35
Deleted: figure
Unknown
Field Code Changed
Unknown
Field Code Changed
Unknown
Field Code Changed

Unknown
Field Code Changed
Ute Daewel 13.3.2019 15:39
Deleted:
Ute Daewel 13.3.2019 15:39
Deleted: s

the model by Radtke et al. (2013) estimated relatively high fish biomasses in contrast to our model, and around Gotland where our model produces fish biomass maxima potentially fostered by the additional availability of macrobenthos as prey, which remains unconsidered by Radtke et al. (2013). [Interestingly, the distribution of fish biomass maxima, estimated by our model, resembles the pattern described as cod nursery areas in the Baltic Sea](#) by Bagge et al. (1994).

5 The structure of modelled North Sea fish biomass is very distinct with maxima in frontal areas such as the tidal mixing front in the southern North Sea and around Dogger Bank and the frontal zone off the German, Danish and British coast, and in the Norwegian Trench. Maxima are also modelled in the Fisher Banks and Oyster Ground and Fladen Ground regions. Minima, in contrast, were estimated in the deeper parts of the western North Sea off the British coast, the German Bight and in the English Channel. They partly resemble the minima estimated for MB, which indicates that fish biomass minima are caused

10 by food shortage in areas with low MB biomass. Especially off the British Coast, studies from Callaway et al. (2002) and Jennings et al. (1999) indicate high fishing effort, indicating that the model likely underestimates the fish biomass in that region. Potential reasons are the model underestimating zooplankton production in that area (Daewel et al., 2015), and the missing impact from the open boundary, where neither zooplankton nor fish were prescribed to enter the model domain.

15 When integrating fish biomass over the North Sea and Baltic Sea regions, it amounts to ca. 0.462 million tonsC and 0.312 million tonsC respectively. Assuming the carbon content of fish being 45% (Huang et al., 2012; Sterner and George, 2000) and the AFDW (ash free dry weight) to wet weight fish ration ranging from 0.1 to 0.2, this corresponds to a simulated total fish biomass in the range of 5.13-10.27 million tons for the North Sea and 3.47-6.93 million tons for the Baltic Sea. Since the AFDW to wet weight ratio is highly variable, even within species, depending on e.g. temperature, season and diet of the fish (Elliott and Hemingway, 2002), it is difficult to determine an exact value for the simulated entire fish assemblage biomass.

20 The modelled estimates of fish biomass are well within the range of what has been estimated for total fish biomass based on observations for the Baltic Sea (Thurow, 1997). Using yield data and age composition data in catches, Thurow (1997) estimated total fish biomass in the Baltic Sea for the time period 1900-1985. His results indicate relatively low fish biomass (<2 million tons) for the first half of the century, but a drastic increase thereafter. For the time period considered here (i.e. 1980-1989) he proposed the fish biomass to be around 7 million tons. [Following ICES \(2018b, 2018a\), fisheries during the 1980's were in the range of 0.7-1 million tons in the Baltic Sea and 2-3 million tons in North Sea. Despite that the model underestimates fish production due to the no horizontal migration assumption and no fish migration over the lateral boundaries, the model's estimates of fish biomass in the North Sea would support the fisheries landing during that time period.](#)

25 Estimates for North Sea total biomass for the time 1983-1985 based on the ICES International young fish survey (IYFS) and the English ground fish survey (EGFS) were published by Sparholt (1990). For the first quarter Sparholt (1990) estimated an average fish biomass of about 8.6 million tons, while for the third quarter the average biomass was estimated to be 13.1 million tons. The discrepancy between first and third quarter was explained by the migration of the western stock of Atlantic mackerel (*Scorpaenopsis scorpaenoides*) and horse mackerel (*Trachurus trachurus*) into the North Sea, which is not considered by our model. Further our results find agreement with output from an Ecopath with Ecosim food-web model of the North Sea as proposed by Mackinson and Daskalov (2007), which estimated that the North Sea total fish biomass was ~11 million tons in

30

Unknown
Field Code Changed

Ute Daewel 13.3.2019 15:40
Deleted: Interestingly, does the distribution of fish biomass maxima, estimated by our model, resemble the pattern that are described as cod nursery areas in the Baltic Sea

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Ute Daewel 7.3.2019 13:18

Deleted: (

Ute Daewel 7.3.2019 13:18

Deleted: ,

Unknown
Field Code Changed

Unknown
Field Code Changed

the year 1991. Indeed, our modelled estimate is well within the range of previously published observations and model results. We suggest that any discrepancies are most likely due to our model neglecting fish migration at large scales. When discussing the spatial variation of estimated fish biomass we need to consider that the model is constrained by the assumption that fish do not move horizontally. Thus, we estimate the production potential for fish in each horizontal grid cell rather than the [actual](#) fish biomass at a given time and location.

3.2 Seasonal dynamics of ecosystem components and diet composition

Values for both MB and fish biomass vary in the course of the year ([Figure 4](#)). While the modelled seasonal amplitude for fish biomass is relatively small (North Sea: $94.7 \text{ mgC m}^{-3} \pm 11\%$; Baltic Sea: $84.8 \text{ mgC m}^{-3} \pm 9.7\%$ of the mean biomass) when compared to the average values, MB seasonality is substantial (North Sea: $4.4 \text{ gC m}^{-3} \pm 222\%$; Baltic Sea: $1.86 \text{ gC m}^{-3} \pm 183\%$ of the mean biomass). Minimum MB and fish biomass is estimated for winter and early spring and the seasonal maximum is modelled for late summer and autumn. The MB maximum lags behind the zooplankton maximum by about 3 months. In contrast to zooplankton, the MB minimum does not [reach](#) values close to zero but the model simulates a significant standing stock for MB also during winter.

In [Figure 4](#) the seasonal cycles for the phytoplankton and zooplankton estimates of the ECOSMO E2E run is presented together with those of the ECOSMO II simulation (Daewel and Schrum, 2013). The seasonal cycles for both phytoplankton and zooplankton are clearly impacted by the consideration of MB and fish. Although the general phytoplankton biomass seasonality and the phenology remain relatively unchanged, the magnitude of the seasonal maximum, especially the diatom bloom, is significantly increased in spring and early summer in both regions NS and BS when the MB/fish groups are included. The consideration of seasonally-variable MB and fish predation on zooplankton imposes a different seasonality on zooplankton mortality compared to the constant mortality rate used in ECOSMO II (Daewel & Schrum, 2013) and hence impacts zooplankton phenology. The reduced zooplankton biomass in the beginning of the season due to MB and fish predation ([Figure 5](#)) consequently leads to a reduction in phytoplankton mortality and to an increase in phytoplankton biomass (top-down process). Additionally, MB competes with zooplankton for resources and thereby changes zooplankton seasonality, especially in autumn in the North Sea when MB biomass is highest and it preys dominantly on dead organic material and phytoplankton ([Figure 5](#)).

An overview on the seasonal feeding dynamics can be obtained by identifying the monthly prey composition for MB ([Figure 5a](#)) and fish ([Figure 5b](#)) in the North Sea and Baltic Sea. For MB, the major food source throughout the year is organic sediments followed by dead organic material, while the percentage of the latter is considerably higher in the North Sea than in the Baltic Sea, presumably due to the fact that a higher percentage of detritus is re-suspended in the [tidally influenced](#), highly turbulent areas of the North Sea. Zooplankton and [phytoplankton are](#) included in the diet when available in spring and summer. The fish prey composition ([Figure 5b](#)) is very similar in both sub-areas; with MB dominating the diet in the autumn and winter months and omnivorous (large) zooplankton dominating in summer. Detritus contributes a significant food source

Ute Daewel 13.2.2019 13:53

Deleted: concrete

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 6.3.2019 10:34

Deleted: .

Ute Daewel 6.3.2019 10:35

Deleted:

Ute Daewel 6.3.2019 10:35

Deleted:)

Ute Daewel 6.3.2019 10:41

Deleted: fall to

Ute Daewel 12.2.2019 13:35

Deleted: figure

Unknown

Field Code Changed

Ute Daewel 13.3.2019 15:41

Deleted: a

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 6.2.2019 17:10

Deleted: P

Ute Daewel 6.2.2019 17:10

Deleted: is

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 15:42

Deleted: es

in March and April, while small zooplankton appears in autumn only in very low amounts. Greenstreet et al., (1997) reviewed food web studies in the North Sea and analyzed the food consumption of fish by guild. When adding up the average MB and zooplankton in the diet of the four in the study [fish guilds considered](#) (demersal piscivores, demersal benthivores, pelagic piscivores, pelagic planktivores), the ratio zooplankton/MB in the diet lies at around 6/4 in summer, which is comparable to our estimated food composition in summer. In contrast to our model results, the estimates from Greenstreet et al. (1997) show no significant seasonal variations in diet composition. Explanations for this disagreement might be found in the model performance of e.g. the zooplankton standing stock. The latter has been estimated to be very low in winter, and hence lead to an intensification of the modelled zooplankton seasonal cycle and to too little zooplankton in the fish diet in winter. Another possible reason for the mismatch between the model and the estimates from Greenstreet et al. (1997) might be related to spatiotemporal differences in the fish biomass and diet.

An EOF analysis on the monthly mean fields for MB and fish biomass reveals the spatial-seasonal pattern. In [Figure 6](#) (MB) and 7 (fish) the first two EOF patterns are shown for MB and fish biomass respectively. Additionally the local explained variance and the related temporal pattern (PC) is given. For MB, the seasonal signal is very homogeneous across the whole area ([Figure 6](#)). With 77%, the first mode explains a significantly large part of the overall variability and the temporal signal resembles the average variability shown in [Figure 4](#). This highlights that the MB seasonality is mainly induced by the seasonal pattern of the system productivity with increased production of fresh organic material in summer and less food availability in winter. [This is in line with observations on seasonality of benthic infauna at three different locations in the North Sea published by Reiss and Krönke \(2005\), who found maximum biomass in late summer. Although the observed seasonality showed the highest magnitude in the German Bight the seasonality was clear at all three locations. The authors concluded that of the potential relevant factors \(food availability/quality, water temperature, predation, hydrodynamic stress\) food quality plays the major role for infauna seasonality, thus is strongly related to primary production. They also suggest food limitation and predation pressure to be the main processes for decrease in abundance during winter. The same authors also looked at seasonality in the epibenthic community \(Reiss and Krönke, 2004\) showing that the epifaunal biomass varies less seasonally, especially in the off shore region, and that the main processes causing seasonal variations are related to migratory behavior, which is not covered by our model. For the Baltic Sea only very local studies in seasonality of MB are available of which some indicate locally strong seasonality \(Anders and Möller, 1983\), while in other regions no seasonal changes in biomass were observed due to the dominance of long-lived species \(Persson, 1983\). In general the comparison to observations indicates that on the one hand the model is able to represent the main seasonality in MB even though epi- and infauna are not separated. On the other hand, in future studies the consideration of an additional functional group encompassing longer-lived species will be required for addressing MB seasonality more correctly.](#)

The second EOF explains about 16% of the overall variability and is especially important in the Gulf of Finland and the Bothnian Bay where it explains up to 80% of the total variability. PC₂ differs from PC₁ by showing a maximum in MB in late autumn and winter with a time lag of about two months compared to PC₁, while the minimum is modelled for July and August. The ecosystem seasonality in the Bothnian Bay and the Gulf of Finland is highly impacted by a relatively long

Unknown
Field Code Changed
Ute Daewel 13.3.2019 15:42
Deleted: s
Ute Daewel 13.3.2019 15:42
Deleted: considered fish guilds

Unknown
Field Code Changed
Ute Daewel 13.3.2019 15:34
Deleted: Figure

Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 12.2.2019 13:36
Deleted: figure

period of winter sea ice cover and the onset of the spring bloom is therefore delayed (see e.g. Andersson et al., 1996; Daewel and Schrum, 2013). This would consequently affect the phenology of MB and fish in that area and explains the difference in the seasonal cycle.

In contrast to the MB pattern, the EOF of the fish biomass seasonality (Figure 7) reveals a clear distinction between seasonal signals in different regions. Together, the first two EOFs explain over 70% of the overall variability whereas the first EOF comprises 44%. This first pattern describes a seasonal cycle with a minimum in March/April and a plateau in maximum fish production between August and December, which dominates the average seasonal cycle shown in Figure 4. It particularly explains the seasonality in the deeper central North Sea, the Norwegian Trench, the Skagerrak and the northern Kattegat region as well as the coastal areas of the central Baltic basins. In all of these regions this mode explains up to 80 % of the variability. The second pattern describes the seasonal variability in the shallow areas of the southern North Sea, including Dogger Bank, the North-Western North Sea and the Belts at the entrance to the Baltic Sea, with a maximum in fish biomass in late spring and summer. The dynamics in these areas are determined by the zooplankton seasonality featuring, unlike the central North Sea, a maximum in summer. The two modes of the estimated fish seasonal cycles clearly indicate two different fish habitats, structured by food availability and temperature. In Figure 8 fish production, partitioned into diet components, in the shallow (Figure 8a) and deeper (Figure 8b) North Sea is illustrated together with the seasonal temperature cycle. The main differences contributing to differences in fish biomass seasonality are the timing of the food resources and the difference in temperature. In the shallow areas of the North Sea, zooplankton forms the major food source for fish in early spring and summer, reaching a maximum in May and June. After that, the MB contribution increases and resumes the role as the major food source in August. In the deeper parts of the NS, the dynamics in fish diet composition are shifted by 1-2 months, and, in contrast to the shallower NS, dead organic material plays an important role throughout the year. Since the seasonality in total fish production is very similar in both regions, this difference in diet composition would per se not lead to a difference in fish biomass seasonality as seen from the EOF analysis (Figure 7). However, in addition to the difference in food resources the two habitats feature very different seasonal temperature cycles. The most likely explanation for the stronger decrease in fish biomass in shallow NS in August and September (Figure 7 PC₂) is the temperature driven higher loss rate (Eq. 11).

Here, we can identify the distinction of NS fish communities approximately at the 50m depth line, which is comparable to the separation line reported for North Sea fish communities in earlier published observational studies (Callaway et al., 2002; Rees et al., 1999). Using data from 270 stations distributed over the whole North Sea, Callaway et al. (2002) separated the NS fish community into several clusters (3 or 5 in dependence of the trawling method) and two main groups. The most conspicuous boundary was defined approximately at the 50m depth contour separating the community in the shallow southern North Sea, which mainly consists of small non-commercial species, from the community in the central North Sea, which was dominated by haddock (*Melanogrammus aeglefinus*), *M. merlangus*, herring (*Clupea harengus*), and plaice (*Pleuronectes platessa*). The authors also suggested the environmental conditions in the region to play a major role in structuring the community. Although our model cannot distinguish between different species and actual communities, the

Unknown
Field Code Changed

Ute Daewel 13.3.2019 15:37
Deleted: Fig.

Ute Daewel 13.3.2019 15:34
Deleted: Figure

Ute Daewel 12.2.2019 13:36
Deleted: figure

Ute Daewel 13.3.2019 15:37
Deleted: Fig.

Ute Daewel 13.3.2019 15:37
Deleted: Fig.

Ute Daewel 13.3.2019 15:44
Deleted: ,

Ute Daewel 13.3.2019 15:44
Deleted: a

Ute Daewel 13.3.2019 15:37
Deleted: Fig.

Ute Daewel 13.3.2019 15:37
Deleted: Fig.

Unknown
Field Code Changed

Unknown
Field Code Changed

Ute Daewel 13.3.2019 15:45
Deleted: for

results indicate a clear distinction between the seasonality and the driving environmental conditions of fish production potential in the shallow southern North Sea and in the central North Sea.

Spatial variations of biomass specific mortality related to zooplankton consumption by fish and MB are given in Figure 9 as an average from 1980-1989. The results were additionally separated into 1st and 2nd quarter of the year to identify potential intra-annual variations as suggested by Maar et al. (2014). The results show a very distinct spatial pattern for fish induced zooplankton mortality (Figure 9a) with increased values in some specific regions in the central North Sea, especially in the vicinity of Dogger Bank, close to the English coast, Oyster ground and in the Fisher Bank (Little and Great Fisher Bank) area. Our model shows furthermore considerable consumption in the Norwegian Trench, along the coast of the southern and central Baltic Sea including the Kattegat/Skagerrak region, in the central basins of the southern Baltic Proper, the Gulf of Gdansk and in the Gulf of Riga. The biomass specific mortality related to MB consumption (Figure 9b), in contrast, is confined to shallow areas with a relatively strong coupling between the benthic and pelagic system. This includes the shallower areas of the southern and central North Sea and the near coastal areas in the Baltic Sea. The difference between the 1st and 2nd half-year is relatively small for the fish induced mortality. However, there is a clear but small increase in the central North Sea and a much stronger change for the Baltic Sea pattern with a higher impact in the second half of the year. Also for MB the impact is substantially stronger for the 2nd half-year, clearly related to the strong seasonal signal in MB biomass.

While we estimate zooplankton predation losses within the model, an earlier study by Maar et al. (2014) proposed spatial-temporal variations in biomass specific mortality of zooplankton based on data of the major zooplanktivorous fish species and on larval distribution in the North Sea (their Figure 10C,D). Our results show clear similarities in magnitude and spatial structure when compared to the results from Maar et al. (2014). The results of Maar et al. (2014) however showed a clear difference between 1st half-year and 2nd half-year with decreased biomass specific mortality in the 2nd half-year in the central North Sea. This intra-annual variation is not evident in our model results. However we found a clear difference in magnitude when comparing winter and summer season (not shown). The reasons for the discrepancies between our model results and Maar et al (2014) are presumably related to interannual variations in fish consumption, which are not considered in a 10-year average, the fact that migration is not considered in the model and thus restrict the spatial variation, and that our functional group cannot resolve species- and stage-specific spatial and temporal variations like e.g. the increase in larval biomass in spring and changes in species composition. On the other hand, the approach from Maar et al. (2014) reveals uncertainties due to the fact that only parts of the North Sea fish assemblage is considered and that the fish biomass is prescribed and not dynamically coupled to zooplankton biomass.

The approach from Maar et al. (2014) provides the possibility to replace the spatially- and temporally-invariant closure term usually used in NPZD type models (Daewel et al., 2014) by a data driven, detailed formulation, and allows for consideration of the predation effects of different fish species and larvae on the zooplankton dynamics. However, the main disadvantage is that, as the authors already pointed out, a huge amount of detailed species specific and potentially undersampled data is required and that the estimated mortality index relies on a number of assumptions concerning e.g. the relevance of the

Ute Daewel 5.3.2019 10:33
Deleted: -
Ute Daewel 5.3.2019 10:33
Deleted: -
Ute Daewel 5.3.2019 11:33
Deleted: The spatial variation of the estimat... [8]
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 5.3.2019 11:35
Deleted: at
Ute Daewel 5.3.2019 16:14
Deleted: and at
Ute Daewel 5.3.2019 16:15
Deleted: zooplankton consumption by
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 5.3.2019 16:15
Deleted: has strong impacts in the
Ute Daewel 5.3.2019 16:16
Deleted: , namely the coastal Region and I... [9]
Ute Daewel 5.3.2019 16:19
Formatted ... [10]
Ute Daewel 5.3.2019 16:19
Formatted ... [11]
Ute Daewel 5.3.2019 16:23
Formatted ... [12]
Unknown
Field Code Changed ... [13]
Ute Daewel 5.3.2019 16:25
Deleted: variable predation rates
Ute Daewel 5.3.2019 16:26
Deleted: n
Ute Daewel 5.3.2019 16:32
Formatted ... [14]
Ute Daewel 5.3.2019 16:32
Formatted ... [15]
Ute Daewel 13.3.2019 15:50
Formatted ... [16]
Ute Daewel 5.3.2019 16:35
Deleted: Such an approach
Ute Daewel 13.3.2019 15:51
Deleted:
Unknown
Field Code Changed ... [17]
Ute Daewel 13.3.2019 15:52
Deleted: ing

individual fish species and spawning time and distribution. Moreover, such a data driven approach has a limited potential for future projections and sensitivity studies on various effects on the ecosystem. However, following a very different, less detailed approach, the spatial variability of our estimates of zooplankton consumption (Figure 9a) by fish compares surprisingly well with the spatial variability of the fish consumption index provided by Maar et al. (2014; their Figure 4), which we consider an implicit validation of our model approach. The consideration of MB-related zooplankton predation mortality is an additional advantage that arises from our modelling approach.

3.3 Model performance and nutrient dynamics

To get a more direct measure for the validity of the modelled fish functional group we used data from the NS-IBTS. In Figure 10, we compared the mean fish functional group biomass distribution to the fish biomass from the NS-IBTS as calculated following the method described in section 2.3. We classified species within the NS fish community into ‘demersal’ (Figure 10a) and ‘pelagic’ groups (Figure 10b) based on life-history characteristics, then summed the biomass of each group to form a ‘combined’ (Figure 10c) group. In contrast to the species-specific differentiation into groups used for the observations, the model results do not provide this level of detail. Here, the differentiation has been performed based on the vertical distribution of the fish biomass. Thus we assigned all biomass in the bottom layer to the ‘demersal’ groups and biomass in the remaining water column to the ‘pelagic’ group. Therefore and because the units in the NS-IBTS data and in the model data differ, the Figures are not quantitatively comparable. When we compare the data for the different fish groups we find the ‘demersal’ fish more strongly increasing with latitude, and, following the North Sea bathymetry, with depth. The ‘pelagic’ fish group biomass, in contrast, shows, in addition to the increase with latitude, a maximum in the south and in the north of the North Sea. Altogether, we find a clear increase in fish biomass with latitude with a maximum at the entrance to the North Sea, but also higher values in the central North Sea and around Dogger Bank. From the data, we can also conclude, that the contribution of pelagic fish biomass to the overall biomass is higher in the south than further north. From a pure qualitative comparison, we find the modelled fish group biomass roughly resembling the observed pattern, with increasing biomass from south to north. The model also represents the maximum in the Kattegat and at the northern shelf edge. However, the model estimates a pronounced minimum off the British coast, which is not evident from the observations and likely stems from the zero boundary condition, meaning that no fish enters or leaves the model area over the lateral boundaries, and missing migration parameterization in the model. In summary, we found that the model is able to represent the spatial fish distribution in the North Sea. However, the differences in fish biomass off the British and partly at the European continental coast and the discrepancy between the observed and modelled ‘pelagic’ group indicate a potential underrepresentation of the pelagic fish stock by the simulated fish functional group.

To understand the effect of changes in the NPZD model closure on model performance with respect to nutrient dynamics, we repeated the nutrient validation for surface nutrients in the North Sea (Figure 11) and for nutrient profiles in the Baltic Sea (Figure 12) as described by Daewel and Schrum (2013). For both surface nitrate (Figure 11a) and phosphate (Figure 11b) the statistics, presented here in a Taylor diagram, indicate an improvement for some regions when MB and fish were considered.

Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Unknown
Field Code Changed
Ute Daewel 13.3.2019 15:52
Deleted:
Ute Daewel 13.3.2019 15:52
Deleted: the here presented
Ute Daewel 12.2.2019 13:36
Deleted: figure
Ute Daewel 6.3.2019 13:30
Deleted: (Fig. 10d)
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 6.3.2019 14:22
Deleted: Note,
Ute Daewel 6.3.2019 14:22
Deleted: that
Ute Daewel 6.3.2019 14:23
Deleted: , and that
Ute Daewel 13.3.2019 15:34
Deleted: figure
Ute Daewel 6.3.2019 14:23
Deleted: therefore
Ute Daewel 6.3.2019 14:24
Deleted: both
Ute Daewel 13.3.2019 15:54
Deleted: ,
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Unknown
Field Code Changed
Ute Daewel 13.3.2019 15:37
Deleted: Fig.
Ute Daewel 13.3.2019 15:37
Deleted: Fig.

Larger improvements occur in regions with relatively high estimated biomass for MB and fish, such as in region E of the English coast, where the correlation coefficient for nitrate improved from under 0.4 to 0.5, and for phosphate from under 0.6 to above 0.7, yet with a stronger bias for both nutrients. Better results were also accomplished in the central North Sea (region K), where the standard deviation moved significantly closer to that of the observations. Small improvements are also shown for regions F and L.

The MB and fish group potentially alters the nutrient dynamics in the Baltic Sea (Figure 12). Although we found only relatively small changes for phosphate relative to the ECOSMO II simulation in both considered locations, clear differences for the nitrogen and oxygen profiles are apparent, especially in the intermediate depth levels between 50 m and 150 m and at the surface. The model indicates a slight upward shift of the oxycline when fish and MB are resolved, which also affects nitrate by relocating the nitrate maximum. This results in decreased model performance with respect to nitrate in the intermediate layer, but improves the performance at the surface by increasing the initially too low modelled surface nutrient concentrations. Ammonium is significantly improved in lower layers at BY15 station for the ECOSMO-E2E model. Several processes interact to determine the changes in vertical nutrient profiles. Possible candidates are, e.g., changes in the oxygen dynamics by including oxygen dependent fish and MB, changes in sediment dynamics, since organic sediments are ingested by MB and subsequently nutrients are released on different time and spatial scales. Additionally, we found that the nutrient dynamics are also sensitive to the parameter choice of zooplankton mortality and the loss rate through fisheries and apex predation (cf. 3.4).

3.4 Ecosystem response to structuring drivers

With the two sets of scenarios we try to evaluate the impact of changes in model closure (fisheries mortality) and zooplankton mortality on ecosystem structure.

In the fisheries-scenarios we would expect a top-down response of the ecosystem dynamics to changes in fisheries, such that reduced fish biomass would relax the predation on the secondary producers (zooplankton and MB), which consequently would increase in biomass and reduce phytoplankton (see e.g. Cury et al. (2003)). Our model results indicate this type of trophic response for the Baltic Sea ecosystem (Figure 13b), but with very little efficiency for the lowest trophic level. The reduction of fish biomass for the highest catch rate scenario is about 98% compared to the control run, and for zooplankton and MB the increase is 13% and 62% respectively. The response of phytoplankton biomass, in contrast, is only in the order of 4% and hence small compared to the interannual variability of phytoplankton biomass, which is in the order of 10%.

The North Sea ecosystem responds less predictably than the Baltic Sea to simulated changes in the fish model closure term. Although the reduction of fish biomass in the North Sea results in an increase in MB, zooplankton biomass does not respond correspondingly (Figure 13a). The introduction of a moderate loss term leads to a comparably strong reduction of zooplankton biomass, while, with further increase in the loss rate zooplankton biomass increases again. As in the Baltic Sea, phytoplankton biomass is reduced with increasing fishing effort but the response is even smaller (~3%). The most likely reason for the more complex response of the North Sea ecosystem is the tighter coupling between MB and zooplankton and

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 6.2.2019 17:30

Deleted: is

Ute Daewel 6.2.2019 17:30

Deleted: for

Ute Daewel 13.3.2019 16:01

Deleted: F

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Unknown

Field Code Changed

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

phytoplankton (see [Figure 7a](#)). Since zooplankton forms a prey group for MB in the North Sea, a major change in MB and fish biomass affects the relevance of the two zooplankton predator groups and the increased predation pressure by MB will counteract (and potentially overrule) the relaxed predation by fish.

The second set of scenario experiments was designed to understand the ecosystem response to changes in the zooplankton natural mortality. In the new E2E model configuration a change of this term cascades up and down the trophic food chain (Fig 13c&d). In both systems, a reduction in zooplankton natural mortality leads consequently to an increase in zooplankton biomass and to a decrease in phytoplankton and an associated decrease in MB. The difference between the systems becomes manifest in the response of the fish group, which is positive in the Baltic Sea ([Figure 13d](#)), but reverses in the North Sea ([Figure 13c](#)) with higher fish biomass in a low zooplankton environment. This response highlights again the major role of the MB in the North Sea ecosystem, which partly competes with zooplankton and forms a major prey item for fish.

Despite the strong changes in the magnitude of phytoplankton and zooplankton biomass, the phenology of the seasonal cycles was almost not impacted by the sensitivity changes (not shown). The only distinct change is a decrease in phytoplankton spring biomass in the Baltic Sea when the model closure term is increased. Almost none other of the phenological changes described in section 3.2 was affected when fish biomass was decreased in the first set of sensitivity experiments, highlighting the dominant role of MB for these changes.

4 Conclusion

We presented here a [3d](#) resolved food web model that is based on a functional group approach ranging from nutrients to fish. Differently to the study by Fennel (2010), we did not distinguish between different fish species to avoid uncertainties associated with the choice of fish species and their contribution, compared to the unconsidered remaining biomass. Our approach integrates the full production potential for fish into one single functional group by defining the feeding pathways via primary and secondary production, including zooplankton and MB. This has certain advantages. For example, we avoid the parameterization of a detailed species dependent food web, and moreover, the adaptability of the model to other ecosystems is independent of the local fish assemblage. [The advantage of the generic functional group approach used in the model for all trophic levels is that we can simplify a complex community structure and reduce the information to the basic common features, thus avoiding a huge parameter set and data requirements. Still the model is able to simulate relevant ecosystem dynamics at high spatial and temporal resolutions with relatively low computational requirements.](#)

Despite the simplicity of the approach, we found the model able to reproduce the observed spatial pattern and magnitude of both macrobenthos and fish biomass in the North Sea and Baltic Sea as described in the literature (see [Figure 3](#) and section 3.1). This highlights the advantage of our approach, and adds weight to the assumption that fish biomass distribution consequently emerges from prey availability and environmental conditions. Furthermore, the model was able to distinguish between the two different fish production areas, separated around the 50-m depth line (compare [Callaway et al., 2002](#)), with

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 7.3.2019 11:33

Formatted: No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Ute Daewel 4.3.2019 16:57

Deleted: -

Ute Daewel 7.3.2019 11:32

Formatted: Font:10 pt, English (UK)

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 13.3.2019 16:03

Deleted:

Unknown

Field Code Changed

differences in the seasonal cycle and the diet composition. Although this differentiation is not based on species composition as in Callaway et al. (2002), it shows that the basic concept that biotic and abiotic conditions determine the composition of the local fish community, as realized in the model, and allows conclusions to be formed about the local fish community, even when it is not explicitly prescribed in the model. This opens up possibilities for additional investigations on, e.g., the inter-annual variability of fish production and biomass through general fish diet composition, and how this compares to observed long-term fish stock variations. Future model developments and applications should particularly address the composition of local fish communities, by classifying fish in two or more functional groups such as planktivores and piscivores, or into pelagic and benthic feeding guilds, to allow for a clearer representation of the food web structure.

However, the simplicity of the model and the related assumptions confines the model interpretation, and some of the simplifications require a revision in future model applications. Besides the redistribution of the MB and fish into several food-web specific functional groups, neglecting fish movements in the model approach is a clear limitation, since we know that fish are mobile and would migrate in response to e.g. food shortage, spawning behaviour or predators. In contrast to the Norwegian Sea, where distinct feeding migrations are observed for the pelagic fish component (Nøttestad et al., 2011) following the northward progressing zooplankton blooms and light conditions, the North Sea and Baltic Sea exhibit a relatively constant spatial pattern of system productivity, with highly productive areas along the coast and less productivity in the central seasonally-stratified regions. Hence, the migratory movements of North Sea and Baltic Sea fish stocks might not be based solely on large feeding migrations, but may also related to temperature and salinity changes and spawning behaviour (Hinrichsen et al., 2016; Hunter et al., 2003; Pinto et al., 2018; Radtke et al., 2013). Additionally, fish migrate into the area from the North Atlantic (e.g. Sparholt, 1990). Two questions arise specifically related to this topic: i) Is including migration strategies on a functional group level effective and reasonable in the North Sea and Baltic Sea environment, considering the variability among species? (ii) Would migration behaviour effectively impact the productivity of the system at the higher trophic level? We would therefore like to highlight the necessity to investigate the impact of specific migration strategies in continuative studies.

One major aim of the model development was to solve the closure term problem that arises with NPZD type models when choosing a fixed zooplankton mortality term (for review see Daewel et al. (2014)). The model results show that the inclusion of a higher trophic functional group can provide a more consistent and dynamic closure term, which produces a realistic but variable mortality field independent of in situ observations, in contrast to observational-based zooplankton predation as used e.g. by Maar et al. (2014). In return, the closure term problem is transferred to the new “end” of the food web, namely the fish group mortality. In future studies, this should be addressed by including dynamic formulations for apex predators and fisheries. This might be accomplished by introducing simple fisheries catch rates as explored in the scenario runs in this study (section 3.4), or by coupling the model to socio-economic models, which allows inclusion of social interests and management decisions in the modelling approach (e.g. Charles, 1989; Schlüter et al., 2014). The latter approach could allow the model to be applied in a fisheries management context – i.e. if the model’s ability to capture local fish community structure with respect to potential production and species composition is further developed.

Unknown
Field Code Changed

Ute Daewel 6.2.2019 17:30
Deleted: for

Ute Daewel 13.3.2019 16:04
Deleted:

Ute Daewel 13.3.2019 16:10
Deleted: food

Unknown
Field Code Changed

Ute Daewel 13.3.2019 16:11
Deleted: (Hinrichsen et al., 2016; Hunter et al., 2003; Radtke et al., 2013).

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Unknown
Field Code Changed

Code and data availability. Model code access and data can be obtained upon request. The code is available in the Helmholtz-Centre-Geesthacht git repository <https://coastgit.hzg.de/udaewel/hamsom-ecosmoe2e/>, and licensed under apache license version 2.

5 Acknowledgements

This work is a contribution to the FP7-SeasERA SEAMAN Collaborative Project financed by the Norwegian Research Council (NRC-227779/E40).

References

- Anders, K. and Möller, H.: Seasonal fluctuations in macrobenthic fauna of the Fucus belt in Kiel Fjord (western Baltic Sea), *Helgoländer Meeresuntersuchungen*, 36, 277–283, 1983.
- Andersson, A., Hajdu, S., Haecky, P., Kuparinen, J. and Wikner, J.: Succession and growth limitation of phytoplankton in the Gulf of Bothnia (Baltic Sea), *Mar. Biol.*, 126(4), 791–801, doi:10.1007/BF00351346, 1996.
- Bagge, O., Thurow, F., Steffensen, E. and Bay, J.: The Baltic cod, *Dana*, 10, 1–28 [online] Available from: <http://agris.fao.org/agris-search/search/display.do?f=1995/DK/DK95002.xml;DK9521335>, 1994.
- Barthel, K., Daewel, U., Pushpadas, D., Schrum, C., Årthun, M. and Wehde, H.: Resolving frontal structures: On the computational costs and pay-off using a less diffusive but computational more expensive advection scheme., *Ocean Dyn.*, doi:10.1007/s10236-012-0578-9, 2012.
- Blackford, J. C., Allen, J. I. and Gilbert, F. J.: Ecosystem dynamics at six contrasting sites : a generic modelling study, *J. Mar. Syst.*, 52, 191–215, doi:10.1016/j.jmarsys.2004.02.004, 2004.
- Butenschön, M., Clark, J., Aldridge, J. N., Icarus Allen, J., Artioli, Y., Blackford, J., Bruggeman, J., Cazenave, P., Ciavatta, S., Kay, S., Lessin, G., Van Leeuwen, S., Van Der Molen, J., De Mora, L., Polimene, L., Sailley, S., Stephens, N. and Torres, R.: ERSEM 15.06: A generic model for marine biogeochemistry and the ecosystem dynamics of the lower trophic levels, *Geosci. Model Dev.*, 9(4), doi:10.5194/gmd-9-1293-2016, 2016.
- Callaway, R., Alsvag, J., de Boois, I., Cotter, J., Ford, a., Hinz, H., Jennings, S., Kröncke, I., Lancaster, J., Piet, G., Prince, P. and Ehrich, S.: Diversity and community structure of epibenthic invertebrates and fish in the North Sea, *ICES J. Mar. Sci.*, 59(6), 1199–1214, doi:10.1006/jmsc.2002.1288, 2002.
- Casini, M., Hjelm, J., Molinero, J.-M., Lövgren, J., Cardinale, M., Bartolino, V., Belgrano, A. and Kornilovs, G.: Trophic cascades promote threshold-like shifts, *Proc. Natl. Acad. Sci.*, 106(1), 197–202, 2009.
- Charles, A. T.: Bio-Socio-Economic Fishery Models: Labour Dyanmics and Multi-Objective Management, *Can. J. Fish. Aquat. Sci.*, 46, 1313–1322, 1989.

Unknown
Field Code Changed

- Christensen, V. and Walters, C. J.: Ecopath with Ecosim: methods, capabilities and limitations, *Ecol. Modell.*, 172(2–4), 109–139, doi:10.1016/j.ecolmodel.2003.09.003, 2004.
- Clarke, A. and Johnston, N. M.: Scaling of metabolic rate with body mass and temperature in teleost fish, *J. Anim. Ecol.*, 68(5), 893–905, doi:10.1046/j.1365-2656.1999.00337.x, 1999.
- 5 Colebrook, J. M., Environment, N., Place, P. and Hoe, T.: Continuous plankton records: relationships between species of phytoplankton and zooplankton in the seasonal cycle, *Mar. Biol.*, 323, 313–323, 1984.
- Coull, K., Jermyn, A., Newton, A., Henderson, G. and Hall, W.: Length/weight Relationships for 88 Species of Fish Encountered in the North East Atlantic, *Scottish Fish. Res. Reports*, (43), 82, 1989.
- Cury, P. and Shannon, L.: Regime shifts in upwelling ecosystems: observed changes and possible mechanisms in the northern and southern Benguela, *Prog. Oceanogr.*, 60(2–4), 223–243, doi:10.1016/j.pocean.2004.02.007, 2004.
- 10 Cury, P., Shannon, L. and Shin, Y.-J.: The functioning of marine ecosystems, in *Responsible Fisheries in the Marine Ecosystem*, edited by M. Sinclair and G. Valdimarsson, pp. 103–123, FAO, Rome and CABI Publishing, Wallingford., 2003.
- Daan, N., Bromley, P. J., Hislop, J. R. G. and Nielson, N. A.: Ecology of North Sea Fish, *Netherlands J. Sea Res.*, 26(2–4), 343–386 [online] Available from: <http://www.mendeley.com/research/no-title-avail/>, 1990.
- 15 Daewel, U. and Schrum, C.: Simulating long-term dynamics of the coupled North Sea and Baltic Sea ecosystem with ECOSMO II: Model description and validation, *J. Mar. Syst.*, 119–120, 30–49, doi:10.1016/j.jmarsys.2013.03.008, 2013.
- Daewel, U., Peck, M. A., Kühn, W., St. John, M. A., Alekseeva, I. and Schrum, C.: Coupling ecosystem and individual-based models to simulate the influence of environmental variability on potential growth and survival of larval sprat (*Sprattus sprattus* L.) in the North Sea, *Fish. Oceanogr.*, 17(5), 333–351, doi:10.1111/j.1365-2419.2008.00482.x, 2008.
- 20 Daewel, U., Hjøllø, S. S., Huret, M., Ji, R., Maar, M., Niiranen, S., Travers-trolet, M., Peck, M. A., Wolfshaar, K. E. Van De and van de Wolfshaar, K. E.: Predation control of zooplankton dynamics: a review of observations and models, *ICES J. Mar. Sci.*, 71(2), 254–271, doi:10.1093/icesjms/fst125, 2014.
- Daewel, U., Schrum, C. and Gupta, A. K.: The predictive potential of early life stage individual-based models (IBMs): an example for Atlantic cod *Gadus morhua* in the North Sea, *Mar. Ecol. Prog. Ser.*, 534, 199–219, doi:10.3354/meps11367, 25 2015.
- Ekeröth, N., Blomqvist, S. and Hall, P. O. J.: Nutrient fluxes from reduced Baltic Sea sediment: effects of oxygenation and macrobenthos, *Mar. Ecol. Prog. Ser.*, 544(February), 77–92, doi:10.3354/meps11592, 2016.
- Elliott, M. and Hemingway, K.: Fishes in Estuaries, *Fishes in Estuaries*, i–xx, doi:10.2134/jeq2003.3750, 2002.
- Fennel, W.: Towards bridging biogeochemical and fish-production models, *J. Mar. Syst.*, 71(1–2), 171–194, 30 doi:10.1016/j.jmarsys.2007.06.008, 2008.
- Fennel, W.: Parameterizations of truncated food web models from the perspective of an end-to-end model approach, *J. Mar. Syst.*, 76(1–2), 171–185, doi:10.1016/j.jmarsys.2008.05.005, 2009.
- Fennel, W.: A nutrient to fish model for the example of the Baltic Sea, *J. Mar. Syst.*, 81(1–2), 184–195, doi:10.1016/j.jmarsys.2009.12.007, 2010.

- Froese, R. and Pauly, D.: FishBase 2000: concepts, design and data sources., ICLARM, Los Baños, Laguna, Philippines., 2000.
- Froese, R., Thorson, J. T. and Reyes Jr, R. B.: A Bayesian approach for estimating length-weight relationships in fishes, *J. Appl. Ichthyol.*, 30(1), 78–85, doi:10.1111/jai.12299, 2014.
- 5 Fulton, E., Smith, A. and Punt, A.: Which ecological indicators can robustly detect effects of fishing?, *ICES J. Mar. Sci.*, 62(3), 540–551, doi:10.1016/j.icesjms.2004.12.012, 2005.
- Fulton, E. A.: Approaches to end-to-end ecosystem models, *J. Mar. Syst.*, 81(1–2), 171–183, doi:10.1016/j.jmarsys.2009.12.012, 2010.
- Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M. and Charnov, E. L.: Effects of size and temperature on metabolic rate., *Science*, 293(5538), 2248–51, doi:10.1126/science.1061967, 2001.
- 10 Gogina, M. and Zettler, M. L.: Diversity and distribution of benthic macrofauna in the Baltic Sea. Data inventory and its use for species distribution modelling and prediction, *J. Sea Res.*, 64(3), 313–321, doi:10.1016/j.seares.2010.04.005, 2010.
- Gogina, M., Glockzin, M. and Zettler, M. L.: Distribution of benthic macrofaunal communities in the western Baltic Sea with regard to near-bottom environmental parameters. 1. Causal analysis, *J. Mar. Syst.*, 80(1–2), 57–70, doi:10.1016/j.jmarsys.2009.10.001, 2010.
- 15 Greenstreet, S.: Seasonal variation in the consumption of food by fish in the North Sea and implications for food web dynamics, *ICES J. Mar. Sci.*, 54(2), 243–266, doi:10.1006/jmsc.1996.0183, 1997.
- Greenstreet, S. P. R., Bryant, A. D., Broekhuizen, N., Hall, S. J. and Heath, M. R.: Seasonal variation in the consumption of food by fish in the North Sea and implications for food web dynamics, *ICES J. Mar. Sci.*, 54(2), 243–266 [online] Available from: <http://dx.doi.org/10.1006/jmsc.1996.0183>, 1997.
- 20 Guerra, A. and Rocha, F.: The life history of *Loligo vulgaris* and *Loligo forbesi* (Cephalopoda: Loliginidae) in Galician waters (NW Spain), *Fish. Res.*, 21(1–2), 43–69, doi:10.1016/0165-7836(94)90095-7, 1994.
- Heath, M. R.: Ecosystem limits to food web fluxes and fisheries yields in the North Sea simulated with an end-to-end food web model, *Prog. Oceanogr.*, 102, 42–66 [online] Available from: <http://www.sciencedirect.com/science/article/pii/S0079661112000213> (Accessed 5 October 2012), 2012.
- 25 Heip, C., Basford, D., Craeymeersch, J. A., Dewarumez, J. M., Dorjes, J., de Wilde, P., Duineveld, G., Eleftheriou, A., Herman, P. M. J., Niermann, U., Kingston, P., Kunitzer, A., Rachor, E., Rumohr, H., Soetaert, K. and Soltwedel, T.: Trends in biomass, density and diversity of North Sea macrofauna, *ICES J. Mar. Sci.*, 49(1), 13–22, doi:10.1093/icesjms/49.1.13, 1992.
- 30 Hinrichsen, H.-H., Lehmann, A., Petereit, C., Nissling, A., Ustups, D., Bergström, U. and Hüsey, K.: Spawning areas of eastern Baltic cod revisited: Using hydrodynamic modelling to reveal spawning habitat suitability, egg survival probability, and connectivity patterns, *Prog. Oceanogr.*, 143, 13–25, doi:10.1016/j.pocean.2016.02.004, 2016.
- Huang, L., Wu, Y., Wan, R. and Zhang, J.: Carbon, nitrogen and phosphorus stoichiometry in Japanese anchovy (*Engraulis japonicus*) from the Huanghai Sea, China, *Acta Oceanol. Sin.*, 31(4), 154–161, doi:10.1007/s13131-012-0229-5, 2012.

- Hunter, E., Metcalfe, J. D. and Reynolds, J. D.: Migration route and spawning area fidelity by North Sea plaice., *Proc. Biol. Sci.*, 270(1529), 2097–2103, doi:10.1098/rspb.2003.2473, 2003.
- ICES: Manual for the International Bottom Trawl Surveys. Series of ICES Survey Protocols. SISP 1-IBTS VIII., Ser. ICES Surv. Protoc. 1-IBTS VIII, 68, 2012.
- 5 ICES: Baltic Sea Ecoregion – Fisheries overview, , (November), 1–23, doi:https://doi.org/10.17895/ices.pub.4648, 2018a.
- ICES: Greater North Sea Ecoregion – Fisheries overview, , (November), 1–31, doi:https://doi.org/10.17895/ices.pub.4647, 2018b.
- Jennings, S., Alvsvåg, J., Cotter, A. J. R., Ehrich, S., Greenstreet, S. P. R., Jarre-Teichmann, A., Mergardt, N., Rijnsdorp, A. D. and Smedstad, O.: Fishing effects in northeast Atlantic shelf seas: patterns in fishing effort, diversity and community structure. III. International trawling effort in the North Sea: An analysis of spatial and temporal trends, *Fish. Res.*, 40(2), 125–134, doi:10.1016/S0165-7836(98)00208-2, 1999.
- 10 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R. and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Meteorol. Soc.*, 77(3), 437–471, doi:10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2, 1996.
- 15 Klaoudatos, D. S., Conides, a. J., Anastasopoulou, a. and Dulčić, J.: Age, growth, mortality and sex ratio of the inshore population of the edible crab, *Cancer pagurus* (Linnaeus 1758) in South Wales (UK), *J. Appl. Ichthyol.*, 29(3), 579–586, doi:10.1111/jai.12122, 2013.
- Kröncke, I. and Bergfeld, C.: North sea benthos: a review, *Senckenbergiana maritima*, 33, 205–268, doi:10.1007/BF03043049, 2003.
- 20 Maar, M., Friis, E., Larsen, J., Skovgaard, K., Wan, Z., She, J., Jonasson, L. and Neumann, T.: Ecosystem modelling across a salinity gradient from the North Sea to the Baltic Sea, *Ecol. Modell.*, 222(10), 1696–1711, doi:10.1016/j.ecolmodel.2011.03.006, 2011.
- Maar, M., Rindorf, A., Möller, E. F., Christensen, A., Madsen, K. S. and van Deurs, M.: Zooplankton mortality in 3D ecosystem modelling considering variable spatial–temporal fish consumptions in the North Sea, *Prog. Oceanogr.*, doi:10.1016/j.pcean.2014.03.002, 2014.
- 25 Mackinson, S. and Daskalov, G.: An ecosystem model of the North Sea to support an ecosystem approach to fisheries management: description and parameterisation, *Sci. Ser. Tech. Rep.*, 142(142), 196 pp, 2007.
- McCully, S., Scott, F. and Ellis, J.: Lengths at maturity and conversion factors for skate (Rajidae) around the British Isles, with an analysis of data in the literature, *ICES J. Mar. Sci.*, 69(10), 1812–1822, doi:10.1093/icesjms/fst048, 2012.
- 30 Megrey, B. A., Rose, K. A., Klumb, R. A., Hay, D. E., Werner, F. E., Eslinger, D. L. and Smith, S. L.: A bioenergetics-based population dynamics model of Pacific herring (*Clupea harengus pallasii*) coupled to a lower trophic level nutrient – phytoplankton – zooplankton model: Description , calibration , and sensitivity analysis, *Ecol. Modell.*, 202, 144–164, doi:10.1016/j.ecolmodel.2006.08.020, 2007.

- Michaelis, L. and Menten, M. L.: Die Kinetik der Invertinwirkung, *Biochem. Zeitschrift Beiträge zur chem. Physiol. u. Pathol.*, 49, 333–369 [online] Available from: <http://publikationen.ub.uni-frankfurt.de/frontdoor/index/index/docId/17273> (Accessed 22 August 2012), 1913.
- Möllmann, C., Kornilovs, G. and Sidrevics, L.: Long-term dynamics of main mesozooplankton species in the central Baltic Sea, *J. Plankton Res.*, 22(11), 2015–2038, 2000.
- Monod, J.: Recherches sur la croissance des cultures bactériennes, Hermann & cie, Paris. [online] Available from: <http://www.worldcat.org/title/recherches-sur-la-croissance-des-cultures-bacteriennes/oclc/6126763> (Accessed 22 August 2012), 1942.
- Müller-Navarra, S. H. and Lange, W.: Modelling tides in the Baltic Sea - A short note on the harmonic analysis of a one-year water level time series., in *Proceedings of the 6th HIROMB Scientific Workshop, St. Petersburg 2003.*, pp. 16–20, St. Petersburg., 2004.
- Neumann, T. and Schernewski, G.: Eutrophication in the Baltic Sea and shifts in nitrogen fixation analyzed with a 3D ecosystem model, *J. Mar. Syst.*, 74(1–2), 592–602, doi:10.1016/j.jmarsys.2008.05.003, 2008.
- Nøttestad, L., Giske, J., Holst, J. C. and Huse, G.: A length-based hypothesis for feeding migrations in pelagic fish, *Can. J. Fish. Aquat. Sci.* [online] Available from: http://www.nrcresearchpress.com/doi/abs/10.1139/f99-222#.VuAkJ_D9TF (Accessed 9 March 2016), 2011.
- Oguz, T., Salihoglu, B. and Fach, B.: A coupled plankton–anchovy population dynamics model assessing nonlinear controls of anchovy and gelatinous biomass in the Black Sea, *Mar. Ecol. Prog. Ser.*, 369, 229–256, 2008.
- Omstedt, A. and Hansson, D.: The Baltic Sea ocean climate system memory and response to changes in the water and heat balance components, *Cont. Shelf Res.*, 26(2), 236–251, doi:10.1016/j.csr.2005.11.003, 2006.
- Peck, M. A., Arvanitidis, C., Butenschön, M., Canu, D. M., Chatzinikolaou, E., Cucco, A., Domenici, P., Fernandes, J. A., Gasche, L., Huebert, K. B., Hufnagl, M., Jones, M. C., Kempf, A., Keyl, F., Maar, M., Mahévas, S., Marchal, P., Nicolas, D., Pinnegar, J. K., Rivot, E., Rochette, S., Sell, A. F., Sinerchia, M., Solidoro, C., Somerfield, P. J., Teal, L. R., Travers-Trolet, M. and van de Wolfshaar, K. E.: Projecting changes in the distribution and productivity of living marine resources: A critical review of the suite of modelling approaches used in the large European project VECTORS, *Estuar. Coast. Shelf Sci.*, doi:10.1016/j.ecss.2016.05.019, 2015.
- Persson, L.-E.: Temporal and spatial variation in coastal macrobenthic community structure, Hanö bay (southern Baltic), *J. Exp. Mar. Biol. Ecol.*, 68, 277–293, 1983.
- Pierce, G. J., Boyle, P. R., Hastie, L. C. and Key, L.: The life history of *Loligo forbesi* (Cephalopoda: Loliginidae) in Scottish waters, *Fish. Res.*, 21(1–2), 17–41, doi:10.1016/0165-7836(94)90094-9, 1994.
- Pinto, C., Travers-Trolet, M., Macdonald, J., Rivot, E. and Vermard, Y.: Combining multiple data sets to unravel the spatio-temporal dynamics of a data-limited fish stock., 2018.
- Politikos, D. V., Curchitser, E. N., Rose, K. A., Checkley, D. M. and Fiechter, J.: Climate variability and sardine recruitment in the California Current: A mechanistic analysis of an ecosystem model, *Fish. Oceanogr.*, (November 2017), 1–21,

- doi:10.1111/fog.12381, 2018.
- Radtke, H., Neumann, T. and Fennel, W.: A Eulerian nutrient to fish model of the Baltic Sea — A feasibility-study, *J. Mar. Syst.*, 125, 61–76, doi:10.1016/j.jmarsys.2012.07.010, 2013.
- Rees, H. L., Pendle, M. a, Waldock, R., Limpenny, D. S. and Boyd, S. E.: A comparison of benthic biodiversity in the North Sea, English Channel, and Celtic Seas\n10.1006/jmsc.1998.0438, *ICES J. Mar. Sci.*, 56(2), 228–246, doi:10.1006/jmsc.1998.0438, 1999.
- Rees, H. L., Eggleton, J. D., Rachor, E. and Vanden Berghe, E.: Structure and dynamics of the North Sea benthos., 2007.
- Reiss, H. and Krönke, I.: Seasonal variability of epibenthic communities in different areas of the southern North Sea, *ICES J. Mar. Sci.*, 61, 882–905, doi:10.1016/j.icesjms.2004.06.020, 2004.
- Reiss, H. and Krönke, I.: Seasonal variability of infaunal community structures in three areas of the North Sea under different environmental conditions, *Estuar. Coast. Shelf Sci.*, 65, 253–274, doi:10.1016/j.ecss.2005.06.008, 2005.
- Rodhe, J., Tett, P. and Wulff, F.: The Baltic and North Seas: A Regional Review of some important Physical-Chemical-Biological Interaction Processes. , in *The Sea*, Vol. 14 B, edited by A. R. Robinson and K. Brink, pp. 1033–1075, Harvard University Press., 2006.
- Rose, K. A., Fiechter, J., Curchitser, E. N., Hedstrom, K., Bernal, M., Creekmore, S., Haynie, A., Ito, S., Lluch-Cota, S., Megrey, B. A., Edwards, C. A., Checkley, D., Koslow, T., McClatchie, S., Werner, F., MacCall, A. and Agostini, V.: Demonstration of a Fully-Coupled End-to-End Model for Small Pelagic Fish Using Sardine and Anchovy in the California Current, *Prog. Oceanogr.*, doi:10.1016/j.pcean.2015.01.012, 2015.
- Schlüter, M., Hinkel, J., Bots, P. W. G. and Arlinghaus, R.: Application of the SES framework for model-based analysis of the dynamics of social-ecological systems, *Ecol. Soc.*, 19(1), doi:10.5751/ES-05782-190136, 2014.
- Schrum, C. and Backhaus, J. O.: Sensitivity of atmosphere-ocean heat exchange and heat content in the North Sea and the Baltic Sea, *Tellus - Ser. A Dyn. Meteorol. Oceanogr.*, 51(4), 526–549, doi:10.1034/j.1600-0870.1992.00006.x, 1999.
- Schrum, C., Alekseeva, I. and St. John, M.: Development of a coupled physical–biological ecosystem model ECOSMO, *J. Mar. Syst.*, 61(1–2), 79–99, doi:10.1016/j.jmarsys.2006.01.005, 2006.
- Seinä, A. and Palosuo, E.: The classification of the maximum annual extent of ice cover in the Baltic Sea 1720-1995, *Meri - Rep. Ser. Finnish Inst. Mar. Res.*, 20, 79–910, 1996.
- Shin, Y.-J. and Cury, P.: Using an individual-based model of fish assemblages to study the response of size spectra to changes in fishing, *Can. J. Fish. Aquat. Sci.*, 61(3), 414–431, doi:10.1139/f03-154, 2004.
- Shin, Y.-J., Travers, M. and Maury, O.: Coupling low and high trophic levels models: Towards a pathways-orientated approach for end-to-end models, *Prog. Oceanogr.*, 84(1), 105–112, 2010.
- Shin, Y. and Cury, P.: Exploring fish community dynamics through size-dependent trophic interactions using a spatialized individual-based model, *Aquat. Living Resour.*, 14(2), 65–80, doi:doi: 10.1016/S0990-7440(01)01106-8, 2001.
- Skogen, M. D., Søiland, H. and Svendsen, E.: Effects of changing nutrient loads to the North Sea, *J. Mar. Syst.*, 46, 23–38, doi:10.1016/j.jmarsys.2003.11.013, 2004.

- Sparholt, H.: An estimate of the total biomass of fish in the North Sea, ICES J. Mar. Sci., 46(2), 200–210, doi:10.1093/icesjms/46.2.200, 1990.
- Sterner, R. W. and George, N. B.: Carbon , Nitrogen , and Phosphorus Stoichiometry of Cyprinid Fishes, Ecology, 81(1), 127–140, 2000.
- 5 Storch, H. Von and Zwiers, F. W.: Statistical Analysis in Climate Research, Cambridge University Press, The Pitt Building, Trumpington Street, Cambridge CB2 1RP., Cambridge. UK., 1999.
- Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, J. Geophys. Res., 106(D7), 7183–7192, doi:10.1029/2000JD900719, 2001.
- Templeman, W.: Length-weight Relationships & Morphometric Characteristics and Thorniness of Thorny Skate (*Raja radiata*) from the Northwest Atlantic, J. Northwest Atl. Fish. Sci., 7, 89–98, 1987.
- 10 Thurow, F.: Estimation of the total fish biomass in the Baltic Sea during the 20th century, ICES J. Mar. Sci., 54(3), 444–461, doi:10.1006/jmsc.1996.0195, 1997.
- Timmermann, K., Norkko, J., Janas, U., Norkko, A., Gustafsson, B. G. and Bonsdorff, E.: Modelling macrofaunal biomass in relation to hypoxia and nutrient loading, J. Mar. Syst., 105–108, 60–69, doi:10.1016/j.jmarsys.2012.06.001, 2012.
- 15 Tomczak, M. T., Niiranen, S., Hjerne, O. and Blenckner, T.: Ecosystem flow dynamics in the Baltic Proper—Using a multi-trophic dataset as a basis for food–web modelling, Ecol. Modell., 230, 123–147, doi:10.1016/j.ecolmodel.2011.12.014, 2012.
- Travers, M., Shin, Y.-J., Jennings, S. and Cury, P.: Towards end-to-end models for investigating the effects of climate and fishing in marine ecosystems, Prog. Oceanogr., 75(4), 751–770, doi:10.1016/j.pocean.2007.08.001, 2007.
- Utne, K. R., Hjøllø, S. S., Huse, G. and Skogen, M.: Estimating the consumption of *Calanus finmarchicus* by planktivorous fish in the Norwegian Sea using a fully coupled 3D model system, Mar. Biol. Res., 8(5–6), 527–547, doi:10.1080/17451000.2011.642804, 2012.
- 20 Vikebø, F., Jørgensen, C., Kristiansen, T. and Fiksen, Ø.: Drift , growth , and survival of larval Northeast Arctic cod with simple rules of behaviour, Mar. Ecol. Prog. Ser., 347, 207–219, doi:10.3354/meps06979, 2007.
- Watson, J. R., Stock, C. A. and Sarmiento, J. L.: Exploring the role of movement in determining the global distribution of marine biomass using a coupled hydrodynamic – Size-based ecosystem model, Prog. Oceanogr., doi:10.1016/j.pocean.2014.09.001, 2014.
- 25

Abbr.	Definition	Value	Units
r_{MB}	MB half saturation constant	0.5	mmol C m ⁻³
m_{MB}	MB mortality rate	0.001	day ⁻¹
ϵ_{MB}	MB excretion rate	0.025	day ⁻¹
γ_{MB}	Assimilation efficiency	0.75	
$\sigma_{MB,X}$	Grazing rate	0.1	day ⁻¹

Table 1. Parameters used for the MB functional group reaction terms

Y	X	P	Z1	Z2	DET	DOM	SED1	MB
MB		0.2	0.2	0.3	0.1	0.1	0.1	-
Fi		-	0.25	0.45	0.05	-	-	0.25

5

Table 2: Feeding preferences for macrobenthos (MB) and fish (FI) ^{a_{y,x}}

Abbr.	Definition	Value	Units
r_{Fi}	Fish half saturation constant	0.7	mmol C m ⁻³
$r_{Fi,MB}$	Fish half saturation constant (MB prey)	0.9	mmol C m ⁻³
m_{Fi}	Fish mortality rate	0.001	day ⁻¹
μ_{Fi}	Fish excretion rate	0.002	day ⁻¹
θ_{Fi}	T control parameter excretion	0.5	
γ_{Fi}	Assimilation efficiency	0.7	
$\sigma_{Fi,X}$	Grazing rates F on MB, Z _{1,2}	0.01	day ⁻¹
$\sigma_{Fi,D}$	Grazing rates F on D	0.005	day ⁻¹
k	Boltzmann's factor	8.6173324*10 ⁻⁵	eV K ⁻¹

Table 3. Parameters used for the fish functional group

Ute Daewel 13.3.2019 16:13

Deleted: Macrobenthos

Ute Daewel 13.3.2019 16:13

Deleted: Fish

State variable	Reaction term
Phytoplankton	$R_{P_{1,2}} = R_{P_{1,2}} - [C_{MB}G_{MB}(C_{P_{1,2}})]_{n=bottom}$
Zooplankton	$R_{Z_{1,2}} = R_{Z_{1,2}} - G_{Fi}(P_{Z_{1,2}}) * \frac{C_{Z_{1,2}}}{P_{Z_{1,2}}} - [C_{MB}G_{MB}(C_X)/\Delta z]_{n=bottom}$
Detritus	$R_{DET} = R_{DET} - P_{Fi}G_{Fi}(P_{DET}) * \frac{C_{DET}}{P_{DET}} - [C_{MB}G_{MB}(C_{DET})]_{n=bottom}$ $+ 0.6((1 - \gamma_{Fi})R_{FiCons} + m_{Fi}C_{Fi})$
Dissolved organic matter	$R_{DOM} = R_{DOM} - [C_{MB}G_{MB}(C_{DOM})]_{n=bottom} + 0.4$ $* (((1 - \gamma_{Fi})R_{FiCons} + m_{Fi}C_{Fi}) + ((1 - \gamma_{MB})R_{MBCons} + m_{MB}C_{MB}))$
Sediments	$R_{SED1} = R_{SED1} - C_{MB}G_{MB}(C_{SED1}) + 0.6 * ((1 - \gamma_{MB})R_{MBCons} + m_{MB}C_{MB})$
Phosphate/Ammonia Oxygen	$R_{PO_4/NH_4} = R_{PO_4/NH_4} + \epsilon_{Fi}C_{Fi} + [\mu_{MB}C_{MB}/\Delta z]_{n=bottom}$ $R_{O_2} = R_{O_2} - c_{C:O_2}[\epsilon_{Fi}C_{Fi} + [\mu_{MB}C_{MB}/\Delta z]_{n=bottom}] \cdot c_{C:O_2:conversion\ factor}$

Table 4. Changes in the biogeochemical reaction terms (R) of ECOSMO due to the macrobenthos (MB) and fish functional groups.

Ute Daewel 7.3.2019 14:30

Formatted Table

Ute Daewel 5.3.2019 10:13

Deleted: μ

Ute Daewel 13.3.2019 16:13

Deleted: F




Figure 1: Model area and bathymetry. Black lines indicate the 30 m and 60 m depth respectively. Insert: area subdivision in ICES-boxes for model comparison to ICES data (see [Figure 10](#)).

Ute Daewel 5.3.2019 10:13

Formatted: Page break before

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 12.2.2019 13:37

Deleted: figure




Figure 2: Schematic diagram of biological-geochemical interactions in ECOSMO E2E.

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)




Figure 3: Simulated spatial pattern of annual mean biomass of macrobenthos (upper panel) and fish (lower panel) (gC m^{-2}).

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)




Figure 4: Average seasonality of ecosystem components. Monthly means averaged for 1980-1989. solid lines: ECOSMO-E2E; dashed lines: ECOSMO II (phytoplankton & zooplankton).

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

5




Figure 5: Prey composition of a) macrobenthos (MB) and b) fish in the North Sea (left) and the Baltic Sea (right). MB feeds on SED (organic material in the sediment), DOM/DET (dead organic material in the water column: dissolved organic matter/detritus), Z (zooplankton), P (phytoplankton). Fish feeds on Zs ("small" herbivorous zooplankton), Zl ("large" omnivorous zooplankton), DET (detritus).

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Ute Daewel 13.3.2019 16:14

Deleted: MB




Figure 6: Empirical Orthogonal Function (EOF) of macrobenthos biomass average seasonality (1980-1989). eof_{1,2}: spatial pattern of the first and second EOF mode; η: global explained variance; η_{local}: local explained variance for the first and second EOF; PC_{1,2}: temporal variability related to eof_{1,2}.

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)




Figure 7: Empirical Orthogonal Function (EOF) of fish biomass average seasonality (1980-1989). eof_{1,2}: spatial pattern of the first and second EOF mode; η : global explained variance; η_{local} : local explained variance for the first and second EOF; PC_{1,2}: temporal variability related to eof_{1,2}.

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)




Figure 8: Seasonal cycle on fish production (primary y-axis) in the shallow (depth<50m) southern North Sea (a) and in the deeper (depth> 50m) northern North Sea, divided into diet components (Zs: herbivorous zooplankton; ZI: omnivorous zooplakton; DET: detritus; MB: macrobenthos). And mean depth averaged temperature in the respective region (solid line; secondary y-axis).

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)





Figure 9) Half-annually averaged (1980-1989) biomass specific mortality (d^{-1}) of zooplankton due to a) fish predation and b) macrobenthos predation.



Deleted:

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Ute Daewel 4.3.2019 16:31

Deleted: Figure 9: Average (1980-1989) production of fish (a) and macrobenthos (b) based on zooplankton consumption only. The measure serves as an indicator for zooplankton predation mortality.

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Ute Daewel 4.3.2019 16:46

Formatted: Tabs: 5.13 cm, Left

Ute Daewel 4.3.2019 16:31

Deleted:

Ute Daewel 4.3.2019 16:54

Formatted





Figure 2: a-c) Mean (1980-1989) total fish biomass from the ICES IBTS survey & fish biomass from ECOSMO E2E for the associated sampling time and area in the first quarter of the year (Jan-Mar) for demersal species (a), pelagic species (b), and combined biomass (c). Left panels: spatial distribution of fish biomass. Right panels: Biomass versus latitude and the mean of biomass at latitude (black line).

Formatted: Font:12 pt, Font color: Text 1, Kern at 12 pt

Ute Daewel 4.3.2019 16:32



Deleted:

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Unknown

Formatted: Font:12 pt, Font color: Text 1, Kern at 12 pt

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Ute Daewel 4.3.2019 16:46

Ute Daewel 4.3.2019 16:54

Formatted

[18]

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:34

[19]

Model Validation ICES Data




Figure 10: Taylor diagram for surface (<10m) nutrients (model versus ICES data) in different areas of the North Sea (area separation in ICES boxes according to [Figure 1](#)) (nitrate (a); phosphate (b)). Arrows indicate regions with relatively large changes in the validation measures.

Ute Daewel 4.3.2019 16:37

Formatted: Page break before

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Ute Daewel 4.3.2019 16:46

Deleted: 11

Unknown

Field Code Changed

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)




Figure 11: Modeled (blue: ECOSMO II; green ECOSMO E2E) and observed (HELCOM data, black) vertical nutrient profiles. Data were averaged over the 10-year period 1980-1989 and mean (full line) and standard deviation (dashed line) are presented at two distinct locations in the Baltic Sea (BY15 (a); BY5 (b) see [Figure 1](#)).

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:46

Deleted: 12

Ute Daewel 13.3.2019 16:15

Deleted:

Ute Daewel 13.3.2019 15:37

Deleted: Fig.

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)




Figure 12: Spider plots showing averaged changes in North Sea and Baltic Sea annual mean phytoplankton, zooplankton, MB, and fish biomass (mgC m⁻³) due to specific changes in the food web components. a,b: scenarios for fisheries mortality (FM); c,d: scenarios for changes in zooplankton natural mortality (ZM) .

Ute Daewel 13.3.2019 15:34

Deleted: Figure

Unknown

Field Code Changed

Ute Daewel 4.3.2019 16:46

Deleted: 13

Ute Daewel 4.3.2019 16:54

Formatted: English (UK)