

## ***Interactive comment on “EcH<sub>2</sub>O-iso 1.0: Water isotopes and age tracking in a process-based, distributed ecohydrological model” by Sylvain Kuppel et al.***

**Sylvain Kuppel et al.**

[sylvain.kuppel@abdn.ac.uk](mailto:sylvain.kuppel@abdn.ac.uk)

Received and published: 4 June 2018

The authors would like to thank the Referee #1 for her/his valuable comments and suggestions to improve the manuscript. They have been taken into account in the revised manuscript, as follows (original referee's comments in bold):

C1

### **General comments**

**This paper introduces the isotopic tracking of the ecohydrological model EcH<sub>2</sub>O. The new model development is evaluated using isotope time series from a montane, low-energy catchment in Scotland. The isotope tracking addition to the model is interesting for the GMD readership and the approach is in general well-documented. The availability of isotope time series in different parts of the study catchment is also very useful for gaining scientific insights. However, . . .**

**1. The paper lacks a clear focus at times, and the writing varies between being very detailed to very general. The authors know the topics very well, and occasionally make jumps or sweeping descriptions that easily lose the reader. (Examples in specific Comments.)**

We thank the referee for this comment. In general, we have tried to make the narrative more consistent as specified in the corresponding specific comments below.

**2. Also, the model development rationale is not entirely clear, which makes it difficult to understand whether the evaluation procedure and criteria are sound, well defined, and in proportion to the goals the model are set to achieve.**

The objective of the paper is to describe and demonstrate the development of a flux/age tracking component built on an existing ecohydrological model. The rationale for this new development is twofold. First, the tracking component is added to a spatially-distributed energy and water balance model with a strong physical base that explicitly simulates the spatio-temporal heterogeneity of the water mixing processes. Then, we evaluate how this ecohydrological model calibrated solely on hydrometric/energy balance data could simulate spatio-temporal isotope variations without any additional calibration of the tracking and fractionation components. Because of the diversity of fluxes and storage dynamics tracked in the model, we put the emphasis on testing the new model with a wide range of isotopic datasets, and use visual inspection and generic

C2

quantitative metrics (such as mean absolute error and model-data correlation, see response to the corresponding specific comment) for a generic evaluation and further discussion. In the future, when moving towards more operational purposes, specific calibrations of the isotopic component using metrics such as KGE or NSE may be beneficial. Note that specific aspects of this discussion relevant to the model development rationale, and the evaluation metrics are further addressed below in specific Comments.

**3. The authors also do not test the sensitivity of neither parameters, mixing assumptions, nor isotope model structure, which limit the insights that could have been generated in the subsequent evaluation process.**

A comprehensive sensitivity analysis of parameters was already performed by Kuppel et al. (2018), along with a description of the ensemble of parameters used in this paper, which were derived from a multi-objective calibration method conducted using constraints from hydrometric and energy balance observations (see Sect. 3.3). Additional parameter sensitivity analysis and calibration using isotopes datasets would provide complementary information to further constrain parameter uncertainty. However, by doing so we would lose an opportunity to assess how the original ECH<sub>2</sub>O structure performs against a dataset that is truly independent from the standard hydrometric information typically used in model calibration/validation exercises. A comparison of the performance of different mixing models is beyond the scope of the paper. The presented model simulates isotope tracking using a simple full mixing assumption, which avoids hard-to-test partial or incomplete mixing hypotheses and therefore permits an in-depth discussion of model strengths and weaknesses for potential applications and hypothesis-driven model developments.

**4. The authors repeatedly refer to Kuppel et al. (2018) and at times assume the reader to have taken part of it. This is a bit unfortunate, as Kuppel et al. (2018) is not open access (and also was not accessible for me during my review). Please consider including key information, if only in Supplementary information.**

C3

It is difficult not to refer extensively to Kuppel et al. (2018) because it gives key details such as a description of improvements to the original ECH<sub>2</sub>O model, a quantification of the model performance on the study site and also describes the basic configuration used to evaluate ECH<sub>2</sub>O-iso. While it is not possible to reproduce all this information in the paper, we strived to include the information relevant for the interpretation of the results of the present study. Nevertheless, we recognize this can be frustrating. To ameliorate this problem we revised the manuscript to add further details regarding model development rationale, key features and limitations, and the range of environments on which the model has been successfully applied. We have also added a figure to the Supplementary Information that indicates the time spans used for calibration and evaluation (Fig. S1) and the list and description of the calibrated parameters (Table S1, adapted from Kuppel et al., 2018):

*"Table S1. Calibrated parameters used in this study, grouped according to their four components: soil units or vegetation types."*

C4

Name	Description
<b>Soil-distributed</b> (Peat, Gley, Podzol, Ranker)	
$D_{soil}$	Total soil depth (m)
$D_{L1}$	Depth of the 1st hydrological layer (m)
$D_{L2}$	Depth of the 2nd hydrological layer (m)
$\phi$	Porosity ( $\text{m}^3 \cdot \text{m}^{-3}$ )
$K_{hx}$	Saturated horizontal hydraulic conductivity ( $\text{m} \cdot \text{s}^{-1}$ )
$K_{hratio}$	Ratio of vertical-to-horizontal hydraulic conductivity (-)
$\lambda_{BC}$	Brooks-Corey exponent parameter (-)
$\Psi_{ae}$	Air-entry pressure head (m)
$\theta_r$	Residual soil moisture ( $\text{m}^3 \cdot \text{m}^{-3}$ )
$k_{root}$	Exponential root profile ( $\text{m}^{-1}$ )
<b>Vegetation-distributed</b> (Pine, Hather, Moss Grass)	
$gs_{max}$	Maximal stomatal conductance ( $\text{m} \cdot \text{s}^{-1}$ )
$CWS_{max}$	Maximum interception storage per unit LAI (m)
$T_{opt}$	Optimal photosynthesis temperature (C)
$\Psi_d$	Soil water potential halving stomatal conductance (-m)
$c$	Sensitivity of stomatal conductance to soil water potential (-)
$K_{beer}$	Light attenuation coefficient (-)

(see Figure at the end of the response) **Figure S1.** *Temporal windows –at daily resolution– covered by each of the datasets (orange) at the different sites (italic font) grouped by observation type (bold font) used to calibrate the ECH<sub>2</sub>O-iso model, while the full simulation period (blue) is used for evaluating the isotopes and age tracking module.*"

**5. The paper is lengthy and readability could be improved by e.g., summarising tables and more condensed graphs that can act as reference, or point the reader to the key results (e.g., notations table, definitions table, and scatterplots etc., more figures like Fig 8).**

C5

Striking a balance between providing sufficient detail while keeping the paper concise is challenging. We moved away much of the methodological details to Supplementary Materials or to Kuppel et al (2018) and much of the bulk of the paper describes and discusses results on the temporal and spatial patterns of water compositions and age, which are key foci of our study. Nonetheless, we have edited the manuscript to reduce verbosity and have moved the section and figures on Ic-excess (Fig. 11) to Supplementary Information, as they offered similar information to Fig. 10.

**6. The authors mention in their literature review and discussions other models ranging from local to global scale, but it's not clear if the authors mean that their modelling procedure can be scaled up.**

The review was meant to contextualize the model within the state of the art, and to indicate that other similar models with different strengths and weaknesses exist. However, one of the features of the ECH<sub>2</sub>O model is that it can be run at a wide range of spatial scales, provided that the necessary inputs are available. Indeed, its spatial domain is constructed and determined by a regular-gridded digital elevation model (DEM) map that defines the topography and the drainage network, and establishes the finite-differences grid on which the governing equations are solved (Maneta Silverman, 2013). Currently applications have been conducted at the plot scale (Maneta Silverman, 2013; Douinot et al., *Plot scale modelling to asses forest effects on water partitioning and flux ages*, in prep.), in small catchments (1-10 km<sup>2</sup>) (Kuppel et al., 2018; Lozano-Parra et al., 2014), in larger watersheds and small regions (10<sup>2</sup>-10<sup>3</sup> km<sup>2</sup>) (Maneta and Silverman, 2013; Simeone, 2018). While these studies obviously did not include isotopic tracking, the hydrologic core is the same. In the revised manuscript, we added the following sentence in the model description (sect. 2.1) in order to emphasize this multi-scaling potential:

*[...] relative humidity, and wind speed). In addition, the flexible definition of the spatial domain in ECH<sub>2</sub>O allows for applications at a range of scales: from the plot (Maneta and Silverman, 2013), to small catchments (1-10 km<sup>2</sup> – Lozano-Parra et al., 2014;*

C6

Kuppel et al., 2018), to larger watersheds ( $10^2$ - $10^3$  km $^2$  – Maneta and Silverman, 2013; Simeone, 2018)."

**7. In the abstract and conclusions, the authors claim that the framework is useful beyond the type of low energy catchment simulated here. However, I feel this statement is misleading and goes well beyond the evidence provided in the paper, and would require e.g. validation in other types of catchments.**

The abstract has been edited to avoid making such claim (see the first specific Comment below). In the conclusions however, we argue that the ECH<sub>2</sub>O-iso was not specifically designed for simulating the kind of catchment here studied and that it has applicability to other regions. In addition, at present the implementation of isotope and age tracking also avoids any location-specific parameterization. From a methodological viewpoint, the specificity of our site lies not so much in its environmental conditions but rather in the richness of available datasets. As a result, there is no reason to think that our methodology (including the ECH<sub>2</sub>O-iso model) could not perform well in other environments. The conclusions have been modified in the revised manuscript to emphasize this aspect (P30L14):

*"Despite some limitations, this isotope-based evaluation suggests a reasonable capture of the velocity fields (i.e., how fast water parcels move) across the catchment, and complements a previous calibration and evaluation mostly using hydrometric observations (water fluxes and storage dynamics) which indicated a good simulation of catchment functioning from a celerity viewpoint (i.e., how fast energy propagates via the hydraulic gradient) (Kuppel et al., 2018). Satisfying this dual velocity-celelity perspective is key to characterising water pathways and quantifying the associated travel times in different ecohydrological compartments of headwater landscapes. Complementing more conceptual approaches, the physical basis of the ECH<sub>2</sub>O-iso model further provides the potential to extrapolate these insights beyond recorded conditions and scales, and to notably project the reciprocal feedbacks between plant water use, hydrological pathways and potential environmental changes. The relatively simple con-*

C7

*ceptualisation of compartment-scale velocities, e.g. assuming complete mixing and without site-specific parameterization, and the absence of isotopic calibration, already make the current results particularly encouraging. It also provides a useful framework for hierachising model development and benchmarking needs. For example, some of the model-data discrepancies in our results stress the necessary incorporation of partial mixing hypotheses, likely to be critical in drier and/or flatter landscapes where diffusive water movement prevails. Second, our model-data analysis of isotope dynamics strongly reflects fractionation effects, be it via soil evaporation or species-specific plant water use. Finally, the versatility of climatic settings in which the original ECH<sub>2</sub>O model has already been evaluated facilitates applying the presented methodology beyond the specifics of a high-latitude, low-energy, wet and steep headwater catchment such as the one simulated here. Further, the flexible spatial domain used by the model will help providing a process-based modelling framework for plot-to-catchment-scale hypothesis testing. This is timely for current challenges in critical zone science, such as exploring the occurrence and mechanisms behind the postulated ecohydrological separation of water fluxes (Berry et al., 2017)."*

**8. Equations: subscripts and superscript should be in upright font when constituting a describing word (e.g., out, in, snow etc.) and only in cursive for variables (e.g., t). Function names such as "max" and "min" should also be in upright font.**

We thank the referee for this suggestion. All subscript and superscript notations, as well as function names, have been formatted accordingly in the revised manuscript.

### Specific Comments

**Abstract: Very sweeping and general, and raises many questions. Please consider to be more specific. E.g., what is meant by "good [. . .] match in most cases", "powerful tool", "some model development"? What kind of cases, why**

C8

is it powerful, what kind of model development? And what is the model development rationale? What can the model be used for? “Celerity” – a term used in the abstract, introduction, discussion and conclusion, but not clearly explained in the analyses and results sections.

We thank the Referee for this. We have edited the abstract to add precision, as well as to add specificity to the rationale and potential model applications. We also addressed the issue of making “celerity” easier to understand / redefined when used here and in other parts of the manuscripts (see other corresponding comments). In the revised manuscript, the abstract now reads as follows:

*“We introduce ECH<sub>2</sub>O-iso, a new development of the physically-based, fully-distributed ecohydrological model ECH<sub>2</sub>O where the tracking of water isotopic tracers (<sup>2</sup>H and <sup>18</sup>O) and age has been incorporated. ECH<sub>2</sub>O-iso is evaluated at a montane, low-energy experimental catchment in northern Scotland using 16 independent isotope time series from various landscape positions and compartments; encompassing soil water, groundwater, stream water, and plant xylem. The results show consistent isotopic ranges and temporal variability (seasonal and higher-frequency) in across the soil profile at most sites (especially on hillslopes), a broad model-data agreement in heather xylems, and consistent deuterium dynamics in stream water and in groundwater. Since ECH<sub>2</sub>O-iso was calibrated only using hydrometric and energy flux datasets, tracking water composition provides a truly independent validation of the physical basis of the model for successfully capturing catchment hydrological functioning, both in terms of celerity of energy propagation shaping the hydrological response (e.g. runoff generation under prevailing hydraulic gradients), and of flow velocities of water molecules (e.g., in consistent tracer concentrations at given locations and times). Additionally, we also show that the spatially-distributed formulation of ECH<sub>2</sub>O-iso provides the possibility to quantitatively link water stores and fluxes with spatio-temporal patterns of isotopes ratios and water ages. However, our study case also highlights model-data discrepancies in some compartments, such as an over-dampened variabil-*

C9

*ity in groundwater and stream water  $l_c$ -excess, and over-fractionated riparian topsoils. The adopted minimalistic framework, without site-specific parameterization of isotopes and age tracking, facilitates the interpretation of these mismatches into model development and benchmarking needs, while taking into account the idiosyncrasies of our study catchment. Notably, we suggest that more advanced conceptualisation of soil water mixing and of plant water use would be needed to reproduce some of the observed patterns. Balancing the need for basic hypothesis testing with that of improved simulations of catchment dynamics for a range of applications (e.g., plant water use under changing environmental conditions, water quality issues, and calibration-derived estimates of landscape characteristics), further works could also benefit from including isotope-based calibration.”*

**Introduction: It could be useful for the authors to explain how such their study is linked to practical and societal meaningful issues.** For example, the authors explains many times how isotopic characterisation could “provide insights into water pathways”, linked to “water flux partitioning”, and understanding “catchment functioning”, but the reader is left to figure out on her own if these topics are interesting and important also in a broader context. E.g., could improving our understanding of catchment functioning also for example be directly linked to our capacity to design models capable of forecasting floods, and works well under a rapidly changing climate? No need to be lengthy, but just to provide a context. Some interesting debates about evaporation partitioning is also not included, among others: (Coenders-Gerrits et al., 2014; Evaristo et al., 2015; Jasechko et al., 2013; Schlesinger and Jasechko, 2014; Wei et al., 2017).

We thank the Referee for bringing this perspective. In the revised manuscript, the end of the first paragraph of the Introduction has been modified in this regard (P2L6):

*“[...] water pathways at scales ranging from the pedon (Sprenger et al., 2018) to the catchment landscape (McGuire and McDonnell, 2006; Birkel and Soulsby, 2015). At larger scales, such approaches can yield global estimates of terrestrial water flux par-*

C10

titioning (Good et al., 2015), where recent scrutiny has been brought upon separating plant transpiration from other source of evaporative losses (e.g., Jasechko et al., 2013; Coenders-Gerrits et al., 2014; Schlesinger and Jasechko, 2014; Wei et al., 2017). At catchment and watershed scales, an understanding of landscape functioning in turn helps designing robust models to predict the impact of climate extremes and environmental changes in society-relevant issues such as water resources management, flood forecasting, and impact assessment of land cover – land use change (e.g., Troy et al., 2015; Zhang et al., 2017)."

**P2L10: What do the authors mean when writing that the “hydrology has remained simplistic” in land surface models? Please specify. And why are dynamic vegetation models and global hydrological models not mentioned?**

Given the scope of the paper, we only refer to land surface models on which (to our knowledge) isotopes tracking have been implemented: JSBACH (Haese et al., 2013), ORCHIDEE (Risi et al., 2016) and CLM (Wong et al., 2017). These land surface model have simplified descriptions of the hydrologic system that do not include explicit laterals water transfers (or is represented using a calibrated residence time for linear storage decrease), shallow and deeper subsurface flows, or channel routing. This is made clearer in the revised manuscript. Some of these models also include vegetation dynamics, and this will be mentioned. To our knowledge, no global hydrological model integrates isotope tracking. The revised manuscript has been modified as follows:

*[...] While the simulation of energy budgets and biogeochemical cycles is increasingly detailed in these land surface models -sometimes including vegetation dynamics- the hydrology has, however, remained somewhat simplistic (or even absent) regarding lateral transfers as overland flow, shallow and deeper subsurface flows and channel routing (Fan, 2015). This makes it difficult to take advantage of isotopes tracking to characterise the role of cascading downstream water redistribution in the spatial patterns of catchment functioning. [...]"*

C11

**P3L9: “evaporative losses in ET”. Please consider “terrestrial evaporation”.**

It has been modified to “terrestrial evaporation” in the revised manuscript. In addition, we now use “E” instead of “ET” to refer to evapotranspiration.

**P3L11: “transpiration (T)”. Please consider using “Et” for transpiration, to avoid the confusion with temperature T.**

“T” has now been replaced by “ $E_t$ ” throughout the revised manuscript.

**P3L19: Key features are described, but the rationale is not explained. E.g., why the model developed is the described way? What are the authors hoping to achieve?**

We have modified the introduction so that the rationale of the original  $ECH_2O$  development explains our choice for developing an isotopes and age tracking module (P3L20): *“Here, we implement isotope and age tracking in the physically-based, fully-distributed model  $ECH_2O$  (Maneta and Silverman, 2013). This model was chosen because of it provides a physically-based, yet computationally-efficient representation of energy-water-ecosystem couplings where intra-catchment connectivity (both vertical and lateral) can be explicitly resolved. In addition,  $ECH_2O$  separately solves the energy balance at the top of the canopy and at the soil surface, allowing a process-based separation of  $E_s$ ,  $E_t$ , and  $E_c$ . The novel isotopic and age [...]”*

**P3L28: Please consider new paragraph for the research questions.**

**P3L30: The research questions could be formulated in a more specifically way. E.g., What are “physics”? Are “mixing assumptions” really investigated in this paper? What kind of “implications and opportunities” do the authors have in mind?**

We answer to these two comments jointly. These are good suggestions, and the research questions are now shown as a list in the revised manuscript, for further clarity. In addition, these questions have been modified as follows: *“We ask the following*

C12

questions:

- To what extent can a hydrometrically-calibrated, physically-based hydrologic model correctly reproduce internal catchment dynamics of isotopes?
- What are the limitations of these isotopic simulations? Do they relate to the underlying model physics and/or to the tracking approach adopted?
- How useful and transferrable is this model framework for simulating spatio-temporal patterns of isotopes and water ages?"

**P4 Sect 2.1: Please describe the key features and main limitations of the ECH<sub>2</sub>O model. Including examples of where and for what kind of purposes the model has been used would also be useful.**

We have extended the first paragraph of Sect. 2.1 to include a further description of ECH<sub>2</sub>O and examples of past applications:

"[...] relative humidity, and wind speed). In addition, the flexible definition of the spatial domain in ECH<sub>2</sub>O allows for applications at a range of scales: from the plot (Maneta and Silverman, 2013), to small catchments (1-10 km<sup>2</sup> – Lozano-Parra et al., 2014; Kuppel et al., 2018), to larger watersheds (10<sup>2</sup>-10<sup>3</sup> km<sup>2</sup> – Maneta and Silverman, 2013; Simeone, 2018). Despite some potential limitations due to the absence of diffusion-driven water redistribution or an explicit biogeochemical cycle providing ecosystem respiration, to date the model yielded satisfactory results and insights across the diversity of climatic settings (semiarid to humid/energy-limited) and scientific focuses (e.g., water balance, energy balance, or plant hydraulics) covered by the aforementioned studies. A comprehensive description of ECH<sub>2</sub>O can be found [...]"

**P5 Fig 1: Please consider illustrating the isotope tracking assumptions within the model chart, e.g., transpiration is not considered fractioning, throughfall is not aging etc.**

C13

This a good suggestion, we have modified Fig. 1 and its caption, so that fractionating processes appears more explicitly but we kept the throughfall assumptions in the main text in order not overload the figure (the revised figure can be found at the end of this document):

*"Figure 1. Water compartments (black rectangles) and fluxes (coloured arrows) as represented in ECH<sub>2</sub>O, with the dashed arrows indicating processes where isotopic fractionation is simulated. The numbers between brackets reflect the sequence of calculation within a time step. Note that water routing (steps [8] to [13]) differs between cells where a stream is present (○) or not (\*)."*

**P6L9: "One exception. . ." Perhaps new paragraph?**

It has been amended in the revised manuscript.

**P6L14 "No spill-over". Not sure what is meant. There is throughfall, right?**

By "no spill-over", we meant that since in the ECH<sub>2</sub>O model "canopy drainage occurs at the rate at which precipitation increases above the maximum canopy storage" (Maneta and Silverman, 2013), and because maximum canopy storage is constant in our simulations, only the precipitation from the current time step can contribute to throughfall. This is the reason why throughfall does not age, as correctly pointed out by Referee1 a few paragraphs above. We made this clearer in the revised manuscript:

*"[...]. Only the same-time-step precipitation can contribute to throughfall in the ECH<sub>2</sub>O model, whenever the resulting canopy storage would exceed the maximum canopy storage capacity (Maneta and Silverman, 2013), the latter being constant in our simulations. As a result, intercepted water eventually evaporates from the canopy and does not interact with the surface/subsurface. [...]"*

**P8L13 "PET" Please consider using Epot, as PET could also be precipitation, evaporation, and temperature.**

We realized that this acronym is not used anywhere else in the manuscript, so was

C14

removed from the revised manuscript.

#### P11L3-4 “Autumn” Lowercase letters

It has been corrected in the revised manuscript

**P13L26-27 “model-to-data ratio of standard deviation and model-data Pearson’s correlation factor”. Please consider discussion the merits and pitfalls of using these evaluation metrics. See for example (Biondi et al., 2012) for review of different validation procedures that might be of relevance.**

We thank the Referee for this suggestion. As stated in our reply to General Comment 2, our approach consists in a generic evaluation of the new model using an ensemble of diverse isotopic datasets across ecohydrological compartments. This is why we rely on visual inspection (recommended in Biondi et al., 2012) as well as on generic metrics of model skill. The mean absolute error gives a generic quantification model-data fit across different type with lower sensitivity to high values within time series, contrary to metrics based on squared differences (such as RMSE or NSE; Krause et al., 2005; Legates and McCabe, 1999) and, to a lesser extent the Kling-Gupta Efficiency (KGE, Kling et al., 2012). In addition, NSE and KGE have been developed primarily for extracting information (and scores) from stream hydrograph for time series with a large number of points and pronounced variability, which is not the case for most isotopic datasets used here. Following the Referee’s comment, in the revised manuscript we have used the mean absolute error (MAE) and Pearson’s correlation factor as reference metrics. First, model-data MAE is shown for all relevant time series (displayed the ensemble median so as not to overload the figures). Second, Fig. 8 has been modified in order to display the normalized MAE (using the range of values of observations) against the Pearson’s correlation factor:

***“Figure 8. Summary of model performance in the dual space of mean absolute error (normalized by the observed range of values) and Pearson’s correlation factor between modelled and observed time series, for (a)  $\delta^{2}\text{H}$  and (b)  $\text{Ic}$ -excess, showing the median***

C15

*and 90%-spread over the ensemble. The size of each symbol is proportional to the logarithm of the number of observation points available. Performances in soil compartments at Forest site A are further separated between periods 2013 and 2015-2016 (the latter indicated with an asterisk), corresponding to two separate field data collection campaigns. Two groundwater wells are presents at the peat site.”*

As pointed out in Biondi et al. (2012) and elsewhere, normalized MAE provides a more balanced evaluation, permits a direct comparison between different types of observables of varying distributions and dynamics, and is sensitive to model biases. The Pearson’s correlation on the other hand captures very well if the model and observables have similar variances, but does not capture biases and is not robust to outliers, especially for time series with few points and/or low variability (groundwater and xylem). In response to this comment these edits have been brought to the main text in Sect 3.4 (P13L26):

*“As outlined in Sect. 1, our model evaluation is meant to test the ability of  $\text{ECH}_2\text{O}$  to generically simulate isotope dynamics across compartments. We used mean absolute error (MAE) to quantify model-data fit for all isotopic outputs, some of which present low temporal variability, have skewed distributions, or have a relatively lower sampling record and resulting in typical hydrograph-oriented efficiency metrics (e.g., Nash-Sutcliffe or Kling-Gupta, Nash and Sutcliffe, 1970; Kling et al., 2012) being less applicable. The median value are shown on corresponding time series (Figs. 3–7). It is then normalized by each dataset range and used in conjunction with Pearson’s correlation factor in Fig. 8 as a summary of model performance. The correlation coefficient axis in this dual model performance space represents the quality of the model in representing the variation of the data, while the normalized MAE axis provides information on the accuracy (bias) of the model.”*

And in section 4.1 (P17L13):

*“A summary of model performance is shown in Fig. 8 for all sites/compartments, using*

C16

*the dual space of normalized MAE (using each dataset range, x-axis) and Pearson's linear correlation factor (y-axis). The vast majority of median normalized MAE were below 1, and more than half of evaluated datasets showed values below 0.5. Values above 0.7 were mostly found for groundwater and xylem compartments, a clustering especially marked for  $\delta^2H$ . In addition, most median model-data correlations were significantly positive between 0.4 and 0.85, noting a tighter clustering around high values for  $\delta^2H$  than  $l_c$ -excess. Insignificant or negative correlations were mostly found where only a few data points were available (xylem) or where seasonal variability was low (e.g. groundwater). Interestingly, median model-data agreement in topsoil at Forest site A significantly differed between 2013 (mobile water sampling via lysimeters) and the 2015-2016 period (bulk water sampling via direct equilibration). This was notable in the dramatic increase of model-data correlation (0.17 to 0.8) and decrease of normalized MAE (0.5 to 0.25) for topsoil  $\delta^2H$  in the latter case, which is consistent with our interpretation that the simulated soil water composition represents that of bulk water."*

**P14 Sect 4.1.** The time series section is detailed and provide considerable amount of information. However, it is also difficult for the reader to quickly get a grasp of the main strength and weaknesses of the model. Please consider including e.g., scatterplots.

Fig. 8 of the revised manuscript provides the recommended scatterplots (see reply to previous referee comment). We have also edited the manuscript to facilitate the interpretation of the figure and guide the reader through the description of the evaluation metrics (at the very end of Sect. 3.4).

**P21 Fig 9, P23 Fig 10: Possibly consider moving some of the maps to the SI, and condense the information by grouping (by e.g., riparian/upstream/downstream etc types of regions).**

Please refer to our reply to General Comment 5.

**P26L9- "By keeping the. . ." Parts of this could also be modelling rationale that**

C17

**could been useful in the introduction section or model set-up.**

Following this suggestion, we have emphasized this aspect in the abstract (see related comment above), and at the end of the introduction just after the list of research questions (also reformulated, see related comment above):

*"These questions are here addressed by testing the new tracer-enhanced model ( $ECH_2O$ -iso, Sect. 2) in a small, low-energy montane catchment (Sect. 3). This site has previously been modelled applying  $ECH_2O$  for calibration, using multiple datasets of long-term ecohydrological fluxes and storage variables (Kuppel et al., 2018). We take advantage of this earlier work as a reference ensemble of calibrated model parameterizations, and no additional isotopic calibration is conducted. In addition to using long-term, high resolution isotopic datasets for rainfall and runoff ( $2H$  and  $18O$ ), we assess the spatio-temporal variations of model-data agreement in soil water, groundwater, and plant xylem at different locations (Sect. 4.1). Following this generic evaluation, the model is used to infer seasonally-varying patterns of water fluxes and isotopes signatures (Sect. 4.2), and water age (Sect. 4.3). Model strengths and weaknesses, insights in processes and potential ways forward are discussed in Sect. 5, before drawing conclusions in Sect. 6."*

**P3019: "ecohydrological feedbacks". A bit general, and not clear what the authors mean. Ecosystem response in terms of CO2 fertilisation and root depth development?**

The term "reciprocal ecohydrological feedbacks" here only encapsulates the reciprocal feedbacks between plant water use and terrestrial water pathways, in the face of environmental change. Given the simplified biogeochemistry used in  $ECH_2O$ , the effect of CO2 fertilization, of changes in nutrient availability, or of rooting depth development cannot be explored at present. We have modified this sentence in the revised manuscript as follows:

*"[...] Complementing more conceptual approaches, the physical basis of the  $ECH_2O$ -*

C18

*iso model further provides the potential to extrapolate these insights beyond recorded conditions and scales, and to notably project the reciprocal feedbacks between plant water use, hydrological pathways and potential environmental changes.”*

## References

Berry, Z. C., Evaristo, J., Moore, G., Poca, M., Steppe, K., Verrot, L., Asbjornsen, H., Borma, L. S., Bretfeld, M., Hervé-Fernández, P., Seyfried, M., Schwendenmann, L., Sinacore, K., De Wispelaere, L., and McDonnell, J.: The two water worlds hypothesis: Addressing multiple working hypotheses and proposing a way forward, *Ecohydrology*, p. e1843, <https://doi.org/10.1002/eco.1843>, 2017.

Biondi, D., Freni, G., Iacobellis, V., Mascaro, G. and Montanari, A.: Validation of hydrological models: Conceptual basis, methodological approaches and a proposal for a code of practice, *Phys. Chem. Earth, Parts A/B/C*, 42–44, 70–76, doi:10.1016/j.pce.2011.07.037, 2012.

Birkel, C. and Soulsby, C.: Advancing tracer-aided rainfall–runoff modelling: a review of progress, problems and unrealised potential, *Hydrol. Process.*, 29(25), 5227–5240, 2015.

Coenders-Gerrits, A. M. J., van der Ent, R. J., Bogaard, T. A., Wang-Erlandsson, L., Hrachowitz, M. and Savenije, H. H. G.: Uncertainties in transpiration estimates, *Nature*, 506(7478), E1–E2, doi:10.1038/nature12925, 2014.

Evaristo, J., Jasechko, S. and McDonnell, J. J.: Global separation of plant transpiration from groundwater and streamflow, *Nature*, 525(7567), 91–94, doi:10.1038/nature14983, 2015.

Good, S. P., Noone, D. and Bowen, G.: Hydrologic connectivity constrains partitioning of global terrestrial water fluxes, *Science*, 349(6244), 175–177, doi:10.1126/science.aaa5931, 2015.

Jasechko, S., Sharp, Z. D., Gibson, J. J., Birks, S. J., Yi, Y. and Fawcett, P. J.: Terrestrial water fluxes dominated by transpiration, *Nature*, 496(7445), 347–50, doi:10.1038/nature11983, 2013.

Kling, H., Fuchs, M., Paulin, M.: Runoff conditions in the upper Danube basin under an ensemble of climate change scenarios. *J. Hydrol.* 424, 264–277, 2012.

Krause, P., Boyle, D.P., Bäse, F.: Comparison of different efficiency criteria for hydrological model assessment. *Adv. Geosci.* 5, 89–97, 2005.

Kuppel, S., Tetzlaff, D., Maneta, M. P., and Soulsby, C.: What can we learn from multi-data calibration of a process-based ecohydrological model?, *Environmental Modelling Software*, 101, 301–316, <https://doi.org/10.1016/j.envsoft.2018.01.001>, 2018.

Legates, D.R., McCabe, G.J.: Evaluating the use of “goodness-of-fit” measures in hydrologic and hydroclimatic model validation. *Water Resour. Res.* 35, 233–241, 1999.

Lozano-Parra, J., Maneta, M. P., and Schnabel, S.: Climate and topographic controls on simulated pasture production in a semiarid Mediterranean watershed with scattered tree cover, *Hydrology and Earth System Sciences*, 18, 1439, 2014.

McGuire, K. J. and McDonnell, J. J.: A review and evaluation of catchment transit time modeling, *J. Hydrol.*, 330(3), 543–563, 2006.

Maneta, M. P. and Silverman, N. L.: A spatially distributed model to simulate water, energy, and vegetation dynamics using information from regional climate models, *Earth Interactions*, 17, 1–44, 2013.

Schlesinger, W. H. and Jasechko, S.: Transpiration in the global water cycle, *Agric. For. Meteorol.*, 189–190, 115–117, doi:10.1016/j.agrformet.2014.01.011, 2014.

Simeone, C.: Coupled ecohydrology and plant hydraulics model predicts Ponderosa seedling mortality and lower treeline in the US Northern Rocky

Mountains, Master Thesis, University of Montana. [online] Available from: <https://scholarworks.umt.edu/etd/11128>, 2018.

Sprenger, M., Tetzlaff, D., Buttle, J., Laudon, H., Leistert, H., Mitchell, C. P., Snelgrove, J., Weiler, M. and Soulsby, C.: Measuring and Modeling Stable Isotopes of Mobile and Bulk Soil Water, Vadose Zone J., 17(1), 2018.


Troy, T. J., Pavao-Zuckerman, M. and Evans, T. P.: Debates-Perspectives on socio-hydrology: Socio-hydrologic modeling: Tradeoffs, hypothesis testing, and validation, Water Resour. Res., 51(6), 4806–4814, 2015.

Wei, Z., Yoshimura, K., Wang, L., Miralles, D. G., Jasechko, S. and Lee, X.: Revisiting the contribution of transpiration to global terrestrial evapotranspiration, Geophys. Res. Lett., 44(6), 2792–2801, doi:10.1002/2016GL072235, 2017.

Zhang, M., Liu, N., Harper, R., Li, Q., Liu, K., Wei, X., Ning, D., Hou, Y. and Liu, S.: A global review on hydrological responses to forest change across multiple spatial scales: Importance of scale, climate, forest type and hydrological regime, J. Hydrol., 546, 44–59, 2017.


Interactive comment on Geosci. Model Dev. Discuss., <https://doi.org/10.5194/gmd-2018-25>, 2018.

C21




**Fig. 1.** New Figure S1.

C22



**Fig. 2.** New Figure 1

C23



**Fig. 3.** New Figure 8

C24