SKRIPS v1.0: A regional coupled ocean–atmosphere modeling framework (MITgcm–WRF) using ESMF/NUOPC, description and preliminary results for the Red Sea

Rui Sun¹, Aneesh C. Subramanian¹, Arthur J. Miller¹, Matthew R. Mazloff¹, Ibrahim Hoteit², and Bruce D. Cornuelle¹

¹Scripps Institution of Oceanography, La Jolla, California, USA

²Physical Sciences and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia

Correspondence: Rui Sun (rus043@ucsd.edu); Aneesh Subramanian (acsubram@ucsd.edu)

Abstract. A new regional coupled ocean–atmosphere model is developed and its implementation is presented in this paper. The coupled model is based on two open-source community model components: (1) MITgcm ocean model; (2) Weather Research and Forecasting (WRF) atmosphere model. The coupling between these components is performed using ESMF (Earth System Modeling Framework) and implemented according to National United Operational Prediction Capability (NUOPC)

- 5 protocols. The coupled model is named the Scripps-KAUST Regional Integrated Prediction System (SKRIPS). The SKRIPS is demonstrated with a real-world example by simulating a 30-day period including a series of extreme heat events occurring on the eastern shore of the Red Sea region in June 2012. The results obtained by using the coupled model, along with those in forced stand-alone oceanic or atmospheric simulations, are compared with observational data and reanalysis products. We show that the coupled model is capable of performing coupled ocean-atmosphere simulation, although all configurations of
- 10 coupled and uncoupled models have good skill in modeling the heat events. In addition, a scalability test is performed to investigate the parallelization of the coupled model. The results indicate that the coupled model code scales well and the ESMF/NUOPC coupler accounts for less than 5% of the total computational resources in the Red Sea test case. The coupled model and documentation are available at https://library.ucsd.edu/dc/collection/bb1847661c, and the source code is maintained at https://github.com/iurnus/scripps_kaust_model.

15 1 Introduction

20

Accurate and efficient forecasting of oceanic and atmospheric circulation is essential for a wide variety of high-impact societal needs, including extreme weather and climate events (Kharin and Zwiers, 2000; Chen et al., 2007), environmental protection and coastal management (Warner et al., 2010), management of fisheries (Roessig et al., 2004), marine conservation (Harley et al., 2006), water resources (Fowler and Ekström, 2009), and renewable energy (Barbariol et al., 2013). Effective forecasting relies on high model fidelity and accurate initialization of the models with the observed state of the coupled ocean–atmosphere system. Although global coupled models are now being implemented with increased resolution, higher-resolution regional

coupled models, if properly driven by the boundary conditions, can provide an affordable way to study air-sea feedback for frontal-scale processes.

A number of regional coupled ocean–atmosphere models have been developed for various goals in the past decades. An early example of building a regional coupled model for realistic simulations focused on accurate weather forecasting in the Baltic

- 5 Sea (Gustafsson et al., 1998; Hagedorn et al., 2000; Doscher et al., 2002), and showed that the coupled model improved the SST (Sea Surface Temperature) and atmospheric circulation forecast. Enhanced numerical stability in the coupled simulation was also observed. These early attempts were followed by other practitioners in ocean-basin-scale climate simulations (e.g. Huang et al., 2004; Aldrian et al., 2005; Xie et al., 2007; Seo et al., 2007; Somot et al., 2008; Fang et al., 2010; Boé et al., 2011; Zou and Zhou, 2012; Gualdi et al., 2013; Van Pham et al., 2014; Chen and Curcic, 2016; Seo, 2017). For example, Huang
- 10 et al. (2004) implemented a regional coupled model to study three major important patterns contributing to the variability and predictability of the Atlantic climate. The study suggested that these patterns originate from air-sea coupling within the Atlantic Ocean or by the oceanic response to atmospheric intrinsic variability. Seo et al. (2007) studied the nature of ocean-atmosphere feedbacks in the presence of oceanic mesoscale eddy fields in the eastern Pacific Ocean sector. The evolving SST fronts were shown to drive an unambiguous response of the atmospheric boundary layer in the coupled model, and lead to model anomalies
- 15 of wind stress curl, wind stress divergence, surface heat flux, and precipitation that resemble observations. This study helped substantiate the importance of ocean-atmosphere feedbacks involving oceanic mesoscale variability features.

In addition to basin-scale climate simulations, regional coupled models are also used to study weather extremes. For example, the COAMPS (Coupled Ocean/Atmosphere Mesoscale Prediction System) was applied to investigate idealized tropical cyclone events (Hodur, 1997). This work was then followed by other realistic extreme weather studies. Another example is the

- 20 investigation of extreme bora wind events in the Adriatic Sea using different regional coupled models (Loglisci et al., 2004; Pullen et al., 2006; Ricchi et al., 2016). The coupled simulation results demonstrated improvements in describing the air-sea interaction processes by taking into account oceanic surface heat fluxes and wind-driven ocean surface wave effects (Loglisci et al., 2004; Ricchi et al., 2016). It was also found in model simulations that SST after bora wind events had a stabilizing effect on the atmosphere, reducing the atmospheric boundary layer mixing and yielding stronger near-surface wind (Pullen et al., 2016).
- 25 2006). Regional coupled models were also used for studying the forecasts of hurricanes, including hurricane path, hurricane intensity, SST variation, and wind speed (Bender and Ginis, 2000; Chen et al., 2007; Warner et al., 2010).

Regional coupled modeling systems also play important roles in studying the effect of surface variables (e.g., surface evaporation, precipitation, surface roughness) in the coupling processes of oceans or lakes. One example is the study conducted by Powers and Stoelinga (2000), who developed a coupled model and investigated the passage of atmospheric fronts over the

30 Lake Erie region. Sensitivity analysis was performed to demonstrate that parameterization of lake surface roughness in the atmosphere model can improve the calculation of wind stress and heat flux. Another example is the investigation of Caspian Sea levels by Turuncoglu et al. (2013), who compared a regional coupled model with uncoupled models and demonstrated the improvement of the coupled model in capturing the response of Caspian Sea levels to climate variability.

In the past ten years, many regional coupled models have been developed using modern model toolkits (Zou and Zhou, 2012; Turuncoglu et al., 2013; Turuncoglu, 2019) and include waves (Warner et al., 2010; Chen and Curcic, 2016), sediment

transport (Warner et al., 2010), sea ice (Van Pham et al., 2014), and chemistry packages (He et al., 2015). However, this work was motivated by the need for a coupled regional ocean–atmosphere model implemented using an efficient coupling framework and with compatible state estimation capabilities in both ocean and atmosphere. The goal of this work is to (1) introduce the design of the newly developed regional coupled ocean–atmosphere modeling system, (2) describe the implementation

- 5 of the modern coupling framework, (3) validate the coupled model using a real-world example, and (4) demonstrate and discuss the parallelization of the coupled model. In the coupled system, the oceanic model component is the MIT general circulation model (MITgcm) (Marshall et al., 1997) and the atmospheric model component is the Weather Research and Forecasting (WRF) model (Skamarock et al., 2019). To couple the model components in the present work, the Earth System Modeling Framework (ESMF) (Hill et al., 2004) is used because of its advantages in conservative re-gridding capability,
- 10 calendar management, logging and error handling, and parallel communications. The National United Operational Prediction Capability (NUOPC) layer in ESMF (Sitz et al., 2017) is also used between model components and ESMF. Using the NUOPC layer can simplify the implementation of component synchronization, execution, and other common tasks in the coupling. The innovations in our work are: (1) we use ESMF/NUOPC, which is a community-supported, computationally-efficient coupling software for earth system models, and (2) we use MITgcm together with WRF, both of which work with the Data Assimilation
- 15 Research Testbed (DART) (Anderson and Collins, 2007; Hoteit et al., 2013). The resulting coupled model is being developed for coupled data assimilation and subseasonal to seasonal (S2S) forecasting. By coupling WRF and MITgcm for the first time with ESMF, we can provide an alternative regional coupled model resource to a wider community of users. These atmospheric and oceanic model components have an active and well-supported user-base.
- After implementing the new coupled model, we demonstrate it on a series of heat events that occurred on the eastern shore of the Red Sea region in June 2012. The simulated surface variables of the Red Sea (e.g., sea surface temperature, 2-m temperature, and surface heat fluxes) are examined and validated against available observational data and reanalysis products. To demonstrate that the coupled model can perform coupled ocean–atmosphere simulations the results are compared with those obtained using stand-alone oceanic or atmospheric models. This paper focuses on the technical aspects of the SKRIPS, and is not a full investigation of the importance of coupling for these extreme events. In addition, a scalability test of the coupled model is performed to investigate its parallel capability.
 - The rest of this paper is organized as follows. The description of the individual modeling components and the design of the coupled modeling system are detailed in Section 2. Section 3 introduces the design of validation experiment and the validation data. Section 4 discusses the preliminary results in the validation test. Section 5 details the parallelization test of the coupled model. The last section concludes the paper and presents an outlook for future work.

30 2 Model Description

The newly developed regional coupled modeling system is introduced in this section. The general design of the coupled model, descriptions of individual components, and ESMF/NUOPC coupling framework are presented below.

2.1 General Design

The schematic description of the coupled model is shown in Fig. 1. The coupled model is comprised of five components: the oceanic component MITgcm, the atmospheric component WRF, the MITgcm–ESMF and WRF–ESMF interfaces, and the ESMF coupler. They are to be detailed in the following sections.

The coupler component runs in both directions: (1) from WRF to MITgcm, and (2) from MITgcm to WRF. From WRF to MITgcm, the coupler collects the atmospheric surface variables (i.e., radiative flux, turbulent heat flux, wind velocity, precipitation, evaporation) from WRF and updates the surface forcing (i.e., net surface heat flux, wind stress, freshwater flux) to drive MITgcm. From MITgcm to WRF, the coupler collects oceanic surface variables (i.e., SST and ocean surface velocity) from MITgcm and updates them in WRF as the bottom boundary condition. Re-gridding the data from either model component

10 is performed by the coupler, in which various coupling intervals and schemes can be specified by ESMF (Hill et al., 2004).



Figure 1. The schematic description of the coupled ocean–atmosphere model. The yellow block is the ESMF/NUOPC coupler; the red blocks are the implemented MITgcm–ESMF and WRF–ESMF interfaces; the white blocks are the oceanic and atmospheric components. From WRF to MITgcm, the coupler collects the atmospheric surface variables (i.e., radiative flux, turbulent heat flux, wind velocity, precipitation, evaporation) and updates the surface forcing (i.e., net surface heat flux, wind stress, freshwater flux) to drive MITgcm. From MITgcm to WRF, the coupler collects oceanic surface variables (i.e., SST and ocean surface velocity) and updates them in WRF as the bottom boundary condition.

2.2 The Oceanic Component (MITgcm)

The MITgcm (Marshall et al., 1997) is a 3-D, finite-volume, general circulation model used by a broad community of researchers for a wide range of applications at various spatial and temporal scales. The model code and documentation, which are under continuous development, are available on the MITgcm webpage (http://mitgcm.org/). The 'Checkpoint 66h' (June 2017) version of MITgcm is used in the present work

5 2017) version of MITgcm is used in the present work.

The MITgcm is designed to run on high-performance computing (HPC) platforms and can run in non-hydrostatic and hydrostatic modes. It integrates the primitive (Navier–Stokes) equations, under the Boussinesq approximation, using finite volume method on a staggered 'Arakawa C-grid'. The MITgcm uses modern physical parameterizations for subgrid-scale horizontal and vertical mixing and tracer properties. The code configuration includes build-time C pre-processor (CPP) options and

10 run-time switches, which allow for great computational modularity in MITgcm to study a variety of oceanic phenomena (Evangelinos and Hill, 2007).

To implement the MITgcm–ESMF interface, we separate the MITgcm main program into three subroutines that handle initialization, running, and finalization, shown in Fig. 2(a). These subroutines are used by the ESMF/NUOPC coupler that controls the oceanic component in the coupled run. The surface boundary fields are exchanged online¹ via the MITgcm–

- 15 ESMF interface during the simulation. The MITgcm oceanic surface variables are the export boundary fields; the atmospheric surface variables are the import boundary fields (see Fig. 2(b)). These boundary fields are registered in the coupler following NUOPC protocols with timestamps² for the coupling. In addition, MITgcm grid information is provided to the coupler in the initialization subroutine for online re-gridding of the exchanged boundary fields. To carry out the coupled simulation on HPC clusters, the MITgcm–ESMF interface runs in parallel via MPI communications. The implementation of the present MITgcm–
- 20 ESMF interface is based on the baseline MITgcm–ESMF interface (Hill, 2005), but updated for compatibility with the modern version of ESMF/NUOPC. We also modify the baseline interface to receive atmospheric surface variables and send oceanic surface variables.

2.3 The Atmospheric Component (WRF)

The Weather Research and Forecasting (WRF) Model (Skamarock et al., 2019) is developed by the NCAR/MMM (Mesoscale and Microscale Meteorology Division). It is a 3-D, finite-difference atmospheric model with a variety of physical parameterizations of sub-grid scale processes for predicting a broad spectrum of applications. WRF is used extensively for operational forecasts (http://www.wrf-model.org/plots/wrfrealtime.php) as well as realistic and idealized dynamical studies. The WRF code and documentation are under continuous development on Github (https://github.com/wrf-model/WRF).

In the present work, the Advanced Research WRF dynamic version (WRF-ARW, version 3.9.1.1, https://github.com/NCAR/ 30 WRFV3/releases/tag/V3.9.1.1) is used. It solves the compressible Euler non-hydrostatic equations, and also includes a run-time

¹In this manuscript, 'online' means the manipulations are performed via subroutine calls during the execution of the simulations; 'offline' means the manipulations are performed when the simulations are not executing.

²In ESMF, 'timestamp' is a sequence of numbers, usually based on the time, to identify ESMF fields. Only the ESMF fields having the correct timestamp will be transferred in the coupling.



Figure 2. The general code structure and run sequence of the coupled ocean–atmosphere model. In panel (a), the black block is the *application driver*; the red block is the *parent gridded component* called by the *application driver*; the green/brown blocks are the *child gridded/coupler components* called by the *parent gridded component*. Panel (b) and (c) shows the sequential and concurrent mode implemented in SKRIPS, respectively. PETs (Persistent Execution Threads) are single processing units (e.g., CPU or GPU cores) defined by ESMF. OCN, ATM, and CON denote oceanic component, atmospheric component and connector component, respectively. The blocks under PETs are the CPU cores in the simulation; the small blocks under OCN or ATM are the small sub-domains in each core; the block under CON is the coupler. The red arrows indicate that the model components are sending data to the connector and the yellow arrows indicate that the model components are coupling time steps.

hydrostatic option. The WRF-ARW uses a terrain-following hydrostatic pressure coordinate system in the vertical direction and utilizes the 'Arakawa C-grid'. WRF incorporates various physical processes including microphysics, cumulus parameterization, planetary boundary layer, surface layer, land surface, and longwave/shortwave radiation, with several options available for each process.

- 5 Similar to the implementations in MITgcm, WRF is also separated into initialization, run, and finalization subroutines to enable the WRF–ESMF interface to control the atmosphere model during the coupled simulation, shown in Fig. 2(a). The implementation of the present WRF–ESMF interface is based on the prototype interface (Henderson and Michalakes, 2005). In the present work, the prototype WRF–ESMF interface is updated to modern versions of WRF-ARW and ESMF, based on the NUOPC layer. This prototype interface is also expanded to interact with the ESMF/NUOPC coupler to receive the oceanic
- 10 surface variables and send the atmospheric surface variables. The surface boundary condition fields are registered in the coupler following the NUOPC protocols with timestamps. The WRF grid information is also provided for online re-gridding by ESMF. To carry out the coupled simulation on HPC clusters, the WRF–ESMF interface also runs in parallel via MPI communications.

2.4 ESMF/NUOPC Coupler

The coupler is implemented using ESMF version 7.0.0. The ESMF is selected because of its high-performance and flexibility for building and coupling weather, climate, and related Earth science applications (Collins et al., 2005; Turuncoglu et al., 2013; Chen and Curcic, 2016; Turuncoglu and Sannino, 2017). It has a superstructure for representing the model and coupler

5 components and an infrastructure of commonly used utilities, including conservative grid remapping, time management, error handling, and data communications.

The general code structure of the coupler is shown in Fig. 2. To build the ESMF/NUOPC driver, a main program is implemented to control an ESMF parent component, which controls the child components. In the present work, three child components are implemented: (1) the oceanic component; (2) the atmospheric component; and (3) the ESMF coupler. The

10 coupler is used here because it performs the two-way interpolation and data transfer (Hill et al., 2004). In ESMF, the model components can be run in parallel as a group of Persistent Execution Threads (PETs), which are single processing units (e.g., CPU or GPU cores) defined by ESMF. In the present work, the PETs are created according to the grid decomposition, and each PET is associated with an MPI process.

The ESMF allows the PETs to run in sequential mode, concurrent mode, or mixed mode (for more than three components).

15 We implemented both sequential and concurrent modes in SKRIPS, shown in Fig. 2(b) and 2(c). In sequential mode, a set of ESMF gridded/coupler components are run in sequence on the same set of PETs. At each coupling time step, the oceanic component is executed when the atmospheric component is completed and vice versa. On the other hand, in concurrent mode, the gridded components are created and run on mutually exclusive sets of PETs. If one component finishes earlier than the other, its PETs are idle and have to wait for the other component, shown in Fig. 2(c). However the PETs can be optimally 20 distributed by the users to best achieve load balance. In this work, all simulations are run in sequential mode.

In ESMF, the gridded components are used to represent models, and coupler components are used to connect these models. The interfaces and data structures in ESMF have few constraints, providing the flexibility to be adapted to many modeling systems. However, the flexibility of the gridded components can limit the interoperability across different modeling systems. To address this issue, the NUOPC layer is developed to provide the coupling conventions and the generic representation

- 25 of the model components (e.g. drivers, models, connectors, mediators). The NUOPC layer in the present coupled model is implemented according to consortium documentations (Hill et al., 2004; Theurich et al., 2016), and the oceanic/atmospheric component each has:
 - 1. Prescribed variables for NUOPC to link the component;
 - 2. The entry point for registration of the component;
- 30 3. An *InitializePhaseMap* which describes a sequence of standard initialization phases, including documenting the fields that a component can provide, checking and mapping the fields to each other, and initializing the fields that will be used;
 - 4. A *RunPhaseMap* that checks the incoming clock of the driver, examines the timestamps of incoming fields, and runs the component;

- 5. Timestamps on exported fields consistent with the internal clock of the component;
- 6. The *finalization method* to clean up all allocations.

The subroutines that handle initialization, running, and finalization in MITgcm and WRF are included in the *InitializePhaseMap*, *RunPhaseMap*, and *finalization method* in the NUOPC layer, respectively.

5 3 Experiment Design and Observational Datasets

We simulate a series of heat events in the Red Sea region, with a focus on validating and assessing the technical aspects of the coupled model. There is a desire for improved and extended forecasts in this region, and future work will investigate whether a coupled framework can advance this goal. The extreme heat events are chosen as a test case due to their societal importance. While these events and the analysis here may not maximize the value of coupled forecasting, these real-world events are

- 10 adequate to demonstrate the performance and physical realism of the coupled model code implementation. The simulation of the Red Sea extends from 0000 UTC 01 June 2012 to 0000 UTC 01 July 2012. We select this month because of the record-high surface air temperature observed in the Makkah region, located 70 km inland from the eastern shore of the Red Sea (Abdou, 2014).
- The computational domain and bathymetry are shown in Fig. 3. The model domain is centered at 20° N and 40° E, and the
 bathymetry is from the 2-minute Gridded Global Relief Data (ETOPO2) (National Geophysical Data Center, 2006). WRF is implemented using a horizontal grid of 256 × 256 points and grid spacing of 0.08°. The cylindrical equidistant map (latitude-longitude) projection is used. There are 40 terrain-following vertical levels, more closely spaced in the atmospheric boundary layer. The time step for atmosphere simulation is 30 seconds, which is to avoid violating the CFL condition. The Morrison 2-moment scheme (Morrison et al., 2009) is used to resolve the microphysics. The updated version of the Kain–Fritsch
- 20 convection scheme (Kain, 2004) is used with the modifications to include the updraft formulation, downdraft formulation, and closure assumption. The Yonsei University (YSU) scheme (Hong et al., 2006) is used for the planetary boundary layer (PBL), and the Rapid Radiation Transfer Model for GCMs (RRTMG; Iacono et al. (2008)) is used for longwave and shortwave radiation transfer through the atmosphere. The Rapid Update Cycle (RUC) land surface model is used for the land surface processes (Benjamin et al., 2004). The MITgcm uses the same horizontal grid spacing as WRF, with 40 vertical z-levels that
- 25 are more closely spaced near the surface. The time step of the ocean model is 120 seconds. The horizontal sub-grid mixing is parameterized using nonlinear Smagorinsky viscosities, and the K-profile parameterization (KPP) (Large et al., 1994) is used for vertical mixing processes.

During coupled execution the ocean model sends SST and ocean surface velocity to the coupler, and they are used directly as the boundary conditions in the atmosphere model. The atmosphere model sends the surface fields to the coupler, including

(1) surface radiative flux (i.e., longwave/shortwave radiation), (2) surface turbulent heat flux (i.e., latent/sensible heat), (3)
 10-m wind speed, (4) precipitation, (5) evaporation. The ocean model uses the atmospheric surface variables to compute the surface forcing, including (1) total net surface heat flux, (2) surface wind stress, (3) freshwater flux. The total net surface heat

flux is computed by adding latent heat flux, sensible heat flux, shortwave radiation flux, and longwave radiation flux. The surface wind stress is computed by using the 10-m wind speed (Large and Yeager, 2004). The freshwater flux is the difference between precipitation and evaporation. The latent and sensible heat fluxes are computed by using COARE 3.0 bulk algorithm in WRF (Fairall et al., 2003). In the coupled code, different bulk formulae in WRF or MITgcm can also be used.



Figure 3. The WRF topography and MITgcm bathymetry in the simulations. Three major cities near the eastern shore of the Red Sea are highlighted. The Hijaz Mountains and Ethiopian Highlands are also highlighted.

- 5 To validate the coupled model, the following sets of simulations using different surface forcings are performed according to the validation tests in the literature (Warner et al., 2010; Turuncoglu et al., 2013; Ricchi et al., 2016):
 - 1. Run CPL: a two-way coupled MITgcm–WRF simulation. The coupling interval is 20 minutes to capture the diurnal cycle (Seo et al., 2014). This run tests the implementation of the two-way coupled ocean–atmosphere model.
 - 2. Run ATM.STA: a stand-alone WRF simulation with its initial SST kept constant throughout the simulation. This run allows assessment of the WRF model behavior with realistic, but persistent SST. This case serves as a benchmark to highlight the difference between coupled and uncoupled runs, and also to demonstrate the impact of evolving SST.

10

- 3. Run ATM.DYN: a stand-alone WRF simulation with a varying, prescribed SST based on HYCOM/NCODA reanalysis data. This allows assessing the WRF model behavior with updated SST and is used to validate the coupled model. It is noted that in practice an accurately evolving SST would not be available for forecasting.
- 4. Run OCN.DYN: a stand-alone MITgcm simulation forced by the ERA5 reanalysis data. The bulk formula in MITgcm is used to derive the turbulent heat fluxes. This run assesses the MITgcm model behavior with prescribed lower-resolution atmospheric surface forcing, and is also used to validate the coupled model.
 - 9

The ocean model uses the HYCOM/NCODA $1/12^{\circ}$ global reanalysis data as initial and boundary conditions for ocean temperature, salinity, and horizontal velocities (https://www.hycom.org/dataserver/gofs-3pt1/reanalysis). The boundary conditions for the ocean are updated on a 3-hourly basis and linearly interpolated between two simulation time steps. A sponge layer is applied at the lateral boundaries, with a thickness of 3 grid cells. The inner and outer boundary relaxation timescales

- 5 of the sponge layer are 10 and 0.5 days, respectively. In CPL, ATM.STA, and ATM.DYN, we use the same initial condition and lateral boundary condition for the atmosphere. The atmosphere is initialized using the ECMWF ERA5 reanalysis data, which have a grid resolution of approximately 30 km (Hersbach, 2016). The same data also provide the boundary conditions for air temperature, wind speed, and air humidity every 3 hours. The atmosphere boundary conditions are also linearly interpolated between two simulation time steps. The lateral boundary values are specified in WRF in the 'specified' zone, and the
- 10 'relaxation' zone is used to nudge the solution from the domain toward the boundary condition value. Here we use the default width of one point for the specific zone and four points for the relaxation zone. The top of the atmosphere is at the 50 hPa pressure level. In ATM.STA, the SST from HYCOM/NCODA at the initial time is used as a constant SST. The time-varying SST in ATM.DYN is also generated using HYCOM/NCODA data. We select HYCOM/NCODA data because the ocean model initial condition and boundary conditions are generated using it. For OCN.DYN we select ERA5 data for the atmospheric state
- 15 because it also provides the atmospheric initial and boundary conditions in CPL. The initial conditions, boundary conditions, and forcing terms of all simulations are summarized in Table 1.

initial and boundary conditions		ocean surface	atmospheric forcings	
		conditions		
CPL	ERA5 (atmosphere)	from MITgom	from WRF	
	HYCOM/NCODA (ocean)	from wirrgem		
ATM.STA	ED A 5	HYCOM/NCODA	N.A.	
	EKAJ	initial condition kept constant		
ATM.DYN	ERA5	HYCOM/NCODA	N.A.	
		updated every 3 hours		
OCN.DYN	HYCOM/NCODA	N.A.	ERA5	

Table 1. The initial conditions, boundary conditions, and forcing terms used in the simulations.

The validation of the coupled model focuses on temperature, heat flux, and surface wind. Our aim is to validate the coupled model and show that the heat and momentum fluxes simulated by the coupled model are comparable to the observations or the reanalysis data. The simulated 2-m air temperature (T2) fields are validated using ERA5. In addition, the simulated T2 for

20

three major cities near the eastern shore of the Red Sea are validated using ERA5 and ground observations from the NOAA National Climate Data Center (NCDC climate data online at https://www.ncdc.noaa.gov/cdo-web/). The simulated SST data are validated against the OSTIA (Operational Sea Surface Temperature and Sea Ice Analysis) system in GHRSST (Group for High Resolution Sea Surface Temperature) (Donlon et al., 2012; Martin et al., 2012). In addition, the simulated SST

fields are validated against HYCOM/NCODA data. Since the simulations are initialized using HYCOM/NCODA data, this aims to show the increase of the differences. Surface heat fluxes (e.g., turbulent heat flux and radiative flux), which drive the oceanic component in the coupled model, are validated using MERRA-2 (Modern-Era Retrospective analysis for Research and Applications, version 2) data (Gelaro et al., 2017). We use MERRA-2 data because (1) it is an independent reanalysis

- 5 data compared to the initial and boundary conditions used in the simulations, and (2) it also provides $0.625^{\circ} \times 0.5^{\circ}$ (lon × lat) resolution reanalysis fields of turbulent heat fluxes (THF). The 10-m wind speed is also compared with MERRA-2 data to validate the momentum flux in the coupled code. The validation of the freshwater flux is shown in the Appendix because (1) the evaporation is proportional to the latent heat in the model and (2) the precipitation is zero in the cities near the coast in Fig. 3. The validation data are summarized in Table 2.
- 10 When comparing T2 with NCDC ground observations, the simulation results and ERA5 are interpolated to the NCDC stations. When interpolating to NCDC stations near the coast, only the data saved on land points are used³. The maximum/minimum T2 every 24 hours from the simulations and ERA5 are compared to the observed daily maximum/minimum T2. On the other hand, when comparing the simulation results with the analysis or reanalysis data (HYCOM, GHRSST, ERA5, and MERRA-2), we interpolate these data onto the model grid to achieve a uniform spatial scale (Maksyutov et al., 2008;
- 15 Torma et al., 2011).

Table 2. The observational data and reanalysis data used to validate the simulation results.

variable	validation data		
sea surface temperature (SST)	GHRSST and HYCOM/NCODA		
2-m air temperature (T2)	ERA5 and NCDC climate data		
turbulent heat fluxes	MERRA-2		
radiative fluxes	MERRA-2		
10-m wind speed	MERRA-2		

4 Results and Discussions

The Red Sea is an elongated basin covering the area between 12-30°N and 32-43°E. The basin is 2250 km long, extending from the Suez and Aqaba gulfs in the north to the strait of Bab el-Mandeb in the south, which connects the Red Sea and the Indian Ocean. In this section, the simulation results obtained by using different model configurations are presented to show that SKRIPS is capable of performing coupled ocean–atmosphere simulations. The T2 from CPL, ATM.STA, and ATM.DYN are compared with the validation data to evaluate the atmospheric component of SKRIPS; the SST obtained from CPL and

²⁰

 $^{^{3}}$ In ATM.STA, ATM.DYN, and CPL, we use land-sea mask=1 as land points because the land-sea mask is either 0 (sea) or 1 (land) in WRF. In ERA5, we use land-sea mask > 0.9 as land points because the land-sea mask is a fractional value between 0 (sea) to 1 (land).

OCN.DYN are compared to validate the atmospheric component of SKRIPS; the surface heat fluxes and 10-m wind are used to assess the coupled system.

4.1 2-m Air Temperature

We begin our analysis by examining the simulated T2 from the model experiments, aiming to validate the atmospheric com-

- 5 ponent of SKRIPS. Since the record-high temperature is observed in the Makkah region on June 2nd, the simulation results on June 2nd (36 or 48 hours after the initialization) are shown in Fig. 4. The ERA5 data, and the difference between CPL and ERA5 are also shown in Fig. 4. It can be seen in Fig. 4(I) that CPL captures the T2 patterns in the Red Sea region on June 2nd compared with ERA5 in Fig. 4(II). Since the ERA5 T2 data are in good agreement with the NCDC ground observation data in the Red Sea region (detailed comparisons of all stations are not shown), we use ERA5 data to validate the simulation results. The
- 10 difference between CPL and ERA5 is shown in Fig. 4(III). The ATM.STA and ATM.DYN results are close to the CPL results and thus are not shown, but their differences with respect to ERA5 are shown in Fig. 4(IV) and 4(V), respectively. Fig. 4(VI) to 4(X) show the nighttime results after 48 hours. It can be seen in Fig. 4 that all simulations reproduce the T2 patterns over the Red Sea region reasonably well compared with ERA5. The mean T2 biases and root mean square errors (RMSEs) over the sea are shown in Table 3. The biases of the T2 are comparable with those reported in other benchmark WRF simulations (Xu
- 15 et al., 2009; Zhang et al., 2013a; Imran et al., 2018).

The simulation results on June 10th and 24th are shown in Fig. 5 to validate the coupled model over longer periods of time. It can be seen in Fig. 5 that the T2 patterns in CPL are generally consistent with ERA5. The differences between the simulations (CPL, ATM.STA and ATM.DYN) and ERA5 shows that the T2 on land are consistent for all three simulations. However, the T2 over the sea in CPL has smaller mean biases and RMSEs compared with ATM.STA, also shown in Table 3. Although the

20 difference in T2 is very small compared with the mean T2 (31.92 °C), the improvement of the coupled run on the 24th (1.02 °C) is comparable to the standard deviation of T2 (1.64 °C). The T2 over the water in CPL is closer to ERA5 because MITgcm in the coupled model provides a dynamic SST which influences T2. On the other hand, when comparing CPL with ATM.DYN, the mean difference is smaller (10th: +0.04 °C; 24th: -0.62 °C). This shows CPL is comparable to ATM.DYN which is driven by an updated warming SST.

Table 3. The biases and RMSEs of T2 simulated in all simulations in comparison with ERA5 data.

	after 36 hours	after 48 hours	after 9.5 days	after 23.5 days
Run CPL	bias: -1.36; RMSE: 1.91	bias: -0.82; RMSE: 1.19	bias: -1.24; RMSE: 1.96	bias: -0.81; RMSE: 1.80
Run ATM.STA	bias: -1.48; RMSE: 2.01	bias: -0.92; RMSE: 1.27	bias: -1.56; RMSE: 2.27	bias: -1.83; RMSE: 2.59
Run ATM.DYN	bias: -1.36; RMSE: 1.90	bias: -0.84; RMSE: 1.28	bias: -1.20; RMSE: 1.93	bias: -1.43; RMSE: 2.14

The mean biases and RMSEs of T2 over the Red Sea during the 30-day simulation are shown in Fig. 6 to demonstrate the evolution of simulation errors. It can be seen that ATM.STA can still capture the T2 patterns in the first week, but it under-



Figure 4. The 2-m air temperature as obtained from the CPL, the ERA5 data, and their difference (CPL–ERA5). The differences between ATM.STA and ATM.DYN with ERA5 (i.e., ATM.STA–ERA5, ATM.DYN–ERA5) are also presented. The simulation initial time is 0000 UTC Jun 01 2012 for both snapshots. Two snapshots are selected: (1) 1200 UTC Jun 02 2012 (36 hours from initial time); (2) 0000 UTC Jun 03 2012 (48 hours from initial time). The results on Jun 02 are presented because the record-high temperature is observed in the Makkah region.

predicts T2 by about 2 °C after 20 days because it has no SST evolution. On the other hand, CPL has smaller bias (-0.60 °C) and RMSE (1.28 °C) compared with those in ATM.STA (bias: -1.19 °C; RMSE: 1.71 °C) during the 30-day simulation as the SST evolution is considered. The ATM.DYN case also has smaller error than ATM.STA and its error is comparable with that in CPL (bias: -0.72 °C; RMSE: 1.31 °C), indicating that the skill of the coupled model is comparable to the stand-alone atmosphere model driven by 3-hourly reanalysis SST. The differences of the mean biases and RMSEs between model outputs and ERA5 data are also plotted in Fig. 6. It can be seen that CPL has smaller error than ATM.STA throughout the simulation. The bias and RMSE between CPL and ATM.DYN are within about 0.5 °C. This shows the capability of the coupled model for performing realistic regional coupled ocean–atmosphere simulations.

5

To validate the diurnal T2 variation of the coupled model in Fig. 4, the time series of T2 in three major cities as simulated in

10 CPL and ATM.STA are plotted in Fig. 7, starting from June 1st. The ERA5 data and the daily observed high/low temperature data from NOAA NCDC are also plotted for validation. Both coupled and uncoupled simulations generally captured the four major heat events (i.e., June 2nd, 10th, 17th, and 24th) and the T2 variations during the 30-day simulation. For the daily high T2, the RMSE in all simulations are close (CPL: 2.09 °C; ATM.STA: 2.16 °C; ATM.DYN: 2.06 °C), and the error does not increase in the 30-day simulation. For the daily low T2, before June 20th (lead time < 19 days), all simulations have consistent</p>

15 RMSEs compared with ground observation (CPL: 4.23 °C; ATM.STA: 4.39 °C; ATM.DYN: 4.01 °C). In Jeddah and Yanbu,



Figure 5. The T2 obtained in CPL, the T2 in ERA5, and their difference (CPL–ERA5). The difference between ATM.STA and ATM.DYN with ERA5 data (i.e., ATM.STA–ERA5, ATM.DYN–ERA5) are also presented. The simulation initial time is 0000 UTC Jun 01 2012 for both snapshots. Two snapshots are selected: (1) 1200 UTC Jun 10 2012 (9.5 days from initial time); (2) 1200 UTC Jun 24 2012 (23.5 days from initial time).



Figure 6. The bias and RMSE between the T2 obtained by the simulations (i.e., ATM.STA, ATM.CPL, and CPL) in comparison with ERA5 data. Only the errors over the Red Sea are considered. The differences between the simulation errors from CPL and stand-alone WRF simulations are presented below the mean bias and the RMSE. The initial time is 0000 UTC Jun 01 2012 for all simulations.

CPL has better captured the daily low T2 after June 20^{th} (Jeddah: 3.95 °C; Yanbu: 3.77 °C) than ATM.STA (Jeddah: 4.98 °C; Yanbu: 4.29 °C) by about 1 °C and 0.5 °C, respectively. However, the T2 difference of Mecca, which is located 70-km from the sea, is negligible (0.05 °C) between all simulations throughout the simulation.



Figure 7. Temporal variation the 2-m air temperature at three major cities near the eastern shore of Red Sea (Jeddah, Mecca, Yanbu) as resulting from CPL, ATM.STA and ATM.DYN. The temperature data are compared with the time series in ERA5 and daily high/low temperature in the NOAA national data center dataset. Note that some gaps exist in the NCDC ground observation dataset. Four representative heat events are highlighted in this figure.

4.2 Sea Surface Temperature

- ⁵ The simulated SST patterns obtained in the simulations are presented to demonstrate that the coupled model can capture the ocean surface state. The snapshots of SST obtained from CPL are shown in Fig. 8(I) and Fig. 8(VI). To validate the coupled model, the SST fields obtained in OCN.DYN are shown in Fig. 8(II) and 8(VII), and the GHRSST data are shown in Fig. 8(II) and 8(VIII). The SST obtained in the model at 0000 UTC (about 3 A.M. local time in the Red Sea region) is presented because the GHRSST is produced with nighttime SST data (Roberts-Jones et al., 2012). It can be seen that both CPL and OCN.DYN
- 10 are able to reproduce the SST patterns reasonably well in comparison with GHRSST for both snapshots. Though CPL uses higher-resolution surface forcing fields, the SST patterns obtained in both simulations are very similar after two days. On June

 24^{th} , the SST patterns are less similar, but both simulation results are still comparable with GHRSST (RMSE < 1° C). Both simulations under-estimate the SST in the northern Red Sea and over-estimate the SST in the central and southern Red Sea on June 24^{th} .



Figure 8. The SST in CPL, the SST in OCN.DYN, and GHRSST. The corresponding differences between the simulations and GHRSST are also plotted. Two snapshots of the model outputs are selected: (1) 0000 UTC Jun 02 2012; (2) 0000 UTC Jun 24 2012. The simulation initial time is 0000 UTC Jun 01 2012 for both snapshots.

- To quantitatively compare the errors in SST, the time histories of the SST in the simulations (i.e., OCN.DYN and CPL) and validation data (i.e., GHRSST and HYCOM/NCODA) are shown in Fig. 9. The mean biases and RMSEs between model outputs and validation data are also plotted. In Fig. 9(a) the snapshots of the simulated SST are compared with available HYCOM/NCODA data every 3 hours. In Fig. 9(b) the snapshots of SST outputs every 24 hours at 0000 UTC (about 3 A.M. local time in the Red Sea region) are compared with GHRSST. Compared with Fig. 9(b), the diurnal SST oscillation can be observed in Fig. 9(a) because the SST is plotted every 3 hours. Generally, OCN.DYN and CPL have a similar range of error compared to both validation datasets in the 30-day simulations. The simulation results are compared with HYCOM/NCODA data to show the increase of RMSE in Fig. 9(a). Compared with HYCOM/NCODA, the mean differences between CPL and OCN.DYN are small (CPL: 0.10 °C; OCN.DYN: 0.03 °C). The RMSE increases in the first week, but does not grow after that. On the other hand, when comparing with the GHRSST, the initial SST patterns in both runs are cooler by about 0.8 °C. This is because the models are initialized by HYCOM/NCODA, which has temperature in the topmost model level cooler
- 15 than the estimated foundation SST reported by GHRSST. After the first 10 days, the difference between GHRSST data and HYCOM/NCODA decreases, and likewise the difference between the simulation results and GHRSST also decreases. It should be noted that the SST simulated by CPL has smaller error (bias: -0.57 °C; RMSE: 0.69 °C) compared with OCN.DYN (bias:

-0.66 °C; RMSE: 0.76 °C) by about 0.1 °C when validated using GHRSST. This indicates the coupled model can adequately simulate the SST evolution compared with the uncoupled model forced by ERA5 reanalysis data.



Figure 9. The bias and RMSE between the SST from the simulations (i.e., OCN.DYN and CPL) in comparison with the validation data. Panel (a) shows the 3-hourly SST obtained in the simulations compared with 3-hourly HYCOM/NCODA data. Panel (b) shows the daily SST at 0000 UTC (about 3 A.M. local time in the Red Sea region) obtained in the simulations compared with GHRSST. Both simulations are initialized at 0000 UTC Jun 01 2012.

4.3 Surface Heat Fluxes

The atmospheric surface heat flux drives the oceanic component in the coupled model, hence we validate the heat fluxes in the 5 coupled model as compared to the stand-alone simulations. Both the turbulent heat fluxes and the net downward heat fluxes are compared to MERRA-2 and their differences are plotted. To validate the coupled ocean–atmosphere model, we only compare the heat fluxes over the sea.

The turbulent heat fluxes (THF; sum of latent and sensible heat fluxes) and their differences with the validation data are shown in Fig. 10 (the snapshots are shown in the Appendix). It can be seen that all simulations have similar mean THF over
the Red Sea compared with MERRA-2 (CPL: 119.4 W/m²; ATM.STA: 103.4 W/m²; ATM.DYN: 117.5 W/m²; MERRA-2: 115.6 W/m²). For the first two weeks, the mean THFs obtained in all simulations are overlapping in Fig. 10. This is because all simulations are initialized in the same way, and the SST in all simulations are similar in the first two weeks. After the second week, CPL has smaller error (bias: -1.8 W/m²; RMSE: 69.9 W/m²) compared with ATM.STA (bias: -25.7 W/m²; RMSE: 76.4 W/m²). This is because the SST is updated in CPL and is warmer compared with ATM.STA. When forced by a warmer

15 SST, the evaporation increases (also see the Appendix) and thus the latent heat fluxes increase. On the other hand, the THFs

in CPL are comparable with ATM.DYN during the 30-day run (bias: 1.9 W/m^2), showing that SKRIPS can capture the THFs over the Red Sea in the coupled simulation.



Figure 10. The turbulent heat fluxes out of the sea obtained in CPL, ATM.STA, and ATM.DYN in comparison with MERRA-2. The top panel shows the mean THF; the middle panel shows the mean bias; the bottom panel shows the RMSE. Only the hourly heat fluxes over the sea are shown .

The net downward heat fluxes (sum of THF and radiative flux) are shown in Fig. 11 (the snapshots are shown in the Appendix). Again, for the first two weeks, the heat fluxes obtained in ATM.STA, ATM.DYN and CPL are overlapping. This is because all simulations are initialized in the same way, and the SST in all simulations are similar in the first two weeks. After the second week, CPL has slightly smaller error (bias: 11.2 W/m²; RMSE: 84.4 W/m²) compared with the ATM.STA simulation (bias: 36.5 W/m²; RMSE: 94.3 W/m²). It should be noted that the mean bias and RMSE of the net downward heat fluxes can be as high as a few hundred W/m² or 40% compared with MERRA-2. This is because WRF over-estimated the shortwave radiation in the daytime (the snapshots are shown in the Appendix). However, the coupled model still captures the mean and standard deviation of the heat flux compared with MERRA-2 data (CPL mean: 110.6 W/m², standard deviation: 350.7 W/m²; MERRA-2 mean 104.7 W/m², standard deviation 342.3 W/m²). The over-estimation of shortwave radiation by the RRTMG scheme is also reported in other validation tests in the literature under all-sky conditions due to the uncertainty of cloud or aerosol (Zempila et al., 2016; Imran et al., 2018). Although the surface heat flux is over-estimated in the daytime, the SST over the Red Sea is not over-estimated (shown in Section 4.2).

15 4.4 10-m Wind Speed

To evaluate the simulation of the surface momentum by the coupled model, the 10-m wind speed patterns obtained from ATM.STA, ATM.DYN, and CPL are compared to the MERRA-2 reanalysis.



Figure 11. The total surface heat fluxes into the sea obtained in CPL, ATM.STA, and ATM.DYN in comparison with MERRA-2. The top panel shows the mean surface heat flux; the middle panel shows the mean bias; the bottom panel shows the RMSE. Only the heat fluxes over the sea are shown.

The simulated 10-m wind velocity fields are shown in Fig. 12. The RMSE of the wind speed between CPL and MERRA-2 data is 2.23 m/s when using the selected WRF physics schemes presented in Section 3. On June 2nd, high wind speeds are observed in the northern and central Red Sea, and both CPL and ATM.STA capture the features of the wind speed patterns. On June 24th, high wind speeds are observed in the central Red Sea and are also captured by both CPL and ATM.STA. 5 The mean 10-m wind speed over the Red Sea in ATM.STA, ATM.DYN, and CPL during the 30-day simulation are shown in Fig. 13. The mean error of CPL (mean bias: -0.23 m/s; RMSE: 2.38 m/s) is slightly smaller than the ATM.STA (mean bias: -0.34 m/s; RMSE: 2.43 m/s) by about 0.1 m/s. Although CPL, ATM.STA, ATM.DYN have different SST fields as the atmospheric boundary condition, the 10-m wind speed fields obtained in the simulations are all consistent with MERRA-2 data. The comparison shows the SKRIPS is capable of simulating the surface wind speed over the Red Sea in the coupled simulation.

Scalability Test 5

10

Parallel efficiency is crucial for coupled ocean-atmosphere models when simulating large and complex problems. In this section, the parallel efficiency in the coupled simulations is investigated. This aims to demonstrate (1) the implemented ESMF/NUOPC driver and model interfaces can simulate parallel cases effectively and (2) the ESMF/NUOPC coupler does

not add a significant computational overhead. The parallel speed-up of the model is investigated to evaluate its performance 15 for a constant size problem simulated using different numbers of CPU cores (i.e. strong scaling). Additionally, the CPU time



Figure 12. The magnitude and direction of the 10-m wind obtained in the CPL, the MERRA-2 data, and their difference (CPL–MERRA-2). The differences between ATM.STA and ATM.DYN with MERRA-2 (i.e., ATM.STA–MERRA-2, ATM.DYN–MERRA-2) are also presented. Two snapshots are selected: (1) 1200 UTC Jun 02 2012; (2) 1200 UTC Jun 24 2012.



Figure 13. The magnitude of the 10-m wind obtained in CPL, ATM.STA, and ATM.DYN in comparison with MERRA-2. The top panel shows the mean 10-m wind; the middle panel shows the mean bias; the bottom panel shows the RMSE. Only the hourly surface wind fields over the sea are shown to validate the coupled model.

spent on oceanic and atmospheric components of the coupled model is detailed. The test cases are run in sequential mode. The parallel efficiency tests are performed on the Shaheen-II cluster at KAUST (https://www.hpc.kaust.edu.sa/). The Shaheen-II cluster is a Cray XC40 system composed of 6174 dual-socket compute nodes based on 16-core Intel Haswell processors running at 2.3GHz. Each node has 128GB DDR4 memory running at 2300MHz. Overall the system has a total of 197,568 CPU

5 cores (6174 nodes $\times 2 \times 16$ CPU cores) and has a theoretical peak speed of 7.2 PetaFLOPS (10¹⁵ floating point operations per second).

The parallel efficiency of the scalability test is $N_{p0}t_{p0}/N_{pn}t_{pn}$, where N_{p0} and N_{pn} are the numbers of CPU cores employed in the baseline case and the test case, respectively; t_{p0} and t_{pn} are the CPU times spent on the baseline case and the test case, respectively. The speed-up is defined as t_{p0}/t_{pn} , which is the relative improvement of the CPU time when solving the problem.

- 10 The scalability tests are performed by running 24-hour simulations for ATM.STA, OCN.DYN, and CPL cases. There are 2.6 million atmosphere cells (256 lat×256 lon×40 vertical levels) and 0.4 million ocean cells (256 lat×256 lon×40 vertical levels, but about 84% of the domain is land and masked out). We started using N_{p0} = 32 because each compute node has 32 CPU cores. The results obtained in the scalability test of the coupled model are shown in Fig. 14. It can be seen that the parallel efficiency of the coupled code is close to 100% when employing less than 128 cores and is still as high as 70% when using 256
- 15 cores. When using 256 cores, there are a maximum of 20480 cells (16 lat×16 lon×80 vertical levels) in each core. The decrease in parallel efficiency results from the increase of communication time, load imbalance, and I/O (read and write) operation per CPU core (Christidis, 2015). From results reported in the literature, the parallel efficiency of the coupled model is comparable to other ocean-alone or atmosphere-alone models with similar numbers of grid points per CPU core (Marshall et al., 1997; Zhang et al., 2013b).



Figure 14. The parallel efficiency test of the coupled model in the Red Sea region, employing up to 512 CPU cores. The simulation using 32 CPU cores is regarded as the baseline case when computing the speed-up. Tests are performed on the Shaheen-II cluster at KAUST.

The CPU time spent on different components of the coupled run is shown in Table. 4. The time spent on the ESMF coupler is obtained by subtracting the time spent on oceanic and atmospheric components from the total time of the coupled run. The most time-consuming process is the atmosphere model integration, which accounts for 85% to 95% of the total. The ocean model integration is the second most time-consuming process, which is 5% to 11% of the total computational cost. If an ocean-only

- 5 region was simulated, the costs of the ocean and atmosphere models would be more equal compared with the Red Sea case. It should be noted that the test cases are run in sequential mode, and the cost breakdown among the components can be used to address load balancing in the concurrent mode. The coupling process takes less than 3% of the total cost in the coupled runs using different numbers of CPU cores in this test. Although the proportion of the coupling process in the total cost increases when using more CPU cores, the total time spent on the coupling process is similar. The CPU time spent on two uncoupled
- 10 runs (i.e., ATM.STA, OCN.DYN) is also shown in Table. 4. Compared with the uncoupled simulations, the ESMF-MITgcm and ESMF-WRF interfaces do not increase the CPU time in the coupled simulation. In summary, the scalability test results suggest that the ESMF/NUOPC coupler does not add significant computational overhead when using SKRIPS in the coupled regional modeling studies.

Table 4. Comparison of CPU time spent on coupled and stand-alone runs. The CPU time presented in the table is normalized by the time spent on the coupled run using 512 CPU cores. The CPU time spent on two stand-alone simulations are presented to show the difference between coupled and stand-alone simulations.

	$N_{p} = 32$	64	128	256	512
CPL run	7.27	4.04	2.02	1.39	1.00
WRF in CPL run	6.88(95%)	3.82(94%)	1.89(93%)	1.25(90%)	0.85(85%)
MITgcm in CPL run	0.37(5%)	0.19(5%)	0.12(6%)	0.11(8%)	0.11(11%)
Coupler in CPL run	0.02(0%)	0.03(1%)	0.02(1%)	0.03(2%)	0.03(3%)
ATM.STA run	6.89	3.80	1.84	1.22	0.83
OCN.DYN run	0.38	0.19	0.13	0.09	0.08

6 Conclusion and Outlook

15 This paper describes the development of the Scripps–KAUST Regional Integrated Prediction System (SKRIPS). To build the coupled model, we implement the coupler using ESMF with its NUOPC wrapper layer. The ocean model MITgcm and the atmosphere model WRF are split into initialize, run, and finalize sections, with each of them called by the coupler as subroutines in the main function.

The coupled model is validated by using a realistic application to simulate the heat events during June 2012 in the Red Sea region. Results from the coupled and stand-alone simulations are compared to a wide variety of available observational and reanalysis data. We focus on the comparison of the surface atmospheric and oceanic variables because they are used to drive the oceanic and atmospheric components in the coupled model. From the comparison, results obtained from various configurations of coupled and stand-alone model simulations all realistically capture the surface atmospheric and oceanic variables in the Red Sea region over a 30-day simulation period. The coupled system gives equal or better results compared with stand-alone model components. The 2-m air temperature in three major cities obtained in CPL and ATM.DYN are comparable and better than

5 ATM.STA. Other surface atmospheric fields (e.g., 2-m air temperature, surface heat fluxes, 10-m wind speed) in CPL are also comparable with ATM.DYN and better than ATM.STA over the simulation period. The SST obtained in CPL is also better than that in OCN.DYN by about 0.1 °C when compared with GHRSST.

The parallel efficiency of the coupled model is examined by simulating the Red Sea region using increasing numbers of CPU cores. The parallel efficiency of the coupled model is consistent with that of the stand-alone ocean and atmosphere models

10 using the same number of cores. The CPU time associated with different components of the coupled simulations shows that the ESMF/NUOPC driver does not add a significant computational overhead. Hence the coupled model can be implemented for coupled regional modeling studies on HPC clusters with comparable performance as that attained by uncoupled stand-alone models.

The results presented here motivate further studies evaluating and improving this new regional coupled ocean–atmosphere model for investigating dynamical processes and forecasting applications. This regional coupled forecasting system can be improved by developing coupled data assimilation capabilities for initializing the forecasts. In addition, the model physics and model uncertainty representation in the coupled system can be enhanced using advanced techniques, such as stochastic physics parameterizations. Future work will involve exploring these and other aspects a regional coupled modeling system suited for forecasting and process understanding.

20 Code and data availability. The coupled model, documentation, and the cases used in this work are available at https://library.ucsd.edu/dc/ collection/bb1847661c, and the source code is maintained on Github https://github.com/iurnus/scripps_kaust_model. ECMWF ERA5 data are used as the atmospheric initial and boundary conditions. The ocean model uses the assimilated HYCOM/NCODA 1/12° global analysis data as initial and boundary conditions. To validate the simulated SST data, we use the OSTIA (Operational Sea Surface Temperature and Sea Ice Analysis) system in GHRSST (Group for High Resolution Sea Surface Temperature). The simulated 2-m air temperature (T2) is validated against the ECMWF ERA5. The observed daily maximum and minimum temperatures from NOAA National Climate Data Center is used to validate the T2 in three major cities. Surface heat fluxes (e.g., latent heat fluxes, sensible heat fluxes, and longwave/shortwave radiation) are compared with MERRA-2 (Modern-Era Retrospective analysis for Research and Applications, version 2).

Appendix A: Snapshots of Heat Fluxes

To examine the modeling of turbulent heat fluxes and radiative fluxes, the snapshots of these heat fluxes obtained from 30 ATM.STA, ATM.DYN, and CPL are presented and validated using the MERRA-2 data.

The snapshots of the THFs at 1200 UTC June 2^{nd} and 24^{th} are presented in Fig. A1. It can be seen that all simulations reproduce the THFs reasonably well in comparison with MERRA-2. On June 2^{nd} , all simulations exhibit similar THF patterns.

This is because all simulations have the same initial conditions, and the SST fields in all simulations are similar within two days. On the other hand, for the heat event on June 24th, CPL and ATM.DYN exhibit more latent heat fluxes coming out of the ocean (170 and 153 W/m²) than those in ATM.STA (138 W/m²). The mean biases in CPL, ATM.DYN, and ATM.STA are 23.1 w/m², 5.1 w/m², and -9.5 w/m², respectively. Although CPL has larger bias at the snapshot, the averaged bias and RMSE

- 5 in CPL are smaller (shown in Fig. 10). Compared with the latent heat fluxes, the sensible heat fluxes in the Red Sea region are much smaller in all simulations (about 20 W/m²). It should be noted that MERRA-2 has unrealistically large sensible heat fluxes in the coastal regions because the land points 'contaminate' the values in the coastal region (Kara et al., 2008; Gelaro et al., 2017), and thus the heat fluxes in the coastal regions are not shown.
- The net downward shortwave and longwave radiation fluxes are shown in Fig. A2. Again, all simulations reproduce the shortwave and longwave radiation fluxes reasonably well. For the shortwave heat fluxes, all simulations show similar patterns on both June 2nd and 24th. The total downward heat fluxes, which is the sum of the results in Figs. A1 and A2, are shown in Fig. A3. It can be seen that the present simulations over-estimate the total downward heat fluxes on June 2nd (CPL: 580 W/m²; ATM.STA: 590 W/m²; ATM.DYN: 582 W/m²) compared with MERRA-2 (525 W/m²), especially in the southern Red Sea because of over-estimating the shortwave radiation. To improve the modeling of shortwave radiation, a better understanding
- 15 of the cloud and aerosol in the Red Sea region is required (Zempila et al., 2016; Imran et al., 2018). Again, the heat fluxes in the coastal regions are not shown because of the inconsistency of land-sea mask. Overall, the comparison shows the coupled model is capable of capturing the surface heat fluxes into the ocean.

Appendix B: Evaporation

To examine the simulation of surface freshwater flux in the coupled model, the surface evaporation fields obtained from 20 ATM.STA, ATM.DYN, and CPL are compared with the MERRA-2 data.

- The surface evaporation fields from CPL are shown in Fig. B1. The MERRA-2 data and the difference between CPL and MERRA-2 are also shown to validate the coupled model. The ATM.STA and ATM.DYN simulation results are not shown, but their differences with CPL are also shown in Fig. B1. It can be seen in Fig. B1(III) and B1(VIII) that CPL reproduces the overall evaporation patterns in the Red Sea. CPL is able to capture the relatively high evaporation in the northern Red Sea and the relatively low evaporation in the southern Red Sea in both snapshots, shown in Fig. B1(I) and B1(VI). After 36-hours, the simulation results are close with each other (e.g., the RMSE between CPL and ATM.STA simulation is smaller than 10 cm/year). However, after 24 days, CPL agrees better with MERRA-2 (bias: 6 cm/year; RMSE: 59 cm/year) than ATM.STA (bias: -25 cm/year; RMSE: 68 cm/year). In addition, the CPL results are consistent with those in ATM.DYN. This shows the coupled ocean–atmosphere simulation can reproduce the realistic evaporation patterns over the Red Sea. Since there is no
- 30 precipitation in three major cities (Mecca, Jeddah, Yanbu) near the eastern shore of the Red Sea during the month according to NCDC climate data, the precipitation results are not shown.



Figure A1. The turbulent heat fluxes out of the sea obtained in CPL, MERRA-2 data, and their difference (CPL–MERRA-2). The differences between ATM.STA and ATM.DYN with MERRA-2 (i.e., ATM.STA–MERRA-2, ATM.DYN–MERRA-2) are also presented. Two snapshots are selected: (1) 1200 UTC Jun 02 2012; (2) 1200 UTC Jun 24 2012. The simulation initial time is 0000 UTC Jun 01 2012 for both snapshots. Only the heat fluxes over the sea are shown to validate the coupled model.



Figure A2. The net downward shortwave and longwave radiation obtained in CPL, MERRA-2 data, and their difference (CPL–MERRA-2). The differences between ATM.STA and ATM.DYN with MERRA-2 (i.e., ATM.STA–MERRA-2, ATM.DYN–MERRA-2) are also presented. Two snapshots are selected: (1) 1200 UTC Jun 02 2012; (2) 1200 UTC Jun 24 2012. The simulation initial time is 0000 UTC Jun 01 2012 for both snapshots. Only the heat fluxes over the sea are shown to validate the coupled model.



Figure A3. Comparison of the total downward heat fluxes obtained in CPL, MERRA-2 data, and their difference (CPL–MERRA-2). The differences between ATM.STA and ATM.DYN with ERA5 (i.e., ATM.STA–MERRA-2, ATM.DYN–MERRA-2) are also presented. Two snapshots are selected: (1) 1200 UTC Jun 02 2012; (2) 1200 UTC Jun 24 2012. The simulation initial time is 0000 UTC Jun 01 2012 for both snapshots. Only the heat fluxes over the sea are shown to validate the coupled model.

Author contributions. RS worked on the coding tasks for coupling WRF with MITgcm using ESMF, wrote the code documentation, and performed the simulations for the numerical experiments. RS and ACS worked on the technical details for debugging the model and drafted the initial manuscript. All authors designed the computational framework and the numerical experiments. All authors discussed the results and contributed to the writing of the final manuscript.

5 Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We appreciate the computational resources provided by COMPAS (Center for Observations, Modeling and Prediction at Scripps) and KAUST used for this project. We gratefully acknowledge the research funding from KAUST (grant number: OSR-2-16-RPP-3268.02). We are immensely grateful to Caroline Papadopoulos for helping with installing software, testing the coupled code, and using the HPC clusters. We appreciate Professor U.U. Turuncoglu sharing part of their ESMF/NUOPC code on GitHub which helps our code

10 development. We wish to thank Dr. Ganesh Gopalakrishnan for setting up the stand-alone MITgcm simulation (OCN.DYN) and providing the external forcings. We thank Drs. Stephanie Dutkiewicz, Jean-Michel Campin, Chris Hill, Dimitris Menemenlis for providing their ESMF– MITgcm interface. We wish to thank Dr. Peng Zhan for discussing the simulations of the Red Sea. We also thank the reviewers for their insightful review suggestions.



Figure B1. The surface evaporation patterns obtained in CPL, the MERRA-2 data, and their difference (CPL–MERRA-2). The differences between uncoupled atmosphere simulations with MERRA-2 (i.e., ATM.STA–MERRA-2, ATM.DYN–MERRA-2) are also presented. Two snapshots are selected: (1) 1200 UTC Jun 02 2012; (2) 1200 UTC Jun 24 2012. Only the evaporation over the sea is shown to validate the coupled ocean–Atmosphere model.

References

Abdou, A. E. A.: Temperature trend on Makkah, Saudi Arabia, Atmospheric and Climate Sciences, 4, 457-481, 2014.

- Aldrian, E., Sein, D., Jacob, D., Gates, L. D., and Podzun, R.: Modelling Indonesian rainfall with a coupled regional model, Climate Dynamics, 25, 1–17, 2005.
- 5 Anderson, J. L. and Collins, N.: Scalable implementations of ensemble filter algorithms for data assimilation, Journal of Atmospheric and Oceanic Technology, 24, 1452–1463, 2007.
 - Barbariol, F., Benetazzo, A., Carniel, S., and Sclavo, M.: Improving the assessment of wave energy resources by means of coupled waveocean numerical modeling, Renewable Energy, 60, 462–471, 2013.

Bender, M. A. and Ginis, I.: Real-case simulations of hurricane-ocean interaction using a high-resolution coupled model: effects on hurricane

10 intensity, Monthly Weather Review, 128, 917–946, 2000.

Benjamin, S. G., Grell, G. A., Brown, J. M., Smirnova, T. G., and Bleck, R.: Mesoscale weather prediction with the RUC hybrid isentropic– terrain-following coordinate model, Monthly Weather Review, 132, 473–494, 2004.

Boé, J., Hall, A., Colas, F., McWilliams, J. C., Qu, X., Kurian, J., and Kapnick, S. B.: What shapes mesoscale wind anomalies in coastal upwelling zones?, Climate Dynamics, 36, 2037–2049, 2011.

15 Chen, S. S. and Curcic, M.: Ocean surface waves in Hurricane Ike (2008) and Superstorm Sandy (2012): Coupled model predictions and observations, Ocean Modelling, 103, 161–176, 2016.

- Chen, S. S., Price, J. F., Zhao, W., Donelan, M. A., and Walsh, E. J.: The CBLAST-Hurricane program and the next-generation fully coupled atmosphere–wave–ocean models for hurricane research and prediction, Bulletin of the American Meteorological Society, 88, 311–318, 2007.
- Christidis, Z.: Performance and Scaling of WRF on Three Different Parallel Supercomputers, in: International Conference on High Performance Computing, pp. 514–528, Springer, 2015.
- Collins, N., Theurich, G., Deluca, C., Suarez, M., Trayanov, A., Balaji, V., Li, P., Yang, W., Hill, C., and Da Silva, A.: Design and implementation of components in the Earth System Modeling Framework, The International Journal of High Performance Computing Applications, 19, 341–350, 2005.
- Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The operational sea surface temperature and sea ice analysis (OSTIA) system. Remote Sensing of Environment, 116, 140–158, 2012.
- Doscher, R., Willén, U., Jones, C., Rutgersson, A., Meier, H. M., Hansson, U., and Graham, L. P.: The development of the regional coupled ocean-atmosphere model RCAO, Boreal Environment Research, 7, 183–192, 2002.
- Evangelinos, C. and Hill, C. N.: A schema based paradigm for facile description and control of a multi-component parallel, coupled atmosphere-ocean model, in: Proceedings of the 2007 Symposium on Component and Framework Technology in High-Performance
- and Scientific Computing, pp. 83–92, ACM, 2007.

5

10

- Fairall, C., Bradley, E. F., Hare, J., Grachev, A., and Edson, J.: Bulk parameterization of air-sea fluxes: Updates and verification for the COARE algorithm, Journal of Climate, 16, 571–591, 2003.
 - Fang, Y., Zhang, Y., Tang, J., and Ren, X.: A regional air-sea coupled model and its application over East Asia in the summer of 2000, Advances in Atmospheric Sciences, 27, 583–593, 2010.
- 20 Fowler, H. and Ekström, M.: Multi-model ensemble estimates of climate change impacts on UK seasonal precipitation extremes, International Journal of Climatology, 29, 385–416, 2009.
 - Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., et al.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2), Journal of Climate, 30, 5419–5454, 2017.
- 25 Gualdi, S., Somot, S., Li, L., Artale, V., Adani, M., Bellucci, A., Braun, A., Calmanti, S., Carillo, A., Dell'Aquila, A., et al.: The CIRCE simulations: regional climate change projections with realistic representation of the Mediterranean Sea, Bulletin of the American Meteorological Society, 94, 65–81, 2013.
 - Gustafsson, N., Nyberg, L., and Omstedt, A.: Coupling of a high-resolution atmospheric model and an ocean model for the Baltic Sea, Monthly Weather Review, 126, 2822–2846, 1998.
- 30 Hagedorn, R., Lehmann, A., and Jacob, D.: A coupled high resolution atmosphere-ocean model for the BALTEX region, Meteorologische Zeitschrift, 9, 7–20, 2000.

Harley, C. D., Randall Hughes, A., Hultgren, K. M., Miner, B. G., Sorte, C. J., Thornber, C. S., Rodriguez, L. F., Tomanek, L., and Williams, S. L.: The impacts of climate change in coastal marine systems, Ecology Letters, 9, 228–241, 2006.

He, J., He, R., and Zhang, Y.: Impacts of air-sea interactions on regional air quality predictions using WRF/Chem v3.6.1 coupled with ROMS

v3.7: southeastern US example, Geoscientific Model Development Discussions, 8, 9965–10009, 2015.
 Henderson, T. and Michalakes, J.: WRF ESMF Development, in: 4th ESMF Community Meeting, Cambridge, USA, Jul 21, 2005.
 Hersbach, H.: The ERA5 Atmospheric Reanalysis., in: AGU Fall Meeting Abstracts, San Francisco, USA, Dec 12-16, 2016.

- Hill, C., DeLuca, C., Balaji, Suarez, M., and Silva, A.: The architecture of the Earth system modeling framework, Computing in Science & Engineering, 6, 18–28, 2004.
- Hill, C. N.: Adoption and field tests of M.I.T General Circulation Model (MITgcm) with ESMF, in: 4th Annual ESMF Community Meeting, Cambridge, USA, Jul 20-21, 2005.
- 5 Hodur, R. M.: The Naval Research Laboratory's coupled ocean/atmosphere mesoscale prediction system (COAMPS), Monthly Weather Review, 125, 1414–1430, 1997.
 - Hong, S.-Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes, Monthly Weather Review, 134, 2318–2341, 2006.
 - Hoteit, I., Hoar, T., Gopalakrishnan, G., Collins, N., Anderson, J., Cornuelle, B., Köhl, A., and Heimbach, P.: A MITgcm/DART ensemble
- analysis and prediction system with application to the Gulf of Mexico, Dynamics of Atmospheres and Oceans, 63, 1–23, 2013.
 Huang, B., Schopf, P. S., and Shukla, J.: Intrinsic ocean–atmosphere variability of the tropical Atlantic Ocean, Journal of Climate, 17, 2058–2077, 2004.
 - Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: calculations with the AER radiative transfer models, Journal of Geophysical Research: Atmospheres, 113, 2008.
- 15 Imran, H., Kala, J., Ng, A., and Muthukumaran, S.: An evaluation of the performance of a WRF multi-physics ensemble for heatwave events over the city of Melbourne in southeast Australia, Climate dynamics, 50, 2553–2586, 2018.

Kain, J. S.: The Kain–Fritsch convective parameterization: an update, Journal of Applied Meteorology, 43, 170–181, 2004.

- Kara, A. B., Wallcraft, A. J., Barron, C. N., Hurlburt, H. E., and Bourassa, M.: Accuracy of 10 m winds from satellites and NWP products near land-sea boundaries, Journal of Geophysical Research: Oceans, 113, 2008.
- 20 Kharin, V. V. and Zwiers, F. W.: Changes in the extremes in an ensemble of transient climate simulations with a coupled atmosphere-ocean GCM, Journal of Climate, 13, 3760–3788, 2000.
 - Large, W. G. and Yeager, S. G.: Diurnal to decadal global forcing for ocean and sea-ice models: the data sets and flux climatologies, Tech. rep., NCAR Technical Note: NCAR/TN-460+STR. CGD Division of the National Center for Atmospheric Research, 2004.

Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parame-

- terization, Reviews of Geophysics, 32, 363–403, 1994.
 - Loglisci, N., Qian, M., Rachev, N., Cassardo, C., Longhetto, A., Purini, R., Trivero, P., Ferrarese, S., and Giraud, C.: Development of an atmosphere-ocean coupled model and its application over the Adriatic Sea during a severe weather event of Bora wind, Journal of Geophysical Research: Atmospheres, 109, 2004.
 - Maksyutov, S., Patra, P. K., Onishi, R., Saeki, T., and Nakazawa, T.: NIES/FRCGC global atmospheric tracer transport model: Description,
- 30 validation, and surface sources and sinks inversion, Earth Simulator, 9, 3–18, 2008.
 - Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, Journal of Geophysical Research: Oceans, 102, 5753–5766, 1997.
 - Martin, M., Dash, P., Ignatov, A., Banzon, V., Beggs, H., Brasnett, B., Cayula, J.-F., Cummings, J., Donlon, C., Gentemann, C., et al.: Group for High Resolution Sea Surface Temperature (GHRSST) analysis fields inter-comparisons. Part 1: A GHRSST multi-product ensemble
- 35 (GMPE), Deep Sea Research Part II: Topical Studies in Oceanography, 77, 21–30, 2012.
 - Morrison, H., Thompson, G., and Tatarskii, V.: Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: Comparison of one-and two-moment schemes, Monthly Weather Review, 137, 991–1007, 2009.
 - National Geophysical Data Center: 2-minute Gridded Global Relief Data (ETOPO2) v2, 2006.

- Powers, J. G. and Stoelinga, M. T.: A coupled air-sea mesoscale model: Experiments in atmospheric sensitivity to marine roughness, Monthly Weather Review, 128, 208–228, 2000.
- Pullen, J., Doyle, J. D., and Signell, R. P.: Two-way air-sea coupling: A study of the Adriatic, Monthly Weather Review, 134, 1465–1483, 2006.
- 5 Ricchi, A., Miglietta, M. M., Falco, P. P., Benetazzo, A., Bonaldo, D., Bergamasco, A., Sclavo, M., and Carniel, S.: On the use of a coupled ocean-atmosphere-wave model during an extreme cold air outbreak over the Adriatic Sea, Atmospheric Research, 172, 48–65, 2016.
 - Roberts-Jones, J., Fiedler, E. K., and Martin, M. J.: Daily, global, high-resolution SST and sea ice reanalysis for 1985–2007 using the OSTIA system, Journal of Climate, 25, 6215–6232, 2012.
 - Roessig, J. M., Woodley, C. M., Cech, J. J., and Hansen, L. J.: Effects of global climate change on marine and estuarine fishes and fisheries, Reviews in Fish Biology and Fisheries, 14, 251–275, 2004.

10

35

- Seo, H.: Distinct influence of air-sea interactions mediated by mesoscale sea surface temperature and surface current in the Arabian Sea, Journal of Climate, 30, 8061–8080, 2017.
- Seo, H., Miller, A. J., and Roads, J. O.: The Scripps Coupled Ocean–Atmosphere Regional (SCOAR) model, with applications in the eastern Pacific sector, Journal of Climate, 20, 381–402, 2007.
- 15 Seo, H., Subramanian, A. C., Miller, A. J., and Cavanaugh, N. R.: Coupled impacts of the diurnal cycle of sea surface temperature on the Madden–Julian oscillation, Journal of Climate, 27, 8422–8443, 2014.
 - Sitz, L., Di Sante, F., Farneti, R., Fuentes-Franco, R., Coppola, E., Mariotti, L., Reale, M., Sannino, G., Barreiro, M., Nogherotto, R., et al.: Description and evaluation of the Earth System Regional Climate Model (RegCM-ES), Journal of Advances in Modeling Earth Systems, 2017.
- 20 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., and Huang, X.-Y.: A description of the Advanced Research WRF Version 4, Tech. rep., NCAR Technical Note: NCAR/TN-556+STR, 145 pp, doi:10.5065/1dfh-6p97, 2019.

Somot, S., Sevault, F., Déqué, M., and Crépon, M.: 21st century climate change scenario for the Mediterranean using a coupled atmosphere– ocean regional climate model, Global and Planetary Change, 63, 112–126, 2008.

- 25 Theurich, G., DeLuca, C., Campbell, T., Liu, F., Saint, K., Vertenstein, M., Chen, J., Oehmke, R., Doyle, J., Whitcomb, T., et al.: The earth system prediction suite: toward a coordinated US modeling capability, Bulletin of the American Meteorological Society, 97, 1229–1247, 2016.
 - Torma, C., Coppola, E., Giorgi, F., Bartholy, J., and Pongrácz, R.: Validation of a high-resolution version of the regional climate model RegCM3 over the Carpathian basin, Journal of Hydrometeorology, 12, 84–100, 2011.
- 30 Turuncoglu, U., Giuliani, G., Elguindi, N., and Giorgi, F.: Modelling the Caspian Sea and its catchment area using a coupled regional atmosphere-ocean model (RegCM4-ROMS): model design and preliminary results, Geoscientific Model Development, 6, 283, 2013.
 - Turuncoglu, U. U.: Toward modular in situ visualization in Earth system models: the regional modeling system RegESM 1.1, Geoscientific Model Development, 12, 233–259, 2019.
 - Turuncoglu, U. U. and Sannino, G.: Validation of newly designed regional earth system model (RegESM) for Mediterranean Basin, Climate Dynamics, 48, 2919–2947, 2017.
 - Van Pham, T., Brauch, J., Dieterich, C., Frueh, B., and Ahrens, B.: New coupled atmosphere-ocean-ice system COSMO-CLM/NEMO: assessing air temperature sensitivity over the North and Baltic Seas, Oceanologia, 56, 167–189, 2014.

- Warner, J. C., Armstrong, B., He, R., and Zambon, J. B.: Development of a coupled ocean-atmosphere-wave-sediment transport (COAWST) modeling system, Ocean Modelling, 35, 230–244, 2010.
- Xie, S.-P., Miyama, T., Wang, Y., Xu, H., De Szoeke, S. P., Small, R. J. O., Richards, K. J., Mochizuki, T., and Awaji, T.: A regional ocean–atmosphere model for eastern Pacific climate: toward reducing tropical biases, Journal of Climate, 20, 1504–1522, 2007.
- 5 Xu, J., Rugg, S., Byerle, L., and Liu, Z.: Weather forecasts by the WRF-ARW model with the GSI data assimilation system in the complex terrain areas of southwest Asia, Weather and Forecasting, 24, 987–1008, 2009.
 - Zempila, M.-M., Giannaros, T. M., Bais, A., Melas, D., and Kazantzidis, A.: Evaluation of WRF shortwave radiation parameterizations in predicting Global Horizontal Irradiance in Greece, Renewable Energy, 86, 831–840, 2016.

Zhang, H., Pu, Z., and Zhang, X.: Examination of errors in near-surface temperature and wind from WRF numerical simulations in regions

10 of complex terrain, Weather and Forecasting, 28, 893–914, 2013a.

Zou, L. and Zhou, T.: Development and evaluation of a regional ocean-atmosphere coupled model with focus on the western North Pacific summer monsoon simulation: Impacts of different atmospheric components, Science China Earth Sciences, 55, 802–815, 2012.

Zhang, X., Huang, X.-Y., and Pan, N.: Development of the upgraded tangent linear and adjoint of the Weather Research and Forecasting (WRF) Model, Journal of Atmospheric and Oceanic Technology, 30, 1180–1188, 2013b.