

# Supplementary material for: An ensemble of AMIP simulations with prescribed land surface temperatures

Duncan Ackerley<sup>1,2</sup>, Robin Chadwick<sup>3</sup>, Dietmar Dommenget<sup>1</sup>, and Paola Petrelli<sup>4</sup>

<sup>1</sup>ARC Centre of Excellence for Climate System Science, School of Earth Atmosphere and Environment, Monash University, Clayton 3800, Victoria, Australia

<sup>2</sup>Met Office, Exeter, UK

<sup>3</sup>Met Office Hadley Centre, Exeter, UK

<sup>4</sup>ARC Centre of Excellence for Climate System Science, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia


**Correspondence:** Duncan Ackerley (duncan.ackerley@metoffice.gov.uk)

*Copyright statement.* The works published in this journal are distributed under the Creative Commons Attribution 4.0 License. This licence does not affect the Crown copyright work, which is re-usable under the Open Government Licence (OGL). The Creative Commons Attribution 4.0 License and the OGL are interoperable and do not conflict with, reduce or limit each other.


© Author(s) and British Crown Copyright 2018

5 This supplementary information contains three figures:


- (1) The change in surface air temperature, precipitation and mean sea level pressure for each of the perturbed (SST+4K, 4xCO<sub>2</sub>, 4xCO<sub>2</sub>rad and solar+3.3%) AMIP free land simulations relative to the AMIP free land control run (Fig. S1) as a reference to show the climatic changes associated with each forcing.
- (2) The change in surface air temperature, precipitation and mean sea level pressure for each of the perturbed (SST+4K, 4xCO<sub>2</sub>, 4xCO<sub>2</sub>rad and solar+3.3%) AMIP prescribed land simulations relative to the AMIP prescribed land control run (Fig. S2) as a reference to show the climatic changes in the prescribed land experiments are comparable to those of their free land counterparts.
- (3) The change in surface latent heat flux over the Amazon for each free land simulation relative to its prescribed land counterpart to show the increases associated with the higher precipitation in the prescribed land simulations (Fig. S3).



**Figure S1.** Differences in surface air temperature (TAS, K) for (a) A4K-A, (b) A4x-A, (c) Arad4x-A, (d) Asc-A. Equivalent differences between like-for-like simulations are given in (e)–(h) and (i)–(l) for precipitation (PR,  $\text{mm day}^{-1}$ ) and mean sea level pressure (PSL, hPa), respectively. The points labelled with an "x" indicate the differences are statistically significant using the Student's t-test ( $p \leq 0.05$ ).



**Figure S2.** Differences in surface air temperature (TAS, K) for (a) A4K<sub>PL</sub>4K-A<sub>PL</sub>, (b) A4x<sub>PL</sub>4x-A<sub>PL</sub>, (c) Arad4x<sub>PL</sub>rad4x-A<sub>PL</sub>, (d) Asc<sub>PL</sub>sc-A<sub>PL</sub>. Equivalent differences between like-for-like simulations are given in (e)-(h) and (i)-(l) for precipitation (PR, mm day<sup>-1</sup>) and mean sea level pressure (PSL, hPa), respectively. The points labelled with an "x" indicate the differences are statistically significant using the Student's t-test ( $p \leq 0.05$ ).



**Figure S3.** The difference in surface latent heat flux ( $\text{W m}^{-2}$ ) for (a)  $A_{PL}$  -  $A$ , (b)  $A4K_{PL4K}$  -  $A4K$ , (c)  $A4x_{PL4x}$  -  $A4x$ , (d)  $Arad4x_{PLrad4x}$  -  $Arad4x$  and (e)  $Asc_{PL}$  -  $Asc$ . The points labelled with an "x" indicate the differences are statistically significant using the Student's t-test ( $p \leq 0.05$ ).