

***Interactive comment on “MOVEIM v1.0:
Development of a bottom-up motor vehicular
emission inventories for the urban area of Manaus
in central Amazon rainforest” by Paulo R. Teixeira
et al.***

Anonymous Referee #2

Received and published: 28 June 2018

- 1) It is important to emphasize the methodology used to extrapolate the VTK information, obtained at peak times, for the others hours (off-peak period).
- 2) Why not use fleet age in the categorization and emissions factors? For example, the CO EF used (1.2 g / km) is very high for the fleet age that the author mentions in the introduction (“...average age of 5 years attributed to the timing of rapid urban growth during the economic expansion between 2009 and 2015...”). This can lead to an overestimation of emissions. The author can use a Top-down approach to age distribution (From a national database like ANFAVEA) or local information from DETRAN.

C1

- 3) The authors considered evaporative emissions?
- 4) With regard to fuel, the authors can estimate from consumption (local information) or from the national inventory (This hypothesis - top-down approach - is more subject to overestimation).
- 5) Please, check the NOx emission factor used to trucks (0.5 g/km). I am thinking there is an underestimation of NOx emission by heavy vehicles.

Interactive comment on Geosci. Model Dev. Discuss., <https://doi.org/10.5194/gmd-2018-81>,
2018.

C2