

## Development and implementation of a new biomass burning emissions injection height scheme (BBEIH v1.0) for the GEOS-Chem model (v9-01-01)

5 Liye Zhu<sup>1,2,3</sup>, Maria Val Martin<sup>4,5</sup>, Luciana V. Gatti<sup>6,7</sup>, Ralph Kahn<sup>8</sup>, Arsineh Hecobian<sup>3</sup>, and Emily V. Fischer<sup>3</sup>

<sup>1</sup> School of Atmospheric Sciences, Sun Yat-sen University, Guangzhou 510275 China

10 <sup>2</sup> Guangdong Province Key Laboratory for Climate Change and Natural Disaster Studies, Sun Yat-sen University, Guangzhou 510275 China

<sup>3</sup> Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA.

<sup>4</sup> Chemical and Biological Engineering Department, The University of Sheffield, Sheffield, UK.

15 <sup>5</sup> Now at Leverhulme Center for Climate Change Mitigation, Animal Plant Sciences Department, The University of Sheffield, Sheffield, UK.

<sup>6</sup> Instituto de Pesquisas Energéticas e Nucleares (IPEN)–Comissão Nacional de Energia Nuclear (CNEN), Cidade Universitária, São Paulo CEP, Brazil.

<sup>7</sup> National Institute for Space Research, INPE/CCST, LaGEE (Greenhouse Gas Laboratory), São José dos Campos-SP, Brazil

20 <sup>8</sup> Climate and Radiation Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.

Correspondence to: Emily V. Fischer (evf@atmos.colostate.edu)

**Abstract.** Biomass burning is a significant source of trace gases and aerosols to the  
25 atmosphere, and the evolution of these species depends acutely on where they are  
injected into the atmosphere. GEOS-Chem is a chemical transport model driven by  
assimilated meteorological data that is used to probe a variety of scientific questions  
related to atmospheric composition, including the role of biomass burning. This paper  
presents the development and implementation of a new global biomass burning emissions  
30 injection scheme in the GEOS-Chem model. The new injection scheme is based on  
monthly gridded Multi-Angle Imaging Spectro Radiometer (MISR) global plume-height  
stereoscopic observations in 2008. To provide specific examples of the impact of the  
model updates, we compare the output from simulations with and without the new MISR-  
based injection height scheme to several sets of observations from regions with active  
35 fires. Our comparisons with ARCTAS aircraft observations show that the updated  
injection height scheme can improve the ability of the model to simulate the vertical  
distribution of peroxyacetyl nitrate (PAN) and carbon monoxide (CO) over North

American boreal regions in summer. We also compare a simulation for October 2010 and 2011 to vertical profiles of CO over the Amazon Basin. When coupled with larger  
40 emission factors for CO, a simulation that includes the new injection scheme also better matches selected observations in this region. Finally, the improved injection height improves the simulation of monthly mean surface CO over California during July 2008, a period with large fires.

## 1 Introduction

45 Properly describing the injection altitude of smoke in the atmosphere is an essential step in predicting the impact of emissions from landscape fires on atmospheric composition (Paugam et al., 2016). Injecting smoke higher in the atmosphere in chemical transport models can extend or reduce the lifetime of trace species, and it can alter the spatial extent of smoke-influence in the atmosphere (Freitas et al., 2006). The impact of  
50 injection height on smoke dispersion is three-fold: 1) winds in the free troposphere are generally stronger than in the boundary layer, thus when smoke is emitted aloft, defined plumes are sometimes detected thousands of kilometers downwind (e.g., Colarco et al. (2004); Damoah et al. (2004); Forster et al. (2001); Val Martín et al. (2006)). 2) Removal processes tend to be more efficient in the boundary layer (e.g. Boy et al. (2008)). 3)  
55 Chemical evolution within the plume can be sensitive to injection height, because altitude impacts plume temperature, ambient relative humidity, smoke-cloud interactions, and photolysis rates (e.g. Freitas et al., 2006). Given the importance for atmospheric composition and air quality predictions (e.g. Stein et al., 2009), substantial efforts have been made to better understand how injection height varies by ecosystem type and season  
60 (e.g., Val Martin et al. (2010); Tosca et al. (2011); Mims et al. (2010)), which environmental drivers of injection height are most important (e.g., Kahn et al. (2007); Val Martin et al. (2012)), and how best to estimate smoke injection height in models (e.g., Paugam et al. (2016) and references therein) to produce improvements in model simulations of trace constituents (e.g., Gonzi et al. (2015)).

65 GEOS-Chem is a global chemical transport model (CTM) ([www.geos-chem.org](http://www.geos-chem.org)) (Bey et al., 2001) that is routinely used to simulate the impacts of biomass burning on atmospheric composition (e.g., Lewis et al. (2013) and Leung et al. (2007)). GEOS-Chem is driven by GEOS assimilated meteorological data from the NASA Global Modeling and

Assimilation Office (GMAO), and it includes a state-of-the-science description of  
70 tropospheric oxidant chemistry, necessary for understanding the chemical and dynamical  
processes controlling the evolution of biomass burning emissions. The public-release  
version of GEOS-Chem emits all biomass burning emissions into the atmospheric  
boundary layer. This may be appropriate for some fire types, but is likely a source of  
error for many regions with active biomass burning (e.g. Leung et al., 2007). The main  
75 objective of the current paper is to introduce a new global biomass burning injection  
height scheme for GEOS-Chem based on Multi-Angle Imaging Spectro Radiometer  
(MISR) plume injection height observations from 2008. The MISR instrument was  
launched into a sun-synchronous, polar orbit aboard the NASA Earth Observing System's  
Terra satellite in December 1999, and acquires global observations at nine viewing angles  
80 in each of four spectral bands about once per week (e.g., Diner et al., 1998). Smoke  
aerosol injection height is derived from source plumes with discernable features in the  
MISR multi-angle views (Kahn et al., 2008).

Though biomass burning impacts atmospheric composition across a suite of  
temporal and geographic scales, this paper presents model-observation comparisons for  
85 specific biomass burning plumes having well-sampled vertical structure. The data  
available to make such important comparisons is limited. However, this is an important  
step toward using the model to address broader aspects of atmospheric composition. To  
the best of our knowledge, this paper represents the first effort at using measured global  
smoke plume injection heights from MISR as constraints on a CTM. There have been  
90 efforts to do this on a regional scale for specific fire seasons (e.g., Chen et al., 2009; Jian  
and Fu, 2014), but we are unaware of similar global implementations. Val Martin et al.  
(2012) studied the performance of one of the most advanced physically based plume-rise  
models. They concluded that given the uncertainties and performance of that approach,  
empirically derived plume injection heights, such as those we use here, provide better  
95 constraints on smoke transport.

Much of the model development presented here was motivated by the persistent  
challenge CTMs appear to face at accurately simulating peroxyacetyl nitrate (PAN) in the  
atmosphere (e.g. Emmons et al. (2015)). This compound plays a central role in oxidant  
chemistry, particularly in remote regions (Moxim et al., 1996). However, it has a

100 temperature dependent lifetime (Singh and Hanst, 1981), and thus its evolution in the  
atmosphere is particularly sensitive to plume injection height. As a first step toward  
validating the revised model, we compare the output from a simulation with improved  
injection heights to multiple sets of observations from regions with active fires, providing  
examples of cases where injecting a substantial percent of biomass burning emissions in  
105 the free troposphere is important for properly simulating PAN as well as carbon  
monoxide (CO).

## **2 Methods**

### **2.1 Overview of Model Development**

Figure 1 illustrates the process of implementing an observationally based injection  
110 scheme into GEOS-Chem. This section describes the details associated with each step in  
the process. The new injection scheme is based on MISR plume injection height  
observations from 2008 (Section 2.2). The model configuration is described in Section  
2.3. We then map the native MISR injection altitude (0-8 km) to emitted percentages of  
total biomass burning emissions to GMAO 47-layer reduced vertical grid and  $2^\circ \times 2.5^\circ$   
115 horizontal grid (Section 2.4).

### **2.2 Analysis of MISR Plume Height Observations**

The new injection scheme is developed based on the MISR plume-height  
stereoscopic observations in 2008 (Val Martin et al., 2018, submitted to Remote  
Sensing). The MISR data we used is part of the MISR Plume Height Project2, which was  
120 derived for the AeroCom multi-model biomass burning experiment. The dataset is  
publicly available from <https://misr.jpl.nasa.gov/getData/accessData/MisrMinxPlumes2/>.  
Briefly, MISR-based injection heights are given by altitude (250 m, from 0 to 8 km above  
ground level), land cover type, season and region. Land cover classifications are based on  
MODIS Level 3 land cover product MOD12Q1 (Friedl et al., 2010). There are twelve  
125 classifications used here: Evergreen Needle Leaf Forest, Evergreen BroadLeaf Forest,  
Deciduous Needle Leaf Forest, Deciduous BroadLeaf Forest, Mixed Forest, Closed  
Shrub, Open Shrub, Woody Savanna, Savanna, Grassland, Wetland and Cropland. We  
define seasons as spring (MAM), summer (JJA), fall (SON) and winter (DJF), and  
considered 8 main fire regions (North America, South America, Africa, Europe, Boreal  
130 Eurasia, South Asia and Australia).



To convert the MISR-based vertical distribution of smoke injection height, Val Martin et al. (2018) first transformed the MISR vertical distribution percentages from 0 to 8 km at 250 m bins into the GEOS-Chem 47 level vertical grid (0.058, 0.189, 0.32, 0.454, 0.589, 0.726, 0.864, 1.004 km, etc.). Second, they determined the largest land cover type coverage in each GEOS-Chem grid. For that, they re-gridded their land cover unit map from  $0.005^\circ \times 0.005^\circ$  to  $2^\circ \times 2.5^\circ$  degree resolution assigning the highest ranked land cover type to each  $2^\circ \times 2.5^\circ$  grid. Finally, they applied the re-gridded vertical distribution of smoke percentages to each  $2^\circ \times 2.5^\circ$  degree grid depending on the defined land cover type and region. An overview of the MISR instrument and standard products is given by Diner et al. (1998), and more details about the MISR plume digitizing tool and the MISR plume database can be found in Nelson et al. (2013) and Val Martin et al. (2018, submitted to Remote Sensing), respectively.

There are several subtleties to the MISR-based plume-height climatology that are worth specifically noting here. MISR equator-crossing time during the day is about 10:30 AM, so the diurnal distribution of emissions is not sampled, and in particular, these data does not represent the mid-to-late afternoon period, when wildfires tend to be most intense. In order to evaluate the impact of the afternoon peaks on the parameterization, a qualitative assessment of the diurnal representativeness of the MISR plume-height record is required, as well as the corresponding 4 micron brightness temperature anomalies (termed Fire Radiative Power, or FRP) data from other satellite instruments (e.g., Ichoku and Kaufman, 2005). Limitations of the parameterization are further discussed in Val Martin et al. (2018, submitted to Remote Sensing), some of which would be worth exploring in the future. Also, the MISR-based plume-height climatology does not include plumes smaller than a certain size, and this size varies with observing conditions. Several factors contribute to this limitation. MODIS thermal anomalies are used to identify fire locations, some fires are smaller than MODIS pixels, others can be obscured by the tree canopy or overlying smoke, and fires for which the emissivity at 4 microns is low (e.g., smoldering fires), are sometimes missed (Kahn et al., 2008). These issues also affect satellite-based smoke emissions inventories such the one used here (see Section 2.4). The other limitation is that small fires may sometimes be missed by the MINX digitizer users, and/or can be digitized with low quality as they have low stereo-height retrieval densities.

To account for these issues, we include an adjustment to the smoke injection height scheme to account for small fires. Specifically, we use Global Fire Emissions Database version 4 (GFED4s) (Randerson et al., 2012) to estimate the fraction of small fires in each region and biome for the study year 2008. As nearly all small fires inject smoke only within the boundary layer, we apply a small-fire correction to the lowest model atmospheric layer as described in Val Martin et al. (2018, submitted to Remote Sensing). Note that aside from the small-fire information in GFED4s, derived separately from the standard satellite retrieval approach of the GFED products, we use GFEDv3 for this study. The emission factors for several species, such as CO for Temperate Forests, are lower in GFEDv4 compared to GFEDv3, which exacerbates known problems of low CO with GFED-initialized models (Akagi et al. 2011; van der Werf et al. 2017). As discussed in Section 2.4 below, we increased and tested the emission factors in GFEDv3 based on the findings in Petrenko et al. (2017).

### 175 **2.3 GEOS-Chem Configuration**

We use the Goddard Earth Observing System-Chemistry (GEOS-Chem) global 3-D chemical transport model including detailed ozone-NO<sub>x</sub>-VOC-aerosol chemistry (version 9.01.01, [www.geos-chem.org](http://www.geos-chem.org)) with modifications to emitted species and the chemical mechanism specifically for PAN as described in Fischer et al. (2014). Most relevant to this work, we use GFEDv3 monthly biomass burning emissions (van der Werf et al., 2010), with updated emission factors for non-methane volatile organic compounds (NMVOCs) and nitrogen oxides (NO<sub>x</sub>) from Akagi et al. (2011). The current work aims at addressing specifically the issue of injection height. Our injection height parameterization could be used with any emission inventory. The version of GEOS-Chem that we chose for developing and implementing the improved injection height scheme includes a number of code updates focused specifically on providing a better representation of PAN chemistry. It includes a more detailed chemical mechanisms related to PAN and a larger suite of precursor NMVOCs emissions. This model version has also been compared to a large suite of aircraft observations. Otherwise, we have used the standard input file settings used in GEOS-Chem. However, we note that choosing a monthly-averaged emission dataset can create biases for specific case studies of biomass burning.

PAN in biomass burning plumes is particularly sensitive to injection altitude because the lifetime of PAN is highly temperature dependent. Thus we focus a substantial portion of our model-measurement comparison on this species. The model experiments in Fischer et al. (2014) were among main motivations for the current paper. Thus, our model configurations are mainly based on the configuration used in this earlier study. However, the current work is focused on understanding potential changes in model performance following the inclusion of the new MISR-based injection height scheme. To keep this focus, there are two differences between the model configuration in Fischer et al. (2014) and our “standard model.” 1) We adjust the biomass burning emissions used in Fischer et al. (2014) to remove the increased biomass burning emissions for northern Asia, originally applied for 2008 in Fischer et al. (2014). These were applied in Fischer et al. (2014) because Kaiser et al. (2012) and Yu et al. (personal communication) found that GFEDv3 underestimates fire emissions at boreal latitudes. 2) We also remove the injection partitioning assumption applied in Fischer et al. (2014), which emitted 35% of total biomass burning emissions above the boundary layer to test the sensitivity of PAN to this choice. Fischer et al. (2014) found improved the PAN simulation, but it is a much coarser approach than what has been done here.

In the following text and figures, we refer to the version of model with the two changes noted above as the “standard model” because the injection of biomass burning is treated as in the public release benchmarked version of GEOS-Chem. We refer to the observationally based injection scheme as the “new injection scheme.” As is discussed later, we then apply different scaling factors for fire emissions following Petrenko et al. (2017) (see below) to the “new injection scheme.” We refer to this final model configuration in our figures as the “new injection scheme with increased CO.”

#### **2.4 GEOS-Chem Implementation**

We associate the native MISR injection altitude (0-8 km) with the emitted percentages of total biomass burning emissions, and map them to the GMAO 47-layer reduced vertical grid and a  $2^\circ \times 2.5^\circ$  horizontal grid. The injection percentages of total column biomass burning emissions for each month for each grid cell are saved in a binary file. (Code modifications to read in the data of percentages and distribute the biomass burning emissions to every grid cell are contained within the setemis.F FORTRAN

module in GEOS-Chem version 9.01.01. The binary file can be updated based on  
225 different analyses (e.g., a more recent year, or a different analysis approach), and little  
effort would be required to update this within the code.

In Figure 2a, we show an example vertical profile of injection percentages at  
56°N, 105°W from the standard model and the new injection scheme. In contrast to a  
blanket approach of emitting all biomass burning emissions within the boundary layer,  
230 the new injection scheme emits a large percentage of these emissions above the boundary  
layer at this location. The global map in Figure 2b shows the injection percentages at 850  
hPa in July 2008 for the globe based on the MISR stereo-height data. The amount of the  
total biomass burning emissions at any given location that are injected into the layer  
encompassing 850 hPa varies substantially. Figure 2a cannot be interpreted as the total  
235 amount of smoke emitted in this layer of the atmosphere; this is a plot of the percent of  
the total column that the model emits at that level. For example, there are regions during  
the month of July 2008 with high percentages of emissions injected at a given level, but  
very small total column emissions overall.

Given the combined limitations in the MISR analysis (Section 2.2) and the GFED  
240 emissions database at representing small fires, our scheme is unlikely to correctly  
represent the fraction of total smoke that is emitted above the boundary layer in places  
where small fires make a significant smoke contribution. Randerson et al. (2012) updated  
the GFEDv4 inventory to include an estimate of the emissions from fires below the  
detection limit of the satellite observations used to construct the standard GFED database,  
245 and most other satellite-based emission inventories. These small fires tend to include  
agricultural and shrubland fires as well as some grassland fires, peat fires, and ground  
fires where the overlying tree canopy is dense. The number of small fires is large in some  
places, their overall contribution to total emissions can be large, and they often produce  
diffuse, smoky haze rather than discrete plumes that are feasible to map from space. They  
250 also tend to inject smoke into the planetary boundary layer rather than above it. These  
fires are not the focus of the MISR injection height analysis or the MODIS FRP analysis,  
and although we have attempted to account for this (Section 2.2), this is a limitation of  
our overall approach.

As a final model experiment, we increase the CO emissions by a factor of 1.5 for

255 burning over savannas, a factor of 1.5 for burning associated with deforestation, and a  
factor of 2 for extra-tropical forests following Petrenko et al. (2017). We use  $2^\circ \times 2.5^\circ$   
horizontal resolution for our global simulations.

## 2.5 Observational Datasets

As a demonstration of the potential impact of the model development and its  
260 relevance to a few example regions, we compare GEOS-Chem output with improved  
injection heights to smoke-impacted trace gas observations from aircraft over boreal  
North America (July 2008) and from aircraft sites in the Amazon basin (2010 – 2011).  
We also compare the model output to monthly mean surface CO observations in regions  
impacted by major fires.

### 265 2.5.1 North America

Boreal North America is an interesting focal region because emissions from  
biomass burning lead to enhancements in high latitude tropospheric ozone during  
summer (Arnold et al., 2015). The representation of injection height has implications for  
inverse studies of emissions from fires in this region and the magnitude of the ozone  
270 enhancement that results from these emissions (Leung et al., 2007). The second portion  
of the NASA Arctic Research of the Composition of the Troposphere from Aircraft and  
Satellites (ARCTAS) mission was conducted over western Canada during June and July  
2008. A complete list of species observed by the NASA DC-8 aircraft during ARCTAS  
can be found in Jacob et al. (2010). In the present study, we use ARCTAS observations  
275 of CO and PAN from July 2008 to illustrate the updated performance of the model with  
the new injection scheme over western North America.

### 2.5.2 Amazon

We highlight the Amazon basin as another interesting region as emissions from  
deforestation fires over Amazonian forests represent a large percent of global emissions  
280 from deforestation (van der Werf et al., 2010). Year-to-year variability in this region has  
been associated with climate extremes (Chen et al., 2013). Thus, it is important to  
understand the fire injection height over this region in order to fully quantify the impact  
of these fires on atmospheric composition, and to better predict how this impact could  
evolve in the future. We use the CO observations from four sites across the Amazon  
285 basin in 2010 and 2011: Alta Floresta (ALF;  $8.80^\circ\text{S}$ ,  $56.75^\circ\text{W}$ ), Rio Branco (RBA;

9.38°S, 67.62°W), Santarém (SAN; 2.86°S; 54.95°W) and Tabatinga (TAB; 5.96°S, 70.06°W). Bi-weekly vertical profiles of CO were measured from just above the forest canopy to 4.4 km above sea level (Gatti et al., 2014). As described in Gatti et al. (2014), samples were collected using a small aircraft. Air samples were collected in flasks that were analyzed using a replica of the NOAA Earth System Research Laboratory (ESRL) trace gas analysis system. The measurements were taken at specific altitude levels on each flight day. Up to six or eight observations are available at each individual altitude level for each month (4 sites with 2 vertical profiles). We also compared model output to aircraft observations from the Balanço Atmosférico Regional de Carbono na Amazônia (BARCA) program, which deployed in 2008 (Andreae et al., 2012). This dataset contains a strong influence of biomass burning emissions. However, when we sampled the model at the locations of the observations, there were no differences in the simulated CO profiles in the two sets of simulations with the different injection schemes. The CO mixing ratios for the regions were biased low, i.e. model mixing ratios were between 80 and 125 ppb, indicating no smoke influence, whereas the corresponding observations were largely > 150 ppb. Andreae et al. (2012) discuss problems with GFEDv3 CO emissions for this region; specifically noting that the emissions in this database could be up to a factor of seven too low for the BARCA period.

### 2.5.3 Surface Observations

Leung et al. (2007) showed that the choice of injection height for boreal fire emissions impacts the simulation of surface CO mixing ratios in the Northern Hemisphere. They compared GEOS-Chem simulated anomalies in CO mixing ratios with surface measurements from the NOAA ESRL Global Monitoring Division (GMD), Carbon Cycle Cooperative Air Sampling Network (Novelli et al., 2003). Therefore, we also performed a comparison with monthly mean observations from 18 sites that may have been impacted by fires during 2008. In most locations (16 of 21) where we conducted comparisons, the model with the MISR-based injection height did not produce notably different surface monthly mean CO mixing ratios (i.e. changes are less than 1 ppb). However, there are four stations where the updated model produces substantially lower monthly mean surface CO mixing ratios than the standard model, and this change produces a better simulation of CO at these locations. We present these results in Section

3.3.

### 3 Results

#### 3.1 North American Boreal Fires

320 Figure 3a shows the total percent of biomass burning emissions for each model  
column emitted above 700 hPa during July 2008 based on MISR observations. We use  
the 700 hPa level to signify an approximate mid-day boundary layer top pressure over  
North America. The percent of emissions injected above 700 hPa in the updated version  
of the model is quite large over boreal regions, exceeding 60% for some locations such as  
325 the one shown in Figure 3b. In boreal regions, the majority of biomass burning emissions  
are produced by a relatively small number of large fires that last days to weeks (Stocks et  
al., 2002; Brey et al., 2018). Figure 3a shows the strong north-south gradient in the  
percent of emissions injected above this atmospheric level. In contrast to boreal regions,  
the new scheme continues to inject nearly all the fire emissions into the boundary layer  
330 over the central U.S. during this month. An example profile of the emitted percent by  
model layer is shown in Figure 3c. There are typically very few fires during July in this  
region; those that do occur are typically short-lived and often involve cropland (Brey et  
al., 2018).

The impact of the new injection scheme on simulated PAN has significant spatial  
335 variability over North America during July 2008, and this is driven by the large spatial  
variability in the fires and the smoke injection level. Figures 4a - 4c present the  
differences in simulated PAN mixing ratios between the updated and the standard model  
at the surface, 850 hPa and 510 hPa on 1 July 2008, respectively. As expected, the new  
injection scheme decreases simulated PAN mixing ratios at the surface and within the  
340 boundary layer over boreal regions. Simulated PAN mixing ratios increase in the mid  
and upper-troposphere. Figure 5 presents a similar example for 4 July 2008.

Figures 4d and 5d (black lines and open circles) show average vertical profiles of  
PAN intercepted by the DC8 during the ARCTAS flights on these particular days. The  
NASA DC8 sampled fresh smoke from the Lake McKay fire (56.5°N, 106.8°W) on 1  
345 July 2008 at several distances downwind (see Alvarado et al. (2010)). We sampled both  
versions of the model along the aircraft pathway at the corresponding observation time,  
and these average profiles are also plotted in Figures 4d. In the lower troposphere, the

standard model largely overestimates PAN on 1 July 2008 (Figure 4d). The new injection scheme decreases the simulated PAN in the boundary layer significantly (~ 300 pptv) and matches better to the ARCTAS observations. The same is not true for the comparison with CO (Figure 5). The DC8 sampled several plumes above 3 km on 4 July. As described in Alvarado et al. (2010), this was a period with strong updrafts, which led to lofting of biomass burning emissions (Fuelberg et al., 2010). We note that time of day could be very important for these comparisons. The aircraft sampled these plumes in the mid-to-late afternoon, so MISR heights are likely underestimates of the actual injection altitudes for the cases shown in Figures 4, 5, 6 and 7.

On 4 July 2008 (Figure 6d), the new injection scheme does not change simulated PAN meaningfully near the surface where the aircraft was located. Figures 6e and 6c show that the aircraft did not fly through the low altitude regions of the model, which showed important changes from the injection scheme. The near surface PAN mixing ratios were not impacted south of the Hudson Bay. However, the new injection scheme does increase PAN by ~130 pptv in the lower to mid free troposphere. This improves the model-measurement comparison substantially between 800 and 500 hPa. The GEOS-Chem simulations presented in Alvarado et al. (2010) substantially underestimated PAN relative to the ARCTAS observations. Here we have included the partitioning of NO<sub>x</sub> immediately to PAN and HNO<sub>3</sub> as originally suggested by Alvarado et al. (2010) and we have updated the injection height. Along with the other updates in Fischer et al. (2014), this appears to greatly improve the ability of the model to simulate the appropriate magnitude of PAN for the cases shown.

Figure 7 presents simulated and observed CO from the 4 July ARCTAS flight. Similar to PAN, for this particular profile the new injection scheme decreases CO in lower troposphere and increases it in middle troposphere (Figure 7d). However, both the standard model and the new injection scheme underestimate CO significantly compared to ARCTAS observations. Both model versions continue to produce a monotonic decrease in CO from the surface to upper levels, and although the new injection scheme increases CO just above 700 hPa, it is not able to simulate the enrichment layer that appears present in the observations. The mean CO underestimate shown in Figure 7d is 15%-56%. The model does not appear to have such a low bias for the 1 July case (Figure



5), but there are very few samples at higher altitudes in this flight. Alvarado et al. (2010)  
380 and Fisher et al. (2010) previously compared a GEOS-Chem simulation to ARCTAS  
observations. The simulation used in those prior studies was based on daily emissions  
from the Fire Locating and Monitoring of Burning Emissions (FLAMBE) emission  
inventory (Reid et al., 2009). Monthly mean GFEDv2 emissions were used for the model  
spin up. The FLAMBE inventory overestimated CO emissions from fires in this region  
385 (Alvarado et al., 2010). In contrast, we find that CO is under-predicted using GFEDv3  
monthly average emissions.

We do not aim to optimize the ability of the model to simulate these specific  
plumes, but rather to show the magnitude of the changes with respect to this well-studied  
set of plumes. Chen et al. (2009) found that switching from monthly to 8-day time  
390 intervals for GFEDv2 in GEOS-Chem had the largest effect on simulating measured day-  
to-day variability in CO for boreal fires during the 2004 fire season. So it is possible that  
this approach (or alternatively using daily or 3-hourly fire fractions) would also improve  
the ability of the model to capture these specific plumes. However, both the daily or 3-  
hourly emissions inventories in our case are still likely to be an underestimate of the true  
395 emissions. Thus, we did not pursue these options. However, to simply show the impact of  
changing the emission factors, we include an additional simulation (pink line in Figure  
7d) with both the updated injection scheme and increased emissions of CO (factor of 2  
for extra-tropical fires and 1.5 for savannahs) following Petrenko et al. (2017), which has  
successfully reproduced the satellite observations of AOD with a series of adjustments to  
400 biomass burning emissions. We also include results from a standard model simulation  
with increased CO emissions (green line in Figure 7d). The green line indicates that this  
model configuration substantially increases the CO mixing ratios within the boundary  
layer as expected. Comparing this simulation (green line in Figure 7d) to the simulation  
incorporating both new injection scheme and increased CO emissions (pink line in Figure  
405 7d), shows the impact of both changes. By comparison, CO is higher at levels above the  
boundary layer and slightly lower in the boundary layer.

### **3.2 Amazon Basin Comparison**

As discussed in Section 2.5.2, we compare the simulated CO profiles to observed  
CO profiles at four Amazon basin sites in each month during 2010 and 2011. We note

410 that we evaluated GEOS-Chem over the Amazon with observations collected in different  
years than the MISR plume height data used to develop the parameterization. We made  
this choice because 2010-2011 CO profiles are available for use in the model-  
measurement comparison, and the MISR smoke-plume-height climatology from 2005-  
415 2012 shows little inter-annual variability over this region (Gonzalez-Alonso et al., in  
preparation). Where the data and the model are clearly smoke-impacted, a simulation that  
includes both the new injection scheme and increased CO emissions does improve the  
simulated CO profiles over this region. Figure 8 shows the total emitted percent of  
biomass burning emission injected above 700 hPa over the Amazon from March to  
November based on the MISR data. We note that 700 hPa is above the boundary layer in  
420 this location. The boundary layer in this region typically extends to 1200 – 1500 m above  
ground level in the morning to early afternoon, which corresponds to 880 – 840 hPa.  
From March to August, the total emitted percent in each grid cell above 700 hPa over the  
Amazon area is generally <15%. Thus, the new injection scheme does not produce a large  
difference in simulated CO profiles compared to the baseline simulation. However, the  
425 total emitted percent above 700 hPa in each grid cell is generally >25% from September  
to November. Figure 9b shows the vertical profiles of the emitted percent of biomass  
burning smoke from the standard model and the new injection scheme at the RBA site,  
for one case in October. Although peak emitted percentages in both simulations are near  
the top of the boundary layer, the new injection scheme has the emissions pushed higher  
430 in the atmosphere. Figure 9c shows comparison of the simulations with the corresponding  
bi-weekly observations at RBA in October of 2010 and 2011. Above the lowermost km,  
the simulated CO from the standard model generally under predicts the observed CO  
mixing ratios. The new injection scheme decreases CO mixing ratios in the boundary  
layer (by up to 45 ppb) and increases CO mixing ratios in troposphere (by up to 12 ppb).  
435 CO mixing ratios are 20 – 75 ppb lower than the RBA observations. With the increased  
CO emissions (pink), simulated CO mixing ratios near the surface provide a better match  
to the observations than the other two model versions. Away from the surface the  
simulation that includes both increased CO and the new injection scheme also performs  
better than the other two model versions, but the model still under predicts CO mixing  
440 ratios in this region of the atmosphere by ~50 ppb.

We also compared the model output to the CO mixing ratio profiles over the other three sites (TAB, ALF and SAN), and the impact of new injection scheme on CO mixing ratios at all levels over all three locations is small. Consistent with Andreae et al. (2012), we also found that the simulated CO mixing ratios are generally under-predicted in all months, especially during the biomass burning seasons. For example, the simulated CO mixing ratios are almost three times lower than observations in September at the SAN site. Gatti et al. (2014) found an emission ratio of 72.8 ppb CO/CO<sub>2</sub> ppm. For comparison the emission ratios used in GFEDv3 as implemented in GEOS-Chem are 97.5 and 59.5 ppb CO/CO<sub>2</sub> ppm for deforestation and savannas respectively. It is possible that either the emission factors themselves may be too low in GFEDv3 or there are fires missing from the inventory, so redistributing them in the atmosphere is not sufficient to better simulate their impact on atmospheric composition.

### 3.3 Averaged Impacts on CO

As the case studies of individual plumes presented in Sections 3.1 and 3.2 show, injecting the emissions of boreal fires higher in the atmosphere often increases the CO mixing ratio in the mid-troposphere above and directly downwind of the fire. For the 4 July smoke plume from ARCTAS (Fig. 7), the new model substantially reduces CO mixing ratios near the surface and at 850 hPa. There is an increase in CO at 510 hPa directly above and directly downwind of the fire as compared to the standard model (Figure 7a). However, Figure 7a also shows a decrease in CO at 510 hPa over much, but not all, of the domain. When viewed hemispherically, the net effect of lofting emissions out of the boundary layer is to produce lower average CO mixing ratios in the mid-upper troposphere because the average lifetime of CO against oxidation by OH is slightly shorter. Annual and globally averaged concentrations of OH increase slightly with altitude from 1000 hPa to 700 hPa (Spivakovsky et al., 2000). Thus when a fraction of the CO emissions are immediately moved out of the boundary layer, this fraction reacts more quickly with OH than in the standard simulation. The same issue applies throughout the atmosphere, and can be visualized for the Amazon region in Figure 9c. The CO mixing ratio decreases with altitude above 650 hPa at a faster rate in the simulation with the new injection scheme than in the standard model. This effect is not local to a given fire, but reflects the cumulative impact of changing the emission altitude for a substantial

quantity of CO emissions. In our model, the resulting changes to monthly mean CO are not large, but there is quite a bit of variability by season and location. Typical monthly mean decreases in CO mixing ratios away from freshly injected biomass burning plumes are <5% in the mid to upper troposphere. The changes in CO between model versions reflects changed injection heights throughout the Northern Hemisphere, not just the fire producing the particular smoke sampled by the aircraft this day. The response of CO is very different than that of PAN. The main loss of PAN is via thermal decomposition, so injecting PAN (or its precursors) higher in the atmosphere will increase PAN in the mid-  
475 to-upper troposphere. Though we highlight CO and PAN here as examples, injection height will impact the chemical evolution of nearly all species emitted from fires.

Figure 10 shows a comparison of our different model versions to monthly mean surface CO mixing ratios from four sites where there are substantial changes in 2008 monthly mean simulated CO with the new injection scheme. The decreases in simulated surface CO can be substantial when the emissions are moved up higher in the atmosphere based on the MISR analysis. Figure 10a and 10b indicate that the standard model over  
485 predicts July 2008 surface CO mixing ratios at two California monitoring sites, Trinidad Head and Point Arena. There were hundreds of wildfires in Northern California in June and July 2008 (Gyawali et al., 2009; Brey et al., 2018). The model with the improved injection height parameterization removes a large CO peak in July that is clearly not  
490 present in the surface observations. The lower panels of Figure 10 indicate that the model overpredicts surface CO abundances during much of the year at these two sites in the Southern Hemisphere, Bukit Kototabang (BKT), Indonesia and Cape Grim (CGO), Tasmania. However, the updated version of the model does reduce the model-  
495 measurement discrepancy at BKT between March and September 2008 by ~50%.

#### **4 Summary**

This paper introduces the development and implementation of a new global biomass burning emissions injection scheme in the GEOS-Chem model. The injection scheme is based on a MISR plume-injection-height climatology for 2008. This  
500 climatology was derived from space-based, multi-angle imagery. Additional (i.e. based on other datasets) or updated (i.e. other years) gridded climatologies of injection height could be implemented with relatively little effort given the code infrastructure that is now

in place. We have completed multi-year simulations with the new injection scheme and compared the model output to three smoke-impacted observational datasets.

505           Based on MISR snapshots, the percentage of total column biomass burning emissions that are typically injected above the boundary layer is relatively high for North American boreal regions. We find that the updated model is better able to simulate observed daytime observed vertical profiles of PAN and CO over boreal regions during the 2008 summer fire season, and including a better representation of injection height is  
510 likely very important for predicting the transport and chemical evolution of smoke plumes originating in this region. However, the version of GEOS-Chem used here has a persistent low bias in CO throughout the atmospheric column. Though our injection height climatology is based on observations from 2008, we also used this to simulate October 2010 and 2011 for the Amazon region. We made this choice because this season  
515 provided access to CO profiles that could be used for model-measurement comparison, and for this region, smoke injection heights do not appear to vary much interannually.

          In testing our model updates, we consistently found that it was important to do model-observation comparisons on specific biomass burning plumes with well-sampled vertical structure. When the model is sampled to match observations with less vertical  
520 information (e.g., MOPITT CO or TES PAN retrievals), the differences between the simulations appeared very small. However, when the model is compared to specific plumes, an improved injection height does produce notable differences in the simulations that can have air quality and possibly climate implications (see also Vernon et al. (2018)). Thus moving forward, we recommend that simulations with improved vertical injection  
525 height schemes for biomass burning plumes be compared to specific plumes, rather than larger-scale observations.

          It is important to note that the MISR plume heights that form the basis for our injection scheme are only snapshots. MISR is in a sun-synchronous orbit, and it crosses the equator at 10:30 local time. Actual wildfire smoke injection heights vary diurnally  
530 and less predictably hour-to-hour or day-to-day as burning progresses. Our scheme provides one consistent, statistically based injection height for each month; however, the ARCTAS aircraft observations also represent daytime measurements. A future development may be to attempt to anchor the model plume height at the MISR overpass

time, rather than assuming a constant plume height. A better comparison would include  
535 plumes observed throughout the diurnal cycle.

Though these model developments offer clear improvements under some  
situations, limitations in this approach should be noted. Most importantly, the MISR  
climatology that underpins this model development is based on snapshots of injection  
height. Thus it may not apply to all fires in a given location at all times of day. The  
540 MISR plume height climatology also may not represent the injection height of small fires  
as well as it does for larger ones. We expect that this approach will be most appropriate in  
regions where the total smoke emissions are dominated by fires large enough to be  
observed by the satellite instrument. However, most small fires inject only into the  
boundary layer, so if the *amount* of small-fire smoke is available, its vertical distribution  
545 can be assumed with some confidence.

**Data Availability:** The GEOS-Chem code used to generate this paper has already been  
passed to the GEOS-Chem model support team and we currently plan to include it as an  
option in the next public version of the model. The anticipated release date will be prior  
550 to publication. The code and data used for this study are published as supplemental  
materials and can be directly applied to the work based on GEOS-Chem v9-01-01. The  
aircraft and surface data used in this paper is already publically available.

**Appendices:** N/A

**Supplemental Information:** N/A

555 **Team List and Author Contributions:**

**Liye Zhu** led the majority of the analysis associated with this manuscript.

**Maria Val Martin** led the analysis of the MISR data and developed the monthly average  
gridded climatology of plume heights for 2008.

**Arsineh Hecobian** provided the smoke designation associated with the ARCTAS aircraft  
560 data.

**Luciana V. Gatti** led the aircraft measurements over the Amazon Basin.

**Ralph Kahn** helped develop the MISR plume-height algorithm, and led or mentored  
much of its application to wildfire smoke and volcanic plumes.

**Emily V. Fischer** led the conception of the work and the writing of this manuscript.

565 **Competing interests:** The authors declare that they have no conflict of interest.

**Disclaimer:** N/A

**Special issue statement:** N/A

**Acknowledgements** This work was supported by NASA Award Numbers NNX14AF14G and NNX14AN47G. PAN data from ARCTAS was provided  
570 by Greg Huey supported by NASA Award Number NNX08AR67G. Amazon vertical profile data were provided by L.V. Gatti supported by NERC (NE/F005806/1) and FAPESP (08/58120-3). We thank Glenn Diskin for the use of the ARCTAS CO data. We thank Paul C. Novelli for the use of the CO data from NOAA ESRL Carbon Cycle  
575 Cooperative Global Air Sampling Network. Maria Val Martin was partially supported by the Leverhulme Trust through a Leverhulme Research Centre Award (RC-2015-029). Ralph Kahn is supported in part by NASA's Climate and Radiation Research and Analysis Program under H. Maring, and NASA's Atmospheric Composition Program under R. Eckman

580

## References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039-4072,  
585 10.5194/acp-11-4039-2011, 2011.
- Alvarado, M. J., Logan, J. A., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R. C., Min, K. E., Perring, A. E., Browne, E. C., Wooldridge, P. J., Diskin, G. S., Sachse, G. W., Fuelberg, H., Sessions, W. R., Harrigan, D. L., Huey, G., Liao, J., Case-Hanks, A., Jimenez, J. L., Cubison, M. J., Vay, S. A., Weinheimer, A. J., Knapp, D. J., Montzka, D.  
590 D., Flocke, F. M., Pollack, I. B., Wennberg, P. O., Kurten, A., Crounse, J., Clair, J. M. S., Wisthaler, A., Mikoviny, T., Yantosca, R. M., Carouge, C. C., and Le Sager, P.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmos. Chem. Phys.*, 10, 9739-9760, 10.5194/acp-10-9739-2010, 2010.
- 595 Andreae, M. O., Artaxo, P., Beck, V., Bela, M., Freitas, S., Gerbig, C., Longo, K., Munger, J. W., Wiedemann, K. T., and Wofsy, S. C.: Carbon monoxide and related trace gases and aerosols over the Amazon Basin during the wet and dry seasons, *Atmos. Chem. Phys.*, 12, 6041-6065, 10.5194/acp-12-6041-2012, 2012.
- Arnold, S. R., Emmons, L. K., Monks, S. A., Law, K. S., Ridley, D. A., Turquety, S.,  
600 Tilmes, S., Thomas, J. L., Bouarar, I., Flemming, J., Huijnen, V., Mao, J., Duncan, B. N., Steenrod, S., Yoshida, Y., Langner, J., and Long, Y.: Biomass burning influence on high-latitude tropospheric ozone and reactive nitrogen in summer 2008: a multi-model analysis based on POLMIP simulations, *Atmos. Chem. Phys.*, 15, 6047-6068, 10.5194/acp-15-6047-2015, 2015.

- 605 Boy, J., Rollenbeck, R., Valarezo, C., and Wilcke, W.: Amazonian biomass burning-derived acid and nutrient deposition in the north Andean montane forest of Ecuador, *Global Biogeochemical Cycles*, 22, n/a-n/a, 10.1029/2007GB003158, 2008.
- Brey, S. J., Ruminski, M., Atwood, S. A., and Fischer, E. V.: Connecting smoke plumes to sources using Hazard Mapping System (HMS) smoke and fire location data over North  
610 America, *Atmos. Chem. Phys. Discuss.*, 2017, 1-29, 10.5194/acp-2017-245, 2017.
- Brey, S. J., Ruminski, M., Atwood, S. A., and Fischer, E. V.: Connecting smoke plumes to sources using Hazard Mapping System (HMS) smoke and fire location data over North America, *Atmos. Chem. Phys.*, 18, 1745-1761, 10.5194/acp-18-1745-2018, 2018.
- Chen, Y., Li, Q., Randerson, J. T., Lyons, E. A., Kahn, R. A., Nelson, D. L., and Diner,  
615 D. J.: The sensitivity of CO and aerosol transport to the temporal and vertical distribution of North American boreal fire emissions, *Atmos. Chem. Phys.*, 9, 6559-6580, 10.5194/acp-9-6559-2009, 2009.
- Chen, Y., Morton, D. C., Jin, Y., Collatz, G. J., Kasibhatla, P. S., van der Werf, G. R.,  
620 DeFries, R. S., and Randerson, J. T.: Long-term trends and interannual variability of forest, savanna and agricultural fires in South America, *Carbon Management*, 4, 617-638, 10.4155/cmt.13.61, 2013.
- Colarco, P. R., Schoeberl, M. R., Doddridge, B. G., Marufu, L. T., Torres, O., and  
Welton, E. J.: Transport of smoke from Canadian forest fires to the surface near Washington, D.C.: Injection height, entrainment, and optical properties, *Journal of  
625 Geophysical Research: Atmospheres*, 109, n/a-n/a, 10.1029/2003JD004248, 2004.
- Damoah, R., Spichtinger, N., Forster, C., James, P., Mattis, I., Wandinger, U., Beirle, S.,  
Wagner, T., and Stohl, A.: Around the world in 17 days - hemispheric-scale transport of forest fire smoke from Russia in May 2003, *Atmos. Chem. Phys.*, 4, 1311-1321, 10.5194/acp-4-1311-2004, 2004.
- 630 Diner, D. J., Beckert, J. C., Reilly, T. H., Bruegge, C. J., Conel, J. E., Kahn, R. A., Martonchik, J. V., Ackerman, T. P., Davies, R., Gerstl, A. W., Gordon, H. R., Muller, J.-P., Myneni, R. B., Sellers, P. J., Pinty, B., and Verstraete, M. M.: Multi-angle imaging spectroradiometer (MISR) instrument description and experiment overview, *IEEE Trans. Geosci. Remote Sens.*, 36, 0196-2892(98)04161-8, 1998.
- 635 Emmons, L. K., Arnold, S. R., Monks, S. A., Huijnen, V., Tilmes, S., Law, K. S., Thomas, J. L., Raut, J. C., Bouarar, I., Turquety, S., Long, Y., Duncan, B., Steenrod, S., Strode, S., Flemming, J., Mao, J., Langner, J., Thompson, A. M., Tarasick, D., Apel, E. C., Blake, D. R., Cohen, R. C., Dibb, J., Diskin, G. S., Fried, A., Hall, S. R., Huey, L. G., Weinheimer, A. J., Wisthaler, A., Mikoviny, T., Nowak, J., Peischl, J., Roberts, J. M.,  
640 Ryerson, T., Warneke, C., and Helmig, D.: The POLARCAT Model Intercomparison Project (POLMIP): overview and evaluation with observations, *Atmos. Chem. Phys.*, 15, 6721-6744, 10.5194/acp-15-6721-2015, 2015.




- 645 Fischer, E. V., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Millet, D. B., Mao, J., Paulot, F., Singh, H. B., Roiger, A., Ries, L., Talbot, R. W., Dzepina, K., and Pandey Deolal, S.: Atmospheric peroxyacetyl nitrate (PAN): a global budget and source attribution, *Atmos. Chem. Phys.*, 14, 2679-2698, 10.5194/acp-14-2679-2014, 2014.
- 650 Fisher, J. A., Jacob, D. J., Purdy, M. T., Kopacz, M., Le Sager, P., Carouge, C., Holmes, C. D., Yantosca, R. M., Batchelor, R. L., Strong, K., Diskin, G. S., Fuelberg, H. E., Holloway, J. S., Hyer, E. J., McMillan, W. W., Warner, J., Streets, D. G., Zhang, Q., Wang, Y., and Wu, S.: Source attribution and interannual variability of Arctic pollution in spring constrained by aircraft (ARCTAS, ARCPAC) and satellite (AIRS) observations of carbon monoxide, *Atmos. Chem. Phys.*, 10, 977-996, 10.5194/acp-10-977-2010, 2010.
- 655 Forster, C., Wandering, U., Wotawa, G., James, P., Mattis, I., Althausen, D., Simmonds, P., O'Doherty, S., Jennings, S. G., Kleefeld, C., Schneider, J., Trickl, T., Kreipl, S., Jäger, H., and Stohl, A.: Transport of boreal forest fire emissions from Canada to Europe, *Journal of Geophysical Research: Atmospheres*, 106, 22887-22906, 10.1029/2001JD900115, 2001.
- 660 Freitas, S. R., Longo, K. M., and Andreae, M. O.: Impact of including the plume rise of vegetation fires in numerical simulations of associated atmospheric pollutants, *Geophysical Research Letters*, 33, n/a-n/a, 10.1029/2006GL026608, 2006.
- Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, A. M.: MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sens. Environ.*, 168-182, 10.1016/j.rse.2009.08.016, 2010.
- 665 Fuelberg, H. E., Harrigan, D. L., and Sessions, W.: A meteorological overview of the ARCTAS 2008 mission, *Atmos. Chem. Phys.*, 10, 817-842, 10.5194/acp-10-817-2010, 2010.
- 670 Gatti, L. V., Gloor, M., Miller, J. B., Doughty, C. E., Malhi, Y., Domingues, L. G., Basso, L. S., Martinewski, A., Correia, C. S. C., Borges, V. F., Freitas, S., Braz, R., Anderson, L. O., Rocha, H., Grace, J., Phillips, O. L., and Lloyd, J.: Drought sensitivity of Amazonian carbon balance revealed by atmospheric measurements, *Nature*, 506, 76, 10.1038/nature12957, 2014.
- 675 Gonzi, S., Palmer, P. I., Paugam, R., Wooster, M., and Deeter, M. N.: Quantifying pyroconvective injection heights using observations of fire energy: sensitivity of spaceborne observations of carbon monoxide, *Atmos. Chem. Phys.*, 15, 4339-4355, 10.5194/acp-15-4339-2015, 2015.
- 680 Gyawali, M., Arnott, W. P., Lewis, K., and Moosmüller, H.: In situ aerosol optics in Reno, NV, USA during and after the summer 2008 California wildfires and the influence of absorbing and non-absorbing organic coatings on spectral light absorption, *Atmos. Chem. Phys.*, 9, 8007-8015, 10.5194/acp-9-8007-2009, 2009.


- 685 Hecobian, A., Liu, Z., Hennigan, C. J., Huey, L. G., Jimenez, J. L., Cubison, M. J., Vay, S., Diskin, G. S., Sachse, G. W., Wisthaler, A., Mikoviny, T., Weinheimer, A. J., Liao, J., Knapp, D. J., Wennberg, P. O., Kürten, A., Crouse, J. D., Clair, J. S., Wang, Y., and Weber, R. J.: Comparison of chemical characteristics of 495 biomass burning plumes intercepted by the NASA DC-8 aircraft during the ARCTAS/CARB-2008 field campaign, *Atmos. Chem. Phys.*, 11, 13325-13337, 10.5194/acp-11-13325-2011, 2011.
- 690 Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A., Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results, *Atmos. Chem. Phys.*, 10, 5191-5212, 10.5194/acp-10-5191-2010, 2010.
- 695 Jian, Y., and Fu, T. M.: Injection heights of springtime biomass-burning plumes over peninsular Southeast Asia and their impacts on long-range pollutant transport, *Atmos. Chem. Phys.*, 14, 3977-3989, 10.5194/acp-14-3977-2014, 2014.
- Kahn, R. A., Chen, Y., Nelson, D. L., Leung, F.-Y., Li, Q., Diner, D. J., and Logan, J. A.: Wildfire smoke injection heights: Two perspectives from space, *Geophysical Research Letters*, 35, n/a-n/a, 10.1029/2007GL032165, 2008.
- 700 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527-554, 10.5194/bg-9-527-2012, 2012.
- 705 Leung, F. T., Logan, J. A., Park, R., Hyer, E., Kasischke, E., Streets, D., and Yurganov, L.: Impacts of enhanced biomass burning in the boreal forests in 1998 on tropospheric chemistry and the sensitivity of model results to the injection height of emissions, *Journal of Geophysical Research: Atmospheres*, 112, doi:10.1029/2006JD008132, 2007.
- 710 Lewis, A. C., Evans, M. J., Hopkins, J. R., Punjabi, S., Read, K. A., Purvis, R. M., Andrews, S. J., Moller, S. J., Carpenter, L. J., Lee, J. D., Rickard, A. R., Palmer, P. I., and Parrington, M.: The influence of biomass burning on the global distribution of selected non-methane organic compounds, *Atmos. Chem. Phys.*, 13, 851-867, 10.5194/acp-13-851-2013, 2013.
- 715 Ichoku, C. and Kaufman, Y.J. A method to derive smoke emission rates from MODIS fire radiative energy measurements. *IEEE transactions on Geoscience and Remote Sensing* 2005, 43, 2636–2649.
- Mims, S. R., Kahn, R. A., Moroney, C. M., Gaitley, B. J., Nelson, D. L., and Garay, M. J.: MISR Stereo-heights of grassland fire smoke plumes in Australia, *IEEE Trans. Geosci. Remote Sens.*, 48, 10.1109/TGRS.2009.2027114, 2010.

- 720 Moxim, W. J., Levy, H., and Kasibhatla, P. S.: Simulated global tropospheric PAN: Its transport and impact on NO<sub>x</sub>, *Journal of Geophysical Research: Atmospheres*, 101, 12621-12638, 10.1029/96JD00338, 1996.
- Nelson, L. D., Garay, J. M., Kahn, A. R., and Dunst, A. B.: Stereoscopic Height and Wind Retrievals for Aerosol Plumes with the MISR Interactive eXplorer (MINX), *Remote Sensing*, 5, 10.3390/rs5094593, 2013.
- 725 Novelli, P. C., Masarie, K. A., Lang, P. M., Hall, B. D., Myers, R. C., and Elkins, J. W.: Reanalysis of tropospheric CO trends: Effects of the 1997–1998 wildfires, *Journal of Geophysical Research: Atmospheres*, 108, 10.1029/2002JD003031, 2003.
- Novelli, P. C., Crotwell, A., Lang, P. M., and Mund, J.: Atmospheric Carbon Monoxide Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1988-2016, Version: 2017-07-28, 2017.
- 730 Paugam, R., Wooster, M., Freitas, S., and Val Martin, M.: A review of approaches to estimate wildfire plume injection height within large-scale atmospheric chemical transport models, *Atmos. Chem. Phys.*, 16, 907-925, 10.5194/acp-16-907-2016, 2016.
- Petrenko, M., Kahn, R., Chin, M., and Limbacher, J.: Refined Use of Satellite Aerosol Optical Depth Snapshots to Constrain Biomass Burning Emissions in the GOCART Model, *Journal of Geophysical Research: Atmospheres*, 122, 10,983-911,004, 10.1002/2017JD026693, 2017.
- 735 Randerson, J. T., Chen, Y., Werf, G. R., Rogers, B. M., and Morton, D. C.: Global burned area and biomass burning emissions from small fires, *J. Geophys. Res.*, 117, 10.1029/2012JG002128, 2012.
- 740 Reid, J. S., Hyer, E. J., Prins, E. M., Westphal, D. L., Wang, J., Christopher, S. A., Curtis, C. A., Schmidt, C. C., Eleuterio, D. P., Richardson, K. A., and Hoffman, J. P.: Global Monitoring and Forecasting of Biomass-Burning Smoke: Description of and Lessons From the Fire Locating and Modeling of Burning Emissions (FLAMBE) Program, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 2, 144-162, 2009.
- 745 Singh, H. B., and Hanst, P. L.: Peroxyacetyl nitrate (PAN) in the unpolluted atmosphere: An important reservoir for nitrogen oxides, *Geophysical Research Letters*, 8, 941-944, 10.1029/GL008i008p00941, 1981.
- 750 Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B. A., Horowitz, L. W., Fusco, A. C., Brenninkmeijer, C. A. M., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Three - dimensional climatological distribution of tropospheric OH: Update and evaluation, *Journal of Geophysical Research: Atmospheres*, 105, 8931-8980, 10.1029/1999JD901006, 2000.

- 755 Stein, A. F., Rolph, G. D., Draxler, R. R., Stunder, B., and Ruminski, M.: Verification of the NOAA Smoke Forecasting System: Model Sensitivity to the Injection Height, *Weather and Forecasting*, 24, 379-394, 10.1175/2008WAF2222166.1, 2009.
- Stocks, B. J., Mason, J. A., Todd, J. B., Bosch, E. M., Wotton, B. M., Amiro, B. D., Flannigan, M. D., Hirsch, K. G., Logan, K. A., Martell, D. L., and Skinner, W. R.: Large forest fires in Canada, 1959–1997, *Journal of Geophysical Research: Atmospheres*, 107, FFR 5-1-FFR 5-12, 10.1029/2001JD000484, 2002.
- 760
- Tosca, M. G., Randerson, J. T., Zender, C. S., Nelson, D. L., Diner, D. J., and Logan, J. A.: Dynamics of fire plumes and smoke clouds associated with peat and deforestation fires in Indonesia, *Journal of Geophysical Research: Atmospheres*, 116, n/a-n/a, 10.1029/2010JD015148, 2011.
- 765
- Val Martin, M., Logan, J. A., Kahn, R. A., Leung, F. Y., Nelson, D. L., and Diner, D. J.: Smoke injection heights from fires in North America: analysis of 5 years of satellite observations, *Atmos. Chem. Phys.*, 10, 1491-1510, 10.5194/acp-10-1491-2010, 2010.
- Val Martin, M., Kahn, R. A., Logan, J. A., Paugam, R., Wooster, M., and Ichoku, C.: Space-based observational constraints for 1-D fire smoke plume-rise models, *Journal of Geophysical Research: Atmospheres*, 117, n/a-n/a, 10.1029/2012JD018370, 2012.
- 770
- Val Martín, M., Honrath, R. E., Owen, R. C., Pfister, G., Fialho, P., and Barata, F.: Significant enhancements of nitrogen oxides, black carbon, and ozone in the North Atlantic lower free troposphere resulting from North American boreal wildfires, *Journal of Geophysical Research: Atmospheres*, 111, n/a-n/a, 10.1029/2006JD007530, 2006.
- 775
- Val Martin, M., Kahn, R. A., and Tosca, M. (2018), A Global Climatology of Wildfire Smoke Injection Height Derived from Space-based Multi-angle Imaging, submitted to *Remote Sensing (MISR Special Issue)*, manuscript ID remotesensing-359296.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707-11735, 10.5194/acp-10-11707-2010, 2010.
- 780
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth Syst. Sci. Data*, 9, 697-720, 10.5194/essd-9-697-2017, 2017.
- 785
- Vernon, C. J., Bolt, R., Canty, T., and Kahn, A. R.: The impact of MISR-derived injection height initialization on wildfire and volcanic plume dispersion, *Atmos. Chem. Phys.*, submitted, 2018.
- 790




795 **Figure 1:** Overview of the implementation of an observationally based scheme to inject biomass burning emissions within GEOS-Chem.



800 **Figure 2:** (a) Vertical profile of the percent of emissions in each model level for a sample location over boreal Canada (56°N, 105°W) from the public release version of GEOS-Chem (blue) and the new observationally-based injection scheme (red). The dashed line indicates the averaged boundary layer top of this month. The solid black line is at 850

hPa, corresponding to the layer shown in (b). (b) Percent of total column biomass burning emissions emitted into the 850 hPa layer in each model grid cell for July 2008.


805



810


**Figure 3:** (a) Percentage of total column biomass burning emissions injected above 700 hPa over North America for July 2008, based on MISR observations. The two example locations shown in (b) and (c) are marked as blue stars. (b) Vertical profile of the percent of emissions in each model level over 56°N, 105°W. (c) Vertical profile of the percent of emissions in each model level over 40°N, 82.5°W. The dashed line indicates the averaged boundary layer top during this month.

815



**Figure 4:** (a)-(c) Differences in simulated PAN mixing ratios between a GEOS-Chem simulation with and without the new observationally based biomass burning injection scheme over North America at three different levels: 510 hPa, 850 hPa, and surface for July 1, 2008. (d) Median vertical profiles of ARCTAS PAN mixing ratios (black), standard model (blue), and new injection scheme (red) on July 1, 2008. The whiskers represent 25% and 75% percentiles of the data in the pressure bins. The numbers on the left are the numbers of observations in different pressure bins. (e) ARCTAS in situ


aircraft observations for July 1, 2008 colored by ambient pressure for the inset black box in (a) - (c). (f) ARCTAS observations for July 1, 2008 colored by PAN mixing ratio.




**Figure 5:** Same as in Figure 4, but for CO. The pink profiles are from a simulation that also increased the emissions of CO from boreal fires as described in Section 2.4. The green profiles are from a simulation of standard model with increased CO emissions as the pink profiles.

830







835 **Figure 6:** Same as Figure 4, but for July 4, 2008.




**Figure 7:** Same as in Figure 5, but for July 4, 2008. The pink profiles are from a simulation that also increased the emissions of CO from boreal fires as described in Section 2.4. The green profiles are from a simulation of standard model with increased CO emissions as the pink profiles.



845 **Figure 8:** Percentage of total column biomass burning emissions injected above 700 hPa over the Amazon from March to November of 2010 and 2011, based on MISR plume-height analysis.



850 **Figure 9:** (a) Map of four measurement sites in the Amazon basin. (b) Vertical profile of  
the percent of emissions in each model level at site RBA from the public release version  
of GEOS-Chem (blue) and the new observationally-based injection scheme (red). The  
dashed line indicates the averaged boundary layer top during this month. (c) Median  
vertical profiles of CO mixing ratios observed at RBA (black), simulated with the  
855 standard model (blue), simulated with the new injection scheme (red), and simulated with  
the new injection scheme and with increased CO (Petrenko et al., 2017) in October of  
2010 and 2011.



860 **Figure 10:** Observed and simulated monthly mean CO mixing ratios at select NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network sites.