

Response to referee comments

The revised manuscript with track changes is provided below.

Anonymous Referee #4

Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)

I very much appreciate the development and implantation of a new individual tree model that is physiology based and can be applied over periods that allow to judge the development of species mixtures under changing environmental conditions. I feel however that considerable improvements in the model description, presentation and discussion could still be achieved.

Model description

The model description in general would also benefit from a better explanation about which processes are run in which time steps. While photosynthesis is calculated hourly, time steps for allocation are not explicitly addressed. They may be done annually, similarly to dimensional calculations. If this is true (I couldn't find any statement about it), it raises some yet unexplained questions: Where is the foliage compartment outside the vegetation period and how does it respire? Is the biomass at the end of the year abruptly changing or are empirical functions used to distribute the growth over the year? Shouldn't the fact that biomass is actually changing during the year be reflected in the maintenance respiration calculations? If, however, allocation is done hourly or daily, how is the first leaf flushing justified in deciduous trees when there is not yet any photosynthesis? In particular since there seems to be no reserve compartment. The best model description is in the caption of figure 1 but I think this (and more) information needs to be in the text too.

Author response (R):

We reviewed the section 2.1 of the paper "Overall operation of the HETEROFOR model" and we specified the time step of each process. Carbon allocation is done once a year at the end of the vegetation period which allows to update tree dimensions for the next yearly time step during which tree size does not change. Outside the vegetation period, the foliage compartment is not considered and does not respire. The small respiration rate of buds during the dormant period is therefore not explicitly accounted but is indirectly taken into account by the calibration of the respiration coefficient.

I also noticed that in many equations, parameters are given as alpha, beta, gamma and delta letters without further indication - which is a bit strange and sincerely against common rules since this kind of notation is not specific. In my opinion, also empirical parameters (including a, b, c...) should be unmistakably identifiable and should also be given and explained (including sources) in a table somewhere.

R:

The value of these parameters is given in Table 2. We recognize however that this information did not appear sufficiently clearly. In the revised Table 2, we clearly identify the parameters, their value and the associated equation (changes highlighted in red).

Model demonstration and discussion

I appreciate that examples for demonstrating the model's performance are based on a number of stands that differ somewhat in their structure. I was a bit disappointed though that only different model approaches (such as different photosynthesis schemes) have been compared and no parameter sensitivity has been done. I admit, however, that this might be a very comprehensive demand and should possibly be restricted to completely new model formulations. On the other hand, the tests are still limited to two tree species and more or less only one climate area. If no other sites will be incorporated (which would be good but which is something I don't really expect), the issue should be carefully acknowledged as a limited demonstration of model performance.

R:

The objective of this paper is first to describe a new model and then to show that it gives interesting results in some case studies. We agree that this first model evaluation is limited and have better acknowledged this in the conclusion (P36L10-11). This first evaluation will be complemented by a larger one at the European scale.

In addition, existing literature should be better considered for the following topics: 1) the linear relationship between NPP and GPP has been recently discussed by Collalti et al. 2019; 2) the allometry depends on competition issues that need to be treated based on individual tree situations, which is a likely explanation for the relatively bad performance of increment calculations (see e.g. del Rio et al. 2019), 3) an hierarchical allocation approach that distributes carbon into different compartments has certain disadvantages, so that source-sink approaches or other approaches needs to be at least discussed as possible alternatives (e.g. Carl et al. 2018, Thurn et al. 2017). These points are particularly important in a single-tree model than in a stand model.

R:

The references suggested by the reviewer helped us to substantially improve the discussion. Collalti et al. (2019) was used to discuss the pros and cons of the *npp* to *gpp* ratio approach for modelling autotrophic respiration (P31L8-25). The way our model accounts for the impact of intra and inter-specific competition on tree allometry was analysed in the light of the study of Del Rio et al. (2019) (P31L34 to P32L3, P33L4-6). The study of Carl et al. (2018) allowed us to improve the discussion on the size-growth relationships and on the competition mode (P33L18-33) while improvement perspectives were opened thanks to Thurn et al. (2017) (P33L15-16).

Besides the evaluations which are basically based on growth measurements that are very integrated outcomes from many underlying processes, I would like to see some curves of how different compartments of different trees or groups of trees are developing over time in dependence on climatic input. Do we see any climate or management impact here? Are different groups have different strategies or are developing differently? Is this development reasonably at all? Please elaborate.

R:

We followed the reviewer suggestion and carry out one-year simulations to highlight the impact of climate conditions (comparison of a dry, normal and wet year) and of management (reduction of stand basal area by 25%) on biomass production and on its allocation to tree compartments. This new simulation experiment was described in the material and methods (P21L17-24), presented in the results (P26L6-16 + Fig. 5) and discussed (P33L7-16).

There are a couple of other more specific things that caught my eye and may be corrected or answered:

P3L13: Another argument why forestry trials are not suitable to derive estimates for future conditions are that these future conditions, i.e. CO₂ concentrations, simply cannot be observed in the present.

R:

Thank you for the suggestion! We have considered it in the revised version (P3L12).

P4L13ff: The BALANCE model is described here as having no soil layers and is missing basic soil processes. This is wrong, as can be easily derived from Figs. 3 and 6 in Grote and Pretzsch 2002. I admit that neither weathering nor nutrients other than nitrogen are included but still the model hosts a full carbon and nitrogen cycle and calculations in different soil layers, considering different nutrient availability with rooting depth. Please correct.

R:

In the revised version, we corrected the description of BALANCE which considers indeed different soil layers (P4L13-14).

P6L14: Does the model run when 'no meteorological measurements' (resp. scenario data) are provided as input? How can this work out reasonably?

R:

When no hourly meteorological measurements are available, the model can be run with a default empirical approach based on the radiation use efficiency approach. In this case, the only input needed by the model is the mean monthly global radiation. This more empirical approach provides good results for conditions similar to those in which the model was calibrated but do not allow extrapolations to future conditions. This option does not account for inter-annual variations in tree growth.

P7L12ff (also in Fig.1, and P9): GPP is either derived by a radiation efficiency approach or with the Farquhar model? This is irritating. There should be a rule when the one or other method is applied. Has this something to do with the age/size or position of the tree similar to the differentiation of respiration?

R:

The user can choose one of the two options (radiation efficiency approach or Farquhar model) depending on his objectives and on meteorological data availability. We rephrased it to avoid misunderstanding (P7L23-25, legend of Fig. 1).

P7L19: The expression 'compartment' or 'tree compartment' is used here and in other places but never defined. In P6L32 it is used for bark and foliage, P8L1 it is soil solution, at P12 it describes foliage and fine roots, and at P26 we learn that 'structural compartments' exist besides foliage and roots. The same holds for the expression 'organ' which is partly used as a synonym to 'compartment'. Figure 1 caption tells us about 'structural organs' that are 'roots, trunk, and branches' and also later 'organ' development is described with equations throughout P11 to P13 without telling the reader what is actually meant.

R:

In order to clarify the scientific terms, we defined 'tree compartment' in P6L11-12 and did not use anymore the term 'organ'. We also restricted the use of the word 'compartment' to 'tree compartment' and modified the sentence where it was used for the soil solution.

P9L7 (and figure 1): Fruits are very interesting but seldomly considered as a separate and dynamic tree compartment that is explicitly modelled. From Fig. 1 and the description in P13 it seems that it is considered but how data on this compartment can actually be used to modify allocation pattern, needs to be described further (taking the carbon from all other compartments equally or selectively?).

R:

As for the other functional tree compartments (foliage, fine roots), the carbon necessary for fruit production is subtracted from npp before the allocation to the structural tree compartments (Eq. 23).

P9L10: This indicates that photosynthesis is driven with daily meteorological data which for the Farquhar approach only makes sense if data are downscaled to hourly (or similar) time steps (also indicated in P10L18). Why is this not possible for the hydrological model then which explicitly requires hourly input?

R:

Hourly (and not daily) meteorological data are used to calculate photosynthesis with the Farquhar approach as mentioned in P11L18.

P10L14: Possible relationships between Farquhar photosynthesis and soil water restrictions have been implemented before (see e.g. Granier, Knauer, van Wijk, Wang in references). I guess, that also the new implementation will be based on one of them which could be indicated here. Otherwise, I fear that a short description of how this is done is indispensable for understanding the carbon assimilation dynamics.

R:

In HETEROFOR, the impact of soil water on Farquhar photosynthesis is described with a decreasing exponential function of the soil water potential. This relationship is presented in the companion paper (see Eq. 56 in de Wergifosse et al., in review).

P11L4ff: I am a bit puzzled about these explanations. Why is it bad that 'the crown to stem diameter ratio changes during the course of the tree development' so that the npp/gpp ratio also changes. Isn't this exactly what you want? And why is this index better although a relation of npp/gpp to this index has not been demonstrated?

R:

The crown to stem diameter ratio characterizes the tree shape reflecting past competition conditions but also changes during the course of the tree development. We standardized it to remove the size effect in order to obtain an index only dependent on the tree shape. This index is particularly useful to account for the large difference in oak crown extension according to the silvicultural system (large crowns in former coppices with standards vs narrow crowns in dense high forests). For oak, we observed that this index provides good results. However, the user can fix the parameter associated to this index to 0 for some tree species (we did for beech) or if he is not convinced by its added value. Fixing this parameter to 0 is equivalent to use a constant npp/gpp ratio.

We rephrase this paragraph in order to clarify the explanations (P12L1-11).

P11L16ff: Maintenance respiration is taken from photosynthesis (according to equation 7), but its calculation is independent of carbon gain. So, the net gain might get negative. What's happening then? Is there any reserve compartment?

R:

The net primary production (npp) is calculated once a year by subtracting maintenance respiration from gross primary production (Eq. 7) and no reserve compartment is considered. Contrary to what frequently happen when npp is calculated daily, the net gain is nearly always positive when calculated on an annual basis. If it turns out that the npp is negative due to exceptional conditions (highly outcompeted trees, water stress), the tree dies. Generally, when it happens, npp decreases progressively and tree decline takes several years.

P13L15/Eq.22: How is this parameterized and even how is the function itself derived? My question arises from the fact that fructification is highly variable and relates to specific climate conditions such as temperature in spring. Also, the parameter given in the appendix seems to be a variable based on this equation.

R:

The inter-annual variation in fruit production is not described by a specific routine. For past conditions, the user can provide an optional input file with the fruit production per year and tree species. For future or unknown conditions, mean value are used by default. The fruit distribution among trees depends on their size and on the light availability according to Eq. 22. In this equation, the parameter α takes a default value or is adapted based on the fruit production of the year (provided in the input file). In the future, we plan to develop a more sophisticated routine to predict the inter-annual variations in fruit production and to distribute it among trees.

In the revised version of the paper, we provided additional explanations on the fruit production modelling (P14L20-24)

P13L30ff/Eq.26-28: How could height and diameter growth be derived from the total above structural carbon gain if diameter and height growth only depend on stem (trunk) development? I would expect that the dimensional growth depends on db_stem rather than db_structural. Given

that the derivation of height and diameter increment is correct, has anybody checked if the biomass increase is actually supporting the dimensional change? Or does it imply a wood density change?

R:

Height and dbh growth are obtained from the aboveground structural biomass gain using an allometric equation predicting aboveground structural biomass from height and dbh (Eq. 26). To do so, this allometric equation was derived with respect to time and then rearranged (Eqs 27 and 28).

The increment in stem biomass is not used to calculate dbh and height since it is still unknown at this stage of the calculation. It is calculated once the dbh increment is known (Eq. 34). As the dimensional change is calculated based on the aboveground structural biomass gain, this biomass increase perfectly corresponds to the dimensional change, without implying any change in wood density. Carbon budgets were achieved to check that no calculation errors were made.

P14L30: Hounzandji et al. 2015 distinguish branches/twigs but not roots into three fractions of different diameter classes. The assumption that roots could be described with the same equations (and parameters?) without any additional data seems very bold to me. As it is, the description gives a false impression of what has been derived from literature. Furthermore, if a root fraction of 0-4 cm is assumed, it would include fine roots, which are, however, separately treated in the model. The differentiation is therefore inconsistent.

R:

The equations of Hounzandji et al. (2015) provide branch proportions and not absolute values. We considered that the mathematical forms of the equations is also suitable for roots as long as the parameters are adjusted with root data. Since we did not have such data, we temporarily used the same parameters than for the branches. However, these distributions in root categories has no impact on the functioning of the model since this information is not used elsewhere. These are just model outputs. If the user does not trust this categorization, he may just consider the whole root compartment without distinguishing it by category. As the proportions are applied to the root compartment, the fraction < 4 cm does not include the fine roots. This is the reason why it is called '< 4 cm' and not '0 - 4 cm'. In the revised version of the paper, we clarified it (P16L6-9).

P17L17ff: The reconstruction is a nice feature but depends on the knowledge about how many trees have died and when they died during the investigated period. Otherwise, npp is underestimated since it only refers to the remaining trees. Could you acknowledge this in the paragraph?

R:

In addition to repeated stand inventories, the reconstruction tool also requires a file listing the trees which were cut or died between the two inventory dates and the last year during which they were present in the stand. We added this information in the revised version (P19L26-28).

References

Carl, C., Biber, P., Veste, M., Landgraf, D. and Pretzsch, H. (2018). Key drivers of competition and growth partitioning among *Robinia pseudoacacia* L. trees. *Forest Ecol. Manage.* 430, 86-93.

Collalti, A. and Prentice, I. C. (2019). Is NPP proportional to GPP? Waring's hypothesis 20 years on. *Tree Physiol.* 39, 1473-1483.

del Río, M., Bravo-Oviedo, A., Ruiz-Peinado, R. and Condés, S. (2019). Tree allometry variation in response to intra- and inter-specific competitions. *Trees* 33, 121-138.

Granier, A., Loustau, D. and Bréda, N. (2000). A generic model of forest canopy conductance dependent on climate, soil water availability and leaf area index. *Annales des Sciences Forestières* 57, 755-765.

Knauer, J., Werner, C. and Zaehle, S. (2015). Evaluating stomatal models and their atmospheric drought response in a land surface scheme: A multibiome analysis. *J. Geophys. Res. - Biogeosci.* 120, 1894–1911

Thurm, E. A., Biber, P. and Pretzsch, H. (2017). Stem growth is favored at expenses of root growth in mixed stands and humid conditions for Douglas-fir (*Pseudotsuga menziesii*) and European beech (*Fagus sylvatica*). *Trees-Struct. Funct.* 31, 349-365.

Van Wijk, M. T., Dekker, S. C., Bouten, W., Bosveld, F. C., Kohsiek, W., Kramer, K., et al. (2000). Modeling daily gas exchange of a Douglas-fir forest: comparison of three stomatal conductance models with and without a soil water stress function. *Tree Physiol.* 20, 115-122.

Wang, Y.-P. and Leuning, R. (1998). A two-leaf model for canopy conductance, photosynthesis and partitioning of available energy I: Model description and comparison with a multi-layered model. *Agric. Forest Meteorol.* 91, 89-111.

HETEROFOR 1.0: a spatially explicit model for exploring the response of structurally complex forests to uncertain future conditions. I. Carbon fluxes and tree dimensional growth.

5

Mathieu Jonard¹, Frédéric André¹, François de Coligny², Louis de Wergifosse¹, Nicolas Beudez², Hendrik Davi³, Gauthier Ligot⁴, Quentin Ponette¹, Caroline Vincke¹

¹Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, 1348, Belgium

10 ²INRA, CIRAD, CNRS, IRD, Univ Montpellier, Montpellier, 34398 France

³Ecologie des Forêts Méditerranéennes (URFM), Institut National de la Recherche Agronomique (INRA), Avignon, 84914, France

⁴Gembloux Agro-Bio Tech, Université de Liège, Gembloux, 5030, Belgium

15 *Correspondence to:* Mathieu Jonard (mathieu.jonard@uclouvain.be)

Abstract.

Given the multiple abiotic and biotic stressors resulting from global changes, management systems and practices must be adapted in order to maintain and reinforce the resilience of forests. Among others, the transformation of monocultures into uneven-aged and mixed stands is an avenue to improve forest resilience. To explore the forest response to these new silvicultural practices under a changing environment, one needs models combining a process-based approach with a detailed spatial representation, which is quite rare.

We therefore decided to develop our own model (HETEROFOR for HETEROgeneous FORest) according to a spatially explicit approach describing individual tree growth based on resource sharing (light, water and nutrients). HETEROFOR was progressively elaborated through the integration of various modules (light interception, phenology, water cycling, photosynthesis and respiration, carbon allocation, mineral nutrition and nutrient cycling) within CAPSIS (Computer-Aided Projection for Strategies in Silviculture), a collaborative modelling platform devoted to tree growth and stand dynamics. The advantage of using such a platform is to use common development environment, model execution system, user-interface and visualization tools and to share data structures, objects, methods and libraries.

This paper describes the carbon-related processes of HETEROFOR (photosynthesis, respiration, carbon allocation and tree dimensional growth) and evaluates the model performances for three broadleaved stands of contrasted species composition (Wallonia, Belgium). This first evaluation showed that HETEROFOR predicts well individual radial growth (Person's correlation of 0.83 and 0.63 for European beech and sessile oak, respectively) and is able to reproduce size-growth relationships. We also noticed that the most empirical option for describing maintenance respiration provides the best results while the process-based and empirical approaches similarly perform for photosynthesis and crown extension. To illustrate how the model can be used to predict climate change impacts on forest ecosystems, the growth dynamics in this stand was simulated according to four IPCC climate scenarios. According to these simulations, the tree growth trends will be governed by the CO₂ fertilization effect with the increase in vegetation period length and in water stress also playing a role but offsetting each other.

1 Introduction

Forest structure and composition result from soil and climate conditions, management and natural disturbances. All these drivers of forest ecosystem functioning are rapidly evolving due to global changes (Aber et al., 2001; Lindner et al., 2010; Campioli et al., 2012). While environmental and societal changes make no doubt, their magnitude and the way they will occur locally remain largely uncertain (Lindner et al., 2014). Designing silvicultural systems and selecting tree species adapted to future conditions seems therefore a risky bet (Ennos et al., 2019). Messier et al. (2015) proposed another approach recognizing that forests are complex adaptive systems whose future dynamics is inherently uncertain. To maintain the ability of forests to provide a large range of goods and services whatever the future conditions, their resilience and adaptability must be improved by favouring uneven-aged structure and tree species mixture (Thompson et al., 2009; Oliver et al., 2015). As the combinations of site conditions, climate projections, stand structures and tree species compositions are nearly infinite, all the management options that could potentially enhance the resilience and adaptive capacity of forests cannot be tested in situ (Cantarello et al., 2017). Furthermore, such silvicultural trials provide results only for present or past conditions~~would provide results and only~~ in the long run given the life span of trees. Scenario analysis based on model simulations are therefore useful to select the most promising management strategies and to evaluate their long-term sustainability. To explore forest response to new silvicultural practices and to unexperienced climate conditions in a realistic way, one needs new process-based models able to deal with mixed and structurally complex stands and to incorporate uncertainties in future conditions (Pretzsch et al., 2015).

In connection with the traditional forestry viewing forests as a stable system that can be controlled, many empirical models were developed to predict tree growth in monocultures considering that past conditions will remain unchanged in the future. On the other hand, scientists developed process-based eco-physiological models to better understand short- and long-term forest ecosystem response to multiple and interacting environmental changes (Dufrêne et al., 2005). This can indeed not be done through direct experimentation because the multisite and multifactorial experiments required for doing so would be too complex and too expensive (Aber et al., 2001; Boisvenue and Running, 2006). Most experiments of environment manipulation focus on single or few factors during a limited period of time, which precludes to properly take into account interactions, feedbacks and acclimation. To simplify the mathematical formalization of eco-physiological processes (e.g., radiation interception) and limit the calculation time, these process-based models were first designed for pure even-aged stands without considering the spatial heterogeneity of stand structure.

With the increasing interest for uneven-aged stands and tree species mixtures, cohort and tree-level models were also developed. Pretzsch et al. (2015) reviewed 54 forest growth models to show how they represent species mixing. Among those models, 36 were process-based with 9 at the stand, 11 at the cohort and 16 at the tree level. While cohort models allow to describe the vertical structure of the stand, tree-level models are generally necessary to consider the spatial heterogeneity in the horizontal dimensions. To represent stand structure in three dimensions, the model must not only operate at the individual level but also consider the tree position. In the review of Pretzsch et al. (2015), 11 process-based models were individual-based and spatially explicit but only three of them accounted simultaneously for radiation transfer, water cycling and phenology (i.e.,

BALANCE, EMILION and MAESPA). Since it describes canopy and water balance processes using a state-of-the-art approach (based on a fine crown discretization), MAESPA is a very useful tool for analysing outcomes of eco-physiological experiments (Duursma and Medlyn, 2012). MAESPA is however not suitable for multi-year simulations since it contains no routine for carbon allocation, respiration and tree dimensional growth. EMILION is also restricted to one-year simulation (no organ emergence) and is specific to pine species with a quite detailed structural approach (Bosc et al., 2000). In contrast, tree dimensional growth is well described in BALANCE which possesses a fine representation of tree structure (Grote and Pretzsch, 2002). In BALANCE, radiation interception by trees and water cycling are based on simpler eco-physiological concepts compared to MAESPA and photosynthesis is calculated with a 10-day time step using the routine of Haxeltine and Prentice (1996). As the Forest v5.1 model (Schwalm and Ek, 2004), BALANCE has the advantage of merging two traditions, conventional growth and yield models together with process-based approaches, providing outputs familiar to foresters (classical tree and stand measurements obtained from forest inventory) as well as carbon fluxes and stocks. Among the three models, BALANCE is the only one that considers mineral nutrition through the impact of nitrogen (N) availability on tree growth. ~~The approach used for modelling nutrient cycling is however very simple. Soil is not partitioned into horizons and the~~Some soil chemistry processes (e.g. ion exchange, mineral weathering) are however not described although they are essential to estimate bioavailability of the major nutrients other than N (P, K, Mg, Ca). Not considered in the review of Pretzsch et al. (2015), iLand is another individual-based model describing the eco-physiological processes with an intermediate level of detail using simplified eco-physiological concepts (such as the radiation use efficiency approach) in order to simulate forest dynamics also at the landscape scale. Later, Simioni et al. (2016) developed the NOTG 3D model to study water and carbon fluxes in Mediterranean forests using an individual-based approach to account for the spatial structure of the stand. This model is more suited for short-term (a few years) rather than long-term (a rotation) simulations since tree dimensions are updated based on fixed empirical relationships between diameter at breast height (*dbh*) and tree height or crown radius.

As the models accounting for both the functional and spatial complexity are rare, we developed a new model (HETEROFOR) using a spatially explicit approach to describe individual tree growth based on resource use (light, water and nutrients) in HETEROgeneous FORests. While the BALANCE and iLand model existed and responded roughly to our expectations, we decided to build a new model for several reasons. First, we thought that another model of this particular type would not be redundant if based on other concepts. Instead of calculating an index of light availability, we chose to estimate radiation interception for all trees using a ray tracing approach. For calculating photosynthesis and tree transpiration, we selected the Farquhar model with shorter time step than in BALANCE in order to account for hourly variations in climate and soil water conditions. While we used a slightly more complex approach for the water balance module (Darcy approach instead of bucket model for soil water dynamics, rainfall partitioning when passing through the canopy), our model rests on a simpler representation of tree structure than BALANCE. Second, we aimed at incorporating a detailed tree nutrition and nutrient cycling module since we realized the necessity to integrate nutritional constraints in forest growth modelling, especially for predicting the response to climate change (Fernandez-Martinez et al., 2014; Jonard et al., 2015). Finally, we wanted to develop the model in a collaborative modelling platform dedicated to tree growth and stand dynamics. Among the various platforms,

CAPSIS was the only one allowing multi-model integration and providing a user-friendly interface (Dufour-Kowalski et al., 2012). HETEROFOR was therefore progressively elaborated through the integration of various modules (light interception, phenology, water cycling, photosynthesis and respiration, carbon allocation, mineral nutrition and nutrient cycling) within CAPSIS. The advantage of such a platform is to use common development environment, model execution system, user-
5 interface and visualization tools and to share data structures, objects, methods and libraries.

To simulate the response of forests to management and changing environmental conditions, integrate and structure the existing knowledge into process-based models is essential but not sufficient. These models must also be documented and evaluated in order to know exactly their strengths and limits when analysing their outputs. The objectives of this paper are (i) to describe the carbon-related processes of HETEROFOR (photosynthesis, respiration, carbon allocation and tree dimensional growth),
10 (ii) evaluate the model ability in reconstructing tree growth in three broadleaved stands of contrasted species composition and compare various options for describing photosynthesis, respiration and crown extension and (iii) illustrate its potentialities by simulating tree growth dynamics in an oak and beech stand under various IPCC climate scenarios.

2. Materials and methods

2.1 Overall operation of the HETEROFOR model

HETEROFOR is a model integrated in the CAPSIS (Computer-Aided Projection for Strategies in Silviculture) platform dedicated to forest growth and dynamics modelling (Dufour-Kowalski et al., 2012). CAPSIS provides to HETEROFOR the execution system and the methods necessary to run simulations and display the results. When running simulations with HETEROFOR, CAPSIS creates a new project in which the variables describing the forest state are stored at a yearly time step, starting from the initial forest characteristics (initial step). Though some data structures and methods are shared with other models integrated in CAPSIS, the initialisation and evolution procedures are specific to HETEROFOR.

For the initialization, HETEROFOR loads a series of files containing tree species parameters, input data on tree (location, dimensions and chemistry), soil (chemical and physical properties) and open field hourly meteorological data. These data are used to create trees and soil horizons at the initial step. The tree is divided in three structural compartments (branch, stem, root) and three functional ones (leaf, fine root and fruit). Then, HETEROFOR predicts tree growth at a yearly time step based on underlying processes modelled at finer time steps and at different spatial levels.

After the initialization step, and at the end of each successive yearly time step, the phenological periods for each deciduous species (leaf development, leaf colouring and shedding) are defined for the next step from meteorological data. When no hourly meteorological measurements are available, the vegetation period is defined by the user who provides the budburst and the leaf shedding dates. Knowing the key phenological dates and the rates of leaf expansion, colouring and falling, the foliage state of the deciduous species is predicted at any timewith a daily time step during the year (de Wergifosse et al., in review). It is characterized by the proportions of leaf biomass and of green leaves relatively to complete leaf development, which are key variables to simulate energy, water and carbon fluxes within the forest ecosystem. The proportion of green leaves impacts photosynthesis, leaf respiration and tree transpiration, as these processes are not active anymore on discoloured leaves which however still intercept solar radiation and rainfall. Based on a ray tracing approach, the SAMSARALIGHT library of CAPSIS (Courbaud et al., 2003) calculates the solar radiation absorbed by the trunk and the crown of each individual tree and the radiation transmitted to the ground. This allows HETEROFOR to estimate the proportions of incident radiation absorbed by the trunk and the crown of each tree and the part transmitted to the ground either on average over the whole vegetation period (simplified budget) or hourly for several key dates (detailed budget). These proportions and the incident radiation measured in the meteorological station are used to compute the hourly global, direct and diffuse radiation absorbed per unit bark or leaf area. Predicting how solar energy is distributed within the forest ecosystem is necessary to estimate foliage, bark and soil evaporation, tree transpiration and leaf photosynthesis.

Every hour, HETEROFOR performs a water balance and updates the water content of each horizon. Rainfall is partitioned in throughfall, stemflow and interception (Andre et al., 2008a; 2008b and 2011). Part of the rainfall reaches directly the ground (throughfall) while the rest is intercepted by foliage and bark. ~~These two tree compartments~~They both have a certain water storage capacity which is regenerated by evaporation. When the foliage is saturated, the overflow joins the throughfall flux

whose proportion increases. As the bark saturates, water flows along the trunk to form stemflow. ~~Foliage and bark storage capacity as well as stemflow proportion are determined at the tree level and then upscaled to the stand level, while evaporation from these surfaces is evaluated at the stand scale. Throughfall is also determined at the stand level as the difference between incident rainfall and the abovementioned fluxes.~~ Throughfall and stemflow supply the first soil horizon (forest floor) with water while soil evaporation and root uptake deplete it. The water evaporation from the soil (as well as from the foliage and the bark) is calculated ~~at stand scale~~ with the Penman-Monteith equation based on the solar radiation absorbed by each component. Using the same equation, individual tree transpiration is estimated by determining the stomatal conductance from tree characteristics, soil extractable water and meteorological conditions. The distribution of root water uptake among the soil horizons is done according to the water potential and the vertical distribution of fine roots. Water exchanges between soil horizons are considered as water inputs (capillary rise) or outputs (drainage). This soil water transfers are calculated based on the water potential gradients according to the Darcy law and using pedotransfer functions to determined soil hydraulic properties. By default, HETEROFOR calculates the water fluxes at the stand scale by aggregating individual fluxes (i.e. tree transpiration) or tree properties (e.g. foliage and bark capacity, stemflow proportion). With this option, all trees are taking up water in the same soil horizons assuming that soil water is redistributed homogeneously between two hourly time steps. However, the user can choose an alternative option to calculate all the water fluxes at the individual level. In this case, the model distributes the total soil volume in individual soil volumes (called pedon) and performs a water balance for each one. Contrary to the default option assuming a homogeneous horizontal water redistribution, the alternative option supposes no water redistribution among pedons (de Wergifosse et al., in review). All these soil water fluxes are considered at the stand level (de Wergifosse et al., in review). ~~Foliage and bark storage capacity as well as stemflow proportion are determined at the tree level and then upscaled to the stand level, while evaporation from these surfaces is evaluated at the stand scale. Throughfall is also determined at the stand level as the difference between incident rainfall and the abovementioned fluxes.~~

The user can choose to calculate ~~The~~ the gross primary production of each tree (*gpp*) ~~is~~ either ~~obtained~~ based on a radiation use efficiency approach distinguishing sunlit and shaded leaves (yearly time step) or ~~calculated hourly~~ using the Farquhar et al. (1980) model (hourly time step). The latter is analytically coupled to the stomatal conductance model proposed by Ball et al. (1987). The photosynthesis is computed using the Library CASTANEA also present in CAPSIS (Dufrêne et al., 2005). This calculation requires the proportions of sunlit and shaded leaves, the direct and diffuse photosynthetically active radiation (*PAR*) absorbed per unit leaf area and the mean soil water potential. At the end of the vegetation period, *gpp* is then converted to net primary production (*npp*) after subtraction of growth and maintenance respiration. Maintenance respiration is either considered as a proportion of *gpp* (depending on the crown to stem diameter ratio) or calculated hourly for each tree compartment by considering the living biomass, the nitrogen concentration and a Q10 function for the temperature dependency following Ryan (1991) as in Dufrêne et al. (2005). Carbon allocation is ~~made~~ done once a year at the end of the vegetation period which allows to update tree dimensions for the next yearly time step during which tree size does not change. Carbon is allocated in priority to foliage and fine roots by ensuring a functional balance between carbon fixation and nutrient uptake through a fine root to

leaf biomass ratio depending on the tree nutritional status (Helmisaari et al., 2007). Allometric relationships are then used to describe carbon allocation to structural components (trunk, branches and structural roots) and to derive tree dimensional growth (diameter at breast height, total height, height to crown base, height of largest crown extension, crown radii in 4 directions) while considering competition with neighbouring trees (Fig. 1).

- 5 Knowing the chemical composition of the tree compartments for a given tree nutrient status, HETEROFOR computes the individual tree nutrient requirements based on the estimated annual growth rate and deduces the tree nutrient demand after subtraction of the amount of re-translocated nutrients. On another hand, the potential nutrient uptake is obtained by calculating the maximum rate of ion transport towards the roots (by diffusion and mass flow). The actual uptake is then determined by adjusting the tree nutrient status and growth rate so that tree nutrient demand matches soil nutrient supply. The nutrient
10 limitation of tree growth is achieved through the regulation of maintenance respiration and through the effect of the tree nutrient status on fine root allocation.

- The soil chemistry is characterized at the stand scale for the various soil horizons defined by the user. In each soil horizon, the chemical composition of the soil solution is in equilibrium with the exchange complex and the secondary minerals. ~~This compartment~~ It receives the nutrients coming from atmospheric deposition, organic matter mineralization and primary mineral
15 weathering, and is depleted by root uptake and immobilization in micro-organisms. The chemical equilibrium within the soil solution, with the exchange complex or the minerals is updated yearly with the PHREEQC geochemical model (Charlton and Parkhurst, 2011) coupled to HETEROFOR through a dynamic link library.

- In this paper, we present a detailed description of the processes regulating the carbon fluxes (Fig. 1) while the phenology and water balance modules are presented in a companion paper (de Wergifosse et al., in review) and the nutrient cycling and tree
20 nutrition module will be described later in a third paper.




Figure 1. Conceptual diagram of the HETEROFOR model. The incident *PAR* radiation is absorbed by individual trees using a ray tracing model (SAMSARALIGHT library). Then, the absorbed *PAR* (*aPAR*) is converted into gross primary production (*gpp*) based on the *PAR* use efficiency concept (first option) or with a biochemical model of photosynthesis (coupling with the CASTANEA library second option). The photosynthesis calculation depends on the soil water content which is updated hourly thanks to the water balance module described in details in de Wergifosse et al. (in review). The net primary production (*npp*) is obtained using a *npp* to *gpp* ratio or by subtracting the growth and maintenance respiration (the latter being temperature dependent). *npp* is first allocated to foliage using an allometric equation function of tree diameter (*dbh*) and crown radius (*cr*). All these processes (radiation interception, photosynthesis and respiration as well as evapotranspiration) depend on the foliage development stage which is determined based on the phenology module. The carbon allocated to fine roots is determined based on a fine root to foliage ratio dependent on the tree nutritional status. Fruit production is calculated with an allometric equation based on *dbh* and on light availability. The remaining carbon is allocated to structural organs-compartments (roots, trunk and branches) using a fixed proportion for the below-ground part. *dbh* and height growth (Δdbh , Δh) are deduced from the change in aboveground biomass by deriving and rearranging an allometric equation. Finally, crown extension is predicted with a distance-dependent or -independent approach.

2.2 Detailed model description

2.2.1 Initialization

To initialize HETEROFOR, the relative position (x, y, z) and the main dimensions of each tree must be provided: girth at breast height (gbh in cm), height (h in m), height of maximum crown extension ($hlce$ in m), height to crown base (hcb in m) and crown radii in the four cardinal directions (cr in m). During the initialization phase, the biomass of each tree compartment is calculated according to the equations used for carbon allocation (see sect. 2.2.4). If available, site-specific allometric equations can also be used to calculate initial biomasses of tree compartments. When data on fruit litterfall are available, a file providing the amount of fruit litterfall per year and per tree species can be loaded and used to adapt the allometric equations predicting fruit production at the individual level. When the water balance module is activated, two additional files must be loaded: a file describing soil horizon properties and another one for the hourly meteorology.

2.2.2 Gross primary production

The annual gross primary production of each tree (gpp in kgC yr^{-1}) is calculated either based on a PAR use efficiency (PUE) approach (Monteith, 1977) or using the photosynthesis method of the CASTANEA model (Dufrêne et al., 2005). Whatever the option retained, a series of variables are needed to calculate gpp .

For the PUE approach, the model uses the solar radiation absorbed by each tree during the vegetation period ($aRAD$ in MJ yr^{-1}). $aRAD$ is then converted in PAR ($aPAR$ in $\text{mol photons yr}^{-1}$) by supposing that 46% of the solar radiation (RAD) is PAR and that 1 MJ is equivalent to 4.55 moles of photons. The diffuse and direct components of $aPAR$ are also considered ($aPAR_{diff}$ and $aPAR_{dir}$ in $\text{mol photons yr}^{-1}$). While all the leaves receive diffuse PAR , only sunlit leaves absorb direct PAR . To estimate the sunlit leaf proportion ($Prop_{sl}$) at the tree level, HETEROFOR uses an adaptation of the classical stand-scale approach based on the Beer-Lambert law (Teh, 2006):

$$Prop_{sl} = \frac{1 - \exp(-k \cdot LAI)}{k} \quad (1)$$

with

k , the extinction coefficient,

LAI , the leaf area index ($\text{m}^2 \text{m}^{-2}$).

At the individual scale, the leaf area index is calculated by dividing the tree leaf area (a_{leaf} in m^2) by the crown projection area (cpa in m^2). The value obtained is then multiplied by the light competition index (LCI in MJ MJ^{-1}) to account for the shading effect of the neighbouring trees:

$$Prop_{sl} = \frac{1 - \exp\left(-k \cdot \frac{a_{leaf}}{cpa}\right)}{k} \cdot LCI \quad (2)$$

where LCI is the ratio between the absorbed radiation calculated with and without neighbouring trees in SAMSARALIGHT. LCI ranges from 1 (no light competition) to 0 (no light reaching the tree).

To adapt the *PAR* use efficiency concept (*PUE*) at the tree level, we considered a distinct *PUE* for sunlit (*sl*) and shaded (*sh*) leaves and calculated an average *PUE* weighted as follows:

$$pue = \frac{aPAR_{diff} \cdot (Prop_{sl} \cdot PUE_{sl} + Prop_{sh} \cdot PUE_{sh}) + aPAR_{dir} \cdot PUE_{sl}}{aPAR} \quad (3)$$

This *pue* is then used to calculate *gpp* based on *aPAR* and a reducer accounting for water stress (*red_{water}*):

$$gpp = aPAR \cdot pue \cdot red_{water} \quad (4)$$

The default value of *red_{water}* is 1 but, when the hydrological module is activated, it is set to the ratio between the actual and the potential (i.e., considering no soil water limitation) tree transpiration (*t_{actual}* and *t_{pot}*, in l per year). This ratio estimates the fraction of the vegetation period during which stomata are partially or totally closed due to limitation in soil water availability. Since this ratio is always lower or equal to 1, a correction factor is applied to avoid introducing a bias.

$$red_{water} = \frac{t_{actual}}{t_{pot}} \cdot corr \quad (5)$$

gpp can also be estimated using the photosynthesis method of CASTANEA (Dufrêne et al., 2005). This method consists in the biochemical model of Farquhar et al. (1980) analytically coupled with the approach of Ball et al. (1987) that linearly relates stomatal conductance to the product of the carbon assimilation rate by the relative humidity. The slope of this relationship varies between 0 and 1 with the soil water availability characterized in HETEROFOR based on a decreasing exponential function of the mean soil water potential (see Eq. 56 in de Wergifosse et al., in review). The formulation of Ball et al. (1987) was slightly adapted to the tree level by accounting for the influence of tree height. Indeed, leaf water potential increases with leaf height and induces a decrease in stomatal conductance (Ryan and Yoder, 1997; Schäfer et al., 2000).

The photosynthesis method requires, at an hourly time step, the direct and diffuse *PAR* absorbed per unit leaf area. The direct *PAR* is intercepted only by sunlit leaves and is obtained by multiplying the hourly incident *PAR* ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$) by the proportion of direct *PAR* absorbed by sunlit leaves. For a tree, this proportion is by default fixed for the whole vegetation period and calculated as the ratio between the direct *PAR* absorbed per unit sunlit leaf area during the vegetation period (in $\text{mol photons.m}^{-2}.\text{yr}^{-1}$) and the incident *PAR* cumulated over the same period (in $\text{mol photons m}^{-2} \text{yr}^{-1}$). A similar procedure is used for the diffuse absorbed *PAR*, except that it is related to the total leaf area. When using the detailed version of SAMSARALIGHT, the proportions of direct/diffuse *PAR* absorbed per unit leaf area change every hour during the day and depending on the phenological stage.

2.2.3 Growth and maintenance respiration

gpp is converted to annual net primary production (*npp* in kgC yr^{-1}) using either a ratio depending on the crown to stem diameter ratio (Eq. 6) or after subtraction of growth (*gr*) and maintenance respiration (*mr*) (Eq. 7) according to the theory of respiration developed by Penning de Vries (1975).

$$npp = gpp \cdot r_{npp_gpp}(DdI) \quad (6)$$

$$npp = gpp - mr - gr \quad (7)$$

Mäkelä and Valentine (2001) showed that the npp to gpp ratio changes with some tree characteristics (tree size, height and age). Based on simulated gpp and npp reconstructed by using the model in reverse mode (see sect. 2.2.7), we tested the impact of several variables characterizing tree dimensions and shape (height, dbh , crown radius, crown volume, crown to stem diameter ratio, aboveground volume or biomass) on the npp to gpp ratio. The best relationship was obtained with the crown to stem diameter ratio (Dd in $m\ m^{-1}$) which had a negative effect on the npp to gpp ratio. This indicates that the proportion of gpp lost by respiration increases for trees with a large crown. Unfortunately, As the crown to stem diameter ratio not only varies with the tree shape reflecting past competition conditions but also changes during the course of the tree development for some tree species. Therefore, we standardized it to remove the size effect in order to obtain an index ($DdIndex$) only characterizing the tree shape, standardized it to obtain a crown to stem diameter index ($DdIndex$). This index is particularly useful to account for the large differences in oak crown extension according to the silvicultural system (large crowns in former coppices with standards vs narrow crowns in dense high forests).

$$r_{npp_gpp} = \alpha + \beta \cdot DdIndex \quad (8)$$

where α and β are parameters and $DdIndex$ is defined as :

$$DdIndex = \frac{Dd}{Dd_{pred}} \quad (9)$$

with

Dd , the crown to stem diameter ratio determined from the tree mean crown radius (cr_{mean} in m) and diameter at breast height (dbh in m),

Dd_{pred} , the crown to stem diameter ratio predicted based on the girth at breast height (gbh in cm):

$$Dd_{pred} = \alpha + \beta \cdot gbh + \gamma \cdot \frac{1}{gbh} + \delta \cdot \frac{1}{gbh^2} \quad (10)$$

In Eq. (7), maintenance respiration is calculated for each tree by summing the maintenance respiration of each organ compartment estimated from the nitrogen content of its living biomass and considering a Q_{10} function for the temperature dependency. During daytime, the inhibition of foliage respiration by light is taken into account by considering that this inhibition reduces respiration by 62% (Villard et al., 1995).

$$mr = \sum_{organcomp} \left(b_{comp,organ} \cdot f_{living} \cdot [N] \cdot R_{T_{ref}} \cdot Q_{10,organ}^{\frac{T-T_{ref}}{10}} \right) \quad (11)$$

with

$b_{organcomp}$, the organ tree compartment biomass (kg of organic matter),

f_{living} , the fraction of living biomass,

$[N]$, the nitrogen concentration ($g\ kg^{-1}$),

$R_{T_{ref}}$, the maintenance respiration per g of N at the reference temperature ($15^{\circ}C$),

T , is the air temperature for aboveground ~~tree organs compartments~~ or the soil horizon temperature for roots (see Appendix A). Root maintenance respiration is estimated for each soil horizon separately.

The fraction of living biomass is fixed to 1 for leaves and fine roots or equals the proportion of sapwood for the structural ~~tree organs compartments~~. The sapwood proportion is derived from the sapwood area ($a_{sapwood}$ in cm^2) determined based on an empirical function of ~~organ-the tree compartment~~ diameter ($\phi_{organcomp.}$ in cm):

$$a_{sapwood} = a + b \cdot \phi_{organcomp.} + c \cdot \phi_{organcomp.}^2 \quad (12)$$

Growth respiration is the sum of the ~~organ-tree compartment~~ growth respiration which is proportional to their ~~organ~~ biomass increment (see sect. 2.2.4):

$$gr = \sum_{organ} (R_{gr} \cdot \Delta b_{organ}) \quad (13)$$

where R_{gr} is the growth respiration per unit biomass increment ($kgC\ kgC^{-1}$).

2.2.4 Carbon allocation and dimensional growth

For each tree, the npp and the carbon retranslocated from leaves and roots (rt_{leaf} and $rt_{fine\ root}$ in $kgC\ yr^{-1}$) are distributed among the various tree compartments at the end of the year. rt_{leaf} and $rt_{fine\ root}$ are determined as follows :

$$rt_{leaf\ or\ fine\ root} = b_{leaf\ or\ fine\ root} \cdot \delta_{leaf\ or\ fine\ root} \cdot rtr_{leaf\ or\ fine\ root} \quad (14)$$

where b_{leaf} and $b_{fine\ root}$ are the tree leaf and fine root biomasses (kgC), δ_{leaf} and $\delta_{fine\ root}$ are the leaf and fine root turnover rates ($kgC\ kgC^{-1}\ yr^{-1}$), and rtr_{leaf} and $rtr_{fine\ root}$ are the leaf and fine root retranslocation rates ($kgC\ kgC^{-1}$).

b_{leaf} is estimated with an allometric equation based on the stem diameter at breast height (dbh in cm) and on the crown to stem diameter ratio (Dd):

$$b_{leaf} = \alpha \cdot dbh^\beta \cdot Dd^\gamma \quad (15)$$

$b_{fine\ root}$ is deduced from the leaf biomass using the fine root to leaf ratio ($r_{fine\ root\ to\ foliage}$):

$$b_{fine\ root} = b_{leaf} \cdot r_{fine\ root_leaf} \quad (16)$$

$r_{fine\ root_leaf}$ takes a value between a minimum ($r_{fine\ root_leaf_min}$) and maximum ($r_{fine\ root_leaf_max}$) ratio depending on the tree nutritional status, in accordance with the concept of functional balance (Mäkelä 1986). This means that a higher ratio is used (more carbon allocation to fine roots) when tree suffers from nutrient deficiency. For each nutrient, a candidate ratio is obtained based on a linear relationship depending on the nutritional status. The ratio increases when the nutritional status deteriorates and this effect is more pronounced for nitrogen (N) > phosphorus (P) > potassium (K) > magnesium (Mg) > calcium (Ca). Among the candidate ratios, the maximum is retained in order to account for the fact that the most limiting nutrient has the dominant effect. For each nutrient, the nutritional status is bounded between 0 and 1 and calculated based on the foliar concentrations (provided in the inventory file) and on the optimum and deficiency thresholds (Mellert and Göttlein, 2012).

$$Status(Nutrient) = \frac{[Foliar\ Nutrient] - Deficiency}{Optimum - Deficiency} \quad (17)$$

The leaf and fine root litter amounts (s_{leaf} and $s_{fine\ root}$ in kgC yr⁻¹) are estimated based on the turnover rate taking into account the retranslocation:

$$s_{leaf\ or\ fine\ root} = b_{leaf\ or\ fine\ root} \cdot \delta_{leaf\ or\ fine\ root} \cdot (1 - rt_{leaf\ or\ fine\ root}) \quad (18)$$

- 5 In the allocation, priority is given to leaves and fine roots. The carbon allocated to leaves corresponds to the annual leaf production (p_{leaf} in kgC yr⁻¹) which is equal to the amount of leaves fallen the previous year plus the leaf biomass change (Δb_{leaf} in kgC yr⁻¹):

$$p_{leaf} = b_{leaf\ t-1} \cdot \delta_{leaf} + \Delta b_{leaf} \quad (19)$$

where Δb_{leaf} is determined by :

$$10 \quad \Delta b_{leaf} = b_{leaf\ t} - b_{leaf\ t-1} \quad (20)$$

with $b_{leaf\ t-1}$ and $b_{leaf\ t}$ being the tree leaf biomasses corresponding to the previous and the current years, respectively.

The fine root production is then estimated according to the same logic:

$$p_{fine\ root} = b_{fine\ root\ t-1} \cdot \delta_{fr} + \Delta b_{fr} \quad (21)$$

- 15 where $b_{fine\ root\ t-1}$ is provided by Eq. (16).

When the carbon allocated to leaf and fine root is higher than the npp plus the retranslocated carbon (suppressed trees with low gpp and npp for their size), the leaf and fine root productions are recalculated so that they do not exceed 90% of the available carbon.

- 20 Then, the fruit production (p_{fruit} in kgC yr⁻¹) is estimated with an allometric equation similar to Eq. (15) and is considered directly proportional to the light competition index since fructification is known to be favoured when tree crowns are exposed to the sun. A threshold dbh ($dbh_{threshold}$ in cm) is fixed below which no fruit production occurs.

$$p_{fruit} = \alpha \cdot LCI \cdot (dbh - dbh_{threshold})^\beta \quad (22)$$

In this equation, the parameter α takes a default value or is adapted based on the fruit production of the year (when the file with the amount of fruit litterfall per year and per tree species is loaded).

- 25 Part of the carbon is also used to compensate for branch and root mortality. The branch mortality- (s_{branch} in kgC yr⁻¹) is described with an equation of the same form as Eq. (15) while the structural root mortality (s_{root} in kgC yr⁻¹) is obtained using a turnover rate similar to that of the branches.

After subtracting the leaf, fine root and fruit productions and the root and branch senescence, the remaining carbon is allocated to structural organ-tree compartment growth:

$$30 \quad \Delta b_{structural} = npp + rt - p_{leaf} - p_{fine\ root} - p_{fruit} - s_{branch} - s_{root} \quad (23)$$

At this stage, the remaining carbon is partitioned between the above- and below-ground parts of the tree according to a fixed root to shoot ratio (r_{root_shoot}):

$$\Delta b_{structural_above} = \frac{\Delta b_{structural}}{(1+r_{root_shoot})} \quad (24)$$

$$\Delta b_{structural_below} = \Delta b_{structural} - \Delta b_{structural_above} \quad (25)$$

The increment in aboveground structural biomass is then used to determine the combined increment in dbh and total height (h in m) based on an allometric equation used to predict aboveground woody biomass (Genet et al., 2011; Hounzandji et al., 2015):

$$b_{structural_above} = \alpha + \beta (dbh^2 \cdot h)^\gamma \quad (26)$$

Deriving this equation and rearranging terms gives:

$$\Delta b_{structural_above} = \beta \gamma (dbh^2 \cdot h)^{\gamma-1} \Delta (dbh^2 \cdot h) \quad (27)$$

$$\Delta (dbh^2 \cdot h) = \frac{\Delta b_{structural_above}}{\beta \gamma (dbh^2 \cdot h)^{\gamma-1}} \quad (28)$$

The development of the left term provides:

$$\Delta (dbh^2 \cdot h) = (dbh + \Delta dbh)^2 \cdot (h + \Delta h) - dbh^2 \cdot h \quad (29)$$

which can be further developed (see Appendix B for details) to isolate Δh :

$$\Delta h \cong \frac{\Delta (dbh^2 \cdot h)}{dbh^2} - \frac{h \cdot \Delta dbh^2}{dbh^2} \quad (30)$$

From Eq. (30), we know that the height increment can be expressed as a function of $\frac{\Delta (dbh^2 \cdot h)}{dbh^2}$. In the following, we refer to it as the height growth potential (Δh_{pot}) since it corresponds to the height increment if all the remaining carbon was allocated to height growth. Contrary to the other term of Eq. (30) $\left(\frac{h \cdot \Delta dbh^2}{dbh^2}\right)$ which is unknown, this height growth potential can be evaluated at this step by dividing the result of Eq. (28) by dbh^2 . However, depending on the level of competition for light and on the tree size, only part of this height growth potential will be effectively realised for height increment. For each tree species, an empirical relationship predicting height growth from the height growth potential, the light competition index and the tree size (dbh or height) was therefore fitted based on successive inventory data (see Appendix E):

$$\Delta h = a + b \cdot dbh + c \cdot h + d \cdot LCI + e \cdot \Delta h_{pot} + f \cdot (\Delta h_{pot})^2 + g \cdot (\Delta h_{pot})^3 \quad (31)$$

The dbh increment is then determined by rearranging Eq. (29):

$$\Delta dbh = \sqrt{\frac{\Delta (dbh^2 \cdot h) + dbh^2 \cdot h}{(h + \Delta h)}} - dbh \quad (32)$$

The increments in root, stem and branch biomass are obtained as follows:

$$\Delta b_{root} = r_{root_shoot} \cdot \Delta b_{structural_above} \quad (33)$$

$$\Delta b_{stem} = f \cdot \rho \cdot ((dbh + \Delta dbh)^2 \cdot (h_{del} + \Delta h_{del}) - dbh^2 \cdot h_{del}) \quad (34)$$

$$\Delta b_{branch} = \Delta b_{structural_above} - \Delta b_{stem} \quad (35)$$

with

f is the form coefficient ($m^3 m^{-3}$),

ρ is the stem volumetric mass (kgC m^{-3}),

h_{del} is the Delevoy height (m) corresponding to the height at which stem diameter is half the diameter at breast height (see Appendix C).

5 The branch and root biomasses are then distributed in 3 categories defined based on the diameter: small branches/roots < 4 cm, medium branches/roots between 4 and 7 cm, coarse branches/roots > 7 cm. The proportions of small, medium and coarse branches/roots are determined based on ~~the~~ equations of the same form as those presented in Hounzandji et al. (2015) for oak branches. Until we can adjust these equations on appropriate data sets, the parameters of Hounzandji et al. (2015) are also used for beech branches and for oak and beech roots. The distribution in root categories has no impact on the functioning of the model since this information is not used elsewhere. This is just a model output that the user can ignore or consider as a whole.-

2.2.5 Crown extension

Depending on whether the competition with the neighbouring trees is taken into account or not, the crown dynamics can be describe by two different approaches. When local competition is not considered (distance-independent approach), change in crown dimensions are derived from dbh or height increment based on empirical relationships:

$$\Delta hlce = hlce\% \cdot \Delta h \quad (36)$$

$$\Delta hcb = hcb\% \cdot \Delta h \quad (37)$$

$$\Delta cr = Dd_{pred} \cdot \frac{\Delta dbh}{200} \quad (38)$$

where $hcb\%$ and $hlce\%$ are the proportions of the total height corresponding to the height to crown base (hcb in m) and to the height of largest crown extension ($hlce$ in m), respectively;

Δcr is the change in crown radius (in m) whatever the direction;

Dd_{pred} is the crown to stem diameter ratio estimated by Eq. (10).

Alternatively, the changes in crown dimensions can be described based on the competition with the neighbouring trees (distance-dependent approach). The space around a target tree is divided into 4 sectors according to the 4 cardinal directions (North between 315° and 45°, East between 45° and 135°, South between 135° and 225°, West between 225° and 315°). In each sector, the tree which is the closest to the target tree is retained as a competitor if its height is higher than the hcb of the target tree. Beyond a certain distance (i.e., two times the maximal crown radius: 10 m), no competitor is considered. For each main direction, the model calculates an $hlce$ at equilibrium ($hlce_{eq}$ in m) for the target tree. This $hlce$ at equilibrium is located between a minimum (hcb in m) and a maximum ($hlce_{max}$ in m). $hlce_{max}$ is obtained by determining the higher intersection between the potential crowns of the target tree and the competitor. The potential crown of a tree is the crown that this tree would have had in absence of competition and is considered as having the shape of a half ellipsoid centred on the tree trunk and with the semi-axis lengths equal to the tree potential crown radius (cr_{pot} in m, see below) and to the crown length ($h - hcb$). $hlce_{eq}$ is positioned between the minimum and the maximum values according to the competition intensity estimated based on the target tree and the competitor heights (h_{target} and h_{comp} in m) as well as the hcb of the target tree (Appendix D):

$$hlce_{eq} = hcb + (hlce_{max} - hcb) \cdot \max\left(0, \min\left(1, \frac{h_{comp} - hcb}{h_{target} - hcb}\right)\right) \quad (39)$$

The four values of $hlce_{eq}$ are then averaged ($hlce_{eq_mean}$).

Finally, the change in $hlce$ is determined as follows:

if $hlce < hlce_{eq_mean}$,

$$\Delta hlce = \min(\Delta hlce_{max}, hlce_{eq_mean} - hlce) \quad (40)$$

else,

$$\Delta hlce = \max(-\Delta hlce_{max}, hlce_{eq_mean} - hlce) \quad (41)$$

where $\Delta hlce_{max}$ is the maximum change in $hlce$ allowed by the model.

The change in hcb is obtained with the same logic:

if $hcb < hcb_{eq_mean}$,

$$\Delta hcb = \min(\Delta hcb_{max}, hcb_{eq_mean} - hcb) \quad (42)$$

else,

$$\Delta hcb = \max(-\Delta hcb_{max}, hcb_{eq_mean} - hcb) \quad (43)$$

where hcb_{eq_mean} is the hcb estimated from the tree height based on $hcb\%$ (Eq. 37).

The change in the four crown radii is calculated based on crown radii at equilibrium (cr_{eq} in m) which are estimated by considering the competitive strength of the target and neighbouring trees. For a given direction, cr_{eq} is calculated based on the potential (free growth) crown radius of the target tree (cr_{pot_target} in m) and of its competitor (cr_{pot_comp} in m), the distance between the two trees (d in m) and the crown overlap ratio ($r_{overlap}$ in m^{-1}):

$$cr_{eq} = \frac{cr_{pot_target}}{cr_{pot_target} + cr_{pot_comp}} \cdot d \cdot r_{overlap_target} \quad (44)$$

The potential crown radius (cr_{pot}) of a tree is determined by:

$$cr_{pot} = \frac{dbh}{200} \cdot Dd_{pred} \cdot sh \quad (45)$$

where Dd_{pred} is the crown to stem diameter ratio estimated by Eq. (10) and sh is a coefficient allowing to shift from the mean to the maximum Dd_{pred} .

The crown overlap ratio is estimated by considering neighbouring trees of the same species two by two and by calculating the ratio between the sum of their crown radii and the distance between the corresponding tree stems. This overlap ratio accounts for the capacity of a tree species to penetrate in neighbouring crowns.

The change in crown radius is then determined as follows for each direction:

if $cr < cr_{eq}$,

$$\Delta cr = \min(\Delta cr_{max}, cr_{eq} - cr) \quad (46)$$

else,

$$\Delta cr = \max(-\Delta cr_{max}, cr_{eq} - cr) \quad (47)$$

with Δcr_{min} and Δcr_{max} being respectively the minimum and the maximum change in cr allowed by the model. They are obtained similarly as cr_{pot} :

$$\Delta cr_{max} = \frac{\Delta dbh}{200} \cdot Dd \cdot sh \quad (48)$$

2.2.6 Tree harvesting and mortality

During the simulation, thinning can be achieved at each annual step either (i) by selecting the trees from a list or a map or according to tree characteristics (tree species, age, *dbh*, height,...), or (ii) by defining the number of trees to be thinned per diameter class using an interactive histogram, or (iii) by loading a file listing the trees that must be thinned. In addition, the thinning methods developed for GYMNOS and QUERGUS are compatible with HETEROFOR. They allow to reach a target basal area, density or relative density index by thinning from below or from above or by creating gaps (Ligot et al., 2014).

When the *npp* of a tree is not sufficient to ensure a normal leaf and fine root development (for suppressed trees and/or after a severe drought), the leaf biomass is reduced and induces a defoliation which is estimated as follows:

$$Def = \frac{b_{leaf} - b_{leaf_corr}}{b_{leaf}} \cdot 100 \quad (49)$$

where b_{leaf} and b_{leaf_corr} are respectively the leaf biomass estimated with Eq. (15) and the leaf biomass corrected to match the available carbon (see sect. 2.2.4).

Tree mortality occurs when trees reach a defoliation of 90%, considering that a tree with less than 10% of its leaves is in an advanced stage of decline and is unlikely to recover (Manion, 1981). Hence, HETEROFOR takes into account the mortality resulting from carbon starvation due to light competition and/or water stress (stomatal closure).

2.2.7 Growth reconstruction

HETEROFOR was adapted to allow the user to run it in reverse mode starting from the known increments in *dbh* and *h* to reconstruct individual *npp* using exactly the same parameters and equations as in the normal mode. To achieve a reconstruction, an inventory file with tree measurements must be loaded to create the initial step. From this initial step, the reconstruction tools can be launched and requires another inventory file with tree measurements achieved one or several years later. Based on these two inventories, HETEROFOR calculates the mean *dbh* and *h* increments for each tree and use the model equations to reconstruct each step and evaluate among other individual *npp*. The *npp* is obtained by re-arranging Eq. (23) in which the carbon allocated to the structural biomass is calculated from the *dbh* and *h* increments using Eq. (27), (25) and (24). The carbon allocated to leaf, fine root and fruit production is determined respectively with Eq. (19), (21) and (22) while the amount retranslocated from leaves and roots before senescence is evaluated with Eq. (14). Finally, the terms of Eq. (23) accounting for the leaf and fine root litter were determined with Eq. (18). In addition to two stand inventories, the reconstruction tool also requires a file listing the trees which were cut or died between the two inventory dates and the last year during which they were present in the stand.

2.3 Input variables and parameter setting for a case study

The model was tested in three stands contrasting in structure and species composition. These stands were located close to each other (< 1km) on the same tableland (300 m elevation) in the western part of the Belgian Ardennes at Baileux (50° 01' N, 4° 24' E). The average annual rainfall is slightly above 1000 mm and the mean annual temperature is 8°C. The forest (60 ha) consists of sessile oak (*Quercus petraea* Liebl.) and European beech (*Fagus sylvatica* L.) and lies on acid brown earth soil (luvisol according to the FAO soil taxonomy) with a moder humus and an A_hB_wC profile. The soil has been developed on a loamy and stony solifluxion sheet in which weathering products of the bedrock (Lower Devonian: sandstone and schist) were mixed with added periglacial loess.

By the end of the 19th century, the Baileux forest was probably an oak coppice with a few standards. Taking advantage of the massive oak regeneration in the 1880s, the forest developed progressively into a high forest and was then invaded by beech. In 2001, the area was covered by even-aged oak trees and heterogeneously sized beech trees. At that time, three experimental plots were installed at the Baileux site in order to study the impact of tree species mixing on ecosystem functioning (Jonard et al., 2006, 2007, 2008; André et al., 2008a, 2008b, 2010, 2011): two plots were located in stands dominated either by sessile oak or by beech and the third one was a mixture of both species (Table 1). In each plot, all trees with a circumference higher than 15 cm were mapped (coordinates) and measured (stem circumference at a height of 1.3 m, total tree height, height of largest crown extension, height to crown base, crown diameters in two directions) at the end of the years 2001 and 2011.

Meteorological data were monitored with an automatic meteorological station located in an open field 300 m away from the forest site. Soil horizon properties were characterized based on the soil profile description and the measurements carried out by Jonard et al. (2011).

To run the simulations, the values of some model parameters were taken directly from the literature. Other parameters involved in empirical relationships were fitted either with data from previous studies or with unpublished monitoring data collected in the study site or in the ICP Forests level II plots of Wallonia (Table 2). Potential explanatory variables of Eq. 31 used to estimate height growth were selected by applying a stepwise forward selection procedure based on the Bayesian Information Criterion (BIC). A multivariate model was then adjusted with the selected variables (Appendix E). The parameters of the *npp* to *gpp* ratio relationship, the maintenance respiration per g of N at 15°C and the *PAR* use efficiency of sunlit and shaded leaves were adjusted with the nlm function of R (R Core Team, 2013) based on observed basal area increments (*BAIs*) using the maximum likelihood approach. This fitting was achieved only based on the data of the mixed stand while the model performances were evaluated on the tree stands.

All the simulations carried out in this study were run with the default option for modelling phenology and water balance (de Wergifosse et al., in review).

2.4 Statistical evaluation of model predictions

The quality of the model was evaluated for various combinations of model options (i.e., photosynthesis model of CASTANEA vs *PUE*, *npp* to *gpp* ratio vs temperature-dependent maintenance respiration, distance-dependent vs -independent crown extension), by comparing predicted and observed BAIs using several statistical indices and tests such as the normalized average error, the *P* value of the paired *t*-test, the regression test, the root mean square error and the Pearson's correlation (Janssens and Heuberger, 1995). For the regression test, the Deming fitting procedure (mcreg function of the mcr package in R) was retained to account for the errors on both the observations and the predictions.

The model quality was also evaluated based on its ability to reconstruct the size - growth relationships for sessile oak and European beech in the three stands of Baileux. The observed and predicted *BAIs* of the trees (calculated for the 2001 – 2011 period) were related to their girth at the beginning of the assessment period. A segmented regression was then applied to observations and predictions to determine the girth threshold beyond which *BAI* linearly increases with girth and to estimate the slope of the linear relationship between *BAI* and initial girth. The heteroscedasticity of the residuals was accounted by modelling their standard deviation with a power function of the initial girth. The fitting was carried out using the nlm function in R.

2.5 Simulation experiment

To assess how the tree biomass production and its allocation to the different tree compartments were affected by climate conditions and management in the model, we simulated the evolution of the mixed stand during a dry (2003 with P=948mm, T°air=9.88°C), a normal (2005 with P=1027mm, T°air=9.67 °C) and a wet year (2012 with P=1117mm, T°air=9.37°C) and we repeated these simulations after thinning this stand by reducing its basal area by 25%. The biomass production and its allocation was assessed at the stand level as well as at the tree level for seven cohorts (four beech cohorts and three oak cohorts) defined based on the tree species and on the girth-class distribution. For this first simulation experiment, we used the following options: photosynthesis model, *npp* to *gpp* ratio and distance-independent crown extension.

A second simulation experiment was performed To illustrate how the model can be used to predict climate change impacts on forest ecosystem functioning, the growth dynamics in the mixed stand of Baileux was simulated according to three IPCC climate scenarios using the following options: photosynthesis model, *npp* to *gpp* ratio and distance-independent crown extension. The climate scenarios retained for this study were obtained from the global circulation model CNRM-CM5 (Voldoire et al., 2013) based on the Representative Concentration Pathways for atmospheric greenhouse gases described in the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (Collin et al., 2013). The Representative Concentration Pathways (RCP2.6, RCP4.5 RCP8.5) are characterized by the radiative forcing in the year 2100 relative to preindustrial levels (+2.6 W m⁻², +4.5 W m⁻², +8.5 W m⁻²). The CNRM-CM5 describes the earth system climate using variables such as air temperature and precipitations on a low-resolution grid (1.4° in latitude and longitude). Although reliable for

estimating global warming, such a model fails to capture the local climate variations. Therefore, these climate projections were downscaled by the Royal Meteorological Institute of Belgium (RMI), using the regional climate model ALARO-0 (Giot et al., 2016). The meteorological files that were received from RMI are hourly values of the longwave and shortwave radiations, air temperature, surface temperature, rainfall, specific humidity, zonal and meridional wind speeds and atmospheric pressure with a 4 km spatial resolution. Specific humidity was converted into relative humidity using the Tetens formula (Tetens, 1930). For a reference period (1976 - 2005), we compared the models predictions with observed meteorological data and detected some biases, especially for precipitations (overestimation of 27%). The biases were corrected by adding to the predictions (or by multiplying them with) a correction factor specific to the month (Maraun and Widmann, 2018). An additive correction factor was used for the bounded variables (radiations, precipitation, relative humidity, wind speed) and a multiplicative one for the other variables (air and surface temperatures).

For the simulations, two 24-year periods (100 years apart) were considered. The period from 1976 to 1999 served as a historical reference while the rest of the simulations based on climate projections were conducted for the 2076-2099 period. The simulations were performed either by keeping the CO₂ concentration of the atmosphere constant (i.e., 380 ppm) or by allowing it to vary according to the years and climate scenarios. Each simulation started with the same initial stand (mixed stand of Baileux in 2001) and lasted 24 years; a thinning operation (25% in basal area) was achieved in 1978 or 2078 and in 1990 or 2090 (12-year cutting cycle). The mean basal area increment obtained with the various climate scenarios were compared using the Tukey multiple comparison test.

3. Results

3.1 Reconstructed *npp* vs predicted *gpp*

Based on two successive stand inventories (2001 and 2011) and using HETEROFOR in reverse mode (see sect. 2.2.7), the individual *npp* was reconstructed and related to the *gpp* predicted with the photosynthesis method of CASTANEA. The linear relationship between *npp* and *gpp* explained 79 and 83 % of the variability for sessile oak and for European beech, respectively (Fig. 2). The intercept was positive and just significantly different from 0 but did not differ between the two trees species. The slope of the relationship was higher for sessile oak (0.50) than for European beech (0.40).




Figure 2. Relationship between the individual *npp* reconstructed based on successive stand inventories (2001 and 2011) and the *gpp* predicted with the process-based option (photosynthesis method of CASTANEA) for the three stands. Values in parentheses are 95% confidence intervals for the intercept and the slope in the equations. The Pearson's correlation between *npp* and *gpp* is indicated on the graph.

3.2 Model performance in predicting individual basal area increment (*BAI*)

HETEROFOR was run with different combinations of options for describing photosynthesis (biochemical model of CASTANEA vs *PUE*), respiration (*npp* to *gpp* ratio vs temperature-dependent maintenance respiration) and crown extension (distance-dependent vs -independent). The predictions carried out using the photosynthesis routine of CASTANEA were generally slightly better correlated to the observations than those obtained with the *PUE* approach which however displayed somewhat lower RMSE (Table 3). For both options of photosynthesis calculation, the use of the maintenance respiration routine provided less accurate predictions (higher NAE and RMSE and lower Person's *r*) than the *npp* to *gpp* ratio approach and the degradation of the model performance due to the maintenance respiration option was more marked for European beech

than for Sessile oak (Table 3). The option for describing crown extension had little effect on prediction quality. Depending on the criterion considered, on the options selected for calculating photosynthesis and respiration and on the tree species, the distance-independent approach was sometimes the best but not systematically (Table 3).

For the simulations using the CASTANEA photosynthesis, we retained the *npp* to *gpp* approach and the distance-dependent crown extension as the best combination of options since the associated predictions were on average not biased for oak and only slightly for beech (Table 3). For this combination of options, the regression of the observed BAIs on the predictions showed however a slight underestimation of the low BAIs and a small overestimation of the high BAIs, which were more pronounced for European beech than for Sessile oak (Fig. 3). For the PUE method, the *npp* to *gpp* ratio and the distance-independent crown extension provided the most accurate predictions (Table 3).




Figure 3. Comparison of observed and predicted basal area increments (BAIs) for the simulation with the photosynthesis method of CASTANEA, the *npp* to *gpp* ratio approach to account for tree respiration and the distance-dependent crown extension (see Table 3). The dashed line represents the Deming regression between observations and predictions with the shaded area indicating the 95% confidence interval and the solid line the 1:1 relationship.

3.3 Reconstructing size – growth relationships

The size - growth relationships were very similar between observations and predictions for the mixed stand on which the model was calibrated (Fig. 4). For the European beech in the beech dominated stand, the predicted increase in BAI with the initial girth was steeper than the observed one revealing a slight overestimation of the tree growth (Fig. 4). The proportion of the *BAI*

variance explained by the size - growth relationship (R^2) was higher for European beech than for sessile oak for both observations and predictions (Fig. 4).




Figure 4. Reconstruction of the size - growth relationships for sessile oak and European beech in the three stands using the photosynthesis method of CASTANEA, the *npp* to *gpp* ratio approach to account for tree respiration and the distance-dependent crown extension. The predicted relationships between the individual *BAI* (calculated for the 2001–2011 period) and the initial girth are compared with observed ones. The solid and dashed lines represent the segmented regression applied respectively to observations


and predictions to determine the girth threshold beyond which radial growth linearly increases with girth and to estimate the slope of the linear relationship between *BAI* and initial girth. The 95% confidence intervals for the intercept and the slope are provided as well as the R^2 of the model. No relationship was fitted for the European beech in the oak dominated stand given the lack of data.

5 3.4 Simulation of climate change impact on tree growth


The first simulation experiment revealed that thinning affected much more the smallest trees than the biggest ones (Fig. 5). The smallest beech cohort (girth of 0 to 61 cm) almost doubled their annual biomass production after the thinning (+85%) while the thinning impact on the biggest oak and beech trees was hardly noticeable (+4% and +2%, respectively). When looking at the different tree compartments, one may notice that the thinning effect was more pronounced on the structural compartments, i.e. roots, stem and branches (+52%) than on the functional ones, i.e. fine roots, leaves and fruits (+22%). While thinning increased the individual biomass production, it decreased the biomass production at the stand level (-15%). Compared to dry climate conditions, the biomass production at the stand level was 11% higher under normal conditions (Fig. 5). This effect was observed for all the cohorts even if, inversely to the thinning effect, dryer conditions affected less the smallest trees (+2% for the 0-61cm beech cohort) than the biggest ones (+13% for oak and beech trees with a girth larger than 140 cm). Whatever the scale considered (tree or stand), there was nearly no difference in biomass production between normal and wet conditions. The climate condition effects were marked only on the structural compartment (+25%).

When the CO₂ concentration of the atmosphere was fixed, no effect of the climate scenario was detected on stand *BAI* but a slight impact was assessed on sessile oak *BAI* which was higher for the RCP2.6 than for the historical scenario (Fig. 56). For the simulations with a variable atmospheric CO₂ concentration, the difference in total, sessile oak and European beech *BAI* were much more pronounced between climate scenarios. For the whole stand as well as for oak and beech separately, *BAI* increased in the order - historical, RCP2.6, RCP4.5 and RCP8.5 -, with the stand *BAI* of these RCP scenarios being between 17 and 72% higher than that of the historical scenario. All scenarios had a *BAI* significantly different from each other, except RCP2.6 and RCP4.5 for the whole stand and the two tree species and historical and RCP2.6 for European beech (Fig. 56).


All cohorts and tree species




European beech (circ. 0-61 cm)




Sessile oak (circ. 61-105 cm)




European beech (circ. 61-105 cm)



Sessile oak (circ. 105-140 cm)



European beech (circ. 105-140 cm)



Sessile oak (circ. 140+ cm)



European beech (circ. 140+ cm)






Figure 5. Effects of climate conditions and thinning on biomass production and on its allocation to tree compartment in the mixed stand of Baileux. The data used to make these graphs were obtained by simulations using the following options: photosynthesis model, npp to gpp ratio and distance-independent crown extension.




Figure 56. Basal area increment (*BAI*) of the mixed stand in Baileux (and of its two main tree species) simulated with climate scenarios produced with the GCM model CNRM-CM5, downscaled with ALARO-0 and corrected empirically for remaining biases. The simulations were performed by using the CASTANEA method to calculate photosynthesis, the *npp* to *gpp* ratio approach and a distance-independent description of crown extension. The CO₂ concentration of the atmosphere was either kept constant (left) or increased with time according to the climate scenario considered (right). Two time periods were considered. 1976-1999 was used as a reference period for running the model with the historical climate scenario while the simulations with future climate scenarios were achieved for the 2076-2099 period. The climate scenarios were based on the representative concentration pathways for atmospheric greenhouse gases described in the fifth assessment report of IPPC. For a given tree species and CO₂ concentration modality, the scenarios with common letters have a *BAI* not significantly different from each other ($\alpha=0.05$).

4. Discussion

Few tree-level, process-based and spatially explicit models have been developed and these often contain only some of the modules necessary to estimate resource availability (solar radiation, water and nutrients). While a description of these models is generally available in the literature, their evaluation by comparison with tree growth measurements is not always accessible or was carried out based on stand-level variables. We have therefore very few information to compare the performances of HETEROFOR at the tree level with those of similar models. Simioni et al. (2016) faced the same problem with the NOTG 3D model.

HETEROFOR first estimates the key phenological dates, the radiation interception by trees and the hourly water balance (de Wergifosse et al., in review). Then, based on the absorbed *PAR* radiation, individual *gpp* is calculated with a *PUE* approach or with the photosynthesis routine of CASTANEA (Dufrêne et al., 2005). Whatever the option retained for calculating tree respiration and crown extension, the photosynthesis routine of CASTANEA and the *PUE* efficiency approach performed similarly (Table 3). This is quite encouraging that the process-based approach for estimating photosynthesis provided predictions of the same quality than the empirical approach fitted with tree growth data taken on the study site. If no extrapolation to future climate is required, the *PUE* approach remains however still valuable, especially when hourly meteorological data are lacking. For the three stands in Baileux, we related the *npp* reconstructed from successive tree inventories with the *gpp* predicted based on the CASTANEA approach (Fig. 2). The good linear relationships (Pearson's correlation > 0.89) obtained for both oak and beech make us confident in the adaptation of the photosynthesis routine of CASTANEA for the tree level. Furthermore, since the parameters of the photosynthesis routine were taken directly from CASTANEA and not calibrated specifically for HETEROFOR, one can expect that the agreement between the predicted *gpp* and the reconstructed *npp* could still be improved.

When comparing the two options available in HETEROFOR for converting *gpp* into *npp*, model performances were systematically better with the *npp* to *gpp* ratio approach than with the temperature-dependent routine for maintenance respiration calculation (Table 3). This can be partly explained since the error in the maintenance respiration calculation results from various sources. At the tree compartment level, uncertainties in the estimation of biomass, sapwood proportion, nitrogen concentration and temperature are multiplied (Eq. 11). Then, the errors made on all tree compartments are summed up. Among these uncertainty sources, the inaccuracy in the estimation of the sapwood proportion could explain why the maintenance respiration routine provided better results for sessile oak than for European beech (Table 3). Since the sapwood of sessile oak can easily be distinguish from the heartwood based on the colour change, we had a lot of sapwood measurements available to fit a relationship. For European beech, this was not the case; instead, we used a sapwood relationship obtained based on sap flow measurements (Jonard et al., 2011). This relationship could certainly be improved by direct measurements of sapwood made after staining the wood to highlight the living parenchyma. Another way to improve these relationships is to consider the social status of the trees since dominant trees have a higher sapwood depth than the suppressed one (Rodriguez-Calcerrada et al., 2015). We tried to account for this by estimating the sapwood area based on the tree growth rate but it did not significantly

increased the quality of the predictions. The poor performances obtained with the maintenance respiration option also indicates that the processes at play are still poorly understood and that further research are needed on this topic.

The process-based approach for estimating maintenance respiration accounts explicitly for the temperature effect through a Q_{10} function. With the npp to gpp ratio approach, temperature is considered more indirectly by assuming that it affects

respiration and photosynthesis in the same proportion, which is valid only in a given range of temperature ($<20^{\circ}\text{C}$) and for non-stressing conditions. Indeed, the optimum temperature for photosynthesis is between 20 and 30°C while the optimum temperature for respiration is just below the temperature of enzyme inactivation ($>45^{\circ}\text{C}$). Therefore, between 30 and 45°C , photosynthetic rates decrease, but respiration rate could continue to increase (Yamori et al., 2013). This reasoning however

does not consider that the base rate of respiration acclimate to new mean temperature conditions and that this acclimation process tends to maintain the npp to gpp ratio more stable (Collalti and Prentice, 2019). ~~In addition~~ On another hand, while

water stress reduces both photosynthesis and respiration, its effect on the two processes is not necessarily equivalent (Rodriguez-Calcerrada et al., 2014). The main argument in favour of ~~Compared to~~ the npp to gpp ratio approach is the tight

coupling between respiration and photosynthesis since the substrate for respiration originates from photosynthesis. The npp to gpp ratio is unfortunately neither universal nor constant. It may vary with tree development stage, climate, soil fertility and

competition conditions (Collalti and Prentice, 2019). The alternative option based on ,the maintenance respiration calculation seems more appropriate to ~~is theoretically more appropriate to~~ simulate the impact of climate change but this is at the expense

of less accurate predictions at the tree level. ~~but this is at the expense of less accurate predictions at the tree level.~~ The ideal is to compare the two options to evaluate the prediction uncertainty associated with the modelling of respiration. ~~Alternatively, one can choose one or the other option~~

~~depending on the aim of the simulation.~~ In the future, ~~we could also implement a third option accounting for thermal acclimation such as in 3D-CMCC (Collalti et al., 2018).~~ the two approaches could be improved. Applying the reconstruction

procedure of HETEROFOR on a large diversity of sites would allow us to estimate the npp to gpp ratio in many different situations, to create a function predicting the npp to gpp ratio based on its main drivers and to subsequently use it in the model.

In parallel, the respiration calculation could be refined by accounting for thermal acclimation such as in 3D-CMCC (Collalti

et al., 2018).

The differences in prediction quality between the two methods of crown extension modelling (distance-dependent vs – independent approach) were quite small, probably because the length of the simulation was not sufficient to drastically affect the crown dimensions which had been initialized based on measurements. ~~In addition, d~~ Describing mechanisms that governs crown development in interaction with neighbours (mechanical abrasion, crown interpenetration) is ~~however indeed~~ crucial to

capture non-additive effect of species mixtures (Pretzsch, 2014). By accounting for crown plasticity, our distance-dependent approach could help better understand how uneven-aged and mixed stands optimize light interception by canopy packing and how they increase productivity (Forester and Albrecht, 2014; Juncker et al., 2015). To fully evaluate the interest of this approach, the predicted crown development should be compared with precise crown measurements repeated over several decades and taken in a large diversity of stand structures. When the model will be calibrated for a larger number of tree species,

long-term simulations could also be performed to evaluate to which extent the model is able to reconstruct the empirical relationships describing tree allometry variations in response to intra- and inter-specific competition. Such relationships were established by del Rio et al. (2019) using data from the Spanish National Forest Inventory.

Based on the current evaluation, the process-based variant perform similarly than the more empirical one for photosynthesis and crown extension but not for respiration, probably because the processes are better known for photosynthesis. For the best combination of options using the CASTANEA photosynthesis (npp to gpp ratio/distance-dependent crown extension), the Pearson's correlation between measurements and predictions of individual basal increment amounted to 0.83 and 0.63 for European beech and sessile oak, respectively. By comparison, Grote and Pretzsch (2002) obtained a correlation of 0.60 for the individual volume of beech trees with the BALANCE model. This lower correlation can partly be explained by the integration of the uncertainty on tree height in the volume estimations. The HETEROFOR performances in terms of tree growth are quite good and could still be improved by a Bayesian calibration of the parameters.

Individual *npp* and retranslocated C are allocated first to foliage and fine roots and then partitioned between above- and below-ground structural compartments. Based on the derivative and rearrangement of a biomass allometric equation, the increment in aboveground structural biomass is used to estimate the combined increment in *dbh* and height. This results in a system of one equation with two unknowns (increment in *dbh* and height). We decided to resolve it by fixing the height growth based on a relationship taking into account tree size (*dbh* or height), the height growth potential (height increment if all the remaining carbon was allocated to height growth) and a light competition index. An intermediate level of sophistication was adopted to describe height growth, between the simple height-*dbh* allometry and the fine description of tree architecture of functional-structural models. Height-*dbh* relationships provides a static picture in which age and neighbour effects are confounded and are not suitable to describe individual growth trajectories (Henry and Aarsen, 1999). More sophisticated relationships considering age and dominant height can be used for even-aged stands (Le Moguédec and Dhôte, 2012) but are hardly applicable in uneven-aged stands for which tree age is unknown. On the other hand, the functional-structural models based on resource availability at organ level and using a short time step can only be applied to a limited number of trees given the long computing times (Letort et al., 2008).

Our individual height growth model was fitted with height data measured ten years apart (Appendix E). A large uncertainty was however associated to these data. First, height measurements were obtained to the nearest meter given the difficulty to clearly identify the top of the trees in closed canopy forests. Second, two errors were added since the height increment was obtained by the difference between two height measurements. Consequently, the uncertainty was more or less of the same order of magnitude than the expected height growth in ten years. Despite these uncertainties, a substantial part of the variability was explained by the model (72% for European beech, 43% for oak). Among the variables tested, the height growth potential had the main effect, which is not surprising since this height growth potential noisily contains the information on height increment. We were also able to highlight the effect of light competition. For a same height growth potential, trees undergoing stronger light competition seem to invest more carbon for height growth than for *dbh* increment (Fig. E1 in Appendix E), which is corroborated by results of other studies (e.g., Lines et al., 2012). This strategy aims at minimizing overtopping by

neighbours and maximizing light interception (Jucker et al., 2015). Trouvé et al. (2015) found similar results and showed the positive effect of stand density on height growth in the allocation between height and diameter increment in even-aged stands of sessile oak. The decrease in the red:far red ratio of incident light promotes apical dominance and internode elongation through the phytochrome system (shade avoidance reaction, Henry and Aarsen, 1999). By considering the light availability effect on height growth at the tree level, HETEROFOR adapts tree allometry to intra- and inter-specific competition, which is crucial to accounts for mixing effects in structurally-complex stands (del Rio et al., 2019).

A first simulation experiment was achieved to assess how tree biomass production and its allocation to tree compartments respond to climate conditions and thinning. What came out of these simulations is in line with the basic principles of silviculture, thinning favoured individual tree growth (especially that of the smaller trees) by redistributing stand biomass production on a smaller number of trees. At the stand level, thinning slightly reduced biomass production since its intensity was substantial and the simulation lasted only one year which was not sufficient to allow the remaining trees to fill the gaps by extending their crown. Drought conditions affected negatively biomass increment of structural components and was more pronounced on big than on small trees. Indeed, when soil water availability decreases, smaller trees maintain a higher stomatal conductivity because of their lower position in the stand canopy. Functional compartments were less influenced by climate because carbon is allocated to them in priority in the model. The allocation routine could be improved by making the fine root to foliage ratio and the root to shoot ratio dependent on the mean soil water availability (Thurm et al., 2017).

We were also quite satisfied to observe that the model was able to reproduce the size-growth relationship. This ~~relationship~~ approach describes the growth partitioning among trees in a stand, which is useful to estimate the mode of competition. For the three studied stands and the two tree species, the competition was partially size asymmetric with a resource partitioning in favour of the larger trees (Carl et al., 2018). Within the studied stands, the European beech trees can be classified in two groups: a group of small suppressed trees whose radial growth was close to 0 and which were just surviving and the rest of the trees (beyond a girth threshold) whose radial growth linearly increased with girth. Regarding sessile oak, nearly all the trees were in the second group, which can be related to the fact that sessile oak is a less shade-tolerant species than European beech. In the mixed stand, the nearly perfect match between the predicted and observed relationships indicates that the model was able to reproduce growth partitioning among trees of different tree species and size. This very good results can be ascribed to the fact that the extinction coefficient and the respiration parameters were calibrated with data of this stand. In the beech dominated stand, the model slightly underestimated the radial growth of the small oak trees and overestimated that of the big beech trees. In this case, the model seems to allocate too much resources to the big beech trees which shade the small oak trees. This could be improved by a model calibration partly specific to this stand (for the *npp* to *gpp* ratio) or by a calibration with data covering a much larger range of stand structures. is characterized by two parameters: the threshold which defines the girth beyond which radial growth linearly increases with girth and the slope providing the theoretical maximum growth efficiency (Le Moguédec and Dhôte, 2012). For European beech, the observed threshold was 49 cm and was easy to detect visually since there were many trees with a girth inferior to that exhibiting nearly no basal area increment. For sessile oak, the observed threshold was

~~more variable than for European beech (41 and 60 cm) and nearly all trees had a higher girth. This can be related to the fact that sessile oak is a less shade tolerant species than European beech.~~

To illustrate one possible application of HETEROFOR, a second simulation experiment was achieved and allowed us to compare the radial growth predicted for 2076-2099 according to three IPCC scenarios with that simulated for an historical period (1976-1999). When atmospheric CO₂ concentration was kept constant (380 ppm), differences among scenarios remained non-significant, except for sessile oak displaying a slightly higher basal area increment for the RCP2.6 than for the historical scenario (Fig. 56). Analyzing in-depth the model outputs, we found that this lack of effects resulted from a balance between negative and positive impacts of climate change. While the increase in air temperature (+0.9 and 3.7°C for RCP2.6 and 8.5) and in the vegetation period length (+8 and 37 days for RCP2.6 and 8.5) favoured photosynthesis, the more frequent and intense water stress negatively affected it (de Wergifosse et al., in review). The positive and negative effects of climate change were of the same magnitude for both tree species and offset each other. For the simulations with a variable atmospheric CO₂ concentration, the differences among scenarios were much larger highlighting a strong CO₂ fertilization effect for both sessile oak and European beech (Fig. 5). These results are in agreement with Reyer et al. (2014) who used the 4C model to predict productivity change in Europe according to a large range of climate change projections. They found NPP increases in most European regions (except a few cases in Mediterranean mountains) when considering persistent CO₂ effects by using variable atmospheric CO₂ concentration. Assuming an acclimation of photosynthesis to CO₂ (by maintaining atmospheric CO₂ constant), they predicted increases in Northern, decreases in Southern and ambivalent responses elsewhere in Europe. Similar response patterns were also obtained by Morales et al. (2007). Rötzer et al. (2013) used the BALANCE model to compare the impact of future and current climate conditions on the productivity of beech in Germany and showed a 30% decrease in NPP without considering the rise in atmospheric CO₂ concentration. After evaluating CASTANEA against eddy covariance and tree growth data in a few highly instrumented sites, Davi et al. (2006) simulated the trend in GGP and net ecosystem productivity (NEP) in these sites from 1960 to 2100. For sessile oak and European beech, they obtained a 53% and 67% increase in GPP and NEP, respectively.

Given the magnitude of the CO₂ fertilization effect (leading to a 72% increase in basal area increment in 100 years for RCP8.5), we conducted retrospective simulations to check that HETEROFOR reproduces well the increase in productivity observed by Bontemps et al. (2011) for beech forests in the north-east of France (data not shown). Based on historical atmospheric CO₂ concentrations, we simulated radial growth during two periods (1879-1910 vs 1979-2010) using the same climate data (obtained by re-analysis for 1979-2010). These simulations showed a productivity increase of 12% over 100 years. By comparison, Bontemps et al. (2011) reported productivity increases ranging from 10 to 70% over 100 years depending on the nitrogen status of the forest. The increase in radial growth simulated with HETEROFOR for the mixed stand in Baileux (Fig. 56) seems therefore plausible but assumes unchanged nutritional status. Increased productivity generates however higher nutrient demand by trees, which is not systematically satisfied by larger soil nutrient supply, especially in the poorest sites. Consequently, the augmentation of forest productivity will most likely be constrained by nutrient availability and give rise to a deterioration of the nutritional status as already observed across Europe (Jonard et al., 2015). To improve our predictions,

nutritional constraints must be taken into account. In this perspective, a mineral nutrition and nutrient cycle module was incorporated in HETEROFOR. As it was developed in parallel to the water balance, some adaptations are needed for a perfect coupling of the two modules (e.g., change from an annual to a monthly time step for soil chemistry update). A complete description and evaluation of the nutrient module will be provided in a future study.

5. Conclusion and future prospects

Our ambition was to develop a model responsive to both management actions and environmental changes that would be particularly well adapted to mixed and uneven-aged stands. We thought that this model had to be tree-level, spatially explicit and process-based. Except BALANCE, iLand and more recently NOTG 3D, no such a model existed in the literature and took simultaneously the radiation transfer, the water balance and the nutrient budget into account. To fill this gap, we elaborated the HETEROFOR model based on concepts quite different from those used for BALANCE, iLAND and NOTG 3D. In this study, a first evaluation of the model performances showed that HETEROFOR predicts well individual radial growth and is able to reproduce size-growth relationships. We also noticed that the most empirical option for describing maintenance respiration provides the best results while the process-based and empirical approaches perform similarly for photosynthesis and crown extension. As this model evaluation was limited to two tree species and one climate, it only provides a first overview of the model potential.

Here, only the core of HETEROFOR was described. The water balance and phenology modules are presented and evaluated in a companion paper (de Wergifosse et al., in review) while the nutrient module will be described later. For the next steps, we plan to couple HETEROFOR with existing libraries such as regeneration, genetics and economics. As HETEROFOR was developed within the CAPSIS platform, it is continually improving thanks to the collaborative dynamics among modellers. A broader assessment of the model performances will be carried out based on forest monitoring plots distributed all over Europe. Indeed, HETEROFOR was designed to be particularly suitable for the level II plots of ICP Forests. The processes were described at a scale that facilitates the comparison between model predictions and observations. Many data collected in these plots can be used to initialize and run the model or to calibrate and evaluate it. HETEROFOR can also be seen as a tool for integrating forest monitoring data and quantifying non-measured processes. While it is now calibrated for oak and beech forests, HETEROFOR will be parameterised for a large range of tree species in order to use it for testing and reproducing identity and diversity effects.

Given all the uncertainties related to climate change impacts, it is an illusion to believe that a model will predict accurately the future dynamics of forest growth. However, models such HETEROFOR can be very useful to compare scenarios. Among others, HETEROFOR can be used to select the management options that maximise ecosystem resilience or to quantify uncertainty in the response of forest ecosystem to climate change.

6. Code availability

The source code of CAPSIS and HETEROFOR is accessible to all the members of the CAPSIS co-development community. Those who want to join this community are welcome but must contact François de Coligny (coligny@cirad.fr) or Nicolas Beudez (nicolas.beudez@inra.fr) and sign the CAPSIS charter (<http://capsis.cirad.fr/capsis/charter>). This charter grants access on all the models to the modellers of the CAPSIS community. The modellers may distribute the CAPSIS platform with their own model but not with the models of the others without their agreement. CAPSIS4 is a free software (LGPL licence) which includes the kernel, the generic pilots, the extensions and the libraries. For HETEROFOR, we also choose an LGPL license and decided to freely distribute it through an installer containing the CAPSIS4 kernel and the latest version (or any previous one) of HETEROFOR upon request from Mathieu Jonard (mathieu.jonard@uclouvain.be). The version 1.0 used for this paper is available at <http://amap-dev.cirad.fr/projects/capsis/files>. The end-users can install CAPSIS from an installer containing only the HETEROFOR model while the modellers who signed the CAPSIS charter can access to the complete version of CAPSIS with all the models. Depending on your status (end-user vs modeller or developer), the instructions to install CAPSIS are given on the CAPSIS website (<http://capsis.cirad.fr/capsis/documentation>).

The source code for the modules published in Geoscientific Model Development can be downloaded from https://github.com/jonard76/HETEROFOR_LGPL_PLUS (DOI 10.5281/zenodo.3403280).

7. Data availability

The data used in this paper are available through the input files for HETEROFOR which are embedded in the installer (see sect. 6).

8. Appendices

8.1 Appendix A – Description of the soil heat transfer routine

The temperature of the mineral soil (T in $^{\circ}\text{C}$) is calculated by soil depth increment (Δz in m) using a simplification of the soil heat transfer equation assuming a constant thermal diffusivity (D in $\text{m}^2 \text{s}^{-1}$) across the soil profile. The thermal diffusivity characterizes the rate of heat transfer within the soil and corresponds to the ratio of the thermal conductivity (K in $\text{W m}^{-1} \text{K}^{-1}$) to the volumetric heat capacity (c_v in $\text{J m}^{-3} \text{K}^{-1}$).

$$\frac{\partial T}{\partial t} = \frac{1}{c_v} \cdot \frac{\partial}{\partial z} \cdot \left(K \frac{\partial T}{\partial z} \right) \Rightarrow \frac{\partial T}{\partial t} = D \cdot \frac{\partial^2 T}{\partial z^2} \quad (50)$$

Eq. (50) can be rewritten as follows according to Anlauf and Liu (1990) and Baker and Don Scott (1998):

$$T_{z,t+\Delta t} = T_{z,t} + D \cdot \frac{\Delta t}{\Delta z^2} \cdot (T_{z+\Delta z,t} + T_{z-\Delta z,t} - 2T_{z,t}) \quad (51)$$

The soil depth increment can be chosen by the user but it has to be smaller than one third of the thiniest horizon. The soil depth increment can be slightly modified by the model to ensure the soil depth is a multiple of the soil depth increment. Then, a stability criterion is checked for each hour and if it is not respected, the temporal step is divided by two.

$$K \cdot \frac{\Delta T}{\Delta z^2} < 0.5 \quad (52)$$

The thermal diffusivity is calculated for each soil horizon based on the thermal conductivity and the volumetric heat capacity and then averaged by weighing according the horizon thickness. The thermal conductivity is obtained with the empirical model of Kersten (1949):

$$K = 0.1442 \cdot (0.9 \cdot \log(\theta) - 0.2) \cdot 10^{0.6243\rho_b} \quad (\text{for silt or clay soils}) \quad (53)$$

$$K = 0.1442 \cdot (0.7 \cdot \log(\theta) + 0.4) \cdot 10^{0.6243\rho_b} \quad (\text{for sandy soils}) \quad (54)$$

with θ , the gravimetric soil water content (g g^{-1}),

ρ_b , the bulk density (kg m^{-3}).

The volumetric heat capacity of soils is approximated through a separation of the soil constituents in solid and liquid phases:

$$c_v \simeq 836 \cdot \rho_b + 4180 \cdot \theta \cdot \rho_b \cdot 1000 \cdot \rho_w \quad (55)$$

with ρ_w , the volumetric mass of water (kg m^{-3}).

To initialize the procedure, the top and bottom temperature during the whole simulation and the initial temperature at each soil depth must be known. The soil temperature at the top of the mineral soil (just under the forest floor) is given by Eq. (56) adapted from van Wijk and de Vries (1963) and Cichota et al. (2004). The bottom temperature is fixed and corresponds to the mean annual air temperature. This assumption can be made as the soil depth largely exceeds 1 meter. The initial temperature is found through a simple interpolation of the temperatures between the soil interface and the bottom.

$$T_t = \bar{T}_y + \frac{(\bar{T}_{d-1} - \bar{T}_y)}{A_{\text{air}}} \cdot A_{\text{soil}} + \frac{a_{\text{air}}}{2} \cdot \text{red}_d \cdot \sin\left(\omega(t - t_{T_{\text{max}}}) + \frac{\pi}{2} - \omega \frac{\Delta z}{\text{Damping}}\right) \quad (56)$$

with \bar{T}_y , mean annual air temperature ($^{\circ}\text{C}$),

\bar{T}_{d-1} , mean air temperature of the previous day ($^{\circ}\text{C}$),

A_{air} , annual air temperature amplitude corresponding to the difference between the maximum and the minimum mean daily temperature over the year ($^{\circ}\text{C}$),

5 A_{soil} , parameter corresponding to the mean annual soil temperature amplitude ($^{\circ}\text{C}$),

a_{air} , daily air temperature amplitude ($T_{\text{max}} - T_{\text{min}}$) calculated over the 24 hour period centered on the considered time ($^{\circ}\text{C}$),

red_d , parameter reducing the daily air temperature amplitude to the daily soil temperature amplitude (fixed to 0.13)

ω , radial frequency (h^{-1}) = $\frac{2\pi}{24}$,

10 $t_{T_{\text{max}}}$, hour of the day at which air temperature is maximal (as the sinusoidal shape of the diurnal soil temperature cycle is not perfectly symmetric, $t_{T_{\text{max}}}$ is adapted so that the period between maximum and minimum soil temperature is exactly 12 hours),

Δz , thickness of organic horizons (m),

Damping, parameter accounting for the phase shift between the diurnal cycle of the air and soil temperature (fixed to 0.0853 after calibration).

15

The temperature of the organic horizons was obtained as the mean between air temperature and the temperature at the interface between organic horizons and mineral soil.

8.2 Appendix B – Development of Eq. (29)

Equation (29) can be developed in order to isolate Δh :

$$\Delta(dbh^2 \cdot h) = (dbh + \Delta dbh)^2 \cdot (h + \Delta h) - dbh^2 \cdot h \quad (57)$$

$$\Delta(dbh^2 \cdot h) = (dbh^2 + 2 \cdot dbh \cdot \Delta dbh + (\Delta dbh)^2) \cdot (h + \Delta h) - dbh^2 \cdot h$$

$$5 \quad \Delta(dbh^2 \cdot h) = dbh^2 \cdot h + 2 \cdot dbh \cdot \Delta dbh \cdot h + (\Delta dbh)^2 \cdot h + dbh^2 \cdot \Delta h + 2 \cdot dbh \cdot \Delta dbh \cdot \Delta h + (\Delta dbh)^2 \cdot \Delta h - dbh^2 \cdot$$

$$h \Delta(dbh^2 \cdot h) = \Delta dbh^2 \cdot (h + \Delta h) + (\Delta dbh)^2 \cdot (h + \Delta h) + dbh^2 \cdot \Delta h$$

$$\Delta(dbh^2 \cdot h) = \Delta h \cdot (\Delta dbh^2 + (\Delta dbh)^2 + dbh^2) + h \cdot (\Delta dbh^2 + (\Delta dbh)^2) \quad (58)$$

Considering $(\Delta dbh)^2 \ll \Delta dbh^2 \ll dbh$, the following approximation can be done:

$$10 \quad \Delta(dbh^2 \cdot h) \cong \Delta h \cdot dbh^2 + h \cdot \Delta dbh^2 \quad (59)$$

$$\Delta h \cdot dbh^2 \cong \Delta(dbh^2 \cdot h) - h \cdot \Delta dbh^2 \quad (60)$$

$$\Delta h \cong \frac{\Delta(dbh^2 \cdot h)}{dbh^2} - \frac{h \cdot \Delta dbh^2}{dbh^2} \quad (61)$$

8.3 Appendix C - Delevoy height estimation

The Delevoy height is the height at which stem diameter is half the diameter at breast height and is calculated as follows from taper (cm m⁻¹):

$$h_{del} = 1.3 + \frac{dbh - \frac{dbh}{2}}{taper} \tag{62}$$

where the taper is obtained based on the girth at 10% of the tree height (G10%) and the relative girth at 60% of the tree height (RG60%) for which empirical equations are provided by Dagnelie *et al.* (1999) for several temperate tree species:

$$taper = \frac{(1 - CR60\%) \cdot C10\%}{0.5 \cdot h \cdot \pi} \tag{63}$$

with

$$C10\% = a + b \cdot \pi \cdot dbh + c \cdot (\pi \cdot dbh)^2 + d \cdot (\pi \cdot dbh)^3 + e \cdot h + f \cdot (\pi \cdot dbh)^2 \cdot h \tag{64}$$

$$CR60\% = a + \frac{b}{C10\%} + \frac{c}{C10\%^2} \tag{65}$$

Table C1. Parameters of Eqs. (64) and (65) for sessile oak and European beech according to Dagnelie et al. (1999)

	a	b	c	d	e	f
Sessile oak						
C10%	3.9330	1.0284	-0.31611 10 ⁻³	0.44036 10 ⁻⁶	-0.33113	-0.28051 10 ⁻⁵
CR60%	0.4838	14.667	-405.67			
European beech						
C10%	3.8541	1.0235	-0.36276 10 ⁻³	0.40063 10 ⁻⁶	-0.30551	-0.20411 10 ⁻⁵
CR60%	0.5286	0	0			

8.4 Appendix D – Estimation of the height of largest crown extension ($hlce$) at equilibrium




Figure D1. Illustration of the routine used to determine the height of largest crown extension at equilibrium ($hlce_{eq}$) of a target tree in three contrasted situations of competition. A first step consists in determining the intersection between the potential crown of the target tree and the competitor. Then, the $hlce_{eq}$ is fixed between the maximum $hlce$ (corresponding to the intersection between potential crowns) and the minimum $hlce$ (which is the height to crown base) based on the relative height of the competitor.

8.5 Appendix E – Height growth modelling results

The main factor explaining the height increment was the so-called height growth potential (Δh_{pot}) with a quadratic effect for beech and a cubic effect for oak (Table E1, Fig. E1). For both tree species, the light competition index (LCI) had a negative effect on height increment, meaning that, for a same height growth potential, trees under stronger competition for light had a higher height growth than trees within better light conditions. For European beech, the variable selection procedure led to select height (which had a negative effect) to account for tree size while dbh was retained for sessile oak and had a positive effect. Even if the root mean square error was slightly higher for European beech (0.094) than for sessile oak (0.083), the height growth model explained a much larger proportion of the variability for European beech (72%) than for sessile oak (43%), partly because the height growth range was higher for European beech.

Table E1. Parameters, R^2 and RMSE of the height growth model (Eq. 31) for European beech and sessile oak.

	European beech	Sessile oak
intercept	0.0233	-0.0562
dbh (in cm)		0.0023
h (in m)	-0.0048	
LCI	-0.2556	-0.1874
$(\Delta dbh^2 h_{tot})/dbh^2$ (in m)	0.6631	0.8183
$[(\Delta dbh^2 h_{tot})/dbh^2]^2$	-0.1777	-0.9178
$[(\Delta dbh^2 h_{tot})/dbh^2]^3$		0.4733
RMSE	0.09397	0.083017
R^2	0.72	0.43




Figure E1. Effect of the height growth potential on oak and beech height growth for two levels of light competition (strong light competition = light competition index ≤ 0.15 , lower light competition = light competition index > 0.15). The solid lines represent the model predictions obtained using Eq. (31) with parameter values of Table E1 and with mean values for dbh , height or the light competition index.

9. Author contribution

MJ, FA, FdC, LdW, NB and HD developed the model code. FA carried out the calibration. MJ~~and~~, FA and LdW performed the simulations and analysed the model outputs. MJ prepared the manuscript with contributions from all co-authors.

10. Competing interests

5 The authors declare that they have no conflict of interest.

11. Acknowledgements

This work was supported by the *Service Public de Wallonie* (SPW/DGO 3/DNF) through the *Accord-Cadre de Recherche et Vulgarisation Forestières 2014–2019* and by the *Fonds de la Recherche Scientifique – FNRS* under the PDR-WISD Grant n°09 (project SustainFor) and the FRIA grant n°1.E005.18 (LdW PhD fellowship). We are also grateful to the RMI (Royal

10 Meteorological Institute of Belgium) for providing us climate projection data for the Baileux site.

12. References

- Aber, J., Neilson, R.P., McNulty, S., Lenihan, J.M., Bachelet, D., and Drapek, R.J.: Forest processes and global environmental change: predicting the effects of individual and multiple stressors. *Bioscience*, 51, 735-751, 2001.
- André, F., Jonard, M., and Ponette, Q.: Precipitation water storage capacity in a temperate mixed oak–beech canopy. *Hydrol. Process.*, 22, 4130-4141, 2008a.
- André, F., Jonard, M., and Ponette, Q.: Influence of species and rain event characteristics on stemflow volume in a temperate mixed oak–beech stand. *Hydrol. Process.*, 22, 4455-4466, 2008b.
- André, F., Jonard, M., and Ponette, Q.: Biomass and nutrient content of sessile oak (*Quercus petraea* (Matt.) Liebl.) and beech (*Fagus sylvatica* L.) stem and branches in a mixed stand in southern Belgium. *Sci. Total Environ.*, 408, 2285-2294, 2010.
- André, F., Jonard, M., Jonard, F., and Ponette, Q.: Spatial and temporal patterns of throughfall volume in a deciduous mixed-species stand. *J. Hydrol.*, 400, 244-255, 2011.
- Anlauf, R. and Liu, Y.P.: Simulation of simple one-dimensional transport processes. *In: Richter, J. (Ed.), Models for Processes in the Soil: Programs and Exercises.* Catena Verlag, Cremlingen-Desdedt, Germany, 1990.
- Baker, D.L. and Don Scott, H.: A limited tutorial on using finite differences in soil physics problems, 1998.
- <http://www.aquarien.com/sptutor/findifa8/sld001.htm>, <http://www.aquarien.com/findif/Findifa4.html>.
- Ball, J.T., Woodrow, I.E., and Berry, J.A.: A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions. *In: Biggins, J. (Ed.), Progress in Photosynthesis Research*, vol. 4, 1987.
- Boisvenue, C. and Running, S.W.: Impacts of climate change on natural forest productivity—evidence since the middle of the 20th century. *Global Change Biol.*, 12, 862-882, 2006.
- Bontemps, J.D., Hervé, J.C., Leban, J.M., and Dhôte, J.F.: Nitrogen footprint in a long-term observation of forest growth over the twentieth century. *Trees*, 25, 237–251, 2011.
- Bosc, A.: EMILION, a tree functional-structural model: presentation and first application to the analysis of branch carbon balance. *Ann. For. Sci.*, 57, 555-569, 2000.
- Cantarello, E., Newton, A.C., Martin, P.A., Evans, P.M., Gosal, A., and Lucash, M.S.: Quantifying resilience of multiple ecosystem services and biodiversity in a temperate forest landscape. *Ecol. and Evol.*, 7, 9661-9675, 2017.
- [Carl, C., Biber, P., Veste, M., Landgraf, D. and Pretzsch, H.: Key drivers of competition and growth partitioning among *Robinia pseudoacacia* L. trees. *Forest Ecol. Manage.*, 430, 86-93, 2018.](#)
- Charlton, S.R. and Parkhurst, D.L.: Modules based on the geochemical model PHREEQC for use in scripting and programming languages. *Comput. and Geosci.*, 37, 1653-1663, 2011.
- Campoli, M., Vincke, C., Jonard, M., Kint, V., Demarée, G. and Ponette, Q.: Current status and predicted impact of climate change on forest production and biogeochemistry in the temperate oceanic European zone: review and prospects for Belgium as a case study. *J. For. Res.*, 17, 1-18, 2012.

- Cichota, R., Elias, E.A., and van Lier, Q.D.J.: Testing a finite-difference model for soil heat transfer by comparing numerical and analytical solutions. *Environ. Model. Softw.*, 19, 495-506, 2004.
- Collalti, A., Trotta, C., Keenan, T. F., Ibrom, A., Bond-Lamberty, B., Grote, R., et al.: Thinning can reduce losses in carbon use efficiency and carbon stocks in managed forests under warmer climate. *J. Adv. Model. Earth Sy.*, 10, 2427–2452, 2018.
- 5 [Collalti, A. and Prentice, I. C.: Is NPP proportional to GPP? Waring's hypothesis 20 years on. *Tree Physiol.* 39, 1473-1483, 2019.](#)
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W.J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A.J., and Wehner, M.: Long-term Climate Change: Projections, Commitments and Irreversibility. *In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., Qin, D., Plattner, G.K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 10 Courbaud, B., de Coligny, F., and Cordonnier, T.: Simulating radiation distribution in a heterogeneous Norway spruce forest on a slope. *Agric. For. Meteorol.*, 116, 1-18, 2003.
- 15 Dagnelie, P., Palm, R., Rondeux, J., Thill, A.: *Tables de cubage des arbres et des peuplements forestiers*. Les Presses Agronomiques de Gembloux, Gembloux, Belgique, 1999.
- Damesin, C., Ceschia, E., Le Goff, N., Ottorini, J.M., and Dufrêne, E.: Stem and branch respiration of beech: from tree measurements to estimations at the stand level. *New Phytol.*, 153, 159–172, 2002.
- 20 Davi, H., Dufrêne, E., Francois, C., Le Maire, G., Loustau, D., Bosc, A., Rambal, S., Granier, A., and Moors, E.: Sensitivity of water and carbon fluxes to climate changes from 1960 to 2100 in European forest ecosystems. *Agric. For. Meteorol.*, 141, 35–56, 2006.
- [del Río, M., Bravo-Oviedo, A., Ruiz-Peinado, R. and Condés, S.: Tree allometry variation in response to intra- and inter-specific competitions. *Trees*, 33, 121-138, 2019.](#)
- 25 de Wergifosse, L., André, F., Beudez, N., de Coligny, F., Goosse, H., Jonard, F., Ponette, Q., Titeux, H., Vincke, C., and Jonard, M.: HETEROFOR 1.0: a spatially explicit model for exploring the response of structurally complex forests to uncertain future conditions. II. Phenology and water cycle, *Geosci. Model Dev. Discuss.*, in review, 2019.
- Dufour-Kowalski, S., Courbaud, B., Dreyfus, P., Meredieu, C., and de Coligny, F.: Capsis: an open software framework and community for forest growth modelling. *Ann. For. Sci.*, 69, 221–233, 2012.
- 30 Dufrêne, E., Davi, H., François, C., Le Maire, G., Le Dantec, V., and Granier, A.: Modelling carbon and water cycles in a beech forest: Part I: Model description and uncertainty analysis on modelled NEE. *Ecol. Model.*, 185, 407-436, 2005.
- Duursma, R.A. and Medlyn, B.E.: MAESPA: a model to study interactions between water limitation, environmental drivers and vegetation function at tree and stand levels, with an example application to [CO₂] × drought interactions. *Geosci. Model Dev.*, 5: 919–940, 2012.

- Ennos, R., Cottrell, J., Hall, J., and O'Brien, D.: Is the introduction of novel exotic forest tree species a rational response to rapid environmental change? – A British perspective. *Forest Ecol. and Manag.*, 432, 718-728, 2019.
- Epron, D., Le Dantec, V., Dufrêne, E., and Granier, A.: Seasonal dynamics of soil carbon efflux and simulated rhizosphere respiration in a beech forest. *Tree Physiol.*, 21, 145–152, 2001.
- 5 Farquhar, G.D., von Caemmerer, S., and Berry, J.A.: A biochemical model of photosynthetic CO₂ assimilation in leaves of C3 species. *Planta*, 149, 78–80., 1980.
- Fernandez-Martinez, M., Vicca, S., Janssens, I.A., Sardans, J., Luyssaert, S., Campioli, M., Chapin III, F.S., Ciais, P., Malhi, Y., Obersteiner, M., Papale, D., Piao, S.L., Reichstein, M., Rodà, F., and Peñuelas, J.: Nutrient availability as the key regulator of global forest carbon balance. *Nat. Clim. Chang.*, 4, 471–476, 2014.
- 10 Forrester, D., I. and Albrecht, A.T.: Light absorption and light-use efficiency in mixtures of *Abies alba* and *Picea abies* along a productivity gradient. *For. Ecol. Manag.*, 328, 94-102, 2014.
- Genet, H., Bréda, N., and Dufrêne, E.: Age-related variation in carbon allocation at tree and stand scales in beech (*Fagus sylvatica* L.) and sessile oak (*Quercus petraea* (Matt.) Liebl.) using a chronosequence approach. *Tree Physiol.*, 30, 177–192, 2010.
- 15 Genet, A., Wernsdörfer, H., Jonard, M., Pretzsch, H., Rauch, M., Ponette, Q., Nys, C., Legout, A., Ranger, J., Vallet P., and Saint-André, L.: Ontogeny partly explains the apparent heterogeneity of published biomass equations for *Fagus sylvatica* in central Europe. *For. Ecol. Manag.*, 261: 1188-1202, 2011.
- Giot, O., Termonia, P., Degrauwe, D., De Troch, R., Caluwaerts, S., Smet, G., Berckmans, J., Deckmyn, A., De Cruz, L., De Meutter, P., Duerinckx, A., Gerard, L., Hamdi, R., Van den Bergh, J., Van Ginderachter, M., and Van Schaeybroeck, B.:
20 Validation of the ALARO-0 model within the EURO-CORDEX framework. *Geosci. Model Dev.*, 9, 1143-1152, 2016.
- Grote, R. and Pretzsch, H., A model for individual tree development based on physiological processes. *Plant Biol.*, 4, 167–180, 2002.
- Haxeltine, A. and Prentice, I.C.: A general model for the light use efficiency of primary production by terrestrial ecosystems. *Funct. Ecol.*, 10, 551-561, 1996.
- 25 Helmsaari, H.S., Derome, J., Nöjd, P., and Kukkola, M.: Fine root biomass in relation to site and stand characteristics in Norway spruce and Scots pine stands. *Tree Physiol.*, 27, 1493-504, 2007.
- Henry, H.A.L. and Aarssen, L.W.: The interpretation of stem diameter-height allometry in trees: biomechanical constraints, neighbour effects, or biased regressions? *Ecol. Lett.*, 2, 89-97, 1999.
- Hounzandji, A.P., Jonard, M., Nys, C., Saint-André, L., and Ponette, Q.: Improving the robustness of biomass functions: from
30 empirical to functional approaches. *Ann. For. Sci.*, 72: 795-810, 2015.
- Janssen, P.H.M. and Heuberger, P.S.C.: Calibration of process-oriented models. *Ecol. Model.*, 83, 55-66, 1995.
- Jonard, M., André, F., and Ponette, Q.: Modeling leaf dispersal in mixed hardwood forests using a ballistic approach. *Ecology*, 87, 2306-2318, 2006.

- Jonard, M., André, F., Jonard, F., Mouton, N., Procès, P., and Ponette, Q.: Soil carbon dioxide efflux in pure and mixed stands of oak and beech, *Ann. For. Sci.*, 64(2), 141-150, 2007.
- Jonard, M., André, F., and Ponette, Q.: Tree species mediated effects on leaf litter dynamics in pure and mixed stands of oak and beech, *Can. J. Forest Res.*, 38(3), 528-538, 2008.
- 5 Jonard, F., André, F., Ponette, Q., Vincke, C., and Jonard, M.: Sap flux density and stomatal conductance of European beech and sessile oak trees in pure and mixed stands during the summer drought of 2003. *J. Hydrol.*, 409, 371-381, 2011.
- Jonard, M., Fürst, A., Verstraeten, A., Thimonier, A., Timmermann, V., Potocic, N., Waldner, P., Benham, S. ; Hansen, K. ; Merila, P. ; Ponette, Q. ; De La Cruz, A. C ; Roskams, P., Nicolas, M., Croisé, L., Ingerslev, M., Matteuci, G., Decinti, B., Bascietto, M., and Rautio, P.: Tree mineral nutrition is deteriorating in Europe. *Global Change Biol.*, 21: 418-430, 2015.
- 10 Jucker, T., Bouriaud, O., Coomes, D.A.: Crown plasticity enables trees to optimize canopy packing in mixed-species forest. *Funct. Ecol.*, 29, 1078-1086, 2015.
- Kersten, M.S.: Laboratory research for the determination of the thermal properties of soils. University of Minnesota, Engineering Experiment Station, 1949.
- Le Moguédec, G. and Dhôte, J.F.: Fagacées: a tree-centered growth and yield model for sessile oak (*Quercus petraea* L.) and
 15 common beech (*Fagus sylvatica* L.). *Ann. For. Sci.*, 69, 257-269, 2012.
- Letort, V., Cournède, P.H., Mathieu, A., de Reffye, P., and Constant, T.: Parametric identification of a functional-structural tree growth model and application to beech trees (*Fagus sylvatica*). *Funct. Plant Biol.*, 35, 951-963, 2008.
- Ligot, G., Balandier, P., Courbaud, B., Jonard, M., and Kneeshaw, D.: Managing understory light to maintain a mixture of species with different shade tolerance. *For. Ecol. Manag.*, 327, 189-200, 2014.
- 20 Lindner, M., Maroschek, M., Netherer, S., Kremer, A., Barbati, A., Garcia-Gonzalo, J., Seidl, R., Delzon, S., Corona, P., Kolström, M., Lexer, M.J., and Marchetti, M.: Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *For. Ecol. Manag.*, 259, 698-709, 2010.
- Lindner, M., Fitzgerald, J.B., Zimmermann, N.E., Reyer, C., Delzon, S., van der Maaten, E., Schelhaas, M-J. Lasch, P., Eggers, J., van der Maaten-Theunissen, M., Suckow, F., Psomas, A., Poulter, B., and Hanewinkel, M.: Climate change and
 25 European forests: What do we know, what are the uncertainties, and what are the implications for forest management? *J. Environ. Manage.*, 146, 69-83, 2014.
- Lines, E.R., Zavala, M.A., Purves, D.W., and Coomes, D.A.: Predictable changes in aboveground allometry of trees along gradients of temperature, aridity and competition. *Glob. Ecol. Biogeogr.*, 21, 1017-1028, 2012.
- Mäkelä, A.: Implications of the pipe model theory on dry matter partitioning and height growth in trees. *J. Theor. Biol.*, 123,
 30 103-120, 1986.
- Mäkelä, A. and Valentine, H.T.: The ratio of NPP to GPP: evidence of change over the course of stand development. *Tree Physiol.*, 21, 1015-1030, 2001.
- Manion, P.D.: *Tree Disease Concepts*. Prentice-Hall, Englewood Cliffs, NJ, 402 pp, 1981.

- Maraun, D. and Widmann, M.: Statistical downscaling and bias correction for climate research. Cambridge University Press, Cambridge, 2018.
- Mellert, K.H. and Göttelein, A.: Comparison of new foliar nutrient thresholds derived from van den Burg's literature compilation with established central European references. *Eur. J. For. Res.*, 131, 1461-1472, 2012.
- 5 Messier, C., Puettmann, K., Chazdon, R., Andersson, K.P., Angers, V.A., Brotons, L., Filotas, E., Tittler, R., Parrott, L., and Levin, S.A.: From management to stewardship: viewing forests as complex adaptive systems in an uncertain world. *Conserv. Lett.*, 8: 368-377, 2015.
- Monteith, J.L.: Climate and the efficiency of crop production in Britain. *Philos. Trans. Royal Soc. B*, 281, 277-294, 1977.
- Morales, P., Hickler, T., Rowell, D.P., Smith, B., and Sykes, M.T.: Changes in European ecosystem productivity and carbon balance driven by regional climate model output. *Global Change Biol.*, 13, 108-122, 2007.
- 10 Oliver, T.H., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A., Orme, C.D.L., Petchey, O.L., Proença, V., Raffaelli, D., Suttle, K.B., Mace, G.M., MartínLópez, B., Woodcock, B.A., and Bullock, J.M.: Biodiversity and resilience of ecosystem functions. *Trends Ecol Evol.*, 30, 673-684, 2015.
- Penning de Vries, F.W.T.: The cost of maintenance processes in plant cells. *Ann. Bot.*, 39, 77-92, 1975.
- 15 Pretzsch, H.: Canopy space filling and tree crown morphology in mixed-species stands compared with monocultures. *For. Ecol. Manag.*, 327, 251-264, 2014.
- Pretzsch, H., Grote, R., Reineking, B., Rötzer, T., and Seifert, S.: Models for forest ecosystem management: a European perspective. *Ann. Bot.*, 101, 1065-1087, 2008.
- Pretzsch, H., Forrester, D.I., and Rötzer, T.: Representation of species mixing in forest growth models. A review and perspective. *Ecol. Model.*, 313, 276-292, 2015.
- 20 Reyer, C., Lasch-Born, P., Suckow, F., Gutsch, M., Murawski, A., and Pilz, T.: Projections of regional changes in forest net primary productivity for different tree species in Europe driven by climate change and carbon dioxide. *Ann. For. Sci.*, 71, 211-225. 2014.
- Rodríguez-Calcerrada, J., Martin-StPaul, N.K., Lempereur, M., Ourcival, J.M., Rey, M. del C. del, Joffre, R., and Rambal, S.: Stem CO₂ efflux and its contribution to ecosystem CO₂ efflux decrease with drought in a Mediterranean forest stand. *Agric. For. Meteorol.*, 195-196, 61-72, 2014.
- 25 Rodríguez-Calcerrada, J., López, R., Salomón, R., Gordaliza, G.G., Valbuena-Carabaña, M., Oleksyn, J., and Gil, L.: Stem CO₂ efflux in six co-occurring tree species: underlying factors and ecological implications. *Plant Cell Environ.*, 38, 1104-1115, 2015.
- 30 Rötzer, T., Liao, Y., Goergen, K., Schüler, G., and Pretzsch, H.: Modelling the impact of climate change on the productivity and water-use efficiency of a central European beech forest. *Clim. Res.*, 58, 81-95, 2013.
- Ryan, M.G.: Effects of climate change on plant respiration. *Ecol. Appl.*, 1, 157-167, 1991.
- Ryan, M.G. and Yoder, B.J.: Hydraulic limits to tree height and tree growth. What keeps trees from growing beyond a certain height? *BioScience*, 47, 235-242, 1997.

- Schäfer, K.V.R., Oren, R., and Tenhunen, J.D.: The effect of tree height on crown level stomatal conductance. *Plant Cell Environ.*, 23, 365–375, 2000.
- Schwalm, C.R., and Ek, A.R.: A process-based model of forest ecosystems driven by meteorology. *Ecol. Model.*, 179, 317–348, 2004.
- 5 Simioni, G., Marie, G., and Huc, R.: Influence of vegetation spatial structure on growth and water fluxes of a mixed forest: Results from the NOTG 3D model. *Ecol. Model.*, 328, 119–135, 2016.
- Team, R.C.: R: A language and environment for statistical computing, 2013.
- Teh, C. B. S.: Introduction to mathematical modeling of crop growth. How the equations are derived and assembled into a computer program, Brown Walker Press, Boca Raton, Florida, USA, 2006.
- 10 Tetens, O.: Über einige meteorologische Begriffe. *z. Geophys.*, 6, 297–309, 1930.
- Thompson, I., Mackey, B., McNulty, S., and Mosseler, A.: Forest resilience, biodiversity, and climate change. A synthesis of the biodiversity/resilience/stability relationship in forest ecosystems. Secretariat of the Convention on Biological Diversity, Montreal. Technical Series no. 43, 2009.
- 15 [Thurm, E. A., Biber, P. and Pretzsch, H.: Stem growth is favored at expenses of root growth in mixed stands and humid conditions for Douglas-fir \(*Pseudotsuga menziesii*\) and European beech \(*Fagus sylvatica*\). *Trees-Struct. Funct.*, 31, 349-365, 2017.](#)
- Trouvé, R., Bontemps, J.D., Seynave, I., Collet, C., and Lebourgeois, F.: Stand density, tree social status and water stress influence allocation in height and diameter growth of *Quercus petraea* (Liebl.). *Tree Physiol.*, 35, 1035–1046, 2015.
- Van Wijk, W.R. and De Vries, D.A.: Periodic temperature variations in a homogenous soil. *In: Van Wijk, W.R. (Ed.), Physics of Plant Environment*. North Holland Publishing Company, Amsterdam, 1963.
- 20 Villar, R., Held, A.A., and Merino, J.: Dark leaf respiration in light and darkness of an evergreen and a deciduous plant species. *Plant Physiol.*, 107, 421–427, 1995.
- Voltaire, A., Sanchez-Gomez, E., Salas y Mélia, D., Decharme, B., Cassou, C., Sénési, S., Valcke, S., Beau, I., Alias, A., Chevallier, M., Déqué, M., Deshayes, J., Douville, H., Fernandez, E., Madec, G., Maisonnave, E., Moine, M.P., Planton, S., Saint-Martin, D., Szopa, S., Tyteca, S., Alkama, R., Belamari, S., Braun, A., Coquart, L., and Chauvin, F.: The CNRM-CM5.1 global climate model: description and basic evaluation. *Clim. Dyn.*, 40: 2091–2121, 2013.
- 25 Vose, J.M. and Bolstad, P.V.: Challenges to modelling NPP in diverse eastern deciduous forests: species-level comparisons of foliar respiration responses to temperature and nitrogen. *Ecol. Model.*, 122, 165–174, 1999.
- Yamori, W., Hikosaka, K., and Way, D.A.: Temperature response of photosynthesis in C3, C4, and CAM plants: temperature acclimation and temperature adaptation. *Photosynth. Res.*, 119, 101–117, 2013.
- 30

Table 1. Stand characteristics for the main tree species derived from stand inventories in 2001. Standard deviation is provided in parentheses.

Stand	Tree species	Tree density (N/ha)	Basal Area (m²/ha)	Gbh¹ (cm)	Dominant height (m)
Oak dominated (0.90 ha)	Sessile oak	187	16.2	100.6 (26.5)	21.9
	European beech	118	4	46.4 (35.6)	15.5
Beech dominated (1.44 ha)	Sessile oak	72	6.4	103.3 (18.1)	23
	European beech	217	16.5	87.5 (41.5)	25
Mixed (1.80 ha)	Sessile oak	118	12.9	115.5 (21.0)	24.5
	European beech	352	17	91.2 (39.3)	25.7

¹ Girth at breast height

Table 2. Description of model parameters for sessile oak and European beech and origin of their value.

Symbol	Description	Units	Value		Origin
			Sessile oak	European beech	
Carbon fixation					
k	extinction coefficient	m ⁻¹		0.53	fitted with tree growth data of the study site
PUE _{sl}	PAR use efficiency of sunlit leaves	kgC mol photons ⁻¹	0.00006	0.000216	fitted with tree growth data of the study site
PUE _{sh}	PAR use efficiency of shaded leaves	kgC mol photons ⁻¹	0.00105	0.000584	fitted with tree growth data of the study site
Respiration					
a _{sapwood}	parameters of the sapwood area function (a/b/c in Eq. 12)		0.00/1.54/0.16	0.00/0.00/0.52	fitted with data from André <i>et al.</i> (2010)
r _{npp_gpp}	parameters of the npp to gpp ratio function (α/β in Eq. 8)		0.997/-0.386	0.959/-0.408	fitted with tree growth data of the study site
R _{Tref}	maintenance respiration per g of N at the reference temperature (15°C)	mole CO ₂ gN ⁻¹ h ⁻¹	0.000079	0.000057	fitted with tree growth data of the study site
R _{gr}	growth respiration per unit biomass increment	kgC kgC ⁻¹		0.2	Dufrène <i>et al.</i> (2005)
Q _{10_leaf or fine root}	temperature dependence coefficient of leaf and fine root respiration	dimensionless		2.1	Vose and Bolstad (1999)
Q _{10_stem and root}	temperature dependence coefficient of stem and root respiration	dimensionless		1.7	Epron <i>et al.</i> (2001)
Q _{10_branch}	temperature dependence coefficient of branch respiration	dimensionless		2.8	Damesin <i>et al.</i> (2002)
Carbon allocation					
b _{leaf}	parameters of the leaf biomass function (a/β/γ in Eq. 15)	kgC	0.0026/1.96/1.96	1.469/2.00/0.00	Jonard <i>et al.</i> (2006)
b _{structural_above}	parameters of the aboveground structural biomass (a/β/γ in Eq. 26)	kgC	0.000/263.4/0.969	0.056/292.8/0.966	Hounzandj <i>et al.</i> (2015) and Genet <i>et al.</i> (2011)
r _{root_shoot}	root to shoot ratio	kgC kgC ⁻¹		0.18	Genet <i>et al.</i> (2010)
r _{fr_leaf_min}	minimum fine root to leaf ratio	kgC kgC ⁻¹		0.5	literature data compilation
r _{fr_leaf_max}	maximum fine root to leaf ratio	kgC kgC ⁻¹		2.5	literature data compilation
δ _{leaf}	leaf relative loss rate	kgC kgC ⁻¹ yr ⁻¹		1	
δ _{fr}	fine root relative loss rate	kgC kgC ⁻¹ yr ⁻¹		1	Grote and Pretzsch (2002)
f	stem form factor	m ³ m ⁻³		0.52	Hounzandj <i>et al.</i> (2015) and Genet <i>et al.</i> (2011)
ρ	stem volumetric mass	kgC m ³	562.17	556	Hounzandj <i>et al.</i> (2015) and Genet <i>et al.</i> (2011)
r _{Tleaf}	leaf retranslocation rate	kgC kgC ⁻¹ yr ⁻¹	0.4	0.45	determined based on tree foliage data collected in the study site
r _{Troot}	fine root retranslocation rate	kgC kgC ⁻¹ yr ⁻¹	0.4	0.45	same values as leaves
S _{branch}	parameters of the branch mortality function (a/β/γ as in Eq. 15)	kgC	6.0E-9/3.064/3.064	5.00E-5/2.681/0.00	fitted with data from André <i>et al.</i> (2010)
p _{fruit}	parameters of the fruit production function (α/β in Eq. 22)	kgC	9.50E-4/2.5	8.00E-4/2.5	fitted with litterfall data from ICP Forests level II plots of Wallonia
dbh _{threshold}	threshold dbh for fruit production	cm		25	field observations
Tree dimension increment					
hlce%	fraction of the total height corresponding to the height of largest crown extension	m m ⁻¹	0.81	0.77	determined based on tree inventory data of the study site
hcb%	fraction of the total height corresponding to the crown base height	m m ⁻¹	0.7	0.61	determined based on tree inventory data of the study site
Dd	parameters of the crown to stem diameter function (α/β/γ/δ in Eq. 10)	m m ⁻¹	16.20/0.0280/0.00/0.00	10.49/0.00/1379/-2881	determined based on tree inventory data of the study site
sh	coefficient used to shift the mean crown to stem diameter ratio relationship to its maximum	dimensionless	1.25	1.5	determined based on tree inventory data of the study site
r _{overlapping}	mean crown overlapping ratio	m m ⁻¹	1	1.2	determined based on tree inventory data of the study site
Δhlc _{max}	maximum annual change in hlce	m yr ⁻¹		0.5	determined based on tree growth data of the study site
Δhcb _{max}	maximum annual change in hcb	m yr ⁻¹		0.5	determined based on tree growth data of the study site

Table 3. Statistical evaluation of predicted basal area increments (vs observations) for various combinations of model options using normalized average error (NAE), paired *t*-test, regression test, root mean square error (RMSE) or Pearson's correlation (Pearson's *r*). Standard deviation or confidence intervals are provided in parentheses.

Model options	NAE	Paired <i>t</i> -test	Orthogonal regression		RMSE	Pearson's <i>r</i>
Tree species		<i>P</i> value	intercept	slope		
<i>Castanea/npp to gpp ratio/distance-independent crown extension</i>						
European beech	0.159	0.00	1.59 (± 1.06)	0.75 (± 0.05)	8.64	0.87
Sessile oak	-0.052	0.18	5.28 (± 2.85)	0.75 (± 0.14)	9.33	0.63
<i>Castanea/npp to gpp ratio/distance-dependent crown extension</i>						
European beech	0.090	0.04	1.75 (± 1.36)	0.79 (± 0.07)	8.95	0.83
Sessile oak	-0.020	0.61	4.54 (± 3.06)	0.77 (± 0.14)	9.11	0.63
<i>Castanea/T° dependent maintenance respiration/distance-independent crown extension</i>						
European beech	0.426	0.00	3.55 (± 2.06)	0.53 (± 0.07)	17.97	0.74
Sessile oak	-0.013	0.79	7.18 (± 3.02)	0.62 (± 0.13)	11.03	0.59
<i>Castanea/T° dependent maintenance respiration/distance-dependent crown extension</i>						
European beech	0.544	0.00	2.89 (± 2.07)	0.53 (± 0.06)	18.60	0.77
Sessile oak	0.054	0.25	6.07 (± 3.42)	0.64 (± 0.14)	11.07	0.58
<i>PUE/npp to gpp ratio/distance-independent crown extension</i>						
European beech	0.007	0.85	1.60 (± 1.19)	0.86 (± 0.06)	7.64	0.85
Sessile oak	-0.181	0.00	2.93 (± 4.14)	1.02 (± 0.24)	9.04	0.54
<i>PUE/npp to gpp ratio/distance-dependent crown extension</i>						
European beech	-0.110	0.01	1.84 (± 1.57)	0.96 (± 0.09)	8.41	0.79
Sessile oak	-0.223	0.00	1.76 (± 5.61)	1.16 (± 0.35)	9.67	0.45
<i>PUE/T° dependent maintenance respiration/distance-independent crown extension</i>						
European beech	0.182	0.01	4.17 (± 2.37)	0.61 (± 0.09)	15.43	0.68
Sessile oak	-0.172	0.00	4.65 (± 3.85)	0.90 (± 0.21)	9.31	0.55
<i>PUE/T° dependent maintenance respiration/distance-dependent crown extension</i>						
European beech	0.223	0.00	3.36 (± 2.47)	0.64 (± 0.09)	14.73	0.71
Sessile oak	-0.176	0.00	4.06 (± 5.02)	0.95 (± 0.28)	9.79	0.47