

**Development and testing of scenarios for implementing land use and land cover changes during the Holocene in Earth System Model Experiments**

Sandy P. Harrison<sup>1</sup>, Marie-José Gaillard<sup>2</sup>, Benjamin D. Stocker<sup>3,4</sup>, Marc Vander Linden<sup>5</sup>, Kees Klein Goldewijk<sup>6,7</sup>, Oliver Boles<sup>8</sup>, Pascale Braconnot<sup>9</sup>, Andria Dawson<sup>10</sup>, Etienne Fluët-Chouinard<sup>11</sup>, Jed O. Kaplan<sup>12,13</sup>, Thomas Kastner<sup>14</sup>, Francesco S.R. Pausata<sup>15</sup>, Erick Robinson<sup>16</sup>, Nicki J. Whitehouse<sup>17</sup>, Marco Madella<sup>18,19,20</sup>, Kathleen D. Morrison<sup>8</sup>

Ms for Geoscientific Model Development (PMIP special issue)

1: Department of Geography and Environmental Science, University of Reading, Reading, UK

2: Department of Biology and Environmental Science, Linnaeus University, Kalmar, Sweden

3: Ecological and Forestry Applications Research Centre, Cerdanyola del Vallès, Spain

4: Department of Earth System Science, Stanford University, Stanford, CA 94305, USA

5: Department of Archaeology, University of Cambridge, UK

6: PBL Netherlands Environmental Assessment Agency, The Hague, The Netherlands

7: Copernicus Institute of Sustainable Development, Utrecht University, The Netherlands

8: University Museum of Archaeology & Anthropology, University of Pennsylvania, Philadelphia, USA

9: Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

10: Department of General Education, Mount Royal University, Calgary, Canada

11: Department of Earth System Science, Stanford University, California, USA

12: Department of Earth Sciences, University of Hong Kong, Hong Kong

13: Institute of Geography, University of Augsburg, Augsburg, Germany

14: Senckenberg Biodiversity and Climate Research Centre, Frankfurt am Main, Germany

15: Centre ESCER, Department of Earth and Atmospheric Sciences, University of Quebec in Montreal, Montreal, Canada

16: Department of Anthropology, University of Wyoming, Laramie, Wyoming, USA

17: School of Geography, Earth and Environmental Science, University of Plymouth, Plymouth, UK

18: Department of Humanities (CaSEs), University Pompeu Fabra, Barcelona, Spain

19: ICREA Passeig Lluís Companys 23 08010 Barcelona, Spain

20: School of Geography, Archaeology and Environmental Studies, University of Witwatersrand, Johannesburg, South Africa


**Abstract:** Anthropogenic changes in land use and land cover (LULC) during the pre-industrial Holocene could have affected regional and global climate. Existing scenarios of LULC changes during the Holocene are based on relatively simple assumptions and highly uncertain estimates of population changes through time. Archaeological and palaeoenvironmental reconstructions have the potential to refine these assumptions and estimates. The Past Global Changes (PAGES) LandCover6k initiative is working towards improved reconstructions of LULC globally. In this paper, we document the types of archaeological data that are being collated and how they will be used to improve LULC reconstructions. Given the large methodological uncertainties involved, both in reconstructing LULC from the archaeological data and in implementing these reconstructions into global scenarios of LULC, we propose a protocol to evaluate the revised scenarios using independent pollen-based reconstructions of land cover and climate. Further evaluation of the revised scenarios involves carbon-cycle model simulations to determine whether the LULC reconstructions are consistent with constraints provided by ice-core records of CO<sub>2</sub> evolution and modern-day LULC. Finally, the protocol outlines how the improved LULC reconstructions will be used in palaeoclimate simulations in the Palaeoclimate Modelling Intercomparison Project to quantify the magnitude of anthropogenic impacts on climate through time and ultimately to improve the realism of Holocene climate simulations.

## 1 Introduction and Motivation

Today, ca 10% the ice-free land surface is estimated to be intensively managed and much of the reminder is under less intense anthropogenic use or influenced by human activities (Arneth et al., 2019). Substantial transformations of natural ecosystems by humans began with the geographically diachronous shift from hunting and gathering characteristic of the Mesolithic to cultivation and more permanent settlement during the Neolithic period (Mazoyer and Roudart, 2006; Zohary et al., 2012; Tauger, 2013; Maezumi et al. 2018), although there is controversy about the relative importance of climate changes and human impact on landscape development both during and since that time. Resolving the uncertainty about the extent and timing of land use is important because changes in land cover as a result of land use (land use land cover: LULC) have the potential to impact climate and the carbon cycle. Direct climate impacts occur through changes in the surface-energy budget resulting from modifications of surface albedo, evapotranspiration, and canopy structure (biophysical impacts, e.g. Pongratz et al., 2010; Myhre et al., 2013; Perugini et al., 2017). LULC affects the carbon cycle through modifications in vegetation and soil carbon storage (biogeochemical impacts, e.g. Pongratz et al., 2010; Mahowald et al., 2017) and turnover times, which changes the C sink/source capacity of the terrestrial biosphere. LULC changes have contributed substantially to the increase in atmospheric greenhouse gases during the industrial period (Le Quéré et al., 2018). It has been suggested that greenhouse gas emissions associated with Neolithic LULC changes were sufficiently large to offset climate cooling after the Mid-Holocene (the overdue-glaciation hypothesis: Ruddiman 2003). Although this has been challenged for several reasons, including inconsistency with the land carbon balance derived from ice-core and peat records (e.g. Joos et al., 2004; Kaplan et al., 2011; Singarayer et al., 2011; Mitchell et al., 2013; Stocker et al. 2017), a LULC impact on climate in more recent millennia appears more plausible.

Climate model simulations have shown that LULC changes have discernible impacts on climate, both in regions with large prescribed changes in LULC and in teleconnected regions with no major local human activity (Vavrus et al., 2008; Pongratz et al., 2010; He et al., 2014; Smith et al., 2016). At the global scale, the biogeophysical effects of the accumulated LULC change during the Holocene which resulted in reconstructed land cover patterns in 1850 CE have been estimated to cause a slight cooling (0.17 °C) that is offset by the biogeochemical warming (0.9 °C), giving a net global warming (0.73 °C) (He et al., 2014). However, in these simulations, biophysical and biogeochemical effects were of comparable magnitude in the most intensively altered landscapes of Europe, Asia, and North America (He et al., 2014). Using parallel simulations, with and without LULC changes, Smith et al. (2016) showed that detectable temperature changes due to LULC could have occurred as early as 7000 years ago (7ka BP) in summer and throughout the year by 3ka BP. All of these conclusions, however, are obviously contingent on the imposed LULC forcing, which is highly uncertain.

There have been several attempts to map LULC changes through time (e.g. Ramankutty and Foley, 1999; Pongratz et al., 2008; Kaplan et al., 2011; Klein Goldewijk et al. 2011; Klein Goldewijk et al. 2017a, b). All of these reconstructions assume that anthropogenic land use is a function of population density and the suitability of land for crops and/or pasture. They then use estimates of regional population trends through time in combination with assumptions about per-capita land use and spatial land use allocation schemes to estimate anthropogenic changes in LULC across time and space. However, differences in the underlying assumptions about land-use per capita, which are generalized from limited and often site-specific data, have resulted in large differences in the final reconstructions (Gaillard et al., 2010; Kaplan et al., 2017). Hence, there are still very large uncertainties about the timing and magnitude of LULC changes, both at a global and at a regional scale (Figure 1).



**Figure 1:** Land use at ca 6000 years ago (6ka BP, 4000 years BCE) from the two widely used global historical land-use scenarios HYDE 3.2 (top panel, Klein Goldewijk et al. 2017a) and KK10 (bottom panel, Kaplan et al. 2011), illustrating the large disagreement between LULC scenarios at a regional scale. In both scenarios, the land-sea mask and lake areas are for the present day.

There is a wealth of archaeological, historical and palaeo-vegetation data that could be used to improve the relatively simple rules used to generate global LULC reconstructions. For example, settlement density and numbers of radiocarbon-dated artifacts can be used to infer population sizes and their temporal dynamics (Rick, 1987; Williams, 2012; Silva and Vander Linden, 2017). Carbonised and waterlogged plant remains and animal bones can be used to infer the occurrence and nature of agriculture at a site, although their presence provides no quantitative information about the area under cultivation (Wright, 2003; Lyman 2008; Orton et al., 2016). Although the record of LULC is likely to be patchy and incomplete, because of preservation and sampling issues, systematic use of archaeological data is one important way to improve current LULC scenarios.

119 The Past Global Change (PAGES, <http://www.pastglobalchanges.org/>) LandCover6k Working  
120 Group (<http://pastglobalchanges.org/ini/wg/landcover6k>) is currently working to develop a rigorous  
121 and robust approach to provide data and data products that can be used to inform the development of  
122 LULC scenarios (Gaillard et al., 2018). LULC changes are taken into account in climate-model  
123 simulations currently being made in the current phase of the Coupled Model Intercomparison Project  
124 (CMIP6) for the historic period and the future scenario runs (Eyring et al., 2016). They are also  
125 included in climate-model simulations of the past millennium (Jungclauss et al., 2017), in order to  
126 ensure that these runs mesh seamlessly with the historic simulations. However, the Land Use  
127 Harmonisation data set (LUH2: Hurtt et al., 2017) only extends back to 850 CE and thus scenarios of  
128 LULC changes are currently not included in the CMIP6 palaeoclimate simulations, including mid-  
129 Holocene simulations, that are used as a test of how well state-of-the-art climate models reproduce  
130 large climate changes. In this paper, we discuss how archaeological data will be used to improve  
131 global LULC scenarios for the Holocene. Given that there are large uncertainties associated with the  
132 primary data and further uncertainties may be introduced when this information is used to modify  
133 existing LULC scenarios, we outline a series of tests that will be used to evaluate whether the revised  
134 LULC scenarios are consistent with the changes implied by independent pollen-based reconstructions  
135 of land cover and whether they produce more realistic estimates of both carbon cycle and climate  
136 changes. Finally, we present a protocol for implementing LULC in Earth System Model simulations  
137 to be carried out in the current phase of the Palaeoclimate Modelling Intercomparison Project (PMIP:  
138 Otto-Bleisner et al., 2017; Kageyama et al., 2018). However, the data sets and protocol will also be  
139 useful in later phases of other CMIP projects, including the Land Use Model Intercomparison Project  
140 (LUMIP) and the Land Surface, Snow and Soil Moisture Model Intercomparison Project (LS3MIP)  
141 (Lawrence et al., 2016; van den Hurk et al., 2016).


142  
143

## 144 **2 LandCover6k Methodology**

145

146 The primary source of information about human exploitation of the landscape comes from  
147 archaeological data. In general, these data are site specific and spatiotemporal coverage is often  
148 patchy, and the types and quality of evidence available vary between sites and regions. Generalising  
149 from site-specific data to landscape or regional scales involves making assumptions about human  
150 behaviour and cultural practices. Because of the inherent uncertainties, we advocate an iterative  
151 approach to incorporate archaeological data into LULC scenarios in LandCover6k (Fig. 2). We  
152 propose to revise the existing LULC scenarios by incorporation of diverse archaeological inputs (Fig.  
153 2, phase 1; see Sections 3 and 4) and to test the revised LULC scenarios for their plausibility and  
154 consistency with other lines of evidence (Fig. 2, phase 2 with iterative testing; see Sections 5-7). As  
155 a first test, the revised LULC scenarios of the extent of cropland and grazing land through time will  
156 be compared with independent data on land-cover changes, specifically pollen-based reconstructions  
157 of the extent of open land (see e.g. Trondman et al., 2015; Kaplan et al., 2017) (Section 5). Further  
158 testing the LULC scenarios involves sensitivity tests using global climate models (Section 6) and  
159 global carbon cycle models (Section 7). While the computational cost of the climate-model  
160 simulations can be minimized using equilibrium time-slice simulations, the carbon cycle constraint  
161 relies on transient simulations, but may be derived from uncoupled, land-only simulations. Simulated  
162 climates at key times can be evaluated against reconstructions of climate variables (e.g. Bartlein et  
163 al., 2011) (Section 6). The parallel evolution of CO<sub>2</sub> and its isotopic composition ( $\delta^{13}\text{C}$ ) can be used  
164 to derive the carbon balance of the terrestrial biosphere and the ocean separately (Elsig et al., 2009)  
165 and, in combination with estimates for other contributors to land carbon changes such as C  
166 sequestration by peat buildup, provides a strong constraint on the evolution of LULC through time.  
167 An under- or over-prediction of anthropogenic LULC-related CO<sub>2</sub> emissions during a specific  
168 interval results in consequences for the dynamics of the atmospheric greenhouse gas burden in

subsequent times (Stocker et al., 2017) (Section 7). Thus, these tests can be used to identify issues in the original archaeological datasets and/or the way these data were incorporated into the LULC scenarios that require further refinement. Phase 3 of the project (Fig. 2) provides a protocol for the implementation of the revised LULC scenarios in Earth System Model simulations (Section 8).



**Figure 2:** Proposed scheme for developing robust LULC scenarios through iterative testing and refinement, as input to Earth System Model (ESM) simulations. The archaeological inputs developed in Phase 1 can be used independently or together to improve the LULC reconstructions; iterative testing of the LULC scenario reconstruction (Phase 2) will ensure that these inputs are reliable before they are used of ESM simulations (Phase 3). The uppermost three LULC simulations capitalize on already planned baseline simulations without LULC; the lowermost two simulations are envisaged as new sensitivity experiments.


### 3 Archaeological data inputs

LandCover6k is creating a number of products that will be used to improve the LULC scenarios (Figure 2). Here, we summarise the important features of these data products before showing how they will be incorporated within a scenario-development framework.

#### 3.1 Population dynamics from $^{14}\text{C}$ data

Radiocarbon is the most routinely used absolute dating technique in archaeology, especially for the Holocene. Many thousands of radiocarbon dates are available from the archaeological literature. A number of regional and pan-regional initiatives are compiling these records through exhaustive survey of the archaeological literature (e.g. the Canadian Archaeological Radiocarbon Database:

194 <https://www.canadianarchaeology.ca/>). Statistical approaches, such as summed probability  
 195 distributions (SPDs), can then be used to infer past demographic fluctuations from these compilations  
 196 (Figure 3). This method assumes that the more people there were, the more remains of their various  
 197 activities they left behind, and that this is directly reflected in the number of samples excavated and  
 198 dated (Rick, 1987; Robinson et al., 2019). There are biases that could affect the expected one-to-one  
 199 relationship between number of people and number of radiocarbon dates on archaeological material,  
 200 including lack of uniform sampling through time and space caused by different archaeological  
 201 research interests and traditions in different regions and increased preservation issues with increasing  
 202 age, but these can be minimised through auditing the datasets. Assessment of the robustness of  
 203 population reconstructions through time can be made statistically, by comparing a null hypothesis of  
 204 demographic growth constructed from an exponential fit to the data with the actual record of number  
 205 of dates through time (Shennan et al., 2013; Timpson et al., 2014). Mathematical simulations show  
 206 that the method is relatively robust for large sample sizes (Williams, 2012). Radiocarbon dates have  
 207 been successfully used in several regions to identify population fluctuations associated with hunter-  
 208 gatherers (Japan: Crema et al., 2016) and the introduction of farming and subsequent changes in  
 209 farming regimes (western Europe: Shennan et al., 2013; Wyoming: Zahid et al., 2016; South Korea:  
 210 Oh et al., 2017; see also Freeman et al., 2018) as well as climatic oscillations (Ireland: Whitehouse et  
 211 al., 2014).  
 212



213  
 214 **Figure 3:** *Reconstruction of changes in population size in the Iberian Peninsula during the Holocene*  
 215 *(9000 to 2000 BP, 9ka to 2ka BP) using summed probability distributions (SPDs) of radiocarbon*  
 216 *dates (data after Balsera et al., 2015). The red line indicates the onset of agriculture in the region.*  
 217 *The lower panels show areas under human use at 6ka (left) and 4ka (right) using kernel density*  
 218 *estimates, where the white dots are actual archaeological sites and the shading shows the implied*  
 219 *density of occupation.*


### 3.2 Date of first agriculture

Radiocarbon dates can also be used to track the timing and process of dispersal events, such as the diffusion of plant and animal domesticates from their initial centres of domestication. Since the distribution of samples is often patchy, geostatistical techniques such as kriging and splines are used to spatially interpolate the information in order to provide quantitative estimates of the timing of spread. Work carried out in Europe (Bocquet-Appel et al., 2009), Asia (Silva et al., 2015), and Africa (Russell et al., 2014) demonstrates that there are different rates of diffusion even within a region, reflecting the possible impact of natural features (e.g. waterways, elevation, ecology) on diffusion rates (Davison et al., 2006; Silva and Steele, 2014). Numerous studies provide robust local estimates for the earliest regional occurrence of agriculture and these are being synthesized to provide a global product within LandCover6k (Figure 2).

### 3.3 Global land-use and livestock maps

Maps of the distribution of archaeological sites or of areas linked to a given food production system have been produced for individual site catchments or small regions (e.g. Zimmermann et al., 2009; Barton et al., 2010; Kay et al., 2019). LandCover6k is developing global land-use maps for specific time windows, using a global hierarchical classification of land-use categories (Morrison et al., 2018) based on land-use types that are widely recognised from the archaeological record. At the highest level, the maps distinguish between areas where there is no (or only limited) evidence of land use, and areas characterized by hunting/foraging/fishing activities, pastoralism, agriculture, and urban/extractive land use (Fig. 4). Except in the cases where land use is minimal (no human land use, extensive/minimal land use), further distinctions are subsequently made to encompass the diversity of land-use activities in each land-use type (Fig. 4). A third level of distinction is made in the case of two categories (agroforestry, wet cultivation) where there are very different levels of intervention in different regions. Explanations of this terminology are given in Morrison et al. (2018). The LandCover6k land-use maps (see e.g. Fig. 5) will be based on different methods ranging from kernel-density estimates to expert assessments depending on the quality and quantity of the archaeological information available from different regions.





**Figure 4:** The hierarchical scheme of land-use classes used for global mapping in LandCover6k (updated from Morrison et al., 2018).


There is considerable variation in how intensely land is used both for crops and for grazing within broad land-use categories both geographically and through time (Ford and Clarke, 2015; Styring et al., 2017). Maps of land-use types do not provide direct information on the intensity of farming practices or how they translate into per-capita land use. Archaeological data about agricultural yields, combined with information from analogous contemporary cultures, historical information (e.g. Pongratz et al., 2008) and theoretical estimates of land use required to meet dietary and energy requirements (e.g. Hughes et al., 2018), can be used to provide regional estimates of per-capita land use for specific land-use categories. LandCover6k will synthesise this information to allow regionally specific estimates of per-capita land use to be derived from the global land-use maps.

Information about the extent of grazing land is an important input for the development of revised LULC scenarios but, from a carbon-cycle modelling perspective, the amount of biomass removed by grazing is also a key parameter. Biomass loss varies not only with population size but also with the type of animal being reared (Herrero et al., 2013; Phelps & Kaplan, 2017) and thus information about what animals were present at a given location and estimates of population sizes are needed for improving the existing LULC scenarios. Although the conditions of bone preservation vary across the globe due to factors such as soil acidity, animal bones are routinely excavated (Lyman, 2008;

272 Reitz & Wing, 2008). Morphometric analysis of bones, along with collateral information such as age-  
 273 related culling patterns, make it possible to determine whether these are the remains of domesticated  
 274 species. We thus have a relatively precise idea of when livestock were introduced into a region, what  
 275 types of animal were being reared at a given time, and can also make informed estimates of population  
 276 size. Although the level of detail will vary geographically, this information can be used to produce  
 277 global livestock maps.

278  
 279 The harvesting of wood for domestic fires, building, and for industrial activities such as  
 280 transportation, pottery-making and metallurgy is an important aspect of human exploitation of the  
 281 landscape in the pre-industrial period (McGrath et al., 2015). It has been argued that even Mesolithic  
 282 hunter-gatherer communities shaped their environment through wood harvesting (Bishop et al.,  
 283 2015). Approaches have been developed to quantifying the wood harvest associated with  
 284 archaeological settlements at specific times based on the evidence of types of wood use, household  
 285 energy requirements, population size, and calorific value of the wood used (see e.g. Marston, 2009;  
 286 Janssen et al., 2017). However, quantitative information on ancient technology and lifestyle is sparse  
 287 and direct estimates of the amount of wood harvest through time are likely to remain highly uncertain  
 288 (Marston et al., 2017; Veal, 2017). Nevertheless, combining evidence-based inferential approaches  
 289 with improved estimates of population size should allow changes in wood harvesting to be taken into  
 290 account in constructing revised LULC scenarios.

291




292  
 293 **Figure 5:** An example of regional land-use mapping. The upper panels show the distribution of known  
 294 archaeological sites superimposed on kernel density estimates of the extent of land-use based on the  
 295 density of observations, and the lower panels show these data superimposed on the LandCover6k

land-use classes for the Middle Neolithic (3600-3400 years BCE, 5600-5400 years BP, 5.6-5.4 ka BP) (left panels) and the Early Neolithic (3750-3600 years BCE, 5750-5600 years BP, 5.7-5.6 ka BP) (right panels) of Ireland. Data points derive from <sup>14</sup>C dated archaeological sites and distributions of settlements and monuments that have been assigned to each archaeological period following the dataset published in McLaughlin et al. (2016). The assigned land-use classes are inferred from archaeological material from one (or more) sites within the grid box. It should not be assumed that the whole gridcell was being used for agriculture during the Middle and Early Neolithic. Informed assessment suggests that agricultural land (crop growing and grazing combined) probably occupied between 10-15% of the total grid area in the low-level food production regions of the eastern and western coastal areas, whilst agricultural land likely represents 5% or less of the total grid cell area in inland areas.

#### 4. Incorporation of archaeological data in LULC scenarios

The existing LULC scenarios are substantially dependent on historical regional population estimates at key times, which are then linearly interpolated to provide a year-by-year estimate of population. Estimates of regional population growth based on suitably screened <sup>14</sup>C data can be used to modify existing population growth curves (Figure 6), both in terms of establishing the initial date of human presence and by modifying a linear growth curve to allow for intervals of population growth and decline.

Information on the timing of the first appearance of agriculture at specific locations can be used to constrain the temporal record of LULC changes in the scenarios. This information can also be used to allocate LULC changes geographically across regions (Figure 6). Global land-use maps can be used to identify areas where there was no permanent agricultural activity at a given time (e.g. either unsettled areas or areas occupied by hunter-gatherer communities) and provide a further constraint on the geographic extent of the LULC changes given by the scenarios (Figure 6). The type of agriculture, including whether the region was predominantly used for tree or annual crops or for pasture, modifies the area of open land specified in the LULC scenarios. Information on the extent of rain-fed versus irrigated agriculture, as indicated by the presence of irrigation structures associated with archaeological sites, can also be used to refine the distribution of these classes in the LULC scenarios. Per-capita land-use estimates and their changes through time (see e.g. Hughes et al., 2018; Weiberg et al., 2019) provide a further refinement of the LULC scenarios, allowing a better characterization of the distinction between e.g. areas given over to extensive versus intensive animal production (rangeland versus pasture in the HYDE 3.2 terminology). There will remain areas of the world for which this kind of fine-grained information is not available. Nevertheless, by incorporating information where this exists, the LandCover6k products will contribute to a systematic refinement of existing LULC scenarios. Iterative testing of the revised scenarios will ensure that they are robust.



**Figure 6:** Schematic illustration of the proposed implementation of  $^{14}\text{C}$ -based population estimates, date of first agriculture, land-use maps, and land-use per capita information in the HYDE model (here indicated as HYDE3.x). The archaeological data are represented as values for a grid cell in geographic space at a given time for date of first agriculture and land use, but as a time series for a specific grid cell for population and land-use per capita. In the case of population estimates, date of first agriculture and land-use per capita data, we show the initial estimate and the revised estimate after taking the archaeological information into account in the HYDE3.x plot. It should be assumed in the case of the land-use mapping that the original estimate was that there was no land use in this region.

## 5. Using pollen-based reconstructions of land cover changes to evaluate LULC scenarios

Pollen-based vegetation reconstructions can be used to corroborate archaeological information on the date of first agriculture from the appearance of cereals and agricultural weeds. These reconstructions can also be used to test the LULC reconstructions, either using relative changes in forest cover or reconstructions of the area occupied by different land cover types. LandCover6k uses the REVEALS pollen source-area model (Sugita, 2007) to estimate vegetation cover from fossil pollen assemblages. REVEALS predicts the relationship between pollen deposition in large lakes and the abundance of individual plant taxa in the surrounding vegetation at a large spatial scale (ca. 100 km x 100 km; Hellman et al., 2008a, b) using models of pollen dispersal and deposition. REVEALS can also be used with pollen records from multiple small lakes or peat bogs (Trondman et al., 2016) although this results in larger uncertainties in the estimated area occupied by individual taxa. The estimates obtained for individual taxa are summed to produce estimates of the area occupied by either plant functional (e.g. summer-green trees, evergreen trees) or land cover (e.g. open land, grazing land, cropland) types.


The geographic distribution of pollen records is uneven. There are also many areas of the world where environments that preserve pollen (i.e. lakes, bogs, forest hollows) are sparse. Site-based reconstructions of land cover are therefore interpolated statistically to produce spatially continuous reconstructions (Nielsen et al., 2012; Pirzamanbein et al., 2014; Pirzamanbein et al., 2018). LandCover6k uses a 1° resolution grid and all available pollen records in each grid cell to produce an estimate of land cover per grid cell through time. The more pollen records per grid cell and pollen counts per time window, the smaller the estimated error on the land-cover reconstruction. The uncertainties on the pollen-based REVEALS reconstructions are partly expressed by their standard errors (SEs). These SEs take into account the SE on the relative pollen productivity (RPP) of each plant taxon included in the REVEALS reconstructions and the variability between the site-specific REVEALS reconstructions (e.g. Trondman et al., 2015). The uncertainties on the pollen-based land cover reconstructions are taken into account when these reconstructions are compared with LULC scenarios (Kaplan et al., 2017).

The REVEALS approach has already been used to produce gridded reconstructions of changes in the amount of open land through time across the northern extratropics (Figure 7; Dawson et al., 2018). These reconstructions provide mean plant cover for time slices of 500 years through the Holocene until 0.7ka BP, and three historical time windows (modern–0.1ka BP, 0.1–0.35ka BP, and 0.35–0.7ka BP). The more pollen samples per time interval and pollen records per grid cell, the more years within the 500 yrs time slice will be represented in the reconstruction. This implies that the number of years represented in a time-slice reconstruction varies in space and time.

A major limitation in applying REVEALS globally is the requirement for information about the relative pollen productivity (RPP) of individual pollen taxa, which is currently largely lacking for the tropics. However, LandCover6k has been collecting RPPs for China, South-East India, Cameroon, Brazil and Argentina and pollen-based land-cover reconstructions will be available for sufficient parts of the tropics to allow testing of the LULC scenarios. Another limitation of the REVEALS reconstructions is that RPP estimates are available for cultivated cereals but not for other cultivars or cropland weeds, so the LandCover6k pollen-based reconstructions will generally underestimate cropland cover (Trondman et al., 2015). It may also be possible to use alternative pollen-based reconstructions of land cover changes, such as the Modern Analogue Approach (MAT: e.g. Tarasov et al., 2007; Zanon et al. 2018); pseudo-biomization (e.g. Fyfe et al., 2014) or STEPPS (Dawson et

al., 2016). While none of these methods require RPPs, MAT and STEPSS can only be applied in regions where the pollen datasets have dense coverage (such as Europe and North America) and pseudo-biomization is affected by the non-linearity of the pollen-vegetation relationship that the REVEALS approach is designed to remove.

Comparison of the reconstructions of the extent of open land with the LULC deforestation scenarios will provide a first evaluation of the realism of the revised LULC scenarios (e.g. Kaplan et al., 2017). Underestimation or overestimation of open land in the LULC scenarios is not necessarily an indication that these scenarios are inaccurate because (a) pollen-based reconstructions cannot distinguish between anthropogenic and climatically determined natural open land (e.g. natural grasslands, steppes, wetlands) and (b) REVEALS underestimates cropland cover because there are no RPP estimates for cultivars other than cereals. However, overestimation of the area of open land in the LULC scenarios might suggest problems either in the archaeological inputs or their implementation, especially for times or regions when other evidence indicates cereals were the major crop. In this sense, despite potential problems, the LandCover6k pollen-based reconstructions of land cover will provide an important independent test of the revised LULC scenarios.




**Figure 7:** Northern extratropical ( $>40^{\circ}\text{N}$ ) mean fractional cover of open land at 6000 years ago (6ka BP: top right panel) and 200 years ago (0.2ka BP: top left panel) estimated using REVEALS, and the



difference in fractional cover between the two periods (lower panel), where red indicates an increase in open land and blue a decrease (after Dawson et al., 2018).

## 6. Testing the reliability of improved scenarios using climate-model simulations

A second test of the realism of the improved LULC scenarios is to examine whether incorporating LULC changes improves the realism of the simulated climate when compared to palaeoclimate reconstructions (Figure 8). The mid-Holocene (6000 years ago, 6 ka BP) is an ideal candidate for such a test because benchmark data sets of quantitative climate reconstructions are available (e.g. Bartlein et al., 2011), the interval has been a focus through multiple phases of PMIP, control simulations with no LULC have already been run, and evaluation of these simulations has identified regions where there are major discrepancies between simulated and reconstructed climates e.g. the observed expansion of northern hemisphere monsoons, climate changes over Europe, the magnitude of high-latitude warming, and wetter conditions in central Eurasia (Mauri et al., 2014; Harrison et al., 2015; Bartlein et al., 2017). There are discernible anthropogenic impacts on the landscape in many of these regions by 6 ka, although they are not as strong as during the later Holocene and they are not present everywhere. Nevertheless, the 6ka BP interval provides a good focus for testing whether improvements to the LULC scenarios produces more realistic simulations of climate. Such an evaluation would need to go beyond the global comparison made here (Figure 8) to regional comparisons to identify whether improvements in simulated climate in regions where there is a large anthropogenic impact on land cover do not result in a degradation in the simulated climate elsewhere.




**Figure 8:** Quantitative comparison of the change in climate between the mid-Holocene (6ka) and the pre-industrial period as shown by pollen-based reconstructions gridded to  $2 \times 2^\circ$  resolution to be compatible with the model resolution (from Bartlein et al., 2011) and in simulations with and without the incorporation of land-use change (from Smith et al., 2016). This figure illustrates the approach that will be taken to evaluate the impact of new LULC scenarios on climate. The imposed land-use changes at 6000 years ago (6ka BP) were derived from the KK10 scenario (Kaplan et al., 2011). The plots show comparisons of mean annual temperature (MAT), mean temperature of the coldest month (MTCO) and mean annual precipitation (MAP) for the northern extratropics (north of  $30^\circ$  N), where each dot represents a model grid cell where comparisons with the pollen-based reconstructions is possible. Although the incorporation of land use produces somewhat warmer and wetter climates in these simulations, overall the incorporation of land-use produces no improvement of the simulated climates at sites with pollen-based reconstructions.

## 7. Testing the reliability of improved scenarios using carbon-cycle models

Carbon-cycle modelling will be used as a further test of the realism of the improved LULC scenarios. Two constraints are available for testing the realism of past LULC scenarios. First, reconstructions of LULC history must converge on the present-day state, which is relatively well constrained by satellite land-cover observations and national statistics on the amount of land under use. Reconstructing the extent of past LULC change thus reduces to allocating a fixed total amount of land conversion from natural to agricultural use over time. More conversion in earlier periods implies less conversion in later periods. At the continental to global scale, cumulative LULC emissions scale linearly with the agricultural area. LULC scenarios that converge to the present-day state also converge to within a small range of cumulative historical emissions (Stocker et al., 2011; Stocker et al., 2017). Deviations from a linear relationship between extent and emissions are due to differences in biomass density in potential natural and agricultural vegetation of different regions affected by anthropogenic LULC. Differences in cumulative emissions for alternative LULC reconstructions with an identical present-day state are due to the long response time of soil carbon content following a change in carbon inputs and soil cultivation. Conserving the total extent of LULC (and allocating a fixed total expansion over time) is thus approximately equivalent to conserving cumulative historical LULC emissions. Thus, more LULC CO<sub>2</sub> emissions in earlier periods imply less CO<sub>2</sub> emissions in more recent periods.

The total C budget of the terrestrial biosphere provides a second constraint on LULC emissions through time. The net C balance of the land biosphere, which reflects the sum of all natural and anthropogenic effects on terrestrial C storage, can be reconstructed from ice-core data of past CO<sub>2</sub> concentrations and  $\delta^{13}\text{C}$  composition (Elsig et al. 2009). Providing that all of the natural contributions to the land C inventory (e.g. the build-up of natural peatlands: Loisel et al., 2014) can be specified from independent evidence, the anthropogenic sources can be estimated as the difference between the total terrestrial C budget and natural contributions (Figure 9) at any specific time.



**Figure 9:** Illustration of the terrestrial C budget approach to evaluate LULC. The total terrestrial C balance (green circle 'total') is constrained by ice core records of CO<sub>2</sub> and its isotopic signature ( $\delta^{13}\text{C}$ ). Estimates for C balance changes of different natural land carbon cycle components (e.g., peatlands, permafrost, forest expansion/retreat, desert greening) can be estimated independently (blue slices 'Natural components') either from empirical upscaling of site-scale observations or from model-based analyses (BGC models forced with varying climate). The remainder (yellow slice 'remainder') is then calculated as the total terrestrial C balance (green circle 'total') minus the sum of the separate estimates of the natural components (blue slices 'Natural components'). The remainder



is effectively the emissions resulting from LULC changes, and can therefore be compared to LULC CO<sub>2</sub> emission estimates by carbon-cycle models.

Transient simulations with a model that simulates CO<sub>2</sub> emissions in response to anthropogenic LULC can be used to test the reliability of the LULC scenarios, by comparing results obtained with prescribed LULC changes through time against a baseline simulation without imposed LULC. This will necessitate making informed decisions about the fraction of land under cultivation that is abandoned or left fallow each year, and the maximum extent of land affected by such episodic cultivation. We envisage using several different offline carbon-cycle models for this purpose in order to take account of uncertainties associated with differences between the carbon-cycle models. The carbon-cycle simulations will be driven by climate outputs (temperature, precipitation and cloud cover) from an existing transient climate simulation made with the ECHAM model (Fischer and Junglaus, 2011) and CO<sub>2</sub> prescribed from ice-core records. The CO<sub>2</sub> emission estimates from these two simulations will then be evaluated using C budget constraints. This evaluation will allow us to pinpoint potential discrepancies between known terrestrial C balance changes and estimated LULC CO<sub>2</sub> emission in given periods over the Holocene.

## 8. Implementation of LULC in Earth System Model simulations

We propose a series of simulations to examine the impact of LULC, using the revised LULC scenarios from LandCover6k and building on climate-model experiments that are currently being run either in CMIP6-PMIP4 (*midHolocene*, *past1000*) or within PMIP although not formally included as CMIP6-PMIP4 experiments.

The *mid-Holocene* (and its corresponding *piControl*) simulation is one of the PMIP entry cards in the CMIP6-PMIP4 experiments (Kageyama et al., 2018; Otto-Bliesner et al., 2017) and it is therefore logical to propose this period for LULC simulations. The LULC sensitivity experiment (*midHoloceneLULC*) should therefore follow the CMIP6-PMIP4 protocol, that is it should be run with the same climate-model components and following the same protocols for implementing external forcings as used in the two CMIP6-PMIP4 experiments (Table 1). Thus, if the *piControl* and *midHolocene* simulations are run with interactive (dynamic) vegetation, then the *midHoloceneLULC* experiment should also be run with dynamic vegetation in regions where there is no LULC change. For most models, this means that the LULC forcing is imposed as a fraction of the grid cell and the remaining fraction of the grid cell has simulated natural vegetation. These new mid-Holocene simulations would allow for a better understanding of the relationship between climate changes and land-surface feedbacks (including snow albedo feedbacks), and the role of water recycling at a regional scale. Thus, modelling groups who are running the *midHolocene* experiment with a fully interactive carbon cycle could also run the LULC experiment allowing atmospheric CO<sub>2</sub> to evolve interactively, subject to the simulated ocean and land C balance.

**Table 1:** Boundary conditions for CMIP6-PMIP4 and the mid-Holocene LULC experiments. The boundary conditions for the CMIP6-PMIP4 *piControl* and *midHolocene* are described in Otto-Bliesner et al. (2017) and are given here for completeness.

Boundary conditions		1850CE (DECK <i>piControl</i> )	6ka ( <i>midHolocene</i> )	6ka LULC ( <i>midHoloceneLULC</i> )
Orbital parameters	Eccentricity	0.016764	0.018682	0.018682
	Obliquity	23.459	24.105	24.105

	Perihelion – 180	100.33	0.87	0.87
	Vernal equinox	Noon, 21 March	Noon, 21 March	Noon, 21 March
Greenhouse gases	Carbon dioxide (ppm)	284.3	264.4	264.4
	Methane (ppb)	808.2	597.0	597.0
	Nitrous oxide (ppb)	273.0	262.0	262.0
	Other GHG	DECK <i>piControl</i>	0	0
Other boundary conditions	Solar constant	TSI: 1360.747	As <i>piControl</i>	As <i>piControl</i>
	Palaeogeography	Modern	As <i>piControl</i>	As <i>piControl</i>
	Ice sheets	Modern	As <i>piControl</i>	As <i>piControl</i>
	Vegetation	Interactive	Interactive	pasture and crop distribution prescribed from a revised scenario
		DECK <i>piControl</i>	As <i>piControl</i>	pasture and crop distribution prescribed from a revised scenario
	Aerosols	interactive	Interactive	Interactive
		DECK <i>piControl</i>	As <i>piControl</i>	As <i>piControl</i>

The real strength of the revised LULC scenarios is to provide boundary conditions for transient climate-model simulations. The CMIP6-PMIP4 simulation of 850-1850 CE (*past1000*) already incorporates LULC changes as a forcing (Jungclaus et al. 2017), based on a harmonized data set that provides LULC changes from 850 through to 2015 CE (Hurtt et al., 2017), which in turn draws on output from the HYDE3.2 scenario (Klein Goldewijk et al., 2017a). The *past1000* protocol (Jungclaus et al., 2017) acknowledges that this default land-use data set is at the lower end of the spread in estimates of early agricultural area indicated by other LULC scenarios and recommends that modelling groups run additional sensitivity experiments using alternative maximum and minimum scenarios. The revised LULC scenarios created by LandCover6k could be used as an alternative to these maximum and minimum scenarios. Other than the substitution of the LandCover6k scenario, the specifications of other forcings would then follow the recommendations for the CMIP6-PMIP4 *past1000* simulation.

A transient climate simulation for a longer period of the Holocene would provide a more stringent test of the impact of LULC on the coupled earth system. We suggest that this transient simulation (*holotrans*) should start from the pre-existing *midHolocene* simulation to capitalise on the fact that the *midHolocene* simulation has been spun up for sufficiently long (Otto-Bleisner et al., 2017) to ensure that the ocean and land carbon cycle is in equilibrium at the start of the transient experiment (Table 2). In order to be consistent with the CMIP6-PMIP4 *midHolocene* protocol (Otto-Bleisner et al., 2017), changes in orbital forcing should be specified from Berger and Loutre (1991) and year-by-year changes in CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O should be specified following Joos and Spahni (2008). LULC changes should be implemented by imposing crop and pasture area through time as specified in the revised LULC scenarios; elsewhere, the simulated vegetation should be active. It will be necessary

to run the Holocene transient climate simulation in two steps. A first simulation (*holotrans\_LULC*) should be run using prescribed atmospheric CO<sub>2</sub> concentration even though the carbon cycle is fully interactive, because this will establish the consistency of the carbon cycle in the land surface model. However, once this is done it will be possible to re-run the simulations with interactive CO<sub>2</sub> emissions. Table 3 provides a summary of the proposed ESM simulations.

**Table 2:** Boundary conditions for baseline PMIP Holocene transient (6 ka BP to 1850 CE) and LULC transient simulations

		Mode	Source/Value	LULC experiment
Orbital parameters		transient		As baseline simulation
Greenhouse gases	CO <sub>2</sub>	transient	Dome C	As baseline simulation
	CH <sub>4</sub>		Combined EPICA & GISP record	As baseline simulation
	N <sub>2</sub> O		Combined EPICA NGRIP, & TALDICE record	As baseline simulation
Solar forcing		transient	Steinhilber et al. (2012)	As baseline simulation
Volcanic forcing		transient	To be determined	As baseline simulation
Palaeogeography		Constant at PI values	Modern	As baseline simulation
Ice sheets		Constant at PI values	Modern	As baseline simulation
Vegetation		interactive		LC6k transient pasture and crop distribution imposed
Aerosols		Constant at PI values		As baseline simulation

Unlike the situation for the mid-Holocene, where there is a global climate benchmark data set (Bartlein et al., 2011) which provides reconstructions of multiple bioclimatic variables of seasonal temperature and moisture, the opportunities for quantitative evaluation of the *holotrans* simulated climate are more limited. Seasonal temperature reconstructions are available for Europe (Davis et al., 2003) and North America (Viau et al, 2006; Viau and Gajewski, 2009). Although there is a new global data set that provides global temperature reconstructions for the Holocene (Kaufman et al., in press), it is based on only 472 terrestrial records worldwide and the results for zonally averaged temperature changes are therefore likely to be more robust than the regional details. There are also time series reconstructions for individual sites outside these two regions (e.g. Nakagawa et al., 2002; Wilmshurst et al., 2007; Ortega-Rosas et al., 2008). Furthermore, the simulated time-course of CO<sub>2</sub> emissions can be compared to the ice core records.

**Table 3:** Summary of proposed simulations.

Name	Mode	Purpose
<i>piControl</i>	equilibrium	Standard CMIP6-PMIP4 simulation

<i>midHolocene</i>	equilibrium	Standard CMIP6-PMIP4 simulation
<i>midHoloceneLULC</i>	equilibrium	Sensitivity to LULC changes
<i>holotrans</i>	transient	Baseline fully transient simulation from 6ka onwards, with no LULC
<i>holotrans_LULC</i>	transient	Fully transient simulation from 6ka onwards, with LULC imposed

The CMIP6-PMIP4 *mid-Holocene* simulations are stylized experiments, lacking several potential forcings (in addition to LULC), including changes in atmospheric dust loading, in solar irradiance, and volcanic forcing. We suggest that additional sensitivity tests could be run to take these additional forcings into account. In the case of solar and volcanic forcing, this would also ensure that the transient *holotrans* simulations mesh seamlessly with the *past1000* simulation. Changes in solar variability during the Holocene should be specified from Steinhilber et al. (2012). There are records of volcanic forcing for the past 2000 years (Sigl et al., 2015; Toohey and Sigl, 2017), and these are used in the *past1000* simulation. Observationally constrained estimates of the volcanic stratospheric aerosol for Holocene are currently under development (M. Sigl, pers comm.) and could be implemented as an additional sensitivity experiment when available. Changes in atmospheric dust loading are not included in the *past1000* simulation but are important during the earlier part of the Holocene (Pausata et al., 2016; Tierney et al., 2017; Messori et al., 2019). Although continuous reconstructions of dust loading through the Holocene are not available, it would be possible to use estimates for particular time-slices (Egerer et al., 2018) to test the sensitivity to this forcing.

## Outcomes and Perspectives

LandCover6k has developed a scheme for using archaeological information to improve existing scenarios of LULC changes during the Holocene, specifically by using archaeological data to provide better estimates of regional population changes through time, better information on the date of initiation of agriculture in a region, more regionally specific information about the type of land use, and more nuanced information about land-use per capita than currently implemented in the LULC scenarios generated by HYDE and KK10. While the final global data set are still in production, fast-track priority products have been created and their impact on current scenarios is being tested.

Although the work of LandCover6k will provide more solid knowledge about anthropogenic modification of the landscape, some information will inevitably be missing and some key regions will be poorly covered. There will still be large uncertainties associated with revised LULC scenarios, even though these will be based on more solid evidence than the existing LULC scenarios. Documenting the uncertainties in the archaeological inputs and their impacts on the revised scenarios is an important goal of the LandCover6k project. We propose using the information about the uncertainties in the archaeological data sources to generate multiple LULC scenarios comparable to the "low-end", "high-end" scenarios used for e.g. in future projections. Furthermore, we have proposed a series of tests that will help to evaluate the realism of the final scenarios, based on independent evidence from pollen-based reconstructions of land cover, reconstructions of climate, and carbon-cycle constraints. These tests should help in identifying which of the potential LULC reconstructions are most realistic and in constraining the sources of uncertainty.

We have proposed the use of offline carbon-cycle simulations solely as a test of the realism of the revised LULC scenarios. Quantifying the LULC contribution to CO<sub>2</sub> emissions during the Holocene would require additional simulations in which other forcings (climate, atmospheric CO<sub>2</sub>, insolation) are kept constant. The difference in simulated total terrestrial C storage between these simulations and LULC simulations provides an estimate of *primary emissions* (Pongratz et al., 2014) and avoids additional model uncertainty regarding the sensitivity of land C storage to atmospheric CO<sub>2</sub> or climate being included in emission estimates. There are other sensitivity tests that would be useful. For example, vegetation-carbon-cycle models differ in their ability to account for gross land use transitions within grid cells (Arneeth et al., 2017). This is critical for simulating effects of non-permanent agriculture where land is simultaneously abandoned and re-claimed within the extent of a model grid cell. Such shifting cultivation-type agriculture implies forest degradation in areas recovering from previous land use and leads to substantially higher LULC emissions compared to model estimates where only net land-use changes are accounted for (Shevliakova et al., 2009). It would therefore be interesting to run additional simulations accounting for net land use change, and indeed separating out the effects of wood harvesting and shifting cultivation.

We anticipate that it will be possible to incorporate realistic LULC scenarios for the mid-Holocene as part of the climate-model sensitivity experiments planned during PMIP4. Such experiments will complement the CMIP6-PMIP4 baseline model experiments, by providing insights into whether discrepancies between simulated and observed 6 ka climate could be the result of incorrect specification of the land-surface boundary conditions. However, the incorporation of archaeological information into LULC scenarios clearly makes it possible to target other interesting periods for such experiments, for example to explore if land-use changes played a role in abrupt events such as the 4.2 ka event, or to examine the impact of population declines in the Americas as a consequence of European colonisation (1500-1750 CE) or the changes in land use globally during the Industrial era (post 1850 CE).

In addition to providing a protocol for the PMIP 6ka sensitivity experiments, we have devised a protocol for implementing the optimal LULC reconstructions for the Holocene in transient climate-model or ESM experiments. The goal here is to provide one of the necessary forcings that could be used for transient simulations in future phases of PMIP. This will allow an assessment of LULC in these simulations, and therefore help address issues that are a focus for other MIPs e.g. LUMIP or LS3MIP. When these new forcings are created, they will be made available through the PMIP4 website ([https://pmip4.lscce.ipsl.fr/doku.php/exp\\_design:lgm](https://pmip4.lscce.ipsl.fr/doku.php/exp_design:lgm), PMIP4 repository, 2017) and the ESGF Input4MIPS repository (<https://esgf-node.llnl.gov/projects/input4mips/>, with details provided in the “input4MIPs summary” link). Modelling groups who run either equilibrium or transient climate-model experiments following this protocol are encouraged to follow the standard CMIP protocol of archiving their simulations through the ESFG.

### Code and Data Availability

The data used for Figure 1 are publicly available. The HYDE3.2 data can be downloaded <https://doi.org/10.17026/dans-25g-gez3>. The KK10 data can be downloaded from <https://doi.org/10.1594/PANGAEA.871369>. The code and data used to generate Figure 1 are available from [https://github.com/jedokaplan/ALCC\\_comparison\\_figure](https://github.com/jedokaplan/ALCC_comparison_figure). The data and code used to generate Figure 3 are available from <https://github.com/mavdlind/GMD>. The data and code used to generate Figure 5 are available from <https://doi.org/10.1594/PANGAEA.897303>. The European pollen-based reconstructions used in Figure 7 are available <https://doi.org/10.1594/PANGAEA.898616>. The pollen data used to generate the Siberian reconstructions is available from <https://doi.org/10.1594/PANGAEA.898616>. An earlier version of this Figure was published in Dawson et al., 2018. The code used to generate 7 Figure is

678 available from <https://doi.org/10.5281/zenodo.3604328>. The pollen-based reconstructions used in the  
679 generation of Figure 8 are available from 10.5281/zenodo.3601028. The climate model outputs used  
680 to generate Figure 8 are available from 10.5281/zenodo.3601040. The code used to generate Figure  
681 8 is available from 10.5281/zenodo.3601011.

682

683

684 **Acknowledgements.** LandCover 6k is a working group of the Past Global Changes (PAGES)  
685 programme, which in turn received support from the Swiss Academy of Sciences. We thank PAGES  
686 for their support for this activity. The land use group also received funding under the Holocene Global  
687 Landuse International Focus Group of INQUA. SPH acknowledges funding from the European  
688 Research Council for “GC2.0: Unlocking the past for a clearer future”. MJG thanks the Swedish  
689 Strategic Research Area MERGE (Modelling the Regional and Global Earth System Model) and  
690 Linnaeus University’s faculty of Health and Life Sciences (Kalmar, Sweden) for financial support.  
691 We thank Joy Singarayer for providing the climate model outputs that were used to generate Figure  
692 8 and Guangqi Li for assistance in producing this figure. BDS was funded by ERC H2020-MSCA-  
693 IF-2015, grant number 701329. The dataset for Figure 5 was generated from the ‘Cultivating  
694 Societies: Assessing the Evidence for Agriculture in Neolithic Ireland’ project, supported by the  
695 Heritage Council, Ireland under the INSTAR programme 2008–2010 (Reference 16682 to  
696 Whitehouse, Schulting, Bogaard and McClatchie).

697

698 **Author Contributions.** SPH, MJG, BDS, MVL, KKG wrote the first draft. SPH, PB, FSRP  
699 contributed to the design of the climate model experiments, BS and TK to the design of the carbon-  
700 cycle simulations. The figures were contributed by JK (Fig. 1), BS (Fig. 2, Fig 9.), MVL (Fig. 3), OB  
701 (Fig. 4), NJW (Fig. 5), KKG (Fig. 6), AD (Fig. 7), SPH (Fig. 8). All authors contributed to the final  
702 version of the paper.

703

704

## 705 References

- 706 Arneth, A., Denton, F., Agus, F., Elbehri, A., Erb, K., Elasha, B.O., Rahimi, M., Rounsevell, M.,  
707 Spence, A., and Valentini, R.: IPCC Special Report on Climate Change, Desertification, Land  
708 Degradation, Sustainable Land Management, Food Security, and Greenhouse gas fluxes in  
709 Terrestrial Ecosystems, 2019.
- 710 Arneth, A., Sitch, S., Pongratz, J., Stocker, B.D., Ciais, P., Poulter, B., Bayer, A.D., Bondeau, A.,  
711 Calle, L., Chini, L.P., Gasser, T., Fader, M., Friedlingstein, P., Kato, E., Li, W., Lindeskog,  
712 M., Nabel, J.E.M.S., Pugh, T.A.M., Robertson, E., Viovy, N., Yue, C., and Zaehle, S.:  
713 Historical carbon dioxide emissions caused by land-use changes are possibly larger than  
714 assumed, *Nature Geosci.*, 10, 79-84, doi: 10.1038/ngeo2882, 2017.
- 715 Balsera, V., Díaz-del-Río, P., Gilman, A., Uriarte, A., and Vicent, J.M.: Approaching the demography  
716 of late prehistoric Iberia through summed calibrated probability distributions (7000-2000 cal  
717 BC), *Quat. Int.*, 208-211, doi:10.106/j.quaint.2015.06.022, 2015.
- 718 Bartlein, P. J., Harrison, S. P., Brewer, S., Connor, S., Davis B. A. S., Gajewski, K., Guiot, J.,  
719 Harrison-Prentice, T. I., Henderson, A., Peyron, O., Prentice, I. C., Scholze, M., Seppä, H.,  
720 Shuman, B., Sugita, S., Thompson, R. S., Vial, A., Williams, J., and Wu, H.: Pollen-based  
721 continental climate reconstructions at 6 and 21 ka: a global synthesis, *Clim. Dyn.*, 37, 775-  
722 802, doi: 10.1007/s00382-010-0904-1, 2011.
- 723 Bartlein, P.J., Harrison, S.P., and Izumi, K.: Underlying causes of Eurasian mid-continental aridity  
724 in simulations of mid-Holocene climate, *Geophys. Res. Lett.*, 44, doi:  
725 10.1002/2017GL074476, 2017.

- Barton, C.M., Ullah, I.I., and Bergin, S.: Land use, water and Mediterranean landscapes: modelling long-term dynamics of complex socio-ecological systems, *Phil. Trans. R. Soc.*, A368, 5275-5297, doi: 10.1098/rsta.2010.0193, 2010.
- Berger, A., and Loutre, M-F.: Insolation values for the climate of the last 10 million of years, *Quat. Sci. Rev.*, 10, 297-317, [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.
- Bishop, R.R., Church, M.J., and Rowley-Conwy, P.A.: Firewood, food and human niche construction: the potential role of Mesolithic hunter-gatherers in actively structuring Scotland's woodlands, *Quat. Sci. Rev.*, 108, 51-75, 2015.
- Bocquet-Appel, J.-P., Naji, S., Vander Linden, M., and Kozłowski, J.K.: Detection of diffusion and contact zones of early farming in Europe from the space-time distribution of  $^{14}\text{C}$  dates, *J. Arch. Sci.*, 36, 807-820, doi: 10.1016/j.jas.2008.11.004, 2009.
- Crema, E.R., Habu, J., Kobayashi, K., and Madella, M.: Summed probability distribution of  $^{14}\text{C}$  dates suggests regional divergences in the population dynamics of the Jomon period in eastern Japan, *PlosOne*, 11, e0154809, doi: 10.1371/journal.pone.0154809, 2016.
- Davis, B.A.S., Brewer, S., Stevenson, A.C., Guiot, J., and Juggins, S.: The temperature of Europe during the Holocene reconstructed from pollen data, *Quat. Sci. Rev.*, 22, 1701-1716, 2003.
- Davison, K., Dolukhanov, P., Sarson, G. R., and Shukurov, P.: The role of waterways in the spread of the Neolithic, *J. Arch. Sci.*, 33, 641-652, doi: 10.106/j.jas.2005.09.017, 2006.
- Dawson, A., Paciorek, C.J., McLachlan, J.S., Goring, S., Williams, J.W., and Jackson, S.T.: Quantifying pollen-vegetation relationships to reconstruct ancient forests using 19th-century forest composition and pollen data, *Quat. Sci. Rev.*, 137, 156-175, doi: 10.1016/j.quascirev.2016.01.012, 2016.
- Dawson, A., Cao, X., Chaput, M., Hopla, E., Li, F., Edwards, M., Fyfe, R., Gajewski, K., Goring, S.J., Herzschuh, U., Mazier, F., Sugita, S., Williams, J.W., Xu, Q., and Gaillard, M-J.: Finding the magnitude of human induced Northern Hemisphere land-cover transformation between 6 and 0.2 ka BP, *PAGES Mag.*, 26, 34-35, <https://doi.org/10.22498/pages.26.1.34>, 2018.
- Egerer, S., Claussen, M., and Reick, C.: Rapid increase in simulated North Atlantic dust deposition due to fast change of northwest African landscape during the Holocene, *Clim. Past*, 14, 1051-1066, <https://doi.org/10.5194/cp-14-1051-2018>, 2018.
- Ellis, E.C., Kaplan, J.O., Fuller, D.Q., Vavrus, S., Klein Goldewijk, K., and Verburg, P. H.: Used planet: A global history, *Proc. Nat. Acad. Sci.*, 110, 7978-7985, 2013.
- Elsig, J., Schmitt, J., Leuenberger, D., Schneider, R., Eyer, M., Leuenberger, M., Joos, F., Fischer, H., and Stocker, T. F.: Stable isotope constraints on Holocene carbon cycle changes from an Antarctic ice core, *Nature*, 461, 507-510, 2009.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937-1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Fischer, N., and Jungclauss, J.H.: Evolution of the seasonal temperature cycle in a transient Holocene simulation: orbital forcing and sea-ice, *Clim. Past*, 7, 1139-1148, <https://doi.org/10.5194/cp-7-1139-2011>, 2011.
- Ford, A., and Clarke, K.C.: Linking the past and present of the ancient Maya: lowland land use, population distribution, and density in the Late Classic Period, in: *The Oxford Handbook of Historical Ecology and Applied Archaeology*, Isendahl, C. and Stump, D. (eds.), doi: 10.1093/oxfordhb/9780199672691.013.33, 2015.
- Freeman, J., Baggio, J.A., Robinson, E., Byers, D.A., Gayo, E., Finley, J.B., Meyer, J.A., Kelly, R.L., and Anderies, J.M.: Synchronisation of energy consumption by human societies throughout the Holocene, *Proc. Nat. Acad. Sci.*, 115, 9962-9967, doi: 10.1073/pnas.1802859115, 2018.
- Fyfe R. M., Woodbridge, J. E., and Roberts, N.: From forest to farmland: pollen-inferred land cover change across Europe using the pseudobiomization approach, *Glob. Change Biol.*, 21: 1197-



1212, doi:10.1111/gcb.12776, 2014.

Gaillard, M.-J., Sugita, S., Mazier, F., Kaplan, J.O., Trondman, A.-K., Brostroem, A., Hickler, T., Kjellstroem, E., Kunes, P., Lemmen, C., Olofsson, J., Smith, B., and Strandberg, G.: Holocene land-cover reconstructions for studies on land-cover feedbacks, *Clim. Past*, 6, 483-499, <https://doi.org/10.5194/cp-6-483-2010>, 2010.

Gaillard, M.-J., Whitehouse, N., Madella, M., Morrison, K., and von Gunten, L.: Past land use and land cover, *PAGES Mag.*, 26, 1-44, doi:10.22498/pages.26.1, 2018.

Harrison, S.P., Bartlein, P.J., Izumi, K., Li, G., Annan, J., Hargreaves, J., Braconnot, P.B., and Kageyama, M.: Evaluation of CMIP5 palaeo-simulations to improve climate projections, *Nature Clim. Change*, 5, 735-743, 2015.

He, F., Vavrus, S.J., Kutzbach, J.E., Ruddiman, W.F., Kaplan, J.O., and Krumhardt, K.M.: Simulating global and local surface temperature changes due to Holocene anthropogenic land cover change, *Geophys. Res. Lett.*, 41, 623–631, 2014.

Hellman, S., Gaillard, M.-J., Broström, A., and Sugita, S.: The REVEALS model, new tool to estimate past regional plant abundance from pollen data in large lakes: validation in southern Sweden, *J. Quat. Sci.*, 22, 1-22, 2008a.

Hellman, S., Gaillard M.-J., Broström, A., and Sugita, S.: Effects of the sampling design and selection of parameter values on pollen-based quantitative reconstructions of regional vegetation: a case study in southern Sweden using the REVEALS model, *Veg. History Archaeobot.*, 17, 445-460, 2008b.

Herrero, M., Havlik, P., Valin, H., Notenbaert, A., Rufino, M.C., Thornton, P.K., Blümmel, M., Weiss, F., Grace, D., and Obersteiner, M.: Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems, *Proc. Nat. Acad. Sci.*, 110, 20888-20893, 2013.

Hughes, R.E., Weiberg, E., Bonnier, A., Finne, M., and Kaplan, J.O.: Quantifying land use in past societies from cultural practice and archaeological data, *Land*, 7, 9, doi.org/10.3390/land/7010009, 2018.

Hurt, G., Chini, L., Sahajpal, R., Frolking, S., Calvin, K., Fujimori, S., Klein Goldewijk, K., Hasegawa, T., Havlik, P., Lawrence, D., Lawrence, P., Popp, A., Stehfest, E., van Vuuren, D., and Zhang, X.: Harmonization of global land-use change and management for the period 850–2100, *Geosci. Model Dev. Discuss.*, 2017.

Janssen, E., Poblome, J., Claeys, J., Kint, V., Degryse, P., Marinova, E., and Muys, B.: Fuel for debating ancient economies. Calculating wood consumption at urban scale in Roman Imperial times, *J. Arch. Sci.: Reports*, 11, 592-599, 2017.

Joos, F., and Spahni, R.: Rates of change in natural and anthropogenic radiative forcing over the past 20,000 years, *Proc. Nat. Acad. Sci.*, 105, 1425–1430, 2008.

Joos, F., Gerber, S., Prentice, I.C., Otto-Bliesner, B.L., and Valdes, P.J.: Transient simulations of Holocene atmospheric carbon dioxide and terrestrial carbon since the last glacial maximum, *Global Biogeochem. Cy.*, 18, GB2002, doi:10.1029/2003GB002156, 2004.

Jungclauss, J.H., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L. P., Egorova, T., Evans, M., González-Rouco, J.F., Goosse, H., Hurt, G.C., Joos, F., Kaplan, J. O., Khodri, M., Klein Goldewijk, K., Krivova, N., LeGrande, A.N., Lorenz, S. J., Luterbacher, J., Man, W., Maycock, A.C., Meinshausen, M., Moberg, A., Muscheler, R., Nehrbass-Ahles, C., Otto-Bliesner, B.I., Phipps, S.J., Pongratz, J., Rozanov, E., Schmidt, G.A., Schmidt, H., Schmutz, W., Schurer, A., Shapiro, A.I., Sigl, M., Smerdon, J.E., Solanki, S.K., Timmreck, C., Toohey, M., Usoskin, I.G., Wagner, S., Wu, C.-J., Yeo, K.L., Zanchettin, D., Zhang, Q., and Zorita, E.: The PMIP4 contribution to CMIP6 – Part 3: The last millennium, scientific objective, and experimental design for the PMIP4 past1000 simulations, *Geosci. Model Dev.*, 10, 4005–4033, <https://doi.org/10.5194/gmd-10-4005-2017>, 2017.

Kageyama, M., Braconnot, P., Harrison, S.P., Haywood, A., Jungclauss, J., Otto-Bliesner, B.,



Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft, P., Ivanovic, R., Lambert, F., Lunt, D., Mahowald, N., Peltier, W.R., Phipps, S., Roche, D., Schmidt, G., Tarasov, L., Valdes, P., Zhang, Q., and Zhou, T.: The PMIP4 contribution to CMIP6 – Part 1: Overview and overarching analysis plan. *Geosci. Model Dev.*, 11: 1033-1057. <https://doi.org/10.5194/gmd-11-1033-2018>, 2018.

Kaplan, J.O., Krumhardt, K.M., Ellis, E.C., Ruddiman, W.F., Lemmen, C., and Klein Goldewijk, K.: Holocene carbon emissions as a result of anthropogenic land cover change, *Holocene*, 21, 775-791, 2011.

Kaplan, J.O., Krumhardt, K.M., Gaillard, M.-J., Sugita, S., Trondman, A.-K., Fyfe, R., Marquer, L., Mazier, F., and Nielsen, A.B.: Constraining the deforestation history of Europe: Evaluation of historical land use scenarios with pollen-based land cover reconstructions, *Land*, 6, 9, doi:10.3390/land6040091, 2017.

Kaufman, D., McKay, N., Routson, C., Erb, M., Davis, B., Heiri, O., Jaccard, S., Tierney, J., Dätwyler, C., et al.: A global database of Holocene paleo-temperature records, *Scientific Data*, in press.

Kay, A. U., Fuller, D. Q., Neumann, K., Eichhorn, B., Höhn, A., Morin-Rivat, J., Champion, L., Linseele, V., Huysecom, E., Ozainne, S., Lespez, L., Biagetti, S., Madella, M., Salzmann, U., and Kaplan, J. O.: Diversification, intensification, and specialization: Changing land use in western Africa from 1800 BC to AD 1500, *J. World Prehistory*, 32, 179–228, doi:10.1007/s10963-019-09131-2. 2019, 2019.

Klein Goldewijk, K., Beusen, A., van Drecht, G., and de Vos, M.: The HYDE 3.1 spatially explicit database of human induced land use change over the past 12,000 years, *Glob. Ecol. Biogeog.*, 20, 73-86, 2011.

Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land-use estimates for the Holocene; HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927-953, <https://doi.org/10.5194/essd-9-927-2017>, 2017a.

Klein Goldewijk, K., Dekker, S.C., and van Zanden, J.L.: Per-capita estimations of long-term historical land use and the consequences for global change research, *J. Land Use Sci.*, 12, 313-337, <https://doi.org/10.1080/1747423X.2017.1354938>, 2017b.

Lawrence, D.M., Hurtt, G.C., Arneth, A., Brovkin, V., Calvin, K.V., Jones, A.D., Jones, C.D., Lawrence, P.J., de Noblet-Ducoudré, N., Pongratz, J., Seneviratne, S.I., and Shevliakova, E.: The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: rationale and experimental design, *Geosci. Model Dev.*, 9, 2973-2998, <https://doi.org/10.5194/gmd-9-2973-2016>, 2016.

Le Quéré, C., Andrew, R.M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P.A., Korsbakken, J.I., Peters, G.P., Canadell, J.G., Arneth, A., Arora, V.K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L.P., Ciais, P., Doney, S.C., Gkritzalis, T., Goll, D.S., Harris, I., Haverd, V., Hoffman, F.M., Hoppema, M., Houghton, R.A., Hurtt, G., Ilyina, T., Jain, A.K., Johannessen, T., Jones, C.D., Kato, E., Keeling, R.F., Goldewijk, K.K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozzi, D., Metzl, N., Munro, D.R., Nabel, J.E.M.S., Nakaoka, S.-I., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P.P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I.T., van der Werf, G.R., Viovy, N., Walker, A.P., Wiltshire, A.J., Wright, R., Zaehle, S., and Zheng, B.: Global Carbon Budget 2018, *Earth Syst. Sci. Data*, 10, 2141-2194, <https://doi.org/10.5194/essd-10-2141-2018>, 2018.

Loisel, J., Yu, Z., Beilman, D.W., Camill, P., Alm, J., Amesbury, M.J., Anderson, D., Andersson, S., Bochicchio, C., Barber, K., Belyea, L.R., Bunbury, J., Chambers, F.M., Charman, D.J., Vleeschouwer, F.D., Fiałkiewicz-Kozieł, B., Finkelstein, S.A., Gałka, M., Garneau, M.,

876 Hammarlund, D., Hinchcliffe, W., Holmquist, J., Hughes, P., Jones, M.C., Klein, E.S., Kokfelt,  
 877 U., Korhola, A., Kuhry, P., Lamarre, A., Lamentowicz, M., Large, D., Lavoie, M., MacDonald,  
 878 G., Magnan, G., Mäkilä, M., Mallon, G., Mathijssen, P., Mauquoy, D., McCarroll, J., Moore,  
 879 T.R., Nichols, J., O'Reilly, B., Oksanen, P., Packalen, M., Peteet, D., Richard, P.J., Robinson,  
 880 S., Ronkainen, T., Rundgren, M., Sannel, A.B.K., Tarnocai, C., Thom, T., Tuittila, E.-S.,  
 881 Turetsky, M., Väliranta, M., and der Linden, M., van Geel B., van Bellen, S., Vitt, D., Zhao,  
 882 Y., and Zhou, W.: A database and synthesis of northern peatland soil properties and Holocene  
 883 carbon and nitrogen accumulation, *Holocene*, 24, 1028-1042, 2014.  
 884 Lyman, R. L.: *Quantitative Paleozoology*. Cambridge University Press, 2008.  
 885 Maezumi, S.Y., Robinson, M., de Souza, J., Urrego, D.H., Schaan, D., Alves, D., and Iriarte, J.: New  
 886 insights from pre-Columbian land use and fire management in Amazonian Dark Earth forests,  
 887 *Front. Ecol. Evol.*, 6, 111, doi: 10.3389/fevo.2018.00111, 2018.  
 888 Mahowald, N.M., Randerson, J.T., Lindsay, K., Munoz, E., Doney, S.C., Lawrence, P.,  
 889 Schlunegger, S., Ward, D.S., Lawrence, D., and Hoffman, F. M.: Interactions between land use  
 890 change and carbon cycle feedbacks, *Glob. Biogeochem. Cy.*, 31, 96–113,  
 891 doi:10.1002/2016GB005374, 2017.  
 892 Marston, J.M.: Modeling wood acquisition strategies from archaeological charcoal remains, *J. Arch.*  
 893 *Sci.*, 36, 2192-2200, 2009.  
 894 Marston, J.M., Holdaway, S.J., and Wendrich, W.: Early- and middle-Holocene wood exploitation in  
 895 the Fayum basin, Egypt, *Holocene*, 27, 1812-1824, 2017.  
 896 Mauri, A., Davis, B. A. S., Collins, P. M., and Kaplan, J. O.: The influence of atmospheric circulation  
 897 on the mid-Holocene climate of Europe: a data–model comparison, *Clim. Past*, 10, 1925–1938,  
 898 <https://doi.org/10.5194/cp-10-1925-2014>, 2014.  
 899 Mazoyer, M., and Roudart, L.: *A History of World Agriculture: From the Neolithic to the Current*  
 900 *Crisis*. Earthscan, UK, 2006.  
 901 McGrath, M. J., Luyssaert, S., Meyfroidt, P., Kaplan, J. O., Burgi, M., Chen, Y., Erb, K., Gimmi, U.,  
 902 McInerney, D., Naudts, K., Otto, J., Pasztor, F., Ryder, J., Schelhaas, M. J., and Valade, A.:  
 903 Reconstructing European forest management from 1600 to 2010, *Biogeosci.*, 12, 4291-4316.  
 904 doi:10.5194/bg-12-4291-2015, 2015.  
 905 McLaughlin, T. R., Whitehouse, N. J., Schulting, R. J., McClatchie, M., Barratt, P. and Bogaard, A.:  
 906 The changing face of Neolithic and Bronze Age Ireland: A big data approach to the  
 907 settlement and burial records. *J. World Prehist.*, 29, 117-153. doi:10.1007/s10963-016-  
 908 9093-0, 2016.  
 909 Messori, G., Gaetani, M., Zhang, Q., Zhang, Q., and Pausata, F. S. R.: The water cycle of the mid-  
 910 Holocene West African monsoon: The role of vegetation and dust emission changes, *Int. J.*  
 911 *Climatol.*, 39, 1927– 1939, <https://doi.org/10.1002/joc.5924>, 2019.  
 912 Mitchell, L., Brook, E., Lee, J., Buizert, C., and Sowers, T.: Constraints on the late Holocene  
 913 anthropogenic contribution to the atmospheric methane budget, *Science*, 342, 964–966,  
 914 doi:10.1126/science.1238920, 2013.  
 915 Morrison, K.D., Hammer, E., Popova, L., Madella, M., Whitehouse, N., Gaillard, M.-J. and  
 916 LandCover6k Land-Use Group Members: Global-scale comparisons of human land use:  
 917 developing shared terminology for land-use practices for global changes, *PAGES Mag.*, 26, 8-  
 918 9, 2018.  
 919 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque,  
 920 J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura T., and Zhang,  
 921 H.: Anthropogenic and natural radiative forcing. in: *Climate Change 2013: The Physical*  
 922 *Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the  
 923 Intergovernmental Panel on Climate Change (T F Stocker, D Qin, G-K Plattner, M Tignor, S  
 924 K Allen, J Boschung, A Nauels, Y Xia, V Bex and P M Midgley (eds), Cambridge: Cambridge  
 925 University Press Cambridge, United Kingdom and New York, NY, USA, 2013.

- 926 Nakagawa, T., Tarasov, P.E., Nishida, K., Gotanda, K., and Yasuda, Y.: Quantitative pollen-based  
 927 climate reconstruction in central Japan: application to surface and Late Quaternary spectra,  
 928 *Quat. Sci. Rev.*, 21, 2099–2113, 2002.
- 929 Nielsen, A.B., Giesecke, T., Theuerkauf, M., Feeser, I., Behre, K.-H., Beug, H.-J., Chen, S.-H.,  
 930 Christiansen, J., Dörfler, W., Endtmann, E., Jahns, S., de Klerk, O., Köhl, N., Latałowa, M.,  
 931 Odgaard, B.V., Rasmussen, P., Stockholm, J.R., Voigt, R., Wiethold, J., and Wolters, S.:  
 932 Quantitative reconstructions of changes in regional openness in north-central Europe reveal  
 933 new insights into old questions, *Quat. Sci. Rev.*, 47, 131–149, 2012.
- 934 Oh, Y., Conte, M., Kang, S., Kim, J., and Hwang, J.: Population fluctuation and the adoption of food  
 935 production in prehistoric Korea: using radiocarbon dates as a proxy for population change,  
 936 *Radiocarbon*, 59, 1761–1770, doi: 10.1017/RDC.2017.1.22, 2017.
- 937 Ortega-Rosas, C.I., Guiot, J., Penalba, M.C., Ortiz-Acosta, M.E.: Biomization and quantitative  
 938 climate reconstruction techniques in northwestern Mexico—with an application to four  
 939 Holocene pollen sequences, *Glob. Planet. Change*, 61, 242–266, 2008.
- 940 Orton, D., Gaastra, J., and Vander Linden, M.: Between the Danube and the Deep Blue Sea:  
 941 zooarchaeological meta-analysis reveals variability in the spread and development of Neolithic  
 942 farming across the western Balkans, *Open Quat.*, 2, doi: 10.5334/oq.28, 2016.
- 943 Otto-Bliesner, B.L., Braconnot, P., Harrison, S.P., Lunt, D.J., Abe-Ouchi, A., Albani, S., Bartlein,  
 944 P.J., Capron, E., Carlson, A.E., Dutton, A., Fischer, H., Goelzer, H., Govin, A., Haywood, A.,  
 945 Joos, F., Legrande, A.N., Lipscomb, W.H., Lohmann, G., Mahowald, N., Nehrbass-Ahles, C.,  
 946 Pausata, F.S.R., Peterschmidt, J.-Y., Phipps, S.J., Renssen, R., and Zhang, Q.: The PMIP4  
 947 contribution to CMIP6 – Part 2: Two interglacials, scientific objective and experimental design  
 948 for Holocene and Last Interglacial simulations. *Geosci. Mod. Dev.*, 10, 3979–4003,  
 949 <https://doi.org/10.5194/gmd-10-1-2017>, 2017.
- 950 Pausata, F.S.R., Messori, G., Zhang, Q.: Impacts of dust reduction on the northward expansion of the  
 951 African monsoon during the Green Sahara period, *Earth Planet. Sci. Lett.*, 434, 298–  
 952 307, <https://doi.org/10.1016/j.epsl.2015.11.049>, 2016.
- 953 Perugini, L., Caporaso, L., Marconi, S., Cescatti, A., Quesada, B., de Noblet-Ducoudré, N., House,  
 954 J.I., and Arneth, A.: Biophysical effects on temperature and precipitation due to land cover  
 955 change, *Environ. Research Lett.*, 12, 053002, <https://doi.org/10.1088/1748-9326/aa6b3f>, 2017.
- 956 Phelps, L.N., and Kaplan, J.O.: Land use for animal production in global change studies: Defining  
 957 and characterizing a framework, *Glob Chang Biol*, 23, 4457–4471, 10.1111/gcb.13732, 2017.
- 958 Pirzamanbein, B., Lindström, J., Poska, A., Sugita, S., Trondman, A., Fyfe, R., Mazier, F., Nielsen,  
 959 A.B., Kaplan, J.O., Bjune, A.E., Birks, H.J.B., Giesecke, T., Kangur, M., Latałowa, M.,  
 960 Marquer, L., Smith, B., and Gaillard, M.-J.: Creating spatially continuous maps of past land  
 961 cover from point estimates: A new statistical approach applied to pollen data, *Ecol. Complexity*,  
 962 20, 127–141, 2014.
- 963 Pirzamanbein, B., Lindström, J., Poska, A., and Gaillard, M.-J.: Modelling spatial compositional data:  
 964 Reconstructions of past land cover and uncertainties, *Spatial Stat.*, 24, 14–31, 2018.
- 965 Pongratz, J., Reick, C., Raddatz, T., and Claussen, M.: A reconstruction of global agricultural areas  
 966 and land cover for the last millennium, *Glob. Biogeochem. Cy.*, 22, 2008.
- 967 Pongratz, J., Reick, C.H., Raddatz, T., and Claussen, M.: Biogeophysical versus biogeochemical  
 968 climate response to historical anthropogenic land cover change, *Geophys. Res. Lett.*, 37,  
 969 L08702, doi:10.1029/2010GL043010, 2010.
- 970 Pongratz, J., Reick, C. H., Houghton, R. A., and J. I. House, J. I.: Terminology as a key uncertainty  
 971 in net land use and land cover change carbon flux estimates, *Earth Syst. Dynam.*, 5, 177–195,  
 972 [www.earth-syst-dynam.net/5/177/2014/](http://www.earth-syst-dynam.net/5/177/2014/) doi:10.5194/esd-5-177-2014, 2014.
- 973 Ramankutty, N., and Foley, J.A.: Estimating historical changes in global land cover: Croplands from  
 974 1700 to 1992, *Glob. Biogeochem. Cy.*, 13, 997–1027, 1999.
- 975 Reitz, E. J., and Wing E.S.: Zooarchaeology. Cambridge University Press, 2008.

976 Rick, J.W.: Dates as data: an examination of the Peruvian Preceramic radiocarbon record, *Am. Antiq.*,  
977 52, 55–73, 1987.

978 Robinson, E., Zahid, H.J., Coddling, B.F., Haas, R., and Kelly, R.L.: Spatiotemporal dynamics of  
979 prehistoric human population growth: radiocarbon ‘dates as data’ and population ecology  
980 models, *J. Arch. Sci.*, 101, 63–71, 2019.

981 Ruddiman, W. F.: The anthropogenic greenhouse era began thousands of years ago, *Clim. Change*,  
982 61, 261–293, doi:10.1023/B:CLIM.00000004577.17928.fa, 2003.

983 Russell, T., Silva, F., and Steele, J.: Modelling the spread of farming in the Bantu-speaking regions  
984 of Africa: an archaeology-based phylogeography, *PlosONE*, 9, e87584, doi:  
985 10.1371/journal.pone.0087584, 2014.

986 Shennan, S., Downey, S.S., Timpson, A., Edinborough, K., Colledge, S., Kerig, T., Manning, K., and  
987 Thomas, M.G.: Regional population collapse followed initial agriculture booms in mid-  
988 Holocene Europe, *Nat. Comms.*, 4, 248, doi: 10.1038/ncomms3486, 2013.

989 Shevliakova, E., Pacala, S. W., Malyshev, S., Hurtt, G. C., Milly, P. C. D., Caspersen, J. P., Sentman,  
990 L. T., Fisk, J. P., Wirth, C., and Crevoisier, C.: Carbon cycling under 300 years of land use  
991 change: Importance of the secondary vegetation sink, *Glob. Biogeochem. Cy.*, 23, GB2022,  
992 doi:10.1029/2007GB003176, 2009.

993 Sigl, M., Winstrup, M., McConnell, J.R., Welten, K.C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee,  
994 M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O.J.,  
995 Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D.R., Pilcher, J.R., Salzer, M., Schüpbach,  
996 S., Steffensen, J.P., Vinther, B.M., and Woodruff, T.E.: Timing and climate forcing of volcanic  
997 eruptions for the past 2,500 years, *Nature*, 523, 543–549, <https://doi.org/10.1038/nature14565>,  
998 2015.

999 Silva, F., and Steele, J.: New methods for reconstructing geographical effects on dispersal rates from  
1000 large-scale radiocarbon databases, *J. Arch. Sci.*, 52, 609–620, doi: 10.1016/j.jas.2014.04.021,  
1001 2014.

1002 Silva, F., and Vander Linden, M.: Amplitude of travelling front as inferred from <sup>14</sup>C predicts levels  
1003 of genetic admixture among European early farmers, *Sci. Reports*, 7, 11985, doi:  
1004 10.1038/s41598-017-12318-2, 2017.

1005 Silva, F., Stevens, C.J., Weisskopf, A., Castillo, C., Qin, L., Bevan, A., and Fuller, D.Q.: Modelling  
1006 the geographical origin of rice cultivation in Asia using the Rice Archaeological Database,  
1007 *PlosOne*, 10, e0137024, 2015.

1008 Singarayer, J.S., Valdes, P.J., Friedlingstein, P., Nelson, S., and Beerling, D.J.: Late Holocene  
1009 methane rise caused by orbitally controlled increase in tropical sources, *Nature*, 470, 82– 85,  
1010 doi:10.1038/nature09739, 2011.

1011 Smith, M.C., Singarayer, J.S., Valdes, P.J., Kaplan, J.O., and Branch, N.P.: The biogeophysical  
1012 climatic impacts of anthropogenic land use change during the Holocene, *Clim. Past*, 12, 923–  
1013 941, doi: <https://doi.org/10.5194/cp-12-923-2016>, 2016.

1014 Steinhilber, F., Abreu, J. A., Beer, J., Brunner, I., Christl, M., Fischer, H., Heikkilä, U., Kubik, P. W.,  
1015 Mann, M., McCracken, K.G., Miller, H., Miyahara, H., Oerter, H., and Wilhelms, F.: 9400  
1016 years of cosmic radiation and solar activity from ice cores and tree rings, *Proc. Natl. Acad. Sci.*,  
1017 109, 5967–5971, 2012.

1018 Stocker, B. D., Strassmann, K., and Joos, F.: Sensitivity of Holocene atmospheric CO<sub>2</sub> and the  
1019 modern carbon budget to early human land use: analyses with a process-based model,  
1020 *Biogeosci.*, 8, 69–88, doi:10.5194/bg-8-69-2011, 2011.

1021 Stocker, B.D., Yu, Z., Massa, C., and Joos, F.: Holocene peatland and ice-core data constraints on  
1022 the timing and magnitude of CO<sub>2</sub> emissions from past land use, *Proc. Natl. Acad. Sci.*, 114,  
1023 1492–1497, doi:10.1073/pnas.1613889114, 2017.

1024 Styring, A., Rösch, M., Stephan, E., Stika, H.-P., Fischer, E., Sillmann, E., and Bogaard, A.:  
1025 Centralisation and long-term change in farming regimes: comparing agricultural practices in

- Neolithic and Iron Age south-west Germany, *Proc. Prehist. Soc.*, 83: 357-381, doi: 10.1017/ppr.2017.3, 2017.
- Sugita, S.: Theory of quantitative reconstruction of vegetation I: pollen from large sites REVEALS regional vegetation composition, *Holocene*, 17, 229–241, 2007.
- Tarasov, P., Williams, J.W., Andreev, A., Nakagawa, T., Bezrukova, E., Herzs Schuh, U., Igarashi, Y., Müller, S., Werner, K., and Zheng, Z.: Satellite- and pollen-based quantitative woody cover reconstructions for northern Asia: Verification and application to late-Quaternary pollen data, *Earth Planet. Sci. Lett.*, 264, 284–298, 2007.
- Tauger, M.B: *Agriculture in World History*, Routledge, 2013.
- Tierney, J.E., Pausata, F.S.R., and deMenocal, P.B.: Rainfall regimes of the Green Sahara, *Sci. Advan.*, 3, e1601503, 2017.
- Timpson, A., Colledge, S., Crema, E., Edinborough, K., Kerig, T., Manning, K., Thomas, M. G. & Shennan, S.: Reconstructing regional population fluctuations in the European Neolithic using radiocarbon dates: a new case-study using an improved method, *J. Arch. Sci.*, 52, 549-557, doi: 10.1016/j.jas.2014.08.011, 2014.
- Toohy, M. and Sigl, M.: Volcanic stratospheric sulphur injections and aerosol optical depth from 500 BCE to 1900 CE, *Earth Syst. Sci. Data*, 9, 809-831, <https://doi.org/10.5194/essd-9-809-2017>, 2017.
- Trondman, A. K., Gaillard, M.-J., Mazier, F., Sugita, S., Fyfe, R., Nielsen, A.B., Twiddle, C., Barratt, P., Birks, H.J.B., Bjune, A. E., Björkman, L., Broström, A., Caseldine, C., David, R., Dodson, J., Dörfler, W., Fischer, E., van Geel, B., Giesecke, T., Hultberg, T., Kalnina, L., Kangur, M., van der Knaap, P., Koff, T., Kuneš, P., Lagerås, P., Latałowa, M., Lechterbeck, J., Leroyer, C., Leydet, M., Lindbladh, M., Marquer, L., Mitchell, F.J. G., Odgaard, B.V., Peglar, S.M., Persson, T., Poska, A., Rösch, M., Seppä, H., Veski, S., and Wick, L.: Pollen-based quantitative reconstructions of Holocene regional vegetation cover (plant-functional types and land-cover types) in Europe suitable for climate modelling, *Glob. Change Biol.*, 21, 676-697, doi:10.1111/gcb.12737, 2015.
- Trondman, A.-K., Gaillard, M.-J., Sugita, S., Björkman, L., Greisman, A., Hultberg, T., Lagerås, P., and Lindbladh, M.: Are pollen records from small sites appropriate for REVEALS model-based quantitative reconstructions of past regional vegetation? An empirical test in southern Sweden, *Veget. Hist. Archaeobot.*, 25, 131–151, doi: 10.1007/s00334-015-0536-9, 2016.
- van den Hurk, B., Kim, H., Krinner, G., Seneviratne, S.I., Derksen, C., Oki, T., Douville, H., Colin, J., Ducharne, A., Cheruy, F., Viovy, N., Puma, M.J., Wada, Y., Li, W., Jia, B., Alessandri, A., Lawrence, D.M., Weedon, G.P., Ellis, R., Hagemann, S., Mao, J., Flanner, M.G., Zampieri, M., Materia, S., Law, R.M., and Sheffield, J.: LS3MIP (v1.0) contribution to CMIP6: the Land Surface, Snow and Soil moisture Model Intercomparison Project – aims, setup and expected outcome, *Geosci. Model Dev.*, 9, 2809-2832, <https://doi.org/10.5194/gmd-9-2809-2016>, 2016.
- Vavrus, S., Ruddiman, W.F., and Kutzbach, J.E.: Climate model tests of the anthropogenic influence on greenhouse-induced climate change: the role of early human agriculture, industrialization, and vegetation feedbacks, *Quat. Sci. Rev.*, 27, 1410-1425, 2008.
- Veal, R., 2017. Wood and charcoal for Rome: towards an understanding of ancient regional fuel economics, In de Haas, T. & Gijs, T. (eds), *Rural communities in a globalizing economy: new perspectives on the economic integration of Roman Italy*, Brill, (New York and Leiden): pp.388-406.
- Viau, A.E., and Gajewski, K.: Reconstructing millennial, regional paleoclimates of boreal Canada during the Holocene, *J. Clim.*, 22, 316–330, 2009.
- Viau, A., Gajewski, K., Sawada, M., and Fines, P.: Mean-continental July temperature variability in North America during the past 14,000 years, *J. Geophys. Res. Atmos.*, 111, D09102, doi:10.1029/2005JD006031, 2006
- Weiberg, E., Hughes, R. E., Finné, M., Bonnier, A., and Kaplan, J. O.: Mediterranean land use



1076 systems from prehistory to antiquity: a case study from Peloponnese (Greece), *J. Land Use Sci.*,  
1077 1-20, doi:10.1080/1747423x.2019.1639836, 2019.

1078 Whitehouse, N., Schulting, R. J., McClatchie, M., Barratt, P., LcLaughlin, T.R., Bogaard, A.,  
1079 Colledge, S., Marchant, R., Gaffrey, J., and Bunting, M.J.: Neolithic agriculture on the  
1080 European western frontier: the boom and bust of early farming in Ireland, *J. Arch. Sci.*, 51, 181-  
1081 205, doi: 10.1016/j.jas.2013.08.009, 2014.

1082 Williams, A.: The use of summed radiocarbon probability distributions in archaeology: a review of  
1083 methods. *J. Archaeol. Sci.*, 39, 578–589, <https://doi.org/10.1016/j.jas.2011.07.014>, 2012.

1084 Wilmshurst, J.M., McGlone, M.S., Leathwick, J.R., and Newnham, R.M.: A pre-deforestation pollen-  
1085 climate calibration model for New Zealand and quantitative temperature reconstructions for the  
1086 past 18000 years BP, *J. Quat. Sci.*, 22, 535–547, 2007.

1087 Wright, P.: Preservation or destruction of plant remains by carbonization. *J.Arch.Sci* 30, 577-583,  
1088 doi: 10.1016/S0305-4403(02)00203-0, 2003.

1089 Zahid, H.J., Robinson, E., and Kelly, R.L.: Agriculture, population growth, and statistical analysis of  
1090 the radiocarbon record, *Proc. Natl. Acad. Sci.*, 113, 931-935, doi: 10.1073/pnas.1517650112,  
1091 2016.

1092 Zanon, M., Davis, B.A.S., Marquer, L., Brewer, S., and Kaplan, J.O.: European forest cover during  
1093 the past 12,000 years: a palynological reconstruction based on modern analogs and remote  
1094 sensing, *Front. Plant Sci.*, 9, 253, doi: 10.3389/fpls.201800253, 2018.

1095 Zimmermann, A., Wendt, K.P., and Hilpert, J.: Landscape archaeology in central Europe. *Proc.*  
1096 *Prehist. Soc.*, 75, 1-53, doi: 10.1017/S007949X00000281, 2009.

1097 Zohary, D., Hopf, M., and Weiss, E.: Domestication of Plants in the Old World: The Origin and  
1098 Spread of Domesticated Plants in South-west Asia, Europe, and the Mediterranean Basin, 4th  
1099 Edn., Oxford University Press, Oxford, 2012.