a point-by-point response to the reviews

Interactive comment on "Mitigation of Model Bias Influences on Wave Data Assimilation with Multiple Assimilation Systems Using WaveWatch III v5.16 and SWAN v41.20" from anonymous referee #1

In general:

The comments and suggestions from Reviewer #1 are excellent, which are very helpful for us to compare the influence of error sources more clearly. In the revision, we have added: (1) section 3.4 to quantify the comparison of the wind forcing error and model error in time and space, (2) clearer description about how to get SWH (significant wave height) bias and bias correction.

The following is the point-by-point reply to address the comments and suggestions of Reviewer #1.

This study investigates the challenge of model bias on wave model data assimilation. It applies a set of twin-model experiments to quantify the different error sources in wave model: initial, boundary, and model error. Based that It provide a simple statistical approach to reduce the impact of model bias and improve the assimilation results.

The topic is interesting and important for wave data assimilation, well fit for GMD. The experiments are well designed and manuscript is in good shape. Here I only have few points to further polish this work. Therefore, I suggest miner revision for current version.

RE: Thanks for reviewer's encouragement.

1 There are three error sources: initial/boundary/model-bias. You have identify them in your biased/unbiased twin experiment. It turns out both boundary and model-bias are important.

In my pointer of view, both error sources could lead to SWH biases/uncertainties for assimilation/simulation. It would be great to separate them and quantify the improvement percentage by your data assimilation from your biased/unbiased twin experiments. The SWH bias and its decrease percentage may give your hind on mitigation the assimilation bias for the real observation. Similar separation should apply to real observation assimilation cases. In your figure 5, mainly represent the

SWH bias. You can recalculate figure 6 (RMSE) after remove the bias in figure 5, which represent the uncertainties related to boundary/model bias.

When you apply average to reduce the assimilation error of SWH. You only reduces the uncertainty part but not the bias part. You have to direct remove the bias from the reanalysis.

RE: Thanks for your thoughtful advice. We have added the quantitative comparison between boundary and model bias, please see lines 443-449 for their performances in the twin experiments. In the experiments of real-data assimilation, we also have replotted the Fig. 6 (Fig. 9 in the revised version) after removing the model bias shown in Fig. 5 (Fig. 6 in the revision). The reviewer is right, as a try, the first method of bias correction is used to reduce the uncertainty part from the reanalysis and the second method is used to reduce the bias part.

2 In your real observation assimilation, the boundary and model-bias are both included. You may compare spatial pattern and the decreasing percentage of SWH bias with those your biased/unbiased twin experiments to speculate which source (boundary/model-bias) has stronger impact in certain area..

RE: Thanks for your excellent suggestion. In the twin experiments, we have compared the spatial pattern of errors caused by wind forcing and model bias, please see lines 458-476 and Fig. 5.

3 You only applied one kind of wind forcing for different models and then use them to do the bias correction. Since the bias also come from the boundary forcing. I encourage you using two kinds of wind forcing to further investigate the bias/uncertainty generated by the boundary forcing.

RE: Thank you for your helpful comment. In the real experiment, it's very difficult to distinguish the error sources. Then, as an understanding, we have tested the influences of these two error sources in the twin experiments. Next, we use a better wind forcing from ERA-Interim reanalysis in this manuscript to drive wave models and conduct bias correction in the real experiments. It is more reasonable and indicative to use bias correction mitigating hybrid errors and finally improve the quality of data assimilation results in the case of assimilating real observations.

4 it is unclear how do you get the SWH bias for bias correction, what is the spatial pattern. I thought it should refer to the figure 5, but I did not find those in the manuscript.

RE: Thanks for your kind reminder. More detailed description about how to get SWH bias has added in the revision, please see lines 556-561 and Fig. 6.

5 Model description in page 4. This part need be more condensed with appropriate reference. Readers would appreciate more on the differences among those models, instead of the comparison to their own previous version. You may highlight the advantage or disadvantage among three modes.

RE: Thanks for your good advice. We simplified the description of this section, please see lines 112-119.

Interactive comment on "Mitigation of Model Bias Influences on Wave Data Assimilation with Multiple Assimilation Systems Using WaveWatch III v5.16 and SWAN v41.20" from anonymous referee #2

In general:

The comments and suggestions from Reviewer 2 are very constructive, which are helpful for us to improve this manuscript. In this revision, we have added: 1) clearer captions of figures, 2) more detailed description for the assumption of equation in section 2.3, 3) more detailed description about how to get SWH bias and bias correction, 4) more discussions on "online-type" bias correction, the improvement of windsea and swell after wind correction etc.

The following is the point-by-point reply to address the comments and suggestions of Reviewer 2.

This paper describes the effect of OI data assimilation on ocean wave simulations, including bias correction. The work is mainly done through running a series of twin experiments. The paper contains scientifically interesting results, clear writing and good quality of figures. There is a potential to use the system to produce a global wave reanalysis (ensemble) product. There are some questions and comments, which are descried as following.

RE: Thanks for your constructive comments, which have been fully addressed in the revision.

In Model configurations: the resolution of wind forcing could cause misleading. For example, 0.125x0.125deg is not the resolution of ERA-Interim itself (in which the model resolution is much coarser), but is the resolution of gridded reanalysis data. Please clarify.

RE: Thanks for your reminder. Clarified. Please see line 131.

AVISO data: along-track data are used in wave simulation and gridded data are used for validation. What will be the possible effect due to difference of these data? Are there any observation-related errors/uncertainties that would influence the validation results? Also, the validation data are not independent to those assimilated. Do author consider using wave buoy data for validation in the future?

RE: Thanks for your great comment. Along-track data is more effective to sample local wave information, while gridded dataset is an integration of multiple satellites, which more focus on the averaging variation over several days. Therefore, it is reasonable absorbing the along-track data into wave simulation in a small assimilation window with the assumption of little wave change in a short time. Nevertheless, it should be mentioned that due to a few satellites available, observations may lack representativeness in global area. At the same time, there is a potential validation error considering the obtained way of gridded data. As the reviewer said, the validation data are not fully independent to those assimilated. In the future, we will add buoy data to the verification, which is more reasonable and powerful to further illustrate the effect of bias correction. We have added discussions in the section 5. Please see lines 657-658.

Following above comment, now some latest observation data contain wave direction information (i.e. peak wave direction, 2D spectra). These obs can be used to assimilate model wave spectra, which will have more advantages than assimilating SWH only. This might be worth mentioning somewhere in the paper.

RE: Thanks for the good suggestion. We have added relevant description about observation variables and more discussions. Please see lines 481-484 and 652-654.

L190: why did authors choose Sigma_M=0.6m and sigma_O=0.25m? Are the same model and observations as used in (Qi and Fan, 2013)? Is sigma varying with time, space and models? There ought to be some assumptions before using these parameters.

RE: Thanks for your good advice. We have added clearer description of sigma. Please see lines 212-216.

In equation 3, sigma_i/sigma_k*r_i,k is for SWH (or wave spectra?) correlation and statistics. I just wonder whether wave covariances will have the same structure as wave error covariances as equation 3 is supposed to be for error covariance. In a storm the high-sea state may have a few hundred km long, but this doesn't necessarily mean the error is propagating in a few hundred km distance. I don't have a solution for this question. But it needs some assumptions on equation 3, with clarifications of potential drawbacks, before using it.

RE: Thanks for your thoughtful question. We have added the description of relevant variables and the assumption of equation 3. Please see lines 224-227.

2.3 section step 4: ocean waves have two components, i.e. windsea waves and swells. The wind should (only) be corrected based on the analysed windsea waves, while analysed swells that are not directly forced by local wind have no impact on wind correction. This concept is described in Lionello et al 1995. Mostly wave models can output windsea and swell SWH. Why did not authors use the windsea SWH (rather than use total SWH) to correct wind forcing? Using analysed total SWH to correct wind, wind could be overly corrected for example when it is a swell dominant event at the DA time.

RE: This is an excellent suggestion. We have tested all corresponding experiments about correcting wind forcing with windsea wave height. However, the results do not show substantial improvement. The possible reason could be attributed to the simple correction method. It seems that in this simple correction method, it is difficult to distinguish the signal by using windsea wave height to correct wind from using the total SWH. In the future, we plan refine the correction method and study this part to enhance the signal-to-noise ratio. Please see lines 433-435 for added discussions on this point.

Fig2: not very clear what b and c are for in these snapshots. SWH difference between what? Which is for increment? Please clarify.

RE: Thank you for your reminder. We have added the detailed caption of Fig. 2. Please see lines 860-865.

Fig3,4: are these statistics for global mean or any regions? In Fig4, what are the correlation coefficients for (spatial correlations)? Same for other figures.

RE: Thanks for your good comments. We have added all relevant captions.

Fig4 shows wind correction does not clearly improve SWH simulation when assimilating J2. What about assimilating all satellite tracks i.e. J2+J3+SA? Does wind correction have a stronger effect?

RE: Thanks for your great suggestion. We have redisplayed the results of wind correction assimilating with three satellite tracks. Please see Fig. 4.

Section 4.1 and Fig7: results show that wind correction only improves wave simulation by certain degrees. 1) Does the wind correction scheme used here have an impact? Can authors show (or suggest) any difference when using the scheme of Lionello et al 1995 (see above comment); 2) How about the spatial distribution of Fig7 red lines? 3) Are there more improvements seen in windsea waves than in swells? (you can simply partition wind sea waves and swell waves from total SWH). I assume wind correction will have a stronger impact in windsea wave simulations at high latitudes with strong wind.

RE: Thanks for your thoughtful question. 1) In the real experiments, wind correction has a positive effect improving the assimilation results. However, different wave models have different improvement magnitudes. Here, SWAN has a weaker performance than WW3. Compared with the scheme in Lionello et al (1995), a big difference is that a coarse adjustment without distinguishing the wave structure (windsea or swell) is conducted to reconstruct the 2-dimensional wave spectrum. We will take this distinction into consideration in the future study. 2) We have displayed the spatial distribution of the red lines in Fig. 7, please see panel d and h of Fig. 9. 3) About the improvement of windsea and swell, we also have added discussions in the revision. Please see lines 582-588.

Section 4.2: Please describe a bit more how the bias is produced and removed in these simulations. It is not very clear to me. How was 'bias correction of model control run' implemented?

RE: Thanks for your great advice. We have added more detailed description about how to get bias and bias correction (please see lines 570-574). And we also have explained the "bias correction of model control run" (please see lines 615-617).

Was bias correction in this paper like the offline-type bias correction? If we can have a long-term historic run, to produce the climatology of wave bias, and then use it as an offline bias correction term before online DA term (simply like some DA procedures in European systems), will this produce similar results as produced in Fig 10? This offline term can potentially be used in forecast as well. One normally won't expect that DA can efficiently correct the long-term and persistent bias, but expects DA is more powerful for correcting the instantons/short-term/flow-dependent errors. It is not simple to have an immediate answer for this question, but it will be useful to have a discussion somewhere in the paper.

RE: Thanks for your excellent suggestion. As the reviewer said, bias correction in this manuscript is an "offline-type". We have 1-year output of model control run and with data assimilation, then do offline-type bias correction and show in Fig. 11. If a longer run is conducted, similar results are supposed to display like Fig. 11. We will verify it in the future study. The effect of data assimilation on correcting the instantons errors is discussed in lines 654-658.

Line 61: to produce=> for producing L337: inaccurate

RE: Thanks for your advice. We have modified this error, please line 375.

a list of all relevant changes made in the manuscript

In this revision, we have added:

(1) more detailed description to make all experiments easily operate and follow, such as: a) the equation assumptions, b) the figure captions, c) the method of obtaining SWH bias and bias correction.

(2) deeper discussion to comprehensive understanding the effect of bias correction on wave data assimilation, such as: a) to quantify the error caused by wind forcing and model bias in time and space, b) to compare the improvement of windsea and swell after wind correction.

a marked-up manuscript version

Mitigation of Model Bias Influences on Wave Data Assimilation with Multiple Assimilation Systems Using WaveWatch III v5.16 and SWAN v41.20

Jiangyu Li^{1,4} and Shaoqing Zhang*1,2,3,4

⁵ ¹Key Laboratory of Physical Oceanography, MOE, China; Ocean University of China, Qingdao, 266100, China ²Pilot National Laboratory for Marine Science and Technology (QNLM), Qingdao, 266100, China

³International Laboratory for High-Resolution Earth System Model and Prediction (iHESP), Qingdao, 266100, China

10 ⁴The College of Oceanic and Atmospheric Sciences, Ocean University of China, Qingdao, 266100, China

Correspondence to: Shaoqing Zhang (szhang@ouc.edu.cn)

Abstract. High-quality wave prediction with a numerical wave model is of societal value. To initialize the wave model, wave data assimilation (WDA) is necessary to combine the model and

- 15 observations. Due to imperfect numerical schemes and approximated physical processes, a wave model is always biased in relation to the real world. In this study, two assimilation systems are first developed using two nearly independent wave models; then, "perfect" and "biased" assimilation frameworks based on the two assimilation systems are designed to reveal the uncertainties of WDA. A series of "biased" assimilation experiments is conducted to systematically examine the adverse
- 20 impact of model bias on WDA. A statistical approach based on the results from multiple assimilation systems is explored to carry out bias correction, by which the final wave analysis is significantly improved with the merits of individual assimilation systems. The framework with multiple assimilation systems provides an effective platform to improve wave analyses and predictions and help identify model deficits, thereby improving the model.

25 1 Introduction

Ocean waves, referring to the ocean surface gravity waves driven by wind, are important physical processes in the study of multiscale coupled systems. Many studies show that ocean waves are necessary for upper ocean mixing processes, whether in small-scale coastal simulations or large-scale global climate simulations (e.g., Babanin et al., 2009; Huang and Qiao, 2010; Qiao et al., 2004, 2010).

- 30 The existence of ocean waves can modify the structures of both atmospheric and marine boundary layers by providing sea surface roughness, wave-induced bottom stress, breaking wave-induced mixing, and so on, which ultimately influence air-sea momentum and heat exchange. Therefore, ocean waves are an important component in atmosphere-ocean interaction flux processes (e.g., Chen et al., 2007; Doyle, 2002; Liu et al., 2011; Warner et al., 2010). In addition, the study of ocean waves can reduce
- 35 and prevent marine disasters and provide guidance for development of the social economy (e.g., Folley and Whittaker, 2009; Rusu, 2015; Wei et al., 2017). Thus, studying ocean waves is of great scientific and social significance.

At present, ocean wave observational techniques are constantly being improved (e.g., Daniel et al., 2011; Hisaki, 2005). Except for traditional buoy observations (e.g., Mitsuyasu et al., 1980; Rapizo et al.,

- 40 2015; Walsh et al., 1989), satellites can provide much near real-time wave observational information, which is beneficial for understanding the state of ocean waves (e.g., Gommenginger et al., 2003; Lzaguirre et al., 2011; Queffeulou P, 2004). However, observations always represent scattered samples in time and space in the real world and therefore do not represent the complete three-dimensional structure and temporal evolution of real world waves.
- 45 Numerical wave models are a powerful tool for studying the physical processes of ocean waves and predicting future wave states. Following the development of the previous two generations, third-generation wave models, such as WAve Modeling (WAM) (WAMDI Group, 1988), WaveWatch III (WW3) (Tolman, 1991), Simulating Waves Nearshore (SWAN) (Booij et al., 1999), and MArine Science and Numerical Modeling (MASNUM) (Yang et al., 2005), integrate the spectral action balance
- 50 equation describing the two-dimensional ocean wave spectrum evolution without additional *ad hoc* assumptions regarding the spectral shape, and these third-generation models are more robust for arbitrary wind fields than previous models. However, there are generally three error sources in wave models. One error source is from an incomplete understanding of the physical processes, approximate expressions of the numerical discretization schemes and so on, which causes systematic errors that are
- 55 usually referred to as wave model bias. The second error source is due to inaccurate wind forcings of wave models. The third error source is from the initial condition uncertainties, which can grow due to nonlinearity of the model equations during model forwarding. In this sense, the model simulated waves do not represent the real world either.

Given the scattering nature of observational information and the approximate characteristics of wave modelling, wave model data assimilation (WDA) is necessary to combine the advantages of both the

60

删除了: modeling

model and observations. WDA optimizes the model initial conditions to produce more accurate wave forecasts and produces more accurate evolution of 3-dimensional wave states to elucidate the underlying mechanisms; this approach dates back to the 1980s (e.g., Esteva, 1988; Janssen et al., 1989).

- 65 Since then, many advanced WDA methods have been developed (e.g., Abdalla et al., 2013; Bauer et al., 1996; Greenslade and Young, 2004; Jesus and Cavaleri, 2015; Lionello et al., 1992; Sun et al., 2017; Vorrips et al., 1999), and their applications have been assessed (e.g., Francis and Stratton, 1990; Heras et al., 1994; Stopa and Cheung, 2014). Furthermore, various observation types, such as buoy, radar and satellite, have been applied to WDA (e.g., Bhatt et al., 2005; Breivik et al., 1998; Feng et al., 2006;
- 70 Greenslade, 2001; Hasselmann et al., 1997; Qi and Cao, 2016; Voorrips, 1999; Waters et al., 2013), and the wave forecasts have also been directly addressed (e.g., Almeida et al., 2016; Emmanouil et al., 2012; Lionello et al., 1995; Qi and Fan, 2013; Sannasiraj et al., 2006; Voorrips, 1999; Wang and Yu, 2009; Zhang et al., 2003).

Due to the approximate nature of the numerical discretization and physical processes, a systematic

- 75 difference between a model and the real world (i.e., model bias) exists. As noted by Zhang et al. (2012), since model bias is not well defined in observational space, the influence of model bias on data assimilation is a challenging research topic. Alternatively, one can simulate model bias using a pair of models and study the adverse impacts on data assimilation. Inspired by previous work (e.g., Dee, 2005; Zhang et al., 2012), here, we use a simple data assimilation scheme with two wave models (WW3 and
- 80 SWAN) to explore the influences of different error sources on WDA. The adverse impacts of wind forcing errors and initial condition uncertainties as well as wave model bias on WDA are studied first, and then two simple statistical methods for bias correction are developed to mitigate assimilation errors and improve wave analysis.

This paper is organized as follows. After the introduction, the methodology is presented in section 2, 85 including a brief description of the employed models and observations, development of the two WDA systems using the WW3 and SWAN models, as well as the design of experiments throughout the study.

Section 3 presents the model bias analysis and the adverse impacts of model bias on WDA. In section 4, the method used to mitigate model bias influences on wave assimilation is explored. Finally, the discussion and conclusion are given in section 5.

删除了: the

2 Methodology

2.1 Models and data

2.1.1 Three models

In the wave models, the variance spectrum or energy density $E(\sigma, \theta)$ is a quantity that represents the 95 wave energy distribution in the radian frequency (σ) and propagation direction(θ). Without ambient ocean currents, the variance or energy of a wave package is conserved. However, if the current is involved, due to the work done by the current on the mean momentum transfer of waves (Longuet-Higgins and Stewart, 1961, 1962), the energy of a spectral component is no longer conserved. In general, an action density spectrum defined as N=E/ σ is considered within the models. Then, the 100 governing equation of the wave model can be written as follows:

$$\frac{\partial N}{\partial t} + \nabla_{\chi} \cdot (\underset{c_g}{\rightarrow} N) + \frac{\partial c_{\sigma} N}{\partial \sigma} + \frac{\partial c_{\theta} N}{\partial \theta} = \frac{S_{tot}}{\sigma}.$$

The left-hand side <u>of this equation</u> is the kinematic <u>process during wave propagation</u>. The second term describes the wave energy propagation in two-dimensional geographical space denoted by $\frac{1}{x}$. The $\frac{1}{c_{r}}$ is

the group velocity that follows the dispersion relation. The third term represents the effect of shifting 105 the radian frequency due to variation in depth. The fourth term represents the depth-induced refraction. c_{σ} and c_{θ} are wave velocities in the frequency σ and direction θ , respectively. On the right-hand side, S_{tot} is the nonconservative source/sink term representing all physical processes that generate, dissipate, or redistribute wave energy. Typically, there are three important physical processes that contribute to S_{tot} , which include the atmosphere-wave interaction, nonlinear wave-wave interaction, and wave-

110 ocean interaction. In a shallow-water case, additional processes must be considered, such as wave-bottom interaction, depth-induced breaking, and triad wave-wave interaction.
In this study, we use three advanced third-generation spectrum models, WAve Modeling (WAM, Cycle 4.5.4) (WAMDI Group, 1988), WaveWatch III (WW3, version 5.16) (Tolman, 1991), and Simulating WAves Nearshore (SWAN, version 41.20) (Booij et al., 1999), which have improved a lot from their

115 predecessors, such as numerical and physical approaches. These wave models have the same form of governing equation, however their numerical implement processes are different due to different consideration. For example, SWAN is more focused on wave propagation processes in shallow water, while WAM and WW3 pay more attention in deep water. For more comparisons, please see section 2.2. 删除了: part 删除了: of this equation

删除了:. The first one is

删除了: developed from the Sea Wave Modelling Project (SWAMP)

设置了格式:英语(美国)

2.1.2 Model configurations

- 125 Three wave models use two-dimensional spectral space containing 29 frequencies that cover from .035 Hz to .555 Hz with a logarithmic distribution and 24 equidistant directions. The geographic space is from 180° W to 180° E in the zonal direction and 75° S to 75° N in the meridional direction with a 1°×1° grid resolution. The topography in this study is taken from the high-resolution ETOPO1 dataset provided by NOAA (website: https://www.ngdc.noaa.gov/mgg/global/). The wind forcing has two
- 130 sources, both of which have 6-hour time intervals. The first dataset is the ERA-Interim reanalysis from European Centre for Medium-Range Weather Forecasts (ECMWF), with a resolution of <u>75°×75°</u> (http://apps.ecmwf.int/datasets/data/interim-full-daily/). The second dataset is the CFSRv2 dataset from NCEP, with a resolution of .205° (longitude) ×.204° (latitude) (https://rda.ucar.edu/datasets/ds094.1/). The time_step of all three models is 15 minutes. All relevant parameters <u>above</u> are set to be identical for

135 every wave model (WW3, SWAN, and WAM) in this study.

2.1.3 Data

The AVISO (Archiving, Validation and Interpolation of Satellite Oceanographic) data (https://www.aviso.altimetry.fr/en/data/products/) are the satellite observational products used in this study. For ocean waves, the AVISO has two satellite altimetry products: along-track data and gridded

- 140 data. The along-track data are used as the observational data source for the wave data in the simulation (sampled from "truth" in the "twin" experiments). The gridded data are used to validate the wave simulation and assimilation (1°×1° resolution with 1-day time intervals). During wave simulation, the significant wave height (SWH) is used as a basic observational variable for data assimilation, which is provided from three ongoing satellites: Jason-2, Jason-3, and Satellite for Argos and ALtiKa (SARAL).
- 145 Figure 1 shows one-cycle ground orbit by taking Jason-2 and SARAL as examples. Jason-3 is the successor of Jason-2, and both satellites share the same orbit.

2.2 Different modeling strategies in WW3 and SWAN

Since the observations are only a sample of real world information, the model bias (i.e., systematic difference between a numerical wave model and the real world) is not well defined against the real

150 world. In this study, we use the systematic difference between the WW3 and SWAN models to simulate the model bias and study the influences on wave data assimilation (WDA). 删除了:. For the first time, WAM creatively overcomes the shortcomings of the first- and second-generation wave models, such as the numerical problems and the restrictions on the spectral shape, and is available for all wind fields and extreme situations. The second model is WaveWatch III (WW3, version 5.16), which is provided online by the National Center for Environmental Prediction (NCEP). In terms of the major aspects such as governing equations, program structures, as well as numerical and physical approaches, WW3 is different from its predecessors (WW2 developed at the NASA Goddard Space Flight Center, and WW1 developed at Delft University of Technology, Netherlands), as WW3 has a more reasonable wind-wave physical mechanism. The third model is Simulating WAves Nearshore (SWAN, version 41.20) provided by Delft University of Technology, Netherlands. SWAN model employs a fully implicit finite differencing scheme. which is unconditionally stable and is more focused on wave propagation processes in shallow water. The model has been successfully applied to simulations of waves in coastal areas, lakes and estuaries, and so on.

删除了:1	25
删除了:1	25
删除了:	

First, let us distinguish the difference in physical and numerical aspects to comprehend the causes of "bias" between these two models. In general, WW3 addresses global scales, and SWAN is more applicable in shallow water. Although the two models have most of the same physical processes, such as the wind input and nonlinear wave-wave interactions, each <u>can provide multiple parameterization</u> scheme<u>s</u> to choose. For example, the nonlinear wave-wave interactions in SWAN include the Discrete Interaction Approximation (DIA) (Hasselmann et al., 1985) and the Webb-Resio-Tracy (Resio and

- 180 Perrie, 1991;Van Vledder, 2006; Webb, 1978), while there are more choices in WW3, such as the Generalized Multiple DIA (Toman, 2004, 2013), the Two-Scale Approximation and Full Boltzmann Integral (Perrie et al., 2013; Perrie and Resio, 2009; Resio et al., 2011; Resio and Perrie, 2008), as well as the Nonlinear Filter scheme (Tolman, 2011). In order to reduce more uncertainties, same scheme is chosen for the same physical process. In numerical aspects, there exist different implementation
- 185 strategies such as the differencing method, which also contributes to bias.

2.3 Two data assimilation systems using WW3 and SWAN

To explore the model bias influences on WDA, we develop two data assimilation systems based on WW3 and SWAN in this study.

Generally, based on the program structure of wave models, we insert the assimilation module between

- 190 calculations of the two-dimensional wave spectrum and outputs of wave parameters so that at the assimilation time, we call on the assimilation module to update the spectrum and SWH. When building the data assimilation systems, we need to consider the different structures of parallelism method, data storage, and information exchange in WW3 and SWAN models as noted in section 2.2.
- To clearly demonstrate the influences of model bias on WDA and minimize its adverse impact, the analysis scheme in both assimilation systems is optimal interpolation (OI), which also is low cost and easy to operate. We implement the OI analysis in three to four steps. The first step uses two Gaussian convolutions of the background and observed SWHs to compute the observational increment of SWH at the observational location. The second step projects the SWH observational increment onto the model grids centered at the observational location but within an impact radius using linear regression.
- 200 The third step transforms the analyzed SWH to update the spectrum of model waves. The fourth step corrects wind forcing using the observational SWH.

Step 1: Computing observational increment by convolution of two Gaussians

删除了: uses

删除了: a different

205 Starting from the idea of the ensemble adjustment Kalman filter (Andersen, 2001), an observational increment at the observational location k, ΔH_k^0 (*H* represents SWH), is computed by the convolution of two Gaussians of the model background and observation, which can usually be obtained from model ensemble members and observational samples. ΔH_k^0 is formulated as follows (Zhang et al, 2007):

$$\Delta H_k^0 = \frac{\frac{1}{(\sigma^M)^2} H^M + \frac{1}{(\sigma^0)^2} H^0}{\frac{1}{(\sigma^M)^2} + \frac{1}{(\sigma^0)^2}} + \frac{\Delta H_k^M}{\left[1 + \left(\frac{\sigma^M}{\sigma^0}\right)^2} - H_k^M.$$
(2)

210 Here, the first and second terms on the right-hand side <u>of the equation</u> adjust the ensemble mean and ensemble spread, respectively, and ΔH_k^M represents the prior model spread. Superscripts *O* and *M* denote the observation and prior quantity estimated by the model, respectively. σ is the corresponding error standard deviation <u>of SWH and varies in time</u>, space and differs in every wave model. The overbar denotes the ensemble mean. In this simplified case, we specify $\sigma^M = 0.6 m$, $\sigma^0 = 0.25 m_a$

215 assuming that σ is constant in all conditions similarly with the previous study (e.g., Qi and Fan, 2013), and use single model and observational values as the ensemble mean. Step 2: Regressing the observational increment onto model grids The second step projects the observational increment ΔH_k^o onto the related model grids using background error covariance, which is a key step in the analysis. To simplify the problem and improve

- 220 the computational efficiency, many studies use a flow-independent distance function to sample the background error covariance for computing the analysis increment at the model grid *i*, ΔH_i^A , as $\Delta H_i^A = (\sigma_i^s)^2 \exp\left(-(\frac{d_{i,k}}{L})\right) \times \Delta H_k^O$. Usually, such an expression is only a symmetrical approximation of the correlation function and cannot represent the spatial structure and propagation characteristics of waves. Here, with the assumption that wave covariance has same structure as wave error covariance, we
- 225 modify the covariance formula to increase its representativeness for wave structure by superimposing a statistical correlation coefficient obtained from the outputted SWH of model simulation into the formula. After analysis, the equation becomes

$$\Delta H_i^A = \frac{\sigma_k^z}{\sigma_k^z} r_{i,k}^z \exp\left(-\left(\frac{d_{i,k}}{L}\right)\right) \times \Delta H_k^O, \qquad \underbrace{if \ d_{i,k} \le R}_{\mathbf{x}}$$
$$\Delta H_i^A = 0, \qquad if \ d_{i,k} > R$$

230 where *L* is the characteristic length and $d_{i,k}$ is the distance between the model grid *i* and observational point *k*. When $d_{i,k}$ is larger than the impact radius *R*, there is no observational impact on this model point from observation *k*. All variables with superscript *s* represent the model statistics from free model

删除了:, 删除了:as in a

删除了: ion

(3)

删除了:, 删除了: 删除了: **设置了格式:** 字体: 倾斜 删除了: 删除了:

control results. For example, $r_{i,k}^s$ is the SWH covariance between the model grid *i* and observation *k*, which is evaluated from the model data time series in corresponding experiments. To ensure the local characteristics of ocean waves, in this study, the characteristics length *L* and impact radius *R* (or the largest $d_{i,k}$) are the same, causing this incremental projection to reach to the e-folding scale. Referring

- 245 to previous studies (e.g., Lionello and Gunther, 1992; Qi and Fan, 2013), we tested different values of L and R as 300 km, 800 km, and 1000 km and found no essential improvement with larger L and R values. Trading-off with computational efficiency, we set L and R as 300 km throughout this study. As shown in Fig. 2, the new covariance represents more wave physics, i.e., the correlation has more asymmetrical and wave-dependent characteristics.
- 250 Step 3: Transforming the SWH to wave spectrum

The assimilation SWH H_i^A is a sum of the prior H_i^M and the analysis increment from step 2 ($H_i^A = H_i^M + \Delta H_i^A$). In the wave model, the form of ocean waves is a two-dimensional wave spectrum that is distributed over frequency and phase. Thus, transforming the assimilation SWH to wave spectrum is necessary to update other wave parameters. Following the previous study (Qi and Fan, 2013), we

assume that the change in wave spectrum is proportional to the energy change that is expressed by the square of SWH. Then, the analyzed spectrum $S_i^A(f, \theta)$ can be written as follows:

$$S_i^A(f,\theta) = \left(\frac{H_i^A}{H_i^M}\right)^2 S_i^M(f,\theta),\tag{4}$$

where f is the wave frequency and θ is the phase direction.

Step 4: Correcting wind forcing using SWH data

260 If the assimilation only adjusts the wave spectrum as described in Step 3, the updated spectral structure may be quickly overwritten by erroneous wind. In this step, we describe a simple scheme using the observed SWH data to correct the wind forcing. Starting from a first guess of wind (the ERA-Interim reanalysis, for instance), the analyzed wind $W_{i,j}^A$ at model grid (i, j) can be written as follows:

$$W_{i,i}^{A} = W_{i,i}^{M} + \Delta W_{i,i}, \tag{5}$$

- 265 where W represents either the u or v component of wind. $\Delta W_{i,j}$ is the corrected wind increment transformed from the updated SWH. While the details of the transformation scheme can be found in Lionello et al. (1992, 1995), we comment on certain aspects relevant to our study. Regardless of boundaries, in general, the energy of ocean waves is determined by the wind speed and duration, which can also be expressed by SWH. In that sense, a function equation can be built, in which the left-hand
- 270 side is an expression of wind speed and duration, while the right-hand side is an expression of SWH,

and they are balanced through wave energy. Then, the analyzed wind speed can be resolved under the assumption that the duration is same in the both prior and analyzed fields.

With respect to the configuration of wave model data assimilation, the model time step is 15 minutes and the assimilation interval is 1 hour. At the assimilation time, we assimilate the along-track

- 275 observations within a 1-hour time window centered at the time. After 10 days, all the observations will cover the global area. The wind data from the reanalysis products (ERA-Interim and NCEP-CFSR in this case) are available every 6 hours. To incorporate the wind correction into the wind forcing of the model, we distribute the wind correction to the adjacent two time-levels of wind data. As the process is looped forward as the wave model state is updated, the wind forcing is adjusted through the SWH 280 assimilation.
 - 2.4 Experimental design

Throughout this study, we use the symbol $MA_{O(s)}^{WF}$ as the name for the assimilation experiment. Here, "*MA*" stands for the "assimilation model" and the subscript "*O*(*s*)" (resp. superscript "*WF*") represents the observing system (resp. wind forcing) in the assimilation. The wind forcing is either the ECMWF

- 285 ERA-Interim (hereafter known as ERAI) or NCEP-CFSR wind (hereafter known as CFSR). Wind forcing can also be corrected by observations of SWH (under this circumstance, the superscript "WF" is replaced by "ASSW"). The observations used in the assimilation could be the model data but are projected on the along-track points of satellite(s) if being used for the twin experiments. Under this circumstance, "O" represents "model that produces observations" and "(s)" represents the used satellite
- 290 tracks (J2-Jason-2, J3-Jason-3, and SA-SARAL, for instance). Otherwise, in the real-data assimilation experiments, the subscript "O(s)" directly lists the satellites that measure the SWH. For example, a symbol named WW3_{SWAND2}^{ERAI} means that the assimilation model is WW3 (here, "MA"="WW3") forced by ERA-Interim wind ("WF"="ERAI"), and the model producing observation is SWAN with Jason-2 satellite track ("O"="SWAN", and "(s)"="(J2)").

295 2.4.1 Twin experiments

Twin experiments refer to a type of Observing System Simulation Experiment (OSSE), in which a model simulation is used to define the "true" solution of a data assimilation problem, and the other model simulation is used to start the assimilation. The "observations" are samples of the "truth" with some white noise to simulate the observational errors. When the "truth" and assimilation are conducted

(设置了格式:英语(美国)

300 by different (resp. identical) models, the framework is a "biased" (resp. "perfect") model twin experiment. Within a twin experiment framework, any aspect of assimilation skills can be measured as the degree to which the "truth" is recovered through the assimilation.
a) Perfect twin experiment

In a perfect twin experiment, we assume that the assimilation model and the observation are unbiased,

305 i.e., both the instrument measuring and numerical modelling processes are sampling the same stochastic dynamical system. Such sampling only has random sampling errors without any systematic difference (bias). We can build this perfect model framework by using the same model to produce the "truth" as the assimilation model but with different initial conditions and wind forcings.

The "observations" from the observational time window (1 hour) centered at the assimilating time can 310 be created by sampling the "truth" SWH with the tracks of the Jason-2, Jason-3 and SARAL satellites, which will cover the global area in 10 days. In this circumstance, if WW3 (resp. SWAN) is used as the assimilation model, the "truth" is produced by the same WW3 (resp. SWAN) model. In the assimilation, we may start the model with different initial conditions and/or wind forcings to examine the influences of initial errors and wind forcing errors on the wave assimilation. Such a perfect twin experiment can

315 be named $WW3_{WW3(s)}^{WF}$ or $SWAN_{SWAN(s)}^{WF}$.

b) Biased twin experiment

To study the impact of model errors on wave assimilation, we use two models to design a "biased" twin experiment. Again, due to the scattering nature of the observations, it is difficult to obtain a complete picture of the model bias against the real world. Given the difference between the WW3 and SWAN

- 320 models described in section 2.2, we use these two models and their assimilation systems here to simulate the model bias and examine its influences on the WDA. We use the ERA-Interim reanalysis wind to force the WW3 (resp. SWAN) to produce the "truth" and "observations" but use the SWAN (resp. WW3) assimilation system to assimilate the "observations." The degree to which the "truth" produced by different model-based assimilation systems is recovered by assimilating the "observations"
- is an assessment of the model bias influences on the WDA. Such a "biased" twin experiment can be named WW3_{SWAN(s)}^{WF} or SWAN_{WW3(s)}^{WF}.
 Under the biased twin experiment framework, we also conduct experiments to examine the impacts of

observing systems on wave assimilations by increasing the observational information based on multiple satellite tracks. For example, we can examine the assimilation results of WW3_{SWAN(J2},^{WF}, 330 WW3_{SWAN(J2+J3})^{WF}, and WW3_{SWAN(J2+J3+S4})^{WF} (resp. SWAN_{WW3(J2})^{WF}, SWAN_{WW3(J2+J3})^{WF}, and

删除了: modeling

 $SWAN_{WW3(J2+J3+SA)}WF$) to understand the impacts of observing systems on different model-based assimilations.

2.4.2 Real-data assimilation experiments

- 335 In this study, we also conduct real-data assimilation experiments using WW3 and SWAN assimilation systems with real track data from the Jason-2, Jason-3 and SARAL satellites. Through real-data assimilation experiments with different model-based assimilation systems, we can 1) increase our understanding of the influences of model errors on the WDA and 2) study the method to reduce the model error influences on the assimilation results. The real-data assimilation experiments can be
- 340 directly named, e.g., $WW3_{J2+J3+SA}^{WF}$ or $SWAN_{J2+J3+SA}^{WF}$.

3 Error sources in wave models and WDA

3.1 Influences of initial and wind forcing errors

Usually, wave numerical simulation can be improved by three methods: 1) reducing the errors in the initial conditions, 2) enhancing the accuracy of the wind forcing, and 3) improving the representation of the wave model and its parameterization.

In this section, we use perfect model twin experiments (as described in section 2.4.1) to exclude model errors and explore the impact of wind forcings and initial conditions on the wave simulations. To compare the performances of the WW3 and SWAN models, we conduct separate experiments with these two models. The "truth" and model control runs are two basic experiments of the perfect twin

- 350 experiment framework. We use the ERA-Interim wind to drive WW3 (resp. SWAN) and generate a long time series of model states as the "truth," which is called *WW3^{ERAI}* (resp. *SWAN^{ERAI}*) for the WW3 (resp. SWAN) perfect model twin experiment. The "observations" are created by interpolating the corresponding "truth" SWH onto the along-track points of satellite orbits. Then, we use the NCEP-CFSR wind to force WW3 (resp. SWAN), called the model control *WW3^{CFSR}* (resp. *SWAN^{CFSR}*), and the
- 355 data assimilation is named WW3_{WW3(s}^{CFSR} (resp. SWAN_{SWAN(s})^{CFSR}). Starting from an independent initial condition produced by the model control, we can conduct the assimilation with the ERA-Interim or NCEP-CFSR wind forcing. The error verification of the assimilation results against the "truth" simulation compared to the error of the model control is an evaluation of the initial error or/and wind forcing error influences on the WDA. All perfect model twin experiments are listed in Table 1.

- 360 First, we conduct two sets of model control experiments WW3^{CFSR} and SWAN^{CFSR} for 80 days (from December 2017 to February 2018). To explore the effect of the initial conditions, we perform the model spin-up for a long time to adequately reach a steady state. Then using the 45th-day model states as the initial conditions, we conduct one more model simulation and data assimilation experiments for each model system as WW3^{ERAI} and WW3_{WW3(2)}^{ERAI} as well as SWAN^{ERAI} and SWAN_{SWAN(2)}^{ERAI}. The root
- 365 mean square errors (RMSEs) of these experiments against the "truth" are shown in Fig. 3 as the red (for WW3^{CFSR}+WW3^{ERAI} and SWAN^{CFSR}+SWAN^{ERAI}) and pink (for WW3^{CFSR} +WW3_{WW3(J2)}^{ERAI} and SWAN^{CFSR}+SWAN_{SWAN(J2)}^{ERAI}) lines.

The SWH RMSE is approximately .34 meters in the WW3 or SWAN model control with the NCEP-CFSR wind. Once the wind forcing is changed to the "perfect" wind (the ERA-Interim) on the 45th day,

- 370 the RMSE quickly drops and is close to zero after approximately 10 days, and the SWAN model takes longer to accomplish this change than the WW3. If data assimilation is added, the RMSE reduces much faster than the model controls (roughly half of the time scale of the correct wind forcing). From the analyses above, we learned, 1) in wave models, the wind forcing plays an important role and an incorrect wind forcing could be a significant error source of WDA; and 2) the WDA can rapidly reduce
- 375 the initial error and improve the predictability of a wave model even when it is forced by an <u>in</u>accurate wind forcing.

3.2 Impact of the observational system

- In this section, using the same model states (at the 45th day) in the corresponding model control as in section 3.1 as the initial conditions, we conduct two sets of assimilation experiments WW3_{WW3(J2})^{CFSR}, 380 WW3_{WW3(J2+J3})^{CFSR}, WW3_{WW3(J2+J3+S4})^{CFSR} and SWAN_{SW4N(J2})^{CFSR}, SWAN_{SW4N(J2+J3+S4})^{CFSR}, SWAN_{SW4N(J2+J3+S4})^{CFSR}, SWAN_{SW4N(J2+J3+S4})^{CFSR}, SWAN_{SW4N(J2+J3+S4})^{CFSR}, Through examining the assimilation quality with one satellite (Jason-2), two satellites (Jason-2+Jason-3) and three satellites (Jason-2+Jason-3+SARAL), we attempt to understand the impact of improving the observing system on the WDA, considering the NCEP-CFSR wind forcing errors against the ECMWF ERA-Interim based on a perfect assimilation model. The RMSEs of all the
- 385 above assimilation experiments are plotted in Fig. 3 as the blue (assimilating Jason-2 only), green (assimilating Jason-2+Jason-3) and cyan (assimilating Jason-2+Jason-3+SARAL) lines. From Fig. 3, we can see that in both models, the assimilation errors are reduced when more observational information is used. The corresponding RMSE reductions in these three experiments from the model control run are 24 %, 32 % and 38 % for WW3 and 26 %, 35 % and 38 % for SWAN,

390 respectively. However, when more satellite observations are assimilated into the model, the magnitude of improvement becomes small (further reduced by 8 % in WW3 and 9 % in SWAN when Jason-3 is added as well as only a 6 % in WW3 and 3 % in SWAN further reduction when SARAL is further added). These results suggest that given wind forcing errors, increasing observational information can help to improve the model behavior, but the improvement is limited.

395 3.3 Adverse impact of model bias

As described in section 2.2, the WW3 and SWAN models discretize the wave action governing equation with different physical processes, parameterization schemes and differencing schemes. These differences result in each wave model having its own distinguished characteristics. To study the adverse impact of the model bias on the wave assimilation, the biased twin experiments described in

- 400 section 2.4.1 are used in this section, where the "truth" model and the assimilation model are different between WW3 and SWAN. For example, the *WW3_{SWAN(J2)}^{ERAI}* (resp. *SWAN_{WW3(J2)}^{ERAI}*) experiment uses WW3 (resp. SWAN) as the assimilation model to assimilate the Jason-2 track point "observations," but the observed values are produced by SWAN (resp. WW3), and all models are forced by the ERA-Interim wind. All related experiments for the biased model framework are described in detail in **Table**
- 405 **2**.

The RMSEs and correlation coefficients produced by the all biased model assimilation experiments are plotted in **Fig. 4**. The black line in each panel represents the result of the WW3 model control forced by the ERA-Interim wind ($WW3^{ERAI}$) against the "truth" simulation by the SWAN model with the same wind forcing ($SWAN^{ERAI}$) (left panels, a and b) (vice versa in the right panels, c and d). Both the

- 410 WW3^{ERAI} and SWAN^{ERAI} experiments are initialized from a cold start by the wind and integrated for 80 days, and the results of the last 40 days are shown in Fig. 4. It is clear that the WW3 and SWAN model simulations are quite different even though both simulations use identical forcings and start from identical initial conditions. The RMSEs of the two model simulations are both 0.58 m, which is much larger than the errors produced by a perfect model but with different wind (~0.34 m, see <u>black lines in</u>
- 415 Fig. 3).

Compared with the model controls $WW3^{ERAI}$ and $SWAN^{ERAI}$, the assimilation experiments $WW3_{SWAN(22)}^{ERAI}$ and $SWAN_{WW3(22)}^{ERAI}$ (pink lines in Fig. 4) can significantly reduce the SWH simulation error (by 24 % and 22 %, respectively) and enhance the correlations (by 3 % and 4 %, respectively) with the "truth" ($SWAN^{ERAI}$ and $WW3^{ERAI}$, respectively). When the "observations" of Jason-3 and

- 420 SARAL are added to the assimilation (i.e., WW3_{SWAN(J2+J3)}^{ERAI} and WW3_{SWAN(J2+J3+SA)}^{ERAI} as well as SWAN_{WW3(J2+J3)}^{ERAI} and SWAN_{WW3(J2+J3+SA)}^{ERAI}) (see the red and blue lines, respectively), the model SWH error (resp. correlation) is further reduced (resp. enhanced), but the amplitude of reduction (resp. enhancement) gradually diminishes (10 % and 5 % for further error reduction and 1 % and 0.8 % for further correlation enhancement in the WW3 assimilation; 10 % and 7 % for further error reduction and
- 425 1.7 % and 0.7 % for further correlation enhancement in SWAN assimilation from the additions of Jason-3 and SARAL, respectively).

The results of two other sets of assimilation experiments called $WW3_{SWAN(J2+J3+S4)}^{ASSW}$ and $SWAN_{WW3(J2+J3+S4)}^{ASSW}$ are also plotted by dotted green lines in **Fig. 4.** The superscript "ASSW" stands for the assimilation-corrected wind, meaning that the wind forcing of the assimilation model is also

430 "corrected" by the "observed" SWH data, as described in Step 4 of section 2.3. We found that in <u>both</u> assimilation systems, using the "observed" SWH data to "correct" the wind can compensate for the model errors to some degree and further reduce the assimilation errors, but the improvement is very limited. The weak improvement could be attributed to the simple wind correction method with total wave height. In the future, a more powerful correction method with windsea wave height may have

435 better performance.

These assimilation results clearly show that even though the wind forcing is perfect, once a biased assimilation model is used, the wave simulation has large errors. Although WDA can greatly reduce the simulation error by assimilating the observational information into the model, due to the existence of the model bias, the error remains at some significant level and cannot be eliminated entirely even by

440 increasing the observational constraints through an improvement in the observational system and constraint of the wind forcing.

3.4 Comparison of the influence of wind forcing with model bias

Comparing the time series of SWH RMSE caused by two important error sources, model bias (0.58 m shown with black lines of panel a/c in Fig. 4) plays a stronger role than wind forcing (0.34 m shown

445 with black lines in Fig. 3). When three satellite observations (Jason-2, Jason-3 and SARAL) are assimilated into twin experiments, the RMSE of SWH reduces 38% in perfect model twin experiments (cyan lines in Fig. 3) and 40% in biased model twin experiments (blue lines of panel a/c in Fig. 4). It is obvious that the error caused by model bias is bigger than by wind forcing and their improvements are almost similar after data assimilation.

删除了: the WW3 model

删除了: In the SWAN model, such wind "correction" cannot compensate for the model error at all....

制除了: Next, with the results of real-data assimilation where both the model and wind forcing have errors, we will analyze and discuss how to mitigate the model bias influences on the WDA
· 设置了格式: 字体: (中文) Times New Roman
• 设置了格式: 字体: (中文) Times New Roman, 英语(英国)
• 设置了格式: 字体: (中文) Times New Roman, 英语(英国)
· (设置了格式: 字体: (中文) Times New Roman
* 带格式的: 标题 2, 右 0.56 字符
• 设置了格式: 字体: (中文) Times New Roman, 英语(英国)
设置了格式: 字体: (中文) Times New Roman, 英语(英国)
删除了: more
设置了格式: 字体颜色: 文字1
删除了:
· 设置了格式: 字体: 加粗
设置了格式: 字体: 加粗

1	A spatial pattern of SWH RMSE is also displayed in Fig. 5 (Due to the similar performance inside twin	设置了格式: 字体: 加粗
	$\underline{\text{experiments}},$ here we only show the results using WW3 model as simulation/assimilation model). In	
460	model control run, it makes sense that the error caused by wind forcing (panel a ($WW3^{CFSR}$). Truth is	
	$\underline{WW3^{ERAI}}$ has distributed in the areas where the wind is strong, such as the area of Antarctic	
	$\underline{\text{Circumpolar Current (ACC) and the north Pacific and Atlantic Ocean, while the error caused by model}$	
	bias (panel c ($WW3^{ERAI}$). Truth is $SWAN^{ERAI}$) is distributed almost globally. After assimilating with	
	three satellite observations, the error spatial distribution of both has improved a lot respectively (panel	
465	b (WW3 <u>WW3(J2+J3+SA)</u> ^{CFSR}), and panel d (WW3 <u>SWAN(J2+J3+SA)</u> ^{ERAI})). Referring to wind distribution, we can	
	divide the global areas roughly into three parts: northern westerly zone (65° N-30° N), trade-wind zone	
1	(30° N-30° S), and circumpolar westerly zone (30° S-65° S). Therefore the error caused by wind	
	forcing (resp. model bias), the decreasing percentages of SWH RMSE in these three areas are 30%	
	(resp. 27%), 45% (resp. 50%), and 46% (resp. 48%), respectively. We can find that the improvement of	
470	both error sources has similar performance in three these areas, weak in the northern westerly zone and	
	almost same strong in the trade-wind zone and circumpolar westerly zone. About the reason why there	
	is a lower improvement in the north Pacific and Atlantic Ocean should be explored in the future. To	
	sum up, the error caused by model bias has larger than wind forcing in the global area generally,	
	especially in the equatorial ocean. However, in the north of Atlantic Ocean, wind forcing has a stronger	
475	impact. After data assimilation, both have improved greatly and have similar spatial pattern (i.e. the	
1	bigger error is still distributed in high latitude). However, their error gap is still existed	设置了格式
	It's worth to mention, that there is a similar performance about the effect of observation system on	删除了: should be
	improving both error sources, whether in time series or spatial distribution. The more observation	删除了: ed
	information absorbed into assimilation systems, the better error improvement in both twin experiments.	
480	However, due to the existence of model bias, this improvement has a limitation and stays at some	
	certain level. If more powerful observation is absorbed (such as: wave direction, wave period, and two-	
	dimensional wave spectra), the limitation maybe stay at a lower level. Next, with the results of real-	
	data assimilation where both the model and wind forcing have errors, we will analyze and discuss how	
1	to mitigate the model bias influences on the WDA	设置了格式: (中文)中文(中国),(其他)英语(美国)

4 Mitigation of model bias influences on wave assimilation

4.1 Bias characteristics of WW3 and SWAN data assimilations

From the above analyses of twin experiment results, we learned that the model bias has a strong 490 adverse impact on WDA. To explore the method of mitigating the model bias influence on the WDA, we conduct the real-data assimilation experiments (same time range as the "twin" experiments) described in section 2.4.2 using the WW3 and SWAN assimilation systems to assimilate the track data of Jason-2 and Jason-2+Jason-3+SARAL. To ensure the performance of the biased model WDA, a longer assimilation (more than two months) is conducted (a total of 70 days). The spatial distributions

495 of the SWH errors (obtained from the difference against the merged gridded AVISO observations over the last 30 days out of 80 days) are shown in Fig. 6. The left (resp. right) panels are for the WW3 (resp. SWAN) simulation and assimilations: WW3^{ERAI}, WW3_{J2}^{ERAI}, and WW3_{J2+J3+SA}^{ERAI} (resp. SWAN^{ERAI}, SWANJ2^{ERAI}, and SWANJ2+J3+SA^{ERAI}).

Comparing panel a with panel d in Fig. 6 reveals that a large difference exists in the simulations of the

- 500 two models. First, the SWAN simulation errors are generally larger than the WW3 simulation errors. Second, the global error distributions are quite different: while the WW3 simulation errors appear negative (resp. positive) over most of the 30° S north (resp. south) area, the SWAN simulation errors appear positive in most of the tropical oceans and negative in the middle latitudes. Both simulations show large errors in the southern ocean coastal area, but over the Antarctic Circumpolar Current area,
- 505 the WW3 error is positive, and the SWAN error is negative. It is interesting that under same wind conditions, two wave models have converse performances. In the future study, it is urgent to find out the detailed approach of two wave models in this area.

The above systematic differences between the two model simulations have significant influences on the results of the WDA. In general, the distribution of assimilation errors shares the same patterns as the

- 510 model simulation errors but with a much smaller magnitude. The net result is that both the WW3 negative (resp. positive) error magnitude over the 30°S north (resp. south) area and the SWAN error magnitude as (+)(-)(+)(-) from south to north are dramatically reduced by the Jason-2 data assimilation (comparing the middle panels with the upper panels), and on this basis, incorporating more observations from Jason-3 and SARAL into the assimilation process, both model error magnitudes are
- 515 further reduced to some degree (comparing the bottom panels with the middle panels). From the corresponding RMSE distributions (it's not shown here), we learned that the large RMSEs mainly

删除了: verified

删除了:5

删除了:5

删除了: Large model errors usually occur in strong wind areas. Although the same parametric scheme of wind input expressed by exponential growth with friction velocity can be activated, the process transforming wind speed provided by users to friction velocity is necessary but different in both models, which may be an important reason why the two models have different performances under the same wind conditions.

删除了: Fig. 6

appear in places where the model bias (time mean error) is large. This finding means that the model bias has a largely adverse impact on the WDA.

- 530 Figure 7 displays the time series of the RMSEs and spatial_correlation coefficients with the global statistics in space. The RMSE (resp. correlation coefficient) of the SWAN model simulation is larger (resp. smaller) than the WW3 model simulation (.66 m RMSE and .806 correlation for SWAN versus .61 m RMSE and .876 correlation for WW3). In the WW3 and SWAN assimilations with the Jason-2 data, the RMSEs are reduced by 8 % and 11 %, respectively, and the time mean correlations
- 535 are enhanced by roughly 1 % and 5 %, respectively. If the data of all three satellites, Jason-2, Jason-3 and SARAL, are assimilated, the RMSEs are reduced by 11 % and 17 % in the WW3 and SWAN assimilations, respectively, and the correlations are enhanced by approximately 2 % and 8 %, respectively. In these real-data assimilation cases, for each model assimilation, both the model and wind forcing have errors. Under this circumstance, the assimilation where the SWH observations are
- 540 used to adjust the model spectrum, the model wind forcing is also corrected and can further reduce the assimilation errors (red lines in Fig. 7). The red lines in Fig. 7 represent the best result of the assimilation given the WW3 and SWAN model biases, which makes full use of the observations from all three satellites to adjust both the model spectrum and wind forcing. Next, we will discuss how to use the results of two assimilation systems to mitigate the wave analysis error.

545 4.2 Mitigation of WDA errors

The mitigation of model bias is a complex issue in which improving the model is a final but longlasting solution. From **Fig.** (), we learned that the WW3 and SWAN assimilation errors have some common (or opposite) characteristics in some locations. For example, while the SWH over the southern ocean coastal area always appears to be overestimated because of the lack of adequate observations to

550 improve in both assimilation systems, the WW3 and SWAN assimilations appear to be the opposite in the Antarctic Circumpolar Current area and the tropical oceans. The WW3 (resp. SWAN) assimilation errors in the Antarctic Circumpolar Current area appear positive (resp. negative), while the WW3 (resp. SWAN) assimilation errors in the tropical oceans appear negative (resp. positive). A question arises: as the first step of mitigating model bias influences on the WDA, can we use a pair of assimilation

555 systems to explore a statistical approach to reduce the wave assimilation errors? Given the opposite behaviors of two assimilation systems existing in certain places, the simplest bias correction could, be conducted by simple average. This assumes that <u>each wave model assimilation</u> (删除了:5

删除了: an

560 system has its own characteristics of systemic error due to deficit physics, and these systemic errors (or model biases) from different wave models follow a Gaussian distribution with trivial expectation. The corresponding results are shown in Fig. 8. Compared with the performance of each individual assimilation system (dashed red lines for WW3 and dotted red lines for SWAN), the results of this bias correction (cyan lines) show that the RMSE is reduced (the left panel) but the spatial correlation is not

- 565 greatly improved (the right panel). It is reasonable that based on the opposite errors deviating from the real world in two assimilation systems, this correction method employing the mathematical average can reduce the RMSE to some extent, but it may not have a significant contribution to improving the correlation coefficient if either the sampling size of model bias is too small (only 2 in this case) or the bias has an asymmetric distribution.
- 570 Considering the potentially asymmetry, of Gaussian distribution of different model biases and small sampling size in practice, as the first step, we calculate the spatial distribution of model bias in every assimilation system, (the time mean of the difference between observations and assimilation results) (Fig. 6) and extract it at each time step in the output, and then calculate the expectation (average) of all assimilation systems as the results after bias correction. The corresponding results are shown with pink
- 575 lines in **Fig. 8**. Both the RMSE and correlation are improved greatly. We also show the spatial distribution of SWH RMSEs after bias correction with the second method. From **Fig. 9**, we can easily find that after bias correction, all the experiments have similar error performances, the smaller in low latitude and the larger in high latitude. As the increasing of satellite observation (one in panel bf, and three in panel cg), the error decreases gradually referred to model control run (panel ae). Based on the
- 580 best assimilation results currently (panel cg), the improvement with a corrected wind is more efficient in WW3 (panel d), especially in high latitude which is the higher error area, but is hardly to find in SWAN (panel h). If we focus on the change without (panel c) and with (panel d) the corrected wind in WW3 model, the RMSE improvement of windsea and swell is 27% and 1% averaged in global. If we divide global area into three parts mentioned as section 3.4, the windsea (resp. swell) improvement of
- 585 regional RMSE mean is 4% (resp. 3%), 3% (resp. 1%), and 6% (resp. 1%) in northern westerly zone (65° N-30° N), trade-wind zone (30° N-30° S), and circumpolar westerly zone (30° S-65° S). It is obvious that the improvement of windsea is better than swell and wind correction has a stronger impact in windsea wave assimilation at high latitude with strong wind. Clearly, this bias correction with physical consideration is more effective to improve the quality of WDA, but it uses observational

删除了: bias

删除了: itself has

1	删际 J:IC
(删除了: (i.e., each wave model has its own characteristics of systematic error due to deficit physics) and small sampling size in
(删除了:
-(设置了格式: 字体颜色: 文字 1
ľ	删除了: i.e., this spatial distribution could be obtained through averaging
Ì	删除了: model
Ì	删除了: long-term mean of the error
Ì	删除了: . In this paper, the corresponding spatial patterns are shown in Fig. 6) for each assimilation system
Ì	删除了: first
(设置了格式: 字体颜色: 绿色
(设置了格式: 字体颜色: 绿色
(设置了格式: 字体: 加粗, 字体颜色: 绿色
(设置了格式: 字体颜色: 绿色
(删除了: for each assimilation system
Ì	

information one more time, while the first method of bias correction processes assimilation results 605 directly without further uses of observational information.

To verify the feasibility and applicability of the bias correction method above, 3 well-known wave models (WW3, SWAN and WAM) with the same data assimilation method are used to conduct longer assimilation and bias correction experiments. The calculation period lasts for 14 months (from November 2016 to December 2017) with sufficient spin-up process to reach a steady assimilation state

- 610 (the 1st month for model spin-up and the 2nd month for assimilation spin-up). The results of the last 12 months (for 2017) are analyzed and presented in Fig. 10, and Fig. 11, for the spatial distributions and time series of RMSEs and spatial correlation coefficients, respectively. From Fig. 10, we can see that both the RMSE and correlation coefficient (panels d and h, respectively) have been improved by the bias correction that combines the advantages of every WDA system (panels a and e for WW3, panels b
- 615 and f for SWAN, panels c and g for WAM). In Fig. 11, the bias correction of model control runs (the second bias correction method is conducted based on the result of model control run from three wave models) shows improvement but is worse than the data assimilation before bias correction (compare green with pink). Compared with the model control (blue), the assimilation results with bias correction (red) can reduce the error by 25 % and significantly enhance the correlation coefficient (from 0.88 to
- 620 0.923). This result confirms that this bias correction based on multiple assimilation systems can effectively enhance the WDA quality.

5 Summary and discussion

Ocean waves cause the sea surface roughness to impact the boundary conditions of the atmosphere and the wind stress of the ocean surface. Wave processes, such as wave-breaking, wave-induced bottom

- 625 stress and so on, have significant effects on ocean mixing. Thus, ocean waves are important physical processes for understanding ocean mixing and air-sea interactions in coupled Earth systems. More accurately predicting ocean waves is of great societal significance. However, multiple error sources exist in wave simulations and predictions, including <u>modelling errors</u>, wind forcing errors and initial condition errors.
- 630 To sort out the source of the errors of wave data assimilation (WDA), a pair of independent WDA systems is first developed using two wave models: Wave Watch III (WW3) and Simulating WAves Nearshore (SWAN). The perfect and biased model "twin" experiment frameworks are designed to

删除了:9		
删除了:0		
删除了:9		

删除了:0

删除了: modeling

clearly identify each error source and examine its influences on WDA. The results show that model bias is a significant error source that has a largely adverse impact. Then, two WDA systems are used to

- 640 design bias correction approaches to mitigate the influences of model bias and improve the assimilation quality. Finally, long-term WDA experiments added by the third WDA system with the WAM model (WAve Modeling) (WW3, SWAN and WAM) are conducted to validate the bias correction method. Three findings are established: 1) When the model is perfect, the initial condition error decays within 10 days, but the WDA can shorten the time scale by half. 2) When the model is biased, despite a
- 645 perfect wind forcing, the wave simulation has large errors and the WDA can only reduce the error to a limited extent. 3) With the results from two assimilation systems, a statistical approach of bias correction significantly improves the quality of final wave analysis by combining the merits from individual assimilation systems.
- Model bias is an obstacle to improving WDA and wave predictions. Using multiple assimilation 650 systems to study the influences of model bias on WDA is an effective approach. As the first step, however, we have used a simple assimilation scheme and simple bias correction method. In follow-up studies, we shall consider <u>powerful observation information (such as: 2-dimensional wave spectra)</u>, advanced assimilation schemes <u>(such as ensemble Kalman Filter)</u>, and more comprehensive correction methods to help improve <u>modelling</u>. For example, the "online" bias correction (the "offline" bias
- 655 correction is used in this paper) during the assimilation process (e.g., Dee, 2005) will be considered to improve the assimilation results and correct the instantons initial condition within individual assimilation systems, after that, a robust observation (such as buoy) is needed to validation the quality of bias correction. In addition, improving the model is an important, inevitable and long-lasting task. In this study, <u>under same wind condition</u>, we find that three models show common bias characteristics in
- 660 the Antarctic Circumpolar Current (ACC) area. If there is a similar performance forcing by other wind, this may suggest that present wave modelling may have deficits in energy spectrum expression for high wind speed areas. In the future, we will further examine the sensitivities of physical processes on high wind speed to mitigate such common modelling bias. All in all, a robust bias correction method with lower model bias and higher representation of wave physical characteristics may further improve wave
- 665 analysis quality. Once a long time series of high-quality wave analyses is available, it is expected that we can improve our understanding of ocean mixing. The physical process of wave-induced mixing is linked with the structure of the ocean mixing layer (Qiao et al., 2010). This process can be expressed as a function of wave number, frequency and wave spectrum and so on, provided by wave analysis. With

删除了: the correct wind forcing can overwrite the initial condition error within a 10-day time scale, but the

删除了: modeling

-(删除了:T
~(删除了: finding
\mathcal{I}	删除了:s
)(删除了: modeling
(删除了: modeling

the framework of multiple WDA systems developed in this study, improved wave predictions can be effectively pursued. How can we further enhance the predictability of ocean waves? The first important step is to understand the physical process of ocean waves better based on a more accurate evolution of 680 wave state from this framework. Answering these questions could be very important and interesting

Code and data availability

Codes, data and scripts used to run the models and produce the figures in this work are available on the Zenodo site: doi:10.5281/zenodo.3445580 or by sending a written request to the corresponding author

685 (Shaoqing Zhang, szhang@ouc.edu.cn).

research topics for the future studies.

Author contribution

Jiangyu Li coded the module of data assimilation and inserted it into the wave models, carried out all the experiments. Shaoqing Zhang designed the experiments, analysed the results with constructive discussions. Both authors wrote the paper together.

690 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

The research is supported by the National Key R&D Program of China (2017YFC1404100 and 2017YFC1404102), the National Natural Science Foundation of China (Grant No. 41775100, 41830964), and the Fundamental Research Funds for the Central Universities (201961001).

删除了: and

References

695

Abdalla, S., Bidlot, J., and Janssen, P.: Assimilation of ERS and ENVISAT wave data at ECMWF, Envisat & Ers Symposium, 572, 2013.

700 Almeida, S., Rusu, L., and Guedes Soares, C.: Data assimilation with the ensemble Kalman filter in a high-resolution wave forecasting model for coastal areas, Journal of Operational Oceanography, 9(2), 103-114, 2016.

Anderson, J. L.: An Ensemble Adjustment Kalman Filter for Data Assimilation, Monthly Weather Review, 129(12), 2884-2903, 2011.

- 705 Babanin, A. V., Ganopolski, A., and Phillips, W. R. C.: Wave-induced upper-ocean mixing in a climate model of intermediate complexity, Ocean Modelling, 29(3), 189-197, 2009. Bauer, E., Hasselmann, K., Young, I. R., and Hasselmann, S.: Assimilation of wave data into the wave model WAM using an impulse response function method, Journal of Geophysical Research, 101(C2), 3801-3816, 1996.
- 710 Bhatt, V., Kumar, R., Basu, S., and Agarwal, V. K.: Assimilation of altimeter significant wave height into a third-generation global spectral wave model, IEEE Transactions on Geoscience and Remote Sensing, 43(1), 110-117, 2005.

Booij, N., Ris, R., and Holthuijsen, L.: A third-generation wave model for coastal regions: 1. Model description and validation, Journal of Geophysical Research, 104(C4), 7649-7666, 1999.

715 Breivik, L. A., Reistad, M., Schyberg, H., Sunde J., Krogstad H. E., and Johnsen, H.: Assimilation of ERS SAR wave spectra in an operational wave model, Journal of Geophysical Research, 103(C4), 7887-7900, 1998.

 Chen, S. S., Price J. F., Zhao, W., Donelan, M. A., and Walsh, E. J.: The CBLAST-Hurricane Program and the Next-Generation Fully Coupled Atmosphere-Wave-Ocean Models for Hurricane Research and Prediction, Bulletin of the American Meteorological Society, 88(3), 311-317, 2007.

- Daniel, T., Manley, J., and Trenaman, N.: The Wave Glider: enabling a new approach to persistent ocean observation and research, Ocean Dynamics, 61(10), 1509-1520, 2011. Dee, D.: Bias and data assimilation, Quarterly Journal of the Royal Meteorological Society, 131(613),
- 725 Doyle, J. D.: Coupled Atmosphere-Ocean Wave Simulations under High Wind Conditions, Monthly Weather Review, 130(12), 3087-3099, 2002.

3323-3343, 2005.

Emmanouil, G., Galanis, G., and Kallos, G.: Combination of statistical Kalman filters and data assimilation for improving ocean waves analysis and forecasting, Ocean Modelling, 59-60(12), 11-23, 2012.

- Forg, H., Vandemark, D., Quilfen, Y., Chapron, B., and Beckley, B.: Assessment of wind-forcing impact on a global wind-wave model using the TOPEX altimeter, Ocean Engineering, 33(11-12), 1431-1461, 2006.
- 735 Folley, M. and Whittaker, T. J. T.: Analysis of the nearshore wave energy resource, Renewable Energy, 34(7), 1709-1715, 2009.

Francis, P. E. and Stratton, R. A.: Some experiments to investigate the assimilation of SEASAT altimeter wave height data into a global wave model, The Quarterly Journal of the Royal Meteorological Society, 116(495), 1225-1251, 1990.

740 Gommenginger, C. P., Strokosz, M. A., Challenor, P. G., and Cotton, P. D.: Measuring ocean wave period with satellite altimeters: A simple empirical model, Geophysical Research Letters, 30(22), 2150, 2003.

Greenslade, D. J. M.: The assimilation of ERS-2 significant wave height data in the Australian region, Journal of Marine Systems, 28, 141-160, 2001.

- 745 Greenslade, D. J. M. and Young, I. R.: Background errors in a global wave model determined from altimeter data, Journal of Geophysical Research, 109(C9), doi: 10.1029/2004JC002324, 2004.
 Hasselmann, S., Hasselmann, K., Allender, J. H., and Barnett, T. P.: Computations and parameterizations of the nonlinear energy transfer in a gravity wave spectrum. Part II: Parameterizations of the nonlinear transfer for application in wave models, Journal of Physical 750 Oceanography, 15(11), 1378-1392, 1985.
- Hasselmann, S., Lionello, P., and Hasselmann, K.: An optimal interpolation scheme for the assimilation of spectral wave data, Journal of Geophysical Research, 102(C7), 15823-15836, 1997. Heras, M. M. D., Burgers, G., and Janssen, P. A. E. M.: Variational Wave Data Assimilation in a Third-Generation Wave Model, Journal of Atmospheric and Oceanic Technology, 11(5), 1350-1369,
- 755 1994.

Hisaki, Y.: Ocean wave directional spectra estimation from an HF ocean radar with a single antenna array: Observation, Journal of Geophysical Research, 110(C11004), doi: 10.1029/2005JC002881, 2005. Huang, C. J. and Qiao, F.: Wave-turbulence interaction and its induced mixing in the upper ocean, Journal of Geophysical Research, 115(C04026), doi: 10.1029/2009JC005853, 2010. 760 Janssen, P. A. E. M., Lionello, P., Reistad, M., and Hollingsworth, A.: Hindcasts and data assimilation studies with the WAM model during the Seasat period, Journal of Geophysical Research, 94(C1), 973:993, 1989.

Jesus, P. Y. and Cavaleri, L.: On the specification of background errors for wave data assimilation systems, Journal of Geophysical Research: Oceans, 121(3), 209-223, 2015.

- 765 Lionello, P., Gunther, H., and Hansen, B.: A sequential assimilation scheme applied to global wave analysis and prediction, Journal of Marine Systems, 6(1-2), 87-107, 1995. Lionello, P., Gunther, H., and Janssen, P. A. E. M.: Assimilation of altimeter data in a global third-generation wave model, Journal of Geophysical Research, 97(C9), 14453-14474, 1992. Liu, B., Liu, H. Q., Xie, L., Guan, C. L., and Zhao, D. L.: A Coupled Atmosphere-Wave-Ocean
- Modeling System: Simulation of the Intensity of an Idealized Tropical Cyclone, Monthly Weather Review, 139(1), 132-152, 2011.

Longuet-Higgins, M. S., and Stewart, R. W.: Radiation stress and mass transport in gravity waves, with application to 'surf beats', Journal of Fluid Mechanics, 13(4), 481-504, 1962.

Longuet-Higgins, M. S., and Stewart, R. W.: The changes in amplitude of short gravity waves on 775 steady non-uniform currents, Journal of Fluid Mechanics Digital Archive, 10(04), 21, 1961.

Lzaguirre, C., Mendez, F. J., Menendez, M., and Losada, I. J.: Global extreme wave height variability based on satellite data, Geophysical Research Letters, 38(10), 415-421, 2011.

Mitsuyasu, H., Tasai, F., Suhara, T., Mizuno, S., Ohkusu, M., Honda, T., and Rikiishi, K.: Observation of the power spectrum of ocean waves using a cloverleaf buoy, Journal of Physical Oceanography, 10, 286-296, 1980.

Perrie, W. and Resio, D. T.: A two-scale approximation for efficient representation of nonlinear energy transfers in a wind wave spectrum. Part II: Application to observed wave spectra, Journal of Physical Oceanography, 39, 2451–2476, 2009.

Perrie, W., Toulany, B., Resio, D. T., Roland, A. and Cuclair, J.: A two-scale approximation for wave-785 wave interactions in an operational wave model, Ocean Modelling, 70, 38-51, 2013.

Qi, P. and Cao, L.: The Assimilation of Jason-2 Significant Wave Height Data in the North Indian Ocean Using the Ensemble Optimal Interpolation, IEEE Transactions on Geoscience and Remote Sensing, 54(1), 287-297, 2016.

Qi, P. and Fan, X.: The impact of assimilation of altimeter wave data on wave forecast model in the 790 north Indian Ocean, Marine Forecasts, 30(4), 70-78, 2013. Qiao, F., Yuan, Y., Ezer, T., Xia, C., Yang, Y., Lv, X., and Song, Z.: A three-dimensional surface wave-ocean circulation coupled model and its initial testing, Ocean Dynamics, 60(5), 1339-1355, 2010. Qiao, F., Yuan, Y., Yang, Y., Zheng, Q., Xia, C., and Ma, J.: Wave-induced mixing in the upper ocean: Distribution and application to a global ocean circulation model, Geophysical Research Letters, 31(11), 202 217 2004

795 293-317, 2004.

Queffeulou, P.: Long-Term Validation of Wave Height Measurements from Altimeters, Marine Geodesy, 27(3-4), 495-510, 2004.

Rapizo, H., Babanin, A. V., Schulz, E., Hemer, M. A., and Durrant, T. H.: Observation of wind-waves from a moored buoy in the Southern Ocean, Ocean Dynamics, 65(9-10), 1275-1288, 2015.

800 Resio, D. T., Long, C. E., and Perrie, W.: The Role of Nonlinear Momentum Fluxes on the Evolution of Directional Wind-Wave Spectra, Journal of Physical Oceanography, 41(4), 781-801, 2011. Resio, D. T. and Perrie, W.: A numerical study of nonlinear energy fluxes due to wave-wave interactions. Part I: Methodology and basic results, Journal of Fluid Mechanics, 223, 603-629, 1991. Resio, D. T. and Perrie, W.: A Two-Scale Approximation for Efficient Representation of Nonlinear

805 Energy Transfers in a Wind Wave Spectrum. Part I: Theoretical Development, Journal of Physical Oceanography, 38(12), 2801-2816, 2008.

Rusu, L.: Assessment of the Wave Energy in the Black Sea Based on a 15-Year Hindcast with Data Assimilation, Energies, 8, 10370-10388, 2015.

Sannasiraj, S. A., Babovic, V., and Chan, E. S.: Wave data assimilation using ensemble error covariances for operational wave forecast, Ocean Modelling, 14(1), 102-121, 2006.

- Stopa, J. E. and Cheung, K. F.: Intercomparison of wind and wave data from the ECMWF Reanalysis
 Interim and the NCEP Climate Forecast System Reanalysis, Ocean Modelling, 75(2), 65-83, 2014.
 Sun, M., Yin, X., Yang, Y., Wu, K., and Sun, B.: On EAKF data assimilation based on MASNUM-WAM, Oceanologia Et Limnologia Sinica, 48(2), 210-220, 2017.
- 815 Tolman, H. L.: A conservative nonlinear filter for the high-frequency range of wind wave spectra, Ocean Modelling, 39(3), 291-300, 2011.

Tolman, H. L.: A Generalized Multiple Discrete Interaction Approximation for resonant four-wave interactions in wind wave models, Ocean Modelling, 70, 11-24, 2013.

Tolman, H. L.: A Third-Generation Model for Wind Waves on Slowly Varying, Unsteady, and Inhomogeneous Depths and Currents, Journal of Physical Oceanography, 21(6), 782-797, 1991. Tolman, H. L.: Inverse modeling of discrete interaction approximations for nonlinear interactions in wind waves, Ocean Modelling, 6(3), 405-422, 2004.

Van Vledder, G.: The WRT method for the computation of non-linear four-wave interactions in discrete spectral wave models, Coastal Engineering, 53, 223-242, 2006.

825 Voorrips, A. C.: Spectral wave data assimilation for the prediction of waves in the North Sea, Coastal Engineering, 37, 455-469, 1999.

Voorrips, A. C., Heemink, A. W., and Komen, G. J.: Wave data assimilation with the Kalman filter, Journal of Marine Systems, 19(4), 267-291, 1999.

Walsh, E. J., Hancock, D. W., and Hines, D. E.: An Observation of the Directional Wave Spectrum

830 Evolution from Shoreline to Fully Developed, Journal of Physical Oceanography, 19(5), 670-690, 1989. WAMDI Group: The WAM Model-A Third Generation Ocean Wave Prediction Model, Journal of Physical Oceanography, 18(12), 1775-1810, 1988.

Wang, Y. and Yu, Z.: Validation of impact of assimilation of altimeter satellite significant wave height on wave forecast in the northwest Pacific, Acta Oceanologica Sinica, 31(6), 1-8, 2009.

835 Warner, J. C., Armstrong, B., He, R., and Zambon, J. B.: Development of a Coupled Ocean-Atmosphere-Wave-Sediment Transport (COASWST) modeling system, Ocean Modelling, 35(3), 230-244, 2010.

Waters, J., Wyatt, L. R., Wolf, J., and Hines, A.: Data assimilation of partitioned HF radar wave data into Wavewatch III, Ocean Modelling, 72, 17-31, 2013.

- 840 Webb, D. J.: Non-linear transfers between sea waves, Deep Sea Research, 25(3), 279-298, 1978. Wei, Y., Gao, Z., Tang, Z., Tang, Z., and Zeng, Y.: Analysis of wave factors in shipwreck accidents based on SAR wave mode data, Journal of Shanghai Ocean University, 26(6), 946-952, 2017. Yang, Y., Qiao, F., Zhao, W., Teng, Y., and Yuan, Y.: MASNUM ocean wave numerical model in spherical coordinates and its application, Acta Oceanologica Sinica, 27(2), 1-7, 2005.
- 845 Zhang, S., Harrison, M. J., Rosati, A., and Wittenbegr, A.: System Design and Evaluation of Coupled Ensemble Data Assimilation for Global Oceanic Climate Studies, Monthly Weather Review, 135(10), 3541-3564, 2007.

Zhang, S., Liu, Z., Rosati, A., and Delworth, T.: A study of enhancive parameter correction with coupled data assimilation for climate estimation and prediction using a simple coupled model, Tellus A:
850 Dynamic Meteorology and Oceanography, 64, doi: 10.3402/tellusa.v64i0.10963, 2012.

Zhang, Z., Qi, Y., Shi, P., Li, Z., and Li, Y.: Preliminary study on assimilation of significant wave heights from T/P altimeter, Acta Oceanologica Sinica, 25(5), 21-28, 2003.



855 Figure 1: The ground projection tracks of satellite Jason-2 (blue line, with an inclination of 66° N(S)) and SARAL (red line, with an inclination of 88° N(S)) in one cycle over approximately 10 days and 35 days, respectively, in the a) global and b) East Asia domains (zoomed out of green box in panel a).



Figure 2: Spatial distributions of a) background correlation coefficients by the empirical correlation model (blue), model data statistics (green) as well as their combination (red) of Eq. (3), b) background adjustment increments of SWH projected from the observational increment with the empirical correction model (blue line) and their combination model (red filled), and c) the difference of background adjustment increments of SWH from panel b by an analysis process given the single observation obtained at 114.09° E, 18.90° N, denoted by the asterisk (unit: meter).

删除了: (red)

(删除了:	SWH
(删除了:	from



- 870 Figure 3: The time series of RMSEs of the a) WW3 and b) SWAN perfect model experiments in the model control run with the NCEP CFSRv2 wind (black, denoted as WW3^{CFSR} in panel a and SWAN^{CFSR} in panel b), assimilating the "observed" data sampled by the tracks of Jason-2 (blue, denoted as WW3_{WW3(J2})^{CFSR} and SWAN_{SWAN(J2})^{CFSR}), Jason-2 & 3 (green, denoted as WW3_{WW3(J2+J3})^{CFSR} and SWAN_{SWAN(J2+J3})^{CFSR}, as well as Jason-2 & 3 and SARAL (cyan, denoted as WW3_{WW3(J2+J3+SA})^{CFSR} and SWAN_{SWAN(J2+J3+SA})^{CFSR}
- against the "truth" simulation forced by the ERA-Interim wind. The red and pink are forced by the NCEP CFSRv2 wind in the first 45 days, but the next 35 days are forced using the ERA-Interim wind (same as "truth") without (denoted as WW3^{CFSR} +WW3^{ERAI} and SWAN^{CFSR} +SWAN^{ERAI}) or with (denoted as WW3^{CFSR} + WW3_{WW3(J2)}).
 ^{ERAI} and SWAN^{CFSR} + SWAN_{SWAN(J2)}) the assimilation of Jason-2 data. The number in parentheses for each color is the corresponding RMSE <u>averaged globally over the verification</u>.
- 880 time period (30 days after the 45-day model spin-up and 5-day assimilation spin-up). The "observed" data are produced by projecting the "truth" SWH onto the satellite orbit.

删除了: summed



0.7 a) 0.6 0.5 0.4 RMSE (m) 0.3 WW3ERAI(.58) 0.2 $WW3_{SWAN(J2)}^{ERAI}(.44)$ WW3_{SWAN(J2+J3)}ERAI(.38) 0.1 $WW3_{SWAN(J2+J3+SA)} = (.36)$ $WW3_{SWAN(J2+J3+SA)} = ERAI(.35)$ $WW3_{SWAN(J2)} = (.43)$ 0 1 **b**) 0.97 0.94 Correlation Coefficient 0.91 0.88 WW3ERAI(.908) WW3_{SWAN(J2)}ERAI(.938) 0.85 WW3_{SWAN(J2+J3)}ERAI(.950) $WW3_{SWAN(J2+J3+SA)}^{ERAI}(.957)$ $WW3_{SWAN(J2+J3+SA)}^{ASSW}(.939)$ 0.82 10 15 20 5 25 30 35 Days 删除了:

Figure 4: The time series of RMSEs (upper) and <u>spatial</u> correlation coefficients (bottom) of the WW3(left) and SWAN(right) biased model experiments in the model control run forced by the ERA-Interim wind (black, denoted as WW3^{ERAI} and SWAN^{ERAI}), assimilations with "observed" data from one (pink, denoted as WW3_{SWAN(21+35})^{ERAI} and SWAN_{WW3(22+35})^{ERAI}, and SWAN_{WW3(22+35})^{ERAI}, and three (blue, denoted as WW3_{SWAN(21+35})^{ERAI} and SWAN_{WW3(22+35})^{ERAI}), and three (blue, denoted as WW3_{SWAN(21+35})^{ERAI} and SWAN_{WW3(21+355})^{ERAI}) and SWAN_{WW3(21+355})^{ERAI} and SWAN_{WW3(21+355})^{ERAI} and SWAN_{WW3(21+355})^{ERAI} and SWAN_{WW3(21+355})^{ERAI} and SWAN_{WW3(21+355})^{ERAI} and SWAN_{WW3(21+355})^{ERAI} and SWAN_{WW3(21+3555})^{ERAI} and SWAN_{W3(21+3555})^{ERAI} and SWAN_{W3(21+3555})^{ERAI} and SWAN_{W3(21+3555})^{ERAI} and SWAN_{W3(21+3555})^{ERAI} and SWAN_{W3(21+3555})^{ERAI} and SWAN_{W3(21+3}











Figure 7: Time series of RMSEs (upper) and <u>spatial</u> correlation coefficients (bottom) of WW3 (left) and ◄ 920 SWAN (right) produced by the model control run (black, denoted as WW3^{ERAI} and SWAN^{ERAI}), assimilation using the data from one (green, denoted as WW3_{J2}^{ERAI} and SWAN_{J2}^{ERAI}) and three satellites (blue, denoted as WW3_{J2+J3+SA}^{ERAI} and SWAN_{J2+J3+SA}^{ASSW} and SWAN_{J2+J3+SA}^{ASSW})_Ψ

带格式的:题注, 右 0.56 字符

(删除了: ←



925

Figure 8: Time series of a) RMSEs and b) <u>spatial</u> correlation coefficients produced by two bias correction schemes (cyan and pink) through a combination of WW3 and SWAN assimilations with the data from three satellites (Jason-2, Jason-3 and SARAL) and wind correction starting from the ERA-Interim wind. The results of the individual assimilation systems are plotted as dotted and dashed red lines (taken from 930 Fig. 7) for reference.

带格式的:题注, 右 0.56 字符

删除了:↩



带格式的:题注, 右 0.56 字符

设置了格式: 字体: (默认) Times New Roman, (中文) Times New Roman, 英语(英国)

(named Bias correction2 in Fig. 8). (unit: meter).









Figure 11: The time series of a) RMSEs and b) spatial correlation coefficients produced by the model control run (blue), data assimilation (pink) and their corresponding bias corrections (green and red) combining three wave model assimilation results (WAM, WW3 and SWAN) globally averaged over 1 year

(改重了 格式: 字体: (中文) Times New Roman
~(删除了:0
\mathcal{I}	设置了格式: 字体: (中文) Times New Roman
X	带格式的: 题注,右 0.56 字符
X	设置了格式: 字体: (中文) Times New Roman
Y,	删除了: . Other information is the same as in Fig. 8
$\langle \rangle$	删除了: ↩
X	设置了格式: 字体: (中文) Times New Roman, 非加粗

Exp. name	Model	Wind force	Assimilation or not	Role
WW3 ^{ERAI}	WW3	ERA-Interim	No	Truth for WW3 assimilation
WW3 ^{CFSR}	WW3	NCEP-CFSR	No	Model control for WW3 assimilation reference
WW3 _{WW3(J2)} CFSR	WW3	NCEP-CFSR	Yes (using Jason-2 track)	
WW3 _{WW3(J2+J3)} CFSR	WW3	NCEP-CFSR	Yes (using tracks of Jason- 2 and Jason-3)	Impact of observational system
WW3 _{WW3(J2+J3+SA)} CFSR	WW3	NCEP-CFSR	Yes (using tracks of Jason- 2, Jason-3 and SARAL)	
SWAN ^{ERAI}	SWAN	ERA-Interim	No	Truth for SWAN assimilation
SWAN ^{CFSR}	SWAN	NCEP-CFSR	No	Model control for SWAN assimilation reference
SWAN _{SWAN(J2)} CFSR	SWAN	NCEP-CFSR	Yes (using Jason-2 track)	
SWAN _{SWAN(J2+J3)} CFSR	SWAN	NCEP-CFSR	Yes (using tracks of Jason- 2 and Jason-3)	Impact of observational system
SWAN _{SWAN(J2+J3+SA}) ^{CFSR}	SWAN	NCEP-CFSR	Yes (using tracks of Jason- 2, Jason-3 and SARAL)	

Table 1: List of perfect model twin experiments.

Exp. name Model Wind source		Assimilation or not	Role	
WW3 ^{ERAI}	WW3	ERA-Interim	No	Truth for SWAN assimilation
SWAN ^{ERAI}	SWAN	ERA-Interim	No	Model control for SWAN assimilation
SWAN _{WW3(J2)} ERAI	SWAN	ERA-Interim	Yes (using Jason-2 track)	
SWAN _{WW3(J2+J3)} ERAI	SWAN	ERA-Interim	Yes (using tracks of Jason-2 and Jason-3)	Impact of observational system
SWANww3(J2+J3+SA) ^{ERAI}	SWAN	ERA-Interim	Yes (using tracks of Jason-2, Jason-3 and SARAL)	
SWANww3(J2 <u>+J3+SA</u>) ^{ASSW}	SWAN	Assimilation-corrected wind based on ERAI	Yes (using tracks of Jason-2. Jason-3 and SARAL)	Impact of assimilation- corrected wind
SWAN ^{ERAI}	SWAN	ERA-Interim	No	Truth for WW3 assimilation
WW3 ^{ERAI}	WW3	ERA-Interim	No	Model control for WW3 assimilation reference
WW3 _{SWAN(J2)} ERAI	WW3	ERA-Interim	Yes (using Jason-2 track)	
WW3 _{SWAN(J2+J3)} ERAI	WW3	ERA-Interim	Yes (using tracks of Jason-2 and Jason-3)	Impact of observational system
WW3 _{SWAN(J2+J3+SA)} ERAI	WW3	ERA-Interim	Yes (using tracks of Jason-2, Jason-3 and SARAL)	

955 Table 2: List of biased model twin experiments.

		WW3 _{SWAN(J2+J3+SA}) ^{ASSW}	WW3	Assimilation-corrected wind based on ERAI	Yes (using tracks of Jason-2, Jason-3 and SARAL)	Impact of assimilation- corrected wind
--	--	--	-----	---	---	---