



## Detecting causality signal in instrumental measurements and climate model simulations: global warming case study

Mikhail Y. Verbitsky<sup>1\*</sup>, Michael E. Mann<sup>2</sup>, Byron A. Steinman<sup>3</sup>, Dmitry M. Volobuev<sup>4</sup>

<sup>1</sup>Gen5 Group, LLC, 275 Grove Street, Suite 2-400, Newton, MA, USA

<sup>2</sup>Department of Meteorology, The Pennsylvania State University, University Park, PA, USA

<sup>3</sup>Large Lakes Observatory and Department of Earth and Environmental Sciences, University of Minnesota Duluth, Duluth, MN, USA

<sup>4</sup>The Central Astronomical Observatory of the Russian Academy of Sciences at Pulkovo, Saint Petersburg, Russia

\*retired

<sup>10</sup>formerly at: Yale University, Department of Geology and Geophysics, New Haven, CT, USA

Correspondence: Mikhail Verbitsky ([verbitskys@gmail.com](mailto:verbitskys@gmail.com))

**Abstract.** Detecting the direction and strength of the causality signal in observed time series is becoming a popular tool for exploration of distributed systems such as Earth's climate system. Here we suggest that in addition to reproducing observed time series of climate variables within required accuracy a model should also exhibit the causality relationship between variables found in nature. Specifically, we propose a novel framework for a comprehensive analysis of climate model responses to external natural and anthropogenic forcing based on the method of conditional dispersion. As an illustration, we assess the causal relationship between anthropogenic forcing (i.e., atmospheric carbon dioxide concentration) and surface temperature anomalies. We demonstrate a strong directional causality between global temperatures and carbon dioxide concentrations (meaning that carbon dioxide affects temperature stronger than temperature affects carbon dioxide) in both the observations and in (CMIP5) climate model simulated temperatures.

### 1. Introduction

The standard approach to attribution of observed global warming employs experiments with climate models. Such “detection and attribution” approaches (e.g., Stocker, 2014) attempt to reproduce observed trends under different external forcing conditions and demonstrate a consistency (or its absence) of simulated climate changes with instrumental observations. A substantial body of detection and attribution studies (e.g., Santer, *et al.*, 2009, 2012; Jones, *et al.*, 2013) spanning the past two decades demonstrates that anthropogenic increases in atmospheric carbon dioxide are very likely the cause of the observed global temperature increase since the mid-19<sup>th</sup> century. Semi-empirical approaches that combine information from model simulations and observations have also proven useful for investigations of modern climate change attribution. Previous work (e.g., Mann *et al.*, 2017) has employed estimates of natural variability derived from a combination of historical simulations and observations to attribute the sequence of record-breaking global temperatures in 2014, 2015, and 2016 to anthropogenic warming by demonstrating that this sequence had a negligible likelihood of occurrence in the absence of anthropogenic warming. Direct investigations of the causal relationship between climate system variables using statistical tools have become recently more common. The most simplistic approach, the Pearson's correlation between two time series, which is often mentioned in the context of causality, does not really measure the causality. While statistically significant correlation quantifies similarity between time series, it does not imply a causality resulting from physical relationships between the natural processes that are expressed by the time series and that can be modeled using differential equations. Instead, it provides a statistical test of a hypothesis that describes a physical link between the two variables (i.e., expressed as time series) without actually testing the plausibility of the physics underlying the hypothesis. The breakthrough Granger developments (Granger, 1969) provided a foundation for several causality measuring techniques based on different hypotheses of data origin. The requirement of the cause leading the effect (but not vice-versa) defines the direction of a causal link if a more general hypothesis of lagged linear connection between noisy autoregressive processes is assumed. Though this hypothesis leads to statistically significant estimates of climate response to the forcing input (e.g., Kaufmann *et al.*, 2006, 2011, Attanasio, 2012, Mokhov *et al.*, 2012, Triacca *et al.*, 2013), it may not be able to reliably detect the direction of causality in the climate system because the potential for non-linearities in the climate system (leading to extreme sensitivity to initial conditions, i.e., deterministic chaos) is not taken into account. For example, Palus *et al.* (2018) demonstrated that coupled chaotic dynamical systems can “violate the first principle of Granger causality that the cause precedes the effect”.

The Shannon information flow approach expands Granger causality to non-linear systems, using transfer entropy as a causality measure. Barnett *et al.* (2009) have shown that transfer entropy is equivalent to Granger causality for



Gaussian processes. The transfer entropy between two probability distributions is typically considered the most general approach for causality detection, and numerous modifications of transfer-entropy-based causality measuring techniques have been developed for different applications (Pearl, 2009), including causality measurements of global warming (e.g., Stips *et al.*, 2016). It should be noted though that all applications require long time series to calculate statistical distributions. The prediction improvement approach is often considered as a generalization of Granger causality for non-linear systems (e.g., Krakovska and Hanzely, 2016). It is highly practical and, besides causality calculations, it may help to improve the prediction accuracy. For pure causality purposes, however, it adds an additional uncertainty because the causality may depend on the chosen prediction method. The convergent cross-mapping approach (Sugihara *et al.*, 2012, Van Nes *et al.*, 2015) has been recently designed to work with relatively short data series, thus addressing the major constraint of transfer-entropy approach. The background hypotheses of the method is more narrow and includes only nonlinear *dynamical* systems, though convergent cross-mapping remains applicable to most natural systems in ecology and geosciences (Sugihara *et al.*, 2012). The approach considers conditional evolution of nearest neighbors in the reconstructed Takens' space so it is sensitive to the noise and may not be applicable to a wide range of time scales. Moreover, Palus *et al.* (2018) have shown that convergent cross-mapping is not capable of determining the directionality of a causal link.

For our case study, we advocate the method of conditional dispersion (MCD) developed by Čenys *et al.* (1991). It has also been designed for non-linear systems and exploits the asymmetry of the conditional dispersion of two variables in Takens' space along all available scales. Therefore, it remains more general and noise resistant than convergent cross-mapping techniques and more general than prediction improvement approaches because it is insensitive to the choice of the prediction method. We propose here to employ the MCD-based causality measurements for a comprehensive analysis of climate model responses to external natural and anthropogenic forcing. While climate models have been tuned in a rough sense to reproduce the observational record, their predictions differ from the observations due to various types of errors and uncertainties. These include: (a) measurement errors in external forcing (e.g., greenhouse gas concentrations, land use, solar variability) used to drive the models; (b) errors in the representation of physical processes in the models (e.g., ocean circulation, cryosphere and biosphere processes, various feedback mechanisms, *etc.*) and incomplete representation of the Earth system (i.e., in many cases lack of representations of dynamic vegetation responses, or oceanic carbon cycle dynamics); (c) errors associated with internal variability in the climate system – for example, models may accurately represent ENSO, The El Niño-Southern Oscillation, but ENSO is an inherently random process and models therefore do not, and should not, reproduce the actual real-world realization of that random process; (d) errors and uncertainties in observational data – for example, surface temperature measurements contain sampling uncertainty due to the irregular sampling in space and time (e.g., lack of data at higher latitudes increasingly back in time). Such sampling uncertainties might lead to model – observational data mismatch that is unrelated to model performance. In addition, there is the potential for biases due, for example, to changes in oceanic and terrestrial measurement platforms over time (e.g., bucket measurements *vs.* intake valves for ocean seawater measurements, or residual urban heat island biases in land-based temperature measurements). The challenge, then, is to determine the best-performing models when all reproduce the observations similarly well. We believe that in addition to reproducing observed time series of climate variables, a model should exhibit the causality relationship between variables found in nature. Since the MCD approach is based on the assumption that each time series is produced by a hypothetical low-dimensional system of dynamical equations, *similarity of causal relationships in both model and observations speaks to the similarities of their parent systems.*

Accordingly, our paper is structured as follows. First, we will briefly describe the method of conditional dispersion. We will illustrate it with several numerical experiments that investigate the causal relationship between surface temperature anomalies for the Northern Hemisphere and atmospheric carbon dioxide concentration measurements. We will show that the causality between carbon dioxide and temperature anomalies is a directional causality, meaning that carbon dioxide affects temperature stronger than temperature affects carbon dioxide. We will then demonstrate that this directional connection cannot be replicated with an independent trend and red noise.

## 2. A glimpse into the method of conditional dispersion.

The MCD approach has been designed for measuring causality between two time series. It is assumed that each time series is a variable produced by its hypothetical low-dimensional system of dynamical equations. The variables contain information about the dynamics of hypothetical parent systems which can be reconstructed using Takens (1981) procedure. Since each of the variables can be used to reconstruct the original parent system manifold, there is



one-to-one correspondence between them. Specifically, if points of one time series are close, the synchronous points of another variable are close as well. Therefore, if two variables ( $u$  and  $x$ ) do not belong to the same or coupled dynamical systems, or in other words, they are independent, then the distance from a reference point to its neighbors of one variable ( $u$ ) does not depend on the distance ( $\varepsilon$ ) between synchronous points of another variable ( $x$ ). In the case of dependency, though, the distance between neighboring points of the controllable variable will be smaller when the distance between points of the driving variable is reduced. Therefore, the dependence of the conditional dispersion  $\sigma(\varepsilon)$  of the variable  $u$  upon the distance  $\varepsilon$  between points of the variable  $x$  becomes a signature of causal relationship between  $u$  and  $x$  (Čenys *et al.*, 1991):

$$\sigma_{xu}^M(\varepsilon) = \left( \frac{\sum_{i \neq j} \|u_i^M - u_j^M\|^2 \Theta(\varepsilon - \|x_i^M - x_j^M\|)}{\sum_{i \neq j} \Theta(\varepsilon - \|x_i^M - x_j^M\|)} \right)^{1/2} \quad (1)$$

Here,  $M$  is dimension of the reconstructed manifold, and  $\Theta$  is the Heaviside function. If variable  $u$  is independent of variable  $x$ , its conditional dispersion  $\sigma_{xu}^M(\varepsilon)$  does not depend upon  $\varepsilon$ . If variable  $x$  is the cause of  $u$ -variability, then conditional dispersion of the variable  $u$  will decline for diminishing  $\varepsilon$ . As an illustration, we show on Fig.1 the conditional dispersion of two variables  $x$  and  $u$  of coupled Henon (1976) maps:


$$\begin{cases} x_{n+1} = 1 + y_n - 1.4x_n x_n \\ y_{n+1} = 0.3x_n + \alpha(y_n - y_n) \end{cases} \quad (2)$$

$$\begin{cases} u_{n+1} = 1 + v_n - 1.4u_n u_n \\ v_{n+1} = 0.3u_n + \beta(y_n - v_n) \end{cases} \quad (3)$$

$$\begin{cases} u_{n+1} = 1 + v_n - 1.4u_n u_n \\ v_{n+1} = 0.3u_n + \beta(y_n - v_n) \end{cases} \quad (4)$$

$$\begin{cases} u_{n+1} = 1 + v_n - 1.4u_n u_n \\ v_{n+1} = 0.3u_n + \beta(y_n - v_n) \end{cases} \quad (5)$$

Here variables  $u$  and  $x$  belong to two dynamical subsystems, (2) - (3) and (4) - (5). The interdependence of these subsystems is defined by coefficients  $\alpha$  and  $\beta$ . When connection is one-directional (for example,  $\alpha=0$ ,  $\beta=0.3$ ), i.e.,  $x$  is the cause of  $u$  but is independent of  $u$ , the conditional dispersion of  $x$ -variable does not depend on  $\varepsilon$  (where  $\varepsilon$  is the distance between synchronous points of  $u$ ) but conditional dispersion of  $u$ -variable declines for diminishing  $\varepsilon$  (where  $\varepsilon$  is the distance between synchronous points of  $x$ ). When the connection is two-directional (for example,  $\alpha=0.1$ ,  $\beta=0.3$ ), the conditional dispersion of both variables declines for smaller  $\varepsilon$ , but a variable which provides a stronger causal force (i.e.,  $x$ ) has a dispersion with a less articulated slope. When variables are equally interdependent (i.e., synchronized), the conditional dispersions of both variables may have the same slope.



**Fig. 1.** The conditional dispersion of coupled dynamical variables  $u$  and  $x$  as described by equations (2) - (5). When connection is one-directional ( $\alpha=0$ ,  $\beta=0.3$ ), i.e.,  $x$  is the cause of  $u$  but is independent of  $u$ , the conditional dispersion of  $x$ -variable does not depend on  $\varepsilon$ , but conditional dispersion of  $u$ -variable declines for diminishing  $\varepsilon$ . When the connection is two-directional ( $\alpha=0.1$ ,  $\beta=0.3$ ), the conditional dispersion of both variables declines for smaller  $\varepsilon$ , but  $x$ -variable which provides a stronger causal force has a dispersion with weaker slope.




### 3. The case study of global warming causality

We will now employ the MCD approach in three numerical experiments for time series of atmospheric carbon dioxide concentration and surface temperature obtained from both direct instrumental measurements and model simulations.

#### 3.1 Detecting causality in instrumental measurements.

First, we investigate the causal relationship between GISTEMP (Hansen *et al.*, 2006) surface temperature assessments for the Northern Hemisphere and atmospheric carbon dioxide concentration measurements (CO<sub>2</sub> NASA GISS Data, 2016) spanning 1880 through 2016. For this purpose, we calculate conditional dispersion of Northern Hemisphere temperature variability (as a function of distance  $\varepsilon$  between synchronous points of the carbon dioxide time series) and conditional dispersion of carbon dioxide (as a function of distance  $\varepsilon$  between synchronous points of the temperature time series). The results are shown in Fig. 2.



**Fig. 2. Left:** GISTEMP (Hansen *et al.*, 2006) surface temperature anomalies and atmospheric carbon dioxide concentration measurements (CO<sub>2</sub> NASA GISS Data, 2016); **Right:** Conditional dispersion of instrumental measurements. Black curve is conditional dispersion of the carbon dioxide concentration. Green curve represents conditional dispersion of the Northern Hemisphere temperature anomalies; its dependence on  $\varepsilon$  is much stronger than that of black curve indicating that carbon dioxide is the cause of temperature changes.

It can be seen that surface temperature and carbon dioxide are interdependent systems (the conditional dispersions of both variables depend upon  $\varepsilon$ ). Nevertheless, *carbon dioxide is the causal force of global warming* because the dependence of the temperature conditional dispersion upon  $\varepsilon$  is much stronger than the same dependence of carbon dioxide conditional dispersion.


#### 3.2 Detecting causality in model simulations. Anthropogenic and natural (volcanic and solar) forcing.

We now apply MCD to the model simulations adopted from the Coupled Model Intercomparison Project Phase 5 (CMIP5) historical simulation experiments (Stocker, 2014). Estimates of the total forced component of Northern Hemisphere mean temperature have been derived by averaging over the full ensemble of CMIP5 multimodel all-forcing historical experiments (Mann *et al.*, 2014, Steinman *et al.*, 2015, Mann *et al.*, 2017). We generated 50 temperature series surrogates using a Monte Carlo resampling approach of Mann *et al.* (2017) and calculated conditional dispersion of Northern Hemisphere temperature variability for each of the 50 surrogates (as a function of distance  $\varepsilon$  between synchronous points of carbon dioxide time series) and conditional dispersion of carbon dioxide (as a function of distance  $\varepsilon$  between synchronous points of every surrogate time series).

The results are presented in Fig.3. It can be seen that the behavior of dispersions derived from multiple simulations' surrogates is quantitatively close to the dispersions obtained from the direct measurements, and therefore that carbon




dioxide is the driver of temperature changes in the model simulations. Though in this experiment we applied MCD-testing network to the full ensemble of CMIP5 models, the same procedure can be applied to any sub-assembly or to individual models if the task is to identify the models that are more consistent with the instrumental data in terms of causality.



**Fig. 3.** Detecting causality in model simulations. Anthropogenic and natural (volcanic and solar) forcing. **Left:** Surrogates of model temperature deviations induced by both natural and anthropogenic forcing together with carbon dioxide concentration measurements (CO<sub>2</sub> NASA GISS Data, 2016). **Right:** Conditional dispersion of the Northern Hemisphere temperature and carbon dioxide concentration. Black curve is conditional dispersion of the carbon dioxide concentration instrumental measurements; green curve represents conditional dispersion of the Northern Hemisphere temperature measurements (same as in the right panel of Fig.2). Blue triangles are mean of 50 multimodel surrogates' conditional dispersions of the Northern Hemisphere temperature; small black dots are mean of 50 multimodel surrogates conditional dispersions of carbon dioxide. Bars represent doubled standard deviation.

### 3.3 Detecting causality in model simulations. Anthropogenic forcing only.

We repeat the analysis described in paragraph 3.2 but for a separate ensemble of anthropogenic-only forcing experiments (Stocker, 2014, Mann *et al.*, 2016 a, 2016 b, 2017). The results are presented in Fig.4.




**Fig. 4.** Same as in Fig.3 but for anthropogenic CO<sub>2</sub> forcing only.

Interestingly, the results of the analysis change minimally when natural forcing (volcanic and solar) are excluded, which implies the dominant causality role of carbon dioxide.



#### 4. Testing boundaries of MCD applicability.

Initially, the MCD approach was applied to causality measurements between deterministic chaotic time series (Čenys *et al.*, 1991). In this study, we expand its applicability to a situation where one of the time series ( $\text{CO}_2$ ) is essentially a regular trend and the history of observations for both  $\text{CO}_2$  and temperature is relatively short. In the next experiment we will test boundaries of MCD applicability and investigate if the MCD approach can distinguish between *interdependent* processes, like  $\text{CO}_2$  and temperature, as well as *independent* but highly autocorrelated processes. For this purpose, we calculate conditional dispersion for two independent but highly autocorrelated time series resembling properties of carbon dioxide series and temperature surrogates. For the carbon dioxide “role” we selected a linear trend. GISS temperature surrogates were replaced by 50 red noise surrogates. Results of the conditional dispersion calculations are shown in Fig.5.



**Fig. 5.** Conditional dispersion of two independent processes having high autocorrelations. **Left:** Example of one-lag autocorrelated (0.92) red noise series simulating statistical properties of GISS temperature record and a linear trend; **Right:** Average conditional dispersion of 50 red noise surrogates. Error bars mark doubled standard deviation.

This example shows that, for relatively short time series, MCD is unable to discriminate cases of independence and very strong interdependence (i.e., synchronization) because spontaneous local correlations may be induced with autocorrelated red noise leading to the same slope of conditional dispersions for both time series. Nevertheless, unlike natural ( $\text{CO}_2$  and temperature) time series, these correlations are not able to induce any directional causality. In other words, were temperature and  $\text{CO}_2$  equally interdependent, MCD would not be able to distinguish this situation from independent trends and red noise. In reality though, carbon dioxide affects temperature stronger than temperature affects  $\text{CO}_2$ , and this situation cannot be replicated by a trend and red noise.

#### 5. Conclusions

In this study we propose an additional climate model validation procedure that assesses whether causality signals between model drivers and responses are consistent with those observed in nature. Specifically, we suggest the method of conditional dispersion (MCD) as the best approach to directly measure the causality between model forcing and response. As an illustration of MCD applicability, we detect the causality signal between atmospheric carbon dioxide concentration and variations of global temperature. Our results suggest that there is a strong causal signal from the carbon dioxide series to the global temperature series or, in other words, that carbon dioxide is the principal *cause* of global warming. This conclusion is applicable to both direct instrumental measurements and multimodel temperature series surrogates. The strength of the causality signal does not significantly change when the additional contribution from natural factors (such as solar and volcanic) are accounted for, implying that increases in carbon dioxide are the main driver of observed warming. It is noteworthy that the causality between carbon dioxide and temperature anomalies is a directional causality: carbon dioxide affects temperature stronger than temperature affects carbon dioxide. This directional connection cannot be replicated using simplistic statistical models for the observed temperature increase (an independent trend and red noise).





**Code and data availability.** The MatLab source code and data (Verbitsky et al., 2019) are available at <https://zenodo.org/record/2605142#.XJirxylzbc> (<http://doi.org/10.5281/zenodo.2605142>). Scripts were tested under MatLab version R2015b (last access: 25 March 2019).

5 **Author contributions.** MV conceived the research. MV and DV wrote the manuscript. All authors contributed equally to the design of the research and to editing the manuscript.

**Competing interests.** The authors declare that they have no conflict of interest.

**Acknowledgements.** Dmitry Volobuev has been funded in part by the Russian Foundation for Basic Research, grant 19-02-00088-a.

## References

- 10 Attanasio, A.: Testing for linear Granger causality from natural/anthropogenic forcings to global temperature anomalies. *Theoretical and applied climatology*, 110 (1-2), 281-289, 2012
- Barnett, L., Barrett, A. B. and Seth, A. K.: Granger causality and transfer entropy are equivalent for Gaussian variables. *Physical review letters*, 103(23), 238701, 2009
- Čenys, A., Lasiene, G. and Pyragas, K.: Estimation of interrelation between chaotic observables. *Physica D: Nonlinear Phenomena*, 52(2-3), 332-337, 1991
- 15 CO2 NASA GISS Data: <https://data.giss.nasa.gov/modelforce/ghgases/Fig1A.ext.txt>, 2012
- Granger, C.W.: Investigating causal relations by econometric models and cross-spectral methods. *Econometrica: Journal of the Econometric Society*, 424-438, 1969
- 20 Hansen, J., Sato, M., Ruedy, R., Lo, K., Lea, D.W. and Medina-Elizade, M.: Global temperature change. *Proceedings of the National Academy of Sciences*, 103(39), 14288-14293, 2006
- Hénon, M.: A two-dimensional mapping with a strange attractor. *The Theory of Chaotic Attractors*. Springer, New York, NY, 94-102, 1976
- Jones, G.S., Stott, P.A. and Christidis, N.: Attribution of observed historical near-surface temperature variations to anthropogenic and natural causes using CMIP5 simulations. *Journal of Geophysical Research: Atmospheres*, 118 (10), 4001-4024, 2013
- 25 Kaufmann, R.K., Kauppi, H. and Stock, J.H.: Emissions, concentrations, and temperature: a time series analysis. *Climatic Change*, 77(3-4), 249-278, 2006
- Kaufmann, R.K., Kauppi, H., Mann, M.L. and Stock, J.H.: Reconciling anthropogenic climate change with observed temperature 1998–2008. *Proceedings of the National Academy of Sciences*, 108(29), 11790-11793, 2011
- 30 Krakovská, A., Hanzely, F.: Testing for causality in reconstructed state spaces by an optimized mixed prediction method. *Physical Review E*, 94(5), 052203, 2016
- Mann, M. E., Steinman, B. A., and Miller, S. K.: On forced temperature changes, internal variability, and the AMO. *Geophys. Res. Lett.* 41, 3211–3219, 2014
- 35 Mann, M. E., Rahmstorf, S., Steinman, B. A., and Miller S. K.: The likelihood of recent record warmth, *Nat. Sci. Rep.*, 6, 19,831, 2016 a



- Mann, M. E., Steinman, B., Miller, S. K., Frankcombe, L., England, M., and Cheung A. H.: Predictability of the recent slowdown and subsequent recovery of large-scale surface warming using statistical methods, *Geophys. Res. Lett.*, 43, 3459–3467, doi:10.1002/2016GL068159, 2016 b
- 5 Mann, M.E., Miller, S.K., Rahmstorf, S., Steinman, B.A. and Tingley, M.: Record temperature streak bears anthropogenic fingerprint. *Geophysical Research Letters*, 44(15), 7936-7944, 2017
- Mokhov, I.I., Smirnov, D.A. and Karpenko, A.A.: March. Assessments of the relationship of changes of the global surface air temperature with different natural and anthropogenic factors based on observations. In *Doklady Earth Sciences*, 443 (1), 381-387, 2012
- 10 Paluš, M., Krakovská, A., Jakubík, J., and Chvosteková, M.: Causality, dynamical systems and the arrow of time. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 28(7), 075307, 2018
- Pearl, J.: *Causality*. Cambridge University Press, 2009
- Santer, B.D., Painter, J., Mears, C.A., Doutriaux, C., Caldwell, P., Arblaster, J., Cameron-Smith, P.J., Gillett, N.P., Gleckler, P.J., Lanzante, J. and Perlwitz, J.: Identifying human influences on atmospheric temperature: Are results robust to uncertainties? *AGU Fall Meeting Abstracts*, 2012
- 15 Santer, B.D., Taylor, K.E., Gleckler, P.J., Bonfils, C., Barnett, T.P., Pierce, D.W., Wigley, T.M.L., Mears, C., Wentz, F.J., Brüggemann, W. and Gillett, N.P.: Incorporating model quality information in climate change detection and attribution studies. *Proceedings of the National Academy of Sciences*, 106 (35), 14778-14783, 2009
- Steinman, B. A., Mann, M. E. and Miller, S. K.: Atlantic and Pacific multidecadal oscillations and Northern Hemisphere temperatures. *Science* 347, 998–991, 2015
- 20 Stips, A., Macias, D., Coughlan, C., Garcia-Goriz, E. and San Liang, X.: On the causal structure between CO<sub>2</sub> and global temperature. *Scientific reports*, 6, 21691, 2016
- Stocker, T. (ed.): *Climate change 2013: the physical science basis: Working Group I contribution to the Fifth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, (Chapter 10: Detection and Attribution of Climate Change: from Global to Regional), 867-952, 2014
- 25 Sugihara, G., May, R., Ye, H., Hsieh, C.H., Deyle, E., Fogarty, M. and Munch, S.: Detecting causality in complex ecosystems. *Science*, 1227079, 2012
- Takens, F.: Detecting strange attractors in turbulence, *Dynamical Systems and Turbulence*, Warwick 1980. Lecture notes in mathematics, 898, 366-381, 1981
- 30 Triacca, U., Attanasio, A. and Pasini, A.: Anthropogenic global warming hypothesis: testing its robustness by Granger causality analysis. *Environmetrics*, 24(4), 260-268, 2013
- Van Nes, E.H., Scheffer, M., Brovkin, V., Lenton, T.M., Ye, H., Deyle, E. and Sugihara, G.: Causal feedbacks in climate change. *Nature Climate Change*, 5(5), 445, 2015
- 35 Verbitsky, M.Y., Mann, M.E., Steinman, B.A., and Volobuev, D.M.: Supplementary code and data to GMD paper "Detecting causality signal in instrumental measurements and climate model simulations: global warming case study" (Version 1.0). *Zenodo*. <http://doi.org/10.5281/zenodo.2605142> (last access: 25 March 2019), 2019