

1  
2  
3  
4 **Observations for Model Intercomparison Project (Obs4MIPs):**  
5 **Status for CMIP6**  
6  
7

8 Duane Waliser<sup>1</sup>, Peter J. Gleckler<sup>2</sup>, Robert Ferraro<sup>1</sup>, Karl E. Taylor<sup>2</sup>, Sasha Ames<sup>3</sup>, James  
9 Biard<sup>4</sup>, Michael G. Bosilovich<sup>5</sup>, Otis Brown<sup>4</sup>, Helene Chepfer<sup>6</sup>, Luca Cinquini<sup>1,7</sup>, Paul J. Durack<sup>2</sup>,  
10 Veronika Erying<sup>8,9</sup>, Pierre-Phillipe Mathieu<sup>10</sup>, Tsengdar Lee<sup>11</sup>, Simon Pinnock<sup>10</sup>, Gerald L.  
11 Potter<sup>5</sup>, Michel Rixen<sup>12</sup>, Roger Saunders<sup>13</sup>, Jörg Schulz<sup>14</sup>, Jean-Noël Thépaut<sup>15</sup>, and Matthias  
12 Tuma<sup>12</sup>  
13  
14

15 <sup>1</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

16 <sup>2</sup>Program for Climate Model Diagnostics and Intercomparison, Lawrence Livermore National  
17 Laboratory, Livermore, California, USA

18 <sup>3</sup> Lawrence Livermore National Laboratory, Livermore, California, USA

19 <sup>4</sup>North Carolina Institute for Climate Studies, North Carolina State University, Asheville, North  
20 Carolina, USA.

21 <sup>5</sup>Goddard Space Flight Center, National Aeronautics and Space Administration, Greenbelt, Maryland,  
22 USA

23 <sup>6</sup>Laboratoire de Météorologie Dynamique / Institut Pierre Simon Laplace, Université Pierre et Marie  
24 Curie, Paris, France.

25 <sup>7</sup>Earth System Research Laboratory, National Ocean and Atmospheric Administration, Boulder,  
26 Colorado, USA.

27 <sup>8</sup>Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen,  
28 Germany.

29 <sup>9</sup>University of Bremen, Institute of Environmental Physics (IUP), Bremen, Germany

30 <sup>10</sup>Climate Office, European Space Agency, Harwell, United Kingdom

31 <sup>11</sup> Earth Science Division, National Aeronautics and Space Administration, Washington, DC, USA

32 <sup>12</sup>World Climate Research Programme, World Meteorological Organization, Geneva, Switzerland

33 <sup>13</sup>MetOffice, Exeter, United Kingdom

34 <sup>14</sup>European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT), Darmstadt,  
35 Germany

36 <sup>15</sup>European Centre of Medium-Range Weather Forecasting, Reading, United Kingdom.  
37

38 *Submitted to the CMIP6 GMD special issue*

39 *September, 2019*

40 *ReSubmitted to the CMIP6 GMD special issue*

41 *April, 2020*  
42

43 *Please email comments to: [duane.waliser@jpl.nasa.gov](mailto:duane.waliser@jpl.nasa.gov) & [gleckler1@llnl.gov](mailto:gleckler1@llnl.gov)*

## Abstract

The Observations for Model Intercomparison Projects (Obs4MIPs) was initiated in 2010 to facilitate the use of observations in climate model evaluation and research, with a particular target being the Coupled Model Intercomparison Project (CMIP), a major initiative of the World Climate Research Programme (WCRP). To this end, Obs4MIPs: 1) targets observed variables that can be compared to CMIP model variables, 2) utilizes dataset formatting specifications and metadata requirements closely aligned with CMIP model output, 3) provides brief technical documentation for each dataset, designed for non-experts and tailored towards relevance for model evaluation, including information on uncertainty, dataset merits and limitations, and 4) disseminates the data through the Earth System Grid Federation (ESGF) platforms, making the observations searchable and accessible via the same portals as the model output. Taken together, these characteristics of the organization and structure of obs4MIPs should entice a more diverse community of researchers to engage in the comparison of model output with observations and to contribute to a more comprehensive evaluation of the climate models.

At present, the number of obs4MIPs datasets has grown to about 80, many undergoing updates, with another 20 or so in preparation, and more than 100 proposed and under consideration. A partial list of current global satellite-based datasets includes: humidity and temperature profiles; a wide range of cloud and aerosol observations; ocean surface wind, temperature, height, and sea ice fraction; surface and top of atmosphere longwave and shortwave radiation; and ozone (O<sub>3</sub>), methane (CH<sub>4</sub>) and carbon dioxide (CO<sub>2</sub>) products. A partial list of proposed products expected to be useful in analyzing CMIP6 results includes: alternative products for the above quantities, and additional products for ocean surface flux and chlorophyll products, a number of vegetation products (e.g. FAPAR, LAI, burnt area fraction), ice sheet mass and height, carbon monoxide (CO) and nitrogen dioxide (NO<sub>2</sub>). While most existing obs4MIPs datasets consist of monthly mean gridded data over the global domain, products with higher time resolution (e.g. daily) and/or regional products are now receiving more attention.

Along with an increasing number of datasets, obs4MIPs has implemented a number of capability upgrades including: 1) an updated obs4MIPs data specifications document that provides for additional search facets and generally improves congruence with CMIP6 specifications for

1 model datasets, 2) a set of six easily understood indicators that help guide users as to a dataset's  
2 maturity and suitability for application, and 3) an option to supply supplemental information about  
3 a dataset beyond what can be found in the standard metadata. With the maturation of the obs4MIPs  
4 framework, the dataset inclusion process, and the dataset formatting guidelines and resources, the  
5 scope of the observations being considered is expected to grow to include gridded in-situ datasets  
6 as well as datasets with a regional focus, and the ultimate intent is to judiciously expand this scope  
7 to any observation dataset that has applicability for evaluation of the types of Earth System models  
8 used in CMIP.

9

# 1. Introduction

State, national and international climate assessment reports are growing in their importance as a scientific resource for climate change understanding and assessment of impacts crucial for economic and political decision-making [WorldBank, 2011; IPCC, 2014; NCA, 2014; EEA, 2015]. A core element of these assessment reports are climate model simulations that not only provide a projection of the future climate but also information relied on in addressing adaptation and mitigation questions. These quantitative projections are the product of extremely complex multi-component, global and regional climate models (GCMs and RCMs). Because of the critical input such models provide to these assessments, and in light of significant systematic biases that potentially impact their reliability [e.g., Meehl et al. 2007; Waliser et al. 2007, 2009; Gleckler et al., 2008; Reichler and Kim, 2008; Eyring and Lamarque, 2012; Whitehall et al., 2012; IPCC, 2013; Stouffer et al. 2017], it is important to expand the scrutiny of them through the systematic application of observations from gridded satellite and reanalysis products as well as in-situ station networks. Enabling such observation-based, multivariate evaluations is needed for assessing model fidelity, performing quantitative model comparison, gauging uncertainty, and constructing defensible multi-model ensemble projections. These capabilities are all necessary to provide a reliable characterization of future climate that can lead to an informed decision-making process.

Optimizing the use of the plethora of observations for model evaluation is a challenge, albeit facilitated to a considerable degree by the vast strides the Coupled Model Intercomparison Project (CMIP) community has made in implementing systematic and coordinated experimentation in support of climate modeling research (Meehl et al., 2007; Taylor et al., 2012; Eyring et al., 2016). CMIP is a flagship project of the World Climate Research Programme and is overseen by its Working Group on Coupled Modelling (WGCM). This architecture includes an increasingly complex set of simulation experiments designed to address specific science questions and to facilitate model evaluation [Meehl et al., 2007; Taylor et al., 2012; Eyring et al., 2016], highly detailed specifications for model output<sup>1</sup> [e.g., Taylor et al., 2009; Juckes et al 2019], and adoption of a distributed approach to manage and disseminate the rapidly increasing data volumes of climate

---

<sup>1</sup> <https://goo.gl/v1drZl>

1 model output [Williams et al. 2016]. The highly collaborative infrastructure framework for CMIP  
2 has been advancing since the first World Climate Research Programme (WCRP) Model  
3 Intercomparison Project [MIP; Gates, 1992], with a payoff that became especially evident during  
4 CMIP3 [Meehl et al., 2007] when the highly organized and readily available model results  
5 facilitated an enormous expansion in the breadth of analysis that could be undertaken [Taylor et  
6 al., 2012; Eyring et al., 2016]. The systematic organization of model results and their archiving  
7 and dissemination was catalytic in developing a similar vision for observations as described in this  
8 article.

9 As the significance of the climate projections has grown in regards to considerations of  
10 adaptation and mitigation measures, so has the need to quantify model uncertainties and identify  
11 and improve model shortcomings. For these purposes, it is essential to maximize the use of  
12 available observations. For instance, observations enable evaluation of a model's climatological  
13 mean state, annual cycle, and variability across timescales, as a partial gauge of model fidelity in  
14 representing different Earth System Processes. The genesis of the obs4MIPs effort stemmed from  
15 the impression that there were many observations that were not being fully exploited for model  
16 evaluation. A notable driver of the early thinking and developments in obs4MIPs was the  
17 recognition – partly from the success of the CMIP experimental architecture in providing greater  
18 model output accessibility – that much of the observation-based model evaluation research was  
19 being conducted by scientists without an expert's understanding of either the observations being  
20 employed or the climate models themselves. Nevertheless, there was a clear imperative, given the  
21 discussion above, to encourage and assist the growing class of climate research scientists who were  
22 beginning to devote considerable effort to the evaluation and analysis of climate model simulations  
23 and projections (left panel of Fig. 1). A sister effort, “CREATE-IP” (initially conceived as  
24 ana4MIPs), has been advanced to make reanalyses data available with a similar objective (Potter,  
25 et al., 2018).

26 While the infrastructure advances made by CMIP had established an obvious precedent, the  
27 daunting prospect of dealing in a similar way with the plethora of observation quantities was  
28 challenging, even when only considering satellite data. Within the NASA holdings, for example,  
29 there have been over 50 Earth observation missions flown, each producing between 1 to nearly  
30 100 quantities, and thus there are likely on the order of 1000 NASA satellite geophysical quantities

1 that might be candidates for migration to obs4MIPs, with many more when accounting for  
2 EUMETSAT, NOAA, ESA, JAXA, etc satellite datasets. Key to making initial progress was the  
3 recognition, illustrated in the right panel of Fig. 1, that only a fraction (perhaps about 10%) of the  
4 available observation variables could be directly compared with the available CMIP output  
5 variables of which there are over a thousand. The aspirations and framework for obs4MIPs were  
6 developed with these considerations in mind. Since the initial implementation of obs4MIPs, there  
7 has been an intention to expand the breadth of datasets including a better match of derived  
8 quantities and model output variables, e.g., through using simulators (e.g., Bodas-Salcedo et al.,  
9 2011) and an increased capacity to host the datasets, as well as to describe and disseminate them.  
10 In addition, for the first time in CMIP, evaluation tools are available that make full use of the  
11 obs4MIPs data for routine evaluation of the models (Eyring et al. 2016) as soon as the output is  
12 published to the ESGF (e.g., the Earth System Model Evaluation Tool, ESMValTool, Eyring et  
13 al., 2019, ; the PCMDI Metrics Package, Gleckler et al., 2016; the NCAR Climate Variability  
14 Diagnostic Package, Phillips et al., 2014; and the International Land Model Benchmarking ILAMB  
15 package, Collier et al., 2018).

16 In the next section, the history and initial objectives of the obs4MIPs project are briefly  
17 summarized. Section 3 describes the needs and efforts to expand the scope of obs4MIPs beyond  
18 its initial objectives, particularly for including a wider range of observational resources in  
19 preparation for CMIP6. Section 4 provides an updated accounting of the obs4MIPs dataset  
20 holdings, descriptions of a number of new features, including updated dataset specifications,  
21 dataset indicators, and accommodation for supplementary material, and a brief description of the  
22 alignment and intersection of obs4MIPs and CMIP model evaluation activities. Section 5  
23 discusses challenges and opportunities for further expansion and improvements to obs4MIPs and  
24 potential pathways for addressing them.

## 25 **2. Background**

26 In late 2009, the Jet Propulsion Laboratory (JPL)/NASA and the Program for Climate Model  
27 Diagnostics and Intercomparison (PCMDI)/DOE began discussions on ways to better utilize global  
28 satellite observations for the systematic evaluation of climate models, with the fifth phase of the

1 WCRP’s CMIP5 in mind. A two-day workshop was held at PCMDI in October 2010 that brought  
2 together experts on satellite observation, modeling, and climate model evaluation [*Gleckler et al.*,  
3 2011]. The objectives of the meeting were to: 1) identify satellite datasets that were well suited to  
4 provide observation reference information for CMIP model evaluation, 2) define a common  
5 template for documentation of observations, particularly with regard to model evaluation, and 3)  
6 begin considerations of how to make the observations and technical documentation readily  
7 available to the CMIP model evaluation community.

8 From the presentations and discussions at the PCMDI workshop and during the months  
9 following, the initial tenets, as well as the name of the activity, were developed [*Teixeira et al.*,  
10 2011]. Consensus was reached on: 1) the use of the CMIP5 model output list of variables [*Taylor*  
11 *et al.*, 2009] as a means to define which satellite variables would be considered for inclusion, 2)  
12 the need for a “technical note” for each variable that would describe the origins, algorithms,  
13 validation/uncertainty, guidance on methodologies for applying the data to model evaluation,  
14 contact information, relevant references, etc. and that would be limited to a few pages targeting  
15 users who might be unfamiliar with satellites and models, 3) having the observation data  
16 technically aligned with the CMIP model output [i.e., CMIP’s specific application of the NetCDF  
17 Climate and Forecast (CF) Metadata Conventions], and 4) hosting the observations on the Earth  
18 System Grid Federation (ESFG) of archive nodes so that they would appear side by side with the  
19 model output. The name “obs4MIPs” was suggested to uniquely identify the data in the ESGF  
20 archive and distinguish it from the diversity of other information hosted there.

21 Along with outlining the initial objectives and tenets of the pilot effort, a first set of about a  
22 dozen NASA satellite observation datasets was identified and deemed particularly appropriate for  
23 climate model evaluations relevant to CMIP and associated IPCC assessment reports, based on  
24 their maturity and long-standing community use. The initial set included temperature and  
25 humidity profiles from the Atmospheric InfraRed Sounder (AIRS) and the Microwave Limb  
26 Sounder (MLS), ozone profiles from the Tropospheric Emission Spectrometer (TES), sea surface  
27 height (SSH) from TOPEX/Jason (joint with CNES - Centre National d’Etudes Spatiales), sea  
28 surface temperature (SST) from the Advanced Microwave Sounder Radiometer-E (AMSR-E, joint  
29 with JAXA – Japanese Aerospace Exploration Agency), shortwave and longwave all-sky and  
30 clear-sky radiation fluxes at the top of the atmosphere from the Cloud and Earth Radiation Budget

1 Experiment Satellite (CERES), and cloud fraction from MODIS, and column water vapor from  
2 the Special Sensor Microwave Imager (SSM/I). All these initial datasets were global, or nearly so,  
3 and had monthly time resolution spanning record lengths between 8 and 19 years. By late 2011  
4 these datasets were archived, with their associated technical notes, on the JPL ESGF node. Further  
5 information on the development and scope of the obs4MIPs effort during this period was captured  
6 in Teixeira et al. [2014].

7 With the success of this pilot effort, NASA and DOE sought to broaden the activity and engage  
8 more satellite teams and agencies by establishing an obs4MIPs Working Group early in 2012 that  
9 included members from DOE, three NASA centers and NOAA. In the subsequent year, this  
10 working group helped identify and shepherd a number of additional datasets into the obs4MIPs  
11 project. These included ocean surface wind vectors and speed from QuikSCAT, precipitation from  
12 the Tropical Rainfall Mapping Mission (TRMM) and the Global Precipitation Climatology Project  
13 (GPCP), aerosol optical depth from the Moderate Resolution Imaging Spectroradiometer  
14 (MODIS) and the Multi-angle Imaging SpectroRadiometer (MISR), aerosol extinction profiles  
15 from Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO), surface  
16 radiation fluxes from CERES, and sea ice from the National Snow and Ice Data Center (NSIDC).  
17 Two of the datasets included higher frequency sampling, with TRMM providing both monthly and  
18 3-hourly values, and GPCP providing both monthly and daily values.

19 All of the datasets contributed to obs4MIPs thus far are gridded products, and many cover a  
20 substantial fraction of the Earth. Most of the data discussed above was provided on a 1 degree x  
21 1 degree (longitude x latitude) grid which was an appropriate target for the CMIP5 generation of  
22 models. More recently, data is being included at the highest gridded resolution available rather  
23 than mapping it to another grid. Calculation of monthly averages, which may be nontrivial  
24 especially for data derived from polar orbiting instruments, is determined on a case-by-case basis  
25 and is described in the Tech Note of each product. Most of the products that have been introduced  
26 into obs4MIPs to-date are based on satellite measurements, but other gridded products based on  
27 in-situ measurements are envisioned to become a part of an expanding set of gridded products  
28 available via obs4MIPs. Ongoing discussions include the possibility of also including some in-  
29 situ data.

30



1

## 2 **3. Expanding the Scope and Contributions**

3 Since its inception, obs4MIPs has continually engaged the climate modeling and model  
4 evaluation communities and endeavored to make them aware of its progress. Awareness and  
5 community support were fostered in part through the publications and workshops mentioned above  
6 [*Gleckler et al.*, 2011; *Teixeira et al.*, 2014; *Ferraro et al.*, 2015], as well as through the WCRP  
7 and the Committee on Earth Observing Satellites (CEOS) and the Coordination Group of  
8 Meteorological Satellites (CGMS) through their Joint Working Group on Climate (JWGC)<sup>2</sup>. The  
9 JWGC published in 2017 an inventory<sup>3</sup> integrating information on available and planned satellite  
10 datasets from all CGMS and CEOS agencies. The inventory is updated annually and serves as one  
11 resource of candidate datasets that might be suitable for obs4MIPs. Based on overlapping interests,  
12 the first international contributions to obs4MIPs were cultivated from the Climate Feedback Model  
13 Intercomparison Project (CFMIP) and the European Space Agency (ESA) through its Climate  
14 Change Initiative (*Hollmann et al.*, 2013) and its Climate Model User Group (CMUG).

15 CFMIP<sup>4</sup> was established, through leadership from the UK Met Office, the Bureau of  
16 Meteorology Research Centre (BMRC) and Le Laboratoire de Météorologie Dynamique (LMD),  
17 in 2003 as a means to bring comprehensive sets of observations on clouds and related parameters  
18 to bear on the understanding of cloud-climate feedback and its representation in climate models.  
19 In addition to the modelling experiments, a deliberate and systematic strategy for archiving the  
20 satellite data relevant to the CFMIP effort was developed and implemented [See *Tsushima et al.*,  
21 2017; *Webb et al.*, 2017 for recent summary information], and it was aligned with the obs4MIPs  
22 strategy and goals. Crucially, this alignment included the use of CF-compliant format, hosting the  
23 data on the ESGF, and having a focus on observed quantities and diagnostics that are fully  
24 consistent with outputs from the CFMIP Observations Simulator Package (COSP ; *Bodas-Salcedo*  
25 *et al.* 2011) for the evaluation of clouds and radiation in numerical models. Based on this relatively

---

<sup>2</sup> <http://ceos.org/ourwork/workinggroups/climate/>

<sup>3</sup> <https://climatemonitoring.info/ecvinventory/>

<sup>4</sup> <http://cfmip.metoffice.com> and <http://climserv.ipsl.polytechnique.fr/cfmip-obs/>

1 close alignment, CFMIP provided over 20 satellite-based observed quantities as contributions to  
2 obs4MIPs. These include a number of cloud and aerosol variables from the CALIPSO, CloudSat  
3 and the Polarization & Anisotropy of Reflectances for Atmospheric Sciences coupled with  
4 Observations from a Lidar (PARASOL) satellite missions as well as the International Satellite  
5 Cloud Climatology Project (ISCCP).

6 ESA established the "Climate Modeling User Group" (CMUG)<sup>5</sup> to provide a climate system  
7 perspective at the center of its Climate Change Initiative (CCI)<sup>6</sup> and to host a dedicated forum  
8 bringing the Earth observation and climate modeling communities together. Having started at  
9 approximately the same time as obs4MIPs with overlapping goals, communication between the  
10 two activities was established at the outset. Through the CCI, a number of global datasets were  
11 being produced that overlapped with the model evaluation goals of obs4MIPs, and CMUG/CCI  
12 succeeded in making early contributions to obs4MIPs. These included an SST product developed  
13 from the Along Track Scanning Radiometers (ATSR) aboard ESA's ERS-1, ERS-2 and Envisat  
14 satellites, specifically the ATSR Reprocessing for Climate (ARC) product, as well as the ESA  
15 GlobVapour project merged MERIS and EUMETSAT's SSM/I water vapor column product.

16 The growing international and multi-agency interest in obs4MIPs and its initial success meant  
17 there was potential to broaden the support structure of obs4MIPs and further expand international  
18 involvement. The establishment of the WCRP Data Advisory Council (WDAC)<sup>7</sup> in late 2011  
19 provided a timely opportunity to foster further development. During 2012, as the WDAC  
20 developed its priorities and identified initial projects to focus on, obs4MIPs was proposed as an  
21 activity that could contribute to the objectives of the WDAC and could be served by WDAC  
22 oversight and promotion. Based on this proposal and ensuing discussions, a WDAC Task Team  
23 on Observations for Model Evaluation (subsequently here, simply "the Task Team") was formed  
24 in early 2013. The terms of reference for the Task Team included: 1) establishing data and  
25 metadata standards for observational and reanalysis datasets consistent with those used in major  
26 climate model intercomparison efforts, 2) encouraging the application of these standards to  
27 observational datasets with demonstrated utility for model evaluation, 3) eliciting community input

---

<sup>5</sup> <http://www.esa-cmug-cci.org>

<sup>6</sup> <http://cci.esa.int>

<sup>7</sup> <http://www.wcrp-climate.org/wdac-overview>

1 and providing guidance and oversight to establish criteria and a process by which candidate  
2 obs4MIPs datasets might be accepted for inclusion, 4) assisting in the coordination of obs4MIPs  
3 and related observation-focused projects (e.g. CFMIP, CREATE-IP – formerly ana4MIPs), 5)  
4 overseeing an obs4MIPs website<sup>8</sup>, 6) recommending enhancements that might be made to ESGF  
5 software to facilitate management of and access to such projects, 7) coordinating the above  
6 activities with major climate model intercomparison efforts (e.g., CMIP) and liaising with other  
7 related WCRP bodies, such as WCRP’s Model Advisory Council (WMAC), including recommend  
8 additions and improvements to CMIP standard model output to facilitate observation-based model  
9 evaluation. Membership of the Task Team<sup>9</sup> draws on international expertise in observations, re-  
10 analyses, and climate modeling and evaluation, as well as program leadership/connections to major  
11 observation-relevant agencies (e.g. ESA, EUMETSAT, NASA, NOAA, DOE).

12 One of the first activities undertaken by the Task Team was to organize a meeting of experts  
13 in satellite data products and global climate modeling for the purpose of planning the evolution of  
14 obs4MIPs in support of CMIP6 [Ferraro *et al.*, 2015]. The meeting, held in late spring of 2014 at  
15 NASA Headquarters, was sponsored by DOE, NASA and WCRP. It brought together over 50  
16 experts in both climate modeling and satellite data from the United States, Europe, Japan, and  
17 Australia. The objectives for the meeting included the following: 1) review and assess the  
18 framework, working guidelines, holdings, and ESGF implementation of obs4MIPs in the context  
19 of CMIP model evaluation, 2) identify underutilized and potentially valuable satellite observations  
20 and reanalysis products for climate model evaluation, in conjunction with a review of CMIP model  
21 output specifications, and recommend changes and additions to datasets and model output to  
22 achieve better alignment, 3) provide recommendations for new observation datasets that target  
23 critical voids in model evaluation capabilities, including important phenomena, subgrid-scale  
24 features, higher temporal sampling, in-situ and regional datasets, and holistic Earth system  
25 considerations (e.g. carbon cycle, composition).

---

<sup>8</sup> <https://www.earthsystemcog.org/projects/obs4mips/>

<sup>9</sup> <https://esgf-node.llnl.gov/projects/obs4mips/governance/>

1        Apart from recommendations of specific datasets to include in obs4MIPs in preparation for  
2 CMIP6, there were several consensus recommendations that have driven subsequent and recent  
3 obs4MIPs developments and expansion activities. These included:

- 4        ● Expand the inventory of datasets hosted by obs4MIPs.
- 5        ● Include higher-frequency datasets and higher-frequency model output.
- 6        ● Develop a capability to accommodate reliable and defensible uncertainty measures.
- 7        ● Include datasets and data specification support for datasets involving offline simulators.
- 8        ● Consider hosting reanalysis datasets in some fashion but with appropriate caveats.
- 9        ● Include gridded in situ datasets and consider other in-situ possibilities.
- 10       ● Provide more information on the degree of correspondence between model and observations.

11 For more details on the discussion and associated recommendations, see Ferraro et al. [2015]. In  
12 the following section, we highlight the considerations and progress that have been made toward  
13 these and other recommendations for expanding and improving obs4MIPs.

## 14        **4. Improvements and Implementation Status for CMIP6**

15        With the recommendations of the planning meeting in hand and with CMIP6 imminent, a  
16 number of actions were taken by the obs4MIPs Task Team and the CMIP Panel (a WCRP group  
17 that oversees CMIP). For the most part, these have provided the means to widen the inventory, to  
18 make the process of contributing datasets to obs4MIPs more straightforward, and to develop  
19 additional features that benefit the users.

### 20        **a) Additional obs4MIPs Data Sets**

21        CMIP6-Endorsed MIPs were required to specify the model output they needed to perform  
22 useful analyses (Eyring et al., 2016), and these formed what is now the CMIP6 data request (Juckes  
23 et al. 2019). The obs4MIPs Task Team responded by encouraging/promoting a wider range of  
24 observation-based datasets and released a solicitation for new datasets in the fall of 2015 that added  
25 emphasis on higher frequency, as well as basin- to global-scale gridded in-situ data. The  
26 solicitation also placed a high priority on data products that might be of direct relevance to the

1 CMIP6-endorsed model intercomparison projects<sup>10</sup>. The outcomes of the solicitation and status  
2 of the obs4MIPs holdings are described below.

3 As of August 2019, the holdings for obs4MIPs<sup>11</sup> include over 80 observational datasets<sup>12</sup>. The  
4 datasets include contributions from NASA, ESA, CNES, JAXA, and NOAA, with the data being  
5 hosted at a number of ESGF data nodes, including LLNL/PCMDI, IPSL, GSFC/NASA,  
6 GFDL/NOAA, British Atmospheric Data Centre (BADC), and German Climate Computing  
7 Center (DKRZ). Along with the previously discussed datasets, there are additional SST and water  
8 vapor products, and outgoing longwave radiation (OLR) and sea ice datasets. Some of these  
9 include both daily and monthly sampled data.

10 There are a number of datasets that have been provided through the ESA CCI effort, including  
11 aerosol optical thickness contribution from the ATSR-2 and AATSR missions, ocean wind speed  
12 from SSM/I, total column methane and CO<sub>2</sub> from ESA, and a near surface, ship-based CO<sub>2</sub> product  
13 from the Surface Ocean CO<sub>2</sub> Atlas (SOCAT); the latter three are particularly important for the  
14 carbon cycle component of Earth System models. A new and somewhat novel dataset is expected  
15 to be contributed which will provide regional OLR data based on the Geostationary Earth  
16 Radiation Budget (GERB) instrument aboard EUMETSAT's geostationary operational weather  
17 satellites. In this case, the data coverage is for Europe and Africa only but with sampling that  
18 resolves the diurnal cycle.

19 In the fall of 2015, the Task Team raised awareness of obs4MIPs by explicitly inviting the  
20 observational community to contribute to obs4MIPs. The call, which was communicated by  
21 WCRP and through other channels, set the end of March 2016 as the deadline for submission. The  
22 call made explicit the desire to include observational datasets that had a regional focus, provided  
23 higher frequency sampling, and in particular were aligned with CMIP6 experimentation and model  
24 output [Eyring *et al.*, 2016]. The response to this call resulted in proposals for nearly 100 new  
25 datasets, with several notable new contribution types. This includes proposals for a number of in-  
26 situ gridded products, merged in-situ and satellite products, and regional datasets. Examples  
27 include global surface temperature, multivariate ocean and land surface fluxes, sea ice and snow,

---

<sup>10</sup> <http://www.wcrp-climate.org/modelling-wgcm-mip-catalogue/modelling-wgcm-cmip6-endorsed-mips>

<sup>11</sup> [www.earthsystemcog.org/projects/obs4mips/](http://www.earthsystemcog.org/projects/obs4mips/)

<sup>12</sup> Not all datasets may be visible on the ESGF unless all nodes are on line.

1 ice sheet mass changes, ozone, complete regional aggregate water and energy budget products,  
2 soil moisture, cloud, aerosol, temperature and humidity profiles, surface radiative flux, and  
3 chlorophyll concentrations.

4 Not long after polling the observational community about possible additions to obs4MIPs,  
5 efforts began in earnest within the CMIP community to dramatically expand the CMIP5 model  
6 output lists for CMIP6. This expansion was primarily driven by the more comprehensive  
7 experimental design for CMIP6 and desire for more in-depth model diagnosis, and secondarily by  
8 the greater availability of observations. It soon became clear that despite risks of slowing the  
9 momentum of obs4MIPs it was better to postpone the inclusion of new datasets until the data  
10 standards for CMIP6 were solidified. This took more than two years (given CMIP6's scope and  
11 complexity), and only when that effort was largely completed in late 2017 was it possible to begin  
12 working to ensure that obs4MIPs data standards would remain technically close to those of CMIP.

### 13 **b) Obs4MIPs Data Specifications (ODS)**

14 The primary purpose of obs4MIPs is to facilitate comparison of observational data to model  
15 output from WCRP intercomparison projects, notably CMIP. To accomplish this, the organization  
16 of CMIP and obs4MIPs data must be closely aligned, including the data structure and metadata  
17 requirements and how they are ingested to the Earth System Grid Federation (ESGF)  
18 infrastructure, which is relied on for searching and accessing the data. The original set of  
19 obs4MIPs dataset contributions adhered to guidelines (ODS V1.0, circa 2012) that were based on  
20 the CMIP5 data specifications. Now, the obs4MIPs data specifications have been refined to be  
21 largely consistent with the CMIP6 data specifications, which will not change until the community  
22 begins to configure a next generation (CMIP7).

23 Updates to the Obs4MIPs Data Specifications (ODS2.1) include accommodation via global  
24 attributes that allow for unique identification of datasets and associated institutions, source types,  
25 and dataset versions (i.e., types of observations)<sup>13</sup>. In addition, the global attributes are constructed  
26 to facilitate organization of the obs4MIPs datasets, and in particular for providing a useful set of  
27 options (or facets) for data exploration via the ESGF search engine.

---

<sup>13</sup> <https://esgf-node.llnl.gov/projects/obs4mips/DataSpecifications>; Gleckler et al., (2019)

1 Meeting the obs4MIPs (or CMIP6) data requirements is facilitated by using the Climate Model  
2 Output Rewriter (CMOR3; Doutriaux et al., 2017)<sup>14</sup>. Use of CMOR3 is not required for producing  
3 obs4MIPs data, but it is strongly recommended because CMOR3 ensures that the necessary  
4 metadata for distributed data searching are included. The version of CMOR used in the initial  
5 phase of obs4MIPs was designed for model output, and some special adaptations were required  
6 when applying it to various gridded observations. Fortunately, during the period while the  
7 CMIP6/obs4MIPs data standards were being developed, important improvements were made to  
8 CMOR3 which included streamlining how it could be used for processing gridded observations.

9 With the updates to ODS2.1 and CMOR3 completed,<sup>15</sup> new and revised datasets are once again  
10 being added to obs4MIPs, and with additional enhancements in place (Section 4c-d), that effort is  
11 expected to be the main priority for obs4MIPs throughout the research phase of CMIP6. For data  
12 providers interested in contributing to obs4MIPs, please see “How to Contribute” on the obs4MIPs  
13 website<sup>16</sup>. Efforts to further improve the process, as well as additional considerations for future  
14 directions are discussed in Section 5.

### 15 **c) Obs4MIPs Dataset Indicators**

16 Obs4MIPs has implemented a set of dataset indicators that provide information on a dataset's  
17 technical compliance with obs4MIPs standards and its suitability for climate model evaluation.  
18 The motivation for including this information is two-fold. First, the indicators provide users with  
19 an overview of key features of a given dataset's suitability for model evaluation. For example, does  
20 the dataset adhere to the key requirements of obs4MIPs (e.g. having a technical note and adhering  
21 to the obs4MIPs data specifications that is required to enable ESGF searching)? Similarly, are  
22 model and observation comparisons expected to be straightforward (e.g. is direct comparison with  
23 model output possible or will it require the use of special coding applied to the model output to  
24 make it comparable)? Another relevant consideration is the degree to which the dataset has  
25 previously been used for model evaluation and whether publications exist that document such use.  
26 Second, the indicators allow for a wider spectrum of observations to be included in obs4MIPs. In

---

<sup>14</sup> <https://cmor.llnl.gov>

<sup>15</sup> <https://github.com/pcmdi/obs4mips-cmor-tables>

<sup>16</sup> <https://esgf-node.llnl.gov/projects/obs4mips/HowToContribute>

1 the initial stages of obs4MIPs, only relatively mature datasets – those already widely adopted by  
2 the climate model evaluation community – were considered acceptable. While this helped ensure  
3 the contributions were relevant for model evaluation, it also limited the opportunity for other or  
4 newer datasets to be exposed for potential use in model evaluation.

5 The establishment of the indicators will facilitate the monitoring and characterization of the  
6 increasingly broad set of obs4MIPs products hosted on the ESGF and will guide users in  
7 determining which observational datasets might be best suited for their purposes. There are six  
8 indicators grouped into three categories: two indicators are associated with obs4MIPs technical  
9 requirements, three indicators are related to measures of dataset maturity and suitability for climate  
10 model evaluation, and one indicator is a measure of the comparison complexity associated with  
11 using the observation for model evaluation. These indicators, grouped by these categories, along  
12 with their potential values are given in Figure 2 (upper). Each of the values is color coded so that  
13 the indicators can be readily shown in a dataset search as illustrated by Figure 2 (lower). In the  
14 present framework, still to be fully exercised, the values of the indicators for a given dataset are  
15 intended to be assigned, in consultation with the dataset provider, by the obs4MIPs Task Team.  
16 Note that the values of the indicators can change over time as a dataset and/or its use for model  
17 evaluation matures or as the degree to which the dataset aligns with obs4MIPs technical  
18 requirements improves. To accommodate this, the values of the indicators will be version-  
19 controlled via the obs4MIPs Github repository. Additional information on the indicators and how  
20 they are assigned can be found on the obs4MIPs website. In brief, these indicators are meant to  
21 serve as an overall summary, using qualitative distinctions, of a dataset’s suitability for climate  
22 model evaluation. They do not represent an authoritative or in-depth scientific evaluation of  
23 particular products as attempted by more ambitious and comprehensive efforts such as the  
24 GEWEX Data and Analysis Panel (GDAP) (e.g. Schroeder et al. 2019).

#### 25 **d) Obs4MIPs Dataset Supplemental Information**

26 As a result of the obs4MIPs-CMIP6 meeting in 2015 [*Ferraro et al. 2015*], many data  
27 providers and users made the case that obs4MIPs should accommodate optional inclusion of  
28 ancillary information with a dataset. Ancillary information might include quantitative uncertainty  
29 information, codes that provide transfer functions or forward models to enable a closer comparison



1 between models and observations, the ability to include data flags, verification data, additional  
2 technical information, etc. Note that with the new obs4MIPs data specifications, “observational  
3 ensembles” (which provide a range of observationally-based estimates of a variable that might  
4 result from reasonable processing choices of actually measured quantities) are accommodated as  
5 a special dataset type and are not relegated to Supplemental Information. The inclusion of  
6 Supplemental Information for an obs4MIPs dataset is optional, and the provision for  
7 accommodating such information is considered a "feature" of the current framework of obs4MIPs  
8 (see Example in Section 4e). In the future, there may be better ways to accommodate such  
9 information, as one particular limitation is that the Supplemental Information is not searchable  
10 from the ESGF search engine, although its existence is readily apparent and accessible once a  
11 particular dataset is located via a search. Additional information for data providers on how to  
12 include supplementary information is available on the obs4MIPs website.

### 13 **e) Example Datasets and Model and Observation Comparison**

14 Here we illustrate how the obs4MIPs conventions and infrastructure are applied using CERES  
15 outgoing longwave radiation and TES ozone. First, following the obs4MIPs data specifications  
16 (*ODS2.1; Section 4b*), data contributors provide some basic “registered content” (RC; see footnote  
17 14) which includes a “source\_id”, identifying the common name of the data set (e.g., CERES) and  
18 version number (e.g., v4.0). The source\_id (CERES-4-0) identifies at a high level the dataset  
19 version, which in some cases (as with CERES) applies for more than one variable. Another  
20 attribute is “region” which for CERES is identified as “global”. Controlled vocabulary (CV)  
21 provides many options for the region attribute as defined by the CF-conventions. Yet another  
22 example is the “Nominal Resolution”, providing an approximate spatial resolution which in the  
23 case of the CERES-4-0 data is “1x1 degree”. These and other attributes defined by ODS2.1 are  
24 included as search facets on the obs4MIPs website. Details of how these and other metadata  
25 definitions are described in detail on the obs4MIPs website.

26 Once the data (uniquely identified via “source\_id”) is registered on the obs4MIPs Github  
27 repository (footnote 15), the obs4MIPs task team works with the data provider to agree on a set of  
28 dataset indicators. In the case of the CERES data, the current status of the obs4MIPs data  
29 indicators is [■ ■ ■ ■ ■ ■ ■ ■ ■ ■]. The color coding is described in Section 4.c and refinements will be

1 posted on the obs4MIPs website<sup>17</sup>. As discussed above, these qualitative indicators provide an  
2 overall summary of a dataset’s suitability for climate model evaluation.

3 As described in Section 4.d, a new feature of obs4MIPs permits data providers to include  
4 Supplemental Information (SI). These data/metadata are “free-form” in that they might not adhere  
5 to any obs4MIPs or other conventions. When a user finds data via an ESGF/CoG search, SI  
6 information, if available, will be accessible adjacent to the data indicators and technical note.  
7 Figure 2 provides an example of this for the TES O3 data. And finally, Figures 3 and 4 show  
8 sample results from two model evaluation packages used in CMIP analyses [*Eyring et al., 2016b*  
9 *and Gleckler et al., 2016*], with other examples of obs4MIPs data being used in the literature (e.g.,  
10 Covey et al., 2016; Tian, B., and X. Dong, 2020)

### 11 **f) Intersection with CMIP6 Model Evaluation Activities**

12 Initially, the primary objective of obs4MIPs was to enable the large and diverse CMIP model  
13 evaluation community to obtain better access to and supporting information on useful  
14 observational datasets. Obs4MIPs as an enabling mechanism continues to be the primary  
15 objective, however it is now evident that there is added value beyond its original intent. In addition  
16 to providing data for researchers, obs4MIPs will be a critical link in support of current community  
17 efforts to develop routine and systematic evaluation [*e.g., Gleckler et al., 2016; Eyring et al.,*  
18 *2016a,b, Righi et al. 2019, Eyring et al., 2019; Phillips et al., 2014; Lee et al. 2018; Collier et al.,*  
19 *2018*]. With the rapid growth in the number of experiments, models and output volumes, these  
20 developing evaluation tools promise to produce a first-look, high-level set of evaluation and  
21 characterization summaries, well ahead of the more in-depth analyses expected to come from the  
22 climate research community. As CMIP6 data volumes are expected to grow to tens of petabytes,  
23 increasingly some model evaluation will likely take place where the data resides. These server-  
24 side evaluation tools will rely on observational data provided via obs4MIPs.

---

<sup>17</sup> <https://esgf-node.llnl.gov/projects/obs4mips/DatasetIndicators>

## 5. Summary and Future Directions

This article summarizes the current status of obs4MIPs in support of CMIP6, including the number and types of new datasets, and the new extensions and capabilities that will facilitate providing and using obs4MIPs datasets. Notable highlights include: 1) the recent contribution of over 20 additional datasets making the total number of datasets about 100, with about 100 or more resulting from the 2016 obs4MIPs data call that are ready for preparation and inclusion, 2) updated obs4MIPs Data Specifications that parallel, for the observations, the changes and extensions made for CMIP6 model data, 3) an updated CMOR3 package to give observation data providers a ready and consistent means for dataset formatting required for publication on the ESG, 4) a set of dataset indicators providing a quick accounting and assessment of a dataset's suitability and maturity for model evaluation, and 5) a provision for including supplementary information for a dataset, information that isn't accommodated by the standard obs4MIPs file conventions (e.g. code, uncertainty information, ancillary data). A number of these capabilities and directions were fostered by the discussions and recommendations in the 2014 obs4MIPs meeting [Ferraro *et al.*, 2015].

It is worth highlighting that a number of the features mentioned above, particularly the dataset indicators, have been implemented to allow a broader variety of observations - in terms of dataset maturity, alternatives for the same geophysical quantity, and immediate relevance for climate model evaluation - to be included. Specifically, in the initial stages of obs4MIPs, the philosophy was to try to identify the "best" dataset for the given variable and/or focus only on observations that had been widely used by the community. More recently, guided by input from the 2015 obs4MIPs meeting and consistent with community model evaluation practices, it was decided that having multiple observation datasets of the same quantity (e.g., datasets derived from different satellites or based on different algorithm approaches) was a virtue. Moreover, as models add complexity and new output variables are produced, and as new observation datasets become available, it may take time to determine how to best use a new observation dataset for model evaluation. In this case, rather than waiting to include a dataset in obs4MIPs while ideas were being explored, it was decided that obs4MIPs could facilitate the maturation process and benefit the model evaluation enterprise better by including any dataset that holds some promise for model

1 evaluation as soon as a data provider is willing and able to accommodate the dataset preparation  
2 and publication steps.

3 Additional considerations being discussed by the obs4MIPs Task Team are the requirements  
4 for assignments of DOIs to the datasets and how to facilitate this process. An important step has  
5 been made as it may be possible to provide DOI's via the same mechanism adopted by CMIP6  
6 [Stockhause and Lautenschlager, 2017] and input4MIPs [Durack et al., 2018]. In addition, there  
7 is discussion about how often to update and/or extend datasets and whether or not to keep old  
8 datasets once new versions have been published. Here, a dataset "extension" is considered as  
9 adding new data to the end of the time series of data with no change in the algorithms, whereas a  
10 dataset "update" involves a revision to the algorithm. At present, the guidance from the Task  
11 Team is to extend the datasets, if feasible, with every new year of data, and if an update is provided,  
12 this would formally represent a new version of the dataset with the previous one(s) remaining a  
13 part of the obs4MIPs archive. The Task Team also has undertaken considerable deliberations on  
14 how to handle reanalysis datasets, given that they often serve as an observational reference for  
15 model evaluation applications. Initially, the archive contained a selected set of variables from the  
16 major reanalysis efforts reformatted to adhere to the same standards as obs4MIPs. This data  
17 remains available in the ESGF archive and is designated analysis for Model Intercomparison  
18 Project (ana4MIPs). The data set is static and not updated as new data become available. A new  
19 initiative called the Collaborative REAnalysis Technical Environment (CREATE) (Potter et al.  
20 2018) is curating recent and updated reanalysis data for intercomparison and model evaluation  
21 purposes. The CREATE project offers an expanded variable list relative to ana4MIPs and is  
22 updated with the newest available data as it is produced by the reanalysis centers. The key variables  
23 are offered for most variables at 6 hour, monthly and for precipitation, daily time resolution. The  
24 service also contains a reanalysis ensemble and spread designated as the Multiple Reanalysis  
25 Ensemble version 3 (MRE3).

26 Finally, obs4MIPs' growing capabilities for accommodating a greater number and broader  
27 range of datasets is pointing towards adoption of the obs4MIPs framework for hosting in-situ  
28 datasets that have value for climate model evaluation. In fact, a likely emphasis of future  
29 obs4MIPs Task Team efforts will be to develop an approach to accommodate in situ data. This  
30 potential widening of scope in turn suggests the possibility for using the obs4MIPs framework to

1 serve the function of curating and providing observation datasets for the monitoring and study of  
2 a more extensive range of environmental processes and phenomena, not specifically focusing on  
3 climate model evaluation.

4

5

1 *Acknowledgements*

2 The authors would like to acknowledge the contributions from the observation providers, and  
3 the associated space and research agencies that have made these contributions possible (see  
4 Sections 2-4), as well as the Earth System Grid Federation (ESGF) for providing the platform for  
5 archive and dissemination. The authors would also like to acknowledge the support of the World  
6 Climate Research Programme (WCRP), specifically the WCRP Data Advisory Council (WDAC)  
7 for facilitating the maintenance and expansion of obs4MIPs and providing a task team framework  
8 to carry out these objectives, as well as the WGCM Infrastructure Panel (WIP) for its fundamental  
9 role in developing data protocols, standards and documentation in support of CMIP. Some work  
10 was supported by the U.S. National Aeronautic and Space Administration (NASA) and the U.S.  
11 Department of Energy by Lawrence Livermore National Laboratory under contract DE-AC52-  
12 07NA27344 under the Office of Science, Climate and Environmental Sciences Division, Regional  
13 and Global Model Analysis Program. DW and RF contributions to this study were carried out on  
14 behalf of the Jet Propulsion Laboratory, California Institute of Technology, under a contract with  
15 the National Aeronautics and Space Administration. OB and JB contributions to this study were  
16 supported by the National Oceanic and Atmospheric Administration through the Cooperative  
17 Institute for Climate and Satellites - North Carolina under Cooperative Agreement  
18 NA14NES432003. We would like to acknowledge the help of Birgit Hassler (DLR) in preparing  
19 the ozone model-observation comparison figures with the ESMValTool, and Jiwoo Lee (PCMDI)  
20 for the comparison figures with the PMP.

21

22 ***Code and data availability.***

23 See <https://esgf-node.llnl.gov/projects/obs4mips/>.

24 ***Author contributions.***

25 DW and PG led the initial drafting of the article. All authors contributed to the development of the  
26 obs4MIPs architecture and implementation progress, as well as the final form of the manuscript.

27 ***Competing interests.***

28 The authors declare that we have no significant competing financial, professional, or personal  
29 interests that might have influenced the performance or presentation of the work described in this  
30 paper.

## 6. References

- 1
- 2 Bodas-Salcedo, A., Webb, M.J., Bony, S., Chepfer, H., Dufresne, J.L., Klein, S.A., Zhang, Y.,  
3 Marchand, R., Haynes, J.M., Pincus, R. and John, V.O., 2011: "COSMOS: Satellite simulation  
4 software for model assessment.", *Bulletin of the American Meteorological Society*, 92(8),  
5 pp.1023-1043, <https://doi.org/10.1175/2011BAMS2856.1>.
- 6 Collier, N., et al., (2018): The International Land Model Benchmarking System (ILAMB): Design  
7 and Theory. *JAMES*, 10, 11, 2731-2754, <https://doi.org/10.1029/2018MS001354>
- 8 Covey, C., P. J. Gleckler, C. Doutriaux, D. N. Williams, A. Dai, J. Fasullo, K. Trenberth, and A.  
9 Berg, 2016: Metrics for the diurnal cycle of precipitation: Toward routine benchmarks for  
10 climate models. *J. Climate*, 29, 4461–4471, <https://doi.org/10.1175/JCLI-D-15-0664.1>.
- 11 Durack, P. J., K. E. Taylor, V. Eyring, S. K. Ames, T. Hoang, D. Nadeau, C. Doutriaux, M.  
12 Stockhause, and P. J. Gleckler (2018), Toward standardized data sets for climate model  
13 experimentation, *Eos*, 99, [doi.org/10.1029/2018EO101751](https://doi.org/10.1029/2018EO101751).
- 14 Doutriaux, C., Taylor, K. E., and D. Nadeau: [CMOR3 and PrePARE Documentation](#), 128 pp,  
15 2017.
- 16 EEA (2015), Overview of reported national policies and measures on climate change mitigation  
17 in Europe in 2015, Information reported by Member States under the European Union  
18 Monitoring Mechanism Regulation Rep., Luxembourg: Publications Office of the European  
19 Union.
- 20 Eyring, V., S. Bony, G. A. Meehl, C. A. Senior, B. Stevens, R. J. Stouffer, and K. E. Taylor (2016),  
21 Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental  
22 design and organization, *Geosci. Model Dev.*, 9, 1937-1958, doi:10.5194/gmd-1939-1937-  
23 2016.
- 24 Eyring, V., and J.-F. Lamarque (2012), Global chemistry-climate modeling and evaluation, *EOS*,  
25 *Transactions of the American Geophysical Union*, 93(51), 539-539.
- 26 Eyring, V., Gleckler, P. J., Heinze, C., Stouffer, R. J., Taylor, K. E., Balaji, V., Guilyardi, E.,  
27 Joussaume, S., Kindermann, S., Lawrence, B. N., Meehl, G. A., Righi, M., and Williams, D.  
28 N., 2016a: Towards improved and more routine Earth system model evaluation in CMIP, *Earth*  
29 *Syst. Dynam.*, 7, 813-830, doi:10.5194/esd-7-813-2016.

1 Eyring, V., Righi, M., Lauer, A., Evaldsson, M., Wenzel, S., Jones, C., Anav, A., Andrews, O.,  
2 Cionni, I., Davin, E. L., Deser, C., Ehbrecht, C., Friedlingstein, P., Gleckler, P., et al. (2016b):  
3 ESMValTool (v1.0) – a community diagnostic and performance metrics tool for routine  
4 evaluation of Earth system models in CMIP, *Geosci. Model Dev.*, **9**, 1747-1802

5 Eyring, V., et al. (2019), ESMValTool (v2.0) – Part 2: an extended set of large-scale diagnostics  
6 for quasi-operational and comprehensive evaluation of Earth system models in CMIP6, GMD,  
7 submitted.

8 Ferraro, R., D. E. Waliser, P. Gleckler, K. E. Taylor, and V. Eyring (2015), Evolving obs4MIPs  
9 To Support Phase 6 Of The Coupled Model Intercomparison Project (CMIP6) Bull. Amer.  
10 Meteor. Soc., Meeting Summary(DOI:10.1175/BAMS-D-14-00216.1), 131-133.

11 Gates, W. L. (1992), AMIP: the Atmospheric Model Intercomparison Project, Bull. Amer. Meteor.  
12 Soc., 73, 1962-1970.

13 Gleckler, P.,J., K. E. Taylor, P. J. Durack, R. Ferraro, J. Baird, S. Finkensieper, S. Stevens, M.  
14 Tuma, and D. Nadeau: The obs4MIPs data specifications 2.1, Manuscript in preparation.

15 Gleckler, P., R. Ferraro, and D. E. Waliser (2011), Better use of satellite data in evaluating climate  
16 models contributing to CMIP and assessed by IPCC: Joint DOE-NASA workshop; LLNL,  
17 October 12-13, 2010, EOS, Vol. 92( No. 20, ), 172.

18 Gleckler, P., K. E. Taylor, and C. Doutriaux (2008), Performance metrics for climate models, J  
19 Geophys Res-Atmos, 113(D6), doi:D06104  
20 10.1029/2007jd008972.

21 Gleckler, P. J., C. Doutriaux, P. J. Durack, K. E. Taylor, Y. Zhang, and D. N. Williams, E. Mason,  
22 and J. Servonnat (2016), A more powerful reality test for climate models, *Eos*, **97**. doi:  
23 [10.1029/2016EO051663](https://doi.org/10.1029/2016EO051663)

24 Hollmann, R., C.J. Merchant, R. Saunders, C. Downy, M. Buchwitz, A. Cazenave, E. Chuvieco,  
25 P. Defourny, G. de Leeuw, R. Forsberg, T. Holzer-Popp, F. Paul, S. Sandven, S.  
26 Sathyendranath, M. van Roozendaal, and W. Wagner, 2013: The ESA Climate Change  
27 Initiative: Satellite Data Records for Essential Climate Variables. *Bull. Amer. Meteor. Soc.*, **94**,  
28 1541–1552, <https://doi.org/10.1175/BAMS-D-11-00254.1>

29 IPCC (2013), Climate Change 2013: The Physical Science BasisRep., 1535 pp, IPCC, Geneva,  
30 Switzerland.



1 IPCC (2014), Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and  
2 III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change Rep.,  
3 151 pp, IPCC, Geneva, Switzerland.

4 Juckes, M., K. E. Taylor, P. Durack, B. Lawrence, M. Mizielinski, A. Pamment, J-Y Peterschmitt,  
5 M. Rixen, and S. Sénésis, 2019: The CMIP6 Data Request (version 01.00.31), Geoscientific  
6 Model Development, in discussion, doi: 10.5194/gmd-2019-219.

7 Lee, H. a. G., A. and McGibbney, L. and Waliser, D. E. and Kim, J. and Loikith, P. C. and Gibson,  
8 P. B. and Massoud, E. C. (2018). "Regional Climate Model Evaluation System powered by  
9 Apache Open Climate Workbench v1.3.0: an enabling tool for facilitating regional climate  
10 studies." Geoscientific Model Development 11: 4435-4449

11 Lee, S., et al. 2015: "Climate Model Diagnostic Analyzer", IEEE International Conference on Big  
12 Data, Santa Clara, CA, October, 2015.

13 Lee, S., et al. 2017: "Scientific and Educational Applications of Climate Model Diagnostic  
14 Analyzer," IEEE Big Data Congress, Honolulu, HI, June, 2017.

15 Meehl, G. A., C. Covey, T. Delworth, M. Latif, B. McAvaney, J. F. B. Mitchell, R. J. Stouffer,  
16 and K. E. Taylor (2007), The WCRP CMIP3 multi-model dataset: A new era in climate change  
17 research, Bulletin of the American Meteorological Society, 88, 1383-1394.

18 NCA (2014), U. S. National Climate Assessment Rep., 1717 Pennsylvania Avenue, NW, Suite  
19 250, Washington, DC.

20 Phillips, A. S., C. Deser, and J. Fasullo (2014), Evaluating modes of variability in climate models,  
21 Eos, Transactions American Geophysical Union, 95(49), 453-455.

22 Potter, G. L., Carriere, L., Hertz, J. D., Bosilovich, M., Duffy, D., Lee, T., and Williams, D. N.:  
23 Enabling reanalysis research using the collaborative reanalysis technical environment  
24 (CREATE), B. Am. Meteorol. Soc., 99, 677–687, [https://doi.org/10.1175/BAMS-D-17-](https://doi.org/10.1175/BAMS-D-17-0174.1)  
25 0174.1, 2018.

26 Reichler, T., and J. Kim (2008), How well do coupled models simulate today's climate?, Bulletin  
27 of the American Meteorological Society, 89(3), 303-+, doi:10.1175/bams-89-3-303.

28 Righi, M., Andela, B., Eyring, V., Lauer, A., Predoi, V., Schlund, M., Vegas-Regidor, J., Bock, L.,  
29 Brötz, B., de Mora, L., Diblen, F., Dreyer, L., Drost, N., Earnshaw, P., Hassler, B., Koldunov,  
30 N., Little, B., Loosveldt Tomas, S., and Zimmermann, K.: ESMValTool v2.0 – Technical

1 overview, *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2019-226>, in  
2 review, 2019.

3

4 Schröder M. et al (2019), The GEWEX Water Vapor Assessment: Overview and Introduction to  
5 Results and Recommendations. *Remote Sens.*, 11(3), 251; <https://doi.org/10.3390/rs11030251>  
6 Stockhause, M. and Lautenschlager, M.: CMIP6 Data Citation of Evolving Data, *Data Science*  
7 *Journal*, 16, 1–13, <https://doi.org/10.5334/dsj-2017-030>, 2017  
8 Stephens, G. L., and D. G. Vane (2007), Cloud remote sensing from space in the era of the A-  
9 Train, *Journal of Applied Remote Sensing*, 1, doi:01350710.1117/1.2709703.

10 Stephens, G. L., et al. (2002), The cloudsat mission and the a-train - A new dimension of space-  
11 based observations of clouds and precipitation, *Bulletin of the American Meteorological*  
12 *Society*, 83(12), 1771-1790.

13 Stouffer, R. J., Eyring, V., Meehl, G. A., Bony, S., Senior, C., Stevens, B., and Taylor, K. E.:  
14 CMIP5 Scientific Gaps and Recommendations for CMIP6, *B Am Meteorol Soc*, 98, 95-105,  
15 2017.

16 Taylor, K. E., R. J. Stouffer, and G. A. Meehl (2009), A Summary of the CMIP5 Experiment  
17 Design, White paper: [http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor\\_CMIP5\\_design.pdf](http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf).

18 Taylor, K. E., R. J. Stouffer, and G. A. Meehl (2012), An Overview of CMIP5 and the Experiment  
19 Design, *Bull. Am. Met. Soc.*, 93, 485–498.

20 Teixeira, J., D. Waliser, R. Ferraro, P. Gleckler, and G. Potter (2011), Satellite observations for  
21 CMIP5 simulations. , *CLIVAR Exchanges. Special Issue on the WCRP Coupled Model*  
22 *Intercomparison Project Phase 5—CMIP5*, 16, 46–47.

23 Teixeira, J., D.E. Waliser, R. Ferraro, P. Gleckler, T. Lee, and G. Potter (2014), Satellite  
24 Observations for CMIP5: The Genesis of Obs4MIPs, *Bull. Amer. Meteor. Soc.*,  
25 <http://dx.doi.org/10.1175/BAMS-D-12-00204.1>.

26 Tian, B., and X. Dong, 2020: The double-ITCZ bias in CMIP3, CMIP5, and CMIP6 models based  
27 on annual mean precipitation. *Geophys. Res. Lett.*, **47**, e2020GL087232,  
28 doi:10.1029/2020GL087232.

29 Tsushima, Y., F. Brient, S. A. Klein, D. Konsta, C. C. Nam, X. Qu, K. D. Williams, S. C.  
30 Sherwood, K. Suzuki, and M. D. Zelinka (2017), The Cloud Feedback Model Intercomparison

1 Project (CFMIP) Diagnostic Codes Catalogue – metrics, diagnostics and methodologies to  
2 evaluate, understand and improve the representation of clouds and cloud feedbacks in climate  
3 models, *Geosci. Model Dev.*, 10(11), 4285-4305, doi:10.5194/gmd-10-4285-2017.

4 Waliser, D., K. W. Seo, S. Schubert and E. Njoku (2007). Global water cycle agreement in the  
5 climate models assessed in the IPCC AR4. *Geophysical Research Letters* 34(16).

6 Waliser, D. E., J. F. Li, C. Woods, R. Austin, J. Bacmeister, J. Chern, A. D. Genio, J. Jiang, Z.  
7 Kuang, H. Meng, P. Minnis, S. Platnick, W. B. Rossow, G. Stephens, S. Sun-Mack, W. K.  
8 Tao, A. Tompkins, D. Vane, C. Walker and D. Wu (2009). Cloud Ice: A Climate Model  
9 Challenge With Signs and Expectations of Progress. *Journal of Geophysical Research*  
10 *CloudSat Special Section*, 114(D00A21).

11 Webb, M. J., T. Andrews, A. Bodas-Salcedo, S. Bony, C. S. Bretherton, R. Chadwick, H. Chepfer,  
12 H. Douville, P. Good, J. E. Kay, S. A. Klein, R. Marchand, B. Medeiros, A. P. Siebesma, C.  
13 B. Skinner, B. Stevens, G. Tselioudis, Y. Tsushima, and M. Watanabe (2017), The Cloud  
14 Feedback Model Intercomparison Project (CFMIP) contribution to CMIP6, *Geosci. Model*  
15 *Dev.*, 10(1), 359-384, doi:10.5194/gmd-10-359-2017.


16 Whitehall, K., et al. (2012), Building model evaluation and decision support capacity for  
17 CORDEX., *WMO Bulletin*, 61, 29-34.

18 Williams, D., V. Balaji, L. Cinquini, S. Denvil, D. Duffy, B. Evans, R. Ferraro, R. Hansen, M  
19 Lautenschlager, and C Trenham, (2016), A Global Repository for Planed-Sized Experiments  
20 and Observations, *Bull. Am. Met. Soc.*, DOI:10.1175/BAMS-D-15-00132.1

21 WorldBank (2011), *Climate change and fiscal policy : A report for APECRep.*, Washington, DC.  
22  
23

# 1 Figures

2  
3



4  
5

6 Figure 1. Two schematics that illustrate key motivations and guiding considerations for obs4MIPs. (left)  
7 Depiction of the large and growing community of scientists undertaking the climate model analysis who  
8 are not necessarily experts in modeling or the details of the observations. (right) Depiction of the large  
9 number of quantities available from model output (e.g. CMIP) and obtained from satellite retrievals,  
10 highlighting that a much smaller subset fall in the intersection but are of greatest relevance to model  
11 evaluation.

12  
13  
14

Technical Requirements		Dataset Suitability and Maturity			Comparison Complexity
Meets obs4MIPs data technical requirements	Includes obs4MIPs technical note information	Closeness or robustness of measurement to observed reference quantity	Maturity with respect to climate model evaluation	Provision for robust uncertainty information	Complexity of Model Observation Comparison
Data suitably processed with CMOR and/or consistent with obs4MIPs standards	Complete technical note information provided	Measurement approach provides a very close relationship to observation quantity	Multiple peer-reviewed examples of application to climate model evaluation	Uncertainty information provided per retrieval/grid point	Comparison can be made directly with CMIP model output variable
Largely complete with minor metadata inconsistencies	Technical note information incomplete and/or could be improved	Measurement approach requires complex and/or non-linear retrieval methods and/or subjective inferences/definitions	One peer-reviewed example of application to climate or component model evaluation.	General uncertainty information given relative to the methodology and dataset as a whole - backed by actual field/in-situ validation exercises	Comparison requires some simple post processing of CMIP output variable(s) (e.g. vertical integral or ratio of two variables)
Non-compliant. Should be removed from database!	Technical note not provided	Measurement approach requires significant use/influence from complex or weakly constrained model and/or has significant ambiguity in definition(s)	No peer-reviewed examples of application to model evaluation	No uncertainty information provided	Comparison requires complex processing of CMIP output (e.g. "simulator", budget calculation)

1

## Obs4MIPs

You are at the CoG-CU node

Home About Us Governance Contact Us Technical Support

**Institute** +

**Instrument** +

**Time Frequency** -

mon (12)

**Realm** -

atmos (12)

**Variable** +

**Variable Long Name** +

**CF Standard Name** +

**Data Node** -

esgf-data.jpl.nasa.gov (12)

Enter Text:

? Search Reset Display 10 results per page [More Search Options](#)

Show All Replicas  
 Show All Versions  
 Search Local Node Only (Including All Replicas)

**Search Constraints:** ✖ esgf-data.jpl.nasa.gov | ✖ atmos | ✖ mon

Total Number of Results: 12  
<< Previous 1 -2-  
Please login to add search results to your Data Cart  
Expert Users: you may display the search URL and return results as XML or return results as JSON

11. **obs4mips.NASA-JPL.TES.tro3.mon**  
Data Node: esgf-data.jpl.nasa.gov  
Version: 20110608  
Total Number of Files (for all variables): 3  
Full Dataset Services: [Show Metadata](#) [List Files](#) [THREDDS Catalog](#) [WGET Script](#) [LAS Visualization](#) [Tech Note](#) [Supplementary Data](#)  
[Globus Download](#) ■ ■ ■ ■ ■


12. **obs4mips.NASA-JPL.QuikSCAT.sfcWind.mon**  
Data Node: esgf-data.jpl.nasa.gov  
Version: 20120411  
Total Number of Files (for all variables): 3  
Full Dataset Services: [Show Metadata](#) [List Files](#) [THREDDS Catalog](#) [WGET Script](#) [LAS Visualization](#) [summary](#) [Globus Download](#)  
■ ■ ■ ■ ■

2

3 Figure 2. (upper) Key to interpretation of obs4MIPs dataset indicators, and(lower) an example of the search  
4 result display of the indicators and links to the [Tech Note] and [Supplementary Data] in the case of datasets  
5 that include those (e.g. TES ozone).


6

# TOA outgoing longwave radiation: DJF (CMIP5 amip: 1981-2005)



1  
2  
3  
4  
5  
6  
7  
8

Figure 3: An illustration of a model-observation comparison using obs4MIPs datasets. This four panel figure shows December-January-February (DJF) climatological mean (1981-2005) results for an individual model (upper left), the CERES-4-0 EBAF dataset (upper right), a difference map of the two upper panels (lower left) and a difference between the CMIP5 Multi-model-mean (MMM) and CERES observations (lower right). The averaging period of the CERES-4-0 DJF mean is 2005-2018. Units are  $W/m^2$ .



1  
2  
3 Figure 4: An illustration of a model-observation comparison using obs4MIPs datasets. Tropospheric ozone  
4 annual cycle calculated from CMIP5 rc4.5 simulations and AURA-TES observations, averaged over the  
5 years 2006-2009, for the NH (left) and SH (right) mid-latitudes (35°-60°) at 250hPa. The individual model  
6 simulations are represented by the different colored lines while AURA-TES is shown as the black line (with  
7 +/- 1 sigma shown in gray).