

Response to editor's comments

We would like to thank you for the in-depth review and constructive comments. Below we provide the point-to-point responses to each comment. Specific comments are given in italic and responses are given in bold. Manuscript with revisions marked using track changes is also provided.

5

Specific Comments

1) Please can you revise your discussions section, to ensure that ALL of the statements you make are backed up with evidence (i.e. at present there are lots of statements which are probably underpinning by your results section, but there is no reference back to the figure etc. which presents that evidence).

10 **Response: Thanks for the comment. The corresponded references and figures are provided.**

2) Please can you also significantly expand the paragraph your have added P25 L9-18. The reviewer sensibly asks for this comparison, but your response is much too light.

15 **Response: Thanks for the comment. This paragraph is a bit reconstructed to include the comparison of biogeochemical fields, the amount of oceanic carbon uptake, and the sensitivity of oceanic carbon uptake. A summary of the comparison of biogeochemical fields between the NESM v3 and IPSL is given and we also add the comparison of oceanic carbon uptake in the historical period between the NESM v3 and CMIP5 models.**

20 *3) Finally, please undertake a (hopefully) final proof-read and edit to ensure that throughout the manuscript the subject of each sentence is completely clear. For example, you have revised the final paragraph of the manuscript to:*

'Overall, compared with both observations and CMIP5 models, the NESM v3 does a good job in simulating ocean biogeochemical fields and oceanic carbon uptake. Despite these model-observation

25 *discrepancies, it is expected that NESM v3 can be used as a useful modeling tool to study interactive feedbacks between the ocean carbon cycle and climate change and the underlying mechanisms.'*

The second sentence says 'Despite these model-observation discrepancies...' but the preceding sentence (and consequently nowhere in the paragraph it) makes no reference to model-observation discrepancies.

Response: Thanks for the comments. We check throughout the manuscript sentence by sentence to make the

writing clear. The publication number in Acknowledgement which is now shown as XXXX, will be given when the manuscript is in final production.

Marine biogeochemical cycling and oceanic CO₂ uptake simulated by the NUIST Earth System Model version 3 (NESM v3)

Yifei Dai^{1, 3}, Long Cao², Bin Wang^{1, 3}

¹ Earth System Modeling Center, and Key Laboratory of Meteorological Disaster of Ministry of Education, Nanjing University of Information Science and Technology, Nanjing 210044, China

² ~~Department~~ Key Laboratory of Atmospheric Sciences GeoScience Big Data and Deep Resource of Zhejiang Province, School of Earth Sciences, Zhejiang University, Hangzhou 310027, China

³ ~~Department of Atmospheric Sciences and~~ China-US joint Atmosphere-Ocean Research Center and International Pacific Research Center, University of Hawaii at Manoa, Honolulu, HI 96822, USA

*Correspondence: Long Cao (longcao@zju.edu.cn)

Abstract. In this study, we evaluate the performance of Nanjing University of Information Science and Technology (NUIST) Earth System Model, version 3 (hereafter NESM v3) in simulating the marine biogeochemical cycle and carbon dioxide (CO₂) uptake. Compared with observations, NESM v3 reproduces reasonably well the large-scale patterns of ~~upper ocean~~ biogeochemical fields in the upper ocean, including nutrients, alkalinity, dissolved inorganic, chlorophyll, and net primary production. Some discrepancies between model simulations and observations are identified and the possible causes are investigated. In the upper ocean, the simulated biases in biogeochemical fields are mainly associated with the shortcoming in simulated ocean circulation. Weak upwelling in the Indian Ocean suppresses the nutrient entrainment to the upper ocean, ~~therefore~~ thus reducing the biological activities and resulting in underestimation of net primary production and chlorophyll concentration. In the Pacific and the Southern Ocean, ~~high nutrient and low chlorophyll~~ nutrients are overestimated as a result from of the strong iron limitation ~~and excessive vertical mixing~~. Alkalinity ~~shows high biases~~ is also overestimated in high-latitude oceans due to ~~the strong~~ excessive convective mixing. The major discrepancy in biogeochemical fields is ~~observed in the deep Northern Pacific. The simulated high concentration of that the model overestimates~~ nutrients, alkalinity, and dissolved inorganic carbon ~~water is too~~ in the deep ~~due to~~ North

Pacific, which is caused by the excessive deep ocean remineralization. The model ~~also~~ reasonably reproduces current-day oceanic CO₂ uptake. Model-simulated ~~total~~ cumulative oceanic CO₂ uptake is 149 PgC ~~from~~ between 1850 ~~to~~ and 2016, which compares well with data-based estimates of 150 ± 20 PgC. In the 1% per year CO₂ increase (1ptCO₂) experiment, the ~~NESM v3 produced~~ diagnosed carbon-climate ($\gamma = -7.9$ PgC/K) and carbon-concentration sensitivity parameters ($\beta = 0.88$ PgC/ppm) in the NESM v3 are comparable with those in Coupled Model Intercomparison Project phase 5 (CMIP5) ~~model results~~ models (β : 0.69 to 0.91 PgC/ppm; γ : -2.4 to -12.1 PgC/K). The ~~nonlinearity of~~ nonlinear interaction between carbon ~~uptake~~ concentration and carbon-climate sensitivity in the NESM v3 accounts for 10.3% of the total carbon uptake, which is within the range of CMIP5 model results (3.6%~10.6%). Overall, the NESM v3 can be employed as a useful modeling tool to investigate large scale interactions between the ocean carbon cycle and climate change.

1 Introduction

The global carbon cycle plays an important role in the climate system. The increase in atmospheric CO₂ is responsible for a large part of the observed increase in global mean surface temperature ([Ciais et al., 2013](#)). From 1750 to 2016, about 645±80 PgC (1 PgC = 10¹⁵ gram carbon) of anthropogenic carbon has been emitted to the atmosphere, including 420±20 PgC from fossil fuels and industry and 225±75 PgC from land-use-change ([Le Quéré et al., 2018](#)). This CO₂ emission caused atmospheric CO₂ concentration to increase by ~~45~~ 48% from an annual mean pre-industrial (PI) value of ~277 parts per million (ppm) ([Joos and Spahni, 2008](#)) to ~~406~~ 409.8 ppm in ~~2017~~ 2019 (~~NOAA ESRL Global Monitoring Division, 2017~~ Dlugokencky and Tans, 2020). As a large carbon reservoir, the global ocean contains more than 50 times the amount of carbon ~~than~~ in the atmosphere ([Denman et al., 2007](#)), and plays a key role in the anthropogenic CO₂ uptake ([Ballantyne et al., 2012](#); [Wanninkhof et al., 2013](#)). Since the year 1870 to 2016, about 25% of anthropogenic CO₂ (about 150±20 PgC) has been absorbed by the ocean ([Le Quéré et al., 2018](#)).

An increase in atmospheric CO₂ perturbs the atmospheric radiative balance and leads to climate change. Changes in atmospheric temperature, precipitation, evaporation, and wind, induce changes in ocean physical properties ~~such as~~ including temperature, salinity, and ocean circulation ([Gregory et al., 2005](#);

[Pierce et al., 2012](#)). These changes in ocean physical properties, in turn, affect the ocean carbon cycle ([Sarmiento and Gruber, 2006](#)). [Friedlingstein et al., \(2006\)](#) proposed that the response of oceanic uptake of atmospheric CO₂ can be represented by the linear sum of two components: 1) carbon-concentration sensitivity, which refers to the response of oceanic CO₂ uptake to increasing atmospheric CO₂; and 2) carbon-climate sensitivity, which refers to the response of oceanic CO₂ uptake to global warming. Adopting this conceptual framework, a number of studies have analyzed the effect of increasing atmospheric CO₂ concentration and global warming on the carbon cycle in terms of the carbon-concentration and carbon-climate sensitivity parameters under different CO₂ emission and concentration scenarios ([Gregory et al., 2009](#); [Boer and Arora, 2009](#); [Tjiputra et al., 2010](#); [Roy et al., 2011](#); [Arora et al., 2013](#); [Arora et al., 2013](#); [Schwinger et al., 2018](#)).

Given the importance of carbon cycle feedback in current and future global climate change, it is necessary to include the representation of the global carbon cycle in climate system models ([Denman et al., 2007](#)). The first and second generations of the NUIST climate system model [have show](#) good skill in simulating internal climate modes, [and](#) global monsoon, ~~and projecting future climate change~~ ([Li et al., 2018](#); [Yang and Wang, 2018a](#); [Yang et al., 2018b](#)). However, the previous generations of NESM do not include an active ocean biogeochemical model and cannot be used to study the ocean carbon cycle ([Cao et al., 2015](#)). Recently, the third version of the NUIST ~~earth system model~~ [Earth System Model](#) (NESM v3) was developed as a newly registered model to [CMIP Coupled Model Intercomparison Project](#) phase 6 (CMIP6, [Cao et al., 2018](#)), ~~which~~; [Eyring et al., 2016](#)). The NESM v3 couples the Pelagic Interactions Scheme for Carbon and Ecosystem Studies [version 2](#) (PISCES v2); [Aumont et al., 2015](#)) to represent the ocean biogeochemical processes ([Aumont et al., 2015](#)).

The objective of this manuscript is to evaluate the performance of NESM v3 in simulating marine carbon-related biogeochemical tracers and oceanic anthropogenic CO₂ uptake. As a newly developed earth system model and a new member of the CMIP community, it is essential to evaluate the model's ability against observational data. First, we analyze the present-day [mean state distributions](#) of macro-nutrients, chlorophyll, net primary production (NPP), sea-air CO₂ flux, dissolved inorganic carbon (DIC), and alkalinity against available observations and ~~observational~~ [observation](#)-based estimates. Then, we evaluate the ~~amount, spatial distribution, and sensitivity parameters of the~~ [model-simulated](#) anthropogenic

~~carbon~~CO₂ uptake. The amount and spatial distribution of oceanic anthropogenic ~~carbon~~CO₂ uptake during the historical period and under future scenarios are compared with observations and CMIP5 ~~model~~models' results. The carbon-concentration and carbon-climate sensitivity ~~parameters and nonlinearity~~ of oceanic CO₂ uptake diagnosed from the NESM v3 are compared with those from CMIP5 models. We also provide a supplementary material that ~~includes comparisons of~~compares biogeochemical fields ~~is~~simulated by the NESM v3 ~~and with that by~~ IPSL-CM5A-LR (hereafter IPSL) ~~that~~, which shares the same marine biogeochemical component (PISCES) as used in NESM v3.

~~The outline of the paper is the following.~~In Section 2, we describe the NESM v3 with a focus on the ocean carbon cycle component, as well as the setup of model simulations. The results of model simulations are analyzed in Section 3. Conclusions and discussions are presented in Section 4.

2 Methods

2.1 Model

2.1.1 Framework of the NESM v3

Detailed descriptions of the physical components, major improvements, and model tuning procedures of the NESM v3 are documented in [Cao et al. \(2018\)](#). Here we give a brief ~~introduction~~review.

NESM v3 consists of three main model components, including European Centre Hamburg Atmospheric Model version 6.3 (ECHAM v6.3) ([Stevens et al., 2012](#); [Giorgetta et al., 2013](#)), Nucleus for European Modeling of the Ocean version 3.4 (NEMO v3.4-revision 3814) ([Madec and NEMO team, 2012](#)) and Los Alamos sea-ice model version 4.1 (CICE v4.1) ([Hunke et al., 2010](#)).

In this study, we use the low-resolution version of ~~the~~NESM v3. The atmospheric resolution is T31L31 ~~which that~~ has a horizontal resolution of $\sim 3.75^\circ$ latitude by 3.75° longitude and 31 layers. The atmospheric model and land surface model are originally adopted from ECHAM v6.3. The detailed ~~information~~description is ~~shown~~provided in [Stevens et al. \(2012\)](#) and [Giorgetta et al. \(2013\)](#). The sea-ice model includes four ice layers and one snow layer with a multi-layer thermodynamic scheme ([Hunke et al., 2010](#); [Cao et al., 2018](#)). ~~Ocean~~The ocean model has the ORCA2 global ocean configuration, ~~which that~~ is a type of tripolar grid. It is based on a 2 degree Mercator mesh and has 31 layers with the thickness

of the ocean layer increasing from 10 m in the upper ocean to 500 m at 5000 m depth. A local transformation is applied in the tropics to refine the resolution to up to 0.5 degree at the equator. In the ocean model, the incoming solar radiation can penetrate to the upper ocean layers as deep as 391 m, and a bio-model penetration parameterization scheme is used to calculate the vertical distribution of solar radiation ([Lengaigne et al., 2009](#)). The ocean background vertical diffusivity is modified in the NESM v3, where the constant value is replaced by latitude-dependent values ([Jochum et al., 2009](#)). The parameterization scheme of ~~the~~ vertical diffusivity is detailed in the supplementary, and supplemental material with the global distribution of vertical diffusivity ~~is also shown (in Fig. S1)~~. Compared to the original vertical diffusivity coefficient constant of 0.12 cm²/s, the coefficients ~~of~~ in the tropical ocean are reduced and that ~~of~~ in the middle subtropical and high latitude oceans are ~~increased, especially in the middle latitude oceans (24°N–33°N and 24°S–33°S)~~ enhanced. Also, the NESM v3 incorporates the parameterization of brine rejection in the ocean model and the reference sea-ice salinity is set as 4 PSU as suggested by [Smith et al \(2010\)](#).

2.1.2 Ocean biogeochemical component

NESM v3 employs the standard PISCES v2 to represent the ocean biogeochemical cycle. The PISCES model is developed from a simple Nutrient-Phytoplankton-Zooplankton-Detritus (NPZD) model ([Aumont et al., 2002](#)). It can be used for both regional and global simulations of lower trophic levels of the marine ecosystem and ocean carbon cycle ([Bopp et al., 2005](#); [Resplandy et al., 2012](#); [Séférian et al., 2013](#)). In the current version, there are 24 prognostic tracers in total, including dissolved inorganic and organic carbon, alkalinity, chlorophyll, and nutrients. We use the same biogeochemical parameters as that used in [Aumont et al. \(2015\)](#). The only exception is the advection scheme for passive tracers. Here we use the Total Variance Dissipation (TVD) formulation instead of the Monotone Upstream Scheme for Conservative Laws (MUSCL) formulation to keep the advection scheme to be consistent with the one used in the physical ocean model. Both TVD and MUSCL schemes have good performance in biogeochemical modeling. The MUSCL scheme has better performance in resolving the small sealesscale processes, while TVD scheme minimizes systematic error through numerical diffusion, and is a better option for coarse-resolution models ([Lévy et al., 2001a](#)).

Two different types of phytoplankton: nanophytoplankton and diatoms, and two size classes of zooplankton: mesozooplankton and microzooplankton, are presented in the model. The life cycle of phytoplankton is regulated by processes of growth, mortality, aggregation, and grazing by zooplankton (Aumont et al., 2015). The growth rate of phytoplankton is determined by temperature, photosynthetically active radiation (PAR), and availability of nutrients, including phosphate, nitrate, silicate, iron, and ammonium. The mortality rate of phytoplankton is set as a constant and is identical for nanophytoplankton and diatoms. The aggregations of nanophytoplankton, which transform the dissolved organic carbon (DOC) to the particular organic matter (POM), only depend on the shear rate, ~~as~~because the main driver of aggregation is the local turbulence. In the NESM v3, this shear rate is set to 1 s^{-1} in the mixed layer and 0.01 s^{-1} below. The same is assumed for diatoms, while the aggregations of diatoms are further enhanced by the nutrients co-limitation. For all species, ~~the~~ phosphate, nitrate, and carbon are linked by a constant Redfield ratio. In ~~the~~ NESM v3, the Redfield ratio of C: N: P is set to be 122:16:1 (Takahashi et al., 1985), and the $-O/C$ ratio is set to 1.34 (Kortzinger et al., 2001). In contrast, the Fe / C, chlorophyll / C, and silicon / C ratio are prognostically simulated by the model based on the external concentrations of the limiting nutrients as in the quota-approach (McCarthy, 1980; Droop, 1983; Aumont et al., 2015). The remineralization of semi-labile DOC can occur in either oxic water or anoxic water depending on the local oxygen concentration, and their degradation rates are specified and identical for oxic respiration and denitrification. Detritus is represented by different types, including POM, calcite, iron particles, and biogenic silica. The POM is partitioned into two size classes: a smaller class (POC: 1-100 μm) and a larger class (GOC: 100-500 μm). The sinking speed of GOC (50-200 m d^{-1}) increases with depth and is much faster than that of POC (3 m d^{-1}). A fraction of phytoplankton would be turned to the POM through the processes of mortality and aggregation. The fate of mortality and aggregation of nanophytoplankton depends on the proportion of the calcifying organisms. For nanophytoplankton, it is assumed that half of the ~~calcifying organisms are associated with the calcifying organisms. Because the density of the calcite is larger than that of organic matter, 50% of the~~ dying calcifiers are routed to the fast-sinking particles. The same is assumed for the mortality of diatoms, and 50% of the dying diatoms are turned to the POM due to the larger density of biogenic silica compared to that of organic matter. The degradation rate of the POM depends on the local temperature with a Q_{10} factor (temperature dependence ratio) of 1.9.

The geochemical boundary condition accounts for the external nutrient supply from five different sources, including atmospheric dust deposition of iron and silicon, river recharge of nutrients, dissolved carbon, and alkalinity, atmospheric deposition of nitrogen, and sediment mobilization of sedimentary iron. In the NESM v3, atmospheric deposition and river recharge are prescribed and sediment mobilization is parameterized. At the bottom of the ocean, different sediment parameterization schemes are applied to biogenic silica, POM, and particulate iron. The amount of permanently buried biogenic silica is assumed to balance the external source, and the burial efficiency of POM is determined by the organic carbon sinking rate at the bottom ~~follows~~ following the algorithm proposed by [Dunne et al. \(2007\)](#). All the particulate iron would be buried into the sediment once they reach the ocean bottom. The amount of the unburied calcite and biogenic silica would dissolve back into the ocean water instantaneously. In this study, the initial conditions of the biogeochemical model have been adopted from World Ocean Atlas 2009 (WOA09, [Garcia, et al., 2010](#)) and Global Ocean Data Analysis Project (GLODAP, [Key et al., 2004](#); [Sabine et al., 2004](#)) dataset.

Carbonate chemistry is formulated based on the Ocean Carbon-Cycle Model Intercomparison Project (OCMIP-2) protocol (more information can be accessed at <http://ocmip5.ipsl.jussieu.fr/OCMIP/>). The quadratic wind-speed formulation proposed by [Wanninkhof \(1992\)](#) is used to compute the air-sea exchange of carbon and oxygen.

2.2 Simulations

In total, there are eight different simulations in this study, including one fully coupled spin-up simulation for 2000 years, one PI-control run (CTRL) for 251 years, three transient runs driven by forcing conditions from historical observational data and shared socio-economic pathway scenarios (SSP5-8.5) -from 1850 to 2100-, ~~(hereafter HistSSP)~~, i.e., fully-coupled HistSSP (FC-HistSSP), biogeochemically-coupled HistSSP (BC-HistSSP), and radiatively-coupled HistSSP (RC-HistSSP), and three idealized 1ptCO₂ runs for 140 years, i.e., FC-~~1%~~1ptCO₂, BC-~~1%~~1ptCO₂, and RC-~~1%~~1ptCO₂. The following is the detailed experimental ~~design~~designs.

First, NESM v3 was spun up for 2000 years with all related parameters set to pre-industrial values (the year 1850), including orbital parameters, land use, aerosol, and greenhouse gas (GHGs) concentration

(284 ppm for CO₂, 790 ppb for CH₄, 275 ppb for N₂O, and 0 ppt for both CFC₁₁ and CFC₁₂). Here, we use PI control simulations of each component model as the initial conditions for the couple model. The atmosphere and sea ice models use the end of a 550-year PI control simulation as the initial conditions, and the ocean model uses the end of a 4000-year PI control simulation as the initial condition. Averaged over). During the last 100 years of the spin-up simulation, the average globally integrated sea-air flux is 1.0 PgC yr⁻¹. The large positive sea-air flux is a result of the 3-dimensional correction of nutrient and alkalinity in the PISCES model (Séférian et al., 2013). The 3-dimensional correction refers to the fact that the global inventory of nutrient and alkalinity are restored toward the observations in January 1st of every year (Aumont et al., 2015). The linear drift of globally integrated sea-air CO₂ flux flux during the last 100 years of the spin-up simulation is 0.0006 PgC yr⁻¹ per year, indicating that a quasi-equilibrium state has been reached for the global ocean carbon cycle. Global mean sea surface temperature (SST) averaged over the last 100 years of ~~2000-year~~the spin-up simulation is 13.1 Celsius (°C) with a linear drift of -0.0001°C per year, and ocean mean temperature is 3.5°C with a linear drift of 0.00016°C per year, indicating that the dynamic ocean component has also reached a quasi-equilibrium state.

Following the protocol of CMIP6 historical and SSP5-8.5 experiment design (Eyring et al., 2016; Jones et al., 2016), starting from the end of the spin-up simulation, the model is further integrated with time-changing conditionsexternal forcings, including GHGs, ozone, aerosol, land use, and solar forcing from 1850 to 2100. From the year 1850 to 2014, GHGs ~~concentration~~concentrations and other forcing conditions are ~~consistent with~~taken from observations, and from the year 2015 to 2100, GHGs concentration and forcing conditions ~~are produced based on~~follow the SSP5-8.5-Meanwhile scenario. Also, we ~~extended the spin-up by~~conducted a 251-years-year (1850-2100) as the PI-control simulation. Also, we performed a similar set of simulations with all forcing conditions to be kept at the pre-industrial levels. Meanwhile, to have a direct comparison with CMIP5 results, ~~i.e., an~~we conducted the idealized 1ptCO₂ ~~run~~simulations, in which atmospheric CO₂ concentration increases at a rate of 1% per year starting from the end state of the ~~pre-industrial~~spin-up simulation with other ~~GHGs concentration~~forcings remaining at the pre-industrial level. ~~The simulation~~These 1ptCO₂ simulations lasted for 140 years until atmospheric CO₂ concentration has quadrupled.

Following Friedlingstein et al., 2006 and Arora et al., 2013, we performed three types of experiments

(biogeochemically-coupled, radiatively-coupled, and fully-coupled) for ~~each scenario~~ [HistSSP and 1ptCO₂ experiments](#) to separate the effect of atmospheric CO₂ and [CO₂-induced](#) global warming on the ocean carbon cycle:

1) BC simulations in which the code of the ocean carbon cycle sees changing atmospheric CO₂, but the code of atmospheric radiation sees a constant pre-industrial concentration of CO₂. In this way, the ocean carbon cycle is only affected by changing atmospheric CO₂, but no direct effect of CO₂-induced warming;

2) RC simulations in which the code of the ocean carbon cycle sees pre-industrial atmospheric CO₂, but the code of atmospheric radiation sees the changing concentrations of atmospheric CO₂. In this way, the ocean carbon cycle is only affected by CO₂-induced warming, but no direct effect of changing atmospheric CO₂.

3) FC simulations in which both the codes of the ocean carbon cycle and atmospheric radiation see the changing concentrations of atmospheric CO₂. In this way, the ocean carbon cycle is affected by changes in both atmospheric CO₂ and CO₂-induced warming.

It is noted that, in the FC-HistSSP, RC-HistSSP, and BC-HistSSP simulations, other forcings (non-CO₂ GHGs, aerosols, and land-use change) still change with time for both the atmospheric and ocean models.

2.3 Evaluation data

Global ocean distribution data of nutrients concentrations, including nitrate, phosphate, and silicate, ~~are is taken~~ from the *World Ocean Atlas 2018* (WOA18, [Garcia, et al., 2018](#)). ~~In this study, we assume the density of the seawater in the model and observations are the same, and then the unit of observations is converted from umol/kg to mmol/m³ by multiplying the modeled density (in the unit of kg/m³).~~

Geographic distributions of DIC, alkalinity, and anthropogenic carbon is taken from the GLODAP v2 ([Key et al., 2015](#); [Lauvset et al., 2016](#)). Both WOA18 and GLODAP v2 data have a horizontal resolution of 1°×1° with 33 levels and represent the climatology in recent decades. In this study, we assume the density of the seawater in the model and observations are the same, and then the unit of observations is converted from umol/kg to mmol/m³ by multiplying the modeled density (in the unit of kg/m³). We compared modeled chlorophyll in recent decades with the SeaWiFS dataset ([NASA Goddard Space Flight Center, 2014](#)), GlobColour merged data ([Maritorena, et al., 2010](#)), and Ocean Colour Climate Change

Initiative (OCCCI) merged data (<http://www.oceancolour.org/>). OCCCI and GlobColour incorporate the same datasets, while their uncertainty information and algorithms are not the same.

~~Mode~~Model-simulated NPP is compared with moderate Resolution Imaging Spectroradiometer (MODIS) estimated marine NPP based on three different algorithms, including the Standard Vertically Generalized Production Model (VGPM), Eppley-VGPM, and the carbon-based Production Model (CbPM). The datasets can be accessed at <http://www.science.oregonstate.edu/ocean.productivity/index.php>. In the VGPM and Eppley-VGPM, NPP is ~~described~~estimated as the product of chlorophyll and photosynthetic efficiencies ([Behrenfeld and Falkowski, 1997a, 1997b](#)), ~~while the~~. Eppley-VGPM emphasizes the photoacclimation effect at high SSTs, i.e., the growth rate is higher at high-temperature regions ([Eppley, 1972](#); [Morel, 1991](#)). In the CbPM, NPP is ~~described~~estimated as the product of carbon biomass and growth rate ([Behrenfeld et al. 2005](#); [Westberry et al. 2008](#)). All ~~the~~ three datasets have a horizontal resolution of $1/12^\circ \times 1/12^\circ$ and cover the period from 2003 to 2014. The distribution of observed surface ocean sea-air CO₂ flux ~~for~~is taken from Takahashi et al (2009) that applies to the reference year of 2000 ~~is taken from Takahashi et al (2009)~~ and has a spatial resolution of 4° latitude by 5° longitude.

To have a direct comparison between the NESM v3 output and corresponding observations, we interpolated all modeled results and observations to a common $1^\circ \times 1^\circ$ grid using the distance-weighted average remapping method, except for the sea-air CO₂ flux. Due to the low resolution of observational sea-air flux, we interpolated the modeled sea-air CO₂ flux to ~~at~~the $4^\circ \times 5^\circ$ grid- used by Takahashi et al. (2009).

2.4 Analysis method

2.4.1 Nutrient decomposition

To ~~figure out whether~~examine the contribution of ocean ~~circulation or dynamics and~~ biogeochemical processes ~~causes the bias of nutrients~~ in the mismatch between model ~~simulation~~results and observations, we ~~decomposing~~decomposed phosphate to its preformed and regenerated components (~~following the~~ method of Weiss et al., 1970; and Duteil et al., 2012). The regenerated phosphate is released through the remineralization processes of organic matter, and the preformed phosphate is the remaining biotically unutilized surface phosphate, which is transported into the ocean interior by ocean circulation. The

regenerated and preformed phosphate are computed as:

$$P_{regenerated} = R_{P:-O_2} \times AOU \quad \text{—————(1)}$$

$$P_{preformed} = P - P_{regenerated} \quad \text{—————(2)}$$

Where AOU is the apparent oxygen utilization, which represents the biological consumption of oxygen.

5 It is computed as the difference between oxygen saturation and simulated oxygen concentration. $R_{P:-O_2}$ represents the oxidation ratio of phosphate and oxygen, which is set to 1/163 in the NESM v3. P represents the simulated phosphate concentration.

2.4.2 Nutrient limitation

10 In the NESM v3, the nutrients limitation coefficient (0~1) is computed from the Michaelis-Menten equation as follow:

$$MM=N / (K+N) \quad \text{—————(3)}$$

Where MM is the Michaelis-Menten coefficient, N is the nutrient concentration, and K is the half-saturation coefficient, which is parameterized based on half-saturation constant and concentrations of nutrients, phytoplankton, and diatoms ([Aumont et al., 2015](#)).

15 We calculated the annual mean nutrient limitation coefficient ~~of~~for each nutrient (phosphate, nitrate, silicate, and iron) and then considered the nutrient with the ~~lowest~~smallest limitation coefficient as the most limiting factor. Temperature and light are assumed to be the most limiting factor when the annual mean nutrient coefficients are greater than 0.9 ~~for all nutrients~~([Moore et al., 2013](#)).

2.4.3 Carbon-concentration and carbon-climate sensitivity parameters

20 Following Arora et al. (2013), we ~~diagnose~~diagnosed the carbon-climate and carbon-concentration sensitivity parameters from two types of experiments performed by a subset of CMIP5 models, i.e., BC simulations and RC simulations.

In the ~~biogeochemically coupled~~BC simulations ~~where the ocean carbon uptake is only affected by changing atmospheric CO₂~~, the relationship between atmospheric CO₂ concentration and sea-air CO₂ flux
25 can be simplified as:

$$\int_0^t F' dt \approx \beta \Delta C_A \quad \text{—————(5+\gamma\Delta T)}$$

(4)

Where F' represents oceanic carbon uptake change in the ~~biogeochemically coupled~~ BC simulation, ΔC_A represents atmospheric CO_2 concentration change, β represents the carbon-concentration sensitivity parameter, γ represents the carbon-climate sensitivity parameter, and ΔT represents surface air temperature change in the BC simulation.

In the ~~radiatively coupled~~ RC simulations where the oceanic carbon uptake is only affected by ~~temperature~~ climate change, the relationship between temperature and sea-air CO_2 flux can be simplified as:

$$\int_0^t F' dt \approx \gamma \Delta T \quad (6) \quad (5)$$

Where F' represents oceanic carbon uptake change in the ~~radiatively coupled~~ RC simulation and ΔT represents surface air temperature change in the RC simulation.

3 Results

3.1 Nutrients

Nutrients play vital roles in the ocean biogeochemical cycle. A lack of nutrients would limit the growth of phytoplankton. Figure 1 compares the model-simulated annual mean spatial distributions of ~~nutrients~~ nutrient concentration averaged over the top 100m ocean with the WOA18 observations from 1985 to 2014, including phosphate (PO_4^{3-}), nitrate (NO_3^-), and silicate (SiO_4^{2-}), ~~averaged over the top 100 m depth from 1985 to 2014 with the WOA18 observations.~~ The model reproduces reasonably well the ~~observed~~ large-scale patterns of upper ocean mean nutrients concentrations with pattern correlation coefficients (PCCs) larger than 0.8 and ~~variability similar to the~~ normalized standard deviations (NSDs, ~~model simulations are normalized by stand deviations of corresponding observations.~~) close to 1.0. The PCCs of nitrate, phosphate, and silicate between model simulation and WOA18 are ~~respectively~~ 0.93, 0.91, and 0.83, ~~respectively~~, and the ~~normalized standard deviations (SDs, normalized to the respective observed standard deviations)~~ NSDs of nitrate, phosphate, and silicate are ~~respectively~~ 1.05, 1.06, and 1.22, ~~respectively~~. For both the NESM v3 simulation and observations, the largest concentrations of phosphate, nitrate, and silicate are observed in the Southern Ocean as a result of strong vertical mixing

and upwelling that bring nutrient-rich deep water to the surface (Whitney, 2011). Also, the strong iron limitation that reduces the biological uptake of the macro-nutrients is one of the main causes of the high macro-nutrients level in the Southern Ocean (de Baar et al., 1990). Relatively high concentrations of nutrients are also simulated in the subarctic Pacific Ocean and the mid-eastern Pacific Ocean. Relatively low concentrations of nutrients are simulated in subtropical regions. The spatial distributions of nutrients in the NESM v3 with high concentrations in the Southern Ocean and low concentrations in the subtropical Pacific are generally consistent with ~~other~~both observations and CMIP5 models results ([Ilyina et al., 2013](#); [Moore et al., 2013](#); [Séférian et al., 2013](#); [Tjiputra et al., 2013](#)). ~~As shown in the supplement, PCCs of nutrients between the NESM v3 and IPSL simulations are all over 0.9.~~

~~For both the NESM v3 simulation and observations, phosphate, nitrate, and silicate in the Southern Ocean have the highest nutrient concentrations. Strong vertical mixing and upwelling bring nutrient rich deep water to the surface (Whitney, 2011). Also, the strong iron limitation (Fig. 4) that reduces the biological uptake of the macro-nutrients is one of the main causes of the high nutrients level in the Southern Ocean. The relatively high concentration of nutrients is simulated in the subarctic Pacific Ocean, and the mid-eastern Pacific Ocean, which is about 50% of the concentration in the Southern Ocean. Relatively low concentration of nutrients, less than 20% of the concentration in the Southern Ocean, is simulated in subtropical regions.~~ Some noticeable discrepancies between model simulations and observations are also found (Fig. ~~a3, b3, c3~~1a3, 1b3, 1c3). Phosphate and nitrate are overestimated in the Southern Ocean and the Pacific Ocean but ~~are~~ underestimated in the Indian Ocean, Subarctic Pacific, and ~~Middle low latitude~~tropical Atlantic. ~~Except for the Indian Ocean and Subarctic Pacific, silicate~~Silicate is overestimated over all of the global ocean, ~~resulting in the high~~ except for the Indian Ocean and Subarctic Pacific. Averaged over the global ~~mean~~upper ocean, model-simulated silicate concentration ~~(is about 50% higher~~greater than ~~that in~~ the WOA18 observations).

Figure 2 shows the ~~zonal mean~~ latitudinal-depth distributions of nutrients from the FC-HistSSP simulation and WOA18 observations in the Pacific, the Atlantic, and the global ocean ~~from 1985 to 2014.~~ Nutrients distributions are reproduced well in the Atlantic by the NESM v3. The deepest penetration of relatively low-nutrients water to a 1000 m depth is simulated in the middle latitude regions. The observed high concentration of nutrients is ~~found~~simulated in the Atlantic south of 45°S, ~~and~~ Also, the equator-ward

~~transport of~~ phosphate and nitrate ~~are equator ward transported~~ by the Antarctic Intermediate Water at near 1000 m depth. ~~is simulated by the model.~~ In the Pacific, the ~~spatial patterns~~ latitude-depth distributions of nutrients broadly agree with observations, but with noticeable positive biases in the deep ~~Northern Pacific. The simulated centers of phosphate and nitrate rich water are too deep.~~ North Pacific.

5 To further analyze the possible reasons for discrepancies in nutrients distributions between the model simulation and observation, we ~~decompose~~ decomposed phosphate to its preformed and regenerated components (Weiss et al., 1970; Duteil et al., 2012) and compare the results with the WOA18 observations (Fig. 3). For the global, Atlantic, and Pacific Ocean, the preformed phosphate diagnosed from the model accounts for 51%, 47%, and 57% of the total phosphate inventory, and the result diagnosed from the WOA18 is 57%, 55%, and 64%, respectively. A relatively small percentage fraction of the preformed phosphate indicates stronger biological activities in the model. Compared to the observations, in the North Atlantic, the model ~~simulates a larger depletion of~~ underestimated the preformed phosphate (~~bias is about 0.2 mmol/m³) in the North Atlantic, which indicates too active,~~ indicating that biological ~~processes~~ uptake in the upper North Atlantic is overestimated by the model. The NESM v3 overestimates the preformed phosphate above ~500-meter depth but underestimates the preformed phosphate at the depth of 500 to 1500 meter. This dipole pattern of preformed phosphate biases is associated with the overestimated vertical mixing that brings excessive nutrient-rich water from the intermediate ocean to the surface (Fig. 3a2, 3a4, and 3a6). In the deep Pacific ~~Deep~~ Ocean, the preformed phosphate concentrations are between 1.3 to 1.5 mmol/m³ for both model simulation and observations.

15
20
25 The ~~NESM v3 simulates a dipole pattern of preformed phosphate bias above 1500 m depth, which is probably associated with the overestimated vertical mixing that brings too much nutrient rich water from the intermediate ocean to the surface (Fig. a2, a4, and a6).~~ The most noticeable ~~positive~~ biases of regenerated phosphate are found in the deep ~~Northern~~ North Pacific. In the model simulation, the high regenerated phosphate ~~water~~ in the deep North Pacific is ~~too deep~~ significantly overestimated, and the biases resemble the difference found in latitudinal-depth distributions of nutrients. In the deep ocean, preformed phosphate is only affected by ocean circulation, ~~while~~ whereas regenerated phosphate is affected by both circulation and remineralization. In the deep ocean, the NESM v3 simulates the preformed phosphate well but overestimates the regenerated phosphate, suggesting indicating that the

overestimated nutrients in the North Pacific deep ocean are mainly caused by biological processes.

Next, we present the NESM v3-simulated patterns of nutrient limitation. As shown in figure 4, the limiting patterns of nanophytoplankton and diatoms are similar in the ~~mid-low~~ tropical and temperate latitude oceans. Iron is the most limiting nutrient for both nanophytoplankton and diatoms in the ~~equatorial~~ eastern Pacific ~~Ocean~~ and the Southern Ocean. Nitrate ~~is the most limiting factor in the subtropical Pacific Ocean,~~ and phosphate ~~is~~ are the most limiting factor in the Indian Ocean, subtropical western Pacific, and ~~middle-low latitude of the~~ tropical Atlantic ~~Ocean~~. At high latitude oceans, nanophytoplankton is mostly limited by the available light and temperature, ~~while~~ whereas diatoms are mostly limited by silicate. The NESM v3 simulated limiting pattern ~~is generally consistent~~ broadly agrees with the results diagnosed from the IPSL-CM4A-LOOP (Schneider et al., 2008) and the Community Earth System Model-Biogeochemistry (CESM1-BGC; Moore et al., 2013), except that the iron limitation diagnosed from the NESM v3 is stronger in the Pacific and the Southern Ocean.

3.2 Biological Production

Figure 5 ~~shows~~ compares the modeled spatial ~~distribution~~ distributions of annual mean surface chlorophyll concentration from 1998 to 2014 ~~compared with~~ that in SeaWiFS observational data, GlobColour merged data, and OCCCI merged data.

In the NESM v3, chlorophyll in both nanophytoplankton and diatoms are parameterized based on the photo-adaptive model (Geider et al., 1997) in which chlorophyll is regulated by the chlorophyll-to-carbon ratio, growth of phytoplankton biomass, mortality, aggregation, and ~~the grazing by~~ zooplankton grazing.

The large-scale pattern of simulated ocean chlorophyll concentration broadly agrees with observations with high levels of chlorophyll in the subarctic Pacific Ocean and North Atlantic (>1 mg Chl m^3) and intermediate levels of chlorophyll in the Southern Ocean (~ 0.5 mg Chl m^3). Also, the observed relatively high chlorophyll concentration in the equatorial Pacific (~ 0.3 mg Chl m^3) surrounded by low chlorophyll concentration seawater over the subtropical oceans (<0.1 mg Chl m^3) are reproduced ~~by the model~~. The relatively high chlorophyll concentrations along the extratropical coastal regions are ~~reproduced,~~ but broadly captured by the NESM v3. However, the model ~~generally~~ underestimates chlorophyll concentration in the tropical coastal regions, especially in the tropical Indian Ocean, maritime

~~continent~~ Maritime Continent, and the tropical Atlantic ~~Ocean~~. This underestimation is partly associated with the deficiencies in modeled coastal dynamics, which is usually not represented well by the relatively coarse global ocean models (Aumont et al., 2015). It is reported that the observed chlorophyll distribution in the coastal region is better reproduced when PISCES is coupled to a higher resolution ocean circulation model (Lee et al., 2000; Hood et al., 2003; Kone et al., 2009). Also, ~~we inspect an underestimation of the~~ model underestimates chlorophyll ~~over~~ concentration in the Northern Indian Ocean. This underestimation of chlorophyll is associated with the underestimation of nutrients over the Indian Ocean (Fig. 1) that ~~increases nutrients limitation and~~ inhibits phytoplankton growth.

In the Southern Ocean where the seawater is typically characterized by high nutrients and low chlorophyll (Lin et al., 2016), noticeable discrepancies in chlorophyll concentrations are seen among different observational-based datasets ~~that, which~~ are associated with different algorithms used for these products. For example, in the Southern Ocean, chlorophyll concentration derived from reflectance by standard algorithms tend to be underestimated by a factor of about 2 to 2.5 (Kahru and Mitchell, 2010). ~~In the Southern Ocean~~ Compared to the three observational data-based estimates, the NESM v3 overestimates the chlorophyll concentration ~~to~~ in the east of 150°E Pacific and ~~underestimates it near the International Date Line (180°E)~~ Indian parts of the Southern Ocean. In the Atlantic part of the Southern Ocean, the modeled chlorophyll concentration is within the range of observational estimates, higher than the SeaWiFS but lower than the GlobColour and OCCCI.

Figure 6 shows the annual mean climatology spatial distributions of vertically integrated NPP from 2003 to 2014. Three different algorithms, including VGPM, Eppley-VGPM, and CbPM, are used to estimate the NPP based on the MODIS observations. Similar to the CbPM, the NESM v3 ~~also~~ simulates the NPP as the product of phytoplankton biomass and growth rate, while the VGPM and Eppley-VGPM describe the NPP as the product of chlorophyll and photosynthetic efficiencies. However, the ~~calculation~~ formulation of growth rate in the NESM v3 is more complex than that in CbPM, which involves chlorophyll, nutrients availability, temperature, respiration, and PAR.

~~There are significant differences between the NESM v3 simulated NPP and VGPM because the~~ The formulation of growth rate in the NESM v3 follows Eppley (1972), i.e., the growth rate is higher at high-temperature regions, while there is no temperature dependence in the VGPM. Therefore, Compared to

VGPM, the NESM v3 estimates more NPP in low latitude oceans and less ~~at NPP in~~ high latitude oceans ~~than that in the VGPM~~ (Fig. 6e). The ~~climatology~~ spatial distribution of the NESM v3 simulated vertically integrated NPP resembles Eppley-VGPM and CbPM estimates (Fig. 6a, c, d). The ~~main~~ NESM v3 broadly reproduces the observed spatial pattern of NPP distribution with high ~~concentrations of~~ NPP in the eastern equatorial Pacific and ~~middle-latitude oceans~~ Maritime Continent and low ~~concentration of~~ NPP in the ~~middle-low latitude oceans, the Southern Ocean,~~ subtropical Pacific and high latitude oceans ~~are reproduced by the NESM v3~~. Also, the NESM v3 ~~reproduced~~ broadly captures the high concentrations of NPP in low-latitude coastal regions ~~to some extent~~. ~~Comparison between three observational data-based estimates and the NESM v3 also suggests that temperature dependence is important to produce the meridional distribution pattern of marine NPP with the high level of NPP over the high-latitude oceans and low level of NPP over the low-latitude oceans.~~

. Although the global pattern of NPP broadly agrees with the ~~observational~~ observation-based estimates, PCC between model simulation and Eppley-VGPM is only 0.5, indicating that some local features are not well described in the NESM v3. Compared to CbPM and Eppley-VGPM, the NESM v3 significantly underestimates the NPP in the Indian Ocean. The NESM v3 also underestimates the NPP in the eastern coastal areas of the United States and the Arctic coastal areas.

Averaged from 2003 to 2014, the globally integrated ocean NPP from the NESM v3 simulation is 45.1 PgC yr⁻¹, compared with the data-based estimates of ~~3738~~ to ~~6765~~ PgC yr⁻¹. (Buitenhuis et al., 2013). The large range of data-based estimates of global NPP is a result of different satellite observations and different algorithms for the NPP estimation (Longhurst et al., 1995; Antoine et al., 1996; Behrenfeld and Falkowski, 1997b; Behrenfeld et al., 2005). Global NPP simulated by CMIP5 models also shows a wide range of values from 30.9 to 78.7 PgC yr⁻¹ (Bopp et al., 2013). NESM v3 simulated global NPP is within the range of data-based estimates and current CMIP5 model estimates. Of the NESM v3 simulated global ocean NPP, 20% is contributed by diatoms, and 80% is contributed by nanophytoplankton. For comparison, from the data-based ~~estimate~~ estimates, 7% to 32% of the total NPP is associated with diatoms (Uitz et al., 2010; Hirata et al., 2011), while ocean biogeochemical models estimate that 15% to 30% of global NPP is from diatoms (Aumont et al., 2003; Dutkiewicz et al., 2005; Yool et al., 2011).

3.3 Dissolved inorganic carbon and alkalinity

Figures 7 and 8 display the modeled and observed spatial distributions of alkalinity and DIC averaged over the upper ocean (0-100 m) and along the zonally averaged section in the Pacific Ocean, Atlantic Ocean, and the global ocean- during the period of 1985 to 2014. The model's skill in simulating upper-ocean alkalinity is moderate (PCC = 0.56). ~~The observed global spatial pattern of alkalinity is generally simulated by the NESM v3.~~ The model reproduced the observed high alkalinity in the subtropical surface oceans ~~and low alkalinity near the maritime continent~~, and the modeled global upper ocean mean alkalinity ~~only~~ has a minor negative bias of 0.45%. The major discrepancies are seen in the Southern Ocean and the subarctic Pacific with a positive bias of more than 80 mmol/m³. In high-latitude oceans, convective mixing of alkalinity-rich deep water is an important factor ~~of changing~~ controlling upper ocean alkalinity, and SST can be used as a proxy of the convective mixing ~~change~~ (Lee et al., 2006). An underestimation of SST of 1°C is simulated at in high latitude oceans (figures not shown), indicating a stronger convective mixing, which may explain the overestimated alkalinity at in high latitude oceans. The alkalinity has a negative bias of more than 60 mmol/m³ near the ~~maritime continent~~ Maritime Continent, where the alkalinity concentration is usually related to salinity (Lee et al., 2006). Cao et al. (2018) found that the NESM v3 simulates excessive precipitation over the ~~maritime continent~~ Maritime Continent, which results in ~~an~~ the underestimation of ~~the~~ salinity of 2 PSU.

NESM v3 simulates well the large-scale pattern of the observed DIC (PCC = 0.78) with high DIC concentration in the middle-high latitude Atlantic and low DIC ~~concentrations~~ concentration in the middle-low latitude Pacific and the Indian Ocean. The model- ~~simulated~~ global mean upper ocean ~~means~~ DIC ~~only~~ has a minor positive bias of 0.27%. Although the global pattern of DIC is different from alkalinity, their ~~bias patterns (model simulation relative to observation)~~ bias patterns are similar (Fig. 7e, f). The largest positive DIC bias of more than 80 mmol C/m³ is simulated in the Southern Ocean and Subarctic Pacific, ~~and while~~ a negative bias of more than 80 mmol C/m³ is simulated in the ~~maritime continent~~ Maritime Continent.

~~The~~ In the Atlantic, the large-scale patterns of the ~~zonal-averaged~~ latitudinal-depth distribution distributions of ~~both~~ DIC and alkalinity ~~are~~ simulated well by the model broadly agree with observations. Both alkalinity and DIC are slightly overestimated in the South Atlantic Ocean and underestimated in the North Atlantic. Apparent biases of DIC and alkalinity are seen in the deep

~~Northern~~North Pacific. One noticeable pattern of the observed DIC and alkalinity is that their maximum concentrations are around 2000-3000 m of the North Pacific Ocean, which the model fails to reproduce. The model also overestimates DIC storage in the deep Pacific Ocean. The mismatches between model simulation and observations, i.e. an underestimation of DIC and alkalinity concentrations in the upper 1000 m depth and an overestimation of their concentrations in the deep ocean, resemble those of nitrate and phosphatenutrients. It indicates that modeled discrepancies of alkalinity and DIC may also be attributed to excessive deep and active remineralization processes, which releases thea large amount of dissolved carbon in the deep ocean.

3.4 Assessment of biogeochemical fields by Taylor diagram

Figure 9 respectively compares the spatial patterns of the NESM v3 and IPSL simulated biogeochemistry-related fields with corresponding observations using a Taylor diagram ([Taylor, 2001](#)). In summary, model-simulated statistical patterns of the nutrients in the upper ocean nutrients compare well with observations, whilewhereas the simulated spatial patterns of chlorophyll, primary production, and alkalinity show larger discrepancies from observations. It is noted that chlorophyll and NPP are not directly observed but diagnosed from the observation-based data, and thus their estimations are subject to considerable uncertainties. Compared with biogeochemical fields in IPSL that shares the same marine biogeochemical cycle component with NESM v3, the NESM v3 has comparable skill in reproducing spatial distributions of nutrients and chlorophyll, but less skill in reproducing DIC and alkalinity with relatively larger SDsNSDs.

We also examine other CMIP5 models results that documented in previous studies ([Moore et al., 2013](#); [Anav et al., 2013](#); [Tjiputra et al., 2013](#); [Séférian et al., 2013](#)). ~~The skill of the models are~~ For different ~~according to the~~ biogeochemical fields ~~examined~~, different models show different skills. For example, PCCs between nutrients and observations in ~~Community Earth System Model (CESM)~~CESM1-BGC are about 0.8, which is lower than that in the NESM v3, but CESM has a better representation of chlorophyll ~~distribution~~ ([Moore et al., 2013](#)). ~~A further study is needed to rank those models, which is beyond the scope of this study~~. Nevertheless, the NESM v3 shows comparable skill in simulating upper ocean biogeochemical fields ~~in the present day~~ with other CMIP5 models.

3.5 Oceanic anthropogenic CO₂ uptake during the historical period

In this section, we ~~compare~~compared the NESM v3 simulated anthropogenic carbon uptake (~~FC-HistSSP simulation~~) during the historical period (~~FC~~) against available ~~observational data~~observation-based estimates.

5 First, we ~~compare~~compared the NESM v3 simulated sea-air CO₂ flux against available observations for the reference year of 2000 ([Takahashi et al., 2009](#)). As shown in Fig. 10, the NESM v3 realistically reproduces the large-scale pattern of the observed sea-air CO₂ flux with CO₂ outgassing in the equatorial oceans and uptake in the mid-to-high latitude oceans (PCC=0.71 and ~~SDs~~NSDs =1.04). For both observation and model results, strong CO₂ uptake is found in the ~~North~~Northern Atlantic where sea surface temperature is low and the formation of deep water is active. Compared to the data-based estimates, modeled sea-air CO₂ flux is overestimated in the tropical Pacific, the Southern ~~ocean~~Ocean, and the ~~Northern~~North Pacific (near 30°N), ~~while~~. Also, the ~~strongest model~~ underestimates ~~of modeled~~ sea-air CO₂ flux ~~are seen~~ in the high-latitude oceans (Fig. 9c and 9d). The globally integrated ocean uptake flux from observation is 1.6 ± 0.9 PgC per year ~~in~~for the reference year of 2000 ([Takahashi et al., 2009](#)), ~~while~~whereas the value is 2.8 PgC per year from the model simulation. The difference between model and observation is mainly originated from the model-simulated positive bias (-1 PgC yr^{-1}) in the pre-industrial steady-state oceanic CO₂ ~~uptake due to the 3-dimensional correction of nutrient and alkalinity in the PISCES model (Séférian et al., 2013). The 3-dimensional correction refers to that the global inventory of nutrient and alkalinity are restored toward the observations every January 1 (Aumont et al.,~~ 2015). In the NESM v3, the pre-industrial steady-state of total oceanic CO₂ uptake is 1.0 PgC PgC per year, compared with the ~~observation~~data-based value of $0.4 \pm 0.2 \text{ PgC PgC}$ per year ([Takahashi et al., 2009](#)). Taking the pre-industrial steady-state into consideration, the total ~~ocean~~oceanic anthropogenic CO₂ uptake flux in the year 2000 is 1.8 PgC from the model simulation, ~~while it is estimated~~compared to ~~be~~the value of 2.0 ± 1.0 PgC from the observation ([Takahashi et al., 2009](#)).

25 WeThen, we compared the NESM v3 simulated anthropogenic CO₂ budget with the data-based ~~estimate~~estimates provided by the Fifth Assessment Report of ~~IPCC~~Intergovernmental Panel on Climate Change (IPCC AR5) (Table 1). The model-simulated ocean uptake of anthropogenic CO₂ is slightly lower than that from the IPCC AR5 but within the estimated uncertainty range. From the pre-industrial ~~time~~era

to the year 2011, NESM v3 simulated cumulative oceanic CO₂ uptake is 137.2 PgC, compared with IPCC data-based estimates of 155 ± 30 PgC (Ciais et al., 2013). The decadal mean oceanic anthropogenic CO₂ uptake diagnosed from the FC-HistSSP run increases from 1.7 to 2.3 PgC yr⁻¹ from the 1980s to 2009, while the observation ranges from 2.0 ± 0.7 to 2.4 ± 0.7 PgC yr⁻¹. Also, compared with recent estimates, from the year 1870 to 2016, the modeled cumulative CO₂ uptake of 149 PgC is within, compared to the observational range of 150 ± 20 PgC (Le Quéré et al., 2018).

The Model-simulated vertically integrated column inventory of modeled ocean storage of anthropogenic DIC (i.e. FC-HistSSP relative to CTRL simulation) from 2000 to 2004 (Fig. 11a) and from 1992 to 1996 (Fig. 11e) are compared with GLODAP v2 (Fig. 11b) and GLODAP v1 (Fig. 11d), respectively. Compared with GLODAP v2, the NESM v3 reasonably captures the large-scale data-based distribution of observed anthropogenic DIC. The largest inventory in the 2000s of more than 100 mol C m⁻² is simulated in the Northern Atlantic where SST is low and deep water formation is active. In the model simulation from 2000 to 2004, the North Atlantic stores 20.8% of the global oceanic anthropogenic carbon, while whereas it is 17.6% in the observation (Fig. 11a and 11b). GLODAP v2. In other oceans ocean basins, the large inventory is mainly found in the middle-latitude areas near 30°N and 30°S, for both model simulations and observations. In the Southern Hemisphere Oceans NESM v3 simulations, from 2000 to 2004, 58.9% of the global oceanic anthropogenic DIC inventory carbon is simulated stored in the Southern Ocean, compared to the value of 62.6% in the observation GLODAP v2.

The most One noticeable discrepancy between the GLODAP v2 and model simulation around 2002 is found in the south of 50°S. Only 8.3% of the global oceanic anthropogenic DIC inventory is simulated, while by the model in this region, whereas the value is 15.5% in the observation GLODAP v2. However, we noticed that the vertically integrated anthropogenic DIC concentration is also low in the southern ocean Southern Ocean (south of 50°S) in the GLODAP v1 and, accounting for only 9.9% of the global inventory is stored in this region. It is noted that anthropogenic. Anthropogenic DIC in the GLODAP is diagnosed by a crude application of the transit time distribution method, and thus the results are subject to considerable uncertainties (Lauvset et al., 2016).

Figure 12 shows the latitudinal-depth distributions of anthropogenic DIC concentration (FC-HistSSP

relative to CTRL simulation) in the Atlantic, Pacific, and the global Ocean from the NESM v3 FC model simulation and GLODAP v2. ~~The anthropogenic CO₂ invades the ocean in the air-sea interface and then penetrates downward.~~ The observed highest high concentrations (more than 51 mmol C m⁻³) in near-surface waters and ~~the observed~~ low concentration (less than 3 mmol C m⁻³) in most of the deep ocean (the Pacific and the middle-low latitude Atlantic) are simulated by the NESM v3. For both data-based estimates and model simulations, a substantial amount of anthropogenic CO₂ has penetrated down to the ocean interior as deep as 1000 m depth with two penetration tongues near 30°N and 40°S, and the deepest penetration of anthropogenic DIC is found in the Northern Atlantic. Deep penetration of anthropogenic DIC is typically associated with convergence zones at temperate ~~latitudes~~ and high latitude oceans where vertical mixing is strong (Sabine et al., 2004). Similar to the vertically integrated inventory of anthropogenic DIC (Fig. 11), the major discrepancy of anthropogenic DIC in the latitudinal-depth distribution is also found in the Southern Atlantic south of 50°S: where the model greatly underestimates the anthropogenic DIC in the entire ocean column.

3.6 Sensitivity of the oceanic CO₂ uptake to increasing atmospheric CO₂ and global warming

The ocean carbon cycle is regulated by changes in atmospheric CO₂ and physical climate (Doney et al., 2004). Increasing atmospheric CO₂ affects oceanic CO₂ uptake directly. Meanwhile, global warming also affects the ocean carbon cycle via changes in ~~climatic fields~~ (climate and biological rates (Gregory et al., 2005; Pierce et al., 2012; Steinacher et al., 2010; Pierce et al., 2012; Olonscheck et al., 2013; Lewandowska et al., 2014). In this section, we first presented the NESM v3 simulated physical climate change and oceanic CO₂ uptake under the historical and SSP5-8.5 ~~scenarios~~ scenarios. Then, we ~~present~~ presented the NESM v3 simulated response of the ocean carbon cycle sensitivity parameters uptake to increasing atmospheric CO₂ and ~~their nonlinearity~~ global warming in the SSP5-8.5 and 1ptCO₂ runs.

3.6.1 NESM v3 simulated physical climate change under historical and SSP5-8.5 scenario

Figure 13 shows the NESM v3 simulated changes (relative to control simulation) in the global annual mean surface air temperature (SAT), mixed layer depth (MLD), and ~~the~~ intensity of Atlantic meridional overturning circulation (AMOC) from 1850 to 2100 under the historical and SSP5-8.5 scenarios. Changes

of SAT, MLD, and AMOC in [the RC-HistSSP](#) and [FC-HistSSP](#) simulations are almost the same, while those changes in the [BC-HistSSP](#) simulation are rather small. In the [FC-HistSSP](#) simulation, the annual ~~global~~-mean SAT anomaly averaged [over the years](#) from 2080 to 2100 (relative to the period of 1986-2005) is 4.6 K, which is at the higher end of the CMIP5 model results (2.6 ~ 4.7 K) under the RCP 8.5 scenario ([Collins and Knutti, 2013](#); [Knutti and Sedláček, 2013](#)). It is noted that the CMIP6 input forcing is used in this study and the atmospheric CO₂ concentration at the end of the 21st century in [the](#) SSP5-8.5 is about 10% higher than ~~the concentration that~~ in the RCP 8.5 scenario. [In the year 2100, Also, we can see a slight increase in SAT change in the BC-HistSSP \(Fig. 13a\), which is 6.3 \(6.7 and 0.8\) K caused by the non-CO₂ GHGs change. Compared with the pre-industrial era, by the end of 21st century \(averaged over the years 2091 to 2100\), the SAT in the FC\(-HistSSP, RC and BC\) simulation. With the increasing atmospheric temperature, the global ocean also becomes warmer in FC and RC-HistSSP, and BC-HistSSP simulations, reducing CO₂ solubility and oceanic CO₂ uptake. increased by 6.0, 6.0, and 0.8 K, respectively.](#)

Modeled MLD decreases since the 1980s. [In](#) [Compared with the year 2100, pre-industrial era, by the end of 21st century, MLD change is 8.5 \(-8 and -1\) meter](#) in the [FC\(-HistSSP, RC-HistSSP, and BC\)-HistSSP](#) simulation [decreased by 7.9, 8.0, and 1.4 meters, respectively](#). The reduction of mixed layer depth indicates a more stratified upper ocean. A substantial weakening of AMOC intensity in the [FC-HistSSP](#) simulation is projected for the 21st century, which is associated with ocean surface warming and increased freshwater input into the North Atlantic ([Gregory et al., 2005](#)). In the pre-industrial period, the model-simulated AMOC index at 30°N is 17.5 Sv (1Sv = 10⁶ m³ s⁻¹), [which is](#) within the range of 14 to 31 Sv from CMIP5 models ([Weaver et al., 2012](#)). ~~The~~ [In the FC-HistSSP simulation, the modeled annual mean of AMOC transport at 30°N averaged from over the years 2004 to 2011 is 17.1 Sv, while whereas the observation record during the same period from RAPID/MOCHA \(Rapid Climate Change programme / Meridional Ocean Circulation and Heatflux Array\) is 17.5 ± 3.8 Sv \(Rhein et al., 2013\). By the year 2100, end of the simulated intensity of AMOC declines 21st century, the NESM v3 simulates a 54% reduction of AMOC \(from 17.5 Sv to 8.0 Sv. In\) from the FC-HistSSP simulation, the simulated 54% weakening of AMOC by whereas the end of this century is at the higher end of what is simulated by AMOC reduction under the RCP8.5 scenario from](#) CMIP5 models [that range ranges](#) from 15% to 60% ~~under the~~

[RCP 8.5 scenario](#) ([Cheng et al., 2013](#)). The higher atmospheric CO₂ concentration at the end of 2100 in the SSP5-8.5 may partly explain the larger AMOC change in this study. Also, [Cao et al. \(2018\)](#) pointed out that the equilibrium climate sensitivity to CO₂ forcing in the NESM v3 is about 10% higher than that of the CMIP5 ensemble.

5 3.6.2 NESM v3 simulated oceanic CO₂ uptakes under historical and SSP5-8.5 scenario

Figure 14 shows the time evolution of the oceanic CO₂ uptake from the BC-[HistSSP](#), RC-[HistSSP](#), FC-[HistSSP](#), and the linear sum of the BC-[HistSSP](#) and RC-[HistSSP](#). In the BC-[HistSSP](#) simulation, the global ocean absorbed a total amount of 662 PgC of anthropogenic CO₂ from the atmosphere by the year 2100. In the RC-[HistSSP](#) simulation, the increased sea surface temperature, enhanced ocean stratification, and ~~the~~ weakened AMOC all act to decrease CO₂ uptake ([Cox et al., 2000](#); [Zickfeld et al., 2008](#); [Roy et al., 2011](#); [Goris et al., 2015](#)). ~~As a result, global warming alone causes the ocean to release CO₂ into the atmosphere by reducing CO₂ solubility, enhancing the oceanic pCO₂ and altering biological rates~~ ([Steinacher et al., 2010](#); [Olonscheck et al., 2013](#); [Lewandowska et al., 2014](#)[Roy et al., 2011](#); [Goris et al., 2015](#); [Cao et al., 2017](#)). By the year 2100, the modeled cumulative CO₂ uptake is -35.9 PgC- in the RC-
10 [HistSSP simulation](#). In the FC-[HistSSP](#) simulation, oceanic CO₂ uptake is affected by both the increase in atmospheric CO₂ and global warming. By the end of the 21st century, in the FC-[HistSSP](#), simulated cumulative oceanic CO₂ uptake since the pre-industrial era is 567 PgC, which is within the ranges ~~from~~of 420 PgC to 600 ~~PgC~~PgC from CMIP5 models results under the RCP 8.5 scenario ([Jones et al., 2013](#)). The sum of the simulated oceanic CO₂ uptake from the BC-[HistSSP](#) and RC-[HistSSP](#) simulations (626
20 PgC) is larger than that from the FC-[HistSSP](#) run (567 PgC), indicating that the effect of increasing atmospheric CO₂ (carbon-concentration sensitivity) and the effect of global warming (carbon-climate sensitivity) on the oceanic CO₂ uptake are not ~~exactly~~perfectly additive. This nonlinearity was also found in previous studies ([Boer and Arora, 2009](#); [Gregory et al., 2009](#); [Schwinger et al., 2014](#)). The NESM v3 simulated nonlinearity (i.e., BC+RC-FC) is 59 PgC by the end of the 21st century, which is larger than the
25 absolute value of the radiative effect on oceanic carbon uptake (-35.9 PgC).

To better understand oceanic CO₂ uptake in response to changing atmospheric CO₂ and global warming ~~in model simulation~~, Figure 15 shows the spatial distribution of anthropogenic sea-air CO₂ flux at the end

of the 21st century (averaged over the ~~year~~years 2091 to 2100) ~~under the SSP5-8.5 scenario~~ from ~~FC-~~HistSSP, ~~RC-~~HistSSP, and ~~BC-~~HistSSP simulations, and the difference between ~~FC-~~HistSSP simulation and the sum of ~~RC-~~HistSSP and ~~BC-~~HistSSP simulations.

In the ~~BC-~~HistSSP simulation, the oceanic anthropogenic CO₂ uptake is 8.0 ~~Pg~~C~~PgC~~ per year at the end of the 21st century. The ocean absorbs atmospheric CO₂ in most regions except for a few scattered grid points at the mid-latitudes with slight CO₂ outgassing. The strongest CO₂ uptake of about 150 g C m⁻² yr⁻¹ is found in the North Atlantic, subarctic Pacific, and the Southern Ocean. Results from the ~~RC-~~HistSSP simulation show CO₂ outgassing in large parts of the global ocean as a result of global warming. In the Arctic Ocean, warming induces a net uptake of CO₂ of 0.07 PgC yr⁻¹ because ~~of~~ the reduced sea-ice extent under global warming ~~that~~ allows more open seawater to absorb atmospheric CO₂. In the ~~Northern~~North Atlantic, the capacity of the ocean uptakes CO₂ is significantly suppressed due to the reduced AMOC.

~~The~~ ~~FC-~~HistSSP simulation shows the ~~combined~~joint effects of ~~the~~ increasing atmospheric CO₂ and global warming (Fig. ~~14e~~15c). ~~Net oceanic~~ CO₂ uptake is simulated in most regions with the strongest uptake in the Southern Ocean, indicating the dominant role of the increasing atmospheric CO₂. CO₂ outgassing is seen in the subtropical Pacific, indicating that the radiative effect dominates the response of oceanic CO₂ uptake in this region.

The nonlinearity of oceanic carbon uptake sensitivity during the 2090s is shown in Figure 15d. In the NESM v3, ~~the oceanic carbon uptake in the FC-HistSSP simulation is lower than the sum of that in the BC-HistSSP and RC-HistSSP simulations.~~ A relatively large nonlinearity is simulated in the ~~Northern~~ Atlantic north of 45°N (19.8% of the total nonlinearity) and the Southern Ocean south of 40°S (35.3% of the total nonlinearity), which is consistent with the findings of previous studies ([Zickfeld et al., 2011](#); [Schwinger et al., 2014](#)). The ~~interactions between different background simulation effects~~oceanic CO₂ content and global warming can partly explain the nonlinearity. Compared with the ~~RC-~~HistSSP simulation, in the ~~FC-~~HistSSP simulation, ~~there is much~~ more carbon ~~in the ocean that~~ is subject to the impact of climate change ~~in the FC simulation.~~ As a consequence, in the ~~FC-~~HistSSP simulation, the increased temperature would have a larger effect on CO₂ solubility and buffer factor ~~that reduces the ocean's capacity to absorb CO₂~~ ([Yi et al., 2001](#)). Also, reduced ocean circulation and increased ocean stratification would slow down the transport of anthropogenic CO₂ from the surface to the deep ocean.

Thus, compared to the BC-HistSSP simulation, slowing ocean ventilation in the FC-HistSSP simulation would cause a larger reduction in oceanic CO₂ uptake. ~~The oceanic carbon uptake in the FC simulation is lower than the sum of the BC and RC simulations, which is consistent with other CMIP5 models (Schwinger et al., 2014).~~

5 The above results also indicate that oceanic CO₂ uptake in high-latitude oceans is more sensitive to both the increasing atmospheric CO₂ concentration and global warming than low-latitude oceans, as well as their nonlinear interactions.

3.6.3 carbon-concentration and carbon-climate sensitivity parameters diagnosed ~~from~~under the SSP5-8.5.

10 In this section, we investigated oceanic CO₂ uptake under the framework of the carbon-concentration and carbon-climate sensitivity parameters.

Figure 16 shows the change in ocean carbon storage against the change in the atmospheric CO₂ concentration (Fig. 16a) and the global annual mean surface temperature (Fig. 16b), respectively. The derived evolution of the carbon-concentration sensitivity parameter β as a function of atmospheric CO₂ concentration and carbon-climate sensitivity parameter γ as a function of the change in temperature is shown in Fig. 16c and 16d, respectively.

As shown in Fig. 16, in the BC-HistSSP and RC-HistSSP simulations, modeled ocean storage of anthropogenic CO₂ scales roughly linearly with atmospheric CO₂ and changes in global mean surface temperature. Increasing atmospheric CO₂ alone increases oceanic CO₂ uptake whereas increasing temperature alone decreases CO₂ uptake. In the year 2100, the carbon-climate sensitivity parameter γ is ~~negative of~~ -5.4 Pg-CPgC/K while the carbon-concentration sensitivity parameter β is ~~positive of~~ 0.79 Pg-CPgC/ppm. ~~From 1850 to 2100, the carbon-climate parameter decreases with the increasing temperature change, indicating that with enhanced warming, each degree increase of surface temperature would induce more CO₂ outgassing from the ocean (Fig. 16d).~~ The Carbon-concentration parameter initially increases with atmospheric CO₂ and then decreases (Fig. 16c). The decreasing trend of β is consistent with the slowdown of the increasing trend of the oceanic CO₂ uptake at the end of the 21st century as a result of decreased oceanic buffer ability ~~due~~(Fig. 16a and 16c). ~~From 1850 to 2100,~~ the carbon-climate parameter becomes more negative with increasing temperature, indicating that the

increase of surface temperature would induce more CO₂ outgassing from the ocean in a warmer world (Fig. 16d). ~~DIC concentration.~~ Similar ~~trends~~ changes of carbon-climate and carbon-concentration sensitivity parameters are also found in CMIP5 ~~models~~ model simulations (Arora et al., 2013). The increased sensitivity of CO₂ outgassing to temperature and the decreased sensitivity of CO₂ uptake to atmospheric CO₂ concentration indicate that the ocean's ability to absorb atmospheric CO₂ would be weakened with increasing atmospheric CO₂ and global warming.

3.6.4 Carbon-concentration and carbon-climate sensitivity parameters from 1ptCO₂ runs.

In this section, we compared the carbon sensitivity parameters diagnosed from the 1ptCO₂ ~~experiment~~ experiments between the NESM v3 and CMIP5 models. The total CO₂ uptake during the 140 years in the FC-1%1ptCO₂ is 636 ~~PgC~~ PgC, while the results from CMIP5 models range from 533 to 676 ~~PgC~~ PgC. The sum of the total CO₂ uptake in the RC-1%1ptCO₂ and ~~the BC-1%1ptCO₂~~ is 65.7 PgC ~~702 PgC~~, which is larger than that in the FC-1%1ptCO₂. The simulated nonlinearity (i.e. BC-1%1ptCO₂ + RC-1%1ptCO₂ - FC-1%1ptCO₂) is about 10.3% of the total CO₂ uptake in the FC-1%1ptCO₂, which is at the higher end of the nonlinearity estimated for CMIP5 models ranging from (3.6%-10.6%) (Schwinger et al., 2014).

~~Then, we compare NESM v3 simulated β and γ parameters with those of CMIP5 results.~~

Figure 17 shows the simulated β and γ ~~parameters~~ in the 1ptCO₂ ~~runs~~ experiments. At the end of 1ptCO₂ ~~runs~~ experiments, the diagnosed value of β from CMIP5 models ranges from 0.69 to 0.91 PgC/ppm with a multi-model mean value of 0.80 PgC/ppm. For comparison, the β diagnosed from the NESM v3 simulations is 0.88 PgC/ppm at. In the end of 1ptCO₂ experiment, the simulation. The declining trend in β is found after simulated by the NESM v3 tends to decrease when the atmospheric CO₂ concentration increases to ~550 ppm, later than in Hist+RCP8.5 experiments (~400 ppm), which is consistent with the results that in CMIP5 models. Compared with β , the At the end of the 1ptCO₂ experiments, the diagnosed value of γ from CMIP5 models has a much larger range, and the value at the end of the simulations ranges from -2.4 to -12.1 PgC/K. The larger spread of γ is associated with the spread of the model-simulated climate change and the dependency of carbon cycle processes on climate change. For comparison, ~~our~~ simulated the diagnosed γ parameter from the NESM v3 simulation is -7.9 PgC/K.

4 Discussion and conclusion

In this study, we ~~evaluate~~evaluated the performance of ~~the~~ NESM v3 in simulating the present-day ocean biogeochemical cycle and historical and future oceanic carbon uptake. We also investigated the response of oceanic CO₂ uptake to the individual and combined effect of increasing atmospheric CO₂ and CO₂-induced ~~global~~-warming under SSP5-8.5 and 1ptCO₂ scenarios. The strengths and limitations of the NESM v3 are analyzed and documented.

The NESM v3 simulates reasonably well the large-scale patterns of observed upper ocean nutrients (~~with~~ PCCs > larger than 0.8 and ~~SDs~~NSDs close to 1.0 (Fig. 1 and 9). The high nutrient concentrations in the Eastern Pacific, subarctic Pacific, and the Southern Ocean are reproduced in the model- (Fig. 1). Also, the simulated global patterns of alkalinity, DIC, chlorophyll, and NPP broadly agree with observations and data-based estimates ~~with their main features reproduced~~ (PCCs: 0.5~0.8; ~~SDs~~NSDs: 0.5~1.6). For example, the high alkalinity concentration in the middle-latitude oceans, ~~the~~ high DIC and chlorophyll concentration in the high-latitude oceans, and ~~the~~ high NPP concentration in the low-latitude oceans are reproduced by the NESM v3- (Fig.5, 6, 7). The ~~global mean concentration of DIC and alkalinity in the upper ocean only has a bias of 0.27% and 0.45%. The integrated model-simulated~~ global ocean NPP averaged over the years from 2003 to 2014 ~~simulated by the NESM v3~~ is 45.1 PgC yr⁻¹, which is comparable with observation-based estimates of ~~37-67~~38-65 PgC yr⁻¹.

~~Compared with observational data~~observation-based estimates, the NESM v3 also does a good job in simulating oceanic CO₂ uptake. The observed global pattern of sea-air CO₂ flux is reproduced ~~in~~by the model- (Fig. 10). In the year 2000, the oceanic anthropogenic CO₂ uptake flux is 1.8 PgC in the NESM v3 simulation, whereas it is 2.0±1.0 PgC in the ~~observations.~~observation (Takahashi et al., 2009). The model-simulated cumulative oceanic anthropogenic CO₂ uptake from the year 1870 to 2016 is 149 PgC, which compares well with data-based estimates of 150 ± 20 PgC (Le Quéré et al., 2018).

~~The behavior of the NESM v3 is also comparable with CMIP5 models in the historical period and future projection. The skills of the models in simulating biogeochemical fields are different according to the variables examined. The integrated global ocean NPP simulated by the NESM v3 (45.1 Pg C) is within the range of CMIP5 models from 30.9 to 78.7 PgC yr⁻¹. At the end of the 21st century of the SSP5-8.5 future scenario, the NESM v3 simulated the total amount of oceanic CO₂ (567 PgC) is at the higher end~~

of CMIP5 results (420–600 PgC). The sensitivity of oceanic CO₂ uptake strongly depends on scenarios. In the 1ptCO₂ experiment, the NESM v3 simulated sensitivities of oceanic CO₂ uptake, i.e., carbon concentration sensitivity ($\beta = 0.88$ PgC/ppm), carbon climate sensitivity ($\gamma = 7.9$ PgC/K), and the nonlinearity (10.3%), compare well with those diagnosed from CMIP5 models (β : 0.69 to 0.91 PgC/ppm, γ : 2.4 to 12.1 PgC/K, and nonlinearity: 3.6% to 10.6%).

Compared with observations, the NESM v3 captures many aspects of the spatial structures of biogeochemical fields and their responses to climate change. However, model-observationsome discrepancies between model simulations and their underlying causes should also be emphasized, which are useful to users of the NESM v3 and also helpful to future model development. In general, observations remain. Our analysis shows that the simulated biases in biogeochemical fields in the upper ocean are primarily associated with the shortcoming in simulated ocean circulation, while the discrepancies in the deep ocean are primarily attributed to excessive remineralization.

In the upper ocean, slight overestimations of nutrients are found in the Pacific and the Southern Ocean (Fig. 1). In our simulations, the regions of overestimated nutrients (Fig.1) in general correspondscorrespond to regions with strong iron limitation (Fig. 4). The strong iron limitation in these areas limits biological activities, therefore reducing the uptake of nutrients by phytoplankton. Also, the overestimated nutrients are probably associated with strong vertical mixing in the Pacific and the Atlantic, which is indicated by the dipole pattern of preformed phosphate bias above the depth of about 1500 m depth (Fig. 3). In the Indian Ocean, the underestimation of nutrients is associated with the weak upwelling (figures not shown) that suppresses the nutrient entrainment to surface water. TheThen, the low-level nutrients nutrient concentrations in the Indian Ocean (Fig.1) reduce the biological activities and then result in underestimationthe underestimations of NPP and chlorophyll- (Fig.5 and 6). Also, in a relatively coarse resolution model, the underestimation of NPP and chlorophyll in the Indian Ocean could be associated with the poor descriptions of mesoscale and submesoscale processes (McGillicuddy et al., 1998; Lévy et al., 2001b). TheAlkalinity is underestimated near the Maritime Continent (Fig. 7), which is related to the underestimation of alkalinity is simulated near the maritime continent, where the model underestimates the surface salinity due to excessive precipitation- (Cao et al., 2018). In the high-latitude ocean, the model underestimates SST of about 1°C- (Cao et al., 2018), indicating strongerstrong

convective mixing, which leads to the overestimation of alkalinity.

As for ~~the~~ vertical profiles of ~~the~~ biogeochemical ~~field~~ fields, the ~~simulated~~ latitudinal-depth ~~distribution~~ distributions of nutrients broadly ~~agrees~~ agree with observations, but the simulated high-concentration centers ~~of nutrients~~ in the ~~Northern~~ North Pacific are too ~~strong and too~~ deep (Fig. 2).

5 Excessive remineralization is the main cause of the overestimated nutrients in the ~~Northern-Deep~~ deep North Pacific ~~according to, as indicated by~~ the ~~results~~ analysis of ~~preformed and regenerated~~ nutrients ~~decomposition~~ (Fig. 3). Similar to the vertical distributions of nutrients, the model ~~simulated high concentration centers of~~ also significantly overestimates alkalinity and DIC ~~are also too strong and too~~

~~deep~~ in the ~~Northern~~ deep North Pacific. (Fig. 8). Excessive remineralization in the deep ocean consumes

10 a large amount of oxygen and releases dissolved organic carbon and nutrients. To better evaluate the NESM v3 simulated ocean dynamics and the ocean carbon cycle, the simulation of natural and bomb $^{14}\text{C}_2$, ~~which are often used to diagnose the behavior of the ocean circulation and carbon cycle model (Levin and Hesshaimer, 2000; Skinner et al., 2017)~~, will be implemented in future versions of NESM.

~~The behavior of the NESM v3 is also comparable with CMIP5 models in terms of simulated marine~~

15 ~~biogeochemical fields and oceanic carbon uptake. As for biogeochemical fields, the skills of models are different for different variables. For example, compared with IPSL, which shares the same ocean~~

~~biogeochemical model (PISCES) with the NESM v3, the NESM v3 has comparable skill in reproducing spatial distributions of the upper-ocean nutrients and chlorophyll, but less skill in reproducing DIC and~~

20 ~~alkalinity (Fig. 9). The integrated global ocean NPP simulated by the NESM v3 (45.1 PgC) is within the range of CMIP5 models that span from 30.9 to 78.7 PgC yr⁻¹ (Bopp et al., 2013). Also, oceanic carbon~~

~~uptake simulated by the NESM v3 compares well with that diagnosed from CMIP5 models. During the historical period from 1850 to 2005, the NESM v3 simulated cumulative ocean carbon uptake is 123 PgC,~~

~~compared to the CMIP5 models' average of 124 ± 30 PgC (Friedlingstein et al., 2014). By the end of the 21st century, accumulative oceanic CO₂ uptake simulated by the NESM v3 under the SSP5-8.5 scenario~~

25 ~~is 567 PgC. For comparison, under the RCP 8.5 scenario, cumulative oceanic CO₂ uptake simulated by CMIP5 models ranges from 420 to 600 PgC (Jones et al., 2013). The sensitivity of oceanic CO₂ uptake~~

~~strongly depends on CO₂ scenarios. In the 1ptCO₂ experiment, the NESM v3 simulated carbon-concentration sensitivity ($\beta = 0.88$ PgC/ppm), carbon-climate sensitivity ($\gamma = -7.9$ PgC/K), and the~~

nonlinearity of oceanic carbon uptake sensitivities (10.3%) compare well with those diagnosed from CMIP5 models (β : 0.69 to 0.91 PgC/ppm, γ : -2.4 to -12.1 PgC/K, and nonlinearity: 3.6% to 10.6%).

Overall, compared with both observations and CMIP5 models, the NESM v3 ~~does a good job~~ shows skills in simulating ocean biogeochemical fields and oceanic carbon uptake. ~~Despite these model-observation discrepancies,~~ With the ongoing CMIP6 project (Eyring et al., 2016), it is expected that a reasonable next step to evaluate NESM v3 with CMIP6 models. The current version of NESM v3 can be used as a useful modeling tool to study interactive feedbacks between the ocean carbon cycle and climate change and the underlying mechanisms.

Code and data availability.

The source code of NESM v3, together with all input data are saved in one compressed file, which can be downloaded from: <https://doi.org/10.5281/zenodo.3524938> after registration. Also, a user guide describing the installation instructions, driver scripts, and software dependencies can be found in the repository at the same link. The simulation results illustrated in this study can be made available upon request to the authors.

Author contributions. Yifei Dai and Long Cao performed the simulations, analyzed the experiments, and made the figures. The NUIST ESM team led by Bin Wang provided the code of NESM v3 used in this study and Bin Wang provided helpful discussions. Yifei Dai, Long Cao, and Bin Wang all contributed to the writing of the manuscript.

Acknowledgement

Long Cao is supported by the National Natural Science Foundation of China (41675063; 41975103). Bin Wang acknowledges the support by the Nanjing University of Information Science and Technology through funding the joint China-US Atmosphere-Ocean Research Center at the University of Hawaii. Yifei Dai acknowledges the support by the China Scholarship Council by providing a scholarship under the State Scholarship Fund. This is the ESMC publication number XXX and IPRC publication number YYYY.

Reference

- Anav, A., Friedlingstein, P., Kidston, M., Boop, L., Ciais, P., Cox, P., Jones, C., Jung, M., Myneni, R., and Zhu, Z.: Evaluating the Land and Ocean Components of the Global Carbon Cycle in the CMIP5 Earth System Models. *Journal of Climate*, 26(18), 6801–6843, 2013.
- 5 Antoine, D., André, J.M., and Morel, A.: Oceanic primary production: 2. Estimation at global scale from satellite (Coastal Zone Color Scanner) chlorophyll. *Global Biogeochemical Cycles*, 10(1), 57-69, 1996.
- Arora, V.K., Boer, G.J., Friedlingstein, P., Eby, M., Jones, C.D., Christian, J.R., Bonan, G., Bopp, L., Brovkin, V., Cadule, P., Hajima, T., Ilyina, T., Lindsay, K., Tjiputra, J.F., and Wu, T.: Carbon–concentration and carbon–climate feedbacks in CMIP5 Earth system models. *Journal of Climate*, 10 26(15), 5289-5314, 2013.
- Aumont, O., Belviso, S., and Monfray, P.: Dimethylsulfoniopropionate (DMSP) and dimethylsulfide (DMS) sea surface distributions simulated from a global three - dimensional ocean carbon cycle model. *Journal of Geophysical Research: Oceans*, 107(C4), 2002.
- 15 Aumont, O., Maier-Reimer, E., Blain, S., and Monfray, P.: An ecosystem model of the global ocean including Fe, Si, P colimitations. *Global Biogeochemical Cycles*, 17(2), 1060, 2003.
- Aumont, O., Éthé, C., Tagliabue, A., Bopp, L., and Gehlen, M.: PISCES-v2: an ocean biogeochemical model for carbon and ecosystem studies. *Geoscientific Model Development*, 8(8), 2465-2513, 2015.
- Ballantyne, A.P., Alden, C.B., Miller, J.B., Tans, P.P., and White, J.W.C.: Increase in observed net 20 carbon dioxide uptake by land and oceans during the past 50 years. *Nature*, 488(7409), 70-72, 2012.
- Behrenfeld, M.J., and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnology and Oceanography*, 42(1), 1-20, 1997a.
- Behrenfeld, M. J., and Falkowski P G.: A consumer's guide to phytoplankton primary productivity models. *Limnology and Oceanography*, 42(7), 1479-1491, 1997b.
- 25 Behrenfeld, M.J., Boss, E., Siegel, D.A., and Shea, D.M.: Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles*, 19, GB1006, 2005.
- Boer, G.J., and Arora, V.: Temperature and concentration feedbacks in the carbon cycle. *Geophysical Research Letters*, 36, L02704, 2009.

- Bopp, L., Aumont, O., Cadule, P., Alvain, S. and Gehlen, M.: Response of diatoms distribution to global warming and potential implications: A global model study. *Geophysical Research Letters*, 32, L19606, 2005.
- Bopp, L., Resplandy, L., Orr, J.C., Doney, S.C., Dunne, J.P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Seferian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences*, 10, 6225-6245, 2013.
- [Buitenhuis, E.T., Hashioka, T. and Quéré, C.L.: Combined constraints on global ocean primary production using observations and models. *Global Biogeochemical Cycles*, 27\(3\), 847-858, 2013.](#)
- Cao, J., Wang, B., Xiang, B., Li, J., Wu, T., Fu, X., Wu, L., and Min, J.: Major modes of short-term climate variability in the newly developed NUIST Earth System Model (NESM), *Adv. Atmos. Sci.*, 32, 585–600, <https://doi.org/10.1007/s00376-014-4200-6>, 2015.
- Cao, J, Wang, B., Yang, Y.-M., Ma, L., Li, J., Sun, B., Bao, Y., He, J., Zhou, X., and Wu, L.: The NUIST Earth System Model (NESM) version 3: Description and preliminary evaluation, *Geosci. Model Dev.* 11(7), 2975-2993, 2018.
- Cao, L., and Zhang, H.: The role of biological rates in the simulated warming effect on oceanic CO₂ uptake. *Journal of Geophysical Research: Biogeosciences*, 122(5), 1098-1106, 2017.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R.B., Piao, S., and Thornton, P.: Carbon and Other Biogeochemical Cycles. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W.J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A.J., and Wehner, M.: Long-term Climate Change: Projections, Commitments and Irreversibility. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Stocker, T.F., D. Qin, G.-K. Plattner, et al., Eds.,

- Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Cox, P.M., Betts, R.A., Jones, C.D., Spall, S.A., and Totterdell, I.J.: Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature*, 408(6809), 184-187, 2000.
- Cheng, W., Chiang, J.C., and Zhang, D.: Atlantic meridional overturning circulation (AMOC) in CMIP5 models: RCP and historical simulations. *Journal of Climate*, 26(18), 7187-7197, 2013.
- Denman, K.L., G. Brasseur, A. Chidthaisong, P. Ciais, P.M. Cox, R.E. Dickinson, D. Hauglustaine, C. Heinze, E. Holland, D. Jacob, U. Lohmann, S Ramachandran, P.L. da Silva Dias, S.C. Wofsy, and X. Zhang.: Couplings Between Changes in the Climate System and Biogeochemistry. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- de Baar, H.J., Buma, A.G., Nolting, R.F., Cadée, G.C., Jacques, G. and Tréguer, P.J.: On iron limitation of the Southern Ocean: experimental observations in the Weddell and Scotia Seas. *Marine ecology progress series*, 105-122, 1990.
- Dlugokencky, E., and Tans, P.: Trends in atmospheric carbon dioxide, National Oceanic & Atmospheric Administration, Earth System Research Laboratory (NOAA/ESRL), available at: https://www.esrl.noaa.gov/gmd/ccgg/trends/gl_data.html, 2020.
- Doney, S.C., Lindsay, K., Caldeira, K., Campin, J.M., Drange, H., Dutay, J.C., Follows, M., Gao, Y., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Madec, G., Maier-Reimer, E., Marshall, J.C., Matear, R.J., Monfray, P., Mouchet, A., Naijar, R., Orr, J.C., Plattner, G.-K., Sarmiento, J., Schlitzer, R., Slater, R., Totterdell, I.J., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Evaluating global ocean carbon models: The importance of realistic physics. *Global Biogeochemical Cycles*, 18, GB3017, 2004.
- Droop, M. R.: 25 years of algal growth kinetics, *Bot. Mar.*, 26, 99–112, 1983.
- Dunne, J.P., Sarmiento, J.L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor. *Global Biogeochemical Cycles*, 21, GB4006, 2007.

- Duteil, O., Koeve, W., Oschlies, A., Aumont, O., Bianchi, D., Bopp, L., Galbraith, E., Matear, R., Moore, J.K., Sarmiento, J.L., and Segschneider, J.: Preformed and regenerated phosphate in ocean general circulation models: can right total concentrations be wrong? *Biogeosciences*, 9(5), 1797-1807, 2012.
- 5 Dutkiewicz, S., Follows, M.J., and Parekh, P.: Interactions of the iron and phosphorus cycles: A three-dimensional model study. *Global Biogeochemical Cycles*, 19, GB1021, 2005.
- Eppley, R.W.: Temperature and phytoplankton growth in the sea. *Fish. Bull.*, 70(4), 1063-1085, 1972.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937-1958, 2016.
- 10 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H.D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A.J., Yoshikawa, C., and Zeng, N.: Climate–carbon cycle feedback analysis: results from the C4MIP model intercomparison. *Journal of Climate*, 19(14), 3337-3353, 2006.
- 15 [Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K., and Knutti, R.: Uncertainties in CMIP5 Climate Projections due to Carbon Cycle Feedbacks. *Journal of Climate*, 27\(2\), 511-526, 2014.](#)
- Garcia, H., Locarnini, R., Boyer, T., Antonov, J., Zweng, M., Baranova, O. and Johnson, D.: World Ocean Atlas 2009, vol. 4, Nutrients (Phosphate, Nitrate, Silicate), edited by: Levitus, S. NOAA Atlas NESDIS, US Gov. Printing Office, Wash., DC, 2010.
- 20 Garcia, H. E., K. Weathers, C. R. Paver, I. Smolyar, T. P. Boyer, R. A. Locarnini, M. M. Zweng, A. V. Mishonov, O. K. Baranova, D. Seidov, and J. R. Reagan.: World Ocean Atlas 2018, Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+nitrite, silicate). A. Mishonov Technical Ed.: NOAA Atlas NESDIS 84, 35 pp, 2018.
- 25 Geider, R.J., MacIntyre, H.L. and Kana, T.M.: Dynamic model of phytoplankton growth and acclimation: responses of the balanced growth rate and the chlorophyll a: carbon ratio to light, nutrient-limitation and temperature. *Marine Ecology Progress Series*, 148, 187-200, 1997.

- Giorgetta, M. A., Roeckner, E., Mauritsen, T., Bader, J., Crueger, T., Esch, M., Rast, S., Kornblueh, L., Schmidt, H., Kinne, S., Hohenegger, C., Möbis, B., Krismer, T., Wieners, K.-H., and Stevens, B.: The Atmospheric General Circulation Model ECHAM6: Model Description, Tech. rep., Max Planck Institute for Meteorology, Hamburg, Germany, 2013.
- 5 Goris, N., Tjiputra, J., Schwinger, J., and Heinze, C.: Responses of carbon uptake and oceanic pCO₂ to climate change in the North Atlantic: A model study with the Bergen Earth System Model, *Global Biogeochem. Cycles*, 29(10), 1567-1583, 2015.
- Gregory, J.M., Dixon, K.W., Stouffer, R.J., Weaver, A.J., Driesschaert, E., Eby, M., Fichet, T., Hasumi, H., Hu, A., Jungclaus, J.H., Kamenkovich, I.V., Levermann, A., Montoya, M., Murakami, S.,
10 Nawrath, S., Oka, A., Sokolov, A.P., and Thorpe, R.B.: A model intercomparison of changes in the Atlantic thermohaline circulation in response to increasing atmospheric CO₂ concentration. *Geophysical Research Letters*, 32, L12703, 2005.
- Gregory, J.M., Jones, C.D., Cadule, P., and Friedlingstein, P.: Quantifying carbon cycle feedbacks. *Journal of Climate*, 22(19), 5232-5250, 2009.
- 15 Hirata, T., Hardman-Mountford, N.J., Brewin, R.J.W., Aiken, J., Barlow, R., Suzuki, K., Isada, T., Howell, E., Hashioka, T., Noguchi-Aita, M., and Yamanaka, Y.: Synoptic relationships between surface Chlorophyll-a and diagnostic pigments specific to phytoplankton functional types. *Biogeosciences*, 8(2), 311-327, 2011.
- Hood, R.R., Kohler, K.E., McCreary, J.P. and Smith, S.L.: A four-dimensional validation of a coupled
20 physical–biological model of the Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography*, 50(22-26), 2917-2945, 2003.
- Hunke, E.C., Lipscomb, W.H., Turner, A.K., Jeffery, N. and Elliott, S.: CICE: the Los Alamos Sea Ice Model Documentation and Software User’s Manual Version 4.1 LA-CC-06-012. T-3 Fluid Dynamics Group, Los Alamos National Laboratory, Los Alamos N.M, 2010.
- 25 Ilyina, Tatiana, Katharina D. Six, Joachim Segschneider, Ernst Maier-Reimer, Hongmei Li, and Ismael Nunez-Riboni: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations. *Journal of Advances in Modeling Earth Systems*, 5, 287-315, 2013.

- Jochum, M.: Impact of latitudinal variations in vertical diffusivity on climate simulations. *Journal of Geophysical Research: Oceans*, 114, C01010, 2009.
- Jones, C., Robertson, E., Arora, V., Friedlingstein, P., Shevliakova, E., Bopp, L., Brovkin, V., Hajima, T., Kato, E., Kawamiya, M. Liddicoat, S., Lindsay, K., Reick, C. H., Roelandt, C., Segschneider, J., and Tjiputra, J.: Twenty-first-century compatible CO₂ emissions and airborne fraction simulated by CMIP5 earth system models under four representative concentration pathways. *Journal of Climate*, 26(13), 4398-4413, 2013.
- Jones, C.D., Arora, V., Friedlingstein, P., Bopp, L., Brovkin, V., Dunne, J., Graven, H., Hoffman, F., Ilyina, T., John, J.G. and Jung, M.: C4MIP–The Coupled Climate–Carbon Cycle Model Intercomparison Project: experimental protocol for CMIP6. *Geoscientific Model Development*, 9(8), 2853-2880, 2016.
- Joos, F., and Spahni, R.: Rates of change in natural and anthropogenic radiative forcing over the past 20,000 years. *Proceedings of the National Academy of Sciences*, 105(5), 1425-1430, 2008.
- Kahru, M., and Mitchell, B.G.: Blending of ocean colour algorithms applied to the Southern Ocean. *Remote Sensing Letters*, 1(2), 119-124, 2010.
- Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP). *Global Biogeochemical Cycles*, 18(4), 357-370, 2004.
- Key, R.M., A. Olsen, S. van Heuven, S. K. Lauvset, A. Velo, X. Lin, C. Schirnick, A. Kozyr, T. Tanhua, M. Hoppema, S. Jutterström, R. Steinfeldt, E. Jeansson, M. Ishi, F. F. Perez, and T. Suzuki.: Global Ocean Data Analysis Project, Version 2 (GLODAPv2), ORNL/CDIAC-162, NDP-P093. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee. 2015.
- Knutti, R., and Sedláček, J.: Robustness and uncertainties in the new CMIP5 climate model projections. *Nature Climate Change*, 3(4), 369-373, 2013.
- Koné, V., Aumont, O., Lévy, M., and Resplandy, L.: Physical and biogeochemical controls of the phytoplankton seasonal cycle in the Indian Ocean: A modeling study. *Indian Ocean Biogeochemical Processes and Ecological Variability*, 185, 147-166, 2009.

- Körtzinger, A., Hedges, J.I., and Quay, P.D.: Redfield ratios revisited: Removing the biasing effect of anthropogenic CO₂. *Limnology and Oceanography*, 46(4), 964-970, 2001.
- Lauvset, S. K., R. M. Key, A. Olsen, S. van Heuven, A. Velo, X. Lin, C. Schirnick, A. Kozyr, T. Tanhua, M. Hoppema, S. Jutterström, R. Steinfeldt, E. Jeansson, M. Ishii, F. F. Pérez, T. Suzuki and S. Watelet.: A new global interior ocean mapped climatology: the 1°x1° GLODAP version 2, *Earth System Science Data*, 8, 325-340, 2016.
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Pongratz, J., Manning, A. C., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Jackson, R. B., Boden, T. A., Tans, P. P., Andrews, O. D., Arora, V. K., Bakker, D. C. E., Barbero, L., Becker, M., Betts, R. A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Cosca, C. E., Cross, J., Currie, K., Gasser, T., Harris, I., Hauck, J., Haverd, V., Houghton, R. A., Hunt, C. W., Hurtt, G., Ilyina, T., Jain, A. K., Kato, E., Kautz, M., Keeling, R. F., Klein Goldewijk, K., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lima, I., Lombardozzi, D., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Nojiri, Y., Padin, X. A., Peregon, A., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Reimer, J., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., van Heuven, S., Viovy, N., Vuichard, N., Walker, A. P., Watson, A. J., Wiltshire, A. J., Zaehle, S., and Zhu, D.: Global Carbon Budget 2017, *Earth Syst. Sci. Data*, 10, 405-448, 2018.
- Lee, C.M., Jones, B.H., Brink, K.H., and Fischer, A.S.: The upper-ocean response to monsoonal forcing in the Arabian Sea: seasonal and spatial variability. *Deep Sea Research Part II: Topical Studies in Oceanography*, 47(7-8), 1177-1226, 2000.
- Lee, K., L. T. Tong, F. J. Millero, C. L. Sabine, A. G. Dickson, C. Goyet, G. H. Park, R. Wanninkhof, R. A. Feely, and R. M. Key: Global relationships of total alkalinity with salinity and temperature in surface waters of the world's oceans, *Geophys. Res. Lett.*, 33, L19605, 2006.
- Lengaigne, M., Madec, G., Bopp, L., Menkes, C., Aumont, O. and Cadule, P.: Bio-physical feedbacks in the Arctic Ocean using an Earth system model. *Geophysical Research Letters*, 36, L21602, 2009.
- [Levin, I. and Hesshaimer, V.: Radiocarbon—a unique tracer of global carbon cycle dynamics. *Radiocarbon*, 42\(1\), 69-80, 2000.](#)

- Lewandowska, A.M., Boyce, D.G., Hofmann, M., Matthiessen, B., Sommer, U. and Worm, B.: Effects of sea surface warming on marine plankton. *Ecology Letters*, 17(5), 614-623, 2014.
- Lévy, M., Estublier, A., and Madec, G.: Choice of an advection scheme for biogeochemical models. *Geophysical Research Letters*, 28(19), 3725-3728, 2001a.
- 5 Lévy, M., Klein, P., and Treguier, A.M.: Impact of sub-mesoscale physics on production and subduction of phytoplankton in an oligotrophic regime. *Journal of Marine Research*, 59(4), 535-565, 2001b.
- Li, J., Y.M. and Wang, B.: Evaluation of NESMv3 and CMIP5 Models' Performance on Simulation of Asian-Australian Monsoon. *Atmosphere*, 9(9), 327, 2018.
- Lin, H., Kuzminov, F.I., Park, J., Lee, S., Falkowski, P.G., and Gorbunov, M.Y.: The fate of photons
10 absorbed by phytoplankton in the global ocean. *Science*, p.aab2213, 2016.
- Longhurst, A., Sathyendranath, S., Platt, T., and Caverhill, C.: An estimate of global primary production in the ocean from satellite radiometer data. *Journal of Plankton Research*, 17(6), 1245-1271, 1995.
- Madec, G.: NEMO ocean engine. Note du pôle de modélisation, No. 27, Institut Pierre-Simon Laplace (IPSL), France, 2012.
- 15 Maritorena, S., d'Andon, O.H.F., Mangin, A. and Siegel, D.A.: Merged satellite ocean color data products using a bio-optical model: Characteristics, benefits and issues. *Remote Sensing of Environment*, 114(8), 1791-1804, 2010.
- McCarthy, J. J.: The kinetics of nutrient utilization, *Can. B. Fish. Aquat. Sci.*, 210, 211–233, 1980.
- McGillicuddy Jr, Robinson, A.R., Siegel, D.A., Jannasch, H.W., Johnson, R., Dickey, T.D., Mcneil, J.,
20 Michaels, A.F., and Knap, A.H.: Influence of mesoscale eddies on new production in the Sargasso Sea. *Nature*, 394(6690), 263-266, 1998.
- Moore, J. K., Lindsay, K., Doney, S. C., Long, M. C., and Misumi, K.: Marine Ecosystem Dynamics and Biogeochemical Cycling in the Community Earth System Model [CESM1(BGC)]: Comparison of the 1990s with the 2090s under the RCP4.5 and RCP8.5 Scenarios. *Journal of Climate*, 26, 9291-
25 9312, 2013.
- Morel, A: Light and marine photosynthesis: a spectral model with geochemical and climatological implications. *Progress in oceanography*, 26(3), 263-306, 1991.
- NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing Group:

- Sea-viewing Wide Field-of-view Sensor (SeaWiFS) Ocean Color Data, NASA OB.DAAC, 2014.
- ~~NOAA ESRL Global Monitoring Division: Atmospheric Carbon Dioxide Dry Air Mole Fractions from quasi-continuous measurements at Mauna Loa, Hawaii. Compiled by K.W. Thoning, D.R. Kitzis, and A. Crotwell. National Oceanic and Atmospheric Administration (NOAA), Earth System Research Laboratory (ESRL), Global Monitoring Division (GMD): Boulder, Colorado, USA. Version 2018-10, 2017. <http://dx.doi.org/10.7289/V54X55RG>.~~
- 5
- Olonscheck, D., Hofmann, M., Worm, B. and Schellnhuber, H.J.: Decomposing the effects of ocean warming on chlorophyll a concentrations into physically and biologically driven contributions. *Environmental Research Letters*, 8(1), p.014043, 2013.
- 10
- Pierce, D.W., Gleckler, P.J., Barnett, T.P., Santer, B.D., and Durack, P.J.: The fingerprint of human-induced changes in the ocean's salinity and temperature fields. *Geophysical Research Letters*, 39, L21704, 2012.
- Resplandy, L., Levy, M., Bopp, L., Echevin, V., Pous, S., Sarma, V.V.S.S., and Kumar, D.: Controlling factors of the oxygen balance in the Arabian Sea's OMZ. *Biogeosciences*, 9, 5095-5109, 2012.
- 15
- Rhein, M., Rintoul, S.R., Aoki, S., Campos, E., Chambers, D., Feely, R.A., Gulev, S., Johnson, G.C., Josey, S.A., Kostianoy, A., Mauritzen, C., Roemmich, D., Talley, L.D., and Wang, F.: Observations: Ocean. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 20
- Roy, T., Bopp, L., Gehlen, M., Schneider, B., Cadule, P., Frölicher, T.L., Segschneider, J., Tjiputra, J., Heinze, C., and Joos, F.: Regional impacts of climate change and atmospheric CO₂ on future ocean carbon uptake: A multimodel linear feedback analysis, *J. Clim.*, 24(9), 2300– 2318, 2011.
- Sabine, C.L., Feely, R.A., Gruber, N., Key, R.M., Lee, K., Bullister, J.L., Wanninkhof, R., Wong, C.S.L., Wallace, D.W., Tilbrook, B., Millero, F.J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A.F.: The oceanic sink for anthropogenic CO₂. *Science*, 305(5682), 367-371, 2004.
- 25
- Sarmiento, J. I., and N. Gruber.: *Ocean Biogeochemical Dynamics*. Princeton University Press, Princeton, NJ, USA, 2006.

- Schneider, B., Bopp, L., Gehlen, M., Segschneider, J., Frölicher, T.L., Cadule, P., Friedlingstein, P., Doney, S.C., Behrenfeld, M.J., and Joos, F.: Climate-induced interannual variability of marine primary and export production in three global coupled climate carbon cycle models. *Biogeosciences*, 5(2), 597-614, 2008.
- 5 Schwinger, J., Tjiputra, J. F., Heinze, C., Bopp, L., Christian, J.R., Gehlen, M., Ilyina, T., Jones, C.D., y Mélia, D.S., Segschneider, J., Séférian, R., and Totterdell, I.: Nonlinearity of ocean carbon cycle feedbacks in CMIP5 Earth system models. *Journal of Climate*, 27(11), 3869-3888, 2014.
- [Schwinger, J. and Tjiputra, J.: Ocean carbon cycle feedbacks under negative emissions. *Geophysical Research Letters*, 45\(10\), 5062-5070, 2018.](#)
- 10 Séférian, Roland, Laurent Bopp, Marion Gehlen, James C. Orr, Christian Ethé, Patricia Cadule, Olivier Aumont, David Salas y Mélia, Aurore Voltaire, and Gurvan Madec.: Skill assessment of three earth system models with common marine biogeochemistry. *Climate Dynamics*, 40(9-10), 2549-2573, 2013.
- [Skinner, L.C., Primeau, F., Freeman, E., de la Fuente, M., Goodwin, P.A., Gottschalk, J., Huang, E., McCave, I.N., Noble, T.L., and Scrivner, A.E.: Radiocarbon constraints on the glacial ocean circulation and its impact on atmospheric CO₂. *Nature Communications*, 8, 16010, 2017.](#)
- 15
- Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., Dukowicz, J., Eden, C., Fox-Kemper, B., Gent, P., Hecht, M., Jayne S., Jochum M., Large W., Lindsay K., Maltrud M., Norton N., Peacock S., Vertenstein M., and Yeager S.: The parallel ocean program (POP) reference manual ocean component of the community climate system model (CCSM) and community earth system model (CESM). Rep. LAUR-01853, 141, 1-140, 2010.
- 20
- Steinacher, M., Joos, F., Frölicher, T.L., Bopp, L., Cadule, P., Cocco, V., Doney, S.C., Gehlen, M., Lindsay, K., Moore, J.K., and Schneider, B.: Projected 21st century decrease in marine productivity: a multi-model analysis. *Biogeosciences*, 7(3), 979-1005, 2010.
- 25
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornblueh, L., Lohmann, U., Pincus, R., Reichler, T., and Roeckner, E.: The atmospheric component of the MPI-M Earth System Model: ECHAM6, *J. Adv. Model. Earth Syst.*, 5, 46–172, 2012.

- Takahashi, T., Broecker, W.S., and Langer, S.: Redfield ratio based on chemical data from isopycnal surfaces. *Journal of Geophysical Research: Oceans*, 90(C4), 6907-6924, 1985.
- Takahashi, T., Sutherland, S.C., Wanninkhof, R., Sweeney, C., Feely, R.A., Chipman, D.W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D.C.E., Schuster, U., Metzl, N.,
5 Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Kortzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T.S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C.S., Delille, B., Bates, N.R., and de Baar, H.J.W.: Climatological mean and decadal change in surface ocean pCO₂, and net sea-air CO₂ flux over the global oceans. *Deep Sea Research Part II: Topical Studies in Oceanography*, 56(8-10), 554-577, 2009.
- 10 Taylor, K.E.: Summarizing multiple aspects of model performance in a single diagram. *Journal of Geophysical Research: Atmospheres*, 106(D7), 7183-7192, 2001.
- Tjiputra, J. F., Assmann, K., Bentsen, M., Bethke, I., Otterå, O.H., Sturm, C., and Heinze, C.: Bergen Earth system model (BCM-C): model description and regional climate-carbon cycle feedbacks assessment. *Geoscientific Model Development*, 3(1), 123-141, 2010.
- 15 Tjiputra, J. F., Roelandt, C., Bentsen, M., Lawrence, D. M., Lorentzen, T., Schwinger, J., Seland, Ø., and Heinze, C.: Evaluation of the carbon cycle components in the Norwegian Earth System Model (NorESM). *Geoscientific Model Development*, 6, 301-325, 2013.
- Uitz, J., Claustre, H., Gentili, B., and Stramski, D.: Phytoplankton class-specific primary production in the world's oceans: seasonal and interannual variability from satellite observations. *Global*
20 *Biogeochemical Cycles*, 24, GB3016, 2010.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean. *Journal of Geophysical Research: Oceans*, 97(C5), 7373-7382, 1992.
- Wanninkhof, R., Park, G.H., Takahashi, T., Sweeney, C., Feely, R.A., Nojiri, Y., Gruber, N., Doney, S.C., McKinley, G.A., Lenton, A., Le Quere, C., Heinze, C., Schwinger, J., Graven, H., and
25 Khatiwala, S.: Global ocean carbon uptake: magnitude, variability and trends. *Biogeosciences*, 10, 1983-2000, 2013.
- Weaver, A. J., Sedláček, J., Eby, M., Alexander, K., Crespin, E., Fichet, T., Philippon - Berthier, G., Joos, F., Kawamiya, M., Matsumoto, K., Steinacher, M., Tachiiri, K., Tokos, K., Yoshimori, M., and

- Zickfeld, K.: Stability of the Atlantic meridional overturning circulation: A model intercomparison. *Geophysical Research Letters*, 39, L20709, 2012.
- Weiss, R.F.: August. The solubility of nitrogen, oxygen and argon in water and seawater. In *Deep Sea Research and Oceanographic Abstracts*, 17(4), 721-735, 1970.
- 5 Westberry, T., Behrenfeld, M.J., Siegel, D.A. and Boss, E.: Carbon-based primary productivity modeling with vertically resolved photoacclimation. *Global Biogeochemical Cycles*, 22, GB2024, 2008.
- Whitney F A.: Nutrient variability in the mixed layer of the subarctic Pacific Ocean, 1987–2010. *Journal of Oceanography*, 67(4), 481-492, 2011.
- 10 Yang, Y.M., and Wang, B.: Improving MJO simulation by enhancing the interaction between boundary layer convergence and lower tropospheric heating. *Climate Dynamics*, 1-23, 2018a.
- Yang, Y.M., Wang, B. and Li, J.: Improving Seasonal Prediction of East Asian Summer Rainfall Using NESM3. 0: Preliminary Results. *Atmosphere*, 9(12), 487, 2018b.
- Yi, C., Gong, P., Xu, M., and Qi, Y.: The effects of buffer and temperature feedback on the oceanic
15 uptake of CO₂. *Geophysical Research Letters*, 28(5), 751-754, 2001.
- Yool, A., Popova, E.E., and Anderson, T. R.: Medusa-1.0: a new intermediate complexity plankton ecosystem model for the global domain. *Geoscientific Model Development*, 4(2), 381-417, 2011.
- Zickfeld, K., Eby, M., and Weaver, A.J.: Carbon-cycle feedbacks of changes in the Atlantic meridional
20 overturning circulation under future atmospheric CO₂. *Global Biogeochemical Cycles*, 22, GB3024, 2008.
- Zickfeld, K., Eby, M., Matthews, H.D., Schmittner, A., and Weaver, A.J.: Nonlinearity of carbon cycle feedbacks. *Journal of Climate*, 24(16), 4255-4275, 2011.

5

10

Table 1. Global ocean anthropogenic CO₂ uptake simulated by NESM v3 during different periods compared against data-based estimates (Ciais et al., 2013). (The uncertainty ranges are given after ±) (It is noted that the pre-industrial time in this study represents the year 1850 while it represents 1750 in IPCC AR5).

	Pre-industrial- 2011 Cumulative PgC	1980-1989 PgC yr ⁻¹	1990-1999 PgC yr ⁻¹	2000-2009 PgC yr ⁻¹	2002-2011 PgC yr ⁻¹
IPCC AR5	155±30	2.0±0.7	2.2±0.7	2.3±0.7	2.4±0.7
NESM v3	137.2	1.7	2.0	2.3	2.3

15

20





Figure 1. Annual mean (averaged over from the year 1985 to 2004) upper ocean (averaged in the upper 100 m) distribution of phosphate (a1, a2), nitrate (b1, b2) and silicate (c1, c2) from the NESM v3 simulations (FC-HistSSP) and the WOA18 observation dataset (in the unit of mmol m^{-3}). The difference between model simulation and observation are also shown (a3, b3, and c3)

5






Figure 2. The latitude-depth distributiondistributions of silicate (a), phosphate (b), and nitrate (c) averaged fromover the years 1985 to 2014 from the NESM v3 simulation (FC-HistSSP) and the WOA18 observation dataset (with in the unit of mmol m^{-3}). a, b, and c represent the silicate, phosphate, and nitrate, respectively. a1, a2, b1, b2, c1, and c2 represent the distributions in the Pacific Ocean, a3, a4, b3, b4, c3, and c4 represent the distributions in the Atlantic Ocean, and a5, a6, b5, b6, c5, and c6 represent the distributions in the Global Ocean.




Figure 3. Latitude-depth ~~distribution~~distributions of preformed and regenerated phosphate concentration (mmol m^{-3}) in the Pacific, Atlantic, and global Ocean averaged over 1985 to 2014. The results diagnosed from the WOA18 observation dataset are shown in the first and third rows (a1, a3, a5, b1, b3, and b5). The difference (model minus observation) between NESM v3 simulations and observation are shown in the second and the last rows (a2, a4, a6, b2, b4, and b6). The panels from a1 to a6 show the preformed component, and the panels from b1 to b6 show the regenerated component.




Figure 4. ~~The diagnosed pattern~~ Diagnosed patterns of nutrients limitation from the NESM v3 simulation (FC-HistSSP). The limitation maps are shown over the annual time scale averaged over the years 1985 to 2014 for phytoplankton (a) and diatoms (b). Different colors represent ~~the factor~~ different factors that most ~~limits~~ limit phytoplankton growth. Replete means nutrient concentrations are sufficient for the phytoplankton growth (growth rate is greater than 90% of their maximal growth rate).

10

15

20

25




Figure 5. Annual mean surface chlorophyll concentration (mg Chl m⁻³) from the NESM v3 FC-HistSSP simulations (a; averaged over 1998 to 2014), the SeaWiFS dataset (b; averaged over 1998 to 2010), the GlobColour merged dataset (c; averaged over 1998 to 2014), and OCCCI merged dataset (d; averaged over 1998 to 2014). The biases between the NESM v3 simulation and observations are shown in the e, f, and g.




Figure 6. Annual mean ~~distribution~~distributions of vertically integrated net primary production (g C m⁻² yr⁻¹) averaged from 2003 to 2014 from the NESM v3 FC-HistSSP simulations (a) and MODIS observation-based estimates (b: VGPM-MODIS; c: CBPM-MODIS; d: Eppley-MODIS). The ~~bias~~biases of model simulations and observations are shown in the right three panels (e, f, g).




Figure 7. Annual mean distributions of upper ocean mean (0-100 m) alkalinity (mmol m^{-3}) (a, b) and DIC (mmol C m^{-3}) (c, d) averaged from 1985 to 2014 from the NESM v3 simulations (a, c) and GLODAP v2 (b, d). The **bias** of the NESM v3 simulations and observations are shown in the last two panels (e, f).




Figure 8. The latitude-depth distributions of the alkalinity (a, in the unit of mmol m^{-3}) and DIC (b, in the unit of mmol C m^{-3}) averaged from 1985 to 2014 from the FC-run-HistSSP simulation are compared with GLODAP v2 observations. The observations are shown in the first and the third rows (a1, a3, a5, b1, b3, b5), whereas the biases (model minus observation) are shown in the second and the last rows (a2, a4, a6, b2, b4, and b6).




Figure 9. Taylor diagram comparing statistical patterns of annual mean biogeochemical fields averaged over the upper 100m ocean between the NESM v3 simulation (FC-HistSSP) and corresponding observations, including upper 100-m ocean nitrate, phosphate, silicate, alkalinity, chlorophyll concentration, and vertically-integrated net primary production. Nutrient concentrations are compared with WOA 18; DIC and alkalinity concentration are compared with GLODAP v2; NPP is compared with the CbPM, and; chlorophyll is compared with SeaWiFS. All fields are normalized by the standard deviation of corresponding observations. Thus, observation fields have a standard deviation of one, which is represented by REF. The distance between the model points and the reference point indicates the root-mean-square (RMS) difference between model simulation and observations. The solid cycle represents the results diagnosed from the NESM v3, and the triangle represents the results from the IPSL, that shares the same marine biogeochemical model as the NESM v3. The NPP of IPSL is not shown here.




Figure 10. Model-simulated sea-air CO₂ flux ($\text{g C m}^{-2} \text{yr}^{-1}$) in the year 2000 compared with data-based observational estimates (Takahashi et al., 2009). Spatial distributions of model simulation (a), observation (b), ~~the difference between model and observation (d), and~~ zonal mean pattern of model simulation and observation (c), ~~and the difference between model and observation (d)~~. Positive values represent CO₂ flux out of the ocean, and negative values represent CO₂ flux into the ocean.




Figure 11. Vertically integrated column inventory of anthropogenic DIC (mol C m^{-2}) from the simulation (a, c), GLODAP v2 (b) and GLODAP v1 and GLODAP v2 observation (b, d). Model simulation results are averaged from 2000 to 2004 (a, represent the period around 2002) and from 1992 to 1996 (c, while the observation is, represent the period around 1994). The anthropogenic DIC from GLODAP v2 are normalized to the year 2002 (b) and), while that from GLODAP v1 are normalized to the year 1994 (d).







Figure 12. Zonal mean latitude-depth ~~distribution~~distributions of anthropogenic DIC (mmol C m^{-3}) from the model simulation (a1: Atlantic, a2: Pacific, and a3: Global) and data-based estimates (GLODAP v2) (b1: Atlantic, b2: Pacific, and b3: Global). Model simulation results are averaged from the year 2000 to 2004 (a), while the observation is normalized to the year 2002 (b).






5 **Figure 13.** Time series of changes in climate changesfields (relative to control simulation) from 1850 to 2100 for the simulation of fully-coupled, biogeochemically-coupled BC-HistSSP, FC-HistSSP, and radiatively-coupled simulations RC-HistSSP. (a) global and annual mean surface air temperature (in the unit of K), (b) global and annual mean mixed layer depth (in the unit of meter, the mixed layer depth is defined to be the depth where the difference in potential density is 0.01 kg m^{-3} relative to the sea surface) and (c) Atlantic meridional overturning circulation index (in the unit of Sv, maximum zonal mean stream function in the Atlantic Ocean at 30°N).



5 **Figure 14.** The NESM v3 simulated (a, in the unit of PgC yr^{-1}) annually anthropogenic oceanic CO₂ uptake change (a, in the unit of PgC yr^{-1}) and (b, in the unit of PgC) cumulative oceanic CO₂ uptake (b, in the unit of PgC) for the simulations RC-HistSSP, BC-HistSSP, FC-HistSSP, and the linear sum of BC-HistSSP and RC-HistSSP from 1850 to 2100.



5 **Figure 15.** Spatial ~~distribution~~distributions of anthropogenic sea-air CO₂ flux (in the unit of g C m⁻² yr⁻¹) at the end of the 21st century (mean of 2091-2100) from the BC-HistSSP (a) BC, RC-HistSSP (b) RC, and FC-HistSSP (c) FC simulations, respectively. Also shown is the difference between FC-HistSSP simulation and the sum of RC-HistSSP and BC-HistSSP simulations (FC-RC-BC). Positive values represent CO₂ flux out of the ocean, and negative values represent CO₂ flux into the ocean.

10




Figure 16. The cumulative oceanic CO₂ uptake against (a) the atmospheric CO₂ in the BC-HistSSP simulation (a) and (b) the global mean surface air temperature change in the RC-HistSSP simulation (b). Also shown is the time evolution of the diagnosed (c) carbon-concentration sensitivity parameter as a function of atmospheric CO₂ (c) and (d) carbon-climate sensitivity parameter as a function of global mean surface air temperature change (d). All the parameters in this figure are diagnosed from HistSSP scenarios

5




Figure 17. Same as Figure 16, but for the 1ptCO₂ runs.