

Dear GMD editor,

Thank you for your comments. We revised the on the manuscript accordingly and below is a list of the revisions we made. Attached to document is a marked-up version of the latest manuscript with change tracking.

1. - page 28, line 90-95: Please elaborate a bit more on recent studies on geographic neural networks (i.e. provide more details on these studies). Otherwise, it is hard to understand why (and on what basis) you compare your method with these studies in the following paragraphs. Also note that the first sentence in this paragraph "With increasing..." is incomplete and does not provide a good introduction to a comparison with neural network models (i.e., this comparison comes out of the blue).

Reply: We revised and extended the introduction to GNNWR and GTNNWR. Please check the updated text at lines 490-498 on page 28.

2. - page 29, line 21: The sentence "Thus, it cannot..." is not complete. Please revise.

Reply: This sentence was revised. Please check lines 524-525 on page 29.

Thanks again for your time.

Sincerely,

Xiaogang (Marshall) Ma

1 A Spatiotemporal Weighted Regression Model (STWR v1.0) for 2 Analyzing Local Non-stationarity in Space and Time

3 Xiang Que^{1,2}, Xiaogang Ma^{2,*}, Chao Ma^{2,*}, Qiyu Chen³

4 ¹ Computer and Information College, Fujian Agriculture and Forestry University, Fuzhou, Fujian, China

5 ² Department of Computer Science, University of Idaho, 875 Perimeter Drive MS 1010, Moscow, ID 83844-1010, USA

6 ³ School of Computer Science, China University of Geosciences (Wuhan), 388 Lumo Road, Wuhan 430074, China

7 *Corresponding author. Email: max@uidaho.edu (Xiaogang Ma); chao@uidaho.edu (Chao Ma)

8
9 **Abstract:** Local spatiotemporal non-stationarity occurs in various natural and socioeconomic processes. Many studies have
10 attempted to introduce time as a new dimension into the geographically weighted regression model (GWR), but the actual
11 results are sometimes not satisfied or even worse than the original GWR model. The core issue here is a mechanism for
12 weighting effects of both temporal variation and spatial variation. In many geographical and temporal weighted regression
13 models (GTWR), the concept of time distance has been inappropriately treated as time interval. Consequently, the combined
14 effect of temporal and spatial variation is often inaccurate in the resulting spatiotemporal kernel function. This limitation
15 restricts the configuration and performance of spatiotemporal weights in many existing GTWR models. To address this
16 issue, we propose a new spatiotemporal weighted regression (STWR) model and the calibration method for it. A highlight of
17 STWR is a new temporal kernel function, in which the method for temporal weighting is based on the degree of impact from
18 each observed point to a regression point. The degree of impact, in turn, is based on the rate of value variation of the nearby
19 observed point during the time interval. The updated spatiotemporal kernel function is based on a weighted combination of
20 the temporal kernel with a commonly used spatial kernel (Gaussian or bi-square) by specifying a linear function of spatial
21 bandwidth versus time. Three simulated datasets of spatiotemporal processes were used to test the performance of GWR,
22 GTWR and STWR. Results show that STWR significantly improves the quality of fit and accuracy. Similar results were

23 obtained by using real-world data for the precipitation hydrogen isotopes ($\delta^2\text{H}$) in Northeastern United States. The Leave-
24 one-out cross-validation (LOOCV) test demonstrates that, comparing with GWR, the total prediction error of STWR is
25 reduced by using recent observed points. Prediction surfaces of models in this case study show that STWR is more localized
26 than GWR. Our research validates the ability of STWR to take full advantage of all the value variation of past observed
27 points. We hope STWR can bring fresh ideas and new capabilities for analyzing and interpreting local spatiotemporal non-
28 stationarity in many disciplines.

29

30 **Key words:** Geographical and temporal weighted regression; Geographically weighted regression; Temporal non-
31 stationarity; Spatial analysis; Spatiotemporal variations; Spatiotemporal weighted regression.

32

33 **1. Introduction**

34 Time, space and attributes are three essential characteristics in geographic entities, and they are recorded to reflect the state
35 and evolution of various real-world phenomena and processes. Because space and time frame all aspects of the discipline of
36 geography (Goodchild, 2013), it is important to observe the spatiotemporal variations and explore appropriate analytical
37 methods to study and reason the internal mechanisms and evolutionary laws. In recent years, new platforms and instruments
38 have brought increasingly massive spatiotemporal data, such as the time- and geo-tagged sensor monitoring records and
39 remote sensing images. Those big data create great opportunities for studying human and environmental dynamics from
40 different perspectives, such as the patterns of human behavior (Chen et al., 2011), environmental risk assessment (Sun et al.,
41 2015), and disease outbreaks (Takahashi et al., 2008). Nevertheless, although spatiotemporal modeling has been a long-term
42 research focus in the field of geographical information science (GIScience) (Cressie, 1991; Cressie and Wike, 2015), the
43 models are not mature yet and challenges still exist (Fotheringham et al., 2015), which call for further work.

44 In this paper, the technological development and discussion focus on modeling local spatiotemporal variations within
45 the framework of geographically weighted regression (GWR). GWR is a method for modeling spatially heterogeneous

46 processes (Brunsdon et al., 1996, 1998; Fotheringham et al., 2003). It has been applied in a variety of areas, such as climate
47 science (Brown et al., 2012), geology (Atkinson et al., 2003), mineral exploration (Wang et al., 2015), transportation analysis
48 (Cardozo et al., 2012), crime studies (Cahill and Mulligan, 2007; Wheeler and Waller, 2009), environmental science (Mennis
49 and Jordan, 2005), and house price modeling (Fotheringham et al., 2015). GWR calibrates a separate regression model at
50 each location through a data-borrowing scheme, in which distance-weights can be calculated by drawing on data from
51 neighboring observations of each regression point (Fraser et al., 2012). This operation complies with Tobler's first law of
52 geography - "Everything is related to everything else, but near things are more related than distant things" (Tobler, 1970).

53 Numerous studies have been devoted to incorporating the temporal dimension into spatial regression (Pace et al., 2000;
54 Gelfand et al., 2004; Crespo et al., 2007; Cressie and Wikle, 2015). However, most of these studies assume that temporal
55 effects are constant over space from a global perspective of modeling (Fotheringham et al., 2015). To address that issue,
56 Crespo et al. (2007) extended GWR by developing spatiotemporal bandwidths that account for varying local spatial effects
57 across time. Huang and Wu (2010, 2014) proposed a geographical and temporal weighted regression model (GTWR) with a
58 method of measuring the spatiotemporal 'closeness' and a parameter ratio τ to deal with different measured units in time
59 and space. Although the approach can address the issue to some extent, Fotheringham et al. (2015) pointed out that a sole
60 measurement of integrated spatial and temporal distances can be misleading as location and time are usually measured at
61 different scales, and he stated that the calculation of distance in three dimensions (time and two-dimensional space) remains
62 a challenge.

63 A spatiotemporal kernel function, which consists of mixed spatial and time-decay bandwidths, was proposed by
64 Fotheringham et al. (2015). Nevertheless, the stepwise strategy applied in this function for bandwidth optimization does not
65 always seem reasonable. In practice, this function needs to first find and fix an optimized spatial bandwidth, then it will find
66 the optimized temporal bandwidth. After that, the spatiotemporal weight will be calculated. This stepwise search process
67 means that the function is not able to optimize both temporal and spatial bandwidths at the same time. However, a more
68 reasonable thought is that the spatiotemporal bandwidth and its weight are simultaneously affected by both spatial and

69 temporal effects of a process. There should be ways to further improve the spatiotemporal kernel function in Fotheringham
70 et al. (2015).

71 The aim of this paper to develop a better methodology for the spatiotemporal kernel function. Following Tobler's first
72 law, we propose an algorithm, the spatiotemporal weighted regression (STWR). In STWR, the velocity of value change is
73 higher related if they were in near time and space. Therefore, STWR can borrow data not only from nearby locations, but
74 also from nearby value variation through time. The latter is what we call as "time distance" in STWR. The time distance is
75 not the concept of time interval, but the rate of value variation through time. It is a kind of value change that reflects the
76 temporal effect of nearby points to the regression point. Accordingly, our local spatiotemporal regression analysis model can
77 take advantage of the variation in data to identify temporal non-stationarity, which is an advantage comparing with GWR
78 and GTWR.

79 Before giving more details about STWR, we can further clarify the meaning of a few concepts. A common issue in the
80 existing GTWR models is that they use the concept of time interval, instead of the above-mentioned "time distance", to
81 calculate temporal and spatiotemporal weights. A time interval is the period between two observed time stages. A time
82 distance, in the context of STWR, is the rate of value variation between an observed point and a regression point through a
83 time interval. We can think about the following scenario for a group of points. The values of some points do not change or
84 change slightly from time A to time B, while a few other points may change greatly in that period. However, many GTWR
85 models ignore the difference in the value changes of observed points during a period of time, and regard that all these points
86 have the same temporal effect to their neighbor regression point. It is hard to believe that some unchanged observations
87 constantly affect their nearby regression points during the observed time interval. Intuitively, different variations of the
88 observed points have different temporal effects. For example, the faster the house price of a point change, the stronger the
89 temporal effect is to the house price at its nearby point. Moreover, the rate of value changes at different observed points
90 (time non-stationary) may also have spatial heterogeneity. The data values observed at different points are results of mixed

91 spatiotemporal effects and some other unknown factors (including errors). Therefore, using only time interval in the
92 calculation of temporal and spatiotemporal weights might interpret local spatiotemporal effect imprecisely.

93 There are other issues in the temporal kernel functions and the multiplication form of spatial and temporal kernels used
94 by the existing GTWR models (Huang et al., 2010; Wu et al., 2014; Fotheringham et al., 2015). When calculating the
95 spatiotemporal effect, these models generally use time intervals and the common kernel functions to calculate temporal
96 weights, such as Gaussian kernel or bi-square kernel. However, an appropriate temporal kernel function should not be the
97 same as the spatial kernel function, because space is in two or three dimensions while time is in one dimension and one
98 direction. Each regression point can borrow observed points from any directions in space but only use points from the past
99 rather than from the future. Moreover, the integrated spatiotemporal weights might be underestimated in these GTWR
00 models by using a multiplication of the spatial and temporal weights. Because both the spatial weights and the temporal
01 weights range from 0 to 1, and the multiplied weight value is never bigger than the smaller one before multiplying, which
02 means that the composite spatiotemporal impacts are never greater than the single spatial impacts and the single temporal
03 impacts. However, the real combined spatiotemporal impacts, may be higher than the single spatial impacts or the temporal
04 impacts, or at least may be higher than the smaller ones. The multiplication formulation of spatiotemporal kernel in GTWR
05 also makes the calculated weight decay faster.

06 The above-mentioned limitations and issues in GWR and GTWR are the driving force behind our development of
07 STWR. The remainder of this article is organized as follows. Section 2 introduces the STWR model formulation, including
08 temporal kernel and spatiotemporal kernel functions. Section 3 describes the methods for bandwidth selection and calibration
09 when STWR is in operation. Section 4 presents results of applying GWR, GTWR and STWR to three sets of simulated data.
10 Section 5 presents experiment results with real-world precipitation hydrogen isotope data. In Section 6, we close the article
11 with a summary of the key findings and a few thoughts for future research.

12

13 **2. The Core Model of STWR**

14 **2.1 The strategy of time distance decay**

15 Since GWR is the background of our work, it is helpful to first give a brief overview of the GWR framework. The basic
16 formulation of GWR can be described in two equations below (Fotheringham et al., 2003).

17
$$y_i = \beta_0(u_i, v_i) + \sum_k \beta_k(u_i, v_i)x_{ik} + \varepsilon_i \quad (1)$$


18
$$\beta_k(u_i, v_i) = (X^T W(u_i, v_i) X)^{-1} X^T W(u_i, v_i) y \quad (2)$$

19 In Equation 1, y_i is a response variable of regression point i at a location with the coordinates (u_i, v_i) . x_{ik} is the k^{th}
20 independent variable, and ε_i denotes the error term for the i^{th} observed point. A key difference between GWR and the
21 traditional global regression method, such as Ordinary Least Squares (OLS), is that GWR allows the coefficient $\beta_k(u_i, v_i)$
22 vary spatially to identify spatial heterogeneity. Equation 2 represents the GWR calibration in a matrix form. $W(u_i, v_i)$ is a
23 diagonal weighting matrix specific to location i , which is calibrated by a specified kernel function with a given bandwidth.
24 Every element w_i in the weighting matrix reflects the impact from another observed point to the regression point. A bigger
25 w_i value means a higher impact.

26 GWR has a strategy of spatial distance decay impact on a regression point (Brunsdon et al., 1998; Fotheringham et al.,
27 2003). A similar “time distance decay” strategy was also discussed in several recent GTWR models (Crespo et al., 2007;
28 Huang et al., 2010; Wu et al., 2014; Fotheringham et al., 2015). Yet, those models did not fully reflect the effect of time
29 distance decay. Sample points are observed at different time stages, and those data points closer in time distance to a
30 regression point have more impact on the regression point than those farther away. The time distance refers to the value
31 variation rate between an observed point and a regression point during a certain time interval. For example, in Fig. 1, there
32 are four time stages from old to new: T-s, T-q, T-p and T. Through a fitting and calibration process, the spatiotemporal
33 bandwidth will be fitted, and the spatiotemporal effects (weights) from observed points to a regression points at time stage T
34 will be calculated by a specific spatiotemporal kernel function. Then, in prediction, the value of a regression point at time
35 stage T can be estimated. Thus, the observed points at time stage T only have spatial effect on the regression point (Fig. 1).
36 There is temporal effect from data points at time stages T-p and T-q (shown as stars, pentagons and triangles in the planes of

37 T-p and T-q in Fig. 1), within a certain spatial bandwidth b_{ST} at each time stage, to the regression point. The time distance
 38 decay should reflect that different variations of the observed points have different temporal effects. However, as mentioned
 39 in the previous section, many existing GTWR models have applied a strategy of time interval decay instead of time distance
 40 decay. Consequently, they regard that all the observed points have the same temporal effect to their neighbor regression
 41 point.

42



43

44 **Fig. 1.** Spatiotemporal impacts of observed points with different rates of value change on a regression point at time stage T.
 45 Temporal bandwidth is the length of time from the intersection point A of the spatiotemporal bandwidth and the time line to
 46 the regression point. Spatial bandwidth and spatiotemporal bandwidth are illustrated in the figure legend.

47

48 Compared to existing GTWR models, the time distance decay strategy of STWR considers the effect of different
 49 variations of observed points through time. For example, some data points may have higher impact on the regression point,
 50 though their spatial distance is farther than other points. Fig. 1 illustrates that the locations of some star-shape points are
 51 farther away from the regression point than some pentagon-shape points at time stage T-p, which denotes that there exist

52 mixed impacts (spatial impact and temporal impact) on the regression point. The temporal impacts depend on the rate of
53 value variation, which is the value difference between the observed point and the regression point divided by a time interval
54 (e.g., [T-p, T] and [T-q, T-p] each is a time interval). If the observed time stage is too long ago or the rate of value variation
55 is too small, and exceeds the limit of optimized temporal bandwidth for the regression point (as shown by observations at
56 time stage T-s), the data points at this time stage may have no impact on the regression point. Even though some of those
57 data points may have huge difference in value and are close to the regression point in space, they are not within the range of
58 the optimized temporal bandwidth. Spatial bandwidths also vary along the time line, and usually the bandwidth gets larger
59 when the observation time is closer to the time stage of the regression point (Fig. 1).

60 2.2 The spatiotemporal kernel function of STWR

61 We assume that a set of observed points $O_{\Delta t} = \{O_{N_t}, O_{N_{t-1}}, \dots, O_{N_{t-q}} | \Delta t = [t - q, t]\}$ are collected during a certain time
62 interval Δt in a study area, where t represents the current time stage and $N_{t-i}, i \in \{0, 1, 2, \dots, q\}$ ($N_t = N_{t-0}$) denotes the
63 number of observed points at each recorded time. As the idea described above, we can borrow neighbor points in space and
64 their value variation during certain recent time intervals, so we can still use Equation 1 to generate local estimates. The
65 weight matrix W in GWR usually depends on the spatial kernel (Fotheringham et al., 2015). In STWR, we need to consider
66 the temporal effect, so the form of W is different from that in GWR. Correspondingly, we should have a spatiotemporal
67 kernel, which can be understood as a temporal extension based on the spatial kernel. However, if we use a multiplication
68 form to combine the temporal kernel and the spatial kernel (Huang et al., 2010; Wu et al., 2014; Fotheringham et al., 2015),
69 we will face the problem of time and space interaction as mentioned above in the Introduction section. To address that issue,
70 we design a weighted average form for the spatiotemporal kernel.

$$71 \quad w_{ijST}^t = (1 - \alpha)k_s(d_{sij}, b_{ST}) + \alpha k_T(d_{tij}, b_T), 0 \leq \alpha \leq 1 \quad (3)$$

72 In Equation 3, w_{ijST}^t is the weight at time t and at the observed location j . k_s and k_T are the spatial and temporal kernel,
73 respectively, and they both have a value range of 0 to 1. α is an adjustable parameter to scale the temporal and spatial
74 effects, which can be optimized with the bandwidth selections. The role of parameter α is different from the scale parameter

75 τ ($\tau = \frac{\alpha}{\lambda}$) in GTWR (Huang et al., 2010). α is introduced here for adjusting the outputs of the spatial kernel k_s and the
 76 temporal kernel k_T , which means measuring the relative strength of the spatial and temporal impacts on the regression point.
 77 But the scale parameter τ is used for adjusting the inconsistency of the time distance and space distance, which cannot
 78 adjust the relative strength of k_s and k_T . d_{sij} and d_{tij} are the spatial (Euclidean) and temporal distance between the
 79 regression point i and an observed data point j , respectively. b_{sT} is the spatial bandwidth b_s at a certain time stage T , and
 80 b_T denotes the temporal bandwidth.

81 The time distance, as mentioned above, is not the time interval but the rate of value variation between an observed point
 82 and a regression point through a time interval. Following the time distance decay strategy in STWR, we can further derive
 83 the temporal kernel k_T as shown below.

$$84 \quad w_{ij\Delta t}^t = \begin{cases} \left[\frac{2}{1 + \exp\left(-\frac{[y_i(t) - y_j(t-q)]/\Delta t}{b_T}\right)} - 1 \right], & \text{if } 0 < \Delta t < b_T \\ 0, & \text{otherwise} \end{cases} \quad (4)$$

85 In Equation 4, $y_i(t) - y_j(t-q)$ is the subtraction of the regression point i 's observed value at t from the point j 's observed
 86 value at $t - q$, which denotes the value change during the time interval Δt . The internal part of the exponential function is
 87 negative, in order to make the weight $w_{ij\Delta t}^t$ range from 0 to 1. The faster the value change rate is, the bigger the weight is,
 88 which means that the time impact is larger. When the time interval Δt is out of the range $(0, b_T)$, the weight will be set to
 89 zero, which denotes that there is no impact because the observed variation is too far to affect the current moment. For
 90 example, if the price of a nearby house has changed a long time ago, it may have little or no impact on the present house
 91 price. But if the house price had a sharp change recently, it will have a big impact on the present house price. Therefore, the
 92 faster the rate of observed value changes and the shorter the time interval is, the greater the impact on the regression point
 93 will be. Compared with GTWR models, the advantage of STWR is that the temporal kernel function k_T can better leverage
 94 the variation data.

95 To calibrate the weight value w_{ijST}^t , we need a spatial kernel function. The most widely used kernel functions are bi-
 96 square and Gaussian (Fotheringham et al., 2003), which are given in Equations 5 and 6, respectively.

$$97 \quad \text{Bi-square: } w_{ijS} = \begin{cases} \left[1 - \left(\frac{d_{stij}}{b_S}\right)^2\right]^2, & \text{if } d_{stij} < b_S \\ 0, & \text{otherwise} \end{cases} \quad (5)$$

$$98 \quad \text{Gaussian: } w_{ijS} = \exp\left[-\frac{1}{2}\left(\frac{d_{stij}}{b_S}\right)^2\right] \quad (6)$$

99 In Equations 5 and 6, b_S is the spatial bandwidth. Derived from b_S and b_{ST} , b_{St} is the initial spatial bandwidth at the given
 100 time stage t of the regression point (i.e., t is the initial time for searching observed points in the past). Many functions can
 101 be specified for the change of spatial bandwidth during the time intervals. Because in most cases it will have smooth change
 102 during a certain short time interval, we assume that the spatial bandwidth changes linearly along with time, as defined
 103 below.

$$104 \quad b_{ST} = b_{St} - \tan\theta * \Delta t, \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2} \quad (7)$$

106 In Equation 7, $\tan\theta$ denotes the slope of spatial bandwidth change in correspondence to Δt , and b_{St} denotes the initial
 107 spatial bandwidth at t . Importing Equations 4 to 7, the calibration of Equation 3 can be further derived into Equations 8 and
 108 9, which are our spatiotemporal kernel functions in STWR. Equations 8 and 9 are based on the bi-square and Gaussian
 109 kernel, respectively. With the STWR spatiotemporal kernel, we only need to optimize the parameters α and θ instead of the
 110 spatial bandwidth b_{ST} . However, we shall traverse all the observed points at the initial time stage t to find the optimized
 111 spatial bandwidth b_{St} . Moreover, we shall also traverse all the time stages to find the optimized temporal bandwidth b_T .

$$w_{yST}^t = \begin{cases} \left[(1-\alpha) * \left[1 - \left(\frac{d_{sij}}{b_{St} - \tan \theta * \Delta t} \right)^2 \right] + \alpha * \left(2 / \left(1 + \exp\left(-\frac{|(y_{i(t)} - y_{j(t-q)}) / y_{j(t-q)}|}{\Delta t / b_T} \right) \right) - 1 \right) \right], \\ \text{if } \Delta t < b_T, \text{ and } d_{sij} < (b_{St} - \tan \theta * \Delta t) \\ 0, \text{ otherwise} \end{cases} \quad (8)$$

$$w_{yST}^t = \begin{cases} \left[(1-\alpha) * \exp\left[-\frac{1}{2} \left(\frac{d_{sij}}{b_{St} - \tan \theta * \Delta t} \right)^2 \right] + \alpha * \left(2 / \left(1 + \exp\left(-\frac{|(y_{i(t)} - y_{j(t-q)}) / y_{j(t-q)}|}{\Delta t / b_T} \right) \right) - 1 \right) \right], \\ \text{if } \Delta t < b_T, \text{ and } d_{sij} < (b_{St} - \tan \theta * \Delta t) \\ 0, \text{ otherwise} \end{cases} \quad (9)$$

14

15 3. STWR in Operation

16 3.1 Bandwidth selection and parameter estimation

17 Some goodness-of-fit diagnostics (Loader, 1999) are widely used in general GWR-based models, such as the cross-
 18 validation (CV) score (Cleveland, 1979; Bowman, 1984) and the Akaike Information Criterion (AIC) (Akaike, 1973;
 19 Akaike, 1998). For STWR, we use cross-validation (CV) as the default searching criteria and we also calculate the value of a
 20 corrected version of AIC (Hurvich et al., 1998), the AICc, which is defined bellow.

$$AIC_c = 2n \ln(\hat{\sigma}) + n \ln(2\pi) + n \left\{ \frac{n + tr(S)}{n - 2 - tr(S)} \right\} \quad (10)$$

22 In Equation 10, n is the sample size, $\hat{\sigma}$ is the estimated standard deviation of the error term, and $tr(S)$ denotes the trace of
 23 the hat matrix S (Hoaglin and Welsch, 1978).

24 Although there is no need to optimize spatial bandwidth b_{ST} of the past time stages in STWR, other parameters such as
 25 α and θ need to be optimized. Also, we should give the b_T and initial b_{St} through trials. For more potential combinations

of these parameters for different spatiotemporal processes, a more reasonable limit and optimization procedure is hence needed.

3.2 Calibration of STWR

Calibration of the STWR models can be conducted by using weighted least squares. The estimator for the coefficients at location (u_i, v_i) is shown below.

$$\beta_t(u_i, v_i) = [(X_{O_{\Delta t}}^T W_{\Delta t}(u_i, v_i) X_{O_{\Delta t}})^{-1} X_{O_{\Delta t}} W_{\Delta t}(u_i, v_i)] y_{O_{\Delta t}} \quad (11)$$

In Equation 11, $X_{O_{\Delta t}}$ and $y_{O_{\Delta t}}$ are observed independent and dependent variables of $O_{\Delta t}$ respectively. $X_{O_{\Delta t}}^T$ is the transpose of $X_{O_{\Delta t}}$. $W_{\Delta t}(u_i, v_i)$ denotes the spatiotemporal weight matrix for observed points at different locations to the regression point (u_i, v_i) at different time stages during Δt . For a better illustration, we show the weight matrix $W_{\Delta t}$ during the time interval Δt in Fig. 2. The matrix $W_{\Delta t}$ here is a bit different form the $W(u_i, v_i)$ in Equation 2. The records in the i^{th} row of $W_{\Delta t}$ are the diagonal elements in $W(u_i, v_i)$, and only no zero values are used to calibrate the coefficients β_k for each regression point. Thus, each row r of this hat matrix is shown below.

$$r_{it} = X_{it} (X_{\Delta t}^T W_{i\Delta t} X_{\Delta t})^{-1} X_{\Delta t} W_{i\Delta t} \quad (12)$$

In Equation 12, X_{it} is the i^{th} row of the matrix of independent variables at t . $X_{\Delta t}$ is the matrix of independent variables during a time interval Δt , and $X_{\Delta t}^T$ is its transpose. Although the $X_{\Delta t}$ in Equation 12 is equal to the $X_{O_{\Delta t}}$ in Equation 11 in the fitting and calibration of STWR, we distinguish $X_{O_{\Delta t}}$ from $X_{\Delta t}$ here. Because $X_{O_{\Delta t}}$ is a specific matrix of independent variables of an observed point set $O_{\Delta t}$ during Δt , while $X_{\Delta t}$ is a general matrix of independent variables of points during Δt . $X_{O_{\Delta t}}$ is only used for fitting and calibration of STWR, while $X_{\Delta t}$ can also be used for prediction in STWR. In other words, we can understand $X_{O_{\Delta t}}$ as a subclass of $X_{\Delta t}$. $W_{i\Delta t}$ is the i^{th} row of the weighted matrix $W_{\Delta t}$.


3.3 Reasonable searching range and procedure of optimization

In order to obtain the optimized α and θ for STWR (Equations 8 and 9), the search range should be limited. Here we use the distance from each regression point $p_i^{(t)}$ to its M^{th} nearest neighbor as the initial spatial bandwidth b_{St} at t . The range of b_{St} is within a finite set of discrete values, because the maximum number of nearest neighbor is limited to N_{t-i} , $i \in$

49 $\{1, 2, \dots, q\}$ for the regression point $p_i^{(t)}$ (N_{t-i} is the total number of observed points at $t-i$). We denote that value set for
50 b_{St} as $BS_{Nt} = \{D_{k+1}, D_{k+2}, \dots, D_{Nt}\}$, in which the element $D_U, U \in \{k+1, k+2, \dots, Nt\}$ denotes the distance from $p_i^{(t)}$ to
51 the U^{th} nearest neighbor, and k equals to the number of independent variables. Moreover, the searching range of the
52 temporal bandwidth b_T is also limited to a finite discrete set $BT_\lambda = \{\Delta t_1, \Delta t_2, \dots, \Delta t_\lambda\}$, in which the element Δt_λ is the time
53 interval from t to $t-\lambda$.

54 The optimization procedure is to traverse the set BT_λ , and for each step we further traverse the set BS_{Nt} to get the
55 optimized α and θ through trials. Some trials of θ may lead to no solution to Equation 11, because there might be less than
56 $(k+1)^{th}$ neighbors within the radius of $b_{St} - \theta \Delta t_\lambda$ from the regression point. Therefore, if it occurs at time stage $t-\lambda$,
57 the spatial bandwidth $b_{St} - \theta \Delta t_\lambda$ needs to be extended to the distance from its $(k+1)^{th}$ nearest neighbor to the regression
58 point, to guarantee the matrix in Equation 11 to be nonsingular.

59



60

61 **Fig. 2.** Weight matrix $W_{\Delta t}$. The symbol $p_k^{(t-i)}, i \in \{0, 1, \dots, q\}, k \in \{1, 2, \dots, N_{t-i}\}$ denotes the k^{th} observed point at $t-i$.

62 The symbol $w_{\Delta t, p_m^{(t)} p_n^{(t)}}^{t-i}, i \in \{0, 1, \dots, q\}, m \in \{1, 2, \dots, N_t\}, n \in \{1, 2, \dots, N_{t-i}\}$ denotes the weight of the n^{th} point $p_n^{(t-i)}$ at

63 $t-i$ to the m^{th} point $p_m^{(t)}$ at t . The symbol $O_{N_{t-i}}, i \in \{0, 1, \dots, q\}$ denotes a set of points observed at $t-i$. Δt denotes all

.64 the time intervals of the weight matrix. In the central and right parts of the figure, the records with background shading
.65 indicate weight values affected by temporal effects.

.66

.67 **3.4 Steps of using STWR for prediction**

.68 In this paper, STWR is used to predicate the current values of regression points with known coordinates. The prediction
.69 formulas of STWR are more complicated than GWR because the spatial distance is calculated directly from the regression
.70 point to each observed data point, while the time distance between the regression point and the data points observed in the
.71 past cannot be calculated directly. Therefore, we specify a few steps for the prediction in STWR. First, we need to have the
.72 optimized initial spatial bandwidth b_{st} , the optimized α and θ , the optimized number of time stages model used and the
.73 fitted weight matrix. Second, all data points within the limited distance of spatial bandwidth at the latest time stage should be
.74 found for the regression point. Third, all the temporal weights of these data points need to be retrieved from the established
.75 weight matrix (Fig. 2). Fourth, we use these retrieved weights to calculate (e.g., use mean value or inverse distance
.76 weighting value) the temporal weight on the regression point. Fifth, by combining with the calculated spatial weight and the
.77 optimized α and θ , we can calculate the spatiotemporal weight on the regression point. Then the value of the regression
.78 point can be calculated.

.79

.80 **4. Experiments with Simulated Data**

.81 **4.1 Simulation design**

.82 To verify the performance of STWR and compare with the results of GWR and GTWR, several groups of simulated data
.83 were used in this study to represent different types of heterogeneity in space and time. All the data and code used in the
.84 experiments are shared on GitHub. Web links are provided at the end of this manuscript.


.85 For GTWR, we only compared with the results generated by algorithms in Huang et al. (2010) and Wu et al. (2014),
.86 because we did not find the software package of Fotheringham et al. (2015). The data generating process (DGP) and the

.87 spatial heterogeneity are introduced here. The basic DGP is a linear model shown in Equation 1 and the study area is a
.88 regular 25×25 lattice. We defined three initial surfaces to represent the spatial heterogeneity of parameters (Fig. 3), which
.89 were generated by Equations 13, 14 and 15, respectively (Fotheringham et al., 2017). Through Equation 1, the two
.90 independent variables x_1 and x_2 were initially generated randomly from the normal distribution $x_1^{initial} \sim N(100, 8)$ and
.91 $x_2^{initial} \sim N(50, 6)$, respectively. They can be set as any other values, and the mean values of both distributions may change
.92 over time. The error term was generated from a normal distribution $\varepsilon \sim N(0, 0.5)$.

$$\beta_{0(zh)}^t = 3 \quad (13)$$

$$\beta_{1(uh)}^t = 1 + \frac{1}{12}(u, v) \quad (14)$$

$$\beta_{2(hh)}^t = 1 + \frac{1}{324}[36 - (6 - u/2)^2][36 - (6 - v/2)^2] \quad (15)$$



.97
.98 **Fig. 3.** Three simulated initial surfaces for representing spatial heterogeneity of parameters.


.99
.00 Several trends were designed to simulate the value change. For a better simulation, we assumed that value variation can
.01 also be spatial heterogeneity. To distinguish from the heterogeneity of the coefficient surface, three other heterogeneity trend
.02 functions were defined by Equations 16, 17 and 18.

$$T_1 V^{t+\Delta t} = V^t + \varphi * \sin(v/4)\Delta t^{npower} \quad (16)$$

04
$$T_2V^{t+\Delta t} = V^t + \varphi * \sin[1/10\pi u]\Delta t^{npower} \quad (17)$$

05
$$T_3V^{t+\Delta t} = V^t + \varphi * \sin[1/6\pi(u + v)]\Delta t^{npower} \quad (18)$$

06 In the above equations, V^t denotes the value at time stage t , φ is used for adjusting the magnitude of change, Δt^{npower}
 07 denotes value change with the n^{th} power of time interval, and $T_iV^{t+\Delta t}$, $i \in \{1,2,3\}$ denotes the V value at time stage $t +$
 08 Δt , which is the result of the i^{th} trend function from the V^t . Fig. 4 shows these trends when φ , V^t , and Δt^{npower} are set to
 09 one.



10
11 **Fig. 4.** Three heterogeneity trend surfaces.

12
13 Our goal of this experiment was to test model performance by using sample data from the simulation process at
 14 different time. Three case studies were designed for different situations. Besides the spatial heterogeneity trends, in our
 15 simulation design we assumed that the mean values of two independent variables x_1 and x_2 also changed over time, which
 16 were generated by Equations 19 and 20, respectively.

17
$$T_1x_m^{t+\Delta t} = x_m^t \pm \eta_1 * \Delta t \quad (19)$$

18
$$T_2x_m^{t+\Delta t} = x_m^t \pm \eta_2 * \Delta t \quad (20)$$

19 In the above two equations, x_m^t denotes the mean of an independent variable x at time stage t , $T_ix_m^{t+\Delta t}$, $i \in \{1,2\}$ denotes
 20 the mean of x at time stage $t + \Delta t$, and η_1 and η_2 are two parameters for adjusting the rate of change. At each time stage
 21 during the simulations, the independent variables x_1 and x_2 are generated by a normal distribution with new means of
 22 $T_1x_m^{t+\Delta t}$ and $T_2x_m^{t+\Delta t}$, respectively.

23 4.2 Results with simulated data

24 We compared the results of OLS, GWR, GTWR, and STWR. A total of 333 random sample points for five time stages
25 (t_0, t_1, t_2, t_3, t_4 from old to new) were collected from the 25×25 lattice generated in the above-mentioned DGP. To simplify
26 the calculation process, we set θ of Equation 7 to zero. Due to the limitation of paper length, in the comparison below the
27 STWR results only include those generated by the spatiotemporal kernel in Equation 8. The objective is to compare the
28 predicted results with the true value at the latest time stage.

29 4.2.1 Case study 1


30 The time interval of observations in case study 1 was one unit, such as one second or one day. The value change of x_1 and
31 x_2 were generated by $\eta_1 = 0.5$ and $\eta_2 = 0.1$, and were affected by T_1V with $\varphi = 0.5$ and $npower = 1$. This means that
32 x_1 and x_2 only changed slightly over time. Table 1 presents the results of the global OLS, GWR, GTWR and STWR at the
33 latest time stage, i.e., stage 5. It shows that the sum of squared errors (SSE) of prediction in STWR is much lower than the
34 other models in at least one magnitude. In addition, the AICc scores (Equation 10) also shows that STWR outperforms
35 GTWR and GWR. As shown in Table 1, the R2 (average R-squared of all regression points) value increases from 13.8% in
36 OLS to 94.2% in GWR, 94.9% in GTWR, and 99.3% in STWR. The estimated standard error, Sigma, reduces to 4.292 in
37 STWR from 23.331 in GTWR. Also, Fig. 5 shows that both the prediction surface (Y_{pred}) and the prediction error surface
38 ($Pred_Error$) of STWR are more accurate than those in GWR. Due to the limitation of the software package in Huang et al.
39 (2010) and Wu et al. (2014), we did not generate images for GTWR in Fig. 5, but the result can be seen from the Sigma
40 value in Table 1.

41
42 **Table 1.** Results of case study 1 at time stage t_4 .

Time stage t_4	SSE	AICc	R2	Sigma
OLS	676366.268	805.455	0.138	
GWR	45674.420	705.529	0.942	33.277

GTWR	40056.823	616.641	0.949	23.331
STWR	5761.109	528.860	0.993	4.293

43



44

45 **Fig. 5.** Comparing prediction results of STWR and GWR in case study 1. Images a1, b1, and c1 are the simulation surfaces
 46 of true Y, the predicted surface of Y by STWR, and the predicted surface of Y by GWR, respectively. Images a2, b2, and c2
 47 are the surface of simulation error, the surface of prediction error of STWR, and the surface of prediction error of GWR,
 48 respectively.

49

50 4.2.2 Case study 2

51 The time interval of observations in case study 2 was 10 units. The value change of x_1 was generated by $\eta_1 = 0.5$ and
 52 affected by T_3V with $\varphi = 0.5$ and $npower=2$. x_2 was generated by $\eta_2=2$ and affected by T_2V with $\varphi = 1$ and $npower$

.53 = 1, which denotes that x_1 and x_2 changed fast over time. Table 2 shows the results of the global OLS, GWR, GTWR and
.54 STWR at the time stage 5. The SSE value in STWR is much lower than other models, and STWR has the highest R2 value
.55 0.995. The Sigma value of STWR is 13.299, which is the lowest and less than one-fifth of the Sigma in GWR and less than
.56 one-sixth of the Sigma in GTWR. Besides, the AICc scores show that STWR significantly outperforms GTWR and GWR.

.57 STWR utilized data from the latest three time stages to calibrate the model. The initial spatial bandwidth b_{st} of STWR
.58 was three nearest neighbors, which was smaller than the one in GWR with 15 nearest neighbors. The optimized α of STWR
.59 was 0.08, which shows that the effect of used observed points to their local regression points was mainly determined by their
.60 spatial distance. In this case, the GWR outperforms GTWR, which may due to the higher ratio of value change. Compared
.61 with the y_{true} surface, the predict surface of STWR is much better than GWR (Fig. 6). For the same reason as mentioned in
.62 case study 1, we did not generate images for GTWR in Fig. 6.


.63

.64

Table 2. Results of case study 2 at time stage t_4 .

Time stage t_4	SSE	AICc	R2	Sigma
OLS	5085961.816	938.610	0.494	
GWR	300088.969	840.178	0.970	87.201
GTWR	627011.021	895.662	0.938	127.821
STWR	52688.545	709.573	0.995	13.299

.65




66
 67 **Fig. 6.** Comparing prediction results of STWR and GWR in case study 2. Images a1, b1, and c1 are the simulation surfaces
 68 of true Y , predicted surface of Y by STWR, and predicted surface of Y by GWR, respectively. Images a2, b2, and c2 are the
 69 surface of simulation error, the surface of prediction error of STWR, and the surface of prediction error of GWR,
 70 respectively.

71

72 4.2.3 Case study 3

73 The time interval of observations in case study 3 was 200 units. In both case studies 1 and 2, the coefficients in Equation 1
 74 were unchanged. In contrast, in case study 3, three surfaces of coefficients changed over time, which were generated by the
 75 trends T_1V , T_2V , and T_3V , respectively. The variations of coefficients were assumed to be slow. The φ and $npower$ in
 76 each trend were set to be 0.2 and 1, respectively. Both η_1 and η_2 were set to be 0.5. The dynamic process of the three
 77 surfaces of coefficients and the y_true surface at each time stage are shown in Fig. 7. The process in case study 3 is more
 78 complicated than a general process, but it may be closer to reality.



80

81 **Fig. 7.** Dynamic process of three surfaces of coefficients and the y_true surface at five different time stages.

82

83 Results of these comparisons in case study 3 show that STWR outperforms both GWR and GTWR in accuracy of

84 model and effectiveness of simulation process (Fig. 8a). Along with the change of the coefficients and the increase of


85 x_1 and x_2 , the R2 values of both GWR and GTWR are consistent in the five time stages, showing an overall downward

.86 trend. But the R2 of STWR is stable and is at a high level among the five time stages. At the beginning stage t_0 , the R2
.87 values of the three models are similar because there are no previous observations that can be used by STWR and GTWR.
.88 The small difference among these models at t_0 may be caused by their different searching range of spatial bandwidth.
.89 Starting from time stage t_1 , STWR and GTWR can borrow points from previous observations. At time stage t_1 , STWR
.90 outperforms both GWR and GTWR, and the advantage of STWR becomes more obvious in the later stages.

.91 It may seem strange that GWR can outperform GTWR (Fig. 8), but that is reasonable for the process in case study 3.
.92 The change of this process is faster; and the time interval of observations is bigger than the previous case studies. STWR is
.93 not only able to deal with time intervals, but also to make full use of the value variation of observed points for calibration. In
.94 contrast, GTWR only uses the time interval information and all the observed points to calibrate, which may cause problems
.95 when the observed values are significantly different in spatial distribution or the time intervals are long. GTWR makes use of
.96 points from previous time stages without considering their variation, but if the actual values are quite different from previous
.97 observations at the current time stage, all the point values for the calibration of GTWR will become smooth. Thus, GWR
.98 outperforms GTWR in this situation because GWR only uses the current data points for model calibration.

.99 STWR is better for estimation than GWR and GTWR because its Sigma value is much smaller. As shown in Fig. 8b,
.00 the Sigma of STWR was half of GWR at time stage t_1 , and even less than a third of GWR at time stage t_4 . The results show
.01 that the advantage of STWR is obvious comparing with GWR and GTWR.

.02




.03
.04 **Fig. 8.** Comparing and evaluating the performance of GWR, GTWR and STWR at five time stages. (a) Comparing the R²
.05 value of different models; (b) Comparing the Sigma value of different models.
.06

.07 At t_4 , STWR used data from all the past time stages to calibrate the model, and its optimized (initial) spatial bandwidth
.08 b_{st} was derived from four nearest neighbors, which was smaller than the one in GWR with 25 nearest neighbors. The
.09 optimized α of STWR was 0, which means that STWR only borrowed points from past time stages without considering
.10 their temporal weights to each regression point at t_4 . The predict surfaces at time stage t_4 is shown in Fig. 9. The Y_{pred}
.11 surface of STWR is much better than GWR, especially in the middle and bottom left parts of the surface. The $Pred_Error$ of
.12 STWR is also much lower than GWR at almost every location. In this case, the α of STWR at each time stage was 0, 0.96,
.13 0, 0.07, and 0, respectively. These values indicate that the temporal effects are different at each stage. They also show that
.14 the value of α can be adaptive to scale the temporal and spatial effects (see Equation 3).
.15


.16 As Fig. 10 shows, the optimized bandwidths are quite different among these models, and the bandwidths of GWR and
.17 GTWR are larger than the initial bandwidth of STWR at each time stage. The optimized bandwidth for each time stage refers
to an optimized number of the nearest neighbors (see Section 3.3). As GTWR considers all the nearest neighbors from

.18 different time stages, the optimized numbers of the nearest neighbors (bandwidth) grow fast, and exceed the GWR model at
.19 time stage t_2 . However, the actual distance from the observed points to the regression points is not necessarily farther. The
.20 initial optimized numbers of the nearest neighbor of STWR are smaller than those in GWR and GTWR, which means that
.21 the initial spatial bandwidth is narrower than the bandwidth of GWR and GTWR. Nevertheless, due to the strategy of
.22 borrowing points from nearby neighbors of past observations, the total points for model calibration in STWR may still be
.23 more than GWR and GTWR. Therefore, the initial optimized numbers of the nearest neighbors in STWR are kept at a lower
.24 level, which means it is more localized than GWR in this sense.
.25



.26
.27 **Fig. 9.** Comparing prediction results of STWR and GWR in case study 3. Images a1, b1, and c1 are the simulation surfaces
.28 of true Y, the predicted surface of Y by STWR, and the predicted surface of Y by GWR, respectively. Images a2, b2, c2 are

.29 the surface of simulation error, the surface of prediction error of STWR, and the surface of prediction error of GWR,
.30 respectively.
.31



.32
.33 **Fig. 10.** Optimized bandwidths (or initial bandwidths) of GWR, GTWR and STWR for the five time stages in case study 3.
.34

.35 **5. Experiments with Real-world Data**

.36 To further test the performance of STWR, we used data of precipitation $\delta^2\text{H}$ isotopes in Northeastern United States in
.37 another case study. We chose $\delta^2\text{H}$ data in three days from October 29 to 31, 2012, which have enough spatiotemporal data
.38 for the test. Here in the comparison the STWR results only include those generated by the spatiotemporal kernel in Equation
.39 8. Data and code used here are shared on Zenodo (See DOI and web links in the 'Code and data availability' section at the
.40 end of the main text of this article).

41 In the experiments, we collected a total of 782 measurements from 116 sites located in Northeastern United States
 42 during the three-day period, and prepared the data on a daily average. The daily precipitation, mean temperature, and
 43 elevation were used as explanatory variables. The model derived from Equation 1 is represented below.

$$y_i = \beta_0 + \beta_1 ppt + \beta_2 tmean + \beta_3 height + \varepsilon_i \quad (21)$$

44
 45 In Equation 21, *ppt* denotes the daily total precipitation (rain + melted snow), *tmean* denotes daily mean temperature, and
 46 *height* is the elevation value. After data preprocessing, there were 272 points for model calibration and 73 points values on
 47 October 31, 2012. For the first day, both GTWR and STWR took no information from the past. Therefore, we only show the
 48 results of SSE, R2 and the optimized initial neighbor (bandwidth) in the model comparisons for the second and third day (D2
 49 and D3) in Tables 3. The SSE of STWR is the lowest at both days. GWR shows a slightly higher SSE than GTWR at D2 and
 50 D3. The R2 of STWR is the highest at both days among these models. GWR has lower R2 than GTWR at D2, and almost the
 51 same R2 as GTWR at D3.

52 Similar to the experiments on three simulation datasets, the result here shows that STWR outperforms GTWR and
 53 GWR. In the experiment, the number of optimized initial neighbors of STWR was smaller than that of GWR and GTWR.
 54 The optimized α of STWR was 0 at both D2 and D3. The optimized temporal bandwidths of STWR (number of time stages
 55 model used) in both D2 and D3 were 2, which means that the STWR in this case only borrowed data points from the latest 2
 56 time stages for D2 and D3. In the result (Table 3), an interesting part to see is that the numbers of optimized initial neighbors
 57 of STWR are smaller than the spatial bandwidths of GWR for D2 and D3. The reason is that STWR borrowed points from
 58 past time stages in the calculation, which led to narrower bandwidths to some extent.

59

60 **Table 3.** Results of model performance with real-world data.


Model	SSE-D2	SSE-D3	R2-D2	R2-D3	Neighbor	Neighbor
					-D2	-D3
OLS	58711.528	52669.399	0.595	0.502		

GWR	33576.400	33043.921	0.769	0.688	52	43
GTWR	32659.808	31967.850	0.775	0.698	37	31
STWR	24022.226	25118.096	0.834	0.763	16	16

.61

.62 We adopted Leave-one-out cross-validation (LOOCV) at D3 for the comparison between STWR and GWR. The
.63 squared errors (SE) of prediction are shown in Fig. 11. The prediction results of STWR are better than GWR for most points.
.64 The mean SE of STWR is smaller than GWR. Moreover, the SE of STWR shows a narrower regional trend, which indicates
.65 that STWR is more robust than GWR. In addition, the total SSE of GWR and STWR are 50216.510 and 39724.995,
.66 respectively. Therefore, the result further validates that the quality of predication in STWR is better than GWR.

.67




.68

.69 **Fig. 11.** LOOCV results of STWR and GWR. (a) Squared error of prediction at each point (leave out); (b) Box plot of the

.70 LOOCV results of GWR and STWR.

.71

.72 In Fig. 12, the predicted $\delta^2\text{H}$ surface at D3 is broadly similar between the GWR and STWR calibrations. The
.73 percentages of explanation of variance in GWR and STWR are similar, which are 68.8% and 76.3%, respectively. However,
.74 like the experiment results with simulated data (Fig. 10), STWR has narrower initial bandwidth, which generates more
.75 localization in the predicted $\delta^2\text{H}$ surface than GWR. For instance, the lower (light yellow and blue parts) or higher (orange
.76 parts) predicted values of $\delta^2\text{H}$ are more concentrated in the $\delta^2\text{H}$ surface of STWR than that of GWR (Fig. 12).
.77



.78
.79 **Fig. 12.** Predicted $\delta^2\text{H}$ surfaces of STWR and GWR at D3.
.80

6. Discussion and Conclusions

Spatiotemporal data analysis is important in many scientific studies. Due to the complexity of spatiotemporal models, spatiotemporal effect may not be fully taken into account when the temporal and spatial information is manipulated simultaneously. In particular, the models for the effect of spatial dynamics should not be simply adapted for modeling the effect of temporal dynamics. Although the GTWR model can borrow points from the near recent, without careful consideration of temporal effect, the performance of GTWR may even be worse than GWR. Increasingly, many scientific issues are not just about spatial non-stationary but involve many spatiotemporal processes. It is necessary to review the limitation of the current spatiotemporal models and make new extensions. The aim of the STWR model developed in this study is to advance the work and discussion in that direction.

Based on a concept similar to GWR, a recently proposed model, named geographically neural network weighted regression (GNNWR) (Du et al., 2020), utilizes both OLS and neural networks to evaluate spatial non-stationarity. It is characterized by a designed spatially weighted neural network (SWNN) which can represent the spatial non-stationary weight matrix in spatial processes. Additionally, a geographically and temporally neural network weighted regression (GTNNWR) model (Wu et al., 2020), which is a temporal extension of GNNWR, was also proposed by the same group for further modelling spatiotemporal non-stationary relationships. GTNNWR can generate a space-time distance by utilizing the so-called spatiotemporal proximity neural network (STPNN), which may address complex non-linear interactions between time and space. Although both STWR and GTNNWR have the potential to handle complex spatiotemporal non-stationarity in various natural and socioeconomic processes, their principles and interpretability are different:

(1) The basic formulation of GNNWR is defined as Equation 22 (Du et al., 2020), which is different from Equation 1 (Fotheringham et al., 2003). The $w_0(u_i, v_i)$ and $w_k(u_i, v_i)$ denote the geographical weight of the constant coefficient β_0 and coefficient β_k , respectively. It assumed that the multiplication of $w_p(u_i, v_i)$ and β_p is equal to $\beta_p(u_i, v_i)$ ($0 \leq p \leq k$). The combined $\beta_p(u_i, v_i)$ is thought as the same as the coefficients of GWR. But in STWR and GWR, the weights and the estimated coefficients are separated. The weights mainly reflect the degree of the influences from the observed points to

Deleted: With increasing combined applications of deep learning and neural network in geospatial non-stationary processes. We first discuss the main differences between STWR and the recently proposed geographic neural network weighted regression (GNNWR) (Du et al., 2020) and geographic and temporal neural network regression (GTNNWR) (Wu et al., 2020). GNNWR is a new attempt to combine the OLS and GWR with Artificial neural networks (ANNs). GTNNWR is based on the GNNWR with combing a new ANNs based method to calculate the spatiotemporal distance. Four main differences between the GTNNWR/GNNWR and STWR are listed below:¶

Formatted: Font: Not Bold

Formatted: Font: Not Bold

16 the regression point, while the coefficient values reflect the relationships between the independent variable and dependent
17 variable.

$$y_i = w_0(u_i, v_i)\beta_0 + \sum_{k=1}^p w_k(u_i, v_i)\beta_k x_{ik} + \varepsilon_i, i = 1, 2, \dots, n \quad (22)$$

18
19
20 (2) GTNNWR and GNNWR use the proposed ANNs based method (Equation 23) (Du et al., 2020) to calculate the
21 weighted matrix, which is quite different from the kernel functions used in GWR and STWR models. Although GTNNWR
22 and GNNWR use the idea of pointwise regression, they do not consider how to "borrow points" from nearby neighbors and
23 do not have the concept of bandwidth. Without spatial bandwidth, all observation points in the study area may have impacts
24 on the regression point, which might violate the Tobler's first law of geography (Tobler, 1970). It may be difficult to
25 understand the relationships between the influence weight and the spatial distances, especially when the study area and the
26 data amounts are large. STWR has spatial bandwidths and follows the Tobler's first law of geography, which can help
27 analyze the affected range of local regression points.

$$W_i = W(u_i, v_i) = SWNN([d_{i1}^s, d_{i2}^s, \dots, d_{in}^s]^T) \quad (23)$$

28
29 (3) The data points will be divided into training set (including validation set) and test set for the GTNNWR and
30 GNNWR, which might require more data points. Thus, it may not be appropriate for analyzing fewer amounts of data points
31 (data acquisitions of many geoscience processes are difficult and costly). STWR and GWR do not need to divide data points
32 into the training set (including validation set) and test set, which requires less data points than GNNWR and GTNNWR.

33 (4) Although GTNNWR utilizing a method named spatiotemporal proximity neural network (STPNN) (Wu et al., 2020)
34 to calculate the spatiotemporal distance, the obtained integrated spatiotemporal distance is lack of explanation, and it is also
35 impossible to tell apart which parts of the calculated weight is affected by time or space. Besides, there is no concept of
36 temporal bandwidth in GTNNWR. Therefore, it fails to inform us the earliest time (stage) since which the observed points
37 start to exert impact on the determination of the regression point. But STWR has temporal bandwidth, and it can distinguish
38 the strength of temporal weight and spatial weight. Therefore, we can analyze the characteristics of the local interaction of
39 time and space according to the temporal bandwidth, spatial bandwidth, and the adjustment parameter α , etc.

Deleted: Thus

Deleted: cannot tell

Deleted: how old

Deleted: historical observation

Deleted: that will have impacts

45 The temporal kernel and the spatiotemporal kernel functions are two important contributions of STWR. The temporal
46 kernel in STWR applies an improved sigmoid form (see Equation 4), which is different from the methods for temporal effect
47 analysis in previous GTWR models. The temporal weight generated by the STWR temporal kernel is limited as a value
48 between 0 and 1. The spatial weight in STWR is also limited as a value between 0 and 1. The STWR spatiotemporal kernel
49 function has a weight adjustment parameter α to scale the temporal and spatial weights (Equation 3). In practice, α can be
50 obtained through optimization. This form of weighted average between temporal and spatial effects in the STWR
51 spatiotemporal kernel is a big improvement comparing with the multiplication form in previous GTWR models. The
52 advantage of the STWR spatiotemporal kernel has been proven in four case studies with both simulated and real-world
53 datasets.

54 Though the performance of STWR is outstanding, the models can still be further extended. A big topic is about the time
55 distance. In the current STWR, the time distance represents the rate of value variation between an observed point and a
56 regression point through a time interval. Nevertheless, we can also use time distance to represent the rate of value variation
57 at each observed point object through time. Note that, from an object-oriented perspective, here we differentiate the point
58 objects from locations, although the point objects have geospatial coordinates as part of their attributes. Following that new
59 definition of time distance, the $y_{i(t)} - y_{j(t-q)}$ in the STWR temporal kernel (Equation 4) can be replaced by $\Delta y_{j(t-q)}$
60 (value variation of an observed point object during Δt). A scenario of interest is that, the observed point objects in the past
61 time stages (such as those shown in Fig. 1) may move to new locations, have no value for a few time stages, or even
62 disappear, so the $\Delta y_{j(t-q)}$ may not exist. We can use object-based methods to address issues caused by that scenario. For
63 example, each point object can be assigned with a unique ID, and then the observed value of the point object at each time
64 stage can be retrieved by using its ID. With this new definition of time distance, the temporal weight on a regression point
65 object is determined by the rate of value variation of its nearby point objects. Several different scenarios for a regression
66 point object at current time stage t are discussed here.

67 (1) The location of an observed point object j is fixed through time (e.g., a fixed sensor). If the value of j is observed
68 at both time stages t and $t - q$, then $\Delta y_{j(t-q)}$ can be calculated directly. If the value of j is observed at t but not
69 observed at $t - q$, we can use interpolation to generate a value for j at $t - q$. If the value of j is not observed at t , but
70 the variation in the past is observed, we can use prediction methods to generate a value for j at t .

71 (2) The location of j is not fixed through time (i.e., j moves). The moving point objects can still have temporal
72 effects to the regression point, then the $\Delta y_{j(t-q)}$ can be calculated. The spatial effect, however, depends on whether j
73 moves out of the spatial bandwidth from the regression point or not.

74 (3) j disappears or appears at a certain time stage. If j does not appear until the current time stage t , the $\Delta y_{j(t-q)}$ can
75 be set to be 0. If j appears in a past time stage (e.g., $t - q$) but it disappears before or at t , we can ignore the impact of j
76 for the regression point object.

77 There are other possibilities for the further improvement of STWR. The first is about the optimization of θ in the
78 spatiotemporal kernel (Equations 8 and 9). The slope θ indicates that the variation of the spatial bandwidth is in a linear
79 form, but it may not be a perfect solution. In many situations, the change of the spatial bandwidth over time may not be
80 linear. The second is about making predications for future time stages. In this paper, we only predict values for points at the
81 current time stage t . Extensions can be made in STWR to predict values for points in future time stages beyond t . The third
82 future work is about exploring multiple spatial and temporal bandwidths of models. Different variables may have different
83 spatial and temporal bandwidths due to their unique characteristics. Correspondingly, we may need more bandwidths to
84 capture the different non-stationarities of those independent variables, to better represent the spatiotemporal heterogeneity.

85 In short, the core contribution of STWR is the clarification of the 'time distance' concept and the new temporal kernel
86 and spatiotemporal kernel functions based on this concept. Our experiments show that STWR outperforms GWR and GTWR
87 in analyzing and interpreting local spatiotemporal non-stationarity. We hope STWR can bring fresh ideas and new
88 capabilities for spatiotemporal data analysis in many disciplines.

89

90 **Code and data availability**

91 The Python source code of STWR v1.0, the data used in the experiments and all the case studies (written in Jupyter
92 Notebook) were archived on Zenodo and made freely accessible via <http://doi.org/10.5281/zenodo.3637689>. Data source of
93 water isotopes $\delta^{2}\text{H}$ is on the website: http://wateriso.utah.edu/waterisotopes/pages/spatial_db/SPATIAL_DB.html. The data
94 of daily precipitation and mean temperature were collected from the PRISM Climate Group
95 (<http://www.prism.oregonstate.edu>), and the elevation data were collected from the GMTED2010
96 (https://topotools.cr.usgs.gov/gmted_viewer/viewer.htm) at U.S. Geological Survey (USGS).

97

98 **Author Contribution.**

99 X.Q., X.M. and C.M. developed the algorithm, X.Q. implemented and coded the algorithm. X.Q. prepared the manuscript
00 with contributions from all co-authors.

01

02 **Competing interests.**

03 The authors declare that they have no conflict of interest.

04

05 **Acknowledgement.**

06 The research presented in this paper was partially supported by the National Science Foundation under Grants No. 1835717
07 and No. 2019609, the China Scholarship Council under Grant No. 201807870006, and the Fujian Provincial Department of
08 Education under Grant No. KLA18025A. The authors thank Prof. Stewart Fotheringham and other colleagues at the Spatial
09 Analysis Research Center (SPARC) of Arizona State University for their insightful comments and suggestions during a
10 seminar about the STWR model.

11

12

13 **References**

- 14 Akaike, H.: Information theory and an extension of the maximum likelihood principle. In: Selected papers of Hirotugu
15 Akaike, Springer, 1998.
- 16 Akaike, H.: Maximum likelihood identification of Gaussian autoregressive moving average models, *Biometrika*, 60, 255-
17 265, 1973.
- 18 Atkinson, P. M., German, S. E., Sear, D. A., and Clark, M. J.: Exploring the relations between riverbank erosion and
19 geomorphological controls using geographically weighted logistic regression, *Geographical Analysis*, 35, 58-82, 2003.
- 20 Bowman, A. W.: An alternative method of cross-validation for the smoothing of density estimates, *Biometrika*, 71, 353-360,
21 1984.
- 22 Brown, S., Versace, V. L., Laurenson, L., Ierodiaconou, D., Fawcett, J., and Salzman, S.: Assessment of spatiotemporal
23 varying relationships between rainfall, land cover and surface water area using geographically weighted regression,
24 *Environmental Modeling & Assessment*, 17, 241-254, 2012.
- 25 Brunson, C., Fotheringham, A. S., and Charlton, M. E.: Geographically weighted regression: a method for exploring spatial
26 nonstationarity, *Geographical analysis*, 28, 281-298, 1996.
- 27 Brunson, C., Fotheringham, S., and Charlton, M.: Geographically weighted regression, *Journal of the Royal Statistical*
28 *Society: Series D (The Statistician)*, 47, 431-443, 1998.
- 29 Cahill, M. and Mulligan, G.: Using geographically weighted regression to explore local crime patterns, *Social Science*
30 *Computer Review*, 25, 174-193, 2007.
- 31 Cardozo, O. D., García-Palomares, J. C., and Gutiérrez, J.: Application of geographically weighted regression to the direct
32 forecasting of transit ridership at station-level, *Applied Geography*, 34, 548-558, 2012.
- 33 Chen, J., Shaw, S.-L., Yu, H., Lu, F., Chai, Y., and Jia, Q.: Exploratory data analysis of activity diary data: a space-time GIS
34 approach, *Journal of Transport Geography*, 19, 394-404, 2011.

-
- 35 Cleveland, W. S.: Robust locally weighted regression and smoothing scatterplots, *Journal of the American statistical*
36 *association*, 74, 829-836, 1979.
- 37 Crespo, R., Fotheringham, S., and Charlton, M.: Application of geographically weighted regression to a 19-year set of house
38 price data in London to calibrate local hedonic price models. In *Proceedings of the 9th International Conference on*
39 *Geocomputation*. National University of Ireland Maynooth, 2007.
- 40 Cressie, N. and Wikle, C. K.: *Statistics for spatio-temporal data*, John Wiley & Sons, 2015.
- 41 Cressie, N. A.: *Statistics for Spatial Data*. New York: John Wiley & Sons, 1993.
- 42 Du, Z., Wang, Z., Wu, S., Zhang, F. and Liu, R.: Geographically neural network weighted regression for the accurate
43 estimation of spatial non-stationarity. *International Journal of Geographical Information Science*, 34:7, 1353-1377,
44 2020.
- 45 Fotheringham, A. S., Brunson, C., and Charlton, M.: *Geographically weighted regression: the analysis of spatially varying*
46 *relationships*, John Wiley & Sons, 2003.
- 47 Fotheringham, A. S., Crespo, R., and Yao, J.: Geographical and temporal weighted regression (GTWR), *Geographical*
48 *Analysis*, 47, 431-452, 2015.
- 49 Fotheringham, A. S., Yang, W., and Kang, W.: Multiscale geographically weighted regression (mgwr), *Annals of the*
50 *American Association of Geographers*, 107, 1247-1265, 2017.
- 51 Fraser, L. K., Clarke, G. P., Cade, J. E., and Edwards, K. L.: Fast food and obesity: a spatial analysis in a large United
52 Kingdom population of children aged 13–15, *American journal of preventive medicine*, 42, e77-e85, 2012.
- 53 Gelfand, A. E., Ecker, M. D., Knight, J. R., and Sirmans, C.: The dynamics of location in home price, *The journal of real*
54 *estate finance and economics*, 29, 149-166, 2004.
- 55 Goodchild, M. F.: Prospects for a space–time GIS: Space–time integration in geography and GIScience, *Annals of the*
56 *Association of American Geographers*, 103, 1072-1077, 2013.
- 57 Hoaglin, D. C. and Welsch, R. E.: The hat matrix in regression and ANOVA, *The American Statistician*, 32, 17-22, 1978.

58 Huang, B., Wu, B., and Barry, M.: Geographically and temporally weighted regression for modeling spatio-temporal
59 variation in house prices, *International Journal of Geographical Information Science*, 24, 383-401, 2010.

60 Hurvich, C. M., Simonoff, J. S., and Tsai, C. L.: Smoothing parameter selection in nonparametric regression using an
61 improved Akaike information criterion, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 60,
62 271-293, 1998.

63 Loader, C. R.: Bandwidth selection: classical or plug-in?, *The Annals of Statistics*, 27, 415-438, 1999.

64 Mennis, J. L. and Jordan, L.: The distribution of environmental equity: Exploring spatial nonstationarity in multivariate
65 models of air toxic releases, *Annals of the Association of American Geographers*, 95, 249-268, 2005.

66 Pace, R. K., Barry, R., Gilley, O. W., and Sirmans, C.: A method for spatial-temporal forecasting with an application to real
67 estate prices, *International Journal of Forecasting*, 16, 229-246, 2000.

68 Sun, T. Y., Conroy, G., Donner, E., Hungerbühler, K., Lombi, E., and Nowack, B.: Probabilistic modelling of engineered
69 nanomaterial emissions to the environment: a spatio-temporal approach, *Environmental Science: Nano*, 2, 340-351,
70 2015.

71 Takahashi, K., Kulldorff, M., Tango, T., and Yih, K.: A flexibly shaped space-time scan statistic for disease outbreak
72 detection and monitoring, *International Journal of Health Geographics*, 7, 14, 2008.

73 Tobler, W. R.: A computer movie simulating urban growth in the Detroit region, *Economic geography*, 46, 234-240, 1970.

74 Wang, W., Zhao, J., Cheng, Q., Carranza, E.J.M.: GIS-based mineral potential modeling by advanced spatial analytical
75 methods in the southeastern Yunnan mineral district, China. *Ore Geology Reviews*, 71, 735-748.
76 <https://doi.org/10.1016/j.oregeorev.2013.08.005>, 2015.

77 Wheeler, D. C. and Waller, L. A.: Comparing spatially varying coefficient models: a case study examining violent crime
78 rates and their relationships to alcohol outlets and illegal drug arrests, *Journal of Geographical Systems*, 11, 1-22, 2009.

79 Wu, B., Li, R., and Huang, B.: A geographically and temporally weighted autoregressive model with application to housing
80 prices, *International Journal of Geographical Information Science*, 28, 1186-1204, 2014.

81 Wu, S., Wang, Z., Du, Z., Huang, B., Zhang, F. and Liu, R.: Geographically and temporally neural network weighted
82 regression for modeling spatiotemporal non-stationary relationships. *International Journal of Geographical Information*
83 *Science*, 1-27. 2020.