

Supplement of "A multirate mass transfer model to represent the interaction of multicomponent biogeochemical processes between surface water and hyporheic zones (SWAT-MRMT-R 1.0)"

Yilin Fang^{1,*}, Xingyuan Chen¹, Jesus Gomez velez², Xuesong Zhang³, Zhuoran Duan¹, Glenn E. Hammond⁴, Amy E. Goldman¹, Vanessa A. Garayburu-Caruso¹, and Emily B. Graham¹

¹Pacific Northwest National Laboratory, Richland, Washington, USA

²Vanderbilt University, Nashville, Tennessee, USA

³Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, Maryland, USA

⁴Sandia National Laboratories, Albuquerque, New Mexico, USA

Contents of this file

1. Table 1

2. Figure S1

Table 1. Vertical and lateral residence times at selected reaches along the Columbia River.

Reach number	Vertical RT (h)	Lateral RT (h)
27	8.98	1380
28	8.98	1380
20	8.98	1380
24	8.99	1380
43	143	46100
53	143	46100
67	143	46100
77	8.49	408
88	7.88	408
93	112	22100
100	258	93800
101	267	93800




Figure 1. Relationship between observed DOC and stream flow discharge.