

RadNet 1.0: Exploring deep learning architectures for longwave radiative transfer

Ying Liu¹, Rodrigo Caballero¹, and Joy Merwin Monteiro¹

¹Department of Meteorology, Stockholm University, Stockholm, Sweden

Correspondence: Ying Liu (ying.liu.sweden@gmail.com)

Abstract. Simulating global and regional climate at high resolution is essential to study the effects of climate change and capture extreme events affecting human populations. To achieve this goal, the scalability of climate models and efficiency of individual model components are both important. Radiative transfer is among the most computationally expensive components in a typical climate model. Here we attempt to model this component using a neural network. We aim to study the feasibility of replacing an explicit, physics-based computation of longwave radiative transfer by a neural network emulator, and assessing the resultant performance gains. We compare multiple neural-network architectures, including a convolutional neural network and our results suggest that the performance loss from the use of conventional convolutional networks is not offset by gains in accuracy. We train the networks with and without noise added to the input profiles and find that adding noise improves the ability of the networks to generalise beyond the training set. Prediction of radiative heating rates using our neural network models achieve up to 370x speedup on a GTX 1080 GPU setup and 11x speedup on a Xeon CPU setup compared to the a state of the art radiative transfer library running on the same Xeon CPU. Furthermore, our neural network models yield less than 0.1 Kelvin per day mean squared error across all pressure levels. Upon introducing this component into a single column model, we find that the time evolution of the temperature and humidity profiles are physically reasonable, though the model is conservative in its prediction of heating rates in regions where the optical depth changes quickly. Differences exist in the equilibrium climate simulated when using the neural network, which are attributed to small systematic errors that accumulate over time. Thus, we find that the accuracy of the neural network in the “offline” mode does not reflect its performance when coupled with other components.

1 Introduction

Computational models of Earth’s climate are essential tools to advance our understanding of the climate system and our ability to predict its response to perturbations such as increased levels of greenhouse gases. Climate models contain algorithmic representations of the various components of the climate system like the atmosphere, ocean, sea ice and land surface. Our ability to predict future changes in climate depends crucially on the accuracy of these models and the extent to which interactions between various components of the climate system are represented.

A basic requirement for increased model fidelity, particularly at the regional scale, is increased spatial resolution. However, the computational burden increases roughly as the fourth power of spatial resolution (since resolution must increase along all

three spatial dimensions, and the time step reduced to ensure numerical stability). To address this problem, various approaches have been used including improved model scalability (Dennis and Loft, 2011) and the use of low-precision floating point operations (Palmer, 2014).

Long simulations using high resolution climate models are needed to explore key questions in climate research, particularly changes in the statistics weather extremes such as windstorms and precipitation events. Radiative transfer (RT) in the atmosphere is among the most computationally burdensome components of such simulations. While the basic equations for calculating RT are straightforward, the complex nature of the absorption bands of greenhouse gases such as carbon dioxide and water vapour requires separate calculation over a very large number of small spectral intervals to obtain accurate results. Since such a line-by-line calculation is extremely computationally intensive and not feasible in a realistic climate model integration, it is necessary to group individual absorption lines into bands or clusters with similar properties as in the correlated-k method (Fu and Liou, 1992). Such methods can dramatically improve the computational performance while retaining adequate accuracy in the computation. Many state-of-the-art climate models use the Rapid Radiative Transfer Model for General circulation models (RRTMG). RRTMG is based on the single-column correlated k-distribution reference model RRTM (Iacono et al., 2008b). RRTMG tries to strike a balance between computational complexity and accuracy by reducing the number of calculations per band while ensuring fidelity with the RRTM code (Iacono et al., 2008a). Nonetheless, even when employing such simplified schemes, RT remains amongst the most numerically expensive components of climate models, and a variety of strategies have been developed to reduce this cost (see for example Pincus and Stevens, 2013, and references therein).

In this paper, we explore the potential performance gains achievable by using a neural network (NN) to calculate radiative transfer. Specifically, we train a variety of alternative NN architectures on a set of radiative heating rate profiles computed using a state-of-the-art RT code (see Sections 2), and compare the computational performance of the NN with that of the RT code itself. Note that this comparison only serves to assess the performance of RT calculation in standalone form. We expect a suitably-trained neural network to be a drop-in replacement for the RT code in a full climate model, and expect that other computational costs—such as data transfer within and between computational nodes—will not change, but we do not explicitly address this issue in this exploratory study. Instead, our focus here is on identifying the most suitable NN architecture in terms of accuracy and computational performance. We also explore the behaviour of the NN in a time-evolving single-column radiative-convective model (Section 4).

Recent advances in NNs have led to rapid progress in the accuracy of pattern and image recognition tasks. In particular, convolutional neural networks (CNNs) (Krizhevsky et al., 2012a) have achieved impressive results for image classification (Krizhevsky et al., 2012b), while recurrent neural networks (RNNs) have made breakthroughs in sequence-to-sequence learning tasks such as machine-translation (Wu et al., 2016). Efforts to use machine learning techniques to model actual physical processes in a climate model have increased recently (Schneider et al., 2017; Gentine et al., 2018; Rasp et al., 2018; O’Gorman and Dwyer, 2018; Scher, 2018; Brenowitz and Bretherton, 2018; San and Maulik, 2018; Brenowitz and Bretherton, 2019; Yuval and O’Gorman, 2020). In particular, it is now being recognized that physical processes whose representation in climate models has usually been inexact and parameterised could potentially be improved by using machine learning techniques. RT, on the other hand, has always been an attractive candidate to optimize in climate models because of the large computational

cost, as discussed above. Optimization has been attempted using traditional optimization, porting to new architectures such as GPUs (Price et al., 2014; Mielikainen et al., 2016; Malik et al., 2017) and using NNs to approximate RT. Initial attempts to retrieve radiative heating profiles used shallow (one hidden layer) networks (Chevallier et al., 1998), and similar NN architectures were successfully used to replace RT in decadal simulations using conventional climate models (Krasnopolsky et al., 2005, 2008, 2009). Recently, a deep NN was used to replace RT in a high resolution GCM, and was successfully used to run the GCM for one year (Pal et al., 2019). These studies show the capability of NNs to accurately approximate radiative heating profiles in a particular climate regime, while raising questions about how generalizable this learning actually is in terms of handling perturbed climate states. Studying the effect of perturbations (in sea-surface temperature, greenhouse gases, aerosols or cloud properties) on the climate of a model is a very typical use-case in climate science, and the performance of NNs in such scenarios has yet to be studied carefully.

In the context of machine learning for climate modelling applications, the following questions are still not well understood (Dueben and Bauer, 2018):

- What NN architectures are most suitable?
- What is the accuracy-efficiency tradeoff between different NN architectures?
- What accuracy loss can we expect when the NN is provided with “non-typical” input values, i.e. values very different from those in the training sample, such as would occur in a perturbed climate experiment?
- What is the speed-up we can expect by replacing a traditional RT scheme with a NN?

Our aim here is to address these four questions. To limit the scope of this exploratory study, we focus on longwave radiative transfer under clear-sky conditions (henceforth, RT thus refers to clear-sky longwave radiative transfer). We use the RRTMG library available within the climt modelling toolkit (Monteiro et al., 2018) to generate radiative cooling profiles to train the NN models. In particular, we compare the accuracy-computational complexity tradeoff between five kinds of NN architectures on both CPU and GPU. We also study the loss in accuracy if perturbations are added to the input. The question of accuracy loss is all the more relevant in RT due to its mathematical structure – since RT is modelled as an integral equation, localised perturbations have global impacts on the profile of radiative heating or cooling obtained.

The paper is organized as follows. The preparation of data for training and validation of the NNs is presented in Section 2. Section 3 presents the NN structures and parameters we have used. Evaluation results are presented in section 4. Finally, we present a brief discussion along with concluding remarks in Section 5.

2 Data and Methods

While radiative transfer is inherently three dimensional, increasing its complexity and computationally cost, it is common to assume horizontal homogeneity (independent column assumption) and retain only a single (vertical) dimension (Meador and Weaver, 1980). This independent column assumption underlies almost all radiative transfer codes used in weather and climate

models, and reduces radiative transfer calculation to an "embarrassingly parallel" 1-dimensional problem in each vertical column of the atmosphere. For a given longitude-latitude point, RT can be represented by a vector whose length is the number of vertical levels into which the column is discretized. The calculation of RT under clear sky (cloud-free) conditions is based on a number of inputs, including vectors of atmospheric pressure, air temperature and specific humidity at each level, while surface temperature and carbon dioxide mixing ratio are represented as scalars. While the clear-sky RT in the atmosphere is affected by other greenhouse gases like methane and aerosols like sulphates, we restrict ourselves to using the above quantities in this exploratory study.

2.1 The ERA-Interim Dataset

We use the ERA-Interim dataset (Dee et al., 2011) to provide temperature and humidity profiles for training the neural network. The horizontal resolution of the data is $0.75^\circ \times 0.75^\circ$ in the horizontal. We use 6 hourly model-level data, which has a higher resolution in the vertical as compared with the pressure level data. The vertical grid is a non-uniform η -coordinate grid with 60 mid-levels from the surface to 0.2 hPa and 61 interface levels from the surface upto 0.1 hPa. This implies that pressure is not a constant and is therefore an additional input to the neural network.

The ERA-Interim dataset consists of 38 years of data spanning the period 1979 to 2016, which amounts to around 6.5 billion sample profiles. We employ the first 7 years of ERA-Interim historical data as the training dataset, i.e., data from 1979 to 1985 and the last 2 years of the ERA-Interim historical data as the validation dataset, i.e., data from 2015 to 2016. Considering the model training time, we have applied random sampling of 1% with respect to each year in the training and validation datasets. This gives around 12 million training samples and 3.5 million validation samples. After sampling, we name the training dataset as $Dataset_1$ and the validation dataset as $Dataset_{1, val}$. The reason for using this data separation schema is because that we would like to examine whether our radiation prediction model is able to generalize to unseen/future data inputs while being trained on the oldest data.

2.2 Perturbed Dataset

Figure 1 shows the mean and variance of ERA-Interim air temperature, humidity and radiative heating rates calculated using RRTMG from 1979 to 2016. Using the above statistics, we have augmented our training data by created a perturbed dataset as follows:

1. Pick an original profile from the historical samples.
2. Generate a random air temperature profile assuming Gaussian distribution at each vertical level using the statistics from Figure 1.
3. Generate a random weight (between -0.2 to 0.2) for the generated air temperature profile.
4. Generate an augmented air temperature profile by adding together the original profile with the weighted random profile vertical level wise.




Figure 1. (a,b) ERA-Interim air temperature (K) and humidity (g kg^{-1}) statistics. (c) Longwave radiative rates (K day^{-1}) calculated using RRTMG. Vertical axis is pressure in Pa.

5. Calculate the maximum humidity given the air temperature and pressure at each vertical level.
6. Calculate the original relative humidity ratio using humidity divided by the maximum humidity at each vertical level.
- 125 7. Calculate the new maximum humidity given the generated air temperature and pressure at each vertical level.
8. Generate the corresponding humidity by multiplying the new maximum humidity and the original relative humidity ratio at each vertical level.
9. We keep the surface temperature and the carbon dioxide mixing ratio the same as the original profile.

The motivation for adding random, vertically uncorrelated perturbations is that the optical properties of the atmosphere (which determine the radiative heating profiles) can be quite noisy in the vertical. This noisiness is due to the presence of clouds, hydrometeors, aerosols and horizontal advection of water vapour at different levels in the atmosphere. Changes in optical depth due to the above factors need not have a strong vertical correlation either. The kind of perturbations we have added represent an extreme case of this physically-motivated reasoning.

Augmented datasets are generated using $Dataset_1$ and $Dataset_{1.val}$. Then, the augmented dataset are 50-50 mixed with $Dataset_1$ and $Dataset_{1.val}$ respectively to create $Dataset_2$ and $Dataset_{2.val}$. The purpose of generating $Dataset_2$ and $Dataset_{2.val}$ is that we would like to use it to investigate the generality of our RT prediction model. The specific evaluation procedures are described in the evaluation section.

2.3 The RT dataset

The calculation of the radiative fields for the historical and perturbed datasets are calculated using the RRTMG component available in the climt modelling toolkit (Monteiro and Caballero, 2016; Monteiro et al., 2018). This component is a python wrapper over the RRTMG fortran library, and provides convenient access to the radiation fields. The statistics of the generated radiative heating profiles are also shown in Figure 1.

3 Neural Network Models

3.1 Neural Network Basics

A neural network is composed of multiple neurons, or even multiple layers of neurons in order to model complex scenarios. A simple neural network is a feed-forward network where information flows only in one direction from input to output. Multilayer perceptron (Gardner and Dorling, 1998) is the most common feed-forward NN. It consists of an input layer that passes the input vector to the network, one or more hidden layers and an output layer. There are usually activation functions applied in each layer. An activation function usually introduces non-linearity in order to allow a NN to tackle with complicated problems and learn complex representations.

Convolutional NN is another type of NN designed for image-focused tasks. It is widely used in many fields such as image classification, object detection and image segmentation (Krizhevsky et al., 2017). CNNs usually consist of three types of layers, convolutional layers, pooling layers and fully-connected layers. A convolutional layer is composed of learnable kernels or filters. The kernel usually considers a small region of input at one time, but covers the entirety of the input. Specifically, it slides over the input spatially and computes dot products between the kernel and the area of input covered by the kernel. With each kernel, a convolutional layer produces an activation map, whose size depends on whether there is a stride or padding. All the activation maps will be stacked together along the depth dimension and passed on to the next layer (O'Shea and Nash, 2015). Neurons in a layer are connected to only a small region of the previous layer instead of everything, which is different

Model Name	Model Structure							Number of Parameters
Model A	input-60x4	fc-512	fc-1024	fc-512	output-60			1202176
Model B	input-60x4	fc-512	fc-1024	fc-2048	fc-1024	fc-512	output-60	5396480
Model C	input-60x4	conv-3x3-128	conv-3x3-256	fc-512	output-60			7666816
Model D	input-60x4	conv-3x3-128	conv-3x3-256	conv-3x3-256	fc-512	output-60		7994496
Model E	input-62x6	conv-3x3-128	conv-3x3-256	fc-512	output-60			15531136
Model F	input-62x5	conv-3x3-128	conv-3x3-256	fc-512	output-60			7928960

Table 1. Neural network models used for predicting RTs. "fc-X" represents a fully connected layer with X number of neurons. "conv-YxY-X" represents a convolutional layer with X number of YxY filters.

from feedforward neural networks. In this way, convolutional layers are better at extracting locality-dependent features, such as shapes and patterns in images.

In the context of RT, we use CNNs to evaluate whether the sensitivity to localised features improves the prediction performance of deep neural networks. In particular, strong local changes in the optical properties of the atmosphere are fairly common due to the presence of clouds or horizontal advection of water vapour. While this work focuses on clear-sky radiation, we study the ability of CNNs to recognise and respond to such local features in the single column simulations.

Table 1 illustrates the structures and parameters of our neural networks. Specifically, we have designed five neural networks, including two feedforward neural networks and three convolutional neural networks (CNN). The input data to the CNNs is prepared by concatenating different variables along the second dimension instead of using different variables as "channels". Therefore, the CNNs use 2D convolutional filters instead of 1D filters. Model A and Model B are implementations of feedforward neural networks with different numbers of layers and of neurons in each layer. Model C is a simplified CNN implementation based on previous work (Simonyan and Zisserman, 2014). The stride of convolutional filters is set to 1 so that the convolutional filters go through the input array with 1 element each step. We have not applied any padding to the input. We have not used pooling layers in between convolutional layers and the convolutional filters are the classic 3x3 filters. Model D is a variant of Model C with one more convolutional layer. Model E has the same neural network structure as Model C. The only difference is that model E has padded the input with an edge of zeros to emphasize on edges of the input. We have used Tensorflow 1.8.0 library for the neural network implementation.

In addition to the above models, we also evaluate a variant of model E denoted as model F. Model F is based on Model E but with a fixed pressure grid. This means that Model F does not take pressure values as input, and interpolates air temperature and humidity from model levels onto a fixed, time-invariant pressure grid. While this configuration reduces the dimensionality of the input, it requires extrapolation of the ERA-Interim data or the calculated/predicted RT to the fixed grid. Specifically, the inputs of a sample profile are B-spline extrapolated according to a fixed pressure grid. We extrapolate the air temperature and humidity values onto the fixed pressure grid based on the profile's pressure range. The inputs corresponding to the rest of the pressure levels are set to 0. After running through model F, the RTs on the static pressure grid are B-spline interpolated back to the original pressure levels, which are the final results. It is important to mention that we constructed the static grid

using 15 equally spaced pressure levels from 1 to 500 Pa, another 15 equally spaced pressure levels from 550 to 50000 Pa, and 30 equally spaced pressure levels from 50300 to 103000 Pa. We made this design choice by observing the distribution of the ERA-Interim data to ensure that our fixed grid encompasses most common pressure profiles in order to achieve a better accuracy on the extrapolation and interpolation. We used model F to run the single-column model simulation presented below, which employs a fixed pressure grid.

3.2 Model training

We trained our five NNs with two datasets, resulting 10 different models. The first dataset is the aforementioned ERA-Interim dataset, namely, $Dataset_1$. The second dataset is the augmented dataset, i.e., $Dataset_2$, in order to generalize the model to a wider operational region beyond $Dataset_1$. Each neural network is trained using the training dataset of either $Dataset_1$ or $Dataset_2$ and validated against either $Dataset_{1.val}$ or $Dataset_{2.val}$.

Each model was trained with 30 epochs under a batch size of 128 starting with a learning rate of 0.001, which then exponentially decays every 10 epochs with base 0.96. This setup was empirically obtained while we observe that all models have converged after the training. Mean squared error is used as the loss function in all models. Parametric Rectified Linear Units (pReLUs) (He et al., 2015a) are used as activation functions in all models since PReLU is able to resolve the problem of vanishing gradient during model training. Adam optimizer (Kingma and Ba, 2014) is employed to compute the gradients.

We present the evaluation results regarding the performance of these models in the next section.

4 Evaluation

4.1 Evaluation Setup

We prepared two datasets, i.e., $Dataset_{1.val}$ and $Dataset_{2.val}$, to evaluate our neural network models. $Dataset_{1.val}$ is used to evaluate the accuracy of the trained models with realistic future data. $Dataset_{2.val}$ is used to evaluate the generality of the trained models as it contains profiles that are perturbed versions of the ERA Interim data.

4.2 Prediction Accuracy

We use vertical level-wise root-mean-squared errors (RMSE) to compare our NN generated radiative cooling rates with those generated by the RRTMG algorithm. Figures 2-5 present results for the different NN models. The RMSE is calculated by taking the difference between NN- and RRTMG-calculated radiative cooling profiles.

Figure 2 presents the RMSE when the NN models are trained using $Dataset_1$ and validated against $Dataset_{1.val}$. These experiments are performed to evaluate the capability of different NN models to predict RT when the atmospheric profiles are sampled from the ERA-Interim dataset itself. We see that a simple 3 layer feedforward neural network (Model A) is able to predict heating rates with a median RMSE of lesser than 0.01 K/day across all pressure ranges. The performance does not improve when more layers of directly connected neurons are added, as shown by Model B. We observe significant RMSE

improvement while using CNNs (Model C,D,E). However, the performance differences among these three CNN models are
215 not substantial except the RMSEs near the surface, which tend to have higher variability as shown in the statistics in Figure 1.
Since surface radiation is particularly important to the climate, efforts have been made to minimize its prediction error. The
input matrix of Model C are padded with zeros in order to allow convolutional filters to put equal emphasis on the edges values
as the middle ones (Innamorati et al., 2018). This creates Model E, which shows much better prediction accuracy on the bottom
and top pressure levels.

220 Figure 3 presents the RMSE when the NN models are trained using $Dataset_2$ and validated against $Dataset_{1, val}$. In this
experiment, we examine whether it is possible to expand the operational region of the NN models without compromising on
their performance on the ERA-Interim dataset. Comparing to Figure 2, we see that the increased generality comes at the cost
of roughly doubled RMSE across all models.

The improvement on generality is suggested by the results shown in Figure 4 and Figure 5 when the NN models are trained
225 using $Dataset_1$ or $Dataset_2$ and validated against $Dataset_{2, val}$. The RMSE increases by almost 100 times across all models
trained with $Dataset_1$ and validated against $Dataset_{2, val}$ (Figure 4) when compared to their validation against $Dataset_{1, val}$
(Figure 2). This suggests that that models trained with $Dataset_1$ cannot really generalize to predict heating profiles from
 $Dataset_{2, val}$. On the other hand, the RMSE increases 10 times when the models are trained using a wider range of data, i.e.,
 $Dataset_2$, as shown in Figure 5. This is mainly because that the model needs to cover a larger operational region.

230 When trained on $Dataset_1$ and validated against $Dataset_{2, val}$ (Figure 4), the RMSE in Model B is significantly higher.
This observation leads us to believe that Model B is more likely to over-fit the training dataset. Given that $Dataset_2$ is more
perturbed than $Dataset_1$ and more parameters and layers in Model B, the nature of feedforward NN (Goodfellow et al., 2016)
makes Model B more deeply coupled with patterns observed in the training data ($Dataset_1$), which leads to larger errors while
evaluating against $Dataset_{2, val}$.

235 Model F displays significant errors on both edges of the pressure levels. This is due to extrapolation errors. Specifically, if
the lowest pressure level in an atmospheric profile is lower than the lowest pressure level of the fixed grid, the profiles need to
be extrapolated. The same issue arises with the highest pressure levels as well. Thus, the errors in model F are mainly due to
these extrapolation based artifacts rather than an issue with the training itself. In fact, this model provides the most stable time
integration of the single column model.

240 In the above evaluations, we have shown that CNN-based models achieve much lower prediction RMSE than feedforward
NN models. However, in the next section, we show that CNN-based models tend to have much slower prediction speed, i.e.,
less speedup as compared with the feedforward models.

4.3 Prediction speed

In this section we compare the computation time of RRTMG and NN models using GPUs and CPUs. These performance
245 evaluations have been performed using Intel Xeon CPU E3-1230 v5 @ 3.40GHz, Nvidia GTX 1060 GPU with 6GB of onboard
memory and Nvidia GTX 1080 GPU with 8GB of onboard memory. Both GPUs we use are commodity hardware, and are easily
available in the market. RRTMG was run in a single-threaded mode for the purposes of this evaluation.




Figure 2. Models are trained with $Dataset_1$ and evaluated against $Dataset_{1.val}$. The plots have 30 linearly spaced levels between 0 and 101300 pascals. The errors from each model are binned to these equally spaced intervals for easier reading. The boxes in the plots present the boxplot of the RT MSEs at every level. Specifically, the boxes describe the 25 (Q1), 50 (Q2) and 75 (Q3) percentile of the MSEs while the two whiskers extend from the edges of box to 1.5 times the interquartile range (Q3-Q1).

Table 2 summarizes the speedups using NN models to predict RT as compared to RRTMG. The calculation time of NN code and RRTMG code are profiled using the python line_profiler based on cProfile. The execution time results are averaged from 10 measurements with execution of 100 000 predictions per measurement. Since RT calculations are embarrassingly parallel, we are able to use batch predictions in our NN models while using a single GPU. The overall results show that the larger the batch size, the larger the speedup observed as long as the CPU or GPU memory is sufficient. In other words, the calculation of M radiative heating profiles is faster than M times the time taken to predict one such profile. This is because of the efficiency of matrix multiplications in NNs while conducting NN forward pass in batches. We note that such a speedup is not possible in a physics-based RT scheme since the calculation of RT for an arbitrary atmospheric profile cannot be expressed as a simple matrix multiplication.

The results show that using only the Xeon CPU, NN model A and B are able to achieve speedups up to 10.88x and 2.82x respectively using a batch size of 1024. When using GTX 1060, we are able to achieve speedups of 123x in model A, 61x




Figure 3. Models are trained with $Dataset_2$ and evaluated against $Dataset_{1.val}$

in model B and 2.8x to 4.5x in CNN-based models (C,D,E). With GTX 1080, which has a larger memory and a faster clock
 260 speed, we observe speedups up to 370x in model A, 123x in model B and 4.4x to 7.7x in CNN-based models (C,D,E).

The results indicate that if the prediction accuracy of Model A is sufficient for a climate simulation, it will provide the greatest
 calculation speedup either using CPU or GPU. Since NNs with comparable or worse accuracy have been used for simulations
 ranging from months to years (Krasnopolsky et al., 2008; Pal et al., 2019), Model A is a promising candidate for modelling
 applications since similar performance gains using a full RT code seems to require a complete rewrite for GPUs (Price et al.,
 265 2014; Mielikainen et al., 2016; Wang et al., 2020). For simulations requiring a higher accuracy, Model C provides significant
 speedups even if a normal GPU is available on the platform.

4.4 Single column model simulation

To explore the ability of the NN model to generalise to new situations, we compare the climate of a single column model when
 RRTMG is replaced by the NN model F (see previous section for a description of model F). The single column model uses a
 270 diffusive boundary layer (Reed and Jablonowski, 2012), a slab surface of 50 meters thickness which behaves like an oceanic




Figure 4. Models are trained with $Dataset_1$ and evaluated against $Dataset_{2.val}$

Baseline RRTMG	0.37 ms									
NN Model Name / Hardware	Xeon CPU E3-1230			GTX 1060			GTX 1080			
NN Batch size	64	256	1024	64	256	1024	64	256	1024	4096
Model A	5.87	10.28	10.88	18.50	61.67	123.33	16.08	61.67	123.33	370.00
Model B	1.87	2.74	2.82	13.70	37.00	61.67	14.80	46.25	74.00	123.33
Model C	0.14	0.14	0.14	3.19	4.11	4.57	4.25	5.52	7.40	7.71
Model D	0.11	0.11	0.11	2.52	3.03	3.33	3.52	4.25	5.44	5.52
Model E	0.09	0.09	0.09	2.16	2.59	2.82	2.98	3.67	4.63	4.40

Table 2. Speedups when using NN models to predict RTs comparing to calculating RTs using RRTMG. Result for RRTMG is shown for the calculation of a sample in units of milliseconds. Results for NN models are shown as speedups for different batch sizes as compared to the RRTMG calculation on the Xeon CPU.

mixed layer, the RRTMG shortwave component, and the Emanuel convection scheme (Emanuel and Zivkovic-Rothman, 1999). The model has no seasonal or diurnal cycle. Carbon dioxide concentration is fixed at 300 ppm, and a fixed ozone concentration




Figure 5. Models are trained with $Dataset_2$ and evaluated against $Dataset_{2.val}$

is prescribed using an observed tropical profile. The model uses pressure as the vertical coordinate and has 60 equally spaced vertical levels between 1013.2 hPa and the model top value at 0 hPa. The model time step is 10 minutes. The tendencies from the various components are stepped forward in time using a third order explicit Adams-Bashforth scheme.

The model is initialised with a dry, isothermal state. We use RRTMG’s longwave component to drive the model until the RMS error between the RRTMG calculated longwave heating rates and those predicted by model F falls below a threshold of 0.5 K/day. Once the errors falls below this value, model F takes over and RRTMG’s longwave component is never used again for the rest of the simulation (shortwave radiation is computed using RRTMG throughout). The switch from RRTMG to model F happens after around 14 days of simulation. This simulation is denoted as “RadNet” in Fig. 6. Another simulation continues to use RRTMG longwave radiation until the end of the simulation and is denoted as “RRTMG” in Fig. 6. As discussed subsequently, the RadNet simulation has a bias in the stratosphere and the temperature profile of the top three levels is constrained to the RRTMG simulation to prevent the simulation from blowing up. Both simulations are run for 2100 days and equilibrium is reached around 1600 days, with constant temperature and humidity profiles afterwards.




Figure 6. Comparison of the vertical profiles of temperature (first row), longwave heating rates (second row) and specific humidity (third row) for the RadNet and RRTMG simulations at three different times.

285 Within the troposphere, both simulations show a realistic moist-adiabatic temperature profile and are in reasonable quanti-
tative agreement. However, there are substantial differences in the stratosphere, and the equilibrium position of the tropopause
seen in Fig. 6c in the RadNet simulation is higher by around 50 hPa as compared to the RRTMG simulation. This is because
Model F has a cooling bias in the upper atmosphere as seen in Fig. 6d, which makes it convectively unstable and therefore the
tropopause shifts upward. The tropospheric temperature profiles are identical since they are set by the convective parameteri-
290 zation in such convectively unstable situations.

As the boundary layer fluxes water vapour into the column from the surface, the atmosphere becomes opaque to longwave
radiation in the lower levels and therefore the longwave cooling is strongest in the level just above the moist, opaque part of
the atmosphere. Figure 6d shows that the cooling peak predicted by model F has a smaller magnitude and is located lower
in the atmosphere. The lower cooling rate peak predicted by the NN results in the slower evolution of the RadNet simulation
295 as compared to the RRTMG simulation, resulting in the difference in height between the two simulations (the cooling peak
rises over time as the convection tries to eliminate the instability produced by radiative cooling). The cooling peak in the
RadNet simulation is situated close to the location of the strongest gradient in water vapour (where the atmosphere transitions
from being opaque to transparent to longwave radiation), which is physically accurate. The differences in magnitude are larger
slightly earlier in the simulation, where the atmospheric profiles are quite unlike the profiles in the training dataset. It seems
300 unlikely that neural nets can predict such “spiky” profiles correctly since the predicted results tend to be smooth in general.
However, the Radnet predicted profiles provide sufficient cooling to make the atmosphere convectively unstable and eventually
mix the entire troposphere of the model.

The NN has a systematic warm bias in the lowest layer of the model, which may be linked to the interpolation errors
discussed previously for model F. This warm bias results in a slightly warmer surface temperature ($\sim 0.5\text{K}$) in the RadNet
305 simulation as seen in Fig. 6c. The warmer profile supports a larger amount of water vapour, and the RadNet simulation has a
moist bias in the lower troposphere as well.

We see that small systematic errors in the predicted heating rates can have a non-trivial effect on the simulated climate
in a single column model, especially in the upper layers of the atmosphere. In particular, errors in radiative heating near the
tropopause can dramatically change the structure of this part of the atmosphere. The neural network tends to cool the upper
310 atmosphere a little more, making it more convectively unstable and pushing the convection and tropopause higher.

To verify the accuracy of the predicted heating profiles, we use the atmospheric profiles from the RadNet simulation to drive
the RRTMG longwave component. The NN and RRTMG heating profiles generated are presented in Fig. 7. The heating profiles
predicted by the NN are fairly accurate, especially in the later parts of the simulation when the atmospheric profiles are similar
to those in the training sample space. The NN predicts the location of the cooling peak accurately even when the atmospheric
315 profiles are unlike those in the training sample space, though it underestimates the magnitude. RRTMG produces fairly noisy
heating profiles in the stratosphere, reflecting the noisy temperature profile simulated by the NN. The noisy stratospheric
temperature profile appears to be a result of the fact that the training data for model F was generated using atmospheric profiles
that had additional noise added to them, which results in noisy heating profiles used for training.




Figure 7. Comparison of the vertical profiles of longwave heating rates predicted by the NN and RRTMG for atmospheric profiles from the RadNet simulation.

5 Discussion and Conclusions

320 Radiative transfer was probably among the first climate model components that neural network models aimed to replace in
 climate simulations. The evolution of NN models has paralleled the evolution of NN architectures themselves, with initial
 attempts using shallow networks while recent attempts (including our own) using deep networks. Since both shallow and deep
 networks seem to perform reasonably well in model simulations (Krasnopolsky et al., 2008; Pal et al., 2019), the question of
 which type of architecture is more suitable inevitably arises.

325 In this paper, we have employed two elementary but widely used classes of neural networks, namely feedforward and con-
 volutional neural networks. We believe that the range of model architectures we have presented in this paper are representative
 within these classes of networks. Our experiments not only explore differences between these two network architectures, but
 they also propose a validation workflow:

- Traditional validation using metrics such a mean-squared error.

330 – Validation against a perturbed dataset, which helps evaluate generalizability of the networks directly

- Validation using a hierarchical climate modelling framework such as climt. Simple climate models such as the single-column model helps climate scientists not familiar with neural networks evaluate the physical consequences of network architecture choices. The neural network could then be deployed in a more complicated setup such as a general circulation model and evaluated again.

335 We believe that different methods of validating a neural network is essential to compare different network architectures in a scientifically relevant manner.

Our experiments show that:

- A larger number of parameters in a neural network lead to slower prediction, as could be expected intuitively
- More parameters do not always lead to a better prediction accuracy

340 Classic convolutional networks provide high accuracy at a higher computational cost. The number of parameters in CNNs can be reduced in multiple ways. For instance, using a 1-D convolutional filter could provide performance gains at the expense of losing the correlation information between different input fields like temperature and specific humidity. Similarly, using a larger stride in convolutional layers, using pooling layers in between convolutional layers or reducing model layers are all ways in which the performance of the classical CNN could be improved. Going beyond the classic architectures we have
345 explored, a variety of architectures have been recently proposed which might increase both accuracy and/or speedup. These architectures include the residual blocks proposed in Resnet (He et al., 2015b), the depth-wise separable convolution used in MobileNets (Howard et al., 2017) and Xception (Chollet, 2016), among others. Furthermore, EfficientNet (Tan and Le, 2019) has shown that an efficient balancing of network depth, width, and resolution can lead to better performance in terms of prediction accuracy and speed. However, any such reduction of model parameters in CNNs or exploring newer architectures
350 must be accompanied with a rigorous validation procedure, which could be similar to the workflow proposed above.

Recent work in NN theory suggests that the mathematical structure of deep neural networks (a series of linear and non-linear operators applied sequentially) is especially suited to capture functions which can be expressed as the composition of other functions (Mhaskar and Poggio, 2016; Lin et al., 2017). Radiative transfer conforms to this structure very well: the total radiative heating rate is the sum of heating rates in each spectral band, and the heating rate in each spectral band requires
355 the calculation of absorption coefficients at each model level, each independent of the other. The two-stream approximation and the independent column assumptions introduce additional locality and symmetry requirements, constraining the problem further. This mathematical structure suggests that deep neural networks are a natural choice to approximate RT. Furthermore, the presence of highly localised scattering and absorbing substances such as clouds and water vapour suggest that RT might benefit from a NN structure which is sensitive to localised patterns. This suggests that convolutional NNs might be a better
360 model for RT, and our results confirm this. However, our results also show that using convolutional NNs reduces performance by 50-100 times as compared to feedforward NNs with only a marginal increase in accuracy. Thus, within our evaluation setup, deep feedforward NNs present the best compromise between accuracy and performance. We note that the performance losses we have observed for CNNs could be reduced using a variety of techniques noted in the previous paragraph. The performance-

accuracy trade off for each of these techniques needs to be evaluated rigorously and provides a promising avenue for future
365 research.

The ability of NNs to generalise to unfamiliar atmospheric profiles seems to be limited as suggested by the cases where the NNs were validated on the perturbed dataset and the single column model comparisons. These results bring to question the applicability of NN based radiative transfer in research configurations where perturbations to the model state or evolution to a wholly new climate state is routinely performed. Thus, NNs seem to work best in an “operational mode” where the state of
370 the climate or weather prediction model is not expected to change dramatically as compared to the training set. The approach of adding of noise to improve NNs’ ability to generalise beyond the training sample has a long history (Sietsma and Dow, 1991). However, our results show that adding noise to the training dataset results in noisy temperature profiles in simulations, especially in the stratosphere where the temperature profile is closer to pure radiative equilibrium.

The dramatic performance gains when using commodity GPUs makes the use of NNs all the more attractive given that
375 most future high-performance computing configurations will include both GPUs and CPUs. NNs allow batching of multiple atmospheric profiles during matrix multiplications, which allows large performance gains. Such batching is not feasible for an actual RT calculation, and each atmospheric profile has to be handled individually. This may be the reason why rewrites of RRTMG for GPUs (Price et al., 2014; Mielikainen et al., 2016; Wang et al., 2020) give similar performance gains to what we have achieved in our setup using NNs. We note that the comparison between RadNet and rewrites of RRTMG for GPUs does
380 not take into consideration differences in GPU architectures and batch sizes, which could change the exact numbers obtained. However, our results highlight the difference that GPUs make in accelerating RadNet.

Another method to assess the ability of NNs to generalise is to actually build a climate model which includes the NN as a component. Since single column models have no diffusion built-in and cannot transport energy horizontally, we believe that they constitute a tougher test case for NNs as compared to GCMs. The lack of dynamics also makes the results easier
385 to interpret. In our test case, we see that the errors in prediction by the NN has a larger impact in the stratosphere than the troposphere due to the tight control of the tropospheric lapse rate by moist convection. The initial atmospheric profile – dry and isothermal – is quite different from the profiles in the training sample space. While the errors in the initial part of the simulation are larger, the NN predicts physically realistic heating profiles with slight differences in location and magnitude. Such physically plausible behaviour in situations quite different from those the NN was trained on gives us confidence that NNs
390 can indeed be used as climate model components in the future. However, it is clear that better strategies for data preparation, selection of NN architecture and testing trained NNs are required to improve NN performance and enable scientists to interpret their impact on climate model simulations.

Code availability. The code used for training RadNet and the Jupyter notebook which simulates the single column model are available at <http://doi.org/10.5281/zenodo.3884964>.

395 The ERA-Interim data can be downloaded from <https://apps.ecmwf.int/datasets/data/interim-full-daily/levtype=sfc/>

Author contributions. YL designed, trained and validated the neural networks, prepared the input data and participated in design of the project and writing the paper. RC participated in design of the project and writing the paper. JMM wrote the climt code to generate the data, interface RadNet to the single column model, analysed the simulations and participated in writing the paper.

Competing interests. The authors declare no competing interests.

400 *Acknowledgements.* All authors were funded by the Swedish e-Science Research Centre (SeRC).

References

- Brenowitz, N. D. and Bretherton, C. S.: Prognostic Validation of a Neural Network Unified Physics Parameterization, *Geophysical Research Letters*, 45, 6289–6298, <https://doi.org/10.1029/2018GL078510>, 2018.
- Brenowitz, N. D. and Bretherton, C. S.: Spatially Extended Tests of a Neural Network Parametrization Trained by Coarse-Graining, *Journal of Advances in Modeling Earth Systems*, 11, 2728–2744, <https://doi.org/10.1029/2019MS001711>, <http://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019MS001711>, 2019.
- Chevallier, F., Chéruy, F., Scott, N. A., and Chédin, A.: A Neural Network Approach for a Fast and Accurate Computation of a Longwave Radiative Budget, *Journal of Applied Meteorology*, 37, 1385–1397, [https://doi.org/10.1175/1520-0450\(1998\)037<1385:ANNAFA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1998)037<1385:ANNAFA>2.0.CO;2), 1998.
- 410 Chollet, F.: Xception: Deep Learning with Depthwise Separable Convolutions, *CoRR*, abs/1610.02357, <http://arxiv.org/abs/1610.02357>, 2016.
- Dee, D. P. et al.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Dennis, J. M. and Loft, R. D.: Refactoring Scientific Applications for Massive Parallelism, in: *Numerical Techniques for Global Atmospheric Models*, *Lecture Notes in Computational Science and Engineering*, pp. 539–556, Springer, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-11640-7_16, 2011.
- Dueben, P. D. and Bauer, P.: Challenges and design choices for global weather and climate models based on machine learning, *Geoscientific Model Development*, 11, 3999–4009, <https://doi.org/https://doi.org/10.5194/gmd-11-3999-2018>, 2018.
- Emanuel, K. A. and Zivkovic-Rothman, M.: Development and Evaluation of a Convection Scheme for Use in Climate Models, *Journal of the Atmospheric Sciences*, 56, 1766–1782, [https://doi.org/10.1175/1520-0469\(1999\)056<1766:DAEOAC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<1766:DAEOAC>2.0.CO;2), 00553, 1999.
- 420 Fu, Q. and Liou, K. N.: On the Correlated k-Distribution Method for Radiative Transfer in Nonhomogeneous Atmospheres, 49, 2139–2156, [https://doi.org/10.1175/1520-0469\(1992\)049<2139:OTCDMF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049<2139:OTCDMF>2.0.CO;2), 1992.
- Gardner, M. and Dorling, S.: *Artificial Neural Networks (The Multilayer Perceptron) – A Review of Applications in the Atmospheric Sciences*, *Atmospheric Environment*, 32, 1998.
- 425 Gentine, P., Pritchard, M., Rasp, S., Reinaudi, G., and Yacalis, G.: Could Machine Learning Break the Convection Parameterization Deadlock?, *Geophysical Research Letters*, 45, 5742–5751, <https://doi.org/10.1029/2018GL078202>, 2018.
- Goodfellow, I., Bengio, Y., and Courville, A.: *Deep learning*, MIT press, 2016.
- He, K., Zhang, X., Ren, S., and Sun, J.: Delving Deep into Rectifiers: Surpassing Human-Level Performance on ImageNet Classification, *CoRR*, abs/1502.01852, <http://arxiv.org/abs/1502.01852>, 2015a.
- 430 He, K., Zhang, X., Ren, S., and Sun, J.: Deep Residual Learning for Image Recognition, *CoRR*, abs/1512.03385, <http://arxiv.org/abs/1512.03385>, 2015b.
- Howard, A. G., Zhu, M., Chen, B., Kalenichenko, D., Wang, W., Weyand, T., Andreetto, M., and Adam, H.: MobileNets: Efficient Convolutional Neural Networks for Mobile Vision Applications, *CoRR*, abs/1704.04861, <http://arxiv.org/abs/1704.04861>, 2017.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2008JD009944>, 2008a.
- 435

- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2008JD009944>, 2008b.
- 440 Innamorati, C., Ritschel, T., Weyrich, T., and Mitra, N. J.: Learning on the Edge: Explicit Boundary Handling in CNNs, *CoRR*, abs/1805.03106, <http://arxiv.org/abs/1805.03106>, 2018.
- Kingma, D. P. and Ba, J.: Adam: A Method for Stochastic Optimization, *CoRR*, abs/1412.6980, <http://arxiv.org/abs/1412.6980>, 2014.
- Krasnopolsky, V. M., Fox-Rabinovitz, M. S., and Chalikov, D. V.: New Approach to Calculation of Atmospheric Model Physics: Accurate and Fast Neural Network Emulation of Longwave Radiation in a Climate Model, *Monthly Weather Review*, 133, 1370–1383, <https://doi.org/10.1175/MWR2923.1>, 2005.
- 445 Krasnopolsky, V. M., Fox-Rabinovitz, M. S., and Belochitski, A. A.: Decadal Climate Simulations Using Accurate and Fast Neural Network Emulation of Full, Longwave and Shortwave, Radiation, *Monthly Weather Review*, 136, 3683–3695, <https://doi.org/10.1175/2008MWR2385.1>, 2008.
- Krasnopolsky, V. M., Fox-Rabinovitz, M. S., Hou, Y. T., Lord, S. J., and Belochitski, A. A.: Accurate and Fast Neural Network Emulations of Model Radiation for the NCEP Coupled Climate Forecast System: Climate Simulations and Seasonal Predictions, *Monthly Weather Review*, 138, 1822–1842, <https://doi.org/10.1175/2009MWR3149.1>, 2009.
- 450 Krizhevsky, A., Sutskever, I., and Hinton, G. E.: ImageNet Classification with Deep Convolutional Neural Networks, in: *Proceedings of the 25th International Conference on Neural Information Processing Systems - Volume 1, NIPS'12*, pp. 1097–1105, Curran Associates Inc., USA, <http://dl.acm.org/citation.cfm?id=2999134.2999257>, 2012a.
- 455 Krizhevsky, A., Sutskever, I., and Hinton, G. E.: Imagenet classification with deep convolutional neural networks, in: *Advances in neural information processing systems*, pp. 1097–1105, 2012b.
- Krizhevsky, A., Sutskever, I., and Hinton, G. E.: ImageNet Classification with Deep Convolutional Neural Networks, *Commun. ACM*, 60, 84–90, 2017.
- Lin, H. W., Tegmark, M., and Rolnick, D.: Why Does Deep and Cheap Learning Work So Well?, *Journal of Statistical Physics*, 168, 1223–1247, <https://doi.org/10.1007/s10955-017-1836-5>, 2017.
- 460 Malik, M., Grosheintz, L., Mendonça, J. M., Grimm, S. L., Lavie, B., Kitzmann, D., Tsai, S.-M., Burrows, A., Kreidberg, L., Bedell, M., Bean, J. L., Stevenson, K. B., and Heng, K.: HELIOS: AN OPEN-SOURCE, GPU-ACCELERATED RADIATIVE TRANSFER CODE FOR SELF-CONSISTENT EXOPLANETARY ATMOSPHERES, *The Astronomical Journal*, 153, 56, <https://doi.org/10.3847/1538-3881/153/2/56>, 2017.
- 465 Meador, W. E. and Weaver, W. R.: Two-Stream Approximations to Radiative Transfer in Planetary Atmospheres: A Unified Description of Existing Methods and a New Improvement, *Journal of the Atmospheric Sciences*, 37, 630–643, [https://doi.org/10.1175/1520-0469\(1980\)037<0630:TSATRT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<0630:TSATRT>2.0.CO;2), 1980.
- Mhaskar, H. N. and Poggio, T.: Deep vs. shallow networks: An approximation theory perspective, *Analysis and Applications*, 14, 829–848, <https://doi.org/10.1142/S0219530516400042>, 2016.
- 470 Mielikainen, J., Price, E., Huang, B., Huang, H. A., and Lee, T.: GPU Compute Unified Device Architecture (CUDA)-based Parallelization of the RRTMG Shortwave Rapid Radiative Transfer Model, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 9, 921–931, <https://doi.org/10.1109/JSTARS.2015.2427652>, 2016.
- Monteiro, J. M. and Caballero, R.: The Climate Modelling Toolkit, in: *Proceedings of the 15th Python in Science Conference*, pp. 69–74, Austin, USA, http://conference.scipy.org/proceedings/scipy2016/joy_monteiro.html, 2016.

- 475 Monteiro, J. M., McGibbon, J., and Caballero, R.: *symp1* (v. 0.3.2) and *climt* (v. 0.11.0) – Towards a flexible framework for building model hierarchies in Python, 2018, 1–21, <https://doi.org/10.5194/gmd-2018-100>, 2018.
- O’Gorman, P. A. and Dwyer, J. G.: Using Machine Learning to Parameterize Moist Convection: Potential for Modeling of Climate, Climate Change, and Extreme Events, *Journal of Advances in Modeling Earth Systems*, 10, 2548–2563, <https://doi.org/10.1029/2018MS001351>, 2018.
- 480 O’Shea, K. and Nash, R.: An Introduction to Convolutional Neural Networks, *CoRR*, abs/1511.08458, 2015.
- Pal, A., Mahajan, S., and Norman, M. R.: Using Deep Neural Networks as Cost-Effective Surrogate Models for Super-Parameterized E3SM Radiative Transfer, *Geophysical Research Letters*, <https://doi.org/10.1029/2018GL081646>, 2019.
- Palmer, T. N.: More reliable forecasts with less precise computations: a fast-track route to cloud-resolved weather and climate simulators?, 372, 20130 391, <https://doi.org/10.1098/rsta.2013.0391>, 2014.
- 485 Pincus, R. and Stevens, B.: Paths to accuracy for radiation parameterizations in atmospheric models, 5, 225–233, <https://doi.org/10.1002/jame.20027>, <http://agupubs.onlinelibrary.wiley.com/doi/full/10.1002/jame.20027>, 2013.
- Price, E., Mielikainen, J., Huang, M., Huang, B., Huang, H. A., and Lee, T.: GPU-Accelerated Longwave Radiation Scheme of the Rapid Radiative Transfer Model for General Circulation Models (RRTMG), *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 7, 3660–3667, <https://doi.org/10.1109/JSTARS.2014.2315771>, 2014.
- 490 Rasp, S., Pritchard, M. S., and Gentine, P.: Deep learning to represent subgrid processes in climate models, *Proceedings of the National Academy of Sciences*, 115, 9684–9689, <https://doi.org/10.1073/pnas.1810286115>, 2018.
- Reed, K. A. and Jablonowski, C.: Idealized tropical cyclone simulations of intermediate complexity: a test case for AGCMs, *Journal of Advances in Modeling Earth Systems*, 4, <https://doi.org/10.1029/2011MS000099>, 2012.
- San, O. and Maulik, R.: Extreme learning machine for reduced order modeling of turbulent geophysical flows, *Physical Review E*, 97, 042 322, <https://doi.org/10.1103/PhysRevE.97.042322>, 2018.
- 495 Scher, S.: Toward Data-Driven Weather and Climate Forecasting: Approximating a Simple General Circulation Model With Deep Learning, *Geophysical Research Letters*, 45, 12,616–12,622, <https://doi.org/10.1029/2018GL080704>, 2018.
- Schneider, T., Lan, S., Stuart, A., and Teixeira, J.: Earth system modeling 2.0: A blueprint for models that learn from observations and targeted high-resolution simulations, *Geophysical Research Letters*, 44, 12–396, 2017.
- 500 Sietsma, J. and Dow, R. J. F.: Creating artificial neural networks that generalize, *Neural Networks*, 4, 67–79, [https://doi.org/10.1016/0893-6080\(91\)90033-2](https://doi.org/10.1016/0893-6080(91)90033-2), 1991.
- Simonyan, K. and Zisserman, A.: Very deep convolutional networks for large-scale image recognition, *arXiv preprint arXiv:1409.1556*, 2014.
- Tan, M. and Le, Q. V.: EfficientNet: Rethinking Model Scaling for Convolutional Neural Networks, *CoRR*, abs/1905.11946, <http://arxiv.org/abs/1905.11946>, 2019.
- 505 Wang, Y., Zhao, Y., Jiang, J., and Zhang, H.: A Novel GPU-Based Acceleration Algorithm for a Longwave Radiative Transfer Model, *Applied Sciences*, 10, 649, <https://doi.org/10.3390/app10020649>, <https://www.mdpi.com/2076-3417/10/2/649>, 2020.
- Wu, Y., Schuster, M., and et al.: Google’s Neural Machine Translation System: Bridging the Gap between Human and Machine Translation, *CoRR*, abs/1609.08144, 2016.
- Yuval, J. and O’Gorman, P. A.: Use of machine learning to improve simulations of climate, *arXiv:2001.03151 [physics]*, [http://arxiv.org/abs/](http://arxiv.org/abs/2001.03151)
- 510 2001.03151, 2020.