



The importance of management information and soil moisture representation for simulating tillage effects on N₂O emissions in LPJmL5.0-tillage

Femke Lutz^{1,2}, Stephen DelGrosso³, Stephen Ogle⁴, Stephen Williams⁴, Sara Minoli¹, Susanne Rolinski¹, Jens Heinke¹, Jetse J. Stoorvogel², and Christoph Müller¹

¹Potsdam Institute for Climate Impact Research (PIK), member of the Leibniz Association, P.O. Box 60 12 03, 14412 Potsdam, Germany

²Wageningen University, Soil Geography and Landscape Group, P.O. Box 47, 6700 AA Wageningen, the Netherlands

³USDA-ARS, Soil management and Sugar Beet Research Unit, 2150 Centre Ave. Bldg. D, Fort Collins, CO 80526, USA

⁴NREL, Colorado State University, Fort Collins, CO 80523, USA

Correspondence: Femke Lutz (femke.lutz@pik-potsdam.de)

Abstract. No-tillage is often suggested as a strategy to reduce greenhouse gas emissions. Modeling tillage effects on nitrous oxide (N₂O) emissions is challenging and subject to large uncertainties, as the processes producing the emissions are complex and strongly non-linear. Previous findings have shown deviations between the LPJmL5.0-tillage model and results from meta-analysis on global estimates of tillage effects on N₂O emissions. Here we tested LPJmL5.0-tillage at four different experimental sites across Europe and the USA, to verify whether deviations in N₂O emissions under different tillage regimes result from a lack of detailed information on agricultural management and/or the representation of soil water dynamics. Model results were compared to observational data and outputs from field-scale DayCent simulations. DayCent has been successfully applied for the simulation of N₂O emissions and provides a richer data base for comparison than non-continuous measurements at the experimental sites. We found that adding information on agricultural management improved the simulation of tillage effects on N₂O emissions in LPJmL. We also found that LPJmL overestimated N₂O emissions as well as the effects of no-tillage on N₂O emissions, whereas DayCent tended to underestimate the emissions of no-tillage treatments. LPJmL showed a general bias to over-estimate soil moisture content. Modifications of hydraulic properties in LPJmL in order to match properties assumed in DayCent, as well as of the parameters related to residue cover, improved the overall simulation of soil water as well as the N₂O emissions simulated under tillage and no-tillage separately. However, the effects of no-tillage (shifting from tillage to no-tillage) did not improve. Advancing the current state of information on agricultural management as well as improvements in soil moisture highlight the potential to improve LPJmL5.0-tillage and global estimates of tillage effects on N₂O emissions.

1 Introduction

Agricultural fields are often tilled to suppress weeds, incorporate crop residues, aerate the soil, prepare the seedbed and improve infiltration. The resulting changes in physical and chemical properties of the soil affect several biochemical processes, including the formation of greenhouse gases (GHG). Many field-scale models and experiments evaluated the effects of tillage and no-



tillage on GHG and soil organic carbon (SOC) (Álvaro-Fuentes et al., 2012; Del Grosso et al., 2009; Jin et al., 2017; Oorts et al., 2007). Nitrous oxide (N₂O) is a very strong GHG and predominantly emitted in agricultural production (Ciais et al., 2014; Smith, 2017). However, studies reported mixed results for the impacts of adapting no-tillage on N₂O emissions from croplands (Deng et al., 2016; Venterea et al., 2011). For instance, no-tillage was found to increase N₂O emissions (Mei et al., 2018; Van Kessel et al., 2013), decrease N₂O emissions (Deng et al., 2016; Plaza-Bonilla et al., 2018; Yoo et al., 2016) or having no significant effects (Alvarez et al., 2012; Boeckx et al., 2011) in comparison to conventional tillage systems.

Soils emit N₂O through a series of processes involving denitrification and nitrification. These processes are driven by microbial activity and strongly respond to soil properties such as moisture, temperature, oxygen, mineral N, and organic carbon (Mosquera et al., 2005; Snyder et al., 2009; Van Kessel et al., 2013). These soil properties are affected by tillage (Lutz et al., 2019a, c) and other management practices (e.g., fertilizer application and residue treatment) (Van Kessel et al., 2013). Due to the complexity of the system, the simulation of tillage effects on N₂O emissions is challenging and subject to large uncertainties.

Lutz et al. (2019a) extended a dynamic global vegetation, hydrology and crop model to explicitly account for the effects of tillage in the simulations of biogeochemical cycles, hydrology and crop yields. This enables simulations of the effects of tillage on crop productivity, the water, carbon and nitrogen cycles, including N₂O emissions at the global scale. However, they found that simulated N₂O emissions from no-tillage exceeded values in most of the climate zones reported in meta-analyses. These deviations between observations and simulations of tillage effects on N₂O emissions can have several different causes, including missing processes and lack of process understanding. Also the parameterization of implemented processes as well as detailed information on management aspects that are explicitly addressed in the model can lead to model deficiencies that could cause the mismatch between observations and simulations.

For example, as detailed information about agricultural management practices is lacking for global-scale applications, assumptions on agricultural management are necessary in these global simulations about e.g., the type, amount and timing of fertilizer applications. Detailed information on fertilization can typically be dealt with in field-scale modeling experiments, whereas at the global scale, there is only general information on fertilization (e.g. Mueller et al., 2012; Potter et al., 2010) which is characterized by gaps and uncertainties (Erb et al., 2017). These generalizations may be a significant contributor to the overall uncertainty for agricultural impact assessments. For instance, Folberth et al. (2019) found that differences in management assumptions (about e.g., growing season, and fertilization) resulted in substantial differences in modeled crop yields using the same crop model.

Second, the formation of N₂O in soils is very sensitive to soil moisture (Butterbach-Bahl et al., 2013). How the effect of tillage on soil moisture is simulated is thus another source of uncertainty that could explain the inaccuracy in modeling tillage effects on N₂O emissions.

In this study, we test the importance of management information as well as the representation of soil water dynamics for the ability to simulate N₂O emissions under different tillage regimes with LPJmL5.0-tillage (Lutz et al., 2019a), for four different experimental sites across Europe and the USA. Simulation results are compared to measurements of N₂O emissions from experimental studies under tillage and no-tillage in different simulation experiments, varying from using observed site-



specific information to using the default assumptions usually applied in the global-scale simulations. Because of the importance of soil moisture for N₂O emissions, we test the accuracy of the simulated soil moisture dynamics and its effects on N₂O emissions against observations. As simulating tillage effects on N₂O emissions is generally challenging, we use the site-specific model DayCent (Del Grosso et al., 2009; Parton et al., 1996), which has previously been applied at the study sites as a benchmark and to provide more detailed information on soil hydrology than the sparse observations. DayCent is a well-established model that has been used for questions related to agricultural impact assessments at various scales (e.g. Begum et al., 2019; Del Grosso et al., 2009; Del Grosso et al., 2002; Gryze et al., 2010). DayCent can be used as a benchmark for which the underlying mechanisms can be analyzed and used for improvements of LPJmL5.0-tillage, even though the performance of DayCent has to be compared to observations first.

65 2 Material and methods

2.1 Overview

In Lutz et al. (2019a), model results deviated from meta-analyses when comparing simulated tillage effects on N₂O emissions. First, we tested whether the deviations are due to a lack of detailed management information. Four experimental sites for which detailed information on management are available were identified. On those sites, LPJmL5.0-tillage was run using management assumptions usually used in a global simulation experiment (LPJmL.G.Orig). To find out if LPJmL5.0-tillage performed better with detailed information on management, we also applied LPJmL5.0-tillage using detailed site-specific management information to derive inputs (LPJmL.D.Orig).

The site-specific DayCent model was used as benchmark to analyze the underlying mechanisms of the N₂O producing processes. For all the simulations of DayCent, detailed information of management was used. Except for the experimental site in Boigneville, DayCent has been used and calibrated for field-scale assessments at the chosen sites (i.e. Campbell et al., 2014; Del Grosso et al., 2009; Yang et al., 2017). Therefore, we expect it to perform better on simulating the effects of tillage on N₂O emissions than LPJmL. We also expect to learn from the underlying mechanisms simulated by DayCent and to use this information for improving process representation and parameterization in LPJmL. All model versions considered here require similar inputs (soil properties, vegetation type, land management information, latitude, daily precipitation, and daily air temperature (minimum and maximum)).

2.2 LPJmL5.0-tillage

LPJmL5.0-tillage is a dynamic global vegetation, hydrology and crop model that simulates nitrogen (N), carbon (C) and water dynamics in natural and agricultural ecosystems. Soils are represented by five hydrologically active layers, with different layer thicknesses.

LPJmL5.0-tillage (in the following referred to as LPJmL) uses three litter pools; representing surface litter, incorporated litter and below-ground litter as well as two soil organic matter (SOM) pools per soil layer, which are characterized by fast



and slow decomposition rates, respectively, and by separate C and N components for each pool. The surface litter pool consists of crop residues which are not removed at harvest or incorporated into the first soil layer through tillage. Residue cover is calculated from the surface litter following Gregory (1982). This residue cover intercepts some rainfall, promotes infiltration
90 into the soil, and limits soil evaporation. Moreover, the presence of a residue cover insulates the soil from air temperature fluctuations. The effects of residue cover on soil water dynamics and soil temperature fluctuations are thoroughly described in Lutz et al. (2019a).

Surface litter decomposes and is incorporated through bioturbation and tillage, forming the incorporated litter pool in the first layer. The below ground litter pool includes crop roots that remain in the soil after harvest. All pools are subjected
95 to decomposition, which is driven by the moisture content and temperature of the soil (for the incorporated litter and below-ground litter pool), or of the moisture content and temperature of the surface litter (surface litter pool). A fixed fraction of the decomposed litter is mineralized and emitted as CO₂, whereas the remaining C is transferred to the SOM pool, where it is then subject to soil C decomposition (see also Von Bloh et al., 2018). The mineralized N is added to the NH₄⁺ pool which is subject to further transformations into other forms of nitrogen (Von Bloh et al., 2018).

100 Nitrification and denitrification are simulated throughout the entire soil profile and are dependent on the water-filled pore space (WFPS), soil temperature, NH₄⁺, pH, SOC (denitrification) and NO₃⁻. The N₂O emissions from denitrification increases exponentially when the WFPS reaches a threshold value of ≥ 90%, as denitrification occurs only in oxygen deficit conditions (see also Krysanova and Wechsung, 2000).

In addition to tillage effects on residues (i.e. incorporating residues into the soil), tillage affects the hydraulic properties
105 of the soil by decreasing the bulk density. Soil hydraulic parameters are calculated through a pedotransfer function (PTF) from Saxton and Rawls (2006) which uses soil texture, SOM, and bulk density changes to calculate field capacity (FC), wilting point (WP), saturation (WSAT) and the saturated hydraulic conductivity (Ksat). The hydraulic parameters determine the water holding capacity- and the water dynamics of the soil. For instance, soil water above WSAT runs off as lateral runoff, while remaining soil water above FC percolates to the next soil layer and generates lateral subsurface runoff or vertical seepage from
110 the soil column.

A full overview of the tillage implementation into LPJmL5.0 as well as affected soil properties and processes can be found in Lutz et al. (2019a), the nitrogen implementation is described by Von Bloh et al. (2018) and a comprehensive description of the LPJmL model is provided by Schaphoff et al. (2018) and Schlüter et al. (2018).

2.3 DayCent

115 The DayCent ecosystem model simulates crop growth, soil water, C and nutrient dynamics (N, P) in natural and agricultural ecosystems (Del Grosso et al., 2009; Parton et al., 1998). The soil is represented by user-specified layers which are hydrologically active. DayCent has two litter pools, representing surface-litter and below-ground litter and three SOM pools (active, slow and passive) characterized by different decomposition rates.

The active and the slow organic matter pools have surface as well as soil components while the passive pool has only
120 a soil component. The litter pools are partitioned into structural and metabolic pools as a function of the lignin to N ratio in



the residue, which are subject to decomposition. Decomposition products of litter supply the SOM pools (surface active, soil active, surface slow and soil slow) and are partitioned among pools based on lignin content. Decomposition of litter and soil organic matter and nutrient mineralization are a function of substrate availability, substrate quality (lignin content, C:N ratio), soil moisture, soil temperature and tillage intensity. N-mineralization, N-fertilization and N-fixation supply the N-pools. NO_3^- is distributed throughout the soil profile, whereas NH_4^+ is confined to the top 10 cm. NO_3^- and NH_4^+ can then be taken up by plants, leached to lower layers (NO_3^- only) or transformed to N gas emissions (e.g. N_2O) through nitrification or denitrification (Del Grosso et al., 2000; Parton et al., 2001). N_2O emissions from nitrification are calculated as a function of soil NH_4^+ concentration, temperature, pH, texture and soil moisture. N_2O from denitrification is calculated as a function of soil NO_3^- concentration, soil moisture, texture and heterotrophic CO_2 respiration rate. N_2O emissions from denitrification increases exponentially when the WFPS exceeds the texture related threshold value and levels off as the soil approaches saturation. The model can simulate different types of tillage (i.e. plowing, tandem disk and field cultivator). Depending on the type of tillage, the decomposition of litter and SOM (active and slow) pools are increased by a specific factor for a period of one month, and a fraction of above-ground residues is transferred to surface litter and top soil layer. Tillage also impacts soil temperature and water dynamics indirectly because the model assumes that precipitation intercepted by surface litter and living biomass evaporates before entering soil. The presence of surface litter insulates the soil from air temperature fluctuations.

If site level measurements of soil hydraulic properties required for DayCent are not available, they are calculated through the PTF from Saxton et al. (1986) and are static throughout the simulations. The PTF uses soil texture to calculate FC, WP, bulk density and Ksat. The soil water model simulates unsaturated water flow using Darcy's equation, runoff, snow dynamics, and the effect of soil freezing on saturated water flow (Pannkuk et al., 1998). DayCent has been shown to reliably model soil water content, N mineralization and N_2O emission rates from different soil types and management practices (Kelly et al., 2000; Parton et al., 2001). Del Grosso et al. (2002) provides an extensive overview of validate results for DayCent.

2.4 Experimental sites

Four experimental sites were selected in which the effects of tillage and no-tillage on N_2O emissions were studied (Table 1 and Table 2). The sites were selected based on the availability of observational data and treatment combination of tillage and no-tillage.

The first study site is located at the Agricultural Research Development and Education Center (ARDEC) near Fort Collins, CO (40° 39'6" N, 104° 59'57" W; 1555 m asl). It was initiated in 1999 on a clay loam soil (fine-loamy, mixed, mesic Aridic Haplustalfs), that was continuously cropped with maize (*Zea mays L.*) for six years. Shortly before sowing, fertilizers (67 kg N ha⁻¹) were applied. The fields were sprinkler irrigated during the growing season. In the tillage treatment, fields were tilled shortly before sowing, and with harvest, followed by tandem disking and then moldboard plowing to a depth of 25 to 30 cm. N_2O emissions were measured three times per week during the growing season (2002-2006) with closed chambers. Soil moisture was measured two to three times per month during the growing season from 2003 to 2006. Soil organic carbon (SOC) was measured once in October 2005. A detailed description of the experimental site can be found in Halvorson et al. (2006).



The second study site is located at the University of Nebraska-Lincoln Agricultural Research and development Center,
155 Ithaca, NE (41° 9'43.3"N, 96° 24'41.4" W; 349 m asl). The experiment was established in 2002 on a silt loam soil that was
previously cropped with rain fed maize, soybean (*Glycine max (L.) Merr.*), oat (*Avena sativa L.*) and alfalfa (*Medicago sativa*
L.). Maize was grown continuously on the field after 2000. During the experiment, N fertilizers were injected to a depth of
10-15 cm, once during the growing season at various rates and compositions (Table 1). The soil in tillage treatments was tilled
before sowing and at harvest to a depth of 15-20 cm. The field was irrigated with varying irrigation amounts. N₂O emissions
160 were measured from April 2011 through May 2016 monthly during the growing season using closed chambers. Soil moisture
was measured at varying intervals from one to five times per month between 2011 and 2015. SOC was measured in May 2001,
November 2010, and November 2014 for different depths (0-0.15, 0.15-0.30, 0.30-0.60, 0.60-0.90, 0.90-1.20 and 1.20-1.50
m). More information regarding the experimental study site is provided by Jin et al. (2017).

The third study site is the W.K. Kellogg Biological Station Long-Term Ecological Research (KBS LTER) experiment
165 located in Southwest Michigan (42° 24' N, 85° 24' W, 288 m asl) on loam soils (Typic Hapludalfs). The experiment was
established in 1988 on an agricultural field that had been tilled for at least 100 years before the experiment. The crop rotation
before 1995 consisted of maize followed by soybean. In 1995, wheat (*Triticum aestivum L.*) was planted after soybean, which
resulted in a maize-soybean-wheat rotation. After the harvest of wheat, the fields stayed bare until the fields were cropped with
maize again. This sequence was followed during the time span analyzed here (1989-2010). Different quantities of N-fertilizers
170 were applied at sowing and/or during the growing season for maize, during the growing season for wheat, and soybean did
not receive fertilizers (Table 1). The tillage treatment was tilled each year with sowing, then during the growing season and at
harvest, to a depth of 20 cm. The fields were not irrigated during the experiment. N₂O emissions were measured once or twice a
month from June 1991 to October 2016 using closed chambers. Soil moisture was measured once per month during the growing
season from 1989 until 2017. SOC was measured annually since 1989 at multiple sampling depths. More information regarding
175 the experimental study site is provided by Grandy et al. (2006) and on the KBS LTER website (<http://lter.kbs.msu.edu>, accessed
November 2018).

The last study site is located in Boigneville, France (48° 33'N, 2° 33'E, altitude unknown) on a silt loam soil (Haplic
Luvisoil) (FAO, 1998). The experiment started in 1970 that had been tilled to 30 cm depth annually. During the experiment, the
site was cropped with a maize-wheat rotation, with maize being sown in April, harvested in October and directly followed by
180 tillage (20 cm for tillage treatments) and sowing of wheat. After harvest of wheat in April, the soil was left bare and was tilled
(20 cm) in November, and left fallow until planting maize in the next growing season. This sequence was followed during the
time span analyzed here (2003-2004). During the experiment, the maize received N-fertilizers in May and wheat in February
and April (Table 1). The fields were irrigated between the end of June and July. N₂O emissions were measured on average
every three weeks using closed chambers. Soil moisture was not measured. Soil organic carbon was measured twice in 2003
185 and once in 2004 at various depths. More information regarding the study site can be found in (Oorts et al., 2007).



Table 1. Overview of experimental sites selected for the study.

Location	Years of Experiment	Soil texture	Land use	Observations	Reference
Boigneville, France	1971–2004	Silt loam	Maize-Wheat	Timespan (average freq. in growing season) N ₂ O: 2003-2004 (three weeks)	Oorts et al. (2007)
Fort Collins, Colorado	1999–2006	Clay loam	Maize	N ₂ O: 2003-2006 (three days)	Halvorson et al. (2006)
Hickory Corners, Michigan	1989–2010	Loam	Maize-Wheat-Soybean	N ₂ O: 1991-2016 (2 weeks)	Grandy et al. (2006)
Mead, Nebraska	2001–2015	Silt loam	Maize	N ₂ O: April 2011-May 2016 (2 weeks)	Jin et al. (2017)



Table 2. Overview of observed input data and LPJmL input data

Site	Observed data						LPJmL data									
	Fertilization		Tillage		Growing Season		Soil Pools		Fertilization		Tillage		Growing Season		Soil Pools	
	Amount (g N m ⁻²)	Day of year	Day of year	Day of year	Day of year	Harvest Day of year	Soil C g C **	Soil N g N **	Amount (g N m ⁻²)	Day of year	Day of year	Day of year	Day of year	Harvest Day of year	Soil C g C **	Soil N g N **
France	15.8	131	301	107	282	282	4553.3	450.3	10.4	122	122	122	297	297	3827.8	297.0
Colorado	6.7	118	30	118	288	288	6092.0	460.4	8	123	123	123	249	249	6267.7	335.7
			109						8	188	270					
			119													
			330													
Michigan	3.3	135	136	128	293	293	9834.2	1148.2	7.155	125	125	125	238	238	6188.4	760.4
	12.3	179	139						7.155	175	251					
Nebraska	20.2	165	114	123	270	270	1762.2	1529.14	8.4	124	124	124	234	234	6267.7	335.7
									8.4	177	131					

* The data are for the years where maize is grown, and vary between years.

** Size of pools are given for soil depth: France, Colorado, Michigan and Nebraska are from 0 to 0.2, 0.2, 1.0 and 1.5 respectively.



2.5 Management information

2.5.1 LPJmL standard setup using global input data

In the LPJmL.G.Orig scenario, all management information as well as soil C and N-pools were used as within the default global simulation of LPJmL (Table 3). The amount of mineral and organic fertilizers was provided by the global gridded crop model intercomparison (Elliott et al., 2015) of the Agricultural Model Intercomparison and Improvement Project (AgMIP, Rosenzweig et al., 2013). It is based on global, gridded data sets for each crop (Mueller et al., 2012; Potter et al., 2010). Fertilizer is assumed to consist of 50% NO_3^- and 50% NH_4^+ . If fertilizer input is low ($\leq 5.0 \text{ gN m}^{-2}$), all is applied at sowing. Otherwise, only half of the fertilizer is applied at sowing and the remainder is applied when the phenological stage fraction (unitless) of the crop reaches 0.4 (Von Bloh et al., 2018). Irrigation events occur when the fractional soil moisture of the water holding capacity (unitless) is below an irrigation threshold value of 0.7 for maize (Jägermeyr et al., 2015).

In the experiments with tillage, tillage occurs twice a year; once at sowing and once at the day of harvest. Sowing dates are calculated internally following Waha et al. (2012). Thereby, the sowing dates are calculated based on a set of rules depending on crop specific thresholds and climate. Here, the sowing date depends on a crop-specific temperature threshold (i.e. $14 \text{ }^\circ\text{C}$ for maize; Waha et al., 2012).

The size of the C and N pools are calculated internally during the spinup (5000 years) of the natural vegetation and land-use history. The land-use history is simulated as with DayCent, in order to establish a comparable starting point when the simulations for the experiments are conducted. Thereby, the spin-up is followed by a simulation of historical land-use change to account for effects on the pools based on the best available information of land management.

2.5.2 LPJmL detailed setup using observed input data

Site-specific observed information for all management inputs as well as soil C and N pools were prescribed for simulation LPJmL.D.Orig (Table 3). For practical reasons, irrigation water was added to precipitation to enable the specification of the amount and the timing of irrigation events. This mimics a sprinkler irrigation technique as part of the irrigation water is intercepted by the canopy. As the current implementation of soil layers and tillage in LPJmL does not allow for distinguishing more detailed tillage types other than conventional tillage and no tillage, we ignored tillage activities that were less intensive (e.g. “shredding”). In order to specify the growing season, phenological heat unit requirements and base temperatures were parameterized so that the simulated harvest dates were matching the reported harvest dates.

The soil C and organic N pools from the simulations were scaled to the observed values. This was done twice, once at the introduction of land-use during spin-up and once at the start of the treatment of the experimental site. If observations were not available for the start of the experiment, the first available observation was taken, assuming that pool sizes remained stable over that time period. The pools (P) at each site were scaled as in equation 1:

$$P_{(cor,l)} = P_{(sim,l)} * \frac{Total_{(obs)}}{Total_{(sim)}} \quad (1)$$



Where $P_{(cor)}$ are the scaled carbon or nitrogen content of the soil pools (g C or N m^{-2}) in layer l of the experimental site and $P_{(sim)}$, the simulated amounts of C or N contained in the soil and litter pools of the different layers l of the experimental site. $Total_{(obs)}$ and $Total_{(sim)}$, are the total of C or N contained in the soil and litter pools summed over the different layers (l) for which observational data of soil organic C and N were available (in g C or g N m^{-2} , respectively) of the experimental site. The differences between simulated and observed input data are depicted in Table 3.

2.6 LPJmL experimental simulations

Agricultural management consists of several practices. To analyze the importance of individual management aspects, we conducted a set of simulations as in LPJmL.D.Orig but ignored one site-specific management practice and replace it with the global assumption as in LPJmL.G.Orig (Table 3). As an example: LPJmL.D.Orig-F, refers to the simulation where all management information are as in the LPJmL.D.Orig, except for the fertilizer information. Instead, the amount, timing and type of fertilizers were used as in LPJmL.G.Orig. Other experimental simulations refer to: LPJmL.D.Orig-I, LPJmL.D.Orig-GS, LPJmL.D.Orig-PS and LPJmL.D.Orig-T, that use the management information as in LPJmL.D.Orig, except for irrigation (I; timing and amount), growing season (GS; sowing- and harvest days), C and N pool sizes (PS) and the timing of tillage (T) respectively. The naming of the simulation consists of three parts: 1) model used (LPJmL), 2) the experiment conducted (e.g. I, GS or PS) and 3) whether it includes modifications (“Mod”; see 2.7) or not (“Orig”).

2.7 Model modifications

Lutz et al. (2019a) found that LPJmL overestimates N_2O emissions. Because of the importance of soil moisture for N_2O emissions, we tested if modifying the simulation of soil moisture can contribute to improving the simulation of N_2O emission. We modified the model with respect to the treatment of the residue cover of the soil in no-tillage systems and with respect to changing the soil parameterization.

As the soil covered by residues under no-tillage practices in LPJmL simulations is very high and thus leads to high soil moisture levels throughout the year (as soil evaporation is reduced and infiltration is enhanced), we tested modifications of the relevant functions for this aspect. To this end, we tested modifications of the parameters that translate litter amounts into soil cover (Gregory, 1982) and those that determine how long the soil is covered with residues. Rather than changing well-established functions on litter decomposition (Schlüter et al., 2018), we modified the parameter on bioturbation that was introduced by Lutz et al. (2019a) and tested its effects on the reduction of the residue cover of the soil.

Lutz et al. (2019a) used an average value of $0.006 (\text{m}^2 \text{g}^{-1})$ (falsely described as 0.004 in their publication, but used so in the code: <https://doi.org/10.5281/zenodo.2652136>) to translate litter biomass into a fraction of soil being covered with residues, which was applied to all litter, neglecting variations in surface litter for different materials. The bioturbation rate was increased from $0.19\% \text{ day}^{-1}$ to $0.63\% \text{ day}^{-1}$ to account for the surface litter being transferred to the incorporated litter pool per day (equivalent to an annual bioturbation rate of 90%, versus 50% as assumed previously).

High N_2O emissions can also result from biases in the parameterization of hydraulic properties. For example, small differences between FC and WSAT lead to frequent triggering of denitrification. To study the role of soil moisture for causing



Table 3. Overview of management data used in LPJmL.D.Orig, LPJmL.G.Orig, Daycent and experimental runs.

Management information	Experimental runs							
	LPJmL.D.	LPJmL.G.	LPJmL.D.Orig					Daycent
	Orig	Orig	-F*	-I*	-GS*	-PS*	-T*	
Fertilizer (amount, type, timing)	Observed data	LPJmL Data	LPJmL data	Observed data	Observed data	Observed data	Observed data	Observed data
Irrigation (amount, timing)	Observed data	LPJmL data	Observed data	LPJmL data	Observed data	Observed data	Observed data	Observed data
Growing season	Observed data	LPJmL data	Observed data	Observed data	LPJmL data	Observed data	Observed data	Observed data
Tillage	Observed data	LPJmL data	Observed data	Observed data	Observed data	Observed data	LPJmL data	Observed data
Soil C and N pool	Observed data	LPJmL data	Observed data	Observed data	Observed data	LPJmL data	Observed data	Observed data

* Experimental runs; all management information is as in the Detail settings, except for one scenario. For example, LPJmL.D-F* excludes fertilization information. The other settings exclude information on irrigation (LPJmL.D-I*), growing season (LPJmL.D-GS*), Pool N and C sizes (LPJmL.D-PS*), and tillage (LPJmL.D-T*)



250 deviations in tillage effects on N₂O emissions, we analyzed if the parameterization of the hydraulic properties causes the overestimation in soil moisture. As observational data on the hydraulic properties are lacking, we here compared the hydraulic properties in relation to soil moisture from DayCent.

2.8 Analyses

2.8.1 N₂O emissions

255 As N₂O emissions are characterized by a high temporal variability, we analyzed two different aggregation levels: annual averages of N₂O emissions and emissions of individual days within the year. We analyzed each tillage type (*tt*, i.e. conventional tillage and no-tillage) separately (N₂O_{*tt*}, equation 2) and differences between the two for both aggregation levels (N₂O_{*diff,year*}; equation 3 and N₂O_{*diff,day*}; equation 4).

$$N_2O_{tt} = \frac{\sum_{day=1}^n N_2O_{day,tt}}{n_{tt}} \quad (2)$$

260 N₂O_{*tt*} is the annual average of simulated and observed N₂O emissions (in g N ha⁻¹ d⁻¹) of *tt* (tillage type: conventional tillage (*till*) or no-tillage (*notill*)), and *n_{tt}* is the number of days with N₂O emissions simulated or observed in the year of *tt*. Thereby, *n_{tt}* equals all 365 days in the simulations and for the observations *n_{tt}* < 365 as observations are not available for every day in the year. We thus assumed that the scarcer observations still represent the full year's dynamics.

The differences in N₂O emissions on annual average (N₂O_{*diff,year*}) were calculated as in equation 3:

$$265 \quad N_2O_{diff,year} = \frac{\sum_{day=1}^n N_2O_{day,notill}}{n_{notill}} - \frac{\sum_{day=1}^n N_2O_{day,till}}{n_{till}} \quad (3)$$

Where N₂O_{*day,notill*} and N₂O_{*day,till*} are daily N₂O emissions in g N ha⁻¹ d⁻¹ for all the days in the year and *n_{notill}* and *n_{till}* the number of days with N₂O emissions simulated or observed in the year for no-tillage and tillage, respectively.

The differences in N₂O emissions for individual days were calculated as in equation 4:

$$N_2O_{diff,day} = N_2O_{notill} - N_2O_{till} \quad (4)$$

270 Where N₂O_{*notill*} and N₂O_{*till*} are daily emissions in all years.

The relative difference (RD in %) of no-tillage to conventional tillage was calculated as in equation 5:

$$RD = \left(\frac{\sum_{day=1}^n N_2O_{notill}}{\sum_{day=1}^n N_2O_{till}} \right) * 100(\%) \quad (5)$$

Where N₂O_{*notill*} and N₂O_{*till*} are daily N₂O emissions in g N ha⁻¹ d⁻¹ for all the days in the year and *n* is the number of days with N₂O emissions simulated or observed.



275 2.8.2 Soil moisture

For the analyses of soil moisture, we focused on the first 0.2 m of the soil, which is the tillage-affected layer. We analyzed the experimental site in Nebraska as this site had the most observations of soil moisture compared to the other experimental sites. As N₂O emissions are regulated by the WFPS in both LPJmL and DayCent, we normalized the soil moisture content and hydraulic properties to porosity (W_{SAT} in mm). The WFPS (fraction) is calculated as in equation 6:

$$280 \quad WFPS = \frac{W}{W_{SAT}} \quad (6)$$

where W is the volumetric soil water content (mm). The $WFPC_{FC}$ (fraction) and $WFPC_{WP}$ (fraction) are the field capacity and wilting point values normalized to WFPS as in equations 7 and 8:

$$WFPC_{FC} = \frac{W_{FC}}{W_{SAT}} \quad (7)$$

$$WFPC_{WP} = \frac{W_{WP}}{W_{SAT}} \quad (8)$$

285 The W_{FC} and W_{WP} are the water content at field capacity and wilting point, respectively.

2.8.3 Evaluation metrics

To quantify the performance of simulated N₂O emissions, we conducted an analyses of coincidence (equation 9) and an analysis of association (equation 10), following Smith and Smith (2007). Therefore, we calculated the deviation between simulated and observed values were by the root mean squared deviation (RMSD in g N ha⁻¹ d⁻¹) for the different sites as in equation 9:

$$290 \quad RMSD = \sqrt{\frac{\sum_{i=1}^n (O_i - S_i)^2}{n}} \quad (9)$$

O_i is the average observed N₂O emission (in g N ha⁻¹ d⁻¹) of year i and S_i the average simulated value of N₂O emission (in g N ha⁻¹ d⁻¹) of year i and n is the total number of valid value pairs for comparison.

To describe how well the dynamics in the observations were captured in the simulations, we calculated the degree of association (r) as in equation 10:

$$295 \quad r = \frac{\sum_{i=1}^n (O_i - \bar{O})(S_i - \bar{S})}{\sqrt{\sum_{i=1}^n (O_i - \bar{O})^2 \sum_{i=1}^n (S_i - \bar{S})^2}} \quad (10)$$

Where \bar{O} and \bar{S} are the average observed and average simulated value respectively over all years (in g N ha⁻¹ d⁻¹). The significance of r corresponds to the tests, null hypothesis: $r=0$.



The mean bias (MB in fraction) was calculated as in equation 11:

$$MB = \frac{\overline{O}}{\overline{S}} \quad (11)$$

300 For soil moisture, the $RMSD$ and r were calculated as well. However, there we focused on one site and calculated the average $RMSD$ and r over all the years, as not much variation in soil moisture is expected between the years.

3 Results and discussion

3.1 Importance of management information

3.1.1 Tillage effects on N_2O emissions

305 Annual averages

The N_2O emissions were overestimated in the LPJmL.G.Orig experiment when analyzing yearly averages of the different sites (Fig. 1 A). This effect was stronger for simulated emissions under no-tillage ($RMSD=36.2 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=-0.07$) than under tillage ($RMSD=23.6 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=-0.31$). DayCent was closer to the observed values for both tillage ($RMSD=7.60 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=0.67$) and no-tillage ($RMSD=4.61 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=0.66$). For the full statistical analyses, we refer to Table A1 in
310 the Appendix.

Using detailed site-specific management information in LPJmL (LPJmL.D.Orig) improved the correlation between the observed and simulated values (Fig. 1 B). The simulated N_2O emissions under no-tillage deviated more from the observed values ($RMSD=38.9 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=0.36$), as the N_2O emissions were still overestimated. This held for the simulated N_2O emissions resulting under conventional tillage as well ($RMSD=31.7 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=0.34$).

315 When analyzing the effect of tillage (difference between no-tillage and tillage), LPJmL.G.Orig showed an increase in emissions with no-tillage (Fig. 2 A), and LPJmL.D.Orig showed both an increase and decrease with no-tillage (Fig. 2 B). On average, no-tillage increased N_2O emissions by 59.5% in LPJmL.G.Orig, and 22.4% in LPJmL.D.Orig across all sites and years. In observations, no-tillage decreased N_2O emissions on average by 16.0% and DayCent shows a reduction of 24.3%. However, observations across the different sites showed, that no-tillage can have very different effects on N_2O emissions.
320 In Boigneville and Michigan, N_2O emissions increased under no-tillage (49.3% and 15.7% respectively), whereas it decreased in Colorado (by 9.01%) and Nebraska (by 29.2%). LPJmL.D.Orig reproduced the observed differences in tillage better ($RMSD=12.0 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=0.48$) than LPJmL.G.Orig ($RMSD=18.0 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=-0.16$), see also Fig. 2. Yet, both versions mainly projected an increase in N_2O emissions from no-tillage practices. DayCent results were closer to the observed values, but slightly underestimated the effects of no-tillage on N_2O emissions ($RMSD=4.96 \text{ g N ha}^{-1} \text{ d}^{-1}$, $r=0.34$).

325 Daily emissions

The simulations with different management information showed that these are relevant for the simulated tillage effects on N_2O emissions on individual days (Fig. 3). On average, more accurate information on management improved the simulations of differences between conventional and no-tillage systems in LPJmL except for the site in Colorado. However, there was




Figure 1. Comparison of observed and simulated yearly averages of N_2O emissions by tillage type and models LPJmL.G.Orig (A), LPJmL.D.Orig (B) and DayCent. Data refer to all four sites and years of the experiments. Each point represents the average of all measured daily values within one year and tillage treatment. Tillage types are indicated by different colors.

no clear pattern between the different experimental runs of LPJmL (Fig. A1 in Appendix A). None of the simulations with
330 partial usage of detailed management information (Table 3) performed clearly better or worse between the LPJmL simulations.
There were only small differences in the distribution of no-tillage effects on N_2O emissions as well as between the averages.
The observations showed that no-tillage both increased (Boigneville, Michigan) and decreased N_2O emissions (Colorado,
Nebraska) on average, as well as on the individual days. The negative effects were reproduced by DayCent in Colorado and
Nebraska. The positive and negative effects were reproduced by LPJmL.D.Orig as well, except in Michigan. LPJmL.G.Orig
335 however, only reproduced the increase in N_2O emissions in Michigan (Fig. 3).

In Colorado, observations showed a decrease in N_2O emissions under no-tillage compared to conventional tillage. In con-
trast, LPJmL.D.Orig and LPJmL.G.Orig showed an increase in emissions with no-tillage, whereas the observed decrease was
well captured by DayCent. In Boigneville, the increase in N_2O emissions under no-tillage was well captured by LPJmL.D.Orig.
DayCent and LPJmL.G.Orig did not capture the increase in N_2O emissions with no-tillage. In Nebraska, LPJmL.D.Orig and
340 DayCent agreed with observations that no-tillage decreases N_2O emission. In Michigan, no-tillage resulted mainly in an in-
crease in emissions in LPJmL, which can also be found in the observations but not in DayCent simulations.

For all sites, LPJmL showed a high variability in N_2O emissions between days (Fig. 3 and Table A1 in Appendix A).
The interquartile ranges from LPJmL simulations were often much wider compared to observations and DayCent simulations.
Hence, the variability of no-tillage effects on daily N_2O emissions was overestimated. DayCent tended to underestimate the
345 variability of N_2O emissions between days (Table A1 in Appendix A).

In LPJmL, the N_2O emissions from no-tillage were entirely caused by changes in denitrification, whereas no-tillage
mainly caused decreases on N_2O emissions from nitrification (Fig. A2 in Appendix A). This can be explained by higher soil




Figure 2. Comparison of observed and simulated effects after converting to no-tillage. The data refer to yearly averages of N_2O emissions and models LPJmL.G.Orig (A), LPJmL.D.Orig (B) and DayCent, of all four sites and years of the experiments.

moisture levels with no-tillage in LPJmL. In general, higher soil moisture levels trigger N_2O emissions from denitrification (anaerobic process), whereas nitrification is decreased (aerobic process). In DayCent, no-tillage mainly decreased N_2O emissions emitted from nitrification and had little effects on denitrification.

3.2 Soil hydrology and model modifications

3.2.1 Soil hydrology

The soil moisture (WFPS) simulated by LPJmL.D.Orig for no-tillage in Nebraska, is high compared to the observed values (RMSD= 0.24 (unitless), $r=0.28$) (Fig. 4). DayCent was closer to the observed values for no-tillage (RMSD= 0.10 (unitless), $r=0.50$) and tillage (RMSD= 0.11 (unitless), $r=0.49$). After modifying the parameters for surface litter and the hydraulic properties, the simulated soil moisture in the experiment LPJmL.D.Mod was closer to the observed values and simulation results from DayCent (Fig. 4). These combined effects showed the best performance for both tillage (RMSD=0.12 (unitless), $r=0.33$) and no-tillage (RMSD=0.14 (unitless), $r=0.48$), compared to implementing the modifications separately (Table 4). The dynamics in soil moisture simulated in the experiment LPJmL.D.Mod better reflected the dynamics simulated by DayCent.

For instance, after October, a decrease in soil moisture was simulated by DayCent (and measured) which was previously not captured by LPJmL.D.Orig. In LPJmL.D.Orig, soil moisture was mostly stationary around FC, which in LPJmL.D.Mod was only the case from April to June.

Although the simulation of soil moisture was improved with the modified settings, LPJmL simulations still overestimated soil moisture in comparison to observations.




Figure 3. Effects of no-tillage on N_2O emissions on individual days (and on average), including the original LPJmL settings, the observations and simulated values by DayCent.

Table 4. Performance of Daycent, and LPJmL compared to soil water observations in Nebraska. The results are shown for both conventional tillage and no-tillage

	<i>RMSE</i>		<i>r</i>	
	Conv. tillage	No tillage	Conv. tillage	No tillage
LPJmL.D.Orig	0.21	0.24	0.10	0.28
Bioturbation	0.20	0.22	0.19	0.40
Parameter residue cover	0.19	0.24	0.20	0.32
Hydraulic properties Daycent	0.15	0.18	0.07	0.23
LPJmL.D.Mod	0.12	0.14	0.33	0.48
Daycent	0.11	0.10	0.49	0.50




Figure 4. Observed and simulated soil moisture of no-tillage in the top soil (0-20 cm) in Nebraska.

365 3.2.2 Tillage effects on N₂O emissions after modifications

Yearly averages

The modifications of the parameters for surface litter and the hydraulic properties improved the yearly tillage and no-tillage effects on N₂O emissions across all the different sites (Fig. 5). The emissions under no-tillage (RSMD=18.1 g N ha⁻¹ d⁻¹, r=0.60) and under tillage (RSMD=16.3 g N ha⁻¹ d⁻¹, r=0.38) were much closer to the observed values than with the original hydrologic parameterization. Although the modifications improved the simulation of tillage and no-tillage, LPJmL.D.Mod still overestimated the changes in emissions when switching from conventional tillage to no-tillage systems (Fig. 5, Table A1). The modifications did not improve the simulation of N₂O emissions after shifting to no-tillage (Fig. 6). Although the deviations of the absolute differences between tillage systems decreased, the correlation with observations was less well captured (RSMD=7.35 g N ha⁻¹ d⁻¹, r=-0.04), negating the improvements achieved through the consideration of detailed management information (LPJmL.G.Orig vs. LPJmL.D.Orig). The conversion to no-tillage systems increased N₂O emissions by 13.0% in LPJmL.D.Mod. The increase in N₂O emissions after shifting to no-tillage in the modified simulations was found across all sites in LPJmL.D.Mod, whereas DayCent showed decreases in N₂O emissions across all sites at the yearly aggregation (Fig. 6). However, the observations showed both increases and decreases in N₂O emissions after shifting to no-tillage for all sites at the yearly aggregation.




Figure 5. Comparison of observed and simulated yearly averages of N_2O emissions by tillage type and models DayCent, LPJmL.D.Mod and LPJmL.D.Orig (in grey). Data refer to all four sites and years of the experiments. Each point represents the average of all measured daily values within one year and tillage treatment. Tillage types are indicated by different colors.




Figure 6. Comparison of observed and simulated effects after converting to no-tillage. The data refer to yearly averages of N_2O emissions and models DayCent, LPJmL.D.Mod and LPJmL.D.Orig (in grey). Data refer to all four sites and years of the experiments.

380 **Daily emissions**

The modified hydrology (LPJmL.D.Mod and LPJmL.G.Mod), decreased the variability of no-tillage effects on N_2O emissions of individual days in most LPJmL simulations (Fig. A3 in Appendix A). The interquartile ranges from daily N_2O emissions simulated by LPJmL were more in agreement compared to the observations and DayCent, as the variability of no-tillage effects on N_2O emissions is declined.

385 In the LPJmL.D.Mod experiment, simulated N_2O emissions from no-tillage are now produced by both denitrification and nitrification (Fig. A2 in Appendix A). The increases in emissions from denitrification were smaller than in the LPJmL.D.Orig experiment and closer to the simulated values by DayCent in Boigneville and Nebraska. The emissions from nitrification increased by switching from conventional tillage to no-tillage systems, whereas they decreased in the LPJmL.D.Orig experiment. However, changes in nitrification remain small, compared to changes in denitrification.

390 **4 General discussion**

Detailed information on agricultural management improved the LPJmL simulation of N_2O emissions produced by tillage and no-tillage, as well as of the effect of switching from conventional tillage to no-tillage systems. However, also with detailed information, LPJmL overestimated the N_2O emissions. The overestimation is caused by too high simulated soil moisture, resulting in high fluxes from denitrification. After correcting for the overestimation in soil moisture, by modifying 1) the
395 parameter that translate litter amounts into soil cover, 2) the parameter that determines the duration of the surface litter layer



and 3) hydraulic properties, the yearly averages of N₂O emissions were closer to the observed values for tillage and no-tillage separately, but not for shifting from conventional tillage to no-tillage. However, the variability of no-tillage effects on N₂O emissions between the days is now reduced in most of the LPJmL simulations and the interquartile ranges from LPJmL simulations are now in better agreement with observations and DayCent.

400 DayCent performed better in simulating tillage and no-tillage effects on N₂O emissions on the yearly averages. However, DayCent tended to underestimate the overall effects and the inter annual variability of no-tillage on the emissions. DayCent mostly simulated a decrease in N₂O emissions upon shifting to no-tillage. A major reason for this is that in DayCent conversion to no-tillage leads to increasing soil organic matter which is associated with decreased availability of mineral N. However, observations showed that no-tillage can also increase N₂O emissions. For example, no-tillage can result in increased soil
405 moisture content which can promote N₂O emissions from denitrification. DayCent simulations showed basically no response in N₂O emissions from denitrification. On the other hand, conventional tillage can increase the decomposition rate of (soil) organic matter, through improved aeration of the soil. Increased decomposition leads to an increase of available N that can be transformed to N₂O through nitrification and denitrification. The higher N₂O emissions with conventional tillage in DayCent, indicates that the increase in decomposition rate of (soil) organic matter due to tillage, is dominant in comparison to the effect
410 of increased soil moisture-driven denitrification rate.

The overall better performance by DayCent likely reflects the years of model development and testing at this scale and previous application at these sites (except the site in Boigneville) (Campbell et al., 2014; Del Grosso et al., 2008; Yang et al., 2017), which enabled more accurate reproduction of observed N₂O emissions. The testing of the model performance as well as improvements to reproduce observed N₂O emissions has been conducted in several studies (Necpálová et al., 2015; Fitton
415 et al., 2014; Del Grosso et al., 2010). For example, model calibration has been conducted to test the model performance based on contributing parameters and key processes that affect N₂O emissions. For instance, the maximum amount of N₂O emissions produced during nitrification as well as the proportion of nitrified N that is lost as N₂O can be specified. LPJmL, is developed for global-scale applications and is therefore usually not calibrated, as suitable calibration targets are typically not available at that scale.

420 The application of LPJmL at the experimental sites provided much insight into the deviations of the tillage effects on N₂O emissions from observations. It enabled to use site-specific information on agricultural management, whereas missing information at global scale has to be supplemented with assumptions. As detailed information improved the simulation of tillage effects on N₂O emissions, advancing the current state of information on agricultural management at the global scale could improve global estimates of tillage effects on N₂O emissions. The study also highlighted the potential of improving the
425 simulation of N₂O emission by improving soil moisture dynamics. Any modification to improve LPJmL5.0-tillage needs to be evaluated at the global scale, as LPJmL is typically applied at that scale (e.g. Heinke et al., 2019; Rolinski et al., 2018; Schaphoff et al., 2018). A first recommendation is to revisit the PTF used in LPJmL5.0-tillage. We saw in this exercise that LPJmL overestimated soil moisture independent of the tillage system. Although the modifications in residue cover improved the results on soil moisture, the most important modification was in the hydraulic properties resulting from the PTF. The modifications
430 still resulted in relatively high soil moisture contents, and therefore possibly still overestimations in N₂O emissions. A reason



for this could be the relatively inefficient percolation of soil moisture to lower soil layers as soon as soil moisture is higher than FC.

N_2O emissions from denitrification increase exponentially when the WFPS exceeds a certain threshold value in LPJmL. This threshold value (which is around 0.8 of WFPS) is a proxy for assuming anaerobic conditions, and is static for all soil texture types. However, finer-textured soils have lower gas diffusivity at a given WFPS than coarser textured soils (e.g. Del Grosso et al., 2000). In soils with lower gas diffusivity, denitrification is assumed to occur at lower levels of WFPS, because atmospheric O_2 may not diffuse into the soil fast enough to fully satisfy microbial demand (Parton et al., 1996). Threshold values for anoxic conditions that are soil texture type specific are currently not accounted for in LPJmL. In DayCent, the effect of gas diffusivity of different soil texture types is taken into account. An index of gas diffusivity is calculated based on the WFPS, bulk density and FC, which is a proxy for pore size distribution and air filled pore space. This index influences the denitrification rate (i.e. lower diffusivity increases denitrification), N_2 to N_2O and NO_x to N_2O ratios. Including such processes in LPJmL might improve simulated N_2O emissions. However, this would require suitable reference data in order to parameterize these processes well.

5 Conclusions

Previous findings have shown deviations between simulations with the LPJmL5.0-tillage model and the results from meta-analyses on global estimates of tillage effects on N_2O emissions. In this study, we tested LPJmL5.0-tillage at different experimental sites to study whether deviations in N_2O emissions result from a lack of detailed information on agricultural management and/or the representation of soil water dynamics. The results were compared to observed values of the experimental sites as well as to results of the field-scale model DayCent.

Adding site-specific information on agricultural management improved the simulation of N_2O emissions under conventional tillage and no-tillage practices, as well as changes in emissions from shifting from conventional tillage to no-tillage in LPJmL5.0-tillage. Although adding information on agricultural management improved the performance of LPJmL5.0-tillage, simulated N_2O emissions remained too high, due to a general bias in over-estimations of soil moisture. By modifying the parameters related to residue cover and the hydraulic properties as used in DayCent, the simulation of soil moisture and N_2O emissions by LPJmL5.0-tillage improved substantially.

Generally, there is substantial uncertainty in simulating the effects of different tillage systems on N_2O emissions. DayCent performed better in simulating N_2O emissions under conventional tillage and no-tillage, but generally showed little response in N_2O emissions on changes in tillage practices. LPJmL5.0-tillage simulations reproduced a broader range of tillage effects on N_2O emissions, but tended to overestimate N_2O emissions in general. Modifications to the detail of management information considered and soil hydrology could always only improve in one deficiency (bias or dynamics) but not in both.

This study confirmed that the deviations in N_2O emissions can be explained by both lacking detailed information on management and relative high soil moisture levels simulated by LPJmL5.0-tillage. Advancing the current state of information on agricultural management can thus improve global estimates of tillage effects on N_2O emissions. Furthermore, the repre-



465 presentation of soil water dynamics and N₂O dynamics highlights the potential to improve LPJmL5.0-tillage. However, given the limited skill to reproduce observed patterns in simulations with LPJmL5.0-tillage, the model currently does not lend itself to evaluating the impacts of different tillage systems on N₂O emissions but requires further research on better representation of soil hydrology and its effects on N₂O emissions.

Code and data availability. The LPJmL source code is publicly available under the GNU AGPL version 3 license. An exact version of the code described here and the R script used for post processing the data from the simulations conducted are archived under
470 <https://doi.org/10.5281/zenodo.3592381> (Lutz et al., 2019b).

<https://doi.org/10.5194/gmd-2019-364>
Preprint. Discussion started: 14 February 2020
© Author(s) 2020. CC BY 4.0 License.



Appendix A




Figure A1. Effects of no-tillage on N₂O emissions on individual days by the different experimental simulations, including the original runs of LPJmL, the observations and simulated values by DayCent.




Figure A2. The relative share of N_2O emissions from nitrification and denitrification on individual days with no-tillage. The simulated values include the original (purple lines) and the modified (black lines) LPJmL settings. The simulated values by DayCent are also shown. Observed values are not available.




Figure A3. Effects of no-tillage on N_2O emissions on individual days by the different experimental simulations, including the original- (purple lines) and the modified (black lines) simulations from LPJmL, the observations and simulated values by DayCent.



Table A1. Performance Daycent and LPJmL over all sites and years

Models	Tillage type	RMSD	r	Sign. ($H_0: r = 0$)	Mean Bias	Standard Deviation
Daycent	Conv. tillage	7.60	0.67	0.00	1.35	3.24
Daycent	No tillage	4.61	0.66	0.00	1.50	2.29
LPJmL.G.Orig	Conv. tillage	23.60	-0.31	0.23	0.31	0.86
LPJmL.G.Orig	No tillage	36.20	-0.07	0.80	0.16	0.51
LPJmL.D.Orig	Conv. tillage	31.70	0.34	0.18	0.22	0.59
LPJmL.D.Orig	No tillage	38.90	0.36	0.16	0.15	0.46
LPJmL.G.Mod	Conv. tillage	14.25	-0.47	0.06	0.46	2.68
LPJmL.G.Mod	No tillage	13.86	-0.07	0.79	0.34	2.41
LPJmL.D.Mod	Conv. tillage	16.30	0.38	0.13	0.37	1.11
LPJmL.D.Mod	No tillage	18.10	0.60	0.01	0.27	0.88
Daycent	No tillage- Conv.tillage	4.96	0.34	0.02	0.88	6.85
LPJmL.G.Orig	No tillage- Conv.tillage	18.00	-0.16	0.55	-0.08	1.17
LPJmL.D.Orig	No tillage- Conv.tillage	12.00	0.48	0.05	-0.16	0.61
LPJmL.G.Mod	No tillage- Conv.tillage	7.17	-0.33	0.19	-0.60	2.21
LPJmL.D.Mod	No tillage- Conv.tillage	7.35	-0.04	0.89	-0.45	1.64



Author contributions. FL and CM designed the study in discussion with SDG and SO. FL conducted all model simulations and wrote the paper with support from CM. FL prepared all figures with support from SM. FL conducted the analyses with inputs from CM and JH. All authors edited the paper text.

475 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. F.L., S.M and S.R. gratefully acknowledge the German Ministry for Education and Research (BMBF) for funding this work, which is part of the MACMIT project (01LN1317A). F.L. also acknowledges the Huub and Julienne Spiertz Fund that enabled to visit the Colorado State University and USDA to collaborate on this research. Support for this research was also provided by the NSF Long-term Ecological Research Program (DEB 1832042) at the Kellogg Biological Station and by Michigan State University AgBioResearch.



480 References

- Alvarez, C., Costantini, A., Alvarez, C. R., Alves, B. J., Jantalia, C. P., Martellotto, E. E., and Urquiaga, S.: Soil nitrous oxide emissions under different management practices in the semiarid region of the Argentinian Pampas, *Nutr. Cycling Agroecosyst.*, 94, 209–220, <https://doi.org/10.1007/s10705-012-9534-9>, 2012.
- Álvaro-Fuentes, J., Morell, F. J., Plaza-Bonilla, D., Arrúe, J. L., and Cantero-Martínez, C.: Modelling tillage and nitrogen fertilization effects on soil organic carbon dynamics, *Soil Tillage Res.*, 120, 32–39, <https://doi.org/10.1016/j.still.2012.01.009>, 2012.
- 485 Begum, K., Kuhnert, M., Yeluripati, J. B., Ogle, S. M., Parton, W. J., Williams, S. A., Pan, G., Cheng, K., Ali, M. A., and Smith, P.: Modelling greenhouse gas emissions and mitigation potentials in fertilized paddy rice fields in Bangladesh, *Geoderma*, 341, 206–215, <https://doi.org/10.1016/j.geoderma.2019.01.047>, 2019.
- Boeckx, P., Van Nieuland, K., and Van Cleemput, O.: Short-term effect of tillage intensity on N₂O and CO₂ emissions, *Agron. Sustainable*
- 490 *Dev.*, 31, 453–461, <https://doi.org/10.1007/s13593-011-0001-9>, 2011.
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: how well do we understand the processes and their controls?, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368, 20130 122, <https://doi.org/10.1098/rstb.2013.0122>, 2013.
- Campbell, E. E., Johnson, J. M., Jin, V. L., Lehman, R. M., Osborne, S. L., Varvel, G. E., and Paustian, K.: Assessing the soil carbon, biomass production, and nitrous oxide emission impact of corn stover management for bioenergy feedstock production using DAYCENT, *Bioenergy research*, 7, 491–502, <https://doi.org/10.1007/s12155-014-9414-z>, 2014.
- 495 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., and Heimann, M.: Carbon and other biogeochemical cycles, pp. 465–570, Cambridge University Press, 2014.
- Del Grosso, S., Parton, W., Mosier, A., Ojima, D., Kulmala, A., and Phongpan, S.: General model for N₂O and N₂ gas emissions from soils due to denitrification, *Global Biogeochem. Cycles*, 14, 1045–1060, <https://doi.org/10.1029/1999GB001225>, 2000.
- 500 Del Grosso, S., Ojima, D., Parton, W., Mosier, A., Peterson, G., and Schimel, D.: Simulated effects of dryland cropping intensification on soil organic matter and greenhouse gas exchanges using the DAYCENT ecosystem model, *Environ. Pollut.*, 116, S75–S83, [https://doi.org/10.1016/S0269-7491\(01\)00260-3](https://doi.org/10.1016/S0269-7491(01)00260-3), 2002.
- Del Grosso, S., Parton, W., Ojima, D., Keough, C., Riley, T., and Mosier, A.: Chapter 18. DAYCENT Simulated Effects of Land Use and Climate on County Level N Loss Vectors in the USA, pp. 1–28, Academic Press/Elsevier, Amsterdam, Boston, 2008.
- 505 Del Grosso, S., Ogle, S., Parton, W., and Breidt, F.: Estimating uncertainty in N₂O emissions from US cropland soils, *Global Biogeochem. Cycles*, 24, <https://doi.org/10.1029/2009GB003544>, 2010.
- Del Grosso, S. J., Parton, W. J., Mosier, A. R., Ojima, D. S., Kulmala, A. E., and Phongpan, S.: General model for N₂O and N₂ gas emissions from soils due to denitrification, *Global Biogeochem. Cycles*, 14, 1045–1060, <https://doi.org/10.1029/1999gb001225>, 2000.
- 510 Del Grosso, S. J., Ojima, D. S., Parton, W. J., Stehfest, E., Heistemann, M., DeAngelo, B., and Rose, S.: Global scale DAYCENT model analysis of greenhouse gas emissions and mitigation strategies for cropped soils, *Global Planet. Change*, 67, 44–50, <https://doi.org/10.1016/j.gloplacha.2008.12.006>, 2009.
- Deng, Q., Hui, D., Wang, J., Yu, C.-L., Li, C., Reddy, K. C., and Dennis, S.: Assessing the impacts of tillage and fertilization management on nitrous oxide emissions in a cornfield using the DNDC model, *Journal of Geophysical Research: Biogeosciences*, 121, 337–349, <https://doi.org/10.1002/2015jg003239>, 2016.
- 515



- Elliott, J., Müller, C., Deryng, D., Chryssanthacopoulos, J., Boote, K. J., Büchner, M., Foster, I., Glotter, M., Heinke, J., Iizumi, T., Izaurralde, R. C., Mueller, N. D., Ray, D. K., Rosenzweig, C., Ruane, A. C., and Sheffield, J.: The Global Gridded Crop Model Intercomparison: data and modeling protocols for Phase 1 (v1.0), *Geosci. Model Dev.*, 8, 261–277, <https://doi.org/10.5194/gmd-8-261-2015>, <https://www.geosci-model-dev.net/8/261/2015/>, 2015.
- 520 Erb, K.-H., Luysaert, S., Meyfroidt, P., Pongratz, J., Don, A., Kloster, S., Kuemmerle, T., Fetzel, T., Fuchs, R., Herold, M., et al.: Land management: data availability and process understanding for global change studies, *Global Change Biol.*, 23, 512–533, <https://doi.org/10.1111/gcb.13443>, 2017.
- FAO: World reference base for soil resources, vol. 3, Food & Agriculture Org., 1998.
- Fitton, N., Datta, A., Hastings, A., Kuhnert, M., Topp, C., Cloy, J., Rees, R., Cardenas, L., Williams, J., Smith, K., et al.: The challenge of
525 modelling nitrogen management at the field scale: simulation and sensitivity analysis of N₂O fluxes across nine experimental sites using DailyDayCent, *Environ. Res. Lett.*, 9, 095 003, <https://doi.org/10.1088/1748-9326/9/9/095003>, 2014.
- Folberth, C., Elliott, J., Müller, C., Balkovič, J., Chryssanthacopoulos, J., Izaurralde, R. C., Jones, C. D., Khabarov, N., Liu, W., Reddy, A., et al.: Parameterization-induced uncertainties and impacts of crop management harmonization in a global gridded crop model ensemble, *PLoS One*, 14, e0221 862, <https://doi.org/10.1371/journal.pone.0221862>, 2019.
- 530 Grandy, A. S., Loecke, T. D., Parr, S., and Robertson, G. P.: Long-term trends in nitrous oxide emissions, soil nitrogen, and crop yields of till and no-till cropping systems, *J. Environ. Qual.*, 35, 1487–1495, <https://doi.org/10.2134/jeq2005.0166>, 2006.
- Gregory, J. M.: Soil cover prediction with various amounts and types of crop residue, *Transactions of the ASAE*, 25, 1333–1337, <https://doi.org/10.13031/2013.33723>, 1982.
- Gryze, S. D., Wolf, A., Kaffka, S. R., Mitchell, J., Rolston, D. E., Temple, S. R., Lee, J., and Six, J.: Simulating greenhouse gas budgets of
535 four California cropping systems under conventional and alternative management, *Ecol. Appl.*, 20, 1805–1819, <https://doi.org/10.1890/09-0772.1>, 2010.
- Halvorson, A. D., Mosier, A. R., Reule, C. A., and Bausch, W. C.: Nitrogen and tillage effects on irrigated continuous corn yields, *Agron. J.*, 98, 63–71, <https://doi.org/10.2134/agronj2005.0174>, 2006.
- Heinke, J., Müller, C., Lannerstad, M., Gerten, D., and Lucht, W.: Freshwater resources under success and failure of the Paris climate
540 agreement, *Earth Syst. Dyn.*, 10, 205–217, <https://doi.org/10.5194/esd-10-205-2019>, 2019.
- Jin, V. L., Schmer, M. R., Stewart, C. E., Sindelar, A. J., Varvel, G. E., and Wienhold, B. J.: Long-term no-till and stover retention each decrease the global warming potential of irrigated continuous corn, *Global Change Biol.*, 23, 2848–2862, <https://doi.org/10.1111/gcb.13637>, 2017.
- Jägermeyr, J., Gerten, D., Heinke, J., Schaphoff, S., Kummu, M., and Lucht, W.: Water savings potentials of irrigation systems: global
545 simulation of processes and linkages, *Hydrol. Earth Syst. Sci.*, 19, 3073–3091, <https://doi.org/10.5194/hess-19-3073-2015>, <https://www.hydrol-earth-syst-sci.net/19/3073/2015/hess-19-3073-2015.html>, 2015.
- Kelly, R., Parton, W., Hartman, M., Stretch, L., Ojima, D., and Schimel, D.: Intra-annual and interannual variability of ecosystem processes in shortgrass steppe, *Journal of Geophysical Research: Atmospheres*, 105, 20 093–20 100, <https://doi.org/10.1029/2000JD900259>, 2000.
- Krysanova, V. and Wechsung, F.: SWIM (Soil and Water Integrated Model) User Manual, <https://www.pik-potsdam.de/members/valen/swim>,
550 accessed on 01.08.2019, 2000.
- Lutz, F., Herzfeld, T., Heinke, J., Rolinski, S., Schaphoff, S., von Bloh, W., Stoorvogel, J. J., and Müller, C.: Simulating the effect of tillage practices with the global ecosystem model LPJmL (version 5.0-tillage), *Geosci. Model Dev.*, 12, 2419–2440, <https://doi.org/10.5194/gmd-12-2419-2019>, 2019a.



- Lutz, F., Müller, C., Heinke, J., Minoli, S., and Rolinski, S.: LPJmL5.0-tillage: Original source code as used in Lutz et al., 2019: submitted
555 to Geosci. Model Dev., <https://doi.org/10.5281/zenodo.3592381>, 2019b.
- Lutz, F., Stoorvogel, J. J., and Müller, C.: Options to model the effects of tillage on N₂O emissions at the global scale, *Ecol. Modell.*, 392,
212–225, <https://doi.org/10.1016/j.ecolmodel.2018.11.015>, 2019c.
- Mei, K., Wang, Z., Huang, H., Zhang, C., Shang, X., Dahlgren, R. A., Zhang, M., and Xia, F.: Stimulation of N₂O emission by conservation
tillage management in agricultural lands: A meta-analysis, *Soil Tillage Res.*, 182, 86–93, <https://doi.org/10.1016/j.still.2018.05.006>, 2018.
- 560 Mosquera, J., ter Beek, C., and Hol, J.: Precise soil management as a tool to reduce CH₄ and N₂O emissions from agricultural soil. II. Field
measurements at arable soils in the Netherlands, Report 9067549851, Agrotechnology & Food Innovations, 2005.
- Mueller, N. D., Gerber, J. S., Johnston, M., Ray, D. K., Ramankutty, N., and Foley, J. A.: Closing yield gaps through nutrient and water
management, *Nature*, 490, 254, <https://doi.org/10.1038/nature11420>, 2012.
- Necpálová, M., Anex, R. P., Fienen, M. N., Del Grosso, S. J., Castellano, M. J., Sawyer, J. E., Iqbal, J., Pantoja, J. L., and Barker, D. W.:
565 Understanding the DayCent model: Calibration, sensitivity, and identifiability through inverse modeling, *Environmental Modelling &
Software*, 66, 110–130, <https://doi.org/10.1016/j.envsoft.2014.12.011>, 2015.
- Oorts, K., Merckx, R., Gréhan, E., Labreuche, J., and Nicolardot, B.: Determinants of annual fluxes of CO₂ and N₂O in long-term no-tillage
and conventional tillage systems in northern France, *Soil Tillage Res.*, 95, 133–148, <https://doi.org/10.1016/j.still.2006.12.002>, 2007.
- Pannkuk, C., Stockle, C., and Papendick, R.: Evaluating CropSyst simulations of wheat management in a wheat-fallow region of the US
570 pacific northwest, *Agric. Syst.*, 57, 121–134, [https://doi.org/10.1016/s0308-521x\(97\)00076-0](https://doi.org/10.1016/s0308-521x(97)00076-0), 1998.
- Parton, W., Mosier, A., Ojima, D., Valentine, D., Schimel, D., Weier, K., and Kulmala, A. E.: Generalized model for N₂ and N₂O production
from nitrification and denitrification, *Global Biogeochem. Cycles*, 10, 401–412, <https://doi.org/10.1029/96GB01455>, 1996.
- Parton, W., Holland, E., Del Grosso, S., Hartman, M., Martin, R., Mosier, A., Ojima, D., and Schimel, D.: Generalized model for NO_x and
N₂O emissions from soils, *Journal of Geophysical Research: Atmospheres*, 106, 17 403–17 419, <https://doi.org/10.1029/2001JD900101>,
575 2001.
- Parton, W. J., Hartman, M., Ojima, D., and Schimel, D.: DAYCENT and its land surface submodel: description and testing, *Global Planet.
Change*, 19, 35–48, [https://doi.org/10.1016/s0921-8181\(98\)00040-x](https://doi.org/10.1016/s0921-8181(98)00040-x), 1998.
- Plaza-Bonilla, D., Álvaro Fuentes, J., Bareche, J., Pareja-Sánchez, E., Justes, É., and Cantero-Martínez, C.: No-tillage reduces long-term
yield-scaled soil nitrous oxide emissions in rainfed Mediterranean agroecosystems: A field and modelling approach, *Agriculture, Ecosys-
580 tems & Environment*, 262, 36–47, <https://doi.org/10.1016/j.agee.2018.04.007>, 2018.
- Potter, P., Ramankutty, N., Bennett, E. M., and Donner, S. D.: Characterizing the spatial patterns of global fertilizer application and manure
production, *Earth Interact.*, 14, 1–22, <https://doi.org/10.1175/2009EI288.1>, 2010.
- Rolinski, S., Müller, C., Heinke, J., Weindl, I., Biewald, A., Bodirsky, B. L., Bondeau, A., Boons-Prins, E., Bouwman, A., and Leffe-
laar, P.: Modeling vegetation and carbon dynamics of managed grasslands at the global scale with LPJmL 3.6, *Geosci. Model Dev.*,
585 <https://doi.org/10.5194/gmd-11-429-2018>, 2018.
- Rosenzweig, C., Jones, J. W., Hatfield, J. L., Ruane, A. C., Boote, K. J., Thorburn, P., Antle, J. M., Nelson, G. C., Porter, C., Janssen, S.,
et al.: The agricultural model intercomparison and improvement project (AgMIP): protocols and pilot studies, *Agric. For. Meteorol.*, 170,
166–182, <https://doi.org/10.1016/j.agrformet.2012.09.011>, 2013.
- Saxton, K., Rawls, W., Romberger, J., and Papendick, R.: Estimating generalized soil-water characteristics from texture, *Soil Sci. Soc. Am.*
590 J., 50, 1031–1036, <https://doi.org/10.2136/sssaj1986.03615995005000040039x>, 1986.



- Saxton, K. E. and Rawls, W. J.: Soil Water Characteristic Estimates by Texture and Organic Matter for Hydrologic Solutions, *Soil Sci. Soc. Am. J.*, 70, 1569–1577, <https://doi.org/10.2136/sssaj2005.0117>, 2006.
- Schaphoff, S., Bloh, W. v., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Knauer, J., Langerwisch, F., Lucht, W., Müller, C., Rolinski, S., and Waha, K.: LPJmL4 – a dynamic global vegetation model with managed land – Part 1: Model description, *Geosci. Model Dev.*, 11, 1343–1375, <https://doi.org/10.5194/gmd-11-1343-2018>, 2018.
- Schlüter, S., Großmann, C., Diel, J., Wu, G.-M., Tischler, S., Deubel, A., and Rücknagel, J.: Long-term effects of conventional and reduced tillage on soil structure, soil ecological and soil hydraulic properties, *Geoderma*, 332, 10–19, <https://doi.org/10.1016/j.geoderma.2018.07.001>, 2018.
- Smith, J. and Smith, P.: *Environmental modelling: an introduction*, Oxford University Press, 2007.
- 600 Smith, K.: Changing views of nitrous oxide emissions from agricultural soil: key controlling processes and assessment at different spatial scales, *European Journal of Soil Science*, 68, 137–155, <https://doi.org/10.1111/ejss.12409>, 2017.
- Snyder, C. S., Bruulsema, T. W., Jensen, T. L., and Fixen, P. E.: Review of greenhouse gas emissions from crop production systems and fertilizer management effects, *Agriculture Ecosystems & Environment*, 133, 247–266, <https://doi.org/10.1016/j.agee.2009.04.021>, 2009.
- Van Kessel, C., Venterea, R., Six, J., Adviento-Borbe, M. A., Linquist, B., and van Groenigen, K. J.: Climate, duration, and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis, *Global Change Biol.*, 19, 33–44, <https://doi.org/10.1111/j.1365-2486.2012.02779.x>, 2013.
- 605 Venterea, R. T., Maharjan, B., and Dolan, M. S.: Fertilizer source and tillage effects on yield-scaled nitrous oxide emissions in a corn cropping system, *Journal of Environmental Quality*, 40, 1521–1531, <https://doi.org/10.2134/jeq2011.0039>, 2011.
- Von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., and Zaehle, S.: Implementing the nitrogen cycle into the dynamic global vegetation, hydrology, and crop growth model LPJmL (version 5.0), *Geosci. Model Dev.*, 11, 2789–2812, <https://doi.org/https://doi.org/10.5194/gmd-11-2789-2018>, 2018.
- 610 Waha, K., Van Bussel, L., Müller, C., and Bondeau, A.: Climate-driven simulation of global crop sowing dates, *Global Ecol. Biogeogr.*, 21, 247–259, <https://doi.org/10.1111/j.1466-8238.2011.00678.x>, 2012.
- Yang, Q., Zhang, X., Abraha, M., Del Grosso, S., Robertson, G., and Chen, J.: Enhancing the soil and water assessment tool model for simulating N₂O emissions of three agricultural systems, *Ecosyst. Health Sustainability*, 3, e01 259, <https://doi.org/10.1002/ehs2.1259>, 2017.
- 615 Yoo, J., Woo, S.-H., Park, K.-D., and Chung, K.-Y.: Effect of no-tillage and conventional tillage practices on the nitrous oxide (N₂O) emissions in an upland soil: soil N₂O emission as affected by the fertilizer applications, *Applied Biological Chemistry*, 59, 787–797, <https://doi.org/10.1007/s13765-016-0226-z>, 2016.