

Dear editor,

referring to the manuscript :

“TheDiaTo (v1.0) – A new diagnostic tool for water, energy and entropy budgets in climate models”
by Valerio Lembo, Frank Lunkeit and Valerio Lucarini (ID: gmd-2019-37)

please find attached the revised draft, with all tracked changes to the original document.

In the following, the comments raised by the two reviewers are reported together with the author's replies and, if applicable, the related changes in the text. Reviewers' comments are reported in plain text. Author's replies and modified text (with quotation marks) in italic.

In addition to that, we have modified Figures 1, 4-a, 6 in order to remove the blank vertical line on the GMT longitude. Labels in Figures 2 and 8 were modified as well, because the definition of enthalpy is more suitable to the computations that are performed here.

Yours Sincerely,

Valerio Lembo

Referee RC1

- Introduction: For energy and water budget studies using observations, please add L'Ecuyer et al, 2015 and Rodell et al, 2015. For studies evaluating the energy and water budgets in climate models, please add Demory et al, 2014; Terai et al, 2017; Vanniere et al, 2018 (e.g. in P2 L29, P3 L22, P4 paragraph 1.2). These studies highlighted the need for more robust evaluation of these budgets in climate models.

Thank you for the suggestions. We have added these references in a new sentence in Sect. 1.2: “Water mass budget has been assessed in observations (L'Ecuyer et al., 2015; Rodell et al., 2015), as well as in climate models, focusing on the hydrological cycle alone (Terai et al., 2018) or evaluating it in conjunction with the energy budgets (Demory et al., 2014; Vannière et al., 2019).”

(l. 15-17, p. 4): “Water mass budget has been assessed in observations (L'Ecuyer et al., 2015; Rodell et al., 2015), as well as in climate models, focusing on the hydrological cycle alone (Terai et al., 2018) or evaluating it in conjunction with the energy budgets (Demory et al., 2014; Vannière et al., 2019).”

- section 2 data and software requirements: are calculations performed on the native grid of the models or on a common grid?

The calculations are performed on the native grid of the model, as provided in the CMIP repository. We specified this in Sect. 2 by adding the following sentence: “For model inter-comparisons (Section 5), computations are performed on the native grid of the model.”

(l. 13-14, p. 6) “ For model inter-comparisons (Section 5), computations are performed on the native grid of the model”.

- section 3: can the tool consider observational data, where available, to validate the models?

The tool can be used to compare models and observational data, as long as the data are global-scale and on a regular grid. The ESMValTool v.2 framework allows for straightforward comparison with Reanalyses (as it will be discussed in the ESMValTool v.2 report that is soon to be submitted on this same journal). Since referee #1 also suggested to mention that, we have added a comment on the range of applicability to observational-based gridded datasets in Sect. 2 as follows: “Therefore, the tool is suitable for the evaluation of any kind of gridded datasets, provided that they contain the necessary variables on a regular grid, including blends of observations and Reanalyses. In our description of the software features, we focus on model evaluation and multi-model intercomparison.”

(l. 5-8, p.g 6) “Therefore, the tool is suitable for the evaluation of any kind of gridded datasets, provided that they contain the necessary variables on a regular grid, including blends of observations and Reanalyses. In our description of the software features, we focus on model evaluation and multi-model intercomparison.”

- section 3: can the tool also work with seasonal means?

Technically, no constraint is given on the usage of a subset of months. Although this would cause in principle no problem on the computations of the budgets and the LEC strength, the material entropy production would be inconsistent, because there is no straightforward argument justifying that the indirect method would provide a reasonable estimate of this quantity (as mentioned when discussing Eq. 11 in Sect. 3.4). A thorough explanation of the assumptions is provided in Lucarini et al. 2011. Furthermore, the implied transports rely on the fact that the differential heating is balanced by the oceans and the atmosphere, but this is not certainly accurate at the smaller than annual timescale (cfr. for instance Rose and Ferreira 2013 on the validity of the Bjerknes compensation mechanism).

- section 3.2 hydrological cycle: Assessing the water budget in climate models is particularly important to determine their conservation error and its evolution throughout the simulation. A model may have a conservation error but that remains stable over time, while another one may constantly lose/gain mass over time (as noted in Liepert and Previdi, 2012). In this study, this would be particularly important to understand what is happening in BNU. Is it possible to see the time evolution of the conservation error with TheDiaTo (Fig. 2 of Liepert and Previdi), and if not, would it be possible to add it? This would be beneficial for a thorough evaluation of the climate models.

The diagnostic tool allows to obtain annual mean time series of energy and water mass budgets for the Northern Hemisphere, the Southern Hemisphere and the global mean. For sake of brevity, we decided not to include them. An example of what the time series for the BNU-ESM model would look like is shown in Figures 1-3. It can be noticed that the conservation error is quite massive, compared to the other models, but relatively constant.

Figure 1: *annual mean time series of the energy budget for BNU-ESM pre-industrial run. Red denotes Southern Hemisphere, blue denotes Northern Hemisphere, black the global mean.*




Figure 2: same as Figure 1, for the last 20 years of the historical experiment.




Figure 3: same as Figure 1, for the last 20 years of the RCP8.5 experiment.

- P15 L21: is there a reason for picking this period: 2441-2460?

The reason for that is that there is no other section of the pre-industrial CanESM2 run for which 20-years of all the needed variables is available on the repository. We explained this by adding the following sentence: “The choice of the sub-period is motivated by the fact that it is the only part of the run for which a 20-years subsequent dataset of the needed variables is available in the repository.”

(l. 14-15, p. 16) “The choice of the sub-period is motivated by the fact that it is the only part of the run for which a 20-years subsequent dataset of the needed variables is available in the repository”.

- P18 L28 & L32: 'bias' generally refers to errors relative to observations. Here, it may be more

robust to refer to 'imbalance'.

Thank you. We replaced “bias” with the more appropriate term “imbalance”.

(l. 11-12, p. 19) “As shown in the first two columns of Table 2, there is a large imbalance in the TOA energy budget under unforced piC conditions.” (and elsewhere in sec. 5.1, l. 11-29)

- Tables 8 and 9: specify that rows are for PiC, hist and rcp85.

We have changed the table captions as follows: “For each model, the first row denotes the estimates from piC, the second row from hist, the third row from rcp8.5”.

(Table 8 & Table 9 captions): “For each model, the first row denotes the estimates from piC, the second row from hist, the third row from rcp8.5. Values are in mW/m²K⁻¹.”

- P1 L5: replace 'Top-of-Atmosphere (TOA)' by TOA, as defined just earlier

Done. Thank you.

(l. 4-5, p. 1) “The routine takes as inputs energy fluxes at surface and at the Top-of-Atmosphere (TOA), which allows for the computation of energy 5 budgets at the TOA, the surface, and in the atmosphere as a residual.”

- P2 L16: remove 'of things'

Done. Thank you.

(l. 16 p. 2): “Difficulties can only increase when looking at the practical side:”

- P2 L21: be more specific with 'the required standards mentioned above' so we don't have to jump back to recall these

We have replaced “prepared according to the required standards mentioned above” with “containing process-based and end-user-relevant diagnostics”.

(l. 20-22, p. 2): “What we propose in this paper is a novel software, containing process-based and end-user-relevant diagnostics, capable of providing an integrated perspective on the problem of models validation and intercomparison.”

- P2 L23: same as above for 'enunciated above ideas'

We have added “reaching a steady state as a result of an interplay of multi-scale processes”.

(l. 22-24, p. 2): “The goal here is to examine models through the lens of their dynamics and thermodynamics, in the view of enunciated above ideas about complex non-equilibrium systems reaching a steady state as a result of an interplay of multi-scale processes.”

- P5 L5: same as above for 'mentioned aspects'

We have added “(i.e. the energy and water mass budgets and meridional transports, the LEC strength and the material entropy production)”.

(9-11, p. 3): “We present here TheDiaTo v1.0, a new software for diagnosing the mentioned aspects of thermodynamics of the climate systems (i.e. the energy and water mass budgets and meridional transports, the LEC strength and the material entropy production) in a broad range of global-scale gridded datasets of the atmosphere.”

- P23 L17: 'effected' -> 'affected'

What we mean here is that baroclinic eddies “effect” baroclinic conversions. They can be both “synoptic-scale” and “planetary-scale” eddies, as shown in a recent paper of us (Lembo et al. 2019, 46, Geophys. Res. Lett.). We have added this reference, since it was not yet accepted by the time the manuscript was submitted.

(l. 29-31 p. 23) “The two quantities are related through the meridional enthalpy transports (cfr. Boschi et al. (2013); Lembo et al. (2019b)), which in mid-latitudes are mainly effected by baroclinic eddies.”

Referee RC2

My main criticism (speaking as an “outsider” - not directly involved in diagnostic studies of state of the art Earth climate models) would be of the description in Section 2 of the data requirements for use of the tool. This is quite short and seems to presume quite a lot of knowledge on the part of the reader (unless they are prepared to spend a lot of time and effort consulting the documentation for the ESMValTool software, referenced as Eyring et al. (2016b)). Table 1 lists the variables required to be presented to the tool for the various calculations, but simply names the variables according to the CMOR convention without explanation. This is not helpful if you don't know that convention, and it is not explained anywhere in the paper except in external references. Why not provide a key for these variables or an explanation in another column of Table 1? Given the general applicability of much of the tool to climates that are not restricted to Earth (as acknowledged in the paper itself), readers would appreciate a bit more help and guidance for novice users. Is there a graphical interface, for example, or is it run via a shell script? Some more guidance here and perhaps an example script would be helpful, either in the text or another appendix.

We thank the reviewer for the constructive criticism. We addressed the issue in two steps. We rearranged Table 1, in order to include a short description of each variable. In addition to that, we substantially rephrased the last part of Section 2, shortly explaining the algorithm. We did not dig into the details of the procedure, because the documentation of the code contains all the necessary information, either one uses it through the ESMValTool v.2, or prefers downloading the stand-alone version of the diagnostic tool (which is now available in a GitHub repository: https://github.com/ValerioLembo/TheDiaTo_v1.0.git)

(l. 3-13, p. 7): “The ESMValTool architecture is implemented as a Python package and the latest version is available at [www.doi.org/10.17874/ac8548f0315](https://doi.org/10.17874/ac8548f0315), where the dependencies and installation requirements are also described. If using TheDiaTo v.1 as part of the ESMValTool architecture, the user is asked to specify the path to input data, work and plot directories, as well as

some details of the local machine. A dedicated namelist (named as "recipe") includes information on the diagnostics, such as a brief description, the reference literature, the developers. The user is asked to specify here the name of the dataset to be analysed, that must be recognised by the ESMValTool preprocessor. The diagnostic tool is thus run calling the ESMValTool interface with the associated recipe.

A stand-alone version of the software is maintained as well, utilizing Python bindings for Climate Data Operators (CDO) (CDO, 2015). It consists of a Python script, reading a namelist with user settings (such as directory paths, model names and options for the usage of the modules). The stand-alone version is publicly available in a GitHub repository (https://github.com/ValerioLembo/TheDiaTo_v1.0), where detailed instructions on the usage of the software are given." and added column in Table 1.

- Figure 5 and associated text: The reservoir terms ASE, ATE, KSE and KTE are not explained or defined very obviously. They presumably refer to stationary and transient components of AE and KE? It would be helpful to state this in the caption and perhaps mention in the text?

Thank you for the comment. We have modified the caption to Figure 5 and we added a short period at Section A2: "The contribution of eddies to APE (EPE) and KE (EKE) consists of two terms, the first one is the term accounting for stationary eddies (ASE and KSE in the diagrams of Figure 5, respectively). The second term accounts for the transient eddies (ATE and KTE in the diagrams of Figure 5, respectively). A discussion on the derivation of these terms is found in Ulbrich and Speth, 1991."

(l. 6-9, p. 27): "The contribution of eddies to APE (EPE) and KE (EKE) consists of two terms, the first one is the term accounting for stationary eddies (ASE and KSE in the diagrams of Figure 5, respectively). The second term accounts for the transient eddies (ATE and KTE in the diagrams of Figure 5, respectively). A discussion on the derivation of these terms is found in Ulbrich and Speth (1991)."

- P.17 line 23 as example of awkward phrasing: should probably read ". . .as a possible reason for the well known cold pole bias. . .". But there are quite a few others – too numerous to list in detail. Perhaps recruit someone whose first language is English to proofread - or the authors should proofread more thoroughly before submission!

We thank the reviewer for the constructive criticism. We have undergone a comprehensive proofread of the manuscript, together with a native speaker. We hope that this effort has substantially improved the quality of the language.

(l. 4-6, p. 18): "This is consistent with previous findings (cfr. Boer and Lambert (2008); Marques et al. (2011)) and is possibly due to the well-known cold Pole bias and the consequently excessive speed of the jet stream." Generally, as evidenced in the document with tracked changes, we performed a careful rewording of the manuscript.

- Presumably this tool could be used on reanalysis datasets too? Perhaps comment further?

Thank you for the suggestion. We have added a sentence in the first paragraph of Section 2, explicitly mentioning that the software is also suitable for the ingestion of Reanalysis datasets: "Therefore, the tool is suitable for the evaluation of any kind of gridded datasets, provided that they contain the necessary variables on a regular grid, including blends of observations and Reanalyses.

In our description of the software feature, we focus on model evaluation and multi-model intercomparison.”

(l. 5-8, p.g 6) “Therefore, the tool is suitable for the evaluation of any kind of gridded datasets, provided that they contain the necessary variables on a regular grid, including blends of observations and Reanalyses. In our description of the software features, we focus on model evaluation and multi-model intercomparison.”

TheDiaTo (v1.0) – A new diagnostic tool for water, energy and entropy budgets in climate models

Valerio Lembo¹, Frank Lunkeit¹, and Valerio Lucarini^{1,2,3}

¹Meteorologisches Institut, Universität Hamburg, Hamburg, Germany

²Department of Mathematics and Statistics, University of Reading, Reading, UK

³Centre for the Mathematics of Planet Earth, Department of Mathematics and Statistics, University of Reading, Reading, UK

Correspondence to: Valerio Lembo (valerio.lembo@uni-hamburg.de)

Abstract. This work presents the Thermodynamic Diagnostic Tool (TheDiaTo), a novel diagnostic tool for studying-investigating the thermodynamics of the climate systems with a wide range of applications, from sensitivity studies to model tuning. It includes a number of modules for assessing the internal energy budget, the hydrological cycle, the Lorenz Energy Cycle and the material entropy production, ~~respectively~~. The routine ~~receives-takes~~ as inputs energy fluxes at surface and at the Top-
5 of-Atmosphere (TOA), which allows for the computation of energy budgets at ~~Top-of-Atmosphere (TOA), at the TOA~~, the surface, and in the atmosphere as a residual. Meridional enthalpy transports are also computed from the divergence of the zonal mean energy ~~budget-fluxes; budgetfrom which the~~ location and intensity of ~~peaks in the two hemispheres are then~~
~~provided-as-outputs~~the maxima in each hemisphere are calculated. Rainfall, snowfall and latent heat fluxes are received as inputs for ~~computing-computation of~~ the water mass and latent energy budgets. If a land-sea mask is provided, the required
10 quantities are separately computed over continents and oceans. The diagnostic tool also computes the annual Lorenz Energy Cycle (LEC) and its storage/conversion terms ~~as annual-mean global-and-hemispheric-values. In order to achieve this, one needs~~
~~to provide-as-input-by~~ hemisphere and as a global mean. This is computed from three-dimensional daily fields of horizontal wind velocity and temperature in the troposphere. Two methods have been implemented for the computation of the material entropy production, one relying on the convergence of radiative heat fluxes in the atmosphere (indirect method) ~~one-and the~~
15 other combining the irreversible processes occurring in the climate system, particularly heat fluxes in the boundary layer, the hydrological cycle and the kinetic energy dissipation as retrieved from the residuals of the LEC (direct method). A version of these diagnostics has been developed as part of the Earth System Model eValuation Tool (ESMValTool) v2.0a1, in order to assess the performances of CMIP6 model simulations, and will be available in the next release~~of the tool~~. The aim of this software is to provide a comprehensive picture of the thermodynamics of the climate system, as reproduced in the state-of-the-
20 art coupled general circulation models. This can prove useful for better understanding of anthropogenic and natural climate change, paleoclimatic climate variability, and climatic tipping points.

1 Introduction

The climate can be ~~seen-viewed~~ as a forced~~and~~, dissipative non-equilibrium system exchanging energy with the external environment, ~~and-setting-a~~. The inhomogeneous absorption of solar radiation is an ongoing source of available potential energy.

The complex mixture of ~~fluid-flows-fluids is then set~~ into motion by ~~converting-the conversion of~~ available potential into mechanical energy ~~through-via~~ a vast range of nonlinear processes. The ~~source-of-available-potential-energy-is-the-inhomogeneous absorption-of-solar-radiation. The~~ kinetic energy is eventually dissipated through viscous stress and converted back into heat. Such processes can be described by taking advantage of the theory of non-equilibrium thermodynamics of continuous multi-
5 phase media, and, in particular, of fluids. ~~Non-equilibrium-systems-are-characterised-by-the-presence-of-, possibly-fluctuating,~~ ~~The presence of (possibly fluctuating)~~ fluxes of matter, chemical species, and energy ~~is a key characteristic of a non-equilibrium system.~~ The steady state is reached as a result of a ~~possibly-potentially~~ complex balance of positive and negative feedbacks, and through the interplay of processes ~~having-with~~ very diverse time scales and physical underpinning mechanisms. ~~Indeed,~~
10 ~~the climate-can-be-viewed-in-such-framework, as its-observed-variability-extends~~ ~~The climate is a prime example of this, with~~ ~~observed variability extending~~ over many orders of magnitude in terms of both spatial and temporal scales, and ~~is-divided-in~~
~~already-with many~~ extremely complex subdomains - the atmosphere, the ocean, the ~~eriosphere~~~~cryosphere~~, the biosphere, the active soil - which have themselves very diverse characteristic internal time-scales and are nonlinearly coupled (Peixoto and Oort, 1992; Lucarini et al., 2014).

It is a major endeavour of contemporary science to improve our understanding of the climate system ~~both~~ in the context of
15 the past, present, and projected future conditions. This is key for understanding, as far as the past goes, the co-evolution of life and of the physico-chemical properties of the ocean, soil, and atmosphere, and for addressing the major challenge faced by our planet as a result of the current anthropogenic climate change. ~~Radically-advancing-the skills-of~~ ~~Improving~~ climate models is key to nearing these goals, and, indeed, efforts aimed in this direction have been widely documented (see the related chapter on the last report, of the Intergovernmental Panel on Climate Change (2013)). Intercomparing and validating climate models is
20 far from being a trivial task, also at a purely conceptual level (Lucarini, 2013; Frigg et al., 2015). Difficulties can only increase when looking at the practical side~~of-things~~: how to choose meaningful metrics to study the performance of climate models? Should they be motivated in terms of basic processes of the climate system, or in terms of relevance for an end-user of climate services? In order to put some order in this conundrum, the community of climate modellers has developed a set of standardized metrics for intercomparing and validating climate models (Eyring et al., 2016a). For obvious reasons, the choice made for
25 such metrics has been biased - for possibly good reasons - in the direction of providing ~~useful-information-for~~ ~~ready-to-use~~ ~~information for end~~ users. What we propose in this paper is a novel software, ~~prepared-according-to-the-required-standards~~ ~~mentioned-above~~~~containing process-based and end-user-relevant diagnostics~~, capable of providing an integrated perspective on the problem of models validation and intercomparison. The goal here is to ~~look-at-examine~~ models through the lens of their dynamics and thermodynamics, in the view of enunciated above ideas about complex non-equilibrium systems ~~reaching~~
30 ~~a steady state as a result of an interplay of multi-scale processes.~~ The metrics that we ~~here-propose~~ ~~propose here~~ are based on the analysis of the energy and water budgets and transports, of the energy transformations, and of the entropy production. We summarize below some of the key concepts behind our work.

1.1 Energy

A requirement for In order to be in steady state, a non-equilibrium system in contact with an external environment ~~to be at steady state is the presence of~~ must have a vanishing - on ~~the~~ average - energy budget. Inconsistencies in the overall energy budget of long-term stationary simulations have been carefully ~~evaluated~~ pointed out (Lucarini and Ragone, 2011; Mauritsen et al., 2012) and various aspects of the radiative and heat transfers within the atmosphere and between the atmosphere and the

5 oceans have been ~~reconsidered~~ evaluated in order to constrain models to a realistic climate (Wild et al., 2013; Loeb et al., 2015). A substantial bias in the energy budget of the atmosphere in particular has been identified in many GCMs, resulting from either the imperfect closure of the kinetic energy budget (Lucarini and Ragone, 2011) or ~~the imperfect closure~~ of the mass balance in the hydrological cycle (Liepert and Previdi, 2012). This picture is made even more complicated by the difficult task of having an accurate observational benchmark of the Earth's energy budget (e.g. Loeb et al. (2009); von Schuckmann et al. (2016)).

10 ~~In general, an increasing consensus is being reached on the fact~~ Many authors have recently suggested that the improvement of climate models requires improving the energetic consistency of the modeled system (Hansen et al., 2011; Lucarini et al., 2011, 2014). Rather than a proxy of a changing climate, surface temperatures and precipitation changes ~~should~~ can be better viewed as a consequence of a non-equilibrium steady state system which is responding to a radiative energy imbalance through ~~a complex interaction of~~ complex interacting feedbacks. A changing climate, under the effect of an external transient forcing,

15 can only be properly addressed if the energy imbalance, and the way it is transported within the system and converted into different forms is taken into account. The ~~models' skill to represent the history of~~ skill of the models at representing historical energy and heat exchanges in the climate system has been assessed by comparing numerical simulations against available observations, where available (Allan et al., 2014; Smith et al., 2015), including the fundamental problem of ocean heat uptake (Exarchou et al., 2015).

20 A key element in defining the steady-state of the climate system is the balance between the convergence of the horizontal (mostly meridional) enthalpy fluxes ~~due to the atmosphere and the ocean by the atmospheric and the oceanic circulations~~ and the radiative imbalance at ~~the top of the atmosphere (TOA). Such TOA. The net radiative~~ imbalance is positive (in-bound) in the low latitudes and negative in the high latitudes, and a compensating horizontal transport must be present in order to ensure steady state. ~~As a result, the transport reduces dramatically~~ This transport dramatically reduces the meridional temperature gradient

25 with respect to what would be set by radiative-convective equilibrium (Manabe and Wetherald, 1967), i.e. in absence of large scale atmospheric and oceanic transport. Differences in the boundary conditions, in the forcing and dissipative processes, and in chemical and physical properties of the atmosphere and ocean lead to a specific partitioning of the enthalpy transport between the two geophysical fluids; a different partitioning associated to the same total transport would lead to ~~rather~~ significantly different climate conditions (Rose and Ferreira, 2013; Knietzsch et al., 2015). The role of meridional heat transports in different

30 paleoclimate scenarios and in relation to different forcing has been addressed in various studies (see e.g. Caballero and Langen (2005); Fischer and Jungclauss (2010)). ~~State-of-the-art climate models have been critically analyzed -- with mixed findings across the available models -- in order to understand whether they~~ The question of whether the current state-of-the-art climate models are able to represent correctly the global picture as well as the details of the atmospheric and oceanic heat transports has been analysed with mixed findings (Lucarini et al., 2011, 2014; Lembo et al., 2016).

In order to understand how the heat is transported by the geophysical fluids, one should clarify what sets them into motion. We focus here on the atmosphere. A comprehensive view of the energetics fuelling the general circulation is given by the Lorenz Energy Cycle (LEC) framework (Lorenz, 1955; Ulbrich and Speth, 1991). This provides a picture of the various processes responsible for conversion of available potential energy (APE) ~~, i.e.,~~ the excess of potential energy with respect to a state of thermodynamic equilibrium (~~cf.~~ see Tailleux (2013) for a review) ~~,~~ into kinetic energy and dissipative heating. Under stationary conditions, the dissipative heating exactly equals the mechanical work performed by the atmosphere. ~~In other words~~ Thus, the LEC formulation allows to constrain the atmosphere to the first law of thermodynamics, and the system as a whole can be ~~seen as a pure thermodynamic~~ interpreted as a heat engine under dissipative non-equilibrium conditions (Ambaum, 2010). The strength of the LEC, or in other words the intensity rate of the conversion of available potential energy (APE) into kinetic energy (KE), has been evaluated in observational-based datasets (Li et al., 2007; Pan et al., 2017) and climate models (Lucarini et al., 2010a; Marques et al., 2011), ~~mostly with estimates generally~~ ranging from 1.5 to 3.5 W/m^2 . A considerable source of uncertainty is the hydrostatic assumption, on which the LEC formulation relies, ~~possibly leading which can lead~~ to significant underestimation of the kinetic energy dissipation (Pauluis and Dias, 2012). ~~In some respects, one can see the~~ The usual formulation of the LEC ~~as suitable for~~ can be seen as describing energy exchanges and transformation at a coarse-grained level, where non-hydrostatic processes are not relevant (~~and the~~ and the system is accurately described by primitive equations). ~~Given its nature of heat engine, an~~ An efficiency can also be attributed to the atmosphere ~~, assuming that the system is as a heat engine, assuming the system as~~ analogous to an engine ~~working operating a warm and a cold operating between a warmer and a colder~~ temperature (Lucarini, 2009). This approach has been generalized by Pauluis (2011), in order to account for the role of water vapor.

1.2 Water

~~Indeed, water~~ Water is an essential ingredient of the climate system, and the hydrological cycle ~~has~~ plays an important role ~~on in~~ the energy pathways of the climate system, ~~because water~~ Water vapour and clouds ~~impact influence~~ the radiative processes inside the system, and water phase exchanges are extremely energy intensive. ~~Similarly to~~ As in the case of energy imbalances, a closed water-mass conserving reproduction of the hydrological cycle ~~has proved to be~~ is essential, not only because of the ~~disparate~~ diverse implications of the hydrological cycle for energy balance and transports in the atmosphere, but also because of its sensitivity to climate change (Held and Soden, 2006) and the ~~extreme relevance~~ importance of the cloud and water vapor feedbacks (Hartmann, 1994). ~~On one hand the~~ The energy budget is ~~relevantly affected~~ influenced by semi-empirical formulations of the water vapor spectrum (Wild et al., 2006), ~~on the other hand the~~ and, similarly, the energy budget influences the moisture budget by means of uncertainties in aerosol-cloud interactions and mechanisms of tropical deep convection (Wild and Liepert, 2010; Liepert and Previdi, 2012). ~~A~~ Water mass budget has been assessed in observations (L'Ecuyer et al., 2015; Rodell et al., 2015), as well as in climate models, focusing on the hydrological cycle alone (Terai et al., 2018) or evaluating it in conjunction with the energy budgets (Demory et al., 2014; Vannière et al., 2019). Therefore, a global scale evaluation of the hydrological cycle, both from a moisture and from an energetic perspective, is ~~thus~~ considered an integral part of an overall diagnostics for the thermodynamics of climate system.

1.3 Entropy

The climate system has long been recognized as featuring irreversible processes through dissipation and mixing in various forms, leading to the production of entropy (Paltridge, 1975), which is ~~one of the key properties~~ key characteristic of non-equilibrium systems (Prigogine, 1962). Several early attempts (Peixoto and Oort, 1992; Johnson, 1997; Goody, 2000) have been made to understand the complex nature of irreversible climatic processes. Recent works (Bannon, 2015; Bannon and Lee, 2017) have proposed an innovative approach by partitioning the system into a control volume made of matter and radiation, which exchanges energy with its surroundings (building on Goody (2000) early works). Raymond (2013) has described the entropy budget of an aggregated dry air + water vapor parcel. From a macroscopic point of view, ~~one usually refers to the~~ "material entropy production" ~~as~~ generally refers to the entropy produced by the geophysical fluids in the climate system, which ~~are is~~ not related to the properties of the radiative fields, but rather to the irreversible processes related to the motion of these fluids. ~~Mainly, this has to do with~~ The material entropy production is dominated by phase changes and water vapor diffusion, as outlined by Pauluis and Held (2002). Lucarini (2009) underlined the link between entropy production and efficiency of the climate engine, which were then ~~used to understand~~ applied to study climatic tipping points, ~~and, in-particular, in particular~~ the snowball/warm Earth critical transition (Lucarini et al., 2010b), ~~to define~~. This allowed the definition of a wider class of climate response metrics (Lucarini et al., 2010a) ~~, and that have been used~~ to study planetary circulation regimes (Boschi et al., 2013). A constraint has also been proposed to the entropy production of the atmospheric heat engine, given by the emerging importance of non-viscous processes in a warming climate (Laliberté et al., 2015).

Given the multiscale properties of the climate system, accurate energy and entropy budgets are affected by subgrid-scale parametrizations (see also Kleidon and Lorenz (2004); Kunz et al. (2008)). These and the discretization of the numerical scheme are generally problematic in terms of conservation principles (Gassmann and Herzog, 2015), and can eventually lead to macroscopic model drifts (Mauritsen et al., 2012; Gupta et al., 2013; Hobbs et al., 2016; Hourdin et al., 2017). See Lucarini and Fraedrich (2009) for a theoretical analysis in a simplified setting.

1.4 This Paper

We present here TheDiaTo v1.0, a new software for diagnosing the mentioned aspects of thermodynamics of the climate systems (i.e. the energy and water mass budgets and meridional transports, the LEC strength and the material entropy production) in a broad range of global-scale gridded datasets of the atmosphere. The diagnostic tool provides global metrics, allowing straightforward comparison of different products. These include:

- Top-of-Atmosphere, atmospheric and surface energy budgets;
- Total, atmospheric and oceanic meridional enthalpy transports;
- Water mass and latent energy budget;
- Strength of the LEC by means of kinetic energy dissipation conversion terms;

- Material entropy production ~~with the~~ by both direct and indirect methods;

The software is structured in terms of independent modules, so that ~~the users are allowed to choose either all or part of~~ users can subset the metrics according to their interest. A version of the tool has been developed as part of the ESMValTool community effort (Eyring et al., 2016b) to provide a standardized set of diagnostics for the evaluation of Coupled Model Intercomparison

- 5 Project Phase 6 (CMIP6) multi-model ensemble simulations (Eyring et al., 2016a). The current version ~~of~~ at time of writing was of ESMValTool is v2.0a1 ~~–A and a~~ new version is soon to be released including TheDiaTo v1.0.

Therefore, our goal is to equip climate modellers and developers with tools for better understanding the strong and weak points of the models of interest. The ~~final~~ aim is to reduce the risk of a model having good outputs for accurate reproductions of quantities of common interest, such as surface temperature or precipitation, but for the wrong dynamical reasons. ~~Clearly,~~

- 10 ~~this is a first necessary~~ This is a necessary first step in the direction of creating a suite of model diagnostics composed of process-oriented metrics. ~~The~~

The paper is structured as follows. The dataset requirements are described ~~(Section 2)~~ in Section 2. We then ~~clarify~~ establish the methods used in each module (Section 3). In Section 4 an example is given of application ~~with one to a~~ 20-years-long model run. In section 5 ~~an extensive evaluation is given of~~ the evolution of ~~various considered the~~ metrics under three different

15 scenarios ~~for which in~~ the CMIP5 ~~(Taylor et al., 2012) provides a~~ multi-model ensemble ~~(Taylor et al., 2012) is evaluated~~. Summary and conclusions are given in Section 6.

2 Data and Software Requirements

The diagnostic tool consists of a set of independent modules, each, except the first one, being triggered by a switch decided by the user: energy budgets and enthalpy transports, hydrological cycle, Lorenz Energy Cycle ~~–and~~

- 20 ~~with via~~ the direct or indirect method.

The software ingests all variables as ~~gridded~~ fields on a regular longitude-latitude grid covering the whole globe. Therefore, the tool is suitable for the evaluation of any kind of gridded datasets, provided that they contain the necessary variables on a regular grid, including blends of observations and Reanalyses. In our description of the software features, we focus on model evaluation and multi-model intercomparison.

- 25 For the LEC computation, 3D fields are required, stored in pressure levels at a daily or finer temporal resolution. If the model does not store data where the surface pressure is lower than the respective pressure levels, daily mean or higher resolution data of near-surface temperatures and horizontal velocity fields are also required for vertical interpolation. For all other computations, 2D fields are required as monthly means at ~~Top-of-the-Atmosphere (TOA)–~~ TOA and at the surface. Input ~~data variables~~ are given as separate ~~NetCDF files – files in NetCDF format. For model inter-comparisons (Section 5), computations are~~
- 30 performed on the native grid of the model. Variables are identified according to their variable names, ~~that~~ Those are required to comply to the Climate and Forecast (CF, <http://cfconventions.org/Data/cf-documents/overview/article.pdf>) and Climate Model Output Rewriter (CMOR, <http://pcmdi.github.io/cmor-site/tables.html>) standards. Datasets that are not complying with the CF-CMOR compliant names are reformatted through ESMValTool built-in preprocessing routines, if ~~known~~ recognized (for

more detail, refer to the dedicated report on ESMValTool ~~v2~~v1.0 Eyring et al. (2016b)). An overview of the required variables is provided in Table 1.

Energy budgets are computed from residuals of instantaneous radiative shortwave (SW) and longwave (LW) fluxes at TOA. At the surface ~~they~~, SW and LW fluxes are combined with instantaneous turbulent latent and sensible heat fluxes (cfr. Section

- 5 3). The radiative fluxes (except the outgoing LW radiation, which is only upwelling) are composed of an upwelling and downwelling component, that are defined positive. The heat fluxes at the surface are computed in climate models from the usual bulk formulas, different models differing in the choice of the parameters, and are positive upwards. Water mass and latent energy budgets are computed from evaporation(implied from latent heat turbulent fluxes at the surface), rainfall and snowfall instantaneous fluxes. ~~In case of recognized cumulative fields, these are converted to instantaneous by the ESMValTool~~
10 ~~preprocessor~~Some models might provide cumulative energy/water mass fields, instead of instantaneous fluxes. If this is the case, the ESMValTool preprocessor performs the conversion to instantaneous fluxes, depending on the analysis timestep.

For the LEC module, 3D fields of the three components of velocity and temperatures ~~, as well as near-surface temperature and horizontal velocity 2D fields~~ are needed at the daily resolution. For the 3D fields, there is no specific constraint on the

- number of pressure levels, although the program has been tested on the standard pressure level vertical discretization used in
15 CMIP5 outputs, consisting on 17 levels from 1000 to 1 hPa. The program then subsets the troposphere between 900 and 100 hPa. LEC computation is performed on Fourier coefficients of the temperature and velocity fields. 2D fields of temperature and horizontal velocity are also required in order to fill gaps in the pressure level discretization by interpolating from the surface on a reference vertical profile. As further discussed later (Sect. 5.2), ~~the vertical interpolation from near-surface fields to fill the empty fields in pressure levels, when necessary, is~~ this technique introduces an inevitable source of uncertainty.

- 20 If explicitly ~~requested~~required by the user, the program is also able to perform computations of energy budgets, enthalpy transports and the hydrological cycle on oceans and continents separately, provided a land-sea mask is supplied. This can either be in the form of land area fraction or ~~of~~ a binary mask.

The ESMValTool architecture is ~~conceived as a library of Python script, whose~~ implemented as a Python package and the latest version is available at www.doi.org/10.17874/ac8548f0315, where the ~~dependency~~dependencies and installation requirements

- 25 are also described. If using TheDiaTo v.1 as part of the ESMValTool architecture, the user is asked to specify the path to input data, work and plot directories, as well as some details of the local machine. A dedicated namelist (named as "recipe") includes information on the diagnostics, such as a brief description, the reference literature, the developers. The user is asked to specify here the name of the dataset to be analysed, that must be recognised by the ESMValTool preprocessor. The diagnostic tool is thus run calling the ESMValTool interface with the associated recipe.

- 30 A stand-alone version of the software is maintained as well, ~~relying on~~utilizing Python bindings for Climate Data Operators (CDO) (CDO, 2015). It consists of a Python script, reading a namelist with user settings (such as directory paths, model names and options for the usage of the modules). The stand-alone version is publicly available in a GitHub repository (https://github.com/ValerioLembo/TheDiaTo_v1.0), where detailed instructions on the usage of the software are given.

3 Methods

3.1 Energy budgets and meridional enthalpy transports

Under the crucial assumptions of neglecting ~~By making the crucial assumption that~~ the heat content of liquid and solid water in the atmosphere, ~~together with~~ the heat associated with the phase transitions in the atmosphere and the effect of salinity and pressure in the oceans ~~are negligible~~, we can write the total specific energy per unit mass for the subdomains constituting the climate system as: $\epsilon = \mathbf{u}^2 + c_v T + \phi + Lq$ for the atmosphere, $\epsilon = \mathbf{u}^2 + c_w T + \phi$ for the ocean, $\epsilon = c_s T + \phi$ for solid earth and ice. Here u is the velocity vector, c_v , c_w and c_s are the specific heat at constant volume of the atmospheric mix at constant volume, water and the solid medium, respectively, L is the latent heat of vaporization of water, q the specific humidity and ϕ the gravitational potential. This leads to an equation for the evolution of the local specific energy in the atmosphere as such (Peixoto and Oort, 1992):

$$\frac{\partial \rho \epsilon}{\partial t} = -\nabla \cdot (\mathbf{J}_h + \mathbf{R} + \mathbf{H}_S + \mathbf{H}_L) - \nabla(\tau \cdot \mathbf{u}), \quad (1)$$

where $\mathbf{J}_h = (\rho E + p)\mathbf{u}$ is the specific enthalpy transport, \mathbf{R} is the net radiative flux, \mathbf{H}_L and \mathbf{H}_S are the turbulent sensible and latent heat fluxes respectively, ~~whereas and~~ τ is the stress tensor. ~~If we neglect~~ By neglecting the kinetic energy component, and vertically integrate integrating Eq. 1, we can write find an equation for the energy tendencies at each latitude ~~, longitude~~ (ϕ) , longitude (λ) and time for the whole climate system, for the atmosphere, and for the surface below it:

$$\begin{cases} \dot{E}_t(\lambda, \phi) = S_t^\downarrow(\lambda, \phi) - S_t^\uparrow(\lambda, \phi) - L_t(\lambda, \phi) - \nabla \cdot J_t(\lambda, \phi) = R_t(\lambda, \phi) - \nabla \cdot J_t(\lambda, \phi) \\ \dot{E}_a(\lambda, \phi) = R_t(\lambda, \phi) - F_s(\lambda, \phi) - \nabla \cdot J_a(\lambda, \phi) = F_a(\lambda, \phi) - \nabla \cdot J_a(\lambda, \phi) \\ \dot{E}_s(\lambda, \phi) = S_s(\lambda, \phi) + L_s(\lambda, \phi) - H_S^\uparrow(\lambda, \phi) - H_L^\uparrow(\lambda, \phi) - \nabla \cdot J_o(\lambda, \phi) = F_s(\lambda, \phi) - \nabla \cdot J_o(\lambda, \phi) \end{cases} \quad (2)$$

where \dot{E}_t , \dot{E}_a , \dot{E}_s denote the total, atmospheric and oceanic surface energy tendencies respectively. $S_s = S_s^\uparrow(\lambda, \phi) - S_s^\downarrow(\lambda, \phi)$, $L_s = L_s^\uparrow(\lambda, \phi) - L_s^\downarrow(\lambda, \phi)$ are the net SW and LW radiative fluxes at the surface, respectively, and S_t^\downarrow , S_s^\downarrow , S_t^\uparrow and S_s^\uparrow the TOA (t) and surface (s) upward (\uparrow) and downward (\downarrow) SW radiative fluxes respectively (and similarly for LW radiative fluxes, denoted by L , provided that ~~the there is no~~ downward LW flux at TOA ~~is neglected~~). R_t , F_a and F_s are the total, atmospheric and surface net energy fluxes respectively. Net fluxes are defined as positive when there is a net energy input, negative when there is a net output. J_t , J_a and J_o are the meridional total, atmospheric and oceanic enthalpy transports. Oceanic transports are assumed to be related to surface EBs, ~~because one assumes that~~ energy budgets, ~~because the~~ long-term enthalpy transports through land are ~~nearly~~ negligible (Trenberth and Solomon, 1994; Trenberth et al., 2001).

When globally averaged, Eq. 2 can be rewritten as (Lucarini et al., 2014):

$$\begin{cases} \dot{E}_t = R_t \\ \dot{E}_a = R_t - F_s = F_a \\ \dot{E}_s = F_s \end{cases} \quad (3)$$

Given the ~~much smaller~~ small thermal inertia of the atmosphere, compared to the oceans, the TOA energy imbalance is expected to be transferred for the most part to the ocean interior ~~in terms of ocean heat uptake, whereas, with~~ F_a is much smaller than

F_s (Levitus et al., 2012; Loeb et al., 2012; Smith et al., 2015). This is usually not ~~found-reproduced~~ in climate models (e.g. ~~Lucarini and Ragone (2011); Loeb et al. (2015); ?~~Lucarini and Ragone (2011); Loeb et al. (2015); Lembo et al. (2019a)).

Under steady-state conditions, the tendency of the internal energy of the system ~~vanishes-will vanish~~ when averaged over sufficiently long ~~timescale~~time periods. We can thus zonally average Eqs. 2 and derive the long-term ~~averages-time means~~ of the meridional ~~enthalpy-transport~~heat transports, or more appropriately, in this formulation, "meridional enthalpy transports":

$$\left\{ \begin{array}{l} T_t(\phi) = 2\pi \int_{\phi}^{\pi/2} a^2 \cos \phi' < \overline{R_t(\phi')} > d\phi' \\ T_a(\phi) = 2\pi \int_{\phi}^{\pi/2} a^2 \cos \phi' < \overline{F_a(\phi')} > d\phi' \\ T_o(\phi) = 2\pi \int_{\phi}^{\pi/2} a^2 \cos \phi' < \overline{F_s(\phi')} > d\phi' \end{array} \right. \quad (4)$$

where T_t , T_a and T_o denote the steady-state total, atmospheric and oceanic meridional enthalpy transports, a the Earth's radius, $<>$ long-term time averaging and overbars the zonal-mean quantities. The stationary condition has to be achieved in the models if the system is unforced. If the system is forced, for instance by ~~means-of-a~~ transient greenhouse gas forcing, a correction is applied in order to ~~prevent-inconsistent~~avoid unphysical cross-polar transports~~-. This correction can be applied~~ as long as the weak non-stationarity condition holds (Carissimo et al., 1985), i.e. the latitudinal variability of the ~~budgets-net energy flux~~ is much larger than the global mean imbalance (cfr. Lucarini and Ragone (2011)). The ~~correction-to-the-energy-fluxes-is-applied~~as-energy fluxes are modified as follows:

$$15 \quad < \overline{B_x}(\phi) >_{co} = < \overline{B_x}(\phi) > - < \overline{B_x}(\phi) > / 2\pi a \quad (5)$$

where B_x refers to either R_t , F_a or F_s .

Peak intensities and ~~latitudinal locations-latitudes~~ are computed as metrics for ~~such-transport~~Comparison among datasets ~~with different resolutions is ensured by interpolating to a common grid. these transports.~~

3.2 Hydrological cycle

20 The atmospheric moisture budget is obtained ~~from-by~~ globally averaging precipitation and evaporation fluxes. The latter are derived from surface latent heat fluxes as:

$$E = \frac{H_L}{L_v} \quad (6)$$

where E from now on denotes the evaporation fluxes and $L_v = 2.5008 \times 10^6$ ~~J/kg is J/kg~~ the latent heat of evaporation (assumed to be constant). If H_L is given in units of ~~W/m² W/m²~~, the implied evaporation flux is ~~coherently-given~~ in units of ~~kg * m⁻² * s⁻¹ kg/m²s⁻¹~~. Rainfall fluxes are derived from total (P) and snowfall (P_s) precipitation as:

$$P_r = P - P_s \quad (7)$$

Under the stationarity assumption, the equation for moisture budget is thus written as:

$$\overline{E} - \overline{P} = \overline{E} - \overline{P_r} - \overline{P_s} \quad (8)$$

where overbars denote [here](#) global averages.

~~For the~~ [The](#) latent energy budget in the atmosphere R_L , ~~the following is used~~ [is then](#):

$$5 \quad \overline{R_L} = \overline{H_L} - L_v \overline{P_r} - (L_v + L_f) \overline{P_s} \quad (9)$$

where $L_f = 3.34 \times 10^5$ ~~J/kg~~ [J/kg](#) is the latent heat of fusion ([which is](#) assumed to be constant). Unlike the water mass budget of Eq. 8, the latent energy budget is not closed, because it does not include the heat captured by snow ~~fusion~~ [melting](#) from the ground. ~~Furthermore, the~~ [The](#) sublimation of ice is also not ~~considered for~~ [accounted for here for](#) sake of simplicity, ~~despite~~ [although](#) it plays a significant role over the Arctic region.

10 3.3 The Lorenz Energy Cycle

The ~~derivation~~ [calculation](#) of the atmospheric LEC in the diagnostic tool follows directly from the general framework proposed by Lorenz (1955) ~~and~~ [revised](#) by Ulbrich and Speth (1991) in order to provide a separation ~~among~~ [between](#) different scales of motion. The LEC allows ~~to investigate~~ [investigation of](#) the general circulation of the atmosphere by looking comprehensively at the energy exchanges between eddies and zonal flow, at the conversion between available potential and kinetic energy, and

15 at the dissipation due to viscous and mixing processes.

[The LEC is depicted diagrammatically in Figure 5, and the equations used to obtain the energy reservoirs and conversion terms are described in the appendices A2-A3.](#) Energy is injected in the ~~reservoir of zonal~~ [reservoirs of](#) APE through differential diabatic heating impacting ~~both~~ [primarily](#) the zonal component (~~primarily~~) ~~and but also~~ the eddy component ~~of the APE~~. The so-called baroclinic instability (mainly occurring in the mid-latitude baroclinic eddies at the synoptic scale) acts in two steps;

20 [: first](#) by transforming zonal mean APE into eddy APE, ~~which is then converted~~ [and then by conversion](#) into eddy kinetic energy (KE) via a lowering of the atmospheric centre of mass. The eddy KE is then converted into smaller scales through the kinetic energy cascade, eventually reaching the dissipative scale, where it is ~~converted~~ [transformed](#) into frictional heating. A remaining part of the eddy KE is converted back into zonal mean flow through the ~~so-called~~ barotropic governor, ensuring the maintenance of the tropospheric and stratospheric jet streams (cfr. Li et al. (2007)).

25 ~~The LEC is depicted diagrammatically in Figure 5, and the equations used to obtain the energy reservoirs and conversion terms are reported in the appendices A2-A3~~ [Other terms, i.e. the generation/dissipation of APE and KE, are computed as residuals of the conversion terms at each reservoir.](#)

In the formulation proposed by Lorenz and widely adopted afterwards, ~~one assumes that motions are~~ [motions are assumed to be](#) quasi-hydrostatic. This assumption is correct as long as one considers sufficiently coarse-grained atmospheric fields. A detailed analysis of non-hydrostatic effects requires dealing with the exchange of available potential and kinetic energy taking

30 place through accelerated vertical motions ~~(cfr. Novak and Tailleux (2018))~~. Furthermore, the ~~energetics refer~~ [approach used here refers](#) to the dry atmosphere (cfr. Appendix A2-A3). In order to ~~take moisture into account, one would have to replace the temperatures with~~ [include moisture effects, temperatures must be replaced by](#) virtual temperatures. ~~The impact of such an~~

~~assumption on the estimate of the~~ A comprehensive moist formulation of the LEC is left for future study.

Following the arguments by Lucarini et al. (2014), we ~~notice note~~ that, under non-stationary ~~state conditions, the requirement for conditions,~~ internal energy conservation ~~holds applies~~ (cfr. Eqs. 4), and the stationarity condition implies that APE and KE tendencies both vanish. Given that the tendency in KE is a balance ~~of between~~ the APE-KE overall conversion (or in other
5 words the mechanical work exerted by the LEC) and the dissipation of KE, we can write:

$$\langle W \rangle = \langle D \rangle = \int_{\Omega} \rho \kappa^2 d\Omega', \quad (10)$$

where W and D denote the LEC work and the dissipation of KE respectively, κ^2 is the specific kinetic energy dissipation rate, ρ is the atmospheric density and Ω the volume of the atmosphere. ~~This justifies the fact that the intensity of the LEC is~~ The LEC can thus be used to obtain the kinetic energy dissipation of the atmosphere. ~~Other terms, i.e. the generation/dissipation of~~
10 ~~APE and KE, are computed as residuals of the conversion terms at each reservoir.~~

3.4 Material entropy production

The total entropy production in the climate system is given by two qualitatively different kinds of processes: ~~on one hand~~ firstly, the irreversible thermalization of photons emitted ~~in the Sun's corona from the Sun~~ at the much lower Earth's surface and atmospheric temperature, ~~on the other hand the~~ and secondly the irreversible processes responsible for mixing and diffusion in
15 the fluids and in the active soil of the climate system. The former accounts for roughly 95% of the total entropy production, the latter is the material entropy production (MEP) ~~in a proper sense~~, and is the quantity of most interest in climate science, because it involves the dynamics of the atmosphere and its interaction with the Earth's surface (see discussion in Kleidon (2009); Lucarini (2009); Lucarini et al. (2011, 2014))

In the long-term mean, assuming that the system is in statistically steady state condition, one can write an equation for the
20 entropy rate of change in the system (Lucarini and Ragone, 2011; Lucarini and Pascale, 2014) as:

$$\int_{\Omega} d\Omega' \left(\overline{\frac{\dot{q}_{rad}}{T}} \right) + \overline{\dot{S}_{mat}} = 0 \quad (11)$$

with \dot{q}_{rad} denoting the local net radiative heating and \dot{S}_{mat} the global material entropy production. ~~From Eq. 11 it is evident that computing~~ It is evident From eq. 11 that the entropy rate of change ~~as a consequence of radiative heating is equivalent to compute~~ can be computed indirectly from the net radiative heating or directly as the sum of the entropy production from all the
25 irreversible processes, both viscous (such as the energy dissipation) and non-viscous (such as the hydrological cycle). These two equivalent methods are here referred to as "indirect method" and "direct method", ~~respectively~~. As long as the volume integral in the first ~~lhs~~ left-hand side member of Eq. 11 is performed on the whole atmospheric domain, the two methods are exactly equivalent (for sake of simplicity we assume that this amounts to the MEP of the climate itself, since the oceanic contribution to the MEP is ~~neglected~~ negligible, accounting for about 2% of the budget ~~, as found by Pascale et al. (2011)~~
30 (Pascale et al., 2011)).

3.4.1 The direct method

The MEP computation with the direct method involves taking into account the viscous processes related to energy cascades toward the dissipative scales, and the non-viscous processes related to sensible heat fluxes (i.e. heat diffusion in the boundary layer, mainly dry air convection (Kleidon, 2009)) and ~~to~~ the hydrological cycle (such as evaporation in unsaturated air, con-

5 condensation in supersaturated air, release of gravitational potential energy due to the fall of the droplet and melting of solid phase water at the ground). ~~Generally speaking, we~~ We can write the MEP equation as:

$$\dot{S}_{mat} = \frac{\kappa^2}{T_v} + \frac{\phi}{T_\phi}, \quad (12)$$

where T_v and T_ϕ denote ~~here~~ the temperatures at which the respective processes occur, ϕ indicates local absorption or emission of heat that is neither radiative nor related to viscous dissipation ~~-In other words,-~~ and ϕ accounts for all indicating the non-

10 viscous irreversible processes.

Overall, the hydrological cycle accounts for about 35 ~~$mW * m^{-2} * K^{-1}$~~ $mW/m^2 K^{-1}$, the sensible heat diffusion for about 2 ~~$mW * m^{-2} * K^{-1}$~~ $mW/m^2 K^{-1}$ and the viscous processes for about 6 to 14 ~~$mW * m^{-2} * K^{-1}$~~ $mW/m^2 K^{-1}$ (cfr. Fraedrich and Lunkeit (2008); Pascale et al. (2011)).

Dealing with direct computation of MEP in climate models has a number of additional implications. ~~The mentioned irreversible~~

15 These processes are dealt with in climate models ~~through the usage of via~~ subgrid-scale parametrizations. The fact that they ~~are~~ should be energy conserving and entropy consistent is far from ~~being~~ trivial (Gassmann, 2013). Further, the numerical scheme adds spurious entropy sources, ~~such as e.g.~~ numerical advection and hyperdiffusion, as ~~thoroughly~~ addressed in an intermediate-complexity model by Pascale et al. (2011) ~~and focusing on or in~~ the dynamical core of a state-of-the-art model by Gassmann and Herzog (2015). The relevance of these non-negligible numerically driven components, and ~~the way how~~ to

20 address them, is strictly model-dependent and ~~a diagnostic tool that is meant to analyze this diagnostic tool which is designed to analyse~~ a potentially diverse ensemble of datasets must come to terms with that limitation.

~~In order to get to our expression for the direct MEP, let us~~ For the direct expression of the MEP, we first explicitly write the non-viscous terms in Eq. 12:

$$\dot{S}_{mat} = \int_{\Omega} d\Omega' \frac{\kappa_s^2}{T} - \int_{\Omega} d\Omega' \frac{\nabla \cdot \mathbf{h}_S}{T} + \dot{S}_{hyd}, \quad (13)$$

25 where $\nabla \cdot \mathbf{h}_S$ denotes the heat diffusion through sensible heat fluxes, and \dot{S}_{hyd} the aggregated MEP related to the hydrological cycle. The specific kinetic energy dissipation rate is here denoted by "s", in order to emphasize that it is now an estimate ~~of such a term~~, as discussed later in this section. As argued by Lucarini and Pascale (2014), there is not an easy way to account for the term related to the hydrological cycle. Pauluis and Held (2002); Pauluis (2011) ~~find point out~~ that the water mass in the atmosphere can be thought of as a "passive tracer", conveying heat until a phase change ~~, i.e. (an irreversible diabatic process;~~

30) allows it to exchange heat with the surrounding, producing material entropy ~~-This means that the atmospheric particle has to be separated in~~ (Pauluis and Held, 2002; Raymond, 2013). For this, each atmospheric parcel must be separated into its dry component and the water mass components in its various phases, and the phase changes ~~have to~~ must be evaluated, in order

to address for the associated MEP (cfr. Pauluis and Held (2002); Raymond (2013)). At in steady-state conditions an indirect estimate of \dot{S}_{hyd} is provided as:

$$\dot{S}_{hyd} = \int_{\Omega_V} d\Omega_V \rho_w \frac{\nabla \cdot \mathbf{h}_L}{T}, \quad (14)$$

where $\nabla \cdot \mathbf{h}_L$ denotes the heat exchange between the water mass particle and the surroundings during ~~its phase transition~~ phase

5 transitions.

In order to express Eq. 13 in terms of climate model outputs, we make some additional assumptions, following the approach by Fraedrich and Lunkeit (2008). First, we assume that the energy dissipation by friction occurs mainly next to the surface, ~~then and~~ we define an operating temperature T_d as an average of surface (T_s) and near-surface temperatures (T_{2m}). Secondly, we estimate the heat diffusion term from sensible turbulent heat fluxes at the surface H_S , transporting heat between the Earth's

10 surface (having temperature T_s) and the boundary layer (having temperature T_{BL} , ~~whose derivation is~~ derivation described below). We then consider the phase exchanges of water mass ~~components as follows~~ components i.e.:

- evaporation at working temperature T_s (~~indirectly derived from latent turbulent heat fluxes at the surface H_L~~ cfr. Section 3.2);
- rain droplet formation through condensation at working temperature T_C , a characteristic temperature of the cloud;
- 15 – snow droplet formation through condensation+solidification at working temperature T_C , ~~a characteristic temperature of the cloud~~;
- snow melting at the ground at working temperature T_s ;

Following from Eq. 4 in Lucarini et al. (2011) we can thus rewrite Eq. 13 as:

$$\begin{aligned} \dot{\Sigma}_{mat} = & \int_A \frac{\kappa_s^2}{T_d} dA' - \int_A H_S \left(\frac{1}{T_s} - \frac{1}{T_{BL}} \right) dA' - \int_A \frac{L_v E}{T_s} dA' + \int_{A_r} \left(\frac{L_v P_r}{T_C} + g \frac{P_r h_{ct}}{T_p} \right) dA'_r \\ & + \int_{A_s} \left(\frac{L_s P_s}{T_C} + g \frac{P_s h_{ct}}{T_p} \right) dA'_s - \int_{A_s} \frac{L_s P_s}{T_s} dA'_s \end{aligned} \quad (15)$$

- 20 Since the latent and sensible heat, rainfall and snowfall precipitation fluxes are given in model outputs as 2D fields, the volume integrals in Eq. 14 reduce to area integrals, with the domain A denoting the Earth's surface and the subdomains A_r and A_s denoting the regions where rainfall and snowfall precipitation occur, respectively. The phase changes associated with snowfall and rainfall precipitation (4th and 5th ~~rhs integrals~~ terms on the right-hand side) are accompanied by a term accounting for the potential to kinetic energy conversion of the falling droplet (with g denoting the gravity acceleration and h_{ct} the distance
- 25 covered by the droplet). ~~One may notice~~ Note that, in principle, this is a viscous term, ~~since as~~ the kinetic energy of the droplet is eventually dissipated into heat at the ground.

The first ~~rhs member~~ term on the right-hand side of Eq. 15 involves the specific kinetic energy dissipation rate (κ_s^2). There is no straightforward way to ~~describe this quantity in climate models~~ extract this quantity from climate model outputs. Pascale et al.

(2011) found that, ~~besides along with~~ the major role of the vertical momentum diffusion ~~in terms of as a~~ frictional stress in the boundary layer, the gravity wave drag and ~~unphysical processes, some unphysical processes~~ - such as horizontal momentum diffusion (hyperdiffusion) ~~, have also - also have~~ a non-negligible role. Each model accounts for these quantities in a different way. Generally, the frictional term is also not present in climate model outputs, thus has to be indirectly estimated from near-

5 surface velocity fields. In order to do so, ~~one has to know~~ the value of the drag coefficient is needed, which is different in every model. ~~In order to tackle this problem, also considering~~ Considering that the kinetic energy dissipation term ~~overall contributes to accounts for~~ less than 10% of the overall MEP, we compute it indirectly, obtaining the kinetic energy dissipation from the intensity of the LEC (cfr. Eq. 10).

The boundary layer temperature T_{BL} is not usually provided as a climate model output, nor the boundary layer thickness is

10 known a-priori. Some manipulations are thus needed to make a working approximation. We start from the definition of a bulk Richardson number:

$$Ri_b = \frac{g}{\theta_{v0}} \frac{(\theta_{vz} - \theta_{v0})z}{u_z^2 + v_z^2} \quad (16)$$

where $g = 9.81 \text{ m} \cdot \text{s}^{-1}$ is the gravity acceleration, θ_{v0} and θ_{vz} are the virtual potential temperatures at the surface and at level z , and u_z and v_z are the zonal and meridional components of the horizontal wind at height z (assumed to be equal to

15 the ~~usually provided~~ near-surface horizontal velocity fields). A critical Richardson number (Ri_{bc}) is defined as the value of the Richardson number at the top of the boundary layer. Its value depends on the nature of the local boundary layer (stable or unstable). In order to distinguish among the stable and unstable boundary layers, a condition on the magnitude of the sensible heat fluxes is imposed (Zhang et al., 2014), so that where H_S is lower than 0.75 W/m^2 , a stable boundary layer is assumed ($Ri_{bc} = 0.39$; boundary layer height z_{BL} : 300 m) , and otherwise an unstable boundary layer is assumed ($Ri_{bc} = 0.28$;

20 boundary layer height z_{BL} : 1000 m). ~~For sake of simplicity, we~~ We approximate the virtual potential temperature as with the dry potential temperature, so that the conversion from temperature to potential temperature and vice versa is given by the basic formula:

$$\theta = T \left(\frac{p}{p_0} \right)^{R_d/c_p} \quad (17)$$

where $R_d = 287.0 \text{ J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$ J/kg K⁻¹ is the gas constant for dry air and $c_p = 1003.5 \text{ J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$ J/kg K⁻¹ is the specific heat of the atmosphere at constant pressure. ~~We can thus obtain~~ Then T_{BL} can be obtained by solving Eq. 16 for $\theta_{z_{BL}}$, imposing Ri_{bc} and z_{BL} ~~as mentioned, and~~ retrieving the value of p at z_{BL} by means of the barometric equation:

$$p_z = p_s e^{-gz/R_d T_s} \quad (18)$$

where p_s the surface air pressure and z is in our case z_{BL} , ~~then using it to obtain~~. This then gives the temperature at the ~~boundary layer top. Here we assume~~ top of the boundary layer. A crucial assumption is here that the boundary layer is approx-

30 imately isothermal.

In Eq. 15, T_C is a temperature representative of the interior of the cloud ~~from~~ where the moist particle drops. In order to define

that, we first retrieve the dewpoint temperature at the surface from the equation:

$$T_d = \frac{1}{1/T_0 - R_v/L_v \log(e/\alpha)} \quad (19)$$

where $T_0 = 273.15 \text{ K}$ is the reference melting temperature, $R_v = 461.51 \text{ J} * \text{kg}^{-1} * \text{K}^{-1}$ is the gas constant for water vapor,

$e = \frac{q_s p_s}{q_s + R_d/R_v}$ is the water vapor pressure (where we have used q_s , i.e. the near-surface specific humidity) ~~;~~ and $\alpha = 610.77$

5 P_a is one of the empirical parameters of the Magnus-Teten formulas for saturation water pressure (cfr. Goff (1957); Buck (1981)). An empirical formula for the computation of the lifting condensation level (LCL) (Lawrence, 2005) can be then used:

$$h_{LCL} = 125(T_s - T_d) \quad (20)$$

If we assume that the moist particle is lifted following a dry adiabatic until it saturates at the LCL, the temperature at such level

10 will be:

$$T_{LCL} = T_s - \Gamma_d h_{LCL} \quad (21)$$

This would be the temperature of the cloud bottom in convective conditions. We hereby assume that similar conditions apply to stratiform clouds. In order to obtain T_C , ~~we average~~ T_{LCL} is averaged with the temperature of the cloud top, which is ~~assumed~~ taken to be the emission temperature T_E at TOA by inversion of the the Stephan-Boltzmann law ~~with~~ applied to the local

15 outgoing longwave (LW) radiation at TOA (i.e. L_t in Sect. 3.1).

The potential energy of the droplets in Eq. 15 is ~~retrieved assuming that the particle starts to fall~~ estimated on the basis that the drop starts from the cloud layer top (h_{ct}). This level is obtained by assuming that the saturated particle, after entering the cloud at the LCL, continues to be lifted in the cloud ~~on~~ following a pseudo-adiabatic path. We thus firstly compute the pseudo-adiabatic lapse rate:

$$20 \quad \Gamma_p = \Gamma_d \left(1 + \frac{L_v q_s}{R_d T_{LCL}} \right) * \left(1 + \frac{\epsilon L_v^2 q_s}{c_p R_d T_{LCL}^2} \right)^{-1} \quad (22)$$

where $\epsilon = 0.622$ is the molecular weight of water vapor/dry air ratio. Once the pseudo-adiabatic lapse rate is known, it is straightforward to compute the height of the cloud top ~~combining it with the~~ by usage of the approximated emission temperature. It can be observed that what we obtain is an upper constraint to the potential energy of the droplets, since we ~~assume have~~ assumed that the particle falls through the whole cloud layer, while the pseudo-adiabatic lapse rate assumes that water vapor

25 gradually precipitates during the ascent. T_p is finally obtained as an average between T_C and T_s .

In conclusion, ~~let us notice~~ note that the MEP budget provided in Eq. 15 is ~~the most~~ a reasonably accurate estimate that can be obtained from ~~largely~~ usually available climate model outputs. ~~Still~~ However, some processes related to intermediate phase transitions in the atmosphere and ~~to~~ heat exchanges at the droplet surface during its coalescence/aggregation stage are not taken into account, because the ~~information provided~~ limited information available in model outputs ~~does not allow for that~~.

30 terms are potentially relevant, as stressed by Pauluis and Held (2002) and Raymond (2013). Furthermore, the ~~contribution from~~ potential energy of the droplet hydrometeors does not usually enters the energetics of a climate model, although ~~it its~~

contribution to the MEP budget is not negligible. Finally, the MEP budget here introduced is focused on the atmosphere. Phase changes in the sea-ice domain ~~provide a potentially large~~ are another potentially significant contribution to the overall MEP of the climate (e.g. Herbert et al. (2011)).

3.4.2 The indirect method

- 5 For the indirect ~~formulation estimation~~ of the entropy budget, we ~~express~~ use a simplified expression of the entropy associated with radiative heat convergence ~~in a simplified formulation~~, following Lucarini et al. (2011). This approach is formally equivalent to the one adopted in Bannon (2015); Bannon and Lee (2017), ~~using which use~~ the definition of a control volume to describe the entropy of the material system, together with the radiation contained in it. ~~Not entering into details of the derivation, we follow~~ Following from Eq. 11 ~~by identifying those we identify the~~ processes responsible for the entropy
- 10 ~~production flux out of the material system~~ through exchanges of radiative energy, as outlined by Ozawa et al. (2003); Fraedrich and Lunkeit (2008). For each process, we thus define an energy flux between two mediums with warmer and colder temperatures. The radiative heat exchange is ~~thus carried on effected~~ locally through vertical exchanges, and on a large scale ~~, mainly~~ through meridional exchanges. ~~If we take,~~ predominantly. Considering SW and LW net radiative fluxes at the surface and at TOA (i.e. the usual output for radiative fluxes in climate models) we can write:

$$15 \quad \dot{\Sigma}_{ind}^{mat} = \int_A \frac{\overline{S_s} + \overline{L_s}}{T_s} dA + \int_A \frac{\overline{S_t} - \overline{S_s}}{T_{A,SW}} dA + \int_A \frac{\overline{L_t} - \overline{L_s}}{T_{A,LW}} dA \quad (23)$$

- where $S_t = S_t^\downarrow - S_t^\uparrow$ is the net SW radiative flux at TOA (~~cfr. see~~ Eq. 2), A is the surface area of the atmosphere, and T_s , $T_{A,SW}$ and $T_{A,LW}$ are characteristic temperatures representative of the surface and of the portion of the atmosphere where LW and SW radiative heat exchanges occur, respectively. Analogously to Eq. 15 the volume integral in Eq. 11 ~~is has been~~ transformed into an area integral, considering that the radiative fluxes ~~occur at the surface are given at the boundaries~~ of the domain and using
- 20 ~~the Gauss~~ Gauss' theorem (Lucarini et al., 2011). This formulation is ~~still~~ an exact expression for the atmospheric MEP ~~, as long as one is able to define~~ SW and LW working temperatures $T_{A,SW}$ and $T_{A,LW}$ can be accurately estimated (cfr. Bannon and Lee (2017) for a discussion on this crucial issue). ~~We follow again the arguments by Lucarini et al. (2011), rewriting~~ Following Lucarini et al. (2011), we rewrite Eq. 23 under the assumption that $T_{A,SW} \approx T_{A,LW} \approx T_E$:

$$\dot{\Sigma}_{ind}^{mat} = \int_A (\overline{S_s} + \overline{L_s}) \left(\frac{1}{T_s} - \frac{1}{T_E} \right) dA + \int_A \frac{\overline{S_t} + \overline{L_t}}{T_E} dA = \overline{\Sigma_{ver}} + \overline{\Sigma_{hor}} \quad (24)$$

- 25 This ~~decisive~~ critical assumption is based on the fact that most of SW and LW radiation is absorbed and emitted in the atmosphere through water vapor into the troposphere (Kiehl and Trenberth, 1997).

- As already stressed by Lucarini et al. (2011), the material entropy budget described in Eq. 24 consists of two terms which have useful physical interpretations. The first term ~~denotes is related to~~ the vertical energy transport between a reservoir at temperature T_s (the surface) and another at temperature T_E (the TOA). For this reason it is referred to as "vertical material entropy production ($\overline{\Sigma_{mat}^{ver}}$). This term is ~~positive almost everywhere almost everywhere positive~~ and accounts for the vertical transport of warm air from the surface, mainly ~~embedded in by~~ moist convection. The second term ~~denotes is related to~~ the
- 30

horizontal energy transport from a warm reservoir at lower latitudes to a cold reservoir at higher latitudes. This is referred to as "horizontal material entropy production" ($\overline{\Sigma_{mat}^{hor}}$) and is associated with the annual mean meridional enthalpy transport setting the ground for the mean meridional circulation (Peixoto and Oort, 1992). ~~One may notice. Note~~ that, while the first term accounts for a local entropy ~~budget~~production, the second ~~has to be rather considered is~~ a horizontal advection of entropy, and

5 should be meaningfully considered only ~~in its as a~~ global integral. Both terms are positive, the first one because the atmosphere is heated from below, the second because the ~~temperature is lower where the~~ heat is transported down-gradient.

Let us finally consider that both the direct and the indirect methods contain crucial approximations. ~~On the other hand, the~~ The 2-layer assumption ~~is critical~~reduces the estimated MEP both with the direct and indirect methods, as shown by Lucarini and Pascale (2014), ~~finding that who investigated~~ the coarse graining of post-processed model data~~always reduces the estimated~~

10 ~~MEP, both with the direct and indirect methods, with the indirect method.~~ We expect that the indirect method will be particularly affected by vertical coarse graining. ~~We expect that~~ Furthermore, the indirect method leads to an overestimation of MEP compared to approximate estimates by Ambaum (2010)), mainly because of the vertical entropy production, as already seen in Lucarini et al. (2014). The impact of considering the 3D radiative fluxes in Eq. 11 is under investigation with a specific intermediate-complexity model,~~and the outcomes will be the topic of a successive work.~~

15 4 Results from a CMIP5 model

Figure 1 about here

A 20-years ~~subset~~extract of a CMIP5 model (CanESM2) simulation under pre-industrial conditions is ~~here analysed,~~ analysed in order to ~~describe~~demonstrate the capabilities of the diagnostic tool. The datasets are retrieved from the Earth System Grid Federation (ESGF) node at Deutsches Klimarechenzentrum (DKRZ). The run used here for the analysis is the one denoted by "r1i1p1". ~~Of~~ From the 995-years (2015-3010) run, ~~we take~~ the 2441-2460 period ~~was used.~~ The choice of the sub-period

20 is motivated by the fact that it is the only part of the run for which a 20-years subsequent dataset of the needed variables is available in the repository.

Figure 1 shows the horizontal distribution of annual mean R_t , F_a and F_s . The TOA energy budget (R_t) is relatively smooth and zonally symmetric, with an area of net energy gain over the tropics and the ~~ocean's subtropics,~~ oceanic subtropics and energy loss elsewhere. A maximum is found over the Eastern Indian - Western Pacific warm pool, where the Indian Monsoons

25 develop and the emission temperature is the lowest ~~, due to the presence of~~ due to deep convection. Interestingly, this pattern is somewhat opposed by the negative values of the TOA ~~EB~~energy budget at similar latitudes over the Sahara, where the highest near-surface temperatures are found. ~~In other words, the~~ The warm and dry conditions characterizing deserted subtropical regions determine the highest thermal emission ~~, largely exceeding which largely exceeds~~ the solar input ~~, thus leading and~~ leads to a net energy loss. The surface energy budget (F_s) is almost vanishing over the continents, given ~~their~~the small thermal

30 inertia of the land surface. The largest absolute values are found in proximity of the main ~~surface~~sub-surface ocean currents. They are mainly negative in ~~coincidence~~the region of the western boundary currents (Gulf Stream, Kuroshio current, Agulhas current), where the ocean's surface transfers heat to the atmosphere. They are negative over the Humboldt Current, extending

deep into the Equatorial Counter Current, and to a lesser extent in proximity of the Antarctic Circumpolar Current, where the ocean's surface takes up heat from the atmosphere. The atmospheric energy budget is clearly by definition a local balance of between the TOA and surface energy budget distribution, the most remarkable feature being the minimum in coincidence of the Equatorial Pacific.

Figure 2 about here

The meridional sections of climatological annual mean total, atmospheric and oceanic northward meridional enthalpy transports are shown in Figure 2. The figure layout follows from the classical-classic approach on meridional transports implied from budgets and their residuals (e.g. Trenberth et al. (2001); Lucarini and Ragone (2011); Lembo et al. (2016)). The transports are vanishing at the Poles-poles by definition, since the Carissimo et al. (1985) correction (cfr. in Equation 5) is applied -accounting to account for the effect of inconsistent model energy biases. The atmospheric transport is slightly asymmetric, being stronger in the SH-Southern Hemisphere (SH) than in the NH-Northern Hemisphere (NH). This is closely related to the asymmetry in the mean meridional circulation, being the latitude where the transport vanishes coincident with the annual mean position of the Inter-tropical Convergence Zone (ITCZ), slightly north of the Equator (Schneider et al., 2014; Adam et al., 2016). The atmospheric transport peaks at about 40 degrees of latitude in both hemispheres, slightly more poleward in the NH. The peaks mark the regions where baroclinic eddies are mainly responsible of become key in transporting energy poleward, and the zonal mean divergence of moist static energy switches sign (positive toward the Equator, negative toward the Poles; Loeb et al. (2015), cfr. Loeb et al. (2015)). The oceanic transport is much less homogeneous than the atmosphere. The two peaks are located in coincidence of near the subtropical and midlatitudinal gyres, the second being smaller than the first. At mid-latitudes of the Southern Hemisphere a relative maximum is found, in some models (cfr. Figure 8) even denoting a counter-transport from the South Pole toward the Equator. This is a critical issue, evidencing that the reproduction of the Southern Ocean circumpolar current is a major source of uncertainty in climate models (Trenberth and Fasullo, 2010).

Figure 3 about here

Figure 3 shows the relation-relationship between atmospheric and oceanic peak magnitudes. CanESM2 exhibits a clear relation between the two quantities in the SH. A stronger oceanic peak corresponds to a weaker atmospheric peak, whereas the relation is less clear in the NH. The anticorrelation of oceanic and atmospheric peaks suggests that the compensation mechanism-theoretised by Stone (1978)-Bjerknes compensation mechanism (Bjerknes, 1969; Stone, 1978) is well reproduced by the model, confirming that the shape of the total meridional heat-enthalpy transports is constrained by geometric and astronomical factors. This is far from being a trivial argument, since changes in the meridional planetary albedo differences-can deeply-can affect the total enthalpy transports (Enderton and Marshall, 2009), as well as the ocean-atmosphere partitioning (Rose and Ferreira, 2013). These plots-facilitate-tool facilitates an evaluation of these arguments-patterns in different models and under different scenarios and forcings.

Figure 4 about here

Figure 4 shows the annual mean horizontal distribution of water vapor in the atmosphere and its zonal mean meridional northward transport. This point-of-view highlights the sources and sinks of humidity in the atmosphere, evidencing that most of the exchanges are effected over the oceans. The water mass (and similarly the latent heat) budget is relatively weak over most of

the continents with significant regional exceptions, notably the Amazon, Bengal and Indonesia, as well as parts of the western coast of the American continent. The water mass budget over these land areas is mainly negative, denoting an excess of precipitation ~~with respect compared~~ to evaporation. The zonal mean water mass transport (Fig. 4(b)) is mainly poleward, except in the SH (30S - Eq) and the NH tropics (10N - 30N), essentially diverging humidity in both directions from ~~both hemispheres~~ ~~oceanic subtropics~~. ~~Clearly, water~~ ~~oceanic subtropics in both hemisphere~~. Water mass (and similarly latent energy, not shown) is primarily advected toward the regions of deep convection, ~~i.e.~~ the ITCZ (slightly north of the Equator), ~~;~~ and secondarily toward both hemispheres extra-tropics, where moisture is provided for the baroclinic eddies (cfr. Cohen et al. (2000)).

Figure 5 about here

The Lorenz Energy Cycle for one year of CanESM2 model run (Figure 5) shows how the energetics of atmospheric dynamics are ~~dealt with in~~ ~~calculated by~~ the diagnostic tool. The reservoirs of Available Potential Energy (APE) and Kinetic Energy (KE) are shown in the blue boxes, separately accounting for the zonal mean, ~~for the~~ stationary eddies (eddies in the time averaged circulation) and ~~for the~~ transient eddies (departure from zonal and time mean). The spectral approach also allows ~~to consider the a~~ partition between planetary wavenumbers, synoptic wavenumbers and higher order eddies, ~~although it is~~ (not shown here).

Most of the energy is stored in the form of APE in the zonal mean flux, and to a lesser extent in the zonal mean kinetic energy. The zonal mean APE is almost instantly converted into eddy potential energy (mainly through meridional advection of sensible heat) and then into eddy kinetic energy (through vertical motions in eddies) by means of mid-latitude baroclinic instability, so the two conversion terms are ~~not surprisingly~~ ~~unsurprisingly~~ qualitatively similar. We also notice that the ~~eddy eddy~~ APE and KE reservoirs have similar magnitudes. As argued before (Li et al., 2007), this is a consequence of the tight relation between temperature perturbations to the zonal mean meridional profile and the eddy synoptic activity. CanESM2 (and other models as well, not shown) agrees well with observational-based datasets ~~on the fact~~ that the stationary eddies play a non-negligible role in the baroclinic-barotropic energy conversion (cfr. Ulbrich and Speth (1991)). As for the KE, the transient eddy reservoirs are about half of the zonal mean, with the stationary eddy playing a more marginal role. The barotropic conversion acts mainly by converting eddy ~~KE~~ into zonal mean KE (i.e. restoring the jet stream), but in part also converting APE into KE (or vice versa) in the zonal mean flow.

Compared to ~~Reanalysis~~ ~~reanalysis~~ datasets (e.g. Ulbrich and Speth (1991); Li et al. (2007); Kim and Kim (2013)), our approach ~~features~~ ~~calculates~~ more energy stored in the zonal mean reservoirs. This is consistent with previous findings (cfr. Boer and Lambert (2008); Marques et al. (2011)) ~~;~~ ~~mentioning as possible reason for that and is possibly due to~~ the well-known cold Pole bias and the consequently excessive speed of the jet stream. Besides the fact that pre-industrial conditions feature different conditions than the present-day, as shown in Table 7, another ~~possible~~ explanation is that, unlike previous results from Reanalysis, we only consider the tropospheric part of the Lorenz Energy Cycle. The conversions of APE to and from stationary eddies also diverge from Reanalyses (Kim and Kim, 2013), although the overall baroclinic conversion is consistent with them. For model inter-comparison in the next ~~Section~~ ~~section~~, we will ~~then~~ consider the sum of the stationary and transient eddies as a single eddy component. The KE-APE conversion in the zonal mean flow is problematic, with CanESM2 having an opposite sign ~~as Reanalyses~~, ~~although also these observational-based products to the measurements made for reanalyses, although they~~

also appear to have certain inconsistencies (Kim and Kim, 2013).

Figure 6 about here

Results obtained from the indirect method for MEP with CanESM2 model are shown in Figure 6, ~~which shows the denoting~~ climatological annual mean maps of the vertical (a) and horizontal ~~components~~ component (b). Two different color maps have
5 been used, in order to emphasize that, ~~despite~~ although the vertical component has smaller maximum values than the horizontal component, it is positive almost everywhere (cfr. Lucarini et al. (2011)). As mentioned in Sect. 3.4.2, the local value of the horizontal component is not meaningful per se, and this component ~~shall~~ should only be addressed globally. Figure 6b is meant to describe an entropy flux divergence from the Tropics, particularly the Indian-Pacific warm pool, toward the high latitudes, roughly reflecting the atmospheric meridional enthalpy transport described in Fig. 2. This ~~allows to provide~~ provides a link
10 between entropy production and the meridional enthalpy transports, with the null isentrope delimiting the areas of enthalpy divergence from those of enthalpy convergence (cfr. Loeb et al. (2015)). The vertical entropy production features its highest values where the evaporation is most intense (cfr. Figure 4(a)). ~~On the contrary,~~ By contrast, the lowest values are found over the continents and the regions of subsidence in the atmospheric meridional circulation. ~~In other words, the~~ The vertical component is indicative of a local budget, in which atmospheric columns are weakly coupled with each other and the mixing occurs
15 on the vertical (Lucarini and Pascale, 2014).

5 Multi-model inter-comparison and changes across different scenarios

We now focus on comparing a 7-members multi-model ensemble from CMIP5 Project under three different scenarios: "piControl" (piC), denoting pre-industrial conditions, "historical" (hist), i.e. a realistic forcing evolution for the 1870-2005 period, and "rcp8.5", representing the 2005-2100 evolution of GHG forcing under business-as-usual emission scenario (in other words, a
20 ~~8.5 W/m^2~~ W/m^2 forcing by the end of the 21th Century). For the "piC scenario, 20-years periods, not necessarily overlapping, have been considered for each model, for hist scenario the 1981-2000 period, for rcp8.5 the 2081-2100 period. The ~~choice of the 7 models and the 20-years periods~~ 20-years periods chosen are motivated by the availability of ~~the CMIP5 model outputs for the three experiments~~ model outputs on the DKRZ ESG node. ~~It can be said that~~ Furthermore, this time length reflects the typical range for decadal climate predictions, as indicated in the Intergovernmental Panel on Climate Change (2013) report, so
25 it ~~well fits~~ aligns with our aim to describe the mean state and inter-annual variability of the climate system ~~with under~~ different conditions. A summary of the main global metrics described in Sect. 3 is reported in Tables 2-4.

Table 2 about here

Table 3 about here

Table 4 about here

30 Figure 7 about here

Figure 8 about here

Figure 9 about here

Figure 10 about here

5.1 Energy and water mass budgets, meridional enthalpy transports

As shown in the first two columns of Table 2, there is a large ~~bias-on-imbalance in~~ the TOA energy budget under unforced piC conditions. Such ~~bias-imbalance~~ sums up with the ~~bias-on~~-atmospheric energy budget ~~imbalance~~ (not shown, see Eq. 3) resulting in the ~~bias-on~~-surface energy budget estimates (the multi-model mean value for R_t being $0.21 \text{ } \cancel{W/m^2} W/m^2$, for F_s $0.73 \text{ } \cancel{W/m^2} W/m^2$). Some clear outliers are found, having either negative (BNU) or positive (MIR-C) values. This ~~bias-imbalance~~ is the signature of the well-known model drift in climate models (Gupta et al., 2013). The fact that ~~the-bias-it~~ is larger on surface budgets is explained by the fact that (cfr. Mauritsen et al. (2012); Hourdin et al. (2017)) models are generally tuned in order to achieve vanishing TOA budgets, whereas surface energy budgets are ~~in-most-cases-left-as-free-parameters-often-un-tuned~~. Panels a-c of Figure 7 emphasize how these biases are relevant with respect to the inter-annual variability of the budgets (computed as the standard deviation of the annual mean values). The atmospheric inter-annual variability is roughly one order of magnitude smaller (about $0.1 \text{ } \cancel{W/m^2} W/m^2$) than the variability ~~on-in-the~~ TOA and surface budgets, emphasizing that the changes in the overall energy imbalance are transferred to a large extent into the ocean (cfr. also Fig. 7d). The ~~model-agreement-inter-model-spread~~ on net energy fluxes is similar at the surface and at TOA, except for two models (BNU and MIR5) exhibiting ~~very-large-biases-a-very-large-imbalance~~ in the atmosphere, which ~~then-reflect-in-TOA-biases-A-closer-look-on-the-is-then-reflected-in-TOA-imbalance~~. There is limited correlation between surface and atmospheric ~~biases-suggests-that-they-are-quite-uncorrelated-with-each-other-imbbalances~~. The inter-annual variability is roughly the same order of magnitude as the ~~biases-imbbalances~~, both in the atmosphere and at the surface/TOA (about $0.20\text{-}0.25 \text{ } \cancel{W/m^2} W/m^2$).

As a consequence of a ~~transient-time-varying~~ GHG forcing, the TOA imbalance ~~is-found-to-increase-increases~~ (cfr. Tables 3 and 4). By the end of the historical period, most models agree on ~~showing~~-a positive imbalance (with respect to unforced biased conditions), ranging between 0.2 and $0.7 \text{ } \cancel{W/m^2} W/m^2$, although still in the range of the inter-annual variability (cfr. Figure 7a). The imbalance is much stronger by the end of the RCP8.5 period, peaking at $2.8 \text{ } \cancel{W/m^2} W/m^2$ (net of the bias, i.e. the value of the imbalance in piC scenario) in MIR-C. The surface imbalance appears to increase consistently with the TOA imbalance.

Figure 9 shows the difference in the ~~zonal mean~~ meridional latent heat transport between the Equator and 10N. As mentioned in Sect. 4 (~~cfr.-see~~ Fig. 4(b)), this is a measure of the moisture convergence toward the ITCZ. There is quite large uncertainty on its value in piC scenario, ranging between 1 and $3 \text{ } \cancel{PW} PW$. In all models the convergence is found to increase by up to $1 \text{ } \cancel{PW} PW$ between piC and RCP8.5. Even though ~~it-is~~ beyond the scope of this work, ~~one-may-argue-that-this-is-this-may-be~~ a robust estimate of the intensity of the uplifting branch, and to some extent of the intensity of the Hadley circulation.

Table 5 denotes a discrepancy of several $\cancel{W/m^2} W/m^2$ in the atmospheric budgets over ocean and land, with a positive imbalance over the former, negative over the latter. Such a well-known imbalance (see Wild et al. (2015) for a review on the model perspective) is key to ~~understand-the-models-capability-probing-the-models-ability~~ to reproduce the hydrological cycle, which ~~can-be-ultimately-seen-as-mediates~~ the convergence of latent energy (cfr. Eq. 9) from oceans (where water mass evaporates) toward the continents (where a large part of it precipitates). Atmospheric ~~energy-and-latent-heat-and-latent-energy-land-ocean~~ asymmetries are compared in the first two columns of Table 5. Models that are relatively well balanced (Can2, IPSL-M, MPI-

LR and MPI-MR) also feature relatively similar ~~atmospheric and latent heat~~ asymmetries. These ~~asymmetries~~ are translated into land-ocean transports, ~~multiplying if~~ ocean and land fluxes ~~are multiplied~~ by their respective surface area. The two transports are ~~obviously required to equal each other, net of the~~ theoretically required to be of equal magnitude but opposite sign (see Table 6) ~~-A reference estimate for such a transport amounts to and are estimated to be close to 2.8 PW (Wild et al., 2015)~~

5 ~~-Only few models are found to stick to PW (Wild et al., 2015). Few models comply to~~ this basic energy conservation ~~rule~~ requirement (with Can2, IPSL-M, MPI-LR and MPI-MR performing better than the others), ~~featuring;~~ others feature differences up to 1.7 ~~PW-PW~~ for BNU. For the better performing models, ~~it can be noticed note~~ that a residual asymmetry holds, which is not attributable to the latent ~~heat energy~~ asymmetry (third column in Table 5). Those transports are directed from land toward the oceans and are interpreted as the land-ocean transports related to asymmetries in the sensible heat fluxes at the

10 surface. The ocean-land latent energy transport is found to increase in RCP8.5 by 0.4 – 0.8 ~~PW-PW~~. This can be interpreted as an increase in the strength of the hydrological cycle, in line with previous findings (e.g. Levang and Schmitt (2015)). Looking at individual components of the hydrological cycle, we find an increase both in evaporation over oceans and precipitation over land.

The mean meridional sections of total, atmospheric ~~vs. and~~ oceanic enthalpy transports for each model are shown in Figure 8 for

15 the piC conditions alone (cfr. Tables 2-4 for an overview of peak magnitude and position values). The choice of not showing hist and rcp85 is motivated by ~~previous arguments on~~ the insensitivity of enthalpy transports to the different forcing (in agreement with the theory by ~~Stone (1978) Bjerknes (1969); Stone (1978)~~, confirmed by previous findings on CMIP models behaviors in disparate forcings representative of ~~nowadays present-day~~ climate or future emission scenarios, e.g. ~~Lucarini et al. (2011); ?~~ Lucarini et al. (2011); Lembo et al. (2019a); Irving et al. (2019)). Models agree on the asymmetry in the atmospheric transport,

20 being stronger in the Southern Hemisphere than in the Northern Hemisphere (see Sect. 3). A significant source of uncertainty is the location of the zero-crossing, which is in some cases very close to the Equator or even displaced in the SH. This is an important metric for the strength and shape of the mean meridional circulation. Compared to the atmosphere, the uncertainty on oceanic enthalpy transports is much larger, especially in the Southern Hemisphere, with some models featuring a counter-transport toward the Equator. As already mentioned (see section 4), a typical source of uncertainty here is the relative maximum

25 in the Southern Hemisphere extra-tropics.

5.2 The intensity of the LEC and its components

Table 7 about here

~~We provide in Table 7 the values of the components for the 7 models in the three scenarios.~~ The eleventh column in Tables 2-4 evidences that the intensity of the Lorenz Energy Cycle is ~~to a large extent relatively~~ constant through the scenarios. Its value ~~is comprised ranges~~ between 2.2 ~~W/m²-W/m²~~ (Can2) and 1.1 ~~W/m²-W/m²~~ (MIR5 in rcp8.5 scenario). As mentioned

30 in Sect. 3.32, a major source of uncertainty is ~~given by~~ the way fields are vertically discretized in pressure levels across the different models. Some models (BNU, Can2, MPI-LR, MPI-MR) internally interpolate the fields ~~, in order to provide~~ meaningful to provide a value in each gridpoint at each level. Others have no values where the surface pressure is lower than the respective pressure level. This normally occurs over Antarctica and mountainous regions (most notably Himalaya and the

Rocky Mountains), where the surface pressure reaches values even lower than 700 hPa. Since the ~~Lorenz Energy Cycle-LEC~~ is computed on spectral fields, the original gridpoint fields have to be continuous, and ~~the gaps, if present, have to~~ any gap must be filled with a vertical interpolation. In order to do so, daily mean near-surface fields of zonal and meridional velocities, and near-surface temperatures are used. Near-surface velocities replace the gaps, whereas near-surface temperatures are interpolated

5 ~~on-the-vertical, considering assuming~~ a vertical profile of temperature reconstructed from barometric equations (cfr. Sect. 3.4.1). Despite the retrieved velocity and temperature profiles ~~are-being~~ qualitatively comparable with those from the internally interpolated models, the stationary eddy conversion terms are unreasonably weaker. This is not entirely surprising, since the interpolation mostly affects mountain regions over the mid-latitudinal continents, i.e. the regions where the orographically driven stationary planetary waves are generated.

10 ~~As mentioned, the~~ The intensity of the LEC is not ~~really-very~~ sensitive to the type of forcing. The result is partially in contrast with ~~with~~ a previous version of MPI-LR (Hernández-Deckers and von Storch, 2010) and ~~Reanalyses-reanalyses~~ (Pan et al., 2017), though the changes/trends in the APE-KE conversion (i.e. the intensity of the LEC) for both studies are not very strong. ~~Nevertheless, Table 7 evidences~~ Table 7 provides values of the reservoirs and conversion terms for the 7 models in the three scenarios, evidencing relevant changes in some of the ~~storage components. Not entering into details of transient vs. stationary~~

15 ~~eddies partitioning, we notice~~ reservoirs across the different scenarios. Note that the zonal mean APE is largely decreased from piC to rcp8.5. This is compensated by an increase of similar magnitude in the zonal mean KE (and, to a lesser extent, ~~of-in~~ the eddy KE). In other words, in the wake of an increasing GHG forcing, the share of kinetic energy to available potential energy is changed in favor of the first. The total (APE+KE) energy contained in all storage terms remains ~~to a large extent stationary~~ approximately constant, in agreement with Pan et al. (2017). These results are consistent with what previously found

20 by Hernández-Deckers and von Storch (2010) with a previous version of MPI-ESM-LR, and Veiga and Ambrizzi (2013) with a state-of-the-art version of MPI-ESM-MR. The zonal-mean APE reduction is consistent with a reduction of the meridional temperature gradient, ~~predominantly as~~ which is predominantly a consequence of high latitude warming amplification (cfr. also Li et al. (2007)). The increase in zonal mean kinetic energy reflects a strengthening of the tropospheric mid-latitude jet stream (~~consistently with what~~ consistent with what was previously found about the SH tropospheric jet, e.g. Wilcox et al. (2012)).

25 Less clear is the impact of climate change on the mid-latitudinal eddy activity. The (slight) increase in eddy kinetic energy may reflect an increased mid-latitudinal baroclinic eddy activity in the Pacific and Atlantic storm tracks (despite large ~~uncertainty~~ still holds on the extent to which models agree on such changes model uncertainty on this aspect, cfr. Intergovernmental Panel on Climate Change (2013)). This, together with the raise of the tropical tropopause (mainly as a consequence of surface warming, cfr. Lin et al. (2017)), may contrast the expected decrease in baroclinic eddy activity associated with a smaller meridional

30 temperature gradient ~~-In other words, this~~ due to polar amplification. This approach allows to straightforwardly associate the different response of the models to the increasing GHG with key aspects of the general circulation of the atmosphere.

5.3 Material entropy production in the two methods

The components of the material entropy production in the indirect and direct methods are closely related to each other and provide further insight into the interpretation of water mass, energy budgets and LEC results.

First, let us consider Table 8, summarizing Table 8 summarizes the main components of the material entropy production in the two methods. The most part of the material entropy production-MEP obtained with the direct method is associated with-related to the hydrological cycle, whereas the vertical component overcomes the horizontal component in the indirect method by an order of magnitude. We have already commented on this in Sect. 3.4.2 and 4. Here we notice that The previous arguments about the changes in intensity of the hydrological cycle and of the atmospheric circulation are here confirmedreflected here as well. The entropy production increases with increasing GHG forcing in all models (except BNU, that we have already noticedbeing strongly water mass and energy unbalanced). The increase in the MEP related to the hydrological cycle ranges between 2.4 $mW \cdot m^{-2} \times K^{-1}$ $mW/m^2 K^{-1}$ (MIR5) and 4.7 $mW \cdot m^{-2} \times K^{-1}$ $mW/m^2 K^{-1}$ (IPSL-M) from piC to RCP8.5, amounting to about a 10% increase. The vertical component increase ranges between 5.6 $mW \cdot m^{-2} \times K^{-1}$ $mW/m^2 K^{-1}$ (MPI-MR) and 7.3 mW/Km^2 $mW/m^2 K^{-1}$ (IPSL-M). Models showing larger increases in the hydrological-cycle-related MEP also feature stronger increase in the vertical component. In other words the vertical component, being which is a signature of MEP related to vertical updraft, mainly through uplift, especially deep tropical convective activity, is closely relate to the strength of the hydrological cycle.

Table 9 provides more insight into the components of the MEP related to the hydrological cycle (cfr. Equation 15). We notice here that the two main changes are on the On one hand the reduction in MEP due to less a decrease of snowfall precipitation (S_s). This reduction ranges between 3 $mW \cdot m^{-2} \times K^{-1}$ $mW/m^2 K^{-1}$ (MIR5) and 8.5 mW/Km^2 $mW/m^2 K^{-1}$ (BNU) from piC to rcp8.5, to which a reduction of less than 1 $mW \cdot m^{-2} \times K^{-1}$ $mW/m^2 K^{-1}$ must be added as a consequence as less snow meltingto reflect a reduction in snow melt (S_m). On the other hand a large increase in MEP due to rainfall precipitation (S_r) is found, ranging between 10.7 $mW \cdot m^{-2} \times K^{-1}$ $mW/m^2 K^{-1}$ (MIR5) and 32.1 $mW \cdot m^{-2} \times K^{-1}$ $mW/m^2 K^{-1}$ (IPSL-M) from piC to rcp8.5, generally overcoming the MEP reduction related to evaporation at the surface ($S_s S_e$). This increase can be interpreted in different ways, either as an increase in water mass which is precipitated, or in terms as of a lower working temperature for rain droplet formation (T_C). As a consequence of the water mass balance, the latent heat associated with evaporation and rainfall precipitation have to precipitation must equal each other (net of less the changes in latent heat related to snowfall precipitation and the marginal contribution by snow melting at the ground). We thus attribute such an increase in S_r to changes in T_C . This might also be indicative of larger rate of convective precipitation on stratiform precipitation, and might be investigated further.

Concerning the other terms of the material entropy productionMEP budget, the one related to the sensible heat fluxes at the surface is slightly reduced, whereas the kinetic energy dissipation term is increased. Given that the LEC intensity, from which the kinetic energy dissipation has been derived, is to a large extent stationary across the scenario, such change is not related to the intensification of the atmospheric circulation (as argued in see Sect. 5.2), rather attributable to the near-surface warming. Finally, the decrease in the horizontal component is in line with the decrease in the APE terms of the LEC (especially the zonal mean term), denoting a weaker heat convergence toward the high latitudes (mainly as a consequence of high latitudes amplified warmingthe polar amplification).

As a wholeIn total, the entropy production is found to increase with increasing GHG forcing (see Tables 2-4), both with via the indirect and the direct method. This is consistent with previous findings (Lucarini et al., 2014, 2011) ,pointing and points

at the role of latent heat release in convective processes ~~-, setting-up-in establishing~~ the response of the climate system. The reduction of the meridional enthalpy transports is also consistent with previous comparisons between dry and moist entropy fluxes (Laliberté and Pauluis, 2010), suggesting ~~a-limiting-role-of-that~~ the hydrological cycle ~~to-that~~ the efficiency of the atmospheric thermal engine (Laliberté et al., 2015).

5 5.4 Baroclinic efficiency and irreversibility

As a wrap-up of the various aspects touched in this section, we introduce two metrics, ~~the-so-called-~~ The first is the "baroclinic efficiency" (Lucarini et al., 2011):

$$\eta = \frac{T_E^> - T_E^<}{T_E^>} \quad (25)$$

, where $T_E^>$ and $T_E^<$ are the emission temperatures averaged in the domains defined by TOA net energy gain and net energy loss (cfr. Figure 1), respectively, ~~and-~~ The second is the "degree of irreversibility" (Lucarini et al., 2011):

$$\alpha = \frac{\overline{\dot{S}_{dir}^{mat}} - \overline{\dot{S}_k^{mat}}}{\overline{\dot{S}_k^{mat}}} \quad (26)$$

, i.e. the ~~ratio-ratio~~ of MEP from irreversible processes others than the kinetic energy dissipation to the MEP related to the kinetic energy dissipation alone. The first parameter accounts for the strength of the mean meridional circulation, driven by the differential thermal gradient. ~~In-other-words,-this-~~ This is a measure of the dry entropy fluxes related to the heat to work conversion associated with the existence of the LEC ~~-,i.e.-and constitutes~~ an upper limit to the efficiency of the atmospheric thermal engine. The second parameter accounts for the relevance of viscous dissipation compared to other non-viscous irreversible processes. This parameter is closely related to the Bejan number, which is widely ~~use-used~~ in thermodynamics for the study of heat transfers in a fluid (Awad, 2016).

Table 10 shows the results from the three scenarios for the seven models that have been considered. The baroclinic efficiency ranges between 0.051 (MIR-C) and 0.066 (IPSL-M) in piC. It undergoes a clear reduction in the wake of increasing forcing, as a consequence of the already discussed reduction in the heat convergence at high latitudes. Remarkably, the last 20 years of hist do not seem to be significantly different from piC, contrary to rcp8.5. The irreversibility, in turn, is generally increased, especially between hist and rcp8.5. As noted before, the hydrological cycle ~~has-an-increasing-importance-~~ plays an increasingly significant role in converting energy into the system in a warmer climate (cfr. also Laliberté and Pauluis (2010); Laliberté et al. (2015)). This reflects a less efficient meridional enthalpy transport from low to high latitudes ~~-,and-consequently-in-and~~ consequently a larger irreversibility of the system, as already argued in Lucarini et al. (2010a, b).

5.5 Links among the metrics

Figure 10 brings up some of the metrics discussed up until now for the piC scenario, ~~to-describe-to-what-extent-they-can-be-related-~~ showing how they relate with each other. The TOA and atmospheric budgets (Figure 10a) are ~~here-clearly-strongly~~ related, with positive/negative outliers determined by positive/negative biases in atmospheric budgets. The other models cluster

around vanishing atmospheric energy imbalances and slightly positive TOA imbalances, likely reflecting the oceanic model drift.

- Figure 10b shows that the uncertainty on the value of baroclinic efficiency is much smaller (10%) than the one related to the LEC (about 50%, even though the treatment of fields in pressure levels is here a critical issue, as discussed in Sect. 5.2). The two quantities are related through the meridional enthalpy transports (cfr. ~~Boschi et al. (2013)~~ Boschi et al. (2013); Lembo et al. (2019b)), which in mid-latitudes are mainly effected ~~through by~~ baroclinic eddies. As mentioned, ~~it can also be noticed that~~ the strength of the baroclinic conversions in the LEC ~~are themselves is itself~~ a measure of the strength of the LEC. One might argue that the baroclinic efficiency peaks for certain values of the LEC intensity, but a larger ensemble would be necessary to prove or disprove such hypothesis.
- Figure 10c shows the relation between horizontal and vertical components of the MEP computed with the indirect methods (cfr. Lucarini et al. (2014)). It can be noticed that larger/smaller values of the vertical components correspond to smaller/larger values of the horizontal component. Indeed, the overall MEP ranges between about 57 and 60 $mW * m^{-2} * K^{-1} mW/m^2 K^{-1}$, while the vertical component alone has a 4 $mW * m^{-2} * K^{-1} mW/m^2 K^{-1}$ uncertainty range, suggesting that ~~somewhat~~ a compensation mechanism somewhat occurs between the vertical and horizontal component, i.e. between the local MEP on the vertical (especially where convection occurs) and the convergence of heat towards the high latitudes.
- Finally, Figure 10d shows the relation between MEP computed via the direct and indirect methods, respectively. Besides one outlier (BNU) all models have values of the direct method about 40 $mW * m^{-2} * K^{-1} mW/m^2 K^{-1}$, whereas through the indirect method they cluster about 57.5 $mW * m^{-2} * K^{-1} mW/m^2 K^{-1}$. Comparison with explicit computations by Pascale et al. (2011) suggest that the discrepancy is mainly attributable to an insufficient representation of the MEP related to the hydrological cycle (which is expected, since some intermediate phase change processes are not taken into account ~~here~~) and to the kinetic energy dissipation (because unphysical processes cannot be accounted for here, given the different numerical schemes of the models, leading to up to 50% reduction of this term).
- Summarizing Across metrics, this analysis emphasizes that the distribution of different models ~~can be quite different from being is quite far from~~ Gaussian. The multi-model ensemble mean and variances ~~here used~~ used here are certainly useful criteria for assessing the model uncertainty, but ~~they shall be carefully used, always paying attention to the care should be taken on the choice of the members and the~~ behavior of each ~~member of the ensemble~~ of them.

6 Summary and Conclusions

- We have ~~here~~ presented TheDiaTo v1.0, a diagnostic tool for the study of different aspects of the thermodynamics of the climate system, with a focus on the atmosphere. The goal of this diagnostic tool is to support the development, ~~the~~ evaluation, and ~~the~~ intercomparison of climate models, and to help the investigation of the properties of the climate in past, current, and projected future conditions. The diagnostic tool ~~consists of few independent~~ is comprised of independent modules, accounting for: (1) the energy budgets and transports in the atmosphere, the oceans, and in the system as a whole, (2) the water mass and latent energy budgets and transports, (3) the Lorenz Energy Cycle (LEC), (4) the material entropy production ~~with the direct or via~~

the direct and indirect methods. Global metrics are provided for immediate comparison ~~among-between~~ different datasets. We provide some examples of practical use of the diagnostic tool. We have presented results obtained from a 20-years subset of CMIP5 model run under unforced pre-industrial conditions, and results from a 20-years multi-model ensemble in three different scenarios: unforced pre-industrial conditions (piC), the end of the historical period (hist) and the last 2 decades of the 21st Century with a business-as-usual scenario (rcp8.5). A summary of the metrics and of the comparisons between the results ~~obtain-obtained~~ across models and scenarios is given in Tables 2-4 and in Figure 10.

The energy and water mass budgets ~~have been computed locally, but from this the transports are~~ are computed locally and from these the transports inferred, providing information about the global scale circulation. Similarly, the material entropy production has been decomposed in a component which essentially accounts for a local budget on the vertical, and another one which accounts for the global meridional enthalpy transport. In other words, the metrics ~~here shown~~ link the local features of the climate to the global energy and mass exchange, allowing for the evaluation of the global impact of localized changes. We have shown how the tool can provide a comprehensive view of the dynamics of the climate system and its response to perturbations. ~~On one hand, it is possible to evaluate the~~ It facilitates the evaluation of the spatial distributions of model biases and their impacts ~~, on the other hand we can interpret the change of the~~ and the interpretation of the changing properties of the system with time in the reduced space defined by the considered metrics.

Apart from the specific - yet important - problem of ~~looking-lookng~~ into climate change scenarios, it is ~~thus also~~ straightforward to use the diagnostic tool ~~for the study of paleoclimate, investigation of tipping points, the to study paleoclimates, investigate tipping points and~~ study of the climate under varied astronomical factors ~~, or~~ chemical composition of the atmosphere. One can envision adapting the model for the analysis of the properties of Earth-like exoplanets.

The requirement of flexibility, which allows the tool to ~~work for~~ be easily applied to a large class of models, inevitably leads to ~~make some simplifying hypotheses on the system~~. The most relevant are the following: a) the quasi-steady-state assumption; b) the hydrostatic assumptions, as background to the LEC framework; c) the identification of the emission temperature as the characteristic temperature in the atmosphere, leading to the 2D formulation of the material entropy production with the indirect method. Other assumptions involve the analysis of the hydrological cycle, ~~as where~~ the latent heat of evaporation and solidification have been assumed ~~as to be~~ constant, even though their value depends on temperature and pressure. Further, it is worth noticing (Loeb et al., 2015) that the latent energy associated with snowfall melting over ~~sea-ice-ocean-free~~ sea-ice-covered regions is not accounted for in CMIP5 models. This accounts for about $0.1 - 0.5 \text{ W/m}^2$. Nevertheless, unlike the water mass budget the latent energy budget is not expected to be closed, since the surface melting is not taken into account (although it is considered for the entropy budget).

Thus far, we have pointed out that the thermodynamic point of view can be ~~clearly~~ linked to fundamental aspects of the atmospheric dynamics. We have ~~linked-related~~ the idea of a baroclinic heat engine (Barry et al., 2002) to the mechanical work carried in a Lorenz Energy Cycle (Lorenz, 1955), along the lines of what proposed by Lucarini et al. (2010a, 2014). A deeper insight into the energetics of the atmospheric dynamics would require an evaluation of the meridional mass streamfunctions. To this ~~aim end~~, an additional diagnostic tool for the streamfunctions is ~~being developed on~~ under development, which uses moist and dry isentropes, rather than ~~on~~ isobaric coordinates, following from Laliberté et al. (2015); Kjellsson (2015), ~~in order~~

to link the Lagrangian and the Eulerian point of view. ~~The extent to which this idea can be pursued obviously depends~~ This approach is contingent on the availability of model outputs. ~~Still, we~~ We have shown here how deeply is the hydrological cycle affected by changes in the mean state of the system, ~~and such so~~ an isentropic approach ~~allows for resolving~~ would allow to resolve the heat exchanges due to moist processes.

- 5 Another open issue is assessing the relevance of coarse graining for the results, not only on the material entropy production (as discussed by Lucarini and Pascale (2014)), but also in terms of LEC and efficiency (when it comes to the type of ~~verteial~~ vertical discretization of the model). On one hand, the method is being tested with 3D fields for the radiative fluxes. On the other hand, the impact of changing the temporal, vertical and horizontal resolution is being assessed through a number of dedicated sensitivity studies ~~with~~ in an intermediate complexity atmospheric model.

10 -

Code and data availability. The diagnostics are part of the ESMValTool community diagnostics (v2.0). The latest release of the tool is available for download at www.doi.org/10.17874/ac8548f0315. CMIP5 data have been gathered from the ESGF node at DKRZ, publicly available upon registration at: <https://esgf-data.dkrz.de/projects/esgf-dkrz/>. The stand-alone version of TheDiaTo, v 1.0, is available on the GitHub repository: https://github.com/ValerioLembo/TheDiaTo_v1.0.git.

15 **Appendix A: Sources, sinks and conversion terms of the Lorenz Energy Cycle**

A1 Symbols and Definitions

- c_p = specific heat at constant pressure
- g = gravity
- p = pressure
- 20 – r = Earth's radius
- t = time
- T = temperature
- T_V = virtual temperature
- u = zonal velocity
- 25 – v = meridional velocity
- ω = vertical velocity

$$- \gamma = -\frac{R}{p} \left(\frac{\partial}{\partial p} [\bar{T}] - \frac{\partial}{\partial p} \frac{[\bar{T}]}{p} \right)^{-1}$$

– \bar{x} = time mean of x

– x' = deviation from time mean

– x = global horizontal mean

5 – $[x]$ = zonal mean

– x^* = deviation from zonal mean

A2 Storage terms

$$- \cancel{ZPE} = \frac{\gamma}{2g} ([\bar{T}] - \{\bar{T}\})^2 \quad \cancel{PZ} = \frac{\gamma}{2g} ([\bar{T}] - \{\bar{T}\})^2$$

$$- EPE = \frac{\gamma}{2g} ([\bar{T}^{*2}] + [\bar{T}'^2])$$

$$10 - \cancel{ZKE} = \frac{1}{2g} ([\bar{u}]^2 + [\bar{v}]^2) \quad \cancel{KZ} = \frac{1}{2g} ([\bar{u}]^2 + [\bar{v}]^2)$$

$$- EKE = \frac{1}{2g} [\bar{u}^{*2} + \bar{v}^{*2} + \bar{u}'^2 + \bar{v}'^2]$$

The contribution of eddies to APE (EPE) and KE (EKE) consists of two terms, the first one is the term accounting for stationary eddies (ASE and KSE in the diagrams of Figure 5, respectively). The second term accounts for the transient eddies (ATE and KTE in the diagrams of Figure 5, respectively). A discussion on the derivation of these terms is found in

15 [Ulbrich and Speth \(1991\)](#).

A3 Conversion terms

–

$$C_A = -\frac{\gamma}{g} \left\{ \frac{\partial [\bar{T}]}{r \partial \phi} [\bar{v}' T' + \bar{v}^* \bar{T}^*] + [\bar{\omega}' T' + \bar{\omega}^* \bar{T}^*] \left(\frac{\partial}{\partial p} ([\bar{T}] - \{\bar{T}\}) - \frac{R}{p c_p} ([\bar{T}] - \{\bar{T}\}) \right) \right\} +$$

$$+ \frac{\gamma}{g} \left\{ + \bar{u}' T'^* \frac{1}{r \cos \phi} \frac{\partial \bar{T}^*}{\partial \lambda} + \bar{v}' T'^* \frac{\partial \bar{T}^*}{r \partial \phi} \right\}$$

$$- C_Z = -\frac{R}{gp} ([\bar{\omega}] - \{\bar{\omega}\}) ([\bar{T}_v] - \{\bar{T}_v\})$$

$$- C_E = -\frac{R}{gp} [\bar{\omega}^* \bar{T}_v^* + \bar{\omega}' T_v']$$

$$\begin{aligned}
C_K = & -\frac{1}{g} \left\{ \left(\frac{\partial [\bar{u}]}{r \partial \phi} + [\bar{u}] \frac{\tan \phi}{r} \right) [\bar{u}^* \bar{v}^* + \overline{u'v'}] + \frac{\partial [\bar{v}]}{r \partial \phi} [\bar{v}^* \bar{v}^* + \overline{v'v'}] \right. \\
& - \frac{\tan \phi}{r} [\bar{v}][\bar{u}^* \bar{u}^* + \overline{u'u'}] + \frac{\partial [\bar{u}]}{\partial p} [\bar{\omega}^* \bar{u}^* + \overline{\omega'u'}] + \frac{\partial [\bar{v}]}{\partial p} [\bar{\omega}^* \bar{v}^* + \overline{\omega'v'}] + \left. \right\} + \\
& + \frac{1}{g} \left\{ \overline{u'u'}^* \frac{1}{r \cos \phi} \frac{\partial \bar{u}^*}{\partial \lambda} + \overline{u'v'}^* \left(\frac{\partial \bar{u}^*}{r \partial \phi} + \bar{u}^* \frac{\tan \phi}{r} + \frac{1}{r \cos \phi} \frac{\partial \bar{v}^*}{\partial \lambda} \right) + \right. \\
& + \left. \overline{v'v'}^* \frac{\partial \bar{v}^*}{\partial \phi} - [\overline{u'u'}]^* \bar{v}^* \frac{\tan \phi}{r} \right\}
\end{aligned}$$

The curly brackets in C_A and C_K emphasize that the diagnostic module is able to distinguish between two components, one dealing with the conversion from/to zonal mean flow to/from eddy flow (first bracket), the other one dealing with the conversion among eddies (second bracket).

The source and sink terms, i.e. the generation/dissipation of APE and KE, are computed as residuals of the conversion terms at each reservoir.

Competing interests. The authors declare that they have no conflict of interest.

Author contributions. Valerio Lembo implemented the new direct method for the material entropy production, wrote the code and the text of this paper. Frank Lunkeit implemented the first version of the LEC computation and supervised the whole code. Valerio Lucarini designed the diagnostic tool with its module partitioning and substantially contributed to the manuscript, in particular in the Introduction and Conclusions sections.

Acknowledgements. We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP. For CMIP the U.S. Department of Energy's Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led the development of software infrastructure in partnership with the Global Organization for Earth System Science Portals. We are also grateful Goodwin Gibbins for a careful proof reading. Valerio Lembo was supported by the Collaborative Research Centre TRR181 "Energy Transfers in Atmosphere and Ocean" funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), project No. 274762653. Valerio Lucarini was partially supported by the SFB/Transregio TRR181 project and by the EU ~~Horizon2020 Blue-Action-Project~~ Horizon 2020 Blue-Action Project (Grant No. 727852) and EU Horizon 2020 CRESCENDO Project (Grant No. 641816).

References

- Adam, O., Schneider, T., Brient, F., and Bischoff, T.: Relation of the double-ITCZ bias to the atmospheric energy budget in climate models, *Geophysical Research Letters*, 43, 7670–7677, <https://doi.org/10.1002/2016GL069465>, 2016.
- 5 Allan, R. P., Liu, C., Loeb, N. G., Palmer, M. D., Roberts, M., Smith, D. M., and Vidale, P. L.: Changes in global net radiative imbalance 1985-2012, *Geophysical Research Letters*, 41, 5588–5598, <https://doi.org/10.1002/2014GL060962>, 2014.
- Ambaum, M. H. P.: *Thermal Physics of the Atmosphere*, Wiley-Blackwell, <https://doi.org/10.1002/9780470710364>, 2010.
- Awad, M. M.: The science and the history of the two Bejan numbers, *International Journal of Heat and Mass Transfer*, 94, 101 – 103, <https://doi.org/https://doi.org/10.1016/j.ijheatmasstransfer.2015.11.073>, 2016.
- 10 Bannon, P. R.: Entropy Production and Climate Efficiency, *Journal of the Atmospheric Sciences*, 72, 3268–3280, <https://doi.org/10.1175/JAS-D-14-0361.1>, 2015.
- Bannon, P. R. and Lee, S.: Toward Quantifying the Climate Heat Engine: Solar Absorption and Terrestrial Emission Temperatures and Material Entropy Production, *Journal of the Atmospheric Sciences*, 74, 1721–1734, <https://doi.org/10.1175/JAS-D-16-0240.1>, 2017.
- Barry, L., Craig, G. C., and Thuburn, J.: Poleward heat transport by the atmospheric heat engine, *Nature*, 415, 774–777, <https://doi.org/10.1038/415774a>, 2002.
- 15 Bjerknes, J.: Atmospheric Teleconnections from the Equatorial Pacific, *Monthly Weather Review*, 97, 163–172, [https://doi.org/10.1175/1520-0493\(1969\)097<0163:ATFTEP>2.3.CO;2](https://doi.org/10.1175/1520-0493(1969)097<0163:ATFTEP>2.3.CO;2), 1969.
- Boer, G. J. and Lambert, S.: The energy cycle in atmospheric models, *Climate Dynamics*, 30, 371–390, <https://doi.org/10.1007/s00382-007-0303-4>, 2008.
- 20 Boschi, R., Lucarini, V., and Pascale, S.: Bistability of the climate around the habitable zone: A thermodynamic investigation, *Icarus*, 226, 1724–1742, <https://doi.org/10.1016/j.icarus.2013.03.017>, 2013.
- Buck, A. L.: New Equations for Computing Vapor Pressure and Enhancement Factor, *Journal of Applied Meteorology*, 20, 1527–1532, [https://doi.org/10.1175/1520-0450\(1981\)020<1527:NEFCVP>2.0.CO;2](https://doi.org/10.1175/1520-0450(1981)020<1527:NEFCVP>2.0.CO;2), 1981.
- Caballero, R. and Langen, P. L.: The dynamic range of poleward energy transport in an atmospheric general circulation model, *Geophysical Research Letters*, 32, L02 705, <https://doi.org/10.1029/2004GL021581>, 2005.
- 25 Carissimo, B. C., Oort, A. H., and Vonder Haar, T. H.: Estimating the Meridional Energy Transports in the Atmosphere and Ocean, [https://doi.org/10.1175/1520-0485\(1985\)015<0082:ETMETI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1985)015<0082:ETMETI>2.0.CO;2), 1985.
- CDO: Climate Data Operators, <http://www.mpimet.mpg.de/cdo>, 2015.
- Cohen, J. L., Salstein, D. A., and Rosen, R. D.: Interannual Variability in the Meridional Transport of Water Vapor, *Journal of Hydrometeorology*, 1, 547–553, [https://doi.org/10.1175/1525-7541\(2000\)001<0547:IVITMT>2.0.CO;2](https://doi.org/10.1175/1525-7541(2000)001<0547:IVITMT>2.0.CO;2), 2000.
- 30 Demory, M.-E., Vidale, P. L., Roberts, M. J., Berrisford, P., Strachan, J., Schiemann, R., and Mizielinski, M. S.: The role of horizontal resolution in simulating drivers of the global hydrological cycle, *Climate Dynamics*, 42, 2201–2225, <https://doi.org/10.1007/s00382-013-1924-4>, 2014.
- Enderton, D. and Marshall, J.: Explorations of Atmosphere–Ocean–Ice Climates on an Aquaplanet and Their Meridional Energy Transports, *Journal of the Atmospheric Sciences*, 66, 1593–1611, <https://doi.org/10.1175/2008JAS2680.1>, 2009.
- Exarchou, E., Kuhlbrodt, T., Gregory, J. M., and Smith, R. S.: Ocean Heat Uptake Processes: A Model Intercomparison, *Journal of Climate*, 28, 887–908, <https://doi.org/10.1175/JCLI-D-14-00235.1>, 2015.

- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016a.
- 5 Eyring, V., Righi, M., Lauer, A., Evaldsson, M., Wenzel, S., Jones, C., Anav, A., Andrews, O., Cionni, I., Davin, E. L., Deser, C., Ehbrecht, C., Friedlingstein, P., Gleckler, P., Gottschaldt, K.-D., Hagemann, S., Juckes, M., Kindermann, S., Krasting, J., Kunert, D., Levine, R., Loew, A., Mäkelä, J., Martin, G., Mason, E., Phillips, A. S., Read, S., Rio, C., Roehrig, R., Senftleben, D., Sterl, A., van Ulft, L. H., Walton, J., Wang, S., and Williams, K. D.: ESMValTool (v1.0) – a community diagnostic and performance metrics tool for routine evaluation of Earth system models in CMIP, *Geoscientific Model Development*, 9, 1747–1802, <https://doi.org/10.5194/gmd-9-1747-2016>, 2016b.
- 10 Fischer, N. and Jungclauss, J. H.: Effects of orbital forcing on atmosphere and ocean heat transports in Holocene and Eemian climate simulations with a comprehensive Earth system model, *Climate of the Past*, 6, 155–168, <https://doi.org/10.5194/cp-6-155-2010>, 2010.
- Fraedrich, K. and Lunkeit, F.: Diagnosing the entropy budget of a climate model, *Tellus A*, 60, 921–931, <https://doi.org/10.1111/j.1600-0870.2008.00338.x>, 2008.
- 15 Frigg, R., Thompson, E., and Werndl, C.: Philosophy of Climate Science Part II: Modelling Climate Change, *Philosophy Compass*, 10, 965–977, <https://doi.org/10.1111/phc3.12297>, 2015.
- Gassmann, A.: A global hexagonal C-grid non-hydrostatic dynamical core (ICON-IAP) designed for energetic consistency, *Quarterly Journal of the Royal Meteorological Society*, 139, 152–175, <https://doi.org/10.1002/qj.1960>, 2013.
- Gassmann, A. and Herzog, H. J.: How is local material entropy production represented in a numerical model?, *Quarterly Journal of the Royal Meteorological Society*, 141, 854–869, <https://doi.org/10.1002/qj.2404>, 2015.
- 20 Goff, A. J.: Saturation pressure of water on the new Kelvin temperature scale, *Transactions of the American Society of Heating and Ventilating Engineers*, pp. 347–354, 1957.
- Goody, R.: Sources and sinks of climate entropy, *Quarterly Journal of the Royal Meteorological Society*, 126, 1953–1970, <https://doi.org/10.1002/qj.49712656619>, <http://doi.wiley.com/10.1002/qj.49712656619>, 2000.
- Gupta, A. S., Jourdain, N. C., Brown, J. N., and Monselesan, D.: Climate drift in the CMIP5 models, *Journal of Climate*, 26, 8597–8615, https://doi.org/10.1175/JCLI_D_12_00521.1, 2013.
- 25 Hansen, J., Sato, M., Kharecha, P., and Von Schuckmann, K.: Earth’s energy imbalance and implications, *Atmospheric Chemistry and Physics*, 11, 13 421–13 449, <https://doi.org/10.5194/acp-11-13421-2011>, 2011.
- Hartmann, D. L.: *Global Physical Climatology*, Academic Press, <https://books.google.com/books?id=Zi1coMyhlHoC{&}pgis=1>, 1994.
- Held, I. M. and Soden, B. J.: Robust Responses of the Hydrological Cycle to Global Warming, *Journal of Climate*, 19, 5686–5699, <https://doi.org/10.1175/JCLI3990.1>, 2006.
- 30 Herbert, C., Paillard, D., and Dubrulle, B.: Entropy production and multiple equilibria: the case of the ice-albedo feedback, *Earth System Dynamics*, 2, 13–23, <https://doi.org/10.5194/esd-2-13-2011>, 2011.
- Hernández-Deckers, D. and von Storch, J.-S.: Energetics Responses to Increases in Greenhouse Gas Concentration, *Journal of Climate*, 23, 3874–3887, <https://doi.org/10.1175/2010JCLI3176.1>, 2010.
- 35 Hobbs, W., Palmer, M. D., and Monselesan, D.: An Energy Conservation Analysis of Ocean Drift in the CMIP5 Global Coupled Models, *Journal of Climate*, 29, 1639–1653, <https://doi.org/10.1175/JCLI-D-15-0477.1>, 2016.
- Hourdin, F., Mauritsen, T., Gettelman, A., Golaz, J.-C., Balaji, V., Duan, Q., Folini, D., Ji, D., Klocke, D., Qian, Y., Rauser, F., Rio, C., Tomassini, L., Watanabe, M., and Williamson, D.: The Art and Science of Climate Model Tuning, *Bulletin of the American Meteorological Society*, 98, 589–602, <https://doi.org/10.1175/BAMS-D-15-00135.1>, 2017.

- Intergovernmental Panel on Climate Change: Climate Change 2013: The Physical Science Basis, Press, Cambridge University, Cambridge Mass., <https://doi.org/10.1017/CBO9781107415324>, 2013.
- Irving, D. B., Wijffels, S., and Church, J. A.: Anthropogenic Aerosols, Greenhouse Gases, and the Uptake, Transport, and Storage of Excess Heat in the Climate System, *Geophysical Research Letters*, 46, 4894–4903, <https://doi.org/10.1029/2019GL082015>, 2019.
- Johnson, D. R.: “General Coldness of Climate Models” and the Second Law: Implications for Modeling the Earth System, *Journal of Climate*, 10, 2826–2846, [https://doi.org/10.1175/1520-0442\(1997\)010<2826:GCOCMA>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<2826:GCOCMA>2.0.CO;2), 1997.
- Kiehl, J. T. and Trenberth, K. E.: Earth’s Annual Global Mean Energy Budget, *Bulletin of the American Meteorological Society*, 78, 197–208, [https://doi.org/10.1175/1520-0477\(1997\)078<0197:EAGMEB>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<0197:EAGMEB>2.0.CO;2), 1997.
- 10 Kim, Y. H. and Kim, M. K.: Examination of the global lorenz energy cycle using MERRA and NCEP-reanalysis 2, *Climate Dynamics*, 40, 1499–1513, <https://doi.org/10.1007/s00382-012-1358-4>, 2013.
- Kjellsson, J.: Weakening of the global atmospheric circulation with global warming, *Climate Dynamics*, 45, 975–988, <https://doi.org/10.1007/s00382-014-2337-8>, 2015.
- Kleidon, A.: Nonequilibrium thermodynamics and maximum entropy production in the Earth system: applications and implications., *Die Naturwissenschaften*, 96, 653–677, <https://doi.org/10.1007/s00114-009-0509-x>, <http://link.springer.com/10.1007/s00114-009-0509-x>, 2009.
- 15 Kleidon, A. and Lorenz, R.: Non-equilibrium Thermodynamics and the Production of Entropy, Springer-Verlag, Berlin/Heidelberg, 1 edn., https://doi.org/10.1007/11672906_1, http://link.springer.com/10.1007/11672906_{_}1, 2004.
- Knietzsch, M.-A., Schröder, A., Lucarini, V., and Lunkeit, F.: The impact of oceanic heat transport on the atmospheric circulation, *Earth System Dynamics*, 6, 591–615, <https://doi.org/10.5194/esd-6-591-2015>, 2015.
- Kunz, T., Fraedrich, K., and Kirk, E.: Optimisation of simplified GCMs using circulation indices and maximum entropy production, *Climate Dynamics*, 30, 803–813, <https://doi.org/10.1007/s00382-007-0325-y>, <https://doi.org/10.1007/s00382-007-0325-y>, 2008.
- Laliberté, F. and Pauluis, O.: Winter intensification of the moist branch of the circulation in simulations of 21st century climate, *Geophysical Research Letters*, 37, 1–6, <https://doi.org/10.1029/2010GL045007>, 2010.
- 25 Laliberté, F., Zika, J., Mudryk, L., Kushner, P. J., Kjellsson, J., and Doos, K.: Constrained work output of the moist atmospheric heat engine in a warming climate, *Science*, 347, 540–543, <https://doi.org/10.1126/science.1257103>, 2015.
- Lawrence, M. G.: The Relationship between Relative Humidity and the Dewpoint Temperature in Moist Air: A Simple Conversion and Applications, *Bulletin of the American Meteorological Society*, 86, 225–234, <https://doi.org/10.1175/BAMS-86-2-225>, 2005.
- L’Ecuyer, T. S., Beaudoin, H. K., Rodell, M., Olson, W., Lin, B., Kato, S., Clayson, C. A., Wood, E., Sheffield, J., Adler, R., Huffman, G., Bosilovich, M., Gu, G., Robertson, F., Houser, P. R., Chambers, D., Famiglietti, J. S., Fetzer, E., Liu, W. T., Gao, X., Schlosser, C. A., Clark, E., Lettenmaier, D. P., Hilburn, K., L’Ecuyer, T. S., Beaudoin, H. K., Rodell, M., Olson, W., Lin, B., Kato, S., Clayson, C. A., Wood, E., Sheffield, J., Adler, R., Huffman, G., Bosilovich, M., Gu, G., Robertson, F., Houser, P. R., Chambers, D., Famiglietti, J. S., Fetzer, E., Liu, W. T., Gao, X., Schlosser, C. A., Clark, E., Lettenmaier, D. P., and Hilburn, K.: The Observed State of the Energy Budget in the Early Twenty-First Century, *Journal of Climate*, 28, 8319–8346, <https://doi.org/10.1175/JCLI-D-14-00556.1>, 2015.
- 30 Lembo, V., Folini, D., Wild, M., and Lionello, P.: Energy budgets and transports : global evolution and spatial patterns during the 20th Century as estimated in two AMIP-like experiments, *Climate Dynamics*, <https://doi.org/10.1007/s00382-016-3173-9>, 2016.
- Lembo, V., Folini, D., Wild, M., and Lionello, P.: Inter-hemispheric differences in energy budgets and cross-equatorial transport anomalies during the 20th century, *Climate Dynamics*, 53, 115–135, <https://doi.org/10.1007/s00382-018-4572-x>, <http://link.springer.com/10.1007/s00382-018-4572-x>, 2019a.

- Lembo, V., Messori, G., Graversen, R., and Lucarini, V.: Spectral Decomposition and Extremes of Atmospheric Meridional Energy Transport in the Northern Hemisphere Midlatitudes, *Geophysical Research Letters*, 46, 2019GL082105, <https://doi.org/10.1029/2019GL082105>, 2019b.
- 5 Levang, S. J. and Schmitt, R. W.: Centennial changes of the global water cycle in CMIP5 models, *Journal of Climate*, 28, 6489–6502, <https://doi.org/10.1175/JCLI-D-15-0143.1>, 2015.
- Levitus, S., Antonov, J. I., Boyer, T. P., Baranova, O. K., Garcia, H. E., Locarnini, R. A., Mishonov, A. V., Reagan, J. R., Seidov, D., Yarosh, E. S., and Zweng, M. M.: World ocean heat content and thermosteric sea level change (0–2000 m), 1955–2010, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051106>, <http://doi.wiley.com/10.1029/2012GL051106>, 2012.
- 10 Li, L., Ingersoll, A. P., Jiang, X., Feldman, D., and Yung, Y. L.: Lorenz energy cycle of the global atmosphere based on reanalysis datasets, *Geophysical Research Letters*, 34, 1–5, <https://doi.org/10.1029/2007GL029985>, 2007.
- Liepert, B. G. and Previdi, M.: Inter-model variability and biases of the global water cycle in CMIP3 coupled climate models, *Environmental Research Letters*, 7, 14006, <https://doi.org/10.1088/1748-9326/7/1/014006>, 2012.
- Lin, P., Paynter, D., Ming, Y., and Ramaswamy, V.: Changes of the Tropical Tropopause Layer under Global Warming, *Journal of Climate*, 30, 1245–1258, <https://doi.org/10.1175/JCLI-D-16-0457.1>, 2017.
- 15 Loeb, N. G., Wielicki, B. a., Doelling, D. R., Smith, G. L., Keyes, D. F., Kato, S., Manalo-Smith, N., and Wong, T.: Toward optimal closure of the Earth’s top-of-atmosphere radiation budget, *Journal of Climate*, 22, 748–766, <https://doi.org/10.1175/2008JCLI2637.1>, 2009.
- Loeb, N. G., Lyman, J. M., Johnson, G. C., Allan, R. P., Doelling, D. R., Wong, T., Soden, B. J., and Stephens, G. L.: Observed changes in top-of-the-atmosphere radiation and upper-ocean heating consistent within uncertainty, *Nature Geoscience*, 5, 1–4, <https://doi.org/10.1038/ngeo1375>, 2012.
- 20 Loeb, N. G., Wang, H., Cheng, A., Kato, S., Fasullo, J. T., Xu, K.-M., and Allan, R. P.: Observational constraints on atmospheric and oceanic cross-equatorial heat transports: revisiting the precipitation asymmetry problem in climate models, *Climate Dynamics*, 46, 3239–3257, <https://doi.org/10.1007/s00382-015-2766-z>, 2015.
- Lorenz, E. N.: Available Potential Energy and the Maintenance of the General Circulation, *Tellus*, 7, 157–167, <https://doi.org/10.1111/j.2153-3490.1955.tb01148.x>, 1955.
- 25 Lucarini, V.: Thermodynamic efficiency and entropy production in the climate system, *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, 80, 1–5, <https://doi.org/10.1103/PhysRevE.80.021118>, 2009.
- Lucarini, V.: Modeling Complexity: The Case of Climate Science, in: *Models, Simulations, and the Reduction of Complexity*, edited by Gähde, U., Hartmann, S., and Wolf, J. H., pp. 229–254, De Gruyter, Berlin, Boston, <https://doi.org/10.1515/9783110313680.229>, 2013.
- 30 Lucarini, V. and Fraedrich, K.: Symmetry breaking, mixing, instability, and low-frequency variability in a minimal Lorenz-like system, *Physical Review E*, 80, 026313, <https://doi.org/10.1103/PhysRevE.80.026313>, <https://link.aps.org/doi/10.1103/PhysRevE.80.026313>, 2009.
- Lucarini, V. and Pascale, S.: Entropy production and coarse graining of the climate fields in a general circulation model, *Climate Dynamics*, 43, 981–1000, <https://doi.org/10.1007/s00382-014-2052-5>, 2014.
- Lucarini, V. and Ragone, F.: Energetics of Climate Models: Net Energy Balance and Meridional Enthalpy Transport, *Reviews of Geophysics*, 49, <https://doi.org/10.1029/2009RG000323>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009RG000323>, 2011.
- 35 Lucarini, V., Fraedrich, K., and Lunkeit, F.: Thermodynamics of climate change: Generalized sensitivities, *Atmospheric Chemistry and Physics*, 10, 9729–9737, <https://doi.org/10.5194/acp-10-9729-2010>, 2010a.
- Lucarini, V., Fraedrich, K., and Lunkeit, F.: Thermodynamic analysis of snowball earth hysteresis experiment: Efficiency, entropy production and irreversibility, *Quarterly Journal of the Royal Meteorological Society*, 136, 2–11, <https://doi.org/10.1002/qj.543>, 2010b.

- Lucarini, V., Fraedrich, K., and Ragone, F.: New Results on the Thermodynamic Properties of the Climate System, *Journal of the Atmospheric Sciences*, 68, 2438–2458, <https://doi.org/10.1175/2011JAS3713.1>, 2011.
- Lucarini, V., Blender, R., Herbert, C., Ragone, F., Pascale, S., and Wouters, J.: Mathematical and physical ideas for climate science, *Reviews of Geophysics*, 52, 809–859, <https://doi.org/10.1002/2013RG000446>, 2014.
- Manabe, S. and Wetherald, R. T.: Thermal Equilibrium of the Atmosphere with a Given Distribution of Relative Humidity, *Journal of the Atmospheric Sciences*, 24, 241–259, [https://doi.org/10.1175/1520-0469\(1967\)024<0241:TEOTAW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1967)024<0241:TEOTAW>2.0.CO;2), 1967.
- Marques, C. A. F., Rocha, A., and Corte-Real, J.: Global diagnostic energetics of five state-of-the-art climate models, *Climate Dynamics*, 36, 1767–1794, <https://doi.org/10.1007/s00382-010-0828-9>, 2011.
- 10 Mauritsen, T., Stevens, B., Roeckner, E., Crueger, T., Esch, M., Giorgetta, M., Haak, H., Jungclaus, J., Klocke, D., Matei, D., Mikolajewicz, U., Notz, D., Pincus, R., Schmidt, H., and Tomassini, L.: Tuning the climate of a global model, *Journal of Advances in Modeling Earth Systems*, 4, <https://doi.org/10.1029/2012MS000154>, 2012.
- Novak, L. and Tailleux, R.: On the Local View of Atmospheric Available Potential Energy, *Journal of the Atmospheric Sciences*, 75, 1891–1907, <https://doi.org/10.1175/JAS-D-17-0330.1>, 2018.
- 15 Ozawa, H., Ohmura, A., Lorenz, R. D., and Pujol, T.: The second law of thermodynamics and the global climate system: A review of the maximum entropy production principle, *Reviews of Geophysics*, 41, 1018, <https://doi.org/10.1029/2002RG000113>, 2003.
- Paltridge, G.: Global dynamics and climate - a system of minimum entropy exchange, *Quarterly Journal of the Royal Meteorological Society*, 101, 475–484, <https://doi.org/10.1002/qj.49710142906>, 1975.
- Pan, Y., Li, L., Jiang, X., Li, G., Zhang, W., Wang, X., and Ingersoll, A. P.: Earth’s changing global atmospheric energy cycle in response to climate change, *Nature Communications*, 8, 14 367, <https://doi.org/10.1038/ncomms14367>, <http://www.nature.com/doi/10.1038/ncomms14367>, 2017.
- 20 Pascale, S., Gregory, J. M., Ambaum, M., and Tailleux, R.: Climate entropy budget of the HadCM3 atmosphere–ocean general circulation model and of FAMOUS, its low-resolution version, *Climate Dynamics*, 36, 1189–1206, <https://doi.org/10.1007/s00382-009-0718-1>, 2011.
- Pauluis, O.: Water Vapor and Mechanical Work: A Comparison of Carnot and Steam Cycles, *Journal of the Atmospheric Sciences*, 68, 91–102, <https://doi.org/10.1175/2010JAS3530.1>, 2011.
- 25 Pauluis, O. and Dias, J.: Satellite estimates of precipitation-induced dissipation in the atmosphere., *Science*, 335, 953–956, <https://doi.org/10.1126/science.1215869>, 2012.
- Pauluis, O. and Held, I. M.: Entropy Budget of an Atmosphere in Radiative–Convective Equilibrium. Part I: Maximum Work and Frictional Dissipation, *Journal of the Atmospheric Sciences*, 59, 125–139, [https://doi.org/10.1175/1520-0469\(2002\)059<0125:EBOAAI>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<0125:EBOAAI>2.0.CO;2), 2002.
- 30 Peixoto, J. P. and Oort, A. H.: *Physics of Climate*, New York, NY (United States); American Institute of Physics, <http://www.osti.gov/scitech/biblio/7287064>, 1992.
- Prigogine, I.: *Non-equilibrium statistical mechanics*, Wiley, New York, 1962.
- Raymond, D. J.: Sources and sinks of entropy in the atmosphere, *Journal of Advances in Modeling Earth Systems*, 5, 755–763, <https://doi.org/10.1002/jame.20050>, 2013.
- 35 Rodell, M., Beaudoin, H. K., L’Ecuyer, T. S., Olson, W. S., Famiglietti, J. S., Houser, P. R., Adler, R., Bosilovich, M. G., Clayson, C. A., Chambers, D., Clark, E., Fetzer, E. J., Gao, X., Gu, G., Hilburn, K., Huffman, G. J., Lettenmaier, D. P., Liu, W. T., Robertson, F. R., Schlosser, C. A., Sheffield, J., Wood, E. F., Rodell, M., Beaudoin, H. K., L’Ecuyer, T. S., Olson, W. S., Famiglietti, J. S., Houser, P. R., Adler, R., Bosilovich, M. G., Clayson, C. A., Chambers, D., Clark, E., Fetzer, E. J., Gao, X., Gu, G., Hilburn, K., Huffman, G. J.,

- Lettenmaier, D. P., Liu, W. T., Robertson, F. R., Schlosser, C. A., Sheffield, J., and Wood, E. F.: The Observed State of the Water Cycle in the Early Twenty-First Century, *Journal of Climate*, 28, 8289–8318, <https://doi.org/10.1175/JCLI-D-14-00555.1>, 2015.
- Rose, B. E. J. and Ferreira, D.: Ocean Heat Transport and Water Vapor Greenhouse in a Warm Equable Climate: A New Look at the Low Gradient Paradox, *Journal of Climate*, 26, 2117–2136, <https://doi.org/10.1175/JCLI-D-11-00547.1>, 2013.
- Schneider, T., Bischoff, T., and Haug, G. H.: Migrations and dynamics of the Intertropical Convergence Zone, *Nature*, 513, 45–53, <https://doi.org/10.1038/nature13636>, 2014.
- Smith, D. M., Allan, R. P., Coward, A. C., Eade, R., Hyder, P., Liu, C., Loeb, N. G., Palmer, M. D., Roberts, C. D., and Scaife, A. a.: Earth’s energy imbalance since 1960 in observations and CMIP5 models, *Geophysical Research Letters*, 42, 2014GL062669, <https://doi.org/10.1002/2014GL062669>, 2015.
- Stone, P. H.: Constraints on dynamical transports of energy on a spherical planet, *Dynamics of Atmospheres and Oceans*, 2, 123–139, [https://doi.org/10.1016/0377-0265\(78\)90006-4](https://doi.org/10.1016/0377-0265(78)90006-4), 1978.
- Tailleux, R.: Available Potential Energy and Exergy in Stratified Fluids, *Annual Review of Fluid Mechanics*, 45, 35–58, <https://doi.org/10.1146/annurev-fluid-011212-140620>, 2013.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- Terai, C. R., Caldwell, P. M., Klein, S. A., Tang, Q., and Branstetter, M. L.: The atmospheric hydrologic cycle in the ACME v0.3 model, *Climate Dynamics*, 50, 3251–3279, <https://doi.org/10.1007/s00382-017-3803-x>, 2018.
- Trenberth, K. E. and Fasullo, J. T.: Simulation of present-day and twenty-first-century energy budgets of the southern oceans, *Journal of Climate*, 23, 440–454, <https://doi.org/10.1175/2009JCLI3152.1>, 2010.
- Trenberth, K. E. and Solomon, A.: The Global Heat-Balance - Heat Transports in the Atmosphere and Ocean, *Climate Dynamics*, 10, 107–134, <https://doi.org/10.1007/Bf00210625>, 1994.
- Trenberth, K. E., Caron, J. M., and Stepaniak, D. P.: The atmospheric energy budget and implications for surface fluxes and ocean heat transports, *Climate Dynamics*, 17, 259–276, <https://doi.org/10.1007/PL00007927>, 2001.
- Ulbrich, U. and Speth, P.: The global energy cycle of stationary and transient atmospheric waves: Results from ECMWF analyses, *Meteorology and Atmospheric Physics*, 45, 125–138, <https://doi.org/10.1007/BF01029650>, 1991.
- Vannière, B., Demory, M.-E., Vidale, P. L., Schiemann, R., Roberts, M. J., Roberts, C. D., Matsueda, M., Terray, L., Koenigk, T., and Senan, R.: Multi-model evaluation of the sensitivity of the global energy budget and hydrological cycle to resolution, *Climate Dynamics*, 52, 6817–6846, <https://doi.org/10.1007/s00382-018-4547-y>, 2019.
- Veiga, J. A. P. and Ambrizzi, T.: A global and Hemispherical Analysis of the Lorenz Energetics Based on the Representative Concentration Pathways Used in CMIP5, *Advances in Meteorology*, 2013, 1–13, <https://doi.org/10.1155/2013/485047>, <http://www.hindawi.com/journals/amete/2013/485047/>, 2013.
- von Schuckmann, K., Palmer, M. D., Trenberth, K. E., Cazenave, A., Chambers, D., Champollion, N., Hansen, J., Josey, S. A., Loeb, N. G., Mathieu, P.-P., Meyssignac, B., and Wild, M.: An imperative to monitor Earth’s energy imbalance, *Nature Climate Change*, 6, 138–144, 2016.
- Wilcox, L. J., Charlton-Perez, A. J., and Gray, L. J.: Trends in Austral jet position in ensembles of high- and low-top CMIP5 models, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD017597>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2012JD017597>, 2012.

- Wild, M. and Liepert, B. G.: The Earth radiation balance as driver of the global hydrological cycle, *Environmental Research Letters*, 5, 25 203, <https://doi.org/10.1088/1748-9326/5/2/025203>, 2010.
- Wild, M., Long, C. N., and Ohmura, A.: Evaluation of clear-sky solar fluxes in GCMs participating in AMIP and IPCC-AR4 from a surface
5 perspective, *Journal of Geophysical Research*, 111, D01 104, <https://doi.org/10.1029/2005JD006118>, 2006.
- Wild, M., Folini, D., Schär, C., Loeb, N. G., Dutton, E. G., and König-Langlo, G.: The global energy balance from a surface perspective, *Climate Dynamics*, 40, 3107–3134, <https://doi.org/10.1007/s00382-012-1569-8>, 2013.
- Wild, M., Folini, D., Hakuba, M. Z., Schär, C., Seneviratne, S. I., Kato, S., Rutan, D., Ammann, C., Wood, E. F., and König-Langlo, G.:
10 The energy balance over land and oceans: an assessment based on direct observations and CMIP5 climate models, *Climate Dynamics*, 44, 3393–3429, <https://doi.org/10.1007/s00382-014-2430-z>, 2015.
- Zhang, Y., Gao, Z., Li, D., Li, Y., Zhang, N., Zhao, X., and Chen, J.: On the computation of planetary boundary-layer height using the bulk Richardson number method, *Geoscientific Model Development*, 7, 2599–2611, <https://doi.org/10.5194/gmd-7-2599-2014>, 2014.




Figure 1. Climatological annual mean maps of (a) TOA, (b) atmospheric and (c) surface energy budgets for CanESM2 model (in W/m^2). The fluxes are positive when entering the domain, negative when exiting the domain.




Figure 2. Climatological annual mean total (blue), atmospheric (orange) and oceanic (green) northward meridional enthalpy transports for 20 years of a pre-industrial CanESM2 model run (in W).




Figure 3. Scatter plots of 20 years pre-industrial CanESM2 annual mean atmospheric vs. oceanic peak magnitudes in the SH (a) and the NH (b) in W .




Figure 4. (a) Climatological annual mean water mass fluxes (in $kg \cdot m^{-2} \cdot s^{-1}$) and (b) annual mean northward meridional water mass transport (in $kg \cdot s^{-1}$) for a 20 years pre-industrial CanESM2 model run.






Figure 5. Diagram of Lorenz Energy Cycle (LEC) annual mean production, dissipation, storage and conversion terms for one year of pre-industrial CanESM2 model run. *AZ* denotes the APE reservoir in the zonal mean flow, *ASE* and *ATE* the APE associated with stationary and transient eddies, respectively. *KZ* denotes the KE associated with the zonal mean flow, *KSE* and *KTE* the KE associated with the stationary and transient eddies (cfr. A2). Reservoirs are displayed in units of $10^5 \text{ J} \cdot \text{m}^{-2} / \text{m}^2$, conversion terms as $\text{W} \cdot \text{m}^{-2}$.



(a)



(b)

Figure 6. Climatological annual mean maps of (a) vertical component of the material entropy production (in $\cancel{W}/m^2 * \cancel{K} \cancel{W}/m^2 K$), (b) horizontal component of the material entropy production (in $\cancel{W}/m^2 * \cancel{K} \cancel{W}/m^2 K$), for a 20 years pre-industrial CanESM2 model run.




Figure 7. Multi-model ensemble scatter plots of annual mean averaged quantities vs. inter-annual variability in the piC scenario for: (a) TOA energy budget, (b) Atmospheric energy budget, (c) Surface energy budget. Panel d shows the atmospheric energy budget vs. the surface energy budget, with whiskers denoting the inter-annual variability as in panels (b) and (c). Blue ellipses denote the σ standard deviation of the multi-model mean (denoted by the red dot). Model IDs are: 1. BNU, 2. Can2, 3. IPSL-M, 4. MIR5, 5. MIR-C, 6. MPI-LR, 7. MPI-MR.




Figure 8. Climatological annual mean (a) total, (b) atmospheric, (c) oceanic northward meridional ~~heat~~enthalpy transports for all models (in W).




Figure 9. Latent heat transport between the Equator and 10N for the 7 models in the three scenarios. The transport increases from piC to RCP8.5. Values are in W and are positive if northward directed.




Figure 10. Multi-model ensemble scatter plots from piC scenario for: (a) Atmospheric energy budget vs. TOA energy budget (both in W/m^2), (b) Baroclinic efficiency vs. LEC intensity (in W/m^2), (c) vertical component of the material entropy production vs. vertical component (together with iso-lines of total material entropy production with the indirect method) (in $\text{W/m}^2 \cdot \text{K}$), (d) Direct vs. indirect method for material entropy production (in $\text{W/m}^2 \cdot \text{K}$). Blue ellipses denote the σ standard deviation of the multi-model mean (denoted by the red dot). Model IDs are: 1. BNU, 2. Can2, 3. IPSL-M, 4. MIR5, 5. MIR-C, 6. MPI-LR, 7. MPI-MR.

Table 1. Variables used in the modules of the diagnostic tool

height	Name (CMOR)	Name (CMOR)	Time res. (min.)	Description	Time res. (min.)	Vert. res.	Energy budget	Energy budget	Hydrological cycle
rsdt		<u>Downward SW radiation at TOA</u>	monthly	2D	X				
rsut		<u>Upward SW radiation at TOA</u>	monthly	2D	X				
rlut		<u>Outgoing LW radiation at TOA</u>	monthly	2D	X				
rl ds		<u>Downward LW radiation at the surface</u>	monthly	2D	X				
rs ds		<u>Downward SW radiation at the surface</u>	monthly	2D	X				
rl us		<u>Upward LW radiation at the surface</u>	monthly	2D	X				
rs us		<u>Upward SW radiation at the surface</u>	monthly	2D	X				
h fls		<u>Latent heat turbulent flux at the surface</u>	monthly	2D	X				X
h fss		<u>Sensible heat turbulent flux at the surface</u>	monthly	2D	X				
pr ^[1]		<u>Rainfall precipitation flux</u>	monthly	2D					X
pr sn ^[1]		<u>Snowfall precipitation flux</u>	monthly	2D					X
ta		<u>Air temperature</u>	day	3D					
ua		<u>Zonal velocity</u>	day	3D					
va		<u>Meridional velocity</u>	day	3D					
w ap		<u>Lagrangian tendency of air pressure</u>	day	3D					
tas		<u>Near-surface air temperature</u>	day	2D					
uas		<u>Near-surface zonal velocity</u>	day	2D					
vas		<u>Near-surface meridional velocity</u>	day	2D					
hus ^[3]		<u>Specific humidity</u>	monthly	3D					
ts		<u>Skin temperature</u>	monthly	2D					
ps		<u>Surface air pressure</u>	monthly	2D					

^[1]: precipitation fluxes are provided in units of $kg/m^2 s$. The program also accepts other units of measure and related fields, depending on the known formats by ESMValTool preprocessor.

^[3]: specific humidity can also be given as near-surface 2-dimensional field (when available), or the lowermost pressure level of the 3-dimensional specific humidity field (variable name: hus s).

Table 2. Annual mean values of a 20-years subset of control runs from 12 models participating to the CMIP5 Project for: TOA and surface energy budgets (B_t and B_s , respectively), maximal and minimal peaks of atmospheric and oceanic meridional **heat enthalpy** transports (with peak locations in latitude degrees specified in brackets) (T_a^{max} , T_a^{min} , T_o^{max} , T_o^{min}), water mass budget ($\overline{E} - \overline{P}$), latent energy budget (\overline{R}_L), mechanical work by the Lorenz Energy Cycle, material entropy production computed with the direct and indirect methods ($\overline{\Sigma}_{dir}^{mat}$ and $\overline{\Sigma}_{ind}^{mat}$, respectively).

	$B_t - R_t$ $(\frac{W}{m^2})$	$B_s - F_s$ $(\frac{W}{m^2})$	T_a^{max} (PW)	T_a^{min} (PW)	T_o^{max} (PW)	T_o^{min} (PW)	$\overline{E} - \overline{P}$ $(\frac{kg}{m^2s} \times 10^{-8})$	\overline{R}_L $(\frac{W}{m^2})$	W $(\frac{W}{m^2})$	$\overline{\Sigma}_{dir}^{mat}$ $(\frac{mW}{m^2K})$	$\overline{\Sigma}_{ind}^{mat}$ $(\frac{mW}{m^2K})$
BNU	2.37	0.79	4.9 (42)	-5.1 (-39)	1.9 (19)	-0.9(-17)	-207.1	-5.89	2.0	64.9	58.7
Can2	0.08	0.19	4.7 (41)	-5.1 (-39)	1.5 (20)	-1.1 (-13)	5.32	-0.55	2.2	42.7	56.6
IPSL-M	0.33	0.32	4.6 (40)	-5.2 (-39)	1.5 (19)	-1.4 (-14)	11.1	-0.48	1.6	38.7	57.9
MIR-C	-3.16	1.50	4.8 (42)	-5.7 (-37)	1.4 (19)	-0.4 (-9)	-1.24	-0.70	1.3	39.8	56.5
MIR5	1.06	1.13	4.2 (42)	-4.6 (-40)	1.3 (18)	-0.6 (-10)	-2.94	-0.71	1.4	43.4	60.3
MPI-LR	0.36	0.58	5.0 (42)	-5.5 (-38)	1.9 (19)	-1.3 (-12)	-4.58	-0.88	1.8	43.4	58.7
MPI-MR	0.45	0.60	5.1 (42)	-5.6 (-39)	1.8 (19)	-1.3 (-11)	-4.03	-0.86	1.7	43.4	58.9

Table 3. Same as in Table 2 for the period 1970-2000 of the historical runs.

	$B_t - R_t$ $(\frac{W}{m^2})$	$B_s - F_s$ $(\frac{W}{m^2})$	T_a^{max} (PW)	T_a^{min} (PW)	T_o^{max} (PW)	T_o^{min} (PW)	$\overline{E} - \overline{P}$ $(\frac{kg}{m^2s} \times 10^{-8})$	\overline{R}_L $(\frac{W}{m^2})$	W $(\frac{W}{m^2})$	$\overline{\Sigma}_{dir}^{mat}$ $(\frac{mW}{m^2K})$	$\overline{\Sigma}_{ind}^{mat}$ $(\frac{mW}{m^2K})$
BNU	2.94	1.41	4.9 (42)	-5.1 (-39)	2.0 (19)	-0.8 (-13)	-199.9	-5.66	1.9	63.9	60.0
Can2	0.50	0.64	4.8 (42)	-5.2 (-39)	1.6 (21)	-1.0 (-12)	5.46	-0.53	2.2	43.2	57.7
IPSL-M	0.92	0.89	4.7 (40)	-5.4 (-39)	1.5 (19)	-1.4 (-13)	10.4	-0.47	1.6	39.6	59.1
MIR-C	-2.71	1.90	4.9 (42)	-5.7 (-37)	1.4 (17)	-0.8 (-11)	-1.12	-0.69	1.2	40.0	57.4
MIR5	1.30	1.32	4.3 (42)	-4.6 (-40)	1.3 (18)	-0.6 (-9)	-1.38	-0.67	1.3	43.4	61.1
MPI-LR	0.95	1.18	5.1 (42)	-5.6 (-38)	1.9 (19)	-1.2 (-11)	-4.91	-0.86	1.8	43.7	59.3
MPI-MR	0.99	1.11	5.1 (42)	-5.7 (-39)	1.8 (19)	-1.2 (-11)	-3.91	-0.82	1.7	43.7	59.5

Table 4. Same as in Table 2 for the period 2071-2100 of the RCP8.5 runs.

	$\overline{B_r - R_L}$ $(\frac{W}{m^2})$	$\overline{B_s - F_s}$ $(\frac{W}{m^2})$	T_a^{max} (PW)	T_a^{min} (PW)	T_o^{max} (PW)	T_o^{min} (PW)	$\overline{E} - \overline{P}$ $(\frac{kg}{m^2 s} \times 10^{-8})$	$\overline{R_L}$ $(\frac{W}{m^2})$	\overline{W} $(\frac{W}{m^2})$	$\overline{\Sigma_{dir}^{mat}}$ $(\frac{mW}{m^2 K})$	$\overline{\Sigma_{ind}^{mat}}$ $(\frac{mW}{m^2 K})$
BNU	4.79	3.19	5.5 (44)	-5.1 (-40)	1.9 (17)	-0.8 (-11)	-146.1	-4.15	1.8	61.7	65.5
Can2	2.36	2.29	5.6 (42)	-5.4 (-39)	1.4 (21)	-0.9 (-11)	6.74	-0.34	2.1	45.6	62.6
IPSL-M	2.79	2.63	4.9 (41)	-5.7 (-40)	1.3 (17)	-1.3 (-12)	7.38	-0.38	1.7	43.4	65.3
MIR-C	-0.29	4.11	4.9 (42)	-6.1 (-37)	1.1 (15)	-0.9 (-8)	-1.13	-0.51	1.1	42.5	63.1
MIR5	3.28	3.11	4.4 (42)	-5.0 (-40)	1.0 (18)	-0.7 (-7)	-0.94	-0.58	1.2	45.0	65.8
MPI-LR	2.68	2.96	5.3 (42)	-6.0 (-37)	1.8 (19)	-1.3 (-11)	-5.87	-0.76	1.6	46.1	64.1
MPI-MR	2.61	2.72	5.4 (44)	-6.1 (-40)	1.7 (17)	-1.3 (-11)	-3.78	-0.68	1.7	46.0	64.3

Table 5. Annual mean land-ocean asymmetries for atmospheric, latent energy budget and the difference of the two, for 20 years of multi-model ensemble simulations under piC conditions. Values are in $\overline{W/m^2}$.

	F_a [W/m ²]		R_L [W/m ²]		$(F_a - R_L)$ [W/m ²]	
	Oc.	Land	Oc.	Land	Oc.	Land
BNU	5.6	-10.7	5.8	-25	-0.2	14.3
Can2	6.3	-21.2	7.4	-18	-1.1	3.2
IPSL-M	4.0	-11	7.3	-15.1	-3.3	4.1
MIR-C	0.2	-16.8	7.8	-21.8	-7.6	5.0
MIR5	3.8	-11	11.8	-25.7	-8.0	14.7
MPI-LR	5.7	-20.5	6.1	-17.9	-0.4	2.6
MPI-MR	5.6	-20.2	6.2	-18.5	-0.6	1.7

Table 6. Evolution of land-ocean asymmetries for latent energy and for the residual of the atmospheric budget, for 20 years of multi-model ensemble simulations under the two extremal scenarios (piC and RCP8.5). Values are in \overline{PW} , and are positive if they are directed toward land. Values in brackets are from spatial integration over land, those not in brackets from integration over oceans.

	R_L [PW]		$(F_a - R_L)$ [PW]	
	piC	rcp8.5	piC	rcp8.5
BNU	2.09 (3.72)	2.91 (3.59)	-0.07 (2.13)	-1.34 (2.29)
Can2	2.67 (2.81)	3.32 (3.34)	-0.39 (-0.47)	-1.17 (0.37)
IPSL-M	2.63 (2.24)	3.07 (2.54)	-1.19 (0.64)	-1.75 (0.91)
MIR-C	2.82 (3.26)	3.21 (3.58)	-2.76 (0.76)	-3.19 (1.51)
MIR5	4.27 (3.83)	4.48 (3.94)	-2.90 (2.19)	-3.58 (2.78)
MPI-LR	2.18 (2.67)	2.63 (3.07)	-0.11 (-0.39)	-0.23 (0.16)
MPI-MR	2.32 (2.76)	2.81 (3.22)	-0.22 (-0.25)	-0.80 (0.26)

Table 7. Annual mean values of APE and KE reservoirs and conversion terms in the LEC. Values are in 10^5 J/m^2 for the reservoirs , in W/m^2 for the conversion terms. For the notation, refer to Appendix A1

	$A_z \text{PZ}$	$A_e \text{EPE}$	$K_z \text{KZ}$	$K_e \text{EKE}$	$\mathcal{C}(A_z, K_z) \text{CZ}$	$\mathcal{C}(A_e, K_e) \text{CE}$	$\mathcal{C}(A_z, A_e) \text{CA}$	$\mathcal{C}(K_e, K_z) \text{CK}$
piC								
BNU	54.9	5.8	9.2	5.6	-0.20	1.7	1.5	0.50
Can2	52.2	5.7	8.8	5.2	-0.20	1.9	1.7	0.40
IPSL-M	52.0	5.7	8.8	5.2	-0.20	1.6	1.4	0.40
MIR-C	53.1	5.7	7.6	5.3	-0.03	1.3	1.2	0.20
MIR5	48.0	5.9	5.9	5.0	-0.05	1.3	1.2	0.20
MPI-LR	46.8	5.6	7.2	5.8	-0.10	1.6	1.5	0.40
MPI-MR	48.8	5.6	7.1	6.1	-0.10	1.6	1.6	0.40
hist								
BNU	52.8	5.7	10.4	5.8	-0.20	1.7	1.5	0.5
Can2	50.7	5.6	10.0	5.4	-0.20	1.9	1.6	0.5
IPSL-M	50.5	5.9	10.5	5.4	-0.20	1.9	1.6	0.50
MIR-C	51.4	5.8	8.1	5.3	-0.02	1.2	1.6	0.40
MIR5	47.2	5.8	6.4	5.1	-0.05	1.3	1.2	0.30
MPI-LR	45.6	5.5	7.6	6.0	-0.15	1.6	1.5	0.50
MPI-MR	46.9	5.6	7.6	6.3	-0.14	1.6	1.5	0.50
rcp8.5								
BNU	46.9	5.3	12.5	6.1	-0.20	1.6	1.3	0.50
Can2	49.1	5.4	13.1	5.9	-0.20	1.9	1.5	0.4
IPSL-M	47.4	5.6	14.8	6.3	0.05	1.8	1.4	0.50
MIR-C	50.0	5.4	11.5	5.6	0.01	1.1	1.3	0.40
MIR5	47.2	5.5	9.1	5.4	-0.03	1.2	1.0	0.3
MPI-LR	44.0	5.3	10.4	6.7	-0.10	1.6	1.4	0.60
MPI-MR	44.8	5.4	9.8	6.8	-0.10	1.6	1.4	0.60

Table 8. Annual mean components of the material entropy production, obtained with the direct and indirect methods. For each model, the first row denotes the estimates from piC, the second row from hist, the third row from rcp8.5. Values are in ~~$mW * m^{-2} * K^{-1}$~~ $mW/m^2 K^{-1}$.

	S_{hyd}	S_{sens}	S_{kin}	S_{ver}	S_{hor}
BNU	54.8	2.67	6.97	51.2	7.5
	54.6	2.59	6.79	53.3	7.4
	52.9	2.40	6.38	58.7	6.8
Can2	32.1	2.97	7.70	50.0	7.5
	32.8	2.85	7.58	51.5	6.6
	35.9	2.64	7.18	56.7	5.96
IPSL-M	30.1	2.92	5.6	49.7	8.2
	31.0	2.84	5.71	51.0	8.3
	34.8	2.64	6.01	57.1	8.2
MIR-C	32.7	2.53	4.53	50.1	6.53
	33.0	2.47	4.44	50.9	6.52
	36.2	2.30	3.96	57.0	6.11
MIR5	36.7	1.84	4.87	54.1	6.19
	36.9	1.80	4.75	55.0	6.18
	39.1	1.71	4.24	59.8	6.0
MPI-LR	34.7	2.51	6.23	51.6	7.05
	35.2	2.41	6.12	52.6	7.00
	38.2	2.23	5.66	57.3	6.82
MPI-MR	34.6	2.52	6.26	51.6	7.32
	35.1	2.41	6.14	52.4	7.27
	37.9	2.23	5.84	57.2	7.15

Table 9. 20-year annual mean material entropy production associated with the hydrological cycle. Each component denotes a different process: (from left to right) evaporation, rainfall precipitation, snowfall precipitation, snow melting at the ground, potential energy of the droplet. For each model, the first row denotes the estimates from piC, the second row from hist, the third row from rcp8.5. Values are in ~~$mW * m^{-2} * K^{-1}$~~ mW/m^2K^{-1} .

	S_e	S_r	S_s	S_m	S_p
BNU	-278.7	307.5	24.1	-2.58	4.53
	-280.7	310.1	23.2	-2.47	4.65
	-296.9	329.3	17.2	-1.83	5.16
Can2	-269.2	276.7	25.9	-2.49	4.21
	-269.5	277.6	22.6	-2.42	4.29
	-282.5	298.0	17.4	-1.86	4.79
IPSL-M	-271.5	274.5	25.9	-2.78	4.02
	-274.8	279.5	24.8	-2.66	4.13
	-293.3	306.2	19.4	-2.08	4.65
MIR-C	-278.6	283.1	22.8	2.46	4.17
	-271.5	280.3	22.4	2.41	4.21
	-286.7	303.5	16.5	-1.77	4.70
MIR5	-315.6	328.0	21.8	-2.35	4.86
	-313.7	326.6	21.4	-2.31	4.91
	-321.7	338.7	18.8	-2.03	4.65
MPI-LR	-288.1	295.2	26.0	-2.80	4.46
	-287.6	296.1	24.9	-2.68	4.53
	-298.2	313.0	20.7	-2.22	4.97
MPI-MR	-291.9	299.1	25.7	-2.76	4.43
	-292.2	300.8	24.6	-2.65	4.52
	-303.2	318.3	20.1	-2.17	4.94

Table 10. 20-year Annual mean irreversibility and baroclinic efficiency for each model and each scenario. Baroclinic efficiency is rescaled as 10^{-2} .

		piC	hist	rcp8.5
BNU	α	8.6	8.8	9.0
	η	6.1	6.1	5.6
Can2	α	4.8	4.9	5.6
	η	5.6	5.7	5.1
IPSL-M	α	6.4	6.4	6.6
	η	6.6	6.7	6.4
MIR-C	α	8.4	8.5	10.1
	η	5.1	5.1	4.8
MIR5	α	8.4	8.6	10.1
	η	5.7	5.8	5.6
MPI-LR	α	6.4	6.3	7.5
	η	6.0	6.1	5.8
MPI-MR	α	6.4	6.2	7.2
	η	6.3	6.4	6.2