

We thank the reviewers for their insightful comments. The reviewers' comments are repeated below in italics, followed by point-by-point responses to each. We also have improved the figures and text covering sea ice after helpful comments from a collaborator (C. Horvat) who has been added as a co-author. All figures were updated to ensure consistent fonts and notations.

The revised manuscript with tracking of all the changes that have been made is appended at the end of this response.

## **Response to Reviewer #1**

*This paper shows the impact of horizontal resolution on ocean-sea-ice model simulations by comparing the outputs from four models at low and high horizontal resolutions. Several biases in the low-resolutions models are improved when the resolution becomes high. The impacts of model resolution on temperature and salinity bias depend on the model. Although the model configuration differences between low and high resolutions differ among the four models, several improvements in high-resolution models compared to low-resolution models seem to be robust. On the one hand, the biases of temperature and salinity fields depend on the model and region. The report in this paper is informative for readers of GMD. I recommend this paper for publication after some minor modifications.*

### Comments

*The time evolution of the global temperature was examined in Section 3.2. Levitus et al. (2005) showed a warming of the world ocean over 1955-2003 based on the observations. The warmings in the models seem to be consistent with the observation including its decadal variations. As well as the sea surface height, the impact of global warming should be discussed.*

A primary focus of Section 3 is on simulation drift, which tends to be much larger than forced variability in the first few forcing cycles of OMIP simulations for many fields, particularly global ocean temperature (Danabasoglu et al. 2014; Griffies et al. 2014). With only one forcing cycle available from the high-resolution OMIP-2 runs, it can be difficult to distinguish forced variability from model drift. The OMIP simulations are initialized from observed climatologies, not from observed 1958 conditions, and so it is not expected that they should accurately simulate the full-depth evolution of the ocean over the 1958-2018 forcing cycle. In the real world, this evolution reflects the cumulative effect of hundreds or thousands of years of surface forcing. Furthermore, the JRA55-do forcing dataset includes adjustments designed to yield zero net climatological heat/freshwater flux into the ocean for the 1988-2007 time period (Tsujino et al. 2018). Thus, an ocean model that perfectly replicates the observed sea surface state should exhibit zero net ocean heating over that 20-year time period when forced with JRA55-do. This is unrealistic, but it is a reasonable choice in the dataset construction given the large uncertainty in the climatological net surface forcing of the ocean (Tsujino et al. 2018). While it is problematic to assess the realism of long-term ocean warming in OMIP simulations for all the reasons outlined above, the JRA55-do forcing should reflect variations in the global energy imbalance that produce interannual to decadal changes in the rate of ocean warming (Trenberth et al. 2014).

In response to the reviewer, we now include two versions of the original Figure 4 (time evolution of global temperature and salinity) in the paper. The original version (showing anomalies relative to initial conditions – Figure 4 in revised manuscript) highlights model drift away from observed ocean climatology. The new version (showing anomalies relative to 2018 – Figure 5 in revised manuscript)

highlights the forced ocean variations of the last few decades of simulation. The comparison with observed heat/salt content change from the WOA18 product in panels c-f of Figure 5 shows that high resolution improves the fidelity and reduces the spread of forced ocean heat content change (particularly 0-2000m heat content) over recent decades. Figures 7-9 in the revised manuscript (time evolution of ocean temperature as a function of depth) are left unchanged because they are intended to illuminate the geographical regions where model drift (i.e., bias growth) is most pronounced, not where forced ocean heat uptake occurs.

**Version 2 of Figure 4, to be included in the manuscript as Figure**

*Figure 9 shows the surface temperature and salinity differences between the simulated fields over 2009-2018 and the initial fields, which discusses about how the simulated field changes from its initial state. Global warming appears to influence much the temperature difference. The model bias compared to the observation should be correctly verified. The simulated surface temperature and salinity are constrained to atmospheric forcing and the restoring to the observation respectively. Because these constrain are not perfect. Therefore, it is worth verifying the surface temperature and salinity biases compared with recent observations such as Argo float in the same period.*

The SST/SSS bias plot (now Fig. 10) has been replotted using concurrent observations (WOA18) spanning 1980-2018 to match Figures 11, 12, and 13 in the revised manuscript. The new plot differs somewhat from the old, but still indicates that model bias is essentially a reflection of model drift away from the initial conditions. This bias generally sets up within a decade or two of initialization (Figures 7-10). The discussion in the paper has been updated accordingly.

Minor comments

*There are not a few typos in the texts. The lead-author should be responsible to make readable paper. In Section 2, the model descriptions of four models are shown. It seems that different authors responsible to each model wrote the texts for each configuration, which makes the texts unreadable. The configuration of each model should be written in the same writing style.*

Agreed. The description of the models was revised to be consistent with each other.

*GMD recommend the unit with negative exponent (e.g.  $m s^{-1}$  instead of m/s).*

Corrected throughout the manuscript.

*L82: 'AMOC' appears first. 'Atlantic meridional overturning circulation' should be shown.*

Done.

*L98:  $\sigma_2$  should be defined.*

Defined.

*L107:  $\Delta x$  should be defined such as 'zonal grid spacing'.*

Now defined. HYCOM uses a regular grid on a Mercator projection.

*L145-146: 'm2 s-1' should be modified by using superscript.*

Corrected throughout the manuscript.

*L313: 'section 2b,c' should be 'section 3.2 and 3.3'.*

Corrected.

*Sec.3.4: The surface temperature and salinity averaged over 2009-2018 are compared to the initial state (Fig. 9). On the one hand, the vertical structures averaged over 1980-2018 are compared to the initial state (Fig. 10, 11, 12). Why are the average periods different?*

Good point. In light of the reviewer's earlier comment on section, Figure 9 (now Figure 10) was revised with a comparison to the 1980-2018 average period.

*Sec.3.7: Please plot the EKE time series along the ACC, if possible. Were the EKE in the models were increased by the zonal wind intensification since the 1970s.*

The difficulty with an evaluation of the EKE response to winds is that there is also an initial condition transient in the buoyancy gradients across the ACC due to the model drift discussed above (see Figure 14c illustrating the buoyancy drift). Since the MOC across the ACC is equilibrating for a significant fraction of the whole time of the high-resolution integration, it is difficult to interpret the mechanisms leading to EKE response and it is somewhat beyond the scope of this paper. Thus, we anticipate that this figure would be quite confusing to the readers. In light of this comment and one from Reviewer 2, we have expanded the discussion.

*L580: I cannot find Ollitrault and Colin de Verdière, 2014 in References.*

The reference has been added.

*Fig. 27: Please add the curves of heat transport of IAP-LICOM high resolution in Fig. 27.*

Done.

*'summer' and 'winter' in the captions of Fig. 30 and 31 should be 'winter' and 'summer' respectively.*

Corrected.

*L242, 553, 1170: Renault et al. (2019) should be Renault et al. (2020).*

Corrected.

*L909-918: Please confirm the sequence of references.*

Corrected.

*L963 includes two references.*

Corrected.

*L1204 includes two references.*

Corrected.

*Table.1: Please show the initial conditions.*

The initial conditions were added to Table 1.

*Table.1: 'Relative wind stress' should be 'Absolute wind stress' for FSU-HYCOM'.*

Corrected.

## **Response to Reviewer #2**

### General Comments

*This is an important paper for the ocean and climate modelling community feeding into wider decisions on the benefits of high resolution ocean components to address climate science questions. The conclusion that the impact of resolution is consistent across model families in terms of circulation but model dependent in terms of temperature and salinity biases is an important one. My main comment is that the high and low resolution versions of the models are not always comparable due to differences in vertical as well as horizontal resolution. It would have been preferable to have changed only the horizontal resolution and associated physics so that the effects of horizontal resolution can be isolated as stated in the abstract. I would suggest that at very least, this issue should be addressed in the model description and discussion in the context of Stewart et al. (Ocean Modelling, 2017) and included as a future recommendation.*

Done. This was an opportunistic study taking advantage of ongoing simulations rather than a dedicated parallel set of simulations. As discussed below, more often than not, the low- and high-resolution were configured independently for different scientific goals and followed different development trajectories, when ideally they should have been designed in tandem.

*I would recommend this paper for publication assuming that this and other specific issues below are addressed.*

### Specific comments:

*L21: See discussion above on isolating the effects of horizontal resolution*

This is now discussed in the introduction of Section 2.



*L86: Title would be better as Model Descriptions*

The title has been changed to “Description of the models”

*L93: A summary table of the four models at both resolutions would be useful for the reader*

The four models are summarized in Table 1. The table has been updated to include the climatologies used to initialize the models.

*L117: Why is FSU model 41 layers in low res and 36 layers in high res? (see above)*

The 36-layer high-resolution configuration was at the time our default configuration and we retained it to compare to previous runs performed with other atmospheric forcings. The 41-layer coarse-resolution runs were performed afterward for inclusion in the Tsujino et al. (2020) using the latest vertical grid with all the additional layers in the upper ocean. This is now addressed in the revised manuscript.

*L157: What is the impact in the NCAR model of two additional levels as well as partial step formulation?*

The additional (250 m thick) vertical levels in high resolution POP increase the maximum ocean depth from 5500 m to 6000 m, allowing for a more realistic representation of deep ocean features resolved by the 0.1° grid. The partial bottom cell formulation improves the representation of deep bathymetric slopes. Neither has been tested in the 1° version, and so the impacts are unknown, but are presumed to be small compared to the horizontal grid refinement which includes overall improved representation of small-scale bathymetric features (even without the minor changes to vertical discretization). It should be noted that the bottom depths in the POP model description (section 2.2) were incorrect and should have been 5500m (1-degree) and 6000m (0.1-degree). This has been corrected in the revised manuscript.

*L207: I couldn't see any mention in the description of the lack of salt flux normalization in IAP?*

This has been added.

*L213: What is the impact of changing from 30 to 55 levels in the high resolution?*

*It seems to me that this is more consistent with resolving baroclinic modes at high resolution (Stewart et al., 2017) but 55 levels is probably insufficient to resolve the first baroclinic mode.*

The main goal was to increase the vertical resolution in the deep ocean to improve the simulation of the deep circulation and the AMOC transport. The AMOC transport is better represented at 26.5°N in the high-resolution CAS-LICOM experiment than in the low-resolution experiment (Figure 26). We do agree with the reviewer that 55 levels may be insufficient to resolve the first baroclinic mode, but the 55 levels were dictated by the computing resources at hand. This is now addressed in the revised manuscript.

*L229: Title would be better as Temporal evolution and model drift*

Agreed. The title has been changed accordingly.

*L249: Is there any comment on why the global average KE is so different between NCAR and IAP (which should be fairly comparable in terms of resolution)?*

This is under investigation by the LICOM team, but it most likely due to the numerics, especially the two-step shape preservation advection scheme used in the momentum equations. The Euler-backward finite difference scheme used in the advection scheme is quite diffusive and the LICOM team is exploring ways to decrease the frequency of which the Euler-backward finite difference scheme is used. In addition, CAS-LICOM uses a relatively large horizontal diffusivity. This is now addressed in the revised manuscript.

*L258: Add this discussion of the salt flux normalization to model description and specifically in description of IAP*

Done.

*L275: Please correct 'all high resolution models warm faster over the upper 2000m and global temperature than their lower-resolution partners'. This doesn't appear to be true for FSU.*

Corrected.

*L284: I understand that including this is beyond the scope of this paper, but the authors could recommend and pass to the discussion, that the vertical heat and salt budgets are compared in future as in Griffies et al. (J. Climate, 2015) and Von Storch et al. (Ocean Modelling, 2016)*

Done.

*L335: Is it possible to hypothesise what leads to reductions or increases in the biases and the different model behaviour?*

We can hypothesize that the thermocline bias is related to the representation of vertical eddy heat flux (Griffies et al. 2015), which tends to be stronger and more realistic in high resolution simulations (see revised Figure 4 in response to reviewer #1). The degradation in thermocline bias in POP high resolution could also be due to missing submesoscale physics, which are parameterized in low resolution, but absent in high resolution, versions of that model. This is now addressed in the revised manuscript.

*L417: is the similarity of the high resolution model and RAPID AMOC evolutions statistically significant or can it be shown to link with reduced deep convection? Are the convection patterns different across resolutions? I think without further discussion, this point is rather speculative.*

Most of the discussion highlights in broad terms the similarity between the model multi-decadal variability and the observations as put forward by Hakkinen and Rhines (2004) and Yashayaev (2007) and does not speculate on the RAPID variability. The RAPID time series is only provided for reference and to show consistency among the observations and the models. The text was slightly modified to better convey this point.

*L445: Should figure 14c be labelled as AABW cell?*

We prefer global MOC transport (see Lumpkin and Speer (2007) and Talley (2013) for observed values). Figure 14c (now Figure 15c) will be relabeled as “global MOC transport at 34°S” and the figure caption modified to reflect that the negative values in global MOC transport are due to the northward-flowing AABW below the southward return flow.

*L502: Since the temperature and salinity fields at depth are still evolving at the end of these experiments, it seems more likely that the ACC changes are related to buoyancy drifts. Can you comment on this?*

There is indeed an initial condition transient in the buoyancy gradients across the ACC (Figure 14c) and the MOC across the ACC is equilibrating for a significant fraction of the whole time of the high-resolution integration. The discussion was extended to reflect this point. See also the response to reviewer #1 regarding the EKE sensitivity to decadal wind stress changes.

*L528: Also comment here about the choice of vertical levels affecting the representation of baroclinic modes (Stewart et al., 2017)*

Done.

*L539: I would argue from figure 16, that there is at least marginal improvement in the representation of the NW corner across the models but it is difficult to see in the multi panel figure*

The representation of the NW corner in these experiments was further discussed in a companion CLIVAR Variations/Exchanges article (Figure 1 of Chassignet et al., 2020). The improvement is very small. This is now referred to in the revised manuscript.

*L555: Please comment on Agulhas eddies in other models*

Done.

*L710: Can you comment on the cause of the increased northward heat transport in the Atlantic by looking at the components? Presumably, it is related to a stronger Gulf Stream transport*

Northward heat transport is usually correlated with higher AMOC transport. In the high-resolution experiments, there is also an overall reduction in subtropical Atlantic upper ocean cold bias (see Figure 23) which is likely to have an impact (Msadek et al., 2013). This has been addressed in the revised manuscript.

*L834: Reference should be Griffies et al. (2015) see below. Include here recommendation on diagnosing vertical heat and salt budgets in future experiments?*

Done.

*L848: I think specific comment is merited here on the aim to maintain traceability across resolution hierarchies in terms of numerics and parameterisations as well as vertical resolution*

Done.

*Technical corrections:*

*L106 (and similar throughout): present as  $3 \times 10^{-5}$*

Corrected.

*L414: extra 'only' in the sentence*

Corrected

*L505: The sentence doesn't make sense as currently written*

Rewritten.

# Impact of horizontal resolution on global ocean-sea-ice model simulations based on the experimental protocols of the Ocean Model Intercomparison Project phase 2 (OMIP-2)

Eric P. Chassignet<sup>1</sup>, Stephen G. Yeager<sup>2</sup>, Baylor Fox-Kemper<sup>3</sup>, Alexandra Bozec<sup>1</sup>, Frederic Castruccio<sup>2</sup>, Gokhan Danabasoglu<sup>2</sup>, [Christopher Horvat](#)<sup>3</sup>, Who M. Kim<sup>2</sup>, Nikolay Koldunov<sup>4</sup>, Yiwen Li<sup>5</sup>, Pengfei Lin<sup>5</sup>, Hailong Liu<sup>5</sup>, Dmitry Sein<sup>4,6</sup>, Dmitry Sidorenko<sup>4</sup>, Qiang Wang<sup>4</sup>, and Xiaobiao Xu<sup>1</sup>

<sup>1</sup>Center for Ocean-Atmospheric Prediction Studies, Florida State University, Tallahassee, FL, USA

<sup>2</sup>National Center for Atmospheric Research, Boulder, CO, USA

<sup>3</sup>Brown University, Providence, RI, USA

<sup>4</sup>Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung (AWI), Bremerhaven, Germany

<sup>5</sup>State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

<sup>6</sup>Shirshov Institute of Oceanology, Russian Academy of Science, Moscow, Russia

*Correspondence to:* Eric P. Chassignet (echassignet@fsu.edu)

**Abstract.** This paper presents global comparisons of fundamental global climate variables from a suite of four pairs of matched low- and high-resolution ocean and sea-ice simulations that are obtained following the OMIP-2 protocol (Griffies et al., 2016) and integrated for one cycle (1958-2018) of the JRA55-do atmospheric state and runoff dataset (Tsujino et al., 2018). Our goal is to assess the robustness of climate-relevant improvements in ocean simulations (mean and variability) associated with moving from coarse ( $\sim 1^\circ$ ) to eddy-resolving ( $\sim 0.1^\circ$ ) horizontal resolutions. The models are diverse in their numerics and parameterizations, but each low-resolution and high-resolution pair of models is matched so as to isolate, to the extent possible, the effects of horizontal resolution. A variety of observational datasets are used to assess the fidelity of simulated temperature and salinity, sea surface height, kinetic energy, heat and volume transports, and sea ice distribution. This paper provides a crucial benchmark for future studies comparing and improving different schemes in any of the models used in this study or similar ones. The biases in the low-resolution simulations are familiar and their gross features – position, strength, and variability of western boundary currents, equatorial currents, and Antarctic Circumpolar Current – are significantly improved in the high-resolution models. However, despite the fact that the high-resolution models “resolve” most of these features, the improvements in temperature or salinity are inconsistent among the different model families and some regions show increased bias over their low-resolution counterparts. Greatly enhanced horizontal resolution does not deliver unambiguous bias improvement in all regions for all models.

## 1 Introduction

A key decision in climate model design is the spatial resolution of different components. The global scope of the integrations and the centennial to millennial timescales and multiple scenarios required to capture changes in the climate set the problem; the spatial resolution is therefore the result of available computing. This decade, computing has become

Formatted: English (United States)

Deleted: New

Formatted: English (United States)

sufficiently powerful to make mesoscale-rich resolution affordable in the ocean and sea-ice models over most of the earth, which allows the ocean model to simulate more intense internal variability than a lower resolution model. As this new regime of coupled modeling is entered, it is important to understand both the behavior of ocean and sea ice models in a controlled framework, and the benefits and challenges that come with higher resolution. This work introduces a set of matched numerical simulations in low- and high-resolution ocean and sea-ice models with the latest forcing protocol. It is anticipated that these results will inform fully coupled modeling studies where ocean resolution varies, and also that follow-on studies will build on our results to examine process and regional detail.

In 2016, an international group of ocean modelers behind the development and analysis of global ocean–sea-ice models used as a component of the earth system models in CMIP6 proposed an Ocean Model Intercomparison Project (OMIP) (Griffies et al., 2016). The essential element behind the OMIP is a common set of atmospheric and river runoff datasets for computing surface boundary fluxes to drive the ocean–sea-ice models, many of which are used as components of coupled climate system models. The OMIP protocol is an outcome of the Coordinated Ocean–sea-ice Reference Experiments (CORE) which assessed the performance of ocean–sea-ice models (Griffies et al., 2009; Danabasoglu et al., 2014; Griffies et al., 2014; Downes et al., 2015; Farneti et al., 2015; Danabasoglu et al., 2016; Wang et al., 2016a, 2016b; Ilıcak et al., 2016; Tseng et al., 2016; Rahaman et al., 2019) using the atmospheric and river runoff dataset of Large and Yeager (2009). However, this dataset has not been updated since 2009 and a new dataset (JRA55-do; Tsujino et al., 2018) has been developed for the OMIP based on the Japanese Reanalysis (JRA-55) product from Kobayashi et al. (2015) to ensure that it is regularly updated. This raw reanalysis product has been substantially adjusted to match reference states based on observations or ensemble mean of other atmospheric reanalysis products as detailed in Tsujino et al. (2018) to create a suitable forcing dataset for ocean and sea-ice models, referred to as JRA55-do. The continental river discharge is provided by a river-routing model forced by input runoff from the land-surface component of JRA-55 adjusted to ensure similar long-term variabilities as in the CORE dataset (Suzuki et al., 2018). Runoff of ice-sheet and glaciers from Greenland (Bamber et al., 2012; Bamber et al., 2018) and Antarctica (Depoorter et al., 2013) are also incorporated. Tsujino et al. (2020) presents an evaluation of the simulations from CMIP6-class global ocean–sea-ice models forced with the JRA55-do datasets. This effort compares CORE-forced (i.e., OMIP-1) and JRA55-do-forced (i.e., OMIP-2) simulations considering metrics commonly used in the evaluation of global ocean–sea-ice models to assess model biases.

Many features are very similar between OMIP-1 and OMIP-2 simulations, but Tsujino et al. (2020) identify many improvements in the simulated fields in transitioning from OMIP-1 to OMIP-2. For example, the sea surface temperature of the OMIP-2 simulations reproduce the global warming hiatus in the 2000s and the recent observed warming, which are absent in OMIP-1 partly because the OMIP-1 forcing stopped in 2009. The low bias in the sea-ice area fraction in summer of both hemispheres in OMIP-1 is significantly improved in OMIP-2. The overall reproducibility of both seasonal and interannual variation in sea surface temperature and sea surface height (dynamic sea level) is also improved in OMIP-2. Tsujino et al. (2020) attributes many of the remaining model biases either to errors in representing important processes in ocean–sea-ice models, some of which are expected to be mitigated by taking finer horizontal and/or vertical resolutions, or to shared biases

70 in the atmospheric forcing. In this paper, we make a first attempt at quantifying the impacts of the models' horizontal resolution on biases.

Our goal is to assess the robustness of climate-relevant improvements in ocean simulations (mean and variability) associated with moving from coarse ( $\sim 1^\circ$ ) to eddy-resolving ( $\sim 0.1^\circ$ ) horizontal resolutions. Using the same atmospheric forcing (JRA55-do) for both low- and high-resolution configurations, we perform a multi-model analysis to identify the robust differences and improvements associated with increased resolution given the same forcing datasets. Within the ocean modeling community, it is usually assumed that high-resolution simulations should in general produce better results than low-resolution ones (Fox-Kemper et al., 2019). While this is clearly the case for surface currents and internal variability, we will show that greatly enhanced horizontal resolution does not necessarily deliver unambiguous bias improvement in temperature and salinity in all regions. It is important to note several caveats when interpreting the results presented in this paper: First, this is based on a limited number of numerical models (four) and, second, because of the large computational cost associated with the high-resolution runs (factor 1000 more expensive), only one JRA55-do cycle (1958-2018) is analyzed in this paper (versus 6 cycles for the coarse-resolution runs of Tsujino et al. (2020)). Also, because of the short integration time, some of the results may not be robust (Atlantic meridional overturning circulation variability, deep ocean circulation, etc.) (Danabasoglu et al., 2016). The layout of the paper is as follows. The models used in the comparison are described in Section 2. Section 3 highlights differences in the magnitude of the models' drift while Section 4 focuses on the detrended interannual to decadal variability and the differences in the modeled ocean climates. The results are summarized and discussed in the final section.

## 2 Description of the models

Because the goal is to identify the robust differences and improvements associated with increased horizontal resolution given the same forcing datasets, most of the participating modeling groups configured their high-resolution configuration with similar parameters to that of the low-resolution configuration (see Table 1 for a detailed description of the parameters used in the low- and high-resolution model configurations). Ideally, only the horizontal resolution and associated physics should be changed to isolate the effects of horizontal resolution (Stewart et al., 2017), but this was not achievable for the present study since many of the low- and high-resolution simulations were often configured independently for distinct scientific goals and followed different development trajectories (e.g., vertical grids). It is also important to note that not all the models used the same climatology for the initial conditions, nor did they use the same wind stress formulation (absolute versus relative winds). When evaluating the drift of a numerical simulation, it is performed with respect to the climatology used to initialize the run.

### 2.1 FSU-HYCOM

The FSU-HYCOM model is a global configuration of the HYbrid Coordinate Ocean Model (HYCOM) (Bleck, 2002; Chassignet et al., 2003; Halliwell, 2004). The sea-ice component is CICE version 4 (CICE4, Hunke and Lipscomb, 2010). The initial conditions in (potential) temperature and salinity are given by the Generalized Digital Environmental Model (GDEM4, Teague et al., 1990; Cames, 2009). The Large and Yeager (2004) bulk formulation is used for turbulent air-sea fluxes except

Deleted: MOC

Formatted: English (United States)

Deleted: Model's d

Formatted: English (United States)

Formatted: Space After: 0 pt

Deleted: coarse

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Moved (insertion) [3]

Moved (insertion) [4]

Moved (insertion) [5]

105 ~~for the surface wind-stress that is calculated without surface currents (absolute wind stress). No restoration is applied on the sea surface temperature. There is no parameterization of the overflows.~~

For the ~~low~~-resolution configuration, ~~FSU-HYCOM uses~~ a tripolar Arakawa C-grid of 0.72° horizontal resolution with refinement to 0.33° at the equator (500 cells in the zonal direction and 382 in the meridional direction). The 2-minute NAVO/Naval Research Laboratory DBDB2 dataset provides the bottom topography. Forty-one hybrid coordinate layers ~~are~~  
 110 ~~used, with potential density  $\sigma_t$  target densities ranging from 17.00 to 37.42 kg m<sup>-3</sup> are used (same configuration as Tsujino et al. (2020)).~~ The vertical discretization combines fixed pressure coordinates in the mixed layer and unstratified regions, isopycnic coordinates in the stratified open ocean, and terrain-following coordinates over shallow coastal regions. ~~The surface salinity is restored to the monthly GDEM4 climatology over the entire domain with a salinity piston velocity of 50 m/60 days and the salinity flux at each time step is adjusted to ensure a net global flux of zero.~~ Vertical mixing is the KPP model (Large  
 115 et al., 1994) with a background diffusion of  $3 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ . Interface height smoothing by a biharmonic operator is used to correspond to Gent and McWilliams (1990), with a mixing coefficient determined by the grid spacing  $\Delta x$  (~~regular grid on a Mercator projection~~) times a velocity scale of  $0.02 \text{ m s}^{-1}$  everywhere, except in the North Pacific and North Atlantic where a Laplacian operator with a velocity scale of  $0.01 \text{ m s}^{-1}$  is used. Gent and McWilliams (1990) is not implemented where the FSU-HYCOM has coordinate surfaces aligned with constant pressure (mostly in the upper ocean mixed layer) and there is no  
 120 rotation of the lateral diffusion along neutral surfaces.

For the ~~high~~-resolution configuration, ~~FSU-HYCOM uses~~ a tripolar Arakawa C-grid of 0.08° (1/12°) horizontal resolution (4500 cells in the zonal direction and 3298 in meridional direction). The model bathymetry is derived from the 30 arc-second GEBCO08 dataset. A vertical resolution of 36 hybrid layers, with  $\sigma_t$  target densities ranging from 26.00 to 37.24 kg m<sup>-3</sup>, is used. ~~The 36-layer high resolution configuration was at the time our default configuration and was retained to compare to previous runs performed with other atmospheric forcing datasets. The 41-layer coarse-resolution runs were performed for inclusion in the Tsujino et al. (2020) using the latest vertical grid with all the additional layers in the upper ocean. While the vertical resolution is lower in both configurations than recommended by Stewart et al. (2017) for z-coordinate models, the statistics of eddy scale and the vertical structure of the resolved eddy motions are well captured with this layer discretization when compared to a z-coordinate model with 300 levels (Ajayi et al., 2020).~~ The surface salinity is restored to  
 125 the monthly GDEM4 climatology over the entire domain with a salinity piston velocity of 50 m/60 days and the salinity flux at each time step is adjusted to ensure a net global flux of zero. The KPP model (Large et al., 1994) with a background diffusivity of  $10^{-5} \text{ m}^2 \text{ s}^{-1}$  provides the vertical mixing. The horizontal advection uses a second-order flux corrected transport scheme. An interface height smoothing is applied through a biharmonic operator (with a velocity scale of  $0.015 \text{ m s}^{-1}$ ).

## 2.2 NCAR-POP

135 The NCAR-POP model is based on the ocean component of the Community Earth System Model version 2 (CESM2; Danabasoglu et al. 2020) and is a global configuration of the Parallel Ocean Program version 2 (POP2; Smith et al. 2010) with several modifications to the model physics and numerics including improved treatment of continental freshwater discharge

Moved (insertion) [6]

Deleted: coarse

Deleted: the grid

Deleted: is

Deleted: whose

Deleted: e

Deleted: /

Deleted: <sup>3</sup>

Moved up [4]: The initial conditions in (potential) temperature and salinity are given by the Generalized Digital Environmental Model (GDEM4, Teague et al., 1990; Carnes, 2009)).

Moved up [3]: The sea-ice component is CICE (Hunke and Lipscomb, 2010).

Moved up [5]: The Large and Yeager (2004) bulk formulation is used for turbulent air-sea fluxes except for the surface wind-stress that is calculated without surface currents (absolute wind stress). No restoration is applied on the sea surface temperature.

Deleted: . T

Deleted: he salinity flux is globally normalized at each time step

Deleted: /

Deleted: The horizontal advection uses a second-order flux corrected transport scheme. A Laplacian diffusion of  $0.03\Delta x$  is applied on temperature and salinity and a combination of Laplacian ( $0.03\Delta x$ ) and biharmonic ( $0.05\Delta x^2$ ) dissipation is applied on the velocities (see Table 1). The model baroclinic and barotropic time steps are 1800s (leap-frog) and 56.25s (explicit) respectively.

Deleted: in m

Moved up [6]: There is no parameterization of the overflows.

Formatted: Indent: First line: 0.38"

Deleted: global

Deleted: the grid

Deleted: is

Deleted: /

Deleted: <sup>3</sup>

Deleted: The initial conditions in temperature and salinity : ... [1]

Deleted: ,

Deleted: /s

Formatted: English (United States)

Deleted: The horizontal diffusion uses a combination of co ... [2]

Deleted: There is no parameterization of the overflows.

Formatted: English (United States)

Moved (insertion) [2]

Deleted: , but

Deleted: features



into unresolved estuaries (Sun et al., 2019) and a new parameterization of Langmuir mixing (Li et al., 2016). The sea-ice component of CESM2 is CICE version 5.1.2 (CICE5; Hunke et al., 2015) which features new mushy-layer thermodynamics (Turner and Hunke, 2015) with prognostic sea-ice salinity and an updated melt pond parameterization (Hunke et al., 2013). CICE5 uses the same horizontal mesh grid as the POP2 configuration to which it is coupled. The initial conditions are given by World Ocean Atlas 2013 (WOA13; Locarnini et al., 2013; Zweng et al., 2013). The surface stress is a function of ocean surface velocity (relative wind stress), and sea surface salinity is restored to WOA13 monthly climatology with a piston velocity of 50 m over one year. Both configurations use a precipitation factor, computed once per year, to prevent salinity drift as discussed in Appendix C of Danabasoglu et al. (2014).

For the low-resolution configuration, NCAR-POP utilizes a dipole mesh grid with the grid northern pole displaced into Greenland. The horizontal resolution (nominal  $1^\circ$ ) is uniform in the zonal direction ( $1.125^\circ$ ), but varies in the meridional direction (from  $0.27^\circ$  at the Equator to  $\sim 0.5^\circ$  at mid-latitudes). The z-coordinate vertical grid has 60 levels, going from 10 m at the surface to 250 m in the deep ocean to a maximum depth of 5500 m. The sub-grid scale closures and parameter settings used in this configuration are well-documented (Danabasoglu et al., 2012, 2014, 2020); some of the details are listed here to facilitate model intercomparison. The low-resolution NCAR-POP employs the skew-flux form of the GM isopycnal transport parameterization (Griffies, 1998), with depth-dependent thickness and isopycnal diffusivities (Ferreira et al., 2005; Danabasoglu and Marshall, 2007) from roughly  $3000 \text{ m}^2 \text{ s}^{-1}$  in the near surface to  $300 \text{ m}^2 \text{ s}^{-1}$  in the deep ocean. Near surface mesoscale diabatic fluxes are also parameterized (Ferrari et al., 2008; Danabasoglu et al., 2008) with diffusivity set to  $3000 \text{ m}^2 \text{ s}^{-1}$ , while the near-surface restratification effects of submesoscale mixed layer eddies are parameterized following Fox-Kemper et al. (2008, 2011). A modified version of the KPP parameterization (Large et al., 1994; Danabasoglu et al., 2006) is used for vertical mixing, with non-uniform background diffusivity that reflects tidal mixing effects (Jayne, 2009). The low-resolution NCAR-POP (but not the high-resolution) uses an overflow parameterization to represent the density-driven flows of the Denmark Strait, Faroe Bank Channel, and the Weddell Sea (Danabasoglu et al., 2010). This configuration uses hourly coupling rather than the daily coupling used in previous CORE publications (e.g., Danabasoglu et al., 2014).

For the high-resolution configuration, NCAR-POP utilizes a tripole grid with the grid northern poles in North America and Asia. It is based on versions documented in McClean et al. (2011) and Small et al. (2014), but has been updated to the CESM2 code base. The horizontal grid (nominal  $0.1^\circ$ ) varies from 11 km at the Equator to 2.5 km at high latitudes, and the vertical grid (62 levels) is the same as that used in the low-resolution NCAR-POP, but extends deeper to 6000 m. The additional two vertical levels (250 m thick) increase the maximum ocean depth from 5500 m to 6000 m, allowing for a more realistic representation of deep ocean features resolved by the  $0.1^\circ$  grid. The high-resolution NCAR-POP uses a partial bottom cell formulation of the vertical grid for more accurate representation of bathymetry. In this configuration, biharmonic horizontal mixing is used for tracers and momentum, but there is otherwise no parameterization of eddy-induced mixing. New features in the CESM2 version include the use of half-hour coupling and a modified virtual salt flux formulation that uses a local reference salinity (the estuary parameterization used in the low-resolution is not used in the high-resolution configuration, but the latter does use new methods for redistributing continental freshwater fluxes over several vertical layers near the surface).

- Moved up [2]: Parallel Ocean Program version 2 (POP2; Smith et al., 2006) contribution is based on the Community Earth System Model (CESM2) version 2.2.2 (Smith et al., 2020) ... [3]
- Deleted: contribution is based on the Community Earth System Model (CESM2) version 2.2.2 (Smith et al., 2020) ... [3]
- Moved (insertion) [8]
- Deleted: For a more detailed description of the CESM2 ocean model configuration, see the CESM2 Ocean Model Configuration ... [4]
- Deleted: ¶
- Deleted: NCAR's
- Deleted: (NCAR-L)
- Deleted: 60
- Deleted: NCAR-L
- Deleted: NCAR-L
- Deleted: s-1
- Formatted ... [5]
- Formatted ... [6]
- Deleted: s-1
- Formatted ... [7]
- Formatted ... [8]
- Formatted ... [9]
- Formatted ... [10]
- Formatted ... [11]
- Formatted ... [12]
- Deleted: NCAR-L
- Deleted: NCAR-H
- Deleted: NCAR-L
- Deleted: NCAR's
- Formatted ... [13]
- Deleted:
- Moved down [7]: (NCAR-H) is based on versions documented in Danabasoglu et al. (2010) and Danabasoglu et al. (2014) ... [7]
- Deleted: (NCAR-H)
- Moved (insertion) [7]
- Deleted: it
- Deleted: It
- Deleted: NCAR-L
- Deleted: 5
- Deleted: with 2 additional levels for a total of 62 levels
- Deleted: NCAR-H (but not NCAR-L)
- Deleted: In NCAR-H, b
- Deleted: i
- Deleted: of NCAR-H
- Deleted: NCAR-L
- Deleted: NCAR-H
- Moved up [8]: In both NCAR-H and NCAR-L simulations, the

275 2.3 AWI-FESOM

~~The AWI-FESOM model is a global configuration of the~~ Finite Element/volume Sea-ice Ocean Model (FESOM) ~~version~~  
1.4 (Wang et al., 2014; Danilov et al., 2015) of the Alfred Wegener Institute Climate Model (AWI-CM, Sidorenko et al., 2015,  
2018; Rackow et al., 2018, 2019; Sein et al., 2018). Both the ocean and sea ice modules work on unstructured triangular meshes  
(Danilov et al., 2004; Wang et al., 2008), thus allowing for multi-resolution simulations. The tracer equations employ a flux-  
corrected advection scheme, as well as the KPP scheme (Large et al., 1994) for vertical mixing. The background vertical  
diffusivity is latitude and depth dependent (Wang et al., 2014). Mesoscale eddies are parameterized by using along-isopycnal  
mixing (Redi, 1982) and Gent-McWilliams advection (Gent and McWilliams, 1990) with vertically varying diffusivity as  
implemented in Danabasoglu et al. (2008). The eddy parameterization is switched on where the first baroclinic Rossby radius  
is not resolved by local grid size. In the momentum equation, the Smagorinsky (1963) viscosity in a biharmonic form is applied.  
The sea ice thermodynamics follow Parkinson and Washington (1979), with a prognostic snow layer to account for snow to  
ice conversion. The Semtner (1976) zero-layer approach, assuming linear temperature profiles in both snow and sea ice, is  
used in this model version. The elastic-viscous-plastic (EVP, Hunke and Dukowicz, 1997) rheology is used with modifications  
that improve the convergence (Danilov et al., 2015, Wang et al., 2016). The sea surface salinity (SSS) is restored to monthly  
WOA13 climatology with a bolus velocity of 50 m over 300 days everywhere. The air-sea turbulence fluxes are calculated  
using the bulk formulation of Large and Yeager (2009). The full ocean surface velocity is used in the calculation of wind stress  
(relative wind stress). The initial conditions are derived from PHC3.0 (Steele et al., 2001).

~~For the low-resolution configuration, AWI-FESOM uses a nominal 1 degree bulk horizontal resolution which has been~~  
~~used in previous CORE-II simulations (e.g., Wang et al., 2016b),~~ with the North Atlantic sub-polar gyre region and Arctic  
Ocean set to 25 km (see Figure 1a) ~~and with a near equatorial resolution of 1/3°.~~

~~For the high-resolution configuration, AWI-FESOM uses the grid introduced by~~ Sein et al. (2016). As the variability of  
sea surface height (SSH) can manifest the variability of mesoscale eddies, the horizontal resolution is scaled with the strength  
of the observed SSH variability on this grid. In particular, the resolution is about 10 km along the western boundary currents,  
the Antarctic Circumpolar Current, and the Agulhas Current region (Figure 1b). Before generating this grid, the field of SSH  
variance is smoothed spatially to make sure that the main currents are within high resolution regions even if their positions  
change to some extent. The resolution is also increased along the coast and where the observed sea ice concentration variability  
is high. This multi-resolution grid has about 1.3M surface cells, similar to the size of a uniform 1/4° mesh. In both setups, 46  
z-levels are used, with 10 m spacing in the upper 100 m. This is slightly less than the recommended 50 levels by Stewart et al.  
(2017) to resolve the first baroclinic mode in a z-coordinate model.

- Deleted: The
- Formatted: English (United States)
- Deleted: configuration
- Deleted: ) is also examined
- Deleted: FESOM version 1.4 (Wang et al., 2014; Danilov et al., 2015) is employed in this study and all the CMIP6 simulations as well.
- Deleted: S
- Deleted: The simulations are driven with the JRA55 forcing following the OMIP protocol.
- Deleted: In
- Formatted: Space After: 0 pt
- Deleted: setup
- Deleted: the
- Deleted: bulk horizontal resolution is
- Moved (insertion) [9]
- Deleted: .
- Deleted: . N
- Deleted: is
- Deleted: This low-resolution grid has been used
- Moved up [9]: in previous CORE-II simulations (e.g., Wang et al., 2016b).
- Deleted: The
- Deleted:
- Deleted: was
- Deleted: in
- Deleted: nodes
- Deleted: layers




Figure 1. Horizontal resolution (km) of the two FESOM grids: (a) Low resolution, (b) High resolution.

## 2.4 IAP-LICOM

The IAP-LICOM model is a global configuration of the LASG/IAP Climate system Ocean Model (LICOM) (Zhang et al. 1989; Liu et al., 2004; Liu et al., 2012; Yu et al., 2018; Lin et al., 2020) developed by the Institute of Atmospheric Physics (IAP) from the Laboratory of Atmospheric Sciences and Geophysical Fluid Dynamics (LASG) of the Chinese Academy of Sciences (CAS). LICOM is the ocean component of the Flexible Global Ocean–Atmosphere–Land System model (FGOALS, e.g., Li et al., 2013, Bao et al., 2013) and of the CAS Earth System Model (CAS-ESM, M. Zhang, private communication). Version 3 of LICOM (LICOM3) is coupled to CICE4 through the NCAR flux coupler 7 (Craig et al., 2012; Lin et al., 2016). The surface salinity is restored to the monthly PHC3.0 climatology over the entire domain with a salinity piston velocity of 50

Formatted: English (United States)

Deleted: is a global ocean general circulation model

Formatted: English (United States)

Deleted: LICOM (

Deleted: ,

Deleted: also

Deleted: both

Deleted: LICOM v

Deleted: with

Deleted: Community Ice Code version 4 (

Deleted: )

Deleted:

Moved down [10]: are employed for both low- and high-resolution experiments following the OMIP protocol. However, only the thermodynamic part of CICE4 (no dynamics) was applied for high resolution.

m /4 years (50 m/30 days for the sea ice regions). ~~However, there is no freshwater flux normalization.~~ The full ocean surface velocity is used in the calculation of wind stress (relative wind stress). ~~The vertical viscosity and diffusion coefficients in the mixed layer have computed by the scheme of Canuto et al. (2001, 2002) with the background values of  $2 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$  and the upper limit of  $2 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$ . Recently, the tidal mixing scheme of St. Laurent et al. (2002) has been implemented in LICOM3 by Yu et al. (2017). The initial conditions are derived from PHC3.0.~~

~~For the low-resolution configuration, IAP-LICOM uses a Murray's (1996) tripolar Arakawa B-grid with two North "poles" at  $65^\circ\text{N}$ ,  $65^\circ\text{E}$  and  $65^\circ\text{N}$ ,  $115^\circ\text{W}$ , with a resolution of approximately 1 degree in both zonal and meridional directions ( $360 \times 218$  grid points). The vertical grid uses the eta-coordinate (Mesinger and Janjic, 1985) with 30 levels. The horizontal viscosity consists of a Laplacian with a coefficient of  $5400 \text{ m}^2 \text{ s}^{-1}$ . The mesoscale eddies are parameterized using the isopycnal tracer diffusion scheme of Redi (1982) and the eddy-induced tracer transport scheme of Gent and McWilliams (1990) with tapering factors as in Large et al. (1997) and the buoyancy frequency ( $N^2$ ) related thickness diffusivity of Ferreira et al. (2005).~~

~~For the high-resolution configuration, IAP-LICOM uses the same Murray's (1996) tripolar grid as in the low-resolution, but with two North "poles" at  $55^\circ\text{N}$ ,  $95^\circ\text{E}$  and  $55^\circ\text{N}$ ,  $85^\circ\text{W}$  and with a resolution of  $1/10^\circ$  (11 km zonally and varying from 11 km at equator to 8 km in mid-latitudes -  $3600 \times 2302$  grid points). There are 55 levels in the vertical which is just above the minimum recommended by Stewart et al. (2017) to resolve the first baroclinic mode. The additional 25 levels from the low-resolution configuration increase resolution in the deep ocean and improve the simulation of the deep circulation and the AMOC transport. The horizontal viscosity consists of a biharmonic operator with a coefficient of  $-2.8 \times 10^{10} \text{ m}^4 \text{ s}^{-1}$ . The Gent and McWilliams (1990) scheme is turned off and the tracers use a biharmonic isopycnal diffusivity with a coefficient of  $-2.8 \times 10^{10} \text{ m}^4 \text{ s}^{-1}$ . It important to note that only the thermodynamic part of CICE4 (no dynamics) was used in the high-resolution IAP-LICOM.~~

### 3 Temporal evolution and drift

#### 3.1 Mean kinetic energy

Figure 2 shows the evolution of the domain-averaged mean kinetic energy for all experiments (solid lines for the high-resolution experiments, dashed lines for low resolution experiments) from 1958 to 2018. The evolution is very similar between different models, all exhibit a quick spin-up of the kinetic energy in the first five years which levels off for the rest of the integration. Not surprisingly, the total kinetic energy is significantly higher for the high-resolution experiments over the low-resolution experiments. For the high-resolution configurations, the FSU-HYCOM has the highest kinetic energy, with a globally averaged value of  $\sim 35 \times 10^{-4} \text{ m}^2 \text{ s}^{-2}$  and the IAP-LICOM has the lowest kinetic energy, with a globally averaged value of  $\sim 15 \times 10^{-4} \text{ m}^2 \text{ s}^{-2}$  in the high-resolution configuration. For comparison, a previous high-resolution  $1/10^\circ$  global simulation, performed with an older version of POP by Maltrud and McClean (2005) using daily NCEP/NCAR reanalysis forcing and absolute winds in wind stress calculations, has a global averaged kinetic energy at  $25\text{-}30 \times 10^{-4} \text{ m}^2 \text{ s}^{-2}$  (see their Figure 1). The higher kinetic energy in FSU-HYCOM can be partially explained by the wind stress formulation which does not take into

**Deleted:** LICOM3 is an ocean model with free sea surface. It uses the primitive equations with Boussinesq and hydrostatic approximations and is configured with the

**Deleted:** for the low-resolution experiment and at  $55^\circ\text{N}$ ,  $95^\circ\text{E}$  and  $55^\circ\text{N}$ ,  $85^\circ\text{W}$  for the high-resolution experiment

**Deleted:** The horizontal grid is the Arakawa B-grid

**Deleted:** for the low-resolution experiment and  $1/10^\circ$  for the high-resolution experiment (11 km zonally and varying from 11 km at equator to 8 km in mid-latitudes)

**Deleted:** and 55

**Deleted:** for the low- and high-resolution experiments, respectively

**Deleted:** The low-resolution and high-resolution LICOM3 have a total of  $360 \times 218$  and  $3600 \times 2302$  grid points in the horizontal, respectively. The two-step preserving shape advection scheme (Yu, 1994; Xiao, 2006) and the implicit vertical viscosity/diffusivity (Yu et al., 2018) were adopted for both momentum and tracer equations. The split-explicit Leapfrog with Asselin filter is used for the time integration of both momentum and tracer.

**Formatted:** Space After: 0 pt

**Deleted:** ¶  
The vertical viscosity and diffusion coefficients in the mixed layer have computed by the scheme of Canuto et al. (2001, 2002) with the background values of  $2 \times 10^{-6} \text{ m}^2/\text{s}$  and the upper limit of  $2 \times 10^{-2} \text{ m}^2/\text{s}$ . Recently, a tidal mixing scheme of St. Laurent et al. (2002) has been adopted in LICOM3 by Yu et al. (2017). The Laplacian form with the coefficient of  $5400 \text{ m}^2/\text{s}$  was chosen for the horizontal viscosity in the low-resolution version and the biharmonic form with [... [14]

**Deleted:** has been

**Deleted:**

**Deleted:** form of the

**Deleted:** the

**Deleted:** Besides, the chlorophyll-a dependent solar pene [... [15]

**Moved (insertion) [10]**

**Deleted:** /s

**Deleted:** was applied

**Deleted:** are employed for both low- and high-resolution [... [16]

**Deleted:** was applied for high resolution

**Deleted:** Models time

**Formatted:** English (United States)

**Deleted:** /s<sup>2</sup>

**Deleted:**

**Deleted:** /s<sup>2</sup>

**Deleted:**

**Deleted:**

**Deleted:** /s<sup>2</sup>

**Deleted:**

account the ocean current velocities (absolute winds) while the other three models do (relative winds). The latter has an eddy killing effect that can reduce the total kinetic energy by as much as 30% (see Renault et al., 2020, for a review). This is roughly the difference that is seen between FSU-HYCOM and NCAR-POP, and POP with absolute winds in the wind stress (Maltrud and McClean, 2005) has a level of kinetic energy that is close to FSU-HYCOM. But even with the highest resolution used here (~0.1°), the total kinetic energy remains significantly lower than what can be inferred from observations and higher resolution models (closer to  $50 \times 10^{-4} \text{ m}^2 \text{ s}^{-2}$ , i.e. Chassignet and Xu (2017)). The increase in total kinetic energy from the low- to the high-resolution configuration is approximately a factor of 4 for all models, except for AWI-FESOM (factor 2 only). This is probably because the high-resolution AWI-FESOM has a variable grid spacing (Figure 1b) and does not resolve the Rossby radius of deformation everywhere. The level of total kinetic energy is substantially lower in IAP-LICOM, most likely because of the two-step shape preservation advection scheme used in the momentum equations and higher diffusivity (Table 1).





Figure 2: Time evolution of the domain-averaged kinetic energy ( $10^{-4} \text{ m}^2 \text{ s}^{-2}$ ) for all experiments.

3.2 Global mean sea level, temperature, and salinity

As stated by Griffies et al. (2014), “[t]he CORE [and OMIP] protocols (Griffies et al., 2009; Danabasoglu et al., 2014; Griffies et al., 2016) introduce a negligible change to the liquid ocean mass (non-Boussinesq) or volume (Boussinesq), and the salt should remain nearly constant (except for relatively small exchanges with the sea ice)”. Changes to the simulated global mean sea level should arise only because of thermosteric effects (i.e., changes in ocean heat content and redistribution of heat) in simulations that preserve salt content (i.e., that have zero net surface freshwater flux). In most models, the global sea level time evolution (Figure 3) is dominated by changes in the global mean ocean temperature (Figure 4a). IAP-LICOM is the exception, in which the global sea level shows a downward trend until ~1990, and then slowly rises. This is due to an increase in global mean salinity (Figure 4b) which dominates the global sea level changes despite a large increase in global volume mean temperature (Figure 4a). This increase is explained by the lack of surface restoring salt-flux normalization in IAP-

Deleted: 19

Deleted: /s<sup>2</sup>



Deleted:

Formatted: English (United States)

Deleted: /

Deleted: 2

Formatted: English (United States)

LICOM (see section 2.4). For details on the salt flux normalization used in the other models, the reader is referred to Appendix B.3 of Griffies et al. (2009) and Appendix C in Danabasoglu et al. (2014), which describe the techniques used to ensure that there is no net salt added to or removed from the ocean-sea ice system (Griffies et al., 2014). One can also note that most models show an increase in global temperature and sea surface height after 1980-90 associated with rising air temperature.





Figure 3: Time evolution from the initial conditions of the global steric sea surface height change for all experiments.

An increase in the horizontal resolution does not necessarily imply a reduction in temperature and salinity drift and no coherent picture emerges from the comparison. If one focuses on the time evolution of the globally averaged temperature in the upper 700 m (Figure 4c), there are only small changes for AWI-FESOM and IAP-LICOM, while FSU-HYCOM shows a significant reduction in the drift and NCAR-POP an increase as resolution is increased. For salinity in the upper 700 m (Figure 4d), the increase in resolution significantly reduces the drift in NCAR-POP and FSU-HYCOM, no changes for IAP-LICOM, and a significant increase for AWI-FESOM. It is important to note that, while the salt flux restoring may differ among the four models, it remains the same for each model as the resolution is increased. When considering the upper 2000 m (Figure 4f), there is a significant reduction in the global salinity drift for NCAR-POP and FSU-HYCOM, less so for IAP-LICOM, and no changes for AWI-FESOM. We note that most high-resolution models (FSU-HYCOM being the exception) warm faster over the upper 2000 m and global temperature than their lower-resolution partners, which is not true for the upper 700 m. Figure 5 contains identical data as in Figure 4, except that, by rebasing anomalies to the final year, it highlights the forced ocean variations of the last few decades of simulation by comparing the modeled global temperature/salinity and heat/salt content change to that of World Ocean Atlas 18 (WOA18). The comparison shows that high resolution improves the fidelity and reduces the spread of forced ocean heat content change (particularly 0-2000 m heat content) over the recent decades. Figure 6 shows in more detail the evolution of the global temperature and salinity from the initial conditions as a function of depth. AWI-FESOM shows the smallest changes in temperature throughout the water column, but the largest in salinity despite having a salt flux adjustment to ensure that the global salinity remains constant (shown in Figure 2). There is a significant freshening in the upper 400 m compensated by a salinification in the deeper ocean. This is more pronounced in the high-resolution experiment. Increasing the resolution significantly improves the drift in both temperature and salinity in FSU-HYCOM, but not so much in the other simulations. One could actually argue that the drift is stronger in NCAR-POP with a

Deleted: The

Deleted: is given in



Deleted:

Formatted: English (United States)


Deleted: all

Deleted: 5




significant warming in the upper 400 m and freshening in the upper 100 m. While it is beyond the scope of this paper, additional insights could be gained by computing vertical heat and salt budgets as in Griffies et al. (2015) and Von Storch et al. (2016).

In the next section, we investigate in more details the evolution of the temperature and salinity as function of depth and time by ocean basin.




Formatted: English (United States)




Deleted:  
Formatted: Check spelling and grammar

515 **Figure 4: Time evolution of global temperature (°C) and salinity (psu) change (relative to initial conditions) for all experiments**  
(depth-integrated, upper 700 m, upper 2000 m).




520 **Figure 5: Time evolution of global temperature (°C) and salinity (psu) change (relative to year 1818) for all experiments and World**  
**Ocean Atlas (WOA18) (depth-integrated, upper 700 m, upper 2000 m).**






**Figure 6:** Time evolution of global temperature (°C) and salinity (psu) change (relative to initial conditions) as a function of depth for all experiments.




**Figure 7:** Time evolution of Atlantic Ocean temperature (°C) and salinity (psu) change (relative to initial conditions) as a function of depth for all experiments.



Deleted:

Formatted: English (United States)


Deleted: 5... Time evolution of the...global temperature (°C) and salinity (psu) change (relative to initial conditions) (... [17])




Deleted:

Formatted: English (United States)


Deleted: 6... Time evolution of the...Atlantic Ocean temperature (°C) and salinity (psu) change (relative to initial conditions) (... [18])



**Figure 8:** Time evolution of Pacific Ocean temperature (°C) and salinity (psu) change (relative to initial conditions) as a function of depth for all experiments.



**Figure 9:** Time evolution of Indian Ocean temperature (°C) and salinity (psu) change (relative to initial conditions) as a function of depth for all experiments.




Deleted:

Formatted: English (United States)

Deleted: 7

Deleted: the

Deleted: s



Deleted:

Formatted: English (United States)

Deleted: 8

Deleted: the

Deleted: s

3.3 Temperature and salinity bias (depth vs. time) by ocean basin

565 Figures 7, 8, and 9 show the time evolutions of the horizontal-mean depth profiles of the temperature and salinity for the Atlantic, Pacific, and Indian Oceans, respectively. The color bar is the same in all figures, including Figure 6 (global), therefore allowing for a qualitative estimate of where the drift is most significant. To a large extent, the time evolution in each of the major ocean basins mimics that of the global, but with some significant differences. For the Atlantic Ocean (Figure 7), there is a surface freshening in the upper 100 to 200 m as resolution is increased in FSU-HYCOM, NCAR-POP, and AWI-FESOM. IAP-LICOM, on the other hand, becomes more saline and warmer in the upper 100 m. The latter is accompanied by a large 570 freshening and cooling below 100 m to approximately 600 m. Overall, the drift is smaller in the Pacific for most models, with a freshening in the upper ocean for all models as resolution is increased. The exception is again IAP-LICOM which shows a significant increase in salinity in the upper 200 m and this could be a consequence of the fact that there is no zero normalization of the surface restoring salt-flux. The temperature bias in the Pacific Ocean is similar to that of the global ocean. In the Indian Ocean (Figure 9), FSU-HYCOM and NCAR-POP exhibit larger biases than AWI-FESOM and IAP-LICOM, but those biases 575 are smaller at the higher resolution.

3.4 Mean temperature and salinity bias

The drift plots from sections 3.2 and 3.3 indicate that temperature and salinity bias structures are well established within the first two decades of the spin up so that time averages computed over the latter decades of the simulations should provide a reasonable estimate of the stationary biases characterizing each model. Figure 10 shows latitude-longitude maps of surface 580 temperature and salinity differences computed over the 1980-2018 time period with respect to WOA18. All models exhibit reductions in SST bias as resolution is increased. The largest bias reductions are seen in the Southern Ocean, in the Agulhas retroflection region, and along the Gulf Stream extension in the North Atlantic. All these changes are presumably primarily related to an improved representation of the pathways of strong surface currents in those regions, which could result not only from differences between resolved and parameterized eddies but also from differences in resolved bathymetry. SST bias in 585 eastern boundary upwelling regions has been shown to be strongly sensitive to atmospheric forcing resolution (Tsujino et al., 2020), somewhat sensitive to the regridding technique used for near-surface winds (Small et al., 2015), and also slightly sensitive to ocean resolution for seasonal means (Kurian et al., 2020). However, Figure 10 shows minimal change in annual mean SST bias in the upwelling zones off the west coasts of Namibia, Peru, and North America. In addition to the Gulf Stream, all models exhibit some degree of bias reduction in the Brazil-Malvinas Confluence zone at high resolution, but no such systematic improvement is evident in the Kuroshio-Oyashio extension region. Sea surface salinity bias is generally reduced 590 globally in the high-resolution simulations, with the exception again being AWI-FESOM which exhibits a more pronounced negative salinity bias in line with the enhanced near-surface salinity drift at high resolution in that model (cf. Figures 6-9). The SSS bias in the Arctic results from the differences in salinity between WOA18 and the climatologies used in the salinity restoring (GDEM, WOA13, or PHC3.0, see section 2 for details).


Deleted: 6  
Deleted: 7  
Deleted: 8  
Formatted: English (United States)  
Deleted: 5  
Deleted: 6

Deleted: 8

Deleted: odel m  
Deleted: versus initial climatology  
Formatted: English (United States)  
Deleted: 2b,c  
Deleted: 9  
Deleted: biases  
Deleted: final 10 years of integration (historical years 2009-2018)  
Deleted: the climatology used to initialized the models (GDEM, WOA13, or PHC3.0, see section 2 for details)  
Deleted: With the possible exception of AWI-FESOM, a  
Deleted: clear


Deleted: 9

Deleted: 5-8



**Figure 10: Modeled surface temperature (°C) and salinity (psu) difference from WOA18 over the 1980-2018 time period.**

The vertical structure of mean temperature and salinity bias is displayed as zonal-averages across the different ocean basins in Figures 11-13. This reveals that the near-surface global drift toward warming seen in FSU-HYCOM and NCAR-POP is partly related to a degradation of the tropical thermocline in all basins. One hypothesis is that the thermocline bias is related to the representation of vertical eddy heat flux (Griffies et al. 2015), which tends to be stronger and more realistic in high-resolution simulations (see Figure 5). In FSU-HYCOM, higher resolution improves the situation, while in NCAR-POP, the tropical thermocline temperature bias actually gets worse with resolution. The degradation in thermocline bias in POP high-resolution could also be due to missing submesoscale physics, which are parameterized in the low-resolution configuration, but absent in the high-resolution. In both FSU-HYCOM and NCAR-POP, and to some extent IAP-LICOM, large extratropical/polar temperature biases associated with intermediate and deep waters decrease with enhanced resolution. In AWI-FESOM, both high and low latitude biases get worse. It turns out that the relatively low near-surface temperature drift seen in the AWI-FESOM model (Figures 6-9) is related to large, but largely compensating, anomalies of opposite sign at each depth level. Cold/warm anomalies that develop in the Tropics and at high latitudes in that model at both resolutions tend to disappear in global or basin-wide means. Such compensation does not occur for salinity which is too fresh in the thermocline and too saline in deeper intermediate waters in AWI-FESOM, a characteristic that gets worse as resolution is increased. The



Deleted:

Formatted: English (United States)

Deleted: 9

Deleted: Modeled s

Deleted: with the climatology used to initialize the model

Deleted: 0

Deleted: 2

Deleted: 5

Deleted: 8



IAP-LICOM model exhibits some large changes in the sign and structure of zonal mean bias as resolution changes, but as with AWI-FESOM, it does not lend strong support to the hypothesis that model temperature and salinity bias can be reduced by increasing model horizontal resolution. As already mentioned, the NCAR-POP model exhibits improved representation of high latitude intermediate and deep waters, but degraded representation of the tropical thermocline, as resolution is enhanced. FSU-HYCOM is the only model that shows near ubiquitous reduction in temperature/salinity bias in all ocean basins in the high-resolution version; for the other models, greatly enhanced horizontal resolution does not deliver unambiguous bias improvement in all regions. This is a rather unexpected result.





Figure 1: Global zonal temperature (°C) and salinity (psu) difference with the climatology used to initialize the model.



660 **Figure 12:** Atlantic zonal temperature (°C) and salinity (psu) difference with the climatology used to initialize the model.

**Deleted:**

**Formatted:** English (United States)

**Deleted:** 1




Figure 13: Indo-Pacific zonal temperature (°C) and salinity (psu) difference with the climatology used to initialize the model.

### 3.5 Northern and southern hemisphere sea ice volume and extent

Due to observational constraints, sea ice is often quantified and monitored by passive microwave satellites in terms of sea ice extent (SIE), defined as the sum of all grid areas with 15% or higher sea-ice concentration. Figures 14a-b display the modeled annual mean northern and southern hemisphere sea-ice extent for all simulations and include comparisons to the latest observations from the National Snow and Ice Data Center (NSIDC) (Fetterer et al., 2017). Similar to the observations of the 5<sup>th</sup> and 6<sup>th</sup> CMIPs (Stroeve et al., 2012; Notz et al., 2020; Shu et al., 2020), the multi-model mean is representative of the passive microwave mean SIE from 1978-2018, with large inter-model differences. All models show a clear decline in SIE in the northern hemisphere and weaker trends in the southern hemisphere. Because observed sea-ice extent is highly correlated with near-surface air temperature (e.g., Olonscheck et al., 2019), the consistency of the temporal variability between different simulations and observations suggests that this consistency is driven by a realistic JRA55-do forcing. In general, in both hemispheres, an increase in resolution reduces SIE bias, with the exception of the IAP-LICOM model.


Indo-Pac Temperature 1980-2018 mean bias FSU-HYCOM
Indo-Pacific Temperature 1980-2018 mean bias NCAR-POP

HIGH RES.
LOW RES.

Indo-Pac Salinity 1980-2018 mean bias FSU-HYCOM
Indo-Pacific Salinity 1980-2018 mean bias NCAR-POP


HIGH RES.
LOW RES.

Deleted:
Formatted: English (United States)
Deleted: 2
Deleted: -
Formatted: English (United States)
Formatted: No underline, English (United States)
Formatted: English (United States)
Formatted: No underline, English (United States)
Formatted: English (United States)
Formatted: No underline, English (United States)
Formatted: English (United States)
Formatted: No underline, English (United States)
Formatted: English (United States)
Formatted: No underline, English (United States)
Formatted: English (United States)
Formatted: No underline, English (United States)
Formatted: English (United States)
Formatted: No underline, English (United States)
Formatted: English (United States)
Formatted: No underline, English (United States)
Deleted: Sea ice is usually quantified and monitored by satellite in terms of sea ice extent, defined as the area with 15% or high ... [19]



**Figure 14:** Modeled time evolution of annual mean Arctic and Antarctic a-b) sea ice extent in  $10^{12} \text{ m}^2$  and c-d) sea ice volume in  $10^{13} \text{ m}^3$ . The black lines are observations from National Snow and Ice Data Center (NSIDC) and results from Pan-Arctic Ice Ocean Modeling and Assimilation System (PIOMAS) (Schweiger et al., 2011).

Sea ice volume (SIV) is a second independent measure of sea ice simulation performance that is representative of sea ice state, determined by the thermodynamical, optical, and dynamical properties of the ice itself and the sensitivity of modeled SIV to model formulation is insightful for understanding the physical realism of the sea ice schemes. The modeled time evolution of annual mean northern/southern hemisphere SIV is displayed in Figures 14c-d. Relative to SIE, there is significant inter-model spread in SIV. With the exception of IAP-LICOM, all simulations exhibit a continuous decline of northern hemisphere sea-ice volume during the satellite era, with trends similar to that of PIOMAS (Schweiger et al., 2011). Changes to SIV also do not appear to be consistent among the models when moving from low- to high-resolution: it increases for FSU-HYCOM and IAP-LICOM and decreases for NCAR-POP and AWI-FESOM. IAP-LICOM is a clear outlier in both hemispheres, which points to issues in the numerical implementation of its sea ice code at higher resolutions: sea ice thickness on average increases as the resolution is increased despite declining SIE. The IAP-LICOM sea-ice model has only thermodynamic processes, and the absence of dynamic sea-ice may lead to an excessive accumulation of volume.



Deleted:

Formatted: English (United States)

Deleted: 3

Deleted: ng

Deleted: s



### 3.6 Meridional overturning circulation

The Atlantic meridional overturning circulation (AMOC) is often quantified by a meridional overturning streamfunction with respect to depth,  $\psi_z$ , at each latitude, defined as the meridional transport (in Sv) across the basin above a constant depth  $z$ :

$$\psi_z(y, z) = \overline{\int_a^0 v(x', y, z', t) dz' dx'},$$

where  $v$  is the 4-d meridional velocity, the overbar denotes a time average (here annual mean), and the  $x$  integration covers the entire span of the Atlantic basin. The magnitude of the AMOC, or the AMOC transport, is then defined as the maximum of the streamfunction  $\psi_z$  with respect to  $z$ , representing the total northward transport above the overturning depth (approximately 1000 m in the Atlantic Ocean). This important measure of the AMOC is quantified and monitored by the RAPID moorings near 26.5°N (e.g., Smeed et al., 2018).

The modeled annual mean AMOC transport across 26.5°N from 1958 to 2018 is displayed in Figure 15a, along with the RAPID results (2005-2017) for reference. The vertical distribution of the time mean zonal streamfunction in the Atlantic Ocean is shown later in Figure 26. For the four models, the high-resolution simulations have a higher AMOC transport than their low-resolution counterparts: the mean AMOC transport in 2004-2018 is 7.8-14.9 Sv in the four low resolution simulations, versus 14.0-19.8 Sv in the four high-resolution (17.2 ± 1.5 Sv as observed by RAPID, e.g. Smeed et al., 2018). However, there is not an obvious difference in the AMOC sensitivity to forcing or in the trend between the low- and high-resolution experiments. The simulated time evolution of the AMOC transports at this latitude depends on the model, its parameterizations, resolution, and on the number of spin-up cycles (Danabasoglu et al., 2016). Despite the fact that the simulations were only integrated over one JRA55-do cycle (1958-2018), all four low-resolution models show a similar multi-decadal variability with a transport decrease from 1958 to late 1970s, an increase from the late 1970s to late 1990s, and a decrease again thereafter. This multi-decadal variability is also present in the CORE-II simulations (Danabasoglu et al., 2016; Xu et al., 2019). In the high-resolution simulations, FSU-HYCOM, NCAR-POP, and AWI-FESOM exhibit a similar multi-decadal variability (of low transports in the late 1970s and high transports in the early 1990s) as seen in previous basin-scale simulations (e.g., Böning et al., 2006; Xu et al., 2013). The decline of the AMOC transport from early 1990s to 2000s may be a consequence of the warming and reduced deep convection in the western subpolar North Atlantic that has been documented quite extensively (e.g., Hakkinen and Rhines, 2004; Yashayaev, 2007). The high-resolution AWI-FESOM model has a strong AMOC decreasing trend (of 4-5 Sv) initially during the first 20 years and then has a time evolution of the AMOC that is very similar to that of FSU-HYCOM and NCAR-POP. The high-resolution IAP-LICOM shows an increasing AMOC transport from 1958 to the early 1990s (by 10 Sv) and decrease thereafter (by 5 Sv), which is quite different from the other three models, but it does appear to capture the observed increase over the last few years.

**Deleted:** Another measure of the sea-ice simulation performance is the sea-ice volume. As there are no reliable long-term sea ice volume observations, we use the model result from the Pan-Arctic Ice Ocean Modeling and Assimilating system (PIOMAS, Schweiger et al., 2011) at the Polar Science Center to assess the northern hemisphere sea-ice volume in our simulations. The modeled time evolution of the northern/southern hemisphere sea-ice volume is displayed in Figures 13c-d. Despite differences in the mean averaged values, all but one simulation exhibit a continuous decline of the northern hemisphere sea-ice volume that is comparable to the PIOMAS observations (Figure 13c) and this result is not significantly influenced by the model's horizontal resolution. The outlier simulation is the high-resolution IAP-LICOM which shows an overall increase of northern hemisphere sea-ice volume (despite a decline in sea ice extent). This because its sea-ice model has only thermodynamic processes and the absence of dynamic sea-ice leads to an excessive accumulation and volume. For the southern hemisphere, the time evolution of the modeled sea-ice volume is also similar for all simulations, except again for the high-resolution IAP-LICOM.

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)


**Formatted:** English (United States)

**Formatted:** English (United States)


**Deleted:** 4

**Deleted:** 5

**Deleted:** only



**Figure 15:** Modeled annual mean Atlantic meridional overturning circulation (AMOC) transports at a) 26.5°N and b) 34°S; and c) global meridional overturning circulation (MOC) transports at 34°S. The global MOC transport is negative because the northward flowing Antarctic bottom water (AABW) is below the southward return flow (see Lumpkin and Speer (2007) and Talley (2013) for observed values). The solid black line in panel a) is derived from the RAPID array measurements (Smeed et al., 2018).



Deleted:

Formatted: English (United States)

Deleted: 4

Formatted

Deleted: Modeled annual mean Atlantic meridional overturning circulation (AMOC) transports at a) 26.5°N and b) 34°S; and c) global Meridional overturning circulation (MOC) transports at 34°S. Solid black line in panel a) are results from RAPID array.¶

795 The AMOC transport in the South Atlantic Ocean has been quantified near 34°S, using several observational techniques including expendable Bathythermograph (XBT), Argo profiles, and moored current meter arrays in the western and eastern boundaries (Baringer and Garzoli, 2007; Dong et al., 2009, 2014, 2015; Garzoli et al., 2013; Goes et al., 2015; Meinen et al., 2013; 2018) and these observations yield a time mean AMOC transport of about 14-20 Sv (see Xu et al. (2020a) for an in-  
800 depth discussion of the circulation in the South Atlantic Ocean). The modeled temporal evolution of the AMOC transports at this latitude are overall similar to that at 26.5°N in the North Atlantic (Figures 15a-b). The range of the modeled time mean AMOC transport at 34°S in the high-resolution simulations, from 14.7 Sv in FSU-HYCOM to 20.1 Sv in IAP-LICOM, is about the same as the observational range mentioned above.

The Antarctic Bottom Water (AABW) cell of the global overturning circulation (Talley, 2013) can be visualized as a streamfunction similar to Equation 1, except that the zonal integration now spans across the full longitude circle. The transport associated with this cell is defined as the minimum of the streamfunction that is typically found near 3500 m (Lumpkin and  
805 Speer, 2007) and is a measure of the northward flow of the near bottom water (AABW and/or Circumpolar Deep Water, CDW) from the Southern Ocean into the Atlantic, Pacific, and Indian Oceans that eventually upwells and returns southward at a shallower depth. Several observational estimates are provided based on hydrographic sections near 32°S, but the uncertainty remains quite high (29±7.6 Sv in Talley (2013)). Figure 15c displays the modeled transport at 34°S. The low-resolution  
810 simulations show a consistently lower transport, with the mean transport value for 2004-2018 ranging from 7.4 Sv in IAP to 15.3 Sv in FSU-HYCOM. The low transport in IAP-LICOM is due to a long-term downward trend in this simulation: from 20 Sv in the early 1960 to 5 Sv in the late 2010s. The other three models, especially FSU-HYCOM, exhibit relatively stable transports in the last 30 years (of the integration). The high-resolution simulations show a much wider range, with time mean transports (2004-2018) ranging from 11.7 Sv in IAP-LICOM to 26.5 Sv in AWI-FESOM. The time evolution of the 34°S  
815 transports in the high-resolution simulations typically shows an increase in the early stage of the integration followed by a gradual decrease afterward. But the timescale of the increase and decrease periods varies significantly between models. As a result, the transport is relatively stable in IAP-LICOM and AWI-FESOM for the last 30 years of the integration but continues to decrease slowly in the NCAR-POP and FSU-HYCOM. Interestingly, despite the large difference in time mean transports, all four high-resolution models exhibit a similar interannual variability at this latitude for the last 30 years of the integration.

820 **3.7 Drake Passage and Indonesian Throughflow transports**

The Antarctic Circumpolar Current (ACC) is a strong oceanic current that flows eastward around Antarctica. It connects the Pacific, Atlantic, and Indian Oceans to its north and is the primary means of inter-basin exchange, enabling a truly global overturning circulation (e.g., Gordon, 1986; Schmitz, 1996; Talley, 2013). An accurate knowledge of the ACC transport is fundamental to understand its influence on the global circulation. Substantial observational efforts have been made toward  
825 quantifying the ACC transport, especially in Drake Passage, where the ACC is constricted between the Antarctica and the southern tip of the South America. The major observations include: 1) the International Southern Ocean Studies (ISOS) program in the late 1970s and early 1980s (Whitworth, 1983; Whitworth and Peterson, 1985); 2) the repeat hydrographic

Formatted: English (United States)

Deleted: 4


Deleted: 4

Deleted: 11

Formatted: English (United States)


Formatted: English (United States)

sections along the World Ocean Circulation Experiment (WOCE) line SR1b (e.g., Cunningham et al., 2003); 3) the repeat shipboard ADCP surveys that directly measure the current in the top 1000 m of the water column (Firing et al., 2011); 4) the DRAKE program that combines a moored current meter array (2006-2009) and satellite observations (e.g., Koenig et al., 2014), and 5) the cDrake program of a high-resolution bottom current meter mooring array and cPIES observations from 2007 to 2011 (e.g., Chidichimo et al., 2014; Donohue et al., 2016). The time mean, full-column transport estimates range from 134±15 Sv based on ISOS, to 141±2.7 Sv based on DRAKE program, and to 173.3±10.7 Sv based on the cDrake results.



**Figure 16:** Modeled annual mean volume transport of the ACC through Drake Passage and of the Indonesian Throughflow (ITF) from Pacific to Indian Ocean. The dotted lines in a) is the observational range between the canonical transport value (134 Sv) based on the ISOS program and 173 Sv based on cDrake observations. The dotted line in b) is the observed ITF transport value (15 Sv) based on INSTANT observations.

The modeled annual mean Drake Passage transport is displayed in Figure 16a together with the mean observational range (134-173 Sv). The transport in the low-resolution simulations differ from each other's: IAP-LICOM and AWI-FESOM show a continuous decrease; FSU-HYCOM shows a continuous increase; whereas NCAR-POP shows an increase in the first 20 years, then levels off for the next 40 years of the integration. It is expected that the low-resolution models will be sensitive to different parameterizations and their ability to simulate eddy saturation (Gent and Danabasoglu, 2011; Munday et al. 2013). As a result, the time mean transport for the last 15 years of these low-resolution simulations are either near the edge or outside of the above observational range: 132, 135, 173, and 190 Sv, respectively, for IAP-LICOM, AWI-FESOM, NCAR-POP, and FSU-HYCOM. The evolution of Drake Passage transport is slightly more similar in high-resolution simulations (compared to the low-resolution counterparts). Even at high resolution, mean ACC transport is sensitive to subgrid schemes (Pearson et al. 2017). Three models show a fast decrease in the first 10-15 years and then a very small decrease thereafter, except that FSU-HYCOM shows a small increase in the first 20-years and then levels off thereafter. The high-resolution models exhibit a higher



Deleted:  
Formatted: English (United States)  
Deleted: 5  
Deleted: 5  
Deleted: based on the most recent observations

interannual variability correlation than in the low-resolution models. The ACC transports in NCAR-POP and AWI-FESOM, 120 and 130 Sv, respectively, are lower than the latest best mean transport estimate (Xu et al., 2020b), but still within the uncertainty range; while the transports in IAP-LICOM and FSU-HYCOM, 145 and 157 Sv, respectively, are within the mean observational range.

The long-term trend in the modeled ACC transport merits some further discussions. Observations indicate that, as shown by a higher Southern Annular Mode index, the westerly winds in the Southern Ocean have been intensified since the 1970s. This increase in westerly winds is also present in the JRA55-do forcing, yet none of the high-resolution simulations exhibit a long-term upward trend (three models even have a slightly weakening trend). The ACC transport time series from DRAKE program (Koenig et al., 2014) as well as from the WOCE line SR1b also show a stable ACC transport from early 1990s. Given that the ACC is wind-driven, a lack of dependence for the ACC transport on the zonal wind at long-term timescale may be at first surprising, but it has also been shown that the strengthening of winds does generate more eddy activity along the path of the ACC, without necessarily changing its total transport (e.g., Hallberg and Gnanadesikan, 2006; Morrison and Hogg, 2013; Munday et al., 2013; Farneti et al. 2015). Following this argument, one may expect to see an increasing ACC transport in low-resolution, non-eddying simulations as long as the coefficient used in the eddy parameterization (i.e., GM) remains constant (Gent and Danabasoglu, 2011). Only FSU-HYCOM shows a long-term increase in the ACC transport and two models (IAP-LICOM, and AWI-FESOM) even show a long-term decrease. This may be a consequence of the eddy parameterization choice made by the different models, assuming that the ACC is quasi-equilibrated and that changes are not related to buoyancy changes in the Southern Ocean (e.g., rate of bottom water formation). There is, however, an initial condition transient in the buoyancy gradients across the ACC (Figure 15c) and the MOC across the ACC is equilibrating for a significant fraction of the integration time: one cannot rule out that the ACC changes result from buoyancy drifts. All models have weaker Drake Passage transports at higher resolution, except for IAP-LICOM.

The Indonesian Throughflow Flow (ITF) connects the tropical Pacific and Indian Oceans and provides a pathway for inter-basin exchange of heat and freshwater. Tropical Pacific and Indian Oceans water masses can go through the ITF and contribute to the upper AMOC limb in the South Atlantic Ocean as part of the global MOC via the Agulhas Current (e.g., Gordon, 1986; Schmitz, 1996; Talley, 2013). A good overview of the Indonesian Sea oceanography and the ITF observations can be found in Gordon (2005) and Gordon et al. (2010). The 3-year mean total ITF transport measured by the International Nusantara Stratification and Transport (INSTANT) program in 2004-2006 is  $15 \pm 2.5$  Sv (Sprintall et al., 2009; Gordon et al., 2010), about 30% greater than the values of non-simultaneous measurements made prior to 2000. The ITF transport variability exhibits a close relationship to the phase of El Niño—Southern Oscillation (Meyers, 1996).

The modeled annual mean ITF transport is displayed in Figure 16b. The low-resolution simulations exhibit a wide range of the ITF transports, ranging from about 5 Sv in the IAP model, to 11 Sv in NCAR-POP model, and to 15 Sv in FSU-HYCOM and AWI-FESOM (30-year mean). The ITF transports exhibit a gradual decrease in AWI-FESOM, but are relatively stable in the other three models. In the high-resolution simulations, however, the models yielded a quite similar ITF transport, especially in the second half of the integration. All four models represented a similar mean transport of 15 Sv that is close to observations

Deleted: I  
Formatted: English (United States)  
Deleted: among these models is more correlated  
Formatted: English (United States)  
Deleted: er  
Formatted: English (United States)  
Deleted: slightly  
Formatted: English (United States)  
Deleted: s from the latest observations, but still  
Formatted: English (United States)

Deleted: is likel  
Deleted: y

Deleted: another key  
Deleted: mean of water mass as well as  
Deleted: It connects the tropical Pacific and Indian Oceans directly and, through  
Deleted: the  
Deleted: Leakage  
Deleted: , eventually feeds into the upper AMOC limb in the South Atlantic Ocean as part of the global MOC scheme

Deleted: 5

and all show a very similar variability. While large-scale forcing mechanisms are thought to set the basic level of ITF transport (Godfrey, 1989), the difference between the low-resolution and high-resolution simulations is likely impacted by the model's ability to represent the small-scale bathymetry feature of the Indonesian Seas (Jochum et al., 2009), after all, the ITF enters the Indian Ocean through a number of narrow straits that are hard to represent in the low-resolution models. IAP-LICOM and NCAR-POP have low ITF transports at low-resolution and appear to have stronger sensitivity to model resolution.

#### 4 Stationary ocean climate

##### 4.1 Sea Surface Height (SSH) and Eddy Kinetic Energy (EKE)

The impact on the circulation of increasing the horizontal resolution is two-fold. First, the solution becomes more nonlinear and allows for a better representation of western boundary currents. Second, the first Rossby radius of deformation is resolved over most of the globe (Hallberg, 2013) and eddies are formed through barotropic and baroclinic instabilities, although higher vertical modes including submesoscale eddies are not often resolved (nor are they resolved by altimetry, i.e. AVISO) (c.f., Stewart et al., 2017). Figure 17 displays the mean sea surface height bias with respect to AVISO and its 5-day variance over the 1993-2018 period. Overall, the large-scale patterns are well represented globally, and one can clearly see the improved representation of the western boundary currents as resolution is increased (Gulf Stream and Kuroshio). In the North Atlantic, most coarse resolution models to date have the tendency to exhibit an overshooting Gulf Stream and a poor representation of the North Atlantic Current (NAC) at the Northwest Corner. This is the case for three out of the four models. Instead of flowing northeastward along the continental rise past the Flemish Cap and continuing northwestward as in the observations (Rossby, 1996), the North Atlantic Current is strongly zonal in NCAR-POP, AWI-FESOM, and IAP-LICOM, and does not turn northeastward at the Flemish Cap. This has been a long-standing issue for many ocean components of the CMIP climate models and it does not necessarily improve as the computational mesh is refined. Increasing the horizontal resolution did improve the Gulf Stream separation (see Chassignet and Marshall (2008) and Chassignet and Xu (2017) for a review) in all models, but not necessarily the representation of the Northwest corner circulation (Figure 17 - see Figure 1 of Chassignet et al. (2020) for details). FSU-HYCOM is the only model that has a good representation of the Northwest corner in both the low- and high-resolution experiments. Since all the models use the same atmospheric forcing dataset, the difference is solely due to the numerical and physical choices made by each modeling group.

As expected, there is a significant increase in the SSH variance as resolution is increased and the eddying solution SSH variance maps are much closer to the observations (top panel) than their low-resolution counterparts. In the high-resolution experiments ( $\sim 0.1^\circ$ ), the variability is however still lower than observed, especially in the gyre interiors and in the experiments that use relative winds (NCAR-POP, AWI-FESOM, and IAP-LICOM). This underestimation is thus partly a consequence of the well-known "eddy-killing" effect which results from taking into account the shear between atmospheric wind and ocean current when computing the wind stress and which can reduce the kinetic energy by as much as 30% (see discussion in section 2a). However, the use of absolute wind in FSU-HYCOM is not sufficient to raise the level of surface variability to that of the


Formatted: English (United States)

Deleted: 6


Deleted: 6



observations and Chassignet and Xu (2017) argues that one actually needs to significantly increase the resolution ( $\sim 0.01^\circ$ ) in order to resolve the submesoscale instabilities that can energize the mesoscale (Callies et al., 2016) and therefore enhance eddy kinetic energy comparable to the mesoscale AVISO observations. It is more physical to take into account the vertical shear between atmospheric winds and ocean currents when computing the wind stress (see Renault et al., 2020, for a review) as it allows for a better representation of western boundary current systems (Ma et al., 2016), especially the Agulhas Current retroflexion and associated eddies (Renault et al., 2017). In FSU-HYCOM which uses absolute wind, the Agulhas eddies are too regular and follow the same pathway. The use of relative winds not only increases the eddy decay, it also impacts the location of the Agulhas retroflexion and where the eddies are formed. The three simulations with relative winds have a better representation of the Agulhas eddies pathway.



Deleted: 19



Deleted:

Formatted: English (United States)

**Figure 17:** Top panel: Mean 1993-2018 SSH and variance AVISO. Middle panel: Difference between the mean modeled SSH and AVISO SSH. Bottom panel: Modeled variance derived from 5-day average outputs. The low-resolution IAP-LICOM SSH variance was not provided.

The eddy kinetic energy maps for the high-resolution simulations are displayed in Figures 18-21 for the surface, at 700 m, and at 1000 m. Observed surface EKE maps are either derived from altimetry (Figure 18a) or drifters (Figure 18b). Because of the irregular sampling, this necessarily implies some type of smoothing in space and time. In the case of the AVISO along-track altimetry observations, they are optimally interpolated on a  $0.25^\circ$  grid which filters scales less than 150 km (due to track separation and measurement noise and errors) and time scales less than 10 days (repeat cycle of the altimeters). There is a significant reduction in EKE when computing it using 5- or 10-day average outputs (as opposed to online at every time step) as shown in Figure 19 for FSU-HYCOM and NCAR-POP. In addition to a significant EKE reduction in the most active regions, the time averaging removes much of the small-scale variability associated with inertial motions and ageostrophic effects. There is not a big difference between the 5-day and the 10-day maps and for the rest of this section, we will use 5-day average model outputs when comparing the EKE to observations. Overall, as for the SSH variability, the EKE is larger in FSU-HYCOM because absolute winds are used to force the models. As already discussed, the use of relative winds does improve the pathway for the Agulhas current eddies, but from Figure 18, there is also a significant reduction of EKE in the ACC, and it also suppresses variability in many areas. This is especially true for the Indian Ocean, in the tropics, and west of the Hawaiian Islands.

Most model-observation comparisons usually focus on the surface fields because of very sparse spatiotemporal sampling at depth (see for example the EKE derived at 700 m from several years of SOFAR float measurements in the Gulf Stream region (Richardson, 1993) or the EKE distribution from ARGO floats at 1000 m (Ollitrault and Colin de Verdière, 2014)). Scott et al. (2010) evaluated the total kinetic energy simulated in eddying ocean models (on the order of  $0.1^\circ$  horizontal resolution) relative to moored current records. They found that the models agreed within a factor of 2 above 3500 m depth and within a factor of 3 below 3500 m depth. Thoppil et al. (2011) show that surface and the abyssal EKE increase with resolution and that better upper-ocean EKE allows strong eddy-driven abyssal circulation. This also means that, while the EKE at depth in the high-resolution experiments is a significant improvement over the quasi non-existent EKE of the low-resolution simulations, it is still significantly less than very limited observations (Richardson, 1993; Ollitrault and Colin de Verdière, 2014). Significantly higher resolution may be necessary in order to obtain a level of EKE close to the observations (see Chassignet and Xu (2017) for a discussion). When using relative winds at this resolution as in NCAR-POP, AWI-FESOM, or IAP-LICOM, there is very little EKE at depth when compared to FSU-HYCOM forced with absolute winds (Figures 20, and 21).

Deleted: 6

Deleted: 7

Deleted: 0

Deleted: 7

Deleted: 7

Deleted: 8

Deleted: 7

Deleted: s

Deleted: clearly

Deleted: .

Deleted: coarse

Deleted: 19

Deleted: 0







Figure 18: Surface EKE from a) AVISO and b) drifters observations. **Modeled** surface EKE calculated from 5-day average fields for the high-resolution c) FSU-HYCOM, d) NCAR-POP, e) AWI-FESOM and f) IAP-LICOM.



Deleted:

Formatted: English (United States)

Deleted: 7

Deleted: S






Figure 19: Comparison of surface EKE calculated from the total integrated kinetic energy, 5-day average fields and 10-day average fields in the high-resolution FSU-HYCOM and NCAR-POP.




Deleted:

Formatted: English (United States)


Deleted: 8



**Figure 20:** EKE at 700 m calculated from 5-day average fields in a) FSU-HYCOM, b) NCAR-POP, c) AWI-FESOM and d) IAP-LICOM.




**Figure 21:** EKE at 1000 m calculated from 5-day average fields in a) FSU-HYCOM, b) NCAR-POP, c) AWI-FESOM and d) IAP-LICOM.



Deleted:

Formatted: English (United States)

Deleted: 19



Deleted:


Formatted: English (United States)

Deleted: 0



## 4.2 Sea Surface Temperature (SST) and Sea Surface Salinity (SSS) deseasonalized variance

Surface temperature and salinity show sensitivities to resolution that are largely consistent with those described above for SSH in terms of both mean state and variability. All models show greatly enhanced SST variance (computed from deseasonalized monthly values spanning 1980-2018) at high resolution in regions of high mesoscale eddy activity, including: the Southern Ocean, the Agulhas Current retroflexion, the Brazil-Malvinas Confluence region, the Kuroshio and its extension across the North Pacific, and the Gulf Stream and its NAC extension (Figure 22). A global, satellite-based SST dataset spanning 1982-2018 provides some measure of ground truth for comparison, but the sampling limitations of microwave and infrared measurements of SST restrict the observed estimate to a ( $1/4^\circ$ ) grid that is 2-3 times coarser than the high-resolution model grids considered here (OISST.v2; Reynolds et al., 2007; Banzon et al., 2014).




**Figure 22: Modeled and observed (OISST) deseasonalized SST variance.**

As was noted for SSH, the high-resolution FSU-HYCOM simulation generally exhibits the highest SST variance in the midlatitude gyre interiors compared to other models, but it has lower SST variance in strongly eddying regions such as the Agulhas retroflexion, Kuroshio, and NAC. Overall, the high-resolution FSU-HYCOM shows the closest match to the observed benchmark. The improved structure of SST variance in the subpolar Atlantic in FSU-HYCOM is related to the improved NAC pathway in that model (discussed above). Compared to OISST.v2, the low-resolution models generally underestimate SST variance while the high-resolution models tend to overestimate the variance, particularly in the western boundary currents, Southern Ocean, and Agulhas retroflexion region. The resolution sensitivity is most dramatic in the NCAR-POP model, whose low-resolution version shows the weakest SST variance of any of the simulations considered here. The overestimation of SST

Formatted: English (United States)


Formatted: English (United States)

Deleted: 1



Deleted: 1

Formatted: Font: Not Bold, Not Italic, English (United States)



Deleted: 1

Deleted: 1

Formatted: English (United States)

variance has been noted before in eddy-resolving coupled simulations (e.g., Small et al. 2014), but it is not clear whether this discrepancy is attributable to deficiencies in the models or in the observation-based estimates which cannot fully resolve ocean mesoscale variability. The absence of a dynamic (and ocean mesoscale-resolving) atmosphere in these simulations may partially explain the overly high SST variance insofar as important eddy-damping processes (Ma et al., 2016) are missing in this experimental framework. It is interesting that all high resolution simulations generate more SST variance in the Kuroshio region than seen in OISST.v2, and yet they all show less variance than observed in the northeast Pacific and along the southeastern branch of the North Pacific gyre (near Hawaii). There is an indication of slightly enhanced ENSO-related tropical Pacific SST variance when going from low to high resolution, but there is not a systematic change in the spatial structure of this variance which appears to depend mainly on model formulation. The representation of high extratropical SST variance along the eastern boundaries of each of the major ocean basins shows robust improvement with resolution across the different models. The high resolution versions all show higher (and more realistic) variance in the following locations: the Benguela Current region in the eastern South Atlantic; the coastal region off the west coast of North America and Baja; the Leeuwin Current region off the west coast of Australia; and the Canary Current region off the west coast of North Africa. Improvements in simulated SST variability in these regions is more apparent than improvements in the mean state (Figure 10). The overall picture of greatly enhanced SST variance at eddying resolution is an important, but not unexpected, result that has significant implications for climate modeling because resolving air-sea interactions at the ocean mesoscale has been shown to result in qualitatively different coupled model behavior (e.g., Bryan et al., 2010; Ma et al., 2016, 2017; Laurindo et al., 2019).




Figure 23: Modeled and observed deseasonalized SSS variance.

While surface salinity generally exhibits globally enhanced variance at high resolution, particularly in the eddy-rich locations already highlighted above, the SSS variance plots are striking in that they show considerably more sensitivity to model formulation than to model horizontal resolution (Figure 23). The increase in SSS variance at both resolutions across model systems—from FSU-HYCOM, to NCAR-POP, to AWI-FESOM, and to IAP-LICOM—is likely related to the steady increase in surface salinity restoring timescale (Table 1). In the absence of reliable long-term global SSS observations, it is

Deleted: 9

Deleted:

Formatted: English (United States)

Deleted: 2

Deleted: 2

difficult to say which model formulation is best or whether high resolution improves the simulation of surface salinity variability.

4.3 Upper ocean heat and salinity content mean biases and deseasonalized variance

The temperature and salinity distributions in the upper 700 m are less constrained to observations than are the SST (which is restored to time-varying observed values through the sensible heat flux) and the SSS (which is restored to observed climatology through the artificial salinity restoring flux). Therefore, vertically-averaged upper ocean heat content (UOHC) and salt content (UOSC) biases tend to be larger than surface biases, particularly at low latitudes (cf. Figures 10, 24, and 25). The UK Met Office EN4 ocean analysis (Good et al., 2013) provides an observational benchmark for UOHC and UOSC mean and variance, but again considerable caution is warranted when interpreting model-observation discrepancies. The spatial resolution of this analysis is only 1°, and it relaxes to climatology in data-sparse regions, which are extensive in the pre-Argo era (prior to 2000).

Both FSU-HYCOM and NCAR-POP show significant and near ubiquitous mean bias reduction for UOHC and UOSC when resolution is increased (Figures 24 and 25). In contrast, AWI-FESOM and IAP-LICOM show mixed results, with bias reduction in some regions (e.g., the Southern Ocean) offset by bias increase elsewhere. Regions of high observed UOHC variance include all the regions of high observed SST variance mentioned above as well as some variance hotspots in the subtropical western Pacific, the Tasman Sea, the subtropical south Indian Ocean, and the western tropical Atlantic (Figure 24). The high-resolution models tend to show improved representation of these subsurface variance hotspots as well as enhanced variance in high EKE regions (western boundary currents, Agulhas, Southern Ocean, etc.). The high resolution IAP-LICOM model stands out for its globally high (perhaps unrealistically high) UOHC variance. This upper ocean variance bias in IAP-LICOM is even more evident in UOSC (Figure 25), although the limited sampling of 0-700m salinity in the EN4 product implies that the observed variance estimate is almost certainly a gross underestimate. There is a hint (especially apparent in NCAR-POP and IAP-LICOM) of higher resolution resulting in enhanced salinity variance along subtropical cell spiciness pathways from the eastern extratropical Pacific to the tropical western Pacific (e.g., Yeager and Large, 2004).

Formatted: English (United States)

Deleted: 9

Deleted: 3

Deleted: 4

Deleted: re


Deleted: 3

Deleted: 4


Deleted: 3

Deleted: 4

Formatted: English (United States)



**Figure 24:** Top panel: Mean 1980-2018 0-700 m temperature and variance from EN4. Middle panel: Difference between the mean modeled 1980-2018 0-700 m temperature and EN4. Bottom panel: Deseasonalized 0-700 m temperature modeled variance.



Deleted:  
Formatted: English (United States)  
Deleted: 3







Figure 25: Top panel: Mean 1980-2018 0-700 m salinity and variance from EN4. Middle panel: Difference between the mean modeled 1980-2018 0-700 m salinity and EN4. Bottom panel: Deseasonalized 0-700 m salinity modeled variance.

#### 4.4 Zonal AMOC mean and variance

Although the magnitude and the temporal evolution of the AMOC transport, as shown in Figures 15a-b, differ significantly between different models and different horizontal resolutions, its time mean spatial structure and variance are quite similar among all the simulations as shown in Figure 26 for the 1980 to 2018 period. All simulations show a positive upper cell and a negative lower cell. In the low-resolution simulations, the upper cell exhibits a maximum transport near 1000 m and 35°-40°N. The lower cell has its maximum transport near 3500-4000 m and is much weaker overall (~2 Sv), except for IAP-LICOM (~6 Sv in the subtropical North Atlantic). In high-resolution simulations, the upper cell extends deeper in all 4 models, which is more consistent with the observations, and the lower cell has a similar magnitude (2-4 Sv). This can be seen more clearly in Figure 27, which compares the model AMOC profiles for 2004-2018 to the RAPID results near 26.5°N. South of about 20°N,



Deleted:

Formatted: English (United States)

Deleted: 4

Formatted: English (United States)


Deleted: 4

Deleted: 5

Deleted: 6



the high-resolution AWI-FESOM exhibits a weak positive cell near the bottom that is unrealistic and is different from other models.



**Figure 26:** (Top panels) Modeled time mean (1980-2018) Atlantic meridional overturning streamfunction  $\psi_m(y, z)$  in the four models (low- and high-resolution). (Lower panels) Modeled standard deviation (1980-2018) of the Atlantic overturning streamfunction  $\psi_m(y, z)$  derived from monthly mean fields.

The AMOC standard deviation shows a similar pattern between different resolutions and different models (Figure 26). The variability is overall highest near the Equator (IAP-LICOM has the strongest variability near 10°N that differs from the other three). This is due to the large seasonal variability of the AMOC, associated with the migration of the Intertropical Convergence Zone (ITCZ) and the changes in wind patterns and Ekman transport (Xu et al., 2014). The standard deviations based on annual means does not show such a maximum near the Equator (see Figure 7 in Hirschi et al. 2019 for results based on different averaging scales). The variability is much weaker beyond the equatorial region. In the low-resolution simulations, the standard deviation is typically 2 Sv and has a slight elevation near 40°N. In the high-resolution simulations, the standard deviation is about 3 Sv and clearly shows a secondary maximum near 40°N. The difference between the low- and high-resolution simulations near the 40°N highlights the impact of the meandering and mesoscale eddy variability of the Gulf Stream to AMOC variability.

FSU-HYCOM Mean Atlantic Stream 1980-2018

NCAR-POP Mean

HIGH RES.

LOW RES.

Latitude

FSU-HYCOM Std Dev. Atlantic Stream, 1980-2018

NCAR-POP Std De

HIGH RES.

LOW RES.

Latitude

Deleted:

Formatted: English (United States)

Deleted: 5

Deleted: (

Formatted: English (United States)

Deleted: Modeled

Deleted: over 1980-2018

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Deleted: with high and

Deleted:

Deleted: s

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Deleted: over 1980-2018 based on

Deleted: results

Deleted: 5

155 Northward heat transport is usually correlated with higher AMOC transport in the Atlantic Ocean (e.g., Msadek et al.,  
 2013; Xu et al., 2016). In the high-resolution experiments, there is also an overall reduction in subtropical Atlantic upper ocean  
 cold bias (Figure 24) which can have an impact as shown by Msadek et al. (2013). Thus, it is not surprising that the high-  
 resolution simulations have a higher heat transport in the Atlantic than the low-resolution counterparts (Figure 28). The  
 maximum northward heat transport in 20-30°N is about 1.0 PW in the high-resolution models, 0.6-0.8 PW in low-resolution  
 160 models, and 1.25-1.3 PW in observations (Johns et al., 2011; Trenberth et al., 2019). In other basins such as the Indo-Pacific  
 and the Southern Oceans, higher resolution does not necessarily lead to higher heat transport (Figure 28a-c). But in general,  
 the meridional heat transport in high-resolution is closer to observations than the low-resolution simulations (Figure 28).





Figure 27. AMOC profile at 26°N.



Deleted: The AMOC dominates the meridional heat transport


Deleted:

Deleted: 7

Deleted: re

Deleted: 6

Deleted: 7




Deleted:

Formatted: English (United States)

Deleted: 6

Deleted:



Deleted:

Formatted: English (United States)

**Figure 28:** Modeled time mean meridional heat transport in PW over 1980-2018: a) global, b) Atlantic Ocean, and c) Indo-Pacific Ocean. Black lines are the latest observational results from Trenberth et al. (2019) with uncertainties.

#### 4.5 Northern and southern hemisphere sea ice mean and variance

Figures 29-32 shows 1980-2018 March and September modeled mean sea ice concentration (SIC) and sea ice thickness (SIT) together with the mean 1980-2018 SIC from passive microwave satellites and the mean 2003-2007 SIT from ICESat. As expected, given the relationship between sea ice extent and atmospheric temperature, the observed spatial pattern of winter SIC (averaged from 1980 to 2018) in the northern hemisphere (Figure 29) is well simulated in all the models at both low- and high-resolutions, though with greater consistency among the high-resolution models. The position of the marginal ice zone (MIZ) – or areas where the SIC is between 15% and 80% - varies between models and resolutions. With the SIC close to 100% in the interior Arctic in winter, the variance in modeled SIC (Figures S1-S8) is small in March. The general pattern of the March northern hemisphere SIT is consistent between models, save for the IAP-LICOM, with increasing SIT towards the Canadian Archipelago. At both resolutions, AWI-FESOM presents slightly thicker sea ice throughout most of the Arctic Basin. The high-resolution IAP-LICOM model is unrealistic, with very high sea ice thicknesses surrounding the pole. Figure 30 repeats Figure 29, but for September. The comparison is similar: there are large inter-model differences in the MIZ, but there is a generally consistent pattern of increasing ice concentrations and thickness from the pole toward the Canadian Archipelago and northern Greenland. The location of the summer MIZ in FSU-HYCOM and NCAR-POP is too far away from the coast when compared to the observations and AWI-FESOM. Overall, there is a broad inter-model disagreement in SIT. The SIT variance (Figures S1-S8) is higher in summer, especially in the shelf regions in experiments at both resolutions. While there is some sensitivity to model resolution for the simulated sea ice thickness, the difference due to changes in the resolution is smaller than the difference between the model results and the observations. Since the SIT observations shown in Figures 29-32 are 2003-2007 averages, one may question the value of comparing the 2003-2007 ICESat observations to 1980-2018 averaged modeled fields. For a more direct comparison, the SIT modeled fields averaged from 2003-2007 are shown in Figures S9 and S10 together with the ICESat observations when available, but they are almost identical to the 1980-2018 SIT fields shown in Figures 29-32.

Figures 31-32 reproduce Figures 29-30 for the southern hemisphere. The geographically simpler observed SIC in winter is well represented by the models, with the exception of FSU-HYCOM, which simulates large regions of low concentration ice within the Antarctic pack. There is a clear difference in the modeled MIZ across the simulations: for example, it is wider in the low-resolution IAP-LICOM experiments when compared to the high-resolution experiment, and vice-versa for NCAR-POP. In the AWI-FESOM simulation, the MIZ is much narrower. The MIZ width is an important observable for explaining southern hemisphere sea ice concentration and thickness, largely determined by the presence of ocean surface waves (Kohout et al., 2014; Horvat et al., 2020). As none of the models allow for the propagation of ocean waves into the ice, the feedback between wave propagation and sea ice cover could not take place. In summer, SIT and SIC are poorly represented in FSU-HYCOM, NCAR-POP, and IAP-LICOM, which, at high resolution, produce a too-wide MIZ (FSU-HYCOM, NCAR-POP).

**Deleted:** 7... Modeled time mean meridional hH...at  
tT...ansport in PW over 1980-2018: for...a) global ocean;... b)  
Atlantic Ocean;; ... [21]

**Formatted:** English (United States)

**Deleted:** -...ce mean and variance and variance ... [22]

**Formatted** ... [23]


**Formatted** ... [24]

1225

1230

or no MIZ (IAP-LICOM), and inaccurately low representations of SIT, with most regions thinner than 20 cm. In general, observations indicate the areas of the most compact sea ice (Weddell Sea) are not those of the thickest ice (western Antarctic sector) in summer. None of the simulations adequately represent the observed relatively thick and noncompact sea ice in the western Antarctic. The high summer sea ice concentration in the Weddell and Ross Seas is only reasonably simulated in AWI-FESOM, but with too thick ice. Although, there is a high sensitivity to changing resolution in the Southern Hemisphere, the low- and high-resolution model pairs tend to agree more than the ensemble of low- or high-resolution models, indicating that sea ice model formulation differences may exceed the effects of resolution in these patterns. The models also show some sensitivity to the model resolution for variance of the sea ice concentration and thickness in the southern hemisphere, especially in FSU-HYCOM. However, the difference between models is considerably larger. As IAP-LICOM significantly overestimates the sea-ice thickness in its high-resolution setup, its variance is also much higher than the other models (Figures S1-S8).

- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)



**Figure 29:** Northern hemisphere winter mean (March) sea ice concentration (SIC) and thickness (SIT). (Top panels) 1980-2018 SIC from passive microwave satellites and 2003-2007 SIT from ICESat. (Lower panels) 1980-2018 modeled SIC and SIT.

**Deleted:** The observed spatial pattern of winter sea-ice concentration (averaged from 1980 to 2018 for March) in the northern hemisphere is well simulated in all the models with both low- and high-resolutions, although there tends to be greater consistency among the high-resolution models. The simulated winter sea-ice spatial distribution is very similar between the models and setups with different resolutions. There is only very minor difference in the exact location of sea-ice outer edges in the sub-Arctic seas (Figures 28 and 29). Wintertime agreement is particularly good, with only low-resolution FSU-HYCOM deviating from the other models. In summer, all the models can reproduce the observed spatial pattern with high sea-ice concentration north of Greenland and low concentration towards the Arctic marginal seas. However, the difference between the models is more significant than in winter, especially in the shelf regions, which is true in both the low- and high-resolution experiments. The sea-ice concentration in summer also does not change much with changing resolution, except for IAP-LICOM, which has higher concentration in most parts of the Arctic Ocean in its high-resolution setup. In both seasons, the northern hemisphere sea-ice thickness is the largest north of the Greenland and decreases towards the Eurasian coast. This pattern is reproduced in the models for both res[25]

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** Centered

**Formatted:** English (United States)

**Formatted:** English (United States)


**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)



**Deleted:**

**Formatted:** English (United States)



**Deleted:** 8

**Deleted:**

**Deleted:** cover

**Formatted:** English (United States)





Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)


Formatted: Justified

Deleted: 29

Deleted: ice cover and thickness

Formatted: English (United States)


335 Figure 30: Northern hemisphere summer mean (September) sea ice concentration (SIC) and thickness (SIT). (Top panel) 1980-2018 SIC from passive microwave satellites (summer SIT not available from ICESat). (Lower panels) 1980-2018 modeled SIC and SIT.



**Figure 31:** Southern hemisphere winter mean (September) sea ice concentration (SIC) and thickness (SIT). (Top panels) 1980-2018 SIC from passive microwave satellites and 2003-2007 SIT from ICESat. (Lower panels) 1980-2018 modeled SIC and SIT.


Formatted: Centered

Ice Con. SSMI/NSIDC SEPTEMBER



Deleted:

Ice Thick. IceSat/NSIDC SEPTEMBER



... [28]

Deleted: summer

Deleted: ice cover and thickness

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Justified

Deleted: 0

Formatted: English (United States)

Formatted: English (United States)







Figure 32: Southern hemisphere summer mean (March) sea ice concentration (SIC) and thickness (SIT), (Top panels) 1980-2018 SIC from passive microwave satellites and 2003-2007 SIT from ICESat. (Lower panels) 1980-2018 modeled SIC and SIT.

Formatted: Centered



Deleted: 1

Deleted: ice cover and thickness

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Deleted: winter

Formatted: English (United States)

## 5 Summary and Discussion

The overall goal of this paper is to assess the robustness of climate-relevant improvements in ocean simulations (mean and variability) associated with moving from coarse ( $\sim 1^\circ$ ) to eddy-resolving ( $\sim 0.1^\circ$ ) horizontal resolutions. It also lays out a set of basic large-scale diagnostics for assessing the relative quality, variability, and sensitivity of high-resolution versus low-resolution ocean and sea-ice models. The emphasis is on the key metrics used in climate modelling – SST, OHC, sea level, salinity, sea ice extent and volume, and circulations that tend to have global impacts (MOC, ACC, ITF) on the modeled climate. Here these metrics are assessed in a suite of 4 pairs of low-resolution, high-resolution ocean and sea-ice models forced with the latest JRA55-do dataset (Tsujino et al., 2019). These results will provide a useful baseline for future process-focused analyses as well as ocean model development activities at diverse resolutions.

On the whole, the biases in the low-resolution variants are familiar, and are similar in these models to those found when assessing sensitivity to forcing products (Tsujino et al., 2020). Gross features of the bias patterns in low resolution models – position, strength, and variability of western boundary currents, equatorial currents, and ACC – are significantly improved in the high-resolution models. However, despite the fact that the high-resolution models “resolve” these features, the improvements in temperature or salinity are inconsistent among the different model families and some regions show increased bias over their low-resolution counterparts. SSH variability and near-surface EKE are significantly – even qualitatively – improved in all high-resolution models over their low-resolution counterparts, although all of these models still underpredict the observed SSH variability and EKE, particularly in the ocean interior, which indicates a need for further refinements in resolution (Chassignet and Xu, 2017), and improvements in less dissipative subgrid schemes for high-resolution models (Pearson et al., 2017) are needed. The results in coupled models in the HighResMIP ensemble (Haarsma et al., 2016) show similar improvements in SSH and SST variability and EKE. Considerable differences in the high-resolution models used here were associated with the use of relative winds versus absolute winds. Renault et al. (2020) show that using relative winds contributes a feedback that tends to reduce EKE, and the models here are consistent with a 30% or so reduction of EKE when relative winds are used.

One interesting aspect of the high-resolution models versus the low-resolution models was that the interannual variability in the ACC and ITF transport and AMOC was more consistent among the high-resolution models than among the low-resolution models. The ITF transport was especially in agreement among the high-resolution models, indicating that better representation of the passageways through the Indonesian archipelago is critical. Consistency in all of these transports potentially indicates that higher-resolution models are needed to represent process variability, which may explain some of the past difficulties in comparing the magnitude of these phenomena across coarse-resolution models. However, the mean ACC transport and MOC strength were not in greater agreement among the high-resolution than the low-resolution models, which means that more work remains in evaluating sensitivity to numerics and subgrid-scale schemes for high-resolution models. Furthermore, Danabasoglu et al. (2016) note that low-resolution models come into greater agreement in AMOC variability after more cycles of the CORE forcing – this comparison was limited by the cost of the high-resolution models to only a single

Deleted: ¶

Formatted: English (United States)

Formatted: English (United States)

Deleted: remains

cycling of the forcing. It is unclear if the high-resolution models are in greater agreement only in the first cycle or generally. The short duration of a single forcing cycle limits the comparison of the decadal changes that are emphasized in Danabasoglu et al. (2016), so the improved agreement discussed here among high-resolution models is year-by-year rather than decade-by-decade. Nonetheless, the high-resolution models had systematically stronger and more variable AMOC, which was in better agreement with observations, both in maximum overturning and profile, than the low-resolution models.

From a climate modeling perspective, ocean heat content, sea level, and sea ice stability are key metrics. There is some indication that the high-resolution models may warm more quickly below 700 m than the low-resolution models, indicating errors in parameterizations of vertical eddy heat transport. Griffies et al. (2016) found similar sensitivity to resolution in a model hierarchy which they explained as resolved eddies versus parameterized eddies affecting vertical heat transport. However, warming between 0-700 m and global mean sea level rise were not systematically different across model families between the low-resolution and high-resolution models. Additional insights could be gained by computing vertical heat and salt budgets as in Griffies et al. (2015) and Von Storch et al. (2016). Regional sea level rise, however, was significantly more variable in the high-resolution models where mesoscale eddies and variability of western boundary and equatorial currents make an impact on regional sea level rise. Thus, it is critical to use high-resolution models to assess the ocean dynamic sea level (Gregory et al., 2019) contribution to regional sea level rise. Generally, there is a highly variable representation of sea ice with the exception of Arctic sea ice concentration – with high inter-model differences in sea ice thickness and volume, and generally inconsistent representation of sea ice in the Southern Hemisphere and sea ice thickness in the Arctic. Inter-model differences are larger than improvements with changing resolution – pointing to the known role of atmospheric forcing in setting modeled sea ice states as well as to the poor representation of sea ice physics and associated parameterizations. Sea ice concentration bias reduction is achievable, for example, through the inclusion of lateral melting and feedbacks between ice fracture and sea ice melting (Horvat and Tziperman, 2015; Roach et al., 2018; Bateson et al., 2020), but the wide spread in sea ice thickness and poor representation of Antarctic sea ice is known in CMIP-class models (Roach et al, 2020; Notz et al, 2020, Shu et al., 2020) and indicates that a fundamental overhaul of sea ice modelling may be needed.

Low-resolution versus high-resolution comparisons are often motivated by identifying persistent patterns across multiple models of bias or improvements with resolution. However, as apparent in many metrics, there is no overall consistency among either the low-resolution or high-resolution ensemble from which to draw simplistic conclusions. This paper therefore does not dwell on the detailed differences among these models' numerics and parameterizations, but is instead intended to serve as a benchmark for future studies comparing and improving different schemes in any of these models or similar ones. The numerics and parameterizations of these models are in continual development in both low-resolution and high-resolution versions. Ideally, however, one should strive to minimize the number of changes as the resolution is increased to ensure traceability. Other modeling centers have expressed interest in participating in this protocol in the future, and this paper provides a basis for comparison.

Finally, the models used in the present study are not coupled to active atmospheres, biospheres, land models, nor land ice models. Thus, many important feedbacks and aspects of climate modeling are not addressed here. However, these same

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

**Deleted:** In terms of sea ice, high-resolution models were better than or equal to low-resolution models in reproducing observations of extent and volume (except for the IAP-LICOM model), and the high-resolution models were more consistent in reproducing observed trends and regional patterns in sea ice.¶

**Deleted:** basis

**Deleted:** The set of tests performed here are a benchmark against which development changes can be assessed.

**Deleted:** T

1450 feedbacks complicate diagnosis of the source of model biases and resolution sensitivity, which is where this study intends to contribute.

#### Code and data availability

The forcing dataset for OMIP-2 is available through input4MIPs (<https://esgf-node.llnl.gov/search/input4mips/>) – see Table 1 in the supplement of a list of the files. An archive of the model outputs and the scripts used to process data and generate figures are available at <https://doi.org/10.5281/zenodo.3685918>. For the observed sea ice concentration, the data are available at <https://nsidc.org/data/NSIDC-0192/versions/2>. For the observed sea ice thickness, the data are available at <https://nsidc.org/data/NSIDC-0393/versions/1> and at <https://earth.gsfc.nasa.gov/index.php/cryo/data/antarctic-sea-ice-thickness> for the Arctic and Antarctic, respectively.

Deleted: are available at .

Formatted: English (United States)

#### Author contribution

1460 EC, SY, and BFK proposed and led this evaluation study. AB processed the model outputs and produced the figures. The following authors are responsible for the individual models, simulations, and diagnostics: AB, XX, and EPC for FSU-HYCOM; SY, FC, WK, and GD for NCAR-POP; QW, SD, NK, and DS for AWI-FESOM; HL, YL, and PL for IAP-LICOM3. All authors contributed to the writing and editing processes.

#### Competing interests

1465 The authors declare that they have no conflict of interest.

#### Acknowledgements

The authors would like to thank the two anonymous reviewers for their constructive comments. The FSU contribution was supported by the National Oceanic and Atmospheric Administration (NOAA) Earth System Prediction Capability Project (Award NA15OAR4320064), NOAA Climate Program Office (CPO) MAPP Program (Award 5NA15OAR4310088), and NSF Physical Oceanography Program (Award 1537136). The NCAR contribution was supported by the NOAA CPO Climate Variability and Predictability (CVP) Program. NCAR is a major facility sponsored by the US National Science Foundation (NSF) under Cooperative Agreement No. 1852977. The AWI contribution was supported by the projects S1 (Diagnosis and Metrics in Climate Models) and S2 (Improved parameterizations and numerics in climate models) of the Collaborative Research Centre TRR 181 “Energy Transfer in Atmosphere and Ocean” funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project no. 274762653, Helmholtz Climate Initiative REKLIM (Regional Climate Change) and European Union’s Horizon 2020 Research & Innovation programme through grant agreement No. 727862 APPLICATE. The IAP contribution was supported by the National Natural Science Foundation of China (Grants No. 41931183 and 41976026). D. Sein was supported by the state assignment of FASO Russia (theme 0149-2019-0015). BFK’s contribution was supported by NSF 1350795, ONR N00014-17-1-2393, and NOAA NA19OAR4310366.

Formatted: English (United States)

Formatted: English (United States)

## References

Ajayi, A., Le Sommer, J., Chassignet, E., Molines, J.-M., Xu, X., Albert, A. and Cosme, E.: Spatial and temporal variability of the North Atlantic eddy field from two kilometeric-resolution ocean models, *J. Geophys. Res. Oceans*, 125, e2019JC015827, doi:10.1029/2019JC015827, 2020.

Bamber, J. L., Tedstone, A. J., King, M. D., Howat, I. M., Enderlin, E. M., van den Broeke, M. R. and Noel, B.: Land Ice Freshwater Budget of the Arctic and North Atlantic Oceans: 1. Data, Methods, and Results, *J. Geophys. Res. Ocean.*, 123(3), 1827–1837, doi:10.1002/2017JC013605, 2018.

Bamber, J., van den Broeke, M., Ettema, J., Lenaerts, J. and Rignot, E.: Recent large increases in freshwater fluxes from Greenland into the North Atlantic, *Geophys. Res. Lett.*, 39(19), doi:10.1029/2012GL052552, 2012.

Banzon, V. F., Reynolds, R. W., Stokes, D. and Xue, Y.: A 1/4°-Spatial-Resolution Daily Sea Surface Temperature Climatology Based on a Blended Satellite and in situ Analysis, *J. Clim.*, 27(21), 8221–8228, doi:10.1175/JCLI-D-14-00293.1, 2014.

Bao, Q., Lin, P., Zhou, T., Liu, Y., Yu, Y., Wu, G., He, B., He, J., Li, L., Li, J., Li, Y., Liu, H., Qiao, F., Song, Z., Wang, B., Wang, J., Wang, P., Wang, X., Wang, Z., Wu, B., Wu, T., Xu, Y., Yu, H., Zhao, W., Zheng, W. and Zhou, L.: The Flexible Global Ocean-Atmosphere-Land system model, Spectral Version 2: FGOALS-s2, *Adv. Atmos. Sci.*, 30(3), 561–576, doi:10.1007/s00376-012-2113-9, 2013.

Baringer, M. O. and Garzoli, S. L.: Meridional heat transport determined with expendable bathythermographs-Part I: Error estimates from model and hydrographic data, *Deep. Res. Part I Oceanogr. Res. Pap.*, 54(8), 1390–1401, doi:10.1016/j.dsr.2007.03.011, 2007.

Bleck, R.: An oceanic general circulation model framed in hybrid isopycnic-Cartesian coordinates, *Ocean Model.*, 4(1), 55–88, doi:10.1016/S1463-5003(01)00012-9, 2002.

Böning, C. W., Scheinert, M., Dengg, J., Biastoch, A. and Funk, A.: Decadal variability of subpolar gyre transport and its reverberation in the North Atlantic overturning, *Geophys. Res. Lett.*, 33(21), L21S01, doi:10.1029/2006GL026906, 2006.

Bryan, F. O., Tomas, R., Dennis, J. M., Chelton, D. B., Loeb, N. G. and McClean, J. L.: Frontal Scale Air–Sea Interaction in High-Resolution Coupled Climate Models, *J. Clim.*, 23(23), 6277–6291, doi:10.1175/2010JCLI3665.1, 2010.

Callies, J., Flierl, G., Ferrari, R. and Fox-Kemper, B.: The role of mixed-layer instabilities in submesoscale turbulence, *J. Fluid Mech.*, 788, 5–41, doi:10.1017/jfm.2015.700, 2015.

Canuto, V. M., Howard, A., Cheng, Y. and Dubovikov, M. S.: Ocean turbulence. Part I: One-point closure model—momentum and heat vertical diffusivities, *J. Phys. Oceanogr.*, 31(6), 1413–1426, doi:10.1175/1520-0485(2001)031<1413:OTPIOP>2.0.CO;2, 2001.

Canuto, V. M., Howard, A., Cheng, Y. and Dubovikov, M. S.: Ocean turbulence. Part II: Vertical diffusivities of momentum, heat, salt, mass, and passive scalars, *J. Phys. Oceanogr.*, 32(1), 240–264, doi:10.1175/1520-0485(2002)032<0240:OTPIVD>2.0.CO;2, 2002.

Chassignet, E. P., Smith, L. T., Halliwell, G. R. and Bleck, R.: North Atlantic simulations with the Hybrid Coordinate Ocean Model (HYCOM): Impact of the vertical coordinate choice, reference pressure, and thermobaricity, *J. Phys. Oceanogr.*, 33(12), 2504–2526, doi:10.1175/1520-0485(2003)033<2504:NASWTH>2.0.CO;2, 2003.

Chassignet, E. P. and Marshall, D. P.: Gulf Stream separation in numerical ocean models, in *Geophysical Monograph Series*, 177, 39–61, 2008.

Chassignet, E. P. and Xu, X.: Impact of horizontal resolution (1/12° to 1/50°) on Gulf Stream separation, penetration, and variability, *J. Phys. Oceanogr.*, 47(8), 1999–2021, doi:10.1175/JPO-D-17-0031.1, 2017.

Chassignet, E. P., Yeager, S. G., Fox-Kemper, B., Bozec, A., Castruccio, F., Danabasoglu, G., Kim, W. M., Koldunov, N., Li, Y., Lin, P., Liu, H., Sein, D., Sidorenko, D., Wang, Q. and Xu, X.: Impact of horizontal resolution on the energetics of global ocean-sea-ice model simulations, *CLIVAR Variations/Exchanges*, 18(1), 23–30, doi:10.5065/g8w0-fy32, 2020.

Chidichimo, M. P., Donohue, K. A., Watts, D. R. and Tracey, K. L.: Baroclinic transport time series of the Antarctic Circumpolar Current measured in Drake Passage, *J. Phys. Oceanogr.*, 44(7), 1829–1853, doi:10.1175/JPO-D-13-071.1, 2014.

Craig, A. P., Vertenstein, M. and Jacob, R.: A new flexible coupler for earth system modeling developed for CCSM4 and CESM1, *Int. J. High Perform. Comput. Appl.*, 26(1), 31–42, doi:10.1177/1094342011428141, 2012.

Formatted: English (United States)

Formatted: Default Paragraph Font, English (United States)

Formatted: English (United States)

Moved down [12]: Canuto, V. M., Howard, A., Cheng, Y. and Dubovikov, M. S.: Ocean turbulence. Part II: Vertical diffusivities of momentum, heat, salt, mass, and passive scalars, *J. Phys. Oceanogr.*, 32(1), 240–264, doi:10.1175/1520-0485(2002)032<0240:OTPIVD>2.0.CO;2, 2002.

Moved (insertion) [12]

Moved down [1]: Chassignet, E. P. and Marshall, D. P.: Gulf stream separation in numerical ocean models, in *Geophysical Monograph Series*, vol. 177, pp. 39–61., 2008.

Formatted: English (United States)

Moved (insertion) [1]

Deleted: s

Deleted: vol.

Deleted: pp.

- Cunningham, S. A.: Transport and variability of the Antarctic Circumpolar Current in Drake Passage, *J. Geophys. Res.*, 108(C5), 8084, doi:10.1029/2001JC001147, 2003.
- Danabasoglu, G., Bates, S. C., Briegleb, B. P., Jayne, S. R., Jochum, M., Large, W. G., Peacock, S. and Yeager, S. G.: The CCSM4 Ocean Component, *J. Clim.*, 25(5), 1361–1389, doi:10.1175/JCLI-D-11-00091.1, 2012.
- 1545 Danabasoglu, G., Ferrari, R. and McWilliams, J. C.: Sensitivity of an ocean general circulation model to a parameterization of near-surface eddy fluxes, *J. Clim.*, 21(6), 1192–1208, doi:10.1175/2007JCLI1508.1, 2008.
- Danabasoglu, G., Large, W. G. and Briegleb, B. P.: Climate impacts of parameterized Nordic Sea overflows, *J. Geophys. Res.*, 115(C11), C11005, doi:10.1029/2010JC006243, 2010.
- Danabasoglu, G., Large, W. G., Tribbia, J. J., Gent, P. R., Briegleb, B. P. and McWilliams, J. C.: Diurnal coupling in the tropical oceans of CCSM3, *J. Clim.*, 19(11), 2347–2365, doi:10.1175/JCLI3739.1, 2006.
- 1550 Danabasoglu, G. and Marshall, J.: Effects of vertical variations of thickness diffusivity in an ocean general circulation model, *Ocean Model.*, 18(2), 122–141, doi:10.1016/j.ocemod.2007.03.006, 2007.
- Danabasoglu, G., Yeager, S. G., Bailey, D., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Böning, C., Bozec, A., Canuto, V. M., Cassou, C., Chassignet, E., Coward, A. C., Danilov, S., Diansky, N., Drange, H., Farneti, R., Fernandez, E., Fogli, P. G., Forget, G., Fujii, Y., Griffies, S. M., Gusev, A., Heimbach, P., Howard, A., Jung, T., Kelley, M., Large, W. G., Leboissetier, A., Lu, J., Madec, G., Marsland, S. J., Masina, S., Navarra, A., George Nurser, A. J., Pirani, A., y Mélia, D. S., Samuels, B. L., Scheinert, M., Sidorenko, D., Treguier, A. M., Tsujino, H., Uotila, P., Valcke, S., Voldoire, A. and Wang, Q.: North Atlantic simulations in Coordinated Ocean-ice Reference Experiments phase II (CORE-II). Part I: Mean states, *Ocean Model.*, 73, 76–107, doi:10.1016/j.ocemod.2013.10.005, 2014.
- 1560 Danabasoglu, G., Yeager, S. G., Kim, W. M., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Bleck, R., Böning, C., Bozec, A., Canuto, V. M., Cassou, C., Chassignet, E., Coward, A. C., Danilov, S., Diansky, N., Drange, H., Farneti, R., Fernandez, E., Fogli, P. G., Forget, G., Fujii, Y., Griffies, S. M., Gusev, A., Heimbach, P., Howard, A., Ilicak, M., Jung, T., Karspeck, A. R., Kelley, M., Large, W. G., Leboissetier, A., Lu, J., Madec, G., Marsland, S. J., Masina, S., Navarra, A., Nurser, A. J. G., Pirani, A., Romanou, A., David, S. y M., Samuels, B. L., Scheinert, M., Sidorenko, D., Sun, S., Treguier, A. M., Tsujino, H., Uotila, P., Valcke, S., Voldoire, A., Wang, Q. and Yashayaev, I.: North Atlantic simulations in Coordinated Ocean-ice Reference Experiments phase II (CORE-II). Part II: Inter-annual to decadal variability, *Ocean Model.*, 97, 65–90, doi:10.1016/j.ocemod.2015.11.007, 2016.
- 1565 Danilov, S., Wang, Q., Timmermann, R., Iakovlev, N., Sidorenko, D., Kimritz, M., Jung, T. and Schröter, J.: Finite-Element Sea Ice Model (FESIM), version 2, *Geosci. Model Dev.*, 8(6), 1747–1761, doi:10.5194/gmd-8-1747-2015, 2015.
- 1570 Danilov, S., Kivman, G. and Schröter, J.: A finite-element ocean model: Principles and evaluation, *Ocean Model.*, 6(2), 125–150, doi:10.1016/S1463-5003(02)00063-X, 2004.
- Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., Van Den Broeke, M. R. and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, 502(7469), 89–92, doi:10.1038/nature12567, 2013.
- 1575 Dong, S., Baringer, M. O., Goni, G. J., Meinen, C. S. and Garzoli, S. L.: Seasonal variations in the South Atlantic Meridional Overturning Circulation from observations and numerical models, *Geophys. Res. Lett.*, 41(13), 4611–4618, doi:10.1002/2014GL060428, 2014.
- Dong, S., Garzoli, S., Baringer, M., Meinen, C. and Goni, G.: Interannual variations in the Atlantic meridional overturning circulation and its relationship with the net northward heat transport in the South Atlantic, *Geophys. Res. Lett.*, 36(20), L20606, doi:10.1029/2009GL039356, 2009.
- 1580 Dong, S., Goni, G. and Bringas, F.: Temporal variability of the South Atlantic Meridional Overturning Circulation between 20°S and 35°S, *Geophys. Res. Lett.*, 42(18), 7655–7662, doi:10.1002/2015GL065603, 2015.
- Donohue, K. A., Tracey, K. L., Watts, D. R., Chidichimo, M. P. and Chereskin, T. K.: Mean Antarctic Circumpolar Current transport measured in Drake Passage, *Geophys. Res. Lett.*, 43(22), 11,760–11,767, doi:10.1002/2016GL070319, 2016.
- 1585 Downes, S. M., Farneti, R., Uotila, P., Griffies, S. M., Marsland, S. J., Bailey, D., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Böning, C., Bozec, A., Canuto, V. M., Chassignet, E., Danabasoglu, G., Danilov, S., Diansky, N., Drange, H., Fogli, P. G., Gusev, A., Howard, A., Ilicak, M., Jung, T., Kelley, M., Large, W. G., Leboissetier, A., Long, M., Lu, J., Masina, S., Mishra, A., Navarra, A., George Nurser, A. J., Patara, L., Samuels, B. L., Sidorenko, D., Spence, P., Tsujino, H., Wang, Q. and Yeager, S. G.: An assessment of Southern Ocean water masses and sea ice during 1988–2007 in a suite of interannual CORE-II simulations, *Ocean Model.*, 94, 67–94, doi:10.1016/j.ocemod.2015.07.022, 2015.
- 1590

- Farneti, R., Downes, S. M., Griffies, S. M., Marsland, S. J., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Böning, C., Bozec, A., Canuto, V. M., Chassignet, E., Danabasoglu, G., Danilov, S., Diansky, N., Drange, H., Fogli, P. G., Gusev, A., Hallberg, R. W., Howard, A., Ilicak, M., Jung, T., Kelley, M., Large, W. G., Leboissetier, A., Long, M., Lu, J., Masina, S., Mishra, A., Navarra, A., George Nurser, A. J., Patara, L., Samuels, B. L., Sidorenko, D., Tsujino, H., Uotila, P., Wang, Q. and Yeager, S. G.: An assessment of Antarctic Circumpolar Current and Southern Ocean meridional overturning circulation during 1958–2007 in a suite of interannual CORE-II simulations, *Ocean Model.*, 93, 84–120, doi:10.1016/j.ocemod.2015.07.009, 2015.
- Ferrari, R., McWilliams, J. C., Canuto, V. M. and Dubovikov, M.: Parameterization of eddy fluxes near oceanic boundaries, *J. Clim.*, 21(12), 2770–2789, doi:10.1175/2007JCLI1510.1, 2008.
- Ferreira, D., Marshall, J. and Heimbach, P.: Estimating Eddy Stresses by Fitting Dynamics to Observations Using a Residual-Mean Ocean Circulation Model and Its Adjoint, *J. Phys. Oceanogr.*, 35(10), 1891–1910, doi:10.1175/JPO2785.1, 2005.
- Fetterer, F., Knowles, K., Meier, W., Savoie, M. and Windnagel, A. K.: Sea Ice Index, Version 3, Boulder, Colorado USA., 2017.
- Firing, Y. L., Chereskin, T. K. and Mazloff, M. R.: Vertical structure and transport of the Antarctic Circumpolar Current in Drake Passage from direct velocity observations, *J. Geophys. Res.*, 116(C8), C08015, doi:10.1029/2011JC006999, 2011.
- Fox-Kemper, B., Danabasoglu, G., Ferrari, R., Griffies, S. M., Hallberg, R. W., Holland, M. M., Maltrud, M. E., Peacock, S. and Samuels, B. L.: Parameterization of mixed layer eddies. III: Implementation and impact in global ocean climate simulations, *Ocean Model.*, 39(1–2), 61–78, doi:10.1016/j.ocemod.2010.09.002, 2011.
- Fox-Kemper, B., Adcroft, A., Böning, C. W., Chassignet, E. P., Curchitser, E., Danabasoglu, G., Eden, C., England, M. H., Gerdes, R., Greatbatch, R. J., Griffies, S. M., Hallberg, R. W., Hanert, E., Heimbach, P., Hewitt, H. T., Hill, C. N., Komuro, Y., Legg, S., Sommer, J. Le, Masina, S., Marsland, S. J., Penny, S. G., Qiao, F., Ringler, T. D., Treguier, A. M., Tsujino, H., Uotila, P. and Yeager, S. G.: Challenges and prospects in ocean circulation models, *Front. Mar. Sci.*, 6(FEB), doi:10.3389/fmars.2019.00065, 2019.
- Fox-Kemper, B., Ferrari, R. and Hallberg, R.: Parameterization of mixed layer eddies. Part I: Theory and diagnosis, *J. Phys. Oceanogr.*, 38(6), 1145–1165, doi:10.1175/2007JPO3792.1, 2008.
- Gent, P. R. and Danabasoglu, G.: Response to increasing southern hemisphere winds in CCSM4, *J. Clim.*, 24(19), 4992–4998, doi:10.1175/JCLI-D-10-05011.1, 2011.
- Gent, P. R. and McWilliams, J. C.: Isopycnal mixing in ocean circulation models, *J. Phys. Oceanogr.*, 20(1), 150–155, doi:10.1175/1520-0485(1990)020<0150:imiocm>2.0.co;2, 1990.
- Godfrey, J. S.: A Sverdrup model of the depth-integrated flow for the world ocean allowing for island circulations, *Geophys. Astrophys. Fluid Dyn.*, 45(1–2), 89–112, doi:10.1080/03091928908208894, 1989.
- Goes, M., Goni, G. and Dong, S.: An optimal XBT-based monitoring system for the South Atlantic meridional overturning circulation at 34°S, *J. Geophys. Res. Ocean.*, 120(1), 161–181, doi:10.1002/2014JC010202, 2015.
- Good, S. A., Martin, M. J. and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *J. Geophys. Res. Ocean.*, 118(12), 6704–6716, doi:10.1002/2013JC009067, 2013.
- Gordon, A. L., Sprintall, J., Van Aken, H. M., Susanto, R. D., Wijffels, S., Molcard, R., Ffield, A., Pranowo, W. and Wirasantosa, S.: The Indonesian throughflow during 2004–2006 as observed by the INSTANT program, *Dyn. Atmos. Ocean.*, 50(2), 115–128, doi:10.1016/j.dynatmoce.2009.12.002, 2010.
- Gordon, A. L.: Oceanography of the Indonesian seas and their throughflow, *Oceanography*, 18(SPL.ISS. 4), 15–27, doi:10.5670/oceanog.2005.01, 2005.
- Gordon, A. L.: Inter-ocean exchange of thermocline water, *J. Geophys. Res.*, 91(C4), 5037, doi:10.1029/jc091ic04p05037, 1986.
- Gregory, J. M., Griffies, S. M., Hughes, C. W., Lowe, J. A., Church, J. A., Fukimori, I., Gomez, N., Kopp, R. E., Landerer, F., Cozannet, G. Le, Ponte, R. M., Stammer, D., Tamsiea, M. E. and van de Wal, R. S. W.: Concepts and Terminology for Sea Level: Mean, Variability and Change, Both Local and Global, *Surv. Geophys.*, 40(6), 1251–1289, doi:10.1007/s10712-019-09525-z, 2019.
- Griffies, S. M.: The Gent–McWilliams Skew Flux, *J. Phys. Oceanogr.*, 28(5), 831–841, doi:10.1175/1520-0485(1998)028<0831:TGMSF>2.0.CO;2, 1998.
- Griffies, S. M., Biastoch, A., Böning, C., Bryan, F., Danabasoglu, G., Chassignet, E. P., England, M. H., Gerdes, R., Haak, H., Hallberg, R. W., Hazeleger, W., Jungclauss, J., Large, W. G., Madec, G., Pirani, A., Samuels, B. L., Scheinert, M., Gupta,



- A. Sen, Severijns, C. A., Simmons, H. L., Treguier, A. M., Winton, M., Yeager, S. and Yin, J.: Coordinated Ocean-ice Reference Experiments (COREs), *Ocean Model.*, 26(1–2), 1–46, doi:10.1016/j.ocemod.2008.08.007, 2009.
- Griffies, S. M., Yin, J., Durack, P. J., Goddard, P., Bates, S. C., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Böning, C. W., Bozec, A., Chassignet, E., Danabasoglu, G., Danilov, S., Domingues, C. M., Drange, H., Farneti, R., Fernandez, E., Greatbatch, R. J., Holland, D. M., Ilicak, M., Large, W. G., Lorbacher, K., Lu, J., Marsland, S. J., Mishra, A., George Nurser, A. J., Salas y Mélia, D., Palter, J. B., Samuels, B. L., Schröter, J., Schwarzkopf, F. U., Sidorenko, D., Treguier, A. M., Tseng, Y. heng, Tsujino, H., Uotila, P., Valcke, S., Voldoire, A., Wang, Q., Winton, M. and Zhang, X.: An assessment of global and regional sea level for years 1993–2007 in a suite of interannual **CORE-II** simulations, *Ocean Model.*, 78, 35–89, doi:10.1016/j.ocemod.2014.03.004, 2014.
- Griffies, S. M., Winton, M., Anderson, W. G., Benson, R., Delworth, T. L., Dufour, C. O., Dunne, J. P., Goddard, P., Morrison, A. K., Rosati, A., Wittenberg, A. T., Yin, J. J. and Zhang, R.: **Impacts on Ocean Heat from Transient Mesoscale Eddies in a Hierarchy of Climate Models**, *J. Clim.*, 28, 952–977, 2015.
- Griffies, S. M., Danabasoglu, G., Durack, P. J., Adcroft, A. J., Balaji, V., Böning, C. W., Chassignet, E. P., Curchitser, E., Deshayes, J., Drange, H., Fox-Kemper, B., Gleckler, P. J., Gregory, J. M., Haak, H., Hallberg, R. W., Heimbach, P., Hewitt, H. T., Holland, D. M., Ilyina, T., Jungclaus, J. H., Komuro, Y., Krasting, J. P., Large, W. G., Marsland, S. J., Masina, S., McDougall, T. J., George Nurser, A. J., Orr, J. C., Pirani, A., Qiao, F., Stouffer, R. J., Taylor, K. E., Treguier, A. M., Tsujino, H., Uotila, P., Valdivieso, M., Wang, Q., Winton, M. and Yeager, S. G.: OMIP contribution to CMIP6: Experimental and diagnostic protocol for the physical component of the Ocean Model Intercomparison Project, *Geosci. Model Dev.*, 9(9), 3231–3296, doi:10.5194/gmd-9-3231-2016, 2016.
- Haarsma, R. J., Roberts, M. J., Vidale, P. L., Senior, C. A., Bellucci, A., Bao, Q., Chang, P., Corti, S., Fučkar, N. S., Guemas, V., von Hardenberg, J., Hazeleger, W., Kodama, C., Koenigk, T., Leung, L. R., Lu, J., Luo, J.-J., Mao, J., Mizielinski, M. S., Mizuta, R., Nobre, P., Satoh, M., Scoccimarro, E., Semmler, T., Small, J. and von Storch, J.-S.: High Resolution Model Intercomparison Project (HighResMIP v1.0) for CMIP6, *Geosci. Model Dev.*, 9(11), 4185–4208, doi:10.5194/gmd-9-4185-2016, 2016.
- Häkkinen, S. and Rhines, P. B.: Decline of Subpolar North Atlantic Circulation during the 1990s, *Science* (80-. ), 304(5670), 555–559, doi:10.1126/science.1094917, 2004.
- Hallberg, R.: Using a resolution function to regulate parameterizations of oceanic mesoscale eddy effects, *Ocean Model.*, 72, 92–103, doi:10.1016/j.ocemod.2013.08.007, 2013.
- Hallberg, R. and Gnanadesikan, A.: The role of Eddies in determining the structure and response of the wind-driven Southern hemisphere overturning: Results from the Modeling Eddies in the Southern Ocean (MESO) project, *J. Phys. Oceanogr.*, 36(12), 2232–2252, doi:10.1175/JPO2980.1, 2006.
- Halliwel, G. R.: Evaluation of vertical coordinate and vertical mixing algorithms in the HYbrid-Coordinate Ocean Model (HYCOM), *Ocean Model.*, 7(3–4), 285–322, doi:10.1016/j.ocemod.2003.10.002, 2004.
- Hirschi, J. J.-M., Barnier, B., Böning, C., Biastoch, A., Blaker, A. T., Coward, A., Danilov, S., Drijfhout, S., Getzlaff, K., Griffies, S. M., Hasumi, H., Hewitt, H., Iovino, D., Kawasaki, T., Kiss, A. E., Koldunov, N., Marzocchi, A., Moat, B., Molines, J.-M., Myers, P. G., Penduff, T., Roberts, M., Treguier, A.-M., Sein, D. V., Sidorenko, D., Small, J., Spence, P., Thompson, L., Weijer, W. and Xu, X.: The Atlantic meridional overturning circulation in high resolution models (in review), *J. Geophys. Res. Ocean.*, 2019.
- Hunke, E. C., Lipscomb, W. H., Turner, A. K., Jeffery, N. and Elliott, S.: CICE: the Los Alamos Sea Ice Model documentation and software user's manual, Version 5.1, LA-CC-06-012, T-3 Fluid Dyn. Group, Los Alamos Natl. Lab., 116, 2015.
- Horvat, C. and Tziperman, E.: **A prognostic model of the sea-ice floe size and thickness distribution**, *Cryosphere*, 6, 2119–2134, doi: 10.5194/tc-9-2119-2015, 2015.
- Horvat, C., Blanchard-Wrigglesworth, E. and Petty, A.: **Observing waves in sea ice with ICESat-2**, *Geophys. Res. Lett.*, doi: 10.1029/2020GL087629, 2020.
- Hunke, E. C. and Dukowicz, J. K.: An elastic-viscous-plastic model for sea ice dynamics, *J. Phys. Oceanogr.*, 27(9), 1849–1867, doi:10.1175/1520-0485(1997)027<1849:AEVPMF>2.0.CO;2, 1997.
- Hunke, E. C., Lipscomb, W. H., Turner, A. K., Jeffery, N. and Elliott, S.: CICE: the Los Alamos Sea Ice Model Documentation and Software User's Manual Version 4.1 LA-CC-06-012., 2010.
- Hunke, E. C., Hebert, D. A. and Lecomte, O.: Level-ice melt ponds in the Los Alamos sea ice model, CICE, *Ocean Model.*, 71, 26–42, doi:10.1016/j.ocemod.2012.11.008, 2013.

Moved (insertion) [11]

Deleted: core

Moved up [11]: Griffies, S. M., Yin, J., Durack, P. J., Goddard, P., Bates, S. C., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Böning, C. W., Bozec, A., Chassignet, E., Danabasoglu, G., Danilov, S., Domingues, C. M., Drange, H., Farneti, R., Fernandez, E., Greatbatch, R. J., Holland, D. M., Ilicak, M., Large, W. G., Lorbacher, K., Lu, J., Marsland, S. J., Mishra, A., George Nurser, A. J., Salas y Mélia, D., Palter, J. B., Samuels, B. L., Schröter, J., Schwarzkopf, F. U., Sidorenko, D., Treguier, A. M., Tseng, Y. heng, Tsujino, H., Uotila, P., Valcke, S., Voldoire, A., Wang, Q., Winton, M. and Zhang, X.: An assessment of global and regional sea level for years 1993–2007 in a suite of interannual core-II simulations, *Ocean Model.*, 78, 35–89, doi:10.1016/j.ocemod.2014.03.004, 2014.¶

Deleted: Hailong, L., Xuehong, Z., Wei, L., Yongqiang, Y. and Rucong, Y.: An eddy-permitting oceanic general circulation model and its preliminary evaluation, *Adv. Atmos. Sci.*, 21(5), 675–690, doi:10.1007/bf02916365, 2004.¶

- Ilicak, M., Drange, H., Wang, Q., Gerdes, R., Aksenov, Y., Bailey, D., Bentsen, M., Biastoch, A., Bozec, A., Böning, C., Cassou, C., Chassignet, E., Coward, A. C., Curry, B., Danabasoglu, G., Danilov, S., Fernandez, E., Fogli, P. G., Fujii, Y., Griffies, S. M., Iovino, D., Jahn, A., Jung, T., Large, W. G., Lee, C., Lique, C., Lu, J., Masina, S., George Nurser, A. J., Roth, C., Salas y Mélia, D., Samuels, B. L., Spence, P., Tsujino, H., Valcke, S., Voldoire, A., Wang, X. and Yeager, S. G.: An assessment of the Arctic Ocean in a suite of interannual CORE-II simulations. Part III: Hydrography and fluxes, *Ocean Model.*, 100, 141–161, doi:10.1016/j.ocemod.2016.02.004, 2016.
- Jayne, S. R.: The Impact of Abyssal Mixing Parameterizations in an Ocean General Circulation Model, *J. Phys. Oceanogr.*, 39(7), 1756–1775, doi:10.1175/2009JPO4085.1, 2009.
- Jochum, M., Fox-Kemper, B., Molnar, P. H. and Shields, C.: Differences in the Indonesian seaway in a coupled climate model and their relevance to Pliocene climate and El Niño, *Paleoceanography*, 24(1), n/a-n/a, doi:10.1029/2008PA001678, 2009.
- Johns, W. E., Baringer, M. O., Beal, L. M., Cunningham, S. A., Kanzow, T., Bryden, H. L., Hirschi, J. J. M., Marotzke, J., Meinen, C. S., Shaw, B. and Curry, R.: Continuous, Array-Based Estimates of Atlantic Ocean Heat Transport at 26.5°N, *J. Clim.*, 24(10), 2429–2449, doi:10.1175/2010JCLI3997.1, 2011.
- Karspeck, A. R., Stammer, D., Köhl, A., Danabasoglu, G., Balmaseda, M., Smith, D. M., Fujii, Y., Zhang, S., Giese, B., Tsujino, H. and Rosati, A.: Comparison of the Atlantic meridional overturning circulation between 1960 and 2007 in six ocean reanalysis products, *Clim. Dyn.*, 49(3), 957–982, doi:10.1007/s00382-015-2787-7, 2017.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K. and Kiyotoshi, T.: The JRA-55 reanalysis: General specifications and basic characteristics, *J. Meteorol. Soc. Japan*, 93(1), 5–48, doi:10.2151/jmsj.2015-001, 2015.
- Koenig, Z., Provost, C., Ferrari, R., Sennéchal, N. and Rio, M.-H.: Volume transport of the Antarctic Circumpolar Current: Production and validation of a 20 year long time series obtained from in situ and satellite observations, *J. Geophys. Res. Ocean.*, 119(8), 5407–5433, doi:10.1002/2014jc009966, 2014.
- Kohout, A., Williams, M., Dean, S. and Meylan, M.H.: Storm-induced sea-ice breakup and the implications for ice extent, *Nature*, 509, 604–607, doi:10.1038/nature13262, 2014
- Kurian, J., P., Li, P., Chang, P., Patricola, C. M. and Small, J.: Impact of the Benguela coastal low-level jet on the Southeast Tropical Atlantic SST bias in a regional ocean model (submitted), *Clim. Dyn.*, 2020.
- Large, W. G. and Yeager, S. G.: Diurnal to decadal global forcing for ocean and sea-ice models: The data sets and flux climatologies, NCAR Tech. Note, TN-460+ST(May), 105pp, doi:10.5065/D6KK98Q6, 2004.
- Large, W. G., McWilliams, J. C. and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization, *Rev. Geophys.*, 32(4), 363, doi:10.1029/94RG01872, 1994.
- Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air - Sea flux data set, *Clim. Dyn.*, 33(2–3), 341–364, doi:10.1007/s00382-008-0441-3, 2009.
- Large, W. G., Danabasoglu, G., Doney, S. C. and McWilliams, J. C.: Sensitivity to Surface Forcing and Boundary Layer Mixing in a Global Ocean Model: Annual-Mean Climatology, *J. Phys. Oceanogr.*, 27(11), 2418–2447, doi:10.1175/1520-0485(1997)027<2418:STSFAB>2.0.CO;2, 1997.
- Laurindo, L. C., Siqueira, L., Mariano, A. J., and Kirtman, B. P.: Cross-spectral analysis of the SST/10-m wind speed coupling resolved by satellite products and climate model simulations, *Clim. Dyn.*, 52, 5071–5098, doi:10.1007/s00382-018-4434-6, 2019
- Li, L., Lin, P., Yu, Y., Wang, B., Zhou, T., Liu, L., Liu, J., Bao, Q., Xu, S., Huang, W., Xia, K., Pu, Y., Dong, L., Shen, S., Liu, Y., Hu, N., Liu, M., Sun, W., Shi, X., Zheng, W., Wu, B., Song, M., Liu, H., Zhang, X., Wu, G., Xue, W., Huang, X., Yang, G., Song, Z. and Qiao, F.: The flexible global ocean-atmosphere-land system model, Grid-point Version 2: FGOALS-g2, *Adv. Atmos. Sci.*, 30(3), 543–560, doi:10.1007/s00376-012-2140-6, 2013.
- Li, Q., Webb, A., Fox-Kemper, B., Craig, A., Danabasoglu, G., Large, W. G. and Vertenstein, M.: Langmuir mixing effects on global climate: WAVEWATCH III in CESM, *Ocean Model.*, 103, 145–160, doi:10.1016/j.ocemod.2015.07.020, 2016.
- Lin, P., Liu, H., Xue, W., Li, H., Jiang, J., Song, M., Song, Y., Wang, F. and Zhang, M.: A coupled experiment with LICOM2 as the ocean component of CESM1, *J. Meteorol. Res.*, 30(1), 76–92, doi:10.1007/s13351-015-5045-3, 2016.
- Lin, P., Liu, H. and Zhang, X.: Sensitivity of the upper ocean temperature and circulation in the equatorial Pacific to solar radiation penetration due to phytoplankton, *Adv. Atmos. Sci.*, 24(5), 765–780, doi:10.1007/s00376-007-0765-7, 2007.

Formatted: English (United States)

Liu, H. L. and Lin, P. F.: A global eddy-resolving ocean forecast system in China -- LICOM Forecast System (LFS) (under review), *J. Oper. Oceanogr.*, 2020.

1760 [Liu, H. L., Zhang, X. H., Li, W., Yu, Y. Q. and Yu R. C.: An eddy-permitting oceanic general circulation model and its preliminary evaluation, \*Adv. Atmos. Sci.\*, 21\(5\), 675–690, doi:10.1007/bf02916365, 2004.](#)

Liu, H., Lin, P., Yu, Y. and Zhang, X.: The baseline evaluation of LASG/IAP Climate system Ocean Model (LICOM) version 2, *Acta Meteorol. Sin.*, 26(3), 318–329, doi:10.1007/s13351-012-0305-y, 2012.

1765 Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., Seidov, D. and Levitus, S.: *World Ocean Atlas 2013, Volume 1, Temperature*, doi:10.7289/V55X26VD, 2013.

Lumpkin, R. and Speer, K.: Global Ocean Meridional Overturning, *J. Phys. Oceanogr.*, 37(10), 2550–2562, doi:10.1175/JPO3130.1, 2007.

1770 Ma, X., Chang, P., Saravanan, R., Montuoro, R., Nakamura, H., Wu, D., Lin, X. and Wu, L.: Importance of Resolving Kuroshio Front and Eddy Influence in Simulating the North Pacific Storm Track, *J. Clim.*, 30(5), 1861–1880, doi:10.1175/JCLI-D-16-0154.1, 2017.

Ma, X., Jing, Z., Chang, P., Liu, X., Montuoro, R., Small, R. J., Bryan, F. O., Greatbatch, R. J., Brandt, P., Wu, D., Lin, X. and Wu, L.: Western boundary currents regulated by interaction between ocean eddies and the atmosphere, *Nature*, 535(7613), 533–537, doi:10.1038/nature18640, 2016.

1775 Maltrud, M. E. and McClean, J. L.: An eddy resolving global 1/10° ocean simulation, *Ocean Model.*, 8(1–2), 31–54, doi:10.1016/j.ocemod.2003.12.001, 2005.

McClean, J. L., Bader, D. C., Bryan, F. O., Maltrud, M. E., Dennis, J. M., Mirin, A. A., Jones, P. W., Kim, Y. Y., Ivanova, D. P., Vertenstein, M., Boyle, J. S., Jacob, R. L., Norton, N., Craig, A. and Worley, P. H.: A prototype two-decade fully-coupled fine-resolution CCSM simulation, *Ocean Model.*, 39(1–2), 10–30, doi:10.1016/j.ocemod.2011.02.011, 2011.

1780 Meinen, C. S., Speich, S., Perez, R. C., Dong, S., Piola, A. R., Garzoli, S. L., Baringer, M. O., Gladyshev, S. and Campos, E. J. D.: Temporal variability of the meridional overturning circulation at 34.5°S: Results from two pilot boundary arrays in the South Atlantic, *J. Geophys. Res. Ocean.*, 118(12), 6461–6478, doi:10.1002/2013JC009228, 2013.

Meinen, C. S., Speich, S., Piola, A. R., Ansong, I., Campos, E., Kersalé, M., Terre, T., Chidichimo, M. P., Lamont, T., Sato, O. T., Perez, R. C., Valla, D., van den Berg, M., Le Hénaff, M., Dong, S. and Garzoli, S. L.: Meridional Overturning Circulation Transport Variability at 34.5°S During 2009–2017: Baroclinic and Barotropic Flows and the Dueling Influence of the Boundaries, *Geophys. Res. Lett.*, 45(9), 4180–4188, doi:10.1029/2018GL077408, 2018.

1785 Mesinger, F. and Janjic, Z. I.: Problems and numerical methods of the incorporation of mountains in atmospheric models, *Lect. Appl. Math.*, 22, 81–120, 1985.

Meyers, G.: Variation of Indonesian throughflow and the El Niño–Southern Oscillation, *J. Geophys. Res. Ocean.*, 101(C5), 12255–12263, doi:10.1029/95JC03729@10.1002/(ISSN)2169-9291.PACLLWBC1, 1996.

1790 Morrison, A. K. and McC. Hogg, A.: On the Relationship between Southern Ocean Overturning and ACC Transport, *J. Phys. Oceanogr.*, 43(1), 140–148, doi:10.1175/JPO-D-12-057.1, 2013.

Msadek, R., Johns, W. E., Yeager, S. G., Danabasoglu, G., Delworth, T. L. and Rosati, A.: The Atlantic Meridional Heat Transport at 26.5°N and Its Relationship with the MOC in the RAPID Array and the GFDL and NCAR Coupled Models, *J. Clim.*, 26(12), 4335–4356, doi:10.1175/JCLI-D-12-00081.1, 2013.

1795 Munday, D. R., Johnson, H. L. and Marshall, D. P.: Eddy Saturation of Equilibrated Circumpolar Currents, *J. Phys. Oceanogr.*, 43(3), 507–532, doi:10.1175/JPO-D-12-095.1, 2013.

Murray, R. J.: Explicit generation of orthogonal grids for ocean models, *J. Comput. Phys.*, 126(2), 251–273, doi:10.1006/jcph.1996.0136, 1996.

[Notz, D., Dörr, J., Bailey, D. A., Blockley, E., Bushuk, M., Debernard, J. B., Dekker, E., DeRepentigny, P., Docquier, D., Fuckar, N. S., Fyfe, J. C., Jahn, A., Holland, M., Hunke, E., Iovino, D., Khosravi, N., Massonnet, F., Madec, G., O'Farrell, S., Petty, A., Rana, A., Roach, L., Rosenblum, E., Rousset, C., Semmler, T., Stroeve, J., Tremblay, B., Toyoda, T., Tsujino, H. and Vancoppenolle, M.: Arctic sea ice in CMIP6, \*Geophys. Res. Lett.\*, 47, e2019GL086749, doi:10.1029/2019GL086749, 2020.](#)

1800 Ohlmann, J. C.: Ocean Radiant Heating in Climate Models, *J. Clim.*, 16(9), 1337–1351, doi:10.1175/1520-0442-16.9.1337, 2003.

Ollitrault, M. and Colin de Verdière, A.: The Ocean General Circulation near 1000-m Depth, *J. Phys. Oceanogr.*, 44(1), 384–409, doi:10.1175/JPO-D-13-030.1, 2014.

1805 [Ollitrault, M. and Colin de Verdière, A.: The ocean general circulation near 1000 m depth, \*J. Phys. Oceanogr.\*, 44, 384–408, doi:10.1175/JPO-D-13-030.1, 2014.](#)

Deleted: 1948-

Deleted: o

Deleted: a

Deleted: ,

Olonscheck, D., Mauritsen, T. and Notz, D.: Arctic sea-ice variability is primarily driven by atmospheric temperature fluctuations, *Nat. Geosci.*, 12(6), 430–434, doi:10.1038/s41561-019-0363-1, 2019.

Parkinson, C. L. and Washington, W. M.: LARGE-SCALE NUMERICAL MODEL OF SEA ICE., *J Geophys Res*, 84(C1), 311–337, doi:10.1029/jc084ic01p00311, 1979.

Pearson, B., Fox-Kemper, B., Bachman, S. and Bryan, F.: Evaluation of scale-aware subgrid mesoscale eddy models in a global eddy-rich model, *Ocean Model.*, 115, 42–58, doi:10.1016/j.ocemod.2017.05.007, 2017.

Rackow, T., Goessling, H. F., Jung, T., Sidorenko, D., Semmler, T., Barbi, D. and Handorf, D.: Towards multi-resolution global climate modeling with ECHAM6-FESOM. Part II: climate variability, *Clim. Dyn.*, 50(7–8), 2369–2394, doi:10.1007/s00382-016-3192-6, 2018.

Rackow, T., Sein, D. V., Semmler, T., Danilov, S., Koldunov, N. V., Sidorenko, D., Wang, Q. and Jung, T.: Sensitivity of deep ocean biases to horizontal resolution in prototype CMIP6 simulations with AWI-CM1.0, *Geosci. Model Dev.*, 12(7), 2635–2656, doi:10.5194/gmd-12-2635-2019, 2019.

Rahaman, H., Srinivasu, U., Panickal, S., Durgadoo, J. V., Griffies, S. M., Ravichandran, M., Bozec, A., Cherchi, A., Voldoire, A., Sidorenko, D., Chassignet, E. P., Danabasoglu, G., Tsujino, H., Getzlaff, K., Ilicak, M., Bentsen, M., Long, M. C., Fogli, P. G., Farneti, R., Danilov, S., Marsland, S. J., Valcke, S., Yeager, S. G. and Wang, Q.: An assessment of the Indian Ocean mean state and seasonal cycle in a suite of interannual CORE-II simulations, *Ocean Model.*, 145, doi:10.1016/j.ocemod.2019.101503, 2020.

Redi, M. H.: Oceanic isopycnal mixing by coordinate rotation, *J. Phys. Oceanogr.*, 12(10), 1154–1158, doi:10.1175/1520-0485(1982)012<1154:OIMBCR>2.0.CO;2, 1982.

Renault, L., Masson, S., Arsouze, T., Madec, G. and McWilliams, J. C.: Recipes for how to force oceanic model dynamics, *J. Adv. Model. Earth Syst.*, 2019MS001715, doi:10.1029/2019MS001715, 2020.

Renault, L., McWilliams, J. C. and Penven, P.: Modulation of the Agulhas Current retroflection and leakage by oceanic current interaction with the atmosphere in coupled simulations, *J. Phys. Oceanogr.*, 47(8), 2077–2100, doi:10.1175/JPO-D-16-0168.1, 2017.

Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S. and Schlax, M. G.: Daily High-Resolution-Blended Analyses for Sea Surface Temperature, *J. Clim.*, 20(22), 5473–5496, doi:10.1175/2007JCLI1824.1, 2007.

Richardson, P. L.: A census of eddies observed in North Atlantic SOFAR float data, *Prog. Oceanogr.*, 31(1), 1–50, doi:10.1016/0079-6611(93)90022-6, 1993.

Roach, L. A., Horvat, C., Dean, S. M. and Bitz, C. M.: An emergent sea ice floe size distribution in a global coupled ocean-sea ice model, *Journal of Geophysical Research: Oceans*, 123, 4322–4337, doi:10.1029/2017JC013692, 2018.

Roach, L. A., Dörr, J., Holmes, C. R., Massonnet, F., Blockley, E. W., Notz, D., Rackow, T., Raphael, M. N., O’Farrell, S. P., Bailey, D. A. and Bitz, C. M.: Antarctic sea ice area in CMIP6, *Geophys. Res. Lett.*, 47, e2019GL086729, doi:10.1029/2019GL086729, 2020.

Rossby, T.: The North Atlantic Current and surrounding waters: At the crossroads, *Rev. Geophys.*, 34(4), 463–481, doi:10.1029/96RG02214, 1996.

Rucong, Y.: A two-step shape-preserving advection scheme, *Adv. Atmos. Sci.*, 11(4), 479–490, doi:10.1007/BF02658169, 1994.

Schmitz Jr, W. J.: On the World Ocean Circulation. Volume 1. Some Global Features/North Atlantic Circulation., 1996.

Schweiger, A., Lindsay, R., Zhang, J., Steele, M., Stern, H. and Kwok, R.: Uncertainty in modeled Arctic sea ice volume, *J. Geophys. Res.*, 116, C00D06, doi:10.1029/2011JC007084, 2011.

Scott, R. B., Arbic, B. K., Chassignet, E. P., Coward, A. C., Maltrud, M., Merryfield, W. J., Srinivasan, A. and Varghese, A.: Total kinetic energy in four global eddying ocean circulation models and over 5000 current meter records, *Ocean Model.*, 32(3–4), 157–169, doi:10.1016/j.ocemod.2010.01.005, 2010.

Sein, D. V., Danilov, S., Biastoch, A., Durgadoo, J. V., Sidorenko, D., Harig, S. and Wang, Q.: Designing variable ocean model resolution based on the observed ocean variability, *J. Adv. Model. Earth Syst.*, 8(2), 904–916, doi:10.1002/2016MS000650, 2016.

Sein, D. V., Koldunov, N. V., Danilov, S., Sidorenko, D., Wekerle, C., Cabos, W., Rackow, T., Scholz, P., Semmler, T., Wang, Q. and Jung, T.: The Relative Influence of Atmospheric and Oceanic Model Resolution on the Circulation of the North Atlantic Ocean in a Coupled Climate Model, *J. Adv. Model. Earth Syst.*, 10(8), 2026–2041, doi:10.1029/2018MS001327, 2018.

Deleted: H

Deleted: F

Deleted: O

Deleted: M

Deleted: D

Deleted: 19

Formatted: Default Paragraph Font, English (United States)

Formatted: English (United States)

- Semtner, A. J.: A Model for the Thermodynamic Growth of Sea Ice in Numerical Investigations of Climate, *J. Phys. Oceanogr.*, 6(3), 379–389, doi:10.1175/1520-0485(1976)006<0379:amftg>2.0.co;2, 1976.
- [Shu, Q., Wang, Q., Song, Z., Qiao, F., Zhao, J., Chu, M. and Li, X.: Assessment of sea ice extent in CMIP6 with comparison to observations and CMIP5, \*Geophys. Res. Lett.\*, 47, e2020GL087965, doi:10.1029/2020GL087965, 2020.](#)
- Sidorenko, D., Koldunov, N. V., Wang, Q., Danilov, S., Goessling, H. F., Gurses, O., Scholz, P., Sein, D. V., Volodin, E., Wekerle, C. and Jung, T.: Influence of a salt plume parameterization in a coupled climate model, *J. Adv. Model. Earth Syst.*, 10(9), 2357–2373, doi:10.1029/2018MS001291, 2018.
- Sidorenko, D., Rackow, T., Jung, T., Semmler, T., Barbi, D., Danilov, S., Dethloff, K., Dorn, W., Fieg, K., Goessling, H. F., Handorf, D., Harig, S., Hiller, W., Juricke, S., Losch, M., Schröter, J., Sein, D. V. and Wang, Q.: Towards multi-resolution global climate modeling with ECHAM6–FESOM. Part I: model formulation and mean climate, *Clim. Dyn.*, 44(3–4), 757–780, doi:10.1007/s00382-014-2290-6, 2014.
- Smagorinsky, J.: General circulation experiments with the primitive equations, *Mon. Weather Rev.*, 91(3), 99–164, doi:10.1175/1520-0493(1963)091<0099:gcwtp>2.3.co;2, 1963.
- Small, R. J., Bacmeister, J., Bailey, D., Baker, A., Bishop, S., Bryan, F., Caron, J., Dennis, J., Gent, P., Hsu, H., Jochum, M., Lawrence, D., Muñoz, E., DiNezio, P., Scheitlin, T., Tomas, R., Tribbia, J., Tseng, Y. and Vertenstein, M.: A new synoptic scale resolving global climate simulation using the Community Earth System Model, *J. Adv. Model. Earth Syst.*, 6(4), 1065–1094, doi:10.1002/2014MS000363, 2014.
- Small, R. J., Curchitser, E., Hedstrom, K., Kauffman, B. and Large, W. G.: The Benguela upwelling system: Quantifying the sensitivity to resolution and coastal wind representation in a global climate model, *J. Clim.*, 28(23), 9409–9432, doi:10.1175/JCLI-D-15-0192.1, 2015.
- Smeed, D. A., Josey, S. A., Beaulieu, C., Johns, W. E., Moat, B. I., Frajka-Williams, E., Rayner, D., Meinen, C. S., Baringer, M. O., Bryden, H. L. and McCarthy, G. D.: The North Atlantic Ocean is in a state of reduced overturning, *Geophys. Res. Lett.*, 45(3), 1527–1533, doi:10.1002/2017GL076350, 2018.
- Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., Dukowicz, J., Eden, C., Fox-Kemper, B., Gent, P. and others: The parallel ocean program (POP) reference manual: ocean component of the community climate system model (CCSM) and community earth system model (CESM), Rep. LAUR-01853, 141, 1–140, 2010.
- Sprattall, J., Wijffels, S. E., Molcard, R. and Jaya, I.: Direct estimates of the Indonesian Throughflow entering the Indian Ocean: 2004–2006, *J. Geophys. Res.*, 114(C7), C07001, doi:10.1029/2008JC005257, 2009.
- St. Laurent, L. C., Simmons, H. L. and Jayne, S. R.: Estimating tidally driven mixing in the deep ocean, *Geophys. Res. Lett.*, 29(23), 21-1-21–4, doi:10.1029/2002gl015633, 2002.
- Steele, M., Morley, R. and Ermold, W.: PHC: A Global Ocean Hydrography with a High-Quality Arctic Ocean, *J. Clim.*, 14(9), 2079–2087, doi:10.1175/1520-0442(2001)014<2079:PAGOHV>2.0.CO;2, 2001.
- [Stewart, K.D., Hogg, A. McC., Griffies, S. M., Heerdegen, A. P., Ward, M. L., Spence, P. and England, M. H.: Vertical resolution of baroclinic modes in global ocean models, \*Ocean Modell.\*, 113, 50–65, doi:10.1016/j.ocemod.2017.03.012, 2017.](#)
- [Stroeve, J. C., Kattsov, V., Barrett, A., Serreze, M., Pavlova, T., Holland, M. and Meier, W. N.: Trends in Arctic sea ice extent from CMIP5, CMIP3 and observations, \*Geophys. Res. Lett.\*, 39, L16502, doi:10.1029/2012GL052676, 2012.](#)
- Sun, Q., Whitney, M. M., Bryan, F. O. and Tseng, Y.: Assessing the skill of the improved treatment of riverine freshwater in the Community Earth System Model (CESM) relative to a new salinity climatology, *J. Adv. Model. Earth Syst.*, 11(5), 1189–1206, doi:10.1029/2018MS001349, 2019.
- Suzuki, T., Yamazaki, D., Tsujino, H., Komuro, Y., Nakano, H. and Urakawa, S.: A dataset of continental river discharge based on JRA-55 for use in a global ocean circulation model, *J. Oceanogr.*, 74(4), 421–429, doi:10.1007/s10872-017-0458-5, 2018.
- Talley, L.: Closure of the Global Overturning Circulation Through the Indian, Pacific, and Southern Oceans: Schematics and Transports, *Oceanography*, 26(1), 80–97, doi:10.5670/oceanog.2013.07, 2013.
- Talley, L. D.: Freshwater transport estimates and the global overturning circulation: Shallow, deep and throughflow components, *Prog. Oceanogr.*, 78(4), 257–303, doi:10.1016/j.pocean.2008.05.001, 2008.
- Teague, W. J., Carron, M. J. and Hogan, P. J.: A comparison between the Generalized Digital Environmental Model and Levitus climatologies, *J. Geophys. Res.*, 95(C5), 7167, doi:10.1029/jc095ic05p07167, 1990.

Formatted: English (United States)

- Thoppil, P. G., Richman, J. G. and Hogan, P. J.: Energetics of a global ocean circulation model compared to observations, *Geophys. Res. Lett.*, 38(15), doi:10.1029/2011GL048347, 2011.
- Trenberth, K. E., Zhang, Y., Fasullo, J. T. and Cheng, L.: Observation-based estimates of global and basin ocean meridional heat transport time series, *J. Clim.*, 32(14), 4567–4583, doi:10.1175/JCLI-D-18-0872.1, 2019.
- 1920 Tseng, Y., Lin, H., Chen, H. ching, Thompson, K., Bentsen, M., Böning, C. W., Bozec, A., Cassou, C., Chassignet, E., Chow, C. H., Danabasoglu, G., Danilov, S., Farneti, R., Fogli, P. G., Fujii, Y., Griffies, S. M., Ilicak, M., Jung, T., Masina, S., Navarra, A., Patara, L., Samuels, B. L., Scheinert, M., Sidorenko, D., Sui, C. H., Tsujino, H., Valcke, S., Voldoire, A., Wang, Q. and Yeager, S. G.: North and equatorial Pacific Ocean circulation in the CORE-II hindcast simulations, *Ocean Model.*, 104, 143–170, doi:10.1016/j.ocemod.2016.06.003, 2016.
- 1925 Tsujino, H., Urakawa, L. S., Griffies, S. M., Danabasoglu, G., Adcroft, A. J., Amaral, A. E., Arsouze, T., Bentsen, M., Bernardello, R., Böning, C. W., Bozec, A., Chassignet, E. P., Danilov, S., Dussin, R., Exarchou, E., Fogli, P. G., Fox-Kemper, B., Guo, C., Ilicak, M., Iovino, D., Kim, W. M., Koldunov, N., Lapin, V., Li, Y., Lin, P., Lindsay, K., Liu, H., Long, M. C., Komuro, Y., Marsland, S. J., Masina, S., Nummelin, A., Rieck, J. K., Ruprich-Robert, Y., Scheinert, M., Sicardi, V., Sidorenko, D., Suzuki, T., Tatebe, H., Wang, Q., Yeager, S. G. and Yu, Z.: Evaluation of global ocean–sea-ice model simulations based on the experimental protocols of the Ocean Model Intercomparison Project phase 2 (OMIP-2) (submitted), *Geosci. Model Dev.*, 2020.
- 1930 Tsujino, H., Urakawa, S., Nakano, H., Small, R. J., Kim, W. M., Yeager, S. G., Danabasoglu, G., Suzuki, T., Bamber, J. L., Bentsen, M., Böning, C. W., Bozec, A., Chassignet, E. P., Curchitser, E., Boeira Dias, F., Durack, P. J., Griffies, S. M., Harada, Y., Ilicak, M., Josey, S. A., Kobayashi, C., Kobayashi, S., Komuro, Y., Large, W. G., Le Sommer, J., Marsland, S. J., Masina, S., Scheinert, M., Tomita, H., Valdivieso, M. and Yamazaki, D.: JRA-55 based surface dataset for driving ocean–sea-ice models (JRA55-do), *Ocean Model.*, 130, 79–139, doi:10.1016/j.ocemod.2018.07.002, 2018.
- 1935 Turner, A. K. and Hunke, E. C.: Impacts of a mushy-layer thermodynamic approach in global sea-ice simulations using the CICE sea-ice model, *J. Geophys. Res. Ocean.*, 120(2), 1253–1275, doi:10.1002/2014JC010358, 2015.
- von Storch, J.-S., Haak, H., Hertwig, E. and Fast, I.: Vertical heat and salt fluxes due to resolved and parameterised meso-scale eddies, *Ocean Model.*, 108, 1–19, doi:10.1016/j.ocemod.2016.10.001, 2016.
- 1940 Wang, Q., Danilov, S., Jung, T., Kaleschke, L. and Wernecke, A.: Sea ice leads in the Arctic Ocean: Model assessment, interannual variability and trends, *Geophys. Res. Lett.*, 43(13), 7019–7027, doi:10.1002/2016GL068696, 2016.
- Wang, Q., Danilov, S. and Schröter, J.: Finite element ocean circulation model based on triangular prismatic elements, with application in studying the effect of topography representation, *J. Geophys. Res.*, 113(C5), C05015, doi:10.1029/2007JC004482, 2008.
- 1945 Wang, Q., Danilov, S., Sidorenko, D., Timmermann, R., Wekerle, C., Wang, X., Jung, T. and Schröter, J.: The Finite Element Sea Ice-Ocean Model (FESOM) v.1.4: formulation of an ocean general circulation model, *Geosci. Model Dev.*, 7(2), 663–693, doi:10.5194/gmd-7-663-2014, 2014.
- 1950 Wang, Q., Ilicak, M., Gerdes, R., Drange, H., Aksenov, Y., Bailey, D. A., Bentsen, M., Biastoch, A., Bozec, A., Böning, C., Cassou, C., Chassignet, E., Coward, A. C., Curry, B., Danabasoglu, G., Danilov, S., Fernandez, E., Fogli, P. G., Fujii, Y., Griffies, S. M., Iovino, D., Jahn, A., Jung, T., Large, W. G., Lee, C., Lique, C., Lu, J., Masina, S., Nurser, A. J. G., Rabe, B., Roth, C., Salas y Mélia, D., Samuels, B. L., Spence, P., Tsujino, H., Valcke, S., Voldoire, A., Wang, X. and Yeager, S. G.: An assessment of the Arctic Ocean in a suite of interannual CORE-II simulations. Part I: Sea ice and solid freshwater, *Ocean Model.*, 99, 110–132, doi:10.1016/j.ocemod.2015.12.008, 2016.
- 1955 Wang, Q., Ilicak, M., Gerdes, R., Drange, H., Aksenov, Y., Bailey, D. A., Bentsen, M., Biastoch, A., Bozec, A., Böning, C., Cassou, C., Chassignet, E., Coward, A. C., Curry, B., Danabasoglu, G., Danilov, S., Fernandez, E., Fogli, P. G., Fujii, Y., Griffies, S. M., Iovino, D., Jahn, A., Jung, T., Large, W. G., Lee, C., Lique, C., Lu, J., Masina, S., Nurser, A. J. G., Rabe, B., Roth, C., Salas y Mélia, D., Samuels, B. L., Spence, P., Tsujino, H., Valcke, S., Voldoire, A., Wang, X. and Yeager, S. G.: An assessment of the Arctic Ocean in a suite of interannual CORE-II simulations. Part II: Liquid freshwater, *Ocean Model.*, 99, 86–109, doi:10.1016/j.ocemod.2015.12.009, 2016.
- 1960 Whitworth, T.: Monitoring the transport of the Antarctic Circumpolar Current at Drake Passage, *J. Phys. Oceanogr.*, 13(11), 2045–2057, doi:10.1175/1520-0485(1983)013<2045:mttota>2.0.co;2, 1983.
- Whitworth, T. and Peterson, R. G.: Volume transport of the Antarctic Circumpolar Current from Bottom pressure measurements, *J. Phys. Oceanogr.*, 15(6), 810–816, doi:10.1175/1520-0485(1985)015<0810:vtotac>2.0.co;2, 1985.



- 1965 Xiao, C.: Adoption of a two-step shape-preserving advection scheme in an OGCM and its coupled experiment, 89pp, MS thesis, Inst. Atmos. Physics, Chinese Acad. Sci. (in Chinese), 2006.
- Xu, X., Chassignet, E. P., Dong, S. and Baringer, M. O.: Transport structure of the South Atlantic Ocean derived from a high-resolution numerical model and observations, *J. Geophys. Res.* (submitted), 2020a.
- 1970 Xu, X., Chassignet, E.P., Firing, Y. L. and Donohue, K.: Antarctic Circumpolar Current transport through Drake Passage: What can we learn from comparing high-resolution model results to observations? *J. Geophys. Res. Ocean.*, doi:10.1029/2020JC016365, 2020b.
- Xu, X., Hurlburt, H. E., Schmitz, W. J., Zantopp, R., Fischer, J. and Hogan, P. J.: On the currents and transports connected with the atlantic meridional overturning circulation in the subpolar North Atlantic, *J. Geophys. Res. Ocean.*, 118(1), 502–516, doi:10.1002/jgrc.20065, 2013.
- 1975 Xu, X., Chassignet, E. P., Johns, W. E., Schmitz, W. J. and Metzger, E. J.: Intraseasonal to interannual variability of the Atlantic meridional overturning circulation from eddy-resolving simulations and observations, *J. Geophys. Res. Ocean.*, 119(8), 5140–5159, doi:10.1002/2014jc009994, 2014.
- Xu, X., Chassignet, E. P. and Wang, F.: On the variability of the Atlantic meridional overturning circulation transports in coupled CMIP5 simulations, *Clim. Dyn.*, 52(11), 6511–6531, doi:10.1007/s00382-018-4529-0, 2019.
- 1980 Xuehong, Z. and Xinzhong, L.: A numerical world ocean general circulation model, *Adv. Atmos. Sci.*, 6(1), 44–61, doi:10.1007/BF02656917, 1989.
- Yashayaev, I.: Hydrographic changes in the Labrador Sea, 1960–2005, *Prog. Oceanogr.*, 73(3–4), 242–276, doi:10.1016/j.pocean.2007.04.015, 2007.
- Yeager, S. G. and Large, W. G.: Late-Winter Generation of Spiciness on Subducted Isopycnals, *J. Phys. Oceanogr.*, 34(7), 1528–1547, doi:10.1175/1520-0485(2004)034<1528:LGOSOS>2.0.CO;2, 2004.
- 1985 Yu, Y. Q., Tang, S. L., Liu, H. L., Lin, P. F. and Li, X. L.: Development and Evaluation of the Dynamic Framework of an Ocean General Circulation Model with Arbitrary Orthogonal Curvilinear Coordinate, *Chinese J. Atmos. Sci.* (in Chinese), 42(4), 877–889, doi:j.issn.1006-9895.1805.17284, 2008.
- Yu, Z. P., Liu, H. L. and Lin, P. F.: A numerical study of the influence of tidal mixing on Atlantic meridional overturning circulation (AMOC) Simulation, *Chinese J. Atmos. Sci.* (in Chinese), 41(5), doi:j.issn.1006-9895.1702.16263, 2017.
- 1990 Zweng, M. M., Reagan, J. R., Antonov, J. I., Locarnini, R. A., Mishonov, A. V., Boyer, T. P., Garcia, H. E., Baranova, O. K., Johnson, D. R., Seidov, D. and others: World ocean atlas 2013. Volume 2, Salinity, 2013.

**Deleted:** (submitted)

**Formatted:** English (United States)

**Formatted:** English (United States)

**Formatted:** English (United States)



Table 1: Model parameters for the low-and high-resolution configurations

	Horizontal grid spacing	Explicit horizontal viscosity	Explicit vertical viscosity	Explicit horizontal diffusivity
<i>FSU-HYCOM low-resolution</i>	0.72°	$A = \max(\text{Smagorinsky}, \text{Laplacian } A_2) + \text{biharmonic } A_4$ with Smagorinsky = $0.1 \Delta x^2$ deformation tensor, $A_2 = 0.03 \Delta x \text{ m}^2 \text{ s}^{-1}$ , and $A_4 = -0.05 \Delta x^3 \text{ m}^4 \text{ s}^{-1}$	Background viscosity of $3 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ in KPP	Laplacian ( $0.03 \Delta x \text{ m}^2 \text{ s}^{-1}$ )
<i>FSU-HYCOM high-resolution</i>	$1/12^\circ$ (8 km at the Equator, 6 km at mid-latitudes)	$A = \max(\text{Smagorinsky}, \text{Laplacian } A_2) + \text{Biharmonic } A_4$ with Smagorinsky = $0.05 \Delta x^2$ deformation tensor, $A_2 = 20 \text{ m}^2 \text{ s}^{-1}$ , and $A_4 = -0.01 \Delta x^3 \text{ m}^4 \text{ s}^{-1}$	Background viscosity of $3 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ in KPP	Laplacian ( $0.005 \Delta x \text{ m}^2 \text{ s}^{-1}$ )
<i>NCAR-POP low-resolution</i>	Nominal $1^\circ$ bipolar grid with tropical refinement down to $1/4^\circ$	Anisotropic horizontal viscosity (see Danabasoglu et al., 2012, for details).	Spatially variable background viscosities between $10^{-5}$ and $3 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ occurring at the equator and at about $30^\circ$ of latitude.	Isopycnal diffusivity with enhanced values that can be as large as $3000 \text{ m}^2/\text{s}$ in the upper ocean and diminish to $300 \text{ m}^2/\text{s}$ by a depth of about 2000 m.
<i>NCAR-POP high-resolution</i>	$1/10^\circ$ tripolar grid	Horizontal biharmonic ( $-2.7 \times 10^{10} \text{ m}^4 \text{ s}^{-1}$ )	Background viscosity of $10^{-4} \text{ m}^2 \text{ s}^{-1}$ at the surface up to $10^{-3} \text{ m}^2 \text{ s}^{-1}$ in the abyss in KPP.	Horizontal biharmonic diffusion ( $-3 \times 10^9 \text{ m}^4/\text{s}$ )
<i>AWI-FESOM low-resolution</i>	Nominal $1^\circ$ ; refined at the equator and around Antarctica; 25 km north of $45^\circ\text{N}$ (see Figure 1a)	Biharmonic-Smagorinsky = $\Delta x^4$ deformation tensor/32	Background viscosity of $10^{-4} \text{ m}^2 \text{ s}^{-1}$ in KPP	0
<i>AWI-FESOM high-resolution</i>	Scaled with the observed sea surface height variance, ranging from about 10 to 50 km (see Figure 1b)	Biharmonic-Smagorinsky = $\Delta x^4$ deformation tensor/32	Background viscosity of $10^{-4} \text{ m}^2 \text{ s}^{-1}$ in KPP	0
<i>IAP-LICOM low-resolution</i>	$1^\circ$ (110 km in longitude, about 110 km at equator and 80 km at mid-latitudes)	Laplacian ( $5400 \text{ m}^2/\text{s}$ )	Background viscosity of $2 \times 10^{-6} \text{ m}^2/\text{s}$ as in Canuto et al. (2001, 2002) with an upper limit of $2 \times 10^{-2} \text{ m}^2/\text{s}$	Laplacian ( $5400 \text{ m}^2/\text{s}$ )
<i>IAP-LICOM high-resolution</i>	$1/10^\circ$ (11 km in longitude, about 11 km at equator and 8 km at mid-latitudes)	Biharmonic ( $-2.8 \times 10^{10} \text{ m}^4/\text{s}$ )	Background viscosity of $2 \times 10^{-6} \text{ m}^2/\text{s}$ as in Canuto et al. (2001, 2002) with an upper limit of $2 \times 10^{-2} \text{ m}^2/\text{s}$	Biharmonic ( $-2.8 \times 10^{10} \text{ m}^2/\text{s}$ )

Formatted: English (United States)

Deleted:

Deleted: /

Deleted: <sup>2</sup>

Deleted: ×

Deleted: /s<sup>2</sup>

Deleted: /s

Deleted: /s

Formatted: English (United States)

Deleted: /s<sup>2</sup>

Deleted: ×

Deleted: /s<sup>2</sup>

Deleted: /s

Deleted: /s

Formatted: English (United States)

Deleted: /s

Formatted: English (United States)

Deleted: /s

Deleted: /s

Deleted: /s

Formatted: English (United States)

Deleted: ×

Deleted: /

Formatted: English (United States), Superscript

Deleted: <sup>2</sup>

Formatted: English (United States)

Formatted: English (United States)

Deleted: ×

Deleted: /

Formatted: English (United States), Superscript

Deleted: <sup>2</sup>

Formatted: English (United States)

Deleted:

Formatted: English (United States)

Formatted: English (United States)

**Table 1: Model parameters for the low-and high-resolution configurations (continued)**

	<i>Isopycnal scheme, e.g. GM</i>	<i>Mixed layer scheme</i>	<i>Momentum advection scheme</i>	<i>Tracer advection scheme</i>	<i>Time stepping scheme</i>
<i>FSU- HYCOM low- resolution</i>	Laplacian $(0.01\Delta x^2 s^{-1}) +$ biharmonic $(-0.02\Delta x^3 m^4 s^{-1})$ thickness diffusion	KPP	2 <sup>nd</sup> order FCT	2 <sup>nd</sup> order FCT	Split-explicit leapfrog with Asselin filter (0.125)
<i>FSU- HYCOM high- resolution</i>	Biharmonic thickness- diffusion $(-0.015 \Delta x^3 m^4 s^{-1})$	KPP	2 <sup>nd</sup> order FCT	2 <sup>nd</sup> order FCT	Split-explicit leapfrog with Asselin filter (0.125)
<i>NCAR- POP low- resolution</i>	GM + submesoscale parameterization	KPP	2 <sup>nd</sup> order centered	2 <sup>nd</sup> order centered	2 <sup>nd</sup> order leapfrog scheme with Asselin filter
<i>NCAR- POP high- resolution</i>	None	KPP	2 <sup>nd</sup> order centered	3 <sup>rd</sup> order upwind	2 <sup>nd</sup> order leapfrog scheme with averaging time step
<i>AWI- FESOM low- resolution</i>	Laplacian Redi and thickness diffusion, diffusivity flow- dependent in the range of 0 to 1500 m <sup>2</sup> as implemented in Danabasoglu et al. (2008).	KPP	Taylor-Galerkin	2 <sup>nd</sup> order FCT	Pressure split; implicit SSH
<i>AWI- FESOM high- resolution</i>	Scaled with the observed sea surface height variance, ranging from about 10 to 50 km (see Figure 1b)	KPP	Taylor-Galerkin	2 <sup>nd</sup> order FCT	Pressure split; implicit SSH
<i>IAP- LICOM low- resolution</i>	Both Redi and GM with coefficient computed as in Ferreira et al. (2005)	Canuto et al. (2001, 2002)	Two step preserving shape (Yu, 1994)	Two step preserving shape (Yu, 1994)	Split-explicit leapfrog with Asselin filter (0.2 for barotropic; 0.43 for baroclinic; 0.43 for tracer)
<i>IAP- LICOM high- resolution</i>	1/10° (11 km in longitude, about 11 km at equator and 8 km at mid-latitudes)	Canuto et al. (2001, 2002)	Two step preserving shape (Yu, 1994)	Two step preserving shape (Yu, 1994)	Split-explicit leapfrog with Asselin filter (0.2 for barotropic; 0.43 for baroclinic; 0.43 for tracer)

Formatted: English (United States)

Formatted Table

Formatted: English (United States)

Deleted: /s

Deleted: /s

Formatted: English (United States)

Deleted: /s

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Table 1: Model parameters for the low-and high-resolution configurations (continued)

	Bottom drag	Surface wind-stress	Vertical coordinates	Initial conditions	SSS restoring
<i>FSU-HYCOM</i> low-resolution	Quadratic bottom drag $C_b( U  + U_{par})U^2$ with $C_b = 1.5 \times 10^{-3}$ and $U_{par} = 0.05 \text{ m s}^{-1}$	Absolute	41 hybrid layers	GDEM4	30 m/60 days to monthly GDEM
<i>FSU-HYCOM</i> high-resolution	Quadratic bottom drag $C_b( U  + U_{par})U^2$ with $C_b = 1.5 \times 10^{-3}$ and $U_{par} = 0.05 \text{ m s}^{-1}$	Absolute	36 hybrid layers	GDEM4	30 m/60 days to monthly GDEM
<i>NCAR-POP</i> low-resolution	Quadratic bottom drag with $C_b = 10^{-3}$	Relative	60 z-levels	WOA13	50 m/1 year to monthly WOA13
<i>NCAR-POP</i> high-resolution	Quadratic bottom drag with $C_b = 10^{-3}$	Relative	62 z-levels with partial-bottom-cell	WOA13	50 m/1 year to monthly WOA13
<i>AWI-FESOM</i> low-resolution	Quadratic bottom drag with $C_b = 2.5 \times 10^{-3}$	Relative	46 z-levels	WOA13	50 m/300 days to monthly PHC3.0
<i>AWI-FESOM</i> high-resolution	Quadratic bottom drag with $C_b = 2.5 \times 10^{-3}$	Relative	46 z-levels	WOA13	50 m/300 days to monthly PHC3.0
<i>IAP-LICOM</i> low-resolution	Quadratic bottom drag with $C_b = 2.6 \times 10^{-3}$	Relative	30 $\eta$ levels	PHC3.0	50 m/4 years to monthly PHC3.0 (50 m/30 days for the sea ice regions)
<i>IAP-LICOM</i> high-resolution	Quadratic bottom drag with $C_b = 2.6 \times 10^{-3}$	Relative	55 $\eta$ levels	PHC3.0	50 m/4 years to monthly PHC3.0 (50 m/30 days for the sea ice regions)

Formatted: Font: Italic, English (United States)

Formatted: English (United States)

Formatted Table

Formatted: English (United States)

Formatted

... [30]

Deleted: Relative ...bsolute wind stress

... [33]

Formatted

... [31]

Deleted: 0.72°

Formatted

... [32]

Formatted: English (United States)

Formatted: English (United States)

Formatted

... [34]

Deleted: Relative...wind stress

... [37]

Formatted

... [35]

Formatted

... [36]

Deleted: /s

Formatted: English (United States)

Deleted: wind stress

Formatted: English (United States)

Deleted: wind stress

Formatted: English (United States)

Deleted: wind stress

Formatted: English (United States)

Deleted: wind stress

Formatted: English (United States)

Deleted: wind stress

Formatted: English (United States)

Deleted: wind stress

Formatted: English (United States)

<b>Page 4: [1] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 4:01:00 PM</b>
The initial conditions in temperature and salinity are given by GDEM4. Similar to the low-resolution configuration, the turbulent air-sea fluxes are computed using the Large and Yeager (2004) bulk formulation and surface wind-stress calculation does not include the surface currents. No restoration is applied on the sea surface temperature.		
<b>Page 4: [2] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 4:47:00 PM</b>
The horizontal diffusion uses a combination of constant Laplacian diffusion of 20 m <sup>2</sup> /s and a biharmonic dissipation of 0.01Δx <sup>3</sup> for velocities; and the same biharmonic diffusion for tracers. The model baroclinic and barotropic time steps are 150s (leap-frog) and 5 s (explicit), respectively.		
<b>Page 5: [3] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 3:55:00 PM</b>
contribution is based on the Community Earth System Model version 2 (CESM2; Danabasoglu et al. 2020). The ocean component is based on the		
<b>Page 5: [4] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 3:57:00 PM</b>
For a more detailed description of the CESM2 ocean and sea-ice components, and changes relative to CESM1, the reader is referred to Danabasoglu et al. (2020). The initial conditions are derived from WOA13 (Locarnini et al., 2013; Zweng et al., 2013).In both NCAR-H and NCAR-L simulations, t		
<b>Page 5: [5] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States), Superscript		
<b>Page 5: [6] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 5: [7] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States), Superscript		
<b>Page 5: [8] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 5: [9] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States), Superscript		
<b>Page 5: [10] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 5: [11] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States), Superscript		
<b>Page 5: [12] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 5: [13] Formatted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 5:59:00 PM</b>
Space After: 0 pt		

The vertical viscosity and diffusion coefficients in the mixed layer have computed by the scheme of Canuto et al. (2001, 2002) with the background values of  $2 \cdot 10^{-6} \text{ m}^2/\text{s}$  and the upper limit of  $2 \cdot 10^{-2} \text{ m}^2/\text{s}$ . Recently, a tidal mixing scheme of St. Laurent et al. (2002) has been adopted in LICOM3 by Yu et al. (2017). The Laplacian form with the coefficient of  $5400 \text{ m}^2/\text{s}$  was chosen for the horizontal viscosity in the low-resolution version and the biharmonic form with the coefficient of  $-2.8 \cdot 10^{10} \text{ m}^4/\text{s}$  for the high-resolution experiment. In the low-resolution LICOM3, the isopycnal tracer diffusion scheme of Redi (1982) and the eddy-induced tracer transport scheme of Gent and McWilliams (1990, GM) with the same coefficients are used to parameterize the effects of mesoscale eddies on the large-scale circulation. Two tapering factors of Large et al. (1997) and a buoyancy frequency ( $N^2$ ) related thickness diffusivity of Ferreira et al. (2005) are also employed. However, for the high resolution, GM

Besides, the chlorophyll-a dependent solar penetration of Ohlmann (2003) introduced by Lin et al. (2007) was also implemented in both simulations.

are employed for both low- and high-resolution experiments following the OMIP protocol. However, onl

5

5

5

6

6

6

Sea ice is usually quantified and monitored by satellite in terms of sea ice extent, defined as the area with 15% or higher sea-ice concentration. Figures 13a-b display the modeled (annual mean) northern and southern hemisphere sea-ice extent for all simulations and are compared to the latest observations from the National Snow and Ice Data Center (NSIDC, Fetterer et al., 2017). Despite a wide range of time mean sea-ice extent, all simulations represent well the observed variability of the sea-ice extent from 1979 to 2018. In the northern hemisphere, the sea-ice extent shows a clear decline since the beginning of the monitored period, whereas, in the southern hemisphere, the sea ice extent remains more or less stable until the last few years of the integration with a small upward trend. Only the low-

resolution version of FSU-HYCOM shows an inconsistent negative trend in the southern hemisphere sea-ice extent. It is known that the observed sea-ice extent is highly correlated with the near-surface air temperature (e.g., Olonscheck et al., 2019). Thus, the consistency of the temporal variability between different simulations and observations suggests that the air-temperature in the JRA55-do is quite realistic. It is also notable that in the northern hemisphere, all of the high-resolution models show less bias in sea-ice extent than their low-resolution counterparts. In the southern hemisphere, all of the high-resolution models show less bias in extent than their low-resolution counterparts, with the exception of IAP-LICOM.

<b>Page 22: [20] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
--------------------------------	------------------------	-----------------------------

English (United States)

<b>Page 22: [20] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
--------------------------------	------------------------	-----------------------------

English (United States)

<b>Page 22: [20] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
--------------------------------	------------------------	-----------------------------

English (United States)

<b>Page 39: [21] Deleted</b>	<b>Eric Chassignet</b>	<b>6/22/2020 3:40:00 PM</b>
------------------------------	------------------------	-----------------------------

7

<b>Page 39: [21] Deleted</b>	<b>Eric Chassignet</b>	<b>6/22/2020 3:40:00 PM</b>
------------------------------	------------------------	-----------------------------

7

<b>Page 39: [21] Deleted</b>	<b>Eric Chassignet</b>	<b>6/22/2020 3:40:00 PM</b>
------------------------------	------------------------	-----------------------------

7

<b>Page 39: [21] Deleted</b>	<b>Eric Chassignet</b>	<b>6/22/2020 3:40:00 PM</b>
------------------------------	------------------------	-----------------------------

7

<b>Page 39: [21] Deleted</b>	<b>Eric Chassignet</b>	<b>6/22/2020 3:40:00 PM</b>
------------------------------	------------------------	-----------------------------

7

<b>Page 39: [21] Deleted</b>	<b>Eric Chassignet</b>	<b>6/22/2020 3:40:00 PM</b>
------------------------------	------------------------	-----------------------------

7

<b>Page 39: [22] Deleted</b>	<b>Eric Chassignet</b>	<b>6/17/2020 4:10:00 PM</b>
------------------------------	------------------------	-----------------------------

-

<b>Page 39: [22] Deleted</b>	<b>Eric Chassignet</b>	<b>6/17/2020 4:10:00 PM</b>
------------------------------	------------------------	-----------------------------

-

<b>Page 39: [23] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
--------------------------------	------------------------	-----------------------------

English (United States)

<b>Page 39: [23] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
--------------------------------	------------------------	-----------------------------

English (United States)







<b>Page 39: [24] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 39: [24] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 39: [24] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 39: [24] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 39: [24] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 41: [25] Deleted</b>	<b>Eric Chassignet</b>	<b>6/17/2020 4:42:00 PM</b>

The observed spatial pattern of winter sea-ice concentration (averaged from 1980 to 2018 for March) in the northern hemisphere is well simulated in all the models with both low- and high-resolutions, although there tends to be greater consistency among the high-resolution models. The simulated winter sea-ice spatial distribution is very similar between the models and setups with different resolutions. There is only very minor difference in the exact location of sea-ice outer edges in the sub-Arctic seas (Figures 28 and 29). Wintertime agreement is particularly good, with only low-resolution FSU-HYCOM deviating from the other models. In summer, all the models can reproduce the observed spatial pattern with high sea-ice concentration north of Greenland and low concentration towards the Arctic marginal seas. However, the difference between the models is more significant than in winter, especially in the shelf regions, which is true in both the low- and high-resolution experiments. The sea-ice concentration in summer also does not change much with changing resolution, except for IAP-LICOM, which has higher concentration in most parts of the Arctic Ocean in its high-resolution setup.

In both seasons, the northern hemisphere sea-ice thickness is the largest north of the Greenland and decreases towards the Eurasian coast. This pattern is reproduced in the models for both resolutions, except for the high-resolution setup of IAP-LICOM, which has its thickest ice centered around the North Pole (Figures 28 and 29). The sea-ice thickness is underestimated for both seasons in FSU-HYCOM and NCAR-POP, especially in summer, while it is overestimated for the winter season in the East Siberian Sea sector in AWI-FESOM. There is some sensitivity to model resolution for the simulated sea-ice thickness, but the difference due to changes in the resolution is smaller than the difference between the model results and the observation.

As the sea-ice concentration is close to 100% in winter inside the Arctic Ocean, its variance (Figures S1-S8) is very small in this season, as consistently shown by all the model simulations. In summer, the variance of sea-ice concentration is high along the outer sea-ice edge. As the summer sea-ice edges in FSU-HYCOM and NCAR-POP are farther from the coast than in AWI-FESOM, the bands of high variance are located at larger distance from the coast in the former two models. The variance does not change much with changing resolution in the models, except for IAP-LICOM, because its sea-ice concentration is quite different when different resolutions are used. The variance of northern hemisphere sea-ice thickness is stronger along the coastlines in the other three models. Along the Canadian


coastline, the thickness variance is relatively high for both seasons in these models. In winter, the thickness variance is also high along the East Siberian Sea coast in FSU-HYCOM and NCAR-POP. In summer, the thickness variance is high over the Eurasian continental shelves in AWI-FESOM, but not in the other models, because these models have too thin summer ice compared to the observation in these regions. Similar to the finding for the mean sea-ice concentration and thickness, their variance is not very sensitive to the model resolution and the difference between models is more pronounced (Figures S1-S8).


In the southern hemisphere (Figures 30 and 31), the observed spatial pattern of sea-ice concentration in winter is well represented by the models. Only FSU-HYCOM has large difference in winter sea-ice concentration in the southern hemisphere when changing the model resolution. The concentration is too low compared to the observation in its low-resolution setup, and it is more realistic when the resolution is increased. In summer, the sea ice concentration in the southern hemisphere is not well reproduced in most models, except for AWI-FESOM. The high summer sea-ice concentration in the Weddell and Ross Seas is only reasonably simulated in AWI-FESOM. The high-resolution setup of IAP-LICOM has too high concentration in a too wide band. The high variability of Antarctic sea-ice by region that obscures the overall Antarctic sea ice change is apparent in the high variance of the models in summertime Antarctic, but there is little agreement about pattern. The low-resolution, high-resolution model pairs tend to agree more than the ensemble of low-resolution or ensemble of high-resolution models, indicating that sea ice model formulation differences may exceed the effects of resolution in these patterns.

Sea ice is thicker along the Antarctic coast and thinner at its outer edge at the north. The models can reasonably capture this spatial pattern (Figures 30 and 31). In winter, the sea-ice is too thin in the low-resolution setup of FSU-HYCOM, which is improved in its high-resolution setup. NCAR-POP and AWI-FESOM has much less sensitivity to model resolution for the Antarctic sea ice thickness than in the other models. In summer, sea-ice thickness in the Southern Ocean is significantly underestimated in most areas by the models. It is closer to the observation in AWI-FESOM, although it is thinner than the observation in most regions. The sea-ice thickness in the high-resolution setup of IAP-LICOM is significantly overestimated in both seasons, which is also manifested in the time series of sea ice volume (Figure 13).


The variance (Figures S1-S8) of winter sea ice concentration in the southern hemisphere shows a circumpolar band of high values, located along the outer sea ice edge. In FSU-HYCOM, there is high variance in the southeastern Weddell Sea, where the mean winter sea ice concentration is too low. This implies formation of large polynya in this region in some years. This feature is eliminated in the high-resolution FSU-HYCOM setup. In summer, the sea-ice concentration variance is quite different between the models. In particular, the variance is very low in the low-resolution FSU-HYCOM setup because it has much less sea ice than in other simulations. The variance of winter sea-ice thickness in the southern hemisphere is the most pronounced along the sea ice outer edge and the eastern Antarctic Peninsula coast, and in the western Antarctic sector. The models are consistent in the location of large variance, although there are differences in the strength of the variance. In summer, strong variance is present mainly in the Weddell Sea and along the coast of the western Antarctic. The models do show some sensitivity to the model resolution for variance of the sea-ice concentration and thickness in the southern hemisphere, especially in FSU-HYCOM.


However, the difference between models is even larger. As IAP-LICOM significantly overestimated the sea-ice thickness in its high-resolution setup, the variance is also much higher than in other models (Figures S1-S8).











<b>Page 60: [30] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [30] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [30] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [30] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [30] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [30] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [31] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [31] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [31] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [32] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [32] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [32] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [32] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [33] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 2:56:00 PM</b>
Relative		
<b>Page 60: [33] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 2:56:00 PM</b>
Relative		
<b>Page 60: [34] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [34] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		

<b>Page 60: [34] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [34] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [34] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [34] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [34] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [35] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [35] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [35] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [36] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [36] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [36] Formatted</b>	<b>Eric Chassignet</b>	<b>6/30/2020 4:16:00 PM</b>
English (United States)		
<b>Page 60: [37] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 2:56:00 PM</b>
Relative		
<b>Page 60: [37] Deleted</b>	<b>Eric Chassignet</b>	<b>6/18/2020 2:56:00 PM</b>
Relative		