



Trends of inorganic and organic aerosols and precursor gases in Europe: insights from the EURODELTA multi-model experiment over the 1990–2010 period

- 5 Giancarlo Ciarelli^{1,2,a,*}, Mark R. Theobald³, Marta G. Vivanco³, Matthias Beekmann¹, Wenche Aas⁴, Camilla Andersson⁵, Robert Bergström^{5,6}, Astrid Manders-Groot⁷, Florian Couvidat², Mihaela Mircea⁸, Svetlana Tsyro⁹, Hilde Fagerli⁹, Kathleen Mar¹⁰, Valentin Raffort¹¹, Yelva Roustan¹¹, Maria-Teresa Pay¹², Martijn Schaap⁷, Richard Kranenburg⁷, Mario Adani⁸, Gino Briganti⁸, Andrea Cappelletti⁸, Massimo D'Isidoro⁸, Cornelis Cuvelier¹³, Arineh Cholakian^{1,2}, Bertrand Bessagnet^{2,b},
10 Peter Wind^{9,14} and Augustin Colette²

¹Laboratoire Inter-Universitaire des Systèmes Atmosphériques (LISA), UMR CNRS 7583, Université Paris Est Créteil et Université Paris Diderot, Institut Pierre Simon Laplace, Créteil, France

15 ²National Institute for Industrial Environment and Risks (INERIS), Parc Technologique ALATA, 60550 Verneuil-en-Halatte, France

³CIEMAT, Research Centre for Energy, Environment and Technology, Madrid, Spain

⁴Norwegian Institute for Air Research (NILU), Box 100, 2027 Kjeller, Norway

⁵Swedish Meteorological and Hydrological Institute, 60176 Norrköping, Sweden

20 ⁶Department of Space, Earth and Environment, Chalmers University of Technology, 41296 Gothenburg, Sweden

⁷Netherlands Organisation for applied scientific research (TNO), P.O. Box 80015, 3508 TA Utrecht, the Netherlands

⁸Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Via Martiri di Monte Sole 4, 40129 Bologna, Italy

⁹Climate Modelling and Air Pollution Division, Research and Development Department, Norwegian Meteorological Institute (MET Norway), Blindern, 0313 Oslo, Norway

25 ¹⁰Institute for Advanced Sustainability Studies (IASS), Postdam, Germany

¹¹CEREA, Joint Laboratory Ecole des Ponts ParisTech – EDF R&D, Champs-Sur-Marne, France

¹²Barcelona Supercomputing Center, Centro Nacional de Supercomputación, Jordi Girona, 29, 08034 Barcelona, Spain

¹³ex European Commission - JRC, Ispra, Italy

¹⁴Faculty of Science and Technology, University of Tromsø, Tromsø, Norway

30 ^aNow at: Department of Chemical Engineering, Carnegie Mellon University, Pittsburgh, USA

^bNow at: Hangzhou Futuris Environmental Technology Co. Ltd, Zhejiang Overseas High-Level Talent Innovation Park, No. 998 WenYi Road, 311121, Hangzhou, Zhejiang, China

*Corresponding author: Giancarlo Ciarelli (Giancarlo.Ciarelli@lisa.u-pec.fr)



Abstract. In the framework of the EURODELTA-Trends (EDT) modeling experiment, several chemical transport models (CTMs) were applied for the 1990–2010 period to investigate air quality changes in Europe as well as the capability of the models to reproduce observed long-term air quality trends. Five CTMs have provided modeled air quality data for twenty-one continuous years in Europe using emission scenarios prepared by IIASA/GAINS and corresponding year-by-year meteorology derived from ERA-interim global reanalysis. For this study, long-term observations of particle sulfate (SO_4^{2-}), total nitrate (TNO_3), total ammonium (TNH_x) as well as sulfur dioxide (SO_2) and nitrogen dioxide (NO_2) for multiple sites in Europe were used to validate the model results. The trends analysis was performed for the full twenty-one years (referred to as PT), but also for two 11-year sub-periods: 1990–2000 (referred to as P1) and 2000–2010 (referred to as P2).

The experiment revealed that the models were able to reproduce the faster decline in observed SO_2 concentrations during the first decade, i.e. 1990–2000, with a 64–76% mean relative reduction in SO_2 concentrations indicated by the EDT experiment (range of all the models) versus an 82% mean relative reduction in observed concentrations. During the second decade, P2, the models estimated a mean relative reduction in SO_2 concentrations of about 34–54%, which was also in line with that observed (47%). Comparisons of observed and modeled NO_2 trends revealed a mean relative decrease of 25% and between 19–23% (range of all the models) during the P1 period, and 12% and between 22–26% (range of all the models) during the P2 period, respectively.

Comparisons of observed and modeled trends in SO_4^{2-} concentrations during the P1 period indicated that the models were able to reproduce the observed trends at most of the sites, with a 42–54% mean relative reduction indicated by the EDT experiment (range of all models) versus a 57% mean relative reduction in observed concentrations, and with good performances also during the P2 and PT periods. Moreover, especially during the P1 period, both modeled and observational data indicated smaller reductions in SO_4^{2-} concentrations compared with its gas-phase precursor (i.e. SO_2), which could be mainly attributed to increased oxidant levels and pH-dependent cloud chemistry.

An analysis of the trends in TNO_3 concentrations indicated a 28–39% and 29% mean relative reduction in TNO_3 concentrations for the full period for model data (range of all the models) and observations, respectively. Further analysis of the trends in modeled HNO_3 and particle nitrate (NO_3^-) concentrations revealed that the relative reduction in HNO_3 was larger than that for NO_3^- during the P1 period, which was mainly attributed to an increased availability of “free-ammonia”. By contrast, trends in modeled HNO_3 and NO_3^- concentrations were more comparable during the P2 period. Also, trends of TNH_x concentrations were, in general, under-predicted by all models, with worst performance for the P1 period than for P2.

Trends in modeled anthropogenic and biogenic secondary organic aerosol (ASOA and BSOA) concentrations together with the trends in available emissions of biogenic volatile organic compounds (BVOCs) were also investigated. A strong decrease in ASOA was indicated by all the models, following the reduction in anthropogenic NMVOCs precursors. Biogenic emission data provided by the modeling teams indicated a few areas with statistically significant increase in isoprene emission and monoterpene emissions during the 1990–2010 period over Fennoscandia and Eastern European regions (i.e. around 14–27%), which was mainly attributed to the increase of surface temperature. However, the modeled BSOA concentrations did not linearly follow the increase in biogenic emissions. Finally, a comprehensive evaluation against positive matrix factorization (PMF) data, available during the second period (P2) at various European sites, revealed a systematic under-estimation of the modeled SOA fractions of between a factor of 3 to 11, on average, most likely because of missing SOA precursors and formation pathways, with reduced biases for the models that accounted for chemical aging of semi-volatile SOA components in the atmosphere.



1. Introduction

Particulate matter (PM) is one of the greatest environmental concerns, affecting climate and visibility, and having deleterious effects on human health (Cohen et al., 2017; Pope and Dockery, 2006; WHO, 2013). Although particulate matter can be directly emitted from different sources, e.g. power plants, industry and transport, PM with an aerodynamic diameter below $2.5 \mu\text{m}$ ($\text{PM}_{2.5}$) is mainly of secondary origin (Crippa et al., 2014), i.e. formed in the atmosphere after various reactions involving gas-phase precursors such as nitrogen oxide (NO_2), sulfur dioxide (SO_2), ammonia (NH_3), volatile organic compounds (VOCs) and several oxidants (e.g. OH, O_3 and NO_3). Particles in this size range can penetrate deeply into the respiratory system leading to respiratory and cardio-vascular problems. The formation mechanisms leading to the formation of secondary aerosols, especially the organic fraction, are complex, non-linear and still not fully understood (Bian et al., 2017; Lachatre et al., 2018; Tsigaridis et al., 2014).

Emissions of SO_2 and NO_2 have largely declined in Europe over the recent decades (Fagerli and Aas, 2008; Tørseth et al., 2012; UNECE LRTAP, 2016). For SO_2 and NO_2 , emissions were reported to have declined by about 65% and 31%, respectively, between 1990 and 2009 whereas emissions of NH_3 were reported to have declined by about 29%, although the emission changes exhibit high spatial variability within the European domain (Tørseth et al., 2012). NH_3 , which is emitted mainly from agricultural activities, is one of the key chemical species involved in the formation of secondary inorganic aerosol. It is the most important base in the atmosphere (Seinfeld and Pandis, 2012) and can react very rapidly with sulfuric acid (H_2SO_4), which is formed from the oxidation of SO_2 with OH (in the gas phase), O_3 and hydrogen peroxide (H_2O_2) in the aqueous phase, to form ammonium sulfate or ammonium bisulfate (Seinfeld and Pandis, 2012). If enough NH_3 is available after the neutralization of H_2SO_4 , it can react with nitric acid (HNO_3), which is mainly formed from the oxidation of NO and NO_2 , to produce the semi-volatile ammonium nitrate. Formation of ammonium nitrate usually occurs when the molar concentration of $\text{NH}_3 + \text{NH}_4^+$ is more than twice the sulfate concentration (i.e. “free ammonia regime”) (Tsimpidi et al., 2007).

Past and future trends in the total PM concentration have recently obtained large attention thanks to the availability of long-term observational data-sets and increased computational power available for long-term CTM simulations. Tørseth et al. (2012) analyzed long-term air quality trends from the European Monitoring and Evaluation Programme (EMEP) during a period of 40 years. Their study showed a substantial reduction in ambient concentrations of sulfur species of about 70-90% since 1980, well in line with emission reductions, and a reduction of about 23% in NO_2 concentrations since the beginning of the 1990s. However, available observations of total nitrate ($\text{TNO}_3 = \text{HNO}_3(\text{g}) + \text{NO}_3^-(\text{p})$) showed only a minor reduction (about 8%), compared to the larger reductions in NO_2 . Aksoyoglu et al. (2014) performed an air quality modeling study with the Comprehensive Air Quality with extension (CAMx) model to evaluate air quality changes due to anthropogenic emission changes in the framework of the revised Gothenburg protocol. They performed air quality simulations for the emission years 1990, 2005 and 2020 with emission scenarios prepared from IIASA/GAINS. Their results indicated that the annual mean $\text{PM}_{2.5}$ concentration in Europe decreased by 20-50% between 1990 and 2005. Moreover, simulated annual mean $\text{PM}_{2.5}$ concentrations were 30% lower in 2020 compared with 2005, with larger decreases for eastern European countries (Aksoyoglu et al., 2014). Similarly, Colette et al. (2011) investigated the capability of six regional and global CTMs for simulating air quality changes between 1998 and 2007 with a focus on NO_2 , O_3 and PM_{10} . Their results indicated that the models could reproduce the trends of primary pollutants, but they had difficulties in reproducing the small observed trends in O_3 , and the year-to-year variability was under-estimated, in general. More recently, Banzhaf et al., 2015 applied the LOTO-EUROS model for the 1990–2009 period to investigate trends of air quality in Europe. They concluded that the model was able to well reproduce the observed trends in primary and secondary produced pollutants. In addition, they also performed a source apportionment study to evaluate the formation efficiency of secondary inorganic species during the 1990–2009 period. Their results



indicated an increase in SO_4^{2-} formation efficiency (between 20–50%) as well as for NO_3^- (up to 20%) compared with 1990.

Organic aerosol (OA) is often a major fraction of $\text{PM}_{2.5}$. OA is a complex mixture of thousands of organic compounds with different chemical and physical properties and volatilities (semi-volatile to low volatile) (Huang et al., 2014; Jimenez et al., 2009). Numerous measurement campaigns performed in different parts of the world and periods of the year, have revealed that only a minor fraction of the observed total OA mass is directly emitted as primary organic aerosol (POA). A more abundant component, referred to as secondary organic aerosol (SOA), was found to often dominate the composition of OA especially in rural areas (Crippa et al., 2014).

The formation of SOA in the atmosphere is mainly initiated by the oxidation of gas-phase organic compounds in different ranges of saturation concentrations, usually referred to as low-volatile, semi-volatile, intermediate-volatile and high-volatile ranges (Donahue et al., 2012, 2011). Some of the resulting gas-phase oxidation products will acquire lower saturation concentration due to the addition of oxygen-containing functional groups and will eventually condense on pre-existing organic particles leading to formation of SOA (depending on temperature and OA concentrations). On the other hand, other organic compounds will obtain lower molecular weight and will fall into higher saturation concentration ranges through fragmentation, and they will likely reside in the gas-phase.

A recent model intercomparison exercise, AeroCom (Tsigaridis et al., 2014), investigated the performance of thirty-one global models with respect to OA revealing large differences between models in terms of SOA formation, mainly because of the assumptions made in the SOA scheme used (e.g. chemical aging, multiphase chemistry and semi-volatile SOA assumptions). In addition, comparison with several observational data-sets revealed that even though the models were able to simulate the secondary nature of OA, they tended to largely underestimate the observed OA, especially in urban areas (Tsigaridis et al., 2014). In Europe, recent applications of CTMs have started to provide a comprehensive picture of the main sources of OAs as well as their temporal variation throughout the year. Bergström et al., 2012 applied the EMEP MSC-W model with the volatility basis set (VBS) model and tested different assumptions on the volatility distribution of POA as well as on the parameterizations of the aging processes. Their studies revealed an underestimation of OA concentrations, especially during winter periods and in northern European countries, most likely as a result of uncertainties in the emissions from the residential sector (mainly wood burning emissions). Summer-time OA levels, on the other hand, were highly influenced by biogenic SOA precursors (isoprene and terpene), also confirmed by more recent studies (Cholakian et al., 2017; Chrit et al., 2017; Ciarelli et al., 2016).

In this study, we investigate the trends in SIA and SOA in Europe during the 1990–2010 period calculated by five CTMs that participated in the EURODELTA-Trends exercise (Colette et al., 2017). The novel multi-model EURODELTA-Trends (EDT) exercise (launched within the Task Force on Measurement and Modelling of the EMEP Programme supporting the Convention on Long Range Transboundary Air pollution (CLRTAP)), provided 21-year of continuous $\text{PM}_{2.5}$ components and their gas-phase precursors concentrations over Europe from the year 1990, and with “real” year-to-year meteorological input data. It provides a base for validating the performance of multiple models over an extended period (i.e. 1990–2010) and to assess the variation of various chemical species not routinely measured in Europe.

The paper is organized as follows: Section 2 provides a general overview of the EURODELTA-Trends experiment, with a description of the models participating in the exercise and the input data used to perform the experiment. The observational data is described in Section 2 along with information regarding the quality control criteria. Results and discussions are presented in section 3. The trends in anthropogenic emissions and inorganic species are discussed in section 3.1 and section 3.2, respectively. An evaluation of the secondary organic aerosol fraction is presented in section 3.3 (for the 2000–2010 period) together with the trends in biogenic emission and anthropogenic and biogenic SOA



concentrations. Finally, conclusions are presented in Section 4.

2. Methods

2.1 Overview of the EURODELTA-Trends experiment

5 The EURODELTA-Trends experiment builds upon the expertise of the previous EURODELTA phases initiated in 2004 (van Loon et al., 2007). In the latest EURODELTA experiments, i.e. EURODELTA III, the performances of several CTMs were investigated for common air quality pollutants, i.e. NO₂, O₃, SO₂, and PM₁₀ and PM_{2.5} at a European scale for specific periods of the EMEP and EUCAARI intensive measurements campaigns (Bessagnet et al., 2016).

10 The follow-up EURODELTA experiments, referred to as EURODELTA-Trends (EDT), aim at investigating the changes in air quality in Europe over the 1990–2010 period. In this framework, state-of-the-art CTMs, were applied over the European domain (Figure 1) with common input data (meteorological fields, anthropogenic emissions and boundary conditions). The participating models carried out extensive sensitivity tests that aimed at disentangling the role of different drivers (e.g. meteorology and emissions) on changes in air quality. The complete list of data available, chemical species and sensitivity tests are reported in detail in Colette et al., 2017.

15 In this study, one tier of simulations was used to investigate the models' capabilities to reproduce gas-phase PM precursors as well as SIA trends over the 1990–2010 period. This tier, referred to as tier 3A, provides twenty-one years of modeled air quality data in Europe driven with “real” meteorology, observation-based boundary conditions and anthropogenic emission scenarios based on the IIASA/GAINS model. Biogenic emissions were calculated separately by the different modeling teams using their own biogenic model driven by the meteorological data (e.g. temperature and radiation).

2.2 Description of the participating models

25 Eight state-of-the-art air quality CTMs delivered their results for the EDT experiments: CHIMERE (Mailler et al., 2017; Menut et al., 2013), CMAQ (Byun and Schere, 2006), EMEP MSC-W (Simpson et al., 2012), LOTOS-EUROS (Manders et al., 2017; Schaap et al., 2008), MATCH (Andersson et al., 2015, 2007; Robertson et al., 1999), MINNI (Mircea, 2016), Polyphemus (Mallet et al., 2007; Sartelet et al., 2012) and WRF-Chem (Grell et al., 2005; Mar et al., 2016). Given the large computational demand of the simulations, only five modeling teams were able to deliver 21 years of continuous air modeled data: CHIMERE, EMEP MSC-W, LOTOS-EUROS, MATCH and MINNI, the results of which are used in this study. Most of the other models provided air quality data for 3 intermediate years: 1990, 2000 and 2010.

The set-up for each participating model is reported in Table 1. The complete list of physical and chemical schemes (including dry and wet deposition parameterizations) can be found in (Colette et al., 2017). The models differ in terms of the adopted gas-phase chemistry mechanisms as well as SIA and SOA formation modules. Here, we briefly describe the main characteristics of the various schemes used by the models.

35 Various gas-phase schemes were used to perform the gas-phase chemistry (Table 1): The Carbon Bond mechanism version 4 (referred to as TNO-CBM-IV), EmChem09, MELCHIOR2 and SAPRC99.

40 - The TNO-CBM-IV gas-phase scheme (Schaap et al., 2009), used by the LOTOS-EUROS model, includes 33 gas-phase species and 9 organic species emitted directly into the atmosphere. Most of the included organic species are lumped according to the carbon-carbon bond type and only a minority of them are explicitly represented (e.g. isoprene and formaldehyde). 104 chemical reactions and 14 photolytic reactions are mapped



in the TNO-CBM-IV mechanism for gas-phase chemistry.

- 5 - The EmChem09 gas-phase scheme (Simpson et al., 2012), used by EMEP MSC-W and MATCH models, include 72 species, 137 chemical reactions and 26 photochemical reactions. The rates and products were designed to be as close as possible to the IUPAC recommendations (<http://www.iupac-kinetic.ch.cam.ac.uk/>) and most of the reaction coefficients were taken from Atkinson et al., 2006, 2004. The MATCH model used a modified version of isoprene chemistry based on the work of Carter, 1996 and Langner et al., 1998.
- 10 - The MELCHIOR2 gas-phase scheme (Derognat et al., 2003), used by the CHIMERE model, is a reduced version of the MELCHIOR1 mechanism and it includes 120 chemical reactions and hydrocarbon degradation as in the EMEP gas-phase mechanism, with a few adaptations included for low-NO_x conditions and NO_x-nitrate chemistry. All rate constants are taken from Atkinson et al., 1997 and De Moore et al., 1994.
- 15 - The SAPRC99 gas-phase scheme (Carter, 2000), used by the MINNI model, includes a detailed speciation of about 400 types of VOCs and with detailed reaction schemes for most of the non-aromatic hydrocarbons and oxygenates in the presence of NO_x. The isoprene photooxidation is explicitly included, the “four-product” condensed isoprene mechanism considers methacrolein, methyl vinyl ketone, lumped C5 unsaturated aldehyde products (ISOPROD), and the methacrolein PAN analogue (MPAN).
- 20 To resolve the composition and phase state of inorganic aerosol, most of the models used the ISORROPIAv2.1 scheme (version 1.7 for the MINNI model and version II for LOTOS-EUROS) which assumes thermodynamic equilibrium with its gas phase precursors (Nenes et al., 1999, 1998). The EMEP MSC-W model adopted the approach proposed by Binkowski and Shankar, 1995, i.e. the MARS equilibrium module, and does not include sodium chloride and dust components, whereas the MATCH model is based on the work of Mozurkewich, 1993. Transformation of HNO₃ to coarse nitrate is included by all the models except MINNI.
- 25 As already mentioned, NH₃ is a key ingredient for the formation of secondary inorganic aerosols. NH₃ compensation points are included in LOTOS-EUROS to account for the presence of NH₃ in the stomata, external leaf surfaces or at the soil surface and partially included in the EMEP MSC-W model by assuming zero NH₃ dry deposition to growing crops.
- 30 Different gas-phase and thermodynamic organic aerosol schemes with various levels of complexity were used by the modeling teams (Table 1): The volatility basis set with and without aging of SOA (Bergström et al., 2012a; Simpson et al., 2012), referred to as VBS-NPAS and VBS-NPNA, respectively, the H₂O mechanism (Couvidat et al., 2012) coupled with the SOAP module (Couvidat and Sartelet, 2015), and the SORGAM mechanism (Schell et al., 2001). None of the models included emission of semi-volatile organic compounds (SVOCs) and/or of intermediate-volatile organic
- 35 compounds (IVOCs). LOTOS-EUROS did not enable any SOA scheme and therefore the organic model description is not included here.
- 40 - The VBS-NPAS and VBS-NPNA organic aerosol modules, used by the EMEP MSC-W and MATCH models respectively, assumes primary organic aerosol (POA) emission to be non-volatile, assuming European emission inventories to consist of inert PM compounds. Semi-volatile SOA is formed from oxidation of anthropogenic and biogenic volatile organic compounds (VOC) (for details regarding the volatility basis set and SOA-yields see Bergström et al., 2012). In the EMEP (VBS-NPAS) model the OH-reaction rate for SOA



aging is set to $4.0 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$; each reaction of the organic compounds in the gas-phase decreases the volatility by one order of magnitude and increases the mass by +7.5 % to account for oxygen-addition (fragmentation processes are not included). SOA aging is not included in the VBS-NPNA scheme.

- 5 - The H₂O organic aerosol module (Couvidat et al., 2012), used by the CHIMERE model, uses different types of surrogate organic species: hydrophilic species (which condense preferentially into an aqueous phase) and hydrophobic species (which condense only into an organic phase). These surrogate species are produced from the oxidation of volatile organic compounds. In H₂O, SOA are formed from 4 classes of precursors: aromatic compounds, isoprene, monoterpenes and sesquiterpenes. For aromatic compounds, toluene and xylene are used
- 10 as SOA precursors when reacting with the OH radical and without accounting for SOA aging. The H₂O mechanism accounts for the effect of nitrogen oxides on SOA formation as well as the dissociation of organic acids in an aqueous phase, the oligomerization of aldehydes. More details of the scheme can be found in Couvidat et al., 2018, 2012.
- 15 - The SORGAM mechanism (Schell et al., 2001), used by the MINNI modelling system, includes 4 SOA precursors classes (alkanes, alkenes, aromatics and monoterpenes) to represent the contributions of anthropogenic precursors and biogenic precursors to SOA formation. The Volatile Organic Compounds (VOCs), are oxidized by reactions with hydroxyl radical (OH), ozone (O₃), and nitrate radical (NO₃). The anthropogenic SOA are formed from aromatics like toluene, xylene, cresol, from internal alkenes and long
- 20 “alkanes” as those grouped together in the ALK5 and OLE2 classes, respectively, in SAPRC99 gas-phase mechanism. Biogenic SOA is produced only by monoterpenes which partitioning parameters are obtained from a weighted average of smog chamber experiments for α -pinene, β -pinene, d₃-carene, sabinene, limonene.

2.3 Emissions

25

2.3.1 Biogenic and natural emissions

Emissions of biogenic volatile organic compounds (BVOCs) were not prescribed by the EDT experiments. Each participating team used their own emission model to calculate biogenic emissions.

- 30 One group of models used the MEGANv2.04 (Guenther et al., 2006) and MEGANv2.1 (Guenther et al., 2012) emission models: CHIMERE and MINNI, respectively. CHIMERE uses highly resolved spatiotemporal data (30 arcsec every 8 days) generated from MODIS for Leaf area index (LAI) inputs. The 30 arcsec USGS (US Geophysical Survey) land-use database is used to provide information on the plant functional type (PFT). The PFT is then combined with the emission factors for each functional type of Guenther et al. (2012) to compute the landscape average emission factors. MINNI
- 35 derived them from the CORINE Land Cover (CLC2006) inventory. The MEGAN model is driven with meteorological variables, such as temperature, wind speed, humidity, solar radiation and soil moisture. The leaf area index retrieved from the TERRA/MODIS satellite is used to simulate the vegetation growth (eight-day and one-month average LAI data at $0.25^\circ \times 0.25^\circ$ degrees resolution for CHIMERE and MINNI, respectively). Common BVOCs species such as isoprene, α -pinene as well as other classes of monoterpenes, are generated for each hour and grid cell of the domain. In
- 40 the CHIMERE model, emissions of sesquiterpenes are also included and used as an input for SOA chemistry. More information on the MEGAN emission algorithms can be found in Guenther et al., 2006, 2012.

The second group of models: LOTOS-EUROS, MATCH, EMEP MSC-W used a detailed tree inventory of 115 species for 30 European countries based on the work of Koeble and Seufert, 2001 and aggregated tree species based on land-



cover types. For this group of models, the environmental factors to derive biogenic emissions include the light correction factor (γ_L) and the temperature correction function (γ_T), which are applied to three types of emission categories: isoprene, pool-dependent monoterpenes and light-dependent monoterpenes based on (Guenther et al., 1993). More information on the EMEP MSC-W BVOC emission algorithm can be found in (Simpson et al., 2012).

- 5 Finally, sea-salt, emitted in water droplets from the sea during high wind speed conditions and as results of breaking of waves and/or bursting of air bubbles, are included in all the models, based on different schemes, as described in Colette et al., 2017. Windblown dust emission were taken into account by all the models except MATCH, while road traffic dust resuspension was only included in the EMEP MSC-W model (Colette et al., 2017).

2.3.2 Anthropogenic emissions

10 Anthropogenic gridded emissions by country and activity, i.e. SNAP (Selected Nomenclature for reporting of Air Pollutants) codes, were estimated using the Greenhouse gas - Air Pollution Interactions and Synergies (GAINS) model (Amann et al., 2011). Emission of SO_x , NO_x , NH_3 , non-methane volatile organic compounds (NMVOCs) as well as primary $\text{PM}_{2.5}$, PM_{10} , black carbon and primary organic aerosol were prepared at a $0.25^\circ \times 0.40^\circ$ resolution (latitude \times longitude). Anthropogenic emissions were calculated for the years 1990, 1995, 2000, 2005 and 2010, and linearly interpolated by country and activity sector for the 5-years periods to obtain the continuous 21-year emission dataset. Data for the different emission sectors were obtained from Eurostat (<http://ec.europa.eu/eurostat>), the International Energy Agency (IEA, 2012) and the UN food and Agriculture Organization (FAO) (<http://www.fao.org/statistics/en/>). Additionally, data from the International Fertilizer Association (IFA) and the COPERT model (Athanasiadis et al., 2009) were used for the agriculture and transportation sectors, respectively. An error in primary particulate emission matter for Russia, North Africa and maritime areas for the period 1991-1999 was identified at the end of the exercise. However, the effect of the error was estimated to be very limited (Theobald et al., 2019).

25 The complete anthropogenic emission dataset accounts for source-specific emission limits as well as for various European air quality directives (e.g. the UNECE Gothenburg Protocol, UNECE, 1999). This emission dataset, referred to as ECLIPSE_V5, was delivered by IIASA as country national totals by activity sector. It was subsequently spatialized by INERIS on the EURODELTA-Trends grid for use in the CTMs using the gridding process described in Terrenoire et al., 2015 and Bessagnet et al., 2016. For the residential heating sector (SNAP2) a proxy based on population density was applied using a bottom-up inventory available for France. More information about the re-gridding can be found in (Colette et al., 2017).

30

2.3.3 Meteorological data

To provide meteorological inputs to the modeling teams, dynamically downscaled regional climate model simulations was used in combination with ERA-interim global reanalysis data (Dee et al., 2011). The Weather Research and Forecast Model (WRF version 3.3.1; Skamarock et al., 2008) was used at a resolution of 0.44° to generate the meteorological parameters. To reduce the uncertainty of the meteorological data, WRF was re-run with ERA-interim reanalysis data in grid-nudging mode as described in Stegehuis et al., 2015 and subsequently interpolated at a 25 km resolution to match the EDT grid, although there were a few differences between the procedures of the modeling team. LOTOS-EUROS used RACMO2-downscaled data and MATCH used HIRLAM-downscaled data. More information on the meteorological inputs can be found in (Colette et al., 2017).

40



2.3.4 Observational data and trend assessment

The observations are reported to EMEP, and the original time series are available in EBAS (<http://ebas.nilu.no>). The datasets chosen for the trend assessment have passed the completeness criteria of 75% of data available over the full 1990–2010 period and had undergone visual screening tests. The secondary dataset with annual and seasonal average concentrations is available from the webpage set up by Task Force on Measurements and Modeling (TFMM) for this study (<https://wiki.met.no/emep/emep-experts/tfmmtrendstations>). The datasets include yearly measurements of long-term air concentrations of sulfur dioxide (SO₂), particle sulfate (SO₄²⁻), nitrogen dioxide (NO₂), total nitrate (TNO₃ = HNO₃(g) + NO₃(p)) and total ammonium (TNH_x = NH₃(g) + NH₄⁺(p)) performed in Europe between 1990 and 2010. Overall, the numbers of observational sites available for each of the species are 30, 20, 25, 13 and 16 for SO₂, SO₄²⁻, NO₂, TNO₃ and TNH_x, respectively. Figure 2 illustrate the geographical distribution of the observational sites for each of the species, all classified as rural background stations. It can be noted that most of the stations are located over the Northern and Central part of the domain. The complete list of the observational sites is reported in Table S1.

NO₂ is mainly sampled with the manual method where NO₂ is selectively absorbed on impregnated glass sinters. Some sites do however use chemiluminescence monitor with molybdenum converter, which is not selective for NO₂, thus these measurements might be biased, and this is especially important in areas with low concentrations (Reed et al., 2016), but it is not assumed that the trends will be largely affected when same method is used during the whole period. The other components are mostly measured using a filterpack sampler with no size cut off in the inlet. The 3-stage filterpack separates gas and aerosol species, but for nitrogen compounds this separation might be biased due to the volatile nature of NH₄NO₃. Therefore TNO₃ and TNH_x are usually used for robust estimate of the atmospheric nitrogen loading (Tørseth et al., 2012). However, it is recommended to report the measurements of all the species since it may give valuable insight into the gas/particle ratio despite possible biases. Details of the method used are found in the annual data report (i.e. EMEP, 2012 for the 2010 data).

The linear trends for each species and observational site were calculated with the Theil-Sen method (Sen, 1968) and their significances were evaluated at 95% confidence level ($p < 0.05$) using the non-parametric Mann-Kendall test (Kendall, 1948; Mann, 1945). Trends in observational data were compared with trends in modeled data calculated with the same methodology. Since anthropogenic emissions did not decline linearly during the full period covered by the experiment (1990–2010, referred to as PT), and larger emissions reduction are expected during the earlier 90s, the trend analysis was performed for two sub-periods: a first period between 1990 and 2000, referred to as P1, and a second period between 2000 and 2010, referred as to P2. In addition, to provide a more comprehensive picture of the trends in the air pollutant concentrations, the trends analysis was also performed for several sub-regions adapted from the commonly used PRUDENCE climatic zones classification (<http://ensemblesrt3.dmi.dk/quicklook/regions.html>). The extension of the sub-regions used in the study is reported in Figure 1.

The evaluation of modeled SOA was performed using an extensive data-set of secondary organic aerosol concentrations retrieved with Positive Matrix Factorization (PMF) analysis (Paatero, 1999) and recently compiled by Tsimpidi et al., 2016. This data-set includes SOA average concentrations at 51 stations in Europe during the P2 period. In order to remove local pollution events, likely not included in emission inventories, stations with average SOA concentrations higher than 7 µg m⁻³ during the measurement period were excluded from the analysis (i.e. 3 stations). Most of the measurements were performed during short campaigns using aerosol mass spectrometers (AMS) in different periods of the years, lasting from about two weeks to one month. The spatial distribution of the stations is presented in Figure S1. The complete list of stations used is reported in Table S2 along with information regarding the year and the seasons during which measurements were made.



3. Results and Discussion

3.1 Trends in anthropogenic emissions

5 Table 2 reports the absolute and relative trends in SO_x, NO_x, NH₃ and NMVOCs emissions for the full 1990–2010 period as well as for the P1 and P2 periods over the entire domain. For the full period, SO_x emissions show a decline of about 69%. SO_x emissions declined faster during the P1 period compared with the P2 with decreases of 54 and 37%, respectively (Table 2). The large reduction in SO_x emissions was largely attributed to emission reductions in the “combustion in energy and transformation industries” sector, largely achieved by the switch to low-sulfur containing
10 fuels (e.g. natural gas) and the adoption of desulphurization technologies in large industries. NO_x emissions were reduced by 25% during the P1 period and by 17% during the P2 period. These reductions were mainly achieved through emission reductions in the road transport sector following the introduction of the new EURO standards for passenger cars. However, in 2010 this sector still represented the most important source of anthropogenic NO_x emissions in Europe (EEA, 2012). Important NO_x emission reductions were also achieved thanks to the adoption
15 of low-NO_x burners and selective and non-selective catalytic reduction measures for the “combustion in energy and transformation industries” sector. NH₃ emissions declined only a little compared to SO_x and NO_x emissions. NH₃ emission mainly arises from agricultural activities, which had less stringent controls compared to SO_x and NO_x emission ceilings. NH₃ emissions declined by 19% over the P1 period but only by 6% over the P2 period.

20 Emissions of NMVOCs showed a decline of 59% over the full 1990–2010 period with similar relative reductions achieved during the P1 and P2 periods: 33% per period. NMVOC emission reductions were mainly driven by the road transport sector, and by the year 2010, most of the NMVOC emissions arise from the use of solvents (EEA, 2012). Huang et al., 2017 compiled a global gridded data set of speciated NMVOC emissions for the 1970–2010 period and analyzed the trends. Among the different world regions, North America and Europe were reported to have reduced their
25 NMVOC emissions since 1970 due to the introduction of EURO emission standards for vehicles. A significant reduction of formaldehyde emissions was reported in 2010 compared with 2000, mainly because of the increasing adoption of EURO standards and the transition from coal to cleaner fuels (e.g. natural gas). The latter resulted in a substantial decrease in the aromatic species and in an increase in the contribution of alkanes and alkanals to the emissions of NMVOCs (Huang et al., 2017).

30



3.2 Trends in inorganic species

3.2.1 Comparison of modeled and observed SO₂ and NO₂ concentrations trends

- 5 Figure 3 and Table 3 report the mean relative trends of all the sites included in the analysis (Table S1). Overall, the observations indicate relative reductions of 25, 12 and 36% in NO₂ concentrations for the P1, P2 and PT, respectively, with the models estimating similar ranges of relative reductions, i.e. 19–23% for the P1 periods, 22–26% for the P2 period and 44–47% for the full period, depending on the model (Table 3). Only about half of the individual observed trends were reproduced within a factor of two by individual models and all models performed worse in the
- 10 second period (P2), overestimating the observed trends (Figure 4 and Table 3). Such behavior could indicate possible difficulties for CTMs in capturing long-term trends at relatively low concentrations with small annual changes, typical of the P2 period, it also could point to overestimated negative trends in national emission data bases.
- For SO₂, the observed relative reductions were 82, 47 and 97% for the P1, P2 and full periods, respectively (Figure 3 and Table 3). The models indicate very similar ranges of SO₂ reductions, i.e. 64–76% for the P1 periods, 34–54% for the
- 15 P2 period and between 84–97% for the full period, depending on the model (Table 3). This is in line with the emission reduction trends presented in Section 3.1 and with previous trend studies for Europe (Tørseth et al., 2012). Table 3 also reports the fraction of model estimates within a factor of 2 of the observed trends. Most of the models were able to reproduce the observed SO₂ trends within a factor of two at most of the individual sites (Figure 3 and Table 3). Overall, model performance was better during the P1 period compared with P2.
- 20 Figure 5 shows the percentage of statistically significant/non-significant increasing/decreasing observed and modelled SO₂ and NO₂ trends:
- For SO₂, most of the stations had significant decreasing trends in concentrations during the P1 period, with only a small fraction of the stations with non-significant decreasing trends. All the models were able to reproduce this pattern, albeit with a slight over-estimation of the significant-decreasing fraction. During the P2 periods, there is a tendency of most of
- 25 the models to over-estimate the number of significant trends, with CHIMERE and LOTOS-EUROS being closer to the fraction of significant/non-significant decreasing trends indicated by the observations. As expected, the agreement between the modeled and observed fractions of significant/non-significant increasing/decreasing trends was improved for the full period (PT), mainly because of the larger number of data-points in the time series, with all the sites indicating significant observed and modeled decreasing trends.
- 30 For NO₂, the models were able to reproduce the observed fraction of significant/non-significant increasing/decreasing trends in the P1 period, with most of the models indicating a significant-decrease in NO₂ concentrations at most of the stations (slightly lower for EMEP MSC-W). The analysis for the P2 period shows that the reduced fraction of observed significant-decreasing trends compared with the P1 period, was not well reproduced by the models, all of them tending to over-estimate the fraction of significantly decreasing trends (Figure 5). Again, a possible explanation for the
- 35 degraded model performance during the P2 period could be related to the relatively low pollutant concentrations, which might be challenging to model at such coarse resolution, as well as to uncertainties in the measurements data (see Section 2.3.4). One site in Ireland (IE0001R) was the only site with significant increasing observed trends during the P1 and PT periods, a result which was not reproduced by any of the models.

40 3.2.2 Comparisons of modeled and observed SO₄²⁻, TNO₃ and TNH_x concentration trends

Figure 6 shows the mean modeled and observed relative trends in SO₄²⁻, TNH_x and TNO₃ for all the sites included in the analysis (Table S1). In consistency with the gas-phase analysis, the trends are reported for the two sub-periods, i.e.



1990–2000 (P1) and 2000–2010 (P2), as well as for the full period (PT).

Overall, the observations indicated that concentrations of SO_4^{2-} declined by 57, 14 and 66% for the P1, P2 and PT periods (mean of all the stations, Figure 6 and Table 4), with the models indicating relative reductions of 42–54% for the P1 periods, 23–35% for the P2 period and 61–78% for the full period, depending on the model. The reductions in SO_4^{2-} concentrations were larger during P1 than during the P2 period and most of the model estimates were within a factor of two of the observed values for all the periods (Figure 7 and Table 4). Two sites, one in Ireland (IE0001R) and one in Poland (PL0003R), showed an increase in SO_4^{2-} concentrations (Figure 7), which none of the models were able to reproduce.

The percentage of statistically significant/non-significant increasing/decreasing trends in the observed and modeled SO_4^{2-} trends is reported in Figure 8, showing a good agreement between the observed and modeled significances (and their direction) for the three periods. Statistically significant increasing trends in SO_4^{2-} concentrations were only observed at the PL0003R site in Poland during the P2 period (Figure 8), a result which none of the models were able to reproduce. Interestingly, observed SO_4^{2-} concentrations declined less than those of SO_2 (Table 3), a behavior also reproduced by all the models. The non-linear dependencies between the reduction in SO_2 and SO_4^{2-} concentrations are influenced by different factors. First, the strong reduction in SO_x emissions will increase the availability of OH radicals, which will directly enhance the homogeneous reaction rate of SO_2 . Second, all the models account for the dependence of the aqueous chemistry of SO_2 on pH levels. Thus, heterogeneous reactions of SO_2 are also expected to proceed more efficiently due to the increase of pH levels over time.

Observations of TNH_x reveal that concentrations declined by 28, 22 and 46% for P1, P2 and PT, respectively (Table 4).

In general, most of the models under-predict the relative changes; the modeled relative reductions for the P1, P2 and PT periods were 15–26%, 14–21% and 27–38%, respectively, with the P1 period showing only a minor fraction of the data-points within a factor of two (Figure 7 and Table 4). Indeed, large uncertainties remain in terms of ammonia emissions, which might affect model performance for TNH_x . Moreover, we would like to underline that none of the participating model accounted for the influence of meteorology (e.g. temperature) on ammonia emissions and relied on static emission profiles provided by the EURODELTA exercise. Recent studies, however, have shown that better agreement in terms of the modeled ammonia concentrations can be achieved when ammonia emissions are modulated with local meteorological conditions (Backes et al., 2016; Hendriks et al., 2016). Compared with the other investigated species, a larger variation in terms of the significance of the trends can be seen in Figure 8, with most of the models tending to over-estimate the fraction of significant decreasing trends. Statistically significant increasing trends in observed TNH_x concentrations were found at one station in Norway (NO0039R) for the full period (PT), with none of the models being able to reproduce this feature.

TNO_3 concentrations, on the other hand, declined to a lesser extent than those of SO_4^{2-} and TNH_x . For all periods, the observed relative changes in TNO_3 concentration were 16, 19 and 29%, for the P1, P2 and PT periods, respectively, with the models estimating similar ranges for the P1 and P2 periods, i.e. 16–19%, 8–27% and 28–39%, for the P1, P2 and PT periods, respectively (Figure 6 and Table 4). Most of the model estimates were more than a factor of two larger than the observed values for the P1 and PT periods (Figure 7 and Table 4). The percentage of statistically significant/non-significant increasing/decreasing observed and modeled TNO_3 trends revealed that most of the models were able to reproduce the large fraction of non-significant decreasing observed trends. The EMEP and MATCH models estimate a larger fraction of significant decreasing trends than the other models in both the P1 and P2 periods, where CHIMERE and MINNI show the largest fraction of non-significant decreases. CHIMERE also shows the largest fraction of non-significant increasing trends during the P2 period (Figure 8).



3.2.3 Trends in modeled HNO_3 and NO_3^- concentrations for different sub-regions

In order to further investigate the trends in TNO_3 concentrations described in the previous paragraph, we also investigated the modeled trends in HNO_3 and NO_3^- concentrations (for the different sub-regions in Figure 1).

Figure 9 illustrates the relative trends in HNO_3 and NO_3^- (sum of the coarse and fine particle fractions) for the P1 (first two columns in Figure 9) and P2 (last two columns in Figure 9) periods, for all the models that participated in the experiment. In general, during the P1 period, the models indicate larger significant decreases in HNO_3 compared with NO_3^- , especially over the Fennoscandia and central European regions. A few differences in the spatial distribution of the modeled HNO_3 and NO_3^- trends can be seen in Figure 9. For instance, for the LOTOS-EUROS model, the significant relative trends in both HNO_3 and NO_3^- concentrations are more comparable during the P1 period whereas MINNI and CHIMERE estimates larger areas of non-significant trends in NO_3^- concentrations, with some significant-increases in few parts of the domain. Figure 10 and Table 5 show the models' average relative changes and standard deviation (over land) of HNO_3 and NO_3^- for all the PRUDENCE zones for the P1 and P2 periods. During the P1 period, the regions classified as Alps (AL), British Island (BI), Benelux area (BX), France (FR) and Mid-Europe (ME), had the largest decrease in HNO_3 concentrations, with average decreases between 36 and 44% (Table 5). A comparison of HNO_3 and NO_3^- relative trends for the same regions shows that NO_3^- concentrations declined to a lesser extent, i.e. around 20%, which is roughly half of the modeled relative reduction in HNO_3 concentrations. The largest difference between the relative reduction in HNO_3 and NO_3^- concentrations occurred over the Scandinavian regions for the P1 period, of 24 and 5%, respectively. On the other hand, the reduction of HNO_3 and NO_3^- concentrations were comparable in the Eastern-European region and over the Iberian Peninsula (Table 5), as well as during the P2 periods.

The non-linear response of HNO_3 concentrations and NO_3^- concentrations, i.e. larger relative reduction in HNO_3 compared to NO_3^- , could be attributed to the shift in the thermodynamic equilibrium of HNO_3 versus particle nitrate NO_3^- . In fact, for specific regions and especially during the P1 period, the large reduction in SO_2 emissions increased the availability of “free-ammonia” and thus the transfer of more HNO_3 into the particle phase, favoring the formation of ammonium nitrate. This also increases the TNO_3 lifetime as dry deposition is much more rapid for HNO_3 than for NO_3^- . This effect could contribute to the reduced TNO_3 decreases with respect to the NO_2 decrease. Figure 11 shows the modeled $\text{HNO}_3/\text{NO}_3^-$ and $\text{NH}_3/\text{NH}_4^+$ molar ratio for the P1 and P2 periods, as predicted by all the models. In general, the models indicate significant decreasing trends in the $\text{HNO}_3/\text{NO}_3^-$ ratio especially over Scandinavian regions (to a lesser extent in the LOTOS-EUROS model) and a strong increase in the $\text{NH}_3/\text{NH}_4^+$ ratio over the whole domain except for some Eastern European areas.

3.2.4 Trends in modeled SO_4^{2-} and SO_2 concentrations for different sub-regions

In this section the trends in modeled SO_4^{2-} and SO_2 for the sub-regions reported in Figure 1, are discussed. Figure 12 illustrates the relative trends in SO_2 and SO_4^{2-} concentrations for the P1 (first two column) and P2 (last two columns) periods for all the models that participated in the experiment. Trends were predicted to be statistically significant over the whole domain during the P1 period, and to a lesser extent over Eastern and Northern European regions during the P2 period. A larger decline in SO_2 concentrations during the P1 period was predicted in the Mid-Europe areas, around 85% relative reductions, compared to the Iberian Peninsula, Mediterranean and Fennoscandia areas (average reduction 24, 48 and 33%, respectively) (Figure 12 and Table 6). During the P2 period, the modeled relative reductions of SO_2 were in the range 33-65%, with the Iberian Peninsula showing a larger reduction compared to the P1 period. As already discussed in the evaluation section, SO_4^{2-} declined to a lesser degree than SO_2 , by 34 and 59% in the P1 period and 30-



49% in the P2 period, likely because of the increased availability of oxidant species and pH-dependent cloud chemistry.



3.3 Trends in organic species

3.3.1 SOA evaluation

5 During the P2 period, various field campaigns were performed in Europe using aerosol mass spectrometer instruments (AMS) to measure ambient OA concentrations at various sites. Using factor-analysis techniques, i.e. positive-matrix factorization analysis (PMF), it was possible to apportion the measured OA to a direct emitted organic factor, i.e. POA, and a more oxidized secondary factor, referred to as SOA. Even though this methodology is affected by various sources of uncertainties, we used this dataset to provide a general benchmark for modeled SOA performance during the P2
10 period.

Figure 14 shows the average modeled and observed SOA concentrations (retrieved from PMF analysis) for all the sites included in the analysis (Table S2 and Figure S1), as well as for campaigns carried out during winter and summer periods (astronomical seasons). All the models underestimate observed SOA concentrations, in general, with larger variabilities between models than for the secondary inorganic species (section 3.2.2). On average, the models
15 underestimated the SOA concentrations by about a factor of 3 to 11 (Table 6, all periods), depending on the specific model (MATCH underestimating the most and EMEP MSC-W the least) and with a larger underestimation during winter periods. The EMEP MSC-W model which accounts for aging of SOA, was closer to the observations, with average SOA concentrations of about $0.7 \mu\text{g m}^{-3}$. The higher SOA mass modeled by the EMEP MSC-W model could be explained by the shift of relatively high-volatile organic compounds toward lower volatility ranges when aging
20 processes are accounted for (with a reaction rate toward OH of $4.0 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ in the case of the EMEP MSC-W model for both ASOA and BSOA). Such processes will increase the SOA mass since low-volatile oxidation products will rapidly condense into the particle-phase. Interestingly, the MATCH model which used the same VBS scheme as the EMEP MSC-W model, but without considering SOA aging processes, tends to under-estimate SOA concentrations substantially (Figure 14). This indicates the importance of these chemical mechanisms in CTMs and
25 their impact on SOA formation. On the other hand, the models based on the two-product scheme and molecular surrogate approach scheme, i.e. MINNI and CHIMERE, respectively, yielded very similar results for the total SOA mass, with SOA concentrations ranging in between the two VBS models (i.e. around 0.4 and $0.6 \mu\text{g m}^{-3}$, averaged over all sites).

Figure S2 and Figure S3 illustrate the modeled and observed (retrieved with PMF analysis) SOA concentrations at the
30 individual sites for winter and summer periods. In general, the models had difficulties in reproducing the SOA concentrations at specific urban sites, such as Paris and Manchester (in both summer and winter periods), and in reproducing high levels of SOA concentrations at few specific sites, e.g. Harkingen and Payerne, where emissions from biomass burning are high. This could be due to missing aerosol precursors (SVOCs emissions) in the resident-heating sectors, which have been shown to have high uncertainties (Denier van der Gon et al., 2015).

35 Figure 15 shows the modeled relative and absolute contributions of anthropogenic and biogenic secondary organic aerosols to SOA concentrations. For most of the models, larger contributions of ASOA to SOA were estimated during winter periods and/or in urban areas (e.g. in Paris, Manchester and Payerne), whereas the BSOA contribution to SOA was largest during warmer periods. Especially in summer, large emissions of biogenic volatile organic compounds can act as an important source of SOA. The CHIMERE model simulated the largest contribution of BSOA, with only minor
40 variations between the stations and periods.



3.3.2 Trends in BVOCs emissions and SOA concentrations

In this section, the trends in BVOCs emissions, i.e. isoprene and terpene, are presented together with the trends in BSOA and ASOA concentrations. The trends analysis for BSOA and ASOA is reported for the full 1990–2010 period, for the different PRUDENCE zones and with the methodology as defined in Section 2.3.4. Note that not all the participants provided biogenic emissions for the full 21 years period, and only the EMEP MSC-W, CHIMERE and MATCH models provided year-by-year emissions of isoprene and monoterpenes for the EDT experiments. Moreover, for the EDT set-up, CHIMERE does not include biogenic emissions for Latitudes north of 65°N.

Figure 16 and Figure 17 illustrate the trends in isoprene and monoterpenes (first two columns) for the full 1990–2010 period. An increase in both isoprene and monoterpene species was found especially over Eastern Europe (EE), with relative increases of a 15–27% for isoprene and 14–18% for monoterpenes, and over the Fennoscandia regions (SC), with relative increases of 12–24% in isoprene emissions and 7–17% in monoterpene emissions, depending on the model (Figure 16 and Figure 17). Interestingly, the increase in biogenic emissions was predicted by all biogenic models (i.e., MEGANv2.1 and the one using vegetation data from Koeble and Seufert, 2001). These increases were mainly attributed to the increase in surface temperature used to drive the different biogenic models (Figure S5), which however were found to be not significant. The increase in surface temperature was found to be larger over Fennoscandia, Mediterranean and Eastern European areas compared to the remaining zones, (i.e. increases around 0.02, 0.02 and 0.03 K y⁻¹, respectively). We want to underline that the surface solar radiation (SSR) could also play an important role for the emission of biogenic precursors (especially for isoprene). The strong reduction in SO₄²⁻ concentrations described in section 3.2.2, especially in the eastern regions of the domain, might have induced an increase in the incoming solar radiation, referred to as brightening-periods (Wild 2009, 2011), which could affect the emission of biogenic species. However, none of the models that participated in the exercise have explicitly accounted for such an interaction and the ERA-interim forcing data rely mainly on climatological aerosol profiles. On the other hand, recent sensitivity studies performed in Europe showed that such effects might be relatively small (Oikonomakis et al., 2018). Figure 16 and Figure 17 also show the relative trends in BSOA and ASOA concentrations for all the models that were able to provide twenty-one years of data. Even though some models indicated few increases in the biogenic SOA concentrations over the Fennoscandia regions, these increases were found to be smaller than the increase in biogenic emissions and in some cases, biogenic SOA concentrations were also estimated to have declined (Figure 16 and Figure 17). This might sound counterintuitive, since one would expect more biogenic SOA to be produced as more biogenic precursors are available and, in general, the increased availability of OH radicals due to the reduction in NO_x and SO_x emissions, thus increasing the oxidation efficiency of biogenic SOA precursors, especially for isoprene. A possible explanation for this non-linear relation between the trends in biogenic emissions and the trends in BSOA concentrations could be due to the trends in the anthropogenic OA concentrations. As shown in Figure 16 and Figure 17 a strong decrease in the ASOA concentrations was found for the entire 1990–2010 period, in line with the reduction in the NMVOC precursors described in Section 3.1 (mainly from transportation sectors). Modeled ASOA concentrations indicate a decline of about 60–70% over the whole domain (considering only land areas). This might have had an effect on the formation of the BSOA fraction: in fact, the strong reduction in ASOA concentrations, and other aerosol organic and inorganic components, will reduce the availability of organic and inorganic material onto which the low-volatile oxidized compounds can condense, directly affecting the formation of the BSOA fraction. Additionally, oxidant levels and thus oxidation pathways could have changed over time, affecting as said before OH, but also NO₃ concentrations.



4. Conclusions

A modelling experiment to evaluate the capability of several chemical transport models (CTMs) to reproduce long-term air quality trends in Europe was initiated within the EURODELTA-Trends (EDT) exercise.

5 Common anthropogenic emissions, meteorological input data and boundary conditions were used by the participants, whereas the chemical and physical parameters varied between the models. Modeled air quality data for the 1990–2010 period was validated against quality-controlled long-term measurements with a focus on several primary and secondary inorganic and organic pollutants.

In general, the experiment revealed that the models were able to reproduce relatively well the observed trends in gas-phase precursors (i.e. SO_2 and NO_2), as well as secondary inorganic species, i.e. sulfate (SO_4^{2-}), total nitrate (TNO_3) and total ammonium (TNH_x), with a few exceptions at some specific sites. The range of modeled trends over 1990–2010 encompass the observed ones for SO_2 , SO_4^{2-} and TNO_3 , but not for TNH_x and NO_2 . The modeled relative declines of NO_2 concentrations were found to be 19–23% during the 1990–2000 period (P1) and 22–26% during the 2000–2010 period (P2), depending on the model. These values were in-line with the relative trends calculated from the observations, around 25 and 12%, for the P1 and P2 periods, respectively (mean values of all sites) even if models did not catch the observed stronger decrease in P1 and weaker decrease in P2. Their difficulty in reproducing the weaker decline over the second period is attributed to the challenge in modelling low NO_2 levels at EMEP background sites during that period.

The large decline in SO_2 and SO_4^{2-} concentrations during the early 90s, due to the switch to low-sulfur fuels (e.g. natural gas) and the adoption of desulfurization technologies, was well reproduced by the models, with most of the absolute trends in observations being reproduced within a factor of 2. As expected, SO_2 decreases faster than SO_x emissions, and SO_4^{2-} decreases less. This is due to the increase in cloud pH that accelerates in-cloud sulfur chemistry that consumes SO_2 to form SO_4^{2-} , constituting a positive retroaction for SO_2 decrease and a negative retroaction for SO_4^{2-} . This effect was well reproduced by the models.

25 TNH_x decreases much faster than NH_3 emissions (respectively 46% and 15% over 1990–2010). This is due to a change of TNH_x partitioning, which shifts towards gas when the atmospheric load of acids (H_2SO_4 and HNO_3) decrease. Consequently, a larger fraction of TNH_x is in NH_3 form, which deposits faster than NH_4^+ , leading to a positive retroaction in enhancing the downward trend of TNH_x . Deposition plays a critical role in TNH_x trends, and it has been noted that the observed decrease in wet deposition in the 1990s was largely driven by a couple of monitoring stations that experienced a sharp drop between 1995 and 1996 (Theobald et al., 2019) which the models fail to capture.

30 The trends in TNH_x concentrations were thus under-estimated, especially during the first decade (1990–2000), with the models exhibiting larger discrepancies compared to the other investigated species. Ammonia emissions certainly play an important role in the model performance for TNH_x . In fact, large uncertainties remain regarding present-day ammonia emissions and higher uncertainties are probably to be expected during the early 1990s.

35 The models estimated relatively lower trends in TNO_3 concentrations compared to other inorganic species (during the P1 periods), which was also indicated by the observations. A further analysis of the modeled HNO_3 and NO_3^- components revealed that HNO_3 declined more than NO_3^- during the 1990–2000 period. We attributed the later to a possible shift in the thermodynamic equilibrium of HNO_3 following the strong reduction in SO_2 concentrations, resulting in more “free-ammonia” available to drive the HNO_3 into the particle phase. Such an effect was particularly enhanced over Fennoscandia regions where differences up to a factor of 5 in the modeled relative reductions of HNO_3 and NO_3^- concentrations were found, 24% relative reduction in HNO_3 and 5% for NO_3^- , respectively (average values of all the models for the P1 period). Because HNO_3 deposits faster than NO_3^- , the shift of TNO_3 partitioning towards particles increases the lifetime of atmospheric nitrogen (as reported by Simpson et al., 2014), which contributes to



explain that TNO_3 decreases less than NO_x emission.

A comprehensive data-set of SOA concentrations retrieved from positive matrix factorization analyses (PMF) was used to investigate the models' capabilities of reproducing the SOA concentrations during the 2000–2010 period. The analysis of modeled SOA concentrations indicated that the models under-estimated the SOA fraction by varying

5 extents, by a factor of 3 to 11, depending on the model, suggesting that large uncertainties in the SOA formation mechanisms as well as in the emissions of SOA precursors remains, and more studies are needed to better elucidate the evolution of the SOA fraction, especially with a long-term prospective. The under-estimation of the SOA fraction seemed to be more pronounced during winter periods, in line with previous studies indicating missing SOA precursors in the residential sector, one of the major contributors to SOA concentrations in Europe during winter periods.

10 Therefore, this experiment confirmed once more the need to improve emission inventories of primary organic aerosol for the residential sector, especially regarding wood-burning emissions.

The analysis of the modeled trends in emissions of biogenic volatile organic compounds (BVOCs), isoprene and monoterpene, revealed an increase in these precursor emissions during the 1990–2010 period, especially in Eastern European regions and in Fennoscandia regions, by about 20%. The increase was independent of the land-use and biogenic model used and was mainly attributed to the increase of the surface temperature during the 1990–2010 period.

15 Modeled trends in ASOA concentrations indicated a strong reduction following emission reductions of non-methane volatile organic precursors, by around 60%, because of the implementation of new EURO standards for passenger cars, among other. However, modeled trends in BSOA concentrations remain less clear. Despite the modeled increase in biogenic emissions, modeled BSOA concentrations showed relatively small increasing trends, or even decreasing

20 trends. A possible explanation was mainly attributed to the reduction in the aerosol mass indicated by all the models. The latter could eventually reduce the condensation sink of low-volatility organic compounds and reduce the capability to form additional organic material from biogenic precursors, despite the increase in BVOCs emissions. Thus, more work is still needed to better characterize the trends in organic aerosol, and especially of the BSOA fraction.



Data and code availability. Technical details of the EURODELTA project simulations that permit the replication of the experiment are available on the wiki of the EMEP Task Force on Measurement and Modelling (<https://wiki.met.no/emep/emep-experts/tfmmtrendeurodelta>, last access: 21 December 2018), which also includes ESGF links to corresponding input forcing data. The EURODELTA-Trends model results are made available for public use on the AeroCom server (<https://wiki.met.no/aerocom/user-server>, last access: 21 December 2018). See Colette et al. (2017) for full terms and conditions for the use of these data. The R code and procedures used for the model evaluation are available from the corresponding author upon request.

Author contributions. ACo coordinated the EURODELTA-Trends (EDT) exercise and WA was responsible for the compilation and quality control of the observations. The following modelling teams set-up, pre-processed, ran and post-processed the simulations for each model: FC, BB, MGV and ACo for CHIMERE; ST, HF and PW for EMEP; AM, MS and RK for LOTO-EUROS; CA and RB for MATCH; MM, MA, GB, ACa and MD for MINNI. Additional post-processing of model output and uploading to the AeroCom server was done by KC. All of the analyses presented in this paper were carried out by GC with assistance from MT, KM, VR, YR, MTP, ACo, ACh and MB.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgments. G. Ciarelli was supported by ADEME in the frame of the MISTRALS/ChArMEx project and the Swiss National Science Foundation (grant no. P2EZP2_175166). The Ineris coordination of the Eurodelta-Trend exercise was supported by the French Ministry in charge of Ecology in the context of the Task Force on Measurement and Modelling of the EMEP Programme of the LRTAP Convention. Meteorological forcing with the WRF model were provided by R. Vautard and A. Stegehuis from LSCE/IPSL. The CHIMERE simulations were performed using the TGCC super computers under GENCI computing allocation. The participation of CIEMAT was financed by the Spanish Ministry of Agriculture and Fishing, Food and Environment. The MATCH participation was partly funded by the Swedish Environmental Protection Agency through the research program Swedish Clean Air and Climate (SCAC) and partly by NordForsk through the research programme Nordic WelfAir (grant no. 75007). The computing resources and the related technical support used for MINNI simulations have been provided by CRESCO/ENEAGRID High Performance Computing infrastructure and its staff. The infrastructure is funded by ENEA, the Italian National Agency for New Technologies, Energy and Sustainable Economic Development and by Italian and European research programmes (<http://www.cresco.enea.it/english>). MINNI participation to this project was supported by the “Cooperation Agreement for support to international Conventions, Protocols and related negotiations on air pollution issues”, funded by the Italian Ministry for Environment and Territory and Sea. The simulations with the EMEP MSC-W model were supported by the Research Council of Norway in the framework of the Programme for Supercomputing: through the EMEP project (grant NN2890K) for CPU, and the Norstore project “European Monitoring and Evaluation Programme” (grant NS9005K) for data storage. The GAINS emission trends were produced as part of the FP7 European Research Project ECLIPSE (Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants); grant no. 282688.



References.

- 5 Aksoyoglu, S., Keller, J., Ciarelli, G., Prévôt, A.S.H., Baltensperger, U., 2014. A model study on changes of European and Swiss particulate matter, ozone and nitrogen deposition between 1990 and 2020 due to the revised Gothenburg protocol. *Atmos Chem Phys* 14, 13081–13095. <https://doi.org/10.5194/acp-14-13081-2014>.
- 10 Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F., Winiwarter, W., 2011. Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications. *Environ. Model. Softw.* 26, 1489–1501. <https://doi.org/10.1016/j.envsoft.2011.07.012>.
- 15 Andersson, C., Bergström, R., Bennet, C., Robertson, L., Thomas, M., Korhonen, H., Lehtinen, K.E.J., Kokkola, H., 2015. MATCH-SALSA – Multi-scale Atmospheric Transport and CHEMistry model coupled to the SALSA aerosol microphysics model – Part 1: Model description and evaluation. *Geosci. Model Dev.* 8, 171–189. <https://doi.org/10.5194/gmd-8-171-2015>.
- 20 Andersson, C., Langner, J., Bergström, R., 2007. Interannual variation and trends in air pollution over Europe due to climate variability during 1958–2001 simulated with a regional CTM coupled to the ERA40 reanalysis. *Tellus B* 59, 77–98. <https://doi.org/10.1111/j.1600-0889.2006.00196.x>.
- 25 Athanasiadis, I.N., Mitkas, P.A., Rizzoli, A.E., Gómez, J.M. (Eds.), 2009. *Information Technologies in Environmental Engineering: Proceedings of the 4th International ICSC Symposium Thessaloniki, Greece, May 28-29, 2009*, Environmental Engineering. Springer-Verlag, Berlin Heidelberg.
- 30 Atkinson, R., Baulch, D.L., Cox, R.A., Crowley, J.N., Hampson, R.F., Hynes, R.G., Jenkin, M.E., Rossi, M.J., Troe, J., 2004. Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x, HO_x, NO_x and SO_x species. *Atmospheric Chem. Phys.* 4, 1461–1738. <https://doi.org/10.5194/acp-4-1461-2004>.
- 35 Atkinson, R., Baulch, D.L., Cox, R.A., Crowley, J.N., Hampson, R.F., Hynes, R.G., Jenkin, M.E., Rossi, M.J., Troe, J., IUPAC Subcommittee, 2006. Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species. *Atmospheric Chem. Phys.* 6, 3625–4055. <https://doi.org/10.5194/acp-6-3625-2006>.
- 40 Atkinson, R., Baulch, D.L., Cox, R.A., Hampson, R.F., Kerr, J.A., Rossi, M.J., Troe, J., 1997. Evaluated Kinetic, Photochemical and Heterogeneous Data for Atmospheric Chemistry: Supplement V. IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry. *J. Phys. Chem. Ref. Data* 26, 521–1011. <https://doi.org/10.1063/1.556011>.
- 45 Backes, A., Aulinger, A., Bieser, J., Matthias, V., Quante, M., 2016. Ammonia emissions in Europe, part I: Development of a dynamical ammonia emission inventory. *Atmos. Environ.* 131, 55–66. <https://doi.org/10.1016/j.atmosenv.2016.01.041>.
- 50 Banzhaf, S., Schaap, M., Kranenburg, R., Manders, A.M.M., Segers, A.J., Visschedijk, A.J.H., Denier van der Gon, H.A.C., Kuenen, J.J.P., van Meijgaard, E., van Ulft, L.H., Cofala, J., Builtjes, P.J.H., 2015. Dynamic model evaluation for secondary inorganic aerosol and its precursors over Europe between 1990 and 2009. *Geosci Model Dev* 8, 1047–1070. <https://doi.org/10.5194/gmd-8-1047-2015>.
- 55 Bergström, R., Denier van der Gon, H.A.C., Prévôt, A.S.H., Yttri, K.E., Simpson, D., 2012a. Modelling of organic aerosols over Europe (2002–2007) using a volatility basis set (VBS) framework: application of different assumptions regarding the formation of secondary organic aerosol. *Atmos Chem Phys* 12, 8499–8527. <https://doi.org/10.5194/acp-12-8499-2012>.
- 60 Bergström, R., Denier van der Gon, H.A.C., Prévôt, A.S.H., Yttri, K.E., Simpson, D., 2012b. Modelling of organic aerosols over Europe (2002–2007) using a volatility basis set (VBS) framework: application of different assumptions regarding the formation of secondary organic aerosol. *Atmos Chem Phys* 12, 8499–8527. <https://doi.org/10.5194/acp-12-8499-2012>.
- Bessagnet, B., Pirovano, G., Mircea, M., Cuvelier, C., Aulinger, A., Calori, G., Ciarelli, G., Manders, A., Stern, R., Tsyro, S., García Vivanco, M., Thunis, P., Pay, M.-T., Colette, A., Couvidat, F., Meleux, F., Rouil, L., Ung, A., Aksoyoglu, S., Baldasano, J.M., Bieser, J., Briganti, G., Cappelletti, A., D’Isidoro, M., Finardi, S., Kranenburg, R., Silibello, C., Carnevale, C., Aas, W., Dupont, J.-C., Fagerli, H., Gonzalez, L., Menut, L., Prévôt, A.S.H., Roberts, P., White, L., 2016. Presentation of the EURODELTA III intercomparison exercise – evaluation of the chemistry transport models’ performance on criteria pollutants and joint analysis with meteorology. *Atmos Chem Phys* 16, 12667–12701.



<https://doi.org/10.5194/acp-16-12667-2016>.

- 5 Bian, H., Chin, M., Hauglustaine, D.A., Schulz, M., Myhre, G., Bauer, S.E., Lund, M.T., Karydis, V.A., Kucsera, T.L., Pan, X., Pozzer, A., Skeie, R.B., Steenrod, S.D., Sudo, K., Tsigaridis, K., Tsimpidi, A.P., Tsyro, S.G., 2017. Investigation of global particulate nitrate from the AeroCom phase III experiment. *Atmos Chem Phys* 17, 12911–12940. <https://doi.org/10.5194/acp-17-12911-2017>.
- 10 Binkowski, F., Shankar, U., 1995. The Regional Particulate Matter Model: 1. Model description and preliminary results - Binkowski - 1995 - *Journal of Geophysical Research: Atmospheres* - Wiley Online Library [WWW Document]. URL <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/95JD02093> (accessed 5.8.18).
- 15 Byun, D., Schere, K.L., 2006. Review of the Governing Equations, Computational Algorithms, and Other Components of the Models-3 Community Multiscale Air Quality (CMAQ) Modeling System. *Appl. Mech. Rev.* 59, 51–77. <https://doi.org/10.1115/1.2128636>.
- 20 Carter, W.P.L., 2000. Documentation of the SAPRC-99 Chemical Mechanism for VOC Reactivity Assessment, Final Report to California Air Resources Board Contract No. 92-329, and 95-308, available at: <http://www.cert.ucr.edu/~carter/absts.htm#saprc99>.
- 20 Carter, W.P.L., 1996. Condensed atmospheric photooxidation mechanisms for isoprene. *Atmos. Environ.* 30, 4275–4290. [https://doi.org/10.1016/1352-2310\(96\)00088-X](https://doi.org/10.1016/1352-2310(96)00088-X).
- 25 Cholakian, A., Beekmann, M., Colette, A., Coll, I., Siour, G., Sciare, J., Marchand, N., Couvidat, F., Pey, J., Gros, V., Sauvage, S., Michoud, V., Sellegri, K., Colomb, A., Sartelet, K., Langley DeWitt, H., Elser, M., Prévot, A.S.H., Szidat, S., Dulac, F., 2017. Simulation of fine organic aerosols in the western Mediterranean area during the ChArMEx 2013 summer campaign. *Atmos Chem Phys Discuss* 2017, 1–45. <https://doi.org/10.5194/acp-2017-697>.
- 30 Chrit, M., Sartelet, K., Sciare, J., Pey, J., Marchand, N., Couvidat, F., Sellegri, K., Beekmann, M., 2017. Modelling organic aerosol concentrations and properties during ChArMEx summer campaigns of 2012 and 2013 in the western Mediterranean region. *Atmos Chem Phys* 17, 12509–12531. <https://doi.org/10.5194/acp-17-12509-2017>.
- 35 Ciarelli, G., Aksoyoglu, S., Crippa, M., Jimenez, J.-L., Nemitz, E., Sellegri, K., Äijälä, M., Carbone, S., Mohr, C., O’Dowd, C., Poulain, L., Baltensperger, U., Prévôt, A.S.H., 2016. Evaluation of European air quality modelled by CAMx including the volatility basis set scheme. *Atmos Chem Phys* 16, 10313–10332. <https://doi.org/10.5194/acp-16-10313-2016>.
- 40 Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope, C.A., Shin, H., Straif, K., Shaddick, G., Thomas, M., Dingenen, R. van, Donkelaar, A. van, Vos, T., Murray, C.J.L., Forouzanfar, M.H., 2017. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *The Lancet* 389, 1907–1918. [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6).
- 45 Colette, A., Andersson, C., Manders, A., Mar, K., Mircea, M., Pay, M.P., Raffort, V., Tsyro, S., Cuvelier, C., Adani, M., Bessagnet, B., Bergström, R., Briganti, G., Butler, T., Cappelletti, A., Couvidat, F., D’Isidoro, M., Doumbia, T., Fagerli, H., Granier, C., Heyes, C., Klimont, Z., Ojha, N., Otero, N., Schaap, M., Sindelarova, K., Stegehuis, A.I., Roustan, Y., Vautard, R., van Meijgaard, E., Vivanco, M.G., Wind, P., 2017a. EURODELTA-Trends, a multi-model experiment of air quality hindcast in Europe over 1990–2010. *Geosci. Model Dev.* 10, 3255–3276. <https://doi.org/10.5194/gmd-10-3255-2017>.
- 50 Colette, A., Andersson, C., Manders, A., Mar, K., Mircea, M., Pay, M.-T., Raffort, V., Tsyro, S., Cuvelier, C., Adani, M., Bessagnet, B., Bergström, R., Briganti, G., Butler, T., Cappelletti, A., Couvidat, F., D’Isidoro, M., Doumbia, T., Fagerli, H., Granier, C., Heyes, C., Klimont, Z., Ojha, N., Otero, N., Schaap, M., Sindelarova, K., Stegehuis, A.I., Roustan, Y., Vautard, R., van Meijgaard, E., Vivanco, M.G., Wind, P., 2017b. EURODELTA-Trends, a multi-model experiment of air quality hindcast in Europe over 1990–2010. *Geosci Model Dev* 10, 3255–3276. <https://doi.org/10.5194/gmd-10-3255-2017>.
- 55 Colette, A., Granier, C., Hodnebrog, Ø., Jakobs, H., Maurizi, A., Nyiri, A., Bessagnet, B., D’Angiola, A., D’Isidoro, M., Gauss, M., Meleux, F., Memmesheimer, M., Mieville, A., Rouïl, L., Russo, F., Solberg, S., Stordal, F., Tampieri, F., 2011. Air quality trends in Europe over the past decade: a first multi-model assessment. *Atmos Chem Phys* 11, 11657–11678. <https://doi.org/10.5194/acp-11-11657-2011>.
- 60 Couvidat, F., Bessagnet, B., Garcia-Vivanco, M., Real, E., Menut, L., Colette, A., 2018. Development of an inorganic



- and organic aerosol model (CHIMERE 2017 β v1.0): seasonal and spatial evaluation over Europe. *Geosci Model Dev* 11, 165–194. <https://doi.org/10.5194/gmd-11-165-2018>.
- 5 Couvidat, F., Debry, É., Sartelet, K., Seigneur, C., 2012. A hydrophilic/hydrophobic organic (H₂O) aerosol model: Development, evaluation and sensitivity analysis: H₂O--A MODEL TO PREDICT ORGANIC AEROSOL. *J. Geophys. Res. Atmospheres* 117, n/a-n/a. <https://doi.org/10.1029/2011JD017214>.
- Couvidat, F., Sartelet, K., 2015. The Secondary Organic Aerosol Processor (SOAP v1.0) model: a unified model with different ranges of complexity based on the molecular surrogate approach. *Geosci Model Dev* 8, 1111–1138. <https://doi.org/10.5194/gmd-8-1111-2015>.
- 10 Crippa, M., Canonaco, F., Lanz, V.A., Äijälä, M., Allan, J.D., Carbone, S., Capes, G., Ceburnis, D., Dall’Osto, M., Day, D.A., DeCarlo, P.F., Ehn, M., Eriksson, A., Freney, E., Hildebrandt Ruiz, L., Hillamo, R., Jimenez, J.L., Junninen, H., Kiendler-Scharr, A., Kortelainen, A.-M., Kulmala, M., Laaksonen, A., Mensah, A.A., Mohr, C., Nemitz, E., O’Dowd, C., Ovadnevaite, J., Pandis, S.N., Petäjä, T., Poulain, L., Saarikoski, S., Sellegri, K., Swietlicki, E., Tiitta, P., Worsnop, D.R., Baltensperger, U., Prévôt, A.S.H., 2014. Organic aerosol components derived from 25 AMS data sets across Europe using a consistent ME-2 based source apportionment approach. *Atmos Chem Phys* 14, 6159–6176. <https://doi.org/10.5194/acp-14-6159-2014>.
- 20 De Moore, W.B., Sandetr, S.P., Golden, D.M., Hampson, R.F., Kurylo, M.J., Howard, C.J., Ravishankara, A.R., Kolb, C.E., Molina, M.J., 1994. Chemical kinetics and photochemical data for use in stratospheric modelling evaluation.
- Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A.P., Monge-Sanz, B.M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., Vitart, F., 2011. The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Q. J. R. Meteorol. Soc.* 137, 553–597. <https://doi.org/10.1002/qj.828>.
- 25 Denier van der Gon, H.A.C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S.N., Simpson, D., Visschedijk, A.J.H., 2015. Particulate emissions from residential wood combustion in Europe – revised estimates and an evaluation. *Atmos Chem Phys* 15, 6503–6519. <https://doi.org/10.5194/acp-15-6503-2015>.
- Derognat, C., Beekmann, M., Baeumle, M., Martin, D., Schmidt, H., 2003. Effect of biogenic volatile organic compound emissions on tropospheric chemistry during the Atmospheric Pollution Over the Paris Area (ESQUIF) campaign in the Ile-de-France region. *J. Geophys. Res.* 108. <https://doi.org/10.1029/2001JD001421>.
- 35 Donahue, N.M., Epstein, S.A., Pandis, S.N., Robinson, A.L., 2011. A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics. *Atmos Chem Phys* 11, 3303–3318. <https://doi.org/10.5194/acp-11-3303-2011>.
- 40 Donahue, N.M., Kroll, J.H., Pandis, S.N., Robinson, A.L., 2012. A two-dimensional volatility basis set – Part 2: Diagnostics of organic-aerosol evolution. *Atmos Chem Phys* 12, 615–634. <https://doi.org/10.5194/acp-12-615-2012>.
- 45 EEA, 2012. EEA Report, No 4/2012. Air quality in Europe - 2010 Report. <https://doi.org/10.2800/55823>.
- EMEP, 2012. Data Report 2010 Acidifying and eutrophying compounds and particulate matter. EMEP/CCC-Report 1/2012. URL: <https://www.nilu.no/projects/ccc/reports/cccr1-2012.pdf>.
- 50 Fagerli, H., Aas, W., 2008. Trends of nitrogen in air and precipitation: Model results and observations at EMEP sites in Europe, 1980–2003. *Environ. Pollut.* 154, 448–461. <https://doi.org/10.1016/j.envpol.2008.01.024>.
- Fountoukis, C., Nenes, A., 2007. ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K⁺–Ca²⁺–Mg²⁺–NH₄⁺–Na⁺–SO₄²⁻–NO₃⁻–Cl⁻–H₂O aerosols. *Atmos Chem Phys* 7, 1153–1165. <https://doi.org/10.5194/acp-7-1153-2007>.
- 55 Grell, G.A., Peckham, S.E., Schmitz, R., McKeen, S.A., Frost, G., Skamarock, W.C., Eder, B., 2005. Fully coupled “online” chemistry within the WRF model. *Atmos. Environ.* 39, 6957–6975. <https://doi.org/10.1016/j.atmosenv.2005.04.027>.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P.I., Geron, C., 2006. Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature). *Atmospheric Chem. Phys.* 6, 3181–3210. <https://doi.org/10.5194/acp-6-3181-2006>.
- 60 Guenther, A.B., Jiang, X., Heald, C.L., Sakulyanontvittaya, T., Duhl, T., Emmons, L.K., Wang, X., 2012. The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for



- modeling biogenic emissions. *Geosci Model Dev* 5, 1471–1492. <https://doi.org/10.5194/gmd-5-1471-2012>.
- 5 Guenther, A.B., Zimmerman, P.R., Harley, P.C., Monson, R.K., Fall, R., 1993. Isoprene and monoterpene emission rate variability: Model evaluations and sensitivity analyses. *J. Geophys. Res.* 98, 12609. <https://doi.org/10.1029/93JD00527>.
- Huang, G., Brook, R., Crippa, M., Janssens-Maenhout, G., Schieberle, C., Dore, C., Guizzardi, D., Muntean, M., Schaaf, E., Friedrich, R., 2017. Speciation of anthropogenic emissions of non-methane volatile organic compounds: a global gridded data set for 1970–2012. *Atmos Chem Phys* 17, 7683–7701. <https://doi.org/10.5194/acp-17-7683-2017>.
- 10 Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K.R., Slowik, J.G., Platt, S.M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S.M., Bruns, E.A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I.E., Prévôt, A.S.H., 2014. High secondary aerosol contribution to particulate pollution during haze events in China. *Nature* 514, 218–222. <https://doi.org/10.1038/nature13774>.
- 15 IEA, 2012. IEA: Energy Statistics of OECD Countries, International Energy Agency Paris, France.
- Jimenez, J.L., Canagaratna, M.R., Donahue, N.M., Prevot, A.S.H., Zhang, Q., Kroll, J.H., DeCarlo, P.F., Allan, J.D., Coe, H., Ng, N.L., Aiken, A.C., Docherty, K.S., Ulbrich, I.M., Grieshop, A.P., Robinson, A.L., Duplissy, J., Smith, J.D., Wilson, K.R., Lanz, V.A., Hueglin, C., Sun, Y.L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J.M., Collins, D.R., Cubison, M.J., E, Dunlea, J., Huffman, J.A., Onasch, T.B., Alfarra, M.R., Williams, P.I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimojo, A., Sun, J.Y., Zhang, Y.M., Dzepina, K., Kimmel, J.R., Sueper, D., Jayne, J.T., Herndon, S.C., Trimborn, A.M., Williams, L.R., Wood, E.C., Middlebrook, A.M., Kolb, C.E., Baltensperger, U., Worsnop, D.R., 2009. Evolution of Organic Aerosols in the Atmosphere. *Science* 326, 1525–1529. <https://doi.org/10.1126/science.1180353>.
- Kendall, M.G., 1948. Rank correlation methods, Rank correlation methods., Griffin, Oxford, England.
- 30 Koeble, R., Seufert, G., 2001. Novel Maps for Forest Tree Species in Europe, A Changing Atmosphere, 8th European Symposium on the Physico-Chemical Behaviour of Atmospheric Pollutants, Torino, Italy, 2001.
- Lachatre, M., Fortems-Cheiney, A., Foret, G., Siour, G., Dufour, G., Clarisse, L., Clerbaux, C., Coheur, P.-F., Van Damme, M., Beekmann, M., 2018. The unintended consequence of SO₂ and NO_x regulations over China: increase of ammonia levels and impact on PM_{2.5} concentrations. *Atmospheric Chem. Phys. Discuss.* 1–25. <https://doi.org/10.5194/acp-2018-1092>.
- 35 Langner, J., Bergström, R., Pleijel, K., 1998. : European scale modeling of sulphur, oxidized nitrogen and photochemical oxidants, Model development and evaluation for the 1994 growing season.
- Mailler, S., Menut, L., Khvorostyanov, D., Valari, M., Couvidat, F., Siour, G., Turquety, S., Briant, R., Tuccella, P., Bessagnet, B., Colette, A., Létinois, L., Markakis, K., Meleux, F., 2017. CHIMERE-2017: from urban to hemispheric chemistry-transport modeling. *Geosci Model Dev* 10, 2397–2423. <https://doi.org/10.5194/gmd-10-2397-2017>.
- 45 Mallet, V., Quélo, D., Sportisse, B., Ahmed de Biasi, M., Debry, é., Korsakissok, I., Wu, L., Roustan, Y., Sartelet, K., Tombette, M., Foudhil, H., 2007. Technical Note: The air quality modeling system Polyphemus. *Atmospheric Chem. Phys.* 7, 5479–5487. <https://doi.org/10.5194/acp-7-5479-2007>.
- 50 Manders, A.M.M., Bultjes, P.J.H., Curier, L., Denier van der Gon, H.A.C., Hendriks, C., Jonkers, S., Kranenburg, R., Kuenen, J.J.P., Segers, A.J., Timmermans, R.M.A., Visschedijk, A.J.H., Wichink Kruit, R.J., van Pul, W.A.J., Sauter, F.J., van der Swaluw, E., Swart, D.P.J., Douros, J., Eskes, H., van Meijgaard, E., van Ulft, B., van Velthoven, P., Banzhaf, S., Mues, A.C., Stern, R., Fu, G., Lu, S., Heemink, A., van Velzen, N., Schaap, M., 2017. Curriculum vitae of the LOTOS–EUROS (v2.0) chemistry transport model. *Geosci Model Dev* 10, 4145–4173. <https://doi.org/10.5194/gmd-10-4145-2017>.
- 55 Mann, H.B., 1945. Nonparametric Tests Against Trend. *Econometrica* 13, 245. <https://doi.org/10.2307/1907187>.
- 60 Mar, K.A., Ojha, N., Pozzer, A., Butler, T.M., 2016. Ozone air quality simulations with WRF-Chem (v3.5.1) over Europe: model evaluation and chemical mechanism comparison. *Geosci Model Dev* 9, 3699–3728. <https://doi.org/10.5194/gmd-9-3699-2016>.
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., Coll, I., Curci, G., Foret, G.,




- Hodzic, A., Mailler, S., Meleux, F., Monge, J.-L., Pison, I., Siour, G., Turquety, S., Valari, M., Vautard, R., Vivanco, M.G., 2013. CHIMERE 2013: a model for regional atmospheric composition modelling. *Geosci Model Dev* 6, 981–1028. <https://doi.org/10.5194/gmd-6-981-2013>.
- 5 Mircea, M., 2016. Impact of Grid Resolution on Aerosol Predictions: A Case Study over Italy. *Aerosol Air Qual. Res.* 16, 1253–1267. <https://doi.org/10.4209/aaqr.2015.02.0058>.
- Mozurkewich, M., 1993. The dissociation constant of ammonium nitrate and its dependence on temperature, relative humidity and particle size. *Atmospheric Environ. Part Gen. Top.* 27, 261–270. [https://doi.org/10.1016/0960-1686\(93\)90356-4](https://doi.org/10.1016/0960-1686(93)90356-4).
- 10 Nees, A., Pandis, S.N., Pilinis, C., 1999. Continued development and testing of a new thermodynamic aerosol module for urban and regional air quality models. *Atmos. Environ.* 33, 1553–1560. [https://doi.org/10.1016/S1352-2310\(98\)00352-5](https://doi.org/10.1016/S1352-2310(98)00352-5).
- 15 Nees, A., Pilinis, C., Pandis, S.N., 1998. ISORROPIA: A new thermodynamic equilibrium model for multiphase multicomponent inorganic aerosols.
- Oikonomakis, E., Aksoyoglu, S., Wild, M., Ciarelli, G., Baltensperger, U., Prévôt, A.S.H., 2018. Modeling the impact of solar “brightening” on summer surface ozone over Europe between 1990 and 2010. *Atmos Chem Phys Discuss* 2018, 1–41. <https://doi.org/10.5194/acp-2017-1182>.
- Paatero, P., 1999. The Multilinear Engine—A Table-Driven, Least Squares Program for Solving Multilinear Problems, Including the n-Way Parallel Factor Analysis Model. *J. Comput. Graph. Stat.* 8, 854–888. <https://doi.org/10.1080/10618600.1999.10474853>.
- 25 Pope, C.A., Dockery, D.W., 2006. Health Effects of Fine Particulate Air Pollution: Lines that Connect. *J. Air Waste Manag. Assoc.* 56, 709–742. <https://doi.org/10.1080/10473289.2006.10464485>.
- 30 Reed, C., Evans, M.J., Di Carlo, P., Lee, J.D., Carpenter, L.J., 2016. Interferences in photolytic NO₂ measurements: explanation for an apparent missing oxidant? *Atmospheric Chem. Phys.* 16, 4707–4724. <https://doi.org/10.5194/acp-16-4707-2016>.
- Robertson, L., Langner, J., Engardt, M., 1999. An Eulerian Limited-Area Atmospheric Transport Model. *J. Appl. Meteorol.* 38, 190–210. [https://doi.org/10.1175/1520-0450\(1999\)038<0190:AELAAT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1999)038<0190:AELAAT>2.0.CO;2).
- 35 Sartelet, K.N., Couvidat, F., Seigneur, C., Roustan, Y., 2012. Impact of biogenic emissions on air quality over Europe and North America. *Atmos. Environ.* 53, 131–141. <https://doi.org/10.1016/j.atmosenv.2011.10.046>.
- 40 Schaap, M., Manders, A.M.M., Hendriks, E., Cnossen, J.M., Segers, A.J., Denier van der Gon, H.A.C., Jozwicka, M., Sauter, F.J., Velders, G.J.M., Beck, J.P., 2009. Regional Modelling of Particulate Matter for the Netherlands, PBL report 500099008.
- Schaap, M., Timmermans, R.M.A., Roemer, M., Boersen, G.A.C., Bultjes, P.J.H., Sauter, F.J., Velders, G.J.M., Beck, J.P., 2008. The LOTOS EUROS model: description, validation and latest developments. *Int. J. Environ. Pollut.* 32, 270. <https://doi.org/10.1504/IJEP.2008.017106>.
- 45 Schell, M., Ackermann, I.J., Hass, H., Binkowski, F.S., Ebel, A., 2001. Modeling the formation of secondary organic aerosol within a comprehensive air quality model system. *J. Geophys. Res. Atmospheres* 106, 28275–28293. <https://doi.org/10.1029/2001JD000384>.
- 50 Seinfeld, J.H., Pandis, S.N., 2012. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*. Wiley.
- Sen, P.K., 1968. Estimates of the Regression Coefficient Based on Kendall’s Tau. *J. Am. Stat. Assoc.* 63, 1379–1389. <https://doi.org/10.1080/01621459.1968.10480934>.
- 55 Simpson, D., Andersson, C., Christensen, J.H., Engardt, M., Geels, C., Nyiri, A., Posch, M., Soares, J., Sofiev, M., Wind, P., Langner, J., 2014. Impacts of climate and emission changes on nitrogen deposition in Europe: a multi-model study. *Atmospheric Chem. Phys.* 14, 6995–7017. <https://doi.org/10.5194/acp-14-6995-2014>.
- 60 Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L.D., Fagerli, H., Flechard, C.R., Hayman, G.D., Gauss, M., Jonson, J.E., Jenkin, M.E., Nyiri, A., Richter, C., Semeena, V.S., Tsyro, S., Tuovinen, J.-P., Valdebenito, Á., Wind, P., 2012. The EMEP MSC-W chemical transport model – technical description. *Atmos Chem Phys* 12, 7825–



7865. <https://doi.org/10.5194/acp-12-7825-2012>.
- 5 Skamarock, W., Klemp, J., Dudhia, J., Gill, D., Barker, D., Wang, W., Huang, X.Y., Duda, M., 2008. A Description of the Advanced Research WRF Version 3. UCAR/NCAR. <https://doi.org/10.5065/D68S4MVH>.
- 5 Stegehuis, A.L., Vautard, R., Ciais, P., Teuling, A.J., Miralles, D.G., Wild, M., 2015. An observation-constrained multi-physics WRF ensemble for simulating European mega heat waves. *Geosci. Model Dev.* 8, 2285–2298. <https://doi.org/10.5194/gmd-8-2285-2015>.
- 10 Terrenoire, E., Bessagnet, B., Rouïl, L., Tognet, F., Pirovano, G., Létinois, L., Beauchamp, M., Colette, A., Thunis, P., Amann, M., Menut, L., 2015. High-resolution air quality simulation over Europe with the chemistry transport model CHIMERE. *Geosci. Model Dev.* 8, 21–42. <https://doi.org/10.5194/gmd-8-21-2015>.
- 15 Theobald, M.R., Vivanco, M.G., Aas, W., Andersson, C., Ciarelli, G., Couvidat, F., Cuvelier, K., Manders, A., Mircea, M., Pay, M.-T., Tsyro, S., Adani, M., Bergström, R., Bessagnet, B., Briganti, G., Cappelletti, A., D'Isidoro, M., Fagerli, H., Mar, K., Otero, N., Raffort, V., Roustan, Y., Schaap, M., Wind, P., Colette, A., 2019. An evaluation of European nitrogen and sulfur wet deposition and their trends estimated by six chemistry transport models for the period 1990–2010. *Atmospheric Chem. Phys.* 19, 379–405. <https://doi.org/10.5194/acp-19-379-2019>.
- 20 Tørseth, K., Aas, W., Breivik, K., Fjæraa, A.M., Fiebig, M., Hjellbrekke, A.G., Lund Myhre, C., Solberg, S., Yttri, K.E., 2012. Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009. *Atmos Chem Phys* 12, 5447–5481. <https://doi.org/10.5194/acp-12-5447-2012>.
- 25 Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P.J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S.E., Bellouin, N., Benedetti, A., Bergman, T., Bernsten, T.K., Beukes, J.P., Bian, H., Carslaw, K.S., Chin, M., Curci, G., Diehl, T., Easter, R.C., Ghan, S.J., Gong, S.L., Hodzic, A., Hoyle, C.R., Iversen, T., Jathar, S., Jimenez, J.L., Kaiser, J.W., Kirkevåg, A., Koch, D., Kokkola, H., Lee, Y.H., Lin, G., Liu, X., Luo, G., Ma, X., Mann, G.W., Mihalopoulos, N., Morcrette, J.-J., Müller, J.-F., Myhre, G., Myriokefalitakis, S., Ng, N.L., O'Donnell, D., Penner, J.E., Pozzoli, L., Pringle, K.J., Russell, L.M., Schulz, M., Sciare, J., Seland, Ø., Shindell, D.T., Sillman, S., Skeie, R.B., Spracklen, D., Stavrou, T., Steenrod, S.D., Takemura, T., Tiitta, P., Tilmes, S., Tost, H., van Noije, T., van Zyl, P.G., von Salzen, K., Yu, F., Wang, Z., Wang, Z., Zaveri, R.A., Zhang, H., Zhang, K., Zhang, Q., Zhang, X., 2014. The AeroCom evaluation and intercomparison of organic aerosol in global models. *Atmos Chem Phys* 14, 10845–10895. <https://doi.org/10.5194/acp-14-10845-2014>.
- 30 Tsimpidi, A.P., Karydis, V.A., Pandis, S.N., 2007. Response of Inorganic Fine Particulate Matter to Emission Changes of Sulfur Dioxide and Ammonia: The Eastern United States as a Case Study. *J. Air Waste Manag. Assoc.* 57, 1489–1498. <https://doi.org/10.3155/1047-3289.57.12.1489>.
- 35 Tsimpidi, A.P., Karydis, V.A., Pandis, S.N., Lelieveld, J., 2016. Global combustion sources of organic aerosols: model comparison with 84 AMS factor-analysis data sets. *Atmos Chem Phys* 16, 8939–8962. <https://doi.org/10.5194/acp-16-8939-2016>.
- 40 UNECE, 1999. UNECE: Protocol to the 1979 Convention on Long-Range Transboundary Air Pollution to abate acidification, eutrophication and ground-level ozone.
- 45 UNECE LRTAP, 2016. Towards Cleaner Air. Scientific Assessment Report 2016. EMEP Steering Body and Working Group on Effects of the Convention on Long-Range Transboundary Air Pollution, Oslo. xx+50pp.
- 50 van Loon, M., Vautard, R., Schaap, M., Bergström, R., Bessagnet, B., Brandt, J., Bultjes, P.J.H., Christensen, J.H., Cuvelier, C., Graff, A., Jonson, J.E., Krol, M., Langner, J., Roberts, P., Rouil, L., Stern, R., Tarrasón, L., Thunis, P., Vignati, E., White, L., Wind, P., 2007. Evaluation of long-term ozone simulations from seven regional air quality models and their ensemble. *Atmos. Environ.* 41, 2083–2097. <https://doi.org/10.1016/j.atmosenv.2006.10.073>.
- 55 WHO, 2013. Health risks of air pollution in Europe – HRAPIE – Summary of recommendations for question D5 on “Identification of concentration-response functions” for cost-effectiveness analysis.



Figures and Tables.



5 **Figure 1:** The extension of the EURODELTA-Trends domain as well as of the sub-regions adapted from the PRUDENCE zones. From South to North: Mediterranean regions (MD), Iberian Peninsula (IP), France (FR), Alps (AL), Mid-Europe (ME), Eastern Europe (EE), Benelux regions (BX), British Isles (BI) and Fennoscandia (SC).







Figure 2: Locations of the observational sites (red dots). The numbers of observational sites available for each species are 30, 20, 25, 13 and 16 for SO₂, SO₄²⁻ (PSO₄), NO₂, TNO₃ and TNH_x, respectively.




5 **Figure 3:** Modeled and observed (Obs) mean relative trends of NO₂ (upper-panels) and SO₂ concentrations (lower-panels) for the P1 (1990–2000), P2 (2000–2010) and PT (1990–2010) periods.



5 **Figure 4:** Modeled and observed NO₂ (upper-panel) and SO₂ (lower-panel) relative trends for the P1 (1990–2000), P2 (2000–2010) and PT (1990–2010) periods (left to right). The continuous line indicates the 1:1 line, and the dotted lines indicate the 1:2 and 1:3 lines (and their reciprocals).




5 **Figure 5:** Percentage of statistically significant/non-significant increasing/decreasing trends in the observations and modeled data for NO₂ (left) and SO₂ (right) for the P1 (1990–2000), P2 (2000–2010) and PT (1990–2010) periods (from top to bottom).



5

Figure 6: Modeled and observed (Obs) mean relative trends in SO_4^{2-} (PSO_4), TNH_x , and TNO_3 concentrations for the P1 (1990–2000), P2 (2000–2010) and PT (1990–2010) periods.



5 **Figure 7:** Modeled and observed SO_4^{2-} (upper-panel), TNH_x (mid-panel) and TNO_3 (lower-panel) relative trends for the P1 (1990–2000), P2 (2000–2010) and PT (1990–2010) periods (left to right). The continuous line indicates the 1:1 line, and the dotted lines indicate the 1:2 and 1:3 lines (and their reciprocals).





Figure 8: Percentage of statistically significant/non-significant increasing/decreasing trends in the observations and modeled data for SO₂⁻² (top-panel) and TNH_x (central-panel) and TN0₃ (bottom-panel) for the P1 (1990–2000), P2 (2000–2010) and PT (1990–2010) periods (from left to right).



5 **Figure 9:** Modeled HNO₃ and NO₃ relative trends over lands for the P1 (1990–2000, first and second columns) and P2 (2000–2010, third and second columns) periods as predicted by all the models (rows; from top to bottom: EMEP MSC-W, CHIMERE, MATCH, LOTOS-EUROS, MINNI). White areas indicate non-significant trends.





Figure 10: Modeled relative trends in HNO₃ and NO₃⁻ concentrations for the different PRUDENCE zones (Figure 1) for the P1 (left) and P2 (right) periods. The columns show the averages (over land) of all the model estimates and the bars show the standard deviation respect to models.

5



5 **Figure 11:** Modeled $\text{HNO}_3/\text{NO}_3^-$ and $\text{NH}_3/\text{NH}_4^+$ molar ratio relative trends over lands for the P1 (1990–2000, first and second columns) and P2 (2000–2010, third and second columns) periods as predicted by all the models (rows; from top to bottom: EMEP, MSC-W, CHIMERE, MATCH, LOTOS-EUROS, MINNI). White areas indicate non-significant trends. Scale was saturate at 100% to facilitate the comprehension of the panel.




Figure 12: Modeled SO_2 and SO_4^{2-} relative trends over lands for the P1 (1990–2000, first and second columns) and P2 (2000–2010, third and second columns) periods as predicted by all the models (rows; from top to bottom: EMEP MSC-W, CHIMERE, MATCH, LOTOS-EUROS, MINNI). White areas indicate non-significant trends. Scale was saturate at 100% to facilitate the comprehension of the panel.

5




Figure 13: Modeled relative trends in SO₂ and SO₄²⁻ concentrations for the different PRUDENCE zones (Figure 1) for the P1 (left) and P2 (right) periods. The columns show the averages (over land) of all the model estimates and the bars show the standard deviation respect to models.

5




Figure 14: Modeled and observed (retrieved from PMF analysis) means and standard deviations of SOA concentrations (a) and for summer campaigns (b) and winter campaigns (c) (Table S2).




Figure 15: Modeled relative and absolute ASOA and BSOA fractions for summer (left-panels) and winter (right-panels) for different years (between 2000 and 2010) and seasons of the year (Table S2).





Figure 16: Modeled relative trends in isoprene and monoterpene emissions for the PT period (1990–2010, first and second column) and biogenic and anthropogenic SOA relative trends for the PT period (1990–2010, third and fourth column) as predicted by all the models (rows; from top to bottom: EMEP MSC-W, CHIMERE, MATCH, LOTOS-EUROS, MINNI). White areas indicate non-significant trends. Scale was saturate at 100% to facilitate the comprehension of the panel. Grey panels indicate missing data.

5



5 **Figure 17:** Modeled relative trends for the biogenic emissions (isoprene and terpene, left panel) and anthropogenic and biogenic SOA concentrations for the different PRUDENCE zones (Figure 1) for the 1990–2010 (PT) period. The columns show the averages (over land) of all the model estimates and the bars show the standard deviation respect to models.



Table 1: Chemical, thermodynamic schemes and biogenic emission models used by the modeling teams in the EURODELTA-Trends experiment.

Model	Gas-phase chemistry	SIA Module	SOA module	VBS for aerosol	Biogenic model
CHIMERE (model version 2017β v1.0)	MELCHIOR2 (Derognat et al., 2003)	ISORROPIA v2.1 (Nenes et al., 1999)	H ₂ O (Couvidat et al., 2012) coupled with SOAP (Couvidat and Sartelet, 2015)	Not used in this study	MEGAN v2.1 (Guenther et al., 2012)
EMEP MSC-W (model rv.4.7, spring 2015)	EmChem09 (Simpson et al., 2012)	MARS (Binkowski and Shankar, 1995)	VBS-NPAS (Simpson et al., 2012)	Yes (Bergström et al., 2012)	Based upon maps of 115 tree species from (Koeble and Seufert, 2001)
LOTOS-EUROS (model version 1.10.005)	TNO-CBM-IV (Schaap et al., 2009)	ISORROPIA II (Fountoukis and Nenes, 2007)	Not used in this study	Not used in this study	Based upon maps of 115 tree species from (Koeble and Seufert, 2001)
MATCH (model version April 2016)	Based on EMEP MSC-W (Simpson et al., 2012) with modified isoprene chemistry (Carter, 1996; Langner et al., 1998)	RH and T dependent equilibrium constant (Mozurkewich, 1993)	Similar to VBS-NPNA (Bergström et al., 2012)	Yes (Bergström et al., 2012)	Based upon maps of 115 tree species from (Koeble and Seufert, 2001)
MINNI (model version 4.7)	SAPRC99 (Carter, 2000)	ISORROPIA v1.7 (Nenes et al., 1998)	SORGAM module (Schell et al., 2001)	None	MEGAN v2.04 (Guenther et al., 2006)



Table 2: Relative and absolute trends in emissions of SO_x, NO_x, NH₃ and NMVOCs in the EURODELTA-Trends exercise (whole domain). Trends are reported for the entire 1990–2010 period as well as for two sub-periods, 1990–2000 and 2000–2010. The linear trends were calculated using the Theil-Sen method (Sen, 1968).

	1990–2000 (P1)		2000–2010 (P2)		1990–2010 (PT)	
	Total relative change (%)	Absolute change per year (ktons y ⁻¹)	Total relative change (%)	Absolute change per year (ktons y ⁻¹)	Total relative change (%)	Absolute change per year (ktons y ⁻¹)
SO _x	-54	-1 952	-37	-668	-69	-1 061
NO _x	-25	-659	-17	-356	-39	-510
NH ₃	-19	-129	-6	-31	-15	-45
NMVOCs	-33	-812	-33	-525	-59	-705



Table 3: Modeled and observed mean relative trends of NO₂ and SO₂ for the P1 (1990–2000), P2 (2000–2010) and PT (1990–2010) periods and percentage of points in Figure 3 within a factor of 2 of the observed trends.

NO ₂	P1 (%)	P2 (%)	PT (%)	P1 (% of points within a factor of 2)	P2 (% of points within a factor of 2)	PT (% of points within a factor of 2)
Obs	-25	-12	-36	-	-	-
EMEP MSC-W	-19	-22	-44	56	56	52
CHIMERE	-23	-25	-47	52	48	56
MATCH	-20	-26	-46	64	52	52
LOTOS-EUROS	-21	-22	-46	48	48	52
MINNI	-19	-24	-44	52	56	56
SO ₂	P1 (%)	P2 (%)	PT (%)	P1 (% of points within a factor of 2)	P2 (% of points within a factor of 2)	PT (% of points within a factor of 2)
Obs	-82	-47	-97	-	-	-
EMEP MSC-W	-76	-54	-97	100	83	100
CHIMERE	-69	-34	-91	97	63	100
MATCH	-67	-48	-88	100	83	100
LOTOS-EUROS	-69	-40	-88	97	67	100
MINNI	-64	-41	-84	97	80	100

**Table 4:** Same as Table 3 but for SO₄²⁻, TNH_x and TNO₃.

SO ₄ ²⁻	P1 (%)	P2 (%)	PT (%)	P1 (% of points within a factor of 2)	P2 (% of points within a factor of 2)	PT (% of points within a factor of 2)
Obs	-57	-14	-66	-	-	-
EMEP MSC-W	-49	-24	-67	95	65	100
CHIMERE	-54	-33	-73	95	60	100
MATCH	-47	-28	-67	95	65	100
LOTOS-EUROS	-42	-23	-61	95	65	100
MINNI	-54	-35	-78	90	65	100
TNH _x	P1 (%)	P2 (%)	PT (%)	P1 (% of points within a factor of 2)	P2 (% of points within a factor of 2)	PT (% of points within a factor of 2)
Obs	-28	-22	-46	-	-	-
EMEP MSC-W	-15	-14	-27	44	75	75
CHIMERE	-23	-14	-31	44	75	75
MATCH	-19	-21	-35	44	81	81
LOTOS-EUROS	-16	-11	-29	25	69	63
MINNI	-26	-13	-38	56	81	75
TNO ₃	P1 (%)	P2 (%)	PT (%)	P1 (% of points within a factor of 2)	P2 (% of points within a factor of 2)	PT (% of points within a factor of 2)
Obs	-16	-19	-29	-	-	-
EMEP MSC-W	-18	-25	-38	62	85	54
CHIMERE	-17	-8	-28	46	38	54
MATCH	-18	-27	-39	46	77	54
LOTOS-EUROS	-19	-17	-37	46	54	54
MINNI	-16	-14	-28	31	54	54



Table 5: Means and standard deviations of the modelled relative changes in HNO₃ and NO₃⁻ concentrations for the P1 (1990–2000) and P2 (2000–2010) periods for all the PRUDENCE regions (Figure 1).

Regions	1990–2000 (P1)		2000–2010 (P2)	
	HNO ₃ Relative change (±SD) (%)	NO ₃ ⁻ Relative change (±SD) (%)	HNO ₃ Relative change (±SD) (%)	NO ₃ ⁻ Relative change (±SD) (%)
AL	-37 (±2)	-19 (±3)	-28 (±4)	-23 (±8)
BI	-44 (±7)	-21 (±12)	-28 (±7)	-19 (±4)
BX	-41 (±5)	-18 (±9)	-20 (±5)	-20 (±7)
EE	-34 (±5)	-30 (±7)	-15 (±4)	-13 (±6)
FR	-36 (±5)	-21 (±4)	-20 (±8)	-12 (±7)
IP	-5 (±3)	-5 (±8)	-26 (±3)	-19 (±9)
MD	-23 (±2)	-10 (±14)	-21 (±2)	-21 (±10)
ME	-42 (±7)	-21 (±6)	-23 (±3)	-26 (±5)
SC	-24 (±4)	-5 (±7)	-19 (±3)	-16 (±12)



Table 6: Means and standard deviations of the modelled relative changes in SO₂ and SO₄²⁻ concentrations for the P1 (1990–2000) and P2 (2000–2010) periods for all the PRUDENCE regions (Figure 1).

Regions	1990–2000 (P1)		2000–2010 (P2)	
	SO ₂ Relative change (±SD) (%)	SO ₄ ²⁻ Relative change (±SD) (%)	SO ₂ Relative change (±SD) (%)	SO ₄ ²⁻ Relative change (±SD) (%)
AL	-69 (±5)	-56 (±4)	-57 (±7)	-41 (±2)
BI	-66 (±4)	-49 (±6)	-65 (±4)	-43 (±6)
BX	-69 (±4)	-52(±9)	-56 (±4)	-37 (±8)
EE	-67 (±4)	-52 (±5)	-31 (±6)	-30 (±4)
FR	-70 (±6)	-53 (±5)	-46 (±7)	-39 (±4)
IP	-24 (±3)	-34 (±5)	-63 (±4)	-49 (±5)
MD	-48 (±5)	-44 (±5)	-38 (±3)	-34 (±3)
ME	-85 (±4)	-59 (±8)	-35 (±8)	-33 (±4)
SC	-65 (±6)	-45 (±5)	-33 (±12)	-36 (±3)



Table 7: Observed and predicted SOA concentrations, averaged over all sites (Table S2 and Figure S1), and for the summer and winter campaigns. Statistics are normalized mean bias (NMB), normalized mean error (NME), Mean Bias (MB) and mean absolute gross error (MAGE).

		Mean Predicted ($\mu\text{g m}^{-3}$)	Mean Observed ($\mu\text{g m}^{-3}$)	NMB (%)	NME (%)	MB ($\mu\text{g m}^{-3}$)	MAGE ($\mu\text{g m}^{-3}$)
Winter	EMEP MSC-W	0.21	1.98	-0.90	0.90	-1.78	1.78
	CHIMERE	0.12	1.98	-0.94	0.94	-1.87	1.87
	MATCH	0.12	1.98	-0.94	0.94	-1.86	1.86
	MINNI	0.29	1.98	-0.85	0.85	-1.69	1.69
Summer	EMEP MSC-W	1.42	2.29	-0.38	0.44	-0.87	1.00
	CHIMERE	0.85	2.29	-0.63	0.63	-1.43	1.43
	MATCH	0.25	2.29	-0.89	0.89	-2.04	2.04
	MINNI	0.52	2.29	-0.77	0.77	-1.76	1.76
All Periods	EMEP MSC-W	0.72	2.03	-0.65	0.73	-1.31	1.48
	CHIMERE	0.39	2.03	-0.81	0.84	-1.64	1.70
	MATCH	0.18	2.03	-0.91	0.91	-1.85	1.85
	MINNI	0.57	2.03	-0.72	0.79	-1.46	1.60