

## 1 Letter to editor:

Dear Editor,

we want to thank you for agreeing to be the editor for our manuscript. After looking at all the reviews, we answered point by point all their questions. We found additional issues in the manuscript in the references (double counted), which we fixed. We added a Figure, Fig.1, which we felt was necessary as pointed out by the reviewers. We also corrected a small typo in Eq.4, where the density of ice and water should be the density of the ocean (note this correction does not appear in the annotated diff manuscript, but in the final revision of the manuscript, as we could not get Latex Diff to handle mathematical formulas). We hope you will find our corrections and comment amenable to publication. Feel free to request further clarifications.

Best regards,

Eric Larour and co-authors.

# 1 Response to Anonymous Referee #1

The authors present a framework to produce probabilistic sea-level projections which allows to incorporate and update contributions from different models of the individual drivers in a flexible, yet consistent, way. It builds on the ISSM-SESAW sea level solver by Adhikari et al. (2016), which was also presented in this journal and which is based on an unstructured mesh, allowing high accuracy at very limited computational costs. The manuscript is clear, it reads well and it presents enough graphics to show the working and capabilities of the software package. Below, I am suggesting a number of minor improvements, listed in the order as they appear in the manuscript.

We thank the referee for the time spent reviewing the manuscript, and for the positive assessment of the manuscript and the methodology implemented in the ISSM-SLPS projection capability. We have tried to address all the concerns raised, as well as present the changes that will be implemented in the amended manuscript to be submitted once the editor requests.

- Line 60: The argument about the time invariancy of fingerprints does not seem very strong, since it is always possible to increase the number of fingerprints for a given source, with the option of using independent scaling factors for each sub-source. Besides, the authors do not seem to demonstrate the effect of time-variable fingerprints on sea level projections. The improvement could be from a computational point of view (due to, e.g., the modularity of the approach), but then it could be explained more clearly.

We agree with the referee that it is always possible to increase the number of fingerprints for a given source, and of using several across time too. However, doing so converges towards a solution that we effectively generalized in our framework, where we can handle any type of thickness change pattern at whichever resolution is required, and at whatever time resolution is optimal. We agree with the reviewer that we have not in the manuscript shown the effect of time variable fingerprints explicitly, though we did compute temporally variable fingerprints to generate Fig. 8. Fig. 1 of the present document shows the impact of temporarily variable fingerprints using an ISMIP6 model projection of ice thickness changes in Greenland. The figure clearly demonstrates the significant impact between normalized RSL fingerprints at 2016 vs 2045 and 2075, with significantly different spatial patterns down to Morocco or Cuba, and over the entire arctic. We will add this figure to the manuscript, as it quantifies clearly the need for temporally variable fingerprints, and we will add the following text in the manuscript:

*An example of the inadequacy of temporally constant fingerprints is shown in Fig. 1 for a projection of Greenland's contribution to RSL at 2016 vs 2045 and 2075. Normalized RSL patterns are clearly different between 2016 and 2045, and the differences are not just local to Greenland, but spill over into North Europe, Alaska, the Canadian arctic, etc...*

- Line 65: Not sure what “profound geometry” means, it is possibly a typo.  
We will replace ”profound geometry” with ”pronounced spatial pattern”, which was the original intended meaning.
- Line 124: Could you add a reference about the “alternative approach” described here?  
Thank you for the suggestion. We will add Thompson et al. (2016) (Fig.3) as an example of this approach.
- Line 148: I guess the term STR refers to a global mean value, not a local one as stated, since  $RSL_{STR}$  only depends on time.  
We thank the reviewer for catching this, indeed STR refers to the global mean value. We will correct the manuscript accordingly.
- Line 168: I am surprised by the fact that  $RSL_{GRD}$  includes viscoelastic deformation of the solid earth, rather than just elastic as in most studies of this kind. If this is the case, it should be highlighted in the introduction.  
We thank the reviewer for spotting this. Indeed it should be explicitly noted that visco-elastic deformation is currently allowed in the projection system. Here, we define visco-elastic deformation any deformation that is not related to GIA, but to smaller time-scale loading processes (50 to 100 years) as observed in West Antarctica Barletta et al. (2018). In practice, this type of deformation is not yet handled in ISSM, but will in the future. We will make the following explicit comment in the manuscript at line 151: *This implies that viscoelastic deformation is split between long-term time scales and short-term fast rebound of the bedrock uplift, such as observed in West Antarctica (Barletta et al., 2018), acting essentially over time scales of 50-100 years.*
- Line 215: I suggest adding that the number of elements is significantly smaller than in the case of an equi-angular 1x1-deg grid, which would already be rather coarse.  
We thank the reviewer for the suggestion, indeed the equi-angular grid at 1x1 deg would require 64,800 vertices, with our mesh standing at only 19,486. We will add this to the manuscript.
- Lines 225-241: I am not sure whether showing a few lines of code is normal in GMD, but, if possible, it would be nice to replace this by a flow chart.  
We understand the reviewer’s point of view. We have debated this extensively, having in the past presented pieces of code in GMD. However, this is the main algorithm core, and we believe it should be explicitly and exactly described. The goal here is to avoid any confusion that could arise from a flowchart. We respectfully ask to keep the code as is, but will defer to the editor’s decision.
- Line 299: It seems a few words are missing, since Fig.8b shows GMSL values, while values for 9 cities around the world are shown in Fig.9. In addition, the caption of Fig.9 does not mention the timeframe for which the projections are generated.  
We thank the reviewer for pointing this redirection to the wrong figure. We

will correct the manuscript accordingly, and add a quick description of Fig 8b in the text as showing the evolution of GMSL's PDF through time. We will also add the time frame in Fig. 9.

- Line 304: I suggest also mentioning that the width of the two PDFs shown in Fig.10 is different, rather than only the mean, because it makes an even stronger point about the benefit of this approach.  
We thank the reviewer for the suggestion. After checking though, the PDFs standard deviations are only different by .4% on a relative basis. The PDFs do indeed appear different at their basis, but we believe this is a visual effect from the color choice. We will put a comment to this effect in the caption of Fig. 10 to dispel any confusion.
- Line 316: “impact” (spelling).  
Thank you for pointing out the typo, we will correct the manuscript accordingly.
- Line 338: The importance of “geodetically compliant patterns”, in spite of being part of the manuscript title, is nowhere really discussed nor explained. From this line, my understanding is that it refers to the possibility of defining the mesh in such a way that it matches the location of specific geodetic observations, hence avoiding unnecessary interpolations. In any case, the issue warrants a more extensive discussion, possibly in the introduction.  
We thank the reviewer for the comment. Geodetically compliant referred to the fact that ISSM-SLPS computes spatio-temporal GRD patterns throughout the projection sampling, which is not always done consistently for other similar frameworks. We will add comments in the introduction referencing this explanation explicitly. In particular, we will add the following at line 117: *“The latter feature builds the basis for a geodetically compliant projection system where GRD patterns and their computations is done systematically and does not introduce biases in the RSL projections.”*

## References

- Barletta, V. R., Bevis, M., Smith, B. E., Wilson, T., Brown, A., Bordoni, A., Willis, M., Khan, S. A., Rovira-Navarro, M., Dalziel, I., Smalley, R., Kendrick, E., Konfal, S., Caccamise, D. J., Aster, R. C., Nyblade, A., and Wiens, D. A.: Observed rapid bedrock uplift in Amundsen Sea Embayment promotes ice-sheet stability, *Science*, 360, 1335–1339, 2018.
- Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., Lipscomb, W. H., Gregory, J., Abe-Ouchi, A., Shepherd, A., Simon, E., Agosta, C., Alexander, P., Aschwanden, A., Barthel, A., Calov, R., Chambers, C., Choi, Y., Cuzzone, J., Dumas, C., Edwards, T., Felikson, D., Fettweis, X., Golledge, N. R., Greve, R., Humbert, A., Huybrechts, P., Le clec'h, S., Lee, V., Leguy, G., Little, C., Lowry, D. P., Morlighem, M., Nias, I., Quiquet, A., Rückamp, M., Schlegel, N.-J., Slater, D., Smith, R., Straneo, F., Tarasov, L., van de Wal, R., and van den Broeke, M.: The future sea-level contribution of the Greenland ice sheet: a multi-model ensemble study of ISMIP6, *The Cryosphere*, <https://doi.org/10.5194/tc-2019-319>, URL <https://www.the-cryosphere-discuss.net/tc-2019-319/>, 2020, accepted.
- Thompson, P. R., Hamlington, B. D., Landerer, F. W., and Adhikari, S.: Are long tide gauge records in the wrong place to measure global mean sea level rise?, *Geophysical research letters*, 2016.




Figure 1: Normalized fingerprints for Greenland at 2016, 2045 and 2075, based on the JPL ISSM experiment 5 simulation contributed to ISMIP6 (Goelzer et al., 2020, accepted). Variations in ice-thickness change patterns are significantly different between the East and West coast of Greenland, and along a South-West gradient too, resulting in significantly different contributions to local RSL at different time snapshots

## 1 Response to Anonymous Referee #2

The authors developed a new method to account for feedbacks between ice sheets, ocean circulation and solid-Earth deformation through dynamically computed sea-level fingerprints for probabilistic projections of local sea-level change. This is a significant and timely contribution because current probabilistic projections use static fingerprints and cannot readily incorporate advances in forward models. The manuscript is well-structured and provides clear examples of how the SLPS works. However, I think the clarity of the manuscript could be improved with some (minor) modifications of the text and figures, see the comments below.

We thank the referee for the time spent reviewing the manuscript and for the positive assessment of the manuscript, in particular the capability of ISSM-SLPS to readily update projections with new forward modeling advances that are tightly coupled with the framework. We address below all the concerns from the referee, and present future changes to the manuscript that will be implemented if the editor moves forwards with accepting corrections.

- L4: For long-term projections (. . .) that provide such probabilistic projectionstive, can this be rewritten?  
*Indeed this is too heavy, we will reformulate to "For a time horizon of 100 years, frameworks have been developed that provide such projections by relying on ..."*
- L8: solid-Earth ‘uplift’ – displacement or deformation would be more complete?  
*We thank the reviewer for the comment, but believe that solid-Earth uplift is a terminology used throughout the Cryosphere/solid-Earth community that has a very specific meaning. Deformation could convey lateral motion too, as well as displacement. Our focus here is on deformation that impacts RSL through VLM in particular. We respectfully would like to keep the terminology as is.*
- L26: why not give examples of the use of projections globally rather than so many references for just the US?  
*We understand the concern of the reviewer, however, here the intent of the manuscript was to point out to how widely the KOPP14 framework has been adopted. For assessment outside the US, the onus is significantly more on the IPCC assessments, which are arguably, given their 4 year cycle, not as responsive to more recent developments in the science community. Given that our intent was focused on the KOPP14 framework we would like to respectfully request to keep ours references as is.*
- L37 & 118: sterodynamic sea-level change is defined incorrectly here, see Gregory et al. (2019). It is the sum of global (not local) thermosteric expansion and ocean dynamics (which include the local steric effect, both thermo- and halosteric) including the IB effect. Can the authors clarify

this?

Another referee #1 pointed to the same issue, and we thank you for spotting this issue. The definition for STR and DSL for our manuscript indeed follows Gregory et al. (2019). The confusion came from the erroneous use of "local" instead of "global" at line 37. The sentence will now read "*which is the sum of globally averaged thermosteric expansion and local sea-level changes due to ocean dynamics (which include the local steric effect, both thermosteric and halosteric).*"

- L38 & 55: Not just ESMs but also AOGCMs  
Dually noted, and the manuscript will be updated accordingly
- L47: Do you mean Kopp et al. (2017) here instead of (2014)?  
Indeed that is what we meant, thank you for spotting the typo, will be reflected in the amended manuscript.
- L64-66: Can the authors comment on the expected importance of geometry changes to 21st century sea-level projections?  
Referee #1 had a similar comment referring to the fact that we did not demonstrate the importance of time variable fingerprints. We will add an example figure (Fig. 1) of the evolution of  $RSL_{GRD}$  for the Greenland Ice Sheet, using a projection from ISMIP6 (Goelzer et al., 2020, accepted) based on the ISSM JPL run for experiment 5. The nominal fingerprints are shown to be significantly different between 2015 and 2100. We refer the reader to the response to referee #1 for the figure, comments, and corresponding changes to the manuscript that address the present comment also.
- L83: 'results such as (. . .) results' - please rewrite  
We will rewrite this sentence to "*In order to be able to account for strong couplings, or to even be able to ingest recent modeling results, one needs to propagate the local mass changes and the associated uncertainties into regional sea-level projections. This is particularly relevant now given new modeling runs that have been carried out within large Modeling Intercomparison Projects (MIPs) such CMIP5 and CMIP6, as well as ISMIP6 or GlacierMIP2.*"
- L91: 'higher-frequency' higher than what? Perhaps use 'high-frequency' instead?  
Thank you for the suggestion. We will replace to "high-frequency", and better define what is meant, with frequencies of interest being daily to monthly.
- L97-103: this paragraph misses a final sentence placing its content in context of the manuscript  
We agree with the reviewer, and will add the following sentence at the end of the paragraph "*All these advances need to be fully integrated into new*



*probabilistic projections of sea-level change, and a new approach therefore needs to be envisioned that will allow for such new processes to be accurately modeled.”*

- L104-107: this sentence doesn't read very well, can you please split this up in smaller sentences?

*We agree with the reviewer. Actually, the whole paragraph will be recast to "Indeed, moving from strategies where continental scale mass changes are sampled and multiplied with the corresponding fingerprint, to actually sampling upstream model inputs is paramount to improving the state of the art. In particular, there is a strong need to fully account for spatial patterns of mass change and their uncertainty (see e.g. Fig. 1b-d), This applies to among others SMB, basal friction, or ice and solid-Earth rheological properties."*

- L114-115: 'and through . . . individual sample' misses a verb, perhaps reverse the order with 'retaining the . . .from Eq. 1'

*We will rephrase the paragraph to "We improve the existing process-based approach by using the Ice-Sheet and Sea-Level System Model (ISSM, Larour et al, 2012c) which allows for inclusion of forward model physics. It also improves the modeling and sampling of covariances between input processes, both temporally and spatially through the computation of high-resolution barostatic-GRD patterns. "*

- Figure 3: the axis labels have fallen off the figure.

*Thank you for spotting this issue, will be fixed in the amended manuscript*

- Figure 5: It is very hard to distinguish colors of cells within the mesh, could this be improved by for example adapting the range of the colorbar?

*We thank the reviewer for the suggestion. By saturating the colorbar, we reach better contrast. We will do so for the manuscript, and provide an improved figure. We will also amend the caption to explain why we saturated the colorbar.*

- L253: has a typo 'approach'

*Thank you for spotting the typo, will be corrected in the manuscript*

- L291-292: "DSL is not sampled, but rather deterministically set to the DSL term of the CMIP5 NorESM-ME runs" – why? please explain

*This is due to the uncertainty in the quality of the CMIP5 runs in terms of global mean thermosteric contribution. We preferred to avoid this uncertainty and concentrate on the geodetically relevant components given the scope of the manuscript.*

- L299: refers to figure 9 instead of 8, and discussion of Figure 9 is missing?

*We refer to referee #1 comments, which also spotted this issue.*

- L316: "inpact"

*Thank you for the typo, will be corrected in the manuscript.*

- L343: 'urther'

Thank you for spotting the typo, will be corrected in the manuscript.:w

## References

- Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., Lipscomb, W. H., Gregory, J., Abe-Ouchi, A., Shepherd, A., Simon, E., Agosta, C., Alexander, P., Aschwanden, A., Barthel, A., Calov, R., Chambers, C., Choi, Y., Cuzzone, J., Dumas, C., Edwards, T., Felikson, D., Fettweis, X., Golledge, N. R., Greve, R., Humbert, A., Huybrechts, P., Le clec'h, S., Lee, V., Leguy, G., Little, C., Lowry, D. P., Morlighem, M., Nias, I., Quiquet, A., Rückamp, M., Schlegel, N.-J., Slater, D., Smith, R., Straneo, F., Tarasov, L., van de Wal, R., and van den Broeke, M.: The future sea-level contribution of the Greenland ice sheet: a multi-model ensemble study of ISMIP6, The Cryosphere, <https://doi.org/10.5194/tc-2019-319>, URL <https://www.the-cryosphere-discuss.net/tc-2019-319/>, 2020, accepted.
- Gregory, J., Griffies, S., Hughes, C., et al.: Concepts and Terminology for Sea Level: Mean, Variability and Change, Both Local and Global, *Surv. Geophys.*, 40, 1251–1289, 2019.

# 1 Response to Anonymous Referee #3

A review of "ISSM-SLPS: geodetically compliant Sea-Level Projection System for the Ice-sheet and Sea-level System Model v4.17" by Larour et al. for possible publication in Geoscientific Model Development.

The authors present a new geodetically compliant approach for modeling future sea-level rise due to ocean thermal expansion, ocean circulation changes, water mass redistribution, and glacial isostatic adjustment. This new approach has the relative advantage compared to previous approaches (e.g., Kopp et al. 2014, 2017) that coupling and interaction between contributors are taken into account. The authors highlight the important result that, by modeling Greenland ice mass loss using 18 basins rather than 1 basin, uncertainties on future sea-level rise are substantially reduced (Figure 11).

I'll confess that, while I study sea level and was invited to review the paper, I'm a physical oceanographer. I don't have the expertise in modeling, geodesy, or glaciology needed to give a thorough review of this paper. I'd strongly recommend the editor to ensure experts in these topical areas weigh in on this paper. That being said, I appreciate the paper. I think it's really valuable that the authors are pushing the envelope and developing flexible, modular, coupled approaches to model the various contributors to future sea-level rise and their uncertainties.

I have no major issues with the manuscript (though, again, I strongly recommend more expert reviewers weigh in). I have a couple editorial remarks, detailed below. My only one real complaint regards terminology. The authors' use of gravitational, rotational, and deformational (GRD), steric, and barystatic sea-level contributions (cf. Equation 1) can be inconsistent with definitions in Gregory et al. (2019). I'd recommend the authors either (1.) adopt the definitions used in Gregory et al. (2019) or (2.) acknowledge where their definitions diverge from Gregory et al. (2019) to avoid confusion.

We thank the reviewer for the time spent on the review and for his valuable insights, especially from the Physical Oceanography point of view. We agree with the reviewer's assessment of the advantage of flexible, modular, coupled approaches to model various contributors to future sea-level rise and their uncertainties. We also agree that our initial explanation of the STR and DSL terms were not compatible with the definitions in Gregory et al. (2019), and will definitely tighten the introduction in this respect. All other referees pointed to the same issue (see in particular referee #1). We below go through all the comments and try and address them, along with modifications to the manuscript that will be carried out if the editor goes forward with requesting a new version.

Specific comments:

- Line 2: paramount to – > important for  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 4: cost and timing – > cost, timing, and risk tolerance

Thank you for the suggestion, we will adopt it in the new manuscript

- Line 23: are summed – > are modeled separately and summed  
Thank you for the suggestion, we will adopt it in the new manuscript
- Equation 1. Comparing to Figure 3 in Gregory et al. (2019), I'm confused by this equation. The equivalent equation in Gregory et al. would be: relative sea level = stereodynamic sea level + gravitation, rotation, deformation (GRD) + barystatic - inverted barometer  
In Gregory et al. (2019), GRD includes GIA, and GRD makes no contribution to global mean sea-level changes. What the authors here call "GRD", Gregory et al. (2019) call "contemporary GRD". Anyway, it's fine that the authors here use slightly different terminology. But they should acknowledge where their definitions diverge from Gregory et al. (2019). Otherwise, readers (i.e., I) will get confused.  
We really appreciate the referee checking against Gregory et al. (2019) and seeing this inconsistency. We will definitely remark in the manuscript on the differences in our approach, and refer to our GRD as contemporary GRD. Here is the new paragraph that will be in the manuscript starting at line 37: *Note here that our definition of GRD is not completely in line with Gregory et al. (2019), as GIA is considered as a separate contributor, and the GRD contribution does contribute to global mean sea-level changes. It is rather in line with the definition of contemporary GRD in Gregory et al. (2019).*
- Line 37: local thermosteric – > global-mean thermosteric  
This was picked up by all referees, and can lead to confusion about the definition of STR and DSL. Thank you for spotting it, we refer to the referee #1 comments on how we addressed this
- Line 53: qualities – > quantities  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 61: "stays constant in time" it's unclear what the authors mean by this phrase  
We will replace by *is constant through time*
- Line 107: paramount to – > important for  
Thank you for the suggestion, we will adopt it in the new manuscript
- Lines 124-126: I'm unfamiliar with studies doing this for projection purposes. Do the authors have a reference in mind for this technique?  
Referre #1 also requested a reference. There is not one involving a projection, but we provided (eg. Thompson et al., 2016, Fig.3) for a good explanation of the approach that could readily be adapted to a projection.
- Line 127: can drive redistribution – > can be coupled to redistribution  
Thank you for the suggestion, we will adopt it in the new manuscript

- Line 128: causes – > manifests in  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 129: cause a change in the load of – > load  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 130: SAL effects. Suggest to reference, e.g., Ray (1998), Stepanov and Hughes (2004), and/or Vinogradova et al. (2015) on these points.  
Thank you for the suggestion, we will adopt it in the new manuscript and reference these studies
- Line 131: Please add ”made by atmosphere-ocean general circulation models (AOGCMs)” after projections and before the Richter et al. (2013) reference  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 148: local – > global-mean  
Thank you for catching this typo that was also important to all three other referees. We have corrected the manuscript accordingly.
- Line 152-153: The authors should clarify whether they remove the global-mean OBP value or not. If not, are the authors making the Greatbatch correction to account for the Boussinesq nature of most CMIP AOGCMs?  
Thank you for the comment. Indeed we remove the global-mean OBP from ocean models, since ocean dynamics don’t add or remove any mass from/to the ocean. The CMIP5 and CMIP6 models all (should) have applied the Greatbatch correction, as confirmed by the fact the global-mean value of the ’zos’ fields is zero, and we use the ’zostoga’ to the models to get the global thermosteric rise. We will add the following paragraph starting at line 155: *”Note also that the global-mean OBP is removed from the ocean models, since ocean dynamics don’t add or remove any mass from/to the ocean. In addition, our projections rely on CMIP5 and CMIP6 fields ’zos’ (the sea-surface height change above geoid, or DSL term) and ’zostoga’ (global average thermosteric sea-level change or STR) where the Greatbatch correction has been applied, resulting in a zero global-mean value of STR.”*
- Line 165: The authors should precisely define the ocean function  $O(\theta, \phi)$  for clarity.  
Thank you for spotting this issue, we will define the ocean function in the manuscript succinctly as  *$O=1$  for oceans and zero otherwise”*
- Line 176: The former – > These  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 193: Please define BAMG on first use  
Thank you for the suggestion, we will define BAMG in the manuscript

- Figure 3 caption: thsi – > this  
Thank you for spotting the type, we will correct it in the new manuscript
- Line 246: alphas – > alpha is  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 256-257: "We display ... the average" I don't understand this sentence, but maybe it's just me.  
This sentence can indeed be clearer, we replace with "*We display the average thinning rate  $\mu$ ,  $\mu + 3\sigma$  and  $\mu - 3\sigma$  (for an arbitrary value of the standard deviation  $\sigma = 5\%$ ).*"
- Line 262: "the KOPP14 ... SLPS framework" Unclear. Are the authors saying that the approach here reduces to and reproduces the Kopp results under certain strong assumptions? Please clarify.  
We are indeed saying that the approach is equivalent to KOPP14 if we use the same partitioning. The assumptions are not so strong, just that the partitioning be the same. However, as demonstrated by Fig. 11, this is not the case anymore once several basins are introduced. We will try and capture this better in the manuscript, with the following statement: "*Once several partitions are adopted however, the refinement in the fingerprint patterns significantly departs from the KOPP14 approach.* "
- Line 289: please add spaces between -1.65, sigma, to, 1.65, sigma  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 291-292: "DSL is not ... CMIP5 NorESM-ME runs" Why only one model and why this model? Variance in model projections of DSL changes can be large and important locally.  
In this demonstration of the capabilities of ISSM-SLPS, we wanted to approach the geodetic angle. Significant variance in model projections are indeed found in the CMIP5 and CMIP6 benchmarks, which completely occultate any other variance from any other inputs. We wanted to avoid this. We ask the reviewer to allow for this exception, as we believe it leads to a better validation of our capability.
- Line 303: "Bayesian exploration approach" The authors reference such an approach several times, but never explain it or give a reference.  
We agree with the reviewer. The GIA statistics relied upon here are from Caron et al. (2018), but the bayesian framework we refer to is described in Caron et al. (2017) and is based on a bayesian inversion method using Simulated Annealing (Kirkpatrick et al., 1983), a variation of the Monte Carlo with Markov chains (MCMC) method (Metropolis and Ulam, 1949; Metropolis et al., 1953). We will better refine the description in the manuscript and give extended citations. The paragraph will now read *These statistics were evaluated using bayesian inversion method based on Simulated Annealing (Kirkpatrick et al., 1983), a variation of the Monte Carlo with Markov chains (MCMC) method (Metropolis and Ulam, 1949;*

*Metropolis et al., 1953). They can be used directly in SLPS, either during a standard probabilistic projection run, or a posteriori as is the case here. These statistics reflect the statistical fitness to a global GIA dataset composed of paleo-RSL indicators and vertical GPS trends.*

- Line 311: "the tails are much larger for the 1 basin scenario" This seems like a very important results, but I don't think the authors have discussed it enough for me to understand physically why this is the case. Suggest to consider adding more of a description.

We agree with the reviewer. The reason for reduced tails is that by multiplying the number of basins, we recompute fingerprints that are more reflective of the true spatial pattern. The example of New York used in Larour et al. (2017) helps in understanding this feature: the entire South-East Greenland contributes zero sea-level change in NY. If a basin is positioned over this entire region, it will contribute zero variance to the PDF distribution for SLR in NY. This leads to a reduction in the tails of the distribution. We will add this explanation in the manuscript too, as suggested. Here is the text we will add starting at line 317: *" This can be visualized better by taking the example of New York, where following Larour et al. (2017) contributions from South Greenland are almost negligible. This implies that all the basins (and corresponding GRD patterns) in South Greenland will contribute zero variance to the PDF for RSL at New York. This will therefore result in smaller tails for projections that rely on more refined basins."*

- Line 311: the "likely" (5-95%) range – > the width of the "likely" (5-95%) range  
Thank you for the suggestion, we will adopt it in the new manuscript
- Line 325: siginificantly – > significantly  
Thank you for spotting the typo, we will correct the manuscript accordingly
- Line 343: urther – > further  
Thank you for spotting the typo, we will correct the manuscript accordingly



## References

- Caron, L., Métivier, L., Greff-Lefftz, M., Fleitout, L., and Rouby, H.: Inverting Glacial Isostatic Adjustment signal using Bayesian framework and two linearly relaxing rheologies, *Geophysical Journal International*, 209, 1126–1147, 2017.
- Caron, L., Ivins, E. R., Larour, E., Adhikari, S., Nilsson, J., and Blewitt, G.: GIA Model Statistics for GRACE Hydrology, Cryosphere, and Ocean Science, *Geophysical research letters*, 45, 2203–2212, 2018.
- Gregory, J., Griffies, S., Hughes, C., et al.: Concepts and Terminology for Sea Level: Mean, Variability and Change, Both Local and Global, *Surv. Geophys.*, 40, 1251–1289, 2019.
- Kirkpatrick, S., Gelatt, C. D., and Vecchi, M. P.: Optimization by Simulated Annealing, *Science*, 220, 671–680, <https://doi.org/10.1126/science.220.4598.671>, 1983.
- Larour, E., Ivins, E. R., and Adhikari, S.: Should coastal planners have concern over where land ice is melting?, *Science Advances*, 3, e1700537, 2017.
- Metropolis, N. and Ulam, S.: The Monte Carlo method, *J. Amer. Stat. Associ.*, 44, 335–341, 1949.
- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., and Teller, E.: Equation of State Calculations by Fast Computing Machines, *The Journal of Chemical Physics*, 21, 1087–1092, <https://doi.org/10.1063/1.1699114>, 1953.
- Thompson, P. R., Hamlington, B. D., Landerer, F. W., and Adhikari, S.: Are long tide gauge records in the wrong place to measure global mean sea level rise?, *Geophysical research letters*, 2016.

# ISSM-SLPS: geodetically compliant Sea-Level Projection System for the Ice-sheet and Sea-level System Model v4.17

Eric Larour<sup>1</sup>, Lambert Caron<sup>1</sup>, Mathieu Morlighem<sup>3</sup>, Surendra Adhikari<sup>1</sup>, Thomas Frederikse<sup>1</sup>, Nicole-Jeanne Schlegel<sup>1</sup>, Erik Ivins<sup>1</sup>, Benjamin Hamlington<sup>1</sup>, Robert Kopp<sup>2</sup>, and Sophie Nowicki<sup>4</sup>

<sup>1</sup>Jet Propulsion Laboratory - California Institute of technology, 4800 Oak Grove Drive MS 300-323, Pasadena, CA 91109-8099, USA

<sup>2</sup>Department of Earth and Planetary Sciences and Institute of Earth, Ocean and Atmospheric Sciences, Rutgers University, New Brunswick, NJ, USA

<sup>3</sup>University of California at Irvine, Department of Earth System Science, Irvine, California, USA

<sup>4</sup>NASA Goddard Space Flight Center, Cryospheric Sciences Lab, Greenbelt, Maryland, USA

**Correspondence:** Eric Larour (eric.larour@jpl.nasa.gov)

## Abstract.

Understanding future impacts of sea-level rise at the local level is [\[..<sup>1</sup>\]](#) important for mitigating its effects. In particular, quantifying the range of sea-level rise outcomes in a probabilistic way enables coastal planners to better adapt strategies, depending on cost [\[..<sup>2</sup>\]](#), [timing and risk tolerance](#). For a time horizon of 100 years, frameworks have been developed that provide such [\[..<sup>3</sup>\]](#) projections by relying on sea-level fingerprints where contributions from different processes are sampled at each individual time step and summed up to create probability distributions of sea-level rise for each desired location. While advantageous, this method does not readily allow for including new physics developed in forward models of each component. For example, couplings and feedbacks between ice sheets, ocean circulation, and solid-Earth uplift cannot easily be represented in such frameworks. Indeed, the main impediment to inclusion of more forward model physics in probabilistic sea-level frameworks is the availability of dynamically computed sea-level fingerprints that can be directly linked to local mass changes. Here, we demonstrate such an approach within the Ice-Sheet and Sea-level System Model (ISSM), where we develop a probabilistic framework that can readily be coupled to forward process models such as those for ice sheets, glacial-isostatic adjustment, hydrology and ocean circulation, among others. Through large scale uncertainty quantification, we demonstrate how this approach enables inclusion of incremental improvements in all forward models and provides fidelity to time-correlated processes. The projection system may readily process input and output quantities that are geodetically consistent with space and terrestrial measurement systems. The approach can also account for numerous improvements in our understanding of sea-level processes.

---

<sup>1</sup>removed: paramount to

<sup>2</sup>removed: and timing. For long-term projections, from present-day to the end of the 21st century

<sup>3</sup>removed: probabilistic projections. They rely

## 20 1 Introduction

Reliable projections of local sea-level change, together with robust uncertainties, are a key quantity for stakeholders to shape adequate and cost-effective mitigation and adaptation measures to sea-level rise (Kopp et al., 2019). Most regional sea-level projections use a process-based approach, in which all relevant processes are [modeled separately and](#) summed up together, including the individual estimates of error, with their spatial signature (Slangen et al., 2012; Church et al., 2013b; Kopp et al., 2014; Jackson and Jevrejeva, 2016; Kopp et al., 2017; Jevrejeva et al., 2019). These projections are widely used by coastal planners and stakeholders, as is for example demonstrated by the impact of Kopp et al. (2014, 2017) on assessment reports across the United States (Gornitz et al., 2019; City of Boston, 2016; Kopp et al., 2016; Kaplan et al., 2016; Callahan et al., 2017; Dalton et al., 2017; Griggs et al., 2017; Miller et al., 2018; Boesch et al., 2018).

In their most simple form, these process-based projections (we generally refer to these as KOPP14, in reference to Kopp et al. (2014)) can be expressed as:

$$RSL_{\text{total}}(\theta, \phi, t) = \sum_{i=1}^n F_{\text{GRD},i}(\theta, \phi) \cdot B_i(t) + RSL_{\text{sterodynamic}}(\theta, \phi, t) + RSL_{\text{GIA}}(\theta, \phi, t). \quad (1)$$

where  $RSL_{\text{total}}(\theta, \phi, t)$  is the total projected relative sea-level (RSL) change at time  $t$ , latitude  $\theta$  and longitude  $\phi$ . For all barystatic processes, or processes that change the total ocean mass, the effects of gravity, rotation and deformation (GRD) on local sea level are computed by multiplying the total barystatic contribution  $B_i(t)$  by the associated barystatic-GRD fingerprint (abbreviated by "fingerprint" from here on), or  $F_{\text{GRD},i}(\theta, \phi)$ , which is computed a priori. This procedure is generally used to include the effects of glacier (GLA) and ice sheet mass loss, as well as for projected changes in terrestrial water storage (TWS). [Note here that our definition of GRD is not completely in line with Gregory et al. \(2019\), as GIA is considered as a separate contributor, and the GRD contribution does contribute to global mean sea-level changes. It is rather in line with the definition of contemporary GRD in Gregory et al. \(2019\).](#) The effects of sterodynamic sea-level change  $RSL_{\text{sterodynamic}}(\theta, \phi, t)$ , which is the sum of [\[..<sup>4</sup>\]global](#) thermosteric expansion and local sea-level changes due to ocean dynamics, is generally included by directly using estimates from Earth System Models (ESMs) and Atmospheric/Oceanic Global Circulation Models (AOGCMs), such as the output of the Coupled Model Intercomparison Project 5 (CMIP5, Taylor et al., 2009). Finally, the glacial isostatic adjustment (GIA) term is generally accounted for using output from a periodically updated global model.

To derive uncertainties for these local projections of sea level change, the barystatic components  $B_i$  are often sampled from a probability distribution found in published probabilistic projections, for example from expert elicitation projects (e.g. Bamber et al., 2019), or other ice-sheet models (DeConto and Pollard, 2016). The sterodynamic contribution often uses the inter-model spread as a source of the uncertainties. While the basis of each probabilistic projection is similar, each group adds additional

---

<sup>4</sup>removed: local

components and physics to Eq. 1. For example, in Kopp et al. (2014) and Kopp et al. (2017), a Gaussian Process Regression model, based on tide-gauge observations is used to account for the effect of non-climatic vertical land motion. Or in Jackson and Jevrejeva (2016) and Kopp et al. (2017), the GRD effects of ocean dynamics (Richter et al., 2013) are explicitly taken into account, with <sup>[..<sup>5</sup>]</sup> Kopp et al. (2017) computing these effects over the entire projection time series.

One of the key strengths of this approach is how simple and transparent it is, as the process from probabilistic estimates of the underlying processes into local sea-level changes is a simple multiplication operation with the respective barystatic-GRD fingerprint. It provides a framework that outputs a probability density function (PDF) for *RSL* at any desired location, from which the expected sea-level change and its confidence intervals can be derived. This provides both efficient calibration/validation <sup>[..<sup>6</sup>]</sup> quantities to projections and streamlines incrementally updated projections. In essence, each modular input may be improved separately, so updates are unencumbered by the queuing up of new modules for incorporation into more complex <sup>[..<sup>7</sup>]</sup> ESMs and AOGCMs.

Yet recently, a growing body of research indicates that additional processes should be considered in this process-based approach. Indeed, inclusion of such processes is critical to improving the quantification of uncertainties in local sea-level change predictions, but they are not directly feasible within the framework of Eq. 1. Below, we highlight some of the key contributors to uncertainty that until now, have not been considered together in large-scale estimates of sea-level change.

First, in Eq. 1, the multiplication of a barystatic mass contributor  $B_i(t)$  with a fingerprint  $F_{GRD,i}(\theta, \phi)$ , assumes that the fingerprint <sup>[..<sup>8</sup>]</sup> is constant through time, which is not always the case (Mitrovica et al., 2011). Instead, a fingerprint results from feedbacks between the geometry of sea-level components. For example local sea level depends on the geometry of ice mass loss, so temporal changes in ice geometry will directly translate into local sea-level changes (e.g. Larour et al., 2017b; Mitrovica et al., 2018). As a result, this temporal variability not only affects the expected local sea-level changes, but also its uncertainties, as the uncertainty of the input mass loss also has a <sup>[..<sup>9</sup>]</sup> pronounced spatial pattern due to relative limitations in measurement and data interpretation. An example of the inadequacy of temporally constant fingerprints is shown in Fig. 1 for a projection of Greenland's contribution to RSL at 2016 vs 2045 and 2075. Normalized RSL patterns are clearly different between 2016 and 2045, and the differences are not just local to Greenland, but spill over into North Europe, Alaska, the Canadian arctic, etc...

Second, covariances in time as well as the co-variances between the individual processes are not always negligibly small, though they are often considered to be or are approximated by a simple relationship (e.g. Church et al., 2013b). Indeed, assuming so could cause a significant misrepresentation of the estimated uncertainties in local sea-level change. For example, Le Bars (2018) showed that most driving factors of sea level are correlated with global-mean temperature changes, and ignoring this inter-process covariance can underestimate uncertainty in local sea-level change. Note that in addition to co-variances between processes, the uncertainty in individual processes may also be correlated temporally. Propagating this full spatio-

---


<sup>5</sup>removed: Kopp et al. (2014)

<sup>6</sup>removed: qualities

<sup>7</sup>removed: Earth System Models

<sup>8</sup>removed: stays constant in

<sup>9</sup>removed: profound geometry



**Figure 1.** Normalized fingerprints for Greenland at 2016, 2045 and 2075, based on the JPL ISSM experiment 5 simulation contributed to ISMIP6 (Goelzer et al., 2020, accepted). Variations in ice-thickness change patterns are significantly different between the East and West coast of Greenland, and along a South-West gradient too, resulting in significantly different contributions to local RSL at different time snapshots

temporal covariance into projections and its uncertainties promotes a better understanding of the spatial and temporal coherence

80 of uncertainties, which could, for example, allow us to assess the likelihood of reaching specific sea levels by 2100 given observed sea-level change during the next 20 years.

Thirdly, recent work on the Antarctic Ice Sheet (AIS) shows a strong coupling between GIA, elastic surface deformation and ice mass loss (Gomez et al., 2018; Barletta et al., 2018; Larour et al., 2019). Such relationships between these processes suggest that any uncertainties in computed ice-sheet histories and solid-Earth properties that propagate into GIA projections 85 (Caron et al., 2018) can also feed back into ice-mass-loss projections, thus considering these processes as independent ignores these couplings. Here, the main problem is that projection frameworks are articulated in terms of changes in mass, while most ice-sheet models, GIA models, TWS evolution models, and glacier models, are explicitly described in terms of local mass change evolution (or thickness changes, in m/yr water equivalent). In order to be able to account for strong couplings, or to even be able to ingest recent modeling results[..<sup>10</sup> ], one needs to propagate the local mass changes and the associated uncertainties 90 into regional sea-level projections. [This is particularly relevant now given new modeling runs that have been carried out within large Modeling Intercomparison Projects \(MIPs\) such CMIP5 and CMIP6, as well as ISMIP6 or GlacierMIP2.](#)


Similarly, additional strong positive feedbacks between ice sheet and ocean dynamics have been evidenced in work from among others, Goldberg et al. (2012, 2018, 2019) and Seroussi et al. (2017). Specifically, these studies suggest that strong coupling between sub ice-shelf ocean circulation (in particular melt rates) and ice-flow dynamics (in particular, grounding 95 line dynamics and mass transport resulting in modifications of an ice-shelf draft) results in significant retreat of ice streams such as Thwaites Glacier and Pine Island Glacier, as Antarctica's warm circumpolar deepwater is advected close to their grounding line. Other [..<sup>11</sup> ][high-frequency processes \(at the daily to monthly level\)](#) such as ocean tides and in particular how tidal currents affect water mass properties at ice sheet marine margins (Padman et al., 2018) are critical in understanding how mass loss rates will evolve. This will significantly impact how melt-rate parameterizations are developed to quantify 100 melt rates, especially in the West Antarctic Ice Sheet area (Seroussi et al., 2017). Significant work remains in calibrating such melt rate parameterizations to correctly account for all afore-mentioned effects. While more work is required in terms of constraining such parameterizations, the impact of such ice/ocean feedbacks have not been assessed in probabilistic sea-level models (PSLMs).

Finally, in the past decade, extensive work has been carried out to probabilistically characterize components like GIA (White- 105 house et al., 2012; Gunter et al., 2014; Caron et al., 2018; Melini and Spada, 2019) or ice-sheet mass balance (Larour et al., 2012b, a; Schlegel et al., 2013, 2015, 2016, 2018). Substantial understanding of the impact of rheological parameters and ice history on the distribution of bedrock uplift and rate of change in geoid rates has been generated through modelling of GIA. Similarly, for ice sheet models, significant knowledge has been generated about how the mass balances of both AIS and Greenland Ice Sheet (GIS) are impacted by surface mass balance (SMB), ice shelf basal melt, ice/bedrock friction, geothermal 110 heat flux, or ice rheology (see e.g. Fig. 2a). [All these advances need to be fully integrated into new probabilistic projections of sea-level change, and a new approach therefore needs to be envisioned that will allow for such new processes to be accurately modeled.](#)

---

<sup>10</sup>removed: such as the large set of Modeling Intercomparison Projects (MIPs) results from the CMIP5 or CMIP6 projects (such as ISMIP6 or GlacierMIP)

<sup>11</sup>removed: higher-frequency processes



**Figure 2.** Contribution to uncertainty in 100-year extreme warming simulations of AIS and three subregions of the AIS, tested for four different model variables independently. Each probability distribution function represents an ensemble of 800 Ice-sheet and Sea-level System Model ice dynamics (ISSM-ICE) runs, sampled using the ISSM-DAKOTA uncertainty quantification framework. (Schlegel et al., 2018)

[..<sup>12</sup>] Indeed, moving from strategies where continental scale mass changes are sampled and multiplied with the corresponding fingerprint, to actually sampling upstream model inputs [..<sup>13</sup>] is important for improving the state of the art. In particular, there is a strong need to fully account for spatial patterns of mass change and [..<sup>14</sup>] their uncertainty (see e.g. Fig. 2b-d), [..<sup>15</sup>] This applies to among others SMB, basal friction, or ice and solid-Earth rheological properties.

For example, Eq. 1 relies on masses that are aggregated at the basin/continental level. However, most ice sheet models compute high-resolution thickness change patterns that are not aggregated. This aggregation greatly reduces the complexity in representation of model physics and uncertainty propagated at the interface between ice-sheet models and PSLMs. A more comprehensive approach that reestablishes interfaces between forward models and PSLMs is therefore necessary, where model outputs are not aggregated or simplified.

<sup>12</sup>removed: Moving

<sup>13</sup>removed: , such as SMB, basal friction or ice and solid-Earth rheological properties among others, with consideration to

<sup>14</sup>removed: its

<sup>15</sup>removed: is paramount to improving the state of the art.

Here we propose a new framework for sea-level projections that is able to account for all terms in Eq. 1. We improve the existing process-based approach by using the Ice-Sheet and Sea-Level System Model (ISSM, Larour et al., 2012c) which allows for inclusion of forward model physics<sup>[..<sup>16</sup> ]</sup>. It also improves the modeling and sampling of covariances between input processes<sup>[..<sup>17</sup> ]</sup>, both temporally and spatially through the computation of high-resolution barystatic-GRD patterns<sup>[..<sup>18</sup> ]</sup>. The latter feature builds the basis for a geodetically compliant projection system where GRD patterns and their computation is done systematically and does not introduce biases in the projections.

## 2 Methods

### 2.1 Theory

130 Sterodynamic sea-level changes form a significant contributor to both global-mean sea-level rise and are responsible for large parts of the regional deviations from the global-mean projected changes (e.g. Slangen et al., 2012; Church et al., 2013b; Slangen et al., 2017). Following the CMIP5 conventions, sterodynamic sea-level changes consist of a global-mean thermosteric contribution (variable name `zostoga`) and a local dynamic contribution (variable name `zos`) with a zero mean over the oceans. Generally, an ensemble of model runs, either based on multiple models (e.g. Church et al., 2013b) or on large-ensemble  
135 experiments based on perturbing a single model (for example Little et al., 2017), can be used to directly sample regional sea-level changes. An alternative approach to generate more samples than model ensemble members is to determine common modes of variability, for example by extracting the largest empirical orthogonal functions from each model and perturbing the associated principal components (eg. Thompson et al., 2016, Fig.3).

While sterodynamic effects do not change the total ocean mass, ocean dynamics can <sup>[..<sup>19</sup> ]</sup>be coupled to redistribution of  
140 ocean mass, which <sup>[..<sup>20</sup> ]</sup>manifests in ocean-bottom pressure changes, particularly on shallow shelf seas (Landerer et al., 2007). These bottom pressure changes <sup>[..<sup>21</sup> ]</sup>load the solid Earth below, and thus result in GRD effects, which are often referred to as self-attraction and loading (SAL) effects (Ray, 1998a; Stepanov and Hughes, 2004; Vinogradova et al., 2015). These SAL effects could cause several centimeters of additional sea-level rise above the sterodynamic signal in century-scale sea-level projections made by atmosphere-ocean general circulation models (AOGCMs) (Richter et al., 2013). By adding the  
145 ocean-bottom pressure changes to the sea-level equation solver, this effect can be incorporated in regional sea-level projections.

As depicted in Eq. 1, in the classical approach, static sea-level fingerprints are computed a priori for each individual process, which typically include glaciers (GLA), the Greenland and Antarctic Ice Sheets and terrestrial water storage (TWS). These fingerprints are subsequently multiplied by the equivalent barystatic contribution, which is often sampled from a PDF, and added, together with the sterodynamic and GIA contribution to obtain local RSL changes and the associated confidence intervals. This

---

<sup>16</sup>removed: , retaining the covariance

<sup>17</sup>removed: from Eq. 1 and through

<sup>18</sup>removed: for each individual sample

<sup>19</sup>removed: drive


<sup>20</sup>removed: causes

<sup>21</sup>removed: cause a change in the load of



150 method is both transparent and simple, while maintaining computational efficiency owing to the fact that the fingerprints are do not have to be computed for each sample or time step.

However, several issues arise from this approach, which can be mitigated using a different method. First, it is assumed that the spatial pattern of mass loss is known a priori and does not vary over time. A common approach is to assume that the mass loss is uniformly distributed over the ice sheet, or that it follows the spatial pattern derived over the GRACE period. Jackson and Jevrejeva (2016) quantified the errors induced by assuming a uniform mass loss, and found that this bias could be up to 155 1 cm and  $\geq 10$  cm for sites distant from and close to centers of mass loss. Furthermore, the approximation of time-invariant fingerprints could lead to biases, when the spatial pattern of mass loss varies over time.



**Figure 3.** Diagram ISSM’s Sea-Level Projection System (ISSM-SLPS) model. The system is driven by requirements from Eq. 2. ISSM-SESAW is the GRD core of the system ( in pink). ISSM-SLPS is a combination of ISSM-SESAW and a layer (in green) that handles STR, DSL and GIA inputs, as well as all uncertainty quantification aspects.

In our approach (Figure 3) ISSM Sea-Level Projection System (ISSM-SLPS) solves for RSL as follows:

$$RSL(\theta, \phi, t) = RSL_{STR}(t) + RSL_{DSL}(\theta, \phi, t) + RSL_{GIA}(\theta, \phi, t) + RSL_{GRD}(\theta, \phi, t). \quad (2)$$

160 The first two terms on the right hand side, i.e.  $RSL_{STR}(t) + RSL_{DSL}(\theta, \phi, t)$ , together represent the sterodynamic sea level change.  $STR$  represents the [.,<sup>22</sup>] global mean thermosteric expansion and  $DSL$  local sea-level changes due to ocean dynamics. These can be obtained from CMIP results. The GIA contribution to ongoing sea level change,  $RSL_{GIA}$ , is given for example by Caron et al. (2018). The last term,  $RSL_{GRD}$  refers to the component of sea level change due to mass induced contemporary GRD response of the solid Earth (Gregory et al., 2019), excluding the GIA processes. This implies that vis-  
165 coelastic deformation is split between long-term time scales and short-term fast rebound of the bedrock uplift, such as

<sup>22</sup>removed: local

observed in West Antarctica (Barletta et al., 2018), acting essentially over time scales of 50-100 years. This includes mass transport between the land and the ocean, as well as that due to dynamic ocean circulation. The latter field is provided by CMIP as the ocean bottom pressure (OBP) products. Note that GRD associated with land-ocean mass transport is usually termed "sea level fingerprint" (e.g., Mitrovica et al., 2009), while the GRD due to OBP variability is termed "self-attraction and loading" phenomenon [..<sup>23</sup>](e.g., Ray, 1998a). As we shall see, we unify both of these elements of contemporary GRD sea level in equation (3). Note also that the global-mean OBP is removed from the ocean models, since ocean dynamics don't add or remove any mass from/to the ocean. In fact, our projections rely on CMIP5 and CMIP6 fields 'zos' (the sea-surface height change above geoid) and 'zostoga' for which the Greatbatch correction has been applied (Greatbatch, 1994).

We compute  $RSL_{GRD}$  using ISSM's Solid Earth and Sea-Level Adjustment module [..<sup>24</sup>](ISSM-SESAW; Adhikari et al., 2016a). Assuming that all of land ice/water mass change directly modulates the ocean mass, we define a global mass conserving loading function,  $M_{global}(\theta, \phi, t)$ , that describes the change in mass per unit area on the solid Earth surface as follows:

$$M_{global}(\theta, \phi, t) = M_{land}(\theta, \phi, t) \left[ 1 - \mathcal{O}(\theta, \phi) \right] + \rho_o \left[ H_{OBP}(\theta, \phi, t) + RSL_{GRD}(\theta, \phi, t) \right] \mathcal{O}(\theta, \phi), \quad (3)$$

where the land loading function (with dimensions of mass per unit area)  $M_{land}(\theta, \phi, t)$  is given by [..<sup>25</sup>]Adhikari et al. (2016a):

$$M_{land}(\theta, \phi, t) = \rho_i \left[ H_{AIS}(\theta, \phi, t) + H_{GIS}(\theta, \phi, t) + H_{GLA}(\theta, \phi, t) \right] + \rho_w H_{TWS}(\theta, \phi, t). \quad (4)$$

Here,  $\rho_i$  is the ice density,  $\rho_w$  is the freshwater density,  $\rho_o$  is the mean density of ocean water, and  $H_{OBP}$  is the (ocean) water equivalent height of the ocean bottom pressure change. Similarly,  $H_{AIS}$ ,  $H_{GIS}$ , and  $H_{GLA}$  are the ice height change in the respective cryospheric domains, and  $H_{TWS}$  is the freshwater height change in the non-cryospheric land domain. Note that we invoke an ocean function  $\mathcal{O}(\theta, \phi)$  in equation (3) to ensure mass conservation in the system. This function is equal to 1 on oceans, 0 everywhere else.

The contemporary mass transport function  $M_{global}(\theta, \phi, t)$  loads the underlying solid Earth that is self-gravitating, rotating, and viscoelastically compressible. The induced spatial pattern of  $RSL_{GRD}(\theta, \phi, t)$  is dictated by the perturbation in Earth's gravitational and rotational potentials and associated viscoelastic deformation of the solid Earth (Farrell and Clark, 1976; Milne and Mitrovica, 1998). In the absence of dynamic sea level and meteorologically induced high-frequency signals, the sea surface height mimics the spatial pattern of geoid (Gregory et al., 2019). Therefore, we may write

$$RSL_{GRD}(\theta, \phi, t) = C(t) + G_{GRD}(\theta, \phi, t) - B_{GRD}(\theta, \phi, t), \quad (5)$$

where  $G_{GRD}(\theta, \phi, t)$  and  $B_{GRD}(\theta, \phi, t)$  represent the change in geoid and bedrock elevation induced by the loading of the solid Earth (equation 3), respectively. Spatial invariant  $C(t)$  is invoked to ensure mass conservation in the Earth system, and it may be readily derived by inserting equation (5) into equation (3) and integrating it over the solid Earth surface.

<sup>23</sup>removed: (e.g., Ray, 1998b)

<sup>24</sup>removed: (ISSM-SESAW; ?)

<sup>25</sup>removed: Adhikari et al. (2016b)

Both  $G_{GRD}(\theta, \phi, t)$  and  $B_{GRD}(\theta, \phi, t)$  appearing in equation (5) may be partitioned into two components each: those related to gravitational potential and those to rotational potential. [..<sup>26</sup> ]These components can be computed by convolving  $M_{global}(\theta, \phi, t)$  with respective Green's functions. These may be defined in terms of surface harmonics with loading Love numbers as coefficients. Given the structure and viscoelastic properties of the solid Earth, these numbers characterize the axisymmetric deformational and gravitational response of Earth to the applied unit surface load. The rotational components depend upon tesseral second-degree loading and tidal Love numbers as well as on the perturbation in Earth's inertia tensor, which in turn depends on  $M_{global}(\theta, \phi, t)$ . In order to solve for  $R_{GRD}(\theta, \phi, t)$ , we require an a priori knowledge of  $M_{global}(\theta, \phi, t)$ , which in turn depends on  $R_{GRD}(\theta, \phi, t)$  itself. The system of equations (3) and (5) is therefore solved iteratively until a desired solution accuracy is achieved. One key feature of this field is that as ice sheets lose mass, the near-field relative sea level drops, and far-field sea level rises at a much larger rate than the barystatic term for the sake of mass conservation. While theoretical/numerical treatments on the topic are found elsewhere (e.g., Farrell and Clark, 1976; Mitrovica and Peltier, 1991; Mitrovica and Milne, 2003; Spada and Stocchi, 2007; Adhikari et al., 2019), version 1.0 of the SESAW algorithm where  $RSL_{GRD}$  is solved for is presented in [..<sup>27</sup> ]Adhikari et al. (2016a).

## 2.2 Meshing

SESAW is a mesh based convolution based on Eq. 2 in Farrell and Clark (1976). As such, it relies on an anisotropic unstructured mesh of the surface of the Earth which is refined according to specific metrics such as distance to the nearest coastline, presence of loads (such as changes in ice thickness or TWS), and the complexity of the coastline. Given the amount of inputs being sampled for in the SLPS system, a systematic approach to refining such a mesh needs to be developed. The main tool for such a refinement is the ISSM implementation of the [..<sup>28</sup> ]Bidimensional Anisotropic Mesh Generator (BAMG) anisotropic mesh refiner (Hecht, 2006a, b). This is a 2D based anisotropic mesher which can refine a mesh according to several constraints at the same time: a metric to specify directions along which the mesh resolution needs to be improved, specific vertex or segment positions, in particular vertex positions of the region outlines, and specified mesh resolutions for user-defined locations. Combining these constraints, we develop an approach based on meshing of a set of 2D continental areas of the Earth, projection of such 2D meshes onto the 3D Earth surface and then stitching of the resulting meshes into one seamless global 3D mesh.

A plot of the 2D regions is given in Fig. 4, which include South America, North America, Australia, Eurasia and the Pacific regions. At the North and South, we have regions defined for Antarctica, and Greenland. Greenland itself has been further refined into 18 regions drawn along the main ice divides of Greenland, following (Zwally et al., 2012, Fig.3). The approach facilitates a direct linkage of models from the existing literature, or potentially from previous ISSM studies such as Seroussi et al. (2017); Schlegel et al. (2018), without having to remesh the entire Earth. This in turn allows for direct comparisons


---

<sup>26</sup>removed: The former

<sup>27</sup>removed: ?

<sup>28</sup>removed: BAMG

between uncertainty quantification projection results where only one specific region is modified, hence allowing an approach  
225 where control runs can be compared against specific variations of an uncertainty quantification projection run.




**Figure 4.** 3D plot of the boundaries used to mesh each continental area of the Earth surface. Regions include South and North America, Australia, Eurasia and the Pacific regions, as well as Greenland and Antarctica. In [..<sup>29</sup>] this particular scenario, Greenland has been subdivided into 18 regions along the boundaries defined in Zwally et al. (2012).

An example mesh of the South-American continent is shown in Fig.5. This mesh relies on defined vertices for the outline, which match the outline vertices for the Pacific, Antarctica, and Eurasian meshes, so that the stitching within a larger 3D mesh can be done without redundancy in vertices along continental boundaries. In addition, GRACE ice mass trends from 2003 to 2016 (Adhikari and Ivins, 2016) are provided as a metric to be used for refinement of the mesh, in particular around the  
230 Patagonian ice fields. The minimum mesh resolution attained for this mesh is 500 m, and the largest is 1,400 km. Finally, the Global Self-consistent, Hierarchical, High-resolution Geography Database *GSHHS\_c\_L1* (Wessel and Smith, 1996) was used as a vertex constraint, so that the final mesh perfectly coincides with the coastline dataset (in black). This allows for the most optimum sea-level solution using the SESAW solver.

Once each region has been meshed in 2D using BAMG, it is projected onto latitude and longitude, and concatenated together  
235 to create a 3D mesh. This is possible because each 2D mesh relies on the same set of boundaries as shown in Fig. 4. The resulting mesh is shown in Fig. 6, and comprises 38944 surface elements for [..<sup>30</sup>]19,486 vertices. For comparison, an equi-angular  $1 \times 1^\circ$  grid would require 64,800 vertices, as much as three times more for a coarse grid resolution.


<sup>30</sup>removed: 19486 vertices.



**Figure 5.** 2D Adaptive mesh of South-America using GRACE observations of ice and hydrological mass change (in cm/yr) from 2003 to 2016. Seismic effects (Richter et al., 2019) are not removed in this rendering of Patagonian ice mass loss that was directly taken from Adhikari and Ivins (2016). The Global Self-consistent, Hierarchical, High-resolution Geography Database (GSHHG) coarse L1 coastline is shown in green. Segments of the triangular mesh are plotted in black. The colorbar for the thickness changes was saturated at  $[-1 \ 1]$  cm/yr in order to improve the contrast of the figure given the high mesh resolution.

### 2.3 Sampling and partitioning

In order to sample variables at each time step, our approach is to use a geographical partitioning of the unstructured mesh. An example is shown in Fig. 7, where a range of values from 1 to 5 has been attributed for each vertex (and element) of the mesh, corresponding respectively to Antarctica (1), Greenland (2), Glaciers (3), the ocean (4) and land (5). For each partition and for each variable that is probabilistically sampled, we define a probability density function (PDF). For normal distributions for example, this will be done through a mean and standard deviation.



**Figure 6.** 3D Earth mesh stitched from 3D projections of 2D regional meshes of the following regions: South and North America, Australia, Eurasia, the Pacific regions as well as Antarctica and Greenland. GRACE observations of ice mass change (in cm/yr) from 2003 to 2016 (Adhikari and Ivins, 2016) are overlaid over the mesh. Left frame azimuth is 30° with elevation of 64°. Right frame azimuth is 205° with elevation of 23°.

The algorithm for sampling through SLPS is explained below, in the generic case where spatial covariances are available  
245 between variables.

```
for t=2019:2100,  
    pdf=PDF(type, pdfspec_arg,t);  
    for j=1:nsamples,  
        alphas = DAKOTA->sample(pdf,j); %alphas array of size [NUM(VAR),RANGE(partition)]  
250    for VAR in (DSL,STR,H_AIS,H_GIS,H_GLA,H_TWS):  
        for VERTEX in MESH:  
            for i=1:range(PARTITION):  
                if PARTITION(VERTEX)==i,  
                    alpha=alphas(VAR,i)  
255                    VAR(VERTEX)=VAR0(VERTEX)*alpha;  
                end  
            end  
        end  
    end  
260    RSL(j,t)=SLPS(DSL,STR,H_AIS,H_GIS,H_GLA,H_TWS);  
end  
end
```

where  $t$  is the time variable (ranging from year 2019 to 2100, at 1 year intervals),  $j$  is the counter for each sample, from 1 to  $nsamples$  (in our case, 10,000),  $VAR$  is the sampled variable (from one of the SESAW inputs, excluding  $RSL_{GLA}$  which is deterministic in our framework),  
265  $VERTEX$  is a counter for all vertices in the mesh  $MESH$ ,  $PARTITION$  is the partition vector (for example ranging from 1 to 5 in Fig. 7),  $PDF$  is the joint probability distribution of variables across all geographical locations,  $DAKOTA$  is the sample generator in ISSM (Eldred et al., 2008; Larour et al., 2012b), [<sup>31</sup>]  $\alpha$  is the  $j^{th}$  sample matrix of scaling factors with size (number of variables, number of partitions),  $VAR0$  the unmodified variable (stored in memory at the beginning of the model run),  $SLPS$  is the sea-level solver, generating  $RSL$  for a specific sample of all the probabilistic variables. In this algorithm, the  $PDF$  distribution is built by specifying its nature and parameters,  
270 e.g. the 'type' argument can indicate the choice of a multivariate Gaussian distribution and 'pdfspec\_arg' specify the vector of means and covariance matrix of  $\alpha$  between each other and between partitions.


For this application, we assume that each variable and each partition is independent, and we set the mean of all distributions to 1. This ensures that values of  $\alpha$  behave as scaling parameters. We use them to directly scale a variable locally, according to which partition area this location geographically belongs. This method is therefore significantly different from the [<sup>32</sup>] approach in KOPPI4, where the entire  
275 mass within a certain partition (for example GIS or AIS) is sampled. Here the sampling is a scaling of a vectorial field, which therefore preserves the local geographical distribution of a given variable. This is shown in Fig. 8 for a scenario where thinning rates of the GIS are sampled using one geographical partition (corresponding to Fig. 7 partition value of 2, in blue). We display [<sup>33</sup>] the average thinning rate

---

<sup>31</sup>removed: alphas

<sup>32</sup>removed: approach

<sup>33</sup>removed: for a standard deviation  $\sigma = 5\%$  the



**Figure 7.** Partition vector (values from 1 to 5, 1 for Antarctica, 2 for Greenland, 3 for Glaciers around the world, 4 for the ocean and 5 for land excepted glaciers). The partition vector is used to sample probabilistic variables in a geographically consistent way, with PDF distributions moments (mean and standard deviation) defined for each partition area.


$\mu$ ,  $\mu + 3\sigma$  and  $\mu - 3\sigma$  (for an arbitrary value of the <sup>34</sup>standard deviation  $\sigma = 5\%$ ). The structure of the thinning rate as it is sampled is kept intact, implying that the spatial covariance of the variable being sampled across the mesh is kept closely similar across samples and within any given partition.

## 2.4 Modularity

The advantage of the partition approach as implemented in SLPS is that various approaches to probabilistic projections can be executed with the same framework. First, as we will show in the next section, the KOPP14 approaches are fully compatible with the SLPS framework. Indeed, fingerprint patterns can be recomputed using local thickness change rate patterns that are spatially constant on the basis of only one partition, such as the entire Greenland or Antarctic ice sheet (contrary to Fig. 12 where Greenland is subdivided). **Once several partitions are adopted however, the refinement in the fingerprint patterns significantly departs from the KOPP14 approach.** Second, existing probabilistic assessments for specific components (such as the impact of changes in surface mass balance or basal friction in Antarctica (Schlegel et al., 2018) on ice thickness changes) can be used directly, using model output (for example for thickness change rates), or PDF distributions from such model outputs. If the uncertainty quantification was done using a Bayesian framework, the model output statistics can be reused directly (using some type of uniform discrete sampling of each model output), hence replicating a Bayesian type exploration

<sup>34</sup>removed: thinning rate samples at three standard deviations from the average






**Figure 8.** Random sampling of thinning rates across Greenland. a) GRACE generated thinning rate pattern at 2005 (in m/yr). b) Thinning rate along the AB profile (from a) (in red, representing the average of the PDF), and samples generated at  $-3\sigma$  (blue) and  $+3\sigma$  (yellow) from the average.

approach of SLPS without incurring any additional computational cost (meaning, not having to rerun the analysis carried out to compute such model outputs). Third, ISSM modules can be activated upstream of the SLPS solver, to push further the boundaries of the uncertainty assessment. For example, an analysis of the impact of SMB variations in one specific region of Antarctica could be carried out using the ice-flow modeling core of ISSM, capable of delivering ice thickness changes directly to the SLPS core. Fourth, these modules can be activated while remaining coupled to other modules. For example, in Larour et al. (2019), it was demonstrated that over centennial time scales, coupling between the elastic uplift of the grounding line, and ice-flow related grounding line migration, are key to controlling the retreat of Thwaites Glacier in West Antarctica. Assessing the uncertainty brought by such processes on sea-level rise (SLR) projections would require this coupling to be activated, which could be done (assuming computational costs are still realistic) without modifications to the SLPS framework. Finally, given how closely ISSM can be integrated within Web Server architectures using its native JavaScript interface (Larour et al., 2017a), SLPS is potentially fully compatible with open source types of collaborative approaches where inputs from the community could be provided directly to Web Servers running ISSM in the background, to generate model projections without significant investment in a computation core and/or an interface to the latter.

### 3 Results and Discussion

SLPS probabilistic projections were validated using model inputs from the Intergovernmental Panel on Climate Change World Fifth Assessment Report (IPCC AR5) (Church et al., 2013a). AR5 supplies several projection components in SLR equivalents: the 'expansion' term ( $STR$ ), the 'glacier' term (which can be converted into an average thickness change rate for  $H_{GLA}$ ), 'antnet' and 'greenet' for net barystatic contribution from the Antarctica and Greenland ice sheets, which can also be converted into an average change rate for  $H_{AIS}$  and  $H_{GIS}$  and the 'landwater' term for TWS contribution to SLR (which can be converted into an average change rate for  $H_{TWS}$ ). For each of these terms, AR5 supplies the mean projection, and the 5-95% percentile confidence interval. We can use this information to calibrate PDF distributions for thickness change rates at each time step, with the mean of each PDF corresponding to the AR5 mean, and the standard-deviation calibrated from the 5-95% interval (corresponding to the  $[-1.65\sigma, 1.65\sigma]$  interval). Because AR5 does not supply spatial patterns, we rely on GRACE 2003-2016 thickness change rate patterns from Adhikari and Ivins (2016) for  $H_{GLA}$ ,  $H_{AIS}$  and  $H_{GIS}$ . For  $H_{TWS}$ , we assume a uniform spatial distribution over all the spatial partitions.  $STR$  is also considered uniform over all the oceans.  $DSL$  is not sampled, but rather deterministically set to the  $DSL$  term of the CMIP5 NorESM-ME runs (Bentsen et al., 2013). GIA is independently sampled (from Caron et al. (2018)) and probabilistically added as an independent PDF. The sampling is carried out on the partitions described in Fig.7 with the notable exception that the GIS is further divided into 18 different basins as defined in Zwally et al. (2012) and as plotted in Fig.12. For each year between 2007 and 2100, 10,000 sample runs of SLPS are carried out (with full geodetic capabilities of the SESAW core). For each partition, samples for the corresponding inputs are generated using a Latin Hypercube Sampling (LHS) algorithm. The runs were carried out on the Pleiades cluster at the NASA Ames Research Center, on 20 Ivy nodes (20 cores per node for an equivalent 400 cores) over 7 hours.



**Figure 9.** ISSM-SLPS projections based on AR5 RCP8.5. For each time step, we sample (10,000 times with Latin Hypercube Sampling, or LHS) the following inputs:  $H_{GIS}$ ,  $H_{AIS}$ , thermal expansion of the ocean ( $STR$ ),  $H_{TWS}$ , and glacier contributions  $H_{GLA}$  (see AR5 WG1 Chapter 13, Church et al. (2013b)). Each input's PDF is calibrated using the AR5 5-95% projection confidence interval, similar to Kopp et al. (2014). The resulting GMSL PDF distribution is shown in a) (in time) and b) (at a sub-set of time steps). The 5-95% confidence interval (likely range, following AR5 definition) is plotted in black in a), along with the temporal mean. Each time step is fully decorrelated from the previous time steps, this test being used to validate against existing an existing AR5 projection

<sup>35</sup>removed:  $-1.65\sigma$  to  $1.65\sigma$

320 Fig. 9 shows projection results for GMSL computed at each time step between 2007 and [..<sup>36</sup> ]2100, and histograms for several time snapshots. We match the mean and 5-95% confidence intervals of AR5 (Fig.9a) as expected. We also show the evolution of RSL in Fig.[..<sup>37</sup> ]10 for nine cities around the world. We provide the mean and standard deviation for each PDF, and show how the sampling of ice-related thickness changes impacts mean and standard deviation. In particular, as expected from the AR5 inputs, we show a marked increase in PDF spreads as time evolves [..<sup>38</sup> ]from 2007 to 2100.

325 Fig. 11 shows the impact of using existing statistics from Caron et al. (2018) to include into SLPS. These statistics were evaluated using [..<sup>40</sup> ]bayesian inversion method based on Simulated Annealing (Kirkpatrick et al., 1983), a variation of the Monte Carlo with Markov chains (MCMC) method (Metropolis and Ulam, 1949; Metropolis et al., 1953). They can be used directly in SLPS, either during a standard probabilistic projection run, or a posteriori as is the case here. These statistics reflect the statistical fitness to a global GIA dataset composed of paleo-RSL indicators and vertical GPS trends. The impact of the migrating Laurentide isostatic bulge on Norfolk, Virginia  
330 is apparent in Fig.11, with an offset of 16 cm in the average projection for the city.

Fig. 12 shows results for a different experiment, in which we quantify the impact of refining the amount of partitions used to sample the uncertainty in ice thickness change rates. For the area of Greenland, we use either one partition (blue boundary), or 18 boundaries (brown basins) from the Zwally et al. (2012) dataset. Each basin is delimited by ice divides, and thus represent a dynamically coherent area, expected to behave (short of ice divides migrating actively) independently from one another. We rerun an SLPS projection using a similar AR5 setup,  
335 and display the contribution of ice-related basins to SLR in New York and Hawaii for 1 and 18 partitions respectively. As expected, the mean in PDF distributions are identical for both 1 and 18 partitions. However, the tails are much larger for the 1 basin scenario. The relative difference in standard deviations between 1 and 18 basins ranges from -23% for New York to -34% for Hawaii. This implies that current probabilistic RSL projections are significantly overestimating (by 20-30%) the width of the "likely" (5-95%) range in ice-melt contribution to RSL.

340 This is understandable because of the fact that in a 1 partition scenario, variations of ice thickness are dictated by scaling of the local ice thickness change rate mean by an identical scalar for the entire partition, which leads to more extreme values for the contribution to RSL. With finer partitions, basins that have low thickness change rates do not [..<sup>41</sup> ]impact RSL as much, and in aggregate the total contribution range varies less. This can be visualized better by taking the example of New York, where following Larour et al. (2017b) contributions from South Greenland are almost negligible. This implies that all the basins (and corresponding GRD patterns) in South Greenland  
345 will contribute zero variance to the PDF for RSL at New York. This will therefore result in smaller tails for projections that rely on more refined basins. A very similar conclusion was found in Schlegel et al. (2018), where Antarctica had to be subdivided in spatially coherent areas, which were not obvious initially and did not mandatorily map into individual basins. The issue is that the error distribution in model inputs had a specific spatial coherence that had to be respected. Assuming this coherence extended to the entire ice sheet led to significantly larger and unrealistic uncertainty ranges in model outputs. Of course, given differing dynamics in each geographical basin, we cannot assume  
350 that the input scaling should be similar (same standard-deviation). This will modify the results in Fig. 12. But our point here is to point out the issue of sub-partitioning as being essential in quantifying the right range of spread in modeled statistical outputs.

This analysis also shows that using SLPS, it is possible to efficiently address the question of how to sample uncertainty in a manner that is consistent with the local behaviour of separate basins, glaciers, ice sheets. In Jackson and Jevrejeva (2016) for example, it is shown that

---


<sup>36</sup>removed: 2100.

<sup>37</sup>removed: 9b.

<sup>38</sup>removed: .

<sup>40</sup>removed: a Bayesian exploration approach, and


<sup>41</sup>removed: impact



**Figure 10.** AR5 calibrated projection of RSL for nine cities around the world [..<sup>39</sup>] from 2007 to 2100. Sampling was carried out for  $H_{AIS}$ ,  $H_{GIS}$ ,  $H_{GLA}$ ,  $STR$  and  $H_{TWS}$  using mean and standard deviations from AR5 (Church et al., 2013a). The patterns for ice thickness are from GRACE 2003-2016 trends (Adhikari and Ivins, 2016).  $DSL$  is fully deterministic, from the CMIP5 NorESM runs (Bentsen et al., 2013).  $RSL_{GIA}$  was deterministically set to 0. Each time step was sampled for using 10,000 LHS samples.

the impact of glacier ice thickness variations around the world is [..<sup>42</sup>] significantly different, and that relying on one fingerprint alone can lead to significant differences in the projection of glacier contribution (up to several percent). Our approach in SLPS ensures that the GRD contribution is systematically reassessed for each sample, at each time step, and the partitioning of our sampling ensures that we correctly

<sup>42</sup>removed: significantly




**Figure 11.** AR5 calibrated projection of RSL for Norfolk (in brown) vs same projection in which GIA statistics from Caron et al. (2018) are used to account for GIA induced RSL (in blue). Note that both PDF distributions have standard deviations that are essentially identical within .4% relative difference.

capture the specificity of each glacier/ice/hydrological area and their unique mass change trends. It is to be noted that a similar approach is currently implemented in new instantiations of the KOPP14 projection system based on sampling of glacier projections across the 19 Randolph Glacier Inventory (RGI) areas used in the GlacierMIP results (Hock et al., 2019). However, these areas can be very large in spatial extent (such as the Low Latitudes or North-Asia areas) and should be broken down. Our approach scales for any barystatic contributor, at any spatial scale (example, sub-basin, or at the glacier level) required by the structure of the error distribution of model inputs.

#### 4 Conclusions

ISSM SLPS is a new sea-level probabilistic projection system which relies on a new partitioning approach to sampling of boundary conditions, forcings and inputs. It is compatible with previous probabilistic frameworks, but allows for a more robust integration of state-of-the-art results in the modeling of ice flow in ice sheets and glaciers, steric sea-level, TWS evolution and GIA. It reestablishes temporal correlation in projections where they were previously lacking, and allows for better constraints on spatial and temporal covariances in the model inputs. In particular, it is capable of systematically computing geodetically compliant patterns of sea-level that are consistent with



**Figure 12.** Impact of sub-sampling the GIS mass (from 1 basin for the whole ice sheet to 18 basins) on barystatic sea-level rise in New York and Hawaii. The distributions are a result of SLPS, where  $H_{GIS}$ ,  $H_{AIS}$  and  $H_{GLA}$  were sampled 10,000 times using an LHS algorithm. The mean in PDF distributions for both scenarios are identical, however the tails are much larger for the 1 basin scenario. The relative difference in standard deviations between 1 and 18 basins ranges from -23% for New York to -34% for Hawaii. This implies that current probabilistic RSL projections could significantly overestimate (20-30%) the "likely" (5-95%) range in ice-melt contribution from glaciers and ice sheets.

space and terrestrial measurement systems. The system relies heavily on the use of high-resolution anisotropic meshes, and allows for a better interfacing with existing modeling frameworks which operate at higher resolutions, and which consistently generate changes in mass density patterns around the globe. SLPS has been validated against previous frameworks and is fully backwards compatible. Differences between SLPS and previous approaches have also been shown both in terms of integration of GIA statistics, and integration of new high-resolution sampling of ice-thickness change patterns in Greenland. This new approach offers a roadmap towards [.,<sup>43</sup>] further increasing the complexity and realism of sea-level probabilistic projection frameworks.

*Code availability.* The ISSM code and its SLPS components are available at <http://issm.jpl.nasa.gov>. The instructions for the compilation of ISSM and SLPS modules are available at <http://issm.jpl.nasa.gov/download>. The public svn repository for the ISSM code can also be found directly at <https://issm.ess.uci.edu/svn/issm/issm/trunk>, and downloaded using user name "anon" and password "anon". The version of the code for this study, corresponding to ISSM release 4.17, is svn version tag number 24683.

*Data availability.* All datasets used in the projections are freely available in the public domain and are referenced in the text.

<sup>43</sup>removed: urther

*Author contributions.* E. Larour carried out all the simulations and implemented SLPS into ISSM. He wrote the bulk of the manuscript. L. Caron contributed computations for GIA. M. Morlighem contributed enhancements to the adaptive meshers in ISSM. All authors contributed to the manuscript both in terms of text, figures and comments.

*Competing interests.* The authors have no competing interests

*Acknowledgements.* This work was supported by the Jet Propulsion Laboratory, California Institute of Technology under a contract with the NASA Sea-level Change Team (N-SLCT), NASA Cryospheric Sciences, NASA Modeling Analysis and Prediction (MAP) and NASA Earth Surface and Interior (ESI) Programs as well as the NASA GRACE and GRACE-FO Science Teams Programs. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center. We would like to acknowledge Dr. Ed. Zaron for his insightful comments on the manuscript.

## References

- Adhikari, S. and Ivins, E. R.: Climate-driven polar motion: 2003–2015, *Science Advances*, 2, <https://doi.org/10.1126/sciadv.1501693>, <http://advances.sciencemag.org/content/2/4/e1501693>, 2016.
- Adhikari, S., Ivins, E. R., and Larour, E.: ISSM-SESAW v1.0: mesh-based computation of gravitationally consistent sea-level and geodetic signatures caused by cryosphere and climate driven mass change, *Geoscientific Model Development*, 9, 1087–1109, <https://doi.org/10.5194/gmd-9-1087-2016>, <https://www.geosci-model-dev.net/9/1087/2016/>, 2016a.
- Adhikari, S., Ivins, E. R., and Larour, E.: ISSM-SESAW v1.0: Mesh-based computation of gravitationally consistent sea level and geodetic signatures caused by cryosphere and climate driven mass change, *Geoscientific Model Development*, 9, 1087–1109, 2016b.
- Adhikari, S., Ivins, E. R., Frederikse, T., Landerer, F. W., and Caron, L.: Sea-level fingerprints emergent from GRACE mission data, *Earth Syst. Sci. Data*, 11, 629–646, 2019.
- Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P., and Cooke, R. M.: Ice Sheet Contributions to Future Sea-Level Rise from Structured Expert Judgment, *Proceedings of the National Academy of Sciences*, 116, 11 195–11 200, <https://doi.org/10.1073/pnas.1817205116>, 2019.
- Barletta, V. R., Bevis, M., Smith, B. E., Wilson, T., Brown, A., Bordoni, A., Willis, M., Khan, S. A., Rovira-Navarro, M., Dalziel, I., Smalley, R., Kendrick, E., Konfal, S., Caccamise, D. J., Aster, R. C., Nyblade, A., and Wiens, D. A.: Observed rapid bedrock uplift in Amundsen Sea Embayment promotes ice-sheet stability, *Science*, 360, 1335–1339, 2018.
- Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevåg, A., Seland, Ø., Drange, H., Roelandt, C., Seierstad, I. A., Hoose, C., and Kristjánsson, J. E.: The Norwegian Earth System Model, NorESM1-M – Part 1: Description and Basic Evaluation of the Physical Climate, *Geoscientific Model Development*, 6, 687–720, <https://doi.org/10.5194/gmd-6-687-2013>, 2013.
- Boesch, D., Boicourt, W., Cullather, R., Ezer, T., G.E. Galloway, J., Johnson, Z., Kilbourne, K., Kirwan, M., Kopp, R., Land, S., Li, M., Nardin, W., Sommerfield, C., and Sweet, W.: Sea-level Rise: Projections for Maryland 2018, University of Maryland Center for Environmental Science, Cambridge, MD, [https://www.umces.edu/sites/default/files/Sea-Level%20Rise%20Projections%20for%20Maryland%202018\\_0.pdf](https://www.umces.edu/sites/default/files/Sea-Level%20Rise%20Projections%20for%20Maryland%202018_0.pdf), 2018.
- Callahan, J. A., Horton, B. P., Nikitina, D. L., Sommerfield, C. K., McKenna, T. E., and Swallow, D.: Recommendation of Sea-Level Rise Planning Scenarios for Delaware: Technical Report, prepared for Delaware Department of Natural Resources and Environmental Control (DNREC) Delaware Coastal Programs, <https://www.dgs.udel.edu/sites/default/files/projects-docs/Delaware%20SLR%20Technical%20Report%202017.pdf>, 2017.
- Caron, L., Ivins, E. R., Larour, E., Adhikari, S., Nilsson, J., and Blewitt, G.: GIA Model Statistics for GRACE Hydrology, Cryosphere, and Ocean Science, *Geophysical research letters*, 45, 2203–2212, 2018.
- Church, J., Clark, P., Cazenave, A., Gregory, J., Jevrejeva, S., Levermann, A., Merrifield, M., Milne, G., Nerem, R., Nunn, P., Payne, A., Pfeffer, W., Stammer, D., and Unnikrishnan, A.: Sea Level Change Supplementary Material. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, [www.climatechange2013.org](http://www.climatechange2013.org), 2013a.
- Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., Payne, A. J., Pfeffer, W. T., Stammer, D., and Unnikrishnan, A. S.: *Climate Change 2013: The Physical Science Basis Change*, chap. 13, pp. 1137–1216, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.026>, <http://www.climatechange2013.org>, 2013b.



- 425 City of Boston, U. S.: Boston Research Advisory Group (BRAG) Climate Projections Full Report, Prepared for Climate Ready Boston, [https://www.boston.gov/sites/default/files/document-file-12-2016/brag\\_report\\_-\\_final.pdf](https://www.boston.gov/sites/default/files/document-file-12-2016/brag_report_-_final.pdf), 2016.
- Dalton, M., Dello, K., Hawkins, L., Mote, P., and Rupp, D.: The Third Oregon Climate Assessment Report, Oregon Climate Change Research Institute, College of Earth, Ocean and Atmospheric Sciences, Oregon State University, Corvallis, OR, <http://www.occri.net/publications-and-reports/third-oregon-climate-assessment-report-2017>, 2017.
- 430 DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591–597, <https://doi.org/10.1038/nature17145>, 2016.
- Eldred, M. S., Adams, B. M., Gay, D. M., Swiler, L. P., Haskell, K., Bohnhoff, W. J., Eddy, J. P., Hart, W. E., Watson, J.-P., Hough, P. D., and Kolda, T. G.: DAKOTA, A Multilevel Parallel Object-Oriented Framework for Design Optimization, Parameter Estimation, Uncertainty Quantification, and Sensitivity Analysis, Version 4.2 User’s Manual, Technical Report SAND 2006-6337, Tech. rep., Sandia National Laboratories, PO Box 5800, Albuquerque, NM 87185, 2008.
- 435 Farrell, W. E. and Clark, J. A.: On Postglacial Sea Level, *Geophysical Journal of the Royal Astronomical Society*, 46, 647–667, 1976.
- Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., Lipscomb, W. H., Gregory, J., Abe-Ouchi, A., Shepherd, A., Simon, E., Agosta, C., Alexander, P., Aschwanden, A., Barthel, A., Calov, R., Chambers, C., Choi, Y., Cuzzone, J., Dumas, C., Edwards, T., Felikson, D., Fettweis, X., Golledge, N. R., Greve, R., Humbert, A., Huybrechts, P., Le clec’h, S., Lee, V., Leguy, G., Little, C., Lowry, D. P., 440 Morlighem, M., Nias, I., Quiquet, A., Rückamp, M., Schlegel, N.-J., Slater, D., Smith, R., Straneo, F., Tarasov, L., van de Wal, R., and van den Broeke, M.: The future sea-level contribution of the Greenland ice sheet: a multi-model ensemble study of ISMIP6, *The Cryosphere*, <https://doi.org/10.5194/tc-2019-319>, <https://www.the-cryosphere-discuss.net/tc-2019-319/>, 2020, accepted.
- Goldberg, D. N., Little, C. M., Sergienko, O. V., Gnanadesikan, A., Hallberg, R., and Oppenheimer, M.: Investigation of land ice-ocean interaction with a fully coupled ice-ocean model: 2. Sensitivity to external forcings, *J. Geophys. Res.*, 117, 1–11, 445 <https://doi.org/10.1029/2011JF002247>, 2012.
- Goldberg, D. N., Snow, K., Holland, P., Jordan, J. R., Campin, J.-M., Heimbach, P., Arthern, R., and Jenkins, A.: Representing grounding line migration in synchronous coupling between a T marine ice sheet model and a z-coordinate ocean model, *Ocean Modelling*, 125, 45 – 60, <https://doi.org/10.1016/j.ocemod.2018.03.005>, 2018.
- Goldberg, D. N., Gourmelen, N., Kimura, S., Millan, R., and Snow, K.: How Accurately Should We Model Ice Shelf Melt Rates?, *Geophys. Res. Lett.*, 46, <https://doi.org/10.1029/2018GL080383>, 2019.
- 450 Gomez, N., Latychev, K., and Pollard, D.: A Coupled Ice Sheet-Sea Level Model Incorporating 3D Earth Structure: Variations in Antarctica during the Last Deglacial Retreat, *Journal of Climate*, 31, 4041–4054, 2018.
- Gornitz, V., Oppenheimer, M., Kopp, R., Orton, P., Buchanan, M., Lin, N., Horton, R., and Bader, D.: New York City Panel on Climate Change 2019 Report Chapter 3: Sea Level Rise, *Annals of the New York Academy of Sciences*, 1439, 71–94, 2019.
- 455 Greatbatch, R. J.: A note on the representation of steric sea level in models that conserve volume rather than mass, *Journal of Geophysical Research*, 99, 12 767, <https://doi.org/10.1029/94jc00847>, 1994.
- Gregory, J., Griffies, S., Hughes, C., et al.: Concepts and Terminology for Sea Level: Mean, Variability and Change, Both Local and Global, *Surv. Geophys.*, 40, 1251–1289, 2019.
- Griggs, G., Arvai, J., Cayan, D., DeConto, R., Fox, J., Fricker, H., Kopp, R., Tebaldi, C., and Whiteman, E.: Rising Seas in 460 California: An Update on Sea-Level Rise Science, California Ocean Science Trust, <http://www.opc.ca.gov/webmaster/ftp/pdf/docs/rising-seas-in-california-an-update-on-sea-level-rise-science.pdf>, 2017.

- Gunter, B., Didova, O., Riva, R., Ligtenberg, S., Lenaerts, J., King, M., Van den Broeke, M., Urban, T., et al.: Empirical estimation of present-day Antarctic glacial isostatic adjustment and ice mass change, *The Cryosphere*, 8, 743–760, 2014.
- Hecht, F.: BAMG: Bi-dimensional Anisotropic Mesh Generator, Tech. rep., FreeFem++, <http://www-rock-inria/gamma/cdrom/www/bamg/eng.htm>, 2006a.
- 465 Hecht, F.: Maillage 2d, 3d, adaptation, Tech. rep., Universite Pierre et Marie Curie, Paris, 2006b.
- Hock, R., Bliss, A., Marzeion, B., Giesen, R. H., Hirabayashi, Y., Huss, M., Radic, V., and Slangen, A. B. A.: GlacierMIP - A model intercomparison of global-scale glacier mass-balance models and projections, *Journal of Glaciology*, 65, 453–467, 2019.
- Jackson, L. P. and Jevrejeva, S.: A probabilistic approach to 21st century regional sea-level projections using RCP and High-end scenarios, *470 Global and Planetary Change*, 146, 179–189, 2016.
- Jevrejeva, S., Frederikse, T., Kopp, R. E., Le Cozannet, G., Jackson, L. P., and van de Wal, R. S. W.: Probabilistic Sea Level Projections at the Coast by 2100, *Surveys in geophysics*, 81, 1–24, 2019.
- Kaplan, M., Campo, M., Auermuller, L., and Herb, J.: Assessing New Jersey’s Exposure to Sea-Level Rise and Coastal Storms: A Companion Report to the New Jersey Climate Adaptation Alliance Science and Technical Advisory Panel Report, Prepared for the New Jersey Climate Adaptation Alliance. New Brunswick, NJ: Rutgers University, <https://njadapt.rutgers.edu/docman-lister/conference-materials/168-crfinal-october-2016/file>, 2016.
- 475 Kirkpatrick, S., Gelatt, C. D., and Vecchi, M. P.: Optimization by Simulated Annealing, *Science*, 220, 671–680, <https://doi.org/10.1126/science.220.4598.671>, 1983.
- Kopp, R., Broccoli, A., Horton, B., Kreeger, D., Leichenko, R., Miller, J., Miller, J., Orton, P., Parris, A., Robinson, D., C.P.Weaver, Campo, M., Kaplan, M., Buchanan, M., Herb, J., Auermuller, L., and Andrews, C.: Assessing New Jersey’s Exposure to Sea-Level Rise and Coastal Storms: Report of the New Jersey Climate Adaptation Alliance Science and Technical Advisory Panel, Prepared for the New Jersey Climate Adaptation Alliance. New Brunswick, New Jersey, <https://njadapt.rutgers.edu/docman-lister/conference-materials/167-njcaa-stap-final-october-2016/file>, 2016.
- 480 Kopp, R. E., Horton, R. M., Little, C. M., Mitrovica, J. X., Oppenheimer, M., Rasmussen, D., Strauss, B. H., and Tebaldi, C.: Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites, *Earth’s future*, 2, 383–406, 2014.
- Kopp, R. E., DeConto, R. M., Bader, D. A., Hay, C. C., Horton, R. M., Kulp, S., Oppenheimer, M., Pollard, D., and Strauss, B. H.: Evolving understanding of Antarctic ice-sheet physics and ambiguity in probabilistic sea-level projections, *arXiv.org*, pp. 1217–1233, 2017.
- Kopp, R. E., Gilmore, E. A., Little, C. M., Lorenzo-Trueba, J., Ramenzoni, V. C., and Sweet, W. V.: Usable Science for Managing the Risks of Sea-Level Rise, *Earth’s Future*, p. 2018EF001145, <https://doi.org/10.1029/2018EF001145>, 2019.
- 490 Landerer, F. W., Jungclaus, J. H., and Marotzke, J.: Ocean Bottom Pressure Changes Lead to a Decreasing Length-of-Day in a Warming Climate, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2006GL029106>, 2007.
- Larour, E., Morlighem, M., Seroussi, H., Schiermeier, J., and Rignot, E.: Ice flow sensitivity to geothermal heat flux of Pine Island Glacier, Antarctica, *J. Geophys. Res. - Earth Surface*, 117, 1–12, <https://doi.org/10.1029/2012JF002371>, 2012a.
- Larour, E., Schiermeier, J., Rignot, E., Seroussi, H., Morlighem, M., and Paden, J.: Sensitivity Analysis of Pine Island Glacier ice flow using ISSM and DAKOTA, *J. Geophys. Res.*, 117, F02009, 1–16, <https://doi.org/10.1029/2011JF002146>, 2012b.
- 495 Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *J. Geophys. Res.*, 117, 1–20, <https://doi.org/10.1029/2011JF002140>, 2012c.

- Larour, E., Cheng, D., Perez, G., Quinn, J., Morlighem, M., Duong, B., Nguyen, L., Petrie, K., Harounian, S., Halkides, D., and Hayes, W.: A JavaScript API for the Ice Sheet System Model: towards an online interactive model for the Cryosphere Community, *Geosci. Model Dev.*, 10, 4393–4403, <https://doi.org/10.5194/gmd-10-4393-2017>, <https://www.geosci-model-dev.net/10/4393/2017/>, 2017a.
- 500 Larour, E., Ivins, E. R., and Adhikari, S.: Should coastal planners have concern over where land ice is melting?, *Science Advances*, 3, e1700537, 2017b.
- Larour, E., Seroussi, H., Adhikari, S., Ivins, E., Caron, L., Morlighem, M., and Schlegel, N.: Slowdown in Antarctic mass loss from solid Earth and sea-level feedbacks, *Science*, 364, eaav7908, 2019.
- 505 Le Bars, D.: Uncertainty in Sea Level Rise Projections Due to the Dependence Between Contributors, *Earth's Future*, <https://doi.org/10.1029/2018EF000849>, 2018.
- Little, C. M., Piecuch, C. G., and Ponte, R. M.: On the Relationship between the Meridional Overturning Circulation, Alongshore Wind Stress, and United States East Coast Sea Level in the Community Earth System Model Large Ensemble, *Journal of Geophysical Research: Oceans*, 122, 4554–4568, <https://doi.org/10.1002/2017JC012713>, 2017.
- 510 Melini, D. and Spada, G.: Some remarks on Glacial Isostatic Adjustment modelling uncertainties, *Geophysical Journal International*, 218, 401–413, 2019.
- Metropolis, N. and Ulam, S.: The Monte Carlo method, *J. Amer. Stat. Associ.*, 44, 335–341, 1949.
- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., and Teller, E.: Equation of State Calculations by Fast Computing Machines, *The Journal of Chemical Physics*, 21, 1087–1092, <https://doi.org/10.1063/1.1699114>, 1953.
- 515 Miller, I., Morgan, H., Mauger, G., Newton, T., Weldon, R., Schmidt, D., Welch, M., and Grossman, E.: Projected Sea Level Rise for Washington State - A 2018 Assessment., A collaboration of Washington Sea Grant, University of Washington Climate Impacts Group, University of Oregon, University of Washington, and US Geological Survey, <http://www.wacoastalnetwork.com/files/theme/wcrp/SLR-Report-Miller-et-al-2018.pdf>, 2018.
- Milne, G. and Mitrovica, J.: Postglacial sea-level change on a rotating Earth, *Geophysical Journal International*, 133, 1–19, 1998.
- 520 Mitrovica, J. and Milne, G.: On post-glacial sea level: I. General theory, *Geophysical Journal International*, 154, 253–267, 2003.
- Mitrovica, J. and Peltier, W.: On post-glacial geoid subsidence over the equatorial ocean, *J. Geophys. Res.*, 96, 20053–20071, 1991.
- Mitrovica, J. X., Gomez, N., and Clark, P. U.: The Sea-Level Fingerprint of West Antarctic Collapse, *Science*, 323, 753, <https://doi.org/10.1126/science.1166510>, 2009.
- Mitrovica, J. X., Gomez, N., Morrow, E., Hay, C., Latychev, K., and Tamisiea, M. E.: On the Robustness of Predictions of Sea Level Fingerprints, *Geophysical Journal International*, 187, 729–742, <https://doi.org/10.1111/j.1365-246X.2011.05090.x>, 2011.
- 525 Mitrovica, J. X., Hay, C. C., Kopp, R. E., Harig, C., and Latychev, K.: Quantifying the Sensitivity of Sea Level Change in Coastal Localities to the Geometry of Polar Ice Mass Flux, *Journal of Climate*, 31, 3701–3709, <https://doi.org/10.1175/JCLI-D-17-0465.1>, 2018.
- Padman, L., Siegfried, M. R., and Fricker, H. A.: Ocean tide influences on the Antarctic and Greenland ice sheets, *Reviews of Geophysics*, 56, 142–184, <https://doi.org/10.1002/2016RG000546>, 2018.
- 530 Ray, R. D.: Ocean self-attraction and loading in numerical tidal models, *Marine Geodesy*, 21, 181–192, <https://doi.org/10.1080/01490419809388134>, 1998a.
- Ray, R. D.: Ocean self-attraction and loading in numerical tidal models, *Marine Geodesy*, 21, 181–192, <https://doi.org/10.1080/01490419809388134>, 1998b.

- Richter, A., Groh, A., Horwath, M., Ivins, E., Marderwald, E., Hormaechea, J. L., Perdomo, R., and Dietrich, R.: The Rapid and Steady Mass  
535 Loss of the Patagonian Icefields throughout the GRACE Era: 2002–2017, *Remote Sensing*, 11, 909, <https://doi.org/10.3390/rs11080909>,  
2019.
- Richter, K., Riva, R., and Drange, H.: Impact of Self-Attraction and Loading Effects Induced by Shelf Mass Loading on Projected Regional  
Sea Level Rise, *Geophysical Research Letters*, 40, 1144–1148, <https://doi.org/10.1002/grl.50265>, 2013.
- Schlegel, N.-J., Larour, E., Seroussi, H., Morlighem, M., and Box, J. E.: Decadal-scale sensitivity of Northeast Greenland ice flow to errors  
540 in surface mass balance using ISSM, *J. Geophys. Res. - Earth Surface*, 118, 1–14, <https://doi.org/10.1002/jgrf.20062>, <http://dx.doi.org/10.1002/jgrf.20062>, 2013.
- Schlegel, N.-J., Larour, E., Seroussi, H., Morlighem, M., and Box, J. E.: Ice discharge uncertainties in Northeast Greenland from boundary  
conditions and climate forcing of an ice flow model, *J. Geophys. Res. - Earth Surface*, 120, 29–54, <https://doi.org/10.1002/2014JF003359>,  
<http://dx.doi.org/10.1002/2014JF003359>, 2015.
- 545 Schlegel, N.-J., Wiese, D. N., Larour, E. Y., Watkins, M. M., Box, J. E., Fettweis, X., and van den Broeke, M. R.: Application of GRACE  
to the assessment of model-based estimates of monthly Greenland Ice Sheet mass balance (2003–2012), *Cryosphere*, 10, 1965–1989,  
<https://doi.org/10.5194/tc-10-1965-2016>, <http://www.the-cryosphere.net/10/1965/2016/>, 2016.
- Schlegel, N.-J., Seroussi, H., Schodlok, M. P., Larour, E. Y., Boening, C., Limonadi, D., Watkins, M. M., Morlighem, M., and van den  
Broeke, M. R.: Exploration of Antarctic Ice Sheet 100-year contribution to sea level rise and associated model uncertainties using the  
550 ISSM framework, *Cryosphere*, 12, 3511–3534, <https://doi.org/10.5194/tc-12-3511-2018>, 2018.
- Seroussi, H., Nakayama, Y., Larour, E., Menemenlis, D., Morlighem, M., Rignot, E., and Khazendar, A.: Continued retreat of  
Thwaites Glacier, West Antarctica, controlled by bed topography and ocean circulation, *Geophys. Res. Lett.*, 44, 6191–6199,  
<https://doi.org/10.1002/2017GL072910>, <http://dx.doi.org/10.1002/2017GL072910>, 2017GL072910, 2017.
- Slangen, A. B. A., Katsman, C. A., van de Wal, R. S. W., Vermeersen, L. L. A., and Riva, R. E. M.: Towards Regional Projections of Twenty-  
555 First Century Sea-Level Change Based on IPCC SRES Scenarios, *Climate Dynamics*, 38, 1191–1209, <https://doi.org/10.1007/s00382-011-1057-6>, 2012.
- Slangen, A. B. A., Adloff, F., Jevrejeva, S., Leclercq, P. W., Marzeion, B., Wada, Y., and Winkelmann, R.: A Review of Recent Updates of  
Sea-Level Projections at Global and Regional Scales, *Surveys in Geophysics*, 38, 385–406, <https://doi.org/10.1007/s10712-016-9374-2>,  
2017.
- 560 Spada, G. and Stocchi, P.: SELEN: A Fortran 90 program for solving the 'sea-level equation?', *Comput. Geosci.*, 33, 538–562, 2007.
- Stepanov, V. N. and Hughes, C. W.: Parameterization of ocean self-attraction and loading in numerical models of the ocean circulation:  
PARAMETERIZATION OF OCEAN LOADING, *Journal of Geophysical Research: Oceans*, 109, <https://doi.org/10.1029/2003jc002034>,  
2004.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: A summary of the CMIP5 experiment design, PCDMI, pp. 1–33, [https://cmip.llnl.gov/cmip5/  
565 docs/Taylor\\_CMIP5\\_design.pdf](https://cmip.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf), 2009.
- Thompson, P. R., Hamlington, B. D., Landerer, F. W., and Adhikari, S.: Are long tide gauge records in the wrong place to measure global  
mean sea level rise?, *Geophysical research letters*, 2016.
- Vinogradova, N. T., Ponte, R. M., Quinn, K. J., Tamisiea, M. E., Campin, J.-M., and Davis, J. L.: Dynamic Adjustment of the Ocean  
Circulation to Self-Attraction and Loading Effects, *Journal of Physical Oceanography*, 45, 678–689, [https://doi.org/10.1175/jpo-d-14-  
570 0150.1](https://doi.org/10.1175/jpo-d-14-0150.1), 2015.

- Wessel, P. and Smith, W. H. F.: A Global, Self-Consistent, Hierarchical, High-Resolution Shoreline Database, *Journal of Geophysical Research: Solid Earth*, 101, 8741–8743, <https://doi.org/10.1029/96JB00104>, 1996.
- Whitehouse, P. L., Bentley, M. J., Milne, G. A., King, M. A., and Thomas, I. D.: A new glacial isostatic adjustment model for Antarctica: calibrated and tested using observations of relative sea-level change and present-day uplift rates, *Geophysical Journal International*, 190, 1464–1482, 2012.
- Zwally, H. J., Giovinetto, M. B., Beckley, M. A., and Saba, J. L.: Antarctic and Greenland Drainage Systems, Tech. rep., GSFC Cryospheric Sciences Laboratory, [http://icesat4.gsfc.nasa.gov/cryo\\_data/ant\\_grn\\_drainage\\_systems.php](http://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php), 2012.