

BFM17 v1.0: A Reduced Biogeochemical Flux Model for Upper Ocean Biophysical Simulations

Katherine M. Smith¹, Skyler Kern¹, Peter E. Hamlington¹, Marco Zavatarelli², Nadia Pinardi², Emily F. Klee³, and Kyle E. Niemeyer³

¹ Paul M. Rady Department of Mechanical Engineering, University of Colorado, Boulder, CO, USA

² Department of Physics and Astronomy, University of Bologna, Bologna, IT

³ School of Mechanical, Industrial, and Manufacturing Engineering, Oregon State University, Corvallis, OR, USA

Correspondence: K. M. Smith (kmsmith@lanl.gov)

Abstract. We present a newly developed upper-thermocline, open-ocean biogeochemical flux model that is complex and flexible enough to capture open-ocean ecosystem dynamics, but reduced enough to incorporate into highly resolved numerical simulations and parameter optimization studies with limited additional computational cost. The model, which is derived from the full 56 state variable Biogeochemical Flux Model (BFM56; Vichi et al. (2007)), follows a biological and chemical functional group approach and allows for the development of critical non-Redfield nutrient ratios. Matter is expressed in units of carbon, nitrogen, and phosphate, following techniques used in more complex models. To reduce the overall computational cost and to focus on upper-thermocline, open-ocean, and non-iron or silicate limited conditions, the reduced model eliminates certain processes, such as benthic, silicate, and iron influences, and parameterizes others, such as the bacterial loop. The model explicitly tracks 17 state variables, divided into phytoplankton, zooplankton, dissolved organic matter, particulate organic matter, and nutrient groups. It is correspondingly called the Biogeochemical Flux Model 17 (BFM17). After describing BFM17, we couple it with the one-dimensional (1D) Princeton Ocean Model (POM) for validation using observational data from the Sargasso Sea. ~~Results show good agreement with the observational data and~~ The results agree closely with observational data, giving correlations above 0.85, except for chlorophyll (0.63) and oxygen (0.37), as well as with corresponding results from BFM56, with correlations above 0.85, except for oxygen (0.56), including the ability to capture the subsurface chlorophyll maximum and bloom intensity. In comparison to previous models of similar size, BFM17 provides improved correlations between several model output and observational data, indicating ~~that significant improvements in the~~ reproduction of *in situ* data can be achieved with a low number of variables, while maintaining the functional group approach. Notable additions to BFM17 over similar complexity models are the explicit tracking of dissolved oxygen, allowance for non-Redfield nutrient ratios, and both dissolved and particulate organic matter, all within the functional group framework.

1 Introduction

Biogeochemical (BGC) tracers and their interactions with upper-ocean physical processes, from basin-scale circulations to millimeter-scale turbulent dissipation, are critical for understanding the role of the ocean in the global carbon cycle. These interactions cause multi-scale spatial and temporal heterogeneity in tracer distributions (Strass, 1992; Yoder et al., 1992; Jr.

et al., 2001; Gower et al., 1980; Denman and Abbott, 1994; Strutton et al., 2012; Clayton, 2013; Abraham, 1998; Bees, 1998; Mahadevan and Archer, 2000; Mahadevan and Campbell, 2002; Levy and Klein, 2015; Powell and Okubo, 1994; Martin et al., 2002; Mahadevan, 2005; Tzella and Haynes, 2007) that can greatly affect carbon exchange rates between the atmosphere and interior ocean, net primary productivity, and carbon export (Lima et al., 2002; Schneider et al., 2007; Hauri et al., 2013; Behrenfeld, 2014; Barton et al., 2015; Boyd et al., 2016). There are still significant gaps, however, in our understanding of how these biophysical interactions develop and evolve, thus limiting our ability to accurately predict critical exchange rates.

Better understanding these interactions requires accurate physical and BGC models that can be coupled together. The exact equations that describe the physics (e.g., the Navier–Stokes or Boussinesq equations) are often known and physically accurate solutions can be obtained given sufficient spatial resolution and computational resources. Due to the vast diversity and complexity of ocean ecology, however, even when only considering the lowest trophic levels, accurately modeling BGC processes can be quite difficult. Put simply, there are no known first-principles governing equations for ocean biology.

As such, two different approaches to modeling BGC processes are often used when faced with this challenge. The first is to increase model complexity and include equations for every known BGC process. Often, these models include species functional types or multiple classes of phytoplankton and/or zooplankton that each serve specific functional roles within the ecosystem, such as calcifiers or nitrogen fixers. The justification for this approach is that particular phytoplankton and zooplankton groups serve as important system feedback pathways, and that without explicit representation of these feedbacks, there is little hope of accurately representing the target ecosystem (Doney, 1999; Anderson, 2005). In many cases, these models also contain variable intra- and extra-cellular nutrient ratios, which are important when accounting for different nutrient regimes within the global ocean and species diversity of non-Redfield nutrient ratio uptake (Dearman et al., 2003).

Although these more complex models are typically highly adaptable and are often able to capture different dynamics than those for which they were calibrated (Blackford et al., 2004; Friedrichs et al., 2007), they for (Blackford et al., 2004; Friedrichs et al., 2007), these more complex models contain many more parameters than their simplified counterparts. Moreover, many of the parameters, such as phytoplankton mortality, zooplankton grazing rates, and bacterial remineralization rates, are inadequately bounded by either observational or experimental data (Denman, 2003). Because of the increased complexity of such models, it is also often difficult to ascertain which processes are responsible for the development of a particular event (e.g., a phytoplankton bloom), and so these models can be ill-suited for process studies. Lastly, while these highly complex models are regularly used within global Earth System Models (ESMs), they are typically prohibitively expensive to integrate within high-fidelity, high-resolution physical models at submesoscales, such as. Examples of such models are those used to enhance fundamental understanding of subgrid-scale (SGS) physics in ESMs and to assist in the development of new SGS parameterizations (Roedel et al., 2012; Hamlington et al., 2014; Suzuki and Fox-Kemper, 2015; Smith et al., 2016, 2018).

In broad terms, the second common BGC modeling approach is focused on substantially decreasing model complexity and severely truncating the number of equations used to describe the dynamics of an ecosystem. Such approaches include the well-known nutrient-phytoplankton-zooplankton-detritus class of models. These models have significantly fewer unknown parameters and can be more easily integrated within complex physical models. Their simplicity also enables greater transparency when attempting to understand the dominant forcing or dynamics underlying a particular event. While they are often capable

of reproducing the overall distributions of chlorophyll, primary production, and nutrients (Anderson, 2005), such simplified models have been shown to under-perform at capturing complex ecosystem dynamics, and often struggle in regions of the ocean for which they were not calibrated (Friedrichs et al., 2007).

Although both of these general BGC modeling approaches have their respective advantages, particularly given their different objectives, the ~~disconnect~~difference between lower-complexity BGC models used in small-scale studies and the more complex BGC models used in global ESMs poses a problem. In particular, the difficulty in directly comparing the two types of models makes the process of “scaling-up” newly developed parameterizations or “downscaling” BGC variables within nested-grid studies much more challenging. This motivates the need for a new BGC model that is reduced enough to be usable within high-resolution, high-fidelity physical simulations for process and parameter optimization studies and parameterization development, but is still complex enough to capture important ecosystem feedback dynamics, as well as the dynamics of vastly different ecosystems throughout the ocean, as required by ESMs.

To begin addressing this need, here we present a new upper-thermocline, open-ocean, 17 state-variable Biogeochemical Flux Model (BFM17) obtained by reducing the larger 56 state variable Biogeochemical Flux Model (BFM56) developed by Vichi et al. (2007). Most high-fidelity, high-resolution physical models are capable of integrating 17 additional tracer equations with limited additional computational cost. Following the approach used in BFM56 (Vichi et al., 2007, 2013), a biological and chemical functional family (CFF) approach underlies BFM17, ~~permitting variable non-Redfield intra- and extra-cellular nutrient ratios, and where~~ matter is exchanged in the model through units of carbon, nitrate, and phosphate. This permits variable non-Redfield intra- and extra-cellular nutrient ratios. Most notably, BFM17 includes a phosphate budget, the importance of which has historically been under-appreciated even though observational data has indicated its potential importance as a limiting nutrient, particularly in the Atlantic Ocean (Ammerman et al., 2003). To reduce model complexity, we parameterize certain processes for which field data are lacking, such as bacterial remineralization.

In the present study, we outline, in detail, the formulation of BFM17 and its development from BFM56. We couple BFM17 to the 1D Princeton Ocean Model (POM) and validate the model for upper-thermocline, open-ocean conditions using observational data from the Sargasso Sea. We also compare results from BFM17 and the larger BFM56 for the same upper-thermocline, open-ocean conditions. As a result of the focus on upper-thermocline, open-ocean conditions, further assumptions have been made in ~~obtaining~~deriving BFM17 from BFM56, such as the exclusion of any representation for the benthic system and the absence of limiting nutrients such as iron and silicate.

It should be noted that the primary focus of the present study is to ~~demonstrate~~introduce the viability of BFM17 as an accurate BGC model for high-resolution, high-fidelity simulations of the upper ocean used in process, parameterization, and parameter optimization studies. This is accomplished here by comparing results from BFM17 to results from observations and BFM56; as such, here we only consider one open-ocean location (i.e., the Sargasso Sea). Although the model must also be applied at other locations to determine its general applicability, its ability ~~, demonstrated herein,~~ to reproduce important and difficult key behaviors in the Sargasso Sea ~~, such as the initial spring bloom and subsequent subsurface chlorophyll maxima,~~ supports its use as a process study model. The correspondence between BFM17 and the more general BFM56 also provides confidence that the reduced model will prove effective at modeling other ocean locations and conditions, and exploring the

range of applicability of BFM17 remains an important direction for future research. We also emphasize that relatively limited calibration of BFM17 parameters has been performed in the present study. Most parameters are set to their values used in the larger BFM56 (Vichi et al., 2007, 2013), and optimization of these parameters over a range of ocean conditions is another important direction of future research, for which BFM17 is ideally suited.

Finally, we note that other similarly complex BGC models have been calibrated using data from the Sargasso Sea, such as those developed in Levy et al. (2005), Ayata et al. (2013), Spitz et al. (2001), Doney et al. (1996), Fasham et al. (1990), Fennel et al. (2001), Hurtt and Armstrong (1996), Hurtt and Armstrong (1999), and Lawson et al. (1996). However, each of these models employs less than 10 species and none uses a CFF approach or includes oxygen, a tracer that is historically difficult to predict. Although some of these models employ data assimilation techniques (e.g., Spitz et al. (2001)) and produce relatively accurate results, most leave room for improvement. With a minimal increase in the number and complexity of the model equations, such as those associated with tracking phosphate in addition to carbon and nitrate, and by including both particulate and dissolved organic nutrient budgets, we ~~postulate-anticipate~~ can-might that a significant increase in model accuracy and applicability can-might be achieved over previous models of similar complexity. Additionally, with this increase in model complexity, the disparate gap between the complexity of BGC models used in small- and global-scale studies is reduced, thereby simplifying up- and down-scaling efforts. This last point is emphasized here by the good agreement between results from BFM17 and BFM56.

In the following, BFM17 is introduced in Section 2, with detailed equations and parameter values provided in Appendix A. Results from a zero-dimensional (0D) test of BFM17 is provided in Appendix B. In Section 3, BFM17 is coupled to the 1D POM physical model. A discussion of the methods used to calibrate and validate the model with observational data collected in the Sargasso Sea is presented in Section 4. Model results, a skill assessment, a comparison to results from BFM56, and a brief comparison to other similar BGC models are discussed in Section 5.

2 Biogeochemical Flux Model 17 (BFM17)

The 17 state equation BFM17 is an upper-thermocline, open-ocean BGC model derived from the original 56 state equation BFM56 (Vichi et al., 2007, 2013), which is based on the CFF approach. In this approach, functional groups are partitioned into living organic, non-living organic, and non-living inorganic CFFs, and exchange of matter occurs through constituent units of carbon, nitrogen, and phosphate. To date, there are no other BGC models with this order of reduced complexity using the CFF approach, making BFM17 unique and able to accurately reproduce complex ecosystem dynamics.

BFM17 is a pelagic model intended for oligotrophic regions that are not iron or silicate limited, and is obtained from the more-complete BFM56 by omitting quantities and processes assumed to be of ~~lesser significance~~ minor importance in these regions, ~~subject to the constraint that variable internal nutrient dynamics are of continued importance~~. We have developed BFM17 primarily for use with high-resolution, high-fidelity numerical simulations, including large eddy simulations (LES) used in process, parameterization, and parameter optimization studies. As such, we do not validate the efficacy of BFM17 as a global BGC model, and note that it is missing potentially important processes for such an application, which we elaborate on

shortly. We also note that we compare BFM17 to the original BFM56 in Section 5 to demonstrate that, although it is reduced in complexity, BFM17 is equally appropriate for use in seasonal process, parameterization, and optimal parameter estimation studies for which a more complex model such as BFM56 may be too computationally expensive. Nevertheless, given the
130 agreement between the BFM17 and BFM56 results in Section 5, there is reason to believe that BFM17 may have potential as a global BGC model, and the examination of the broader applicability of BFM17 is an important direction for future research.

In BFM17, the living organic CFF is comprised of single phytoplankton and zooplankton living functional groups (LFGs); these two groups are the bare minimum needed within a BGC model and already account for six state equations (corresponding to carbon, nitrogen, and phosphate constituents of both groups). The baseline parameters used in BFM17 are those detailed
135 in Vichi et al. (2007), and a complete list of the model parameters is provided in Appendix A. Parameters used in the representation of phytoplankton loosely correspond to the flagellate LFG in BFM56, while the zooplankton parameters correspond to the micro-zooplankton LFG. The only relevant difference with respect to Vichi et al. (2007) is related to the choice of the phytoplankton specific photosynthetic rate ($r_P^{(0)}$ in Table A3 of Appendix A); in this case, the new value was chosen according to the control laboratory cultures of Fiori et al. (2012).

140 Within BFM17, we track chlorophyll, dissolved oxygen, phosphate, nitrate, and ammonium, since their distributions and availability can greatly enhance or hinder important biological and chemical processes. Dissolved oxygen is of particular interest, because it is historically difficult to predict using BGC models of any complexity. This is likely due, in part, to missing physical processes in the mixing parameterizations used in global and column models. This provides motivation for the present study, since a primary goal in the development of BFM17 is to create a BGC model that can be used in combination with
145 high-resolution, high-fidelity physical models (e.g., those found in LES) to understand the effects of these physical processes and how they can be more accurately represented in mixing parameterizations.

Dissolved and particulate organic matter, each with their own partitions of carbon, nitrogen, and phosphate, are also included in BFM17 to account for nutrient recycling and carbon export due to particle sinking. Another primary goal of developing BFM17 is to explore how spatially decoupled (or “patchy”) processes, such as the sinking of organic matter and the subsequent
150 upwelling of multiple recycled nutrients (not just nitrate) affect the fate and distribution of a phytoplankton bloom.

Lastly, remineralization of nutrients is provided by parameterized bacterial closure terms, thereby reducing complexity while still maintaining critical nutrient recycling. The related parameter values (see Table A5 in Appendix A) were chosen according to Mussap et al. (2016), who carried out sensitivity tests to evaluate the many parameters values found in the literature.

Iron is omitted from BFM17, limiting the applicability of the model in regions where iron components are important, such
155 as the Southern Ocean and the tropical Pacific. Thus, if used in such regions, at least a fixed concentration of iron may be needed (although this method has not yet been validated within BFM17). Top-down control of the ecosystem [in the form of explicit predation of zooplankton](#) is also not included. Instead, a simple constant zooplankton mortality is used, as this is a complicated process and understanding where to add this closure and where to feed the particulate and dissolved nutrients from this process in a lower-complexity model is not well understood. However, the addition of a top-down closure term was tested,
160 and no major differences were observed in the model results. Consequently, it was assumed that the constant mortality term was sufficient for this model, similar to other models of this complexity (Fasham et al., 1990; Lawson et al., 1996; Clainche et al.,

Table 1. Notation used for the 17 state variables in the BFM17 model, as well as the chemical functional family (CFF), units, description, and rate equation reference for each state variable. CFFs are divided into living organic (LO), non-living organic (NO), and inorganic (IO) families.

Symbol	CFF	Units	Description	Equation
P_C	LO	mg C m^{-3}	Phytoplankton carbon	(A5)
P_N	LO	mmol N m^{-3}	Phytoplankton nitrogen	(A6)
P_P	LO	mmol P m^{-3}	Phytoplankton phosphorus	(A7)
P_{chl}	LO	$\text{mg Chl-}a \text{ m}^{-3}$	Phytoplankton chlorophyll	(A8)
Z_C	LO	mg C m^{-3}	Zooplankton carbon	(A31)
Z_N	LO	mmol N m^{-3}	Zooplankton nitrogen	(A32)
Z_P	LO	mmol P m^{-3}	Zooplankton phosphorus	(A33)
$R_C^{(1)}$	NO	mg C m^{-3}	Dissolved organic carbon	(A41)
$R_N^{(1)}$	NO	mmol N m^{-3}	Dissolved organic nitrogen	(A42)
$R_P^{(1)}$	NO	mmol P m^{-3}	Dissolved organic phosphorus	(A43)
$R_C^{(2)}$	NO	mg C m^{-3}	Particulate organic carbon	(A44)
$R_N^{(2)}$	NO	mmol N m^{-3}	Particulate organic nitrogen	(A45)
$R_P^{(2)}$	NO	mmol P m^{-3}	Particulate organic phosphorus	(A46)
O	IO	$\text{mmol O}_2 \text{ m}^{-3}$	Dissolved oxygen	(A47)
$N^{(1)}$	IO	mmol P m^{-3}	Phosphate	(A48)
$N^{(2)}$	IO	mmol N m^{-3}	Nitrate	(A49)
$N^{(3)}$	IO	mmol N m^{-3}	Ammonium	(A50)

2004). Additionally, the benthic system within BFM56 (Mussap et al., 2016) has been removed. It is assumed that within the upper thermocline of the open ocean, the ecosystem is not substantially influenced by a benthic system and any water-column influences from depth can be taken into account using boundary conditions (such as those discussed in Section 4). As such, we cannot attest to the accuracy of BFM17 in shelf or coastal regions.

In summary, notable novel attributes of BFM17, in comparison to other models of comparable complexity, are the use of (i) CFFs for living organisms, including two LFGs for phytoplankton and zooplankton, (ii) CFFs for both particulate and dissolved organic matter, (iii) a full nutrient profile (i.e., phosphate, nitrate, and ammonium), and (iv) the tracking of dissolved oxygen. A summary of the 17 state variables tracked in BFM17 is provided in Table 1, and a schematic of the CFFs and LFGs used in BFM17, along with their interactions, is shown in Figure 1. The detailed equations comprising BFM17, as well as all associated parameter values, are presented in Appendix A. Results from an initial 0D test of BFM17 are provided in Appendix B.




Figure 1. Schematic of the 17 state equation BFM17 model. The dissolved organic matter, particulate organic matter, and living organic matter chemical functional families (CFFs) are each comprised of three chemical constituents (i.e., carbon, nitrogen, and phosphorus). The living organic CFF is further subdivided into phytoplankton and zooplankton living functional groups (LFGs).

3 Coupled Physical-Biogeochemical Flux Model

As a demonstration of BFM17 for predicting ocean biogeochemistry in oligotrophic pelagic zones, here we couple the model to a 1D physical mixing parameterization and make comparisons with available observational data in the Sargasso Sea. In order to focus on the upper-thermocline, open-ocean regime for which BFM17 was developed, the physical model only extends 150 m in depth and diagnostically calculates diffusivity terms based upon prescribed temperature and salinity profiles from the observations. While a 1D physical model is unlikely to resolve all processes relevant for biogeochemistry in the upper thermocline, we have made additions, such as large-scale general circulation and mesoscale eddy vertical velocities, as well as relaxation bottom boundary conditions for nutrient upwelling, to better represent missing processes.

For all equations here and in Appendix A, we adopt the same notation style used for BFM56 in Vichi et al. (2007), Mussap et al. (2016), and the BFM User Manual (Vichi et al., 2013) for consistency and clarity. The coupled physical and BGC model is a time-

Table 2. Values, units, and descriptions for parameters used in the combined physical–BFM17 model.

Symbol	Value	Units	Description
$v^{(\text{set})}$	-1.00	m d ⁻¹	Settling velocity of particulate detritus
W	-0.02 – 0	m d ⁻¹	Imposed general circulation vertical velocity
W_E	0 – 0.1	m d ⁻¹	Imposed mesoscale circulation vertical velocity
λ_O	0.06	m d ⁻¹	Relaxation constant for oxygen at bottom
$\lambda_{N(1)}$	0.06	m d ⁻¹	Relaxation constant for phosphate at bottom
$\lambda_{N(2)}$	0.06	m d ⁻¹	Relaxation constant for nitrate at bottom

depth model that integrates in time the generic equation for all biological state variables, denoted A_j , given by

$$\frac{\partial A_j}{\partial t} = \frac{\partial A_j}{\partial t} \Big|_{\text{bio}} - \left[W + W_E + v^{(\text{set})} \right] \frac{\partial A_j}{\partial z} + \frac{\partial}{\partial z} \left(K_H \frac{\partial A_j}{\partial z} \right), \quad (1)$$

where the A_j are the 17 state variables of BFM17, the first term on the right-hand side accounts for sources and sinks within
185 each species due to biological and chemical reactions ~~;~~ (as represented by the equations comprising BFM17 ~~;~~ and outlined in
Appendix A), W and W_E are the vertical velocities due to large-scale general circulation and mesoscale eddies, respectively,
 $v^{(\text{set})}$ is the settling velocity, and K_H is the vertical eddy diffusivity. Although the BFM17 formulation and model results are
the primary focus of the present study, we also perform coupled physical-BGC simulations using BFM56 for comparison.
Equation (1) applies to all 17 state variables in BFM17, as well as to all 56 state variables in BFM56. Consequently, the only
190 differences between the biophysical models with BFM17 and BFM56 are the number of state variables being tracked and the
equations used to calculate the biological forcing terms. The specific forms of Eq. (1) for each of the 17 species in BFM17 are
discussed in Appendix A, and the specific forms of this equation for each of the 56 species in BFM56 were previously discussed
in Vichi et al. (2007). The parameters used in BFM56 correspond to the values provided in Tables A3–A5 of Appendix A, with
the remaining undefined parameters (since BFM56 includes many more model parameters than BFM17) based on values from
195 Mussap et al. (2016).

The ~~vertical velocities range of values for W and W_E in Eq. (1) are the large-scale general circulation and mesoscale eddy~~
~~vertical velocities, respectively. The range of values for each of these velocities are included in Table 2 and the corresponding~~
depth profiles are discussed in Section 4.3. The settling velocity, $v^{(\text{set})}$, in Eq. (1) is only non-zero for the three constituents of
particulate organic matter, and its value is given in Table 2. We assume $v^{(\text{set})} = 0$ for zooplankton, since zooplankton actively
200 swim and oppose their own sinking velocity. Finally, K_H in Eq. (1) is ~~the vertical eddy diffusivity term~~ calculated by the
model, and is described in more detail later in this section.

To obtain the complete 1D biophysical model, BFM17 has been coupled with a modification of the three-dimensional (3D)
Princeton Ocean Model (POM) (Blumberg and Mellor, 1987) that considers only the vertical (specifically, the upper 150 m of
the water column) and time dimensions; that is, the evolution of the system in the (z, t) space. It is well known that the primary

205 calibration dimension in marine ocean biogeochemistry is along the vertical direction, as shown in several previous calibration and validation exercises (Vichi et al., 2003; Triantafyllou et al., 2003; Mussap et al., 2016).

The 1D POM solver (POM-1D) is used to calculate the vertical structure of the two horizontal velocity components, denoted U and V , the potential temperature, T , salinity, S , density, ρ , turbulent kinetic energy, $q^2/2$, and mixing length scale, ℓ . In this model adaptation, vertical temperature and salinity profiles are imposed from given climatological monthly profiles [obtained from observations](#), as previously done in Mussap et al. (2016) and Bianchi et al. (2005). ~~The model computes only POM-1D directly computes~~ the time evolution of the horizontal velocity components, the turbulent kinetic energy and the mixing length scale, all of which are used to compute the turbulent diffusivity term, K_H , required in Eq. (1). In this configuration, POM-1D is called “diagnostic” since temperature and salinity are prescribed. Furthermore, pressure effects are neglected in the density equation and the buoyancy gradients and temperature are used in place of potential temperature since we consider only the upper water column.

[A detailed description of POM can be found in Blumberg and Mellor \(1987\), and in the following we simply provide a description of the physical model and equations solved in POM-1D.](#) In diagnostic mode, [as used in the present study](#), POM-1D solves the momentum equations for U and V given by

$$\frac{\partial U}{\partial t} - fV = \frac{\partial}{\partial z} \left(K_M \frac{\partial U}{\partial z} \right), \quad (2)$$

$$220 \quad \frac{\partial V}{\partial t} + fU = \frac{\partial}{\partial z} \left(K_M \frac{\partial V}{\partial z} \right), \quad (3)$$

where $f = 2\Omega \sin \phi$ is the Coriolis force, Ω is the angular velocity of the Earth, and ϕ is the latitude. The vertical viscosity K_M and diffusivity K_H are calculated using the closure hypothesis of Mellor and Yamada (1982) as

$$K_M = ql S_M, \quad (4)$$

$$K_H = ql S_H, \quad (5)$$

225 where q is the turbulent velocity and S_H and S_M are stability functions written as

$$S_M [1 - 9A_1 A_2 G_H] - S_H [(18A_1^2 + 9A_1 A_2) G_H] = A_1 [1 - 3C_1 - 6A_1/B_1], \quad (6)$$

$$S_H [1 - (3A_2 B_2 + 18A_1 A_2) G_H] = A_2 [1 - 6A_1/B_1]. \quad (7)$$

The coefficients in the above expressions are $(A_1, B_1, A_2, B_2, C_1) = (0.92, 16.6, 0.74, 10.1, 0.08)$, with

$$G_H = \frac{l^2}{q^2} \frac{g}{\rho_0} \frac{\partial \rho}{\partial z}, \quad (8)$$

230 where $\rho_0 = 1025 \text{ kg m}^{-3}$, $g = 9.81 \text{ m s}^{-2}$. Following Mellor (2001), G_H is limited to have a maximum value of 0.028. The equation of state relating ρ to T and S is nonlinear (Mellor, 1991) and given by

$$\begin{aligned} \rho = & 999.8 + (6.8 \times 10^{-2} - 9.1 \times 10^{-3}T + 1.0 \times 10^{-4}T^2 - 1.1 \times 10^{-6}T^3 + 6.5 \times 10^{-9}T^4) T \\ & + (0.8 - 4.1 \times 10^{-3}T + 7.6 \times 10^{-5}T^2 - 8.3 \times 10^{-7}T^3 + 5.4 \times 10^{-9}T^4) S \\ & + (-5.7 \times 10^{-3} + 1.0 \times 10^{-4}T - 1.6 \times 10^{-6}T^2) S^{1.5} + 4.8 \times 10^{-4} S^2, \end{aligned} \quad (9)$$

235 where the polynomial constants have been written only up to the first digit. For a more precise reproduction of these constants, the reader is referred to Mellor (1991). Finally, the governing equations solved to obtain the turbulence variables $q^2/2$ and ℓ are

$$\frac{\partial}{\partial t} \left(\frac{q^2}{2} \right) = \frac{\partial}{\partial z} \left[K_q \frac{\partial}{\partial z} \left(\frac{q^2}{2} \right) \right] + K_M \left[\left(\frac{\partial U}{\partial z} \right)^2 + \left(\frac{\partial V}{\partial z} \right)^2 \right] + \frac{g}{\rho_0} K_H \frac{\partial \rho}{\partial z} - \frac{q^3}{B_1 \ell}, \quad (10)$$

$$\frac{\partial}{\partial t} (q^2 \ell) = \frac{\partial}{\partial z} \left[K_q \frac{\partial}{\partial z} (q^2 \ell) \right] + E_1 \ell K_M \left[\left(\frac{\partial U}{\partial z} \right)^2 + \left(\frac{\partial V}{\partial z} \right)^2 \right] + E_1 \ell \frac{g}{\rho_0} K_H \frac{\partial \rho}{\partial z} - \frac{q^3}{B_1} \widetilde{W}, \quad (11)$$

240 where $K_q = \kappa K_H$ is the vertical diffusivity for turbulence variables, $\kappa = 0.4$ is the von Karman constant, and $\widetilde{W} = [1 + E_2 \ell^2 / \kappa^2 (1/|z| + 1/|z - H|)^2]$ with $(E_1, E_2) = (1.8, 1.33)$. In Eqs. (10) and (11), the time rate of change of the turbulence quantities is equal to the diffusion of turbulence (the first term on the right hand side of both equations), the shear and buoyancy turbulence production (second and third terms), and the dissipation (the fourth term). This is a second-order turbulence closure model that was formulated by Mellor (2001) as a particular case of the Mellor and Yamada (1982) model
245 for upper ocean mixing.

Boundary conditions for the horizontal velocities $\mathbf{U} = (U, V)$ and the turbulence quantities are

$$K_M \frac{\partial \mathbf{U}}{\partial z} \Big|_{z=0} = \boldsymbol{\tau}_w, \quad (12)$$

$$K_M \frac{\partial \mathbf{U}}{\partial z} \Big|_{z=z_{\text{end}}} = 0, \quad (13)$$

$$(q^2, q^2 \ell) \Big|_{z=0} = \left(B_1^{2/3} \frac{|\boldsymbol{\tau}_w|}{C_d}, 0 \right), \quad (14)$$

$$250 \quad (q^2, q^2 \ell) \Big|_{z=z_{\text{end}}} = 0, \quad (15)$$

where $\boldsymbol{\tau}_w = C_d |\mathbf{u}_w| \mathbf{u}_w$ is the surface wind stress, \mathbf{u}_w is the surface wind vector, C_d is a constant drag coefficient chosen to be 2.5×10^{-3} , and $z = 0$ and $z = z_{\text{end}}$ denote the locations of the surface and the greatest depth modeled, respectively.

For all variables except oxygen, surface boundary conditions for the coupled model variable A_j are

$$K_H \frac{\partial A_j}{\partial z} \Big|_{z=0} = 0. \quad (16)$$

255 By contrast, the surface boundary condition for oxygen has the form

$$K_H \frac{\partial O}{\partial z} \Big|_{z=0} = \Phi_O, \quad (17)$$

where Φ_O is the air-sea interface flux of oxygen computed according to Wanninkhof (1992, 2014). The bottom (i.e., greatest depth) boundary conditions for phytoplankton, zooplankton, dissolved organic matter, and particulate organic matter are

$$K_H \frac{\partial A_j}{\partial z} \Big|_{z=z_{\text{end}}} = 0. \quad (18)$$

260 This boundary condition was chosen since it allows removal of the scalar quantity A_j through the bottom boundary of the domain. This can be seen by integrating Eq. (1) over the boundary layer depth using the boundary condition above, giving

$$\frac{\partial}{\partial t} \int_{z=z_{\text{end}}}^{z=0} A_j dz = \left[W + W_E + v^{(\text{set})} \right] A_j \Big|_{z=z_{\text{end}}}, \quad (19)$$

where the biological part of Eq. (1) has been neglected and the resulting temporal change in the integrated scalar A_j is negative since $|(W + W_E)| < |v^{(\text{set})}|$, as shown in Table 2. For oxygen, phosphate, and nitrate, the bottom boundary conditions are

$$265 \quad K_H \frac{\partial A_j}{\partial z} \bigg|_{z=z_{\text{end}}} = \lambda_j \left(A_j|_{z=z_{\text{end}}} - A_j^* \right), \quad (20)$$

where λ_j and A_j^* are the corresponding relaxation velocity and observed at-bottom boundary climatological field data value, respectively, of that species. Base values for the relaxation velocities are included in Table 2. Lastly, the bottom boundary condition for ammonium is

$$K_H \frac{\partial N^{(3)}}{\partial z} \bigg|_{z=z_{\text{end}}} = 0. \quad (21)$$

270 Since observations of ammonium concentration in the observational area are not available, this choice is based on the assumption that the nitrogen diffusive flux from depth to the surface (euphotic) layers occurs mostly in the form of a nitrate flux, consistent with the concepts of “new” and “regenerated” production, as described by Dugdale and Goering (1967) and Mulholland and Lomas (2008).

4 Field Validation and Calibration Data

275 4.1 Study Site Description

Field data for calibration and validation of BFM17 are taken from the Bermuda Atlantic Time-series Study (BATS) (Steinberg et al., 2001) and the Bermuda Testbed Mooring (BTM) (Dickey et al., 2001) sites, which are located in the Sargasso Sea (31°40' N, 64°10' W) in the North Atlantic subtropical gyre. Both sites are a part of the US Joint Global Ocean Flux Study (JGOFS) program. Data has been collected from the BATS site since 1988 and from the BTM site since 1994.

280 Steinberg et al. (2001) provide an overview of the biogeochemistry in the general BATS and BTM area. Winter mixing allows nutrients to be brought up into the mixed layer, producing a phytoplankton bloom between January and March (winter mixed layer depth is typically 150-300 m). As thermal stratification intensifies over the summer months, this nutrient supply is cut off (summer mixed layer depth is typically 20 m). At this point, a subsurface chlorophyll maximum is observed near a depth of 100 m. Stoichiometric ratios of carbon, nitrate, and phosphate are often non-Redfield and, in contrast to many
285 oligotrophic regimes, phosphate is the dominant limiting nutrient (Fanning, 1992; Michaels et al., 1993; Cavender-Bares et al., 2001; Steinberg et al., 2001; Ammerman et al., 2003; Martiny et al., 2013; Singh et al., 2015).

4.2 Data Processing

The region encompassing the BATS and BTM sites is characterized as an open ocean, oligotrophic region that is phosphate limited. This region has thus been chosen for initial ~~calibration and validation testing~~ of BFM17 due to the prevalence of
290 oligotrophic regimes in the open ocean and to demonstrate the ability of BFM17 to capture difficult non-Redfield ratio regimes (which occur in phosphate-limited regions). The BATS/BTM data have also been collected over many years, providing long time series for model calibration and validation.

Data from the BATS/BTM area is used in the present study for two purposes: (i) as initial, boundary, and forcing conditions for the POM-1D biophysical simulations with BFM17 and BFM56, and (ii) as target fields for validation of the simulations.

295 In addition to the subsurface BATS data, we also use BTM surface data, such as the 10 m wind speed and PAR. For each observational quantity, we compute monthly averages over 27 years for the BATS data and 23 years (not continuous) for the BTM data. Additionally, we interpolate the BATS data to a vertical grid with 1 m resolution. We subsequently smooth the interpolated data, using a robust locally estimated scatterplot smoothing (LOESS) method, to maintain a positive buoyancy gradient, thereby eliminating any spurious buoyancy-driven mixing due to interpolation and averaging.

300 Figure 2 shows the monthly climatological profiles of temperature and salinity from the BATS data (maximum mixed layer depth from the climatology is approximately ~~125-149 m~~, which was calculated based upon a ~~$\Delta 0.02-0.2$ kg/m³ criteria~~ increase in density from the surface value), as well as the photosynthetically available radiation (PAR) and 10 m wind speed from the BTM data. Similar processing-The same monthly averaging, vertical interpolations, and smoothing used for the physical variables is also performed ~~on~~ for biological variables, which largely serve as target fields for the validation of BFM17.

305 4.3 Inputs to the Physical Model

The physical model computes density from the prescribed temperature and salinity, and surface wind stress from the 10 m wind speed; temperature, salinity, and wind speed are all provided by the BATS/BTM data. The model also uses this data in the turbulence closure to compute the turbulent viscosity and diffusivity. This diagnostic approach eliminates any drifts in temperature and salinity that might occur due to improper parameterizations of lateral mixing in a 1D model, therefore

310 providing greater reliability. In addition to the 10 m wind speed, temperature, and salinity, BFM requires monthly varying PAR at the surface. For all the monthly mean input data sets, a correction (Killworth, 1995) is applied. This correction is applied to the monthly averages to ~~account for monthly mean errors due to linear interpolation to the~~ reduce the errors incurred by linearly interpolating monthly averages to the much shorter model time step.

We imposed both general circulation, W , and mesoscale eddy, W_E , vertical velocities in the simulations. The imposed

315 vertical profiles of these velocities have been adapted from Bianchi et al. (2005), where the velocities are assumed to be zero at the surface and reach their maxima near the base of the Ekman layer, which is assumed to be at or below the bottom boundary of the simulations. The general large-scale upwelling/downwelling circulation, W , is due to Ekman pumping and is correspondingly given as

$$W = \hat{k} \cdot \nabla \times \left(\frac{\tau_w}{\rho f} \right), \quad (22)$$

320 where \hat{k} denotes the unit vector in the vertical direction. The monthly average value and sign of the wind stress curl, $\nabla \times \tau_w$, for the general BATS/BTM region was taken from the Scatterometer Climatology of Ocean Winds database (Risien and Chelton, 2008, 2011). The monthly value of W from Eq. (22) is then assumed to be the maximum, occurring at the base of the Ekman layer, for that particular month. Given the sign of the wind stress curl for the BATS/BTM region, a negative W was calculated, indicating general downwelling processes in this region. Seasonal profiles of W are shown in Figure 2(e).




Figure 2. Sargasso Sea physical variables, showing climatological monthly averaged (a) temperature, (b) 10 m surface wind speed, (c) surface PAR, and (d) salinity. Panel (e) shows the mean seasonal general circulation velocity, W , and panel (f) shows the bimonthly maximum value of the mesoscale eddy velocity W_E . Monthly averaged mixed layer depths (defined as the depth at which the density is 0.2 kg/m^3 greater than the surface density) are shown as black lines in panel (a).

Due to the prevalence of mesoscale eddies within the BATS/BTM region (Hua et al., 1985), which can provide episodic upwelling of nutrients to the upper water column, we also include an additional positive upwelling vertical velocity, W_E , which has a timescale of 15 days. The general profile of W_E is assumed to be the same as for W , with a value of zero at the surface and a maximum value at depth. However, there is no linear interpolation between each 15-day period and the maximum magnitude of W_E is randomized between 0 and 0.1 m d^{-1} , as shown in Figure 2(f) for each 15-day period.

4.4 Initial and Boundary Conditions

Although the BATS/BTM data includes information on many biological variables, initial conditions for only 5 of the 17 species within BFM17 could be extracted from the data. Similarly to the temperature and salinity, the initial chlorophyll, particulate organic nitrogen, oxygen, nitrate, and phosphate were interpolated to a mesh with 1 m vertical grid spacing, averaged over the initial month of January, and smoothed vertically in space to give the initial profiles seen in Figure 3(a). The remaining 12 state




Figure 3. Sargasso Sea initial and boundary conditions showing (a) initial profiles of nitrate, phosphate, particulate organic carbon, chlorophyll, and oxygen, where each profile, denoted $\phi_i(z)$, is normalized by its depth averaged value, $\langle\phi_i\rangle_z$, and (b) monthly bottom boundary conditions for nitrate, phosphate, and oxygen, where each quantity, $\phi_i(t)$, is normalized by its annual average value $\langle\phi_i\rangle_t$. The depth and annual averaged values are shown in parentheses in the legends of each panel. Units are mmol N/m³ for nitrate, mmol P/m³ for phosphate, mg C/m³ for particulate organic carbon, mg Chl/m³ for chlorophyll, and mmol O/m³ for oxygen.

335 variable initial conditions were determined either through the adoption of the Redfield ratio C:N:P \equiv 106:16:1 (Redfield et al., 2005), or assuming a reasonably low initial value. Since the 1D simulations were run to steady state over 10 years, memory of these initial states was assumed to be lost, with little effect on the results.

For the comparison of BFM17 to BFM56, the initial conditions for the additional state variables were calculated by splitting the total initial phytoplankton and zooplankton carbon values into additional-equal amounts for all phytoplankton and zoo-
 340 plankton groups. The other state variables for each group were again calculated using the Redfield ratio. The initial bacteria distribution was defined by setting the column equal to a constant value.

In both simulations, the bottom boundary conditions for oxygen, nitrate, and phosphate species are based on observed BATS data. Values are taken at the next closest data point below the bottom boundary (at 150 m) and then averaged over the month. Figure 3(b) shows the monthly average bottom boundary conditions for each of the three species.

345 5 Validation-Model Assessment Results

The coupled BFM17-POM1D model was run using the parameter values from Tables 2, A1, and A3-A5, which were decided on the basis of standard literature values (Vichi et al., 2007, 2003, 2013; Fiori et al., 2012). The simulations were allowed to run out to steady-state and multi-year monthly means were calculated as functions of depth for chlorophyll, oxygen, nitrate, phosphate, particulate organic nitrogen (PON), and net primary production (NPP), each of which were measured at the BATS/BTM site.




Figure 4. Comparison of target BATS fields (top row) to BFM17 simulation results (middle row) and BFM56 simulation results (bottom row) for (a,g,m) chlorophyll (mg Chl-a/m^3), (b,h,n) oxygen (mmol O/m^3), (c,i,o) nitrate (mmol N/m^3), (d,j,p) phosphate (mmol P/m^3), (e,k,q) particulate organic nitrogen (PON - mg N/m^3), (f,l,r) and net primary production (NPP - $\text{mg C/m}^3/\text{day}$). Simulation plots are multi-year, monthly averages of the last 3 years of a 10 year integration.

350 The model PON is defined as the sum of nitrogen contained within the phytoplankton, zooplankton, and particulate detritus, and NPP is defined as the net phytoplankton carbon uptake (or gross primary production) minus phytoplankton respiration.

Figure 4 qualitatively compares the BATS data (top row) with the results of from BFM17 (middle row). The model is able to capture the initial spring bloom between January and March brought on by physical entrainment of nutrients, the corresponding peak in net primary production and PON around the same time, and the subsequent subsurface chlorophyll maxima during the summer (evident in Figure 4 as a larger chlorophyll concentration at depths close to 100 m during the summer months). The predicted oxygen levels are lower than observed values, however, the overall structure predicted by BFM17 is quite similar-not completely dissimilar to that of the BATS oxygen field. These trends-results are consistent with the trends-seen-in-the-results-those from BFM56 (bottom row of Figure 4), suggesting that the two models are in generally close agreement. Correlation coefficients between the two models are 0.85 for chlorophyll, 0.56 for oxygen, 0.99 for nitrate,

355




Figure 5. Taylor diagram showing the normalized standard deviation, correlation coefficient, and normalized root mean squared differences between the BFM17 output and the BATS target fields. Observations lie at (1,0). Radial deviations from observations corresponds to the normalized root mean square error (RMSE), radial deviation from the origin correspond to the normalized standard deviation, and angular deviations from the vertical axis correspond to the correlation coefficient. BFM17 and BFM56 results are shown as colored circles and triangles, respectively (chlorophyll = blue, oxygen = orange, nitrate = yellow, phosphate = purple, PON = green, NPP = cyan). Note that BFM56 nitrate and phosphate data points fall on top of one another (yellow and purple triangles).

0.99 for phosphate, 0.95 for PON, and 0.97 for NPP. Differences in chlorophyll and oxygen are likely due to the removal in BFM17 of specific phytoplankton and zooplankton species in favor of general LFGs, to the removal of denitrification, and to the parameterization of remineralization using new closure terms that were calibrated to give reasonable agreement with the observational data.

As mentioned previously, oxygen is historically difficult to predict using BGC models of any complexity. It is likely that this is due, in part, to inaccuracies in the mixing parameterizations used in POM 1D and other physical models. For example, BFM17 struggles to accurately predict oxygen, in part, because the second-order mixing scheme of Mellor and Yamada (1982) lacks sufficient resolution of the winter mixing using just the monthly mean temperature and salinity. However, since it is often not included or presented at all in models of similar complexity to BFM17 (i.e., models reduced enough to reasonably couple to a high-fidelity, high-resolution physical model), studies that explore this hypothesis have been difficult to undertake. Thus, we include oxygen in BFM17 and present our results here to illustrate this exact point, and to lend motivation to developing and using a model such as BFM17 to study the effects of physical processes missing from mixing parameterizations and how they can be better represented.

To ~~quantitatively evaluate~~ obtain a first indication of the performance of BFM17, a model ~~skill~~ assessment was performed for each target field. The same ~~skill~~ assessment was performed for BFM56 to compare the two models. The results are summarized
 375 by the Taylor diagram in Figure 5. This diagram can be used to assess the extent of misfit between the models and observations by showing the normalized root mean square (RMS) errors, normalized standard deviation, and the correlation coefficient between each of the model outputs and the BATS target fields.

The normalized RMS errors were calculated as $\varepsilon_{\text{rms}}/\sigma_{\text{obs}}$, where ε_{rms} is the RMS error between the model and the obser-
 380 $\sigma_{\text{mod}}/\sigma_{\text{obs}}$ where σ_{mod} is the standard deviation of the model fields. The normalized RMS errors, normalized standard deviation, and the correlation coefficients each give an indication of the relative similarities in amplitude, variations in amplitude, and structure of each modeled field compared to the BATS target fields, respectively. For each variable, these statistics were calculated over all months and all depths shown in Figure 4.

The Taylor diagram in Figure 5 shows that BFM17 and BFM56 produce similar results. For most variables, errors in the
 385 amplitudes are within roughly one standard deviation of the observations. Additionally, the structure of the model fields for chlorophyll, nitrate, phosphate, PON, and NPP have high correlations with that of the BATS target fields. The correlation values range from 0.63 for chlorophyll to 0.94 for nitrate in BFM17 and from 0.60 for chlorophyll to 0.93 for phosphate and nitrate in BFM56. For BFM17, variability in amplitude for nitrate, phosphate, oxygen, and NPP are closest to that of the corresponding BATS target fields, while the chlorophyll and PON have too much variability. For BFM56, nitrate, phosphate,
 390 oxygen, and chlorophyll have similar variability in amplitude to the BATS data, while NPP and PON have too little and too much variability, respectively.

Table 3 provides a comparison of correlation coefficients and un-normalized RMS errors, calculated with respect to the observational fields, from BFM17 and BFM56, as well as from other models. Comparisons were only made to models that were calibrated using the same BATS/BTM data, employed some kind of parameter estimation technique, and reported correlation
 395 and RMS errors. Ayata et al. (2013) included six biological tracers, while both Fasham et al. (1990) and Spitz et al. (2001) included seven. The Spitz et al. (2001) study used data assimilation, while the Ayata et al. (2013) and Fasham et al. (1990) studies used only optimization to determine a select set of parameters. All models used climatological monthly mean forcing from the BATS region and reported climatological monthly means for their results. Care was taken to ensure that the same variable definition was compared between all models. Ayata et al. (2013) used a similar 1D physical model as was used here,
 400 while Spitz et al. (2001) and Fasham et al. (1990) used a time-dependent box model of the upper-ocean mixed layer. As such, correlations and RMS error values for comparison to Ayata et al. (2013) were computed over the entire domain (Ayata et al. (2013) calculated their metrics over the top 168 m of their domain). For comparison to Spitz et al. (2001) and Fasham et al. (1990), correlations and RMS errors were calculated only within the mixed layer (defined as the depth at which the density is 0.02-0.2 kg/m^3 greater than the surface density) and are shown as separate columns in Table 3.

~~Table 3 shows that BFM17 and BFM56 give comparable results, with BFM17 producing similar correlation coefficients and RMS error values to those from BFM56. The largest differences compared to the BATS data for both BFM17 and BFM56 are in the oxygen values. The overall slightly better correlation coefficient agreement of BFM17 with the BATS data results from~~

Table 3. Correlation coefficients (and RMS error in parenthesis) between BATS target fields and model data for BFM17, BFM56, and several example models of similar complexity. The first set of BFM columns is calculated over the entire water column, while the second set (denoted with “ML Only”) is calculated over the monthly mixed layer depth only (defined as the depth at which the density is ~~0.02~~0.2 kg/m³ greater than the surface density).

Variable	BFM17	BFM56	Ayata <i>et al.</i> (2013)	BFM17 (ML Only)	BFM56 (ML Only)	Fasham <i>et al.</i> (1990)	Spitz <i>et al.</i> (2001)
Chlorophyll	0.63 (0.08)	0.60 (0.10)	0.60 (0.06)	0.63 (0.07)	0.60 (0.09)	-0.33 (0.34)	0.86 (0.04)
Oxygen	0.37 (31.18)	0.18 (21.84)	-	0.29 (29.53)	-0.09 (20.22)	-	-
Nitrate	0.94 (0.22)	0.93 (0.16)	0.80 (0.33)	0.94 (0.22)	0.93 (0.15)	0.87 (0.28)	0.98 (0.05)
Phosphate	0.91 (0.01)	0.93 (0.005)	-	0.91 (0.01)	0.93 (0.005)	-	-
PON	0.85 (0.15)	0.85 (0.11)	0.45 (0.08)	0.86 (0.14)	0.86 (0.10)	0.48 (0.6)	0.76 (0.12)
NPP	0.93 (0.26)	0.87 (0.63)	0.50 (0.14)	0.94 (0.21)	0.89 (0.5)	-0.47 (0.021)	0.69 (0.016)

~~the adjustment of some model parameters in BFM17 due to the removal of specific phytoplankton and zooplankton species in favor of general LFGs, and to the parameterization of remineralization using new closure terms that were calibrated to give reasonable agreement with the observational data. By contrast, BFM56 was run using the baseline parameter values that were not adjusted to improve agreement with the observations.~~

~~The correlation~~The correlation coefficients and RMS errors for both BFM17 and BFM56 are ~~also~~ comparable with the Ayata et al. (2013) study for chlorophyll, while out-performing this study for nitrate, PON, and NPP. The Spitz et al. (2001) study, which used data assimilation and is therefore naturally more likely to perform better, does in fact do so for predictions of chlorophyll and nitrate. However, the nitrate correlation values for BFM17 and the Spitz et al. (2001) model are both high, although the latter model does have a lower RMS error value. As compared to the Spitz et al. (2001) model, BFM17 has higher correlation values for both PON and NPP, but a larger RMS error for NPP. Lastly, both BFM17 and BFM56 out-perform the Fasham et al. (1990) study for all fields for both correlation coefficient and RMS error values.

These results show that, with a relatively small increase in the number of biological tracers as compared to similar models, BFM17 is generally able to increase correlation coefficient values and decrease RMS error values for ~~each~~many of the target field in comparison to similar models. Moreover, BFM17 approaches the accuracy of models that use data assimilation to improve agreement with the observations, such as the Spitz et al. (2001) model. The extra biological tracers in BFM17, as compared to the Ayata et al. (2013) and Fasham et al. (1990) models, account for variable intra- and extra-cellular nutrient ratios with the addition of phosphorus

Finally, a key benefit of the chemical functional family approach used by BFM17 is the ability of the model to predict non-Redfield nutrient ratios. Figure 6 shows the constituent component ratios normalized by the respective Redfield ratios for BFM17. The figure includes the component ratios of carbon to nitrogen, carbon to phosphorous, and nitrogen to phosphorous for phytoplankton, DOM, and POM. Zooplankton nutrient ratios were not included because the parameterization of the zooplankton relaxes the nutrient ratio back to a constant value. The normalized ratio values are uniform non-unity valued fields.




Figure 6. Fields of BFM17 constituent component ratios of carbon to nitrogen (top row), carbon to phosphorous (middle row), and nitrogen to phosphorous (bottom row) for phytoplankton (first column), dissolved organic detritus (second column), and particulate organic detritus (third column). Each field is normalized by the respective Redfield ratio.

430 Ultimately, Figure 6 shows that BFM17 is able to capture the phosphate-limited dynamics that characterize the BATS/BTM region (Fanning, 1992; Michaels et al., 1993; Cavender-Bares et al., 2001; Steinberg et al., 2001; Ammerman et al., 2003; Martiny et al., 2013; Singh et al., 2015). In particular, Figure 6 shows that all results comparing carbon or nitrogen to phosphorous for BFM17 produce normalized values greater than 1, where the normalization is carried out using the Redfield ratio (i.e., a normalized value greater than 1 indicates that the field is denominator limited). Figure 6 also shows that the ratios are
435 not uniform for phytoplankton, DOM, and POM, with the ratios decreasing with depth as a result of the increased availability of nitrogen and phosphate.

6 Conclusions

In this study, we have presented a new upper-thermocline, open-ocean BGC model that is complex enough to capture open-ocean ecosystem dynamics within the Sargasso Sea region, yet reduced enough to integrate with a physical model with limited additional computational cost. The new model, named the Biogeochemical Flux Model 17 (BFM17) includes 17 state variables and expands upon more reduced BGC models by incorporating a phosphate equation and by tracking dissolved oxygen, as well as variable intra- and extra-cellular nutrient ratios. BFM17 was developed primarily for use within high-resolution, high-fidelity 3D physical models, such as LES, for process, parameterization, and parameter optimization studies, applications for which its more complex counterpart BFM56 would be much too costly.

To calibrate and test the model, it was coupled to the 1D Princeton Ocean Model (POM-1D) and forced using field data from the Bermuda Atlantic test site area. The full 56 state variable Biogeochemical Flux Model (BFM56) was also run using the same forcing. Results were compared between the two models and all six of the BATS target fields—chlorophyll, oxygen, nitrate, phosphate, PON, and NPP—and a model skill assessment was performed, concluding that the BFM17 captures the subsurface chlorophyll maximum and bloom intensity observed in the BATS data and produces comparable results to BFM56. In comparison with similar studies using slightly less complex models, BFM17-POM1D performs on par with, or better than, those studies.

In the future, a sensitivity study is necessary to assess the most sensitive model parameters, both in BFM17 as well as in the 1D physical model. After identification of these most sensitive model parameters, an optimization can be performed to reduce discrepancies between the BATS observation biology fields and the corresponding model output fields. Additionally, it would be useful to study the efficacy of using BFM17 in a global context, to reproduce the ecology in other regions of the ocean, and its sensitivity under various physical forcing scenarios. Finally, BFM17 is now of a size that it can be efficiently integrated in high-resolution, high-fidelity 3D simulations of the upper ocean, and future work will examine model results in this context.

Appendix A: BFM17 Model Equations

In the following, the detailed equations and their parameter values for each of the 17 state variables that comprise BFM17 are outlined. A summary of the 17 state variables is provided in Table 1 and a schematic of the CFFs and LFGs used in BFM17, along with their interactions, is shown in Figure 1. It should be noted that for all BFM17 equations here, we adopt the same notation style used for BFM56 in Vichi et al. (2007), Mussap et al. (2016), and the BFM User Manual (Vichi et al., 2013) for consistency and clarity.

A1 Environmental parameters

BFM17 ~~interacts with~~ is influenced by the environment through temperature and irradiance ~~inputs~~. Temperature directly affects all physiological processes and is represented in the model by introducing the non-dimensional parameter $f_j^{(T)}$ defined as

$$f_j^{(T)} = Q_{10,j}^{(T-T^*)/T^*}, \quad j = P, Z, \quad (\text{A1})$$

where T^* is a base temperature and $Q_{10,j}$ is a coefficient that may differ for the phytoplankton and zooplankton LFGs, denoted P_i and Z_i , respectively. Here, the subscript i is used to denote different chemical constituents (i.e., C, N, and P) and j is used to denote different LFGs. Base values used for T^* and $Q_{10,j}$ are shown in Table A1. The model additionally employs a temperature-dependent nitrification parameter $f_N^{(T)}$, which is defined similarly to Eq. (A1) as

$$f_N^{(T)} = Q_{10,N}^{(T-T^*)/T^*}, \tag{A2}$$

where $Q_{10,N}$ is given in Table A1

In contrast to temperature, irradiance only directly affects phytoplankton, serving as the primary energy source for phytoplankton growth and maintenance. Irradiance is a function of the incident solar radiation at the sea surface. Within BFM17, the amount of photosynthetically active radiation (PAR) at any given location z is parameterized according to the Lambert–Beer model as

$$E_{\text{PAR}}(z) = \varepsilon_{\text{PAR}} Q_S \exp \left[\lambda_w z + \int_z^0 \lambda_{\text{bio}}(z') dz' \right], \tag{A3}$$

where Q_S is the short-wave surface irradiance flux, which is typically obtained from real-world measurements of the atmospheric radiative transfer, ε_{PAR} is the fraction of PAR within Q_S , λ_w is the background light extinction due to water, and λ_{bio} is the light extinction due to suspended biological particles. Values for ε_{PAR} and λ_w are given in Table A1. The extinction coefficient due to particulate matter, λ_{bio} , is dependent on phytoplankton chlorophyll, P_{chl} , and particulate detritus, $R_C^{(2)}$, and is written as

$$\lambda_{\text{bio}} = c_P P_{\text{chl}} + c_{R^{(2)}} R_C^{(2)}, \tag{A4}$$

where c_P and $c_{R^{(2)}}$ are the specific absorption coefficients of phytoplankton chlorophyll and particulate detritus, respectively, with values given in Table A1.

Table A1. Symbols, values, units, and descriptions for environmental parameters within the BFM17 pelagic model.

Symbol	Value	Units	Description
$Q_{10,P}$	2.00	-	Phytoplankton Q_{10} coefficient
$Q_{10,Z}$	2.00	-	Zooplankton Q_{10} coefficient
$Q_{10,N}$	2.00	-	Nitrification Q_{10} coefficient
T^*	10.0	°C	Base temperature
c_P	0.03	$\text{m}^2 (\text{mg chl})^{-1}$	Chlorophyll-specific light absorption coefficient
ε_{PAR}	0.40	-	Fraction of photosynthetically active radiation
λ_w	0.0435	m^{-1}	Background attenuation coefficient
$c_{R^{(2)}}$	0.1×10^{-3}	$\text{m}^2 (\text{mg C})^{-1}$	C-specific attenuation coefficient of particulate detritus

A2 Phytoplankton equations

The phytoplankton LFG in BFM17 is part of the living organic CFF and is composed of separate state variables for the constituents carbon, nitrogen, phosphorous, and chlorophyll, denoted P_C , P_N , P_P , and P_{chl} respectively (see also Table 1). The governing equations for the constituent state variables are given by:

1. Phytoplankton functional group in the living organic CFF, carbon constituent (state variable P_C):

$$\left. \frac{\partial P_C}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}} - \left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{lys}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(2)}}^{\text{lys}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{exu}} - \left. \frac{\partial P_C}{\partial t} \right|_{Z_C}^{\text{prd}}, \quad (\text{A5})$$

2. Phytoplankton functional group in the living organic CFF, nitrogen constituent (state variable P_N):

$$\left. \frac{\partial P_N}{\partial t} \right|_{\text{bio}} = \max \left[0, \left. \frac{\partial P_N}{\partial t} \right|_{N^{(2)}}^{\text{upt}} + \left. \frac{\partial P_N}{\partial t} \right|_{N^{(3)}}^{\text{upt}} \right] - \left. \frac{\partial P_N}{\partial t} \right|_{R_N^{(1)}}^{\text{lys}} - \left. \frac{\partial P_N}{\partial t} \right|_{R_N^{(2)}}^{\text{lys}} - \left. \frac{\partial P_N}{\partial t} \right|_{Z_N}^{\text{prd}}, \quad (\text{A6})$$

3. Phytoplankton functional group in the living organic CFF, phosphorus constituent (state variable P_P):

$$\left. \frac{\partial P_P}{\partial t} \right|_{\text{bio}} = \max \left[0, \left. \frac{\partial P_P}{\partial t} \right|_{N^{(1)}}^{\text{upt}} \right] - \left. \frac{\partial P_P}{\partial t} \right|_{R_P^{(1)}}^{\text{lys}} - \left. \frac{\partial P_P}{\partial t} \right|_{R_P^{(2)}}^{\text{lys}} - \left. \frac{\partial P_P}{\partial t} \right|_{Z_P}^{\text{prd}}, \quad (\text{A7})$$

4. Phytoplankton functional group in the living organic CFF, chlorophyll constituent (state variable P_{chl}):

$$\left. \frac{\partial P_{chl}}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_{chl}}{\partial t} \right|_{\text{syn}}^{\text{syn}} - \left. \frac{\partial P_{chl}}{\partial t} \right|_{\text{loss}}^{\text{loss}}, \quad (\text{A8})$$

where the descriptions of each of the source and sink terms are provided in Table A2. The subscript “bio” on the left-hand side terms indicates that these are the total rate expressions associated with all biological processes.

For the evolution of the phytoplankton carbon constituent given by Eq. (A5), gross primary production depends on the non-dimensional regulation factors for temperature and light as well as on the maximum photosynthetic growth rate and the phytoplankton carbon instantaneous concentration. This then gives

$$\left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}} = r_P^{(0)} f_P^{(T)} f_P^{(E)} P_C, \quad (\text{A9})$$

where $r_P^{(0)}$ is the maximum photosynthetic rate for phytoplankton (reported in Table A3) and $f_P^{(T)}$ is the temperature regulation factor for phytoplankton given by Eq. (A1). The term $f_P^{(E)}$ is the light regulation factor for phytoplankton, which is defined following (Jassby and Platt, 1976) as

$$f_P^{(E)} = 1 - \exp \left(-\frac{E_{\text{PAR}}}{E_K} \right), \quad (\text{A10})$$

where E_{PAR} is defined in Eq. (A3) and E_K (the “optimal” irradiance) is given by

$$E_K = \left[\frac{r_P^{(0)}}{\alpha_{\text{chl}}^{(0)}} \right] \left(\frac{P_C}{P_{\text{chl}}} \right). \quad (\text{A11})$$

Table A2. List of abbreviations used to indicate physiological and ecological processes in the equations comprising the BFM17 pelagic model.

Abbreviation	Process
gpp	Gross primary production
rsp	Respiration
prd	Predation
rel	Biological release: egestion, excretion, mortality
exu	Exudation
upt	Uptake
lys	Lysis
syn	Biochemical synthesis
loss	Biochemical loss
nit	Nitrification

The parameter $\alpha_{\text{chl}}^{(0)}$ is the maximum light utilization coefficient and is defined in Table A3.

Phytoplankton respiration is parameterized in Eq. (A5) as the sum of the basal respiration and activity respiration rates, namely

$$\left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} = b_P f_P^{(T)} P_C + \gamma_P \left[\left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{exu}} \right], \quad (\text{A12})$$

515 where b_P is the basal specific respiration rate, γ_P is the respired fraction of the gross primary production, the gross primary production term is given by Eq. (A9), and the exudation term is defined below in Eq. (A18). Values and descriptions for b_P and γ_P are given in Table A3.

Both phytoplankton exudation and lysis, defined below, depend on a multiple nutrient limitation term $f_P^{(N,P)}$. This term allows for the internal storage of nutrients and depends on the respective nutrient limitation terms for both nitrate and phosphate. It is
520 given by $f_P^{(N,P)} = \min \left[f_P^{(N)}, f_P^{(P)} \right]$, where

$$f_P^{(N)} = \min \left\{ 1, \max \left[0, \frac{P_N/P_C - \phi_N^{(\min)}}{\phi_N^{(\text{opt})} - \phi_N^{(\min)}} \right] \right\}, \quad (\text{A13})$$

$$f_P^{(P)} = \min \left\{ 1, \max \left[0, \frac{P_P/P_C - \phi_P^{(\min)}}{\phi_P^{(\text{opt})} - \phi_P^{(\min)}} \right] \right\}. \quad (\text{A14})$$

The parameters $\phi_N^{(\text{opt})}$ and $\phi_P^{(\text{opt})}$ are the optimal phytoplankton quotas for nitrogen and phosphorus, respectively, while $\phi_N^{(\min)}$ and $\phi_P^{(\min)}$ are the minimum possible quotas, below which $f_P^{(N)}$ and $f_P^{(P)}$ are zero. Values for each of these parameters are
525 included in Table A3.

Phytoplankton lysis includes all mortality due to mechanical, viral, and yeast cell disruption processes, and is partitioned between particulate and dissolved detritus. The internal cytoplasm of the cell is released to dissolved detritus, denoted by $R_i^{(1)}$,

Table A3. Phytoplankton parameters, values, units, and descriptions within the BFM17 pelagic model.

Symbol	Value	Units	Description
$r_P^{(0)}$	1.60	d^{-1}	Maximum specific photosynthetic rate
b_P	0.05	d^{-1}	Basal specific respiration rate
$d_P^{(0)}$	0.05	d^{-1}	Maximum specific nutrient-stress lysis rate
$h_P^{(N,P)}$	0.10	-	Nutrient stress threshold
β_P	0.05	-	Excreted fraction of primary production
γ_P	0.05	-	Activity respiration fraction
$a_P^{(N)}$	0.025	$\text{m}^3 (\text{mg C})^{-1} \text{d}^{-1}$	Specific affinity constant for nitrogen
$h_P^{(N)}$	1.50	$\text{mmol N-NH}_4 \text{m}^{-3}$	Half saturation constant for ammonium uptake
$\phi_N^{(\min)}$	6.87×10^{-3}	$\text{mmolN} (\text{mg C})^{-1}$	Minimum nitrogen quota
$\phi_N^{(\text{opt})}$	1.26×10^{-2}	$\text{mmolN} (\text{mg C})^{-1}$	Optimal nitrogen quota
$\phi_N^{(\max)}$	$1.0\phi_N^{(\text{opt})}$	$\text{mmolN} (\text{mg C})^{-1}$	Maximum nitrogen quota
$a_P^{(P)}$	2.5×10^{-3}	$\text{m}^3 (\text{mg C})^{-1} \text{d}^{-1}$	Specific affinity constant for phosphorus
$\phi_P^{(\min)}$	4.29×10^{-4}	$\text{mmolP} (\text{mg C})^{-1}$	Minimum phosphorus quota
$\phi_P^{(\text{opt})}$	7.86×10^{-4}	$\text{mmolP} (\text{mg C})^{-1}$	Optimal phosphorus quota
$\phi_P^{(\max)}$	$1.0\phi_P^{(\text{opt})}$	$\text{mmolP} (\text{mg C})^{-1}$	Maximum phosphorus quota
$\alpha_{\text{chl}}^{(0)}$	1.52×10^{-5}	$\text{mgC} (\text{mg chl})^{-1} (\mu\text{E})^{-1} \text{m}^2$	Maximum light utilization coefficient
$\theta_{\text{chl}}^{(0)}$	0.016	$\text{mg chl} (\text{mg C})^{-1}$	Maximum chlorophyll to carbon quota

while structural parts of the cell are released to particulate detritus, denoted by the state variable $R_i^{(2)}$, where $i = \text{C, N, P}$ (see also Table 1). The resulting lysis terms in Eqs. (A5)–(A7) are then given by

$$530 \quad \left. \frac{\partial P_i}{\partial t} \right|_{R_i^{(1)}}^{\text{lys}} = \left[1 - \varepsilon_P^{(N,P)} \right] \left[\frac{h_P^{(N,P)}}{f_P^{(N,P)} + h_P^{(N,P)}} d_P^{(0)} P_i \right], \quad i = \text{C, N, P}, \quad (\text{A15})$$

$$\left. \frac{\partial P_i}{\partial t} \right|_{R_i^{(2)}}^{\text{lys}} = \varepsilon_P^{(N,P)} \left[\frac{h_P^{(N,P)}}{f_P^{(N,P)} + h_P^{(N,P)}} d_P^{(0)} P_i \right], \quad i = \text{C, N, P}, \quad (\text{A16})$$

where $h_P^{(N,P)}$ is the nutrient stress threshold and $d_P^{(0)}$ is the maximum specific nutrient-stress lysis rate, both of which are given in Table A3. The term $\varepsilon_P^{(N,P)}$ is a fraction that ensures nutrients within the structural parts of the cell, which are less degradable, are always released as particulate detritus. This fraction is determined by the expression

$$535 \quad \varepsilon_P^{(N,P)} = \min \left[1, \frac{\phi_N^{(\min)}}{P_N/P_C}, \frac{\phi_P^{(\min)}}{P_P/P_C} \right], \quad (\text{A17})$$

where $\phi_N^{(\min)}$ and $\phi_P^{(\min)}$ are given in Table A3.

If phytoplankton cannot equilibrate their fixed carbon with sufficient nutrients, this carbon is not assimilated and is instead released in the form of dissolved carbon, denoted by state variable $R_C^{(1)}$, in a process known as exudation. The exudation term

in Eq. (A5) is parameterized as

$$540 \quad \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{exu}} = \left\{ \beta_P + (1 - \beta_P) \left[1 - f_P^{(N,P)} \right] \right\} \left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}}, \quad (\text{A18})$$

where β_P is the excreted fraction of gross primary production, defined in Table A3, and the gross primary production term is again given by Eq. (A9).

The nutrient uptake of Eqs. (A6) and (A7) combines both the intracellular quota (i.e., Droop) and external concentration (i.e., Monod) approaches Baretta-Bekker et al. (1997). The total phytoplankton uptake of nitrogen, represented by the combination
545 of the two uptake terms in Eq. (A6), is the minimum of a diffusion-dependent uptake rate when internal nutrient quotas are low and a rate that is based upon balanced growth needs and any excess uptake, namely

$$\left. \frac{\partial P_N}{\partial t} \right|_{N^{(2,3)}}^{\text{upt}} = \min \left\{ a_P^{(N)} \left[\frac{h_P^{(N)}}{h_P^{(N)} + N^{(3)}} N^{(2)} + N^{(3)} \right] P_C, \phi_N^{(\max)} G_P + \nu_P \left[\phi_N^{(\max)} - \frac{P_N}{P_C} \right] P_C \right\}, \quad (\text{A19})$$

where $a_P^{(N)}$ is the specific affinity for nitrogen, $h_P^{(N)}$ is the half saturation constant for ammonium uptake, and $\phi_N^{(\max)}$ is the maximum nitrogen quota; base values for these three parameters are given in Table A3. The net primary productivity G_P in
550 Eq. (A19) is given as

$$G_P = \max \left[0, \left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{exu}} - \left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{lys}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(2)}}^{\text{lys}} \right]. \quad (\text{A20})$$

The specific uptake rate ν_P appearing in Eq. (A19) is given by

$$\nu_P = f_P^{(T)} r_P^{(0)}. \quad (\text{A21})$$

It should be noted that only the sum of the two uptake terms, represented by Eq. (A19), is required in the governing equation
555 for P_N given by Eq. (A6). However, in the governing equations for nitrate and ammonium, denoted $N^{(2)}$ and $N^{(3)}$ (see Table 1) that will be presented later, expressions are required for the individual uptake portions from nitrate and ammonium. When the total phytoplankton nitrogen uptake rate from Eq. (A19) is positive, the individual portions from nitrate and ammonium are determined by

$$\left. \frac{\partial P_N}{\partial t} \right|_{N^{(2)}}^{\text{upt}} = \varepsilon_P \left. \frac{\partial P_N}{\partial t} \right|_{N^{(2,3)}}^{\text{upt}}, \quad (\text{A22})$$

$$560 \quad \left. \frac{\partial P_N}{\partial t} \right|_{N^{(3)}}^{\text{upt}} = (1 - \varepsilon_P) \left. \frac{\partial P_N}{\partial t} \right|_{N^{(2,3)}}^{\text{upt}}, \quad (\text{A23})$$

where the rates on the right-hand sides are obtained from Eq. (A19), and ε_P is given as

$$\varepsilon_P = \frac{s_N N^{(2)}}{N^{(3)} + s_N N^{(2)}}. \quad (\text{A24})$$

The preference for ammonium is defined by the saturation function s_N and is given by

$$s_N = \frac{h_P^{(N)}}{h_P^{(N)} + N^{(3)}}. \quad (\text{A25})$$

565 When the phytoplankton nitrogen uptake rate from Eq. (A19) is negative, however, the entire nitrogen uptake goes to the dissolved organic nitrogen pool, $R_N^{(1)}$ [see Eq. (A42)].

As with the uptake of nitrogen, phytoplankton uptake of phosphorus in Eq. (A7) is the minimum of a diffusion-dependent rate and a balanced growth/excess uptake rate. This uptake comes entirely from one pool and the uptake term in Eq. (A7) is correspondingly given by

$$570 \quad \left. \frac{\partial P_P}{\partial t} \right|_{N^{(1)}}^{\text{upt}} = \min \left\{ a_P^{(P)} N^{(1)} P_C, \phi_P^{(\max)} G_P + \nu_P \left[\phi_P^{(\max)} P_C - P_P \right] \right\}, \quad (\text{A26})$$

where $a_P^{(P)}$ is the specific affinity constant for phosphorous and $\phi_P^{(\max)}$ is the maximum phosphorous quota. Values for both parameters are given in Table A3. If the uptake rate is negative, the entire phosphorus uptake goes to the dissolved organic phosphorus pool, $R_P^{(1)}$.

Predation of phytoplankton within BFM17 is solely performed by zooplankton, and each of the predation terms appearing
575 in Eqs. (A5)–(A7) are equal and opposite to the zooplankton predation terms, namely

$$\left. \frac{\partial P_i}{\partial t} \right|_{Z_i}^{\text{prd}} = - \left. \frac{\partial Z_i}{\partial t} \right|_{P_i}^{\text{prd}}, \quad i = \text{C, N, P}. \quad (\text{A27})$$

Equations for the zooplankton predation terms are given in the next section.

Finally, phytoplankton chlorophyll, denoted P_{chl} with the rate equation given by Eq. (A8), contributes to the definition of the optimal irradiance value in Eq. (A11) and of the phytoplankton contribution to the extinction coefficient in Eq. (A4).

580 The phytoplankton chlorophyll source term in Eq. (A8) is made up of only two terms: chlorophyll synthesis and loss. Net chlorophyll synthesis is a function of acclimation to light n conditions, availability of nutrients, and turnover rate, and is given by

$$\left. \frac{\partial P_{\text{chl}}}{\partial t} \right|^{\text{syn}} = \rho_{\text{chl}} (1 - \gamma_P) \left[\left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{exu}} \right] - \frac{P_{\text{chl}}}{P_C} \left[\left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{lys}} + \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(2)}}^{\text{lys}} + \left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} \right], \quad (\text{A28})$$

where ρ_{chl} regulates the amount of chlorophyll in the phytoplankton cell and all other terms in the above expression have been
585 defined previously. The term ρ_{chl} is computed according to a ratio between the realized photosynthetic rate (i.e., gross primary production) and the maximum potential photosynthesis Geider et al. (1997), and is correspondingly given as

$$\rho_{\text{chl}} = \theta_{\text{chl}}^{(0)} \min \left\{ 1, \frac{(1 - \gamma_P)}{\alpha_{\text{chl}}^{(0)} E_{\text{PAR}} P_{\text{chl}}} \left[\left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}} - \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{exu}} \right] \right\}, \quad (\text{A29})$$

where $\theta_{\text{chl}}^{(0)}$ is the maximum chlorophyll to carbon quota and $\alpha_{\text{chl}}^{(0)}$ is the maximum light utilization coefficient, both of which can be found in Table A3. Chlorophyll loss in Eq. (A8) is simpler and is just a function of predation, where the amount of
590 chlorophyll transferred back to the infinite sink is proportional to the carbon predated by zooplankton, giving

$$\left. \frac{\partial P_{\text{chl}}}{\partial t} \right|^{\text{loss}} = \frac{P_{\text{chl}}}{P_C} \left. \frac{\partial P_C}{\partial t} \right|_{Z_C}^{\text{prd}}. \quad (\text{A30})$$

A3 Zooplankton equations

The zooplankton LFG group in BFM17 is part of the living organic CFF and is composed of separate state variables for carbon, nitrogen, and phosphorous, denoted Z_C , Z_N , and Z_P , respectively (see also Table 1). The governing equations for the constituent state variables are given by:

5. Zooplankton functional group in the living organic CFF, carbon constituent (state variable Z_C):

$$\left. \frac{\partial Z_C}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial Z_C}{\partial t} \right|_{P_C}^{\text{prd}} - \left. \frac{\partial Z_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} - \left. \frac{\partial Z_C}{\partial t} \right|_{R_C^{(1)}}^{\text{rel}} - \left. \frac{\partial Z_C}{\partial t} \right|_{R_C^{(2)}}^{\text{rel}}, \quad (\text{A31})$$

6. Zooplankton functional group in the living organic CFF, nitrogen constituent (state variable Z_N):

$$\left. \frac{\partial Z_N}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial Z_N}{\partial t} \right|_{P_N}^{\text{prd}} - \left. \frac{\partial Z_N}{\partial t} \right|_{R_N^{(1)}}^{\text{rel}} - \left. \frac{\partial Z_N}{\partial t} \right|_{R_N^{(2)}}^{\text{rel}} - \left. \frac{\partial Z_N}{\partial t} \right|_{N^{(3)}}^{\text{rel}}, \quad (\text{A32})$$

7. Zooplankton functional group in the living organic CFF, phosphorus constituent (state variable Z_P):

$$\left. \frac{\partial Z_P}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial Z_P}{\partial t} \right|_{P_P}^{\text{prd}} - \left. \frac{\partial Z_P}{\partial t} \right|_{R_P^{(1)}}^{\text{rel}} - \left. \frac{\partial Z_P}{\partial t} \right|_{R_P^{(2)}}^{\text{rel}} - \left. \frac{\partial Z_P}{\partial t} \right|_{N^{(1)}}^{\text{rel}}, \quad (\text{A33})$$

where, once more, descriptions of each of the source and sink terms are provided in Table A2.

Zooplankton predation of phytoplankton, which appears as the first term in each of Eqs. (A31)–(A33), primarily depends on the availability of phytoplankton and their capture efficiency, and is expressed as

$$\left. \frac{\partial Z_i}{\partial t} \right|_{P_i}^{\text{prd}} = \frac{P_i}{P_C} \left[f_Z^{(T)} r_Z^{(0)} \delta_{Z,P} \frac{P_C}{P_C + h_Z^{(F)}} Z_C \right], \quad i = \text{C, N, P}, \quad (\text{A34})$$

where $r_Z^{(0)}$ is the potential specific growth rate and $h_Z^{(F)}$ is the Michaelis constant for total food ingestion. These parameters and their base values are included in Table A4. Here, $f_Z^{(T)}$ is the temperature regulating factor for zooplankton growth given by Eq. (A1). The total food availability can be expressed as $\delta_{Z,P} P_C$, where $\delta_{Z,P}$ is the prey availability of phytoplankton and is included in Table A4.

Zooplankton respiration is the sum of active and basal metabolism rates, where active respiration is the cost of nutrient ingestion, or predation. The resulting respiration rate is given by

$$\left. \frac{\partial Z_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} = (1 - \eta_Z - \beta_Z) \left. \frac{\partial Z_C}{\partial t} \right|_{P_C}^{\text{prd}} + b_Z f_Z^{(T)} Z_C, \quad (\text{A35})$$

where η_Z is the assimilation efficiency, β_Z is the excreted fraction uptake, and b_Z is the basal specific respiration rate. All three parameters are included in Table A4.

The biological release terms in Eqs. (A31)–(A33) are the sum of zooplankton excretion, egestion, and mortality. Excretion and egestion are the portions of ingested nutrients, resulting from predation, that have not been assimilated or used for respiration. Zooplankton mortality is parameterized as the sum of a constant mortality rate and an oxygen-dependent regulation factor

Table A4. Zooplankton parameters, values, units, and descriptions within the BFM17 pelagic model.

Symbol	Value	Unit	Description
b_Z	0.02	d^{-1}	Basal specific respiration rate
$r_Z^{(0)}$	2.00	d^{-1}	Potential specific growth rate
$d_Z^{(0)}$	0.25	d^{-1}	Oxygen dependent specific mortality rate
d_Z	0.05	d^{-1}	Specific mortality rate
η_Z	0.50	-	Assimilation efficiency
β_Z	0.25	-	Fraction of activity excretion
ε_Z^C	0.60	-	Partition between dissolved and particulate excretion of C
ε_Z^N	0.72	-	Partition between dissolved and particulate excretion of N
ε_Z^P	0.832	-	Partition between dissolved and particulate excretion of P
$h_Z^{(F)}$	200.0	mg C m^{-3}	Michaelis constant for total food ingestion
$\delta_{Z,P}$	1.00	-	Availability of phytoplankton to zooplankton
$\nu_Z^{(P)}$	1.0	d^{-1}	Specific rate constant for phosphorous excretion
$\nu_Z^{(N)}$	1.0	d^{-1}	Specific rate constant for nitrogen excretion
$\varphi_P^{(\text{opt})}$	7.86×10^{-4}	mmolP (mg C)^{-1}	Optimal phosphorous quota
$\varphi_N^{(\text{opt})}$	0.0126	mmolN (mg C)^{-1}	Optimal nitrogen quota

given by

$$f_Z^{(0)} = \frac{O}{O + h_O}, \quad (\text{A36})$$

620 where O represents the oxygen constituent of the dissolved gas in the inorganic CFF and h_O is the half saturation coefficient for chemical processes given in Table A5. The total biological release is then partitioned into particulate and dissolved organic matter, giving

$$\frac{\partial Z_i}{\partial t} \Big|_{R_i^{(1)}}^{\text{rel}} = \varepsilon_Z^{(i)} \left\{ \beta_Z \frac{\partial Z_i}{\partial t} \Big|_{P_i}^{\text{prd}} + d_Z + d_Z^{(0)} \left[1 - f_Z^{(0)} \right] f_Z^{(T)} Z_i \right\}, \quad i = \text{C, N, P}, \quad (\text{A37})$$

$$\frac{\partial Z_i}{\partial t} \Big|_{R_i^{(2)}}^{\text{rel}} = \left[1 - \varepsilon_Z^{(i)} \right] \frac{\partial Z_i}{\partial t} \Big|_{R_i^{(1)}}^{\text{rel}}, \quad i = \text{C, N, P}, \quad (\text{A38})$$

625 where $\varepsilon_Z^{(i)}$ is the fraction excreted to the dissolved pool, d_Z is the specific mortality rate, and $d_Z^{(0)}$ is the oxygen dependent specific mortality rate. Base values for each parameter are given in Table A4.

Table A5. Values, units, and descriptions for dissolved organic matter, particulate organic matter, and nutrient parameters within the BFM17 pelagic model.

Symbol	Value	Units	Description
$\alpha_{R^{(1)}}^{(\text{sink}_C)}$	0.05	d^{-1}	Specific remineralization rate of dissolved carbon
$\zeta_{N^{(1)}}$	0.05	d^{-1}	Specific remineralization rate of dissolved phosphorus
$\zeta_{N^{(3)}}$	0.05	d^{-1}	Specific remineralization rate of dissolved nitrogen
$\alpha_{R^{(2)}}^{(\text{sink}_C)}$	0.1	d^{-1}	Specific remineralization rate of particulate carbon
$\xi_{N^{(1)}}$	0.1	d^{-1}	Specific remineralization rate of particulate phosphorus
$\xi_{N^{(3)}}$	0.1	d^{-1}	Specific remineralization rate of particulate nitrogen
$\Lambda_{N^{(3)}}^{(\text{nit})}$	0.01	d^{-1}	Specific nitrification rate at 10 °C
h_O	10.0	$\text{mmolO}_2 \text{ m}^{-3}$	Half saturation for chemical processes
$\Omega_C^{(O)}$	12.0	$\text{mmolO}_2 \text{ mgC}^{-1}$	Stoichiometric coefficient for oxygen reaction
$\Omega_N^{(O)}$	2.0	$\text{mmolO}_2 \text{ mmolN}^{-1}$	Stoichiometric coefficient for nitrification reaction

The zooplankton also excrete into the nutrient pools of phosphate, $N^{(1)}$, and ammonium, $N^{(3)}$. These effects are represented by the final terms of Eqs. (A32) and (A33), which are parameterized by

$$\left. \frac{\partial Z_N}{\partial t} \right|_{N^{(3)}}^{\text{rel}} = \nu_Z^{(N)} \max \left[0, \frac{Z_N}{Z_C} - \varphi_N^{(\text{opt})} \right] Z_N, \quad (\text{A39})$$

$$630 \quad \left. \frac{\partial Z_P}{\partial t} \right|_{N^{(1)}}^{\text{rel}} = \nu_Z^{(P)} \max \left[0, \frac{Z_P}{Z_C} - \varphi_P^{(\text{opt})} \right] Z_P, \quad (\text{A40})$$

where $\nu_Z^{(N)}$ and $\nu_Z^{(P)}$ are specific rate constants and $\varphi_N^{(\text{opt})}$ and $\varphi_P^{(\text{opt})}$ are the optimal zooplankton quotas for nitrogen and phosphorous, respectively. All four parameters are included in Table A4.

A4 Dissolved organic matter equations

The governing equations for the three constituents of dissolved organic matter are given by:

635 8. Dissolved matter in non-living organic CFF, carbon constituent [state variable $R_C^{(1)}$]:

$$\left. \frac{\partial R_C^{(1)}}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{lys}} + \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(1)}}^{\text{exu}} + \left. \frac{\partial Z_C}{\partial t} \right|_{R_C^{(1)}}^{\text{rel}} - \alpha_{R^{(1)}}^{(\text{sink}_C)} R_C^{(1)}, \quad (\text{A41})$$

9. Dissolved matter in non-living organic CFF, nitrogen constituent [state variable $R_N^{(1)}$]:

$$\left. \frac{\partial R_N^{(1)}}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_N}{\partial t} \right|_{R_N^{(1)}}^{\text{lys}} + \left. \frac{\partial Z_N}{\partial t} \right|_{R_N^{(1)}}^{\text{rel}} - \min \left[0, \left. \frac{\partial P_N}{\partial t} \right|_{N^{(2)}}^{\text{upt}} + \left. \frac{\partial P_N}{\partial t} \right|_{N^{(3)}}^{\text{upt}} \right] - \zeta_{N^{(3)}} R_N^{(1)}, \quad (\text{A42})$$

10. Dissolved matter in non-living organic CFF, phosphorus constituent [state variable $R_p^{(1)}$]:

$$\left. \frac{\partial R_p^{(1)}}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_p}{\partial t} \right|_{R_p^{(1)}}^{\text{lys}} + \left. \frac{\partial Z_p}{\partial t} \right|_{R_p^{(1)}}^{\text{rel}} - \min \left[0, \left. \frac{\partial P_p}{\partial t} \right|_{N^{(1)}}^{\text{upt}} \right] - \zeta_{N^{(1)}} R_p^{(1)}. \quad (\text{A43})$$

All terms except for the last terms in each of these equations, representing remineralization, have been defined in previous sections. Remineralization of dissolved organic matter by bacteria is parameterized within BFM17 as a rate that is proportional to the local concentration of that dissolved constituent. In Eq. (A41), remineralization is parameterized as $\alpha_{R^{(1)}}^{(\text{sink}_C)} R_C^{(1)}$, where $\alpha_{R^{(1)}}^{(\text{sink}_C)}$ is a constant that controls the rate at which dissolved carbon is remineralized and returned to the pool of carbon; this constant is given in Table A5. In Eqs. (A42) and (A43), remineralization is represented by the parameters $\zeta_{N^{(3)}}$ and $\zeta_{N^{(1)}}$, which are the specific remineralization rates of dissolved ammonium and phosphate, respectively. These rates are also included in Table A5

A5 Particulate organic matter equations

The governing equations for the three constituents of particulate organic matter are given by:

11. Particulate matter in non-living organic CFF, carbon constituent [state variable $R_C^{(2)}$]:

$$\left. \frac{\partial R_C^{(2)}}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_C}{\partial t} \right|_{R_C^{(2)}}^{\text{lys}} + \left. \frac{\partial Z_C}{\partial t} \right|_{R_C^{(2)}}^{\text{rel}} - \alpha_{R^{(2)}}^{(\text{sink}_C)} R_C^{(2)}, \quad (\text{A44})$$

12. Particulate matter in non-living organic CFF, nitrogen constituent [state variable $R_N^{(2)}$]:

$$\left. \frac{\partial R_N^{(2)}}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_N}{\partial t} \right|_{R_N^{(2)}}^{\text{lys}} + \left. \frac{\partial Z_N}{\partial t} \right|_{R_N^{(2)}}^{\text{rel}} - \xi_{N^{(3)}} R_N^{(2)}, \quad (\text{A45})$$

13. Particulate matter in non-living organic CFF, phosphorus constituent [state variable $R_p^{(2)}$]:

$$\left. \frac{\partial R_p^{(2)}}{\partial t} \right|_{\text{bio}} = \left. \frac{\partial P_p}{\partial t} \right|_{R_p^{(2)}}^{\text{lys}} + \left. \frac{\partial Z_p}{\partial t} \right|_{R_p^{(2)}}^{\text{rel}} - \xi_{N^{(1)}} R_p^{(2)}. \quad (\text{A46})$$

Once again, all terms except for the final remineralization terms in each equation have been defined in previous sections. Remineralization of particulate organic matter by bacteria is parameterized within BFM17 as a rate that is proportional to the local concentration of that particulate constituent. In Eq. (A44), remineralization is parameterized by $\alpha_{R^{(2)}}^{(\text{sink}_C)} R_C^{(2)}$, where $\alpha_{R^{(2)}}^{(\text{sink}_C)}$ is a constant that controls the rate at which the particulate carbon is remineralized. The base value for this constant is provided in Table A5. The parameters $\xi_{N^{(3)}}$ and $\xi_{N^{(1)}}$ are the specific remineralization rates of particulate ammonium and phosphate, respectively. The specific remineralization rates for particulate organic matter are also presented in Table A5.

A6 Dissolved gas and nutrient equations

The only dissolved gas resolved by BFM17 is oxygen, O , (carbon dioxide is treated as an infinite source/sink) and the dissolved nutrients in the model are phosphate, $N^{(1)}$, nitrate, $N^{(2)}$, and ammonium, $N^{(3)}$ (see also Table 1). Governing equations for each of these state variables are given by:

14. Dissolved gas in the inorganic CFF, oxygen constituent (state variable O):

$$\begin{aligned} \left. \frac{\partial O}{\partial t} \right|_{\text{bio}} = & \left. \frac{\partial O}{\partial t} \right|_{\text{wind}} + \Omega_C^{(O)} \left[\left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{gpp}} - \left. \frac{\partial P_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} - \left. \frac{\partial Z_C}{\partial t} \right|_{\text{CO}_2}^{\text{rsp}} - \alpha_{R^{(2)}_{\text{C}}}^{(\text{sink}_{\text{C}})} R_{\text{C}}^{(2)} - \alpha_{R^{(1)}_{\text{C}}}^{(\text{sink}_{\text{C}})} R_{\text{C}}^{(1)} \right] \\ & - \Omega_N^{(O)} \left. \frac{\partial N^{(3)}}{\partial t} \right|_{N^{(2)}}^{\text{nit}}, \end{aligned} \quad (\text{A47})$$

15. Dissolved nutrient in the inorganic CFF, phosphate constituent (state variable $N^{(1)}$):

$$\left. \frac{\partial N^{(1)}}{\partial t} \right|_{\text{bio}} = - \left. \frac{\partial P_P}{\partial t} \right|_{N^{(1)}}^{\text{upt}} + \zeta_{N^{(1)}} R_P^{(1)} + \xi_{N^{(1)}} R_P^{(2)} + \left. \frac{\partial Z_P}{\partial t} \right|_{N^{(1)}}^{\text{rel}}, \quad (\text{A48})$$

16. Dissolved nutrient in the inorganic CFF, nitrate constituent (state variable $N^{(2)}$):

$$\left. \frac{\partial N^{(2)}}{\partial t} \right|_{\text{bio}} = - \left. \frac{\partial P_N}{\partial t} \right|_{N^{(2)}}^{\text{upt}} + \left. \frac{\partial N^{(2)}}{\partial t} \right|_{N^{(3)}}^{\text{nit}}, \quad (\text{A49})$$

17. Dissolved nutrient in the inorganic CFF, ammonium constituent (state variable $N^{(3)}$):

$$\left. \frac{\partial N^{(3)}}{\partial t} \right|_{\text{bio}} = - \left. \frac{\partial P_N}{\partial t} \right|_{N^{(3)}}^{\text{upt}} + \zeta_{N^{(3)}} R_N^{(1)} + \xi_{N^{(3)}} R_N^{(2)} + \left. \frac{\partial Z_N}{\partial t} \right|_{N^{(3)}}^{\text{rel}} - \left. \frac{\partial N^{(3)}}{\partial t} \right|_{N^{(2)}}^{\text{nit}}. \quad (\text{A50})$$

675 Aeration of the surface layer by wind, $\partial O / \partial t|_{\text{wind}}$, is parameterized as described in Wanninkhof (1992, 2014). In a 0D model it is a source term for dissolved oxygen and so belongs in Eq. (A47). However, in any model of one dimension or more it should be treated as a surface boundary condition for dissolved oxygen and so belongs in Eq. (17) and should be omitted from Eq. (A47). The parameters $\Omega_C^{(O)}$ and $\Omega_N^{(O)}$ are stoichiometric coefficients used to convert units of carbon to units of oxygen and nitrogen, respectively. All terms in the above equations have been defined in previous sections, except for nitrification.

680 Nitrification is a source term for nitrate and is parameterized as a sink of ammonium and oxygen as

$$\left. \frac{\partial N^{(2)}}{\partial t} \right|_{N^{(3)}}^{\text{nit}} = \left. \frac{\partial N^{(3)}}{\partial t} \right|_{N^{(2)}}^{\text{nit}} = \Lambda_{N^{(3)}}^{(\text{nit})} f_N^{(T)} f_Z^{(O)} N^{(3)}, \quad (\text{A51})$$

where $\Lambda_{N^{(3)}}^{(\text{nit})}$ is the specific nitrification rate, given in Table A5. The terms $f_N^{(T)}$ and $f_Z^{(O)}$ are defined in Eqs. (A2) and (A36), respectively.

Appendix B: Zero-Dimensional Test of the BFM17

685 As a simple test of BFM17 without the influence of any particular physical model, the BGC model (as discussed previously, the choice of physical model can greatly affect the results) BFM17 was integrated in a 0D (i.e., time only) test for 10 years using sinusoidal forcing for the temperature (in units of °C), salinity (psu), 10 m wind-speed (m s⁻¹), and PAR (W m⁻²) cycles. This forcing is implemented as

$$F^{(\text{var})}(t) = \left[F_s^{(\text{var})} + F_w^{(\text{var})} \right] - 0.5 \left[F_s^{(\text{var})} - F_w^{(\text{var})} \right] \cos(tR), \quad (\text{B1})$$




Figure B1. Seasonal cycle of surface (a) chlorophyll, (b) zooplankton carbon, and (c) nitrate from the 0D test of BFM17. Results are shown for the last 4 years of the 10-year simulation. Panels (a) and (c) show monthly averaged values taken from the observational data described in Section 4.

690 where $F^{(\text{var})}$ is the annually varying forcing term, ‘var’ indicates the variable of interest, corresponding to temperature (‘temp’), salinity (‘sal’), wind speed (‘wind’), and PAR. In Eq. (B1), $F_w^{(\text{var})}$ and $F_s^{(\text{var})}$ are, respectively, the winter and summer extreme values for the forcing term considered, $0 \leq t \leq 360$ is the time, and $R = \pi/180$. The winter and summer values were chosen to be similar to those found in the observational data described in Section 4, with $[F_w^{(\text{temp})}, F_s^{(\text{temp})}] = [10^\circ\text{C}, 30^\circ\text{C}]$, $[F_w^{(\text{sal})}, F_s^{(\text{sal})}] = [37 \text{ psu}, 36.5 \text{ psu}]$, $[F_w^{(\text{wind})}, F_s^{(\text{wind})}] = [6 \text{ m s}^{-1}, 2 \text{ m s}^{-1}]$, and $[F_w^{(\text{PAR})}, F_s^{(\text{PAR})}] = [10 \text{ W m}^{-2}, 120 \text{ W m}^{-2}]$.

695 The exact observational data was not used, and instead we used an idealized version of the data for simplicity and because there is no physical variable in the 0D framework to properly apply the exact observations. Note that, in the 0D framework, the wind forcing does not constrain the biogeochemical dynamics, but does play a role in oxygen exchange with the atmosphere, defined according to Wanninkhof (1992, 2014).

Initial values for chlorophyll, oxygen, phosphate, and nitrate were taken to be similar to values from the BATS/BTM observational data, with $P_{\text{chl}} = 0.2 \text{ mg Chl-}a \text{ m}^{-3}$, $O = 230 \text{ mmol O}_2 \text{ m}^{-3}$, $N^{(1)} = 0.06 \text{ mmol P m}^{-3}$, and $N^{(2)} = 1.0 \text{ mmol N m}^{-3}$. Phytoplankton carbon was calculated using the maximum chlorophyll to carbon ratio, $\theta_{\text{chl}}^{(0)}$ in Table A3. Initial values for zooplankton carbon, dissolved carbon, and particulate organic carbon were assumed to be the same as the phytoplankton carbon. Ammonium was assumed to have the same initial concentration as phosphate. All other constituents were calculated using their respective optimal ratios in Tables A3 and A4.

705 Figure B1 shows the seasonal cycle of surface chlorophyll, zooplankton carbon, and nitrate over the last 4 years of the 10-year simulation period, indicating that a ~~self-consistent and~~ stable seasonal cycle with reasonable ecosystem values can be attained by the model, regardless of its coupling to a physical model. Figures B1(a) and (c) also show monthly averaged values taken from the observational data described in Section 4. Although the agreement between the 0D BFM17 model and the observations is not perfect, both are qualitatively similar and close in magnitude, providing confidence in the accuracy of

710 the model despite the lower fidelity of the 0D test.

Code and data availability. Current versions of BFM17 and BFM56 are at <https://github.com/marco-zavatarelli/BFM17-56/tree/BFM17-56> under the GNU General Public License version 3. The exact versions of the models used to produce the results used in this paper are archived on Zenodo at <http://doi.org/10.5281/zenodo.3839984>. Data and scripts used to produce all figures in this paper are archived on Zenodo at <http://doi.org/10.5281/zenodo.3840562>.

715 *Author contributions.* KMS, SK, PEH, MZ, and NP formulated the reduced BFM order model, EFK and KEN verified and debugged the model, KMS and SK ran the model simulations and produced the results presented in the paper, KMS and PEH prepared the initial draft of the paper, and all authors edited the paper to produce the final version.

Competing interests. The authors declare that they have no competing interests or other conflicts of interest.

720 *Acknowledgements.* KMS and PEH were supported by NSF OCE-1258995 and PEH was supported, in part, by NSF OCE-1924636. SK was supported by an ANSEP Alaska Grown Fellowship and by an NSF Graduate Research Fellowship. EFK and KEN were supported by NSF OAC-1535065. The data analyzed in this paper are available from Mendeley Data (<https://data.mendeley.com>). This work utilized the RMACC Summit supercomputer supported by NSF (ACI-1532235, ACI-1532236), CU-Boulder, and CSU, as well as the Yellowstone (ark:/85065/d7wd3xhc) and Cheyenne (doi:10.5065/D6RX99HX) supercomputers provided by NCAR CISL, sponsored by NSF.

References

- 725 Abraham, E. R.: The generation of plankton patchiness by turbulent stirring, *Nature*, 391, 577–580, 1998.
- Ammerman, J. W., Hood, R. R., Case, D. A., and Cotner, J. B.: Phosphorus Deficiency in the Atlantic: An Emerging Paradigm in Oceanography, *EOS*, 84, 165–170, 2003.
- Anderson, T. R.: Plankton functional type modelling: running before we can walk?, *Journal of Plankton Research*, 27, 1073–1081, 2005.
- Ayata, S. D., Levy, M., Aumont, O., Siandra, A., Sainte-Marie, J., Tagliabue, A., and Bernard, O.: Phytoplankton growth formulation in
730 marine ecosystem models: Should we take into account photo-acclimation and variable stoichiometry in oligotrophic areas?, *Journal of Marine Systems*, 125, 29–40, 2013.
- Baretta-Bekker, J. G., Baretta, J. W., and Ebenhoh, W.: Microbial dynamics in the marine ecosystem model ERSEM II with decoupled carbon assimilation and nutrient uptake, *Journal of Sea Research*, 38, 195–211, 1997.
- Barton, A. D., Lozier, M. S., and Williams, R. G.: Physical controls of variability in North Atlantic phytoplankton communities, *Limnology and Oceanography*, 60, 181–197, 2015.
- 735 Bees, M. A.: Plankton Communities and Chaotic Advection in Dynamical Models of Langmuir Circulation, *Applied Scientific Research*, 59, 141–158, 1998.
- Behrenfeld, M. J.: Climate-mediated dance of the plankton, *Nature Climate Change*, 4, 880–887, 2014.
- Bianchi, D., Zavatarelli, M., Pinardi, N., Capozzi, R., Capotondi, L., Corselli, C., and Masina, S.: Simulations of ecosystem response during
740 the sporel S1 deposition event, *Paleogeography, Palaeoclimatology, Palaeoecology*, 235, 265–287, 2005.
- Blackford, J. C., Allen, J. I., and Gilbert, F. J.: Ecosystem dynamics at six contrasting sites: A generic modelling study, *Journal of Marine Systems*, 52, 191–215, 2004.
- Blumberg, A. F. and Mellor, G. L.: A Description of a Three-Dimensional Coastal Ocean Circulation Model, *Three Dimensional Coastal Ocean Models*, 1987.
- 745 Boyd, P. W., Cornwall, C. E., Davison, A., Doney, S. C., Fourquez, M., Hurd, C. L., Lima, I. D., and McMin, A.: Biological responses to environmental heterogeneity under future ocean conditions, *Global Change Biology*, 22, 2633–2650, 2016.
- Cavender-Bares, K. K., Karl, D. M., and Chisholm, S. W.: Nutrient gradients in the western North Atlantic Ocean: Relationship to microbial community structure and comparison to patterns in the Pacific Ocean, *Deep-Sea Research I*, 48, 2373–2395, 2001.
- Clainche, Y. L., Levasseur, M., Vezina, A., and Dacey, J. W. H.: Behavior of the ocean DMS(P) pools in the Sargasso Sea viewed in a coupled
750 physical-biogeochemical ocean model, *Can. J. Fish. Aquat. Sci.*, 61, 788–803, 2004.
- Clayton, S. A.: Physical Influences on Phytoplankton Ecology: Models and Observations, Ph.D. thesis, Massachusetts Institute of Technology, 2013.
- Dearman, J. R., Taylor, A. H., and Davidson, K.: Influence of autotroph model complexity on simulations of microbial communities in marine mesocosms, *Marine Ecology Progress Series*, 250, 13–28, 2003.
- 755 Denman, K. L.: Modelling planktonic ecosystems: parameterizing complexity, *Progress in Oceanography*, 57, 429–452, 2003.
- Denman, K. L. and Abbott, M. R.: Time scales of pattern evolution from cross-spectrum analysis of advanced very high resolution radiometer and coastal zone color scanner imagery, *Journal of Geophysical Research*, 99, 7433–7442, 1994.
- Dickey, T., Zedler, S., Yu, X., Doney, S., Frye, D., Jannasch, H., Manov, D., Sigurdson, D., McNeil, J., Dobeck, L., Gilboy, T., Bravo, C., Siegel, D., and Nelson, N.: Physical and biogeochemical variability from hours to years at the Bermuda Testbed Mooring site:

- 760 June 1994–March 1998, *Deep Sea Research Part II: Topical Studies in Oceanography*, 48, 2105–2140, [https://doi.org/10.1016/s0967-0645\(00\)00173-9](https://doi.org/10.1016/s0967-0645(00)00173-9), 2001.
- Doney, S., Glover, D. M., and Najjar, R. G.: A new coupled, one-dimensional biological-physical model for the upper ocean: Applications to the JGOFS Bermuda Atlantic Time-series Study (BATS) site, *Deep-Sea Research II*, 43, 591–624, 1996.
- Doney, S. C.: Major challenges confronting marine biogeochemical modeling, *Global Biogeochemical Cycles*, 13, 705–714, 1999.
- 765 Dugdale, R. C. and Goering, J. J.: Uptake of new and regenerated forms of nitrogen in primary productivity., *Limnology and Oceanography*, 12, 196–206, 1967.
- Fanning, K. A.: Nutrient Proveniences in the Sea: Concentration Ratios, Reaction Rate Ratios, and Ideal Covariation, *Journal of Geophysical Research*, 97, 5693–5712, 1992.
- Fasham, M. J. R., Ducklow, H. W., and McKelie, S. M.: A nitrogen-based model of plankton dynamics in the oceanic mixed layer, *Journal*
- 770 *of Marine Research*, 48, 1990.
- Fennel, K., Losch, M., Schroter, J., and Wenzel, M.: Testing a marine ecosystem model: sensitivity analysis and parameter optimization, *Journal of Marine Systems*, 28, 45–63, 2001.
- Fiori, E., Mazzotti, M., Guerrini, F., and Pistocchi, R.: Combined effects of the herbicide terbuthylazine and temperature on different flagellates of the Northern Adriatic Sea, *Aquatic Toxicology*, 128–129, 79–90, <https://doi.org/10.1016/j.aquatox.2012.12.001>, 2012.
- 775 Friedrichs, M. A. M., Dusenberry, J. A., Anderson, L. A., Armstrong, R. A., Chai, F., Christian, J. R., Doney, S. C., Dunne, J., Fujii, M., Hood, R., Jr, D., Moore, J. K., Schartau, M., Spitz, Y. H., and Wiggert, J. D.: Assessment of skill and portability in regional marine biogeochemical models: Role of multiple planktonic groups, *Journal of Geophysical Research*, 112, 1–22, 2007.
- Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic model of phytoplankton growth and acclimation: responses of the balanced growth rate and chlorophyll a:carbon ratio to light, nutrient limitation and temperature, *Marine Ecology Progress Series*, 148, 187–200,
- 780 1997.
- Gower, J. F. R., Denman, K. L., and Holyer, R. J.: Phytoplankton patchiness indicates the fluctuation spectrum of mesoscale oceanic structure, *Nature*, 288, 157–159, 1980.
- Hamlington, P. E., Roedel, L. P. V., Fox-Kemper, B., Julien, K., and Chini, G. P.: Langmuir-Submesoscale Interactions: Descriptive Analysis of Multiscale Frontal Spin-down Simulations, *Journal of Physical Oceanography*, 44, 2249–2272, 2014.
- 785 Hauri, C., Gruber, N., Vogt, M., Doney, S. C., Feely, R. A., Lachkar, Z., Leinweber, A., McDonnell, A. M. P., Munnich, M., and Plattner, G. K.: Spatiotemporal variability and long-term trends of ocean acidification in the California Current System, *Biogeosciences*, 10, 193–216, 2013.
- Hua, B. L., McWilliams, J. C., and Owens, W. B.: An Objective Analysis of the POLYMODE Local Dynamics Experiment. Part II: Streamfunction and Potential Vorticity Fields during the Intensive Period, *Journal of Physical Oceanography*, 16, 506–522, 1985.
- 790 Hurtt, G. C. and Armstrong, R. A.: A pelagic ecosystem model calibrated with BATS data, *Deep-Sea Research II*, 43, 653–683, 1996.
- Hurtt, G. C. and Armstrong, R. A.: A pelagic ecosystem model calibrated with BATS and OWSI data, *Deep-Sea Research I*, 46, 27–61, 1999.
- Jassby, A. D. and Platt, T.: Mathematical formulation of the relationship between photosynthesis and light for phytoplankton, *Limnology and Oceanography*, 21, 540–547, 1976.
- Jr., D. J. M., Kosnyrev, V. K., Ryan, J. P., and Yoder, J. A.: Covariation of mesoscale ocean color and sea-surface temperature patterns in the
- 795 Sargasso Sea, *Deep-Sea Research II*, 48, 1823–1836, 2001.
- Killworth, P. D.: Time Interpolation of Forcing Fields in Ocean Models, *Journal of Physical Oceanography*, 26, 136–143, 1995.

- Lawson, L. M., Hofmann, E. E., and Spitz, Y. H.: Time series sampling and data assimilation in a simple marine ecosystem model, *Deep-Sea Research II*, 43, 625–651, 1996.
- Levy, M. and Klein, P.: Does the low frequency variability of mesoscale dynamics explain a part of the phytoplankton and zooplankton spectral variability?, *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 460, 1673–1687, 2015.
- Levy, M., Gavart, M., Memery, L., Caniaux, G., and Paci, A.: A four-dimensional mesoscale map of the springbloom in the northeast Atlantic (POMMEexperiment): Results of a prognostic model, *Journal of Geophysical Research*, 110, 2005.
- Lima, I. D., Olson, D. B., and Doney, S. C.: Biological response to frontal dynamics and mesoscale variability in oligotrophic environments: Biological production and community structure, *Journal of Geophysical Research*, 107, 3111, 2002.
- Mahadevan, A.: Spatial heterogeneity and its relation to processes in the upper ocean, pp. 165–182, Springer-Verlag, 2005.
- Mahadevan, A. and Archer, D.: Modeling the impact of fronts and mesoscale circulation on the nutrient supply and biogeochemistry of the upper ocean, *Journal of Geophysical Research*, 105, 1209–1225, 2000.
- Mahadevan, A. and Campbell, J. W.: Biogeochemical patchiness at the sea surface, *Geophysical Research Letters*, 29, 32–1–32–4, 2002.
- Martin, A. P., Richards, K. J., Bracco, A., and Provenzale, A.: Patchy productivity in the open ocean, *Global Biogeochemical Cycles*, 16, 1025, 2002.
- Martiny, A. C., Pham, C. T. A., Primeau, F. W., Vrugt, J. A., Moore, J. K., Levin, S. A., and Lomas, M. W.: Strong latitudinal patterns in the elemental ratios of marine plankton and organic matter, *Nature Geoscience*, 2013.
- Mellor, G. L.: An equation of state for numerical models of oceans and estuaries, *Journal of Atmos. Oceanic Tech.*, 8, 609–611, 1991.
- Mellor, G. L.: One-Dimensional, Ocean Surface Layer Modeling: A Problem and a Solution, *Journal of Physical Oceanography*, 31, 790–809, 2001.
- Mellor, G. L. and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems, *Review of Geophysical Space Physics*, 20, 851–875, 1982.
- Michaels, A. F., Knap, A. H., Dow, R. L., Gundersen, K., Johnson, R. J., Sorensen, J., Close, A., Knauer, G. A., Lohrenz, S. E., Asper, V. A., Tuel, M., and Bidigare, R.: Seasonal pattern of ocean biogeochemistry at the U. S. JGOFS Bermuda Atlantic Time-series Study site, *Deep-Sea Research I*, 41, 1013–1038, 1993.
- Mulholland, M. R. and Lomas, M. W.: Nitrogen uptake and assimilation., pp. 303–384, Academic Press, 2008.
- Mussap, G., Zavatarelli, M., Pinardi, N., and Celio, M.: A management oriented 1-D ecosystem model: Implementation in the Gulf of Trieste (Adriatic Sea), *Regional Studies in Marine Science*, 6, 109–123, 2016.
- Powell, T. M. and Okubo, A.: Turbulence, diffusion and patchiness in the sea, *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 343, 11–18, 1994.
- Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The influence of organisms on the composition of sea water, pp. 26–77, Interscience, 2005.
- Risien, C. M. and Chelton, D. B.: A Global Climatology of Surface Wind and Wind Stress Fields from Eight Years of QuikSCAT Scatterometer Data, *Journal of Physical Oceanography*, 38, 2379–2413, <https://doi.org/10.1175/2008JPO3881.1>, 2008.
- Risien, C. M. and Chelton, D. B.: Scatterometer Climatology of Ocean Winds (SCOW), <http://cioss.coas.oregonstate.edu/scow/>, 2011.
- Roekel, L. P. V., Fox-Kemper, B., Sullivan, P. P., E., P., and Haney, S. R.: The form and orientation of Langmuir cells for misaligned winds and waves, *Journal of Geophysical Research - Oceans*, 117, 2012.

- Schneider, B., Bopp, L., Gehlen, M., Segschneider, J., Frolicher, T. L., Joos, F., Cadule, P., Friedlingstein, P., Doney, S. C., and Behrenfeld, M. J.: Spatio-temporal variability of marine primary and export production in three global coupled climate carbon cycle models, *Biogeosciences Discussions*, 4, 1877–1921, 2007.
- Singh, A., Baer, S. E., Riebesell, U., Martiny, A. C., and Lomas, M. W.: C:N:P stoichiometry at the Bermuda Atlantic Time-series Study station in the North Atlantic Ocean, *Biogeosciences*, 12, 6389–6403, 2015.
- Smith, K. M., Hamlington, P. E., and Fox-Kemper, B.: Effects of submesoscale turbulence on ocean tracers, *Journal of Geophysical Research: Oceans*, 121, 908–933, 2016.
- Smith, K. M., Hamlington, P. E., Niemeier, K. E., Fox-Kemper, B., and Lovenduski, N. S.: Effects of Langmuir Turbulence on Upper Ocean Carbonate Chemistry, *Journal of Advances in Modeling Earth Systems*, 10, <https://doi.org/10.1029/2018MS001486>, 2018.
- Spitz, Y. H., Moisan, J. R., and Abbott, M. R.: Configuring an ecosystem model using data from the Bermuda Atlantic Time Series (BATS), *Deep-Sea Research II*, 48, 1733–1768, 2001.
- Steinberg, D. K., Carlson, C. A., Bates, N. R., Johnson, R. J., Michaels, A. F., and Knap, A. H.: Overview of the US JGOFS Bermuda Atlantic Time-series Study (BATS): a decade-scale look at ocean biology and biogeochemistry, *Deep-Sea Research II*, 48, 1405–1447, 2001.
- Strass, V.: Chlorophyll patchiness caused by mesoscale upwelling at fronts, *Oceanographic Research Papers*, 39, 75–96, 1992.
- Strutton, P. G., Lovenduski, N. S., Mongin, M., and Metear, R.: Quantification of Southern Ocean phytoplankton biomass and primary productivity via satellite observations and biogeochemical models, *CCAMLR Science*, 19, 247–265, 2012.
- Suzuki, N. and Fox-Kemper, B.: Understanding Stokes forces in the wave-averaged equations, *Journal of Geophysical Research - Oceans*, 121, 3579–3596, 2015.
- Triantafyllou, G., Petihakis, G., and Allen, I. J.: Assessing the performance of the Cretan Sea ecosystem model with the use of high frequency M3A buoy data set, *Ann. Geophys.*, 21, 365–375, 2003.
- Tzella, A. and Haynes, P. H.: Small-scale spatial structure in plankton distributions., *Biogeosciences*, 4, 173–179, 2007.
- Vichi, M., Oddo, P., Zavatarelli, M., Coluccelli, A., Coppini, G., Celio, M., Umani, S. F., and Pinardi, N.: Calibration and validation of a one-dimensional complex marine biogeochemical flux model in different areas of the northern Adriatic shelf, *Ann. Geophys.*, 21, 413–436, 2003.
- Vichi, M., Pinardi, N., and Masina, S.: A generalized model of pelagic biogeochemistry for the global ocean ecosystem. Part I: Theory., *Journal of Marine Systems*, 64, 89–109, 2007.
- Vichi, M., Cossarini, G., Mlot, E. G., P, P. L., Lovato, T., Mattia, G., Masina, S., McKiver, W., Pinardi, N., Solidoro, C., and Zavatarelli, M.: The Biogeochemical Flux Model (BFM): Equation Description and User Manual, BFM version 5 (BFM-V5); <http://bfm-community.eu>, Bologna, Italy, 1 edn., release 1.0; BFM Report series N. 1; <http://bfm-community.eu>, 2013.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *Journal of Geophysical Research*, 97, 7373–7382, 1992.
- Wanninkhof, R.: Relationship between windspeed and gas exchange over the ocean revisited, *Limnology and Oceanography Methods*, 12, 351–362, 2014.
- Yoder, J. A., Aiken, J., Swift, R. N., H., F. E., and Stegmann, P. M.: Spatial variability in near-surface chlorophyll a fluorescence measured by the Airborne Oceanographic Lidar (AOL), *Deep-Sea Research II*, 40, 37–53, 1992.