

Discussion of “Snow profile alignment and similarity assessment for aggregating, clustering, and evaluating of snowpack model output for avalanche forecasting”

AUTHOR RESPONSE 1

Herla et al.

October 28, 2020

We thank both reviewers for their thoughtful comments that helped us improve the manuscript. This document provides our responses to the referee comments before highlighting the changes made to our manuscript.

Responses to Referee #1 (TG)

General Comment

Referee General Comment: *In their paper, Herla et al. develop a model to compare snow packing profiles based on the alignment of their stratigraphy. The alignment procedure is based on the dynamic time warping (DTW) algorithm; in short, it enables the flexible comparison and clustering of snow layer profiles. The paper is well written and it makes a convincing case for the use of DTW. Notably, it introduces non-trivial domain-specific adaptations: for one, a cost metric built from a weighted combination of heterogeneous discrete parameters (grain type, hardness, age). Each variable is preprocessed carefully in a knowledge-based way (e.g. grain type was addressed through a suitable similarity matrix). Another adaptation is the selection of open-ended alignments coupled with slope-limited patterns, with suitable weights and windows sizes serving as hyper-parameters. The description of the DTW algorithm and the construction of the local cost matrix D are appropriate and clear.*

Author General Comment: We thank TG for his constructive review and helpful comments. We appreciate the encouraging comments about our domain-specific adaptations of DTW. Please see below for our responses to specific comments and suggestions from Referee #1. Additions to the manuscript are included in our responses in quotes, where page- and line-numbers refer to the revised manuscript that includes the highlighted and marked-up changes.

Specific Comments

Referee Comment (RC) 1.1: *Page 9. "In fig. 1..." . Figure 1 is illustrative, but it uses different warping parameters than your final set (e.g. constraints). Hence, consider adding "*As an illustration,* in Figure 1..."*

Author Response (AR) 1.1: Thank you! Done. (P10 L16)

RC 1.2: *'no "stopping" or "going back" is allowed.'* To be more precise: *'no "going back in time" is allowed'.* Actually some patterns, such as your illustration of Figure 1, do allow local "stopping", as you note.

AR 1.2: Thank you! Changed it. (P9 L16)

RC 1.3: *In 2.2.4. (and possibly other places) "The symmetric Sakoe-Chiba local slope constraint" is mentioned. However Sakoe-Chiba's paper introduced *several* possible recursions (see their Table I). Specifically, the one you adopt appears to be "P=1, symmetric". This should be made explicit. You may also mention that it's "symmetricP1" in R.*

AR 1.3: We agree and made this more explicit. (P14 L9ff, P24 L4)

RC 1.4: *Please mention the dtw R package used and the corresponding version.*

AR 1.4: Done. We added a brief note in the Discussion (P22 L5) and a more detailed explanation to the Code Availability section (P23 L28).

RC 1.5: *"Mirroring D about its anti-diagonal". An alternative, possibly simpler way to express this is to say that one reverses both time series.*

AR 1.5: Good suggestion! Changed. (P14 L26)

RC 1.6: *Appendix A: "(the weighting factor 2 indicates that both indices i and j have been incremented)". Weighting factors, or slope weights, ensure that the normalisation prefactor in A2 is path-independent and has that specific form. [This is well explained in Rabiner-Juang's book on speech recognition.] Either omit the sentence or rephrase slightly, e.g., "the slope weights depend on the local indices' increments and ensure that alignments can be compared".*

AR 1.6: Thanks, we rephrased the explanation. (P24 L13f)

RC 1.7: *"(expressed as Manhattan distance)" to clarify: "(expressed as Manhattan distance *from the matrix origin, for symmetric recursions*)"*

AR 1.7: Done. (P25 L14)

RC 1.8: *Possible material for SI: - A figure with step patterns and weights, e.g. subfigure A) plot(symmetrict2) because it is used in Figure 1, and subfigure B) plot(symmetrictP1), used by your alignment, for comparison.*

AR 1.8: Thanks for the suggestion. We included the figure and its explanation in an Appendix B. (P25 L23ff)

Responses to Referee #2 (Matthieu Lafaysse)

General Comment

Referee General Comment RC 2.1: *Herla et al. present a matching algorithm between snowpack profiles able to classify and evaluate snow cover simulations designed for avalanche hazard forecasting. I fully agree with the need of new tools to extend the evaluations of these models and facilitate their use by forecasters. The question is very well introduced in the paper. I am also glad to see that the proposal of Hagenmuller and Pilloix, 2016 to use such matching algorithms for that purpose found an echo in other modelling teams and is applied in new contexts and with other similarity metrics. I also like the discussion about a possible concrete use of the method in an operational forecasting system. The results shown by the authors in a clustering application are interesting and a number of new advances in this paper are valuable for the community (new metrics, classification and synthesis methods).*

However, I am a bit uncomfortable with the way the authors present their work in this context. Indeed their unique reference to Hagenmuller and Pilloix, 2016 is: "Hagenmuller and Pilloix (2016) and Hagenmuller (2018) were the first to align, cluster and aggregate one-dimensional snow hardness profiles from ram resistance field measurements using Dynamic Time Warping (DTW), a method from the fields of time series analysis and data mining. While their contribution demonstrates the usefulness of DTW for snow profile comparisons, their method is not general enough to allow for meaningful comparisons of snow profiles from different sources and varying levels of details." Then, they present their work as "a new approach for computationally comparing, grouping and summarizing snow profiles" and they present the algorithm in details in Section 2.2.1 as it could be expected for the first publication of an algorithm. They also state in the discussion "we are the first to present an alignment algorithm and similarity measure for generic snow profiles". However, the method proposed in this paper can not be told as new. Indeed, the algorithm is extremely similar to the one developed by our colleague P. Hagenmuller and which is already used in several publications (with or without a detailed description of DTW depending on the references): Hagenmuller and Pilloix, 2016; Teich et al., 2019; Hagenmuller et al. 2018a; Hagenmuller et al. 2018b; Viallon et al. (accepted). Note also that DTW was also already applied on snow by Schaller et al., 2016. The innovation of Herla et al. is mainly the distance the authors introduce in the application of this algorithm, in order to use a criteria more focused on mechanical stability, but the general philosophy of the method is the same. Furthermore, the application of such algorithm for clustering also already appears in Hagenmuller et al., 2018b and also include a selection method of a representative profile. Although this reference is not a publication in a peer-review journal but a conference proceedings, the authors are aware of this contribution.

Author General Comment AC 2.1: We thank Matthieu Lafaysse for his review, in particular for the encouraging comment related to our attempt of creating tools of operational value for avalanche forecasting, as well as for the honest presentation of his concerns. In his review we identified three main overarching themes that we will address first before responding to the more technical comments in a point-by-point manner further below.

The main concerns of the referee are centered around proper accreditation, clear communication of our innovations as well as clear separation towards other contributions. We addressed these issues altering the Introduction, the Discussion, and the Abstract, including the additional references suggested by the referee, highlighting how we identified research needs, and more explicitly communicating the innovations of our contribution. Our main modifications include the following (new additions and edits are printed in black fonts, while text from our first submission is printed in gray; page- and line-numbers refer to the revised manuscript with the marked-up changes):

- We added an explanation of the data type we are working with:

“When observing a snow profile, traditionally the most commonly recorded layer characteristics are snow grain type, grain size and layer hardness. [...] Snow profiles that contain information about these layer characteristics are referred to as generic snow profiles hereafter and represent the main source of snow stratigraphy information in operational contexts.” (P2 L15ff)

- We provide a more detailed summary of how DTW has been introduced to the snow community, and we illustrate in more detail how the layer matching algorithm by Hagenmuller and Pilloix (2016) has been applied in the snow community. Furthermore, we made the research gap more specific:
“Hagenmuller and Pilloix (2016) and Hagenmuller et al. (2018b), as well as Schaller et al. (2016) introduced Dynamic Time Warping (DTW), a longstanding method from the fields of time series analysis and data mining, to the snow community. Both implemented a layer matching algorithm to align, cluster, and aggregate one-dimensional snow hardness (or density) profiles from field measurements and thereby demonstrated the usefulness of DTW for snow profile comparisons. In the subsequent years, the layer matching algorithm by Hagenmuller and Pilloix (2016) has been applied to evaluate snow penetrometers (Hagenmuller et al. 2018a) or to characterize spatial variability of the snow cover from ram resistance field measurements (Teich et al. 2019). Consequently, their approach has focused on one-dimensional, continuous, numerical sequences and is not readily applicable to operational snowpack observations by avalanche forecasters.” (P4 L4ff)
- We then explicitly formulate our research goals:
“The objective of this study is to introduce an approach for computationally comparing, grouping and summarizing generic snow profiles that consist of multi-dimensional, discrete sequences of categorical, numerical, and ordinal data types. To maximize the value for avalanche forecasting, our methods focus on structural elements in the profiles that are particularly important for avalanche hazard assessments and can handle both simulated profiles and manual observations with different levels of detail.” (P4 L14ff)
- We re-iterate the innovations of our contribution in the Discussion, where we also include the recently published paper by the referee’s group (Viallon-Galinier et al. 2020). Since the efforts behind that paper were not published or otherwise publicly accessible until mid to end September 2020, we could not have included them in our original submission.

“Building on the well-established and longstanding concept of Dynamic Time Warping (DTW), we developed a snow profile alignment algorithm that combines multiple layer characteristics of categorical, numerical, and ordinal format into a weighted metric and feeds into existing DTW algorithms such as the open-source R package dtw. Moreover, we reviewed and derived useful DTW configurations and hyper-parameter settings for snow profile applications. Since these applications rely on operationally available profile observations that typically focus on information relevant for current avalanche conditions only, our approach is able to handle missing data and take advantage of select layer date tags. To maximize the layer matching performance for profiles with limited details, we implemented a scheme for preferential layer matching based on domain knowledge. In parallel, Viallon-Galinier et al. (2020) extended the layer matching algorithm from Hagenmuller and Pilloix (2016) to conduct a detailed, process-related evaluation of the snowpack model Crocus based on high-quality snow profile observations with a large variety of observed variables that are sampled at specific study sites at regular intervals. Since their goal is the correction of deviating model states with a direct insertion assimilation scheme based on point scale simulations and observations, their evaluation targets not only each individual layer, but also each individual layer characteristic separately. To address operational

avalanche forecasting needs, we additionally developed a similarity measure that focuses on avalanche forecasting specific considerations where certain layers are considered more important. Moreover, combining information from individual layers and their characteristics into a scalar measure allows for clustering and aggregating of sets of profiles to characterize and evaluate the regional scale avalanche hazard conditions.” (P22 L1ff)

Specific Comments

RC 2.2: *The introduction should better emphasize the own innovations of this paper and avoid general statements presenting the whole method as a new algorithm.*

AR 2.2: Done, see general comment above. Additionally, we removed all ambiguous statements.

RC 2.3: *Section 2.2.1 should better emphasize what is common and what differs between this algorithm and the algorithm of Hagenmuller and Pilloix, 2016 and following papers. The details of DTW already described in previous literature can be moved to an Appendix in order to be more focused on the novelties in the main text.*

AR 2.3: We improved upon the distinction between the different contributions, see general comment above. However, we prefer not to move the background section on DTW (i.e., Section 2.2.1) to an appendix. First, we believe it is appropriate and necessary for a methods paper to transparently communicate the presented method. Second, in our opinion, other publications in the snow community that present or use a matching algorithm based on DTW have so far not presented the method in an accessible and reproducible manner. And third, the journal addresses a wider audience of geophysicists to whom a matching algorithm for multi-dimensional, mixed-type sequences or time series might be of relevance. Demonstrating the modification and application of DTW based on an illustrative example such as snow profiles, while relating to the rich body of literature in the DTW community, will allow readers of *gmd* to develop similar methods for other geophysical disciplines.

RC 2.4: *Then, it is unclear what is unsatisfying in the algorithm of P. Hagenmuller. What do the author mean by “not general enough to allow for meaningful comparisons of snow profiles”? Hagenmuller and Pilloix, 2016, discuss the possibility to modify the distance criteria depending on applications: “The metrics D and V between profiles whose definition involve the mean square difference of logarithmic hardness can be adapted to incorporate other snow properties. [...]” with examples. In Viallon et al., accepted, we applied DTW with a more general distance combining density, liquid water content, grain shape and depth, better suited for an overall model evaluation and the algorithm behaves well in this context. The distance presented here by the authors is probably better suited to their application but can not pretend to be more “general”. Do the authors refer only to the management of missing values in snow profiles? Or is there something else? I think it is important to be more specific on the issues in previous references to better justify and emphasize the innovations they want to publish here.*

AR 2.4: We addressed this concern in the modifications described in the general comment above.

RC 2.5: *A number of choices in the distance definition are model-dependent relatively to the SNOWPACK model. For instance, neither layer hardness or layer date are diagnostics of the Crocus model (similar to SNOWPACK in terms of complexity). Furthermore, a number of considerations in the distance definition for grain type are based on considerations about the current typical behaviour of SNOWPACK (lines 5-6 and 23-31 page 5). These considerations might not apply to another model or even to a future version of SNOWPACK with for instance new parameterizations of snow metamorphism. This option can be arguable if there is added value to make this choice but the limitation should be clearly discussed. Will it be necessary to*

modify the distance definition if significant changes are implemented in the model?

AR 2.5: We appreciate that the referee highlighted the close tie of our algorithm to the SNOWPACK model and is interested in limitations with respect to future version of SNOWPACK and their parametrizations of snow metamorphism.

To explain how the simulated variables are calculated and to explain the relation of our algorithm to SNOWPACK, we added the following paragraph at the beginning of Section 2.1: “*While grain types are computed by snow models based on parametrizations of snow metamorphism, and simulated burial date information can easily be derived from simulated deposition date or age of the layer, layer hardness is only a diagnostic variable provided by the model SNOWPACK, but not by Crocus. Therefore the following distance functions are presented in the light of SNOWPACK, and the application to other snow model output may require some modifications.*” (P5 L8ff) More details about the conversion of layer age or date information is then provided in Sect. 2.1.3.

To communicate our reasoning behind the modifications of grain type similarities more clearly, we rewrote the following paragraph: “*SH and DH layers are formed by very different processes—by the deposition of hoar onto the snow surface versus by kinetic growth of crystals within the snowpack. Consequently, Lehning et al. (2001) evaluated the similarity of the two grain types as completely dissimilar. However, both SH and DH represent hazardous weak layers and are of comparable importance in avalanche hazard assessments (Schweizer and Jamieson 2001). Furthermore, practical experience with the current version of SNOWPACK shows that SH layers are often converted to DH layers once buried. To account for both of these aspects, we raised their similarity from 0 to 0.9 for both tasks (Table 1A, B).*” (P7 L1ff)

To summarize: While our modifications to the grain type similarities accommodate peculiarities of SNOWPACK, they are all rooted in considerations of avalanche hazard assessment. We do therefore not expect any necessary modifications for future versions of snowpack that include new parametrizations of snow metamorphism.

RC 2.6: *Finally, it is questionable whereas there is really added value for the community in the future to provide two separate codes from two research teams based on a similar algorithm. The authors requested the code of P. Hagenmuller last winter and they received it with a number of explanations from P. Hagenmuller. This is surprisingly not mentioned in the Acknowledgements section. I think the authors should justify in the paper the need for another code by describing the reasons which have probably prevent them using directly the code from P. Hagenmuller. A better understanding of these limitations might help to avoid further work duplication and hopefully allow more shared developments in the post-processing of snow models in the future, although I am fully aware of the difficulties of such collaborative developments.*

AR 2.6: We apologize for this oversight and we now thank P. Hagenmuller in the Acknowledgements for an email exchange at the outset of this research project. Furthermore, we agree that it is desirable to reduce redundant work and combine efforts, especially in a niche research field like snowpack modeling and avalanche forecasting. However, we also believe that there is value in developing multiple approaches and letting the research community decide what works best for their specific objectives. Furthermore, we strongly believe that the open access approach is most valuable for the research community and therefore built our algorithm on the open source R package dtw, which provides excellent documentation and is also implemented in other programming languages, instead of Hagenmuller’s code.

RC 2.7: *Is there a limitation of snow height differences to apply the rescaling? Is it meaningful to rescale profiles even when the relative difference of snow height reach for instance 500% or more?*

AR 2.7: If the two profiles stem from the same forecast area, or in other words from an area with homogeneous (albeit elevation and aspect dependent) meteorological conditions, rescaling of the profiles is reasonable independently from the relative difference of the snow heights.

Especially when the relative difference of snow heights is extremely large, the two profiles have been subject to systematic variations in their meteorological forcing (e.g., increase of snowfall amounts with elevation). It is exactly these situations, where the rescaling is of particular importance.

Since that question is also related to clustering and aggregating applications, we added the following clarification to the Discussion. *“The similarity measure, and consequently the clustering and aggregating applications are purely based on the agreement of the snowpack structure. Hence, snow depth is not a driver of our similarity assessment unless it leads to deviations in the snow stratigraphy. If combined with monitoring of the snow depths distribution, a clustering or aggregation application can provide a comprehensive picture of the conditions within a specific forecast area.”* (P23 L6ff)

RC 2.8: *Page 10 Line 21: The formulation is a bit ambiguous. Was the value of 0.5 cm indeed chosen for grid resampling in this work?*

AR 2.8: We added an explanation, so that the relevant paragraph reads: *“Once the profiles have been rescaled, each of the two profiles consists of a series of discrete layers along an irregular height grid. To equalize the two different height grids, we resample the profiles onto a regular grid with a constant sampling rate, which represents the final resolution for the alignment procedure. While our algorithm allows users to flexibly set the sampling rate, a resolution of about half a centimeter ensures that typically thin, hazardous weak layers are being captured. Hence, snow profiles in the presented examples have been resampled to 0.5 cm.”* (P11 L15ff)

RC 2.9: *The variables chosen to compute the similarity measure are not prognostic variables of the model but rather indirect diagnostics (grain type, hardness) involving very uncertain parameterizations from prognostic variables. Beyond the clustering application presented in this paper, the possible use of this criteria for model evaluation raises some questions. Indeed, it can not be established if this metric would be really representative of the real skill of the physical model or if these transformation functions would not prevail in the obtained metric. I fully understand the motivation relative to the avalanche hazard application but this topic should be discussed because (1) a perfect model in terms of physical evolution laws might not provide a perfect similarity measure due to errors in the variables transformation and (2) this criteria might not be recommended for a process-related model evaluation where I think a metric based on density and temperature profiles for instance should be preferred. In spite of this comment, I acknowledge that the detailed thinking about an appropriate weighting of layers in the final similarity metric is really interesting and useful (although a bit complex) from the mechanical point of view.*

AR 2.9: We agree with the referee in that a model evaluation based on indirect diagnostics skews the obtained results and does not provide a direct evaluation of the physical model skill. Since our research focuses on making snowpack simulations more accessible for avalanche forecasters, and since those indirect diagnostics are the relevant variables for avalanche hazard applications, an evaluation limited to the prognostic variables is not informative and relevant in our context. To convey this idea more strongly, we extended the relevant paragraph in the Introduction:

“Furthermore, the ability to operationally compare simulated snow profiles against observed ones provides an avenue for continuously monitoring the quality of the simulations and correct them if necessary. To judge the operational value of snowpack models for avalanche forecasting, it is particularly important to focus on snowpack features and layer characteristics that are of direct relevance for avalanche hazard assessments. Since operational snowpack observations and relevant layer characteristics are expressed by variables (such as grain type and layer hardness) that are only indirectly diagnosed by models, the parametrization from prognostic variables introduces another layer of uncertainty. The evaluation of these models for practical purposes

therefore needs to take all of these uncertainties into account.“ (P3 L26ff)

References

- Hagenmuller, P. and T. Pilloix, 2016: A New Method for Comparing and Matching Snow Profiles, Application for Profiles Measured by Penetrometers. *Frontiers in Earth Science*, **4**, doi:10.3389/feart.2016.00052.
- Hagenmuller, P., A. van Herwijnen, C. Pielmeier, and H.-P. Marshall, 2018a: Evaluation of the snow penetrometer Avatech SP2. *Cold Regions Science and Technology*, **149**, 83–94, doi: 10.1016/j.coldregions.2018.02.006.
- Hagenmuller, P., L. Viallon, C. Bouchayer, M. Teich, M. Lafaysse, and V. Vionnet, 2018b: Quantitative Comparison of Snow Profiles. *Proceedings of the 2018 international snow science workshop, Innsbruck, AUT*, 876—879, URL <https://arc.lib.montana.edu/snow-science/item/2668>.
- Lehning, M., C. Fierz, and C. Lundy, 2001: An objective snow profile comparison method and its application to SNOWPACK. *Cold Regions Science and Technology*, **33 (2-3)**, 253–261, doi:10.1016/S0165-232X(01)00044-1.
- Schaller, C. F., J. Freitag, S. Kipfstuhl, T. Laepple, H. Christian Steen-Larsen, and O. Eisen, 2016: A representative density profile of the North Greenland snowpack. *Cryosphere*, **10 (5)**, 1991–2002, doi:10.5194/tc-10-1991-2016, URL <https://tc.copernicus.org/articles/10/1991/2016/>.
- Schweizer, J. and J. B. Jamieson, 2001: Snow cover properties for skier triggering of avalanches. *Cold Regions Science and Technology*, **33 (2-3)**, 207–221, doi:10.1016/S0165-232X(01)00039-8.
- Teich, M., A. D. Giunta, P. Hagenmuller, P. Bebi, M. Schneebeli, and M. J. Jenkins, 2019: Effects of bark beetle attacks on forest snowpack and avalanche formation — Implications for protection forest management. *Forest Ecology and Management*, **438**, 186–203, doi:10.1016/j.foreco.2019.01.052.
- Viallon-Galinier, L., P. Hagenmuller, and M. Lafaysse, 2020: Forcing and evaluating detailed snow cover models with stratigraphy observations. *Cold Regions Science and Technology*, **180**, 103–163, doi:10.1016/j.coldregions.2020.103163.

Snow profile alignment and similarity assessment for aggregating, clustering, and evaluating of snowpack model output for avalanche forecasting

Florian Herla¹, Simon Horton^{1,2}, Patrick Mair³, and Pascal Haegeli¹

¹Simon Fraser University, Burnaby, BC, Canada

²Avalanche Canada, Revelstoke, BC, Canada

³Harvard University, Cambridge, MA, USA

Correspondence: Florian Herla (fherla@sfu.ca)

Abstract. Snowpack models simulate the evolution of the snow stratigraphy based on meteorological inputs and have the potential to support avalanche risk management operations with complementary information relevant to their avalanche hazard assessment, especially in data-sparse regions or at times of unfavorable weather and hazard conditions. However, the adoption of snowpack models in operational avalanche forecasting has been limited, predominantly due to missing data processing algorithms and uncertainty around model validity. Thus, to enhance the usefulness of snowpack models for the avalanche industry, numerical methods are required that evaluate and summarize snowpack model output in accessible and relevant ways. We present algorithms that compare and assess generic snowpack data from both human observations and models, [which consist of multi-dimensional sequences describing the snow characteristics of grain type, hardness, and age](#). Our approach exploits Dynamic Time Warping, a well-established method in the data sciences, to match layers between snow profiles and thereby align them. The similarity of the aligned profiles is then evaluated by our independent similarity measure based on characteristics relevant for avalanche hazard assessment. Since our methods provide the necessary quantitative link to data clustering and aggregating methods, we demonstrate how snowpack model output can be grouped and summarized according to similar hazard conditions. Through emulating a human avalanche hazard assessment approach, our methods aim to promote the operational application of snowpack models so that avalanche forecasters can begin to build understanding in how to interpret and when to trust operational snowpack simulations.

Copyright statement.

1 Introduction

Snow avalanches are a serious mountain hazard, whose risk is managed through a combination of long- and short-term mitigation measures, depending on the character of the exposed elements-at-risk. Avalanche forecasting—the prediction of avalanche hazard over a specific area of terrain (Campbell et al., 2016)—is a critical prerequisite for choosing effective short-term mitiga-

tion measures and timing them properly (e.g., publication of advisories, temporary closures, proactive triggering of avalanches). The task of avalanche forecasters¹ is to integrate the available weather, snowpack and avalanche observations into a coherent mental model of the hazard conditions across their area of interest (LaChapelle, 1966, 1980; McClung, 2002). Statham et al. (2018) describe the essence of avalanche forecasting as answering four sequential questions: 1) *What* type of avalanche problem(s) exist? 2) *Where* are these problems located in the terrain? 3) *How likely* are avalanches to occur? and 4) *How big* will these avalanches be? Because of the complexity of the avalanche system and the large uncertainty due to the spatial and temporal variability of snowpack properties (Schweizer et al., 2007), avalanche forecasts are subjective expert judgments that are expressed in qualitative degrees of belief (Vick, 2002).

Snow profiles describing the stratigraphy of the snowpack and the characteristics of the individual layers (McClung and Schaerer, 2006) are an important source of information for avalanche forecasting. While avalanche observations offer direct evidence of unstable conditions, the information on structural weaknesses and slab properties contained in snow profiles is crucial for developing a more complete understanding of the nature of the avalanche hazard, its spatial distribution, as well as making predictions about the likelihood of avalanches and their expected size. To adequately capture the conditions in an area of interest, it is common practice for avalanche forecasters to collect snow profile information from a variety of informative locations and employ targeted sampling to address specific hypotheses. When observing a snow profile, traditionally the most commonly recorded layer characteristics are snow grain type, grain size and layer hardness. In addition, layers representing critical structural weaknesses are often labeled with their burial date to facilitate tracking and simplify communication (Canadian Avalanche Association, 2016). Snow profiles that contain information about these layer characteristics are referred to as generic snow profiles hereafter and represent the main source of snow stratigraphy information in operational contexts. Snowpack tests might be performed next to profile locations to examine the potential for fracture initiation and failure propagation along specific layers of interest.

As a winter progresses, avalanche forecasters continuously synthesize the collected snow profile information into a comprehensive picture of existing hazard conditions across the terrain. While experienced forecasters can process snow profile information intuitively and effortlessly, the process is actually a challenging exercise in multi-dimensional pattern recognition, pattern matching, and data assimilation, which requires several advanced skills. These include matching key features between profiles, assessing the similarity or dissimilarity of profiles, combining the information of several profiles into an overall perspective, as well as extrapolating the identified patterns across terrain based on knowledge of snowpack processes and how they are affected by terrain. In North America, it is common practice among avalanche forecasters to document their understanding of the local snowpack by sketching synthesized snow profiles for different areas of interest (e.g., elevation or aspect specific).

Since the late 1980s, physically-based numerical snowpack models have been developed to expand the available information sources for avalanche forecasters beyond traditional field observations. The most commonly used snowpack models are Crocus (Brun et al., 1989; Vionnet et al., 2012), and SNOWPACK (Lehning et al., 1999; Bartelt et al., 2002; Lehning et al., 2002b, a). Both of these models simulate the stratigraphy of the snowpack at a point location by integrating meteorological input data

¹We use the term avalanche forecaster to describe anybody who assesses avalanche hazard conditions to make decisions about short-term mitigation options. This can include public avalanche forecasters, avalanche safety technicians, ski patrollers, mountain guides, and private recreationists.

over a winter season. The source for the meteorological forcing can be time series of in-situ observations, outputs of numerical weather prediction models, or assimilation products that integrate both. The physical properties of the individual snow layers (e.g., grain type, grain size, hardness) are simulated using empirical representations of the key snowpack processes (e.g., snow metamorphism, water percolation, settlement) that are tied together by conservation of mass and energy.

5 Over the last 20 years, extensive amounts of research have been conducted to improve the capabilities of snowpack models, and explore their application for avalanche forecasting. Many contributions evaluated or improved the skill of the models with respect to hazardous weak layer formation (Fierz, 1998; Bellaire et al., 2011, 2013; Horton et al., 2014, 2015; Horton and Jamieson, 2016; Van Peurseem et al., 2016), or weak layer detection (Monti et al., 2014a). Others tried to assess snow stability from model outputs (Schweizer et al., 2006; Schirmer et al., 2010; Monti et al., 2014b) and estimate danger levels (Lehning et al., 2004; Schirmer et al., 2009; Bellaire and Jamieson, 2013). Vionnet et al. (2016) and Vionnet et al. (2018) specifically
10 evaluated and improved meteorological data from a weather prediction model serving as input to a snow cover model. While all studies agree that snowpack modeling has the potential to add value to avalanche forecasting, the understanding of under which circumstances and to which degree these models can add value (especially when coupled with weather prediction models) seems to be limited. This knowledge gap is a major hurdle for developing necessary trust for the operational use of
15 these models.

In Canada, the combination of numerical weather and snowpack models offers a tremendous opportunity for providing avalanche forecasters with useful information on snowpack conditions in otherwise data-sparse regions (Storm, 2012). However, the integration of physical snowpack models into operational avalanche forecasting has so far been limited. Informal conversations with forecasters highlight two main issues: 1) the overwhelming volume of data produced by the models, and
20 2) validity concerns due to cumulative impact of potentially inaccurate weather inputs. Morin et al. (2020) provide a more detailed discussion of the challenges around the operational use of snowpack models, which the authors classify into four main categories: issues of accessibility, interpretability, relevance and integrity.

Addressing the two issues effectively requires the development of computer-based methods that can process large numbers of snow profiles. An algorithm for objectively assessing the similarity of simulated snow profiles is the necessary foundation
25 for computationally emulating the snowpack data synthesis process of avalanche forecasters and meaningfully reducing the data volume to a manageable level. Furthermore, the ability to operationally compare simulated snow profiles against observed ones provides an avenue for continuously monitoring the quality of the simulations and correct them if necessary. To judge the operational value of snowpack models for avalanche forecasting, it is particularly important to focus on snowpack features and layer characteristics that are of direct relevance for avalanche hazard assessments. Since operational snowpack observations and relevant layer characteristics are expressed by variables (such as grain type and layer hardness) that are only indirectly diagnosed by models, the parametrization from prognostic variables introduces another layer of uncertainty. The evaluation of these models for practical purposes therefore needs to take all of these uncertainties into account.

While numerical methods for comparing simulated snow profiles exist, they are unable to address the operational needs described above. To evaluate the performance of SNOWPACK, Lehning et al. (2001) developed an algorithm for comparing
35 modeled profiles against manual observations. Since their approach is only concerned with finding manually observed layers

in a specific depth range of the modeled profile, it is not suitable for subsequent clustering and aggregating. Moreover, their agreement score for snow profile pairs is focused on providing insight for model improvements, which has different similarity assessment needs than comparing snow profiles for avalanche hazard assessment purposes. Hagenmuller and Pilloix (2016) and Hagenmuller et al. (2018b) ~~were the first to align, cluster and aggregate one-dimensional snow hardness profiles from ram-resistance field measurements using~~, as well as Schaller et al. (2016) introduced Dynamic Time Warping (DTW), a longstanding method from the fields of time series analysis and data mining. ~~While their contribution demonstrates~~, to the snow community. Both implemented a layer matching algorithm to align, cluster, and aggregate one-dimensional snow hardness (or density) profiles from field measurements and thereby demonstrated the usefulness of DTW for snow profile comparisons; ~~their method is not general enough to allow for meaningful comparisons of snow profiles from different sources and varying levels of details.~~ In the subsequent years, the layer matching algorithm by Hagenmuller and Pilloix (2016) has been applied to evaluate snow penetrometers (Hagenmuller et al., 2018a) or to characterize spatial variability of the snow cover from ram resistance field measurements (Teich et al., 2019). Consequently, their approach has focused on one-dimensional, continuous, numerical sequences and is not readily applicable to operational snowpack observations by avalanche forecasters.

The objective of this study is to introduce a new an approach for computationally comparing, grouping and summarizing generic snow profiles that ~~can handle both simulated profiles and manual observations with different levels of detail~~ consist of multi-dimensional, discrete sequences of categorical, numerical, and ordinal data types. To maximize the value for avalanche forecasting, our methods focus on structural elements in the profiles that are particularly important for avalanche hazard assessments and can handle both simulated profiles and manual observations with different levels of detail. We approach the task by numerically emulating the cognitive process of human forecasters: We first present a layer matching algorithm that aligns profiles in a way that a similarity measure can evaluate their agreement. We then exploit the resulting similarity score between pairs of snow profiles to cluster snow profiles into distinct groups and aggregate them into a representative profile. The derivation of the snow profile alignment algorithm and the similarity measure is presented in Sect. 2, whereas the new methods are valued through practical aggregation and clustering applications as described in Sect. 3. We discuss the implications of our approach alongside its limitations, and conclude with future perspectives in Sect. 4. We believe that the algorithms presented in this paper provide an important step for the development of operational data aggregation and validation algorithms that can make large-scale snowpack simulations more accessible and relevant for avalanche forecasters.

2 Derivation of the snow profile alignment algorithm and similarity measure

In this section we describe how snow profiles can be aligned by matching layers between them (Sect. 2.2), and define a similarity measure to evaluate the agreement between aligned profiles (Sect. 2.3). Since both of these tasks require a method for assessing differences between individual snowpack layers, we start with that in Sect. 2.1.

2.1 Assessing differences between individual snow layers

To align snow profiles and determine the similarity of snow profiles as a whole, we need a method for assessing the similarity of individual layers. While snowpack models provide a wide range of layer properties, the most commonly used characteristics by practitioners are snow grain type, layer hardness, and burial date. To assess the similarity between individual layers that incorporate all three layer characteristics, we first define distance functions for these characteristics, which are normalized to the interval $[0, 1]$ to make them comparable. A distance of zero means that two layer characteristics are identical, whereas a distance of one represents complete dissimilarity.

While grain types are computed by snow models based on parametrizations of snow metamorphism, and simulated burial date information can easily be derived from simulated deposition date or age of the layer, layer hardness is only a diagnostic variable provided by the model SNOWPACK, but not by Crocus. Therefore the following distance functions are presented in the light of SNOWPACK, and the application to other snow model output may require some modifications.

2.1.1 Distance function for grain type

The international classification for snow on the ground (Fierz et al., 2009) organizes snow grain types (also known as grain shapes) into main grain type classes, which can in turn be broken down into more nuanced sub-classes. The following grain types are typical in avalanche forecasting contexts: precipitation particles (PP), decomposing and fragmented particles (DF), round grains (RG), faceted crystals (FC) (including the sub-class rounding facets, FCxr), surface and depth hoar (SH and DH), and melt forms (MF) (including the sub-class melt-freeze crusts, MFcr). New snow layers mostly consist of PP and DF, SH and DH are prototypical persistent weak layers, and MFcr often promote the faceting of adjacent grains, which weakens the interface (Jamieson, 2006). FC, including FCxr, can also be considered persistent weak layers, even though not every faceted layer is a layer of concern according to common snow profile analysis techniques that consider combinations of grain type and grain size (amongst other properties) to identify structural weaknesses in the snowpack (Schweizer and Jamieson, 2007). Following that concept, we use the term *bulk layers* for layers that constitute a large proportion of the snowpack without being structurally weak. While avalanche forecasters typically regard FC indicative of weaker snowpack layers, simulated layers of FC tend to be associated with smaller faceted crystals and thicker layers that represent bulk layers rather than weak layers since SNOWPACK classifies any faceted grain with a size greater than 1.5 mm as DH.

Since grain type is a categorical variable, calculating distances between non-identical grain types is non-trivial. Our approach builds on the original method developed by Lehning et al. (2001), who defined a matrix of normalized grain type similarities between all possible pairs of grain types based on the physics of their formation and metamorphosis. Their approach evaluates the modeled grain type stratigraphy to identify model deficiencies and offer insight for model improvements. By contrast, our focus is on matching layers between snow profiles and assessing their similarity for avalanche hazard assessments. For the layer matching task, the similarity between grain types should indeed be evaluated partly based on their formation processes but also on the knowledge of snowpack model quirks and differences between modeled and observed profiles. For assessing the similarity of profiles, however, the similarity between grain types should be evaluated based on their implications for the

Table 1. Similarities between snow grain types as used for the layer alignment of snow profiles (A), and as used for the similarity assessment between snow profiles (B). Gray cells highlight modifications to the matrix used in Lehning et al. (2001), and bold fonts highlight differences between the two tables A and B.

A) Grain type similarity (snow profile alignments)										
	PP	DF	RG	FC	DH	SH	MF	FCxr	MFcr	na
PP	1.00									0.60
DF	0.80	1.00								0.60
RG	0.50	0.80	1.00							0.60
FC	0.20	0.40	0.40	1.00						0.50
DH	0.00	0.00	0.10	0.80	1.00					0.40
SH	0.00	0.00	0.00	0.60	0.90	1.00				0.40
MF	0.00	0.00	0.00	0.00	0.00	0.00	1.00			0.50
FCxr	0.20	0.40	0.50	0.80	0.70	0.00	0.00	1.00		0.60
MFcr	0.00	0.00	0.00	0.00	0.00	0.00	0.20	0.00	1.00	0.40

B) Grain type similarity (snow profile similarity assessments)										
	PP	DF	RG	FC	DH	SH	MF	FCxr	MFcr	na
PP	1.00									0.50
DF	0.80	1.00								0.50
RG	0.50	0.80	1.00							0.50
FC	0.20	0.40	0.40	1.00						0.50
DH	0.00	0.00	0.10	0.50	1.00					0.50
SH	0.00	0.00	0.00	0.30	0.90	1.00				0.50
MF	0.00	0.00	0.00	0.00	0.00	0.00	1.00			0.50
FCxr	0.20	0.40	0.50	0.60	0.40	0.00	0.00	1.00		0.50
MFcr	0.00	0.00	0.00	0.00	0.00	0.00	0.20	0.00	1.00	0.50

hazard conditions rather than their formation processes. Thus, to make the approach more suitable for aligning snow profiles and assessing their similarity, we adapt the grain type similarity matrix of Lehning et al. (2001) for each of the two tasks separately (Table 1A, B). The matrices contain values between $[0, 1]$ that represent the similarity between two grain types g_1 and g_2 . Values > 0.5 indicate similarity, 0.5 implies indifference, and values < 0.5 indicate dissimilarity. The similarity between two grain types can be converted into a distance function $d_g(g_1, g_2)$ by subtracting the similarity from 1 (i.e., the similarity between identical grain types is 1, their distance is 0) to make it comparable to the other distance functions for hardness and layer date. Table 1A and B are modified from the grain type similarity matrix of Lehning et al. (2001) in the following ways (i.e., gray cells):

1. ~~Practical experience with SNOWPACK shows that buried SH layers are often labeled as DH layers. Hence, the original similarity matrix of Lehning et al. (2001) is unable to recognize the similarity of a buried SH layer in a human observed profile and the equivalent layer in a simulated profile. To address this issue, we raised the similarities between these SH and DH layers are formed by very different processes—by the deposition of hoar onto the snow surface versus~~
5 ~~by kinetic growth of crystals within the snowpack. Consequently, Lehning et al. (2001) evaluated the similarity of the two grain types from 0 to 0.9 for both tasks (Table 1A, B), despite the differences in their formation process. The increased similarity is also in line with the fact that as completely dissimilar. However,~~ both SH and DH represent hazardous weak layers and are of comparable importance in avalanche hazard assessments ~~-(Schweizer and Jamieson, 2001).~~
10 ~~Furthermore, practical experience with the current version of SNOWPACK shows that SH layers are often converted to DH layers once buried. To account for both of these aspects, we raised their similarity from 0 to 0.9 for both tasks (Table 1A, B).~~

Following the same line of logic, we also raised the similarity between SH and FC, another common weak layer grain type. However, to acknowledge the enhanced seriousness of buried SH layers while simultaneously facilitate the weak layer matching, the similarity between SH and FC is higher when aligning snow profiles (Table 1A) than when assessing their similarities (Table 1B)

2. Since human profiles sometimes lack grain type information for certain layers, an automated alignment algorithm needs to be able to cope with missing data in a meaningful way. Since it is more common for bulk layers to have missing grain type information, it is more desirable to match unknown grain types to bulk layers than weak layers. We therefore expanded the similarity matrix of Lehning et al. (2001) with an additional column for unknown grain types and filled it with values that are centered around indifference (i.e., 0.5) (Table 1B), but with slight preference towards bulk layers (Table 1A).

3. While we acknowledge the physical similarity between DH, FC, and FCxr, we want to emphasize their slightly different implications for the hazard conditions. We therefore classified DH layers—often representing buried SH layers, or layers with large FC grains in simulated profiles—as first order weak layers, whereas FC and FCxr were classified as second and third order weak layers, respectively. This hierarchy is implemented in Table 1B: similarities DH–FC and DH–FCxr are equal to and slightly lower than 0.5, respectively, and thus represent indifference or a slight mis-match; the similarity FC–FCxr is slightly greater than 0.5 and thus represents a weak match. These modifications lead to a pronounced distinction between the weak layer grains SH and DH from the less distinct weak layer grains FC and FCxr.

2.1.2 Distance function for layer hardness

30 The second important layer characteristic to consider is hardness, which characterizes the resistance of snow to penetration. In operational field observations the layer hardness is expressed on an ordinal hand hardness scale (Fierz et al., 2009). Hardness observations are taken by gently pushing different objects into snow layers from a snow pit wall. The hardness of a layer is expressed by the largest object that can be pushed into the snow layer with a consistent force of approx. 10–15 N. The ordinal

levels of the hand hardness index are fist (F), four fingers (4F), one finger (1F), pencil (P), knife blade (K), and ice (I). Sub-classifications like $4F+$, $4F-1F$, $1F-$ are possible and refer to *just harder than 4F*, *between 4F and 1F*, and *just softer than 1F*, respectively.

5 A translation of the ordinal index into a numerical scale is straightforward by assuming that *fist* equals to a numerical value of 0, *ice* equals to 6, and the ordinal levels are equidistant (Schweizer and Jamieson, 2007). The normalized distance d_h can then be written as $d_h(h_1, h_2) = \frac{|h_1 - h_2|}{5}$, where h_1 and h_2 represent numerical translations of two hand hardness values and the normalization factor of 5 refers to the largest distance possible (F–I). Hence, only layers with hardness difference of fist to ice are considered completely dissimilar, while all other hardness combinations exhibit some degree of similarity.

2.1.3 Distance function for layer date

10 Snow layers are commonly labeled with either their deposition date (i.e., the date when a specific layer was formed) or their burial date (i.e., the date when a specific layer was buried). While snowpack models predominantly work with deposition dates, practitioners mainly use burial dates. One reason for practitioners' preference for burial dates is the fact that layers can form over several days, which makes assigning a deposition date challenging. However, it is straightforward to derive the burial date of a simulated snowpack layer based on the deposition date of the overlying layer. Hence, layer dates of simulated snow
15 profiles can easily be compared with layer dates recorded by practitioners.

The distance function d_t between two dates t_1 and t_2 becomes trivial as soon as the dates represent the same type of date (deposition or burial) and are converted into Julian dates: $d_t(t_1, t_2) = \frac{|t_1 - t_2|}{c}$. In this case, the normalization factor c determines the time lag when the dates are considered to be completely dissimilar. A normalization factor of $c = 5$, for example, means that date differences from zero to four days become increasingly dissimilar (i.e., $d_t \in [0, 1]$), whereas date differences equal
20 to or greater than five are considered completely dissimilar (i.e., $d_t \geq 1$; d_t typically does not exceed 2, but the exact limit depends on c and the length of the season or the size of the DTW warping window constraint). A normalization factor of $c = 1$ means that only identical dates are considered to have any similarity. Hence, c can be used to account for small deviations in reported burial dates as well as in short time lags of weather patterns across geographic regions. In cases when layer dates are not available for one or both layers, the distance d_t defaults to *indifference* (i.e. $d_t = 0.5$). This makes it possible to label only
25 important layers with their date.

2.2 Aligning snow profiles with Dynamic Time Warping

In this section, we present a numerical algorithm that addresses the challenge of matching corresponding layers in snow profiles. Our method is based on Dynamic Time Warping (DTW), a longstanding algorithm, which was originally designed for speech recognition in the 1970s (Sakoe and Chiba, 1970; Sakoe, 1971; Sakoe and Chiba, 1978). Soon thereafter it was
30 adopted in time series analyses (e.g., Berndt and Clifford, 1994), and it remains a state-of-the-art component in data mining methods such as clustering and classification (e.g., Keogh and Ratanamahatana, 2005; Petitjean et al., 2011; Wang et al., 2013; Paparrizos and Gravano, 2015). Our discussion of DTW starts with a general background section, which is followed by four sections that explain our application of DTW to snow profiles in more details. This includes snow profile pre-processing steps




Figure 1. Illustrative example of the DTW alignment of two sine wave sequences. (a) each element of the first sequence is mapped to one or many elements of the second sequence; the corresponding sequence elements can be found by (b) the (warping) path of least resistance through a local cost matrix that stores the the distances between every possible pair of sequence elements.

as well as inputs to the DTW algorithm. We then discuss suitable DTW parameter choices for snow profile alignments, and make recommendations on how to use the alignment algorithm.

2.2.1 Background on Dynamic Time Warping

The following brief summary of DTW is based on Sakoe and Chiba (1978), Rabiner and Juang (1993), Keogh and Ratanamahatana (2005), and Giorgino (2009).

DTW is an *elastic* distance measure for time series or, more generally, sequences that calculates the dissimilarity between two sequences while allowing for distortions in "time". The distortions are accommodated by mapping each element of one sequence to one or many elements of the other sequence (Fig. 1a). Once mapped, the dissimilarities between each of the matched sequence elements can be computed and combined into one single distance value that quantifies the dissimilarity between the two sequences.

To link corresponding sequence elements, a local cost matrix \mathbf{D} is calculated that stores the distances between every possible pair of sequence elements. Hence, the dimensions of \mathbf{D} are the lengths of the two sequences. Then, the best alignment of the two sequences is represented by the path of least resistance through \mathbf{D} , which is referred to as the optimal warping path \mathbf{P} (Fig. 1b). To ensure meaningful matching between sequence elements, the warping path is subject to the following constraints:

- 15 **Monotonicity and continuity** Subsequent elements of the warping path are contiguous in a sense, that they are horizontally, vertically or diagonally adjacent cells of the local cost matrix \mathbf{D} , while no "~~stopping~~ or ~~going back~~ going back (in time)" is allowed.

Warping window It is common practice to restrict the search for the optimal warping path to the bounds of a warping window around the main diagonal of the local cost matrix \mathbf{D} . That main diagonal represents the lock-step alignment of the two sequences, where the i th element of the first sequence is mapped onto the i th element of the second sequence. Thus, the farther the warping path deviates from the main diagonal, the more extreme the warping gets, and sequence elements that are farther apart from each other get matched. To prevent excessive warping, the warping path is often constrained to a warping window. The shape and size of the optimal warping window changes with the domain and the target data set. A slanted band of constant width (*Sakoe–Chiba band*) or a parallelogram between the sequences’ start and end points (*Itakura parallelogram*) are popular choices. For detailed visualizations see Ratanamahatana and Keogh (2004). DTW with a constrained warping window is commonly referred to as *constrained* Dynamic Time Warping (cDTW).

Local slope constraint While the warping window constrains the envelope of the warping path *globally*, the so-called local slope constraint of the warping path ensures reasonable warping *locally*. More specifically, the local slope constraint controls how many subsequent elements of one sequence can be mapped onto one element of the other sequence. That is an important control, because it regulates how much stretching and compressing of individual sequence elements is allowed. For time series, that means stretching and compressing with respect to time; in the case of snow profiles, it refers to the stretching and compressing of snow layer thicknesses.

Many different local slope constraints have been suggested in the literature. ~~In~~ [As an illustration, in](#) Fig. 1, we use a non-restrictive local slope constraint, which allows arbitrarily many elements to be mapped onto one corresponding element. In this case, the first element of sequence 2 is mapped onto almost 20 elements of sequence 1 (Fig. 1a), which requires a vertical start slope of the warping path (Fig. 1b). [Find an explicit illustration of the non-restrictive local slope constraint in Fig. B1a.](#)

Boundary conditions For a *global* alignment, both the sequences’ start and end points need to be elements of the warping path (Fig. 1b). *Partial* alignments can be computed by relaxing one or both of these constraints. This results in three different options: alignments where the start but not the end points are matched (*open-end* alignment), alignments where the end but not the start points are matched (*open-begin* alignment), or alignments where subsequences of the two sequences that neither include the start or end points are matched. Further details on partial DTW matching can be found in Appendix A and in the review by Tormene et al. (2009).

While there are many potential warping paths through the local cost matrix \mathbf{D} that satisfy these constraints, the objective is to find the *optimal* warping path \mathbf{P} that accumulates the least cost while stepping through \mathbf{D} . This is an optimization problem that can be solved by dynamic programming. We refer the interested reader to Appendix A for more details.

30 2.2.2 Preprocessing of snow profiles: uniform scaling and resampling

The specific nature of snow profile data requires some preprocessing before DTW can be applied in a meaningful way. Variabilities in snowpack structures can be divided into systematic differences due to systematic variations in the meteorological

forcing (e.g., location: a wind scoured ridgeline next to a wind loaded slope; elevation: increase of snowfall amounts with elevation), and random differences due to natural variations in the meteorological forcing (e.g., peculiar patterns of individual storms; small-scale variations) (Schweizer et al., 2007). The former can result in substantial differences in the snow profiles due to accumulation of different forcings over time. (See Sect. S1 in the supplementary material for more details and visualizations of idealized stratigraphic snowpack variability.) While the DTW algorithm is well suited to deal with random differences, it was not designed to cope with systematic differences. Since systematic differences are common in snow profiles, it is necessary to preprocess them for a meaningful application of DTW. Fu et al. (2007) suggest Uniform Scaling—another optimization technique—which minimizes systematic differences by determining an optimal, global scaling factor. For efficiency reasons, we simply scale the snow profiles to identical snow heights instead. We thereby assume that the offset corresponds to approximately the magnitude of the systematic differences, and that the rescaled profiles are characterized predominantly by random differences that can be handled by DTW. A tentative evaluation of this assumption can be found in the supplementary material (Sect. S2.2).

Once the profiles have been rescaled, each of the two profiles consists of a series of discrete layers along an irregular height grid. To equalize the two different height grids, we resample the profiles onto a regular grid with a constant sampling rate, which represents the final resolution for the alignment procedure. ~~A typical sampling rate might be of the order of~~ While our algorithm allows users to flexibly set the sampling rate, a resolution of about half a centimeter ~~to capture hazardous, thin weak layers ensures that typically thin, hazardous weak layers are being captured. Hence, snow profiles in the presented examples have been resampled to 0.5~~ cm. To preserve the discrete layer character of each profile, we do *not* interpolate between the grid points during the resampling process.

2.2.3 Computing a weighted local cost matrix from multiple layer characteristics

Section 2.2.1 introduced how DTW exploits the local cost matrix \mathbf{D} to find the best alignment of two sequences. We will now show how to compute this local cost matrix for snow profiles. Therefore, the following presentation focuses on the layer characteristics of categorical grain type, ordinal layer hardness, and numerical layer date, which are the layer characteristics most commonly recorded by practitioners. Note, however, that the layer date contribution can be omitted if the information is not available, and our algorithm can easily be expanded to include other layer properties.

First, we combine the distances of the individual layer characteristics (defined in Sect. 2.1) into one scalar distance by weighted averaging. The resulting scalar distance fills one cell of the local cost matrix \mathbf{D} , and thus controls how the alignment algorithm matches the corresponding layers in the snow profiles.

Second, we use a weighting scheme for preferential layer matching (Fig. 2) to ensure that the algorithm prioritizes the alignment of snowpack features that are relevant for avalanche hazard assessment when calculating the scalar distance. The intent of the weighting scheme is to create anchor points for key layers by artificially introducing penalties for non-key layers. This is especially advantageous when no date information is available. Similar to our approach with the distance function for grain types, we focus on avalanche hazard assessment priorities and try to emulate a human alignment approach: First priority should be given to the alignment of first-order persistent weak layers (SH and DH), followed by crusts. Second priority




Figure 2. Visualization of a local cost matrix \mathbf{D} , which stores the distance between individual layers of two snow profiles Q and R; Q and R contain the layer characteristics grain type q^g and r^g , as well as hardness q^h and r^h . \mathbf{D} is only filled with values around its diagonal, the rest is clipped by a warping window and a local slope constraint; the back line represents the optimal warping path \mathbf{P} , which accumulates the least cost while stepping through \mathbf{D} , and which defines the alignment of the two profiles; the preferential layer matching implementation becomes apparent in the yellow and orange cells, which represent smaller than average distances and thus will be matched more easily; see text for further explanation.

should be given to the alignment of faceted grain types with first-order persistent weak layers. This cascade of priorities is expressed numerically by the relative differences of the weighting coefficients ν presented in Table 2. We determined these values experimentally by testing numerous snow profile alignments and found that they yield the wanted result without any unwanted side effects.

- 5 To compute \mathbf{D} from two generic snow profiles, we introduce the following notation. Q and R are two rescaled and resampled snow profiles with I number of layers—typically on the order of hundreds of layers, depending on the sampling rate and the time of the season. The indices $i = 1, \dots, I$ and $j = 1, \dots, I$ refer to these layers. Each layer of the snow profiles Q and R contains

Table 2. Weighting scheme for preferential layer matching; the preferential layer matching coefficient $\nu(g_1, g_2)$ depends on the combination of two grain types g_1 and g_2 ; the darker the shading, the more preferably the grain type combinations get matched. Values in the upper triangle are symmetric to values in the lower triangle.

	PP	DF	RG	FC	DH	SH	MF	FCxr	MFcr
PP	5								
DF	5	5							
RG	5	5	5						
FC	5	5	5	5					
DH	5	5	5	4.5	0				
SH	5	5	5	4.5	0	0			
MF	5	5	5	5	5	5	5		
FCxr	5	5	5	5	5	5	5	5	
MFcr	5	5	5	5	5	5	5	5	2.5

Table 3. An illustrative example of a resampled snow profile Q that contains information about the vertical position of the profile layers (at their top interfaces), the grain types q^g , the hardnesses q^h , and the (burial) dates q^t .

Height (cm)	i	q^g	q^h	q^t
0.5	1	DH	1.0	2018-12-15
1.0	2	DH	1.0	2018-12-15
		\vdots		
119.5	239	DF	2.25	—
120.0	240	PP	1.5	—

information about the grain type, the hardness, the burial date, and the vertical position of the layer in the profile (i.e., height or depth). Those characteristics are denoted by q^g, r^g for grain type; q^h, r^h for hardness; and q^t, r^t for date (see Table 3 for an example). Each element D_{ij} of the local cost matrix \mathbf{D} can then be written as

$$D_{ij} = w_g d_g(q_i^g, r_j^g) + w_h d_h(q_i^h, r_j^h) + w_t d_t(q_i^t, r_j^t) + \nu(q_i^g, r_j^g), \quad (1)$$

- 5 where $w_g, w_h,$ and w_t are averaging weights that sum up to 1 ($w_g + w_h + w_t = 1$). Those weights need to be estimated (see supplementary material, Sect. S2.1), but specific values are recommended in Sect. 2.2.5. Since d_g and d_h range within $[0, 1]$, d_t typically within $[0, 2]$, and ν within $[0, 5]$, the distance D_{ij} typically ranges within $[0, 7]$, where a distance of zero ($D_{ij} = 0$) refers to the two layers being identical. Note that—in Equ. (1)— d_g is calculated based on the similarity matrix in Table 1A, which is geared towards snow profile alignments. Figure 2 visualizes a local cost matrix derived from two generic snow profiles.

2.2.4 Obtaining the optimal alignment of the snow profiles

After calculating the local cost matrix \mathbf{D} , there are several constraints on the warping path that need to be specified to tailor DTW to snow profile alignments; those constraints are the warping window, the local slope constraint, and the boundary conditions.

- 5 **Warping window** We use a slanted band of constant width around the main diagonal of \mathbf{D} to constrain the warping path. This so-called *Sakoe-Chiba* band is quantified by the window size ε . See Sect. 2.2.5 for a recommendation on which value of ε to use for snow profile alignments.

Local slope constraint We require the local slope constraint to prevent excessive stretching or compressing of the snow layers in either of the two profiles. The *symmetric Sakoe-Chiba local slope constraint (Sakoe and Chiba, 1978) does constraints (Sakoe and Chiba, 1978) do* exactly that: ~~it limits~~ they limit the amount of stretching or compressing to a specific factor, which is identical for both profiles. We chose a factor that limits the amount of stretching to double the layer thickness, and the compression to half the layer thickness. On the technical level, that means that while stepping through \mathbf{D} , a horizontal or vertical step is only allowed if following a diagonal step (Fig. A1), ~~B1~~. The specific local slope constraint we refer to has been termed by Sakoe and Chiba (1978) as symmetric, $P=1$.

15 Note that the local slope constraint results in a funnel-shaped restriction of the warping window close to the starting point of the alignment, which is more restrictive than the slanted band window (Fig. 2). See Appendix A for a more detailed explanation.

Boundary conditions In cases when defining features of the snowpack at the very bottom or top only exist in one of the two profiles (e.g., lack of an early season snowfall event or missing of the most recent storm at one of the two profile locations), partial snow profile alignments are more suitable than global alignments. In those situations the alignment benefits from relaxing the boundary conditions to accommodate those partial alignments (Fig. 2 and 3). We therefore implement symmetric open-end alignments, where an *entire* profile is mapped onto the other profile with the start points matched but the end points not.

25 Since *open-begin* alignments cannot be calculated with our chosen local slope constraint (Tormene et al., 2009), we developed a workaround by aligning snow profiles both *bottom-up* and *top-down*. The top-down alignment can be calculated straightforwardly by ~~mirroring \mathbf{D} about its anti-diagonal,~~ reversing both profiles and re-running the DTW algorithm ~~on the mirrored matrix~~. However, since the DTW distance (Sect. 2.2.1, Appendix A) cannot be used to effectively identify the better alignment of the two, we use our independent, more nuanced similarity measure of snow profiles (addressed in Sect. 2.3) to assess the quality of the alignment.

30 Finding the optimal warping path P implies matching the corresponding layers between the two profiles Q and R . That in turn allows *warping* one profile onto the other one by optimally stretching and compressing its individual layer thicknesses, so that the *warped profile* is optimally aligned with the other profile. For example, warping profile Q onto profile R produces the




Figure 3. The open-begin alignment of two snow profiles Q and R: the line segments match the corresponding layers between Q and R. Adjusting the layer thicknesses of profile Q yields the warped profile Q_w , which is optimally aligned to profile R.

warped profile Q_w , which contains the same layer sequences as profile Q, but the adjusted layer thicknesses have aligned Q_w to the corresponding layers of R (Fig. 3). The optimally aligned profiles R and Q_w can now be used as input to an independent similarity measure.

2.2.5 Application cases and usage recommendations

- 5 There are two main application cases for the snow profile alignment algorithm: (i) aligning snow profiles from different locations (or sources) but at the same points in time, and (ii) aligning snow profiles from the same location at different points in time.

In the first case, we recommend to use open-end alignments when the optimal alignment of *all individual* layers is required—e.g., in applications that compare individual layers, or in clustering applications that search for regions of similar snowpack conditions. We recommend to use global alignments when modeled profiles are evaluated versus human profiles: even though open-end alignments might yield better alignment results, the explicit mismatch of layers at the very bottom or top of the

profiles represents an important element of the model evaluation. Furthermore, mind that open-end alignments increase the scope of the alignment algorithm, which can sometimes lead to surprising layer matches if the algorithm is used unsupervised.

In the second case, aligning profiles from the same location at different points in time (i.e., layer tracking), one of the two profiles typically has more layers and a higher snow depth. Since the smaller profile should be contained in the taller profile in a similar form, the alignment needs to be open-end and bottom-up. Furthermore, it is *not* necessary to rescale the profiles as recommended in Sect. 2.2.2, but only to resample them. In this case, rescaling actually moves corresponding layers farther apart, which would have to be compensated by increasing the window size ε . To increase your control on which parts of the profiles get matched, we recommend that you label some of the key layers with their date information.

In the supplementary material (Sect. S2), we derive the optimal values for the averaging weights w_g and w_h , and the window size ε by simulating experiments. Based on the results of these experiments, we recommend to use the following default settings: $\varepsilon \approx 0.3$ in conjunction with a bottom-up/top-down approach²; a ratio of $w_h/w_g = 1/4$; the value of w_d depends heavily on how similar the meteorological processes are that shape the snowpack at the two locations; e.g. two profiles from the same elevation and the same aspect that are in close proximity can be aligned based on layer date alone; however, two profiles from opposite aspects and different elevation bands, may be aligned predominantly based on grain type and hardness.

2.3 Assessing the similarity of snow profiles

A measure that quantifies the similarity of snow profiles as a whole is best designed independently from a profile alignment or layer matching routine. Such an independent approach allows for *matching of layers based on physical similarity* (i.e., processes like grain formation and metamorphism, or knowledge about snow cover models), whereas *the similarity of the aligned profiles can be assessed based on characteristics relevant for avalanche hazard assessment*. Therefore, we define a similarity measure Φ for generic snow profiles, based on the layer characteristics grain type and hardness, that can be used to numerically compare, evaluate, and group snow profiles.

Let us again consider two snow profiles. Some of their layers have been matched, while others have not. The non-matched layers are located either at the very bottom or very top of *one* of the two profiles. For example, the two profiles could be the profiles R and Q_w from the previous section. Our goal is to compute a scalar number that expresses the similarity between these two profiles on a scale from 0–1. To do so, we start again by computing the (dis)similarities between the corresponding layers analogously to the previous sections. However, in a similarity measure that is geared towards hazard assessment not every layer is equally important. Furthermore, since important weak layers are often much thinner than the bulk layers, they would be dramatically underrepresented in a measure that computes a standard average across all layers. Thus, we bin all layers according to four major grain type classes relevant for avalanche hazard assessments: (1) new snow crystals (PP and DF) that are commonly associated with surface problems, (2) weak layers (SH and DH) and (3) crusts (MFcr) that are typically related to persistent avalanche problems, and (4) all other grain types that represent bulk layers. We calculate separate similarity values for every class (a scalar value between $[0, 1]$), and the overall similarity between the two profiles is the average similarity derived from the classes.

²Note that this value is much higher than typically recommended in the literature (Ratanamahatana and Keogh, 2004).

To calculate the similarity for a grain type class, we first distinguish between matched and non-matched layers. All non-matched layers are treated as *indifferent*, and are therefore assigned a similarity value of 0.5. Such a strategy makes the measure robust against a varying number of non-matched layers. Next, we calculate the similarities of all matched layers. That can be done with the distance functions from Sect. 2.1, which compute the dissimilarity between two layers based on grain type or hardness. Note that in this context, the grain type distance is calculated based on Table 1B to ensure the derived similarity is most useful for avalanche forecasting. The resulting distance is converted into a similarity by subtracting the distance from 1 (i.e., a distance of 0.8 becomes a similarity of 0.2). If the grain type class is new snow crystals or bulk grains, the similarity of a matched layer is computed as the product between the associated similarity of grain type and hardness. The emerging similarity for the entire class is then the average over all (matched and non-matched) layers within.

While the above approach works well for new snow crystals and bulk layers, weak layers and crusts require additional considerations to be integrated in a meaningful way.

1. Given weak layers or crusts, an identical match of the grain type is arguably more important than a hardness evaluation: many weak layers and crusts are thin, and often melt-freeze crust laminates are characterized by an inhomogeneous hardness. These circumstances challenge precise hardness measurements of these layers and make them prone to error. Additionally, crusts play an important role in avalanches not as a weak layer themselves, but as a layer favoring adjacent weak layer growth (Jamieson, 2006). That in turn makes the grain type of a crust much more important than its hardness when evaluating the similarity between two layers. In summary, a hardness evaluation might introduce more error than benefit—especially when comparing human versus modeled profiles. Therefore, we compute the similarity of a matched weak layer or crust as the associated similarity of grain type alone, thereby neglecting hardness information.

2. For weak layers and crusts, it is specifically important where in the profile the layers are located. Consider a snow profile with two DH layers close to the ground and one SH layer buried under new snow. A second, almost identical profile lacks the buried SH layer. While the likelihood of triggering and potential size of avalanches are similar with respect to the two matched DH layers, they are not with respect to the buried SH layer, which is missing in the second profile. Even though the two profiles are visually almost identical, they require different avalanche risk management approaches. If we calculated the similarity for the weak layer class of those two profiles as described above (i.e., as average over all layers), the thin SH layer would be heavily underrepresented among the thicker DH layers. As a consequence, the weak layer class would exhibit a high similarity value. For example, if the two DH layers are each 10 cm thick, and the SH layer 2 cm, the sampling rate were 1 cm, then the similarity for the weak layer class would be $(20 \cdot 1 + 2 \cdot 0)/22 = 0.9$. Given that the two snowpack conditions demand different risk management approaches, such a high similarity is not meaningful. To mitigate those situations, we divide the two profiles into sections of equal thickness and evaluate the similarity of the weak layers and crusts within those sections separately. The number of sections is determined by the maximum number of weak layers (or crusts, respectively) in either of the two profiles. By evaluating similarities of adjacent weak layers or crusts, we introduce a basic weighting scheme for the position of those layers in the profile. It is based on the idea that avalanche likelihood and size, as well as resulting risk management are rather similar for adjacent

weak layers or crusts, but rather different for weak layers or crusts in opposing depths of the profile. In our example, there are three weak layers, hence, the two profiles are divided into three sections of equal thickness. We assume that all weak layers that are in the same section require a similar risk management approach. So, the similarity for each section is the average similarity of the weak layers within, and each section is equally important for the hazard assessment, so the similarity for the weak layer class is the average similarity with respect to the sections. In our example, the lower section contains the two DH layers, the middle section contains no weak layers, and the upper section contains the SH layer in one profile. Hence the similarity for the weak layer class is $(\frac{20 \cdot 1}{20} + \frac{2 \cdot 0}{2}) / 2 = 0.5$. A weak layer similarity of 0.5 evaluates the two snowpack conditions much better than a similarity of 0.9.

In summary, the resulting similarity measure Φ between two snow profiles expresses the similarity between these two profiles in a scalar value within $[0, 1]$. A similarity of 1 corresponds to the two profiles being identical. Note that the measure is symmetric to the two profiles, so that the first profile is as similar to the second profile as vice versa. By treating non-matched layers as indifferent the measure is able to cope with varying numbers of missing layers at the bottom or top of the profiles, which is important for assessing the similarity of snow profiles from different times and/or locations. To make the similarity measure useful for avalanche forecasting, it weighs hazardous thin layers, crusts, as well as storm snow layers more heavily than bulk layers, and it considers the relative depth of those layers.

3 Aggregation and clustering applications—a practical valuation

Sections 2.2 and 2.3 detail how to match layers between snow profiles, and how to assess the similarity of the aligned profiles for applications in avalanche hazard assessment. Both of these steps are fundamental prerequisites to automate forecaster tasks, such as grouping similar profiles and finding the representative profile of a group. In the data sciences, these tasks are called data clustering and data aggregation. In this section, we demonstrate how to apply simple data clustering and data aggregation methods to snow profiles based on their prior alignment and similarity assessment. We use these application examples as face validation of our methods.

3.1 Clustering of snow profiles

Avalanche professionals need to group snow profiles to understand how snowpack conditions and avalanche hazard vary across space. The snow profile alignment algorithm and similarity measure described in the previous section enable the automation of this task by providing the necessary quantitative link to the well-established field of numerical clustering methods (e.g., James et al., 2013; Sarda-Espinosa, 2019).

To showcase the clustering of snow profiles based on our similarity measure, we are using a set of twelve snow profiles that exhibit both pronounced and subtle differences in their snowpack features (Fig. 4). The alignment and similarity assessment of the snow profiles is based on grain type and hardness information, even though they are visualized in Fig. 4 solely by their grain type sequences. After computing a total $12 \cdot 11 = 132$ profile alignments and similarity assessments, the clustering is carried out by an agglomerative hierarchical clustering algorithm that iteratively fuses individual profiles to clusters based on the




Figure 4. Hierarchical clustering of twelve snow profiles based on prior snow profile alignment and similarity assessment. The colors represent the grain types of distinct layers, the bold black lines separate the four most distinct clusters.

similarity of the profiles³. The resulting cluster hierarchy is best analyzed from the top to the bottom, allowing to separate the set of twelve profiles into up to twelve clusters. The higher a split occurs in the hierarchy, the more distinct the corresponding clusters are to each other.

The first, most distinct split separates the heavily faceted profiles 1–4 from the remaining profiles. The remaining profiles 5 are generally quite similar, except for some subtle, but important features that still require different risk management strategies: Profiles 5–7 show weak layers both in the middle of the snowpack as well as below the new snow (i.e., second split); profiles 8 and 9 have a weak layer sandwiched between two crusts at the bottom of the snowpack (i.e., third split), and profiles 10–12 either have a weak layer only in the middle of the snowpack or none at all. Additionally to examining those most distinct, four clusters, the similarities within the clusters can be investigated further. For example, profile 1 is the most dissimilar profile 10 within the first cluster, being the only profile with a crust below the new snow and the only profile with a pronounced weak layer in mid snow height. Or profile 10 is the outsider in cluster 4, having no weak layer at all.

The relationships among the profiles as established by the alignment algorithm and similarity measure yield a sound clustering result that looks similar to how a human avalanche forecaster would group the profiles according to different strategies on how to manage snowpack conditions and the related avalanche hazard. This example demonstrates that our approach can 15 differentiate between both very different as well as subtly different snowpack conditions.

³We use complete linkage to fuse individual profiles to clusters (James et al., 2013).

3.2 Finding a representative snow profile: The medoid

Avalanche forecasters often draw a representative snow profile that summarizes the most important snowpack features within a group of profiles. In the data mining community, this type of generalization is typically called the *average sequence*, or *aggregate*, and the most sophisticated methods for computing that aggregate are closely tied to the used alignment algorithm and similarity measure (e.g., Petitjean et al., 2011; Paparrizos and Gravano, 2015). In the following, we use the simple approach of identifying the one profile within the group that is most similar to all other profiles, called the *medoid* profile. Visually, the medoid profile can be thought of as the member of the group that is closest to the geometric center of the group. Mathematically, that means that the medoid profile minimizes the accumulated distances to all other profiles. To identify the medoid, we compute the accumulated distances to all other profiles for every profile. As distance δ between two profiles Q and R we use the similarity measure Φ after converting it to a distance by subtracting it from 1 (cf., Sect. 2.3). We apply the similarity measure to both profile pairs (Q, R_w) and (Q_w , R) to account for missing layers. Hence,

$$\delta(Q, R) = \max\{1 - \Phi(Q, R_w), 1 - \Phi(Q_w, R)\}. \quad (2)$$

The pairwise distances between the profiles of the group can be translated into a configuration plot, which gathers similar profiles close to each other, and dissimilar ones further away from each other⁴ (Fig. 5). The medoid profile, being most similar to all other profiles, is the member of the group that represents the group the best. We use the same set of twelve profiles as in Sect. 3.1 to demonstrate the profile aggregation. As the snowpack conditions within the set are too different to meaningfully represent them by one representative profile (Fig. 5a), we can combine clustering and aggregating to draw the representative profiles of the most distinct clusters within the set (Fig. 5b—the three clusters consist of profiles 1–4, 5–7, and 8–12, as depicted in Fig. 4).

While our proof of concept demonstrates a sound and reliable workflow, using the medoid profile may be computationally too expensive to deal efficiently with data volumes beyond the order of tens to hundreds of profiles on an operational basis. However, Paparrizos and Gravano (2015) show that the medoid approach performs just better than any other sequence aggregation method. *Dynamic Time Warping Barycenter Averaging* (Petitjean et al., 2011) might be an alternative aggregating method that could be evaluated in future studies.

4 Discussion and conclusions

The snow profile alignment algorithm and the similarity measure presented in this paper aim to address two of the main reasons that have limited the adoption of snowpack models to support avalanche warning services and practitioners. First, the methods provide the foundation for numerical grouping and summarizing of snow profile data, and thus can help to make snowpack model output more accessible by addressing any avalanche operations' fundamental questions *Where in the terrain do we find which conditions?*. Second, our methods have the potential to make snowpack models more relevant to avalanche forecasters by providing a means for model evaluation against human observations.

⁴We use an ordinal multi-dimensional scaling approach to create the configuration plot for the group of snow profiles (e.g., Mair, 2018).




Figure 5. Configuration plots of a group of twelve snow profiles, where similar profiles are gathered close to each other; (a) shows the geometric center of the whole group, which identifies the one, most representative profile; (b) shows the three most distinct subsets of the whole group and highlights their associated representative profiles, cf. Fig. 4.

Since we are the first to present an alignment algorithm and similarity measure for generic snow profiles, there is no objective approach to evaluate the methods against existing benchmark methods. Unlike in Building on the well-established and longstanding concept of Dynamic Time Warping (DTW), we developed a snow profile alignment algorithm that combines multiple layer characteristics of categorical, numerical, and ordinal format into a weighted metric and feeds into existing DTW algorithms such as the open-source R package *dtw*. Moreover, we reviewed and derived useful DTW configurations and hyper-parameter settings for snow profile applications. Since these applications rely on operationally available profile observations that typically focus on information relevant for current avalanche conditions only, our approach is able to handle missing data and take advantage of select layer date tags. To maximize the layer matching performance for profiles with limited details, we implemented a scheme for preferential layer matching based on domain knowledge. In parallel, Viallon-Galinier et al. (2020) extended the layer matching algorithm from Hagenmuller and Pilloix (2016) to conduct a detailed, process-related evaluation of the snowpack model Crocus based on high-quality snow profile observations with a large variety of observed variables that are sampled at specific study sites at regular intervals. Since their goal is the correction of deviating model states with a direct insertion assimilation scheme based on point scale simulations and observations, their evaluation targets not only each individual layer, but also each individual layer characteristic separately. To address operational avalanche forecasting needs, we additionally developed a similarity measure that focuses on avalanche forecasting specific considerations where certain layers are considered more important. Moreover, combining information from individual layers and their characteristics into a scalar measure allows for clustering and aggregating of sets of profiles to characterize and evaluate the regional scale avalanche hazard conditions.

In the data mining community, ~~where distance~~ similarity (or distance) measures are evaluated through classification applications (Wang et al., 2013). ~~Since~~ snow profile data sets that represent ground truth of alignments or groupings do not (yet) exist. ~~Consequently~~, the evaluation of our methods needs to rely on expert judgement or application valuation. During the development of our algorithm, we therefore manually evaluated the alignment of many profile pairs, a few of which are shown in the supplementary material (Sect. S3) to demonstrate its behavior. Furthermore, we evaluated the interplay of the alignment algorithm and similarity measure through clustering and aggregation applications. Those applications involve many individual profile assessments and therefore represent a meaningful valuation approach, which shows that the alignment algorithm and similarity measure are capable of distinguishing between subtle differences in the snow stratigraphy and yield a sound grouping that could have been carried out by a human avalanche professional.

Although our methods have been designed to accommodate a wide range of requirements and to cope with a variety of scenarios, the following limitations should be considered. First, while the layer matching algorithm can easily be applied to large data sets in an unsupervised manner, not all scenarios within such a data set might be best served with the same parameter settings. Getting meaningful results in a highly diverse data set requires the algorithm to be less constrained. However, this can also result in unrealistic alignments. Second, when alignments are based on grain type and hardness alone, the matching of layers can sometimes be ambiguous—even for human experts. In those cases, labeling a few key layers with their burial date can greatly improve the alignment accuracy with little extra effort, especially as the labeling of weak layers is already established practise in North America (Canadian Avalanche Association, 2016). Note that our algorithm can easily be expanded to include

other layer properties, especially if they are of ordinal or numeric data types such as e.g., grain size or specific surface area. ~~And third~~ Third, it is important to recognize that any numerical approach that condenses the complexity of these similarity assessments to a one dimensional scale between [0, 1] is unable to capture the full expertise and situational flexibility of human forecasters. However, it is a critical prerequisite for algorithmically grouping profiles and establishing ranks among them. As such our similarity measure offers a consistent evaluation of snow profiles that is based on the most commonly available layer characteristics. And lastly, the similarity measure, and consequently the clustering and aggregating applications are purely based on the congruency of the snowpack structure. Hence, snow depth is not a driver of our similarity assessment unless it leads to deviations in the snow stratigraphy. If combined with monitoring of the snow depths distribution, a clustering or aggregation application can provide a comprehensive picture of the conditions within a specific forecast area.

10 Since our methods aim to offer direct benefit in helping to overcome the operational challenges of summarizing snowpack model data and evaluating that data, we imagine its integration into operational avalanche forecasting as follows. Traditionally, an avalanche risk management operation, such as a public warning service or a backcountry guiding operation, obtains vital information about the snowpack through manual snow pit observations at select point locations. Simulated snow profiles across a mountain drainage can sample the snowpack conditions similarly to field observations, except with a higher spatiotemporal coverage and independent from external circumstances. Our methods could group the simulated profiles according to similar conditions, potentially uncovering different avalanche problems. Furthermore, that grouping could quantify the prevalence of those conditions, and the conditions could be linked to their specific location, elevation, and aspect. Then, the different conditions could be summarized by their representative profile to present the data in a familiar way. The human forecaster can use that simulated data as additional data source complementing the field observations, or deploy targeted field observations to verify the model output. Through a continuous evaluation of the model output against human observations or human assessments (e.g., synthesized snow profiles), the momentary validity of the simulations could potentially be extrapolated into data sparse regions. More generally, a continuous evaluation of operational snowpack simulations provides an opportunity to better understand strengths and weaknesses of the involved model chain for its application in avalanche forecasting. Through this line of research, we hope that snowpack models will be further incorporated into operational avalanche hazard assessment routines, so that avalanche forecasters can begin to build understanding in how to interpret and when to trust snowpack simulations.

Code and data availability. The snow profile alignment algorithm and similarity measure, as well as the data and the according analysis scripts to reproduce the results presented in this paper are available from an Open Science Framework registration. All code and data can be accessed through <https://doi.org/10.17605/OSF.IO/9V8AD>. Our alignment algorithm builds upon the R package *dtw* (v1.21-3) (by Giorgino, 2009), which belongs to “the most complete, freely-available (GPL) implementation of Dynamic Time Warping-type (DTW) algorithms up to date”.




Figure A1. Sketch of a cost matrix \mathbf{G} that stores the smallest accumulated cost of visiting a matrix element G_{ij} . In an open-end alignment, the optimal warping path \mathbf{P} starts at G_{11} . The steps of the warping path are governed by the chosen local slope constraint (black arrows), such that only the grey matrix cells can be visited. The warping path ends in the last row or column of the matrix where the accumulated cost is smallest (i.e. at $G_{65} = 6$). See text for more details.

Appendix A: An exemplary solution to the DTW optimization problem

In this section we explain the concept of solving the DTW optimization problem and thereby finding the optimal warping path \mathbf{P} through the local cost matrix \mathbf{D} . We do that exemplarily by applying some of the same constraints that we also use in the snow profile alignment algorithm; most notably, those are the Sakoe-Chiba local slope constraint ([symmetric, \$P=1\$, Fig. B1b](#)) and (symmetric) open-end boundary conditions (cf., Sect. 2.2.4).

As mentioned in Sect. 2.2.1, the DTW optimization problem can be solved recursively with the aid of dynamic programming. Imagine a local cost matrix \mathbf{D} (i -by- j) with individual elements D_{ij} . Another, yet empty matrix \mathbf{G} —the *accumulated* cost matrix—has the same dimension as \mathbf{D} . From the boundary conditions, we know that the first items of the two sequences need to be matched. Thus the optimal warping path \mathbf{P} starts at G_{11} , which holds the same value as D_{11} . From the local slope constraint, we know that a horizontal or vertical step is only allowed if following a diagonal step. Therefore, as we are about to do our first step, we have to do a diagonal one to G_{22} . G_{22} is the second element of the optimal warping path \mathbf{P} , and its value can be calculated by the accumulated cost one step before (i.e., G_{11}) plus twice the *local* cost of the current step (i.e., D_{22}); hence $G_{22} = G_{11} + 2D_{22}$ (where the weighting factor 2 ~~indicates that both indices i and j have been incremented~~).

As represents a slope weight, cf. Appendix B). Since we just did a diagonal step, we are now allowed to step up vertically, diagonally, or horizontally to fields G_{23} , G_{33} , or G_{32} , respectively.

More generally, any element G_{ij} can be calculated by the recursion

$$G_{ij} = \min \begin{bmatrix} G_{i-1j-2} + 2D_{ij-1} + D_{ij} \\ G_{i-1j-1} + 2D_{ij} \\ G_{i-2j-1} + 2D_{i-1j} + D_{ij} \end{bmatrix}. \quad (\text{A1})$$

5 Figure A1 sketches that concept. Each individual step of the optimal warping path \mathbf{P} is governed by the local slope constraint, such that only a limited number of matrix elements can be visited by the warping path. Each of those elements of the cost matrix \mathbf{G} in turn stores the *smallest* accumulated cost that is necessary to arrive at that cell. In a symmetric open-end alignment, the final element of the optimal warping path \mathbf{P} is the one element in the last column or row of \mathbf{G} that has accumulated the least cost.

10 If the optimal warping path \mathbf{P} has K elements from $k = 1, \dots, K$, then each element \mathbf{p}_k stores its location in the cost matrix by $\mathbf{p}_k = (p_k^i, p_k^j)$. Consequently, the accumulated cost of the optimal warping path \mathbf{P} can be expressed as $G_{p_K^i p_K^j}$. For example, for the warping path implied by Fig. A1, $\mathbf{P} = \{(1, 1), (2, 2), \dots, (6, 5)\}$ and $G_{p_K^i p_K^j} = G_{65} = 6$. Finally, the DTW distance between two sequences, d_{DTW} , is expressed by the accumulated cost of the optimal warping path \mathbf{P} normalized by the length of the path (expressed as Manhattan distance from the matrix origin, for symmetric recursions), i.e.

$$15 \quad d_{\text{DTW}} = \frac{1}{p_K^i + p_K^j} G_{p_K^i p_K^j}. \quad (\text{A2})$$

In Sect. 2.2.4 we introduced the *warped* snow profile Q_w . With the insight gained from the current section, we can now precisely define the warped profile Q_w . Therefore we adopt the notation introduced in Sect. 2.2.3, and additionally denote the layer height of the snow profiles Q_w and R as \mathbf{q}_w^{Ht} and \mathbf{r}^{Ht} , respectively. Then the warped profile Q_w can be constructed with the indices \mathbf{p}^i and \mathbf{p}^j , which are given by the warping path \mathbf{P} , by

$$20 \quad \mathbf{q}_w^{\text{Ht}} = \mathbf{r}^{\text{Ht}}_{\mathbf{p}^j}, \quad (\text{A3})$$

$$\mathbf{q}_w^{\text{g}} = \mathbf{q}^{\text{g}}_{\mathbf{p}^i}. \quad (\text{A4})$$

The vectors \mathbf{q}_w^h and \mathbf{q}_w^t can be calculated analogously to Equ. (A4).

Appendix B: Exemplary DTW step patterns

DTW step patterns describe the local slope constraint and the slope weights associated with each step of the warping path.

25 While the local slope constraint limits the warping path to physically meaningful steps, the slope weights, which depend on the local indices' increments, ensure that different alignments can be compared. Both step patterns used in this paper use slope weights that are symmetric to the query and reference sequences, but asymmetric slope weights that favor the advance of a certain sequence also exist. (Rabiner and Juang, 1993; Sakoe and Chiba, 1978; Giorgino, 2009).




Figure B1. Two exemplary DTW step patterns that illustrate different, symmetric local slope constraints and their associated slope weights. While (a) shows an unconstrained pattern, (b) shows a pattern that limits the stretching of either sequence to a factor of 2 by enforcing a diagonal step before a horizontal or vertical one.

While the symmetric, unconstrained step pattern allows the warping path to advance diagonally, vertically, or horizontally at any time of the alignment (Fig. B1a, used in the alignment of sine waves shown in Fig. 1), the Sakoe-Chiba symmetric, $P=1$ step pattern enforces a diagonal step preceding each horizontal or vertical step to limit the stretching and compressing of the sequences to a factor of 2 and $1/2$, respectively (Fig. B1b, used for snow profile alignments as in Fig. 2, 3).

- 5 *Author contributions.* All authors conceptualized the research; SH provided the snowpack modeling infrastructure and the snowpack model output; FH derived the methods and implemented both code and simulations; all authors contributed to writing the manuscript; PH acquired the funding.

Competing interests. The authors declare that they have no conflict of interest.

- 10 *Acknowledgements.* FH thanks T. Giorgino for a valuable exchange on open-end DTW, as well as S. Mayer and B. Richter for exchanging research ideas and providing additional snow profile data for methods testing. We thank P. Hagenmuller for an email exchange at the outset of the project. We also thank Fabien Maussion for supervising the review process and Matthieu Lafaysse, as well as an anonymous reviewer for valuable comments to improve this manuscript.

References

- Bartelt, P., Lehning, M., Bartelt, P., Brown, B., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss avalanche warning: Part I: Numerical model, *Cold Regions Science and Technology*, 35, 123–145, [https://doi.org/10.1016/s0165-232x\(02\)00074-5](https://doi.org/10.1016/s0165-232x(02)00074-5), <https://linkinghub.elsevier.com/retrieve/pii/S0165232X02000733>, 2002.
- 5 Bellaire, S. and Jamieson, J. B.: On estimating avalanche danger from simulated snow profiles, in: *Proceedings of the International Snow Science Workshop Grenoble–Chamonix Mont-Blanc*, pp. 7–11, 2013.
- Bellaire, S., Jamieson, J. B., and Fierz, C.: Forcing the snow-cover model SNOWPACK with forecasted weather data, *The Cryosphere*, 5, 1115–1125, 2011.
- Bellaire, S., Jamieson, J. B., and Jamieson, B.: Forecasting the formation of critical snow layers using a coupled snow cover and weather
10 model, *Cold Regions Science and Technology*, 94, 37–44, <https://doi.org/10.1016/j.coldregions.2013.06.007>, <https://www.sciencedirect.com/science/article/pii/S0165232X13000840><http://dx.doi.org/10.1016/j.coldregions.2013.06.007>, 2013.
- Berndt, D. J. and Clifford, J.: Using dynamic time warping to find patterns in time series., in: *KDD workshop*, vol. 10, pp. 359–370, Seattle, WA, 1994.
- Brun, E., Martin, E., Simon, V., Gendre, C., and Coléou, C.: An energy and mass model of snow cover suitable for operational avalanche
15 forecasting, *Journal of Glaciology*, 35, 333–342, <https://doi.org/10.1017/S0022143000009254>, 1989.
- Campbell, C., Conger, S., Gould, B., Haegeli, P., Jamieson, J. B., and Statham, G.: *Technical Aspects of Snow Avalanche Risk Management–Resources and Guidelines for Avalanche Practitioners in Canada*, Revelstoke, BC, Canada, 2016.
- Canadian Avalanche Association: *Observation Guidelines and Recording Standards for Weather, Snowpack, and Avalanches*, Tech. rep., Revelstoke, BC, Canada, 2016.
- 20 Fierz, C.: Field observation and modelling of weak-layer evolution, *Annals of Glaciology*, 26, 7–13, <https://doi.org/10.3189/1998AoG26-1-7-13>, https://www.cambridge.org/core/product/identifier/S0260305500014452/type/journal_article, 1998.
- Fierz, C., Armstrong, R. L., Durand, Y., Etchevers, P., Greene, E., McClung, D. M., Nishimura, K., Satyawali, P., and Sokratov, S. A.: *The International Classification for Seasonal Snow on the Ground*, vol. 5, UNESCO/IHP, <http://www.unesco.org/water/ihp>, 2009.
- Fu, A. W.-C., Keogh, E. J., Lau, L. Y. H., Ratanamahatana, C. A., and Wong, R. C.-W. W.: Scaling and time warping in time series querying,
25 *The VLDB Journal*, 17, 899–921, <https://doi.org/10.1007/s00778-006-0040-z>, 2007.
- Giorgino, T.: Computing and Visualizing Dynamic Time Warping Alignments in R: The dtw Package, *Journal of Statistical Software*, 31, <https://doi.org/10.18637/jss.v031.i07>, 2009.
- Hagenmuller, P. and Pilloix, T.: A New Method for Comparing and Matching Snow Profiles, Application for Profiles Measured by Penetrometers, *Frontiers in Earth Science*, 4, <https://doi.org/10.3389/feart.2016.00052>, 2016.
- 30 Hagenmuller, P., van Herwijnen, A., Pielmeier, C., and Marshall, H.-P.: Evaluation of the snow penetrometer Avatech SP2, *Cold Regions Science and Technology*, 149, 83–94, <https://doi.org/10.1016/j.coldregions.2018.02.006>, 2018a.
- Hagenmuller, P., Viallon, L., Bouchayer, C., Teich, M., Lafaysse, M., and Vionnet, V.: Quantitative Comparison of Snow Profiles, in: *Proceedings of the 2018 international snow science workshop*, Innsbruck, AUT, pp. 876—879, <https://arc.lib.montana.edu/snow-science/item/2668>, 2018b.
- 35 Horton, S. and Jamieson, J. B.: Modelling hazardous surface hoar layers across western Canada with a coupled weather and snow cover model, *Cold Regions Science and Technology*, 128, 22–31, <https://doi.org/10.1016/j.coldregions.2016.05.002>, 2016.

- Horton, S., Bellaire, S., and Jamieson, J. B.: Modelling the formation of surface hoar layers and tracking post-burial changes for avalanche forecasting, *Cold Regions Science and Technology*, 97, 81–89, <https://doi.org/10.1016/j.coldregions.2013.06.012>, <http://dx.doi.org/10.1016/j.coldregions.2013.06.012>, 2014.
- Horton, S., Schirmer, M., and Jamieson, J. B.: Meteorological, elevation, and slope effects on surface hoar formation, *The Cryosphere*, 9, 1523–1533, <https://doi.org/10.5194/tc-9-1523-2015>, 2015.
- James, G., Witten, D., Hastie, T., and Tibshirani, R.: *Statistical Learning*, 103, <https://doi.org/10.1007/978-1-4614-7138-7>, <http://link.springer.com/10.1007/978-1-4614-7138-7>, 2013.
- Jamieson, B.: Formation of refrozen snowpack layers and their role in slab avalanche release, *Reviews of Geophysics*, 44, RG2001, <https://doi.org/10.1029/2005RG000176>, <http://doi.wiley.com/10.1029/2005RG000176>, 2006.
- 10 Keogh, E. J. and Ratanamahatana, C. A.: Exact indexing of dynamic time warping, *Knowledge and Information Systems*, 7, 358–386, <https://doi.org/10.1007/s10115-004-0154-9>, 2005.
- LaChapelle, E. R.: *Avalanche Forecasting—A Modern Synthesis*, in: *International Association of Scientific Hydrology, Publication*, vol. 69, pp. 410—417, 1966.
- LaChapelle, E. R.: The fundamental processes in conventional avalanche forecasting., *Journal of Glaciology*, 26, 75–84, <https://doi.org/10.3189/s0022143000010601>, 1980.
- 15 Lehning, M., Bartelt, P., Brown, B., Russi, T., Stöckli, U., and Zimmerli, M.: SNOWPACK model calculations for avalanche warning based upon a new network of weather and snow stations, in: *Cold Regions Science and Technology*, vol. 30, pp. 145–157, Elsevier, [https://doi.org/10.1016/S0165-232X\(99\)00022-1](https://doi.org/10.1016/S0165-232X(99)00022-1), <https://www.sciencedirect-com.proxy.lib.sfu.ca/science/article/pii/S0165232X99000221>, 1999.
- 20 Lehning, M., Fierz, C., and Lundy, C.: An objective snow profile comparison method and its application to SNOWPACK, *Cold Regions Science and Technology*, 33, 253–261, [https://doi.org/10.1016/s0165-232x\(01\)00044-1](https://doi.org/10.1016/s0165-232x(01)00044-1), 2001.
- Lehning, M., Bartelt, P., Brown, B., and Fierz, C.: A physical SNOWPACK model for the Swiss avalanche warning Part III: Meteorological forcing, thin layer formation and evaluation, *Cold Regions Science and Technology*, 35, 169–184, [https://doi.org/10.1016/S0165-232X\(02\)00072-1](https://doi.org/10.1016/S0165-232X(02)00072-1), 2002a.
- 25 Lehning, M., Bartelt, P., Brown, B., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss avalanche warning Part II. Snow microstructure, *Cold Regions Science and Technology*, 35, 147–167, [https://doi.org/10.1016/S0165-232X\(02\)00073-3](https://doi.org/10.1016/S0165-232X(02)00073-3), 2002b.
- Lehning, M., Fierz, C., Brown, B., and Jamieson, J. B.: Modeling snow instability with the snow-cover model SNOWPACK, *Annals of Glaciology*, 38, 331–338, <https://doi.org/10.3189/172756404781815220>, 2004.
- Mair, P.: *Modern Psychometrics with R, Use R!*, Springer International Publishing, Cham, <https://doi.org/10.1007/978-3-319-93177-7>, <http://link.springer.com/10.1007/978-3-319-93177-7>, 2018.
- 30 McClung, D. M.: The Elements of Applied Avalanche Forecasting, Part II: The Physical Issues and the Rules of Applied Avalanche Forecasting, *Natural Hazards*, 26, 131–146, <https://doi.org/10.1023/a:1015604600361>, 2002.
- McClung, D. M. and Schaerer, P.: *The avalanche handbook*, Mountaineers Books, 2006.
- Monti, F., Schweizer, J., and Fierz, C.: Hardness estimation and weak layer detection in simulated snow stratigraphy, *Cold Regions Science and Technology*, 103, 82–90, <https://doi.org/10.1016/j.coldregions.2014.03.009>, 2014a.
- 35 Monti, F., Schweizer, J., Gaume, J., and Fierz, C.: Deriving snow stability information from simulated snow cover stratigraphy, in: *Proceedings of the 2014 international snow science workshop, Banff, AB*, pp. 465–469, 2014b.

- Morin, S., Fierz, C., Horton, S., Bavay, M., Dumont, M., Hagenmuller, P., Lafaysse, M., Mitterer, C., Monti, F., Olefs, M., Snook, J. S., Techel, F., Van Herwijnen, A., and Vionnet, V.: Application of physical snowpack models in support of operational avalanche hazard forecasting : A status report on current implementations and prospects for the future, *Cold Regions Science and Technology*, 170, 1098–1107, <https://doi.org/10.1016/J.COLDREGIONS.2019.102910>, <https://www.sciencedirect.com/science/article/pii/S0165232X19302071>, 2020.
- Paparrizos, J. and Gravano, L.: k-shape: Efficient and accurate clustering of time series, in: *Proceedings of the 2015 ACM SIGMOD International Conference on Management of Data*, pp. 1855–1870, ACM, 2015.
- Petitjean, F., Ketterlin, A., and Gançarski, P.: A global averaging method for dynamic time warping, with applications to clustering, *Pattern Recognition*, 44, 678–693, <https://doi.org/10.1016/j.patcog.2010.09.013>, 2011.
- Rabiner, L. and Juang, B.-H.: *Fundamentals of speech processing*, Prentice Hall, NJ, USA, 1993.
- Ratanamahatana, C. A. and Keogh, E. J.: Everything you know about dynamic time warping is wrong, in: *Third workshop on mining temporal and sequential data*, vol. 32, Citeseer, 2004.
- Sakoe, H.: Dynamic-programming approach to continuous speech recognition, in: *1971 Proc. the International Congress of Acoustics*, Budapest, 1971.
- Sakoe, H. and Chiba, S.: A similarity evaluation of speech patterns by dynamic programming, in: *Nat. Meeting of Institute of Electronic Communications Engineers of Japan*, p. 136, 1970.
- Sakoe, H. and Chiba, S.: Dynamic programming algorithm optimization for spoken word recognition, *IEEE Transactions on Acoustics, Speech, and Signal Processing*, 26, 43–49, <https://doi.org/10.1109/tassp.1978.1163055>, 1978.
- Sarda-Espinosa, A.: dtwclust: Time Series Clustering Along with Optimizations for the Dynamic Time Warping Distance, <https://cran.r-project.org/package=dtwclust>, 2019.
- Schaller, C. F., Freitag, J., Kipfstuhl, S., Laepple, T., Christian Steen-Larsen, H., and Eisen, O.: A representative density profile of the North Greenland snowpack, *Cryosphere*, 10, 1991–2002, <https://doi.org/10.5194/tc-10-1991-2016>, <https://tc.copernicus.org/articles/10/1991/2016/>, 2016.
- Schirmer, M., Lehning, M., and Schweizer, J.: Statistical forecasting of regional avalanche danger using simulated snow-cover data, *Journal of Glaciology*, 55, 761–768, 2009.
- Schirmer, M., Schweizer, J., and Lehning, M.: Statistical evaluation of local to regional snowpack stability using simulated snow-cover data, *Cold Regions Science and Technology*, 64, 110–118, <https://doi.org/10.1016/j.coldregions.2010.04.012>, <http://dx.doi.org/10.1016/j.coldregions.2010.04.012>, 2010.
- Schweizer, J. and Jamieson, J. B.: Snow cover properties for skier triggering of avalanches, *Cold Regions Science and Technology*, 33, 207–221, [https://doi.org/10.1016/S0165-232X\(01\)00039-8](https://doi.org/10.1016/S0165-232X(01)00039-8), 2001.
- Schweizer, J. and Jamieson, J. B.: A threshold sum approach to stability evaluation of manual snow profiles, *Cold Regions Science and Technology*, 47, 50–59, <https://doi.org/10.1016/j.coldregions.2006.08.011>, 2007.
- Schweizer, J., Bellaire, S., Fierz, C., Lehning, M., and Pielmeier, C.: Evaluating and improving the stability predictions of the snow cover model SNOWPACK, *Cold Regions Science and Technology*, 46, 52–59, <https://doi.org/10.1016/j.coldregions.2006.05.007>, 2006.
- Schweizer, J., Kronholm, K., Jamieson, J. B., and Birkeland, K. W.: Review of spatial variability of snowpack properties and its importance for avalanche formation, *Cold Regions Science and Technology*, 51, 253–272, <https://doi.org/10.1016/j.coldregions.2007.04.009>, 2007.
- Statham, G., Haegeli, P., Greene, E., Birkeland, K. W., Israelson, C., Tremper, B., Stethem, C., McMahon, B., White, B., and Kelly, J.: A conceptual model of avalanche hazard, *Natural Hazards*, 90, 663–691, <https://doi.org/10.1007/s11069-017-3070-5>, 2018.

- Storm, I.: Public Avalanche Forecast Challenges: Canada's Large Data-Sparse Regions, in: Proceedings, 2012 International Snow Science Workshop, Anchorage, Alaska, pp. 908–912, 2012.
- Teich, M., Giunta, A. D., Hagenmuller, P., Bebi, P., Schneebeli, M., and Jenkins, M. J.: Effects of bark beetle attacks on forest snowpack and avalanche formation — Implications for protection forest management, *Forest Ecology and Management*, 438, 186–203, <https://doi.org/10.1016/j.foreco.2019.01.052>, 2019.
- 5 Tormene, P., Giorgino, T., Quaglini, S., and Stefanelli, M.: Matching incomplete time series with dynamic time warping: an algorithm and an application to post-stroke rehabilitation, *Artificial Intelligence in Medicine*, 45, 11–34, <https://doi.org/10.1016/j.artmed.2008.11.007>, 2009.
- Van Peurse, K., Hendrikx, J., Birkeland, K. W., Miller, D., and Gibson, C.: Validation of a coupled weather and snowpack model across western montana, in: Proceedings of the 2016 international snow science workshop, Breckenridge, Montana, Breckenridge, CO, 2016.
- 10 Viallon-Galinier, L., Hagenmuller, P., and Lafaysse, M.: Forcing and evaluating detailed snow cover models with stratigraphy observations, *Cold Regions Science and Technology*, 180, 103–163, <https://doi.org/10.1016/j.coldregions.2020.103163>, 2020.
- Vick, S. G.: Degrees of belief: Subjective probability and engineering judgment, ASCE Publications, 2002.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geoscientific Model Development*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, <https://www.geosci-model-dev.net/5/773/2012/>, 2012.
- 15 Vionnet, V., Dombrowski-Etchevers, I., Lafaysse, M., Quéno, L., Seity, Y., and Bazile, E.: Numerical Weather Forecasts at Kilometer Scale in the French Alps: Evaluation and Application for Snowpack Modeling, *Journal of Hydrometeorology*, 17, 2591–2614, <https://doi.org/10.1175/jhm-d-15-0241.1>, 2016.
- 20 Vionnet, V., Guyomarc'h, G., Lafaysse, M., Naaim-Bouvet, F., Giraud, G., and Deliot, Y.: Operational implementation and evaluation of a blowing snow scheme for avalanche hazard forecasting, *Cold Regions Science and Technology*, 147, 1–10, <https://doi.org/10.1016/j.coldregions.2017.12.006>, 2018.
- Wang, X., Mueen, A., Ding, H., Trajcevski, G., Scheuermann, P., and Keogh, E. J.: Experimental comparison of representation methods and distance measures for time series data, *Data Mining and Knowledge Discovery*, 26, 275–309, <https://doi.org/10.1007/s10618-012-0250-5>, 2013.
- 25