

# ***Interactive comment on “Modeling of land-surface interactions in the PALM model system 6.0: Land surface model description, first evaluation, and sensitivity to model parameters” by Katrin Frieda Gehrke et al.***

**Katrin Frieda Gehrke et al.**

gehrke@muk.uni-hannover.de

Received and published: 1 February 2021

Summary:

Reviewer Comment (RC): This paper describes the land-surface model in the PALM model system. The subject matter is appropriate for the GMD journal and the text is well written. The details about the model are provided in appropriate amount of detail, equations provided are also appropriate. As part of the paper, there is a "first evaluation" which I think could/should be improved upon, as detailed in my comments below.

Printer-friendly version

Discussion paper



Even though this is a "first evaluation" I think it can be done with more rigor and there are, in my opinion, a few technical flaws. While I think the description of the model is ready to be published, the evaluation of the model (and comparison with observations) needs improvement before that portion of the paper should be published.

Author's reply (AR): In the name of all authors, I would like to thank the anonymous referee for the time and effort that was devoted to reviewing the paper. The valuable comments helped to significantly improve the original manuscript.

General Comments:

RC: 1. Most land-surface models vs observation comparisons I am aware of use the model run in "single-point" mode with the observed tower data driving the model. However, the model output from the PALM LSM has been averaged over the spatial domain. This spatial seems to confound the comparison (e.g., l.337-340). Is there a reason the spatial averaging of the model data is necessary?

AR: In (LES) modeling, it is common practice to employ the spatial average instead of a temporal average as it is done for single-point observations. It would be possible to mimic observations by using a time-averaging window to remove/reduce the turbulent signal from the LES data. Nonetheless, according to Taylor's Hypothesis, a spatial signal in our LES model should relate to a temporal signal in the observations as long as the surface is homogeneous. From a practical point of view, using the spatial average at one point in time is rather convenient and requires much less memory (it can be calculated on-the-fly during the simulation). Physically, the spatial average is in general superior over a temporal average, because Taylor's hypothesis is affected by changes of the mean wind direction, non-stationarity of the flow, and self-correlation. Because a spatial average over homogeneous terrain is generally equivalent to the single-point temporal average we decided to use the standard output of PALM (which is the horizontally-averaged data). This issue is not present in RANS simulations with horizontally homogeneous surface, as the RANS model only provides the

[Printer-friendly version](#)

[Discussion paper](#)



time-averaged flow so that there are no spatial and temporal variations due to turbulence. In order to add information about the spatial variability of the shown LES data, we have included the range (minimum to maximum value) in Figs. 3-6.

RC: 2. The 2-day period seems too short to do a thorough evaluation of the model. There are many decades of Cabauw data, but only a short 2-day period is used. Even for a "first evaluation" this seems like a weakness of the paper. According to the authors, this particular period was chosen because (l.236) "...the forcing from the surface was dominant and larger-scale advection played a minor role." However, many times later in the paper they attribute problem with the model-observation comparison to largerscale issues (e.g., l.333, "...which could be caused by advection processes in reality modifying the residual layer"). A much stronger statement would look at many days when the surface forcing is dominant and then contrast this with many other days when the surface-forcing is not dominant. Then the authors could actually show the model does better (or worse) when surface forcing dominates rather than simply making a vague statements about it.

AR: Even though the Cabauw site is one of Europe's most frequently used sites for this kind of comparison, it is not easy to find a period where there is little advection, no clouds and all measuring instruments are running. The period we chose was suggested by the person in charge for the Cabauw site (Fred Bosveld), who has great experience in the data quality and availability. A major reason for this particular period was that in May 2008 radiosondes were launched three times daily as part of the IMPACT-EUCAARI campaign (Kulmala et al., 2009), which we used for initialization. Overall, the period in question did have a few more days, but as we would have had to incorporate some kind of nudging to the forcing and/or data assimilation to avoid model drift, we decided to simulate a two-day period only. This gives the model the possibility to study the behavior of the model over a full diurnal cycle and also look at how the first day affects the following day(s). In general, it is not possible to derive the height-dependent boundary layer advection from observation needed to drive the

[Printer-friendly version](#)[Discussion paper](#)

LES model, particularly within the boundary layer, where turbulence dominates, such large-scale tendencies are difficult or impossible to obtain. We thus think that for a "first evaluation" the chosen period is acceptable. We have added a brief discussion about this to the manuscript which points out why we have chosen exactly this period.

RC: 3. The smaller observed H and LE fluxes than modeled flux during the daytime (ie, Fig. 7) is almost certainly related to the choice of a 10-min averaging period to calculate the turbulent fluctuations. The authors acknowledge that there are low-frequency issues with the fluxes (ie, l.247-250) and during the daytime the time-scale involved for the fluxes are longer than 10 minutes. Perhaps the Kaimal correction they describe fixes this issue, but using a 10-min window to calculate the fluxes is certainly a problem in the daytime (probably ok for nighttime). Why use a model to try and fix a methodology shortcoming? If a longer time window is used (e.g., either 30-min or an hour), it will make daytime obs H and LE larger and in closer agreement to the modeled fluxes.

AR: We agree with the reviewer that 10-minute intervals averaging time during daytime are certainly too short and the natural choice would be 30 minutes, though even 30-minute intervals are often not sufficient under convective conditions. For the current Cabauw data, 10 minute intervals are used and processed/corrected according to the Cabauw standard. Unfortunately we do not have access to the raw data (i.e. the raw data is not stored permanently) to calculate fluxes based on a longer time window. However we discuss the shortcomings of the method in the manuscript.

RC: 4. p.3, l.71, why is the heat capacity assumed to be zero for vegetation-covered surfaces? Heat capacity has recently been shown to be an important consideration in land-surface models (e.g., Swenson, et al 2019). Getting the heat capacity of the storage terms (soil, biomass) correct is an important consideration to properly close the SEB (e.g., Lindroth, et al 2010, Leuning, et al 2012). These so-called "smaller" terms are important because they tend to have a phase shift (in terms of the diurnal cycle) relative to the other Rnet/H/LE flux terms. The authors appear to focus on the issue of low-frequency contributions to the fluxes, and do not talk about the heat storage terms

[Printer-friendly version](#)[Discussion paper](#)

as a possible problem (in fact, since heat capacity of the vegetation biomass is set to zero can the biomass storage term even be considered?).

AR: The implementation of vegetated surfaces in PALM is based on the parameterization used in the LSM of the Integrated Forecast System (IFS). Accordingly, the skin layer has no heat capacity (IFS Documentation). This is because the heat storage of the vegetation layer is difficult to estimate and would introduce another parameter of uncertainty. Please also note that the vegetation in the simulated domain is short grass (homogeneous), whereas Swenson, et al 2019 and Lindroth, et al 2010 study heat stored in forests, where the heat capacity of the canopy is much much higher than for short grass. The assumption to have zero heat capacity for short grass is not unrealistic and thus a valid approach (we have added a brief discussion about this to the manuscript). The heat capacity of the soil is treated properly by the soil model.

RC: 5. The model has been designed to have many options and work with many different land-surface types (e.g., Table 1)...however, the evaluation is only done for one specific land-surface type. This is a very limited test of the validity of the model over the parameter space—realize article length is an issue—but, what about evaluations of other surface types? At least maybe cover more than just one?

AR: The evaluation is done for 21 cases of which 11 affect land-surface parameters (ALBE\_24, ALBE\_44, CAP\_2e4, COND\_2, COND\_6, EMIS\_95, EMIS\_100, LAI\_05, LAI\_3, ROUGH\_01, ROUGH\_001), and thereby the land-surface type “short grass” is implicitly altered. In our opinion, the parameter space thus appears to be systematic. We acknowledge that adding some more of the pre-set land-surface types (Table 1) may be in the interest of the user, however, this requires more observational sites over the respective homogeneous surface types, which are not easily available with similar data quality as of the Cabauw site. In particular, it is difficult to find locations that are horizontally homogenous. This becomes particularly true for artificial surfaces such as pavements. Nevertheless, we are planning to evaluate the land surface model also for pavements based on observational near-surface data obtained during a measurement

[Printer-friendly version](#)[Discussion paper](#)

campaign on a disused movement area of an airport in Berlin, Germany.

RC: 6. Though there is good information in Section 5.2, it seems like this section would benefit from subsections that guide the reader a bit better. As it is, I find it difficult to extract the key points the authors want to make from the comparison.

AR: Thank you, we have included subsections accordingly and restructured the text in some parts to better point-out the key points.

Specific Comments:

\* RC: does PALM stand for anything? Is this an acronym? If so, it should be stated when first introduced...

AR: Even though the PALM developers do not want to use the long name (abbreviation for Parallelized Large-eddy Simulation Model) anymore and this paper is part of a special issue featuring PALM, we have included a note in the revised manuscript, because readers, who are unfamiliar with the model are expecting some kind of explanation.

\* RC: p.4, l.98, remove parentheses with Duynkerke, 1999 reference

AR: Thank you for spotting this.

\* RC: p.6, l.159, is "high" vegetation, tall vegetation? such as trees?

AR: Yes, we changed it accordingly.

\* RC: p.8, Table 1, why is C<sub>0</sub> set to 0.00 for all vegetation types?

AR: The surface heat capacity is set to 0 by default, because this is, how it is done in the IFS code. Even though this is definitely unsuitable for tall vegetation such as forests, we do not give explicit values, because, to meet our standard, they must be comprehensively tested to not mislead the user. Nevertheless, the user has the possibility to change the value. We added a note to the revised manuscript that this value should be carefully adjusted, if e.g. a forest is simulated.

Printer-friendly version

Discussion paper



\* RC: p.11, l.220, do waves have any effect on the transfer coefficients over water?

AR: In the case of inland water, say for small lakes, rivers, ponds, etc., the transfer coefficients are not altered. For ocean, the roughness lengths vary to account for sub-grid scale waves. Here we use a Charnock parameterization. This information was given in line 94-96. We added another note to the section about treatment of the water surfaces to improve readability.

\* RC: p.12, l.244, "...but means of a Fourier extrapolation." Is there a reference for this method?

AR: A reference (Bosveld, 2020) was added to the revised version.

\* RC: p.12, l.244, Was a soil heat flux plate also used at some depth below or near the temperature measurements? If so, this is not clearly stated. Flux plates are typically used for measuring the soil flux while the soil temperature profile is used for the heat stored in the soil (e.g. see Eq. 7 and discussion in Leuning, et al 2012). [I now see this discussed on p.21, l.405-406]. Perhaps I don't fully understand this, but it seems like the comparison of the modeled and observed soil heat flux needs further consideration. Are the same quantities actually being compared in Fig. 6?

AR: Surface soil heat flux can also be derived from the soil heat flux observations alone. We resolve the 24h time series of G05 and G10 in its Fourier components. Corresponding components can then be extrapolated to the surface in the same way as described above for the diurnal Fourier component. Subsequently an inverse Fourier transformation is performed on the extrapolated components to construct the time series of the surface soil heat flux. The penetration depth for short time scales becomes small. This means that for these high frequency components the signal may be hidden in distortions of the observations, either noise or deviations because the time series is not a response to a perfect cyclic forcing with a period of 24 hours. In the current implementation the first 9 Fourier components are used, thus the fastest resolved cycle has a length of 2h40m. Extrapolation of component is done when the amplitude of the

Printer-friendly version

Discussion paper



10 cm Fourier component is  $> 1 \text{ W m}^{-2}$  and when the amplitude of the 5cm sensor is less than 3 times the amplitude of the 10 cm sensor. If this conditions are not met the amplitude of the 5 cm sensor is used.

\* RC: p.12, l.260, Fig. 2 is mentioned before Fig. 1.

AR: We established the correct order.

\* RC: p.13, l.282, are the root fraction values based on measurements or assumed?

AR: Root density is based on the study of Jager (1976) and assumed for model layers that are in-between the observed layers. We clarified this accordingly.

\* RC: p.16, Table 5, how were the specific values for each variable selected? For example, LAI has values of 0.5 to 3  $\text{m}^2/\text{m}^2$ . Are these realistic or reasonable values? Furthermore, if you want to truly look at the sensitivity to LAI (or other variables), why not vary them between the endpoints, e.g., in steps of 0.1  $\text{m}^2/\text{m}^2$  between 0.5 and 3?

AR: In the revised manuscript we have specified our choices. Indeed, starting from the reference case which was setup as a best guess based on reports of the Cabauw site, we have varied the respective parameter in a reasonable range. We agree with the reviewer that we do not show a full parameter study but rather a small part of the parameter range, which we now make more clear. With this study we did not intend to provide a comprehensive parameter study but an idea on how sensitive the model reacts on specific parameters. This is mainly motivated by the fact that in many cases these input data is either not available and need to be estimated, or often only roughly available. In both cases the uncertainty in the estimated parameters is high. We now try to make this more clear at the end of the setup description.

\* RC: p.18, l.345-374, I understand there is a difference in Rnet which is presumably due to an incorrect modeled surface temperature. But, I'm not sure what to take away from the discussion following this—is the suggestion that the LAI should really be 0.5  $\text{m}^2/\text{m}^2$ ? Is the problem with the observations since the radiative flux divergence is not

[Printer-friendly version](#)

[Discussion paper](#)





included?

AR: The paragraph provides a discussion on how sensitive the surface net radiation reacts on changes of specific model parameters. We do not intend to give the impression that there is one truth and one perfect parameter combination, so thanks for pointing this out. Instead our intention is to outline the sensitivity of the energy balance components, here surface net radiation, on specific parameter variations. In the revised manuscript we try to make this more clear what the intention is.

\* RC: p.21, l.410, "It strikes", should be "It is striking"?

AR: Thanks for this hint. In the revised version we have changed it to "remarkably".

\* RC: p.21, l.413-416, are you suggesting that the observed LAI is incorrect? If you increase LAI, LE should increase at the expense of H..this is not surprising.

AR: Removed sentence to avoid confusion.

\* RC: p.24, l.472, for more info on grid spacing of models in stable conditions, see Sullivan, et al 2016.

AR: Thank you, the reference has been added.

\* RC: p.28, l.575, "differences of up to 50% are possible.". Differences in which variable?

AR: H and LE. The sentence has been rearranged for clarification.

\* RC: p.29, l.602-603, I didn't see how step-like orography is implemented in the LSM? Was this described somewhere in the paper?

AR: The reviewer is right, these information was given out of the context. We have now outlined the issue to make our point more clear.

References:

Bosveld, F.: The Cabauw In-situ Observational Program 2000 – Present: Instruments,

Calibrations and Set-up, Tech. report 384, KNMI, De Bilt, The Netherlands, regularly updated, 2020. IFS Documentation, Physical Processes, Chapter 8 Surface parameterisation, 2016, <http://www.ecmwf.int/en/elibrary/16648-part-iv-physical-processes>

Jager, C., Nakken, C., and Palland, C.: Bodemkundig Onderzoek van twee Graslandpercelen Nabij Cabauw, NV Heidemaatschappij Beheer, in Dutch, 1976.

Kulmala, M., Asmi, A., Lappalainen, H. K., Carslaw, K. S., Pöschl, U., Baltensperger, U., Hov, Ø., Brenquier, J.-L., Pandis, S. N., Facchini, 775 M. C., Hansson, H.-C., Wiedensohler, A., and O'Dowd, C. D.: Introduction: European Integrated Project on Aerosol Cloud Climate and Air Quality interactions (EUCAARI) – integrating aerosol research from nano to global scales, *Atmospheric Chemistry and Physics*, 9, 2825–2841, <https://doi.org/10.5194/acp-9-2825-2009>, 2009

Lindroth, A., Molder, M., and Lagergren, F., 2010: Heat storage in forest biomass improves energy balance closure, *Biogeosciences*, 7, 301-313, doi:10.5194/bg-7-301-2010

Swenson, S. C., Burns, S.P. , and D.M. Lawrence, 2019: The impact of biomass heat storage on the canopy energy balance and atmospheric stability in the Community Land Model. *Journal of Advances in Modeling Earth Systems (JAMES)*, 11, 83-98, doi:10.1029/2018MS001476

---

Interactive comment on *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2020-197>, 2020.

Printer-friendly version

Discussion paper

