

1 **Sensitivity of surface solar radiation to aerosol-radiation and aerosol-** 2 **cloud interactions over Europe in WRFv3.6.1 climatic runs with fully** 3 **interactive aerosols**

4 Sonia Jerez^{1*}, Laura Palacios-Peña¹, Claudia Gutiérrez², Pedro Jiménez-Guerrero¹, Jose María
5 López-Romero¹, E. Pravia-Sarabia¹, Juan Pedro Montávez¹

6 (1) Department of Physics, Regional Atmospheric Modeling group, Regional Campus of
7 International Excellence "Campus Mare Nostrum", University of Murcia, 30100 Murcia, Spain

8 (2) Environmental Sciences Institute, University of Castilla-La Mancha, 45071 Toledo, Spain

9 *Correspondence to: sonia.jerez@gmail.com

10 **Abstract**

11 The amount of solar radiation reaching the Earth's surface can be highly determined by atmospheric
12 aerosols, ~~which the IPCC has called~~ **pointed** as the most uncertain climate forcing agents; through
13 their direct (scattering **and absorption**), semi-direct (absorption **implying a thermodynamic effect**
14 **on clouds**) and indirect (cloud properties modification **when aerosols act as cloud condensation**
15 **nuclei**) effects. Nonetheless, Regional Climate Models hardly ever dynamically model the
16 atmospheric concentration of aerosols and their interactions with radiation (ARI) and clouds (ACI).
17 The objective of this work is to evince the role of modeling ARI and ACI in Weather Research and
18 Forecast (WRF) model simulations with fully interactive aerosols (online resolved concentrations)
19 with a focus on summer mean surface downward solar radiation (RSDS) over Europe. Under
20 **present-day historical** conditions (1991-2010), both ARI and ACI reduce RSDS by a few
21 percentage points over central and northern regions. This reduction is larger when only ARI are
22 resolved, while ACI counteract the effect of the former by up to half. The response of RSDS to the
23 activation of ARI and ACI is mainly led by the aerosol effect on the cloud coverage, while the
24 aerosol effect on the atmospheric optical depth plays a very minor role, **which evinces the**
25 **importance of the.** ~~This suggests that~~ semi-direct and indirect aerosol effects ~~prevail over the~~
26 ~~direct effect.~~ **In fact Consistently**, differences in RSDS among experiments with and without
27 aerosols are softer under clear-sky conditions. In terms of future projections (2031-2050 vs. 1991-

28 2010), the baseline pattern (from an experiment without aerosols) shows positive signals southward
29 and negative signals northward. While ARI enhance the former and reduce the latter, ACI work in
30 the opposite direction and provide a flatter RSDS change pattern, further evincing the opposite
31 impact from semi-direct and indirect effects and the non-banal influence of the latter.

32 **1 – Introduction**

33 Regional Climate Models (RCMs) are powerful tools providing high-resolution climate information
34 by dynamically downscaling coarser datasets, e.g. from Global Circulation Models (GCMs). Their
35 added value comes not only from the increased resolution, but also from the fact that such an
36 increased resolution allows modeling and considering fine scale processes and features that are
37 missed or misrepresented otherwise, e.g. local circulations and land uses (Rummukainen 2010,
38 Jacob et al 2014, 2020, Schewe et al 2019). Still, certain phenomena need to be parametrized, e.g.
39 the turbulence within the planetary boundary layer, microphysics processes and convective
40 phenomena. However, there are relevant processes that GCMs usually model dynamically, but
41 which ~~RCMs usually do not~~ **are not usually included in RCMs runs**. This is the case of the
42 atmospheric aerosol concentration and their multiple non-linear interactions (e.g. Taylor et al 2012
43 vs. Ruti et al 2016), the so-called aerosol-radiation and aerosol-cloud interactions (ARI and ACI
44 respectively; Boucher 2015).

45 Depending on their nature and the ambient conditions, aerosols can act to scatter and/or absorb the
46 solar radiation through ARI, which may result in less or more solar radiation reaching the surface
47 **through direct and semi-direct effects. Direct effects might involve that less solar radiation**
48 **reaches the surface due to its scattering and absorption (Giorgi et al 2002, Nabat et al 2015a,**
49 **Li et al 2017, Kinne 2019), or more if, for instance, absorption warms aloft atmospheric**
50 **layers, thereby leading to more stable atmospheric situations (lower surface temperatures**
51 **than upward) and thus to the inhibition of clouds formation via convective phenomena**
52 **(Giorgi et al 2002, Nabat et al 2015a). Absorption itself can also lead to clouds inhibition**
53 **and/or burn-off through thermodynamic effects, i.e. by heating the air (semi-direct effects),**
54 **thus increasing the amount of solar radiation reaching the surface (Allen and Sherwood**
55 **2010). ~~less because of its scattering (direct effect), more if absorption (semi-direct effect) leads~~
56 ~~to clouds burn-off and/or inhibition (Giorgi et al 2002, Nabat et al 2015a, Li et al 2017, Kinne~~
57 ~~2019).~~ **Besides, a**Aerosols **also** act as cloud condensation nuclei (indirect effect or ACI), which may
58 also result in less or more solar radiation reaching the surface. Abundance of cloud condensation**

59 nuclei rebounds on enhanced scattering via whitened clouds of smaller drops with increased size
60 and lifetime, and on drizzle suppression which reduces below-cloud wet deposition processes
61 (Seinfeld et al 2016, Kinne 2019). On the contrary, in-cloud aerosol scavenging processes lead to
62 out-of-cloud cleaner atmospheres (Croft et al 2012). All these processes can potentially alter local
63 and regional circulations, therefore impacting beyond the radiative balance (Kloster et al 2010,
64 Wilcox et al 2013, Nabat et al 2014, Wang et al 2016, Pavlidis et al 2020).

65 In the current context of climate crisis, the scientific challenge is becoming twofold: (1) to gain a
66 good understanding of the processes that occur in the atmosphere and of what will occur in the
67 future, because this is crucial (IPCC 2013) in order (2) to advance effective measures both at global
68 and regional scales (IPCC 2014). In particular, climate change mitigation strategies require low-
69 carbon energies to grow rapidly in the coming decades (Rohrig et al 2019, IRENA 2019). This rapid
70 transition of the energy sector towards renewable-powered decarbonized systems makes energy
71 production, transmission and distribution increasingly sensitive to weather and climate variability
72 (Bloomfield et al 2016, Collins et al 2018, Kozarcenin et al 2018, Jerez et al 2019). Thus, several
73 works have been devoted to assessing this issue through the use of climate modeling tools. In
74 particular, for the solar resource, Crook et al (2011), Gaetani et al (2014), Wild et al (2015) and
75 Müller et al (2019) showed a generalized increase in Europe by making use of GCM simulations,
76 while Jerez et al (2015), Gil et al (2019) and Tobin et al (2018) reported a different behavior, with
77 RCM simulations projecting a slight general decrease in the amount of solar radiation reaching the
78 surface over Europe.

79 From the previous literature, we point out here three key features that motivated the present work.
80 First, the increasing use of RCM to evaluate the renewable energy resources and their supply
81 potential (e.g. Jerez et al 2013, 2015, 2019, Gil et al 2019, Soares et al 2019, van der Wiel et al
82 2019). Second, the key role of aerosols regarding the accuracy of the simulated solar resource by
83 climate models (Gaetani et al 2014, Nabat et al 2015b, Pavlidis et al 2020), particularly attributed to
84 their direct and semi-direct effects, which would help to explain the aforementioned discrepancy
85 between the GCM and RCM future projections (Boé et al 2020, Gutiérrez et al 2020). Third, none
86 of the previous studies has so far dealt with the non-evident RCM sensitivity to interactively
87 modeled atmospheric aerosol concentrations and the resulting aerosol-radiation and aerosol-cloud
88 interactions in order to simulate the solar resource under **present historical** and future climate
89 scenarios.

90 Hence, our objective here is to shed light on the third point above by assessing the sensitivity of
91 long-term RCM simulations to the inclusion of ARI and ACI using fully interactive (online
92 diagnosed) aerosols. For this, we made use of a widely applied RCM, the Weather Research and
93 Forecasting (WRF) model (Skamarock et al 2008) and its coupled form with Chemistry (WRF-
94 Chem; Grell et al 2005), to perform sets of **present historical** (period 1991-2010) and future
95 (period 2031-2050) simulations over Europe in three ways: (1) without including atmospheric
96 aerosols, (2) with dynamic aerosols and aerosol-radiation interactions activated, and (3) with
97 dynamic aerosols and both aerosol-radiation and aerosol-cloud interactions activated.

98 Section 2 describes experiments and methods; section 3 presents the results; the discussion and
99 conclusions are provided in Section 4.

100 **2 – Experiments, data and methods**

101 **2.1 – General description of the WRF simulations**

102 We performed three experiments using the WRF model version 3.6.1 (Skamarock et al 2008;
103 available at <https://www.mmm.ucar.edu/weather-research-and-forecasting-model>). In all cases, the
104 simulated periods were 1991-2010 (**present historical**) and 2031-2050 (future). Initial and
105 boundary conditions were taken from GCM simulations: the r1i1p1 MPI-ESM-LR historical and
106 RCP8.5-forced runs (Giorgetta et al 2012a,b; available at <https://cera-www.dkrz.de>) from the
107 Coupled Model Intercomparison Project Phase 5 (CMIP5; <https://pcmdi.llnl.gov/mips/cmip5/>;
108 Taylor et al 2012). The Representative Concentration Pathway RCP8.5 (Moss et al 2010) depicts
109 the highest radiative forcing along the XXI century among all RCPs, with doubled CO₂, CH₄, and
110 N₂O concentrations by 2050 compared to the last record of the historical period. Both the observed
111 (past) and estimated (future) temporal evolution of the concentration of these species was
112 appropriately considered in the WRF executions (Jerez et al 2018).

113 The three experiments consisted of, and are named as:

114 BASE: aerosols are not considered in the simulations. No aerosol climatology is used, and no
115 aerosol interactions are taken into account by the model. WRF-alone considers a constant number

116 of cloud condensation nuclei (250 per cm³, set in the model by default) to enable the formation of
117 clouds.

118 ARI: aerosols are **estimated online** ~~treated interactively (see below)~~ and aerosol-radiation
119 interactions are activated in the model (**both direct and semi-direct effects are included in the**
120 **simulations**).

121 ARCI: aerosols are **estimated online** ~~treated interactively (see below)~~ and both aerosol-radiation
122 and aerosol-cloud interactions are activated in the model (**direct, semi-direct and indirect effects**
123 **are included in the simulations**).

124 The WRF spatial configuration consisted of two one-way nested domains (Supp Fig 1). The inner
125 one (target domain) is an Euro-Cordex (<https://www.euro-cordex.net/>; Jacob et al 2014, 2020)
126 compliant domain covering Europe with a horizontal resolution of 0.44° in latitude and longitude.
127 The outer one has a horizontal resolution of 1.32° and covers the most important areas of Saharan
128 dust emission as in Palacios-Peña et al 2019a. This configuration was necessary to generate and
129 include the information of the Saharan dust intrusions through the boundaries of our target domain
130 for the ARI and ARCI experiments, because the boundary conditions from the GCM do not provide
131 this information. In the vertical dimension, 29 unevenly spaced eta levels were specified in the two
132 domains, with more levels near the surface than upward, and the model top was set to 50 hPa. The
133 physics configuration of the WRF model consisted of the Lin microphysics scheme (Lin et al 1983),
134 the RRTM **long- and short-wave** radiative scheme (Iacono et al 2008), the Grell 3D ensemble
135 cumulus scheme (Grell 1993, Grell and Dévényi 2002), the University of Yonsei boundary layer
136 scheme (Hong et al 2006) and the Noah land surface model (Chen & Dudhia 2001, Tewari et al
137 2004). Boundary conditions from the GCM were updated every 6 hours, including the low
138 boundary condition for the sea surface temperature. Nudging was applied to the outer domain, but
139 not to the target domain.

140 **2.2 – Including aerosols in WRF**

141 To perform the ARI and ARCI experiments, we used the WRF model coupled with Chemistry
142 (WRF-Chem) version 3.6.1 (Grell et al 2005, Chin et al 2002). WRF-Chem runs with GOCART
143 aerosol module (Ginoux et al 2001). This scheme includes five species, namely sulfate, mineral
144 dust, sea salt aerosol, organic matter and black carbon, and was coupled with RACM-KPP

145 (Stockwell et al 1997, Geiger et al 2003) as chemistry option. Chemical reactions in the GOCART
146 model include several oxidation processes by the three main oxidants in the troposphere: OH,
147 NO₃, and O₃. The OH radical dominates oxidation during the daytime, but at night its
148 concentration drops and NO₃ becomes the primary oxidant (Archer-Nicholls et al 2014). So, the
149 oxidation pathways represented in GOCART include: (a) dimethyl sulfide (DMS) oxidation by the
150 hydroxyl radical (OH) during the day to form sulfur dioxide (SO₂) and methanesulfonic acid
151 (MSA); (b) oxidation by nitrate radicals (NO₃) at night to form SO₂; and (c) SO₂ oxidation by OH
152 in air and by H₂O₂ and tropospheric ozone (O₃) in clouds (aqueous chemistry) to form sulfate (Chin
153 et al 2000). Henceforth, the skilful characterization of gas-phase radicals such as OH and NO₃ or
154 compounds like O₃ is essential for the representation of oxidation pathways in the atmosphere
155 leading to the formation of secondary aerosols (Jiménez et al 2003). Therefore, in this contribution
156 the RACM (Stockwell et al 1997, Geiger et al 2003) mechanism was coupled to GOCART through
157 the kinetics pre-processor (KPP) in WRF-Chem in order to provide the concentrations of radical
158 and gas-phase pollutants needed by the GOCART aerosol model. The Fast-J module (Wild et al
159 2000) was used as photolysis option. Biogenic emissions were calculated using the Guenther
160 scheme (Guenther et al 2006). Anthropogenic emissions coming from the Atmospheric Chemistry
161 and Climate Model Intercomparison Project (ACCMIP; Lamarque et al 2010) were kept unchanged
162 in the simulation periods (we considered the 2010 monthly values). Natural emissions depend on
163 ambient conditions and varied accordingly in our simulations following Ginoux et al 2001 for dust
164 and Chin et al 2002 for sea salt.

165 The inclusion of aerosol-radiation interactions in the called ARI simulations follows Fast et al
166 (2006) and Chapman et al (2009). The overall refractive index for a given size bin was determined
167 by volume averaging associating each chemical constituent of aerosol with a complex index of
168 refraction. The Mie theory and the summation over all size bins were used to determine the
169 composite aerosol optical properties assuming wet particle diameters, **taking into account the**
170 **humidity variations to allow variations of optical properties**. Finally, aerosol optical properties
171 were transferred to the shortwave radiation scheme. Aerosol-cloud interactions were implemented
172 by linking the simulated cloud droplet number with the microphysics schemes (Chapman et al
173 2009) affecting both the calculated droplet mean radius and the cloud optical depth. Although this
174 WRF-Chem version (3.6.1) does not allow a full coupling with aerosol-cloud interactions **that**
175 **includes the aerosols exerting the highest influence from a climatic point point of view, i.e. sea**
176 **salt and desert dust**, the microphysics implemented here is **a modified version of** a single
177 moment scheme that turns **it** into a two-moment scheme in the simulations denoted ARCI. One-

178 moment microphysical schemes are unsuitable for assessing the aerosol-cloud interactions as they
179 only predict the mass of cloud droplets and do not represent the number or concentration of cloud
180 droplets (Li et al 2008). The prediction of two moments provides a more robust treatment of the
181 particle size distributions, which is key for computing the microphysical process rates and
182 cloud/precipitation evolution. In this sense, **although the Lin microphysics is originally presented
183 as a single moment scheme (Lin et al 1983), a modified Lin double-moment microphysical
184 scheme is implemented in WRF-Chem (Lin et al 2008) and used here to conduct the ARCI
185 simulations. In this scheme, both the mass and the total number of cloud droplets are
186 predicted. The prognostic treatment of cloud droplet number involves water vapor, cloud
187 water, rain, cloud ice, snow and graupel (Ghan et al 1997), and is activated through the
188 “mixactivate” module of WRF-Chem. In that module, WRF-Chem calculates the aerosol
189 number per volume concentration by using, for each aerosol type, the information about the
190 size (the mean volume-diameter of each aerosol mode, obtained from the aerosol mechanism
191 implemented in the simulation), and fixed densities and molecular weight of each type of
192 aerosols. With all this information and the total mass, WRF-Chem estimates the aerosol
193 number for each mode assuming spherical particles. The autoconversion of cloud droplets to
194 rain droplets depends on droplet number (Liu et al 2005). Droplet-number nucleation and
195 (complete) evaporation rates correspond to the aerosol activation and resuspension rates. Ice
196 nuclei based on predicted particulates are not treated. However, ice clouds are included via
197 the prescribed ice nuclei distribution, following the Lin et al (2008) scheme. Thus, the droplet
198 number will affect both the calculated droplet mean radius and cloud optical depth. Finally,
199 the interactions of clouds and incoming solar radiation were implemented by linking the
200 simulated cloud droplet number with the Goddard shortwave radiation scheme, representing
201 the first indirect effect (i.e. increase in droplet number associated with increases in aerosols),
202 and with the Lin microphysics, representing the second indirect effect (i.e. decrease in
203 precipitation efficiency associated with increases in aerosols). ~~although the Lin microphysics is
204 presented as a single moment scheme, the WRF-Chem model makes it possible to transform
205 the single into a double-moment scheme. A prognostic treatment of cloud droplet number was
206 added (Ghan et al 1997), which treats water vapor and cloud water, rain, cloud ice, snow, and
207 graupel. The autoconversion of cloud droplets to rain droplets depends on droplet number
208 (Liu et al 2005). Droplet-number nucleation and (complete) evaporation rates correspond to
209 the aerosol activation and resuspension rates. Ice nuclei based on predicted particulates are
210 not treated. However, ice clouds are included via the prescribed ice nuclei distribution,
211 following the Lin scheme. Finally, the interactions of clouds and incoming solar radiation~~**

212 ~~were implemented by linking the simulated cloud droplet number with the Goddard~~
213 ~~shortwave radiation scheme, representing the first indirect effect, and with the Lin~~
214 ~~microphysics, representing the second indirect effect (Skamarock et al 2008). Thus, the~~
215 ~~droplet number will affect both the calculated droplet mean radius and cloud optical depth.~~

216 An important aspect of the differences in the model setup between experiments is that the
217 autoconversion scheme necessarily changes in the ARCI simulations as compared to the model
218 configuration used for ARI and BASE. The flag *progn* of the WRF namelist should be set to 0
219 for running ARI experiments in order to keep disabled the interaction of the online-estimated
220 aerosols with cloud microphysics, hence ensuring the use of prescribed aerosols (as in the case
221 of the BASE simulations) as this regards. Conversely, *progn* should be set to 1 for running
222 ARCI experiments in order to feed the cloud microphysics scheme with the online-estimated
223 number and physico-chemical properties of aerosols (this effectively turns the Lin scheme into
224 a second-moment microphysical scheme).

225 2.3 – Data and methods

226 The WRF and WRF-Chem outputs were recorded every hour for surface downward solar radiation
227 (RSDS), total cloud cover (CCT) and the concentrations of various aerosol species (dust, black
228 carbon, organic carbon and sea salt). The concentration of sulfates was indirectly computed from
229 the recorded concentrations of SO₂ and OH using the same kinetic reaction implemented in the
230 RACM-KPP module. From the concentrations of the five aerosol species, the atmospheric optical
231 depth (AOD) at 550 nm was estimated using the reconstructed mass-extinction method (Malm et al
232 1994), as in Palacios-Peña et al (2020). The RSDS and CCT data simulated by the driving GCM
233 runs were used for comparison purposes. We also retrieved the AOD at 550 nm as seen by the GCM
234 from the MACv2 data (Kinne et al 2019), whose anthropogenic changes are in accordance with the
235 RCP8.5 while its coarse mode (of natural origin) was not allowed to change. Also, RSDS values
236 from the ERA5 reanalysis (Hersbach et al 2020) were used for validation purposes. Seasonal means
237 means of all the variables were used in the analysis. These means involve all the records within
238 each season in the series.

239 We also studied the sensitivity to resolving aerosol interactions of RSDS and AOD under clear-sky
240 conditions. The analysis in absence of cloudiness will tell us more about the relevance of the direct
241 radiative effect of aerosols. RSDS and AOD clear-sky (RSDS_{cs} and AOD_{cs}, respectively) mean

242 seasonal series were constructed as follows. First, hourly series of CCT, RSDS and AOD were time
243 averaged up to the daily timescale. Second, days with CCT values lower than 1% were retained
244 (this criterion is applied at the grid-box level, for each grid-box individually); otherwise we put a
245 missing value. These clear-sky daily series were then time averaged up to the seasonal time-scale.
246 When pairs of experiments were compared, only coincident clear-sky dates (days) in the series were
247 selected (missing values were also assigned in this case to the non-coincident dates with clear-sky
248 conditions) before performing the seasonal time average. This restriction aims to avoid the
249 masking effect of Earth orbit related issues, of large scale climate drivers and/or local forcings such
250 as water vapor content, since different days may have different daytime lengths and different
251 atmospheric compositions (different atmospheric optical depth or atmospheric transmissivity) that
252 may mask the AOD effect under clear-sky conditions. The analysis involving $RSDS_{cs}$ and AOD_{cs}
253 was carried out only over those grid points where at least 75% of the summer mean values in the
254 series (i.e. at least 15 records per period) were not missing (which, according to our methodology,
255 would occur only if all days within a summer season had CCT values $\geq 1\%$).

256 Spatial correlations between climatological patterns were computed excluding sea grid points,
257 considering absolute values in case they involved differences (while these were depicted in the
258 Figures in relative terms, i.e. in %), using the CDO *fldcor* function
259 (<https://code.mpimet.mpg.de/projects/cdo/embedded/cdo.pdf>). Temporal correlations were
260 computed at the grid point level between the seasonal series, considering absolute values in case
261 they involved differences, using the R *cor* function
262 (<https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/cor>; *Pearson* correlation
263 coefficient selected). The statistical significance of any signal was assessed with a t-test **imposing**
264 **$p < 0.05$** .

265 We focus on the summer season (JJA), when solar energy is at its maximum, AOD typically reaches
266 high values and the aerosol radiative effect has been proven to be strongest (Pavlidis et al 2020).

267 In order to investigate the underlying mechanisms explaining the signals found in RSDS and CCT,
268 additional variables and statistics were used, namely: JJA-mean top-of-the-atmosphere outgoing
269 short-wave radiation (RSOT), surface (2 m height) air temperature (TAS), surface (1000 hPa
270 pressure level) relative humidity (RH), total precipitation (PR) and convective precipitation (PRC);
271 number of cloudy days (CLD, defined as days with mean CCT > 75%) in the summer series; 90th
272 percentile of the JJA day-mean PR series; and number of rainy days (RD, defined as days with

273 mean precipitation > 1 mm) in the JJA daily PR series. **Vertical profiles of air temperature (T)**
274 **and cloud fraction (CLFR) were also considered.**

275 **3 – Results**

276 **3.1 – ~~Present-day climatologies~~ Historical patterns**

277 *Brief validation of the simulated RSDS patterns*

278 As a first test, Supp Fig 2 provides the GCM, ERA5 BASE, ARI and ARCI JJA climatologies of
279 RSDS in the **present historical** period and the results of a brief validation exercise. Although the
280 five patterns depict similar structures (Supp Fig 2a,b,d-f), Supp Fig 2g-i reveals significant
281 deviations of the climatologies from the WRF experiments with respect to the GCM: positive values
282 (higher RSDS values in the RCM experiments) south and northward (up to 20 and 30%
283 respectively), and negative values in between (10-15%, eventually up to 25%). These differences
284 are very similar to those obtained when WRF climatologies are compared with the ERA5 pattern
285 (Supp Fig 2j-l), with a notable exception over the Scandinavian region where the agreement
286 between the WRF experiments and ERA5 is higher than between the WRF experiments and the
287 GCM. In fact, the GCM pattern strongly underestimates RSDS over such a region (over 30%; Supp
288 Fig 2c), while showing a better agreement with ERA5 elsewhere as compared to the WRF
289 simulations.

290 *Aerosols impact on the simulated RSDS patterns*

291 Although the three WRF experiments (BASE, ARI and ARCI) perform similarly when compared to
292 the GCM or ERA5, there are still noticeable differences between them (Fig 1a-c and Supp Fig 3a-
293 c), and it is there that this research focuses. The inclusion of aerosols (ARI and ARCI experiments)
294 reduces the JJA mean values of RSDS in central and northern parts of our domain by a few
295 percentage points (i.e. by $\sim 10 \text{ Wm}^{-2}$) as compared to the BASE experiment (Fig 1a,b and Supp Fig
296 3a-b). This reduction is generally stronger in ARI than in ARCI. Consequently, the ARCI minus
297 ARI pattern (Fig 1c and Supp Fig 3c) depicts mostly positive values (by $\sim 5 \text{ Wm}^{-2}$) over central and
298 southern regions. This result **already** indicates that the indirect aerosols effects tend to counteract
299 the **joint** direct **and semi-direct** effects **seen in the ARI minus BASE pattern** ~~of lower RSDS~~

300 ~~with higher aerosol concentrations over most of the domain~~, reducing it by up to a half **over**
301 **most of the domain**, which is in agreement with previously reported findings (Pavlidis et al 2020).

302 In order to better understand the patterns of differences in RSDS between experiments, Fig 1 (and
303 Supp Fig 3) also provides differences in CCT and AOD (panels d to f and g to i, respectively) and
304 the spatial correlations (*s_corr*) between these patterns and those of RSDS differences.

305 *The role of CCT*

306 Compared to BASE, both ARI and ARCI lead to more cloudiness in central and northern regions
307 (albeit quite slight increases, well below 5%). **This could respond to the direct effect of the**
308 **scattering of the solar radiation due to the high presence of sea salt, dust and sulfate over**
309 **these areas (Fig 2), as an increase in RSOT over these areas is also appreciated in both ARI**
310 **and ARCI simulations (Fig 3a-b). In addition, this direct effect could be triggering the**
311 **following feedback mechanism: the cooling effect downward (where less solar radiation is**
312 **received because of its scattering) cools down surface temperatures (Fig 3d-e), thus increasing**
313 **relative humidity (Fig 3g-h), which may favor the formation of clouds (these should be non-**
314 **convective, mostly low-level, clouds as the decrease in TAS leads to more stable atmospheric**
315 **layers; Fig 4a,b), thus less radiation reaches the surface, thus lower surface temperatures, and**
316 **so on. Noteworthy, both the reduction in RSDS and the accompanying increase in RSOT is**
317 **more marked in ARI than in ARCI over central regions (Fig 1c and Fig 3c), where the indirect**
318 **effects included in the ARCI simulation, such as in-cloud aerosol scavenging processes, could**
319 **lead to cleaner atmospheres than ARI simulates. ~~This could be explained through the~~**
320 **~~following feedback mechanism: the cooling effect of the scattering of radiation by the high~~**
321 **~~presence of sea salt, dust and sulfate over these areas (Supp Fig 4a-j and Supp Fig 5a-b) cools~~**
322 **~~down surface temperatures (Supp Fig 5d-e), thus increasing relative humidity (Supp Fig 5g-h)~~**
323 **~~and favoring the formation of clouds, which leads to less radiation reaching the surface, thus~~**
324 **~~lower surface temperatures, and so on. Nonetheless, these signals would simply indicate that~~**
325 **~~this enhancing mechanism prevails over others. For instance, the semi-direct effect, which acts~~**
326 **~~to suppress cloudiness due to the thermodynamic effect of dark particles, that could explain~~**
327 **~~that this CCT reduction was more evident in ARCI than in ARI (Fig 1c, Supp Fig 3c); or the~~**
328 **~~wet deposition and in-cloud aerosol scavenging processes leading to cleaner atmospheres,~~**
329 **~~which could explain why RSOT was larger in ARI than in ARCI in spite of the former (Supp~~**
330 **~~Fig 5c,f,i).~~**

331 Conversely, both ARI and ARCI lead to less cloudiness southward **as compared to BASE**,
332 especially ARCI (reductions up to 10% in Mediterranean regions; Fig 1d-e). Consistently, the ARCI
333 minus ARI pattern (Fig 1f) depicts negative values (around 5%) along the Mediterranean strip.
334 Therefore, both semi-direct and indirect aerosol effects would tend to diminish cloudiness
335 southward, with the latter (indirect effect) having the greatest impact. This could be due to the fact
336 that a high presence of large aerosols over southern Europe, both in form of dust or sulfate in our
337 case (**Supp Fig 2 4a-j**), ~~hampers the formation of clouds (e.g. Xue et al 2008) and may even~~
338 ~~shortens their lifetime by enhancing precipitation~~ **can accelerate collision-coalescence**
339 **processes fastening that precipitation occurs and thus shortening the lifetime of clouds (e.g.**
340 **Lee et al 2008)**, which is most plausible in the warm season over warm areas (**e.g.** Yin et al 2000),
341 as long as aerosol-cloud interactions are resolved by the model. However, we did not find such an
342 enhanced precipitation effect in our simulations (maybe the signal does not hold at the climatic
343 scales assessed here), only a decrease in both mean cloudiness and number of cloudy days (**Supp**
344 **Fig 35j-l**) together with consistent pictures of lower mean precipitation, lower mean convective
345 precipitation, fewer rainy days and lower extreme precipitation values emerging over those areas
346 where the aerosol effects diminish cloudiness (**Supp Fig 56**). **The reduction in convective**
347 **precipitation (the prevailing form of precipitation over this area during the summer season)**
348 **suggests that absorption might be creating more stable atmospheric situations (by heating**
349 **aloft layers) and thus preventing clouds formation via convective phenomena and increasing**
350 **the incoming surface solar radiation. But we did not find any clear evidence of that either (Fig**
351 **4c). So the thermodynamic effect of aerosols on clouds inhibition and burn-off might justify**
352 **the reduction in CCT (mainly at low levels; Fig 4d) and the accompanying increase in RSDS**
353 **in the southernmost areas. These signals are intensified when we add the indirect aerosols**
354 **effects, likely due to the removal of aerosols via scavenging processes, which cleans the**
355 **atmosphere favoring that the solar radiation reaches the surface.**

356 Whatever the underlying mechanisms are, the patterns of differences between experiments in CCT
357 are well correlated with the corresponding patterns of differences in RSDS, thus indicating a key
358 role of CCT in driving the latter. Indeed, the temporal correlation at the grid point level between the
359 seasonal series of RSDS and CCT differences is above 0.8 (negative) in most of the domain (**Supp**
360 **Fig 47a-c**).

361 *The role of AOD*

362 The inclusion of aerosols also leads to differences of a few percentage points (2-5%) in the AOD
363 values between ARCI and ARI simulations over western areas (Fig 1i), and the AOD climatologies
364 from these two experiments provide a consistently non-null picture (Fig 1g,h; null values can be
365 considered for BASE). However, the patterns for AOD do not correlate with those for RSDS and
366 the seasonal series of differences in AOD hardly correlates with the seasonal series of differences in
367 RSDS except for certain locations of central and southeast Europe (Supp Fig 47d-f). Interestingly,
368 over these locations, the temporal correlation between differences in RSDS and differences in AOD
369 are positive, **further** indicating the secondary role of the direct radiative effect of the aerosols there:
370 if the larger the AOD, the larger the RSDS, it is because semi-direct and indirect effects counteract
371 the impact of the direct scattering effect.

372 *Clear-sky analysis*

373 ~~Thus, there is a~~An overall predominant link between the aerosol effect on cloudiness and its
374 impact on the amount of solar radiation reaching the surface, that totally masks any other
375 mechanism related to the variation in AOD and its direct impact on RSDS, **has been detected so**
376 **far**. On the contrary, as expected, under clear-sky conditions, both the negative spatial correlations
377 between the patterns of AOD_{cs} and RSDS_{cs} differences between experiments (Fig 62), and the
378 negative temporal correlations between the respective series computed at the grid point level (Supp
379 Fig 47g-i), support the relevant role of the AOD_{cs} variable for the simulation of RSDS_{cs}. The
380 differences in RSDS_{cs} between ARI or ARCI and BASE are negative (around 5 Wm⁻²; Fig 62 and
381 Supp Fig 58) over the study area (restricted to the southern half of the domain since the clear-sky
382 series northward lack of sufficient records to perform a robust statistical analysis), illustrating the
383 direct radiative effect of aerosols and further supporting **the important role its smaller impact at**
384 ~~the time-scales considered here as compared to the overall impact~~ of semi-direct and indirect
385 effects (that make the negative clear-sky signals softer and even positive **over some southern**
386 **locations**, as shown in Fig 1a,b). ARCI minus ARI differences in RSDS_{cs} are basically null since
387 semi-direct and indirect effects are largely irrelevant in the absence of cloudiness.

388 **3.2 – Future projections**

389 *Future climatologies*

390 The overall results described above also hold under future climate conditions, while some
391 differences were identified and deserve mention. The inclusion of aerosols reduces RSDS over most
392 of the domain due to direct, **semi-direct** and indirect effects (Supp Fig 69a-c). In particular, this
393 occurs significantly southward, along the Mediterranean strip (~~Supp Fig 9a-c~~), in contrast to the
394 previous results. Over some locations, mainly in central Europe, this reduction is stronger in ARI
395 than in ARCI, as detected under **present-day historical** conditions. However, the opposite (larger
396 RSDS reduction in ARCI than in ARI) occurs elsewhere, interestingly over the Mediterranean strip,
397 which also contrasts with the results found under **present-day historical** conditions (~~Supp Fig 9a-~~
398 ~~e~~). These results further support the sensitivity of the simulations to both aerosol-radiation and
399 aerosol-cloud interactions under changed climates, in such a way that cloudiness still appears to be
400 the most important explanatory variable for the differences in RSDS between experiments, although
401 the role of AOD gains much relevance as compared to the analysis under **present-day historical**
402 conditions (**see the spatial and temporal correlation values in** Supp Fig 69d-i and Supp Fig 712a-
403 f, **respectively**). Under clear-sky conditions (Supp Fig 712g-i and Supp Fig 813), the results are
404 identical to those reported in the previous section.

405 Therefore, what contrasts most with the previous results is that (1) both ARI and ARCI simulations
406 provide diminished values of RSDS (of a few percentage points but statistically significant) over
407 southern locations as compared to BASE (Supp Fig 69a,b), which should primarily respond to the
408 direct aerosol effect of scattering the radiation (enhanced RSOT can be appreciated in Supp Fig
409 910a,b) since it occurs, in particular, in spite of the diminished CCT values simulated by the ARI
410 experiment there (Supp Fig 69d); and (2) such a reduction in RSDS over such southern locations is
411 reinforced when indirect effects are included (Supp Fig 69c), as these do cause higher CCT values
412 than BASE (Supp Fig 69e) and, consequently, higher RSOT values there than ARI (Supp Fig 910a-
413 c). This latter could also respond to the added role of aerosols in modifying the optical properties of
414 clouds. When ACI are considered, aerosols act as cloud condensation nuclei, which can lead to
415 whiter clouds with higher albedo. Interestingly, but out of the scope of this study, different PR shifts
416 east and west across Mediterranean Europe were detected when ARCI and ARI experiments were
417 compared between them, and then ARCI and ARI with BASE (Supp Fig 1011). Over the Balkan
418 Peninsula (south-east of the domain), ACI enhances precipitation, whether in the form of convective
419 precipitation, total precipitation, intense precipitation or number of rainy days, more than ARI does,
420 whereas over the Iberian Peninsula (south-west of the domain), ARI leads to higher precipitation
421 rates and intensity, while reducing the frequency of rainy days as compared to ARCI. These signals

422 suggest that the fact that different aerosol species prevail in these areas (the concentration of sulfate
423 is larger eastward, while the concentration of dust particles is larger westward; Supp Fig 114), and
424 how this affects the ratio between large and fine particulate matter, **might should also** have an
425 impact along with the aforementioned mechanisms **in this case** (López-Romero et al 2020).

426 Since the patterns of differences in the analyzed variables show different structures under **present**
427 **historical** and future climate conditions, the RSDS change patterns vary when ARI and ACI are
428 taken into account by the model, as described below.

429 *Future projections*

430 The change patterns for RSDS are similar in both the BASE and ARI experiments (Fig 73b,c and
431 Supp Fig 1214b,c), showing negative signals in northernmost regions (up to 10%, $\sim 15 \text{ Wm}^{-2}$) and
432 positive signals southward (up to 5%, again $\sim 15 \text{ Wm}^{-2}$). The latter are more widespread in ARI than
433 in BASE, which makes the ARI pattern the most similar to the change pattern from the GCM (Fig
434 73a and Supp Fig 1214a). However, when aerosols-cloud interactions are included in the WRF runs,
435 such a positive RSDS change signals mostly disappear, while the northern negative ones reinforce
436 in some parts as compared to the ARI pattern (Fig 73d and Supp Fig 1214d). These results are in
437 quite good agreement with the corresponding change patterns for CCT (Fig 73e-h and Supp Fig
438 1214e-h) – including the fact that the negative change signals for CCT appearing southward in the
439 GCM, BASE and ARI experiments are much less evident in ARCI – and occur in spite of two
440 constraining facts regarding the AOD simulation approach in our WRF experiments: (1) AOD
441 remains unchanged in the BASE experiment (as illustrated by Fig 73j), and (2) AOD changes from
442 the ARI and ARCI experiments are hardly realistic because their anthropogenic component is
443 disregarded (as specified in Section 2), and thus depict patterns (Fig 73k,l) that have nothing to do
444 with the GCM projection in Fig 73i (which does consider time evolving anthropogenic aerosols). In
445 fact, the spatial correlation between the patterns of AOD and RSDS changes is lower than between
446 those of CTT and RSDS changes. Therefore, direct and semi-direct aerosol effects have **a limited**
447 **little** impact on the RSDS future projections here, while indirect effects play a major role by
448 reducing the future decrease in CCT southward within our domain and thereby dispelling the future
449 increase in RSDS in this region.

450 The change signals for RSDS_{cs} and AOD_{cs} (Fig 84 and Supp Fig 1315) depict different spatial
451 structures to those for RSDS and AOD, turning mostly negative southward and positive northward

452 for RSDS_{cs} (with negative signals around 5% and positive up to 10%, in both cases implying
453 changes up to 20 Wm^{-2}). Although this occurs similarly in the three experiments (BASE, ARI and
454 ARCI), BASE provides the softest signals, which does evince a certain role of the direct aerosol
455 effect. However, there is not a clear relationship between AOD_{cs} change patterns and RSDS_{cs}
456 changes (low spatial correlation), except for some local signals in the north-east where the direct
457 aerosol effect enhances RSDS_{cs} in areas with reduced AOD_{cs} . However, as discussed above, the role
458 of retaining, or not, coincident clear-sky dates between pairs of experiments is important in filtering
459 out the true role of AOD_{cs} on RSDS_{cs} . Thus, the fact that change patterns are constructed over
460 different dates could partially explain the apparently negligible role of AOD_{cs} on RSDS_{cs} in this
461 case. But only partially, as the BASE change pattern for RSDS_{cs} (simulated on the ground of nule
462 AOD_{cs} changes) resembles the respective patterns from ARI and ARCI experiments.

463 **4 - Discussion and conclusions**

464 We presented here a research on the role of dynamically modeled atmospheric aerosols in regional
465 climate simulations with a focus on the impacts on the solar resource during the summer season
466 from a climatic perspective, including projected changes to a medium-range horizon and analysis
467 under clear-sky conditions. For this, we evaluated a set of 20-yr long runs (spanning both **present**
468 **historical** and future periods) without including aerosols and with resolved aerosol-radiation and
469 aerosol-radiation-cloud interactive (two-way) interactions performed with the WRF model (BASE,
470 ARI and ARCI experiments, respectively).

471 We interpreted the signals on the basis that the differences between ARI and BASE can be
472 attributed to direct and semi-direct aerosol effects and the differences between ARCI and ACI to the
473 indirect aerosol effect. Nonetheless, we should acknowledge that the autoconversion scheme called
474 so that cloud droplets can turn into rain droplets in the ARCI simulations is different to the
475 autoconversion scheme activated in the ARI (and BASE) simulations. This change in the WRF-
476 Chem configuration can lead to differences between ARCI and ARI experiments that do not come
477 necessarily from the aerosol-cloud interactions from a physical point of view (Liu et al 2005). In
478 fact, the activation of the aerosol-cloud interactions requires further changes in the model
479 configuration (as compared to the configuration used for the simulations labeled ARI) beyond the
480 autoconversion scheme, such as the activation of aqueous chemistry processes, which could also
481 have an added impact to the effect that can be strictly attributed to the aerosol-cloud interactions.
482 However, technically, the encoding of the WRF-Chem model hampers better isolation of the effect

483 of the aerosol-cloud interactions (the mentioned aspects necessarily change between ARI and ARCI
484 run modes). Therefore, ARCI-ARI differences can not be attributed to the aerosol-cloud interactions
485 from a purely physical point of view, but to the activation of the aerosol-cloud interactions from a
486 modeling point of view. It should also be borne in mind that the set of experiments performed
487 allows any attribution to the interactive aerosol modeling approach adopted here to be made, while
488 it is a distinct feature with respect to previous studies aimed at providing more consistent signals
489 from a physical point of view. **Besides Last**, and more general, internal variability plays a role in
490 the simulations (e.g. Gómez-Navarro et al 2012), and a single member **with a single physics**
491 **configuration**, as was used for the sensitivity experiment, may not be sufficient to obtain generally
492 occurring responses. **Last, we kept the anthropogenic aerosol emissions unchanged throughout**
493 **the simulation period. This approach permits to better isolate the signals from the aerosol-**
494 **radiation-cloud interactions due to the climate variability alone and the so-called climate**
495 **change penalty alone, but at the expense of the reliability of the simulated patterns.**
496 **Anthropogenic emissions have been dramatically reduced since the 1980s and are expected to**
497 **continue in that pathway to the future (IPCC 2013, 2014), so keeping 2010 values (as we did)**
498 **could lead to an underestimation of AOD in the historical period (in fact, it does; reference**
499 **AOD climatologies can be found in Pavlidis et al. 2020) and to its overestimation in the future**
500 **period.** Under these constraints, we draw the following conclusions.

501 The inclusion of aerosols in the WRF simulations reduces in general the amount of solar radiation
502 reaching the surface by a few percentage points (~5%) under both **present historical** and future
503 climate scenarios, as expected (Nabat et al 2015a, Gutiérrez et al 2018, Pavlidis et al 2020). Under
504 **present-day historical** conditions, this effect is larger when the aerosol-cloud interaction remains
505 turned off, because its activation leads to less cloudiness (over Mediterranean Europe) and lower
506 AOD values (over Atlantic Europe), as evidenced when ARCI and ARI simulations were compared.
507 The differences in RSDS between experiments are in overall good agreement with those found in
508 cloudiness, while they seem to be unlinked with the differences in AOD in many parts of the
509 domain. In agreement with Pavlidis et al (2020), AOD plays its major role under clear-sky
510 conditions. However, the differences in JJA-mean values of RSDS under clear skies between
511 experiments with and without dynamic aerosols are hardly about 1%, while still significant in some
512 of the southernmost parts of our European domain, and almost null between ARCI and ARI.

513 Our results suggested a variety of drivers underlying the mechanisms to explain the signals
514 obtained, depending on the region (and season; winter plots are provided in Supp Fig **14-17****16-19** as

515 an example for interested readers), and varying under future climate conditions. These involve the
516 ~~effect of large aerosols in hampering the formation of clouds, increased~~ scattering of solar
517 radiation with the consequent cooling downward, suppression of cloudiness due to thermodynamic
518 effects, modification of the clouds' optical properties, or in-cloud scavenging processes. As these
519 prevailing mechanisms change (up to a point) in the future, the sensitivity of the WRF simulations
520 under future climate conditions, represented through the patterns of differences in RSDS, is
521 somehow depicted differently than under ~~present-day~~ **historical** conditions. Therefore, the future
522 projections also show sensitivity to the way the model considers aerosols.

523 The patterns of change for RSDS and CCT again show high spatial correlations in all the GCM and
524 RCM (BASE, ARI and ARCI) projections. Although lower, still high spatial correlations define the
525 match between the RSDS change patterns and those for AOD in the GCM, while this is not the case
526 in either the ARI or ARCI experiments. The GCM, BASE and ARI experiments agree in projecting
527 positive RSDS change signals in southern and eastern areas (around 5%), while clear differences
528 are found between the GCM and the BASE or ARI RSDS change patterns (with the latter two very
529 similar) in central and northeastern areas, where the positive signals from the GCM turn notably
530 negative in both BASE and ARI. ARCI provides the most singular and negative picture of RSDS
531 changes among all those shown, with widespread decreasing signals of a few percentage points,
532 further reinforcing the fact that the indirect effect tends to counteract the direct and semi-direct
533 effect of aerosols and enlarges the distance between the RCM and the GCM projections.

534 Previous works (Jerez et al 2015, Sørland et al 2018) had already detected inconsistencies in the
535 change signals between RCM projections and those from their driving GCM, which have been
536 related to the way aerosols had been represented in the RCM through their impact on the simulated
537 AOD (Gutiérrez et al 2020, Boé et al 2020), and in particular to their direct and semi-direct effects
538 and their reduced concentrations in the future as long as anthropogenic emissions are projected to
539 decrease. In agreement with these previous findings, insofar as we kept the anthropogenic aerosol
540 emissions unchanged throughout the simulation period, our projections differ from those obtained
541 with the GCM. Nevertheless, the ARI experiment brings our results slightly nearer to those of the
542 GCM as compared to the BASE experiment, perhaps also indicating the key role of the direct and
543 semi-direct aerosol effects for reducing the GCM-RCM discrepancies, as reported in these previous
544 works. However, pushing our understanding further, by turning off the already reported effect of
545 reduced aerosol concentrations in the future via the direct and semi-direct effects, our approach
546 made it possible to identify the prevailing role of CCT changes (over the dynamically simulated

547 natural changes in AOD) to explain our signals of change in RSDS, and the capacity of the aerosol-
548 radiation-cloud interactions to significantly alter our RSDS change patterns (much more than
549 aerosol-radiation interactions alone do). Thus, although change patterns for RSDS certainly look
550 uniform among experiments under clear-sky conditions (likely because we suppressed the
551 anthropogenic component for the changes in AOD, which was identified by Boé et al (2020) as a
552 main factor for these signals indeed), the results presented here further indicate that the joint effect
553 of aerosol-radiation-cloud interactions should be considered in the RCM simulations that serve to
554 build up action-oriented messages in the challenging context of current climate change, calling for
555 caution otherwise and for future research efforts in this line.

556 **Author contribution**

557 S. J. conceived this study. L. P.-P., P. J.-G. and J. P. M. designed the experiments and J. M. L.-R. and
558 E. Pravia-Sarabia carried them out. S. J. performed the analysis and prepared the manuscript with
559 contributions from all co-authors.

560 **Acknowledgments**

561 This study was supported by the Spanish Ministry of Science, Innovation and Universities/*Agencia*
562 *Estatal de Investigación* and the European Regional Development Fund through the projects EASE
563 (RTI2018-100870-A-I00, MCI/AEI/FEDER, UE) and ACEX (CGL2017-87921-R,
564 MINECO/AEI/FEDER, UE), and by the *Fundación Séneca – Agencia de Ciencia y Tecnología de la*
565 *Región de Murcia* through the project CLIMAX (20642/JLI/18). S. Jerez receive funding from the
566 *Plan Propio de Investigación* of the University of Murcia (grant No. UMU-2017-10604). L. Palacios-Peña
567 thanks the FPU14/05505 scholarship and the Spanish Ministry of Education, Culture and Sports. J. M.
568 López-Romero acknowledges the FPI-BES-2015-074062 grant from the Spanish Ministry of Science.

569 **Code and data availability**

570 All data (netcdf files) and relevant codes (scripts for data analysis and WRF namelists) for reproducing this
571 study are publicly accessible at
572 <http://doi.org/10.23728/b2share.a65d25c2b3ba49e1a46e970783e9476e>.

573 **References**

- 574 **Allen, R. J., & Sherwood, S. C. (2010). Aerosol cloud semi direct effect and land sea**
575 **temperature contrast in a GCM. *Geophysical Research Letters*, 37(7).**
- 576 Archer-Nicholls, S., Lowe, D., Utembe, S., Allan, J., Zaveri, R. A., Fast, J. D., Hodnebrog, Ø.,
577 Denier van der Gon, H., McFiggans, G. (2014). Gaseous chemistry and aerosol mechanism
578 developments for version 3.5.1 of the online regional model, WRF-Chem. *Geoscientific Model*
579 *Development*, 7, 2557–2579.
- 580 Bartók, B., Wild, M., Folini, D., Lüthi, D., Kotlarski, S., Schär, C., Vautard, R., Jerez, S., & Imecs,
581 Z. (2017). Projected changes in surface solar radiation in CMIP5 global climate models and in
582 EURO-CORDEX regional climate models for Europe. *Climate Dynamics*, 49(7-8), 2665-2683.
- 583 Bloomfield, H. C., Brayshaw, D. J., Shaffrey, L. C., Coker, P. J., & Thornton, H. E. (2016).
584 Quantifying the increasing sensitivity of power systems to climate variability. *Environmental*
585 *Research Letters*, 11(12).
- 586 Boé, J., Somot, S., Corre, L., & Nabat, P. (2020). Large discrepancies in summer climate change
587 over Europe as projected by global and regional climate models: causes and consequences. *Climate*
588 *Dynamics*, 54(5), 2981-3002.
- 589 Boucher, O. (2015). Atmospheric aerosols. In *Atmospheric Aerosols* (pp. 9-24). Springer,
590 Dordrecht.
- 591 Chapman, E.G., Gustafson, W.I., Easter, R.C., Barnard, J.C., Ghan, S.J., Pekour, M.S., & Fast, J.D.
592 (2009). Coupling aerosol-cloud-radiative processes in the WRF-Chem model: Investigating the
593 radiative impact of elevated point sources. *Atmospheric Chemistry and Physics*, 9, 945–964.
- 594 Chen, F., & Dudhia, J. (2001). Coupling an advanced land surface–hydrology model with the Penn
595 State–NCAR MM5 modeling system. Part I: Model implementation and sensitivity. *Monthly*
596 *Weather Review*, 129(4), 569-585.

- 597 Chenni, R., Makhlof, M., Kerbache, T., & Bouzid, A. (2007). A detailed modeling method for
598 photovoltaic cells. *Energy*, 32(9), 1724-1730.
- 599 Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J.
600 A., Higurashi, A., & Nakajima, T. (2002). Tropospheric aerosol optical thickness from the
601 GOCART model and comparisons with satellite and Sun photometer measurements. *Journal of the*
602 *Atmospheric Sciences*, 59, 461 – 483.
- 603 Chin, M., Rood, R.B., Lin. S.-J., Müller, J.-F., Thompson, M. (2000). Atmospheric sulfur cycle
604 simulated in the global model GOCART: Model description and global properties. *Journal of*
605 *Geophysical Research*, 105(D20), 24671-24687.
- 606 Collins, S., Deane, P., Gallachóir, B. Ó., Pfenninger, S., & Staffell, I. (2018). Impacts of inter-
607 annual wind and solar variations on the European power system. *Joule*, 2(10), 2076-2090.
- 608 Croft, B., Pierce, J. R., Martin, R. V., Hoose, C., & Lohmann, U. (2012). Uncertainty associated
609 with convective wet removal of entrained aerosols in a global climate model. *Atmospheric*
610 *Chemistry and Physics*, 12(22), 10725-10748.
- 611 Crook, J. A., Jones, L. A., Forster, P. M., & Crook, R. (2011). Climate change impacts on future
612 photovoltaic and concentrated solar power energy output. *Energy & Environmental Science*, 4(9),
613 3101-3109.
- 614 Dubey, S., Sarvaiya, J. N., & Seshadri, B. (2013). Temperature dependent photovoltaic (PV)
615 efficiency and its effect on PV production in the world—a review. *Energy Procedia*, 33, 311-321.
- 616 Fast, J. D., Gustafson Jr, W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G.
617 A., & Peckham, S. E. (2006). Evolution of ozone, particulates, and aerosol direct radiative forcing
618 in the vicinity of Houston using a fully coupled meteorology chemistry aerosol model. *Journal of*
619 *Geophysical Research: Atmospheres*, 111(D21).
- 620 Gaetani, M., Huld, T., Vignati, E., Monforti-Ferrario, F., Dosio, A., & Raes, F. (2014). The near
621 future availability of photovoltaic energy in Europe and Africa in climate-aerosol modeling
622 experiments. *Renewable and Sustainable Energy Reviews*, 38, 706-716.

623 Geiger, H., Barnes, I., Bejan, I., Benter, T., & Spittler, M. (2003). The tropospheric degradation of
624 isoprene: an updated module for the regional atmospheric chemistry mechanism. *Atmospheric*
625 *Environment*, 37, 1503 - 1519.

626 Ghan, S. J., Leung, L. R., Easter, R. C., & Abdul Razzak, H. (1997). Prediction of cloud droplet
627 number in a general circulation model. *Journal of Geophysical Research: Atmospheres*, 102(D18),
628 21777-21794.

629 Gil, V., Gaertner, M. A., Gutierrez, C., & Losada, T. (2019). Impact of climate change on solar
630 irradiation and variability over the Iberian Peninsula using regional climate models. *International*
631 *Journal of Climatology*, 39(3), 1733-1747.

632 Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., & Lin, S. J. (2001).
633 Sources and distributions of dust aerosols simulated with the GOCART model. *Journal of*
634 *Geophysical Research: Atmospheres*, 106(D17), 20255-20273.

635 Giorgetta, M., Jungclaus, J., Reick, C., Legutke, S., Brovkin, V., Crueger, T., Esch, M., Fieg, K.,
636 Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Kinne, S., Kornblueh, L., Matei, D., Mauritsen,
637 T., Mikolajewicz, U., Müller, W., Notz, D., Raddatz, T., Rast, S., Roeckner, E., Salzmann, M.,
638 Schmidt, H., Schnur, R., Segschneider, J., Six, K., Stockhause, M., Wegner, J., Widmann, H.,
639 Wieners, K.-H., Claussen, M., Marotzke, J., & Stevens, B. (2012a). Forcing Data for Regional
640 Climate Models Based on the MPI-ESM-LR Model of the Max Planck Institute for Meteorology
641 (MPI-M): The CMIP5 Historical Experiment, World Data Center for Climate (WDCC) at DKRZ,
642 https://doi.org/10.1594/WDCC/RCM_CMIP5_historical-LR.

643 Giorgetta, M., Jungclaus, J., Reick, C., Legutke, S., Brovkin, V., Crueger, T., Esch, M., Fieg, K.,
644 Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Kinne, S., Kornblueh, L., Matei, D., Mauritsen,
645 T., Mikolajewicz, U., Müller, W., Notz, D., Raddatz, T., Rast, S., Roeckner, E., Salzmann, M.,
646 Schmidt, H., Schnur, R., Segschneider, J., Six, K., Stockhause, M., Wegner, J., Widmann, H.,
647 Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B. (2012b). Forcing data for Regional
648 Climate Models Based on the MPI-ESM-LR model of the Max Planck Institute for Meteorology
649 (MPI-M): The CMIP5rcp85 experiment, World Data Center for Climate (WDCC) at DKRZ, [https://](https://doi.org/10.1594/WDCC/RCM_CMIP5_rcp85-LR)
650 doi.org/10.1594/WDCC/RCM_CMIP5_rcp85-LR.

651 Giorgi, F., Bi, X., & Qian, Y. (2002). Direct radiative forcing and regional climatic effects of
652 anthropogenic aerosols over East Asia: A regional coupled climate chemistry/aerosol model study.
653 *Journal of Geophysical Research: Atmospheres*, 107(D20), AAC-7.

654 Gómez-Navarro, J. J., Montávez, J. P., Jiménez-Guerrero, P., Jerez, S., Lorente-Plazas, R.,
655 González-Rouco, J. F., & Zorita, E. (2012). Internal and external variability in regional simulations
656 of the Iberian Peninsula climate over the last millennium.

657 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., & Geron, C. (2006). Estimates of
658 global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols
659 from Nature). *Atmospheric Chemistry and Physics*, 6(11), 3181-3210.

660 Gutiérrez, C., Somot, S., Nabat, P., Mallet, M., Gaertner, M. Á., & Perpiñán, O. (2018). Impact of
661 aerosols on the spatiotemporal variability of photovoltaic energy production in the Euro-
662 Mediterranean area. *Solar Energy*, 174, 1142-1152.

663 Gutiérrez, C., Somot, S., Nabat, P., Mallet, M., Corre, L., van Meijgaard, E., Perpiñán, O., &
664 Gaertner, M. Á. (2020). Future evolution of surface solar radiation and photovoltaic potential in
665 Europe: investigating the role of aerosols. *Environmental Research Letters*, 15(3), 034035.

666 Grell, G. A. (1993). Prognostic evaluation of assumptions used by cumulus parameterizations.
667 *Monthly Weather Review*, 121(3), 764-787.

668 Grell, G. A., & Dévényi, D. (2002). A generalized approach to parameterizing convection
669 combining ensemble and data assimilation techniques. *Geophysical Research Letters*, 29(14), 38-1.

670 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., & Eder, B.
671 (2005). Fully coupled “online” chemistry within the WRF model. *Atmospheric Environment*,
672 39(37), 6957-6975.

673 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz Sabater, J., ... & Simmons,
674 A. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*,
675 146(730), 1999-2049.

676 Hong, S. Y., Noh, Y., & Dudhia, J. (2006). A new vertical diffusion package with an explicit
677 treatment of entrainment processes. *Monthly weather review*, 134(9), 2318-2341.

678 Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., & Collins, W. D.
679 (2008). Radiative forcing by long lived greenhouse gases: Calculations with the AER radiative
680 transfer models. *Journal of Geophysical Research: Atmospheres*, 113(D13).

681 IPCC (2013). *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I*
682 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. T. F. Stocker, D.
683 Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P. M.
684 Midgley (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY,
685 USA, 1535 pp.

686 IPCC (2014). *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group*
687 *III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. O.
688 Edenhofer, R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S.
689 Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T. Zwickel and
690 J.C. Minx (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY,
691 USA.

692 IRENA (2019). *Global energy transformation: A roadmap to 2050 (2019 edition)*. International
693 Renewable Energy Agency, Abu Dhabi.

694 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., et al. (2014).
695 EURO-CORDEX: new high-resolution climate change projections for European impact research.
696 *Regional environmental change*, 14(2), 563-578.

697 Jacob, D., Teichmann, C., Sobolowski, S., Katragkou, E., Anders, I., Belda, M., et al. (2020).
698 Regional climate downscaling over Europe: perspectives from the EURO-CORDEX community.
699 *Regional Environmental Change*, 20(2).

700 Jerez, S., Trigo, R. M., Vicente-Serrano, S. M., Pozo-Vázquez, D., Lorente-Plazas, R., Lorenzo-
701 Lacruz, J., Santos-Alamillos, F., & Montávez, J. P. (2013). The impact of the North Atlantic

702 Oscillation on renewable energy resources in southwestern Europe. *Journal of Applied Meteorology*
703 *and Climatology*, 52(10), 2204-2225.

704 Jerez, S., Tobin, I., Vautard, R., Montávez, J. P., López-Romero, J. M., Thais, F., Bartok, B.,
705 Christensen, O. B., Colette, A., Déqué, M., Nikulin, G., Kotlarski, S., van Meijgaard, E., teichmann,
706 C., & Wild, M. (2015). The impact of climate change on photovoltaic power generation in Europe.
707 *Nature Communications*, 6, 10014.

708 Jerez, S., López-Romero, J. M., Turco, M., Jiménez-Guerrero, P., Vautard, R., & Montávez, J. P.
709 (2018). Impact of evolving greenhouse gas forcing on the warming signal in regional climate model
710 experiments. *Nature Communications*, 9(1), 1304.

711 Jerez, S., Tobin, I., Turco, M., Jiménez-Guerrero, P., Vautard, R., & Montávez, J. P. (2019). Future
712 changes, or lack thereof, in the temporal variability of the combined wind-plus-solar power
713 production in Europe. *Renewable Energy*, 139, 251-260.

714 Jiménez, P., Baldasano, J.M., Dabdub, D. (2003). Comparison of photochemical mechanisms for air
715 quality modeling. *Atmospheric Environment*, 37, 4179-4194.

716 Kinne, S. (2019). Aerosol radiative effects with MACv2. *Atmospheric Chemistry and Physics*,
717 19(16), 10919-10959.

718 Kloster, S., Dentener, F., Feichter, J., Raes, F., Lohmann, U., Roeckner, E., & Fischer-Bruns, I.
719 (2010). A GCM study of future climate response to aerosol pollution reductions. *Climate Dynamics*,
720 34(7-8), 1177-1194.

721 Kozarcanin, S., Andresen, G. B., & Greiner, M. (2018). Impact of Climate Change on the Backup
722 Infrastructure of Highly Renewable Electricity Systems. *Journal of Sustainable Development of*
723 *Energy, Water and Environment Systems*, 6(4), 710-724.

724 Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., et al. (2010). Historical
725 (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols:
726 methodology and application. *Atmospheric Chemistry and Physics*, 10(15), 7017-7039.

727 Lee, S. S., Donner, L. J., Phillips, V. T., & Ming, Y. (2008). Examination of aerosol effects on
728 precipitation in deep convective clouds during the 1997 ARM summer experiment. *Quarterly*
729 *Journal of the Royal Meteorological Society*, 134(634), 1201-1220.

730 Li, G., Wang, Y., & Zhang, R. (2008). Implementation of a two moment bulk microphysics scheme
731 to the WRF model to investigate aerosol cloud interaction. *Journal of Geophysical Research:*
732 *Atmospheres*, 113(D15).

733 Li, X., Wagner, F., Peng, W., Yang, J., & Mauzerall, D. L. (2017). Reduction of solar photovoltaic
734 resources due to air pollution in China. *Proceedings of the National Academy of Sciences*, 114(45),
735 11867-11872.

736 Lin, Y. L., Farley, R. D., & Orville, H. D. (1983). Bulk parameterization of the snow field in a cloud
737 model. *Journal of Climate and Applied Meteorology*, 22(6), 1065-1092.

738 Liu, Y., Daum, P. H., & McGraw, R. L. (2005). Size truncation effect, threshold behavior, and a new
739 type of autoconversion parameterization. *Geophysical research letters*, 32(11).

740 López-Romero, J. M., Montávez, J. P., Jerez, S., Lorente-Plazas, R., Palacios-Peña, L., & Jiménez-
741 Guerrero, P. (2020). Precipitation response to Aerosol-Radiation and Aerosol-Cloud Interactions in
742 Regional Climate Simulations over Europe. *Atmospheric Chemistry and Physics Discussions*,
743 <https://doi.org/10.5194/acp-2020-381>, in review.

744 Malm, W. C., Sisler, J. F., Huffman, D., Eldred, R. A., & Cahill, T. A. (1994). Spatial and seasonal
745 trends in particle concentration and optical extinction in the United States. *Journal of Geophysical*
746 *Research: Atmospheres*, 99(D1), 1347-1370.

747 Mitchell, D. L., Rasch, P., Ivanova, D., McFarquhar, G., & Nousiainen, T. (2008). Impact of small
748 ice crystal assumptions on ice sedimentation rates in cirrus clouds and GCM simulations.
749 *Geophysical research letters*, 35(9).

750 Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Van Vuuren, D. P., et al.
751 (2010). The next generation of scenarios for climate change research and assessment. *Nature*,
752 463(7282), 747.

753 Müller, J., Folini, D., Wild, M., & Pfenninger, S. (2019). CMIP-5 models project photovoltaics are a
754 no-regrets investment in Europe irrespective of climate change. *Energy*, *171*, 135-148.

755 Nabat, P., Somot, S., Mallet, M., Sanchez Lorenzo, A., & Wild, M. (2014). Contribution of
756 anthropogenic sulfate aerosols to the changing Euro Mediterranean climate since 1980.
757 *Geophysical Research Letters*, *41*(15), 5605-5611.

758 Nabat, P., Somot, S., Mallet, M., Sevault, F., Chiacchio, M., & Wild, M. (2015a). Direct and semi-
759 direct aerosol radiative effect on the Mediterranean climate variability using a coupled regional
760 climate system model. *Climate Dynamics*, *44*(3-4), 1127-1155.

761 Nabat, P., Somot, S., Mallet, M., Michou, M., Sevault, F., Driouech, F., Meloni, D., di Sarra, G., di
762 Biagio, C., Formenti, P., Sicard, M., Leon, J. F. & Bouin, M. N. (2015b). Dust aerosol radiative
763 effects during summer 2012 simulated with a coupled regional aerosol–atmosphere–ocean model
764 over the Mediterranean. *Atmospheric Chemistry and Physics*, *15*(6), 3303-3326.

765 Palacios-Peña, L., Lorente-Plazas, R., Montávez, J. P. and Jiménez-Guerrero, P. (2019a) Saharan
766 Dust Modeling Over the Mediterranean Basin and Central Europe: Does the Resolution Matter?.
767 *Frontiers in Earth Science*, *7*, 290.

768 Palacios-Peña, L., Jiménez-Guerrero, P., Baró, R., Balzarini, A., Bianconi, R., Curci, G., Landi, T.
769 C., Pirovano, G., Prank, M., Riccio, A., Tuccella, P., & Galmarini, S. (2019b). Aerosol optical
770 properties over Europe: an evaluation of the AQMEII Phase 3 simulations against satellite
771 observations. *Atmospheric Chemistry and Physics*, *19*(5), 2965-2990.

772 Palacios-Peña, L., Montávez, J.P., López-Romero, J.M., Jerez, S., Gómez-Navarro, J.J., Lorente-
773 Plazas, R., Ruiz, J., & Jiménez-Guerrero, P. (2020). Added Value of Aerosol-Cloud Interactions for
774 Representing Aerosol Optical Depth in an Online Coupled Climate-Chemistry Model over Europe.
775 *Atmosphere*, *11*, 360.

776 Pavlidis, V., Katragkou, E., Prein, A., Georgoulas, A. K., Kartsios, S., Zanis, P., & Karacostas, T.
777 (2020). Investigating the sensitivity to resolving aerosol interactions in downscaling regional model
778 experiments with WRFv3.8.1 over Europe. *Geoscience Model Development*, *13*, 2511–2532

779 Rohrig, K., Berkhout, V., Callies, D., Durstewitz, M., Faulstich, S., Hahn, B., et al. (2019).
780 Powering the 21st century by wind energy – Options, facts, figures. *Applied Physics Reviews*, 6(3),
781 031303.

782 Rummukainen, M. (2010). State of the art with regional climate models. *Wiley Interdisciplinary*
783 *Reviews: Climate Change*, 1(1), 82-96.

784 Ruti, P. M., Somot, S., Giorgi, F., Dubois, C., Flaounas, E., Obermann, A., et al. (2016). MED-
785 CORDEX initiative for Mediterranean climate studies. *Bulletin of the American Meteorological*
786 *Society*, 97(7), 1187-1208.

787 Rutledge, S. A., & Hobbs, P. V. (1984). The Mesoscale and Microscale Structure and Organization
788 of Clouds and Precipitation in Midlatitude Cyclones. XII: A Diagnostic Modeling Study of
789 Precipitation Development in Narrow Cold-Frontal Rainbands. *Journal of Atmospheric Science*, 41,
790 2949–2972.

791 Schewe, J., Gosling, S. N., Reyer, C., Zhao, F., Ciais, P., Elliott, J., Francois, L., Huber, V., Lotze,
792 H. K., Seneviratne, S. I., van Vliet, M. T. H., Vautard, R., Wada, Y., Breuer, L., Büchner, M.,
793 Carozza, D. A., Chang, J., Coll, M., Deryng, D., de Wit, A., Eddy, T. D., Folberth, C., Frieler, K.,
794 Friend, A. D., Gerten, D., Gudmundsson, L., Hanasaki, N., Ito, A., Khabarov, N., Kim, H.,
795 Lawrence, P., Morfopoulos, C., Müller, C., Schmied, H. M., Orth, R., Ostberg, S., Pokhrel, Y.,
796 Pugh, T. A. M., Sakurai, G., Satoh, Y., Schmid, E., Stacke, T., Steenbeek, J., Steinkamp, J., Tang,
797 Q., Tian, H., Tittensor, D.P., Volkholz, J., Wang, X., & Warszawski, L. (2019). State-of-the-art
798 global models underestimate impacts from climate extremes. *Nature Communications*, 10(1), 1005.

799 Sørland, S. L., Schär, C., Lüthi, D., & Kjellström, E. (2018). Bias patterns and climate change
800 signals in GCM-RCM model chains. *Environmental Research Letters*, 13(7), 074017.

801 Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., et al. (2016).
802 Improving our fundamental understanding of the role of aerosol– cloud interactions in the climate
803 system. *Proceedings of the National Academy of Sciences*, 113(21), 5781-5790.

804 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Wang, W., & Powers, J. G.
805 (2008). A description of the Advanced Research WRF version 3. Technical report, NCAR Tech.
806 Note TN-475+STR, doi:10.5065/D68S4MVH.

807 Soares, P. M., Brito, M. C., & Careto, J. A. (2019). Persistence of the high solar potential in Africa
808 in a changing climate. *Environmental Research Letters*.

809 Stockwell, W. R., Kirchner, F., Kuhn, M., & Seefeld, S. (1997). A new mechanism for regional
810 atmospheric chemistry modeling. *Journal of Geophysical Research: Atmospheres*, 102, 25 847 - 25
811 879.

812 Tao, W. K., Simpson, J., & McCumber, M. (1989). An ice-water saturation adjustment. *Monthly*
813 *Weather Review*, 117(1), 231-235.

814 Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 and the experiment
815 design. *Bulletin of the American Meteorological Society*, 93(4), 485-498.

816 Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M. A., Mitchell, K., Ek, M., Gayno, G.,
817 Wegiel, J. W., & Cuenca, R. H. (2004, January). Implementation and verification of the unified
818 NOAA land surface model in the WRF model. In 20th conference on weather analysis and
819 forecasting/16th conference on numerical weather prediction (Vol. 1115). Seattle, WA: American
820 Meteorological Society.

821 Tobin, I., Greuell, W., Jerez, S., Ludwig, F., Vautard, R., Van Vliet, M. T. H., & Breón, F. M. (2018).
822 Vulnerabilities and resilience of European power generation to 1.5 C, 2 C and 3 C warming.
823 *Environmental Research Letters*, 13(4), 044024.

824 van der Wiel, K., Bloomfield, H. C., Lee, R. W., Stoop, L. P., Blackport, R., Screen, J. A., & Selten,
825 F. M. (2019). The influence of weather regimes on European renewable energy production and
826 demand. *Environmental Research Letters*, 14(9), 094010.

827 Wang, H., Xie, S. P., Tokinaga, H., Liu, Q., & Kosaka, Y. (2016). Detecting cross equatorial wind
828 change as a fingerprint of climate response to anthropogenic aerosol forcing. *Geophysical Research*
829 *Letters*, 43(7), 3444-3450.

- 830 Wilcox, L. J., Highwood, E. J., & Dunstone, N. J. (2013). The influence of anthropogenic aerosol
831 on multi-decadal variations of historical global climate. *Environmental Research Letters*, 8(2),
832 024033.
- 833 Wild, M., Folini, D., Henschel, F., Fischer, N., & Müller, B. (2015). Projections of long-term
834 changes in solar radiation based on CMIP5 climate models and their influence on energy yields of
835 photovoltaic systems. *Solar Energy*, 116, 12-24.
- 836 Wild, O., Zhu, X., Prather, M. J., & Fast, J. (2002). Accurate Simulation of In- and Below-Cloud
837 Photolysis in Tropospheric Chemical Models. *Journal of Atmospheric Chemistry*, 37, 245 – 282.
- 838 ~~Xue, H., Feingold, G., & Stevens, B. (2008). Aerosol effects on clouds, precipitation, and the~~
839 ~~organization of shallow cumulus convection. *Journal of the Atmospheric Sciences*, 65(2), 392-~~
840 ~~406.~~
- 841 Yin, Y., Levin, Z., Reisin, T., & Tzivion, S. (2000). Seeding convective clouds with hygroscopic
842 flares: Numerical simulations using a cloud model with detailed microphysics. *Journal of Applied*
843 *Meteorology*, 39(9), 1460-1472.

844 **Figure caption**

845 **Figure 1.** Relative differences between the WRF simulations in the RSDS (a to c), CCT (d to f) and
846 AOD at 550 nm (g to i) summer (JJA) climatologies in the **present historical** period (1991-2010),
847 squared if statistically significant ($p < 0.05$); units: %. Note that panels g and h are referred to the
848 horizontal colorbar just below them and simply represent the AOD summer climatologies in ARI
849 and ARCI respectively. Spatial correlations (s_corr) between the patterns in the second and third
850 rows and the respective patterns in the first row are indicated in the headers.

851 **Figure 2.** Contribution of each aerosol species (BC: black carbon, DUST, OC: organic carbon,
852 SEAS: sea salt, and SULF: sulfate) to the JJA-mean total surface aerosol mass concentration
853 in ARI and ARCI simulations in the period 1991-2010. Units: %.

854 **Figure 3.** Relative differences between the WRF simulations in the top-of-the-atmosphere
855 outgoing short-wave radiation (RSOT, a to c), surface (2 m height) air temperature (TAS, d to
856 f), surface (1000 hPa pressure level) relative humidity (RH, g to I), and number of cloudy days
857 (CLD; defined as days with mean CCT > 75%, j to l) summer (JJA) climatologies in the
858 historical period (1991-2010), squared if statistically significant ($p < 0.05$). Units: K for TAS, %
859 for RSOT, RH and CLD.

860 **Figure 4.** Vertical profiles of the spatial mean differences in summer (JJA) mean air
861 temperature (T, left panels) and cloud fraction (CLFR, right panels) in the historical period
862 (1991-2010) between experiments over two small areas: a northern one (Region N; top panels)
863 and a southern one (Region S; bottom panels), gray shaded in the respective maps. These are
864 plain differences, which units are K for T and % for CLFR.

865 **Figure 5.** Relative differences between the WRF simulations in the summer (JJA)
866 climatologies of various precipitation (PR) statistics in the historical period (1991-2010),
867 squared if statistically significant ($p < 0.05$): mean PR (a to c), 90th percentile of the JJA daily
868 PR series (d to f), number of rainy days (RD) in the JJA daily PR series (defined as days with
869 mean precipitation > 1 mm, g to I), and mean convective precipitation (PRC, j to l). Units: %.

870 **Figure 62.** Relative differences between the WRF simulations in the RSDS_{cs} (a to c) and AOD_{cs} at
871 550 nm (d to f) summer (JJA) climatologies, this is under clear-sky conditions, in the **present**

872 **historical** period (1991-2010), squared if statistically significant ($p < 0.05$); units: %. Note that
873 panels d and e are referred to the horizontal colorbar just below them and simply represent the AOD
874 summer climatologies in ARI and ARCI, respectively. Gray shaded areas depict grid points where
875 less than 75% of the summer mean values in the time series of RSDS_{cs} and AOD_{cs} were not missing.
876 Spatial correlations (s_corr) between the patterns in the second row and the respective patterns in
877 the first row are indicated in the headers.

878 **Figure 73.** Projected changes for the RSDS (a to d), CCT (e to h) and AOD at 550nm (i to l)
879 summer (JJA) climatologies by the GCM (first column) and the WRF experiments (second to fourth
880 columns); units: %. Squares highlight statistically significant signals ($p < 0.05$). Note that panel i is
881 referred to the horizontal colorbar just below it. Spatial correlations (s_corr) between the patterns in
882 the second and third rows and the respective patterns in the first row are indicated in the headers.

883 **Figure 84.** Projected changes for the RSDS_{cs} (a to c) and AOD_{cs} at 550nm (d to f) summer (JJA)
884 climatologies, this is under clear-sky conditions, by the WRF experiments, squared if statistically
885 significant ($p < 0.05$); units: %. Gray shaded areas depict grid points where less than 75% of the
886 summer mean values in the time series of RSDS_{cs} and AOD_{cs} were not missing in either the **present**
887 **historical** or the future period. Spatial correlations (s_corr) between the patterns in the second row
888 and the respective patterns in the first row are indicated in the headers.